

K-Moduli Spaces of Del Pezzo Pairs with Sum of Lines and Wall Crossings

Junyan Zhao

Abstract

In this paper, we study the K-moduli spaces of pairs $(X, c\sum L_i)$, where X is a del Pezzo surface of degree d , L_i 's are the lines on it counted with multiplicity, and $c > 0$ any coefficient such that this pair is log Fano. We prove that for $d = 2, 3, 4$, there are no walls for the K-moduli spaces. For $d = 1$, we exhibit examples of walls.

Contents

1	Introduction	1
2	Preliminaries	2
3	Cubic and quartic del Pezzo pairs	4
3.1	The cubic case	5
3.2	The quartic case	6
4	Degree two case	7
4.1	Singularities of plane curves	8
4.2	Singularities of curves in $\mathbb{P}(1, 1, 4)$	10
5	Discussion on degree one case	10
5.1	Degree one del Pezzo with an A_7 -singularity	10
5.2	Degree one del Pezzo with two D_4 -singularities	11
5.3	Degree one del Pezzo with an A_8 -singularity two $\frac{1}{9}(1, 2)$ -singularities	12

1 Introduction

The moduli spaces of del Pezzo surfaces have been extensively studied thanks to the development of different approaches to construct moduli spaces.

A weighted marked del Pezzo surface is a pair (S, B) with S a del Pezzo surface and $B = \sum b_i B_i$ a boundary divisor, where $b_i \in [0, 1]$ and B_i 's are all the finitely many lines on S . The KSBA moduli spaces parameterizes all such pairs such that (S, B) is semi log canonical (slc) and $K_S + B$ is ample. For example, when $b_i = 1$ for each i , the moduli of such quartic del Pezzo pairs is isomorphic to the moduli $\overline{M}_{0,5}$ of rational curves with 5 marked points (cf. [HKT09, HKT13]). Another interesting case is when $b_i = b$, where b is a positive number such that (S, B) is close to a Calabi-Yau pair (cf. [HKT09, HKT13, GKS21]).

The development of K-stability provides a moduli theory for log Fano pairs, called K-moduli spaces. The general K-moduli theory was established by a group of people in the past decade

([ABHLX20, BLX22, BX19, Jia20, LWX21, LXZ22, XZ20]). Roughly speaking, for fixed numerical invariants, we have a projective scheme parameterizing the K -polystable log Fano pairs with these invariants. Moreover, in the surface case, there is a wall crossing structure when we vary the coefficients of the boundary divisors (cf. [ADL19, ADL21]). See Section 2 for definitions and more details. We can ask the similar question in the K -moduli spaces setting: can we describe the moduli of K -polystable log Fano pairs $(X, c \sum L_i)$ and determine the walls when we vary c , where X is a del Pezzo surface, L_i 's are all the lines on X and $c > 0$ such that $-(K_X + c \sum L_i)$ is ample. In particular, when $c = 0$, then the K -moduli space parameterizes all the del Pezzo surfaces of a fixed degree, which admit a Kähler-Einstein metric (cf. [OSS16, MM93]).

We prove in this article that for $d = 2, 3, 4$, there are no walls for the K -moduli. For $d = 1$, we exhibit examples to see that there is indeed some walls.

Theorem 1.1. (Cor.3.5, Prop.3.7, Thm.4.10) *Let X be a K -semistable degeneration of smooth K -semistable del Pezzo surfaces of degree $d \in \{2, 3, 4\}$, and L_i 's are degeneration of all the lines on it. Then the pair $(X, c \sum L_i)$ is K -semistable for any $c > 0$ such that $-(K_X + c \sum L_i)$ is ample. In other words, there are no walls for $M_{d,c}^K$ when c varies from 0 to the log Calabi-Yau threshold.*

Theorem 1.2. (Proposition 5.2) *Let $0 < c < \frac{1}{240}$ be a number, and $M_{1,c}^K$ be the K -moduli space of K -polystable log Fano pairs $(X, c \sum L_i)$, where X is a del Pezzo surface of degree 1 and L_i 's all the lines on X counted with multiplicities. Then there is a wall $c = c' < \frac{1}{288}$, which is given by the destabilization of the surface pair with an A_7 -singularity.*

Acknowledgements This problem was motivated when talking with Schock Nolan. I would also like to express my gratitude to Yuchen Liu for sharing some primary ideas. The author would also like to thank Izzet Coskun and Schock Nolan for many stimulating discussions.

2 Preliminaries

Convention. In this paper, we work over the field of complex numbers \mathbb{C} . By a surface, we mean a complex connected projective algebraic surface. We keep the notions on singularities of surface pairs the same as [KM98] Chapter 2 and Chapter 4.

For the reader's convenience, we state the definitions and results in simple versions, especially in dimension 2. Most of them can be generalized to higher dimension.

Definition 2.1. *Let X be a normal projective variety, and D be an effective \mathbb{Q} -divisor. Then (X, D) is called a log Fano pair if $K_X + D$ is \mathbb{Q} -Cartier and $-(K_X + D)$ is ample. A normal projective variety X is called a \mathbb{Q} -Fano variety if $(X, 0)$ is a klt log Fano pair.*

Definition 2.2. *Let (X, D) be an n -dimensional log Fano pair, and E a prime divisor on a normal projective variety Y , where $\pi : Y \rightarrow X$ is a birational morphism. Then the log discrepancy of (X, E) with respect to E is*

$$A_{(X,D)}(E) := 1 + \text{coeff}_E(K_Y - \pi^*(K_X + D)).$$

We define the S-invariant of (X, D) with respect to E to be

$$S_{(X,D)}(E) := \frac{1}{(-K_X - D)^n} \int_0^\infty \text{vol}_Y(\pi^*(-K_X - D) - tE) dt,$$

and the β -invariant of (X, D) with respect to E to be

$$\beta_{(X,D)}(E) := A_{(X,D)}(E) - S_{(X,D)}(E)$$

The first definition of K-(poly/semi)stability of log Fano pairs used test configurations. We refer the reader to [ADL19] and [Xu21]. There is an equivalent definition using valuations, which is called *valuative criterion* for K-stability. The advantage of this definition is that it is easier to check.

Theorem 2.3. ([Fuj19, Li17, BX19]) *A log Fano pair (X, D) is*

1. *K-semistable if and only if $\beta_{(X,D)}(E) \geq 0$ for any prime divisor E over X ;*
2. *K-stable if and only if $\beta_{(X,D)}(E) > 0$ for any prime divisor E over X .*

The following powerful result is called *interpolation* of K-stability. We only state a version that we will use later. For a more general statement, see for example [ADL19] Proposition 2.13 or [Der16] Lemma 2.6.

Theorem 2.4. ([ADL19] Proposition 2.13, [Der16] Lemma 2.6) *Let X be a K-semistable \mathbb{Q} -Fano variety, and $D \sim_{\mathbb{Q}} -rK_X$ be an effective divisor.*

- (1) *If $(X, \frac{1}{r}D)$ is klt, then (X, cD) is K-stable for any $c \in (0, \frac{1}{r})$;*
- (2) *If $(X, \frac{1}{r}D)$ is log canonical, then (X, cD) is K-semistable for any $c \in (0, \frac{1}{r})$.*

The following result on openness of K-stability is useful in our analysis of K-moduli spaces.

Theorem 2.5. ([BLX22, Xu20]) *Let (X, D) be a klt log Fano pair which degenerates to a K-semistable log Fano pair (X_0, D_0) . Then (X, D) is also K-semistable.*

Recall that the volume of a divisor D on an n -dimensional normal projective variety Y is

$$\text{vol}_Y(D) := \lim_{m \rightarrow \infty} \frac{\dim H^0(Y, mD)}{m^n/n!}.$$

The divisor D is big by definition if and only if $\text{vol}_Y(D) > 0$.

Definition 2.6. *Let $x \in (X, D)$ be an n -dimensional klt singularity. Let $\pi : Y \rightarrow X$ be a birational morphism such that $E \subseteq Y$ is an exceptional divisor whose center on X is $\{x\}$. Then the volume of $(x \in X)$ with respect to E is*

$$\text{vol}_{x,X,D}(E) := \lim_{m \rightarrow \infty} \frac{\dim \mathcal{O}_{X,x}/\{f : \text{ord}_E(f) \geq m\}}{m^n/n!},$$

and the normalized volume of $(x \in X)$ with respect to E is

$$\widehat{\text{vol}}_{x,X,D} := A_{(X,D)}(D) \cdot \text{vol}_{x,X,D}(E).$$

We define the local volume of $x \in (X, D)$ to be

$$\widehat{\text{vol}}(x, X, D) := \inf_E \widehat{\text{vol}}_{x,X,D},$$

where E runs through all the prime divisor over X whose center on X is $\{x\}$.

Theorem 2.7. ([Fuj18, LL19, Liu18]) *Let (X, D) be an n -dimensional K-semistable log Fano pair. Then for any $x \in X$, we have*

$$(-K_X - D)^n \leq \left(1 + \frac{1}{n}\right)^n \widehat{\text{vol}}(x, X, D).$$

Now let us briefly review some results on the K-moduli spaces of log Fano pairs. We mainly state the results in our setting. For more general statements, see [ADL19] Theorem 3.1, Remark 3.25 or [ADL21] Theorem 2.21.

Definition 2.8. *Let $f : (\mathcal{X}, \mathcal{D}) \rightarrow B$ be a proper flat morphism to a reduced scheme with normal, geometrically connected fibers of pure dimension n , where \mathcal{D} is an effective \mathbb{Q} -divisor on \mathcal{X} which does not contain any fiber of f . Then f is called a \mathbb{Q} -Gorenstein flat family of log Fano pairs if $-(K_{\mathcal{X}/B} + \mathcal{D})$ is \mathbb{Q} -Cartier and ample over B .*

Definition 2.9. *Let $0 < c < 1/r$ be a rational number and (X, cD) be a log Fano surface pair such that $D \sim -rK_X$. A \mathbb{Q} -Gorenstein flat family of log Fano pairs $f : (\mathcal{X}, c\mathcal{D}) \rightarrow C$ over a pointed smooth curve $(0 \in C)$ is called a \mathbb{Q} -Gorenstein smoothing of (X, D) if*

- (1) *the divisors \mathcal{D} and $K_{\mathcal{X}/C}$ are both \mathbb{Q} -Cartier, f -ample, and $\mathcal{D} \sim_{\mathbb{Q},f} -rK_{\mathcal{X}/B}$;*
- (2) *both f and $f|_{\mathcal{D}}$ are smooth over $C \setminus \{0\}$, and*
- (3) *$(\mathcal{X}_0, c\mathcal{D}_0) \simeq (X, cD)$.*

Theorem 2.10. ([ADL19]) *Let χ be the Hilbert polynomial of an anti-canonically polarized smooth del Pezzo surface X of degree $d \in \{1, 2, 3, 4\}$. Let L_i 's be the lines on X and r be the positive integer such that $\sum L_i \sim -rK_X$. Let $c \in (0, 1/r)$ be a rational number. Consider the moduli pseudo-functor sending a reduced base S to*

$$\mathcal{M}_{d,c}^K(S) = \left\{ (\mathcal{X}, \mathcal{D})/S \left| \begin{array}{l} (\mathcal{X}, c\mathcal{D})/S \text{ is a } \mathbb{Q}\text{-Gorenstein smoothable log Fano family,} \\ \mathcal{D} \sim_{S,\mathbb{Q}} -rK_{\mathcal{X}/S}, \text{ each fiber } (\mathcal{X}_s, c\mathcal{D}_s) \text{ is a } K\text{-semistable} \\ \text{pair with sum of lines as boundary, and} \\ \chi(\mathcal{X}_s, \mathcal{O}_{\mathcal{X}_s}(-mK_{\mathcal{X}_s})) = \chi(m) \text{ for } m \text{ sufficiently divisible.} \end{array} \right. \right\}.$$

Then there is a smooth quotient stack $\mathcal{M}_{d,c}^K$ of a smooth scheme by a projective general linear group which represents this pseudo-functor. The \mathbb{C} -points of $\mathcal{M}_{d,c}^K$ parameterize K -semistable \mathbb{Q} -Gorenstein smoothable log Fano pairs (X, cD) with Hilbert polynomial $\chi(X, \mathcal{O}_X(-mK_X)) = \chi(m)$ for sufficiently divisible $m \gg 0$ and $D \sim_{\mathbb{Q}} -rK_X$. Moreover, the stack $\mathcal{M}_{d,c}^K$ admits a good moduli space $M_{d,c}^K$, which is a normal projective reduced scheme of finite type over \mathbb{C} , whose \mathbb{C} -points parameterize K -polystable log Fano pairs.

Theorem 2.11. ([ADL19] Theorem 1.2) *Keep the notation as in the last theorem. There are rational numbers*

$$0 = c_0 < c_1 < c_2 < \cdots < c_n = \frac{1}{r}$$

such that for every $0 \leq j < n$, the K -moduli stacks $\mathcal{M}_{d,c}^K$ are independent of the choice of $c \in (c_j, c_{j+1})$. Moreover, for every $0 \leq j < n$ and $0 < \varepsilon \ll 1$, one has open immersions

$$\mathcal{M}_{d,c_j-\varepsilon}^K \hookrightarrow \mathcal{M}_{d,c_j}^K \hookleftarrow \mathcal{M}_{d,c_j+\varepsilon}^K,$$

which descend to projective birational morphisms

$$M_{d,c_j-\varepsilon}^K \rightarrow M_{d,c_j}^K \leftarrow M_{d,c_j+\varepsilon}^K.$$

3 Cubic and quartic del Pezzo pairs

In this section, we prove that the moduli spaces of cubic and quartic del Pezzo pairs do not have wall crossings.

3.1 The cubic case

In [OSS16], the authors essentially proved that for cubic del Pezzo surfaces, the GIT-(semi/poly)stability is equivalent to the K-(semi/poly)stability. As a consequence, the two moduli spaces are isomorphic. However, the K-moduli space was constructed only in recent years. Therefore we give a modern proof here.

Theorem 3.1. ([OSS16]) *A cubic surface is K-semistable if and only if it is GIT-semistable. In particular, the K-moduli space of cubic surfaces is isomorphic to the GIT moduli space.*

Proof. We know that the Fermat cubic surface is K-stable (cf. [Tia87]). By the openness of the K-stability (cf. Theorem 2.5), a general cubic surface is K-stable. Denote by M_3^K by the K-moduli space of cubic del Pezzo surfaces, which parameterizes K-stable smooth cubic surfaces and their K-polystable limits. Let $X \in M_3^K$ be the limit of a \mathbb{Q} -Gorenstein degeneration of smooth cubics $X_t \in M_3^K$.

Let $(x \in X) \simeq (0 \in \mathbb{A}^2/G_x)$ be a singular point (if it exists), then by Theorem 2.7, we have that

$$3 = (-K_X)^2 \leq \frac{9}{4} \widehat{\text{vol}}(x, X) = \frac{9}{4} \cdot \frac{4}{|G_x|} = \frac{9}{|G_x|},$$

which implies that $|G_x| \leq 3$. If $|G_x| = 2$, then x is an A_1 -singularity. If $|G_x| = 3$, then x is either an A_2 -singularity or a $\frac{1}{3}(1, 1)$ -singularity. The latter case is ruled out by [KSB88] Proposition 3.10. We thus conclude that X has at worst A_1 - or A_2 -singularities. It follows from [Fuj90] Section 2 that X can be embedded by $|-K_X|$ into \mathbb{P}^3 as a cubic surface.

Since for hypersurfaces the K-stability implies GIT-stability (cf. [PT09]), then we get a birational and injective morphism $\Phi : M_3^K \rightarrow M_3^{\text{GIT}}$. As both of these two moduli spaces are proper, then Φ is a finite map, and thus Φ is an isomorphism by Zariski Main Theorem. □

In the proof of the theorem, we deduce that a K-polystable cubic del Pezzo surface has at worst A_1 - or A_2 -singularities. This partially recovers the following classical result of Hilbert.

Theorem 3.2. ([MFK94]) *A cubic surface $X \subseteq \mathbb{P}^3$ is*

- (i) *GIT-stable if and only if it has at worst A_1 -singularities;*
- (ii) *strictly polystable if and only if it is isomorphic to the cubic X_0 defined by $x_0^3 = x_1x_2x_3$;*
- (iii) *semistable if and only if it has at worst A_1 - or A_2 -singularities.*

For the stable cubic surfaces, we know how the lines degenerate and the singularities of the pairs we care about are mild.

Lemma 3.3. ([GKS21] Lemma 5.9, 5.10) *Let X be a stable cubic surface and L_i 's be the 27 lines on it. Then the pair $(X, c \sum L_i)$ is log canonical for $0 < c < \frac{1}{9}$.*

Proof. It is proven in [GKS21] that the log canonical threshold is larger than $\frac{1}{9}$. □

We remark here that to make the pair $(X, c \sum L_i)$ log Fano, the coefficient c has to be less than $\frac{1}{9}$. Now let us deal with the surfaces with A_2 -singularities.

Proposition 3.4. *Let $(X, c \sum L_i)$ be a log Fano pair, where X is a K-semistable cubic del Pezzo with A_2 -singularities and L_i 's are the 27 lines on it. Then $(X, c \sum L_i)$ is log canonical for $0 < c < \frac{1}{9}$.*

Proof. Observe that $X_0 = \{x_0^3 = x_1x_2x_3\}$ is a toric surface which is obtained as follows. Let $(x : y : z)$ be the homogeneous coordinates of \mathbb{P}^2 , then blow up the tangent vector given by $\{y = 0\}$ at $(1 : 0 : 0)$, the tangent vector given by $\{z = 0\}$ at $(0 : 1 : 0)$, and the tangent vector given by $\{x = 0\}$ at $(0 : 0 : 1)$. Denote by E_i and F_i the exceptional divisors on the blow-up with self-intersection -2 and -1 respectively, and H_i the (-2) -curves in the class

$$H - E_1 - 2F_1 - E_2 - F_2, \quad H - E_2 - 2F_2 - E_3 - F_3, \quad H - E_3 - 2F_3 - E_1 - F_1,$$

respectively, where $i = 1, 2, 3$. Denote this surface by \tilde{X}_0 . Finally contract the three pairs of (-2) -curves to get X_0 . Let $\pi : \tilde{X}_0 \rightarrow X_0$ be the contraction map. This surface has three A_2 -singularities, and the 27 lines degenerate to the images of F_1, F_2, F_3 , denoted by G_1, G_2, G_3 , each of which is of multiplicity 9. Then by we deduce that

$$\pi^*G_1 = F_1 + \frac{1}{3}E_1 + \frac{2}{3}H_1 + \frac{1}{3}H_2 + \frac{2}{3}E_2,$$

and the other two relations. As a result, we obtain that

$$A_{(X_0, c \sum L_j)}(F_i) = A_{(X_0, c \sum L_j)}(E_i) = A_{(X_0, c \sum L_j)}(H_i) = 1 - 9c > 0$$

for $0 < c < \frac{1}{9}$. Thus the pair $(X_0, c \sum L_j)$ is log canonical for $0 < c < \frac{1}{9}$. The other pairs whose surfaces have A_2 -singularities degenerate to this case, so we get the result we desire. \square

Corollary 3.5. *Let X be a K -semistable cubic surface and L_i 's are the 27 lines on it. Then the pair $(X, c \sum L_i)$ is K -semistable for $0 < c < \frac{1}{9}$. In other words, there are no walls for $M_{3,c}^K$ when c varies from 0 to $\frac{1}{9}$.*

Proof. This follows immediately from Theorem 2.4 and the discussion of the singularities. \square

3.2 The quartic case

The K -stability of quartic surfaces was studied in [MM93, OSS16].

Theorem 3.6. ([MM93]) *A K -semistable quartic del Pezzo surface has at worst A_1 -singularities.*

By [HW81] Theorem 3.4, we know that every ADE del Pezzo surface is obtained by blowing up points on \mathbb{P}^2 in almost general position.

Proposition 3.7. *Let X be a K -semistable quartic del Pezzo surface and L_i 's the 16 lines on it. Then the pair $(X, c \sum L_i)$ is K -semistable for $0 < c < \frac{1}{4}$.*

Proof. By interpolation of K -stability (cf. Theorem 2.4), it suffices to check that these pairs are log canonical for $0 < c < \frac{1}{4}$.

- (i) If X is smooth, then the 16 lines are distinct and $\sum L_i$ is normal crossing. Thus the pair is log canonical automatically.
- (ii) If X has one A_1 -singularity p , then there exist exactly four lines of multiplicity two and eight lines of multiplicity one. These double lines pass through p , while the remaining eight lines avoid it. Let E be the exceptional divisor of the blow-up at p . Then we have that

$$A_{(X, c \sum L_i)}(E) = 1 - 4c > 0$$

for any $0 < c < \frac{1}{4}$. As the weight of the lines at smooth points is at most three, then the pair is log canonical.

- (iii) If X has two A_1 -singularities p and q , then there exist one line of multiplicity four, four lines of multiplicity two and four lines of multiplicity one. For each singularity, there are exactly eight lines (counted with multiplicities) passing through it. Let E be the exceptional divisor of the blow-up at p . Then we have that

$$A_{(X, c \sum L_i)}(E) = 1 - 4c > 0$$

for any $0 < c < \frac{1}{4}$. As the weight of the lines at smooth points is at most four, then the pair is log canonical.

- (iv) Suppose that X has three A_1 -singularities. Notice that X is obtained as follows: blow up \mathbb{P}^2 at a general tangent vector and at three curvilinear points on a general line, where two of them collide, then take the ample model. There exist two lines of multiplicity four, and four lines of multiplicity two. For each singularity, there are exactly eight lines (counted with multiplicities) passing through it. For the same reason as in (ii), the pair is log canonical.
- (v) If X has four A_1 -singularities, then it is obtained as follows: blow up \mathbb{P}^2 at a general point P and at two tangent vectors whose supporting lines pass through P , then take the ample model. There are four lines of multiplicity four. For each singularity, there are exactly eight lines (counted with multiplicities) passing through it. For the same reason as in (ii), the pair is log canonical.

□

Remark 3.8. Using the same argument as in Theorem 3.1, we can prove that the K-moduli space of quartic del Pezzo surfaces is isomorphic to the GIT-moduli space

$$\mathrm{Gr}(2, H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(2)))^{ss} / \mathrm{PGL}(5).$$

As a consequence, each K-moduli space $M_{4,c}^K$ is isomorphic to this GIT-moduli space.

4 Degree two case

In this section, we prove that there are no walls for the K-moduli spaces $M_{2,c}^K$ when we vary the coefficient c from 0 to $\frac{1}{28}$.

Recall that a smooth del Pezzo surface of degree 2 is a double cover of \mathbb{P}^2 branched along a quartic curve, and the 56 lines are sent pairwise to the 28 bitangent lines of the quartic. Thus the K-stability of a degree 2 del Pezzo surface X is equivalent to the K-stability of a del Pezzo pair $(\mathbb{P}^2, \frac{1}{2}C_4)$, where C_4 is the quartic plane curve along which the double cover $X \rightarrow \mathbb{P}^2$ is branched.

In [OSS16], the authors give a description of the K-moduli space of del Pezzo surfaces of degree 2. It is diffeomorphic to the blow-up of $\mathbb{P}H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(4))^{ss} / \mathrm{PGL}(3)$ at the point parameterizing the double conic. Moreover, each point $[s]$ on the exceptional divisor E represents a surface which is a double cover of $\mathbb{P}(1, 1, 4)$ branched along a hyperelliptic curve $z^2 = f_8(x, y)$, where f_8 is a GIT-polystable octic binary form. In [ADL19], the authors study the wall crossing of the K-moduli of pairs (\mathbb{P}^2, cC_4) when c varies from 0 to $\frac{3}{4}$. They proved that there is a unique wall $c = \frac{3}{8}$. As a result, the K-moduli of degree 2 del Pezzo surfaces is isomorphic to a weighted blow-up of the GIT moduli space $\mathbb{P}H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(4))^{ss} / \mathrm{PGL}(3)$ at the point parameterizing the double conic.

For us, to show that there are no walls, by interpolation of K-semistability, it suffices to check that the $(X, \frac{1}{28} \sum E_i)$ is log canonical, where X is a K-polystable degree quadric del

Pezzo surface and E_i 's are the lines on it. It further reduces to checking that $(\mathbb{P}^2, \frac{1}{2}C + \frac{1}{28} \sum L_i)$ or $(\mathbb{P}^2, \frac{1}{2}C + \frac{1}{28} \sum L_i)$ is log canonical (cf. [KM98] Proposition 5.20), where C is the branched curve and L_i 's are the bitangent lines of it.

The classification of the semistable plane quartics is well-known to the experts. For the reader's convenience, we state the result here.

Lemma 4.1. ([MFK94]) *Let $G = \mathrm{PGL}(4)$ act on the space of plane quartic. A plane quartic curve $C_4 \subseteq \mathbb{P}^2$ is*

- (i) *stable if and only if it has at worst A_1 or A_2 singularities;*
- (ii) *strictly polystable if and only if it is either a double conic, or two reduced conics tangential at two points, where at least one of them is irreducible.*

Remark 4.2. In the case (ii), if both of the (distinct) conics are smooth, then we call it *cateye*; if there is a singular one, we call it *ox*.

4.1 Singularities of plane curves

Let us first compute the special case. First notice that if C_4 is the sum of four lines (intersecting transversely), then each L_i coincides with one of the four lines, and the pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical clearly.

Lemma 4.3. *The pair $(\mathbb{P}^2, \frac{1}{2}(C_2 + C'_2) + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$, where C_2 and C'_2 are two conics tangential at two points p and q .*

Proof. Notice that each line L_i is either a tangent line at p or q , or the line $L = \overline{pq}$. Let multiplicity of the line L in L_i 's be $2m$, and those of the two tangent lines be $14 - m$. By symmetry, we only need to look at the log resolution at q . Let E and F be the exceptional divisors with self-intersection -2 and -1 respectively. Then one has $A(E) = 1 - (14 + m)c$ and $A(F) = 1 - 28c$, which are positive when $0 < c < \frac{1}{28}$. \square

If C_4 is the sum of two lines and a conic or two distinct conic which are GIT-polystable, then the pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ degenerates to one of two cases involving cateye or ox. Thus $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for these cases.

Now we deal with the general case: a plane quartic C_4 with at worst A_1 or A_2 singularities. We first assume that C_4 is irreducible. Then the Table 3.2 and Lemma 3.3.1 in [CS03] describe how these bitangent lines degenerate. We claim that all these curves give log canonical pairs $(\mathbb{P}^2, \frac{1}{2}C_4 + \sum L_i)$.

Lemma 4.4. *The pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$, where C_4 has three nodes.*

Proof. We know from [CS03] that the 28 lines degenerate to 4 bitangent lines of multiplicity 1 not passing through the nodes, 6 tangent lines of multiplicity 2 passing through exactly one node, and 3 lines of multiplicity 4 containing two nodes. Thus for each node p , there are 12 lines passing through it, and at most 4 lines tangential to it. One can blow up at p twice to get a log resolution. We have $A(E) = 1 - 16c > 0$ and $A(F) \geq \frac{3}{2} - 20c > 0$ for $0 < c < 1/28$, where E and F are the two exceptional divisors over p with self-intersection -2 and -1 respectively. The singularities at other points are milder, thus $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical as desired. \square

Lemma 4.5. *The pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$, where C_4 has three cusps.*

Proof. We know from [CS03] that the 28 lines degenerate to 4 bitangent lines of multiplicity 1 not passing through the nodes, and 3 lines of multiplicity 9 containing two cusps. Thus for each cusp p , there are 18 lines passing through it, and none of them are tangential to it. One can blow up at p three times to get a log resolution. Let E, F, G be the exceptional divisors with self-intersection $-3, -2, -1$ respectively. We have $A(E) = 1 - 18c > 0$, $A(F) = \frac{3}{2} - 18c > 0$, and $A(G) = 2 - 36c > 0$ for $0 < c < 1/28$. The singularities at other points are milder, thus $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical as desired. \square

Lemma 4.6. *The pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$, where C_4 has one node and two cusps.*

Proof. By [CS03], we see that the 28 lines degenerate to one bitangent line of multiplicity 1 not passing through the singularities, 2 tangent lines of multiplicity 3 passing through exactly one cusp, 1 line of multiplicity 9 containing two cusps, and 2 lines of multiplicity 6 passing through one node and one cusp.

Thus for the node p , there are 12 lines passing through it, and none of them are tangential to it. One can blow up at p to get a log resolution and the log discrepancy with respect to the exceptional divisor is $1 - 12c > 0$.

For each cusp q , there are 18 lines passing through it, and at most 3 lines tangential to it. Resolving q by blowing up three times, one gets three exceptional divisors E, F, G with self-intersection $-3, -2, -1$ respectively. We have $A(E) = 1 - 18c > 0$, $A(F) \geq \frac{3}{2} - 21c > 0$, and $A(G) \geq 2 - 39c > 0$ for $0 < c < 1/28$. The singularities at other points are milder, thus $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical as desired. \square

Lemma 4.7. *The pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$, where C_4 has one cusp and two nodes.*

Proof. By [CS03], we see that the 28 lines degenerate to 2 bitangent lines of multiplicity 1 not passing through the singularities, 2 tangent lines of multiplicity 3 passing through exactly the cusp, 2 tangent lines of multiplicity 2 passing through exactly one node, 1 line of multiplicity 4 containing two nodes, and 2 lines of multiplicity 6 passing through one node and one cusp.

Thus for the cusp p , there are 18 lines passing through it, and at most 6 of them are tangential to it. Resolving p by blowing up three times, one gets three exceptional divisors E, F, G with self-intersection $-3, -2, -1$ respectively. We have $A(E) = 1 - 18c > 0$, $A(F) \geq \frac{3}{2} - 24c > 0$, and $A(G) \geq 2 - 42c > 0$ for $0 < c < 1/28$.

For each node q , there are 12 lines passing through it, and at most 2 lines tangential to it. One can blow up at p twice to get a log resolution. We have $A(E') = 1 - 12c > 0$ and $A(F') \geq \frac{3}{2} - 14c > 0$ for $0 < c < 1/28$, where E' and F' are the two exceptional divisors over p with self-intersection -2 and -1 respectively. The singularities at other points are milder, thus $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical as desired. \square

Notice that the other pairs where C_4 is irreducible degenerate to one of these four. For the case where C_4 is an irreducible cubic union a line, there is also a description on how the bitangent lines degenerate in [CS03] Section 3.4. Applying the same argument to these pairs, we conclude the following.

Proposition 4.8. *Let C_4 be a GIT-polystable plane quartic which is not a double conic. Then the pair $(\mathbb{P}^2, \frac{1}{2}C_4 + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$. In particular, these pairs are K -polystable for $0 < c < 1/28$.*

4.2 Singularities of curves in $\mathbb{P}(1, 1, 4)$

Recall that an octic binary form $f_8(x, y)$ is GIT-semistable if and only if each of its zeros has multiplicity at most 4 (cf. [MFK94] Section 4.1). Also notice that if the point $(x : y : z)$ is an singularity of the curve $C = \{z^2 = f_8(x, y)\}$, then $z = 0$ and $(x : y)$ is a multiple root of $f_8(x, y)$.

Let p_i , $i = 1, \dots, 8$, be the eight intersection points of C and $\{z = 0\}$. Another important observation is that a bitangent line of a plane quartic degenerate to two rulings of $\mathbb{P}(1, 1, 4)$, each of which passes through some p_i , and there are $\binom{8}{2} = 28$ pairs in total. Thus we only need to focus on the vertex p of $\mathbb{P}(1, 1, 4)$ and the singularities of C .

Proposition 4.9. *The pair $(\mathbb{P}(1, 1, 4), \frac{1}{2}C + c \sum L_i)$ is log canonical for $0 < c < \frac{1}{28}$.*

Proof. Notice that C does not pass through the vertex p . Let E be the exceptional divisor of $\text{Bl}_p \mathbb{P}(1, 1, 4) \rightarrow \mathbb{P}(1, 1, 4)$. Then $A(E) = \frac{1}{2} - 14c > 0$ for $0 < c < \frac{1}{28}$.

Suppose that $p_i = (x : y : 0)$ is a singular point of C . We may assume that $x = 0$ and $y = 1$, and locally the equation of C is $z^2 = x^t$, where $t \in \{2, 3, 4\}$. When $t = 4$, p_i is a tacnode, and there are 28 lines passing through it. Let F and G be the exceptional divisors of the minimal resolution of p_i with self-intersection -2 and -1 respectively. Then $A(F) = 1 - 28c > 0$ and $A(G) = 1 - 28c > 0$ for $0 < c < \frac{1}{28}$. This is also true for the cases $t = 2$ and $t = 3$. □

To sum up, we conclude the following by the same argument as in Corollary 3.5.

Theorem 4.10. *Let M_2^K be the K -moduli space of degree 2 del Pezzo surfaces. Then*

$$M_{2,c}^K \simeq M_2^K$$

for any $0 < c < \frac{1}{28}$. In other words, there is no wall crossing when $0 < c < \frac{1}{28}$.

5 Discussion on degree one case

In this section, we display some examples to show that there exist some walls for $M_{1,c}^K$ when c varies from 0 to $\frac{1}{240}$. The following result describes the K -polystable del Pezzo surfaces with at worst ADE singularities. For the analytic description of the K -moduli space $M_{1,0}^K$, see [OSS16] Section 5.

Proposition 5.1. ([OSS16]) *A nodal del Pezzo surface X of degree 1 is K -polystable if and only if it has either only A_k -singularities with $k \leq 7$, or exactly two D_4 -singularities, and X is not isomorphic to one of the surfaces parameterized by $(a_1 : a_2) \in \mathbb{P}^1$, which are hypersurfaces in $\mathbb{P}(1, 1, 2, 3)$ defined by the equations*

$$w^2 = a_1 z^3 + z^2 x^2 + z y^4 + a_2 z^2 y^2.$$

5.1 Degree one del Pezzo with an A_7 -singularity

We aim to prove the following statement.

Proposition 5.2. *Let Y be a del Pezzo surface of degree one with an A_7 -singularity, and L_i 's be the degeneration of the 240 lines. Then the pair $(Y, c \sum L_i)$ is K -unstable for $c \geq \frac{1}{288}$.*

Proof. Let Y be a degree one del Pezzo surface with an A_7 -singularity. Then Y is obtained by blowing up \mathbb{P}^2 at eight curvilinear points on a smooth cubic curve supported at a single point

and taking the ample model. Let E_1, \dots, E_8 be the eight exceptional divisor on the blow-up \tilde{Y} in the order of blow-up, and H the class of the pull-back of $\mathcal{O}_{\mathbb{P}^2}(1)$. Then the proper transform to \tilde{Y} of the 240 lines on Y are of the following types:

- (a) E_8 , intersecting E_7 , of multiplicity 8;
- (b) $H - E_1 - 2E_2 - 2E_3 - \dots - 2E_8$, intersecting E_2 , of multiplicity 28;
- (c) $2H - E_1 - 2E_2 - 3E_3 - 4E_4 - 5\sum_{i \geq 5} E_i$, intersecting E_5 , of multiplicity 56;
- (d) $3H - 2E_1 - 3E_2 - \dots - 8E_7 - 8E_8$, intersecting E_1 and E_7 , of multiplicity 56;
- (e) $4H - 2E_1 - 4E_2 - 6E_3 - \sum_{i \geq 4} (i+3)E_i$, intersecting E_3 and E_8 , of multiplicity 56;
- (f) $5H - \sum_{i=1}^6 2iE_i - 13E_7 - 14E_8$, intersecting E_6 and E_8 , of multiplicity 28;
- (g) $6H - 3E_1 - \sum_{i \geq 2} (2i+1)E_i$, intersecting E_1 and intersecting E_8 twice, of multiplicity 8;

Denote by L_i and l_i the line and its image on Y of one of the above types, and a_i be the multiplicity of the line L_i , where $i = 0, 1, \dots, 6$ is the coefficient of H in the class. Let $\pi : \tilde{Y} \rightarrow Y$ be the blow-down morphism. Then we have

$$\begin{aligned}\pi^*l_0 &= E_8 + \frac{7}{8}E_7 + \frac{6}{8}E_6 + \frac{5}{8}E_5 + \frac{4}{8}E_4 + \frac{3}{8}E_3 + \frac{2}{8}E_2 + \frac{1}{8}E_1, \\ \pi^*l_1 &= L_1 + \frac{1}{4}E_7 + \frac{2}{4}E_6 + \frac{3}{4}E_5 + \frac{4}{4}E_4 + \frac{5}{4}E_3 + \frac{6}{4}E_2 + \frac{3}{4}E_1, \\ \pi^*l_2 &= L_2 + \frac{5}{8}E_7 + \frac{10}{8}E_6 + \frac{15}{8}E_5 + \frac{12}{8}E_4 + \frac{9}{8}E_3 + \frac{6}{8}E_2 + \frac{3}{8}E_1, \\ \pi^*l_3 &= L_3 + E_7 + E_6 + E_5 + E_4 + E_3 + E_2 + E_1, \\ \pi^*l_4 &= L_4 + \frac{3}{8}E_7 + \frac{6}{8}E_6 + \frac{9}{8}E_5 + \frac{12}{8}E_4 + \frac{15}{8}E_3 + \frac{10}{8}E_2 + \frac{5}{8}E_1, \\ \pi^*l_5 &= L_5 + \frac{3}{4}E_7 + \frac{6}{4}E_6 + \frac{5}{4}E_5 + \frac{4}{4}E_4 + \frac{3}{4}E_3 + \frac{2}{4}E_2 + \frac{1}{4}E_1, \\ \pi^*l_6 &= L_6 + \frac{1}{8}E_7 + \frac{2}{8}E_6 + \frac{3}{8}E_5 + \frac{4}{8}E_4 + \frac{5}{8}E_3 + \frac{6}{8}E_2 + \frac{7}{8}E_1.\end{aligned}$$

In particular, one gets that the log discrepancy

$$A_{(Y, c \sum a_i L_i)}(E_3) = A_{(Y, c \sum a_i L_i)}(E_4) = A_{(Y, c \sum a_i L_i)}(E_5) = 1 - 288c < 0$$

when $\frac{1}{288} < c < \frac{1}{240}$. Thus there is a wall $0 < c < \frac{1}{288}$ given by the degeneration of such pairs. \square

5.2 Degree one del Pezzo with two D_4 -singularities

Proposition 5.3. *Let Z be a del Pezzo surface of degree one with exactly two D_4 -singularities, and L_i 's be the degeneration of the 240 lines. Then the pair $(Z, c \sum L_i)$ is K -stable for any $0 < c < \frac{1}{240}$.*

Proof. Let Z be a degree one del Pezzo surface with exactly D_4 -singularities. Such surfaces are weighted hypersurfaces in $\mathbb{P}(1, 1, 2, 3)$ and there is a one-parameter family parameterizing them (cf. [OSS16] Example 5.19). For us, a description by blowing up projective plane is more useful, since we need to figure out the degeneration of the 240 lines.




Figure 1: Blow-up of \mathbb{P}^2 to get \tilde{Z}

We see that Z can be obtained by blowing up \mathbb{P}^2 in the following way. Fix four distinct points p_1, p_2, p_3, p_4 on a line L in \mathbb{P}^2 , and another point $p \in \mathbb{P}^2 \setminus L$. Blow up \mathbb{P}^2 at p_1 along the tangent direction to p with exceptional divisors E_1, F_1 ; at p_2 along the tangent direction to p with exceptional divisors E_2, F_2 ; and at the point p_3 with exceptional divisors E . Finally blow up the length 3 0-dimensional curvilinear subscheme supported on the line $\overline{pp_4}$ and concentrated at p with exceptional divisors E_3, F_3, G_3 (see Figure 1 for the blow-up procedure and Figure 2 for the configuration of the (-1) -curves and (-2) -curves on \tilde{Z}). Denote this surface by \tilde{Z} , and the ample model by Z .

Let H be the class of the pull-back of $\mathcal{O}_{\mathbb{P}^2}(1)$. Then the proper transform to \tilde{Z} of the 240 lines on Y are of the following types:

- (a) E , of multiplicity 24;
- (b) $H - E - E_3 - F_3 - G_3$, of multiplicity 24;
- (c) F_1 , of multiplicity 64;
- (d) F_2 , of multiplicity 64;
- (e) G_3 , of multiplicity 64.

Notice that the configurations of the curves in Figure 2 is symmetric: the black curves denote the (-2) -curves and the red lines denote the (-1) -curves. Using the same computation as in Section 5.1, one sees that the minimal log discrepancy of the pair $(Z, c \sum a_i L_i)$ is $1 - 240c > 0$, when $0 < c < \frac{1}{240}$. Thus the surfaces with exactly two D_4 -singularities do not contribute any wall. □

Remark 5.4. Notice that the four points p_1, \dots, p_4 on \mathbb{P}^1 have a cross-ratio. This also explains why we have a one-dimensional family of surfaces with exactly two D_4 -singularities.

5.3 Degree one del Pezzo with an A_8 -singularity two $\frac{1}{9}(1, 2)$ -singularities

Among all K-polystable degeneration of smooth degree 1 del Pezzo surface, there is a special one X_∞ with an A_8 -singularity and two $\frac{1}{9}(1, 2)$ -singularities. The surface X_∞ can be viewed as




Figure 2: Configuration of the lines on \tilde{Z}

a degree 18 hypersurface in $\mathbb{P}(1, 2, 9, 9)$ given by the equation $z_3 z_4 = z_2^9$, where $(z_1 : z_2 : z_3 : z_4)$ is the coordinates of weights 1, 2, 9, 9 respectively. Moreover, projecting to $\mathbb{P}(1, 2, 9)$ by

$$(z_1 : z_2 : z_3 : z_4) \mapsto (z_1 : z_2 : z_3 + z_4)$$

realizes X_∞ as a double cover of $\mathbb{P}(1, 2, 9)$ branched along the curve $C_\infty = \{v^9 = w^2\}$, where $(u : v : w)$ is the coordinate of $\mathbb{P}(1, 2, 9)$. In fact, this double cover map is given by the linear system $|-2K_{X_\infty}|$.

Recall that for a smooth del Pezzo surface X_t of degree one, the map given by the linear system $|-2K_{X_t}|$ is a double cover to $\mathbb{P}(1, 1, 2) \subseteq \mathbb{P}^3$ branched along a sextic curve C_t . The 240 lines on X_t are sent pairwise to the 120 conics on $\mathbb{P}(1, 1, 2)$ obtained by intersecting the $\mathbb{P}(1, 1, 2)$ with the 120 tritangent planes of C_t in \mathbb{P}^3 . See [KRSNS18] for details.

The degeneration of $\mathbb{P}(1, 1, 2)$ to $\mathbb{P}(1, 2, 9)$ can be observed in \mathbb{P}^{15} : they are embedded into \mathbb{P}^{15} by the complete linear series $|\mathcal{O}(6)|$ and $|\mathcal{O}(18)|$, respectively. In particular, the 120 conics degenerate to curves in $\mathbb{P}(1, 2, 9)$ of degree 6. Notice that the pair $(\mathbb{P}(1, 2, 9), \frac{1}{2}C_\infty)$ is a \mathbb{T} -variety of complexity one: there is a \mathbb{G}_m -action given by

$$\lambda \cdot (u : v : w) = (u : \lambda^2 v : \lambda^9 w).$$

In particular, each of the 120 sextic curves is \mathbb{G}_m -invariant, and hence is defined by one of the following four equations:

$$u^6 = 0, \quad u^4 v = 0, \quad u^2 v^2 = 0, \quad v^3 = 0.$$

However, the multiplicity of them is not clear to us. We have the following partial result.

Proposition 5.5. *Let X_∞ be the degree 1 del Pezzo surface as above, and L_i 's be the 240 lines on it counted with multiplicities. Then the pair $(X_\infty, c \sum L_i)$ is either K -polystable for any*

$0 < c < \frac{1}{240}$ or K -unstable for any $0 < c < \frac{1}{240}$. Moreover, it is K -polystable if and only if

$$\text{ord}_{u=0} \left(\sum l_j \right) = 240 \quad \text{and} \quad \text{ord}_{u=0} \left(\sum l_j \right) = 240,$$

where l_i 's are the 120 sextic curves in $\mathbb{P}(1, 2, 9)$ given by the images of L_i 's.

Proof. Assume we are in the case when $\text{ord}_{u=0}(\sum l_j) = 240$ and $\text{ord}_{u=0}(\sum l_j) = 240$. By [LZ22] Theorem 1.2, it suffices to check the K -stability of the pair $(\mathbb{P}(1, 2, 9), \frac{1}{2}C_\infty + c \sum l_i)$, which is a \mathbb{T} -pair of complexity one. By [ACC⁺21] Theorem 1.3.9, we only need to compute the β -invariant of the pair with respect to all \mathbb{G}_m -invariant divisors on $\mathbb{P}(1, 2, 9)$ (see also [Zha22] Theorem 2.9 for the statement for pairs). The divisor $\{u = 0\}$ is the unique horizontal divisor on X_∞ . We have that

$$A_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(\{u = 0\}) = 1 - 240c$$

and that

$$S_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(\{u = 0\}) = \frac{(3 - 720c)}{\mathcal{O}(1)^2} \int_0^1 \mathcal{O}(1 - t)^2 dt = 1 - 240c,$$

and thus $\beta_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(\{u = 0\}) = 0$. For the vertical divisor $\{v = 0\}$ we have that

$$A_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(\{v = 0\}) = 1 - 240c$$

and that

$$S_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(\{v = 0\}) = \frac{(3 - 720c)}{\mathcal{O}(1)^2} \int_0^{\frac{1}{2}} \mathcal{O}(1 - 2t)^2 dt = \frac{1}{2}(1 - 240c).$$

For other vertical divisors $D_t = \{v^9 = tw^2\}$ with $\lambda \in \mathbb{C}^*$, similarly, we have that that $\beta_{(\mathbb{P}(1,2,9), \frac{1}{2}C_\infty + c \sum l_i)}(D_t) > 0$. This conclude for the case $\text{ord}_{u=0}(\sum l_j) = \text{ord}_{u=0}(\sum l_j) = 240$. For the other case, the β -invariant for the horizontal divisor $\{u = 0\}$ is non-zero when $0 < c < \frac{1}{240}$ so the pair is K -unstable. □

References

- [ABHLX20] Jarod Alper, Harold Blum, Daniel Halpern-Leistner, and Chenyang Xu. Reductivity of the automorphism group of K -polystable Fano varieties. *Inventiones mathematicae*, 222(3):995–1032, 2020.
- [ACC⁺21] Carolina Araujo, Ana-Maria Castravet, Ivan Cheltsov, Kento Fujita, Anne-Sophie Kaloghiros, Jesus Martinez-Garcia, Constantin Shramov, Nivedita Viswanathan, et al. The Calabi problem for Fano threefolds. 2021.
- [ADL19] Kenneth Ascher, Kristin DeVleming, and Yuchen Liu. Wall crossing for K -moduli spaces of plane curves. *arXiv preprint arXiv:1909.04576*, 2019.
- [ADL21] Kenneth Ascher, Kristin DeVleming, and Yuchen Liu. K -moduli of curves on a quadric surface and $K3$ surfaces. *Journal of the Institute of Mathematics of Jussieu*, pages 1–41, 2021.
- [BLX22] Harold Blum, Yuchen Liu, and Chenyang Xu. Openness of K -semistability for Fano varieties. *Duke Mathematical Journal*, 171(13):2753–2797, 2022.

- [BX19] Harold Blum and Chenyang Xu. Uniqueness of K-polystable degenerations of Fano varieties. *Annals of Mathematics*, 190(2):609–656, 2019.
- [CS03] Lucia Caporaso and Edoardo Sernesi. Recovering plane curves from their bitangents. *J. Algebraic Geom.*, 12:225–244, 2003.
- [Der16] Ruadháí Dervan. On K-stability of finite covers. *Bulletin of the London Mathematical Society*, 48(4):717–728, 2016.
- [Fuj90] Takao Fujita. On singular del Pezzo varieties. In *Algebraic geometry*, pages 117–128. Springer, 1990.
- [Fuj18] Kento Fujita. Optimal bounds for the volumes of Kähler-Einstein Fano manifolds. *American Journal of Mathematics*, 140(2):391–414, 2018.
- [Fuj19] Kento Fujita. A valuative criterion for uniform K-stability of \mathbb{Q} -Fano varieties. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2019(751):309–338, 2019.
- [GKS21] Patricio Gallardo, Matt Kerr, and Luca Schaffler. Geometric interpretation of toroidal compactifications of moduli of points in the line and cubic surfaces. *Advances in Mathematics*, 381:107632, 2021.
- [HKT09] Paul Hacking, Sean Keel, and Jenia Tevelev. Stable pair, tropical, and log canonical compactifications of moduli spaces of del pezzo surfaces. *Inventiones mathematicae*, 178(1):173–227, 2009.
- [HKT13] Brendan Hassett, Andrew Kresch, and Yuri Tschinkel. On the moduli of degree 4 del Pezzo surfaces. *Development of moduli theory (Kyoto, 2013)*, 69:349–386, 2013.
- [HW81] Fumio Hidaka and Kei-ichi Watanabe. Normal Gorenstein surfaces with ample anti-canonical divisor. *Tokyo Journal of Mathematics*, 4(2):319–330, 1981.
- [Jia20] Chen Jiang. Boundedness of \mathbb{Q} -Fano varieties with degrees and alpha-invariants bounded from below. *Ann. Sci. Éc. Norm. Supér (4)*, 53(5):1235–1248, 2020.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134. Cambridge university press, 1998.
- [KRSNS18] Avinash Kulkarni, Yue Ren, Mahsa Sayyary Namin, and Bernd Sturmfels. Real space sextics and their tritangents. In *Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*, pages 247–254, 2018.
- [KSB88] János Kollár and Nicholas I Shepherd-Barron. Threefolds and deformations of surface singularities. *Inventiones mathematicae*, 91(2):299–338, 1988.
- [Li17] Chi Li. K-semistability is equivariant volume minimization. *Duke Mathematical Journal*, 166(16):3147–3218, 2017.
- [Liu18] Yuchen Liu. The volume of singular Kähler-Einstein Fano varieties. *Compositio Mathematica*, 154(6):1131–1158, 2018.
- [LL19] Chi Li and Yuchen Liu. Kähler-Einstein metrics and volume minimization. *Advances in Mathematics*, 341:440–492, 2019.

- [LWX21] Chi Li, Xiaowei Wang, and Chenyang Xu. Algebraicity of the metric tangent cones and equivariant K-stability. *Journal of the American Mathematical Society*, 34(4):1175–1214, 2021.
- [LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to K-stability. *Annals of Mathematics*, 196(2):507–566, 2022.
- [LZ22] Yuchen Liu and Ziwen Zhu. Equivariant k-stability under finite group action. *International Journal of Mathematics*, 33(01):2250007, 2022.
- [MFK94] David Mumford, John Fogarty, and Frances Kirwan. *Geometric invariant theory*, volume 34. Springer Science & Business Media, 1994.
- [MM93] Toshiaki Mabuchi and Shigeru Mukai. Stability and Einstein-Kähler metric of a quartic del Pezzo surface. In *Einstein metrics and Yang-Mills connections*, pages 133–160. CRC Press, 1993.
- [OSS16] Yuji Odaka, Cristiano Spotti, and Song Sun. Compact moduli spaces of del Pezzo surfaces and Kähler-Einstein metrics. *Journal of Differential Geometry*, 102(1):127–172, 2016.
- [PT09] S. T. Paul and G. Tian. CM stability and the generalized Futaki invariant II. *Astérisque*, 328:339–354, 2009.
- [Tia87] Gang Tian. On Kähler-Einstein metrics on certain Kähler manifolds with $c_1(m) > 0$. *Inventiones mathematicae*, 89(2):225–246, 1987.
- [Xu20] Chenyang Xu. A minimizing valuation is quasi-monomial. *Annals of Mathematics*, 191(3):1003–1030, 2020.
- [Xu21] Chenyang Xu. K-stability of Fano varieties: an algebro-geometric approach. *EMS Surv. Math. Sci.*, 8(1):265–354, 2021.
- [XZ20] Chenyang Xu and Ziquan Zhuang. On positivity of the CM line bundle on K-moduli spaces. *Annals of mathematics*, 192(3):1005–1068, 2020.
- [Zha22] Junyan Zhao. K-moduli of quintic del Pezzo pairs and moduli of genus six curves. *arXiv preprint arXiv:2212.06992*, 2022.