

## AN ALMOST STRONG RELATION

SHIMON GARTI AND ANDRÉS VILLAVECES

ABSTRACT. Let  $\mu$  be a strong limit singular cardinal. We prove that if  $2^\mu > \mu^+$  then  $\binom{\mu^+}{\mu} \rightarrow \binom{\tau}{\mu}_{<cf(\mu)}$  for every ordinal  $\tau < \mu^+$ . We obtain an optimal positive relation under  $2^\mu = \mu^+$ , as after collapsing  $2^\mu$  to  $\mu^+$  this positive relation is preserved.

---

2010 *Mathematics Subject Classification.* 03E04, 03E02, 03E10.

*Key words and phrases.* Polarized relations, elementary submodels, pcf theory.

## 0. INTRODUCTION

The *strong polarized partition relation*  $\binom{\lambda}{\kappa} \rightarrow \binom{\lambda}{\kappa}_\theta$  says that for every coloring  $c : \lambda \times \kappa \rightarrow \theta$  one can find  $A \in [\lambda]^\lambda, B \in [\kappa]^\kappa$  such that  $c''(A \times B)$  is constant. The *almost strong polarized partition relation*  $\binom{\lambda}{\kappa} \rightarrow \binom{\tau}{\kappa}_\theta$  for every  $\tau < \lambda$  asserts that for every coloring  $c : \lambda \times \kappa \rightarrow \theta$  and any ordinal  $\tau < \lambda$  one can find a color  $i < \theta$  and a pair of sets  $A \subseteq \lambda, B \subseteq \kappa$  such that  $\text{otp}(A) = \tau, |B| = \kappa$  and  $c''(A \times B) = \{i\}$ . The purpose of this paper is to prove such a relation at strong limit singular cardinals. The main theorem of the paper improves both a result of Erdős-Hajnal-Rado for strong limit singular cardinals with countable cofinality and a result of Shelah for strong limit singular cardinals with uncountable cofinality.

In Theorem 42 of [3] by Erdős, Hajnal and Rado it is shown that  $\binom{\mu^+}{\mu} \rightarrow \binom{\mu}{\mu}_2$  whenever  $\text{cf}(\mu) = \omega$ . Assuming that  $2^\mu > \mu^+$  and  $\mu$  is a strong limit cardinal with countable cofinality we increase the order type of the first coordinate, namely  $\text{otp}(A) = \tau$  for every ordinal  $\tau < \mu^+$ . Similarly, Shelah proved in [13] that  $\binom{\mu^+}{\mu} \rightarrow \binom{\mu^+}{\mu}_{<\text{cf}(\mu)}$  whenever  $\mu > \text{cf}(\mu) > \omega$ ,  $\mu$  is strong limit and  $2^\mu > \mu^+$ . Actually, Shelah's theorem applies to singular cardinals of countable cofinality as well, but in these cardinals it is less interesting since it holds even if without the assumption that  $2^\mu > \mu^+$ . Anyway, we show that the ordinal  $\mu + 1$  can be replaced by every ordinal  $\tau < \mu^+$  for singular cardinals of uncountable cofinality under the same assumptions of  $2^\mu > \mu^+$ . These results give a positive answer to Question 4.8 from [7]. Our proof follows in the footsteps of Shelah's proof in [13], but we add a feature which enables us to stretch monochromatic sets of size  $\mu$  and to obtain sets of the same cardinality but a larger order type.

Let us indicate that the strong relation  $\binom{\mu^+}{\mu} \rightarrow \binom{\mu^+}{\mu}_2$  is consistent for singular cardinals (see [6], [7]), using several assumptions about the structure of cardinal arithmetic. For this relation the assumption  $2^\mu > \mu^+$  is necessary since  $2^\mu = \mu^+$  implies  $\binom{\mu^+}{\mu} \not\rightarrow \binom{\mu^+}{\mu}_2$  as proved in [3], and usually the positive strong relation requires more than just  $2^\mu > \mu^+$ . But for our result, it suffices to assume  $2^\mu > \mu^+$  in order to obtain the almost strong relation at strong limit singular cardinals. Moreover, although collapsing  $2^\mu$  to  $\mu^+$  destroys the strong relation  $\binom{\mu^+}{\mu} \rightarrow \binom{\mu^+}{\mu}_2$  it still preserves the almost strong relation, as will be indicated at the end of the paper.

The result of Shelah was introduced in an expository article of Menachem Kojman, [11], who simplified the proof. The presentation here owes a lot to Kojman's paper, and in particular we follow the notation of that paper. We divide the rest of the paper into two sections. In the first one we discuss the general concept of pcf arrays of elementary submodels. In the second section we prove the combinatorial theorem. The main reason for this separation is that pcf arrays seem to represent a general method. Shelah remarked that

tentatively this concept might lead to other mathematical proofs, though no such one has been discovered so far.

Our notation is standard. When we use elementary submodels of some  $\mathcal{H}(\chi)$  we mean that  $\chi$  is a sufficiently large regular cardinal and this structure can be augmented by any finite number of additional predicates (like a well-ordering of the elements of  $\mathcal{H}(\chi)$ ). If  $\kappa = \text{cf}(\kappa) < \lambda$  then  $S_\kappa^\lambda = \{\delta \in \lambda : \text{cf}(\delta) = \kappa\}$ . We shall use this notation in most cases when  $\lambda$  is a regular cardinal, in which case  $S_\kappa^\lambda$  is a stationary subset of  $\lambda$ . An ordinal  $\eta$  is indecomposable iff it has the form  $\omega^\beta$ , where this denotes ordinal exponentiation. Indecomposable ordinals behave like cardinals in the sense that if  $\alpha < \eta$  then the order type of  $\eta - \alpha$  is still  $\eta$ . The collection of all indecomposable ordinals of some  $\lambda = \text{cf}(\lambda) > \aleph_0$  is a club subset of  $\lambda$ . If  $E$  is a club subset of  $\lambda$  then  $\text{acc}(E)$  is the set of accumulation points of  $E$ , and it is a club of  $\lambda$  as well.

For basic background in pcf theory we suggest [1] and [2]. For advanced theorems, including the main tool used in this paper, we refer to [12]. We thank Saharon Shelah for a very helpful discussion concerning several issues in pcf theory. A good background about polarized relations appears in [14] and in [9].

## 1. PCF ARRAYS OF ELEMENTARY SUBMODELS

We define in this section the notion of a *pcf array of elementary submodels*. We provide sufficient conditions for the existence of this kind of arrays. This will be useful in the proof of the almost strong polarized relation later in the paper.

Let  $\mu$  be a singular cardinal,  $\kappa = \text{cf}(\mu)$ . Let  $\langle \mu_\alpha : \alpha < \kappa \rangle$  be an increasing sequence of regular cardinals such that  $\mu = \bigcup_{\alpha < \kappa} \mu_\alpha$ . Let  $J$  be an ideal over  $\kappa$  which contains the ideal  $J_\kappa^{\text{bd}}$  of all bounded subsets of  $\kappa$ .

The relation  $f <_J g \Leftrightarrow \{\alpha \in \kappa : g(\alpha) \leq f(\alpha)\} \in J$  defined on elements from the product  $\prod_{\alpha < \kappa} \mu_\alpha$  is usually just a partial order. As such, it may possess a  $J$ -increasing cofinal sequence of functions or may not. If there is a cofinal  $J$ -increasing sequence then the minimal length of such a  $J$ -increasing sequence is denoted by  $\text{tcf}(\prod_{\alpha < \kappa} \mu_\alpha, J)$  and called *the true cofinality* of the product.

If  $J$  is a prime ideal (equivalently, if the dual of  $J$  is an ultrafilter) then the partial order defined above is actually a linear order and hence a cofinal sequence exists. The spectrum of possible cofinalities is the most basic concept of pcf theory. Formally, if  $\mathfrak{a}$  is a set of regular cardinals then  $\text{pcf}(\mathfrak{a}) = \{\text{tcf}(\prod \mathfrak{a}, \mathcal{U}) : \mathcal{U} \text{ is an ultrafilter over } \mathfrak{a}\}$ . For the majority of pcf theorems,  $\mathfrak{a}$  has to be progressive, where a set of regular cardinals  $\mathfrak{a}$  is called *progressive* iff  $|\mathfrak{a}| < \min(\mathfrak{a})$ . In many cases the set  $\mathfrak{a}$  is an end-segment of the set of all regular cardinals below a given singular cardinal  $\mu$ . For this set to be progressive one has to assume that  $\mu$  is not a fixed point of the  $\aleph$ -function.

A fundamental theorem of pcf theory is that if  $\mu > \text{cf}(\mu) = \kappa$  then there exists a sequence  $\langle \mu_\alpha : \alpha \in \kappa \rangle$  such that  $\text{tcf}(\prod_{\alpha \in \kappa} \mu_\alpha, J_\kappa^{\text{bd}}) = \mu^+$ . According to this theorem we can realize  $\mu^+$  as a true cofinality using the ideal of bounded subsets of  $\kappa$ . We indicate that it might happen that  $\lambda \in \text{pcf}(\mathfrak{a})$  where  $\mathfrak{a} \subseteq \text{Reg} \cap \mu$  and  $\lambda$  cannot be expressed as a true cofinality with  $J_\kappa^{\text{bd}}$ . But in the case of  $\lambda = \mu^{++}$  we have the following result from [12, Chapter VIII, Theorem 1.1].

**Theorem 1.1.** *Assume that  $\mu > \text{cf}(\mu) = \kappa \geq \aleph_0$ , and  $\mu$  is a strong limit cardinal.*

*Then there exists an increasing sequence of regular cardinals  $\langle \lambda_i : i < \kappa \rangle$  such that  $\mu = \bigcup_{i < \kappa} \lambda_i$  and  $\text{tcf}(\prod_{i < \kappa} \lambda_i, J_\kappa^{\text{bd}}) = \mu^{++}$ .*

□<sub>1.1</sub>

One of the virtues of  $J_\kappa^{\text{bd}}$  is that if we take any unbounded subsequence  $\langle \lambda_{i_\beta} : \beta \in \kappa \rangle$  of the above sequence then  $\text{tcf}(\prod_{\beta \in \kappa} \lambda_{i_\beta}, J_\kappa^{\text{bd}}) = \mu^{++}$  as well. Indeed, if  $(f_\alpha : \alpha \in \mu^{++})$  witnesses  $\text{tcf}(\prod_{i < \kappa} \lambda_i, J_\kappa^{\text{bd}}) = \mu^{++}$  then the restriction of the scale to the subsequence  $\langle \lambda_{i_\beta} : \beta \in \kappa \rangle$  witnesses  $\text{tcf}(\prod_{\beta \in \kappa} \lambda_{i_\beta}, J_\kappa^{\text{bd}}) = \mu^{++}$ .

In general, it is possible that  $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i, J)$  for some  $J \supsetneq J_\kappa^{\text{bd}}$  and  $\lambda$  cannot be realized as the true cofinality of some sequence of regular cardinals




FIGURE 1. A pcf-array

below  $\mu$  with the ideal  $J_\kappa^{\text{bd}}$ . In such cases we do not know how to carry out our argument. The problematic point is that by thinning-out the sequence to  $\langle \lambda_{i_\beta} : \beta \in \kappa \rangle$  we may change the true cofinality. However, in some cases one can ensure that the true cofinality is obtained by the ideal  $J_\kappa^{\text{bd}}$ . A notable example is when  $\mu > \text{cf}(\mu) = \kappa > \aleph_0$  and  $\kappa$  is weakly compact, as proved in [8]. Another case, crucial for our proof, is the case in which  $\lambda = \mu^{++}$ , see [13, Theorem 1.4].

In what follows, we will define matrices of models  $(M_{\alpha i})_{\alpha < \lambda, i < \kappa}$ , where  $\mu > \text{cf}(\mu) = \kappa$  and  $\lambda \geq \mu^+$ ; furthermore, for some increasing sequence of regular cardinals  $\langle \lambda_i : i < \kappa \rangle$ ,  $\mu = \bigcup_{i < \kappa} \lambda_i$ . This last part connects the  $\lambda \times \kappa$ -matrix to the statement  $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i, J)$ .

**Definition 1.2** (Pcf array). Let  $\mu$  be a singular cardinal,  $\kappa = \text{cf}(\mu)$  and  $\chi$  a sufficiently large regular cardinal above  $2^\mu$ .

Assume that  $\mu < \lambda < \text{tcf}(\prod_{i < \kappa} \lambda_i, J)$ ,  $J \supseteq J_\kappa^{\text{bd}}$  and for some increasing sequence of regular cardinals  $\bar{\lambda} = \langle \lambda_i \mid i < \kappa \rangle$ ,  $\mu = \bigcup_{i < \kappa} \lambda_i$ .

A  $\lambda \times \kappa$  pcf array of models  $\mathcal{M}$  is a sequence of elementary submodels of  $\mathcal{H}(\chi)$  of the form  $\langle M_{\alpha i} : \alpha < \lambda, i < \kappa \rangle$  such that (see Figure 1):

- ( $\aleph$ ) For every  $\alpha \in \lambda$ ,  $M_{\alpha i} \prec M_{\alpha j}$  whenever  $i < j$ .
- ( $\beth$ ) For every  $\alpha < \beta < \lambda$  there exists  $u \in J$  such that  $M_{\alpha j} \prec M_{\beta j}$  whenever  $j \in \kappa - u$ .
- ( $\beth$ )  $|M_{\alpha i}| < \lambda_i$  for every  $\alpha \in \lambda, i \in \kappa$ .

We shall say that  $\mathcal{M}$  is based on the sequence  $\bar{\lambda}$  and the ideal  $J$ . The following figure demonstrates the idea in the case where  $J = J_\kappa^{\text{bd}}$ ; in this case, for every  $\alpha < \beta < \lambda$  there exists an ordinal  $i_{\alpha\beta} < \kappa$  such that  $M_{\alpha j} \prec M_{\beta j}$  whenever  $i_{\alpha\beta} \leq j < \kappa$ .

A central concept related to pcf arrays is the following:

**Definition 1.3** (The characteristic sequence). Let  $\mathcal{M} = \langle M_{\alpha i} : \alpha < \lambda, i < \kappa \rangle$  be a pcf array of models based on  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ . For every  $\alpha < \lambda, i < \kappa$  let  $f_\alpha(i) = \sup(M_{\alpha i} \cap \lambda_i)$ . Each function  $f_\alpha \in \prod_{i < \kappa} \lambda_i$  is called the *characteristic function* of  $\mathcal{M}$  at stage  $\alpha$ , and  $(f_\alpha)_{\alpha < \lambda}$  is called the *characteristic sequence*.

Notice that each  $f_\alpha$  is an element of  $\prod_{i < \kappa} \lambda_i$  since  $|M_{\alpha i}| < \lambda_i$ . The main point of the proof in the next section is that for a pcf array  $\mathcal{M}$  and  $\lambda$ -many functions  $f_\alpha$  as above there is a single function  $h \in \prod_{i < \kappa} \lambda_i$  which bounds them all. The reason is that  $\text{tcf}(\prod_{i < \kappa} \lambda_i, J) > \lambda$ . Now the values of  $h$  will be ordinals outside the pertinent models; later in the proof we will use the fact that any first order property of them will be reflected to many ordinals below.

**Claim 1.4.** *Let  $\mu$  be a strong limit singular cardinal,  $\kappa = \text{cf}(\mu) > \omega$  and  $2^\mu > \mu^+$ . Let  $\bar{\mu} = \langle \mu_i : i < \kappa \rangle$  be an increasing continuous sequence of cardinals such that  $\mu = \bigcup_{i < \kappa} \mu_i$  and  $\mu_0 > \kappa$ . Fix also another increasing sequence of cardinals  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$  such that  $\mu = \bigcup_{i < \kappa} \lambda_i$ . There exists a pcf array  $\mathcal{M} = \langle M_{\alpha i} : \alpha < \mu^+, i < \kappa \rangle$  based on  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$  and  $J_\kappa^{\text{bd}}$  such that  $\mu_i^+ M_{\alpha i} \subseteq M_{\alpha i}$  for every  $\alpha \in \mu^+, i < \kappa$  and  $\alpha \subseteq \bigcup_{i < \kappa} M_{\alpha i}$  for every  $\alpha \in \mu^+$ .*

*Proof.*

We may assume, without loss of generality, that  $2^{(\mu_i^+)} < \lambda_i$  for every  $i < \kappa$  by taking a subsequence of  $\bar{\lambda}$  if needed. We use here the fact that  $\mu$  is a strong limit cardinal. Choose for every  $\alpha \in \mu^+, i < \kappa$  some  $M_{\alpha i} \prec \mathcal{H}(\chi)$  such that:

- (a)  $|M_{\alpha i}| = 2^{(\mu_i^+)}$  and  $2^{(\mu_i^+)} \subseteq M_{\alpha i}$ .
- (b)  $\mu_i^+ M_{\alpha i} \subseteq M_{\alpha i}$ .
- (c)  $\alpha \in M_{\alpha i}$  and  $\alpha \subseteq \bigcup_{i < \kappa} M_{\alpha i}$ .
- (d)  $\{M_{\alpha i} : (\alpha, i) <_{\text{lex}} (\beta, j)\} \in M_{\beta j}$ .

The only non-trivial requirement in the construction is  $\alpha \subseteq \bigcup_{i < \kappa} M_{\alpha i}$ , since typically  $\mu < \alpha$  and  $|M_{\alpha i}| < \mu$  for every  $i \in \kappa$ . However,  $\alpha < \mu^+$  so  $\alpha$  is expressible as  $\bigcup_{i < \kappa} \alpha_i$  such that  $|\alpha_i| \leq 2^{(\mu_i^+)}$  for every  $i \in \kappa$ , and the construction follows.

We now explain why the construction works. Requirement  $(\beth)$  of Definition 1.2 is guaranteed by (a) and our assumption on the  $\lambda_i$ 's, so we check  $(\aleph)$  and  $(\beth)$ . We first note that if  $(\alpha, i) <_{\text{lex}} (\beta, j)$  and  $\alpha \in M_{\beta j}$  then  $M_{\alpha i} \in M_{\beta j}$ . This follows from (d), and from the fact that  $i \in M_{\beta j}$  as  $\kappa \in 2^{(\mu_i^+)} \subseteq M_{\beta j}$ . Second, if  $M_{\alpha i} \in M_{\beta j}$  and  $j \leq i$  then  $M_{\alpha i} \subseteq M_{\beta j}$ . Indeed,  $\mathcal{H}(\chi) \models |M_{\alpha i}| = 2^{(\mu_i^+)}$  so there is a function  $g : 2^{(\mu_i^+)} \rightarrow M_{\alpha i}$  which enumerates its elements. Since  $M_{\beta j} \prec \mathcal{H}(\chi)$  there is such a function in  $M_{\beta j}$  and since  $2^{(\mu_i^+)} \leq 2^{(\mu_j^+)} \subseteq M_{\beta j}$  we see that the range of this function is a

subset of  $M_{\beta j}$ , namely  $M_{\alpha i} \subseteq M_{\beta j}$ . Finally, if  $M_{\alpha i} \subseteq M_{\beta j}$  then  $M_{\alpha i} \prec M_{\beta j}$  since both are elementary submodels of the same  $\mathcal{H}(\chi)$ .

Fix any  $\alpha \in \mu^+$ . If  $i < j$  then  $(\alpha, i) <_{\text{lex}} (\alpha, j)$ . Since  $\alpha \in M_{\alpha j}$  by (c), we infer that  $M_{\alpha i} \prec M_{\alpha j}$  so  $(\aleph)$  holds. To check  $(\beth)$ , assume that  $\alpha < \beta$ . Since  $\beta \subseteq \bigcup_{i \in \kappa} M_{\beta i}$  and  $\alpha < \beta$  there is  $i_{\alpha\beta} \in \kappa$  such that  $\alpha \in M_{\beta i_{\alpha\beta}}$ . If  $j \geq i_{\alpha\beta}$  then  $M_{\beta i_{\alpha\beta}} \subseteq M_{\beta j}$  and hence  $\alpha \in M_{\beta j}$ . As  $(\alpha, j) <_{\text{lex}} (\beta, j)$  we have  $M_{\alpha j} \prec M_{\beta j}$  thus proving  $(\beth)$ .

□<sub>1.4</sub>

## 2. ALMOST STRONG RELATIONS

In this section we shall prove the combinatorial results concerning the polarized relation. We first provide a useful lemma which says, roughly, that if a coloring of pairs depends only on one coordinate then it has large monochromatic products. This observation is pertinent for both cases of countable and uncountable cofinality.

**Lemma 2.1.** *Assume that  $\mu > \text{cf}(\mu)$ ,  $\tau < \mu^+$  and  $\theta < \text{cf}(\mu)$ .*

*Let  $c : \mu^+ \times \mu \rightarrow \theta$  be any coloring.*

*If  $A \subseteq \mu^+$ ,  $\text{otp}(A) = \tau$ ,  $B_0 \in [\mu]^\mu$  and  $c \upharpoonright (A \times B_0)$  depends only on the second coordinate then there is a set  $B \in [B_0]^\mu$  such that  $c \upharpoonright (A \times B)$  is constant.*

*Proof.*

Define a function  $d : B_0 \rightarrow \theta$  as follows. Fix any  $\alpha \in A$  and let  $d(\beta) = c(\alpha, \beta)$  for every  $\beta \in B_0$ . The choice of  $\alpha$  is unimportant by the assumption of the lemma. Since  $|B_0| = \mu$  and  $\theta < \text{cf}(\mu)$  one can find  $j \in \theta$  and  $B \in [B_0]^\mu$  such that  $\beta \in B \Rightarrow d(\beta) = j$ . It follows that  $c(\alpha, \beta) = j$  for every  $\alpha \in A, \beta \in B$ , so we are done. □<sub>2.1</sub>

The above lemma will be used within the proof of the main result of the paper, which reads as follows.

**Theorem 2.2.** *Assume that  $\mu > \text{cf}(\mu) = \kappa \geq \aleph_0$ ,  $\mu$  is a strong limit cardinal and  $2^\mu > \mu^+$ . Then  $\binom{\mu^+}{\mu} \rightarrow \binom{\tau}{\mu}_{< \text{cf}(\mu)}$  for every  $\tau < \mu^+$ .*

*Proof.*

Fix two sequences of cardinals,  $\bar{\mu} = \langle \mu_i : i < \kappa \rangle$  and  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ , with the following properties:

- (a)  $\bar{\mu}$  is increasing and continuous,  $\kappa < \mu_0$  and  $\mu = \bigcup_{i < \kappa} \mu_i$ .
- (b)  $\bar{\lambda}$  is an increasing sequence of regular cardinals,  $\mu = \bigcup_{i < \kappa} \lambda_i$ .
- (c) For every  $i < \kappa$ ,  $2^{\mu_i^+} < \lambda_i$ .
- (d)  $\text{tcf}(\prod_{i < \kappa} \lambda_i, J_\kappa^{\text{bd}}) = \mu^{++}$ .

Part (d) is possible by Theorem 1.1. Part (c) is possible since  $\mu$  is a strong limit cardinal and  $\bar{\lambda}$  can be replaced by any unbounded subsequence while keeping (d). The fact that we use  $J_\kappa^{\text{bd}}$  in the representation of  $\mu^{++}$  at (d) is crucial here.

Fix any  $\theta < \kappa$  and a coloring  $c : \mu^+ \times \mu \rightarrow \theta$ . Let  $\tau < \mu^+$  be an ordinal. By the monotonicity of the arrow notation we may increase  $\tau$  and therefore assume that  $\text{cf}(\tau) = \kappa$  and  $\tau = \bigcup_{i < \kappa} \delta_i$  where each  $\delta_i$  is indecomposable and  $\delta_i > \bigcup_{j < i} \delta_j$  for every  $i < \kappa$ . We also choose another sequence of indecomposable ordinals  $\langle \nu_i : i \in \kappa \rangle$  so that  $\delta_i < \nu_i < \delta_{i+1}$  for every  $i \in \kappa$ . These assumptions are possible since  $S_\kappa^{\mu^+}$  is stationary and the set of indecomposable ordinals below  $\mu^+$  is a club.

Let  $\chi$  be a sufficiently large regular cardinal in which the arguments of our proof will be carried out. Let  $\mathcal{M} = \langle M_{\alpha i} : \alpha < \mu^+, i < \kappa \rangle$  be a pcf



array of submodels of  $\mathcal{H}(\chi)$ , based on the ideal  $J_\kappa^{\text{bd}}$  and the sequences  $\bar{\mu}, \bar{\lambda}$  and let  $\langle f_\alpha : \alpha < \mu^+ \rangle$  be its characteristic sequence. We may assume that  ${}^\kappa\theta \subseteq M_{\alpha i}$  for every  $\alpha \in \mu^+, i \in \kappa$ . Since  $\text{tcf}(\prod_{i < \kappa} \lambda_i, J_\kappa^{\text{bd}}) = \mu^{++}$ , we can fix a function  $f^* \in \prod_{i < \kappa} \lambda_i$  such that  $\alpha \in \mu^+ \Rightarrow f_\alpha <_{J_\kappa^{\text{bd}}} f^*$ .

We shall use  $f^*$  as a *translation* of the coloring  $c$ , defined on pairs, to a function defined on singletons in the following sense: for every  $\alpha \in \mu^+$  define  $g_\alpha \in {}^\kappa\theta$  by letting  $g_\alpha(i) = c(\alpha, f^*(i))$ . Here we still depend on both coordinates, but since  $\theta^\kappa < \mu$  there is a fixed  $g^* \in {}^\kappa\theta$  such that the set  $H = \{\alpha \in \mu^+ : g_\alpha = g^*\}$  is of size  $\mu^+$ . Now  $c(\alpha, f^*(i))$  depends only on the righthand coordinate when we restrict ourselves to  $H$  in the lefthand coordinate: if  $\alpha_1, \alpha_2 \in H$  then  $c(\alpha_1, f^*(i)) = g_{\alpha_1}(i) = g^*(i) = g_{\alpha_2}(i) = c(\alpha_2, f^*(i))$ .

The role of  $\mathcal{M}$  and  $f^*$  is to create the small component of the monochromatic product. For the large component we choose a continuous increasing sequence  $\langle N_\zeta : \zeta \in \mu^+ \rangle$  of submodels of  $\mathcal{H}(\chi)$  so that for every  $\zeta \in \mu^+$  the following requirements are met:

- ( $\aleph$ )  $\mu \subseteq N_\zeta, |N_\zeta| = \mu$ .
- ( $\beth$ )  $c, H, g^*, \mathcal{M} \in N_\zeta$ .
- ( $\beth$ )  $N_\zeta \in N_{\zeta+1}$ .

Notice that the set  $E = \{\zeta \in \mu^+ : N_\zeta \cap \mu^+ = \zeta\}$  is a club subset of  $\mu^+$ . It would be helpful to concentrate on members of  $E$ . Since  $E$  is a club we may assume, without loss of generality, that every  $\delta_i$  belongs to  $\text{acc}(E)$ . For every  $i < \kappa$  we choose an ordinal  $\alpha(i) \in \delta_i$  such that  $\alpha(i) > \bigcup\{\nu_j : j < i\}$ . Also, pick  $\alpha(*) \in H - \tau$ .

The monochromatic product will be created now by induction. For every  $i < \kappa$  we build sets  $A_i, B_i$  which will approximate the large and the small components of the product respectively. Likewise, we choose an ordinal  $j(i) \in \kappa$  and we try to keep the following requirements:

- (a)  $j(i) > i$ ; also,  $i_0 < i_1 \Rightarrow \lambda_{j(i_0)} < \mu_{j(i_1)}$ .
- (b) If  $\sigma_0, \sigma_1 \in \{\delta_\ell : \ell \leq i\} \cup \{\alpha(\ell) : \ell \leq i\} \cup \{\alpha(*)\}$  and  $\sigma_0 < \sigma_1$  then  $f_{\sigma_0}(j) < f_{\sigma_1}(j)$  for every  $j \in [j(i), \kappa)$ .
- (c)  $A_i \subseteq H \cap \delta_i, \text{otp}(A_i) = \delta_i$  and  $A_i \in M_{\nu_i j(i)}$ .
- (d)  $B_i \subseteq \lambda_{j(i)} - \bigcup\{\lambda_{j(\ell)} : \ell < i\}, \text{otp}(B_i) = \lambda_{j(i)}$  and  $B_\ell \in M_{\nu_i j(i)}$  for every  $\ell < i$ .
- (e) If  $\alpha \in \bigcup\{A_\ell : \ell \leq i\} \cup \{\alpha(*)\}$  and  $\beta \in B_\ell$  for some  $\ell \leq i$  then  $c(\alpha, \beta) = g^*(j(\ell))$ .

Suppose that we can carry the induction. Define  $A = \bigcup_{i < \kappa} A_i \cup \{\alpha(*)\}$ ,  $B = \bigcup_{i < \kappa} B_i$ . Notice that  $|B| = \bigcup\{\lambda_{j(i)} : i < \kappa\} = \mu$  and  $\text{otp}(A) = \tau + 1$ . Choose any pair of ordinals  $\alpha \in A, \beta \in B$ . Let  $i < \kappa$  be the first ordinal so that  $\beta < \lambda_{j(i)}$  and  $\alpha \in \bigcup\{A_\ell : \ell \leq i\} \cup \{\alpha(*)\}$ . Requirement (d) implies that the  $B_\ell$ s are mutually disjoint. By the disjointness of the  $B_\ell$ s there is a unique  $\ell \leq i$  such that  $\beta \in B_\ell$ , and then  $c(\alpha, \beta) = g^*(j(\ell))$ . This means that  $c(\alpha, \beta)$  depends only on the righthand coordinate (as  $A \subseteq H$ ), so by Lemma 2.1 we are done.

It remains to carry out the induction. Suppose that  $i < \kappa$ , and assume that  $A_\ell, B_\ell, j(\ell)$  have been defined for every  $\ell < i$ . We shall choose  $j(i)$  and build  $A_i$  together, and then we shall describe the construction of  $B_i$ . For every  $\ell < i$  we pick an ordinal  $\eta_\ell \in \kappa$  such that  $A_\ell, B_\ell, j(\ell) \in M_{\alpha(i)\eta_\ell}$ . Notice that for every  $j \in [\eta_\ell, \kappa)$  we will have  $A_\ell, B_\ell, j(\ell) \in M_{\alpha(i)j}$  by the properties of the array  $\mathcal{M}$ . The choice of  $\eta_\ell$  is possible since  $\nu_\ell < \alpha(i)$  and hence there exists  $\eta_\ell \in \kappa$  such that  $M_{\nu_\ell j} \prec M_{\alpha(i)j}$  for every  $j \in [\eta_\ell, \kappa)$ . Now  $A_\ell, B_\ell, j(\ell) \in M_{\nu_\ell j(\ell)}$  by the induction hypothesis, so they belong to  $M_{\nu_\ell j}$  for every  $j \geq j(\ell)$ . It follows that for every sufficiently large  $j < \kappa$  we have  $A_\ell, B_\ell, j(\ell) \in M_{\alpha(i)j}$ . The above argument works separately for every  $\ell < i$ , and we can choose now  $j_0 < \kappa$  such that  $A_\ell, B_\ell, j(\ell) \in M_{\alpha(i)j_0}$  simultaneously for every  $\ell < i$ . This is possible since  $i < \kappa$ .

We choose an increasing sequence  $\langle \zeta_\varepsilon : \varepsilon < \text{cf}(\delta_i) \rangle$  of elements of  $E$  such that  $\zeta_0 > \alpha(i)$  and  $\delta_i = \bigcup \{ \zeta_\varepsilon : \varepsilon < \text{cf}(\delta_i) \}$ . Notice that  $M_{\alpha(i)j_0} \prec N_{\zeta_\varepsilon}$  for every  $\varepsilon < \text{cf}(\delta_i)$ : fixing  $\varepsilon < \text{cf}(\delta_i)$ , we first note that  $M_{\alpha(i)j_0} \in N_{\zeta_\varepsilon}$ , using that both ordinals  $\alpha(i)$  and  $j_0$  belong to  $N_{\zeta_\varepsilon}$ , as  $\mathcal{M} \in N_{\zeta_\varepsilon}$  and  $j_0 \in \kappa \subseteq N_{\zeta_\varepsilon}$ . Now, since  $|M_{\alpha(i)j_0}| < \lambda_{j_0}$  (by the definition of a pcf array), we have in particular  $|M_{\alpha(i)j_0}| < \mu$ ; since  $\mu \subseteq N_{\zeta_\varepsilon}$ ,  $M_{\alpha(i)j_0} \subseteq N_{\zeta_\varepsilon}$ . Finally, as both  $M_{\alpha(i)j_0}$  and  $N_{\zeta_\varepsilon}$  are elementary submodels of  $\mathcal{H}(\chi)$  we may conclude that  $M_{\alpha(i)j_0} \prec N_{\zeta_\varepsilon}$ .

Fix any  $\varepsilon < \text{cf}(\delta_i)$ . We define a first order formula  $\varphi(x)$  with parameters ( $x$  is the free variable), as follows:

$$x \in H \wedge x > \zeta_\varepsilon \wedge (\forall \ell < i)(\beta \in B_\ell \rightarrow c(x, \beta) = g^*(j(\ell))).$$

Remark that  $\mathcal{H}(\chi) \models \varphi(\alpha(*))$  using the induction hypothesis. Notice that  $\varphi(x)$  is definable in  $N_{\zeta_{\varepsilon+1}}$ , so by elementarity there is a set  $\Gamma_\varepsilon \subseteq N_{\zeta_{\varepsilon+1}} - N_{\zeta_\varepsilon}$  such that  $\text{otp}(\Gamma_\varepsilon) = \zeta_{\varepsilon+1}$  and  $\gamma \in \Gamma_\varepsilon \Rightarrow \mathcal{H}(\chi) \models \varphi(\gamma)$ . Let  $A_i = \bigcup \{ \Gamma_\varepsilon : \varepsilon < \text{cf}(\delta_i) \}$ . Observe that  $\text{otp}(A_i) = \delta_i$ .

We can choose now  $j(i) \in \kappa$  by setting the following requirements. First we find  $j_1 \in \kappa$  such that  $A_i \subseteq M_{\nu_i j_1}$  and then we choose  $j(i) \geq \max\{j_0, j_1, i\}$ . Now we increase  $j(i)$  if needed, by considering a few more requirements. We make sure that  $A_i \in M_{\nu_i j(i)}$ . This can be done since  $\delta_i < \nu_i$  and hence every subset of  $\delta_i$  is bounded in  $\bigcup_{j \in \kappa} M_{\nu_i j}$ . In particular,  $A_i \subseteq \delta_i$ , so for some  $j_1 \in \kappa$  we see that  $A_i \subseteq M_{\nu_i j_1} \subseteq M_{\nu_i j(i)}$  and by increasing the above chosen  $j(i)$  we arrive at a sufficiently closed submodel so that inclusion implies membership. If still needed we increase  $j(i)$  once more to get  $M_{\nu_i j(i)} \prec M_{\alpha(*)j(i)}$  and  $f_{\delta_i}(j) < f^*(j)$  for every  $j \geq j(i)$ .

Finally, we define the set  $B_i$ . Let  $\Gamma = \{ \beta \in \lambda_{j(i)} : \forall \alpha \in \bigcup_{\ell \leq i} A_\ell \cup \{ \alpha(*) \}, c(\alpha, \beta) = g^*(j(i)) \}$ . Notice that  $\Gamma$  is definable in  $M_{\alpha(*)j(i)}$ , since all the parameters are in this model. This is true, in particular, for  $g^*$ , since all the functions from  $\kappa$  to  $\theta$  are in every  $M_{\alpha i}$ . The focal point here is that  $f^*(j(i)) \in \Gamma$ . Indeed,  $f^*(j(i)) < \lambda_{j(i)}$  and for every  $\alpha \in H$  we have  $c(\alpha, f^*(j(i))) = g^*(j(i))$ . This is true, in particular, for every  $\alpha \in \bigcup_{\ell \leq i} A_\ell \cup \{ \alpha(*) \}$  as all these ordinals are from  $H$ .

Since  $f^*(j(i)) > f_{\nu_i}(j(i)) = \sup(M_{\nu_i j(i)} \cap \lambda_{j(i)})$  we infer that  $f^*(j(i)) \notin M_{\nu_i j(i)}$ . By elementarity there is an unbounded set of ordinals below  $\lambda_{j(i)}$  in  $M_{\nu_i j(i)}$  which belong to  $\Gamma$ . Truncate the part of this set below  $\bigcup\{\lambda_{j(\ell)} : \ell < i\}$ , and let  $B_i$  be the remainder. One can verify that  $A_i, B_i$  and  $j(i)$  satisfy the requirements of the inductive process, so we are done.

□<sub>2.2</sub>

In this paper we focused on the so-called *balanced* polarized relation, in which the required size of the monochromatic product is identical in both colors. Let us add a few words with regard to the unbalanced relation. The central assumption in our paper, as well as in [13], is  $2^\mu > \mu^+$ . However, if  $(\mu^+)_\mu \rightarrow (\tau)_\theta$  for every  $\tau \in \mu^+$  and one collapses  $2^\mu$  to  $\mu^+$  then this instance of an almost strong relation is preserved by virtue of the completeness of the collapse.

Consider now the parallel unbalanced relation  $(\mu^+)_\mu \rightarrow (\mu^+ \ \tau)_\mu$ . It was shown in [5] that if  $\mu > \text{cf}(\mu) = \omega$  and  $2^\mu = \mu^+$  then  $(\mu^+)_\mu \not\rightarrow (\mu^+ \ \omega_2)_\mu$ , and it was shown in [4] that consistently  $(\mu^+)_\mu \not\rightarrow (\mu^+ \ \omega_1)_\mu$ . For singular cardinals of countable cofinality this negative relation is optimal, since for such cardinals  $(\mu^+)_\mu \rightarrow (\mu^+ \ \tau)_\mu$  for every  $\tau \in \omega_1$ , as proved in [10]. If  $\text{cf}(\mu) > \omega$  then the negative relation is even sharper, as  $(\mu^+)_\mu \not\rightarrow (\mu^+ \ \omega)_\mu$ , a result of Erdős, Hajnal and Rado. Thus, assuming  $2^\mu = \mu^+$  there is a meaningful difference between the balanced and the unbalanced almost strong polarized relations. This is reflected by the fact that at the balanced relation one can obtain  $(\mu^+)_\mu \rightarrow (\tau)_\theta$  for every  $\tau \in \mu^+, \theta \in \text{cf}(\mu)$ , and this is optimal. Indeed,  $2^\mu = \mu^+$  implies the failure of the strong relation, that is  $(\mu^+)_\mu \not\rightarrow (\mu^+)_\mu$ .

We conclude the paper with the following natural problem:

**Question 2.3.** Let  $\mu$  be a strong limit singular cardinal and suppose that  $2^\mu > \mu^+$ . Is it consistent that  $(\mu^+)_\mu \not\rightarrow (\mu^+)_\mu$ ?

## REFERENCES

1. Uri Abraham and Menachem Magidor, *Cardinal arithmetic*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1149–1227. MR 2768693
2. Maxim R. Burke and Menachem Magidor, *Shelah’s pcf theory and its applications*, Ann. Pure Appl. Logic **50** (1990), no. 3, 207–254. MR 1086455
3. P. Erdős, A. Hajnal, and R. Rado, *Partition relations for cardinal numbers*, Acta Math. Acad. Sci. Hungar. **16** (1965), 93–196. MR MR0202613 (34 #2475)
4. Shimon Garti, *Unbalanced polarized relations*, Proc. Amer. Math. Soc. **149** (2021), no. 3, 1281–1287. MR 4211881
5. Shimon Garti, Menachem Magidor, and Saharon Shelah, *Infinite monochromatic paths and a theorem of Erdos-Hajnal-Rado*, Electron. J. Combin. **27** (2020), no. 2, Paper No. 2.8, 11. MR 4245063
6. Shimon Garti and Saharon Shelah, *A strong polarized relation*, J. Symbolic Logic **77** (2012), no. 3, 766–776. MR 2987137
7. ———, *Open and solved problems concerning polarized partition relations*, Fund. Math. **234** (2016), no. 1, 1–14. MR 3509813
8. Moti Gitik and Saharon Shelah, *Applications of pcf for mild large cardinals to elementary embeddings*, Ann. Pure Appl. Logic **164** (2013), no. 9, 855–865. MR 3056300
9. András Hajnal and Jean A. Larson, *Partition relations*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 129–213. MR 2768681
10. Albin L. Jones, *A polarized partition relation for cardinals of countable cofinality*, Proc. Amer. Math. Soc. **136** (2008), no. 4, 1445–1449. MR 2367118
11. Menachem Kojman, *A proof of Shelah’s partition theorem*, Arch. Math. Logic **34** (1995), no. 4, 263–268. MR 1356541
12. Saharon Shelah, *Cardinal arithmetic*, Oxford Logic Guides, vol. 29, The Clarendon Press, Oxford University Press, New York, 1994, Oxford Science Publications. MR 1318912
13. ———, *A polarized partition relation and failure of GCH at singular strong limit*, Fund. Math. **155** (1998), no. 2, 153–160. MR MR1606515 (99b:03064)
14. Neil H. Williams, *Combinatorial set theory*, Studies in Logic and the Foundations of Mathematics, vol. 91, North-Holland Publishing Co., Amsterdam, 1977. MR 3075383

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 91904, ISRAEL

*Email address:* shimon.garty@mail.huji.ac.il

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, BOGOTÁ, COLOMBIA

*Email address:* avillavecesn@gmail.com