

BASIC COMMUTATORS IN n -LIE ALGEBRAS

SEYEDEH NAFISEH AKBAROSSADAT

Abstract. In this paper, we give the structure of free n -Lie algebras. Next, we introduce basic commutators in n -Lie algebras and generalize the Witt formula to calculate the number of the basic commutators. Also, we prove that the set of all of the basic commutators of weight w and length $n + (w - 2)(n - 1)$ is a basis for F^w , where F^w is the w th term of the lower central series in the free n -Lie algebra F .

Keywords: n -Lie algebra, Basic commutators, Free n -Lie algebras, Basic product, String.

MSC: 17B05, 17B10, 17B99.

1. INTRODUCTION AND HISTORY

Lie polynomials appeared at the end of the 19th century and the beginning of the 20th century in the work of Campbell [3], Baker [2], and Hausdorff [4] on exponential mappings in a Lie group, which has led to the so-called Campbell–Baker–Hausdorff formula. Around 1930, Witt introduced the Lie algebra of Lie polynomials. He showed that the Lie algebra of Lie polynomials is actually the free Lie algebra and that its enveloping algebra is the associative algebra of noncommutative polynomials. He proved what is now called the Poincaré–Birkhoff–Witt theorem and showed how a free Lie algebra is related lower central series of the free group. About at the same time, Hall and Magnus, with their commutator calculus, opened the way to the bases of the free Lie algebra. For more details about a historical review of free Lie algebras, we refer the reader to the reference [8] and its references.

The concept of basic commutators is defined in groups and Lie algebras, and there is also a way to construct and identify them. Moreover, a formula for calculating their number is obtained.

In 1962, Shirshov [10] gave a method that generalizes Hall’s method for choosing a basis in a free Lie algebra.

Basic commutators are of particular importance in calculating the dimensions of different spaces and are, therefore, highly regarded. Niroomand and Parvizi [7] investigated some more results about the 2-nilpotent multiplier $\mathcal{M}^{(2)}(L)$ of a finite-dimensional nilpotent Lie algebra L , and by using the Witt formula, they calculated its dimension. Moreover, Salemkar, Edalatzadeh, and Araskhan [9] introduced the concept of c -nilpotent multiplier $\mathcal{M}^{(c)}(L)$ of a finite-dimensional Lie algebra L and obtained some bounds for $\mathcal{M}^{(c)}(L)$ by using the Witt formula and basic commutators.

In 1985, Filippov [5] introduced the concept of n -Lie algebras as an n -ary multilinear and skew-symmetric operation $[x_1, \dots, x_n]$, which satisfies the following generalized Jacobi identity:

$$[[x_1, \dots, x_n], y_2, \dots, y_n] = \sum_{i=1}^n [x_1, \dots, [x_i, y_2, \dots, y_n], \dots, x_n].$$

Clearly, such an algebra becomes an ordinary Lie algebra when $n = 2$.

In this paper, we are going to define basic commutators in n -Lie algebras and give a formula to count the number of them.

2. FREE n -LIE ALGEBRAS

For the first time, Akbarossadat and Saeedi [1] introduced the concept of free n -Lie algebras. In the following subsection, we will review the definition of this concept and some of its properties.

Free n -Lie Algebras. Given a set X , a free n -Lie algebra on X over a field \mathbb{F} is an n -Lie algebra L over \mathbb{F} , together with a mapping $i : X \rightarrow L$, with the following universal property:

For each n -Lie algebra, such as M , and each mapping $f : X \rightarrow M$, there exists a unique n -Lie algebra homomorphism $F : L \rightarrow M$ such that $f = F \circ i$.

A standard argument shows that a free n -Lie algebra on X is necessarily unique, up to n -Lie algebra isomorphism. Also, its existence is shown as follows:

Let X be a set. We construct a Lie algebra generated by X , satisfying no relations other than these:

$$(2.1) \quad [x, x, \dots, x] = 0, \quad \text{for all } x \in X,$$

$$(2.2) \quad \sum_{i=1}^n [x_1, \dots, x_{i-1}, [x_i, x'_1, \dots, x'_{n-1}], x_{i+1}, \dots, x_n] = 0,$$

for all $x_i, x'_j \in X, \quad 1 \leq i \leq n, \quad 1 \leq j \leq n - 1.$

Let $M(X)$ be the set inductively defined as follows:

- (1) $X \subseteq M(X)$.
- (2) If $x_1, \dots, x_n \in M(X)$, then also $(x_1, \dots, x_n) \in M(X)$.

The set $M(X)$ is called the free magma on X . It consists of all bracketed expressions on the elements of X . On $M(X)$ we define an n -ary operation

$$\begin{aligned} M(X) \times \dots \times M(X) &\longrightarrow M(X) \\ x_1 \cdot \dots \cdot x_n &\longmapsto (x_1, \dots, x_n). \end{aligned}$$

For $m \subseteq M(X)$, we define its degree recursively: $deg(m) = 1$ if $m \in X$ and $deg(m) = \sum_{i=1}^n m_i$ if $m = (m_1, \dots, m_n)$. So the degree of an element $m \in M(X)$ is just the number

of elements of X that occur in m (counted with multiplicities). For an integer $d \geq 1$, we set $M_d(X)$ as the subset of $M(X)$ consisting of all $m \in M(X)$ of degree d . Then

$$M(X) = \bigcup_{d \geq 1} M_d(X).$$

Let \mathbb{F} be a field and let $A(X)$ be the vector space over \mathbb{F} spanned by $M(X)$. If we extend the n -ary operation on $M(X)$ n -linearly to $A(X)$, then $A(X)$ becomes an (nonassociative) algebra; it is called the free n -algebra over \mathbb{F} on X . Let $f \in A(X)$; if also $f \in M(X)$, then f is said to be a monomial. For $f \in A(X)$, we define $\deg(f)$ to be the maximum of the degrees of m , where m runs over all $m \in M(X)$ that occur in f with nonzero coefficient.

Let I_0 be the ideal of $A(X)$ generated by all elements

$$\begin{aligned} & (m, m, \dots, m), \quad \text{for all } m \in M(X), \\ & (m_1, \dots, m_i, \dots, m_j, \dots, m_n) + (m_1, \dots, m_j, \dots, m_i, \dots, m_n) \\ & \quad \text{for all } m_i \in M(X), \text{ for all } 1 \leq i, j \leq n \\ & \sum_{i=1}^n (m_1, \dots, m_{i-1}, (m_i, m'_1, \dots, m'_{n-1}), m_{i+1}, \dots, m_n), \quad \text{for all } m_i, m'_i \in M(X). \end{aligned}$$

Set $L(X) = A(X)/I_0$. Let \mathcal{B} be a basis of $L(X)$ consisting of (images of) elements of $M(X)$. Then it is immediate that we have

$$(m, m, \dots, m) = 0,$$

$$(m_1, \dots, m_i, \dots, m_j, \dots, m_n) + (m_1, \dots, m_j, \dots, m_i, \dots, m_n) = 0,$$

and

$$\sum_{i=1}^n (m_1, \dots, m_{i-1}, (m_i, m'_1, \dots, m'_{n-1}), m_{i+1}, \dots, m_n) = 0,$$

for all $m, m_i, m'_i \in \mathcal{B}$. It is easy to check that the relations (2.1) and (2.2) hold for all elements of $L(X)$, so $L(X)$ is an n -Lie algebra. Therefore we will use a bracket to denote the product in $L(X)$ and say $L(X)$ is the free Lie algebra on X .

In other words, an n -Lie algebra L is free on X if and only if it is generated by all possible n -Lie bracketings of all elements of X . The only existing and possible relations among these bracketing generators are the consequence of n -linearity bracketings, generalized Jacobi identity, and

$$[x_1, x_2, \dots, x_{i-1}, x_i, \dots, x_{j-1}, x_j, \dots, x_n] = -[x_1, x_2, \dots, x_{i-1}, x_j, \dots, x_{j-1}, x_i, \dots, x_n],$$

for all $1 \leq i, j \leq n$.

The following lemma is a main tool for the proof of main theorem of this section.

Lemma 2.1. *Let F be a free n -Lie algebra and let $i, j \in \mathbb{N}$. Then $[F^i, F^j, \underbrace{F, \dots, F}_{n\text{-times}}] \subseteq F^{i+j}$.*

Proof. We prove this by induction on i and j . For each $i \geq 1$ and $j = 1$, by the definition of lower central series terms, we have

$$[F^i, F^j, F, \dots, F] = [F^i, F^1, F, \dots, F] = [F^i, F, F, \dots, F] = F^{i+1}.$$

Now, let i be a fixed and arbitrary natural number and let the statement be true for $j - 1$; that is,

$$[F^i, F^{j-1}, F, \dots, F] \subseteq F^{i+1} \quad (\text{induction hypothesis}).$$

By the definition of lower central series terms F^i 's, we know that $F^i = [F^{i-1}, F, F, \dots, F]$. Hence every element of F^i such an element a is $a = [[a_0, f'_2, \dots, f'_n], f_2, \dots, f_n]$, where $a_0 \in F^{i-1}$ and $f_s, f'_r \in F$, for each $2 \leq r, s \leq n$. Now, let $x = [a, b, f_2, \dots, f_n]$ be an arbitrary element of $[F^i, F^j, \underbrace{F, \dots, F}_{n\text{-times}}]$, where $a = [a_0, f'_2, \dots, f'_n] \in F^i$, $b \in F^j$, and

$f'_s \in F$, for all $2 \leq s \leq n$. By the Jacobi identity, we have

$$\begin{aligned} x &= [a, b, f_2, \dots, f_n] = [[a_0, f'_2, \dots, f'_n], b, f_2, \dots, f_n] \\ &= [[a_0, b, f_2, \dots, f_n], f'_2, \dots, f'_n] + [a_0, [f'_2, b, f_2, \dots, f_n], f'_3, \dots, f'_n] \\ &\quad + \dots + [a_0, f'_2, \dots, f'_{n-1}, [f'_n, b, f_2, \dots, f_n]] \\ &= - \underbrace{[[b, a_0, f_2, \dots, f_n], f'_2, \dots, f'_n]}_{\in F^{j+i-1}} - \underbrace{[[f'_2, b, f_2, \dots, f_n], a_0, f'_3, \dots, f'_n]}_{\in F^{j+i-1}} \\ &\quad + \dots + \underbrace{[a_0, f'_2, \dots, f'_{n-1}, [f'_n, b, f_2, \dots, f_n]]}_{\in F^{j+i-1}} \\ &= \sum_{s=2}^n [a_0, f'_2, \dots, [f'_s, b, f_2, \dots, f_n], f'_{s+1}, \dots, f'_n] \in F^{i+j}. \end{aligned}$$

Thus $[F^i, F^j, \underbrace{F, \dots, F}_{n\text{-times}}] \subseteq F^{i+j}$. □

An ideal $I \trianglelefteq L$ is called abelian, if $[I, I, L, \dots, L] = 0$. Using this fact, we prove the following theorem, which is an important result.

Theorem 2.2. *Let F be a free n -Lie algebra and let F^i be the i th term of the lower central series of F , for each $i \in \mathbb{N}$. Then $\frac{F^i}{F^{i+c}}$ is abelian, where $c = 0, 1, \dots, i$.*

Proof. It is obvious for $c = 0$, since $\frac{F^i}{F^i} = 0_{\frac{F^i}{F^i}} = F^i$. Let $c = 1$. Since $[F^i, F^i, F, \dots, F] \subseteq [F^i, F, F, \dots, F] = F^{i+1}$, so $Z\left(\frac{F^i}{F^{i+1}}\right) = \frac{F^i}{F^{i+1}}$, and hence $\frac{F^i}{F^{i+1}}$ is abelian.

Now suppose that $0 \leq c \leq i$. We show that $\frac{F^i}{F^{i+c}}$ is abelian. In this regard, it is enough to show that $\frac{F^i}{F^{i+c}}$ is an abelian ideal of $\frac{F}{F^{i+c}}$, which is equivalent to

$$[F^i, F^i, F, \dots, F] + F^{i+c} = 0_{\frac{F^i}{F^{i+c}}} = F^{i+c}, \quad \text{or} \quad [F^i, F^i, F, \dots, F] \subseteq F^{i+c}.$$

By Lemma 2.1, we know that $[F^i, F^i, F, \dots, F] \subseteq F^{2i}$. Since $0 \leq c \leq i$, so $c + i \leq 2i$, and thus $F^{2i} \subseteq F^{i+c}$. Therefore, $[F^i, F^i, F, \dots, F] \subseteq F^{i+c}$. \square

3. AN INTRODUCTION TO BASIC COMMUTATORS

In this section, our aim is to define basic commutators of n -Lie algebras. For this, we need to introduce some concepts.

Strings and collected parts. Let X be a given set of noncommuting variables (also called letters). We consider formal words or strings $b_1 b_2 \dots b_n$, where each b_i is one of the letters $x_1, x_2, \dots, x_r, \dots$. In other words, the word is a finite sequence of elements of X or, equivalently, a noncommutative monomial. We include the empty word denoted 1. We denote by X^* the set of all words on X . With the product defined by the concatenation of words, X^* becomes a monoid, which is the free monoid on X . Indeed, for any mapping $f : X \rightarrow M$, where M is any monoid, there is a unique homomorphism of monoids $F : X^* \rightarrow M$ such that $f = F \circ i$, where i is the natural injection $X \rightarrow X^*$.

A noncommutative polynomial is a linear combination of words with coefficients in the field \mathbb{F} . The set of all these noncommutative polynomials becomes an algebra over \mathbb{F} with the product inherited from the free monoid X^* , which is denoted by $\mathbb{F}\langle X \rangle$ and is the associative algebra X . Indeed, for any mapping $f : X \rightarrow A$, where A is any \mathbb{F} -algebra, there is a unique homomorphism of algebras as $F : \mathbb{F}\langle X \rangle \rightarrow A$, such that $f = F \circ i$, where i is the natural injection from X into $\mathbb{F}\langle X \rangle$.

Since $\mathbb{F}\langle X \rangle$ is an algebra, so from the product, we can define a bracket on it and turn it into a Lie algebra. For this, let P and Q be two polynomials in $\mathbb{F}\langle X \rangle$. Define $[-, -] : \mathbb{F}\langle X \rangle \times \mathbb{F}\langle X \rangle \rightarrow \mathbb{F}\langle X \rangle$ by $[P, Q] := PQ - QP$. It is easy to check the bilinear and skew-symmetric properties and Jacobi identity. Hence this defines a structure of Lie algebra on $\mathbb{F}\langle X \rangle$.

Now, let $\mathcal{B}_{\mathbb{F}}(X)$ denote the smallest Lie subalgebra of $\mathbb{F}\langle X \rangle$ that contains each letter in X . It is easy to prove that $\mathcal{B}_{\mathbb{F}}(X)$ is a free Lie algebra. In what follows, we show that every element of the Lie subalgebra $\mathcal{B}_{\mathbb{F}}(X)$ is a Lie polynomial and that the set of all basic commutators on X is a basis for $\mathcal{B}_{\mathbb{F}}(X)$. Hence the dimension of it is the number of all basic commutators on X . Indeed, for this purpose, we first need to introduce the concept of basic commutators and some of their properties. In what follows, we discuss this topic.

Now, we introduce formal commutators c_i and weights $\omega(c_i)$ by the rules:

- 1) $c_i = x_i$, $i = 1, \dots, r$, are the commutators of weight 1; that is, $\omega(x_i) = 1$.

2) If c_{i_1}, \dots, c_{i_n} are commutators, then $c_k = (c_{i_1}, \dots, c_{i_n})$ is a commutator and $\omega(c_k) = \sum_{j=1}^n \omega(c_{i_j})$.

Note that these definitions yield only a finite number of commutators of any given weight. We shall order the commutators by their subscripts, counting $c_i = x_i, i = 1, \dots, r$, and listing in order of weight, but giving an arbitrary ordering to commutators of the same weight.

A string $c_{i_1} \dots c_{i_r}$ of commutators is said to be in collected form if $i_1 \leq \dots \leq i_r$, that is, if the commutators are in order read from left to right. An arbitrary string of commutators $c_{i_1} \dots c_{i_r} c_{i_{r+1}} \dots c_{i_s}$ will, in general, has a collected part $c_{i_1} \dots c_{i_r}$ if $i_1 \leq \dots \leq i_r$ and if $i_r \leq i_j, j = r + 1, \dots, s$. Moreover, this sting will have an uncollected part $c_{i_{r+1}} \dots c_{i_s}$, where i_{r+1} is not the least of $i_j, j = r + 1, \dots, s$. The collected part of a string $c_{i_1} \dots c_{i_s}$ will be void unless i_1 is the least of the subscripts.

Collecting process in strings. In this part, we define a collecting process for strings of commutators. If c_u is the earliest commutator in the uncollected part and if $c_{i_j} = c_u$ is the leftmost uncollected c_u , then we replace $c_{i_1} \dots c_{i_m} \dots c_{i_{j-1}} c_{i_j} \dots c_{i_n}$ with $-c_{i_1} \dots c_{i_m} \dots c_{i_j} c_{i_{j-1}} \dots c_{i_n}$. This has the effect of moving c_{i_j} to the left as the formal coefficient of (-1) . Thus c_{i_j} is still the earliest commutator in the uncollected part. Alter enough steps, c_{i_j} will be moved to the $(m + 1)$ st position and will become a part of the collection part. Note that in every step of moving c_{i_j} to the left, the coefficient of (-1) is created that can be abridged.

Basic commutators and basic product. Suppose that sequences of basic commutators c_1, c_2, \dots formed from the generators x_1, x_2, \dots, x_d are given. We call a product of basic commutators $c_{i_1} c_{i_2} \dots c_{i_s}$, a *basic product* if it is in collected order; that is, $i_1 \leq i_2 \leq \dots \leq i_s$. The number of all basic products of weight 2 and length n formed from generators x_1, x_2, \dots, x_d , is d^n .

In applying the collecting process to a positive word, not all commutators will arise. Thus $(x_n, \dots, x_3, x_2, x_1)$ may arise but not $(x_n, \dots, x_1, x_2, x_3)$, since x_3 is collected before x_1 . The commutators that may actually arise are called *basic commutators*. We give a formal definition of the basic commutators for n -Lie algebras generated by x_1, \dots, x_d .

Brief introduction of basic commutators in n -Lie algebras. In the previous sections, we completely and precisely define the basic commutators by using the words. Indeed in this section, we are going to briefly introduce the basic commutators for n -Lie algebra L (especially) with the free representation $\frac{F}{R}$ over the field \mathbb{F} , in the bracket form. In other words, we introduce a method to determine the basic commutators in n -Lie algebras. Note that two properties of n -Lie algebras, as “skew-symmetric” and “Jacobian identity”, are important and necessary to determine the basic commutators.

In general, it can be said that the set of all basic commutators A is the smallest subset of the set of all commutators of L , named W , such that A can be produced W .

The basic commutators are characterized by two concepts as “*weight*” and “*length*”. In the case when $n = 2$, these two concepts are the same. The basic commutator $0 \neq c \in F$ has weight w_c , if $c \in F^{w_c}$ and $c \notin F^{w_c+1}$, where F^{w_c} is w_c th terms of lower central

series. Also, the length of c is the number of its components that are in X , and we denote it by m_c and calculate it as follows:

$$m_c = n + (w_c - 2)(n - 1) = n + w_c n - 2n - w_c + 2 = (w_c - 1)n - (w_c - 2).$$

In the general case, weight and length are denoted by w and m , respectively. Thus, the general form of basic commutators with different weights and lengths is shown in Table 1.

TABLE 1. Generally form of basic commutators

Domain	weight(w)	length (m)	generally form
X	1	1	x_i
F^2	2	n	$[x_{i_1}, \dots, x_{i_n}]$
F^3	3	$2n - 1$	$[[x_{i_1}, \dots, x_{i_n}], x_{j_2}, \dots, x_{j_n}]$
\vdots	\vdots	\vdots	\vdots

With the above discussion, we have Table 2.

TABLE 2. The length of the basic commutators of weight w in n -Lie algebras

w	1	2	3	4	5	6	7	8	...
$n = 2$	1	2	3	4	5	6	7	8	...
$n = 3$	1	3	5	7	9	11	13	15	...
$n = 4$	1	4	7	10	13	16	19	22	...
$n = 5$	1	5	9	13	17	21	25	29	...
$n = 6$	1	6	11	16	21	26	31	36	...
$n = 7$	1	7	13	19	25	31	37	43	...
$n = 8$	1	8	15	22	29	36	43	50	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots

If in any n -Lie algebras (for each $n \in \mathbb{N}$), we denote the number of length of the basic commutator of weight w_0 by $m_{w_0}^n$, then we can obtain the following results:

- $m_{w_0}^n = m_n^{w_0}$, $n \geq 2$, and $w_0 \geq 2$.
- $m_{w_0}^3 = m_{w_0}^2 + m_{w_0-1}^2$.

Let $X = \{x_1, x_2, \dots, x_n, \dots\}$ be a basis set of n -Lie algebra L . We define the relation " $<$ " on X as $x_1 < x_2 < \dots < x_n < \dots$. In what follows, we define the basic commutators in L as follows:

- (1) Every basic element $c_i^1 = x_i$ in X is the basic commutator of weight 1.
- (2) The element $c_s^2 = [x_{i_1}, x_{i_2}, \dots, x_{i_n}]$ (where $x_{i_j} \in X$, for all $j = 1, \dots, n$) is a basic commutator of length n and weight 2. It belongs to F^2 , if $x_{i_1} > x_{i_2} > \dots > x_{i_n}$; in other words, $i_1 > i_2 > \dots > i_n$. We can compare every two basic commutators c_s^2

and c_k^2 with weight 2 and length n . Suppose that $c_s^2 = [x_{i_1}, x_{i_2}, \dots, x_{i_n}]$ and that $c_k^2 = [x'_{j_1}, x'_{j_2}, \dots, x'_{j_n}]$, and let r_0 be the first index (in moving from right to left) of x_{i_r} 's and x'_{j_r} 's which $x_{i_{r_0}} \neq x'_{j_{r_0}}$. If $x_{i_{r_0}} > x'_{j_{r_0}}$ or, equivalently, $i_{r_0} > j_{r_0}$, then we say $c_s^2 > c_k^2$, and otherwise, we say $c_s^2 < c_k^2$.

(3) Let $c_t^w = [c_{j_1}^{w_1}, \dots, c_{j_n}^{w_n}]$, where all of $c_{j_1}^{w_1}, \dots, c_{j_n}^{w_n}$ are basic commutators of weights w_1, \dots, w_n and lengths $n + (w_1 - 2)(n - 1), \dots, n + (w_n - 2)(n - 1)$, respectively. Then c_t^w is called a basic commutator if the following conditions hold:

- $w > w_1 \geq w_2 \geq \dots \geq w_n$.
- Whenever $w_s = w_{s+1}$, then $c_{j_s}^{w_s} > c_{j_{s+1}}^{w_{s+1}}$, equivalently, $j_s > j_{s+1}$.
- Whenever $w_s > w_{s+1}$ and $c_{j_s}^{w_s} = [c_{j_{s_1}}^{w'_1}, \dots, c_{j_{s_n}}^{w'_n}]$, then $c_{j_{s_n}}^{w'_n} \leq c_{j_n}^{w_n} < \dots < c_{j_{s+2}}^{w_{s+2}} < c_{j_{s+1}}^{w_{s+1}}$.

In this case, c_t^w is an basic element of F^w of weight w and length $n + (w - 2)(n - 1)$.

Note that every basic commutator of weight w is smaller than every basic commutator of weight larger than w . Hence $c_i^1 < c_j^2 < c_k^3 < \dots$, for all indices i, j, k, \dots .

Collecting process in commutators of n -Lie algebras. It is obvious that there exist many commutators in generated free n -Lie algebra F over X such that they does not have a basic form. Indeed it is possible to choose an equivalent basic form for them and characterize them by basic commutators. For this, we have to make the necessary changes to them by the properties of n -Lie algebras.

Since F is an n -Lie algebra, so every bracket with at least two same components is equal to zero. In fact, there is no nonzero bracket such that it has at least two same components in F , and all of the brackets in F have n different components.

Suppose that $c = [c_{j_1}^{w_1}, c_{j_2}^{w_2}, \dots, c_{j_r}^{w_r}, c_{j_{r+1}}^{w_{r+1}}, \dots, c_{j_n}^{w_n}]$ is a nonbasic commutator in F . Then there are the following two cases:

(A) Let $c_{j_r}^{w_r} < c_{j_{r+1}}^{w_{r+1}}$. Then it is enough to write it as follows:

$$c = [c_{j_1}^{w_1}, c_{j_2}^{w_2}, \dots, c_{j_r}^{w_r}, c_{j_{r+1}}^{w_{r+1}}, \dots, c_{j_n}^{w_n}] = -[c_{j_1}^{w_1}, c_{j_2}^{w_2}, \dots, c_{j_{r+1}}^{w_{r+1}}, c_{j_r}^{w_r}, \dots, c_{j_n}^{w_n}].$$

(B) Let $c_{j_1}^{w_1} = [c_{r_1}^{w'_1}, c_{r_2}^{w'_2}, \dots, c_{r_n}^{w'_n}]$ be in the basic form. Then c reads as follows:

$$c = [c_{j_1}^{w_1}, c_{j_2}^{w_2}, \dots, c_{j_n}^{w_n}] = [[c_{r_1}^{w'_1}, c_{r_2}^{w'_2}, \dots, c_{r_n}^{w'_n}], c_{j_2}^{w_2}, \dots, c_{j_n}^{w_n}].$$

We know that c is a basic commutator if

$$c_{r_n}^{w'_n} \leq c_{j_n}^{w_n} < c_{j_{n-1}}^{w_{n-1}} < \dots < c_{j_3}^{w_3} < c_{j_2}^{w_2}.$$

Otherwise, c is not in a basic form. Thus we replace c by using the Jacobi identity and case (A).

Now, we state the theorems that are among the main results and our goals in this paper.

Theorem 3.1. *Let F be a free n -Lie algebra with the ordered basis set $X = \{x_1, x_2, x_3, \dots\}$, where $x_1 < x_2 < x_3 < \dots$. The set of all defined basic commutators of weight w (denoted by $B(w)$) on X is linearly independent.*

Proof. It is obvious by the collecting process in commutators of n -Lie algebras. Since F is the generated free n -Lie algebra on X , then it has no other relations except n -linearity, skew-symmetric properties, and Jacobi identity. Therefore, all of the brackets in F apply to these three properties. On the other hand, in the process of determining and identifying the basic commutators, we remove the commutators that are linearly dependent under these three properties (n -linearity, skew-symmetric properties, and Jacobi identity), and we separate only its linearly independent elements. Hence, all of the basic commutators of arbitrary and fixed weight w are linearly independent. \square

The following theorem is the main result of this section, which is similarly discussed in group theory and Lie algebras.

Theorem 3.2. *Let F be a free n -Lie algebra. Then $\frac{F^i}{F^{i+c}}$ is abelian n -Lie algebras whose basis is the set of all basic commutators of weights $i, i + 1, i + 2, i + 3, \dots, i + c - 1$ and lengths $n + (i - 2)(n - 1), n + (i - 1)(n - 1), \dots, n + (i + c - 3)(n - 1)$.*

Proof. This theorem is proved similar to the Lie algebra case. \square

4. COUNTING OF BASIC COMMUTATORS


It is necessary to note that counting of basic commutators on every arbitrary basis set X is not possible, in general. In fact, although it is possible to define the concept of basic commutators for any countable set, counting their number is only possible if the set X is finite. The following formula, known as the *Witt formula*, is provided in group theory and Lie algebras for counting of all of the basic commutators:

$$(4.3) \quad l_d(w) = \frac{1}{w} \sum_{r|w} \mu(r) d^{\frac{w}{r}},$$

where d is the number of generators of given group/Lie algebra (the number of members of X), μ is the *Möbius function*, which is defined for positive integers by the rule $\mu(1) = 1$, and for $w = p_1^{a_1} p_2^{a_2} \dots p_s^{a_s}$ that p_1, p_2, \dots, p_s are distinct primes numbers, by $\mu(p_1 p_2 \dots p_s) = (-1)^s$ and $\mu(p_1^{a_1} p_2^{a_2} \dots p_s^{a_s}) = 0$, if any $a_i > 1$.

In group theory, the strategy of the proof of the above formula is to remove nonbasic commutators from all the commutators. That is, it first counts all possible commutators on X and then removes nonbasic members and their periods. Indeed in n -Lie algebras, with this method, only a bound can be found for the number of basic commutators.

Now, let $X = \{x_1, x_2, \dots, x_d\}$ with the ordered relation $x_d > x_{d-1} > \dots > x_2 > x_1$. We are going to determine the number of all of basic commutators of weight w and length $m = n + (w - 2)(n - 1)$. Now, consider the following string of circles:



We want to put x_i 's in each circle such that the final result is a basic commutator. First, we consider all of the cases for putting them. It is easy to see that the number of these

cases is d^m (every circle has d choices). Obviously, all of them are not basic, and hence we remove nonbasic cases. For this, we remove duplicate and linear dependent commutators by using two properties, skew-symmetric and Jacobi identity of n -Lie algebras. This method is similar to the method used in the group theory (see [6]), and also, Shirshov [10] provided it for Lie algebras. We know that

$$m = n + (w - 2)(n - 1) = n + wn - 2n - w + 2 = (w - 1)n - (w - 2),$$

and so n is a divisor of m if $w \equiv 2 \pmod{n}$; in other words, $w = qn + 2$, for some integer positive q . Thus

$$m = (w - 1)n - (qn + 2 - 2) = (w - 1)n + qn = (w + q - 1)n,$$

and hence $\frac{m}{n} = w + q - 1$ is integer. Therefore, in general, $n \nmid m$, but $n - 1 \mid m - 1$, because

$$m = n + (w - 2)(n - 1) = n + wn - 2n - w + 2,$$

and so

$$\begin{aligned} m - 1 &= n + wn - 2n - w + 2 - 1 = \underline{n - 2n + 1} + \underline{wn - w} \\ &= -(n - 1) + w(n - 1) \\ &= (w - 1)(n - 1). \end{aligned}$$

We call a word $a_1 a_2 \dots a_s$ circular if a_1 is regarded as a_s , and also, $a_1 a_2 \dots a_s, a_2 a_3 \dots a_s a_1, a_3 \dots a_{n-1} a_n a_2, \dots, a_n a_1 a_2 \dots a_{n-1}$ are regarded as the same word. A circular word C of weight w and length $m = n + (w - 2)(n - 1)$ may conceivably be given by repeating a segment of q letters m/q times, where q is some divisor of m . We say that C is of period q if we have

$$\underbrace{\underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}} \underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}} \dots \underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}}}_{m\text{-circles and } m/q\text{-parts}}.$$

So by the above, circular commutators must be removed from all of the generated commutators on X . Also, in weights that smaller than w , we have

$$\underbrace{\underbrace{(\otimes \otimes \otimes \dots \otimes)}_{i\text{-times}} \underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}} \underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}} \dots \underbrace{(\bigcirc \bigcirc \dots \bigcirc)}_{q\text{-times}}}_{(m-i)\text{-circles and } (m-i)/q\text{-parts}}$$

$$(4.4) \quad l_d^n(w) \leq \frac{1}{m} \left(\sum_{r|m} \mu(r) d^{\frac{m}{r}} \right).$$

Note that the equality holds if and only if $n = 2$.

In the next section, we introduce the recursive formula.

5. NEW METHOD OF COUNTING OF BASIC COMMUTATORS

In the previous section, we provide the generalized Witt formula for counting basic commutators in n -Lie algebras with finite basis set X . Indeed, it could not discuss the relationship between the numbers of basic commutators of different weights. Now, in this section, we introduce some formulas in special cases with the new method and show their relationship.

First, note the following theorem in which we give the number of all of the basic commutators of weight 2, in general n -Lie algebras with d generators.

Theorem 5.1. *Let L be an n -Lie algebra of dimension d . Then*

$$l_d^n(2) = \binom{d}{n}.$$

Proof. Let L be an n -Lie algebra of dimension d with the basis $X = \{x_1, x_2, \dots, x_d\}$ and put $w = 2$. We are going to count the number of all basic commutators of weight 2 on d letters. It is clear that the number of all commutators of weight 2 is equal to $d^n = n^n$. By the skew-symmetric property in n -Lie algebras, we have

$$[x_{i_1}, x_{i_2}, \dots, x_{i_j}, \dots, x_{i_k}, \dots, x_{i_n}] = -[x_{i_1}, x_{i_2}, \dots, x_{i_k}, \dots, x_{i_j}, \dots, x_{i_n}],$$

and hence $[x_{i_1}, x_{i_2}, \dots, x_{i_{j-1}}, x, x_{i_{j+1}}, \dots, x_{i_{k-1}}, x, x_{i_{k+1}}, \dots, x_{i_n}] = 0$. Thus just independent and nonzero commutators of weight 2 are as $[x_{i_1}, x_{i_2}, \dots, x_{i_n}]$, where all of x_{i_j} 's are distinct. So $l_d^n(2) = \binom{d}{n}$. \square

Remark. Since every n -Lie algebra has the skew-symmetric property, it is not possible to exist at least two same components. Hence the number of basic commutators of weights greater than or equal to 2 is equal to zero, if $d < n$.

Corollary 5.2. *If $n = d$, then the following properties hold:*

- (1) $l_n^n(1) = n$.
- (2) $l_n^n(2) = \binom{n}{n} = 1$.
- (3) $l_n^n(3) = \binom{n}{n-1} = n$.
- (4) $l_n^n(4) = \binom{n}{2} + n$.
- (5) $l_n^n(5) = \binom{n}{3} + 2\binom{n}{2} + n$.
- (6) $l_n^n(6) = \binom{n}{4} + 3\binom{n}{3} + 3\binom{n}{2} + n$.
- (7) $l_n^n(7) = \binom{n}{5} + 4\binom{n}{4} + 6\binom{n}{3} + 4\binom{n}{2} + n$.
- (8) $l_n^n(8) = \binom{n}{6} + 5\binom{n}{5} + 10\binom{n}{4} + 10\binom{n}{3} + 5\binom{n}{2} + n$.
- (9) $l_n^n(9) = \binom{n}{7} + 6\binom{n}{6} + 15\binom{n}{5} + 20\binom{n}{4} + 15\binom{n}{3} + 6\binom{n}{2} + n$.
- (10) $l_n^n(10) = \binom{n}{8} + 7\binom{n}{7} + 21\binom{n}{6} + 35\binom{n}{5} + 35\binom{n}{4} + 21\binom{n}{3} + 7\binom{n}{2} + n$.

Proof. Let L be an n -Lie algebra of dimension $d = n$ with the basis $X = \{x_1, x_2, \dots, x_n\}$ such that $x_n > x_{n-1} > \dots > x_2 > x_1$.

- (1). It is clear from the definition of basic commutators of weight 1.

(2). By Theorem 5.1, we have

$$l_d^n(2) = l_n^n(2) = \binom{n}{n} = \frac{n!}{(n-n)! \times n!} = 1.$$

(3). We know that the general form of the basic commutators of weight 3 is as follows:

$$[[x_{11}, \dots, x_{1n}], x_{22}, \dots, x_{2n}] \in L^3,$$

where $x_{ij} \in X$, for all $1 \leq j \leq n$ and $i = 1, 2$. By part (2), we know that at $n = d$, there is only one basic commutator of weight 2 as $[x_n, x_{n-1}, \dots, x_2, x_1]$. Therefore, it is enough to determine only the status of the outer components of the second bracket. Since $n = d$, so $(n-1)$ -components x_{22}, \dots, x_{2n} are equal to $(n-1)$ -components of inner bracket. Without reducing the totality, assume that $x_{1j} = x_{2j}$, for all $2 \leq j \leq n$. Hence in this situation, according to the Jacobi identity and skew-symmetric property of n -Lie algebras, we have

$$\begin{aligned} & [[x_{11}, \dots, x_{1n}], x_{22}, \dots, x_{2n}] \\ &= [[x_{11}, \dots, x_{1n}], x_{12}, \dots, x_{1n}] \\ &= [[x_{11}, \dots, x_{1n}], x_{12}, \dots, x_{1n}] + [x_{11}, \cancel{[x_{12}, x_{12} \dots, x_{1n}]} \rightarrow 0, x_{13}, \dots, x_{1n}] \\ &\quad + [x_{11}, x_{12}, \cancel{[x_{13}, x_{12} \dots, x_{1n}]} \rightarrow 0, x_{14}, \dots, x_{1n}] + \dots + [x_{11}, x_{12}, \dots, \cancel{[x_{1n}, x_{12} \dots, x_{1n}]} \rightarrow 0] \\ &= [[x_{11}, \dots, x_{1n}], x_{12}, \dots, x_{1n}]. \end{aligned}$$

Thus, to count the number of all basic commutators of weight 3 on d letters (which $d = n$), it is enough to calculate the number of all possible modes to select $(n-1)$ distinct letters from n letters (in other words, it is equal to the number of all basic commutators of weight 2 on d letters in $(n-1)$ -Lie algebras), which according to Theorem 5.1, is equal to

$$l_n^d(3) = l_n^n(3) = \binom{n}{n-1} = n.$$

So the number of all basic commutators of weight $w = 3$, where $n = d$, is equal to n .

(4). We know that the general form of basic commutators of weight 4 can be considered as $[[[x_{11}, x_{12}, \dots, x_{1n}], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}] \in L^4$. Note that x_{22}, \dots, x_{2n} and x_{32}, \dots, x_{3n} can be considered as strings such as basic commutators of weight 2 on n letters and with length $n-1$. In this case, according to the definition of the basic commutators, $(x_{12}, \dots, x_{1n}) \leq (x_{22}, \dots, x_{2n}) \leq (x_{32}, \dots, x_{3n})$ should be hold. Given that at $n = d$, there is only one basic commutator of weight 2 as $[x_1, x_2, \dots, x_n]$, it suffices to consider the possible choices for (x_{22}, \dots, x_{2n}) and (x_{32}, \dots, x_{3n}) . Obviously, for each $2 \leq j \leq n$, x_{2j} and x_{3j} are equal to one of the $[x_1, x_2, \dots, x_n]$ components, and (x_2, \dots, x_n) is smaller than or equal to (x_{22}, \dots, x_{2n}) and (x_{32}, \dots, x_{3n}) . At $n = d$, the general form of the base shifters of weight 4 can be rewritten as follows:

$$[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}] \in L^4, \quad x_{2j}, x_{3j} \neq x_1, \text{ for all } 2 \leq j \leq n.$$

According to the above description, it can be divided into the following two categories:

$$(5.5) \quad \begin{aligned} & [[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{22}, \dots, x_{2n}], (x_{22}, \dots, x_{2n}) = (x_{32}, \dots, x_{3n}) \\ & \implies \text{the number of all basic commutators in this form is equal to } n. \end{aligned}$$

$$(5.6) \quad \begin{aligned} & [[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}], (x_{22}, \dots, x_{2n}) \neq (x_{32}, \dots, x_{3n}), \\ & \implies \text{the number of all basic commutators in this form is equal to } \binom{n}{2}. \end{aligned}$$

Note that the number of all basic commutator of weight 2 on $d = n$ letters in $(n - 1)$ -Lie algebras is equal to n . Therefore, the number of all basic commutators of weight 4 on $d = n$ letters is equal to the sum of the above two cases (5.5) and (5.6); that is,

$$l_n^n(4) = \binom{l_n^{n-1}(2)}{2} + n = \binom{\binom{n}{2}}{2} + n = \binom{n}{2} + n.$$

(5). Similar to part (4), we have the following categories:

$$[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{22}, \dots, x_{2n}], x_{22}, \dots, x_{2n}],$$

where $(x_{22}, \dots, x_{2n}) = (x_{i2}, \dots, x_{in})$, $i = 3, 4$,

(5.7)

\implies the number of all basic commutators in this form is equal to n .

$$[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}] \quad \text{and}$$

$$[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}], x_{32}, \dots, x_{3n}],$$

where $(x_{22}, \dots, x_{2n}) \neq (x_{32}, \dots, x_{3n})$,

(5.8)

\implies the number of all basic commutators in this form is equal to $2 \times \binom{n}{2}$.

$$[[[x_1, x_2, \dots, x_n], x_{22}, \dots, x_{2n}], x_{32}, \dots, x_{3n}], x_{42}, \dots, x_{4n}],$$

where $(x_{i2}, \dots, x_{in}) \neq (x_{j2}, \dots, x_{jn})$, $i, j = 2, 3, 4$,

(5.9)

\implies the number of all basic commutators in this form is equal to $\binom{n}{3}$.

Hence, $l_n^n(5)$ is equal to the sum of the three cases (5.7), (5.8), and (5.9) as follows:

$$\begin{aligned} l_n^n(5) &= \binom{l_n^{n-1}(2)}{3} + 2\binom{l_n^{n-1}(2)}{2} + n \\ &= \binom{\binom{n}{n-1}}{2} + 2\binom{\binom{n}{n-1}}{2} + n \\ &= \binom{n}{3} + 2\binom{n}{2} + n. \end{aligned}$$

By a similar process to the previous one, it can be proved that

$$l_n^n(6) = \binom{\binom{n}{n-1}}{4} + 3\binom{\binom{n}{n-1}}{3} + 3\binom{\binom{n}{n-1}}{2} + n = \binom{n}{4} + 3\binom{n}{3} + 3\binom{n}{2} + n,$$

$$\begin{aligned} l_n^n(7) &= \binom{\binom{n}{n-1}}{5} + 4\binom{\binom{n}{n-1}}{4} + 6\binom{\binom{n}{n-1}}{3} + 4\binom{\binom{n}{n-1}}{2} + n \\ &= \binom{n}{5} + 4\binom{n}{4} + 6\binom{n}{3} + 4\binom{n}{2} + n, \end{aligned}$$

$$\begin{aligned} l_n^n(8) &= \binom{\binom{n}{n-1}}{6} + 5\binom{\binom{n}{n-1}}{5} + 10\binom{\binom{n}{n-1}}{4} + 10\binom{\binom{n}{n-1}}{3} + 5\binom{\binom{n}{n-1}}{2} + n \\ &= \binom{n}{6} + 5\binom{n}{5} + 10\binom{n}{4} + 10\binom{n}{3} + 5\binom{n}{2} + n, \end{aligned}$$

$$\begin{aligned} l_n^n(9) &= \binom{\binom{n}{n-1}}{7} + 6\binom{\binom{n}{n-1}}{6} + 15\binom{\binom{n}{n-1}}{5} + 20\binom{\binom{n}{n-1}}{4} + 15\binom{\binom{n}{n-1}}{3} + 6\binom{\binom{n}{n-1}}{2} + n \\ &= \binom{n}{7} + 6\binom{n}{6} + 15\binom{n}{5} + 20\binom{n}{4} + 15\binom{n}{3} + 6\binom{n}{2} + n, \end{aligned}$$

$$\begin{aligned} l_n^n(10) &= \binom{\binom{n}{n-1}}{8} + 7\binom{\binom{n}{n-1}}{7} + 21\binom{\binom{n}{n-1}}{6} + 35\binom{\binom{n}{n-1}}{5} + 35\binom{\binom{n}{n-1}}{4} + 21\binom{\binom{n}{n-1}}{3} \\ &\quad + 7\binom{\binom{n}{n-1}}{2} + n \\ &= \binom{n}{8} + 7\binom{n}{7} + 21\binom{n}{6} + 35\binom{n}{3} + 35\binom{n}{4} + 21\binom{n}{3} + 7\binom{n}{2} + n. \end{aligned}$$

□

According to Corollary 5.2, we obtain $l_n^n(w) = \sum_{i=1}^{w-2} a_i \binom{n}{i}$, and in Table 3 (for $i \geq 1$), see the order between the coefficients.

TABLE 3. coefficients $\binom{n}{i}$'s

	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8
$l_n^n(4)$	1	1	–	–	–	–	–	–
$l_n^n(5)$	1	2	1	–	–	–	–	–
$l_n^n(6)$	1	3	3	1	–	–	–	–
$l_n^n(7)$	1	4	6	4	1	–	–	–
$l_n^n(8)$	1	5	10	10	5	1	–	–
$l_n^n(9)$	1	6	15	20	15	6	1	–
$l_n^n(10)$	1	7	21	35	35	21	7	1

According to Table 3, we assume that the following relations are established.

- $a_1 = a_{w-2} = 1$, $a_2 = a_{w-3} = w - 3$, and $a_i = a_{w-i-1}$, for each i .
- Comparing the coefficients in Table 3 with the values of the Khayyam triangle, it is observed that to calculate the coefficients a_i in $l_n^n(w)$, it is sufficient to obtain the $(w - 3)$ th power coefficients of the binomial, that is, $(x + y)^{w-3}$. In other words, a_i is the coefficient $x^{w-3-(i-1)}y^{i-1}$. Each a_i , for $1 \leq i \leq w - 2$, can also be considered the i th digit in the answer of $(11)^{w-3}$.

Table 4 denotes the number of basic commutators in different weights with $n = d$, which can be obtained by the formulas in Corollary 5.2 and manual calculations.

TABLE 4. Number of basic commutators in different weights

$n = d$	2	3	4	5	6	7	8	9	10
$w = 1$	2	3	4	5	6	7	8	9	10
$w = 2$	1	1	1	1	1	1	1	1	1
$w = 3$	2	3	4	5	6	7	8	9	10
$w = 4$	3	6	10	15	21	28	36	45	55
$w = 5$	6	10	20	35	56	84	120	165	220
$w = 6$	9	15	35	70	126	210	330	495	715
$w = 7$	18	21	56	126	252	462	792	1287	2002
$w = 8$	30	28	84	210	462	924	1716	3003	5005
$w = 9$	56	36	120	330	792	1716	3432	6435	11440
$w = 10$	99	45	165	495	1287	3003	6435	12870	24310

We obtain the following simple formulas that are easily proved by induction:

- $l_n^n(4) = l_n^n(3) + (l_n^n(3) - 1) + (l_n^n(3) - 2) + \cdots + 2 + 1 = \sum_{i=0}^{l_n^n(3)-1} (l_n^n(3) - i) = \left(l_n^n(3) \times l_n^n(3) \right) + \sum_{i=1}^{l_n^n(3)} i = \left(l_n^n(3) \right)^2 + \frac{l_n^n(3)(l_n^n(3) - 1)}{2} = \left(l_n^n(3) \right)^2 + \binom{l_n^n(3)}{2}$, where $n \geq 2$.
- $l_n^n(w) = \sum_{w'=2}^w l_{n-1}^{n-1}(w') = l_n^n(w-1) + l_{n-1}^{n-1}(w)$, where $n > 3$.
- $l_3^3(w) = l_3^3(w-1) + (w-1)$, where $w \geq 3$, $l_3^3(1) = 3$, and $l_3^3(2) = 1$.
- $l_4^4(w) = \begin{cases} l_4^4(w-1) + (w-1)r, & w = 2r, \\ l_4^4(w-1) + wr, & w = 2r+1. \end{cases}$

We know that the number of all commutators (all of the basic and nonbasic commutators) with weight w is d^{mw} . Hence,

$$\underbrace{\begin{matrix} d & d & & d & d \\ \circ & \circ & \dots & \circ & \circ \end{matrix}}_{n\text{-times}}.$$

Also, it is obvious that the number of all nonbasic commutators of weight w , denoted by $L_d^n(w)$, is equal to

$$(5.10) \quad L_d^n(w) = d^{mw} - l_d^n(w).$$

On the other hand, by Theorem 5.1, we have $l_d^n(2) = \binom{d}{n}$, and so the number of all nonbasic commutators of weight 2 is equal to

$$(5.11) \quad L_d^n(2) = d^{m_2} - l_d^n(2) = d^n - \binom{d}{n}.$$

Now, we calculate the number of all basic commutators of weight 3. Every commutator of weight 3 has $n+(n-1) = 2n-1$ components, in which n of them form the commutators of weight 2, and others are weight 1 and form the exterior part in the bracket. It is obvious that every nonbasic commutator of weight 2 forms a nonbasic commutator of weight 3, independent of the selection of the $n-1$ components in external parts. So these must be removed, too. We denote the number of all of such cases by $L_d^m(3)$, which is equal to

$$(5.12) \quad L_d^m(3) = L_d^n(2) \times d^{n-1}.$$

On the other hand, note that if a basic commutator of weight 2 is arranged in a nonbasic form way with the external part, then the result will be nonbasic again. So we should remove such cases. The number of these, denoted by $L_d^m(3)$, is calculated as follows:

$$(5.13) \quad L_d^m(3) = l_d^n(2) \times L_d^{n-1}(2)$$

$$(5.14) \quad = \binom{d}{n} \times \left(d^{n-1} - \binom{d}{n-1} \right).$$

Thus,

$$(5.15) \quad L_d^n(3) = L_d^m(3) + L_d^{m_n}(3).$$

Moreover, it may be that two parts are basic, but the third condition of the definition of basic commutators is not true. Hence commutators in this situation must be replaced by terms with the basic form, and the number of them must be decreased. In this case, we can replace it by using the Jacobi identity, such that every term of the Jacobi identity has the basic form, or otherwise, repeat this replacing for nonbasic terms in the Jacobi identity. Note that if any term of Jacobi identity does not have a basic form, then we use the Jacobi identity next-handed to replace it (i.e., if the first time, we write the right Jacobi identity, then next time, must be used the left Jacobi identity to replace the other terms). Note that by attention to the collecting process, all of the above terms can be assumed. We denote the number of them with $L_w^*(n, d, m_{w-1}, n-1)$, where $m_w = m_{w-1} + n - 1$. It is important that if $n = d$, then each product of two basic commutators as ab , in which a is a basic commutator of weight $w - 1$ and b is a basic string of length $n - 1$, has a basic form with weight w . Thus

$$(5.16) \quad L_w^*(n, n, m_{w-1}, n-1) = 0.$$

In general,

$$(5.17) \quad L_w^*(n, d, m_{w-1}, n-1) = l_d^n(w-1) \times l_d^{n-1}(2) - \kappa = l_d^n(w-1) \times \binom{d}{n-1} - \kappa.$$

We are going to find κ by induction on w .

Let us first show that the number of all basic commutators of weight w is equal to κ . According to the above explanation, it is clear that the number of all of the nonbasic commutators is equal to the sum of both $L_d^n(w)$ and $L_w^*(n, d, m_{w-1}, n-1)$, and so other commutators are basic. Thus the number of all basic commutators of weight w is

$$\begin{aligned} l_d^n(w) &= d^{m_w} - L_d^n(w) - L_w^*(n, d, m_{w-1}, n-1) \\ &= d^{m_w} - (L_d^m(w) + L_d^{m_n}(w)) - L_w^*(n, d, m_{w-1}, n-1) \\ &= d^{m_w} - (L_d^n(w-1) \times d^{n-1}) - (l_d^n(w-1) \times L_d^{n-1}(2)) - (l_d^n(w-1) \times l_d^{n-1}(2) - \kappa) \\ &= d^{m_w} - ((d^{m_{w-1}} - l_d^n(w-1)) \times d^{n-1}) - (l_d^n(w-1) \times (d^{n-1} - l_d^{n-1}(2))) \\ &\quad - (l_d^n(w-1) \times l_d^{n-1}(2) - \kappa) \\ &= d^{m_w} - \underbrace{(d^{m_{w-1}} \times d^{n-1})}_{d^{m_w}} + (l_d^n(w-1) \times d^{n-1}) - (l_d^n(w-1) \times d^{n-1}) \\ &\quad + (l_d^n(w-1) \times l_d^{n-1}(2)) - (l_d^n(w-1) \times l_d^{n-1}(2)) + \kappa = \kappa. \end{aligned}$$

Now, we calculate κ by induction on w . It is obvious that the smallest value of w must be 3, and so, for $w = 3$, we have $l_d^n(2) = \binom{d}{n}$, $l_d^{n-1}(2) = \binom{d}{n-1}$, and

$$(5.18) \quad L_3^*(n, d, m_2, n-1) = \binom{d}{n} \binom{d}{n-1} - \sum_{i=1}^{d-n+1} \sum_{j=i+1}^{d-1} (d-j) \left[\binom{d-i+1}{n-1} - j + i + 1 \right].$$

Therefore, by (5.11), (5.12), (5.14), and (5.18), we have

$$l_d^n(3) = \kappa = \sum_{i=1}^{d-n+1} \sum_{j=i+1}^{d-1} (d-j) \left[\binom{d-i+1}{n-1} - j + i + 1 \right].$$

A comparison of two equations (5.19) and (5.18) shows that there are no other basic commutators of weight 3 except equation (5.19). Therefore, we obtain a formula to count basic commutators of weight 3 in any given n -Lie algebras with arbitrary dimension $d \geq n$, and hence the following theorem is proved.

Theorem 5.3. *Let F be a free n -Lie algebra over the ordered set $X = \{x_i | x_{i+1} > x_i, i = 1, 2, \dots, d\}$. Then the number of basic commutators of weight 3 is*

$$(5.19) \quad l_d^n(3) = \sum_{i=1}^{d-n+1} \sum_{j=i+1}^{d-1} (d-j) \left[\binom{d-i+1}{n-1} - j + i + 1 \right].$$

By a similar way and induction, we can rewrite (5.12), (5.13), and (5.15) and obtain the following equations:

$$L_d^n(w) = L_d^n(w-1) \times d^{n-1}, \quad L''^n(w) = l_d^n(w-1) \times L_d^{n-1}(2), \quad L_d^n(w) = L_d^n(w) + L''^n(w).$$

Now, in another way, we calculate the number of basic commutators of weights 4 and 5 on $d \geq 4$ letters. Then we generalize it to the state of arbitrary w and obtain the final and general formula for the various arbitrary values of n , d , and w .

Theorem 5.4. *Let F be a free n -Lie algebra over the ordered set $X = \{x_i | x_{i+1} > x_i, i = 1, 2, \dots, d\}$. Then the number of basic commutators of weight 4 is*

$$(5.20) \quad l_d^n(4) = \sum_{j=1}^{\alpha_0} \beta_{j^*} \left(\binom{\binom{d}{n-1}}{2} + \binom{d}{n-1} \right),$$

where $\alpha_0 = \binom{d-1}{n-1}$, and if $\binom{k-1}{n-1} + 1 \leq j \leq \binom{k}{n-1}$ (for $k = n-1, n, n+1, n+2, \dots, d-1$), then $j^* = \binom{k-1}{n-1} + 1$ and $\beta_{j^*} = (d - n - j^* + 2)$.

Proof. Let $d \geq n$, let $X = \{x_i | x_i > x_{i-1}, i = 1, 2, 3, 4, \dots, d\}$, and let $w = 4$. We know that $l_{d-1}^{n-1}(2) = \binom{d-1}{n-1}$ is the number of collected strings of length $n-1$, too. We calculate the collected strings of length $n-1$ and arrange them as follows. Then we number them from right to left (We use the j counter to count them. Obviously, $j = 1, 2, \dots, l_{d-1}^{n-1}(2)$).

Note that here we display $n = 3$ just to make it easier to display the components, but the same procedure applies to the general case n . Hence

$$(5.21) \quad \begin{array}{cccccccccc} x_{d-1}x_{d-2} & x_{d-1}x_{d-3} & \dots & x_4x_3 & x_4x_2 & x_4x_1 & x_3x_2 & x_3x_1 & x_2x_1 \\ \downarrow & \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ j = \binom{d-1}{n-1} & j = \binom{d-1}{n-1} - 1 & \dots & j = 6 & j = 5 & j = 4 & j = 3 & j = 2 & j = 1 \end{array}$$

The general form of the basic commutators of weight 4 is as follows:

$$(((a_1, a_2, a_3), b_2, b_3), c_2, c_3), \quad a_r, b_s, c_q \in X.$$

First, we select the first collected string from the chain (5.21) (for $j = 1$) and replace a_2, a_3 with it; see Figure 1. We have to calculate the number of all basic commutators by

FIGURE 1.

$$\left(\left(\left(\square \ 2 \ 1 \right) \begin{array}{c} y_1 \ y_2 \\ \square \ \square \\ \downarrow \\ y \end{array} \right) \begin{array}{c} z_1 \ z_2 \\ \square \ \square \\ \downarrow \\ z \end{array} \right)$$

counting the number of allowed modes for cells such as x, y , and z . Since the corresponding term $j = 1$ in the chain (5.21) is the first and smallest sentence, the allowed states for selecting x are x_3, x_4, \dots, x_{d-1} , and x_d , so the number of these states is equal to $d - (n - 1) = d - 2$.

Now, we want to form a chain of all basic commutators of weight 2 (with length $n - 1$) on d letters $x_1, x_1, \dots, x_d \in X$. We know that $l_{d-1}^{n-1}(2) = \binom{d-1}{n-1}$ is the number of all basic commutators of weight 2 and length $n - 1$. We calculate the basic commutators of weight 2 and then arrange them by the defined relation “ $<$ ” on basic commutators as follows. Then we label them from right to left (We use the counter i , to count them. Obviously, $i = 1, 2, \dots, l_d^{n-1}(2)$). Hence

$$(5.22) \quad \underbrace{x_d x_{d-1}}_{i=\binom{d}{n-1}} > \underbrace{x_d x_{d-2}}_{i=\binom{d}{n-1}-1} > \underbrace{x_{d-1} x_{d-2}}_{i=\binom{d}{n-1}-2} > \dots > \underbrace{x_d x_2}_{i=2d-3} > \dots > \underbrace{x_4 x_2}_{i=d+1} > \underbrace{x_3 x_2}_{i=d} > \underbrace{x_d x_1}_{i=d-1} > \dots > \underbrace{x_3 x_1}_{i=2} > \underbrace{x_2 x_1}_{i=1}.$$

Also, since the corresponding sentence $j = i = 1$ in the chain (5.22) is the smallest sentence, all the sentences in the chain (5.22) can be placed after it. On the other hand, both $z \geq y$ and $y \geq x_2 x_1$ conditions must be met. Obviously, both sentences we choose from the chain (5.22) for y and z , one of which is smaller than the other, which can be considered as y and another as z . Therefore, the number of choice states y and z , provided that y and z are distinct, is equal to $\binom{d}{n-1} = \binom{d}{n-1}$. If $y = z$, then we have $\binom{d}{n-1}$ choices for them. Thus, in this case $j = 1$, the number of basic commutators is equal to $(d - 2) \times \left[\binom{d}{n-1} + \binom{d}{n-1} \right]$.

By the same process, we also calculate the number of basic commutators in the cases $j = 2, 3, 4, \dots, \binom{d-1}{n-1}$, and the sum of the number of basic commutators in the various cases j is equal to the number of all the basic commutators of weight 4 on d letters. Moreover,

$$l_d^n(4) = \sum_{j=1}^{\alpha_0} \beta_{j^*} \left(\binom{\binom{d}{n-1}}{2} + \binom{d}{n-1} \right),$$

where $\alpha_0 = \binom{d-1}{n-1}$, and if $\binom{k-1}{n-1} + 1 \leq j \leq \binom{k}{n-1}$, (for $k = n-1, n, n+1, n+2, \dots, d-1$), then put $j^* = \binom{k-1}{n-1} + 1$ and $\beta_{j^*} = (d - n - j^* + 2)$. Therefore it is clear that $1 \leq j^* \leq d - n + 1$. \square

In the following, we want to get a formula for calculating the number of basic commutators of any weight w in every d -dimensional n -Lie algebras. For this purpose, it is necessary to note that in the case of duplicate selection, the placement of strings can also have several different modes, the number of which must also be calculated. For example, suppose $w = 5$. We have to count the number of allowed choices for x , y , and z . If all three strings x , y , and z are distinct, then they are arranged as follows:

$$\left(\left(\left(\underbrace{a_1, a_2, \dots, a_n}_a, \underbrace{x_2, x_3, \dots, x_n}_x, \underbrace{y_2, y_3, \dots, y_n}_y, \underbrace{z_2, z_3, \dots, z_n}_z \right) \right) \right), \quad z > y > x \geq a.$$

Indeed if two of the strings x , y , and z are the same, then the following two states must be considered and counted:

$$\left(\left(\left(\underbrace{a_1, a_2, \dots, a_n}_a, \underbrace{x_2, x_3, \dots, x_n}_x, \underbrace{x_2, x_3, \dots, x_n}_y, \underbrace{z_2, z_3, \dots, z_n}_z \right) \right) \right), \quad z > y = x \geq a,$$

$$\left(\left(\left(\underbrace{a_1, a_2, \dots, a_n}_a, \underbrace{x_2, x_3, \dots, x_n}_x, \underbrace{y_2, y_3, \dots, y_n}_y, \underbrace{y_2, y_3, \dots, y_n}_z \right) \right) \right), \quad z = y > x \geq a.$$

Also, if $x = y = z$, then

$$\left(\left(\left(\underbrace{a_1, a_2, \dots, a_n}_a, \underbrace{x_2, x_3, \dots, x_n}_x, \underbrace{x_2, x_3, \dots, x_n}_y, \underbrace{x_2, x_3, \dots, x_n}_z \right) \right) \right), \quad z = y = x \geq a.$$

In the weight $w = 4$, there exist two outer strings x and y , and so there were two choice modes for x and y , which are as follows: (1)-A state where both x and y are distinct (Equivalently, x and y are replaced by exactly two distinct strings). (2)-The state that x and y are equal (Equivalently, x and y are replaced by only one string). Also, in the case $w = 5$, there exist three outer strings x , y , and z . Hence, there were three choice modes for x , y , and z , which are as follows: (1)-A state that all three strings x , y , and z are distinct (Equivalently, x , y , and z are replaced by exactly three distinct strings, in which case there is only one arrangement for these three strings as $z > y > x$). (2)-The state that two strings of x , y , and z are equal (Equivalently, x , y , and z are replaced by exactly two distinct strings, in which case there is only two arrangement for these strings as $z = y > x$ and $z > y = x$). (3)-The state where all of the three strings are equal together (Equivalently, x , y , and z are replaced by only one string, in which case there is only one arrangement for strings as $z = y = x$).

In general, for an arbitrary w , the number of outer strings is equal to $w - 2$, called $y_1, y_2, y_3, \dots, y_{w-2}$ (where it can be replaced by basic strings of length $n - 1$, or in other words, by basic commutators of weight 2 in $(n - 1)$ -Lie algebras) on d letters, in which the number of all them is equal to $l_d^{n-1}(2) = \binom{d}{n-1}$. If we want to replace them with $w - 2$ distinct strings (a state in which they are all distinct), then we have only one arrangement as $y_1 < y_2 < y_3 < \dots < y_{w-2}$. If we replace all of them with just one string (a state in which they are all repetitive), then we have one arrangement $y_1 = y_2 = y_3 = \dots = y_{w-2}$. If we replace them with $w - 3$ distinct strings (a state in which exactly two strings are same) or replace them with 2 distinct strings (a state in which some of the y_i are equal to string 1 and others are equal to string 2), then the number of arrangements of them is equal to $\alpha_2 = \alpha_{w-1} = \binom{w-3}{0} = \binom{w-3}{w-3}$.

Continuing this process, we can say that if we want to replace them with i distinct strings or replace them with $w - i - 1$ (a state in which exactly i strings are the same), then we have α_i different arrangements, where $\alpha_i = \alpha_{w-i+1} = \binom{w-3}{i-2} = \binom{w-3}{w-i+1}$ and $2 \leq i \leq w - 1$.

Now, considering the above explanations and the proof of the previous theorem, we are ready to introduce the formula for calculating the number of basic commutators.

Theorem 5.5. *Let F be a free n -Lie algebra over the ordered set $X = \{x_i | x_{i+1} > x_i : i = 1, 2, \dots, d\}$ and let w be a positive integer number. Then the number of basic commutators of weight w is*

$$(5.23) \quad l_d^n(w) = \sum_{j=1}^{\alpha_0} \beta_{j^*} \left(\sum_{i=2}^{w-1} \alpha_i \binom{\binom{d}{n-1}}{w-i} \right),$$

where $\alpha_0 = \binom{d-1}{n-1}$, α_i , ($2 \leq i \leq w - 1$) is the coefficient of the $(i - 2)$ th sentence in Newton's binomial expansion $(a + b)^{w-3}$, and if $\binom{k-1}{n-1} + 1 \leq j \leq \binom{k}{n-1}$, (for $k = n - 1, n, n + 1, n + 2, \dots, d - 1$), then $j^* = \binom{k-1}{n-1} + 1$ and $\beta_{j^*} = (d - n - j^* + 2)$.

Proof. According to the recent explanation, it is proved by a similar process of the proof of Theorem 5.20. \square

6. RELATIONSHIP BETWEEN THE NUMBER OF BASIC COMMUTATORS OF n -LIE ALGEBRAS AND LIE ALGEBRAS

We know that the n -Lie algebras are independent of the selection of the bracket, and the concept of basic commutators is also related to the free n -Lie algebras. In addition, if L is a Lie algebra over the field \mathbb{F} with the bracket $[-, -] : L \times L \rightarrow L$, then for each natural number $n \in \mathbb{N}$, L can be equipped with an n -linear function $[-, \dots, -] : \underbrace{L \times L \times \dots \times L}_{n\text{-times}} \rightarrow L$, such that L forms an n -Lie algebra with the bracket $[-, \dots, -]$

on the field \mathbb{F} in which

$$[-, \dots, -] : \underbrace{L \times L \times \dots \times L}_{n\text{-times}} \longrightarrow L$$

$$(x_1, x_2, \dots, x_n) \longmapsto [x_1, x_2, \dots, x_n] := [[\dots [[x_1, x_2], x_3], \dots]].$$

The above explanation may raise the question of whether there is a connection between $l_d(w)$ and $l_d^n(w)$, which are the number of basic commutators of weight w and on d generators in Lie algebras and n -Lie algebras?

In this section, we intend to answer this question and examine the relationship between them. By using (4.3), we have

$$l_d^2(w) = l_d(w) = \frac{1}{w} \sum_{s|w} \mu(s) d^{\frac{w}{s}},$$

for all $w, d \in \mathbb{N}$.

On the other hand, we know that

$$\binom{d}{2} = \frac{d(d-1)}{2!} = \frac{d^2 - d}{2!} = \frac{1}{2} \sum_{s|2} \mu(s) d^{\frac{d}{s}} = l_d^2(2).$$

Now, let $n = 3$. Then

$$\begin{aligned} \binom{d}{3} &= \frac{d^3 - d}{3!} - \frac{3d^2 - 3d}{3!} \\ &= \frac{d^3 - d}{3 \times 2!} - \frac{d^2 - d}{2!} \\ &= \frac{1}{2!} \left(\frac{1}{3} \sum_{s|3} \mu(s) d^{\frac{d}{s}} \right) - \frac{1}{2} \sum_{s|2} \mu(s) d^{\frac{d}{s}} \\ &= \frac{1}{2!} l_d(3) - l_d(2). \end{aligned}$$

By continuing this process, Table 5 can be obtained for different values of $n \in \mathbb{N}$.

Now, we assume that the following equation is true for each natural number smaller than or equal to n_0 :

$$(6.24) \quad \binom{d}{n} = \sum_{s=2}^n (-1)^{n-s} \left(\frac{a_s}{(s-1)! \times (s+1)(s+2) \dots (n-1)n} \right) l_d(s),$$

where, $a_s > 0$, for all $2 \leq s \leq n$. Then for $n_0 + 1$, we have

$$\binom{d}{n_0 + 1} = \frac{\prod_{i=0}^{d-1} (d-i)}{(d - (n_0 + 1))! \times (n_0 + 1)!}$$

$$\begin{aligned}
 &= \frac{\prod_{i=0}^{n_0} (d-i)}{(n_0+1)!} \\
 &= \frac{(d-n_0) \prod_{i=0}^{n_0-1} (d-i)}{(n_0+1) \times n_0!} \\
 &= \frac{(d-n_0)}{n_0+1} \times \frac{\prod_{i=0}^{n_0-1} (d-i)}{n_0!} \\
 &= \frac{(d-n_0)}{n_0+1} \times \left(\sum_{s=2}^{n_0} (-1)^{n_0-s} \left(\frac{a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0} \right) l_d(s) \right) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} (d-n_0) a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} l_d(s) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} (d-n_0) a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(r) d^{s/r} \right) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} (d-n_0) a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r) d^r \right) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{(d-n_0)}{s} \sum_{r|s} \mu(s/r) d^r \right) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r) (d-n_0) d^r \right) \\
 &= \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r) d^{r+1} \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r)(-n_0)d^r \right) \\
 & = \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r)d^{r+1} \right) \\
 & + \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} (-n_0)a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r)d^r \right) \\
 & = \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} \left(\frac{1}{s} \sum_{r|s} \mu(s/r)d^{r+1} \right) \\
 & - \sum_{s=2}^{n_0} \frac{(-1)^{n_0-s} n_0 a_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} l_d(s) \\
 & = \sum_{s=2}^{n_0+1} \frac{(-1)^{n_0-s+1} b_s}{(s-1)! \times (s+1)(s+2) \dots (n_0-1)n_0(n_0+1)} l_d(s).
 \end{aligned}$$

Thus equation (6.24) holds for each natural number n , and we have

$$(6.25) \quad \binom{d}{n} = \sum_{s=2}^n (-1)^{n-s} \left(\frac{a_s}{(s+1)(s+2) \dots (n-1)n} \right) l_d(s).$$

Moreover,

$$\begin{aligned}
 (6.26) \quad \binom{d}{n} & = \sum_{s=2}^n (-1)^{n-s} \left(\frac{a_s}{3 \times 4 \times \dots \times (s-1)(s+1)(s+2) \dots (n-1)n} \right) l_d(s) \\
 & = \sum_{s=2}^n (-1)^{n-s} \left(\frac{a_s}{\prod_{\substack{r=3 \\ r \neq s}}^n r} \right) l_d(s).
 \end{aligned}$$

	$l_d(2)$	$l_d(3)$	$l_d(4)$	$l_d(5)$	$l_d(6)$	$l_d(7)$	$l_d(8)$	$l_d(9)$	$l_d(10)$
$\binom{d}{2} = \frac{1}{2!} \prod_{i=0}^1 (d-i)$	$\frac{1}{1}$	-	-	-	-	-	-	-	-
$\binom{d}{3} = \frac{1}{3!} \prod_{i=0}^2 (d-i)$	$\frac{-3}{3}$	$\frac{1}{2!}$	-	-	-	-	-	-	-
$\binom{d}{4} = \frac{1}{4!} \prod_{i=0}^3 (d-i)$	$\frac{12}{4 \times 3}$	$\frac{-6}{4 \times 2!}$	$\frac{1}{3!}$	-	-	-	-	-	-
$\binom{d}{5} = \frac{1}{5!} \prod_{i=0}^4 (d-i)$	$\frac{-60}{5 \times 4 \times 3}$	$\frac{35}{5 \times 4 \times 2!}$	$\frac{-10}{5 \times 3!}$	$\frac{1}{4!}$	-	-	-	-	-
$\binom{d}{6} = \frac{1}{6!} \prod_{i=0}^5 (d-i)$	$\frac{360}{6 \times 5 \times 4 \times 3}$	$\frac{-224}{6 \times 5 \times 4 \times 2!}$	$\frac{85}{6 \times 5 \times 3!}$	$\frac{-15}{6 \times 4!}$	$\frac{1}{5!}$	-	-	-	-
$\binom{d}{7} = \frac{1}{7!} \prod_{i=0}^6 (d-i)$	$\frac{-2520}{7 \times 6 \times \dots \times 3}$	$\frac{1603}{7 \times 6 \times 5 \times 4 \times 2!}$	$\frac{-735}{7 \times 6 \times 5 \times 3!}$	$\frac{175}{7 \times 6 \times 4!}$	$\frac{-21}{7 \times 5!}$	$\frac{1}{6!}$	-	-	-
$\binom{d}{8} = \frac{1}{8!} \prod_{i=0}^7 (d-i)$	$\frac{20160}{8 \times 7 \times \dots \times 3}$	$\frac{-12810}{8 \times 7 \times \dots \times 4 \times 2!}$	$\frac{6770}{8 \times 7 \times 6 \times 5 \times 3!}$	$\frac{-1960}{8 \times 7 \times 6 \times 4!}$	$\frac{322}{8 \times 7 \times 5!}$	$\frac{-28}{8 \times 6!}$	$\frac{1}{7!}$	-	-
$\binom{d}{9} = \frac{1}{9!} \prod_{i=0}^8 (d-i)$	$\frac{-181440}{9 \times 8 \times \dots \times 3}$	$\frac{113589}{9 \times 8 \times \dots \times 4 \times 2!}$	$\frac{-67320}{9 \times 8 \times \dots \times 5 \times 3!}$	$\frac{22449}{9 \times 8 \times 7 \times 6 \times 4!}$	$\frac{-4536}{9 \times 8 \times 7 \times 5!}$	$\frac{546}{9 \times 8 \times 6!}$	$\frac{-36}{9 \times 7!}$	$\frac{1}{8!}$	-
$\binom{d}{10} = \frac{1}{10!} \prod_{i=0}^9 (d-i)$	$\frac{1814400}{10 \times 9 \times \dots \times 3}$	$\frac{-1109472}{10 \times 9 \times \dots \times 4 \times 2!}$	$\frac{724550}{10 \times 9 \times \dots \times 5 \times 3!}$	$\frac{-269324}{10 \times 9 \times \dots \times 6 \times 4!}$	$\frac{63273}{10 \times 9 \times 8 \times 7 \times 5!}$	$\frac{-9450}{10 \times 9 \times 8 \times 6!}$	$\frac{870}{10 \times 9 \times 7!}$	$\frac{-45}{10 \times 8!}$	$\frac{1}{9!}$

TABLE 5. The connection between $l_d^2(w)$ and $l_d^n(2) = \binom{d}{n}$

According to the above description, the following theorem is proved easily.

Theorem 6.1. *The number of basic commutators of weight 2 on d generators in n -Lie algebras can be concluded by the number of basic commutators of weights $w \leq n$ in Lie algebras.*

Proof. It is easy to check from equation (6.25) and Theorem 5.1. □

The following corollary expresses the relationship between the number of basic commutators in Lie algebras and n -Lie algebras. The following formula is Witt's generalized formula.

Corollary 6.2. *The number of basic commutators of arbitrary weight w on d generators in n -Lie algebras, can be concluded by the number of basic commutators in Lie algebra. We have*

$$l_d^n(w) = \sum_{j=1}^{\alpha_0} \beta_{j^*} \left(\sum_{i=2}^{w-1} \alpha_i \left(\sum_{s=2}^{w-i} (-1)^{w-i-s} \left(\frac{a_s}{\prod_{\substack{r=3 \\ r \neq s}}^{w-i}} r \right) l_{d^*}^*(s) \right) \right),$$

where $d^* = \binom{d}{n-1}$, $\alpha_0 = \binom{d-1}{n-1}$, α_i ($2 \leq i \leq w-1$) is the coefficient of the $(i-2)$ th sentence in Newton's binomial expansion $(a+b)^{w-3}$. If $\binom{k-1}{n-1} + 1 \leq j \leq \binom{k}{n-1}$, (for $k = n-1, n, n+1, n+2, \dots, d-1$), then $j^* = \binom{k-1}{n-1} + 1$ and $\beta_{j^*} = (d-n-j^*+2)$.

Proof. It is proved from equation (5.23) in Theorem 5.5 and Theorem 6.1. □

The following theorem is the most important and practical result of this paper.

Theorem 6.3. *Let F be a free n -Lie algebra and let F^i be the i th term of the lower central series of F , for each $i \in \mathbb{N}$. Then $\frac{F^i}{F^{i+c}}$ is abelian of dimension $l_d^n(i) + l_d^n(i+1) + \dots + l_d^n(i+c-1)$, where $c = 0, 1, 2, \dots, i$.*

Proof. It is proved similar to its version in the theory of Lie algebras. □

REFERENCES

- [1] S.N. Akbarossadat and F. Saeedi, On the dimension of the Schur multiplier of n -Lie algebras, *Linear Multilinear Algebra*, **68**(7), (2020), 1465–1499.
- [2] H.F. Baker, Alternants and continuous groups, *Proc. London Math. Soc.*, **3**, (1905), 24–47.
- [3] J.E. Campbell, On a law of combination of operators bearing of the theory of continuous transformation groups, *Proc. London Math. Soc.*, **28**, (1897), 381–390.
- [4] F. Hausdroff, Die symbolische Exponential formel in der Gruppen theorie, *Leipziger Ber.*, **58**, (1906), 19–48.
- [5] V.T. Filippov, n -Lie algebras, *Sib. Math. Zh.*, **26**(6), (1985), 126–140.
- [6] Marshall Hall, Jr., *The Theory of Groups*, The Macmillan Company, New York, 1959.

- [7] Niroomand, P. and Parvizi, M., 2-capability and 2-nilpotent multiplier of finite dimensional nilpotent Lie algebras, *J. Geom. Phys.*, **121**, (2017), 180–185.
- [8] Reutenauer, Ch., *Free Lie algebras*, Oxford University Press, New York(US), 1993.
- [9] Salemkar, A., Edalatzadeh, B., and Araskhan, M., Some inequalities for the dimension of the c -nilpotent multiplier of Lie algebras, *J. Algebra*, **322**, (2009), 1575–1585.
- [10] Shirshov, A.I., On the bases of a free Lie algebra, *Algebra Log.*, **1**(1), (1962), 14–19.

Seyedeh Nafiseh Akbarossadat

Department of Mathematics, Mashhad Branch, Islamic Azad University, Mashhad, Iran.

E-mail address: n.akbarossadat@gmail.com