

PASCAL DETERMINANTAL ARRAYS AND A GENERALIZATION OF RAHIMPOUR'S DETERMINANTAL IDENTITY

H. TEIMOORI AND H. KHODAKARAMI

ABSTRACT. In this paper, we shall first introduce the Pascal determinantal array of order k as a generalization of the standard Pascal array. We also present an algorithm to produce determinantal arrays of any order. Then we investigate geometric properties of these new determinantal arrays. As a by product, we give a proof of the correctness of our proposed algorithm. Then we give a geometric interpretation of the determinant of any k by k subarray of the Pascal array. Finally, we will give a generalization of Rahimpour's determinantal identity, using the above geometric interpretation.

1. INTRODUCTION

In [1], the first author and M. Bayat proved a conjecture by Rahimpour about determinants inside the Pascal array. After more investigations around the previous project, the second author of this paper made a generalization of the Rahimpour's determinantal identity. Here, we first introduce a new infinite class of squared arrays which are called the Pascal determinantal arrays as a generalization of the standard Pascal array and then we recall the Rahimpour's identity. We then state it's generalization. In section 3, we present an algorithm to produce a Pascal determinantal array of order k from the given Pascal array. In section 4, we quickly review Dodgson's condensation formula for determinants. We also give another recursive formula for Pascal determinantal arrays based on this formula. The weighted version of the star of David rule will be the subject of section 5. We also show that this rule is true for the class of Pascal determinantal arrays. In section 6, we give a proof of the correctness of our algorithm, using the weighted version of the star of David rule and the condensation formula for determinants. Finally in the last section, after giving a geometric interpretation of the entries of the Pascal determinantal array of order k , we present a simple geometric proof of our main result.

2. THE CLASS OF PASCAL DETERMINANTAL ARRAY

Starting from the Pascal infinite array, $P = (P_{i,j} = \binom{i+j}{i})_{i,j \geq 0}$, we can define a new class of infinite arrays which we call them *Pascal determinantal* arrays of order k ($k \geq 1$) denoted by $PD_k = (P_{i,j}^{(k)})_{i,j \geq 0}$, where $P_{i,j}^{(k)}$ is the

Key words and phrases. Determinantal Array, Pascal Array, Dodgson's Condensation, Star of David.

determinant of a k by k sub-array of the Pascal array, starting from it's (i, j) -entry, as follows

1	1	1	1	1	1	1	1
1	2	3	4	5	6	7	8
1	3	6	10	15	21	28	36
1	4	10	20	35	56	84	120
1	5	15	35	70	126	210	330
1	6	21	56	126	252	462	792
1	7	28	84	210	462	924	1716
1	8	36	120	330	792	1716	3432

FIGURE 1. The Pascal array.

$$P_{i,j}^{(k)} := \begin{vmatrix} P_{i,j} & \cdots & P_{i,j+k-1} \\ \vdots & \ddots & \vdots \\ P_{i+k-1,j} & \cdots & P_{i+k-1,j+k-1} \end{vmatrix}.$$

Clearly, the determinantal array of order 1 is exactly the Pascal array (see Figure 1). In Figure 2, we see a Pascal determinantal array of order 2. This is a well-known array which is the squared-form of the so-called Narayana triangle (see A001263 in [5]).

2.1. The Rahimpour Determinantal Identity. Consider the Pascal infinite array, $PD_1 = \binom{i+j}{i}$, $i, j \geq 0$, in Figure 1. Now consider the sub-arrays, shown by squares in Figure 1, of which the left edges lie in the first column (starting from zero) and the other edges are free to lie in any rows or columns of this array of numbers.

She conjectured that the determinants of this square sub-arrays are always equal to the top right entry of that square (see Figure 2). More precisely, we have the following result.

Theorem 1 (See [1]). *Let $P = [P_{i,j} = \binom{i+j}{i}]_{i,j \geq 0}$ be the Pascal infinite array. Then, we have the following determinantal identity*

$$(1) \quad P_{i,j}^{(1)} = P_{i,1}^{(j)}, \quad (i, j \geq 0).$$

The following question naturally arises that, if we let the left edges of square sub-arrays are lie in the k -th column ($k \geq 1$) what will be the extension of the above conjecture?

While working on Rahimpour's determinantal identity, the second author of this paper came up with the following conjecture.

1	1	1	1	1	1	1
1	3	6	10	15	21	28
1	6	20	50	105	196	336
1	10	50	175	490	1176	2520
1	15	105	490	1764	5292	13860
1	21	196	1176	5292	19404	60984
1	28	336	2520	13860	60984	226512

FIGURE 2. Pascal determinantal array of order 2.

Conjecture 2. The determinant of the $n \times n$ square sub-array in Pascal array where the left edge lie in the k -th column is equal to the determinant of the top right k by k square sub-array (see Figure 3). More precisely, we have

$$(2) \quad P_{i,j}^{(k)} = P_{i,k}^{(j)}, \quad (i, j \geq 0, k \geq 1).$$

k	0	1	2	3	4	5	6	7
n	1	1	1	1	1	1	1	1
	1	2	3	4	5	6	7	8
	1	3	6	10	15	21	28	36
	1	4	10	20	35	56	84	120
	1	5	15	35	70	126	210	330
	1	6	21	56	126	252	462	792
	1	7	28	84	210	462	924	1716
	1	8	36	120	330	792	1716	3432

FIGURE 3. The Pascal array and the second author’s conjecture

The main result of our paper is to give a proof of Conjecture 2. A proof of Theorem 1 in [1] is based on the linear algebra properties of Pascal functional matrices [2] and the well-known Cramer’s rule for the inverse of a nonsingular matrix.

Here we will use a completely different approach. We will present an *geometric proof* which relies heavily on the geometric properties of the Pascal array [3], besides using the Dodgson's condensation formula for determinants [4].

3. A RECURSIVE ALGORITHM TO PRODUCE PASCAL DETERMINANTAL ARRAYS

In this section, we give a simple recursive algorithm to produce Pascal determinantal arrays from the standard Pascal array. The algorithm works, as follows

1. Set $PD_1 :=$ The Pascal Array.
 2. Remove the zeroth row and the column of PD_{k-1} and rename the remaining infinite array $RD_k = (R_{i,j}^{(k)})_{i,j \geq 0}$.

3. Set $QD_{i,j}^{(k)} := \frac{R_{i,j}^{(k)}}{R_{0,i+j}^{(k)}}$, $i, j = 0, 1, 2, \dots$

4. Put $P_{i,j}^{(k)} := Q_{i,j}^{(k)} \cdot P_{i,j}^{(1)}$, $i, j = 0, 1, 2, \dots$

Now, $PD_k = (P_{i,j}^{(k)})_{i,j \geq 0}$, is the desired Pascal determinantal array of order k .

Alternatively, we can give the following recursive definition for the entries of the Pascal determinantal array $PD_k = [P_{i,j}^{(k)}]_{i,j \geq 0}$.

$$(1) P_{i,j}^{(1)} = \binom{i+j}{i},$$

$$(2) P_{i,j}^{(k+1)} := \frac{P_{i+1,j+1}^{(k)} P_{i,j}^{(1)}}{P_{1,i+j+1}^{(k)}}, \quad (k \geq 1).$$

Example 3. *The following steps show how to produce PD_3 .*

Step1.

3	6	10	15	21
6	20	50	105	196
10	50	175	490	1176
15	105	490	1764	5292
21	196	1176	5292	19404

Step2.

1	1	1	1	1
1	2	10/3	5	7
1	10/3	25/3	35/2	98/3
1	5	35/2	49	588/5
1	7	98/3	588/5	1764/5

=

1/1	1/1	1/1	1/1	1/1
1/1	4/2	10/3	20/4	35/5
1/1	10/3	50/6	175/10	490/15
1/1	20/4	175/10	9800/20	4116/35
1/1	35/5	490/15	4116/35	24696/70

Step3.

1	1	1	1	1
1	4	10	20	35
1	10	50	175	490
1	20	175	980	4116
1	35	490	4116	24696

Of course, we need to show the correctness of the above algorithm. To do this, we first investigate some algebraic and geometric properties of these determinantal arrays.

4. DODGSON'S CONDENSATION OF DETERMINANTS

Condensation of determinants is a method of computing the determinant of a square matrix due to Charles Dodgson (1866) [4]. For a matrix A of order k , Dodgson's condensation formula states that:

$$A_{k-2}(2,2)A_k(1,1) = A_{k-1}(1,1)A_{k-1}(2,2) - A_{k-1}(1,2)A_{k-1}(2,1), \quad (k \geq 3)$$

where $A_r(i, j)$ denote the r by r minor consisting of r contiguous rows and columns of A , beginning with row i and column j . Note that $A_k(1,1) = \det(A)$, $A_{k-2}(2,2)$ is the central minor; $A_{k-1}(1,1)$, $A_{k-1}(2,2)$, $A_{k-1}(1,2)$, $A_{k-1}(2,1)$ are the northwest, southeast, northeast and southwest minors, respectively. For example, for the matrix

$$A = \begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \\ d_1 & d_2 & d_3 & d_4 \end{bmatrix},$$

we get

$$\begin{aligned} A_2(2,2) &= \begin{bmatrix} b_2 & b_3 \\ c_2 & c_3 \end{bmatrix}, & A_3(1,1) &= \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} \\ A_3(2,2) &= \begin{bmatrix} b_2 & b_3 & b_4 \\ c_2 & c_3 & c_4 \\ d_2 & d_3 & d_4 \end{bmatrix}, & A_3(1,2) &= \begin{bmatrix} a_2 & a_3 & a_4 \\ b_2 & b_3 & b_4 \\ c_2 & c_3 & c_4 \end{bmatrix}, & A_3(2,1) &= \begin{bmatrix} b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \\ d_1 & d_2 & d_3 \end{bmatrix}. \end{aligned}$$

In other words, every k by k minor can be computed in terms of $k-1$ by $k-1$ and $k-2$ by $k-2$ minors. As an immediate consequence this formula, we have the following recurrence relation between the entries of Pascal determinantal arrays of order k ($k \geq 1$):

$$(3) \quad P_{i,j}^{(k)} := \frac{P_{i+1,j+1}^{(k-1)} \cdot P_{i,j}^{(k-1)} - P_{i+1,j}^{(k-1)} \cdot P_{i,j+1}^{(k-1)}}{P_{i+1,j+1}^{(k-2)}}, \quad (i, j = 0, 1, 2, \dots),$$

with the convention that the initial values are $PD_0 := J$ and $PD_1 := P$ in which J and P are all-ones matrix and the Pascal array, respectively.

5. THE WEIGHTED VERSION OF STAR OF DAVID RULE

Consider the Pascal array (see Figure 4). We can draw an arbitrary rectangle whose vertices are entries of the Pascal array. We identify a vertex (the circled vertex) as the anchor of this rectangle. Now we define a weight for this rectangle denoted by W , as follows





FIGURE 4. A rectangle with the Pascal array entries

$$W := \frac{P_{i+m,j+l} \cdot P_{i,j}}{P_{i+m,j} \cdot P_{i,j+l}}, \quad m, l \geq 1, \quad i, j \geq 0.$$

Hilton and Pedersen proved [4] that when we move the anchor of the rectangle through the diagonal of the Pascal array (indicated by the arrow d in Figure 4), the weight remains constant. We will call this property as the weighted version of star of David rule. We can define a symmetric cross of size k in the Pascal array (See Figure 5) as an $2k$ -tuples $(c_1, \dots, c_k, r_1, \dots, r_k)$, where c_i, r_i , $(1 \leq i \leq k)$ are the entries of the Pascal array. We then define the weight of such a cross, W_c , as


$$W_c := \frac{c_1 \cdot c_2 \cdot \dots \cdot c_k}{r_1 \cdot r_2 \cdot \dots \cdot r_k}.$$

Remark 4. An immediate consequence of the weighted version of star of David rule in the Pascal array is that if we move the center of a cross through the diagonal d , then the weight W_c remains constant. We will call this property the *sliding cross rule* in the Pascal array. We also note that if the sliding cross rule holds for $m = l = 1$, then by an straight forward induction on m and l , it can be easily seen that it holds in general for every positive integers m and l .


FIGURE 5. A cross of size k

Our next step is to show that the above sliding rule is also true for any Pascal determinantal array PD_k ($k \geq 1$). To do this, we apply mathematical induction on k . The basis step, $k = 1$, is already true. For the inductive step, we assume that every Pascal determinantal array of order less than $k + 1$ has the sliding property and we show that it is also true for PD_{k+1} .

Consider the Pascal determinantal array PD_k , as in Figure 6, where the circled-points show the entries of this array and the squared entries show the corresponding entries of the Pascal determinantal array of order $k - 1$.

FIGURE 6. Pascal determinantal array PD_k

Consider the previous discussions, the entries of PD_{k+1} can be obtained, as follows (see Figure 7).


 FIGURE 7. Pascal determinantal array PD_{k+1}

$$(4) \quad \begin{aligned} \alpha &= \frac{ae - bd}{A}, \quad \beta = \frac{bf - ec}{B}, \quad \alpha' = \frac{a'e' - b'd'}{A'}, \quad \beta' = \frac{b'f' - e'c'}{B'} \\ \gamma &= \frac{dh - ge}{C}, \quad \theta = \frac{ek - fh}{D}, \quad \gamma' = \frac{d'h' - g'e'}{C'}, \quad \theta' = \frac{e'k' - f'h'}{D'}. \end{aligned}$$

Therefore, it is sufficient to show that $\frac{\alpha\theta}{\beta\gamma} = \frac{\alpha'\theta'}{\beta'\gamma'}$. This is equivalent to

$$(5) \quad \frac{\frac{(ae-bd)(ek-fh)}{AD}}{\frac{(bf-ec)(dh-ge)}{BC}} = \frac{\frac{(a'e'-b'd')(e'k'-f'h')}{A'D'}}{\frac{(b'f'-e'c')(d'h'-g'e')}{B'C'}}$$

By induction hypothesis, we have

$$\frac{ae}{bd} = \frac{a'e'}{b'd'}, \quad \frac{bf}{ec} = \frac{b'f'}{e'c'}, \quad \frac{dh}{ge} = \frac{d'h'}{g'e'}, \quad \frac{ek}{fh} = \frac{e'k'}{f'h'}, \quad \frac{AD}{BC} = \frac{A'D'}{B'C'}$$

or equivalently

$$\frac{ae - bd}{bd} = \frac{a'e' - b'd'}{b'd'}, \quad \frac{ek - fh}{fh} = \frac{e'k' - f'h'}{f'h'},$$

$$\frac{bf - ec}{ec} = \frac{b'f' - e'c'}{e'c'}, \quad \frac{dh - ge}{ge} = \frac{d'h' - g'e'}{g'e'},$$

$$\frac{AD}{BC} = \frac{A'D'}{B'C'}$$

which can be simply written as

$$\frac{\frac{(ae-bd)(ek-fh)}{(bdfh)AD}}{\frac{(bf-ec)(dh-ge)}{(ecge)BC}} = \frac{\frac{(a'e'-b'd')(e'k'-f'h')}{(b'd'f'h')A'D'}}{\frac{(b'f'-e'c')(d'h'-g'e')}{(e'c'g'e')B'C'}}.$$

Therefore by (2), we only need to show that

$$\frac{ecge}{bdfh} = \frac{e'c'g'e'}{b'd'f'h'}, \text{ or equivalently}$$

$$\frac{baecge}{babdfh} = \frac{b'a'e'c'g'e'}{b'a'b'd'f'h'}$$

But we already know, by induction hypothesis, that

$$\frac{ae}{bd} = \frac{a'e'}{b'd'}, \quad \frac{ce}{bf} = \frac{c'e'}{b'f'}, \quad \frac{bg}{ah} = \frac{b'g'}{a'h'}.$$

Thus considering the above remark, we complete the the proof by induction.

6. THE CORRECTNESS OF THE ALGORITHM

Now we are at the position to prove that the algorithm mentioned in the section 2 will produce every Pascal determinantal array of order k . To do this, we apply mathematical induction on k . The basis step, PD_1 , is obviously true. Assume that the algorithm is true for PD_k , as inductive step. Then, we have

$$(6) \quad P_{i,j}^{(k)} = \frac{P_{i+1,j+1}^{(k-1)} P_{i,j}^{(1)}}{P_{1,i+j+1}^{(k-1)}}.$$

Now we show that the algorithm is also true for the Pascal determinantal array PD_{k+1} . By the sliding cross property for PD_k , we have (see Figure 8):

$$(7) \quad \frac{P_{i+1,j}^{(k)} P_{i,j+1}^{(k)}}{P_{i,j}^{(k)} P_{i+1,j+1}^{(k)}} = \frac{P_{1,i+j}^{(k)} P_{0,i+j+1}^{(k)}}{P_{0,i+j}^{(k)} P_{1,i+j+1}^{(k)}},$$

Considering the fact that $P_{0,i+j+1}^{(k)} = P_{0,i+j}^{(k)} = 1$, it is not hard to see that the equality in (7) is equivalent to


$$(8) \quad \frac{P_{1,i+j}^{(k)}}{P_{i+1,j}^{(k)} P_{i,j+1}^{(k)}} = \frac{P_{1,i+j+1}^{(k-1)}}{P_{i,j}^{(k+1)} P_{i+1,j+1}^{(k-1)}},$$

Now, by reversing the both fraction in (8) and multiply them by $P_{i,j}^{(1)}$, we obtain

$$(9) \quad \frac{P_{i,j}^{(1)} P_{i+1,j}^{(k)} P_{i,j+1}^{(k)}}{P_{1,i+j}^{(k)}} = \frac{P_{i,j}^{(1)} P_{i,j}^{(k+1)} P_{i+1,j+1}^{(k-1)}}{P_{1,i+j+1}^{(k-1)}}.$$

Using relation (6), we immediately get

$$(10) \quad \frac{P_{i,j}^{(1)} P_{i+1,j}^{(k)} P_{i,j+1}^{(k)}}{P_{1,i+j}^{(k)}} = P_{i,j}^{(k+1)} P_{i,j}^{(k)}.$$


 FIGURE 8. The sliding property for PD_k

Next, by dividing both sides of (10) to $P_{i,j}^{(k)}$ and also multiplying the left-hand side by $\frac{P_{i+1,j+1}^{(k)}}{P_{i+1,j+1}^{(k)}}$, we have

$$(11) \quad \frac{P_{i+1,j+1}^{(k)} P_{i,j}^{(1)} P_{i+1,j}^{(k)} P_{i,j+1}^{(k)}}{P_{i+1,j+1}^{(k)} P_{i,j}^{(k)} P_{1,i+j}^{(k)}} = P_{i,j}^{(k+1)}.$$

By using the equation (6) one more time, we obtain

$$(12) \quad \frac{P_{i+1,j+1}^{(k)} P_{i,j}^{(1)} P_{1,i+j}^{(k)} P_{0,i+j+1}^{(k)}}{P_{0,i+j}^{(k+1)} P_{1,i+j+1}^{(k)} P_{1,i+j}^{(k)}} = P_{i,j}^{(k+1)}.$$

Finally, after simplifications based on the fact $P_{0,i+j+1}^{(k)} = P_{0,i+j}^{(k+1)} = 1$ and canceling out the similar terms, we thus obtain

$$\frac{P_{i+1,j+1}^{(k)} P_{i,j}^{(1)}}{P_{1,i+j+1}^{(k)}} = P_{i,j}^{(k+1)},$$

as required.

7. A GEOMETRIC INTERPRETATION OF $P_{i,j}^{(k)}$

In this section we present a geometric interpretation of (i, j) -entry of the Pascal determinantal array of order k . To do this, we first need to state a new identity for Pascal array entries which is a direct consequence of the sliding cross rule in Pascal array. Indeed for every positive k and nonnegative j , we have

$$P_{0,j+2k+1} \cdot P_{1,j+2k-1} \cdots P_{k-1,j+3} \cdot P_{k,j+1} = P_{0,j} \cdot P_{1,j+1} \cdots P_{k-1,j+k-1} \cdot P_{k,j+k}.$$

The corresponding entries are shown in Figure 9. For simplicity, we show the entries on left-hand side (LHS) of the above identity with a_i , ($1 \leq i \leq k+1$) and the entries on the right-hand side (RHS) with A_i , ($1 \leq i \leq k+1$). It is worth to note that $a_1 = A_1 = 1$.

The LHS entries are on the line with slope $-\frac{1}{1}$, the line L_a , and the RHS entries are on the line with slope $+\frac{1}{2}$, the line L_A .




FIGURE 9. A geometric interpretation of the new identity

Now we give a geometric proof of the above identity based on induction on k and sliding cross rule in Pascal array. The basis step is obviously true. By induction hypothesis the identity holds for entries b_1, b_2, \dots, b_k and A_1, A_2, \dots, A_k . So, we have

$$(13) \quad b_1 b_2 \dots b_k = A_1 A_2 \dots A_k.$$

On the other hand, considering two parallel crosses $(b_1, b_2, \dots, b_k, r_1, r_2, \dots, r_k)$ and $(a_2, \dots, a_{k+1}, r_2, \dots, r_{k+1})$ of size k , by sliding cross rule, we get

$$(14) \quad \frac{b_1 b_2 \dots b_k}{r_1 r_2 \dots r_k} = \frac{a_2 a_3 \dots a_{k+1}}{r_2 r_3 \dots r_{k+1}},$$

or equivalently

$$(15) \quad b_1 b_2 \dots b_k A_{k+1} = a_1 a_2 \dots a_{k+1}.$$

Now, (13) and (15) imply

$$a_1 a_2 \dots a_{k+1} = A_1 A_2 \dots A_{k+1},$$

as required.

In the next step, we use the algorithm of section 2, to find a closed-form formula for $P_{i,j}^{(k)}$ and it's geometric interpretation. We already know that

$$P_{i,j}^{(k+1)} = \frac{P_{i+1,j+1}^{(k)} P_{i,j}^{(1)}}{P_{1,i+j+1}^{(k)}}.$$

By repeated application of this formula and substitution in the right hand side, after $k - 1$ iterations, we get

$$P_{i,j}^{(k+1)} = \frac{P_{i+k,j+k}^{(1)} P_{i+(k-1),j+(k-1)}^{(1)} \cdots P_{i+1,j+1}^{(1)} P_{i,j}^{(1)}}{P_{1,i+j+2k-1}^{(1)} P_{1,i+j+2k-3}^{(2)} \cdots P_{1,i+j+3}^{(k-1)} P_{1,i+j+1}^{(k)}}.$$

On the other hand, by Rahimpour's theorem [1], we have

$$P_{1,j}^{(k)} = P_{k,j}^{(1)} \quad (j \geq 0, k \geq 1).$$

Thus, we obtain

$$P_{i,j}^{(k+1)} = \frac{P_{i+k,j+k}^{(1)} P_{i+(k-1),j+(k-1)}^{(1)} \cdots P_{i+1,j+1}^{(1)} P_{i,j}^{(1)}}{P_{1,i+j+2k-1}^{(1)} P_{2,i+j+2k-3}^{(1)} \cdots P_{k-1,i+j+3}^{(1)} P_{k,i+j+1}^{(1)}},$$

or equivalently,

$$P_{i,j}^{(k+1)} = \frac{P_{i+k,j+k} \cdot P_{i+(k-1),j+(k-1)} \cdots P_{i+1,j+1} \cdot P_{i,j}}{P_{1,i+j+2k-1} \cdot P_{2,i+j+2k-3} \cdots P_{k-1,i+j+3} \cdot P_{k,i+j+1}}.$$

Now, using the new identity with j replaced by $i + j$, we can alternatively obtain the following formula

$$P_{i,j}^{(k+1)} = \frac{P_{i+k,j+k} \cdot P_{i+(k-1),j+(k-1)} \cdots P_{i+1,j+1} \cdot P_{i,j}}{P_{1,i+j+1} \cdot P_{2,i+j+2} \cdots P_{k-1,i+j+(k-1)} \cdot P_{k,i+j+k}}.$$

Considering the two parallel symmetric crosses $(P_{i,j}, \dots, P_{i+k,j+k}, P_{i,j+k}, \dots, P_{i+k,j})$ and

$(P_{0,i+j}, \dots, P_{k,i+j+k}, P_{k,i+j}, \dots, P_{0,i+j+k})$ of size $k+1$, using the sliding property, we obtain

$$P_{i,j}^{(k+1)} = \frac{P_{i+k,j} \cdot P_{i+(k-1),j+1} \cdots P_{i+1,j+(k-1)} \cdot P_{i,j+k}}{P_{0,i+j+k} \cdot P_{1,i+j+(k-1)} \cdots P_{k-1,i+j+1} \cdot P_{k,i+j}}.$$

Finally, by replacing k with $k - 1$, we get

$$P_{i,j}^{(k)} = \frac{P_{i+(k-1),j} \cdot P_{i+(k-2),j+1} \cdots P_{i+1,j+(k-2)} \cdot P_{i,j+(k-1)}}{P_{i+j+(k-1),0} \cdot P_{i+j+(k-2),1} \cdots P_{i+j+1,k-2} \cdot P_{i+j,k-1}}.$$

Considering the above closed-form formula for $P_{i,j}^{(k)}$, we give the following geometric interpretation of k by k determinants in Pascal array.




FIGURE 10. A geometric interpretation of $P_{i,j}^{(k)}$

We will show a double stick (see Figure 10) by $(b_1, b_2, \dots, b_k | r_1, r_2, \dots, r_k)$ where b_i, r_i ($1 \leq i \leq k$) are entries of Pascal array which lie on the discrete line $x + y = i + j + (k - 1)$, the line L , where

$$b_1 := P_{i+(k-1),j}, b_k := P_{i,j+(k-1)},$$

and

$$r_1 := P_{i+j+(k-1),0}, r_k := P_{i+j,k-1}.$$

We define the weight of the double stick, $W_{i,j}^k$, as follows

$$W_{i,j}^{(k)} = \frac{b_1 b_2 \dots b_k}{r_1 r_2 \dots r_k}.$$

Now, clearly $P_{i,j}^{(k)}$ can be interpreted as the weight of double stick, i.e. $W_{i,j}^{(k)}$. Indeed the following simple fact that the overlapping area of the double stick will increase by increasing the value of k , is the essence of the proof of our main conjecture. Now, we are at the position to give a geometric proof of our conjecture, besides its algebraic proof, based on the above geometric interpretation of k by k determinants in Pascal array.

Proof of the conjecture. Based on the language of Pascal determinantal arrays, the conjecture is equivalent to the following statement (see Figure 11).

For any positive integers i, k and n , $n \geq k$, we have

$$P_{i,k}^{(n)} = P_{i,n}^{(k)}$$




FIGURE 11. Sub-arrays related to the main conjecture

Now let's look at the geometric meaning of $P_{i,k}^{(n)}$. It is equal to $W_{i,k}^{(n)}$, the weight of the double stick in Figure 12 (solid line). Clearly, it lies on the line $L : x + y = i + k + (n - 1)$. So we get

$$P_{i,k}^{(n)} := W_{i,k}^{(n)} = \frac{P_{i+(n-1),k} \cdots P_{i,k+(n-1)}}{P_{i+k+(n-1),0} \cdots P_{i+k,n-1}},$$

or equivalently,

$$P_{i,k}^{(n)} := W_{i,k}^{(n)} = \frac{(P_{i+(n-1),k} \cdots P_{i+k,n-1})(P_{i+(k-1),n} \cdots P_{i,n+(k-1)})}{(P_{i+n+(k-1),0} \cdots P_{i+n,k-1})(P_{i+(n-1),k} \cdots P_{i+k,n-1})}.$$




FIGURE 12. A geometric proof of the conjecture

Now, by canceling the common factor out of numerator and denominator of the above fraction which is geometrically equivalent to omit the overlapping area of the original double stick, we obtain

$$P_{i,k}^{(n)} := \frac{P_{i+(k-1),n} \cdots P_{i,n+(k-1)}}{P_{i+n+(k-1),0} \cdots P_{i+n,k-1}},$$

which is clearly the weight of the new double stick shown by the dash line. As it is easily seen from Figure 12. and the above algebraic formula, this is equal to $W_{i,n}^{(k)}$ or $P_{i,n}^{(k)}$, and this completes the proof.

REFERENCES

- [1] H. Teimoori and M. Bayat, Fermat Row-Eliminated Matrices and Some Binomial Determinants, *J. Math. Gazette.*, 87 (2003), pp. 114-118.
- [2] M. Bayat and H. Teimoori, Pascal k-eliminated Functional Matrix and It's Property, *Linear Algebra Appl.*, 308 (2000), pp.65-75.
- [3] P. Hilton and J. Pedersen, Relation Geometry and Algebra in the Pascal Triangle, Hexagon, Tetrahedron, and Cuboctahedron Part I: Binomial Coefficients, Extended Binomial Coefficients and Prepration for further Work, *The College Math. J.*, 30 (3)(1999), pp. 170-186.
- [4] C. L. Dodgson, Condensation of Determinants: Being a New and Brief Method for Computing their Arithmetic Values, *Proc. Roy. Soc. Ser. A*, 15 (1866)150-155.
- [5] N.J.A.Sloane, Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequence,2008>.
- [6] K. Robin McLean, Determinants of binomial coefficients, *J. Math. Gazette.*, 89 (2005), pp. 247-250.

DEPARTMENT OF COMPUTER SCIENCE, FACULTY OF MATHEMATICAL AND COMPUTER SCIENCES, ALLAMEH TABATABA'I UNIVERSITY, TEHRAN, IRAN.,
Email address: `hossein.teimoori@atu.ac.ir`

MINISTRY OF INFORMATION AND COMMUNICATION TECHNOLOGY (ICT), ZANJAN, IRAN.
Email address: `hasan.khodakarami@gmail.com`