

Random points are good for universal discretization

F. Dai and V. Temlyakov *

Abstract

Recently, there was a big progress in studying sampling discretization of integral norms of finite dimensional subspaces and collections of such subspaces (universal discretization). It was established that sampling discretization results are useful in a number of applications. In particular, they turn out to be useful in sampling recovery. Typically, recent sampling discretization results provide existence of good points for discretization. The main goal of this paper is to show that in the problem of universal discretization the independent random points on a given domain that are identically distributed according to the given probabilistic measure provide good points with high probability. Also, we demonstrate that a simple greedy type algorithm based on good points for universal discretization provide good recovery in the square norm.

Keywords and phrases: Sampling discretization, universality, recovery.

MSC classification 2000: Primary 65J05; Secondary 42A05, 65D30, 41A63.

1 Introduction

This paper is a follow up to [5]. We begin with a brief description of the new ingredients of this paper. In [5] we discussed in detail universal discretization of integral norms for collections of v -dimensional subspaces spanned by

*The first named author's research was partially supported by NSERC of Canada Discovery Grant RGPIN-2020-03909. The second named author's research was supported by the Russian Federation Government Grant No. 14.W03.31.0031.

elements of a given dictionary \mathcal{D}_N . One of the main results of [5] states that under certain conditions on \mathcal{D}_N we can guarantee good universal discretization of the L_p norm, $1 \leq p \leq 2$, with the number of points m of the order $v(\log N)^2(\log v)^2$. Note that this bound on m is much better than previously known bounds which are quadratic in v . Also, the following simple lower bound $m \geq v$ is well known. Thus, the bound $m \ll v(\log N)^2(\log v)^2$ is a reasonably good upper bound. In [5] we only discuss the problem of existence of good points for discretization. In the very recent paper [6] we showed that points of good universal discretization can be used for sparse recovery of multivariate smooth functions. This motivated us to address the problem of finding points for good universal discretization. This is the main new ingredient of this paper. We demonstrate that in a number of existence type results from [5] we can guarantee a similar result with high probability for randomly chosen points. We use the technique developed in [5]. However, the step from existence theorems to statements “with high probability for randomly chosen points” turns out to be nontrivial. In addition to this step we also give a somewhat better bound on m . We prove $m \ll v(\log N)(\log(2v))^2(\log(2v) + \log \log N)$. Roughly speaking, we replace one of $\log N$ by $\log(2v)$. It seems like a very minor improvement, however we prove in Section 5 that we cannot get rid of the factor $\log N$ in the bound for universal discretization. From this point of view we improve the dependence on N from $(\log N)^2$ to $\log N \log \log N$, which is very close to the optimal.

In the very recent paper [6] we used the universal discretization results for the L_2 norm for analyzing a theoretical sparse recovery algorithm, based on a special dictionary \mathcal{D}_N . We showed that in the sense of accuracy it is a very good algorithm – we proved the corresponding Lebesgue-type inequalities for it. In Section 4 we study a very simple greedy-type algorithm for sparse recovery with respect to a special dictionary \mathcal{D}_N and prove the Lebesgue-type inequalities for it.

We now proceed to a detailed description of the main results. We refer the reader to [5], [4], and [10] for historical comments. Let Ω be a nonempty set (usually a compact set) equipped with a probability measure μ . For $1 \leq p \leq \infty$, we let $L_p(\Omega) := L_p(\Omega, \mu)$ denote the Lebesgue space L_p defined with respect to the measure μ on Ω and let $\|\cdot\|_p$ the norm of $L_p(\Omega)$. By discretization of the L_p norm we understand a replacement of the measure μ by a discrete measure μ_m with support on a set $\xi = \{\xi^j\}_{j=1}^m \subset \Omega$. This means that integration with respect to the measure μ is replaced by an appropriate cubature formula. Thus, integration is replaced by evaluation of a

function f at a finite set ξ of points. This is why this way of discretization is called *sampling discretization*. We now give explicit formulations of the sampling discretization problem (also known as the Marcinkiewicz discretization problem) and of the problem of universal discretization.

The sampling discretization problem. Let (Ω, μ) be a probability space and $X_N \subset L_q$ be an N -dimensional subspace of $L_q(\Omega, \mu)$ with $1 \leq q \leq \infty$ (the index N here, usually, stands for the dimension of X_N). We shall always assume that every function in X_N is defined everywhere on Ω , and

$$f \in X_N, \|f\|_q = 0 \implies f = 0 \in X_N.$$

We say that X_N admits the Marcinkiewicz-type discretization theorem with parameters $m \in \mathbb{N}$ and q and positive constants $C_1 \leq C_2$ if there exists a set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ such that for any $f \in X_N$ we have in the case $1 \leq q < \infty$

$$C_1 \|f\|_q^q \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \leq C_2 \|f\|_q^q \quad (1.1)$$

and in the case $q = \infty$

$$C_1 \|f\|_\infty \leq \max_{1 \leq j \leq m} |f(\xi^j)| \leq \|f\|_\infty.$$

Here and throughout the paper, we will use a slight abuse of the notation that $L_\infty(\Omega)$ denotes the space of all uniformly bounded measurable functions on Ω with norm $\|f\|_\infty = \sup_{x \in \Omega} |f(x)|$.

The problem of universal discretization. Let $\mathcal{X} := \{X(n)\}_{n=1}^k$ be a collection of finite-dimensional linear subspaces $X(n)$ of the $L_q(\Omega)$, $1 \leq q \leq \infty$. We say that a set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ provides *universal discretization* for the collection \mathcal{X} if, in the case $1 \leq q < \infty$, there are two positive constants C_i , $i = 1, 2$, such that for each $n \in \{1, \dots, k\}$ and any $f \in X(n)$ we have

$$C_1 \|f\|_q^q \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \leq C_2 \|f\|_q^q.$$

In the case $q = \infty$ for each $n \in \{1, \dots, k\}$ and any $f \in X(n)$ we have

$$C_1 \|f\|_\infty \leq \max_{1 \leq j \leq m} |f(\xi^j)| \leq \|f\|_\infty. \quad (1.2)$$

Note that the problem of universal discretization for the collection $\mathcal{X} := \{X(n)\}_{n=1}^k$ is the sampling discretization problem for the set $\cup_{n=1}^k X(n)$. Also,

we point out that the concept of universality is well known in approximation theory. For instance, the reader can find a discussion of universal cubature formulas in [16], Section 6.8.

In this paper we focus on special collections of subspaces generated by a given dictionary. We begin our discussion with the concept of v -term approximation with respect to a given dictionary $\mathcal{D}_N = \{g_i\}_{i=1}^N$. Given an integer $1 \leq v \leq N$, we denote by $\mathcal{X}_v(\mathcal{D}_N)$ the collection of all linear spaces spanned by g_j , $j \in J$ with $J \subset [1, N] \cap \mathbb{N}$ and $|J| = v$, and denote by $\Sigma_v(\mathcal{D}_N)$ the set of all v -term approximants with respect to \mathcal{D}_N :

$$\Sigma_v(\mathcal{D}_N) := \bigcup_{V \in \mathcal{X}_v(\mathcal{D}_N)} V.$$

Denote

$$\Sigma_v^q(\mathcal{D}_N) := \{f \in \Sigma_v(\mathcal{D}_N) : \|f\|_q \leq 1\}.$$

We focus on special type of dictionaries. We assume that $\Phi_N := \{\varphi_j\}_{j=1}^N$ is a dictionary of N uniformly bounded functions on $\Omega \subset \mathbb{R}^d$ satisfying

$$\sup_{\mathbf{x} \in \Omega} |\varphi_j(\mathbf{x})| \leq 1, \quad 1 \leq j \leq N, \quad (1.3)$$

and that there exists a constant $K > 0$ such that for any $(a_1, \dots, a_N) \in \mathbb{C}^N$,

$$\sum_{j=1}^N |a_j|^2 \leq K \left\| \sum_{j=1}^N a_j \varphi_j \right\|_2^2. \quad (1.4)$$

Theorem 1.1. *Let $1 \leq p \leq 2$. Assume that $\Phi_N = \{\varphi_j\}_{j=1}^N \subset L_\infty(\Omega)$ is a dictionary satisfying the conditions (1.3) and (1.4) for some constant $K \geq 1$. Let ξ^1, \dots, ξ^m be independent random points on Ω that are identically distributed according to μ . Then there exist constants $C = C(p) > 1$ and $c = c(p) > 0$ such that given any integers $1 \leq v \leq N$ and*

$$m \geq CKv \log N \cdot (\log(2Kv))^2 \cdot (\log(2Kv) + \log \log N),$$

the inequalities

$$\frac{1}{2} \|f\|_p^p \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p \leq \frac{3}{2} \|f\|_p^p, \quad \forall f \in \Sigma_v(\Phi_N) \quad (1.5)$$

hold with probability $\geq 1 - 2 \exp\left(-\frac{cm}{Kv \log^2(2Kv)}\right)$.

Remark 1.1. *If the functions $\varphi_1, \dots, \varphi_N$ are linearly independent in $L_2(\Omega, \mu)$, then the condition (1.4) holds automatically with K^{-1} being the smallest eigenvalue of the $N \times N$ matrix $\left[\langle \varphi_i, \varphi_j \rangle_{L_2(\mu)} \right]_{1 \leq i, j \leq N}$.*

We now discuss sparse sampling recovery in the case Φ_N is a uniformly bounded Riesz basis satisfying (1.3) and

$$R_1 \left(\sum_{j=1}^N |a_j|^2 \right)^{1/2} \leq \left\| \sum_{j=1}^N a_j \varphi_j \right\|_2 \leq R_2 \left(\sum_{j=1}^N |a_j|^2 \right)^{1/2}, \quad (1.6)$$

where $0 < R_1 \leq R_2 < \infty$.

Define for a Banach space X

$$\sigma_v(f, \mathcal{D}_N)_X := \inf_{g \in \Sigma_v(\mathcal{D}_N)} \|f - g\|_X$$

to be the best v -term approximation of $f \in X$ in the X -norm with respect to \mathcal{D}_N , and define

$$\sigma_v(\mathbf{F}, \mathcal{D}_N)_X := \sup_{f \in \mathbf{F}} \sigma_v(f, \mathcal{D}_N)_X, \quad \sigma_0(\mathbf{F}, \mathcal{D}_N)_X := \sup_{f \in \mathbf{F}} \|f\|_X$$

for a function class $\mathbf{F} \subset X$.

Definition 1.1. *We say that a set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ provides universal discretization for the collection $\mathcal{X} := \{X(n)\}_{n=1}^k$ of finite-dimensional linear subspaces $X(n)$ if we have*

$$\frac{1}{2} \|f\|_2^2 \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^2 \leq \frac{3}{2} \|f\|_2^2 \quad \text{for any } f \in \bigcup_{n=1}^k X(n). \quad (1.7)$$

We denote by $m(\mathcal{X})$ the minimal m such that there exists a set ξ of m points, which provides universal discretization (1.7) for the collection \mathcal{X} .

The following Theorem 1.2 is a conditional result, which guarantees that a simple greedy type algorithm – Weak Orthogonal Matching Pursuit (WOMP) (see Section 4 for the definition) – provides good sparse recovery by using points of good universal discretization. Theorem 1.1 shows that we can choose those points randomly with high probability. Instead of the space $L_2(\Omega, \mu)$ we consider the space $L_2(\Omega_m, \mu_m)$ where $\Omega_m = \{\xi^\nu\}_{\nu=1}^m$ is from Definition 4.1 and $\mu_m(\xi^\nu) = 1/m$, $\nu = 1, \dots, m$. Let $\Phi_N(\Omega_m)$ be the restriction of Φ_N onto Ω_m .

Theorem 1.2. *Assume that Φ_N is a uniformly bounded Riesz basis of $[\Phi_N]$ satisfying (1.6) for some constants $0 < R_1 \leq R_2 < \infty$. For given parameters $t \in (0, 1]$ and R_1, R_2 from above there exists a constant $c = C(t, R_1, R_2)$ with the following property. Let for integer v such that $u := (1 + c)v \leq N$ and a number $m \geq m(\mathcal{X}_u(\Phi_N))$ the set ξ of m points $\xi^1, \dots, \xi^m \in \Omega$ provide universal discretization for the collection $\mathcal{X}_u(\Phi_N)$. Then for any $f_0 \in \mathcal{C}(\Omega)$ the WOMP with weakness parameter t applied to f_0 with respect to the normalized $\Phi_N(\Omega_m)$ in the space $L_2(\Omega_m, \mu_m)$ provides*

$$\|f_{cv}\|_{L_2(\Omega_m, \mu_m)} \leq C \sigma_v(f_0, \Phi_N(\Omega_m))_{L_2(\Omega_m, \mu_m)}, \quad (1.8)$$

and

$$\|f_{cv}\|_{L_2(\Omega, \mu)} \leq C' \sigma_v(f_0, \Phi_N)_\infty \quad (1.9)$$

with absolute constants C and C' .

Corollary 1.1. *Under assumptions of Theorem 1.2 the WOMP with weakness parameter t applied to $f_0 \in \Sigma_v(\Phi_N)$ with respect to the normalized $\Phi_N(\Omega_m)$ in the space $L_2(\Omega_m, \mu_m)$ recovers it exactly.*

We now discuss an application of Theorem 1.2 to the optimal sampling recovery. The reader can find further discussion in Section 4. We allow any mapping $\Psi : \mathbb{C}^m \rightarrow L_p(\Omega, \mu)$ and for a function class $\mathbf{F} \subset \mathcal{C}(\Omega)$ define

$$\varrho_m^o(\mathbf{F}, L_p) := \inf_{\Psi; \xi} \sup_{f \in \mathbf{F}} \|f - \Psi(f(\xi^1), \dots, f(\xi^m))\|_p.$$

Here, index o stands for optimal. The following Theorem 1.3 is a direct corollary of Theorem 1.2.

Theorem 1.3. *Assume that Φ_N and c are from Theorem 1.2. Then for any $\mathbf{F} \subset \mathcal{C}(\Omega)$ we have*

$$\varrho_m^o(\mathbf{F}, L_2) \leq C \sigma_v(\mathbf{F}, \Phi_N)_\infty \quad (1.10)$$

provided $m \geq m(\mathcal{X}_u(\Phi_N))$ with an absolute constant C .

The first inequality of the kind of (1.10) was obtained in the very recent paper by T. Jahn, T. Ullrich, and F. Voigtlaender [9]. They used deep known results from compressed sensing and applied the ℓ_1 minimization algorithm. In addition to function values at m points their algorithm uses the fact $f \in \mathbf{F}$ and the quantity $\sigma_v(\mathbf{F}, \Phi_N)_\infty$. They proved an inequality similar to (1.10) for the uniformly bounded orthonormal system Φ_N with an extra term (not

important one) in the right hand side of (1.10). Their algorithm does not provide a sparse with respect to Φ_N approximant. We improve the result from [9] in a number of ways. First, we get rid of the extra term in the right hand side. Second, our algorithm (WOMP) only uses the function values at m points and does not use any information about the class \mathbf{F} . Thus, it is universal in that sense. Third, we prove (1.10) for a wider class of systems Φ_N . Forth, our algorithm provides cv -sparse with respect to Φ_N approximant. Finally, for our algorithm we prove the Lebesgue-type inequality (1.9) for individual functions.

2 Some general entropy bounds

We first recall the definition of entropy numbers in Banach spaces. Let A be a compact subset of a Banach space $(X, \|\cdot\|)$. Given a positive number ε , the covering number $N_\varepsilon(A, X)$ of the set A in the space X is defined to be the smallest positive integer n for which there exist n elements $g_1, \dots, g_n \in A$ such that

$$\max_{f \in A} \min_{1 \leq j \leq n} \|f - g_j\| \leq \varepsilon.$$

We denote by $\mathcal{N}_\varepsilon(A, X)$ the corresponding minimal ε -net of the set A in X . The ε -entropy $\mathcal{H}_\varepsilon(A, X)$ of the compact set A in X is then defined as

$$\mathcal{H}_\varepsilon(A, X) = \log_2 N_\varepsilon(A, X), \quad \varepsilon > 0,$$

whereas the entropy numbers of A in the space X are defined as

$$\varepsilon_k(A, X) := \inf\{t > 0 : \mathcal{H}_t(A, X) \leq k\}, \quad k = 0, 1, 2, \dots$$

The following theorem is a direct consequence of a more general conditional theorem that is proved in Section 6 (see Theorem 6.1).

Theorem 2.1. *Let $1 \leq q < \infty$ and $1 \leq v \leq N$. Suppose that \mathcal{D}_N is a dictionary of N functions from $L_\infty(\Omega)$ such that the set $\Sigma_v^q(\mathcal{D}_N)$ satisfies*

$$\varepsilon_k(\Sigma_v^q(\mathcal{D}_N), L_\infty) \leq B_1(v/k)^{1/q}, \quad 1 \leq k < \infty, \quad (2.1)$$

and

$$\|f\|_\infty \leq B_2 v^{1/q} \|f\|_q, \quad \forall f \in \Sigma_v(\mathcal{D}_N) \quad (2.2)$$

for some constants $B_1, B_2 \geq 1$. Let $\xi^1, \dots, \xi^m \in \Omega$ be independent random points satisfying

$$\frac{1}{m} \sum_{k=1}^m \mu_{\xi^k} = \mu, \quad (2.3)$$

where μ_{ξ^j} denotes the probability distribution of ξ^j . If

$$m \geq C(q) B_1^q v (\log(2B_2 v))^2,$$

for some large constant $C(q) > 1$ depending on q , then the inequalities

$$\frac{3}{4} \|f\|_q^q \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^q \leq \frac{5}{4} \|f\|_q^q, \quad \forall f \in \Sigma_v(\mathcal{D}_N)$$

hold with probability

$$\geq 1 - \exp\left[-\frac{c_q m}{B_2^q v \log^2(2B_2 v)}\right].$$

Clearly, the condition (2.3) is a direct consequence of either of the following two conditions:

- (i) ξ^1, \dots, ξ^m are identically distributed according to μ ;
- (ii) there exists a partition $\{\Omega_1, \dots, \Omega_m\}$ of Ω such that $\mu(\Omega_j) = \frac{1}{m}$ and $\xi^j \in \Omega_j$ is distributed according to $m \cdot \mu|_{\Omega_j}$ for each $1 \leq j \leq m$.

Next, we recall the following simple remark from [5].

Remark 2.1. *We point out that (2.1) implies*

$$\|f\|_\infty \leq 3B_1 v^{1/q}, \quad \forall f \in \Sigma_v^q(\mathcal{D}_N). \quad (2.4)$$

Therefore, assumption (2.2) can be dropped with B_2 replaced by $3B_1$ in the bound on m . However, in applications, the constant B_2 in (2.2) may be significantly smaller than $3B_1$. For example, if \mathcal{D}_N is a uniformly bounded orthonormal system with $\max_{f \in \mathcal{D}_N} \|f\|_\infty = 1$, then we can take $B_2 = 1$ in the case $q = 2$.

Theorem 2.1 motivates us to estimate the characteristics $\varepsilon_k(\Sigma_v^q(\mathcal{D}_N), L_\infty)$. In this section we prove that in the case when a system Φ_N satisfies (1.3) and (1.4) the estimates

$$\varepsilon_k(\Sigma_v^2(\Phi_N), L_\infty) \leq Cq_v \left(\frac{v}{k}\right)^{1/2}, \quad k = 1, 2, \dots, \quad (2.5)$$

hold under the additional assumption on Φ_N that there exist a number $q_v := q_{N,v} > 2$ and a universal constant $C_0 > 1$ such that

$$\|f\|_\infty \leq C_0 \|f\|_{q_v}, \quad \forall f \in \Sigma_{2v}(\Phi_N). \quad (2.6)$$

We now recall some known general results, which turn out to be useful for that purpose. The following Theorem 2.2 was proved in [14] (see also [16], p.331, Theorem 7.4.3).

Theorem 2.2. *Let W be a compact subset of a Banach space X for which there exist a system $\mathcal{D}_N \subset X$ with $|\mathcal{D}_N| = N$, and a number $r > 0$ such that*

$$\sigma_m(W, \mathcal{D}_N)_X \leq (m+1)^{-r}, \quad m = 0, 1, \dots, N.$$

Then for $k \leq N$

$$\varepsilon_k(W, X) \leq C(r) \left(\frac{\log(2N/k)}{k}\right)^r. \quad (2.7)$$

For a given set $\mathcal{D}_N = \{g_j\}_{j=1}^N$ of elements we introduce the octahedron (generalized octahedron)

$$A_1(\mathcal{D}_N) := \left\{ f : f = \sum_{j=1}^N c_j g_j, \quad \sum_{j=1}^N |c_j| \leq 1 \right\} \quad (2.8)$$

and the norm $\|\cdot\|_A$ on X_N

$$\|f\|_A := \inf \left\{ \sum_{j=1}^N |c_j| : f = \sum_{j=1}^N c_j g_j \right\}.$$

We now use a known general result for a smooth Banach space. For a Banach space X we define the modulus of smoothness

$$\rho(X, u) := \sup_{\|x\|=\|y\|=1} \left(\frac{1}{2} (\|x+uy\| + \|x-uy\|) - 1 \right).$$

The uniformly smooth Banach space is the one with the property

$$\lim_{u \rightarrow 0} \rho(X, u)/u = 0.$$

A Banach space X is called s -smooth for some parameter $1 < s \leq 2$ and constant $\gamma > 0$ if $\rho(X, u) \leq \gamma u^s$ for all $u > 0$. The following bound is a corollary of greedy approximation results (see, for instance [16], p.455).

Theorem 2.3. *Let X be an s -smooth Banach space for some parameter $1 < s \leq 2$ and constant $\gamma > 0$. Let $\mathcal{D}_N = \{g_j\}_{j=1}^N \subset X$ be a dictionary normalized so that $\|g_j\|_X \leq 1$ for all $1 \leq j \leq N$. Then for any $1 \leq m \leq N$,*

$$\sigma_m(A_1(\mathcal{D}_N), X) \leq C(s)\gamma^{1/s}m^{1/s-1}.$$

We now proceed to a special case of $X = L_p$, where it is known that

$$\rho(L_p, u) \leq (p-1)u^2/2, \quad 2 \leq p < \infty, \quad u > 0. \quad (2.9)$$

Theorem 2.4. *Let $\Phi_N = \{\varphi_j\}_{j=1}^N \subset L_\infty(\Omega)$ be a dictionary satisfying the conditions (1.3), (1.4) and (2.6) for some constants $K \geq 1$ and $q_v := q_{N,v} > 2$. Then we have*

$$\varepsilon_k(\Sigma_v^2(\Phi_N), L_\infty) \leq C\sqrt{Kq_v \cdot \log N} \left(\frac{v}{k}\right)^{1/2}, \quad k = 1, 2, \dots \quad (2.10)$$

Proof. First of all, for any $f = \sum_{j \in G} a_j \varphi_j$ with $|G| = v$, we have

$$\|f\|_A \leq \sum_{j \in G} |a_j| \leq v^{1/2} \left(\sum_{j \in G} |a_j|^2 \right)^{1/2} \leq (Kv)^{1/2} \|f\|_2, \quad (2.11)$$

implying

$$\Sigma_v^2(\Phi_N) \subset (Kv)^{1/2} \Sigma_v^A(\Phi_N),$$

where

$$R\Sigma_v^A(\Phi_N) := \{f \in \Sigma_v(\Phi_N) : \|f\|_A \leq R\}.$$

By Theorem 2.3 with $s = 2$ and by (2.9), we have that for $p \in [2, \infty)$

$$\sigma_m(\Sigma_v^2(\Phi_N), L_p) \leq C(Kv)^{1/2} \sqrt{pm}^{-\frac{1}{2}}, \quad m = 1, 2, \dots, N. \quad (2.12)$$

Thus, Theorem 2.2 implies that for $p \in [2, \infty)$

$$\varepsilon_k(\Sigma_v^2(\Phi_N), L_p) \leq C(Kp \log(2N/k))^{1/2} (v/k)^{1/2}, \quad k = 1, 2, \dots, N. \quad (2.13)$$

Second, by (2.6) we obtain

$$\varepsilon_k(\Sigma_v^2(\Phi_N), L_\infty) \leq C_0 \varepsilon_k(\Sigma_v^2(\Phi_N), L_{q_v}). \quad (2.14)$$

Combining (2.13) and (2.14) we get

$$\varepsilon_k(\Sigma_v^2(\Phi_N), L_\infty) \leq C \sqrt{K q_v \cdot \log N} (v/k)^{1/2}, \quad k = 1, 2, \dots, N. \quad (2.15)$$

Finally, for $k > N$ we use the inequalities

$$\varepsilon_k(W, L_\infty) \leq \varepsilon_N(W, L_\infty) \varepsilon_{k-N}(X_N^\infty, L_\infty)$$

and

$$\varepsilon_n(X_N^\infty, L_\infty) \leq 3 \cdot 2^{-n/N}, \quad 2^{-x} \leq 1/x, \quad x \geq 1,$$

to obtain (2.15) for all k . This completes the proof. \square

In the same way as in Section 3 of [5] we derive from Theorem 2.4, which holds for $p = 2$, the following Theorem 2.5, which holds for $1 \leq p \leq 2$. We do not present this proof here.

Theorem 2.5. *Assume that $\Phi_N = \{\varphi_j\}_{j=1}^N \subset L_\infty(\Omega)$ is a dictionary satisfying the conditions (1.3), (1.4) and (2.6) for some constants $K \geq 1$ and $q_v := q_{N,v} > 2$. Then for $1 \leq p \leq 2$, we have*

$$\varepsilon_k(\Sigma_v^p(\Phi_N), L_\infty) \leq C_p \left(\frac{K q_v \cdot \log N \cdot v}{k} \right)^{1/p}, \quad k = 1, 2, \dots \quad (2.16)$$

3 Proof of Theorem 1.1

Theorem 2.5 provides bounds on the entropy numbers $\varepsilon_k(\Sigma_v^p(\Phi_N), L_\infty)$ under the additional assumption (2.6)

$$\|f\|_\infty \leq C_0 \|f\|_{q_v}, \quad \forall f \in \Sigma_{2v}(\Phi_N).$$

Thus, a combination of Theorem 2.5 with Theorem 2.1 implies the conclusion of Theorem 1.1 for

$$m \geq C(p)(K q_v \cdot \log N)v(\log(2Kv))^2$$

under the additional condition (2.6), which is not assumed in Theorem 1.1. In this section, we will show how to drop this additional assumption.

Recall that $\mathcal{D}_N = \Phi_N := \{\varphi_j\}_{j=1}^N \subset L_\infty(\Omega)$ is a dictionary satisfying the conditions (1.3) and (1.4) for some constant $K \geq 1$. For a set $J \subset [N] := \{1, 2, \dots, N\}$, we define

$$V_J := \text{span} \left\{ \varphi_j : j \in J \right\} \quad \text{and} \quad V_J^p := \{f \in V_J : \|f\|_p \leq 1\}.$$

Then $\Sigma_v(\Phi_N) := \bigcup_{\substack{J \subset [N] \\ |J|=v}} V_J$ for $1 \leq v \leq N$. Furthermore, by (1.3) and (1.4),

we have

$$\|f\|_\infty \leq (Kv)^{\frac{1}{2}} \|f\|_2, \quad \forall f \in \Sigma_v(\Phi_N),$$

which in turn implies that

$$\|f\|_\infty \leq (Kv)^{\frac{1}{p}} \|f\|_p, \quad \forall f \in \Sigma_v(\Phi_N), \quad 1 \leq p \leq 2. \quad (3.1)$$

We need several lemmas. The first three lemmas can be found in [1].

Lemma 3.1. [1, Lemma 2.4] *Let X denote the space \mathbb{R}^N endowed with a norm $\|\cdot\|_X$, and let $B_X := \{x \in X : \|x\|_X \leq 1\}$. Then for any $0 < \varepsilon \leq 1$,*

$$N \log \frac{1}{\varepsilon} \leq \mathcal{H}_\varepsilon(B_X, X) \leq N \log\left(1 + \frac{2}{\varepsilon}\right). \quad (3.2)$$

Lemma 3.2. [1, Lemma 2.1] *Let $\{g_j\}_{j=1}^m$ be independent random variables with mean 0 on some probability space (Ω, μ) , which satisfy*

$$\max_{1 \leq j \leq m} \|g_j\|_{L_1(d\mu)} \leq M_1, \quad \max_{1 \leq j \leq m} \|g_j\|_{L_\infty(d\mu)} \leq M_\infty$$

for some constants M_1 and M_∞ . Then for any $0 < \varepsilon < 1$ we have the inequality

$$\mu \left\{ \omega \in \Omega : \left| \frac{1}{m} \sum_{j=1}^m g_j(\omega) \right| \geq \varepsilon \right\} \leq 2e^{-\frac{m\varepsilon^2}{4M_1M_\infty}}.$$

Lemma 3.3. [1, Lemma 2.5] *Let $T : X \rightarrow Y$ be a bounded linear map from a normed linear space $(X, \|\cdot\|_X)$ into another normed linear space $(Y, \|\cdot\|_Y)$. Let \mathcal{F} be an ε -net of the unit ball $B_X := \{x \in X : \|x\|_X \leq 1\}$ for some constant $\varepsilon \in (0, 1)$. Assume that there exist constants $C_1, C_2 > 0$ such that*

$$C_1 \|x\|_X \leq \|Tx\|_Y \leq C_2 \|x\|_X, \quad \forall x \in \mathcal{F}.$$

Then

$$C_1(\varepsilon)\|z\|_X \leq \|Tz\|_Y \leq C_2(\varepsilon)\|z\|_X, \quad \forall z \in X,$$

where

$$C_1(\varepsilon) := C_1(1 - \varepsilon) - C_2\varepsilon\frac{1 + \varepsilon}{1 - \varepsilon}, \quad C_2(\varepsilon) := C_2\frac{1 + \varepsilon}{1 - \varepsilon}.$$

From the above three lemmas, we can deduce

Lemma 3.4. *Let $1 \leq p \leq 2$ be a fixed number. Let $\{\xi^j\}_{j=1}^\infty$ be a sequence of independent random points identically distributed according to μ . Then for any $0 < \varepsilon \leq \frac{1}{8}$ and any integer*

$$m \geq 16K\varepsilon^{-2}v^2 \log\left(\frac{4e}{\varepsilon} \cdot \frac{N}{v}\right), \quad (3.3)$$

the inequalities

$$(1 - 4\varepsilon)\|f\|_p^p \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p \leq (1 + 4\varepsilon)\|f\|_p^p, \quad \forall f \in \Sigma_v(\Phi_N) \quad (3.4)$$

hold with probability $\geq 1 - 2\exp\left(-\frac{m\varepsilon^2}{16Kv}\right)$.

Proof. By Lemma 3.1, for each $J \subset [N]$ with $|J| = v$, there exists an ε -net $\mathcal{F}_J \subset V_J^p$ of V_J^p in the space L_p such that $|\mathcal{F}_J| \leq \left(1 + \frac{2}{\varepsilon}\right)^v$. Let $\mathcal{F} := \bigcup_{J \subset [N], |J|=v} \mathcal{F}_J$. Then

$$\begin{aligned} \log |\mathcal{F}| &\leq \log \left[\binom{N}{v} \left(1 + \frac{2}{\varepsilon}\right)^v \right] \leq \log \left[\left(\frac{eN}{v}\right)^v \left(1 + \frac{2}{\varepsilon}\right)^v \right] \\ &\leq v \log \left(\frac{4eN}{v\varepsilon} \right). \end{aligned} \quad (3.5)$$

On the other hand, (3.1) implies

$$\|f\|_\infty^p \leq Kv, \quad \forall f \in \mathcal{F}.$$

Thus, using (3.5) and Lemma 3.2, we conclude that the inequalities

$$\left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p - \int_\Omega |f|^p d\mu \right| \leq \varepsilon \|f\|_p^p, \quad \forall f \in \mathcal{F} \quad (3.6)$$

hold with probability

$$\begin{aligned} &\geq 1 - 2|\mathcal{F}| \exp\left(-\frac{m\varepsilon^2}{8Kv}\right) = 1 - 2 \exp\left(\log |\mathcal{F}| - \frac{m\varepsilon^2}{8Kv}\right) \\ &\geq 1 - 2 \exp\left(v \log\left(\frac{4eN}{v\varepsilon}\right) - \frac{m\varepsilon^2}{8Kv}\right). \end{aligned}$$

Under the condition (3.3), we have

$$\frac{m\varepsilon^2}{16Kv} \geq v \log\left(\frac{4eN}{v\varepsilon}\right),$$

implying that (3.6) holds with probability

$$\geq 1 - 2 \exp\left(-\frac{m\varepsilon^2}{16Kv}\right).$$

To complete the proof, we just need to note that by Lemma 3.3, (3.6) implies (3.4). □

Given $\mathbf{x} = (x_1, x_2, \dots, x_m) \in \Omega^m$ and $f : \Omega \rightarrow \mathbb{R}$, we define

$$S(f, \mathbf{x}) := (f(x_1), f(x_2), \dots, f(x_m)) \in \mathbb{R}^m.$$

We denote by L_p^m the space \mathbb{R}^m equipped with the discrete norm

$$\|\mathbf{z}\|_p := \begin{cases} \left(\frac{1}{m} \sum_{j=1}^m |z_j|^p\right)^{1/p}, & 1 \leq p < \infty, \\ \max_{1 \leq j \leq m} |z_j|, & p = \infty, \end{cases}, \quad \mathbf{z} = (z_1, \dots, z_m) \in \mathbb{R}^m.$$

Thus,

$$\|S(f, \mathbf{x})\|_p = \left(\frac{1}{m} \sum_{j=1}^m |f(x_j)|^p\right)^{1/p}, \quad 1 \leq p < \infty,$$

and

$$\|S(f, \mathbf{x})\|_\infty = \max_{1 \leq j \leq m} |f(x_j)|.$$

As usual, we identify a vector in \mathbb{R}^m with a function on the set $[m] := \{1, 2, \dots, m\}$. Under this identification, $\text{span}\{S(\varphi_i, \mathbf{x}) : 1 \leq i \leq N\}$ is a subspace of L_p^m .

Now we are in a position to prove Theorem 1.1. Without loss of generality, we may assume that $m \leq CKv^2 \log\left(\frac{eN}{v}\right)$ for some large constant C , since otherwise Theorem 1.1 follows directly from Lemma 3.4. Let ℓ_v denote the smallest integer $\geq CKv^2 \log\left(\frac{eN}{v}\right)$. Let $m_1 = m\ell_v$. Let $x_1, \dots, x_{m_1} \in \Omega$ be independent random points that are identically distributed according to μ . Let $\mathbf{x} = (x_1, \dots, x_{m_1})$. By Lemma 3.4, the inequalities

$$\frac{4}{5}\|f\|_p^p \leq \|S(f, \mathbf{x})\|_p^p \leq \frac{6}{5}\|f\|_p^p, \quad \forall f \in \Sigma_v(\Phi_N), \quad (3.7)$$

$$\text{and } \frac{4}{5}\|f\|_2^2 \leq \|S(f, \mathbf{x})\|_2^2 \leq \frac{6}{5}\|f\|_2^2, \quad \forall f \in \Sigma_v(\Phi_N) \quad (3.8)$$

hold simultaneously with probability

$$\geq 1 - 4 \exp\left(-\frac{cm_1}{Kv}\right) \geq 1 - 4e^{-mv}.$$

Next, we fix the random points x_1, \dots, x_{m_1} such that both (3.7) and (3.8) hold. Let $\{\Lambda_1, \dots, \Lambda_m\}$ be a partition of the set $[m_1]$ such that $|\Lambda_j| = \ell_v$ for $1 \leq j \leq m$. Let $n_j \in \Lambda_j$ denote a random variable that is uniformly distributed in the finite set Λ_j . Assume in addition that $x_1, \dots, x_{m_1}, n_1, \dots, n_m$ are independent. Define $\xi^j = x_{n_j}$ for $j = 1, 2, \dots, m$. It is easily seen that $\xi^1, \dots, \xi^m \in \Omega$ are independent random points identically distributed according to μ .

Now we consider the discrete $L_p^{m_1}$ -norm instead of the initial norm $\|\cdot\|_{L_p(\Omega)}$. Let

$$S(\Phi_N, \mathbf{x}) := \left\{ S(\varphi_i, \mathbf{x}) : 1 \leq i \leq N \right\} \subset L_2^{m_1}.$$

(3.7) and (3.8) imply that $S(\Phi_N, \mathbf{x})$ is a dictionary in $L_2^{m_1}$ satisfying (1.3) and (1.4) with constant $\frac{5}{4}K$ for the normalized counting measure on $[m_1]$. Moreover,

$$\|S(f, \mathbf{x})\|_\infty \leq \left(\frac{5}{4}Kv\right)^{1/p} \|S(f, \mathbf{x})\|_p, \quad \forall f \in \Sigma_v(\Phi_N).$$

Note that

$$\log m_1 \leq q_v := C \log(2Kv) + \log \log N.$$

Thus, the regular Nikolskii's inequality of the discrete norms $L_q^{m_1}$ implies

$$\|S(f, \mathbf{x})\|_\infty \leq e \cdot \|S(f, \mathbf{x})\|_{q_v}, \quad \forall f \in \Sigma_v(\Phi_N).$$

By Theorem 2.1 and Theorem 2.5 applied to the discrete norm of $L_p^{m_1}$, we conclude that for any integer

$$m \geq C_p K v (\log N) (\log(2vK))^2 (\log(2vK) + \log \log N),$$

the inequalities

$$\frac{3}{4} \|S(f, \mathbf{x})\|_p^p \leq \|S(f, \xi)\|_p^p \leq \frac{5}{4} \|S(f, \mathbf{x})\|_p^p, \quad \forall f \in \Sigma_v(\Phi_N) \quad (3.9)$$

hold with probability

$$\geq 1 - 2 \exp\left(-\frac{cm}{Kv \log^2(2Kv)}\right),$$

where

$$\xi = (\xi^1, \dots, \xi^m) = (x_{n_1}, x_{n_2}, \dots, x_{n_m}).$$

Combining (3.9) with (3.7), we obtain

$$\frac{3}{5} \|f\|_p^p \leq \|S(f, \xi)\|_p^p \leq \frac{3}{2} \|f\|_p^p, \quad \forall f \in \Sigma_v(\Phi_N). \quad (3.10)$$

Finally, we estimate the event E that (3.10) holds. Let E_1 denote the event that both (3.7) and (3.8) hold. The above argument then shows that $\mathbf{P}(E_1) \geq 1 - 4e^{-mv}$ and

$$\begin{aligned} \chi_{E_1} \cdot \mathbf{E}[\chi_E | x_1, \dots, x_{m_1}] &= \chi_{E_1} \cdot \mathbf{E}_{n_1, \dots, n_m}(\chi_E) \\ &\geq \chi_{E_1} \left(1 - 2 \exp\left(-\frac{cm}{Kv \log^2(2Kv)}\right)\right), \end{aligned}$$

where $\mathbf{E}_{n_1, \dots, n_m}(\chi_E) = \mathbf{E}[\chi_E | x_1, \dots, x_{m_1}]$ denotes the conditional expectation given x_1, \dots, x_{m_1} ; that is, the expectation computed with respect to the random variables n_1, \dots, n_m , fixing the variables x_1, \dots, x_{m_1} . Since χ_{E_1} is a function of $(x_1, x_2, \dots, x_{m_1})$, it follows that

$$\begin{aligned} \mathbf{P}(E) &\geq \mathbf{P}(E \cap E_1) = \mathbf{E}[\chi_{E \cap E_1}] = \mathbf{E}_{x_1, \dots, x_n}[\chi_{E_1} \cdot \mathbf{E}_{n_1, \dots, n_m}(\chi_E)] \\ &\geq \left(1 - 2 \exp\left(-\frac{cm}{Kv \log^2(2Kv)}\right)\right) \cdot \mathbf{E}[\chi_{E_1}] \\ &\geq \left(1 - 2 \exp\left(-\frac{cm}{Kv \log^2(2Kv)}\right)\right) (1 - 4e^{-cmv}) \\ &\geq 1 - 2 \exp\left(-\frac{c'm}{Kv \log^2(2Kv)}\right). \end{aligned}$$

This completes the proof.

4 Sparse sampling recovery

We begin this section with the observation that the universal discretization of the L_2 norm is closely connected with the concept of Restricted Isometry Property (RIP), which is very important in compressed sensing.

Let $U = \{\mathbf{u}^j\}_{j=1}^N$ be a system of column vectors from \mathbb{C}^m . We form a matrix $\mathbf{U} := [\mathbf{u}^1 \ \mathbf{u}^2 \ \dots \ \mathbf{u}^N]$ with vectors \mathbf{u}^j being the columns of this matrix. We say that matrix \mathbf{U} or the system U has the RIP property with parameters D and $\delta \in (0, 1)$ if for any subset J of indexes from $\{1, 2, \dots, N\}$ with cardinality $|J| \leq D$ we have for any vector $\mathbf{a} = (a_1, \dots, a_N) \in \mathbb{C}^N$ of coefficients such that $a_j = 0, j \notin J$ the following inequalities

$$(1 - \delta)\|\mathbf{a}\|_2^2 \leq \left\| \sum_{j \in J} a_j \mathbf{u}^j \right\|_2^2 \leq (1 + \delta)\|\mathbf{a}\|_2^2.$$

Let Ω be a compact subset of \mathbb{R}^d with the probability measure μ . For a dictionary $\mathcal{D}_N = \{g_i\}_{i=1}^N$ and a set of points $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ consider the system of vectors $G_N(\xi) = \{\bar{g}_i(\xi)\}_{i=1}^N$ where

$$\bar{g}_i(\xi) := m^{-1/2}(g_i(\xi^1), \dots, g_i(\xi^m))^T, \quad i = 1, \dots, N.$$

Suppose that \mathcal{D}_N is an orthonormal system and that the system $G_N(\xi)$ has the RIP property with parameters v and $\delta \in (0, 1)$. Then for any $f \in \Sigma_v(\mathcal{D}_N)$ we have on one hand

$$f = \sum_{i \in I} a_i g_i, \quad \|f\|_2^2 = \|\mathbf{a}\|_2^2, \quad \bar{f} := (f(\xi^1), \dots, f(\xi^m))^T = m^{1/2} \sum_{i \in I} a_i \bar{g}_i.$$

On the other hand

$$\left\| \sum_{i \in I} a_i \bar{g}_i \right\|_2^2 = \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^2.$$

It is clear that this means that the set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ provides *universal discretization* of the L_2 norm for the collection $\mathcal{X}_v(\mathcal{D}_N)$ with $C_1 = 1 - \delta$ and $C_2 = 1 + \delta$. The inverse statement is also true. Thus, in the case of orthonormal system \mathcal{D}_N the problem of finding a set ξ of points for universal discretization for the collection $\mathcal{X}_v(\mathcal{D}_N)$ and for the RIP property (with $D = v$) of the system $G_N(\xi)$ are closely related.

The reader can find results on the RIP properties of systems $G_N(\xi)$ associated with uniformly bounded orthonormal systems \mathcal{D}_N in the book [7]. For illustration we formulate one result from [7] (see Theorem 12.31 on page 405).

Theorem 4.1. *There exists a universal constant C with the following properties. Let \mathcal{D}_N be uniformly bounded orthonormal system such that $\|g_i\|_\infty \leq K$, $i = 1, 2, \dots, N$. Assume that ξ^1, \dots, ξ^m are independent random variables distributed on Ω according to μ . Then with probability at least $1 - N^{-(\ln N)^3}$ the system $G_N(\xi)$ has the RIP property with parameters s and δ provided*

$$m \geq CK^{-2}\delta^{-2}s(\ln N)^4. \quad (4.1)$$

In this section we discuss sampling recovery of periodic functions where the trigonometric system plays an important role. Let $M \in \mathbb{N}$ and $d \in \mathbb{N}$. Denote $\Pi(M) := [-M, M]^d$ to be the d -dimensional cube. Consider the system $\mathcal{T}(M, d) := \mathcal{T}(\Pi(M))$ of functions $e^{i(\mathbf{k}, \mathbf{x})}$, $\mathbf{k} \in \Pi(M)$ defined on $\mathbb{T}^d = [0, 2\pi)^d$. Then $\mathcal{T}(M, d)$ is an orthonormal system in $L_2(\mathbb{T}^d, \mu)$ with μ being the normalized Lebesgue measure on \mathbb{T}^d . The cardinality of this system is $N(M) := |\mathcal{T}(M, d)| = (2M + 1)^d$. In our further applications we are interested in bounds on $m(\mathcal{X}_v(\mathcal{T}(M, d)))$ in a special case, when $M \leq v^c$ with some constant c , which may depend on d . Then Theorem 1.1 gives

$$m(\mathcal{X}_v(\mathcal{T}(v^c, d))) \leq C(c, d)v(\log(2v))^4. \quad (4.2)$$

The RIP Theorem 4.1 and the above argument in the beginning of this section give (4.2) as well. It is known that in the case of system $\mathcal{T}(M, d)$ the bound (4.1) in Theorem 4.1 can be improved. This allows us to improve the bound (4.2). Namely, results of [8] and [2] combined with the above argument in the beginning of this section give

$$m(\mathcal{X}_v(\mathcal{T}(v^c, d))) \leq C(c, d)v(\log(2v))^3. \quad (4.3)$$

We recall some results from greedy approximation theory using our notations. For notational convenience we consider here a countable dictionary $\mathcal{D} = \{g_i\}_{i=1}^\infty$ of a Hilbert space H with the inner product $\langle \cdot, \cdot \rangle$ and the norm $\|x\| := \langle x, x \rangle^{1/2}$. We say a set \mathcal{D} of functions (elements) from H is a dictionary if each $g \in \mathcal{D}$ has a unit norm ($\|g\| = 1$) and the closure of $\text{span } \mathcal{D}$ is H . An element (function, signal) $f \in H$ is said to be v -sparse with respect to \mathcal{D} if it has a representation $f = \sum_{i \in T} x_i g_i$, $|T| \leq v$. The set of all v -sparse

elements is denoted by $\Sigma_v(\mathcal{D})$. For a given element f_0 we introduce the error of best v -term approximation

$$\sigma_v(f_0, \mathcal{D}) := \inf_{f \in \Sigma_v(\mathcal{D})} \|f_0 - f\|.$$

Let a sequence $\tau = \{t_k\}_{k=1}^\infty$, $0 \leq t_k \leq 1$, be given. The following greedy algorithm was defined in [12] under the name Weak Orthogonal Greedy Algorithm (WOGA).

Weak Orthogonal Matching Pursuit (WOMP). Let f_0 be given. Then for each $m \geq 1$ we inductively define:

- (1) $\psi_m \in \mathcal{D}$ is any element satisfying

$$|\langle f_{m-1}, \psi_m \rangle| \geq t_m \sup_{g \in \mathcal{D}} |\langle f_{m-1}, g \rangle|.$$

- (2) Let $H_m := \text{span}(\psi_1, \dots, \psi_m)$ and let $P_{H_m}(\cdot)$ denote an operator of orthogonal projection onto H_m . Define

$$G_m(f_0, \mathcal{D}) := P_{H_m}(f_0).$$

- (3) Define the residual after m th iteration of the algorithm

$$f_m := f_0 - G_m(f_0, \mathcal{D}).$$

In the case $t_k = 1$, $k = 1, 2, \dots$, WOMP is called the Orthogonal Matching Pursuit (OMP). In this paper we only consider the case $t_k = t$, $k = 1, 2, \dots$, $t \in (0, 1]$. The term *weak* in the definition of the WOMP means that at step (1) we do not shoot for the optimal element of the dictionary which realizes the corresponding supremum but we are satisfied with a weaker property than being optimal. The obvious reason for this is that we do not know in general that the optimal element exists. Another practical reason is that the weaker the assumption the easier it is to satisfy it and, therefore, easier to realize in practice. Clearly, ψ_m may not be unique. Our results apply for any realization (any choice of ψ_m) of the WOMP.

We now formulate a result from [11] (see also [16], Section 8.7). We use the following assumption.

UP(v, D). (v, D)-**unconditional property.** We say that a dictionary \mathcal{D} is (v, D)-unconditional with a constant U if for any v -sparse element $f = \sum_{i \in T} x_i g_i$, for any $A \subset T$, and any Λ , such that $A \cap \Lambda = \emptyset$ and $|A| + |\Lambda| \leq D$, we have for any $\{c_i\}$

$$\left\| \sum_{i \in A} x_i g_i - \sum_{i \in \Lambda} c_i g_i \right\| \geq U^{-1} \left\| \sum_{i \in A} x_i g_i \right\|. \quad (4.4)$$

Theorem 4.2 ([11, Corollary I.1]). *Let H be a Hilbert space. Suppose that the dictionary \mathcal{D} has property $\mathbf{UP}(v, D)$ with a constant U . Then for any $f_0 \in H$ the WOMP with weakness parameter t applied to f_0 provides*

$$\|f_{c(t,U)v}\| \leq C\sigma_v(f_0, \mathcal{D}) \quad \text{for } v + c(t, U)v \leq D$$

with an absolute constant C .

We now discuss sparse sampling recovery in the case Φ_N is a uniformly bounded Riesz basis satisfying (1.3) and

$$R_1 \left(\sum_{j=1}^N |a_j|^2 \right)^{1/2} \leq \left\| \sum_{j=1}^N a_j \varphi_j \right\|_2 \leq R_2 \left(\sum_{j=1}^N |a_j|^2 \right)^{1/2}, \quad (4.5)$$

where $0 < R_1 \leq R_2 < \infty$.

Proof of Theorem 1.2. We use the discretization properties (1.7) in order to establish property $\mathbf{UP}(v, D)$ for the dictionary $\mathcal{D} := \Phi_N(\Omega_m)$ being the restriction of Φ_N onto Ω_m . For any A and Λ such that $A \cap \Lambda = \emptyset$ the inequalities (4.5) imply

$$\left\| \sum_{i \in A} x_i \varphi_i - \sum_{i \in \Lambda} c_i \varphi_i \right\|_{L_2(\Omega, \mu)}^2 \geq R_1^2 \sum_{i \in A} |x_i|^2 \geq R_2^{-2} R_1^2 \left\| \sum_{i \in A} x_i \varphi_i \right\|_{L_2(\Omega, \mu)}^2.$$

This means that the system Φ_N has the $\mathbf{UP}(v, D)$ in the $L_2(\Omega, \mu)$ with any parameters $v \leq N$ and $v \leq D \leq N$ and the constant $U_1 = R_2/R_1$. Inequalities (1.7) imply that the discretized system $\Phi_N(\Omega_m)$ has the $\mathbf{UP}(u, N)$ in the $L_2(\Omega_m, \mu_m)$ with the constant $U_2 = U_1 3^{1/2}$.

We now define a dictionary \mathcal{D}_N in the $L_2(\Omega_m, \mu_m)$ as the normalized system $\Phi_N(\Omega_m)$. Clearly, property \mathbf{UP} does not depend on the normalization. We plan to apply the WOMP algorithm with the weakness parameter t . We now set $c = c(t, U_2)$ and for a given v set $u = v + c(t, U_2)v$ with the constant $c(t, U_2)$ from Theorem 4.2. It remains to apply Theorem 4.2. This proves (1.8) in Theorem 1.2.

We now derive (1.9) from (1.8). Clearly,

$$\sigma_v(f_0, \Phi_N(\Omega_m))_{L_2(\Omega_m, \mu_m)} \leq \sigma_v(f_0, \Phi_N)_\infty.$$

Let $f \in \Sigma_v(\Phi_N)$ be such that $\|f_0 - f\|_\infty \leq 2\sigma_v(f_0, \Phi_N)_\infty$. Then (1.8) implies

$$\|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega_m, \mu_m)} \leq \|f - f_0\|_{L_2(\Omega_m, \mu_m)} + \|f_{cv}\|_{L_2(\Omega_m, \mu_m)}$$

$$\leq (2 + C)\sigma_v(f_0, \Phi_N)_\infty.$$

Using that $f - G_{cv}(f_0, \Phi_N(\Omega_m)) \in \Sigma_u(\Phi_N)$, by discretization (1.7) we conclude that

$$\|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega, \mu)} \leq 2^{1/2}(2 + C)\sigma_v(f_0, \Phi_N)_\infty. \quad (4.6)$$

Finally,

$$\|f_{cv}\|_{L_2(\Omega, \mu)} \leq \|f - f_0\|_{L_2(\Omega, \mu)} + \|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega, \mu)}.$$

This and (4.6) prove (1.9). □

Theorem 1.2 guarantees that the simple greedy algorithm WOMP provides good sparse recovery. Namely, it gives the corresponding Lebesgue-type inequality in the norm $L_2(\Omega_m, \mu_m)$. We now discuss some results from [6], which provide good error of sparse sampling recovery in the norm $L_2(\Omega, \mu)$.

In addition to $\sigma_v(\mathbf{F}, \mathcal{D}_N)_X$ with X being a standard space $L_p(\Omega, \mu)$, $1 \leq p \leq \infty$, we also use the following characteristic, which is convenient for us. For $\xi = (\xi^1, \dots, \xi^m) \in \Omega^m$ let μ_ξ denote the probability measure

$$\mu_\xi := \frac{1}{2}\mu + \frac{1}{2m} \sum_{j=1}^m \delta_{\xi^j},$$

where $\delta_{\mathbf{x}}$ denotes the Dirac measure supported at a point \mathbf{x} .

We proved in [6] the conditional Theorem 4.3. Recall that given a finite dimensional subspace X of continuous functions on Ω , and a vector $\xi = (\xi^1, \dots, \xi^m) \in \Omega^m$, the classical least squares recovery operator (algorithm) (see, for instance, [3]) is defined as

$$LS(\xi, X_N)(f) := \arg \min_{u \in X} \|S(f - u, \xi)\|_2,$$

where $S(g, \xi) := (g(\xi^1), \dots, g(\xi^m))$, and

$$\|S(g, \xi)\|_2 := \left(\frac{1}{m} \sum_{\nu=1}^m |g(\xi^\nu)|^2 \right)^{1/2}.$$

With the help of the classical least squares algorithms, we define a new non-linear algorithm for a given collection $\mathcal{X} = \{X(n)\}_{n=1}^k$ of finite-dimensional subspaces of $\mathcal{C}(\Omega)$:

$$n(\xi, f) := \arg \min_{1 \leq n \leq k} \|f - LS(\xi, X(n))(f)\|_2,$$

$$LS(\xi, \mathcal{X})(f) := LS(\xi, X(n(\xi, f)))(f). \quad (4.7)$$

Definition 4.1. We say that a set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$ provides one-sided universal discretization with constant C_1 for the collection $\mathcal{X} := \{X(n)\}_{n=1}^k$ of finite-dimensional linear subspaces $X(n)$ if we have

$$C_1 \|f\|_2^2 \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^2 \quad \text{for any } f \in \bigcup_{n=1}^k X(n).$$

We denote by $m(\mathcal{X}, C_1)$ the minimal m such that there exists a set ξ of m points, which provides one-sided universal discretization with constant C_1 for the collection \mathcal{X} .

Theorem 4.3 ([6], Theorem 1.2). Let m, v, N be given natural numbers such that $v \leq N$. Let $\mathcal{D}_N \subset \mathcal{C}(\Omega)$ be a dictionary of N elements. Assume that there exists a set $\xi := \{\xi^j\}_{j=1}^m \subset \Omega$, which provides one-sided universal discretization with constant C_1 for the collection $\mathcal{X}_v(\mathcal{D}_N)$. Then for any function $f \in \mathcal{C}(\Omega)$ we have

$$\|f - LS(\xi, \mathcal{X}_v(\mathcal{D}_N))(f)\|_2 \leq 2^{1/2}(2C_1^{-1} + 1)\sigma_v(f, \mathcal{D}_N)_{L_2(\Omega, \mu_\xi)} \quad (4.8)$$

and

$$\|f - LS(\xi, \mathcal{X}_v(\mathcal{D}_N))(f)\|_2 \leq (2C_1^{-1} + 1)\sigma_v(f, \mathcal{D}_N)_\infty. \quad (4.9)$$

Comment 4.1. An advantage of the algorithm $LS(\xi, \mathcal{X}_v(\mathcal{D}_N))$ over the WOMP is that it provides in (4.8) and (4.9) the error in the $L_2(\Omega, \mu)$ norm and the WOMP provides in (1.8) the error in the discrete $L_2(\Omega_m, \mu_m)$ norm. However, a big advantage of the WOMP over $LS(\xi, \mathcal{X}_v(\mathcal{D}_N))$ is that it is a simple algorithm, which is known for easy practical implementation, and the $LS(\xi, \mathcal{X}_v(\mathcal{D}_N))$ is only a theoretical algorithm, which has the step (4.7), which is difficult to realize directly. Both algorithms WOMP and $LS(\xi, \mathcal{X}_v(\mathcal{D}_N))$ only use information from f and provide the error close to best (in the certain sense) error. They do not use the information that $f \in \mathbf{F}$ and automatically provide the error bound in terms of the best class \mathbf{F} , to which f belongs. This makes these algorithms *universal*.

Remark 4.1. In the same way as we derived (1.9) from (1.8) above in the proof of Theorem 1.2 we can prove the following inequality

$$\|f_{cv}\|_{L_2(\Omega, \mu)} \leq C'' \sigma_v(f, \Phi_N)_{L_2(\Omega, \mu_\xi)}. \quad (4.10)$$

Proof. It is clear that the space $L_2(\Omega_m, \mu_m)$ can be seen as the $L_2(\Omega, \mu_m)$ with

$$\mu_m = \frac{1}{m} \sum_{j=1}^m \delta_{\xi^j}.$$

Thus, we have $\mu_\xi = (\mu + \mu_m)/2$. We use discretization assumption (1.7) (we only need the left inequality in (1.7)) and the inequalities

$$\|h\|_{L_2(\Omega, \mu)} \leq 2^{1/2} \|h\|_{L_2(\Omega, \mu_\xi)},$$

$$\|h\|_{L_2(\Omega_m, \mu_m)} = \|h\|_{L_2(\Omega, \mu_m)} \leq 2^{1/2} \|h\|_{L_2(\Omega, \mu_\xi)}.$$

Then

$$\sigma_v(f_0, \Phi_N(\Omega_m))_{L_2(\Omega_m, \mu_m)} \leq 2^{1/2} \sigma_v(f_0, \Phi_N)_{L_2(\Omega, \mu_\xi)}.$$

Let $f \in \Sigma_v(\Phi_N)$ be such that $\|f_0 - f\|_{L_2(\Omega, \mu_\xi)} \leq 2\sigma_v(f_0, \Phi_N)_{L_2(\Omega, \mu_\xi)}$. Then (1.8) implies

$$\begin{aligned} \|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega_m, \mu_m)} &\leq \|f - f_0\|_{L_2(\Omega_m, \mu_m)} + \|f_{cv}\|_{L_2(\Omega_m, \mu_m)} \\ &\leq 2^{1/2}(2 + C)\sigma_v(f_0, \Phi_N)_{L_2(\Omega, \mu_\xi)}. \end{aligned}$$

Using that $f - G_{cv}(f_0, \Phi_N(\Omega_m)) \in \Sigma_u(\Phi_N)$, by discretization (1.7) we conclude that

$$\|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega, \mu)} \leq 2(2 + C)\sigma_v(f_0, \Phi_N)_{L_2(\Omega, \mu_\xi)}. \quad (4.11)$$

Finally,

$$\|f_{cv}\|_{L_2(\Omega, \mu)} \leq \|f - f_0\|_{L_2(\Omega, \mu)} + \|f - G_{cv}(f_0, \Phi_N(\Omega_m))\|_{L_2(\Omega, \mu)}.$$

This and (4.11) prove (4.10). □

We now discuss application of Remark 4.1 to optimal sampling recovery of periodic functions. This discussion complements the one from [6], Section 5. Let $\mathbf{s} = (s_1, \dots, s_d)$ be a vector whose coordinates are nonnegative integers

$$\rho(\mathbf{s}) := \{\mathbf{k} \in \mathbb{Z}^d : [2^{s_j-1}] \leq |k_j| < 2^{s_j}, \quad j = 1, \dots, d\},$$

$$\Gamma(N) := \{\mathbf{k} \in \mathbb{Z}^d : \prod_{j=1}^d \max(|k_j|, 1) \leq N\} \quad - \quad \text{a hyperbolic cross.}$$

For $f \in L_1(\mathbb{T}^d)$

$$\delta_{\mathbf{s}}(f, \mathbf{x}) := \sum_{\mathbf{k} \in \rho(\mathbf{s})} \hat{f}(\mathbf{k}) e^{i(\mathbf{k}, \mathbf{x})}, \quad \hat{f}(\mathbf{k}) := (2\pi)^{-d} \int_{\mathbb{T}^d} f(\mathbf{x}) e^{-i(\mathbf{k}, \mathbf{x})} d\mathbf{x}.$$

Let Q be a finite set of points in \mathbb{Z}^d , we denote

$$\mathcal{T}(Q) := \left\{ t : t(\mathbf{x}) = \sum_{\mathbf{k} \in Q} c_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})} \right\}.$$

Consider the Wiener norm (the A -norm or the ℓ_1 -norm)

$$\|f\|_A := \sum_{\mathbf{k}} |\hat{f}(\mathbf{k})|.$$

The following classes, which are convenient in studying sparse approximation with respect to the trigonometric system, were introduced and studied in [15]. Define for $f \in L_1(\mathbb{T}^d)$

$$f_j := \sum_{\|\mathbf{s}\|_1=j} \delta_{\mathbf{s}}(f), \quad j \in \mathbb{N}_0, \quad \mathbb{N}_0 := \mathbb{N} \cup \{0\}.$$

For parameters $a \in \mathbb{R}_+$, $b \in \mathbb{R}$ define the class

$$\mathbf{W}_A^{a,b} := \{f : \|f_j\|_A \leq 2^{-aj} (\bar{j})^{(d-1)b}, \quad \bar{j} := \max(j, 1), \quad j \in \mathbb{N}\}.$$

The following result is from [6]. It is given there without proof. For completeness we present at the end of this section a proof of a somewhat more general statement.

Lemma 4.1. *There exist two constants $c(a, d)$ and $C(a, b, d)$ such that for any $\xi \in (\mathbb{T}^d)^m$ and $v \in \mathbb{N}$ there is a constructive method $A_{v, \xi}$ based on greedy algorithms, which provides a v -term approximant from $\mathcal{T}(\Gamma(M))$, $|\Gamma(M)| \leq v^{c(a, d)}$, with the bound for $f \in \mathbf{W}_A^{a,b}$*

$$\|f - A_{v, \xi}(f)\|_{L_2(\mathbb{T}^d, \mu_{\xi})} \leq C(a, b, d) v^{-a-1/2} (\log v)^{(d-1)(a+b)}.$$

Let $v \in \mathbb{N}$ be given and let M be from Lemma 4.1. Consider the orthonormal basis $\{e^{i(\mathbf{k}, \mathbf{x})}\}_{\mathbf{k} \in \Gamma(M)}$ of the $\mathcal{T}(\Gamma(M))$ to be the system Φ_N . Then Lemma 4.1, Remark 4.1 and the bound (4.3) imply the following Theorem 4.4.

Theorem 4.4. *There exist two constants $c'(a, d)$ and $C'(a, b, d)$ such that for any $v \in \mathbb{N}$ we have the bound for $a > 0, b \in \mathbb{R}$,*

$$\varrho_m^o(\mathbf{W}_A^{a,b}, L_2(\mathbb{T}^d)) \leq C'(a, b, d)v^{-a-1/2}(\log v)^{(d-1)(a+b)} \quad (4.12)$$

for any m satisfying

$$m \geq c'(a, d)v(\log(2v))^3.$$

Theorem 4.4 gives the following bound for the classical classes \mathbf{W}_p^r of functions with bounded mixed derivative (see [16], p.130, for their definition and [6] for some recovery results).

Theorem 4.5. *There exist two constants $c'(r, d, p)$ and $C'(r, d, p)$ such that for any $v \in \mathbb{N}$ we have the bound for $r > 1/p, 1 < p < 2$,*

$$\varrho_m^o(\mathbf{W}_p^r, L_2(\mathbb{T}^d)) \leq C'(r, d, p)v^{-r+1/p-1/2}(\log v)^{(d-1)(r+1-2/p)} \quad (4.13)$$

for any m satisfying

$$m \geq c'(r, d, p)v(\log(2v))^3.$$

The authors of [9] (see Corollary 4.16 in v2) proved the following interesting bound for $1 < p < 2, r > 1/p$ for $m \geq c(r, d, p)v(\log(2v))^3$

$$\varrho_m^o(\mathbf{W}_p^r, L_2(\mathbb{T}^d)) \leq C(r, d, p)v^{-r+1/p-1/2}(\log v)^{(d-1)(r+1-2/p)+1/2} \quad (4.14)$$

The bound (4.13) is better than (4.14) by the $(\log v)^{-1/2}$ factor. The proof of (4.14) in [9] is based on known results from [15] on sparse trigonometric approximation in the uniform norm. We proved (4.13) by using inequality (4.10) and estimating the sparse trigonometric approximation in the norm $L_2(\mathbb{T}^d, \mu_\xi)$, which is weaker than the uniform norm.

Let $\mathcal{D} = \{g_{\mathbf{k}}\}_{\mathbf{k} \in \mathbb{Z}^d}$ be a system of uniformly bounded functions

$$|g_{\mathbf{k}}(\mathbf{x})| \leq 1, \quad \mathbf{x} \in \Omega, \quad \mathbf{k} \in \mathbb{Z}^d. \quad (4.15)$$

For

$$f = \sum_{\mathbf{k}} a_{\mathbf{k}} g_{\mathbf{k}} \quad (4.16)$$

denote

$$\delta_{\mathbf{s}}(f, \mathcal{D}) := \sum_{\mathbf{k} \in \rho(\mathbf{s})} a_{\mathbf{k}} g_{\mathbf{k}}, \quad f_j := \sum_{\|\mathbf{s}\|_1=j} \delta_{\mathbf{s}}(f, \mathcal{D}), \quad j \in \mathbb{N}_0, \quad \mathbb{N}_0 := \mathbb{N} \cup \{0\}$$

and

$$\|f\|_{A(\mathcal{D})} := \sum_{\mathbf{k}} |a_{\mathbf{k}}|.$$

For parameters $a \in \mathbb{R}_+$, $b \in \mathbb{R}$ define the class $\mathbf{W}_A^{a,b}(\mathcal{D})$ of functions f for which there exists a representation (4.16) satisfying

$$\|f_j\|_{A(\mathcal{D})} \leq 2^{-aj}(\bar{j})^{(d-1)b}, \quad \bar{j} := \max(j, 1), \quad j \in \mathbb{N}. \quad (4.17)$$

Let Q be a finite set of points in \mathbb{Z}^d , we denote

$$\mathcal{D}(Q) := \left\{ t : t(\mathbf{x}) = \sum_{\mathbf{k} \in Q} a_{\mathbf{k}} g_{\mathbf{k}} \right\}.$$

Lemma 4.2. *There exist two constants $c(a, d)$ and $C(a, b, d)$ such that for any $\xi \in \Omega^m$ and $v \in \mathbb{N}$ there is a constructive method $A_{v,\xi}$ based on greedy algorithms, which provides a v -term approximant from $\mathcal{D}(\Gamma(M))$, $|\Gamma(M)| \leq v^{c(a,d)}$, with the bound for $f \in \mathbf{W}_A^{a,b}(\mathcal{D})$*

$$\|f - A_{v,\xi}(f)\|_{L_2(\Omega, \mu_\xi)} \leq C(a, b, d) v^{-a-1/2} (\log v)^{(d-1)(a+b)}.$$

Proof. We prove the lemma for $v \asymp 2^n n^{d-1}$, $n \in \mathbb{N}$. Let $f \in \mathbf{W}_A^{a,b}(\mathcal{D})$ and

$$f = \sum_{\mathbf{k}} a_{\mathbf{k}} g_{\mathbf{k}}$$

be its representation satisfying (4.17). We approximate f_j in $L_2(\Omega, \mu_\xi)$ by a v_j -term approximant from $\mathcal{D}(Q_j)$, $\mathcal{D}(Q_j) := \cup_{\|\mathbf{s}\|_1=j} \rho(\mathbf{s})$. The sequence $\{v_j\}$ will be defined later. Since $L_2(\Omega, \mu_\xi)$ is a Hilbert space and by (4.15)

$$\|g_{\mathbf{k}}\|_{L_2(\Omega, \mu_\xi)} \leq \sup_{\mathbf{x} \in \Omega} |g_{\mathbf{k}}(\mathbf{x})| \leq 1, \quad \mathbf{k} \in Q_j, \quad (4.18)$$

using Theorem 2.19 of [13, p. 93], we deduce from (4.17) that for every j there exists an $h_j \in \Sigma_{v_j}(\mathcal{D}(Q_j))$ (provided by a greedy algorithm – Orthogonal Matching Pursuit) such that

$$\|f_j - h_j\|_{L_2(\Omega, \mu_\xi)} \leq v_j^{-1/2} 2^{-aj} (\bar{j})^{(d-1)b}.$$

We take $\beta \in (0, a)$ (for instance, $\beta = a/2$) and specify

$$v_j := \lceil 2^{n-\beta(j-n)} j^{d-1} \rceil, \quad j = n, n+1, \dots$$

It is clear that there exists $j(\beta, d)$ such that $v_j = 0$ for $j \geq j(\beta, d)n$. For $j \geq j(\beta, d)n$ we set $h_j = 0$. In addition to $\{h_j\}_{j=n}^{j(\beta, d)n}$ we include in the approximant

$$S_n(f, \mathcal{D}) := \sum_{\|\mathbf{s}\|_1 < n} \delta_{\mathbf{s}}(f, \mathcal{D}).$$

Define

$$A_v(f, \beta) := S_n(f, \mathcal{D}) + \sum_{j \geq n} h_j = S_n(f, \mathcal{D}) + \sum_{j=n}^{j(\beta, d)n} h_j.$$

Then, we have built a v -term approximant of f with

$$v \ll 2^n n^{d-1} + \sum_{j \geq n} v_j \ll 2^n n^{d-1}.$$

The error of this approximation in $L_2(\Omega, \mu_\xi)$ is bounded from above by

$$\begin{aligned} \|f - A_m(f, p, \beta)\|_{L_2(\Omega, \mu_\xi)} &\leq \sum_{j \geq n} \|f_j - h_j\|_{L_2(\Omega, \mu_\xi)} \ll \sum_{j \geq n} (\bar{v}_j)^{-1/2} 2^{-aj} j^{(d-1)b} \\ &\ll \sum_{j \geq n} 2^{-1/2(n-\beta(j-n))} j^{-(d-1)/2} 2^{-aj} j^{(d-1)b} \ll 2^{-n(a+1/2)} n^{(d-1)(b-1/2)}. \end{aligned}$$

This completes the proof of lemma. □

5 A lower bound for universal discretization

In this section we discuss one result from [4] (see Theorem 7.4 there) on universal discretization of the L_2 norm of trigonometric polynomials. For $n \in \mathbb{N}$ denote $\Pi_n := \Pi(\mathbf{N}) \cap \mathbb{Z}^d$ with $\mathbf{N} = (2^{n-1} - 1, \dots, 2^{n-1} - 1)$, where,

$$\Pi(\mathbf{N}) := [-N_1, N_1] \times \dots \times [-N_d, N_d].$$

Let $v \in \mathbb{N}$ and $v \leq |\Pi_n|$. Consider

$$\mathcal{S}(v, n) := \{Q \subset \Pi_n : |Q| = v\}.$$

Let

$$\mathcal{T}(Q) := \{f : f = \sum_{\mathbf{k} \in Q} c_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})}, \quad c_{\mathbf{k}} \in \mathbb{C}\}.$$

The following Theorem 5.1 was proved in [4] (see Theorem 7.4 there).

Theorem 5.1. *There exist three positive constants $C_i(d)$, $i = 1, 2, 3$, such that for any $n, v \in \mathbb{N}$ and $v \leq |\Pi_n|$ there is a set $\xi = \{\xi^\nu\}_{\nu=1}^m \subset \mathbb{T}^d$, with $m \leq C_1(d)v^2n$, which provides universal discretization in L_2 for the collection $\mathcal{S}(v, n)$: For any $f \in \cup_{Q \in \mathcal{S}(v, n)} \mathcal{T}(Q)$*

$$C_2(d)\|f\|_2^2 \leq \frac{1}{m} \sum_{\nu=1}^m |f(\xi^\nu)|^2 \leq C_3(d)\|f\|_2^2.$$

Claim 5.1. We prove in this section that the factor n in the bound $m \leq C_1(d)v^2n$ in Theorem 5.1 cannot be replaced by a slower growing in n factor.

We begin with a direct corollary of Theorem 5.1. For $N \in \mathbb{N}$ consider the system $\mathcal{T}_N^s := \{\sin(\pi kx)\}_{k=1}^N$ defined on $[0, 1]$. It is easy to derive from Theorem 5.1 that there exist two positive absolute constants C_i , $i = 1, 2$, such that for any $N \in \mathbb{N}$ there is a set $\xi = \{\xi^\nu\}_{\nu=1}^m \subset [0, 1]$, with $m \leq C_1 \log(N+1)$, which provides the following discretization in L_2 : For any $g \in \mathcal{T}_N^s$ we have

$$C_2\|g\|_2^2 \leq \frac{1}{m} \sum_{\nu=1}^m |g(\xi^\nu)|^2. \quad (5.1)$$

We now show that the replacement of the bound $m \leq C_1 \log(N+1)$ by $m = o(\log N)$ brings a contradiction. For that we use the well known generalization of the Dirichlet Theorem on diophantine approximation: For a given set $\xi = \{\xi^\nu\}_{\nu=1}^m \subset [0, 1]$ there exist a natural number $k \leq N$ and integer numbers a_1, \dots, a_m such that

$$|k\xi^\nu - a_\nu| \leq N^{-1/m}, \quad \nu = 1, \dots, m. \quad (5.2)$$

Then, (5.1) with $g(x) = \sin(\pi kx)$ implies

$$C_2/2 \leq \pi^2 N^{-2/m},$$

which gives a contradiction if $m = o(\log N)$.

6 A more general version of the conditional theorem

Our goal in this section is to prove the following conditional theorem, which is the random version of Theorem 5.1 of [5].

Theorem 6.1. *Let $\mathcal{W} \subset L_\infty(\Omega)$ and let $\mathcal{W}_p := \{f \in \mathcal{W} : \|f\|_p \leq 1\}$ for some $1 \leq p < \infty$. Assume that $\lambda \cdot \mathcal{W} = \mathcal{W}$ for any $\lambda > 0$, and \mathcal{W}_p is a compact subset of L_∞ . Let*

$$R := \sup_{f \in \mathcal{W}_p} \sup_{x \in \Omega} |f(x)|.$$

Let ξ^1, \dots, ξ^m be independent random points on Ω satisfying

$$\frac{1}{m} \sum_{k=1}^m \mu_{\xi^k} = \mu, \quad (6.1)$$

where μ_{ξ^j} denotes the probability distribution of ξ^j . Then there exist positive constants C_p, c_p depending only on p such that for any $\varepsilon \in (0, 1)$ and any integer

$$m \geq C_p \varepsilon^{-5} \left(\int_{c_p \varepsilon^{1/p}}^R u^{\frac{p}{2}-1} \left(\int_u^R \frac{\mathcal{H}_{c_p \varepsilon t}(\mathcal{W}_p, L_\infty)}{t} dt \right)^{\frac{1}{2}} du \right)^2, \quad (6.2)$$

the inequalities,

$$(1 - \varepsilon) \|f\|_p^p \leq \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p \leq (1 + \varepsilon) \|f\|_p^p, \quad \forall f \in \mathcal{W}_p, \quad (6.3)$$

hold with probability

$$\geq 1 - \varepsilon \exp\left(-\frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}\right).$$

It can be easily seen that either of the following two conditions implies the condition (6.1):

- (i) ξ^1, \dots, ξ^m are identically distributed according to μ ;
- (ii) there exists a partition $\{\Omega_1, \dots, \Omega_m\}$ of Ω such that $\mu(\Omega_j) = \frac{1}{m}$ and $\xi^j \in \Omega_j$ is distributed according to $m \cdot \mu \Big|_{\Omega_j}$ for each $1 \leq j \leq m$.

The proof of Theorem 6.1 follows along the same line as that of Theorem 5.1 of [5]. We sketch it as follows.

Proof of Theorem 6.1. Let

$$\mathcal{S}_p := \{f/\|f\|_p : f \in \mathcal{W}, \|f\|_p > 0\} \subset \mathcal{W}_p.$$

Let $c^* = c_p^* \in (0, \frac{1}{2})$ be a sufficiently small constant depending only on p . Let $a := c^* \varepsilon$. Let J, j_0 be two integers such that $j_0 < 0 \leq J$,

$$(1+a)^{J-1} \leq R < (1+a)^J \quad \text{and} \quad (1+a)^{j_0 p} \leq \frac{1}{10} \varepsilon \leq (1+a)^{(j_0+1)p}. \quad (6.4)$$

For $j \in \mathbb{Z}$, let

$$\mathcal{A}_j := \mathcal{N}_{2a(1+a)^j}(\mathcal{S}_p, L_\infty) \subset \mathcal{S}_p$$

denote the minimal $2a(1+a)^j$ -net of \mathcal{S}_p in the norm of L_∞ . For $f \in \mathcal{S}_p$, we define $A_j(f)$ to be the function in \mathcal{A}_j such that $\|A_j(f) - f\|_\infty \leq 2a(1+a)^j$.

For $f \in \mathcal{S}_p$ and $j > j_0$, let $U_j(f) := \{\mathbf{x} \in \Omega : |A_j(f)(\mathbf{x})| \geq (1+a)^{j-1}\}$, and $D_j(f) := U_j(f) \setminus \bigcup_{k>j} U_k(f)$. Define

$$h(f, \mathbf{x}) := \sum_{j=j_0+1}^J (1+a)^j \chi_{D_j(f)}(\mathbf{x}). \quad (6.5)$$

It can be easily verified that

$$(1 - \frac{\varepsilon}{8})|f(\mathbf{x})|^p \leq |h(f, \mathbf{x})|^p \leq (1 + \frac{\varepsilon}{8})|f(\mathbf{x})|^p, \quad (6.6)$$

if $\mathbf{x} \in \bigcup_{j_0 < j \leq J} D_j(f)$, and $|f(\mathbf{x})|^p \leq \frac{\varepsilon}{8}$ otherwise¹. As a result, for any probability measure ν on Ω , and any $f \in \mathcal{S}_p$, we have

$$\left| \|h(f)\|_{L_p(\nu)}^p - \|f\|_{L_p(\nu)}^p \right| \leq \frac{\varepsilon}{8} \|f\|_{L_p(\nu)}^p + \frac{\varepsilon}{8}. \quad (6.7)$$

Now for $j_0 + 1 \leq j \leq J$, let

$$\mathcal{F}_j^p := \{M_j \cdot \chi_{D_j(f)} : f \in \mathcal{S}_p\}, \quad \text{where } M_j := (1+a)^{pj}.$$

Let $\{\varepsilon_j\}_{j=j_0+1}^J$ be a sequence of positive numbers satisfying $\sum_{j=j_0+1}^J \varepsilon_j \leq \varepsilon/4$, which will be specified later. By (6.6), for each $j_0 < j \leq J$,

$$\|M_j \chi_{D_j(f)}\|_1 \leq \|h(f)\|_p^p \leq (1 + \frac{\varepsilon}{8}) \|f\|_p^p \leq 2.$$

¹For details, we refer to the proof of Theorem 5.1 in [5].

Thus, using (6.1) and Lemma 3.2, we conclude that the inequalities ,

$$\sup_{\varphi \in \mathcal{F}_j^p} \left| \frac{1}{m} \sum_{k=1}^m \varphi(\xi^k) - \int_{\Omega} \varphi(x) d\mu(x) \right| \leq \varepsilon_j, \quad j_0 < j \leq J, \quad (6.8)$$

hold simultaneously with probability

$$\geq 1 - \sum_{j=j_0+1}^J |\mathcal{F}_j^p| \exp\left(-\frac{m\varepsilon_j^2}{16M_j}\right). \quad (6.9)$$

Now we fix a set of random points ξ^1, \dots, ξ^m for which (6.8) is satisfied. Using (6.5), (6.8) and the triangle inequality, we then deduce

$$\sup_{f \in \mathcal{S}_p} \left| \frac{1}{m} \sum_{j=1}^m |h(f, \xi^j)|^p - \|h(f)\|_p^p \right| \leq \sum_{j=j_0+1}^J \varepsilon_j < \frac{\varepsilon}{4}.$$

This together with (6.7) implies that for any $f \in \mathcal{S}_p$,

$$\left| \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p - 1 \right| \leq \frac{3}{8}\varepsilon + \frac{\varepsilon}{8} \cdot \frac{1}{m} \sum_{j=1}^m |f(\xi^j)|^p,$$

from which the desired inequality (6.3) follows.

Thus, to complete the proof, it remains to construct a sequence of positive numbers $\{\varepsilon_j\}_{j=j_0+1}^J$ such that

$$\sum_{j=j_0+1}^J \varepsilon_j \leq \varepsilon/4, \quad (6.10)$$

and

$$\sum_{j=j_0+1}^J |\mathcal{F}_j^p| \exp\left(-\frac{m\varepsilon_j^2}{16M_j}\right) \leq C\varepsilon \exp\left(-\frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}\right). \quad (6.11)$$

Since $\mathcal{S}_p \subset \mathcal{W}_p$, we can estimate the cardinalities of the sets \mathcal{F}_j^p as follows:

$$|\mathcal{F}_j^p| \leq |\mathcal{A}_j| \times \dots \times |\mathcal{A}_J| \leq \prod_{k=j}^J N_{a(1+a)^k}(\mathcal{W}_p, L_\infty) =: L_j. \quad (6.12)$$

A straightforward calculation shows that

$$\log L_j = \sum_{k=j}^J \mathcal{H}_{a(1+a)^k}(\mathcal{W}_p, L_\infty) \leq C\varepsilon^{-1} \int_{(1+a)^{j-1}}^R \mathcal{H}_{at}(\mathcal{W}_p, L_\infty) \frac{dt}{t}. \quad (6.13)$$

Now we specify the numbers ε_j as follows:

$$\varepsilon_j := \delta + \tau_j, \quad j_0 < j \leq J,$$

where $\delta := \frac{\varepsilon}{8(J-j_0)}$, and $\tau_j > 0$ is defined by

$$2 \log L_j = \frac{m\tau_j^2}{16M_j}, \quad \text{that is } \tau_j := 4\sqrt{2} \sqrt{\frac{M_j \cdot \log L_j}{m}}. \quad (6.14)$$

We have

$$\sum_{j=j_0+1}^J \varepsilon_j = 4\sqrt{2}m^{-1/2} \sum_{j=j_0+1}^J (M_j \log L_j)^{\frac{1}{2}} + \frac{\varepsilon}{8},$$

which, by (6.13) and straightforward calculations, is estimated above by

$$\leq \frac{\varepsilon}{8} + Cm^{-\frac{1}{2}}\varepsilon^{-\frac{3}{2}} \int_{4^{-1}(10^{-1}\varepsilon)^{1/p}}^R u^{\frac{p}{2}-1} \left(\int_u^R \mathcal{H}_{c_p \varepsilon t}(\mathcal{W}_p, L_\infty) \frac{dt}{t} \right)^{\frac{1}{2}} du.$$

Thus, to ensure (6.10), it is enough to assume that

$$m \geq C\varepsilon^{-5} \left[\int_{4^{-1}(10^{-1}\varepsilon)^{1/p}}^R u^{\frac{p}{2}-1} \left(\int_u^R \mathcal{H}_{c_p \varepsilon t}(\mathcal{W}_p, L_\infty) \frac{dt}{t} \right)^{\frac{1}{2}} du \right]^2$$

with $C = C_p$ being a large constant depending only on p .

Finally, we estimate the sum

$$S := \sum_{j=j_0+1}^J |\mathcal{F}_j^p| \exp\left(-\frac{m\varepsilon_j^2}{16M_j}\right).$$

Since $\varepsilon_j^2 \geq \tau_j^2 + \delta^2$, we obtain from (6.12) that

$$S \leq \sum_{j=j_0+1}^J \exp\left(\log L_j - \frac{m\tau_j^2}{16M_j} - \frac{m\delta^2}{16M_j}\right) \leq \exp\left(-\frac{m\delta^2}{16M_J}\right) \sum_{j=j_0+1}^J \frac{1}{L_j}.$$

Using (6.4), we have

$$\frac{m\delta^2}{16M_J} = m \cdot \left(\frac{\varepsilon}{8(J-j_0)} \right)^2 \cdot \frac{1}{16M_J} \geq \frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}.$$

Since

$$L_j \geq N_{a(1+a)^j}(\mathcal{W}_p, L_\infty),$$

this implies

$$S \leq \exp\left(-\frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}\right) \sum_{j=j_0+1}^J \frac{1}{N_{a(1+a)^j}(\mathcal{W}_p, L_\infty)}.$$

On the other hand, from the proof of Theorem 5.1 in [5], we know that

$$N_t(\mathcal{W}_p, L_\infty) \geq \frac{R}{4t}, \quad \forall 0 < t < R. \quad (6.15)$$

It follows that

$$S \leq \exp\left(-\frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}\right) \cdot 4c_*R^{-1}\varepsilon \sum_{j=j_0+1}^J (1+a)^j \leq \varepsilon \exp\left(-\frac{cm\varepsilon^4}{R^p(\log \frac{R}{\varepsilon})^2}\right),$$

which is as desired. This completes the proof. \square

Acknowledgement. The authors thank Tino Ullrich for bringing to their attention the paper [8].

References

- [1] J. Bourgain, J. Lindenstrauss and V. Milman, Approximation of zonoids by zonotopes, *Acta Math.*, **162** (1989), 73–141.
- [2] J. Bourgain, An improved estimate in the restricted isometry problem, In *Geometric Aspects of Functional Analysis*, volume 2116 of *Lecture Notes in Mathematics*, pages 65–70. Springer, 2014.
- [3] A. Cohen and G. Migliorati, Optimal weighted least-squares methods, *SMAI J. Computational Mathematics* **3** (2017), 181–203.

- [4] F. Dai, A. Prymak, V.N. Temlyakov, and S.U. Tikhonov, Integral norm discretization and related problems, *Russian Math. Surveys* **74:4** (2019), 579–630. Translation from *Uspekhi Mat. Nauk* **74:4(448)** (2019), 3–58; arXiv:1807.01353v1.
- [5] F. Dai and V. Temlyakov, Universal sampling discretization, arXiv:2107.11476v1 [math.FA] 23 Jul 2021.
- [6] Dai and V. Temlyakov, Universal discretization and sparse sampling recovery, arXiv:2301.05962v1 [math.NA] 14 Jan 2023.
- [7] S. Foucart and H. Rauhut, *A Mathematical Introduction to Compressive Sensing*, Birkhäuser, 2013.
- [8] I. Haviv and O. Regev, The restricted isometry property of subsampled Fourier matrices, In *Geometric aspects of functional analysis*, volume 2169 of *Lecture Notes in Math.*, pages 163–179. Springer, Cham, 2017.
- [9] T. Jahn, T. Ullrich, and F. Voigtlaender, Sampling numbers of smoothness classes via ℓ^1 -minimization, arXiv:2212.00445v1 [math.NA] 1 Dec 2022.
- [10] B. Kashin, E. Kosov, I. Limonova, and V. Temlyakov, Sampling discretization and related problems, *Journal of Complexity*, **71** (2022), 101653; arXiv:2109.07567v1 [math.FA] 15 Sep 2021.
- [11] E. Livshitz and V. Temlyakov, Sparse approximation and recovery by greedy algorithms, *IEEE Transactions on Information Theory*, **60** (2014), 3989–4000; arXiv: 1303.3595v1 [math.NA] 14 Mar 2013.
Constructive Approximation (Advanced Problems), Springer, 1996.
- [12] V.N. Temlyakov, Weak greedy algorithms, *Adv. Comput. Math.* **12** (2000), 213–227.
- [13] V.N. Temlyakov, *Greedy Approximation*, Cambridge University Press, 2011.
- [14] V.N. Temlyakov, An inequality for the entropy numbers and its application, *J. Approx. Theory*, **173** (2013), 110–121.

- [15] V.N. Temlyakov, Constructive sparse trigonometric approximation and other problems for functions with mixed smoothness, arXiv: 1412.8647v1 [math.NA] 24 Dec 2014, 1–37; *Matem. Sb.*, **206** (2015), 131–160.
- [16] V. Temlyakov, *Multivariate Approximation*, Cambridge University Press, 2018.

F. Dai, DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES
UNIVERSITY OF ALBERTA
EDMONTON, ALBERTA T6G 2G1, CANADA
E-MAIL: fdai@ualberta.ca

V.N. Temlyakov, UNIVERSITY OF SOUTH CAROLINA,
STEKLOV INSTITUTE OF MATHEMATICS,
LOMONOSOV MOSCOW STATE UNIVERSITY,
AND MOSCOW CENTER FOR FUNDAMENTAL AND APPLIED MATHEMATICS.
E-MAIL: temlyak@math.sc.edu