

**ON VARIETIES OF LIE ALGEBRAS
WITH INFINITE BASIS PROPERTY**

VESSELIN S. DRENSKY

ABSTRACT. Over an arbitrary field of positive characteristic we construct an example of a locally finite variety of Lie algebras which does not have a finite basis of its polynomial identities. As a consequence we construct varieties of Lie algebras with prescribed properties.

We shall use the notation in [1] and [2]. In particular, all products are left normed: $xyz = (xy)z$ and $xy^n = \underbrace{xy \cdots y}_n$. In [1] and [2] M. R. Vaughan-Lee and the author constructed examples of infinitely based varieties of Lie algebras over a field of positive characteristic. A similar example constructed also Yu. G. Kleiman (unpublished). In this regard, Yu. A. Bahturin raised the question about the existence of locally finite infinitely based varieties of algebras. Quite recently I. B. Volichenko [3] constructed such an example in the case of a field of characteristic 2. Using the ideas from [1] and [2] we give the answer in the case of an arbitrary field of positive characteristic.

Theorem 1. *Let K be a field of positive characteristic p . Then the variety of Lie algebras over K defined by the identities*

$$(1) \quad (x_1x_2)(x_3x_4) \cdots (x_{2p-1}x_{2p})x_{2p+1} = 0,$$

$$(2) \quad ((x_1x_2)(x_3x_4))((x_5x_6)(x_7x_8)) = 0,$$

$$(3) \quad x_1x_2^{p^2+2} = 0,$$

$$(4) \quad (x_1x_2 \cdots x_{k+2})(x_1x_2)^{p-1} = 0, \quad k = 1, 2, \dots,$$

is locally finite and infinitely based.

We shall prove the theorem in several steps. We shall follow the exposition of [2, §2].

1. Let T_n be the set of all subsets of the set $\{1, 2, \dots, 2n-1\}$ and let A_n be an algebra with basis over K

$$\{a_{\sigma s}^{(1)}, a_{\tau s}^{(k)}, b_s \mid \sigma, \tau \in T_n, |\tau| \text{ is odd}, k = 2, 3, \dots, p-1, s = 0, 1, \dots, p\}.$$

For all $\tau \in T_n$, where $|\tau|$ is odd, we set $a_{\tau s}^{(p)} = b_s$ if $|\tau| = 2n-1$ and $a_{\tau s}^{(p)} = 0$ if $|\tau| < 2n-1$. We define the multiplication in A_n in the following way:

Published as Vesselin S. Drensky, Infinitely based varieties of Lie algebras (Russian), *Serdica* 9 (1983), No. 1, 79-82. Zbl 0525.17006, MR0725813.
Available online at <http://www.math.bas.bg/serdica/1983/1983-079-082.pdf>.

$$a_{\sigma s}^{(k)} a_{\tau t}^{(l)} = (-1)^{|\sigma|} a_{\pi s+t}^{(k+l)},$$

where $\pi = \sigma \cup \tau$, if $k+l \leq p$, $s+t \leq p$, $\sigma \cap \tau = \emptyset$, $|\sigma| + |\tau| \equiv 1 \pmod{2}$,

$$a_{\sigma s}^{(k)} a_{\tau t}^{(l)} = 0 \text{ in all other cases.}$$

Let us consider the set of linear transformations of A_n

$$(5) \quad \{g_{\lambda s}, h_{\mu s} \mid \lambda, \mu \in T_n, |\lambda| \text{ is odd, } |\mu| \text{ is even, } s = 0, 1, \dots, p\},$$

defined by the following equalities:

$$a_{\sigma s}^{(k)} g_{\lambda t} = a_{\pi s+t}^{(k)}, \text{ where } \pi = \sigma \cup \lambda, \text{ if } s+t \leq p, \sigma \cap \lambda = \emptyset, |\sigma| \text{ is odd,}$$

$$a_{\sigma s}^{(k)} h_{\mu t} = k a_{\pi s+t}^{(k)}, \text{ where } \pi = \sigma \cup \mu, \text{ if } s+t \leq p, \sigma \cap \mu = \emptyset,$$

$$a_{\sigma s}^{(k)} g_{\lambda t} = a_{\sigma s}^{(k)} h_{\mu t} = 0 \text{ in all other cases.}$$

As in [2] we can prove that the algebra A_n is a Lie algebra in the variety $\mathfrak{A}^2 \cap \mathfrak{N}_p$ and the transformations (5) are derivations of the algebra A_n which pairwise commute. Let us denote by B_n the split extension of the algebra A_n by the abelian algebra of derivations D_n generated by the transformations (5). Following [2] it is easy to show that B_n satisfies the identities (1) and (2).

2. Let us denote by C_n the subalgebra of B_n generated by the set

$$\{a_{\sigma 0}^{(1)}, g_{\lambda 0}, h_{\sigma 1}, h_{\mu 0} \mid \sigma = \emptyset, \lambda = \{1\}, \mu = \{2, 3\}, \{4, 5\}, \dots, \{2n-2, 2n-1\}\}.$$

The algebra C_n has a basis as a K -vector space

$$(6) \quad \{a_{\sigma s}^{(k)}, g_{\lambda 0}, h_{\mu 0}, h_{\nu 1} \mid \sigma \in T_n \text{ with } a_{\sigma s}^{(k)} \in C_n, k = 1, 2, \dots, p, s = 0, 1, \dots, p, \\ \lambda = \{1\}, \mu = \{2, 3\}, \{4, 5\}, \dots, \{2n-2, 2n-1\}, \nu = \emptyset\}.$$

We shall prove that C_n is nilpotent of class $2p+n$ and satisfies the identity (3):

In any nonzero product $l_1 l_2 \cdots l_m$ of elements in (6) the elements $a_{\sigma s}^{(k)}$ appear not more than p times, the element $h_{\nu 1}$ appears not more than p times and every element $g_{\lambda 0}, h_{\mu 0}$ appears not more than once. Hence $m \leq 2p+n$ and $C_n \in \mathfrak{N}_{2p+n}$.

Let us assume that $m \geq p^2 + 2$. We shall prove that $x_1 x_2^m = 0$ in C_n . Let $l_1, l_2 \in C_n$, $l_2 = a + d$, where $a \in A_n$, $d \in D_n$. Then $l_1 l_2 = a_1 \in A_n$. We shall unwrap parentheses in the product $l_1 l_2^m = (l_1 l_2) l_2^{m-1} = a_1 (a + d)^{m-1}$:

$$(7) \quad l_1 l_2^m = \sum a_1 c_1 c_2 \cdots c_{m-1},$$

where $c_i = a$ or $c_i = d$, $i = 1, 2, \dots, m-1$. If $a_1 c_1 c_2 \cdots c_{m-1} \neq 0$, then a appears among c_1, c_2, \dots, c_{m-1} not more than $p-1$ times; if a appears exactly $p-1$ times, then $c_{m-1} = a$ (because $a_{\sigma t}^{(p)}$ belongs to the center of C_n). Therefore the nonzero products in the sum (7) are of the form

$$(8) \quad a_1 d^{k_1} a d^{k_2} a \cdots a d^{k_s} a^\varepsilon, \quad s \leq p-1, \varepsilon = 0, 1,$$

and $k_1 + k_2 + \cdots + k_s + (s-1) + \varepsilon = m-1$. Hence

$$(9) \quad m = k_1 + k_2 + \cdots + k_s + s + \varepsilon \leq k_1 + k_2 + \cdots + k_s + p.$$

The element d is a linear combination of derivations in (6), i.e. $d = \sum \alpha_i d_i$, $\alpha_i \in K$, and the d_i 's commute pairwise. Then ($\text{char} K = p$)

$$(10) \quad \left(\text{ad} \sum \alpha_i d_i \right)^p = \sum \alpha_i^p (\text{ad} d_i)^p,$$

$$(11) \quad (\text{ad} d_i)^p = \begin{cases} 0, & \text{if } d_i = g_{\lambda 0}, h_{\mu 0}, \mu \neq \emptyset, \\ \text{ad} h_{\nu p}, & \text{if } d_i = h_{\nu 1}, \nu = \emptyset, \end{cases}$$

because $a_{\sigma s}^{(k)} h_{\nu 1}^p = k^p a_{\sigma s+p}^{(k)} = k a_{\sigma s+p}^{(k)} = a_{\sigma s}^{(k)} h_{\nu p}$, $\nu = \emptyset$. Let us consider the product $a_1 d^{k_1} \cdots a d^{k_s} a^\varepsilon$ in (8). Let

$$(12) \quad k_i = \max(k_1, \dots, k_s), \quad k_j = \max(\{k_1, \dots, k_s\} \setminus \{k_i\}).$$

If $k_i \leq 2p - 1$, $k_j \leq p - 1$, then ($s \leq p - 1$ from (12))

$$k_1 + \cdots + k_s \leq (2p - 1) + (s - 1)(p - 1) \leq (2p - 1) + (p - 2)(p - 1) = p^2 - p + 1$$

and by (9) $m \leq p^2 + 1$ which is impossible. Hence there are two cases: $k_j \geq p$ or $k_i \geq 2p$. For convenience of the notation we assume that $i = 1$, $j = 2$ (with the same considerations in the general case).

(i) Let $k_2 \geq p$. Then, in virtue of (10) and (11),

$$\begin{aligned} a_1 d^{k_1} a d^{k_2} &= a_1 d^p d^{k_1-p} a d^p d^{k_2-p} = a_1 \left(\sum \alpha_i d_i \right)^p d^{k_1-p} a \left(\sum \alpha_i d_i \right)^p d^{k_2-p} \\ &= a_1 \left(\sum \alpha_i^p \text{ad}^p d_i \right) d^{k_1-p} a \left(\sum \alpha_i^p \text{ad}^p d_i \right) d^{k_2-p} = \alpha^{2p} a_1 h_{\nu 1}^p d^{k_1-p} a h_{\nu 1}^p d^{k_2-p} \\ &= \alpha^{2p} a_1 h_{\nu p} d^{k_1-p} a h_{\nu p} d^{k_2-p} = 0, \quad \nu = \emptyset. \end{aligned}$$

(Here α is the coefficient of $d_{i_0} = h_{\nu 1}$ in $\sum \alpha_i d_i$.) Hence in this case the product (8) is equal to zero.

(ii) $k_1 \geq 2p$. Then it follows from (10) and (11) that

$$\begin{aligned} a_1 d^{k_1} &= a_1 d^p d^p d^{k_1-2p} = a_1 \left(\sum \alpha_i d_i \right)^p \left(\sum \alpha_i d_i \right)^p d^{k_1-2p} \\ &= a_1 \left(\sum \alpha_i^p \text{ad}^p d_i \right)^2 d^{k_1-2p} = \alpha^{2p} a_1 \text{ad}^{2p} h_{\nu 1} d^{k_1-2p} = \alpha^{2p} a_1 h_{\nu p}^2 d^{k_1-2p} = 0, \quad \nu = \emptyset. \end{aligned}$$

Again, the product (8) is equal to zero. Therefore, the algebra C_n satisfies the identity (3).

3. The variety defined by the identities (1), (2), and (3) is solvable and satisfies the Engel identity. By the theorem of K. W. Gruenberg [4] it is locally nilpotent and hence locally finite.

4. As in [2, Lemmas 2.1 and 2.2 and Theorem 2.3] we can prove that C_n satisfies all identities (4) for $k < n$ and $(x_1 x_2 \cdots x_{n+2})(x_1 x_2)^{p-1} \neq 0$ in C_n . Since C_n is nilpotent of class $2p + n$ it satisfies also (4) for $k > n$, i.e., the identities (4) are independent modulo the identities (1), (2), and (3). This completes the proof of the theorem.

Using ideas of A. Yu. Ol'shanskii [6] and [5] and Theorem 1 we can construct some examples of varieties of Lie algebras with prescribed properties.

Theorem 2. *Over any field of positive characteristic there exists a continuum of locally finite varieties of Lie algebras which form a strictly increasing chain with respect to the inclusion.*

Proof. We order the set \mathbb{Q} of the rational numbers:

$$(13) \quad r_1, r_2, r_3, \dots,$$

and consider the set of Lie algebras $\{L_r \mid r \in \mathbb{Q}\}$. Here $L_r \cong C_m$, where m is the number of r in (13) and C_m is the algebra from the proof of Theorem 1. For any real α we define the variety

$$(14) \quad \mathfrak{W}_\alpha = \text{var}\{L_r \mid r \leq \alpha\}.$$

It follows from Part 4 of the proof of Theorem 1 that if $\alpha < \beta$, then $\mathfrak{W}_\alpha \subsetneq \mathfrak{W}_\beta$. \square

Corollary 3. *Over any field of positive characteristic there exists a continuum of locally finite varieties of Lie algebras with pairwise different ordinal functions. (The ordinal function $f_{\mathfrak{V}} : \mathbb{N} \rightarrow \mathbb{N}$ of the locally finite variety \mathfrak{V} is defined by the equality $f_{\mathfrak{V}}(n) = \dim F_n(\mathfrak{V})$, $n = 1, 2, \dots$, where $F_n(\mathfrak{V})$ is the relatively free algebra of rank n in \mathfrak{V} .)*

Proof. Let us consider the varieties (14). If $\alpha > \beta$ then

$$(15) \quad \mathfrak{V}_{\beta} \subset \mathfrak{V}_{\alpha}$$

and for every n there is a canonical epimorphism $\psi_n : F_n(\mathfrak{V}_{\alpha}) \rightarrow F_n(\mathfrak{V}_{\beta})$. Since the inclusion (15) is strict, there is an n such that $\ker \psi_n \neq 0$, which implies that $f_{\mathfrak{V}_{\alpha}}(n) \neq f_{\mathfrak{V}_{\beta}}(n)$. \square

Remark 4. As in [6] we can prove that over a field of characteristic $p > 0$ there exist two different varieties with the same ordinal functions. For example, let $k = p^3 + p^2 + p + 2$, $\mathfrak{M} = \mathfrak{A}^2 \cap \mathfrak{N}_k$. We consider the subvarieties \mathfrak{V} and \mathfrak{W} of \mathfrak{M} defined, respectively, by the identities

$$v(x_1, x_2, y_0, y_1, \dots, y_p, y_{p+1}) = x_1 x_2 y_0^p y_1^p \cdots y_p^p y_{p+1}^p = 0,$$

$$w(x_1, x_2, y_0, y_1, \dots, y_p, y_{p+1}) = x_1 x_2 y_0^{p^2} y_1^{p^2} \cdots y_p^{p^2} y_{p+1}^p = 0.$$

Then $f_{\mathfrak{V}}(n) = f_{\mathfrak{W}}(n)$.

Remark 5. Using the ideas from his paper [7] Yu. N. Mal'tsev (unpublished) constructed an example of two just non-commutative varieties of associative algebras over a finite field with the same ordinal functions. Using the ideas from [6] we can construct the following simple example. Let K be a field of characteristic $p > 0$ and let $k = p^2 + p + 1$. Define the variety \mathfrak{M} of associative algebras $xy - yx = x_1 x_2 \cdots x_{k+1} = 0$ and its subvarieties \mathfrak{V} and \mathfrak{W} defined, respectively by the identities

$$v(x_0, x_1, \dots, x_p, y) = x_0 x_1 \cdots x_p y^{p^2} = 0, \quad w(x_0, x_1, \dots, x_p, y) = (x_0 x_1 \cdots x_p)^p y = 0.$$

Then $\mathfrak{V} \neq \mathfrak{W}$ but \mathfrak{V} and \mathfrak{W} have the same ordinal functions.

The results of this paper are included in the Ph.D. thesis of the author [8]. The author expresses his sincere gratitude to Yu. A. Bahturin for the stating of the problem and attention to the work.

REFERENCES

- [1] M. R. Vaughan-Lee, Varieties of Lie algebras, Quart. J. Math. Oxford (2), 21 (1970), 297-308.
- [2] V. S. Drensky, Identities in Lie algebras (Russian), Algebra i Logika 13 (1974), 265-290. Translation: Algebra and Logic 13 (1974), 150-165.
- [3] I. B. Volichenko, On varieties of center-by-metabelian Lie algebras, Preprint No. 16, 1980, Institute of Mathematics of National Academy of Sciences of Belarus, Minsk.
- [4] K. W. Gruenberg, Two theorems on Engel groups, Proc. Cambridge Philos. Soc. 49 (1953), 377-380.
- [5] A. Yu. Ol'shanskii, On some infinite systems of identities (Russian), Tr. Semin. Im. I. G. Petrovskogo 3 (1978), 139-146. Translation: Transl., Ser. 2, Am. Math. Soc. 119 (1983), 81-88.
- [6] A. Yu. Ol'shanskii, On the orders of free groups of locally finite varieties (Russian), Izv. Akad. Nauk SSSR, Ser. Mat. 37 (1973), 89-94. Translation: Math. USSR, Izv. 7 (1973), 85-90.
- [7] Yu. N. Mal'tsev, Almost commutative varieties of associative rings (Russian), Sib. Mat. Zh. 17 (1976), 1086-1096. Translation: Sib. Math. J. 17 (1976), 803-811.
- [8] V. S. Drensky, Solvable Varieties of Lie Algebras (Russian), Ph. D. thesis, Dept. of Mechanics and Math., Moscow State Univ., Moscow, 1979.