

Endomorphisms of a certain ring of rational functions

by Milo Moses

January 31, 2023

Contents

1	Introduction	1
2	General Methods	2
3	Endomorphisms of $\mathcal{M}_{\mathbb{Q}}$	8
4	Endomorphisms of \mathcal{M}_K, K/\mathbb{Q} algebraic	14

1 Introduction

For any field K , define the ring \mathcal{M}_K as follows:

$$\left(\begin{array}{l} \text{elements: ratios of monic polynomials in } K(x) \\ \text{addition: multiplication of rational functions} \\ \text{multiplication: } \left[\prod_i (x - \alpha_i) \right] \cdot \left[\prod_j (x - \beta_j) \right] = \left[\prod_{i,j} (x - \alpha_i \beta_j) \right] \end{array} \right)$$

This sort of ring can naturally arise in the study of vector spaces. Endomorphisms of vector spaces have associated characteristic polynomials, and the direct sum/tensor product of endomorphisms corresponds to addition/multiplication in \mathcal{M}_K . We will put a topology on \mathcal{M}_K by letting Cauchy sequences be those in which the leading coefficients converge. We will later show that it makes \mathcal{M}_K into a topological ring. The following is the main theorem proved in this text:

Theorem 1.1. *Let K/\mathbb{Q} be an algebraic extension, and let $\phi : \mathcal{M}_K \rightarrow \mathcal{M}_K$ be a continuous ring endomorphism. Then, either*

1. *There exist integers $k, s \geq 0$ and $\sigma \in \text{Gal}(K/\mathbb{Q})$ such that*

$$\phi \left(\prod_i (x - \alpha_i) \right) = x^{\sum_i \alpha_i^s - 1} \cdot \prod_i (x - \sigma(\alpha_i)^k)$$

2. All roots of rational functions in the image of ϕ are roots of unity. If K/\mathbb{Q} is finite, then there exists N such that all the roots are N th roots of unity.

The constant N appearing in part (2) of Theorem 1.1 is only effective when $K = \mathbb{Q}$. In that case one can easily compute N given (say,) $\phi([x - 2])$ using the methods of the proof. This can be turned into precise results, such as $N < (\# \text{ roots of } \phi([x - 2]))^{1.8}$.¹

2 General Methods

In this section, we will establish some general results about \mathcal{M}_K , with K being an arbitrary field. Once convenient, we will specialize to $K \subseteq \mathbb{C}$. Our main tool will be the ring morphisms $t_k : \mathcal{M}_K \rightarrow K$ defined on elements of the form $[x - \alpha]$ by

$$t_k([x - \alpha]) = \alpha^k,$$

and extended to general elements by linearity. Note that we use the convention $0^0 = 1$, so t_0 is the degree map. A first property of these maps is that they can be used to compute polynomial coefficients:

Theorem 2.1 (Newton's Formulas). *For any polynomial $f \in K[x]$ of degree d and $k \leq d$, we have that*

$$ke_k(f) = \sum_{n=1}^k (-1)^{n+1} e_{k-n}(f) t_n([f]).$$

Here, $e_n(f)$ are the elementary symmetric polynomial coefficients defined implicitly by

$$f(x) = \sum_{n=0}^d (-1)^{d-n} e_{d-n}(f) x^n.$$

Proof. These are classical. □

Corollary 2.1. *If K has characteristic 0, then for any monic polynomials $f, g \in K[x]$, $e_k(f) = e_k(g)$ for all $k < N$ if and only if $t_k([f]) = t_k([g])$ for all $k < N$.*

Proof. This is immediate from the Theorem. □

In particular, we have that the t_k define an injection

$$\mathcal{M}_K \xrightarrow{\prod_{k=0}^{\infty} t_k} \prod_{k=0}^{\infty} K.$$

¹The result here is not proved explicitly, but it is a simple corollary of our methods. The constant 1.8 is not optimal, and can be readily improved to $1/\ln(2) + \epsilon$

This allows us to define the topology on \mathcal{M}_K as the subspace topology, with $\prod_{k=1}^{\infty} K$ given the product topology, and K always given the discrete topology. It is clear that convergence in \mathcal{M}_K is equivalent to the convergence of the t_k , which in turn is equivalent to the convergence of the e_k , which are the leading coefficients of a polynomial. We will now show to what extent t_k are the only continuous ring morphisms $\mathcal{M}_K \rightarrow K$.

Lemma 2.1. *If S is a subring of K^N for some natural number N , then the nonzero morphisms in $\text{Hom}(S, K)$ are projections of S onto a single index, followed by composition with some ring morphism $K \rightarrow K$.*

Proof. We proceed by induction on N . If $N = 1$, then S is a subring of K . Thus, the map $S \rightarrow K$ is automatically projection of S onto a single index followed by a morphism. Additionally, it is standard application of Zorn's lemma that every map $S \rightarrow K$ can be lifted to a map $K \rightarrow K$.

For our inductive step, suppose the result is proved for $N - 1$. Fix a map $\phi : S \rightarrow \mathbb{Z}$. It is clear that ϕ doesn't factor through the projection $L \rightarrow L'$ forgetting the 1st index if and only if there are two elements which are equal outside of the 1st index, but give different elements under ϕ . Equivalently, this means that there is a nonzero element $(a_1, 0, 0\dots)$ on which ϕ does not act by 0.

If ϕ factors through the projection forgetting the 1st or 2nd index, then the induction hypothesis gives us the conclusion. Suppose it does not factor through forgetting either. This means there are elements $(a_1, 0, 0\dots)$ and $(0, a_2, 0\dots)$ under which ϕ does not act by 0. However, these elements multiply to 0. This implies there exist nonzero elements of K which multiply to 0, which is a contradiction. This completes the proof. \square

Proposition 2.1. *The only continuous ring morphisms $\phi : \mathcal{M}_K \rightarrow K$ are of the form $\sigma \circ t_k$, where $\sigma : K \rightarrow K$ is a ring morphism.*

Proof. Since K is discrete, the kernel of ϕ contains an open neighborhood of 0. The ideals

$$I_N = \{r \text{ st } t_k(r) = 0 \forall k \geq N\}$$

form a basis of open neighborhoods of 0 under the topology for \mathcal{M}_K . Hence, we have that $I_N \in \ker \phi$ for some N . Hence, any continuous map $\mathcal{M}_K \rightarrow K$ factors through projection onto \mathcal{M}_K/I_N . This is a subring of K^N , hence Lemma 2.1 gives the desired result. \square

This is extremely powerful, as it lets us define the following:

Definition 2.1. For any continuous endomorphisms $\phi : \mathcal{M}_K \rightarrow \mathcal{M}_K$, we define the functions $\phi^* : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$ and $\sigma^\phi : \mathbb{Z}_{\geq 0} \rightarrow (\text{morphisms } K \rightarrow K)$ to be the unique values such that

$$t_n(\phi(r)) = \sigma_n^\phi(t_{\phi^*(n)}(r))$$

for all $r \in \mathcal{M}_K$.

We now define the most important endomorphisms of \mathcal{M}_K :

Proposition 2.2. *For every $n \geq 1$, the map*

$$\lambda_n : \mathcal{M}_K \rightarrow \mathcal{M}_K$$

sending $\prod_i (x - \alpha_i)$ to $\prod_i (x - \alpha_i^n)$ is a continuous endomorphism of \mathcal{M}_K . We have that $\lambda_k^(n) = kn$ for all k, n , and $\sigma_n^{\lambda_k} = 1$.*

Proof. This is a straightforward computation. \square

The following formula will underpin our use of cyclotomic methods throughout the proof.

Corollary 2.2. *If K has characteristic 0, then for any $f \in \overline{K}[x]$,*

$$x^d f(1/x) = \prod_{n \geq 1} (1 - x^n)^{c_n}$$

where

$$c_n = \frac{1}{n} \sum_{d|n} \mu(n/d) t_d([f]).$$

The quantities $(1+x)^z$ are defined using Newton's generalized binomial theorem.

Proof. This is the so-called Cyclotomic identity, and can be found in many references. \square

Let $\mathcal{M}_{\mathbb{Z}}$ be the subring of $\mathcal{M}_{\mathbb{Q}}$ consisting of rational functions with integer coefficients.

Proposition 2.3. *For any $r \in \mathcal{M}_{\mathbb{Z}}$ and $n \in \mathbb{Z}_{\geq 1}$,*

$$\frac{1}{n} \sum_{d|n} \mu(n/d) t_d(r) \in \mathbb{Z}.$$

Proof. We prove the result for $r = [f]$, $f \in \mathbb{Z}[x]$ monic, and extend to a general element by linearity. Let N be the smallest number for which $c_N = \frac{1}{N} \sum_{d|N} \mu(n/d) t_d(r)$ is not in \mathbb{Z} . Now, we look at the coefficient of x^N on both sides of

$$x^d f(1/x) = \prod_{n \geq 1} (1 - x^n)^{c_n}.$$

The left hand side's coefficient is an element of \mathbb{Z} by assumption. On the right hand side, terms with $n > N$ do not contribute since they are too large. By assumption, all terms with $n < N$ contribute an element of \mathbb{Z} . Thus, the final answer is an element of \mathbb{Z} if and only if the coefficient of x^N in $(1 - x^N)^{c_N}$ is in \mathbb{Z} . The coefficient is exactly $-c_N$ and hence $c_N \in \mathbb{Z}$ which is a contradiction. This yields the result. \square

The following lemmas are necessary to continue:

Proposition 2.4. *Let s_k be any sequence of complex numbers given explicitly by*

$$s_k = \sum_j c_j \theta_j^k$$

where j runs over a finite set of values, c_j are complex, and θ_j are on the unit circle. There are infinitely many $q \geq 1$ such that

$$|s_k - s_{k+q}| \leq \frac{4\pi}{q^{1/J}} \sum_i |c_i|$$

for every $k \geq 0$, where J is the number of values j runs over. For every k_0 , s_{k_0} is a limit point of s_k . In particular, if s_k converges then it must be constant. The only way for s_k to be constant is for the coefficients of all roots other than 1 to be zero.

Proof. To begin, we assume without loss of generality that all the θ_j are distinct. Let $\theta_j = e^{2\pi i x_j}$. By Dirichlet's Simultaneous Approximation theorem, we can find infinitely many values q for which there exists p_j such that

$$\left| x_j - \frac{p_j}{q} \right| \leq \frac{1}{q^{1+1/J}}.$$

Using the estimate $|e^x - 1| \leq 2|x|$ for $|x| < 1$, we get

$$|\exp(2\pi i q x_j) - 1| \leq \frac{4\pi}{q^{1/J}}.$$

Choosing any $k \geq 1$ and summing over j , this immediately gives

$$|s_k - s_{k+q}| \leq \frac{4\pi}{q^{1/J}} \sum_i |c_i|.$$

Since this tends to 0 as q gets large, it is clear that every value of s_k must be a limit point.

Now, suppose that s_k is a constant. By subtracting off the coefficient of 1, we can assume $s_k = 0$ uniformly. The average value of $|s_k|^2$ is $\sum_j |c_j|^2$, and hence we must have $c_j = 0$ for all j , as desired. \square

Sometimes the following will also be useful:

Proposition 2.5. *If $r \in \mathcal{M}_{\mathbb{C}}$ is such that $t_k(r)$ goes over finitely many values, then all roots of r are roots of unity.*

Proof. If r has any roots greater than 1, then Proposition 2.4 gives that $t_k(r)$ tends to infinity. Hence, we must have that all roots of r are in absolute value less than or equal to 1. We thus have

$$r = \prod_i (x - \theta_i)^{c_i} \cdot r',$$

where θ_i are distinct complex numbers of absolute value 1, and all roots of r' are less than 1. Letting $E < 1$ be the largest root of r' , we get that

$$t_k(r) = S_k + O(E^k)$$

where $S_k = \sum_i c_i \cdot \theta_i^k$, and O denotes big-Oh notation. Choose q large so that $|S_{k+q} - S_k|$ is uniformly small, which is possible by Proposition 2.4. Then, we would have that

$$t_{k+q}(r) - t_k(r)$$

is uniformly small as q gets large, for large enough k . However, since $t_k(r)$ goes over finitely values, this must imply that $t_k(r) = t_{k+q}(r)$ for all large enough k . In particular, we must have that $S_{k+q} - S_k$ grows small at an exponentially small rate. Proposition 2.4 says this means that S_k is uniformly zero. Hence, we get that $\theta_i^q = 1$ for all q .

Now, consider the roots of r smaller than 1. Since S_k is periodic, we must have that the t_k of the smaller roots must go over finitely many values. However, the t_k of the smaller roots also tend to zero. The only way to go over finitely many values and tend to zero is if the t_k is zero for all large enough roots. The only way to be zero for large enough roots is if there are no roots of absolute value less than 1, by repeating the same as above. \square

The above work motivates the following definition:

Proposition 2.6. *The following conditions on an element $r \in \mathcal{M}_{\mathbb{Z}}$ are equivalent:*

1. $r = [f/g]$, where the roots of f and g are all of absolute value 1 or 0
2. $r = [f/g]$, where the roots of f and g are all of absolute value ≤ 1
3. $\lim_{k \rightarrow \infty} t_k(r) / (1 + \epsilon)^k = 0$ for all $\epsilon > 0$
4. $r = c_0[x] + \sum_{n \geq 1} c_n [1 - x^{-n}]$, where $c_n \in \mathbb{Z}$ are constants only finitely of which are non-zero

The subring of elements $r \in \mathcal{M}_{\mathbb{Z}}$ with any (all) of these equivalent properties is called $\mathcal{M}_{\mathbb{Z}, \Phi}$.

Proof. We begin by showing that 2 \implies 1. To do this, we note that the smallest non-zero coefficient of f is equal to the product of its non-zero roots. This product must be a non-zero integer, and since none of the roots are larger than 1 they must all be of absolute value 1.

Now we show that 3 \implies 2. Let C be the absolute value of the largest root of r . If $C \leq 1$, we are done. If $C > 1$, write

$$t_n(r) = S_n \cdot C^n + O(E^n)$$

where $S_n = \sum_i c_i \cdot \theta_i^n$, $|\theta_i| = 1$, and $E < C$. Then,

$$S_n = \frac{t_n(r)}{C^n} + O((E/C)^n),$$

so $S_n \rightarrow 0$. Thus S_n meets the conditions of Lemma 2.4, and hence is zero constant. This contradicts the fact that there is at least one root of absolute value C , since otherwise the sum would be non-trivial.

Now we show that 4 \implies 3. To do this, we note that

$$t_k \left(c_0[x] + \sum_{n \geq 1} c_n [1 - x^{-n}] \right)$$

is periodic and hence tends to zero when divided by arbitrarily large values.

Finally we show that 1 \implies 4. We induct on $\deg(f) + \deg(g)$. If $\deg(f) \geq 1$ with root ζ , then ζ is an algebraic number all of whose Galois conjugates are also of absolute value one and hence must be a primitive root of unity of degree n , $\zeta = \zeta_n$, by a classical theorem of Kronecker. Thus, $\Phi_n(x) \mid \deg(f)$ where Φ_n is the n th cyclotomic polynomial and hence

$$[f/g] = [(f/\Phi_n(x))/g] + \sum_{d \mid n} \mu(n/d) [x^d - 1].$$

Continuing until the only roots of f and g are all zero, we are done. \square

We define $\widehat{\mathcal{M}}_K$ to be the topological completion of \mathcal{M}_K .

Proposition 2.7. *The subring $\mathcal{M}_{\mathbb{Z}, \Phi}$ is dense in $\mathcal{M}_{\mathbb{Z}}$. Every element $r \in \widehat{\mathcal{M}}_{\mathbb{Z}, \Phi} = \widehat{\mathcal{M}}_{\mathbb{Z}}$ can be written uniquely in the form*

$$r = c_0[x] + \sum_{n \geq 1} c_n [1 - x^{-n}],$$

for arbitrary c_n . Moreover, for $k \in \mathbb{Z}_{\geq 1}$

$$t_k(r) = \sum_{d \mid k} c_d$$

and $t_0(r) = c_0$.

Proof. To begin, we show that every element of the form $r = c_0[x] + \sum_{n \geq 1} c_n [1 - x^{-n}]$ is in $\widehat{\mathcal{M}_{\mathbb{Z}, \Phi}}$. That follows from the fact that

$$r = \lim_{N \rightarrow \infty} \left(c_0[x] + \sum_{n=1}^N c_n [1 - x^{-n}] \right),$$

where each term in the limit is in $\mathcal{M}_{\mathbb{Z}, \Phi}$ by Proposition 2.6 part 3. Next, we show that these sums follow the t_k formula stated. Namely, we begin by checking

$$\begin{aligned} t_k([x^n - 1]) &= \sum_{j=0}^{n-1} e^{2\pi j i/k} \\ &= \begin{cases} n & n|k \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Hence, dividing through by x^n , we get the same formula for $t_k([1 - x^{-n}])$ except that it has no contribution at 0. Extending by linearity we get the formula. Uniqueness follows from Möbius inversion.

For the last part of the proposition, we check that every element of $\widehat{\mathcal{M}_{\mathbb{Z}}}$ really can be written in this form. let r_N be a sequence in $\mathcal{M}_{\mathbb{Z}}$ with r as a limit point. Let

$$c_n = \lim_{N \rightarrow \infty} \frac{1}{n} \sum_{d|n} \mu(n/d) t_d(r_N),$$

which converges since t_k are continuous. Moreover, $c_n \in \mathbb{Z}$ since by Proposition 2.3 they are at every step in the limit. It is now clear that $r = c_0[x] + \sum_{n \geq 1} c_n [1 - x^{-n}]$ by Möbius inversion. \square

We now prove an important multiplication formula on $\mathcal{M}_{\mathbb{Z}, \Phi}$:

Proposition 2.8. *For any n, m it holds that*

$$[x^n - 1][x^m - 1] = \gcd(n, m) [x^{\text{lcm}(n, m)} - 1]$$

Proof. This is a trivial computation that follows directly from the definition of multiplication in $\mathcal{M}_{\mathbb{Z}}$. \square

3 Endomorphisms of $\mathcal{M}_{\mathbb{Q}}$

In this section, we study $\text{End}(\mathcal{M}_{\mathbb{Q}})$. A natural place to start is to study maps on the smaller domain $\mathcal{M}_{\mathbb{Z}, \Phi}$:

Proposition 3.1. *If $\phi \in \mathcal{M}_{\mathbb{Z}, \Phi} \rightarrow \mathcal{M}_{\mathbb{C}}$, is continuous then for all rational prime powers p^α we have that*

$$\phi\left([x^{p^\alpha} - 1]\right) = s_{-\infty}[x] + \sum_{\beta=0}^{\alpha} s_\beta \cdot p^{\alpha-\beta} \cdot [x^{p^\beta} - 1],$$

where $s_\beta \in \{-1, 0, 1\}$.

Proof. Seeing as $t_k(\phi(r))$ must go over finitely values, Proposition 2.5 gives us that $\phi(r)$ lies in $\mathcal{M}_{\mathbb{Z}, \Phi}$. Let $r = \phi([x^{p^\alpha} - 1])$, and let $c_n = \frac{1}{n} \sum_{d|n} \mu(n/d) t_d(r)$. Choose N such that $c_N \neq 0$. Write $N = p^\gamma N'$, with p not dividing N' . We begin by showing that $\gamma \leq \alpha$. Namely we compute

$$\begin{aligned} p^\gamma N' \leq p^\gamma N' |c_N| &= \left| \sum_{d|p^\gamma N'} \mu(p^\gamma N'/d) t_d(r) \right| \\ &= \left| \sum_{d|N'} \mu(N'/d) (t_{p^\gamma d}(r) - t_{p^{\gamma-1}d}(r)) \right| \\ &\leq \sum_{d|N'} p^\alpha \leq p^\alpha N'. \end{aligned}$$

Hence, $\gamma \leq \alpha$ as desired. Note that in the case $\gamma = 0$ the formula written is not strictly correct (i.e, there is not $p^{\gamma-1}d$ correction term), but since $\alpha \geq 1$ our proposition is true a posteriori in this case. Now, we have that Nc_N is a multiple of p^α . In particular, c_N must be a multiple of $p^{\alpha-\gamma}$. We use this to bound N' . Namely, from before we have the inequality

$$p^\gamma N' |c_N| \leq \sum_{d|N'} p^\alpha.$$

Seeing as $|c_N| \geq p^{\alpha-\gamma}$, we have equality if and only if $c_N = \pm p^{\alpha-\gamma}$ and $N' = \sum_{d|N'} 1$. This means $N' = 1$ or $N' = 2$.

We now show that $N' = 2$ is impossible. Let γ now be the smallest value (if it exists) such that $c_{2p^\gamma} \neq 0$. Then,

$$\begin{aligned} t_{2p^\gamma} &= \sum_{d|2p^\gamma} dc_d \\ &= \sum_{d|p^\gamma} dc_d + 2p^\gamma c_{2p^\gamma} \\ &= t_{p^\gamma} \pm 2p^\alpha. \end{aligned}$$

Since t_{p^γ} and t_{2p^γ} both equal 0 or p^α , it is impossible. Hence, $c_{2p^\gamma} = 0$ for all γ . This is the desired result. \square

An important element to study in endomorphisms is $[x]$. We classify the possible places it can go:

Proposition 3.2. *For all $\phi \in \text{End}(\mathcal{M}_{\mathbb{Q}})$, we must have that $\phi([x])$ is one of the following: $[x]$, 0 , $[x - 1]$, or $[(x - 1)/x]$.*

Proof. As before, $t_k([x])$ takes on only the values 0 or 1. If $\phi([x]) = c_0[x] + \sum_{n=1}^{\infty} c_n [x^n - 1]$, then

$$|nc_n| = \left| \sum_{d|n} \mu(n/d) t_d(\phi([x])) \right| \leq \sum_{d|n} 1$$

and so if $c_n \neq 0$ then $n \in \{0, 1, 2\}$. Moreover, in those cases we must have $c_n = 1$ is in those cases. Looking over the finite number of options, we get the desired result \square

The cases $\phi([x]) = [x - 1]$ or $\phi([x]) = [(x - 1)/x]$ can be treated with ease:

Proposition 3.3. *The following properties of an endomorphism in $\text{End}(\mathcal{M}_{\mathbb{Q}})$ are equivalent:*

1. $\forall k \geq 1, \phi^*(k) = 0$
2. $\exists k \geq 1$ s.t $\phi^*(k) = 0$
3. $\phi([x]) = [x - 1]$ or $[(x - 1)/x]$
4. $\exists k$ s.t $\phi(r) = t_0(r)[x - 1] + (t_k(r) - t_0(r)) [x] \forall r \in \mathcal{M}_{\mathbb{Q}}$

Proof.

(1) \implies (2). Trivial.

(2) \implies (3). Let $k \geq 1$ be the value such that $\phi^*(k) = 0$. Then, $t_k(\phi([x])) = t_{\phi^*(k)}([x]) = 1$. Hence, we cannot have $\phi([x])$ be 0 or $[x]$ and so by Proposition 3.2 we get the result.

(3) \implies (4). If $\phi([x]) = [(x - 1)/x]$, then

$$\begin{aligned} \phi([x]r) &= \phi(t_0(r)[x]) \\ &= t_0(r) [(x - 1)/x] = t_0(r)[x - 1] - t_0(r)[x] \\ &= \phi(r)[(x - 1)/x] = \phi(r) - t_0(\phi(r))[x] \end{aligned}$$

so collecting we get that

$$\phi(r) = t_0(r)[x - 1] + (t_{\phi^*(0)}(r) - t_0(r)) [x].$$

We proceed similarly when $\phi([x]) = [x - 1]$.

(4) \implies (1). It is an unenlightening computation that every $\phi \in \text{End}(\mathcal{M}_{\mathbb{Q}})$ of the form in (4) is an endomorphism. For any $j \geq 1$,

$$t_j(t_0(r)[x-1] + (t_{\phi^*(0)}(r) - t_0(r))[x]) = t_0(r)$$

so $\phi^*(j) = 0$ and hence (1) is implied. Combining these implications, we obtain our result. \square

We now analyze the remaining case, $\phi^*(1) \neq 0$. The first step is to study the behavior of ϕ^* more closely.

Lemma 3.1. *If $\phi \in \text{End}(\mathcal{M}_{\mathbb{Q}})$ with $\phi^*(1) \neq 0$, then $v_p(\phi^*(k))$ depends only on $v_p(k)$.*

Proof. By Proposition 3.3, $\phi^*(1) \neq 0$ implies $\phi^*(k) \neq 0$ for all k and hence $v_p(\phi^*(k))$ will all be well defined. By the classification given in Lemma 3.1, we get that $t_k(\phi([x^{p^\alpha} - 1]))$ depends only on $v_p(k)$. In turn, $t_k(\phi([x^{p^\alpha} - 1]))$ as α varies determines $v_p(\phi^*(k))$ and hence we are done. \square

Proposition 3.4. *If $\phi^*(1) \neq 0$, $\phi^*(n)/\phi^*(1)$ is multiplicative, and sends powers of p to powers of p .*

Proof. If n is a power of p , then $v_q(n) = v_q(n)$ for all $q \neq 0$, and hence $v_q(\phi^*(n)) = v_q(\phi^*(1))$ for all $q \neq p$ and hence $\phi^*(n)/\phi^*(1)$ is a power of p . Thus, in general,

$$\begin{aligned} \phi^*(n) &= \prod_p p^{v_p(\phi^*(n))} \\ &= \prod_{p^\alpha || n} p^{v_p(\phi^*(p^\alpha))} \cdot \prod_{\substack{p^\alpha || \phi^*(1) \\ p \nmid n}} p^\alpha \\ &= \phi^*(1) \cdot \prod_{p^\alpha || n} p^{v_p(\phi^*(p^\alpha)) - v_p(\phi^*(1))} \end{aligned}$$

and hence $\phi^*(n)/\phi^*(1)$ is multiplicative. \square

Lemma 3.2. *Suppose $\phi^*(1) \neq 0$, and $\phi([x - k])$ has a root of absolute value greater than 1 for some $k \geq 2$. Then,*

$$\limsup_{N \rightarrow \infty} \frac{\phi^*(N)}{N}$$

is a nonzero rational number.

Proof. Let C be the absolute value of the largest root of $r = \phi([x - k])$. By assumption, $C > 1$. Letting $E < C$ be the absolute value of the second largest root of $\phi([x - k])$, we find that

$$\begin{aligned} k^{\phi^*(n)} &= t_{\phi^*(n)}([x - k]) \\ &= t_n(r) \\ &= S_n \cdot C^n + O(E^n), \end{aligned}$$

where $S_n = \sum_j c_j \theta_j^n$, $c_j \in \mathbb{Z} \neq 0$, and $|\theta_j| = 1$ are all distinct. If S_N were negative or complex for some N , then S_N would be a limit point of S_n by Lemma 2.4. This is a contradiction since then we would have that $k^{\phi^*(n)}$ is negative or complex for some n . We can thus take logarithms, and get that

$$\phi^*(n) = \log(C)n + \log(S_n + O((E/C)^n)).$$

Here, by log we mean logarithm base k . It is standard that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n < N} |S_n|^2 = \sum_j |c_j|^2.$$

Hence, $S_n \geq 1$ for a positive proportion of n since S_n is bounded above. In particular, we find that

$$\limsup_{n \rightarrow \infty} \frac{\phi^*(n)}{n} = \log(C) > 0.$$

Choose $N \geq 1$ such that $S_N \neq 0$. Choose q large such that

$$|S_n - S_{n+q}| \ll \frac{1}{q^{1/J}},$$

where $J \geq 1$ is a fixed integer. There are infinitely many such q by Lemma 2.4. We find uniformly over $m \ll q^{1/J}$

$$\begin{aligned} & |\phi^*(N + mq) - \log(C) \cdot mq| \\ & \ll_N |\phi^*(N + mq) - \log(C) \cdot (N + mq) + \log(S_{N+mq})| \\ & \quad + |\log(S_{N+mq}) - \log(S_N)| \\ & \ll_N 1. \end{aligned}$$

We now manipulate to find that when $N^2 m < q^{1/J}$,

$$\begin{aligned} & \frac{1}{\phi^*(N)} |\phi^*(N + (N^2 m)q) - \log(C) \cdot (N^2 m)q| \\ & = \left| \frac{\phi^*(1 + m(Nq))}{\phi^*(1)} - \frac{N \log(C)}{\phi^*(N)} \cdot m(Nq) \right| \\ & \ll_N 1. \end{aligned}$$

Now, we observe the following. By Proposition 3.4, $\phi^*(p^a)/\phi^*(1)$ is a power of p for each prime p and $a \geq 1$. If $\phi^*(p^a)/\phi^*(1) \geq p^{a+1}$, then $\phi^*(p^a)/p^a \geq \phi^*(1)p$. Since $\phi^*(n) \leq \log(C)n + O(1)$, this can only happen for finitely many p . In particular, let m be the product of all primes for which $\phi^*(p^a)/\phi^*(1) \geq p^{a+1}$ for some $a \geq 1$. Then $1 + m(Nq)$ is relatively prime to all of these primes, and hence

$$\frac{\phi^*(1 + m(Nq))}{\phi^*(1)(1 + m(Nq))}$$

will be the reciprocal of an integer. Hence, since

$$\left| \frac{\phi^*(1 + m(Nq))}{\phi^*(1)(1 + m(Nq))} - \frac{N \log(C)}{\phi^*(N)} \right| \ll_N \frac{1}{q}$$

for large enough q , we have that $\frac{N \log(C)}{\phi^*(N)}$ can be approximated arbitrarily well by reciprocals of integers. Seeing as the set of reciprocals of integers is discrete, this means that $\frac{N \log(C)}{\phi^*(N)}$ itself is the reciprocal of an integer. In particular, $\log(C)$ is rational and we get our result. \square

Lemma 3.3. *In the language of Lemma 3.2, $S_n = 1$ for all n .*

Proof. By the discussion of Lemma 3.2, $S_n \geq \sqrt{\sum_j |c_j|^2}$ for a positive proportion of n . Suppose that $\sum_j |c_j|^2 = 1$. Then, clearly, $S_n = \theta^n$ for a single value θ . To make S_n a positive real number for each n , we must have $\theta = 1$. This gives the desired result.

The other case is that $\sum_j |c_j|^2 \neq 1$. Here, we get that $S_n \geq \sqrt{2}$ for a positive proportion of n . For these values, we have

$$\phi^*(n) = \log(C)n + \log(S_n) + O((E/C)^n).$$

Let D be the denominator of $\log(C)$, which exists by Lemma 3.2. We have

$$D\phi^*(n) = D \log(C)n + D \log(S_n) + O((E/C)^n).$$

Since $S_n \geq \sqrt{2}$, $D \log(S_n)$ is bounded away from zero. Hence,

$$D\phi^*(n) = D \log(C)n + O(1),$$

where $O(1)$ is a bounded size non-zero integer. Every prime dividing $\phi^*(n)$ must either divide $\phi^*(1)$ or n by Proposition 3.4. If a prime divides both $\phi^*(n)$ and n it must also divide $O(1)$, and hence be of bounded size. Hence, we get that there exists $B > 0$ such that $\phi^*(n)$ is B -smooth for all n . The set of B -smooth numbers is of proportion zero in the naturals. In particular, $n/\log(C)$ is always a bounded distance from a B -smooth number and hence must be running over a set of proportion zero in the naturals. However, we assumed that n was running over a set of positive density. This is a contradiction. Hence $S_n = 1$ for all n and we are done. \square

Proposition 3.5. *If $\phi^*(1) \neq 0$ and $\phi([x - k])$ has a root of absolute value greater than 1 for some $k \geq 2$, then $\phi = \lambda_{\phi^*(1)} + (t_{\phi^*(0)} - t_0)[x]$.*

Proof. In the languages of Lemmas 3.2 and 3.3, we have that

$$\phi^*(n) = \log(C)n + O((E/C)^n).$$

Seeing as $\phi^*(n)$ and $\log(C)n$ are rational, this means that $\phi^*(n) = \log(C)n$ for large enough n . By the multiplicativity of $\phi^*(n)/\phi^*(1)$, this means that $\log(C) = \phi^*(1)$, so $\phi^*(n) = \phi^*(1)n$ for large enough n . Again by the multiplicativity of $\phi^*(n)/\phi^*(1)$, this means that $\phi^*(n) = \phi^*(1)n$ for all $n \geq 1$. Introducing the $\phi^*(0) = 0$, and hence $\phi^*(n) = \phi^*(1)n$ for all $n \geq 1$.

It is clear that $\lambda_{\phi^*(1)}^*(n) = \phi^*(1)n$ for all $n \geq 0$. Introducing the $t_{\phi^*(0)} - t_0$ to get the correct value of $\phi^*(0)$, we conclude our proof. \square

Lemma 3.4. *If $\phi^*(1) \neq 0$ and all the roots of $\phi([x - k])$ are of absolute value less than or equal to 1 for some $k \geq 2$, then $\phi([1 - x^{-n}]) = 0$ for all but finitely many n .*

Proof. We find that $t_k(\phi([x - k]))$ is bounded. Seeing as it is integral, we get that it ranges over finitely many values. Hence, $\phi^*(k)$ ranges over finitely many values. Thus, we have that $N \nmid \phi^*(k)$ for all $k \geq 1$, for all but finitely many N , by Proposition 3.3. For these N , we find that $t_k(\phi([x^N - 1]))$ is zero for all $k \geq 1$. Hence, we must have $\phi([x^N - 1]) = N[x]$ for all but finitely many N , giving the result. \square

Theorem 3.1. *If $\phi \in \text{End}(\mathcal{M}_{\mathbb{Q}})$, then either $\phi = \lambda_k + (t_j - t_0)$ for some k, j or $\phi([1 - x^{-n}]) = 0$ for all but finitely many n .*

Proof. If $\phi^*(1) = 0$, then $\phi([1 - x^{-n}]) = 0$ for all but finitely many n by the (2) \iff (3) equivalence in Proposition 2.7. Since every $r \in \mathcal{M}_{\mathbb{Q}}$ can be written as an infinite sum of elements of the form $[1 - x^{-n}]$ by Proposition [ref], this means that $\phi(r)$ will be an N th root of unity, where N is the least common multiple of those n such that $\phi([1 - x^{-n}])$ is nonzero.

Alternatively, suppose $\phi^*(1) \neq 0$. If $\phi([x - 2]) \in \mathcal{M}_{\mathbb{Z}, \Phi}$ then by Lemma 3.4 we have that $\phi([1 - x^{-n}]) = 0$ for all but finitely many n , and hence we can conclude by the above argument. Otherwise, by Lemma 3.5 we have that ϕ is of the desired form. \square

4 Endomorphisms of \mathcal{M}_K , K/\mathbb{Q} algebraic

Let $\phi : \mathcal{M}_K \rightarrow \mathcal{M}_K$ be a continuous ring endomorphism. By Proposition 2.1, for every n there exist $\sigma_n^\phi \in \text{Gal}(K/\mathbb{Q})$ and $\phi^*(n) \in \mathbb{Z}_{\geq 0}$ so that $t_n \circ \phi = \sigma_n^\phi \circ t_{\phi^*(n)}$. In particular, the property of t_n being rational for every n is preserved by ϕ , and hence in turn the property of having rational coefficients is preserved, by Newton's formulas.

Thus, ϕ restricts to a map $\mathcal{M}_{\mathbb{Q}} \rightarrow \mathcal{M}_{\mathbb{Q}}$. Theorem 3.1 says then that either $\phi^*(k)$ is bounded uniformly with respect to k , or it is linear of the form $\phi^*(1)k$ for $k \geq 1$. We now treat these two cases:

Proposition 4.1. *Let $\phi : \mathcal{M}_K \rightarrow \mathcal{M}_K$ be a continuous ring endomorphism for which $\phi^*(k)$ is bounded. Then all roots of elements in the image of ϕ are roots of unity. If K/\mathbb{Q} is finite, then all the roots are N th roots for a fixed N .*

Proof. To begin, choose $r \in \mathcal{M}_K$. Then $t_k(\phi(r)) = \sigma_k(t_{\phi^*(k)}(r))$ runs over finitely many values, since $t_{\phi^*(k)}(r)$ runs over finitely many values and $\text{Gal}(K/\mathbb{Q})$ can send those values to some subset of their (finitely many) Galois conjugates. By Proposition 2.5, we get that all the roots of $\phi(r)$ are roots of unity.

Suppose K/\mathbb{Q} is finite. Then, $t_k(\phi(r))$ goes over finitely many values, and that finite quantity is now uniform as r varies. Suppose there were no N such that every $\phi(r)$ were periodic mod N . Then, by taking weighted linear combinations of these values we could create elements r for which $\phi(r)$ takes on arbitrarily many values. This is a contradiction, hence $\phi(r)$ is periodic mod N for some uniform period N .

We now show that r being periodic mod N implies all of its nonzero roots are N th roots of unity, which in turn will imply our result. Write $t_k(r) = \sum_i c_i \alpha_i^k$. Seeing as this is periodic mod N , we find that

$$\sum_i c_i \alpha_i^{k_0} (\alpha_i^N)^k$$

is constant with respect to k for every k_0 . Hence, we find that

$$\sum_{\alpha_i^N = \alpha} c_i \alpha_i^{k_0} = 0$$

for all $\alpha \neq 1$. Thus, again by Proposition 2.5, we get that $c_i = 0$ whenever $\alpha_i^N \neq 1$. In particular, all α_i are N th roots of unity, so we are done. \square

Lemma 4.1. *Let $\alpha \in \overline{\mathbb{Q}}$ be an algebraic number. For all large enough integers N , none of the Galois conjugates of $\alpha' = N\alpha + 1$ differ in ratio by a root of unity.*

Proof. Suppose that there are infinitely many N for which $N\alpha + 1$ and $N\sigma(\alpha) + 1$ differ in ratio by a root of unity, for some automorphism σ . Then there is an infinite sequence $\frac{\alpha + 1/N}{\sigma(\alpha) + 1/N}$ of roots of unity. All of these terms have bounded algebraic degree over \mathbb{Q} , and hence they must all be J th roots of unity for some J . The set of J th roots of unity is discrete, and hence a non-constant convergent sequence of J th roots of unity is impossible. Thus, we get a contradiction and the desired result is obtained. \square

Proposition 4.2. *Let $\phi : \mathcal{M}_K \rightarrow \mathcal{M}_{\mathbb{C}}$ be a continuous ring endomorphism for which $\phi^*(n) = \phi^*(1)n$ for all $n \geq 1$. Then, $\sigma_n^\phi = \sigma_1^\phi$ for all $n \geq 1$.*

Proof. Choose $\alpha \in K$. We wish to show that $\sigma_n^\phi(\alpha) = \sigma_1^\phi(\alpha)$. By Proposition 4.1, we can assume without loss of generality no Galois conjugates of α differ in ratio by root of unity. Considering where $\phi([x - \alpha])$ goes, we find that

$$\sigma_n^\phi(\alpha)^{\phi^*(1)n} = t_n(\phi([x - \alpha])) = S_n \cdot C^n + O(E^n),$$

where C is a real number, $E < C$, and $S_n = \sum_i c_i \cdot \theta_i^n$ is a weighted sum of complex numbers of the unit circle. Taking absolute values and a logarithm, we get that uniformly for n such that $|S_n| \geq 6/5$

$$\log |\sigma_n^\phi(\alpha)| \cdot \phi^*(1)n = \log(C) \cdot n + \log |S_n| + O((E/C)^n).$$

Taking ratios, we get

$$\log |\sigma_n^\phi(\alpha)| = \frac{\log(C)}{\phi^*(1)} + \frac{\log |S_n|}{\phi^*(1)n} + O((E/C)^n).$$

Seeing as $\sigma_n^\phi(\alpha)$ go over a finite set as n varies, the only way to be arbitrarily well approximated by $\log(C)/\phi^*(1)$ is if $\log |\sigma_n^\phi(\alpha)| = \frac{\log(C)}{\phi^*(1)}$ for large enough n . Hence, for large enough n with $|S_n| \geq 6/5$ we have

$$\frac{\log |S_n|}{\phi^*(1)n} = O((E/C)^n).$$

Seeing as $\log |S_n|$ is bounded away from 0, this can only happen for finitely many n . Hence, $|S_n| \geq 6/5$ for only finitely many n . Hence,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N |S_n|^2 = \sum_i |c_i|^2 \leq (6/5)^2 < 2.$$

Seeing as the $|c_i|$ are integers, for $\sum_i |c_i|^2$ to be less than 2 we must have $c_i = 1$ for some i , and $c_i = 0$ for all other i . Hence, $S_n = \theta^n$ for some $|\theta| = 1$. In particular, we find that

$$\sigma_n^\phi(\alpha)^{\phi^*(1)n} = (\theta C)^n + O(E^n)$$

for all large enough n . Hence,

$$\sigma_n^\phi(\alpha)^{\phi^*(1)n} = (\theta C)^n$$

for all $n \geq 1$. Hence,

$$\left(\frac{\sigma_n^\phi(\alpha)}{\sigma_1^\phi(\alpha)} \right)^{\phi^*(1)n} = 1$$

for all $n \geq 1$. By our assumption on α , we thus get that $\sigma_n^\phi(\alpha) = \sigma_1^\phi(\alpha)$. We get the desired conclusion. \square

Combining the two cases, we get the desired result.