

Multiple Weighted Estimates for Multilinear Commutators of Multilinear Singular Integrals with Generalized Kernels

LIWEN GAO, YAN LIN AND SHUHUI YANG

ABSTRACT. In this paper, the weighted L^p boundedness of multilinear commutators and iterated commutators of multilinear singular integral operators with generalized kernels is established, where the weight is multiple weight. Our results are generalizations of the corresponding results for multilinear singular integral operators with standard kernels and Dini kernels.

1 Introduction

The multilinear singular integral operator theory is the most brilliant part of modern harmonic analysis. It originated from the work [1]-[3] of Meyer and Coifman in the 1970s. The theory mainly study the operator as follows:

Definition 1.1. Suppose $m \in \mathbb{N}^+$ and a function $K(y_0, y_1, \dots, y_m)$ is defined apart from the diagonal $y_0 = y_1 = \dots = y_m$ in $(\mathbb{R}^n)^{m+1}$. T is an m -linear operator defined on product of test functions that relates to K , then its integral representation is as follow:

$$T(f_1, \dots, f_m)(x) = \int_{\mathbb{R}^n} \cdots \int_{\mathbb{R}^n} K(x, y_1, \dots, y_m) \prod_{j=1}^m f_j(y_j) dy_1 \cdots dy_m, \quad (1.1)$$

where $f_j (j = 1, \dots, m) \in C_0^\infty(\mathbb{R}^n)$ and $x \notin \bigcap_{j=1}^m \text{supp} f_j$.

In particular, we call K a standard m -linear Calderón-Zygmund kernel if it has the following size estimates:

$$|K(y_0, y_1, \dots, y_m)| \leq \frac{A}{(\sum_{k,l=0}^m |y_k - y_l|)^{mn}}, \quad (1.2)$$

for some $A > 0$ and all $(y_0, y_1, \dots, y_m) \in (\mathbb{R}^n)^{m+1}$ are away from the diagonal, and smooth estimates:

$$|K(y_0, \dots, y_j, \dots, y_m) - K(y_0, \dots, y'_j, \dots, y_m)| \leq \frac{C|y_j - y'_j|^\varepsilon}{(\sum_{k,l=0}^m |y_k - y_l|)^{mn+\varepsilon}}, \quad (1.3)$$

2010 Mathematics Subject Classification. Primary 42B25, 42B35.

Key words and phrases. Multilinear singular integral with generalized kernel, Multiple weights, Iterated commutator, BMO function.

This work was financially supported by the National Natural Science Foundation of China (No. 12071052).

for some $\varepsilon > 0$, whenever $0 \leq j \leq m$ and $|y_j - y'_j| \leq \frac{1}{2} \max_{0 \leq k \leq m} |y_j - y_k|$.

Definition 1.2. Let T be an m -linear operator defined by (1.1) with a standard m -linear Calderón-Zygmund kernel K , we say that T is a standard m -linear Calderón-Zygmund operator if it satisfies either of the following condition.

Given a group of numbers t_1, t_2, \dots, t_m meeting $1 \leq t_1, t_2, \dots, t_m \leq \infty$ and $1/t = 1/t_1 + \dots + 1/t_m$,

(1) T maps $L^{t_1,1} \times \dots \times L^{t_m,1}$ into $L^{t,\infty}$ if $t > 1$,

(2) T maps $L^{t_1,1} \times \dots \times L^{t_m,1}$ into L^1 if $t = 1$,

where the signs $L^{t_1,1}, L^{t_2,1}, \dots, L^{t_m,1}$ and $L^{t_m,\infty}$ all stand for Lorentz spaces.

The initial interest in the study of commutators is related to the generalization of the classical factorization theorem for Hardy space. Then Pérez and Torres generalized the definition to the commutator generated by multilinear singular integral operator and BMO function in [15]. Now we recall the definition of the multilinear commutators and iterated commutators as follows.

Definition 1.3. Let $\vec{b} = (b_1, \dots, b_m)$ be a family of locally integrable functions. The m -linear commutator generated by \vec{b} and the m -linear operator T is defined by

$$T_{\vec{b}}(f_1, \dots, f_m) = \sum_{j=1}^m T_{\vec{b}}^j(\vec{f}),$$

whenever each term is the commutator of b_j and the j th entry of T , that is

$$T_{\vec{b}}^j = b_j T(f_1, \dots, f_j, \dots, f_m) - T(f_1, \dots, b_j f_j, \dots, f_m).$$

Definition 1.4. Suppose that T is an m -linear singular integral operator and $\vec{b} = (b_1, \dots, b_m)$ is a family of locally integrable functions, then the m -linear iterated commutator generated by T and \vec{b} is defined to be

$$T_{\prod \vec{b}}(f_1, \dots, f_m) = [b_1, [b_2, \dots, [b_{m-1}, [b_m, T]_m]_{m-1} \dots]_2]_1(\vec{f}),$$

where $\vec{f} = (f_1, \dots, f_m)$. If T is linked in the usual way to the kernel K studied in this paper, then we can write

$$\begin{aligned} T_{\prod \vec{b}}(f_1, \dots, f_m)(x) &= \int_{(\mathbb{R}^n)^m} \prod_{j=1}^m (b_j(x) - b_j(y_j)) K(x, y_1, \dots, y_m) f_1(y_1) \cdots f_m(y_m) \\ &\quad \times dy_1 \cdots dy_m. \end{aligned}$$

In 2002, Grafakos-Kalton in [5] and Grafakos-Torres in [7]-[9] gave a systematic statement of the theory. In [7], the authors showed the boundedness of classical singular integral operator. Grafakos-Martell made some work about weighted estimates for standard m -linear Calderón-Zygmund operator in [6]. The weight that they discuss is the Muckenhoupt weight.

They obtained that T is bounded from $L^{p_1}(w_1) \times \cdots \times L^{p_m}(w_m)$ to $L^p(w)$ for any $1 < p_1, \dots, p_m < \infty$ and $1/p = 1/p_1 + \cdots + 1/p_m$.

In [9], Grafakos and Torres put forwards a question: Is there a multiple weight theory? The most appropriate multilinear maximal function or multiple weights to work with in this direction are not yet clear. Therefore, to solve the question the multiple weight was given by Lerner, et al in [10]. The definition of the multiple weight is as follows.

Definition 1.5. Let $1 \leq p_1, \dots, p_m < \infty$. Given $\vec{w} = (w_1, \dots, w_m)$, set

$$v_{\vec{w}} = \prod_{j=1}^m w_j^{\frac{p}{p_j}}.$$

\vec{w} satisfies the $A_{\vec{p}}$ condition if

$$\sup_Q \left(\frac{1}{|Q|} \int_Q v_{\vec{w}} \right)^{\frac{1}{p}} \prod_{j=1}^m \left(\frac{1}{|Q|} \int_Q w_j^{1-p'_j} \right)^{\frac{1}{p'_j}} < \infty.$$

When $p_j = 1$, the condition $\left(\inf_Q w_j \right)^{-1}$ takes place of the condition $\left(\frac{1}{|Q|} \int_Q w_j^{1-p'_j} \right)^{\frac{1}{p'_j}}$.

In [10], the authors not only put forwards the definition of the multiple weight, but also presented a new maximal function

$$\mathcal{M}(\vec{f})(x) = \sup_{Q \ni x} \prod_{j=1}^m \frac{1}{|Q|} \int_Q |f_j(y_j)| dy_j,$$

and denoted $\mathcal{M}_q(\vec{f})$ as

$$\mathcal{M}_q(\vec{f})(x) = \sup_{Q \ni x} \left(\prod_{j=1}^m \frac{1}{|Q|} \int_Q |f_j(y_j)|^q dy_j \right)^{\frac{1}{q}}.$$

This kind of maximal function will become an key tool in our paper in order to match the multiple weight. In [10], by using the key tools the authors obtained a range of results. Set \vec{w} under the assumption of Definition 1.5 and with regard to the standard multilinear Calderón-Zygmund operator T , for any $1 < p_1, \dots, p_m < \infty$ with $1/p = 1/p_1 + \cdots + 1/p_m$, T and $T_{\vec{b}}$ are both bounded from $L^{p_1}(w_1) \times \cdots \times L^{p_m}(w_m)$ into $L^p(v_{\vec{w}})$.

In 2014, Zhang-Lu introduced multilinear singular integral operators with kernels of Dinis type in [14]. The operators that they studied weakened the condition (1.3) in the foundation of standard multilinear Calderón-Zygmund operator into the following condition,

$$\begin{aligned} |K(y_0, \dots, y_j, \dots, y_m) - K(y_0, \dots, y'_j, \dots, y_m)| &\leq \frac{C}{(|y_0 - y_1| + \cdots + |y_0 - y_m|)^{mn}} \\ w \left(\frac{|y_j - y'_j|}{|y_0 - y_1| + \cdots + |y_0 - y_m|} \right), \end{aligned} \quad (1.4)$$

whenever $|y_j - y'_j| \leq \frac{1}{2} \max_{1 \leq j \leq m} |y_0 - y_j|$ and $w(t)$ is non-negative and non-decreasing function on \mathbb{R}^+ .

We easily know that the condition (1.4) can become the condition (1.3) by letting $w(t) = t^\varepsilon$. Thus the standard multilinear Calderón-Zygmund operator is a special case of multilinear singular integral operators with kernels of Dinis type.

In [14], Zhang and Lu mainly studied the multilinear singular integrals with kernels of Dinis type and the multilinear commutators with multiple weight.

In [12], Lin-Xiao gave a kind of multilinear singular integrals with generalized kernels which changed the condition (1.4) into a weaker condition.

For any positive integers k_1, \dots, k_m ,

$$\begin{aligned} & \left(\int_{2^{k_m}|y_0-y'_0| \leq |y_m-y_0| < 2^{k_m+1}|y_0-y'_0|} \cdots \int_{2^{k_1}|y_0-y'_0| \leq |y_m-y_0| < 2^{k_1+1}|y_0-y'_0|} \right. \\ & \left. |K(y_0, y_1, \dots, y_m) - K(y'_0, y_1, \dots, y_m)|^q dy_1 \dots dy_m \right)^{\frac{1}{q}} \\ & \leq C |y_0 - y'_0|^{-\frac{mn}{q'}} \prod_{i=1}^m C_{k_i} 2^{-\frac{n}{q'} k_i}, \end{aligned} \quad (1.5)$$

where $1 < q < \infty$, C_{k_i} is a positive constant about k_i and q' is the conjugate exponent of q .

Motivated by [6, 10, 12, 13, 14], we now discuss the weighted boundedness of multilinear commutators and iterated commutators of multilinear singular integral operators with generalized kernels, where the weight is multiple weight. However, the condition (1.5) can not be suit for the multiple weight condition. This fact forces us to search for another kind of generalized kernels as follows. For $k \in \mathbb{N}$,

$$\begin{aligned} & \left(\int_{2^k \sqrt{m}|y_0-y'_0| \leq |\vec{y}-\vec{y}_0| < 2^{k+1} \sqrt{m}|y_0-y'_0|} |K(y_0, y_1, \dots, y_m) - K(y'_0, y_1, \dots, y_m)|^q dy_1 \dots dy_m \right)^{\frac{1}{q}} \\ & \leq C C_k |y_0 - y'_0|^{-\frac{mn}{q'}} 2^{-\frac{kmn}{q'}}, \end{aligned} \quad (1.6)$$

whenever $\vec{y}_0 = (y_0, y_0, \dots, y_0)$, $\vec{y} = (y_1, y_2, \dots, y_m)$, (q, q') is a fixed pair of positive numbers satisfying $\frac{1}{q} + \frac{1}{q'} = 1$ and C_k is a positive constant about k .

Remark 1.6. By calculation, we can find that condition (1.4) implies condition (1.6) by putting $C_k = w(2^{-k})$. According to the contents that we mentioned in the foregoing, the work in [13] includes the classical case. So we can deduce naturally that our work contains both the cases of standard kernel and Dini type kernel.

This paper is organized as follows. In Section 2, we list the necessary definition and lemmas. In Section 3, we establish the weighted estimates of $T_{\vec{b}}$ and $T_{\prod \vec{b}}$ by their sharp estimation. In Section 4, we will give the proof of all theorems.

2 Necessary Definition and Lemma

Meanwhile, we also give some definition and lemmas used in this paper.

Definition 2.1. Fix positive integers k and m satisfying $1 \leq k < m$ and we suppose C_k^m be a family of all finite subsets $\sigma = \{\sigma(1), \dots, \sigma(k)\}$ of $\{1, 2, \dots, m\}$ with k different elements. If $j < l$, then $\sigma(j) < \sigma(l)$. For any $\sigma \in C_k^m$, let $\sigma' = \{1, 2, \dots, m\} \setminus \sigma$ be the complementary sequence. In particular, $C_0^m = \emptyset$. For an m -tuple \vec{b} and $\sigma \in C_k^m$, the k -tuple $\vec{b}_\sigma = (b_{\sigma(1)}, \dots, b_{\sigma(k)})$ is a finite subset of $\vec{b} = (b_1, \dots, b_m)$. Given T the m -linear operator and $\sigma \in C_j^m$, the multilinear iterated commutator is given by

$$T_{\prod b_\sigma}^-(f_1, \dots, f_m) = [b_{\sigma(1)}, [b_{\sigma(2)}, \dots, [b_{\sigma(j-1)}, [b_{\sigma(j)}, T]_{\sigma(j)}]_{\sigma(j-1)} \dots]_{\sigma(2)}]_{\sigma(1)}(\vec{f}).$$

According to (1.1), we can also represent the multilinear iterated commutator by the form.

$$\begin{aligned} T_{\prod b_\sigma}^-(f_1, \dots, f_m)(x) &= \int_{(\mathbb{R}^n)^m} \prod_{j=1}^m \left(b_{\sigma(i)}(x) - b_{\sigma(i)}(y_{\sigma(i)}) \right) K(x, y_1, \dots, y_m) \\ &\quad \times f_1(y_1) \cdots f_m(y_m) dy_1 \cdots dy_m. \end{aligned}$$

Obviously, $T_{\prod b_\sigma}^- = T_{\prod \vec{b}}$ when $\sigma = \{1, 2, \dots, m\}$ and $T_{\prod b_\sigma}^- = T_{b_j}^j$ when $\sigma = \{j\}$.

Definition 2.2. The Hardy-Littlewood maximal operator M is defined by

$$Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy,$$

where the supremum is taken over all balls which contain x . We can also define the operator $M_s(f)(x)$ by $M_s(f) = [M(|f|^s)]^{1/s}$, $s > 0$. The sharp maximal operator M^\sharp is defined by

$$M^\sharp(f)(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y) - f_B| dy \sim \sup_{B \ni x} \inf_{a \in \mathbb{C}} \frac{1}{|B|} \int_B |f(y) - a| dx.$$

We denote the l -sharp maximal operator by $M_l^\sharp(f) = [M^\sharp(|f|^l)]^{1/l}$, $l > 0$.

Lemma 2.3(see [10]). *Suppose that p, q satisfy $0 < p < q < \infty$. Then there is a positive constant $C = C_{p,q}$ so as to for any measurable function f , the following equality*

$$|Q|^{-1/p} \|f\|_{L^p(Q)} \leq C |Q|^{-1/q} \|f\|_{L^{q,\infty}(Q)}$$

holds for any Q .

Lemma 2.4(see [11]). *Suppose that f is a function in $BMO(\mathbb{R}^n)$, $1 \leq p < \infty$, $x \in \mathbb{R}^n$, and $r_1, r_2 > 0$. Then there is a positive constant C independent of f, x, r_1 and r_2 such that*

$$\left(\frac{1}{|B(x, r_1)|} \int_{B(x, r_1)} |f(y) - f_{B(x, r_2)}|^p dy \right)^{1/p} \leq C \left(1 + \left| \ln \frac{r_1}{r_2} \right| \right) \|f\|_{BMO}.$$

Lemma 2.5(see [10], [4]). *Let $0 < p, \delta < \infty$ and $w \in A_\infty$. Then there exists a constant $C > 0$ depending only on the A_∞ constant of w such that*

$$\int_{\mathbb{R}^n} [M_\delta(f)(x)]^p w(x) dx \leq C \int_{\mathbb{R}^n} [M_\delta^\sharp(f)(x)]^p w(x) dx,$$

for every function f such that the left-hand side is finite.

Lemma 2.6(see [10]). Let $\vec{w} = (w_1, \dots, w_m)$ and $1 \leq p_1, \dots, p_m < \infty$. Then $w \in A_{\vec{p}}$ if and only if

$$\begin{cases} w_j^{1-p'_j} \in A_{mp'_j}, & j = 1, 2, \dots, m, \\ v_{\vec{w}} \in A_{mp}. \end{cases}$$

The condition $w_j^{1-p'_j} \in A_{mp'_j}$ in the case $p_j = 1$ is replaced by $w_j^{\frac{1}{m}} \in A_1$.

Lemma 2.7(see [10]). Let $1 < p_j < \infty$, $j = 1, 2, \dots, m$, and $1/p = 1/p_1 + \dots + 1/p_m$. Then the inequality

$$\|\mathcal{M}(\vec{f})\|_{L^p(v_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}$$

holds for any \vec{f} if and only if \vec{w} is content with $A_{\vec{p}}$ condition.

Lemma 2.8(see [18]). Let $m \geq 2$ and T be an m -linear singular integral operator defined by (1.1) with generalized kernel satisfying (1.6) and $\sum_{k=1}^{\infty} C_k < \infty$. Suppose for fixed $1 \leq r_1, \dots, r_m \leq q'$ with $1/r = 1/r_1 + \dots + 1/r_m$, T is bounded from $L^{r_1} \times \dots \times L^{r_m}$ into $L^{r, \infty}$. If $0 < \delta < 1/m$, then we have

$$M_{\delta}^{\sharp}(T(\vec{f}))(x) \leq C \mathcal{M}_{q'}(\vec{f})(x),$$

for all m -tuples $\vec{f} = (f_1, \dots, f_m)$ of bounded measurable functions with compact support.

3 Main results

Theorem 3.1. Let $m \geq 2$ and T be an m -linear singular integral operator defined by (1.1) with generalized kernel satisfying (1.6) and $\sum_{k=1}^{\infty} kC_k < \infty$. Suppose for fixed $1 \leq r_1, \dots, r_m \leq q'$ with $1/r = 1/r_1 + \dots + 1/r_m$, T is bounded from $L^{r_1} \times \dots \times L^{r_m}$ into $L^{r, \infty}$. If $\vec{b} \in BMO^m$, $0 < \delta < 1/m$, $\delta < \varepsilon_0 < \infty$ and $q' < s < \infty$, then for all m -tuples $\vec{f} = (f_1, \dots, f_m)$ of bounded measurable functions with compact support, there is

$$M_{\delta}^{\sharp}(T_{\vec{b}}(\vec{f}))(x) \leq C \|\vec{b}\|_{BMO^m} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right),$$

where $\|\vec{b}\|_{BMO^m} = \max_{1 \leq j \leq m} \|b_j\|_{BMO}$.

Theorem 3.2. Let $m \geq 2$ and T be an m -linear singular integral operator defined by (1.1) with generalized kernel satisfying (1.6) and $\sum_{k=1}^{\infty} kC_k < \infty$. Suppose for fixed $1 \leq r_1, \dots, r_m \leq q'$ with $1/r = 1/r_1 + \dots + 1/r_m$, T is bounded from $L^{r_1} \times \dots \times L^{r_m}$ into $L^{r, \infty}$. If $\vec{b} \in BMO^m$, then for $q' < p_1, \dots, p_m < \infty$ with $1/p = 1/p_1 + \dots + 1/p_m$, $T_{\vec{b}}$ is bounded from $L^{p_1}(w_1) \times \dots \times L^{p_m}(w_m)$ into $L^p(v_{\vec{w}})$, where $\vec{w} = (w_1, \dots, w_m) \in A_{\vec{p}/q'}$ and

$$v_{\vec{w}} = \prod_{j=1}^m w_j^{\frac{p}{p_j}}.$$

Theorem 3.3. Let $m \geq 2$ and T be an m -linear singular integral operator defined by (1.1) with generalized kernel satisfying (1.6) and $\sum_{k=1}^{\infty} k^m C_k < \infty$. Suppose for fixed $1 \leq r_1, \dots, r_m \leq q'$ with $1/r = 1/r_1 + \dots + 1/r_m$, T is bounded from $L^{r_1} \times \dots \times L^{r_m}$ into $L^{r, \infty}$. If $\vec{b} \in BMO^m$, $0 < \delta < 1/m$, $\delta < \varepsilon_0 < \infty$ and $q' < s < \infty$, then for all m -tuples $\vec{f} = (f_1, \dots, f_m)$ of bounded measurable functions with compact support,

$$\begin{aligned} M_{\delta}^{\sharp}(T_{\prod \vec{b}}(\vec{f}))(x) &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right) \\ &\quad + C \sum_{j=1}^{m-1} \sum_{\sigma \in C_j^m} \prod_{i=1}^j \|b_{\sigma(i)}\|_{BMO} M_{\varepsilon_0}(T_{\prod \vec{b}_{\sigma'}}(\vec{f}))(x). \end{aligned}$$

Theorem 3.4. Let $m \geq 2$ and T be an m -linear singular integral operator defined by (1.1) with generalized kernel satisfying (1.6) and $\sum_{k=1}^{\infty} k^m C_k < \infty$. Suppose for fixed $1 \leq r_1, \dots, r_m \leq q'$ with $1/r = 1/r_1 + \dots + 1/r_m$, T is bounded from $L^{r_1} \times \dots \times L^{r_m}$ into $L^{r, \infty}$. If $\vec{b} \in BMO^m$, $q' < p_1, \dots, p_m < \infty$ with $1/p = 1/p_1 + \dots + 1/p_m$, then $T_{\prod \vec{b}}$ is bounded from $L^{p_1}(w_1) \times \dots \times L^{p_m}(w_m)$ into $L^p(v_{\vec{w}})$, where $\vec{w} = (w_1, \dots, w_m) \in A_{\vec{P}/q'}$ and $v_{\vec{w}} = \prod_{j=1}^m w_j^{\frac{p}{p_j}}$.

4 Proof of the main results

Proof of Theorem 3.1. Fix $x \in \mathbb{R}^n$. For any ball $B \ni x$, we first consider the commutator.

$$T_{b_1}^1(f_1, \dots, f_m)(z) = b_1(z)T(f_1, \dots, f_m)(z) - T(b_1 f_1, \dots, f_m)(z).$$

Let $B^* = 16\sqrt{m}B$, then for any $z \in B$,

$$T_{b_1}^1(f_1, \dots, f_m)(z) = (b_1(z) - b_{1B^*})T(f_1, \dots, f_m)(z) - T((b_1 - b_{1B^*})f_1, \dots, f_m)(z),$$

where $b_{1B^*} = \frac{1}{|B^*|} \int_{B^*} b_1(z) dz$. Since $0 < \delta < 1$, then for any $c \in \mathbb{C}$, we have

$$\begin{aligned} &\left(\frac{1}{|B|} \int_B |T_{b_1}^1(\vec{f})(z)|^{\delta} - |c|^{\delta} dz \right)^{\frac{1}{\delta}} \\ &\leq C \left(\frac{1}{|B|} \int_B |(b_1(z) - b_{1B^*})T(\vec{f})(z)|^{\delta} dz \right)^{\frac{1}{\delta}} \\ &\quad + C \left(\frac{1}{|B|} \int_B |T((b_1 - b_{1B^*})f_1, \dots, f_m)(z) - c|^{\delta} dz \right)^{\frac{1}{\delta}} \\ &:= I + II. \end{aligned}$$

We can find an l such that $1 < l < \min\{\frac{\varepsilon_0}{\delta}, \frac{1}{1-\delta}\}$, then $l\delta < \varepsilon_0$ and $l'\delta > 1$. By Hölder's inequality and Lemma 2.4, we obtain

$$I \leq C \left(\frac{1}{|B|} \int_B |b_1(z) - b_{1B^*}|^{\delta l'} dz \right)^{\frac{1}{\delta l'}} \left(\frac{1}{|B|} \int_B |T(\vec{f})(z)|^{\delta l} dz \right)^{\frac{1}{\delta l}}$$

$$\begin{aligned}
&\leq C(1 + \ln 16\sqrt{m})\|b_1\|_{BMO}M_{\delta l}(T(\vec{f}))(x) \\
&\leq C\|b_1\|_{BMO}M_{\varepsilon_0}(T(\vec{f}))(x).
\end{aligned}$$

For each j , $f_j = f_j^0 + f_j^\infty$ where $f_j^0 = f_j\chi_{B^*}$. Then

$$\prod_{j=1}^m f_j(y_j) = \prod_{j=1}^m f_j^0(y_j) + \sum_{(\alpha_1, \dots, \alpha_m) \in \Gamma} f_1^{\alpha_1}(y_1) \cdots f_m^{\alpha_m}(y_m),$$

where $\Gamma = \{(\alpha_1, \alpha_2, \dots, \alpha_m) : \text{there is at least one } \alpha_j \neq 0\}$. Denote $\vec{z}_0 = (z_0, z_0, \dots, z_0)$, $\vec{y} = (y_1, y_2, \dots, y_m)$, the center of B by x_0 and the radius of B by r_B . We choose $z_0 \in 4B \setminus 3B$. Let

$$c = \sum_{(\alpha_1, \dots, \alpha_m) \in \Gamma} T((b_1 - b_{1B^*})f_1^{\alpha_1}, f_2^{\alpha_2}, \dots, f_m^{\alpha_m})(z_0).$$

Then

$$\begin{aligned}
II &\leq C \left(\frac{1}{|B|} \int_B |T((b_1 - b_{1B^*})f_1^0, f_2^0, \dots, f_m^0)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\
&\quad + C \sum_{(\alpha_1, \dots, \alpha_m) \in \Gamma} \left(\frac{1}{|B|} \int_B |T((b_1 - b_{1B^*})f_1^{\alpha_1}, f_2^{\alpha_2}, \dots, f_m^{\alpha_m})(z) \right. \\
&\quad \left. - T((b_1 - b_{1B^*})f_1^{\alpha_1}, f_2^{\alpha_2}, \dots, f_m^{\alpha_m})(z_0)|^\delta dz \right)^{\frac{1}{\delta}} \\
&:= II_0 + \sum_{(\alpha_1, \dots, \alpha_m) \in \Gamma} II_{\alpha_1, \dots, \alpha_m}.
\end{aligned}$$

Let $t = s/q'$, then it follows from $s > q'$ that $t > 1$. Due to for any j , $r_j \leq q'$, we can get $r_j t \leq s$. By Lemma 2.3 and Lemma 2.4, we have

$$\begin{aligned}
II_0 &\leq C|B|^{-1/\delta} \|T((b_1 - b_{1B^*})f_1^0, f_2^0, \dots, f_m^0)\|_{L^\delta(B)} \\
&\leq C|B|^{-1/r} \|T((b_1 - b_{1B^*})f_1^0, f_2^0, \dots, f_m^0)\|_{L^{r, \infty}(\mathbb{R}^n)} \\
&\leq C \left(\frac{1}{|B^*|} \int_{B^*} |(b_1(z) - b_{1B^*})f_1(y_1)|^{r_1} dy_1 \right)^{\frac{1}{r_1}} \prod_{j=2}^m \left(\frac{1}{|B^*|} \int_{B^*} |f_j(y_j)|^{r_j} dy_j \right)^{\frac{1}{r_j}} \\
&\leq C \left(\frac{1}{|B^*|} \int_{B^*} |b_1(z) - b_{1B^*}|^{r_1 t'} dy_1 \right)^{\frac{1}{r_1 t'}} \left(\frac{1}{|B^*|} \int_{B^*} |f_1(y_1)|^{r_1 t} dy_1 \right)^{\frac{1}{r_1 t}} \\
&\quad \times \prod_{j=2}^m \left(\frac{1}{|B^*|} \int_{B^*} |f_j(y_j)|^{r_j} dy_j \right)^{\frac{1}{r_j}} \\
&\leq C\|b_1\|_{BMO} \prod_{j=1}^m \left(\frac{1}{|B^*|} \int_{B^*} |f_j(y_j)|^s dy_j \right)^{\frac{1}{s}} \\
&\leq C\|b_1\|_{BMO} \mathcal{M}_s(\vec{f})(x).
\end{aligned}$$

For any $(\alpha_1, \alpha_2, \dots, \alpha_m) \in \Gamma$, we can find a $j \in \{1, \dots, m\}$ that makes $\alpha_j = \infty$. Then for any $\vec{y} = (y_1, y_2, \dots, y_m) \in \text{supp}f_1^{\alpha_1} \times \dots \times \text{supp}f_m^{\alpha_m}$ and $z \in B$, $|\vec{y} - \vec{z}_0| \geq 2\sqrt{m}|z - z_0|$, and $2r_B \leq |z - z_0| \leq 5r_B$. Thus

$$\begin{aligned}
II_{\alpha_1, \alpha_2, \dots, \alpha_m} &\leq C \frac{1}{|B|} \int_B \int_{|\vec{y} - \vec{z}_0| \geq 2\sqrt{m}|z - z_0|} |K(z, \vec{y}) - K(z_0, \vec{y})| |b_1(y_1) - b_{1B^*}| \\
&\quad \times \prod_{j=1}^m |f_j(y_j)| d\vec{y} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \int_{2^k \sqrt{m}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1} \sqrt{m}|z - z_0|} |K(z, \vec{y}) - K(z_0, \vec{y})| \\
&\quad \times |b_1(y_1) - b_{1B^*}| \prod_{j=1}^m |f_j(y_j)| d\vec{y} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k \sqrt{m}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1} \sqrt{m}|z - z_0|} |K(z, \vec{y}) - K(z_0, \vec{y})|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \left(\int_{|\vec{y} - \vec{z}_0| < 2^{k+1} \sqrt{m}|z - z_0|} |b_1(y_1) - b_{1B^*}|^{q'} \prod_{j=1}^m |f_j(y_j)|^{q'} d\vec{y} \right)^{\frac{1}{q'}} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k \sqrt{m}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1} \sqrt{m}|z - z_0|} |K(z, \vec{y}) - K(z_0, \vec{y})|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \left(\frac{1}{|B(z_0, 2^{k+1} \sqrt{m}|z - z_0)|} \int_{|y_1 - z_0| < 2^{k+1} \sqrt{m}|z - z_0|} |(b_1(y_1) - b_{1B^*})| \right. \\
&\quad \times \left. f_1(y_1) \right)^{\frac{1}{q'}} \\
&\quad \times \prod_{j=2}^m \left(\frac{1}{|B(z_0, 2^{k+1} \sqrt{m}|z - z_0)|} \int_{|y_j - z_0| < 2^{k+1} \sqrt{m}|z - z_0|} |f_j(y_j)|^{q'} dy_j \right)^{\frac{1}{q'}} \\
&\quad \times (2^{k+1} \sqrt{m}|z - z_0|)^{\frac{mn}{q'}} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k \sqrt{m}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1} \sqrt{m}|z - z_0|} |K(z, \vec{y}) - K(z_0, \vec{y})|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \left(\frac{1}{|B(x_0, 2^{k+2} \sqrt{m}|z - z_0)|} \int_{|y_1 - x_0| < 2^{k+2} \sqrt{m}|z - z_0|} |b_1(y_1) - b_{1B^*}|^{q't'} dy_1 \right)^{\frac{1}{q't'}} \\
&\quad \times \left(\frac{1}{|B(z_0, 2^{k+1} \sqrt{m}|z - z_0)|} \int_{|y_1 - z_0| < 2^{k+1} \sqrt{m}|z - z_0|} |f_1(y_1)|^{q't} dy_1 \right)^{\frac{1}{q't}} \\
&\quad \times \prod_{j=2}^m \left(\frac{1}{|B(z_0, 2^{k+1} \sqrt{m}|z - z_0)|} \int_{|y_j - z_0| < 2^{k+1} \sqrt{m}|z - z_0|} |f_j(y_j)|^{q't} dy_j \right)^{\frac{1}{q't}} \\
&\quad \times (2^{k+1} \sqrt{m}|z - z_0|)^{\frac{mn}{q'}} dz
\end{aligned}$$

$$\begin{aligned}
&\leq C \|b_1\|_{BMO} \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} |z - z_0|^{-\frac{mn}{q'}} C_k 2^{-\frac{kmn}{q'}} k (2^{k+1} \sqrt{m} |z - z_0|)^{\frac{mn}{q'}} \\
&\quad \times \prod_{j=1}^m \left(\frac{1}{|B(z_0, 2^{k+1} \sqrt{m} |z - z_0|)|} \int_{|y_j - z_0| < 2^{k+1} \sqrt{m} |z - z_0|} |f_j(y_j)|^s dy_j \right)^{\frac{1}{s}} dz \\
&\leq C \|b_1\|_{BMO} \mathcal{M}_s(\vec{f})(x).
\end{aligned}$$

Thus

$$\sum_{(\alpha_1, \dots, \alpha_m) \in \Gamma} II_{\alpha_1, \dots, \alpha_m} \leq C \|b_1\|_{BMO} \mathcal{M}_s(\vec{f})(x).$$

Through the above estimates we get the result

$$\begin{aligned}
M_{\delta}^{\sharp}(T_b^1(\vec{f}))(x) &\leq C \|b_1\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right) \\
&\leq C \|\vec{b}\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right).
\end{aligned}$$

Similarly, we can get to for any $j = 1, 2, \dots, m$,

$$M_{\delta}^{\sharp}(T_b^j(\vec{f}))(x) \leq C \|\vec{b}\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right).$$

So we have

$$\begin{aligned}
M_{\delta}^{\sharp}(T_b(\vec{f}))(x) &= M_{\delta}^{\sharp} \left(\sum_{j=1}^m T_b^j(\vec{f}) \right) (x) \\
&\leq \sum_{j=1}^m M_{\delta}^{\sharp}(T_b^j(\vec{f}))(x) \\
&\leq C \|\vec{b}\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right),
\end{aligned}$$

which completes the proof of the theorem. \square

Proof of Theorem 3.2. Let $\vec{q} = \frac{\vec{P}}{q'} = (\frac{p_1}{q'}, \dots, \frac{p_m}{q'})$. Due to $w \in A_{\vec{q}}$, by Lemma 2.6, each $\psi_j = w_j^{-\frac{1}{q_j-1}}$ belongs to A_{∞} where $q_j = \frac{p_j}{q'}$, $j = 1, \dots, m$. By inverse Hölder's inequality, we can find constants $c_j, t_j > 1$ depending on the A_{∞} constant of ψ_j such that

$$\left(\frac{1}{|B|} \int_B w_j^{-\frac{t_j}{q_j-1}} \right)^{\frac{1}{t_j}} \leq \frac{c_j}{|B|} \int_B w_j^{-\frac{1}{q_j-1}},$$

for any ball B . Pick a $d_j > 1$ that makes

$$\frac{t_j}{q_j - 1} = \frac{1}{\frac{q_j}{d_j} - 1}.$$

Then $q_j > d_j > 1$, $j = 1, \dots, m$.

Let $d = \min\{d_1, \dots, d_m\}$ and $c = \max\{c_1, \dots, c_m\}$. We have for $q_0 = \frac{p}{q}$

$$\begin{aligned}
& \left(\frac{1}{|B|} \int_B v_{\vec{w}} \right)^{1/\frac{q_0}{d}} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{\frac{q_j}{d}-1}} \right)^{1-1/\frac{q_j}{d}} \\
&= \left(\frac{1}{|B|} \int_B v_{\vec{w}} \right)^{\frac{d}{q_0}} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{\frac{q_j}{d}-1}} \right)^{\left(\frac{q_j}{d}-1\right)\frac{d}{q_j}} \\
&\leq \left(\frac{1}{|B|} \int_B v_{\vec{w}} \right)^{\frac{d}{q_0}} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{\frac{q_j}{d_j}-1}} \right)^{\left(\frac{q_j}{d_j}-1\right)\frac{d}{q_j}} \\
&= \left(\frac{1}{|B|} \int_B v_{\vec{w}} \right)^{\frac{d}{q_0}} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{t_j}{q_j-1}} \right)^{\left(\frac{q_j-1}{t_j}\right)\frac{d}{q_j}} \\
&\leq c^{dm} \left(\frac{1}{|B|} \int_B v_{\vec{w}} \right)^{\frac{d}{q_0}} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{q_j-1}} \right)^{(q_j-1)\frac{d}{q_j}} \\
&\leq c^{dm} [w]_{A_{\vec{q}}}^d.
\end{aligned}$$

Thus $\vec{w} \in A_{\vec{q}/d}$. Let $s = q'd$, then we can find that $s > q'$ and $s < p_j$, $j = 1, \dots, m$ from $d > 1$. We can get the result

$$\vec{w} \in A_{\vec{q}/d} = A_{\vec{p}/s}.$$

Take ε_0, δ so that $0 < \delta < \varepsilon_0 < \frac{1}{m}$. We can deduce the following conclusion from Lemma 2.5, Theorem 3.1, Lemma 2.8, Lemma 2.7 and $\vec{w} \in A_{\vec{p}/s}$.

$$\begin{aligned}
\|T_{\vec{b}}(\vec{f})\|_{L^p(v_{\vec{w}})} &\leq \|M_\delta(T_{\vec{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} \\
&\leq \|M_\delta^\sharp(T_{\vec{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} \\
&\leq C\|\vec{b}\|_{BMO^m} \|M_{\varepsilon_0}(T(\vec{f})) + \mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} \\
&\leq C\|\vec{b}\|_{BMO^m} (\|M_{\varepsilon_0}^\sharp(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})}) \\
&\leq C\|\vec{b}\|_{BMO^m} \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} = C\|\vec{b}\|_{BMO^m} \|\mathcal{M}(\vec{f}^s)\|_{L^{\frac{p}{s}}(v_{\vec{w}})}^{\frac{1}{s}} \\
&\leq C\|\vec{b}\|_{BMO^m} \prod_{j=1}^m \| |f_j|^s \|_{L^{\frac{p_j}{s}(w_j)}}^{\frac{1}{s}} = C\|\vec{b}\|_{BMO^m} \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.
\end{aligned}$$

This finishes the proof of the Theorem 3.2. \square

Proof of Theorem 3.3. For the sake of simplicity, we only consider the case $m = 2$. The proof of other cases is similar. Let f_1, f_2 be bounded measurable functions with compact support and $b_1, b_2 \in BMO$. Then for any constants λ_1 and λ_2 ,

$$T_{\prod \vec{b}}(\vec{f})(z) = (b_1(z) - \lambda_1)(b_2(z) - \lambda_2)T(f_1, f_2)(z) - (b_1(z) - \lambda_1)T(f_1, (b_2 - \lambda_2)f_2)(z)$$

$$\begin{aligned}
& -(b_2(z) - \lambda_2)T((b_1 - \lambda_1)f_1, f_2)(z) + T((b_1 - \lambda_1)f_1, (b_2 - \lambda_2)f_2)(z) \\
= & -(b_1(z) - \lambda_1)(b_2(z) - \lambda_2)T(f_1, f_2)(z) + (b_1(z) - \lambda_1)T_{b_2 - \lambda_2}^2(f_1, f_2)(z) \\
& + (b_2(z) - \lambda_2)T_{b_1 - \lambda_1}^1(f_1, f_2)(z) + T((b_1 - \lambda_1)f_1, (b_2 - \lambda_2)f_2)(z).
\end{aligned}$$

Let C_0 be a constant determined later. Fixed $x \in \mathbb{R}^n$, for any ball $B(x_0, r_B)$ containing x and $0 < \delta < \frac{1}{2}$, we have

$$\begin{aligned}
& \left(\frac{1}{|B|} \int_B |T_{\Pi \vec{b}}(\vec{f})(z)|^\delta - |C_0|^\delta dz \right)^{\frac{1}{\delta}} \\
\leq & \left(\frac{1}{|B|} \int_B |T_{\Pi \vec{b}}(\vec{f})(z) - C_0|^\delta dz \right)^{\frac{1}{\delta}} \\
\leq & C \left(\frac{1}{|B|} \int_B |(b_1(z) - \lambda_1)(b_2(z) - \lambda_2)T(f_1, f_2)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\
& + C \left(\frac{1}{|B|} \int_B |(b_1(z) - \lambda_1)T_{b_2 - \lambda_2}^2(f_1, f_2)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\
& + C \left(\frac{1}{|B|} \int_B |(b_2(z) - \lambda_2)T_{b_1 - \lambda_1}^1(f_1, f_2)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\
& + C \left(\frac{1}{|B|} \int_B |T((b_1 - \lambda_1)f_1, (b_2 - \lambda_2)f_2)(z) - C_0|^\delta dz \right)^{\frac{1}{\delta}} \\
:= & I + II + III + IV.
\end{aligned}$$

Then we calculate each term respectively. Denote $B^* = 16\sqrt{2}B$ and $\lambda_j = (b_j)_{B^*} = \frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} b_j(x) dx$, $j = 1, 2$. Since $0 < \delta < \frac{1}{2}$ and $0 < \delta < \varepsilon_0 < \infty$, it is easy to seek an l such that $1 < l < \min\{\frac{\varepsilon_0}{\delta}, \frac{1}{1-\delta}\}$. We can deduce that $l\delta < \varepsilon_0$ and $l'\delta > 1$. Choose $q_1, q_2 \in (1, \infty)$ satisfying $\frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{l'}$, then $\frac{1}{q_1} + \frac{1}{q_2} + \frac{1}{l} = 1$, $q_1\delta > 1$ and $q_2\delta > 1$. By Lemma 2.4, we have

$$\begin{aligned}
I & \leq C \left(\frac{1}{|B|} \int_B |T(f_1, f_2)(z)|^{l\delta} dz \right)^{\frac{1}{l\delta}} \left(\frac{1}{|B|} \int_B |b_1(z) - \lambda_1|^{q_1\delta} dz \right)^{\frac{1}{q_1\delta}} \\
& \quad \times \left(\frac{1}{|B|} \int_B |b_2(z) - \lambda_2|^{q_2\delta} dz \right)^{\frac{1}{q_2\delta}} \\
& \leq C \|b_1\|_{BMO} \|b_2\|_{BMO} M_{\delta l}(T(f_1, f_2))(x) \\
& \leq C \|b_1\|_{BMO} \|b_2\|_{BMO} M_{\varepsilon_0}(T(f_1, f_2))(x).
\end{aligned}$$

It follow from Hölder's inequality that

$$\begin{aligned}
II & \leq C \left(\frac{1}{|B|} \int_B |T_{b_2 - \lambda_2}^2(f_1, f_2)(z)|^{l\delta} dz \right)^{\frac{1}{l\delta}} \left(\frac{1}{|B|} \int_B |b_1(z) - \lambda_1|^{l'\delta} dz \right)^{\frac{1}{l'\delta}} \\
& \leq C \|b_1\|_{BMO} M_{\delta l}(T_{b_2 - \lambda_2}^2(f_1, f_2))(x) \\
& \leq C \|b_1\|_{BMO} M_{\varepsilon_0}(T_{b_2}^2(f_1, f_2))(x).
\end{aligned}$$

Similarly we can get

$$III \leq C \|b_2\|_{BMO} M_{\varepsilon_0}(T_{b_1}^1(f_1, f_2))(x).$$

In the next, we discuss *IV*. Split f_i into two parts $f_i = f_i^0 + f_i^\infty$, where $f_i^0 = f \chi_{B^*}$ and $f_i^\infty = f_i - f_i^0$, $i = 1, 2$. Choose $z_0 \in 4B \setminus 3B$. Denote $\vec{z}_0 = (z_0, z_0)$ and $\vec{y} = (y_1, y_2)$. Let

$$C_0 = \sum_{(\alpha_1, \alpha_2) \in \Gamma} T((b_1 - \lambda_1)f_1^{\alpha_1}, (b_2 - \lambda_2)f_2^{\alpha_2})(z_0),$$

where $\Gamma = \{(\alpha_1, \alpha_2): \text{there is at least one } \alpha_j \neq 0, j = 1, 2\}$, then

$$\begin{aligned} IV &\leq C \left(\frac{1}{|B|} \int_B |T((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^0)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\ &\quad + C \sum_{(\alpha_1, \alpha_2) \in \Gamma} \left(\frac{1}{|B|} \int_B |T((b_1 - \lambda_1)f_1^{\alpha_1}, (b_2 - \lambda_2)f_2^{\alpha_2})(z) \right. \\ &\quad \left. - T((b_1 - \lambda_1)f_1^{\alpha_1}, (b_2 - \lambda_2)f_2^{\alpha_2})(z_0)|^\delta dz \right)^{\frac{1}{\delta}} \\ &:= IV_0 + \sum_{(\alpha_1, \alpha_2) \in \Gamma} IV_{\alpha_1, \alpha_2}. \end{aligned}$$

Let $t = s/q'$, then it follows from $s > q'$ that $t > 1$. Due to for any $j = 1, 2$, $r_j \leq q' < s$, we can get $r_j t \leq s$. By Lemma 2.3 and Hölder's inequality,

$$\begin{aligned} IV_0 &\leq C |B|^{-1/\delta} \|T((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^0)\|_{L^\delta(B)} \\ &\leq C |B|^{-1/r} \|T((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^0)\|_{L^{r, \infty}(B)} \\ &\leq C \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |b_1(y_1) - \lambda_1|^{r_1} |f_1(y_1)|^{r_1} dy_1 \right)^{\frac{1}{r_1}} \\ &\quad \times \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |b_2(y_2) - \lambda_2|^{r_2} |f_2(y_2)|^{r_2} dy_2 \right)^{\frac{1}{r_2}} \\ &\leq C \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |f_1(y_1)|^{r_1 t} dy_1 \right)^{\frac{1}{r_1 t}} \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |b_1(y_1) - \lambda_1|^{r_1 t'} dy_1 \right)^{\frac{1}{r_1 t'}} \\ &\quad \times \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |f_2(y_2)|^{r_2 t} dy_2 \right)^{\frac{1}{r_2 t}} \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |b_2(y_2) - \lambda_2|^{r_2 t'} dy_2 \right)^{\frac{1}{r_2 t'}} \\ &\leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |f_1(y_1)|^s dy_1 \right)^{\frac{1}{s}} \\ &\quad \times \left(\frac{1}{|16\sqrt{2}B|} \int_{16\sqrt{2}B} |f_2(y_2)|^s dy_2 \right)^{\frac{1}{s}} \\ &\leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \mathcal{M}_s(\vec{f})(x). \end{aligned}$$

For any $(\alpha_1, \alpha_2) \in \Gamma$, we can find a $j \in \{1, 2\}$ that makes $\alpha_j = \infty$. Then for any $\vec{y} = (y_1, y_2) \in \text{supp}f_1^{\alpha_1} \times \text{supp}f_2^{\alpha_2}$ and $z \in B$, $|\vec{y} - \vec{z}_0| \geq 2\sqrt{2}|z - z_0|$ and $2r_B \leq |z - z_0| \leq 5r_B$. Thus

$$\begin{aligned}
IV_{\alpha_1, \alpha_2} &\leq C \frac{1}{|B|} \int_B \int_{|\vec{y} - \vec{z}_0| \geq 2\sqrt{2}|z - z_0|} |K(z, y_1, y_2) - K(z_0, y_1, y_2)| \\
&\quad \times \prod_{j=1}^2 |b_j(y_j) - \lambda_j| |f_j(y_j)| d\vec{y} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \int_{2^k\sqrt{2}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1}\sqrt{2}|z - z_0|} |K(z, y_1, y_2) - K(z_0, y_1, y_2)| \\
&\quad \times \prod_{j=1}^2 |b_j(y_j) - \lambda_j| |f_j(y_j)| d\vec{y} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k\sqrt{2}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1}\sqrt{2}|z - z_0|} |K(z, y_1, y_2) - K(z_0, y_1, y_2)|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \left(\int_{|\vec{y} - \vec{z}_0| < 2^{k+1}\sqrt{2}|z - z_0|} \prod_{j=1}^2 |b_j(y_j) - \lambda_j|^{q'} |f_j(y_j)|^{q'} d\vec{y} \right)^{\frac{1}{q'}} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k\sqrt{2}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1}\sqrt{2}|z - z_0|} |K(z, y_1, y_2) - K(z_0, y_1, y_2)|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \prod_{j=1}^2 \left(\frac{1}{|B(z_0, 2^{k+1}\sqrt{2}|z - z_0)|} \int_{|y_j - z_0| < 2^{k+1}\sqrt{2}|z - z_0|} |b_j(y_j) - \lambda_j|^{q'} |f_j(y_j)|^{q'} dy_j \right)^{\frac{1}{q'}} \\
&\quad \times |B(z_0, 2^{k+1}\sqrt{2}|z - z_0)|^{\frac{2}{q'}} dz \\
&\leq C \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} \left(\int_{2^k\sqrt{2}|z - z_0| \leq |\vec{y} - \vec{z}_0| < 2^{k+1}\sqrt{2}|z - z_0|} |K(z, y_1, y_2) - K(z_0, y_1, y_2)|^q d\vec{y} \right)^{\frac{1}{q}} \\
&\quad \times \prod_{j=1}^2 \left(\frac{1}{|B(x_0, 2^{k+2}\sqrt{2}|z - z_0)|} \int_{|y_j - x_0| < 2^{k+2}\sqrt{2}|z - z_0|} |b_j(y_j) - \lambda_j|^{q't'} |dy_j| \right)^{\frac{1}{q't'}} \\
&\quad \times \prod_{j=1}^2 \left(\frac{1}{|B(z_0, 2^{k+1}\sqrt{2}|z - z_0)|} \int_{|y_j - z_0| < 2^{k+1}\sqrt{2}|z - z_0|} |f_j(y_j)|^{q't} dy_j \right)^{\frac{1}{q't}} \\
&\quad \times |B(z_0, 2^{k+1}\sqrt{2}|z - z_0)|^{\frac{2}{q'}} dz \\
&\leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \frac{1}{|B|} \int_B \sum_{k=1}^{\infty} |z - z_0|^{-\frac{2n}{q'}} C_k 2^{-\frac{2kn}{q'}} k^2 (2^{k+1}\sqrt{2}|z - z_0|)^{\frac{2n}{q'}} \\
&\quad \times \prod_{j=1}^2 \left(\frac{1}{|B(z_0, 2^{k+1}\sqrt{2}|z - z_0)|} \int_{|y_j - z_0| < 2^{k+1}\sqrt{2}|z - z_0|} |f_j(y_j)|^s dy_j \right)^{\frac{1}{s}} dz \\
&\leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \mathcal{M}_s(\vec{f})(x).
\end{aligned}$$

Thus

$$\sum_{(\alpha_1, \alpha_2) \in \Gamma} IV_{\alpha_1, \alpha_2} \leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \mathcal{M}_s(\vec{f})(x).$$

So

$$IV \leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \mathcal{M}_s(\vec{f})(x).$$

Finally, we can get the result,

$$\begin{aligned} M_\delta^\sharp(T_{\prod \bar{b}}(\vec{f}))(x) &= M^\sharp(|T_{\prod \bar{b}}(\vec{f})|^\delta)^{\frac{1}{\delta}}(x) \\ &\leq C \sup_{B \ni x} \left(\frac{1}{|B|} \int_B \|T_{\prod \bar{b}}(\vec{f})(z)^\delta - |C_0|^\delta dz \right)^{\frac{1}{\delta}} \\ &\leq C \|b_1\|_{BMO} \|b_2\|_{BMO} \left(M_{\varepsilon_0}(T(\vec{f}))(x) + \mathcal{M}_s(\vec{f})(x) \right) \\ &\quad + C (\|b_1\|_{BMO} M_{\varepsilon_0}(T_{b_2}^2(\vec{f}))(x) + \|b_2\|_{BMO} M_{\varepsilon_0}(T_{b_1}^1(\vec{f}))(x)). \end{aligned}$$

This completes the proof of Theorem 3.3. \square

Proof of Theorem 3.4.

Proceeding as in the proof of Theorem 3.2, we can find a s such that $q' < s$, $s < p_j$, $j = 1, \dots, m$ and $\vec{w} \in A_{\vec{p}/s}$.

Choose $\delta, \varepsilon_1, \varepsilon_2, \dots, \varepsilon_m$ satisfying $0 < \delta < \varepsilon_1 < \varepsilon_2 < \dots < \varepsilon_m < \frac{1}{m}$. We can deduce the conclusion by Lemma 2.5 and Lemma 2.8.

$$\|M_{\varepsilon_j}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} \leq C \|M_{\varepsilon_j}^\sharp(T(\vec{f}))\|_{L^p(v_{\vec{w}})} \leq C \|\mathcal{M}_{q'}(\vec{f})\|_{L^p(v_{\vec{w}})} \leq C \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})}.$$

By Theorem 3.3, we have

$$\begin{aligned} \|M_\delta^\sharp(T_{\prod \bar{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \left(\|M_{\varepsilon_1}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} \right) \\ &\quad + C \sum_{j=1}^{m-1} \sum_{\sigma \in C_j^m} \prod_{i=1}^j \|b_{\sigma(i)}\|_{BMO} \|M_{\varepsilon_1}(T_{\prod \bar{b}_{\sigma'}}(\vec{f}))\|_{L^p(v_{\vec{w}})} \\ &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \left(\|M_{\varepsilon_1}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} \right) \\ &\quad + C \sum_{j=1}^{m-1} \sum_{\sigma \in C_j^m} \prod_{i=1}^j \|b_{\sigma(i)}\|_{BMO} \|M_{\varepsilon_1}^\sharp(T_{\prod \bar{b}_{\sigma'}}(\vec{f}))\|_{L^p(v_{\vec{w}})}. \end{aligned}$$

For the purpose of reducing the dimension of BMO functions in the commutators, we apply the Theorem 3.3 to $\|M_{\varepsilon_1}^\sharp(T_{\prod \bar{b}_{\sigma'}}(\vec{f}))(x)\|_{L^p(v_{\vec{w}})}$.

Let $\sigma = \{\sigma(1), \dots, \sigma(j)\}$ and $\sigma' = \{\sigma(j+1), \dots, \sigma(m)\}$, $A_h = \{\sigma_1: \text{any finite subset of } \sigma' \text{ with different elements}\}$ and $\sigma'_1 = \sigma' - \sigma_1$,

$$\begin{aligned} \|M_{\varepsilon_1}^\sharp(T_{\prod \vec{b}_{\sigma'}}(\vec{f}))\|_{L^p(v_{\vec{w}})} &\leq C \prod_{i=j+1}^m \|b_{\sigma(i)}\|_{BMO} \left(\|M_{\varepsilon_2}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} \right) \\ &\quad + C \sum_{h=1}^{m-j-1} \sum_{\sigma_1 \in A_h} \prod_{i=1}^h \|b_{\sigma_1(i)}\|_{BMO} \|M_{\varepsilon_2}(T_{\prod \vec{b}_{\sigma_1'}}(\vec{f}))\|_{L^p(v_{\vec{w}})}. \end{aligned}$$

Repeating the process above and using Theorem 3.1, we can get

$$\begin{aligned} \|M_{\delta}^\sharp(T_{\prod \vec{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \left(A_{m+1}(m, n) \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} + A_1(m, n) \right. \\ &\quad \times \|M_{\varepsilon_1}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + A_2(m, n) \|M_{\varepsilon_2}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \dots \\ &\quad \left. + A_m(m, n) \|M_{\varepsilon_m}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} \right), \end{aligned}$$

where $A_1(m, n), A_2(m, n), \dots, A_{m+1}(m, n)$ are finite real numbers related to m and n .

We can deduce the following conclusion from Lemma 2.5, Lemma 2.7 and $\vec{w} \in A_{\vec{p}/s}$.

$$\begin{aligned} \|T_{\prod \vec{b}}(\vec{f})\|_{L^p(v_{\vec{w}})} &\leq \|M_{\delta}(T_{\prod \vec{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} \\ &\leq C \|M_{\delta}^\sharp(T_{\prod \vec{b}}(\vec{f}))\|_{L^p(v_{\vec{w}})} \\ &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \left(A_{m+1}(m, n) \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} + A_1(m, n) \right. \\ &\quad \times \|M_{\varepsilon_1}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + A_2(m, n) \|M_{\varepsilon_2}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} + \dots \\ &\quad \left. + A_m(m, n) \|M_{\varepsilon_m}(T(\vec{f}))\|_{L^p(v_{\vec{w}})} \right) \\ &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \|\mathcal{M}_s(\vec{f})\|_{L^p(v_{\vec{w}})} \\ &\leq C \prod_{j=1}^m \|b_j\|_{BMO} \|f_j\|_{L^p(w_j)}. \end{aligned}$$

The proof is completed. \square

Remark 3.5. It should be pointed out that the corresponding results of multilinear commutators and iterated commutators of multilinear singular integral operators with Dinis type kernels in [13, 16, 17] and the standard kernels can be deduced as special cases of our results in this paper.

References

- [1] R. R. Coifman and Y. Meyer, *On commutators of singular integrals and bilinear singular integrals*, Trans. Amer. Math. Soc. **212** (1975), 315-331.

- [2] R. R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque **57**, Paris, 1978.
- [3] R. R. Coifman and Y. Meyer, *Commutateurs d'intégrales singulières et opérateurs multilinéaires*, Ann. Inst. Fourier. (Grenoble) **28** (1978), no. 3, 177-202.
- [4] C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), no. 3-4, 137-193.
- [5] L. Grafakos and N. Kalton, *Multilinear Calderón-Zygmund operators on Hardy spaces*, Collect. Math. **52** (2001), no. 2, 169-179.
- [6] L. Grafakos and J. M. Martell, *Extrapolation of weighted norm inequalities for multivariable operators and applications*, J. Geom. Anal. **14** (2004), no. 1, 19-46.
- [7] L. Grafakos and R. H. Torres, *Multilinear Calderón-Zygmund theory*, Adv. Math. **165** (2002), no. 1, 124-164.
- [8] L. Grafakos and R. H. Torres, *Discrete decompositions for bilinear operators and almost diagonal conditions*, Trans. Amer. Math. Soc. **354** (2002), no. 3, 1153-1176.
- [9] L. Grafakos and R. H. Torres, *On multilinear singular integrals of Calderón-Zygmund type*, Publ. Mat.(2002), 57-91.
- [10] A. K. Lerner, S. Ombrosi, C. Pérez, R. H. Torres and R. Trujillo-González, *New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory*, Adv. Math. **220** (2009), no. 4, 1222-1264.
- [11] Y. Lin and S. Z. Lu, *Strongly singular CalderónZygmund operators and their commutators*, Jordan J. Math. Stat. **1** (2008), 31-49.
- [12] Y. Lin and Y. Y. Xiao, *Multilinear singular integral operators with generalized kernels and their multilinear commutators*, Acta Math. Sin. (Engl. Ser.) **33** (2017), no. 11, 1443-1462.
- [13] Y. Lin and N. Zhang, *Sharp maximal and weighted estimates for multilinear iterated commutators of multilinear integrals with generalized kernels*, J. Inequal. Appl. (2017).
- [14] G. Z. Lu and P. Zhang, *Multilinear CalderónZygmund operators with kernels of Dini's type and applications*, Nonlinear Anal. **107** (2014), 92-117.
- [15] C. Pérez and R. H. Torres, *Sharp maximal function estimates for multilinear singular integrals*, Contemp. Math. **320** (2003), 323-331.
- [16] J. Sun and P. Zhang, *Commutators of multilinear CalderónZygmund operators with kernels of Dini type kernels on some function spaces*, J. Nonlinear Sci. Appl. **10** (2017), no. 9, 5002-5019.
- [17] J. Sun and P. Zhang, *Commutators of multilinear CalderónZygmund operators with kernels of Dini's type and applications*, J. Math. Inequal. **13** (2019), no. 4, 1071-1093.

- [18] S. H. Yang, P. F. Li and Y. Lin, *Multiple weight inequalities for multilinear singular integral operators with generalized kernels*, Adv. Math. (China), accepted, to appear.

LIWEN GAO

SCHOOL OF SCIENCE

CHINA UNIVERSITY OF MINING AND TECHNOLOGY, BEIJING

BEIJING 100083, P. R. CHINA

Email address: **gaoliwen1206@163.com**

YAN LIN

SCHOOL OF SCIENCE

CHINA UNIVERSITY OF MINING AND TECHNOLOGY, BEIJING

BEIJING 100083, P. R. CHINA

Email address: **linyan@cumtb.edu.cn**

SHUHUI YANG

SCHOOL OF SCIENCE

CHINA UNIVERSITY OF MINING AND TECHNOLOGY, BEIJING

BEIJING 100083, P. R. CHINA

Email address: **yangshuhui0601@163.com**