

ON THE SUMMABILITY OF RANDOM FOURIER–JACOBI SERIES

Partiswari Maharana¹ and Sabita Sahoo²¹ Department of Mathematics, Sambalpur University, Odisha, India² Department of Mathematics, Sambalpur University, Odisha, India.¹ partiswarimath1@suniv.ac.in² sabitamath@suniv.ac.in

ABSTRACT. This article is a study on the summability of random Fourier–Jacobi series of some functions in different spaces. We consider the random series $\sum_{n=0}^{\infty} a_n A_n(\omega) p_n^{(\gamma, \delta)}(y)$, where $p_n^{(\gamma, \delta)}(y)$, $\gamma, \delta > -1$ are orthonormal Jacobi polynomials, the scalars a_n are Fourier–Jacobi coefficients of a function f and the random variables $A_n(\omega)$ are Fourier–Jacobi coefficients of the symmetric stable process $X(t, \omega)$ of index $\alpha \in [1, 2]$. It is established that the random Fourier–Jacobi series is Θ –summable in probability, if a_n are the Fourier–Jacobi coefficients of function f in the space $C_{[-1,1]}^{(\eta, \tau)}$. The Cesàro (C, ϕ) , $\phi \geq 1$ summability of random Fourier–Jacobi series is shown, for the symmetric stable process $X(t, \omega)$ of index $\alpha \in [1, 2]$ under different conditions on the parameters γ, δ, η and τ . The other cases of summability, such as Riesz, Rogosinski, etc., are also discussed. Further, the Nörlund summability, generalized Nörlund summability, and lower triangular summability of random Fourier–Jacobi series are proved if a_n are the Fourier–Jacobi coefficients of a function $f \in L_{[-1,1]}^{1,(\gamma, \delta)}$, and $A_n(\omega)$ are associated with the symmetric stable process $X(t, \omega)$ of index one. It is observed that the conditions on the parameters γ, δ differ from that of the conditions on γ, δ for the Fourier–Jacobi series of functions f in $L_{[-1,1]}^{1,(\gamma, \delta)}$.

2020 MSC Classification: 60G99, 40G15.

Key words: Cesàro summability; Convergence in probability; Fourier–Jacobi series; Jacobi polynomials; Lower triangular summability; Nörlund summability; Random variables; Stochastic integral; Symmetric stable process.

1. INTRODUCTION

The foremost aspect in the study of Fourier analysis is the convergence of Fourier series of functions. In the general case, the convergence of the Fourier series of all functions is not always possible. However, using some summability methods, convergence is possible in some sense, like in Cesàro, Riesz, Nörlund, etc. The summability of Fourier series in orthogonal polynomials is investigated by many researchers. Also, the weighted convergence of certain sums in the Fourier–Jacobi series was investigated by Lubinsky, Totik [7], and many others.

Further, Nayak, Pattanayak, Mishra [11], Towghi [14], and others studied the summability of random Fourier series. Motivated by the works of Liu and Liu [5, 6], we investigated the convergence of random Fourier–Jacobi series of functions f in various continuous spaces [9] and the space $L_{[-1,1]}^p$ [10].

In this article, we have studied the summability of random Fourier series

$$\sum_{n=0}^{\infty} a_n A_n(\omega) p_n^{(\gamma, \delta)}(y) \quad (1.1)$$

in orthonormal Jacobi polynomial $p_n^{(\gamma, \delta)}(y)$, where a_n are scalars, $A_n(\omega)$ are random variables. The scalars a_n are the Fourier–Jacobi coefficients of a function f defined as

$$a_n := \int_{-1}^1 f(t) p_n^{(\gamma, \delta)}(t) \rho^{(\gamma, \delta)}(t) dt, \quad \gamma, \delta > -1. \quad (1.2)$$

The functions f chosen here are from the continuous space $C_{[-1,1]}^{(\eta,\tau)}$ and the space $L_{[-1,1]}^{p,(\gamma,\delta)}$. In relation to the space $C_{[-1,1]}^{(\eta,\tau)}$, the Θ -summability, Cesàro (C, ϕ) summability, and some other summability of random series (1.1) is established. The Nörlund summability, generalized Nörlund summability, and lower triangular summability of random series (1.1) are established in connection to the space $L_{[-1,1]}^{1,(\gamma,\delta)}$.

The class of continuous functions

$$C_{[-1,1]}^{(\eta,\tau)} := \left\{ f \in C(-1, 1) \mid \lim_{|y| \rightarrow 1} (f\rho^{(\eta,\tau)})(y) = 0 \right\}$$

is a linear space of real-valued continuous functions defined on the interval $(-1, 1)$ equipped with the norm

$$\|f\|_{\rho^{(\eta,\tau)}} := \|f\rho^{(\eta,\tau)}\| := \max_{y \in [-1, 1]} \left\{ |(f\rho^{(\eta,\tau)})(y)| \right\},$$

where

$$\rho^{(\eta,\tau)}(y) := (1-y)^\eta(1+y)^\tau, (y \in [-1, 1])$$

is the Jacobi weight with parameters $\eta, \tau \geq 0$.

The space $L_{[-1,1]}^{p,(\gamma,\delta)}$ with the weight function $\rho^{(\gamma,\delta)}(y) := (1-y)^\gamma(1+y)^\delta$, $\gamma, \delta > -1$, is the class of all measurable functions f on the segment $[-1, 1]$, such that

$$\int_{-1}^1 |f(y)\rho^{(\gamma,\delta)}(y)|^p dy < \infty.$$

This space is equipped with the norm

$$\|f\|_{L_{[-1,1]}^{p,(\gamma,\delta)}} = \left\{ \int_{-1}^1 |f(y)\rho^{(\gamma,\delta)}(y)|^p dy \right\}^{\frac{1}{p}}, \text{ for } p \geq 1.$$

In random series (1.1), the random variables $A_n(\omega)$ are the Fourier–Jacobi coefficients of the symmetric stable process $X(t, \omega)$ of index $\alpha \in [1, 2]$. We know that, the stochastic integral

$$\int_a^b f(t) dX(t, \omega) \quad (1.3)$$

is defined in the sense of probability and is a random variable, if f is a continuous function in $[a, b]$ and $X(t, \omega)$, for $t \in \mathbb{R}$, is a continuous stochastic process with independent increments [8]. Moreover, the stochastic integral (1.3) is defined in the sense of probability, for $f \in L_{[a,b]}^p$, $p \geq \alpha \geq 1$ and $X(t, \omega)$ is a symmetric stable process of index $\alpha \in [1, 2]$ [11]. Accordingly, if $f \in L_{[-1,1]}^{p,(\eta,\tau)}$, $p \geq 1, \eta, \tau > -1$ i.e. $f\rho^{(\eta,\tau)} \in L_{[-1,1]}^p$, then the stochastic integral

$$\int_{-1}^1 f(t)\rho^{(\eta,\tau)}(t) dX(t, \omega)$$

exists in probability, for $p \geq \alpha \geq 1$. A sequence of random variables X_n is said to converge in probability to a random variable X , if $\lim_{n \rightarrow \infty} P(|X_n - X| > \epsilon) = 0$, for $\epsilon > 0$. In particular, if $f(t)$ is the orthonormal Jacobi polynomial $p_n^{(\gamma,\delta)}(t)$, $\gamma, \delta > -1$, then $p_n^{(\gamma,\delta)}(t)\rho^{(\eta,\tau)}(t)$ is continuous in $[-1, 1]$, for all $\eta, \tau \geq 0$ and is in both the spaces $C_{[-1,1]}^{(\eta,\tau)}$ and $L_{[-1,1]}^{p,(\eta,\tau)}$. Hence the integrals

$$A_n(\omega) := \int_{-1}^1 p_n^{(\gamma,\delta)}(t)\rho^{(\eta,\tau)}(t) dX(t, \omega) \quad (1.4)$$

exist and are random variables, which are not independent, for each $n = 1, 2, \dots$. The series (1.1) is called the random Fourier–Jacobi series of function f . The summability of random Fourier–Jacobi series (1.1) of functions f in the space $C_{[-1,1]}^{(\eta,\tau)}$ and $L_{[-1,1]}^{p,(\gamma,\delta)}$, $p \geq 1$ are studied in this article.

The summability in the space $C_{[-1,1]}^{(\eta,\tau)}$ is discussed by using the summation matrix Θ which is a lower triangular infinite matrix (see equation (2.2)). We find different summability methods for different values of the entries of the matrix Θ . The various summability of the random series (1.1) depend on different conditions on the summation matrix Θ . The Nörlund summability and some other variants of it are established, for the random Fourier–Jacobi series of functions in the space $L_{[-1,1]}^{p,(\gamma,\delta)}$, $p = 1$. Each of these summability methods has conditions on the parameters γ, δ that differ from the conditions on these parameters in the case of the Fourier–Jacobi series of functions $f \in L_{[-1,1]}^{1,(\gamma,\delta)}$. If $0 \leq \gamma < 1/2$ and $\delta \geq 0$, the random Fourier–Jacobi series is Nörlund summable. In this case, however the Fourier–Jacobi series is summable, for $-1/2 \leq \gamma < 1/2$ and $\delta > -1/2$. The generalized Nörlund summability of random Fourier–Jacobi series holds for $\gamma \geq 0$, $\delta - \gamma > 1$ and $\delta + \gamma > 0$, but it is observed for the Fourier–Jacobi series if $\gamma \geq 0$, $\delta - \gamma > 1$ and $\delta + \gamma \geq -1$. Furthermore, the random Fourier–Jacobi series is lower triangular summable for $0 \geq \gamma \geq 1/2$ and $\delta \geq 0$. The Fourier–Jacobi series, on the other hand, is lower triangular summable if $-1/2 \geq \gamma \geq 1/2$ and $\delta > -1/2$.

This article is organized as follows. The preliminaries and some results relevant to this work are stated in Section 2. The Θ -summability in general of the random Fourier–Jacobi series (1.1) associated with symmetric stable process of index $\alpha \in [1, 2]$ under some conditions on γ, δ, η and τ is discussed in Section 3. Further, the other summability of the random Fourier–Jacobi series (1.1), such as Cesáro, Riesz, Rogosinski, etc., for different set of values of entries in the matrix Θ are established. Section 4 investigates the summability of random Fourier–Jacobi series (1.1) of functions $f \in L_{[-1,1]}^{1,(\gamma,\delta)}$. We prove the Nörlund summability, generalized Nörlund summability, and lower triangular matrix summability of the random series (1.1) associated with the symmetric stable process of index $\alpha = 1$, for the parameter $\gamma, \delta \geq 0$ at a particular point only.

2. PRELIMINARIES AND RESULTS ON SUMMABILITY OF FOURIER–JACOBI SERIES

Some definitions and results from the work of Chripkó, Raghuvanshi, Thorpe, and Dhakal [3, 4, 12, 13] are presented here which will be of use in proving our results in next sections.

Chripkó [3] discussed the Θ -summability of Fourier–Jacobi series

$$\sum_{n=0}^{\infty} a_n p_n^{(\gamma,\delta)}(y) \quad (2.1)$$

in the weighted space $C_{[-1,1]}^{(\eta,\tau)}$ of continuous functions. He considered the summation matrix Θ as

$$\Theta := \begin{bmatrix} \theta_{0,1} & & & \\ \theta_{0,2} & \theta_{1,2} & & \\ \theta_{0,3} & \theta_{1,3} & \theta_{2,3} & \\ \vdots & \vdots & \vdots & \\ \vdots & \vdots & \ddots & \ddots \end{bmatrix}, \quad (2.2)$$

where $\theta_{k,n}$'s are the real entries. The Θ -sum of a Fourier–Jacobi series (2.1) is defined as

$$\mathbf{s}_n^{\Theta,(\gamma,\delta)}(f, y) := \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma,\delta)}(y), \quad (y \in [-1, 1], n \in \mathbb{N}), \quad (2.3)$$

where a_k are the Fourier–Jacobi coefficients of a function $f \in C_{[-1,1]}^{(\eta,\tau)}$.

Chripkó considered some choices of the summation matrix Θ , which are stated below.

Case-1

If $\theta_{k,n} := 1$ ($k = 0, 1, 2, \dots, n-1, n \in \mathbb{N}$), then the n th Fourier–Jacobi series (2.3) of f is equal to the n th partial sum of the Fourier–Jacobi series (2.1).

Case-2

Let

$$\theta_{k,n} := \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} \quad (\phi \geq 0, k = 0, \dots, n-1, n \in \mathbb{N}),$$

where $B_0^{(\phi)} := 1$, $B_m^{(\phi)} := \binom{m+\phi}{m} = \frac{(\phi+1)\dots(\phi+m)}{m!}$, ($m \in \mathbb{N}$). Then $\Theta := (\theta_{k,n})$ is the Cesàro summation matrix and the Cesàro means of the Fourier–Jacobi series (2.1) is defined as

$$\begin{aligned} \sigma_n^{(\phi),\gamma,\delta}(f, y) &:= \sum_{k=0}^{n-1} \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} a_k p_k^{(\gamma,\delta)}(y), \quad y \in [-1, 1], \quad n \in \mathbb{N} \\ &= \int_{-1}^1 \mathbf{F}_n^{(\phi),\gamma,\delta}(y, t) \rho^{(\gamma,\delta)}(t) dt, \end{aligned} \quad (2.4)$$

where

$$\mathbf{F}_n^{(\phi),\gamma,\delta}(y, t) := \sum_{k=0}^{n-1} \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} p_k^{(\gamma,\delta)}(t) p_k^{(\gamma,\delta)}(y).$$

If $\phi = 0$, then it is the same as in Case-1. For $\phi \geq 1$, the sum (2.4) is known as Cesàro (C, ϕ) means of the Fourier–Jacobi series (2.1). In particular, for $\phi = 1$, we obtain the Fejer summation of the Fourier–Jacobi series (2.1).

Case-3

Let $\theta_{k,n} := \left[1 - \left(\frac{k}{n}\right)^\nu\right]^\mu$, $k = 0, 1, \dots, n-1, n \in \mathbb{N}$, where $\nu, \mu \geq 0$ are constant real numbers. Then

$$\mathbf{s}_n^{(\gamma,\delta)\nu,\mu}(f, y) := \sum_{k=0}^{n-1} \left[1 - \left(\frac{k}{n}\right)^\nu\right]^\mu a_k p_k^{(\gamma,\delta)}(y), \quad (y \in [-1, 1])$$

is the Riesz (R, ν, μ) sum of the Fourier–Jacobi series (2.1).

Case-4

Let

$$\theta_{k,n} := \begin{pmatrix} 1, & \text{if } 0 \leq \frac{k}{n} \leq s \\ \frac{\frac{k}{n} - 1}{s - 1}, & \text{if } s < \frac{k}{n} \leq 1 \end{pmatrix},$$

where $k \in 0, 1, 2, \dots, n-1, n \in \mathbb{N}$ and $s \in (0, 1)$. Then

$$\mathbf{s}_n^{(\gamma,\delta),s}(f, y) := \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma,\delta)}(y)$$

is the de la Vallée Poussin means of the Fourier–Jacobi series (2.1).

Case-5

Let $\theta_{k,n} := \cos \frac{\pi k}{2n}$, for $k = 0, 1, \dots, n-1, n \in \mathbb{N}$, then

$$\mathbf{s}_n^{(\gamma,\delta),R}(f, y) := \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma,\delta)}(y), \quad y \in [-1, 1]$$

is known as the Rogosinski means of the Fourier–Jacobi series (2.1).

Choudhary [2] discussed the regular Nörlund summability (also known as Nörlund summability) of Fourier–Jacobi series (2.1). The regular Nörlund sum of a series is defined below:

Let $\sum_{n=0}^{\infty} a_n$ be a given infinite series with the sequence of partial sums $\{s_n\}$. Let p_n be a sequence of constants, real or complex and let $P_n = p_0 + p_1 + \dots + p_n$. Define the sequence to sequence transformation h_n as

$$h_n := \frac{1}{P_n} \sum_{v=0}^n p_{n-v} s_v = \frac{1}{P_n} \sum_{v=0}^n p_v s_{n-v}, \quad P_n \neq 0$$

The sequence $\{h_n\}$ is said to be Nörlund means of $\{s_n\}$, generated by the constants $\{p_n\}$. The series $\sum_{n=0}^{\infty} a_n$ is Nörlund summable to the sum s if $\lim_{n \rightarrow \infty} h_n$ exists and equal to s .

The Nörlund sum of Fourier–Jacobi series at the point $y = 1$ is

$$\mathbf{h}_n^{(\gamma, \delta)}(f, y) := \frac{1}{P_n} \sum_{k=0}^n p_{n-k} \mathbf{s}_k^{(\gamma, \delta)}(f, 1),$$

where $\mathbf{s}_k^{(\gamma, \delta)}(f, 1)$ is the n th partial sum of the Fourier–Jacobi series (2.1) at the point $y = 1$. We use the symbol (\mathbf{N}, p_n) for it. The following theorem is established by Choudhary.

Theorem 1. [2] *Let (\mathbf{N}, p_n) be a Nörlund method defined by a real non–negative monotonic non–increasing sequence of coefficients $\{p_n\}$ such that*

$$P_n \rightarrow \infty \text{ as } n \rightarrow \infty. \quad (2.5)$$

If

$$\Psi(t) = O\left[\frac{p(1/t)}{P(1/t)} t^{(2\gamma+1)}\right], \text{ as } t \rightarrow 0, \quad (2.6)$$

and

$$\sum_{n=0}^{\infty} \frac{n^{\gamma+1/2}}{P_n} < \infty, \quad (2.7)$$

then the series (2.1) of some function $f \in L_{[-1,1]}^{1,(\gamma, \delta)}$ is (\mathbf{N}, p_n) summable at the point $y = 1$ to the sum A provided $-1/2 \leq \gamma < 1/2$, $\delta > -1/2$ and the antipole condition

$$\int_{-1}^b (1+x)^{\delta/2-3/4} |f(x)| dx < \infty, \quad (2.8)$$

is satisfied, for fixed b .

Raghuvanshi [12] discussed the generalized Nörlund summability of Fourier–Jacobi series (2.1). The generalized Nörlund summability of a series is defined as below:

Let $\{s_n\}$ be the sequence of partial sums of an infinite series $\sum_{n=0}^{\infty} \mathbf{a}_n$. Let $\{r_n\}$ and $\{q_n\}$ be any two sequences of positive real constants with R_n and Q_n as their n th partial sums respectively, and let

$$(q * r)_n := \sum_{k=0}^n q_{n-k} r_k = \sum_{k=0}^n q_k r_{n-k}$$

tends to infinity as $n \rightarrow \infty$. As in [1], define the sequence to sequence transformation as

$$t_n^{q,r} := \frac{1}{(q * r)_n} \sum_{k=0}^n q_{n-k} r_k s_k.$$

If $t_n^{q,r} \rightarrow s$ as $n \rightarrow \infty$, then the infinite series $\sum_{n=0}^{\infty} \mathbf{a}_n$ is said to be generalized Nörlund summable to s . We use the notation $|\mathbf{N}, q_n, r_n|$ for generalized Nörlund summation. The $|\mathbf{N}, q_n, r_n|$ sum of the Fourier–Jacobi series (2.1) at the point $y = 1$ is defined as

$$\mathbf{u}_n^{(\gamma, \delta)}(f, 1) := \frac{1}{(q * r)_n} \sum_{k=0}^n q_k r_{n-k} \mathbf{s}_k^{(\gamma, \delta)}(f, 1),$$

where $\mathbf{s}_k^{(\gamma, \delta)}(f, 1)$ is the n th partial sum of the Fourier–Jacobi series (2.1) at the point $y = 1$. The following result is established by Raghuvanshi.

Theorem 2. [12] Let $|\mathbf{N}, q_n, r_n|$ be the generalized summability method defined by a non-negative real constants sequences $\{q_n\}$, $\{r_n\}$ and let $\gamma > -1/2$, $\delta - \gamma > 1$, $\delta + \gamma \geq -1$ such that

$$\sum_{k=2}^n \frac{(q * r)_k}{k^{\gamma+(1/2)} \log k} = O\left(\frac{(q * r)_n}{n^{\gamma+(1/2)}}\right) \text{ as } n \rightarrow \infty. \quad (2.9)$$

Also suppose that

$$\int_{1-t}^t |f(u) - A| du = O\left(\frac{t}{\log(1/t)}\right) \text{ as } t \rightarrow 0, \quad (2.10)$$

and the antipole condition

$$\int_{-1}^b (1+y)^{(\delta-\gamma-1)/2} |f(y)| dy < \infty, \text{ } b \text{ fixed}, \quad (2.11)$$

are satisfied, then the series (2.1) is $|\mathbf{N}, q_n, r_n|$ summable to the sum A at the point $y = 1$.

Dhakar [4] studied the summability of Jacobi series in lower triangular matrix method. The lower triangular matrix method of a series is defined as below:

Let $M = (\mathbf{b}_{n,k})$ be an infinite lower triangular matrix satisfying the Silverman-Töeplitz conditions [13] of regularity i.e.

$$\sum_{k=0}^n \mathbf{b}_{n,k} \rightarrow 1 \text{ as } n \rightarrow \infty,$$

where $\mathbf{b}_{n,k} = 0$, for $k > n$ and $\sum_{k=0}^n |\mathbf{b}_{n,k}| \leq M$, for M a finite +ve constant.

Consider s_n be the n th partial sum of an infinite series $\sum_{n=0}^{\infty} \mathbf{a}_n$. Let $\{t_n\}$ be the lower triangular matrix sums of $\{s_n\}$, generated by the sequence of coefficient $(\mathbf{b}_{n,k})$ i.e. the sequence-to-sequence transformation is

$$t_n := \sum_{k=0}^n \mathbf{b}_{n,k} s_k.$$

If $t_n \rightarrow s$ as $n \rightarrow \infty$, then the series $\sum_{n=0}^{\infty} \mathbf{a}_n$ is said to be summable by the lower triangular matrix method to s . We use the notation (\mathbf{T}) for lower triangular matrix summation.

The (\mathbf{T}) summation of Jacobi series (2.1) at the point $y = 1$ is

$$\mathbf{t}_n^{(\gamma, \delta)}(f, 1) = \sum_{k=0}^n \mathbf{b}_{n,k} \mathbf{s}_k^{(\gamma, \delta)}(f, 1),$$

where $\mathbf{s}_k^{(\gamma, \delta)}(f, 1)$ is the n th partial sum of the Fourier-Jacobi series (2.1) at the point one. The following result is established by Dhakar on (\mathbf{T}) summability of Fourier-Jacobi series (2.1) of the function $f \in L_{[-1,1]}^{1,(\gamma, \delta)}$.

Theorem 3. [4] Let $M = (\mathbf{b}_{n,k})$ be an infinite lower triangular regular matrix such that the element $(\mathbf{b}_{n,k})$ is positive, monotonic increasing in k , $0 \leq k \leq n$,

$$D_{n,\tau} = \sum_{k=n-\tau}^n \mathbf{b}_{n,k}, \quad D_{n,n} = 1, \text{ for all } n, \quad (2.12)$$

and

$$n^{\gamma+1/2} D_{n, [1/\beta]} = o(1), \quad 0 < \beta < \pi \text{ as } n \rightarrow \infty, \quad (2.13)$$

where $\tau = \text{Integral part of } \frac{1}{\phi} = [\frac{1}{\phi}]$ and $0 \leq \phi \leq \pi$. For $-1/2 \leq \gamma \leq 1/2$, $\delta > -1/2$, if

$$\int_{1-t}^1 |f(u) - A| du = o\left(\frac{t}{\xi(1/t) \log(1/t)}\right) \text{ as } t \rightarrow 0, \quad (2.14)$$

then the Jacobi series (2.1) is (\mathbf{T}) summable to the sum A at $y = 1$ provided $\xi(t)$ is positive monotonic non-decreasing function of t such that

$$\sum_{k=a}^n \frac{D_{n,k}}{k^{(2\gamma+3)/2} \xi(k) \log k} = O\left(\frac{1}{n^{(2\gamma+1)/2}}\right), \quad (2.15)$$

and satisfy the antipole condition

$$\int_0^{1/n} t^{\delta-1/2} |f(-cost) - A| dt = o(1) \text{ as } n \rightarrow \infty. \quad (2.16)$$

3. SUMMABILITY OF RANDOM FOURIER–JACOBI SERIES IN THE SPACE $C_{[-1,1]}^{(\eta,\tau)}$

Consider the random Fourier–Jacobi series

$$\sum_{n=0}^{\infty} a_n A_n(\omega) p_n^{(\gamma,\delta)}(y), \quad (3.1)$$

where the scalars a_n are the Fourier–Jacobi coefficients of a function $f \in C_{[-1,1]}^{(\eta,\tau)}$, $\eta, \tau \geq 0$ defined as in (1.2) and the random variables $A_n(\omega)$ are defined as in (1.4). The Θ –sum of the random Fourier–Jacobi series (3.1) is

$$\mathbf{S}_n^{\Theta,(\gamma,\delta)}(f, y, \omega) := \sum_{k=0}^{n-1} \theta_{k,n} a_k A_k(\omega) p_k^{(\gamma,\delta)}(y). \quad (3.2)$$

We will use the following notations on Θ –summation matrix same as in Chripkó [3].

$$T_1 : \lim_{n \rightarrow \infty} (1 - \theta_{k,n}) = 0, \text{ for all fixed } k = 0, 1, 2, \dots$$

$$T_2 : \theta_{n-1,n} = O\left(\frac{1}{n}\right), \quad (n \in \mathbb{N}).$$

$$T_3 : \Delta^2 \theta_{k-1,n} = O\left(\frac{1}{n^2}\right), \quad (k = 1, 2, \dots, n-1, n \in \mathbb{N}).$$

$$T_4 : \Delta^2 \theta_{k-1,n} \quad (k = 1, 2, \dots, n-1, n \in \mathbb{N}) \text{ is of constant sign.}$$

$$T_5 : \operatorname{sgn} \Delta^2 \theta_{k-1,n} = \operatorname{sgn} \theta_{n-1,n} \quad (k = 1, 2, \dots, n-1, n \in \mathbb{N}),$$

where $\Delta^2 \theta_{k,n} := \Delta \theta_{k+1,n} - \Delta \theta_{k,n}$, $\Delta \theta_{k,n} := \theta_{k+1,n} - \theta_{k,n}$, $(\theta_{n,n} := 0)$.

Furthermore, the following notations will be used.

(A) if (T_1) , (T_2) and (T_3) hold,

(B) if (T_1) , (T_2) and (T_4) hold,

(C) if (T_1) , (T_5) hold.

The convergence of Θ –sum (3.2) of the random Fourier–Jacobi series (3.1) is proved in Theorem 5 by imposing some restrictions on $\gamma, \delta, \eta, \tau$ and the entries of summation matrix Θ . To prove this theorem, we need the following result.

Lemma 4. [11] *Let $X(t, \omega)$ be a symmetric stable process of index α , $1 \leq \alpha \leq 2$ and $f(t)$ be any function in $L_{[a,b]}^p$, $p \geq 1$, then for all $\epsilon > 0$,*

$$P\left(\left|\int_a^b f(t) dX(t, \omega)\right| > \epsilon\right) \leq \frac{C 2^{\alpha+1}}{(\alpha+1)\epsilon'^{\alpha}} \int_a^b |f(t)|^{\alpha} dt,$$

where C is a positive constant and $\epsilon' < \epsilon$.

Theorem 5. *The Θ –sum (3.2) of random Fourier–Jacobi series (3.1) converges in probability to the stochastic integral*

$$\int_{-1}^1 f(y, t) \rho^{(\eta,\tau)}(t) dX(t, \omega), \quad (3.3)$$

if the parameters $\gamma, \delta \geq -1/2, \eta, \tau \geq 0$ satisfy the following conditions

$$\frac{\gamma}{2} - \frac{1}{4} < \eta < \frac{\gamma}{2} + \frac{3}{4}, \text{ and } \frac{\delta}{2} - \frac{1}{4} < \tau < \frac{\delta}{2} + \frac{3}{4}, \quad (3.4)$$

with the entries of Θ -matrix satisfying either (A), (B) or (C).

Proof. The integral form of the n th partial Θ -sum of the random Fourier–Jacobi series (3.1) involving Jacobi polynomials is

$$\begin{aligned} \mathbf{S}_n^{\Theta, (\gamma, \delta)}(f, y, \omega) &:= \sum_{k=0}^{n-1} \theta_{k,n} a_k A_k(\omega) p_k^{(\gamma, \delta)}(y) \\ &= \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma, \delta)}(y) \int_{-1}^1 p_k^{(\gamma, \delta)}(t) \rho^{(\eta, \tau)}(t) dX(t, \omega) \\ &= \int_{-1}^1 \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma, \delta)}(y) p_k^{(\gamma, \delta)}(t) \rho^{(\eta, \tau)}(t) dX(t, \omega) \\ &= \int_{-1}^1 \mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) \rho^{(\eta, \tau)}(t) dX(t, \omega), \end{aligned}$$

where

$$\mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) := \sum_{k=0}^{n-1} \theta_{k,n} a_k p_k^{(\gamma, \delta)}(t) p_k^{(\gamma, \delta)}(y).$$

Now

$$\begin{aligned} &\mathbf{S}_n^{\Theta, (\gamma, \delta)}(f, y, \omega) - \mathbf{S}_m^{\Theta, (\gamma, \delta)}(f, y, \omega) \\ &= \int_{-1}^1 \left(\mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) - \mathbf{s}_m^{\Theta, (\gamma, \delta)}(f, y, t) \right) \rho^{(\eta, \tau)}(t) dX(t, \omega). \end{aligned}$$

By the inequality in Lemma 4, we obtain

$$\begin{aligned} &P\left(\left| \mathbf{S}_n^{\Theta, (\gamma, \delta)}(f, y, \omega) - \mathbf{S}_m^{\Theta, (\gamma, \delta)}(f, y, \omega) \right| > \epsilon \right) \\ &= P\left(\left| \int_{-1}^1 \left(\mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) - \mathbf{s}_m^{\Theta, (\gamma, \delta)}(f, y, t) \right) \rho^{(\eta, \tau)}(t) dX(t, \omega) \right| > \epsilon \right) \\ &\leq \frac{C2^{\alpha+1}}{(\alpha+1)\epsilon'^{\alpha}} \int_{-1}^1 \left| \left(\mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) - \mathbf{s}_m^{\Theta, (\gamma, \delta)}(f, y, t) \right) \rho^{(\eta, \tau)}(t) \right|^{\alpha} dt, \text{ for } \epsilon' < \epsilon. \end{aligned}$$

Invoking Chripkó result (see [3, Theorem 3.2]) on the convergence of Fourier–Jacobi series (2.1), we know that $\mathbf{s}_n^{\Theta, (\gamma, \delta)}$ converges uniformly in $C_{[-1,1]}^{(\eta, \tau)}$, if a_n are the Fourier–Jacobi coefficients of a function f in $C_{[-1,1]}^{(\eta, \tau)}$ and $\gamma, \delta, \eta, \tau$ satisfy the conditions in equation (3.4). Hence $\mathbf{s}_n^{\Theta, (\gamma, \delta)}$ is a Cauchy sequence.

This yields, for $\alpha \in [1, 2]$ and $f \in C_{[-1,1]}^{(\eta, \tau)}$,

$$\lim_{n, m \rightarrow \infty} \int_{-1}^1 \left| \left(\mathbf{s}_n^{\Theta, (\gamma, \delta)}(f, y, t) - \mathbf{s}_m^{\Theta, (\gamma, \delta)}(f, y, t) \right) \rho^{(\eta, \tau)}(t) \right|^{\alpha} dt = 0.$$

This result holds uniformly for all y, t in the range $[-1, 1]$, implying that $\mathbf{S}_n^{\Theta, (\gamma, \delta)}(f, y, \omega)$ is a Cauchy sequence in the sense that it converges in probability. Hence, it will converge to a random variable. Now we will show that the sequence $\mathbf{S}_n^{\Theta, (\gamma, \delta)}(f, y, \omega)$ converges to the stochastic integral $\int_{-1}^1 f(y, t) \rho^{(\eta, \tau)}(t) dX(t, \omega)$

in probability. By Lemma 4,

$$\begin{aligned} & P\left(\left|\mathbf{S}_n^{\Theta,(\gamma,\delta)}(f, y, \omega) - \int_{-1}^1 f(y, t)\rho^{(\eta,\tau)}(t)dX(t, \omega)\right| > \epsilon\right) \\ &= P\left(\left|\int_{-1}^1 \left(\mathbf{s}_n^{\Theta,(\gamma,\delta)}(f, y, t) - f(y, t)\right)\rho^{(\eta,\tau)}(t)dX(t, \omega)\right| > \epsilon\right) \\ &\leq \frac{C2^{\alpha+1}}{(\alpha+1)\epsilon'^{\alpha}} \int_{-1}^1 \left|\left(\mathbf{s}_n^{\Theta,(\gamma,\delta)}(f, y, t) - f(y, t)\right)\rho^{(\eta,\tau)}(t)\right|^{\alpha} dt, \text{ for } \epsilon' < \epsilon. \end{aligned}$$

If $\gamma, \delta, \eta, \tau$ satisfy the conditions in (3.4), then by Chripkó [3, Theorem 3.2],

$$\lim_{n \rightarrow \infty} \int_{-1}^1 \left|\left(\mathbf{s}_n^{\Theta,(\gamma,\delta)}(f, y, t) - f(y, t)\right)\rho^{(\eta,\tau)}(t)\right| dt = 0.$$

Hence, for $\alpha \in [1, 2]$,

$$\lim_{n \rightarrow \infty} \int_{-1}^1 \left|\left(\mathbf{s}_n^{\Theta,(\gamma,\delta)}(f, y, t) - f(y, t)\right)\rho^{(\eta,\tau)}(t)\right|^{\alpha} dt = 0.$$

This proves that the Θ –sum (3.2) of random Fourier–Jacobi series (3.1) converges in probability to the stochastic integral (3.3). \square

The following theorem demonstrates that the random Fourier–Jacobi series (3.1) is Cesàro (C, ϕ) , $\phi \geq 1$ summable, if a_n are the Fourier–Jacobi coefficients of some function f in the space $C_{[-1,1]}^{(\eta,\tau)}$ under the same conditions on γ, δ, η and τ as stated in previous theorem.

Theorem 6. *The random Fourier–Jacobi series (3.1) is Cesàro (C, ϕ) , $\phi \geq 1$ summable in probability to the stochastic integral (3.3), if $\eta, \tau \geq 0, \gamma, \delta \geq -1/2$ and the entries of Θ –matrix satisfy the conditions (3.4) in Theorem 5.*

Proof. The n th (C, ϕ) , $\phi \geq 1$ means of random Fourier–Jacobi series (3.1) is

$$\mathbf{Q}_n^{(\phi),\gamma,\delta}(f, y, \omega) := \sum_{k=0}^{n-1} \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} \mathbf{S}_k^{(\gamma,\delta)}(f, y, \omega), \quad (3.5)$$

where $B_m^n = \frac{\Gamma(m+n+1)}{\Gamma(m+1)\Gamma(n+1)}$ and $\mathbf{S}_k^{(\gamma,\delta)}(f, y, \omega)$ is the n th partial sum of random Fourier–Jacobi series (3.1). The (C, ϕ) means (3.5) can be expressed in the integral form as

$$\begin{aligned} \mathbf{Q}_n^{(\phi),\gamma,\delta}(f, y, \omega) &:= \sum_{k=0}^{n-1} \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} \mathbf{S}_k^{(\gamma,\delta)}(f, y, \omega) \\ &= \int_{-1}^1 \sigma_n^{(\phi),\gamma,\delta}(f, y, t)\rho^{(\eta,\tau)}(t)dX(t, \omega), \end{aligned}$$

where

$$\sigma_n^{(\phi),\gamma,\delta}(f, y, t) := \sum_{k=0}^{n-1} \frac{B_{n-k-1}^{(\phi)}}{B_{n-1}^{(\phi)}} \mathbf{s}_k^{(\gamma,\delta)}(f, y, t),$$

and $\mathbf{s}_n^{(\gamma,\delta)}(f, y)$ is the n th partial sum of Fourier–Jacobi series (2.1).

Now

$$\begin{aligned} & \mathbf{Q}_n^{(\phi),\gamma,\delta}(f, y, \omega) - \mathbf{Q}_m^{(\phi),\gamma,\delta}(f, y, \omega) \\ &= \int_{-1}^1 \left(\sigma_n^{(\phi),\gamma,\delta}(f, y, t) - \sigma_m^{(\phi),\gamma,\delta}(f, y, t)\right)\rho^{(\eta,\tau)}(t)dX(t, \omega). \end{aligned}$$

By the inequality in Lemma 4, we obtain $\mathbf{Q}_n^{(\phi), \gamma, \delta}(f, y, \omega)$ is a Cauchy sequence. With the help of Chripkó result (see [3, Corollary 3.4]) on (C, ϕ) , $\phi \geq 1$ summation and the Lemma 4, the Cesàro means of the random series (3.1) converges in probability to the stochastic integral $\int_{-1}^1 f(y, t) \rho^{(\eta, \tau)}(t) dX(t, \omega)$ in probability. \square

The following theorem states the summability of random Fourier–Jacobi series (3.1) in some other means under the same conditions (3.4) in Theorem 5 by using the Lemma 4, for f in $C_{[-1,1]}^{(\eta, \tau)}$, $\eta, \tau \geq 0$.

Theorem 7. *The random Fourier–Jacobi series (3.1) is*

- (a) *Riesz* (R, μ, ν) , for $\mu, \nu \geq 1$,
- (b) *de la Vallée Poussin*, for every $s \in (0, 1)$,
- (c) *Rogosinski summable in probability to the stochastic integral (3.3)*, if $\eta, \tau \geq 0, \gamma, \delta \geq -1/2$ satisfy the conditions in Theorem 5.

4. SUMMABILITY OF RANDOM FOURIER–JACOBI SERIES IN $L_{[-1,1]}^{1,(\gamma, \delta)}$ SPACE

Let $f \in L_{[-1,1]}^{1,(\gamma, \delta)}$, $\gamma, \delta \geq 0$ and a_n be the Fourier–Jacobi coefficients of a function f . Consider the random Fourier–Jacobi series as

$$\sum_{n=0}^{\infty} a_n A_n(\omega) p_n^{(\gamma, \delta)}(y), \quad (4.1)$$

where a_n are Fourier–Jacobi coefficients of a function $f \in L_{[-1,1]}^{1,(\gamma, \delta)}$ and $A_n(\omega)$ defined as

$$\int_{-1}^1 p_n^{(\gamma, \delta)}(t) \rho^{(\gamma, \delta)}(t) dX(t, \omega) \quad (4.2)$$

are the Fourier–Jacobi coefficients of symmetric stable process $X(t, \omega)$ of index $\alpha = 1$. It is shown in the following theorem that the Nörlund (\mathbf{N}, p_n) sum of the random Fourier–Jacobi series (4.1) converges in probability to the stochastic integral

$$\int_{-1}^1 A \rho^{(\gamma, \delta)}(t) dX(t, \omega), \quad (4.3)$$

where A is a positive fixed constant.

Theorem 8. *If p_n is the non-negative non-increasing sequences satisfy the conditions (2.5), (2.6), (2.7) along with the antipole condition (2.8) stated in the Theorem 1 and $0 \leq \gamma < 1/2, \delta \geq 0$, then the random Fourier–Jacobi series (4.1) is (\mathbf{N}, p_n) summable in probability to the stochastic integral (4.3) at the point one.*

Proof. Let

$$\mathbf{S}_n^{(\gamma, \delta)}(f, y, \omega) := \sum_{k=0}^n a_k A_k(\omega) p_k^{(\gamma, \delta)}(y) \quad (4.4)$$

be the n th partial sum of the random Fourier–Jacobi series (4.1). For $y = 1$, the integral form of $\mathbf{S}_n^{(\gamma, \delta)}(f, 1, \omega)$ is

$$\begin{aligned} \mathbf{S}_n^{(\gamma, \delta)}(f, 1, \omega) &:= \sum_{k=0}^n a_k A_k(\omega) p_k^{(\gamma, \delta)}(1) \\ &= \int_{-1}^1 \mathbf{s}_n^{(\gamma, \delta)}(1, t) \rho^{(\gamma, \delta)}(t) dX(t, \omega), \end{aligned} \quad (4.5)$$

where

$$\mathbf{s}_n^{(\gamma, \delta)}(f, 1, t) := \sum_{k=0}^n a_k p_k^{(\gamma, \delta)}(t) p_k^{(\gamma, \delta)}(1).$$

The (\mathbf{N}, p_n) means of the random Fourier–Jacobi series (4.1) at the point $y = 1$ is

$$\mathbf{H}_n^{(\gamma, \delta)}(f, 1, \omega) := \frac{1}{P_n} \sum_{k=0}^n p_{n-k} \mathbf{S}_k^{(\gamma, \delta)}(f, 1, \omega). \quad (4.6)$$

The integral form of (\mathbf{N}, p_n) means (4.6) is

$$\begin{aligned} \mathbf{H}_n^{(\gamma, \delta)}(f, 1, \omega) &:= \frac{1}{P_n} \sum_{k=0}^n p_{n-k} \mathbf{S}_n^{(\gamma, \delta)}(f, 1, \omega) \\ &= \int_{-1}^1 \mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) \rho^{(\gamma, \delta)}(t) dX(t, \omega), \end{aligned}$$

where

$$\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) := \frac{1}{P_n} \sum_{k=0}^n p_{n-k} \mathbf{s}_n^{(\gamma, \delta)}(f, 1, t).$$

By Lemma 4, for $\alpha = 1$,

$$\begin{aligned} &P\left(\left|\mathbf{H}_n^{(\gamma, \delta)}(f, 1, \omega) - \mathbf{H}_m^{(\gamma, \delta)}(f, 1, \omega)\right| > \epsilon\right) \\ &\leq \frac{C2^{\alpha+1}}{\epsilon^\alpha(\alpha+1)} \int_{-1}^1 \left| \left(\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) - \mathbf{h}_m^{(\gamma, \delta)}(f, 1, t) \right) \rho^{(\gamma, \delta)}(t) \right| dt. \end{aligned}$$

If the equations (2.5), (2.6) and (2.7) are satisfied by P_n with the antipole condition (2.8), then the Nörlund (\mathbf{N}, p_n) sum of Fourier–Jacobi series (2.1) at the point $y = 1$ converges to A . So, $\{\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t)\}$ forms a Cauchy sequence. Thus

$$\lim_{n \rightarrow \infty} \int_{-1}^1 \left| \left(\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) - \mathbf{h}_m^{(\gamma, \delta)}(f, 1, t) \right) \rho^{(\gamma, \delta)}(t) \right| dt = 0.$$

This gives that $\mathbf{H}_n^{(\gamma, \delta)}(f, 1, \omega)$ is a Cauchy sequence and converges in probability to a random variable. With the help of Lemma 4,

$$\begin{aligned} &P\left(\left|\mathbf{H}_n^{(\gamma, \delta)}(f, 1, \omega) - \int_{-1}^1 A \rho^{(\gamma, \delta)}(\psi) dX(\psi, \omega)\right| > \epsilon\right) \\ &\leq \frac{C2^{\alpha+1}}{\epsilon^\alpha(\alpha+1)} \int_{-1}^1 \left| \left(\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) - A \right) \rho^{(\gamma, \delta)}(t) \right| dt. \end{aligned}$$

Hence by the Theorem 1,

$$\lim_{n \rightarrow \infty} \int_{-1}^1 \left| \left(\mathbf{h}_n^{(\gamma, \delta)}(f, 1, t) - A \right) \rho^{(\gamma, \delta)}(t) \right| dt = 0.$$

This completes the proof of theorem. \square

The following two theorems state about the generalized Nörlund summability and lower triangular summability of random Fourier–Jacobi series (4.1) which can be established by following the same steps in Theorem 8.

Theorem 9. *If $\gamma \geq 0$, $\delta - \gamma > 1$, $\delta + \gamma > 0$ and $\{q_n\}$, $\{r_n\}$ are the non-negative real constants sequences satisfy the conditions (2.9), (2.10) and the antipole condition (2.11) stated in the Theorem 2, then the random Fourier–Jacobi series (4.1) is $|\mathbf{N}, q_n, r_n|$ summable in probability to the stochastic integral (4.3) at the point $y = 1$.*

Theorem 10. *If $0 \leq \gamma \leq 1/2$, $\delta \geq 0$ and the infinite triangular regular matrix $M = (\mathbf{b}_{n,k})$ be such that, its entries $(\mathbf{b}_{n,k})$ are positive, monotonic increasing in k , $0 \leq k \leq n$, and satisfy the conditions (2.12), (2.13), (2.14), (2.15) with the antipole condition (2.16) stated in the Theorem 3, then the random Fourier–Jacobi series (4.1) is (\mathbf{T}) summable to the stochastic integral (4.3) at the point $y = 1$.*

ACKNOWLEDGMENTS

This research work was supported by University Grant Commission (National Fellowship with letter no-F./2015-16/NFO-2015-17-OBC-ORI-33062).

REFERENCES

- [1] Borwein, D.: On product of sequences. *J. Lond. Math. Soc.* 33 (1958) 352-357.
- [2] Choudhary, R. S.: On Nörlund summability of Jacobi series. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend.* 52 (5) (1972) 644-652.
- [3] Chripkó, Á.: Weighted approximation via Θ -summations of Fourier–Jacobi series. *Studia Sci. Math. Hungar.* 47 (2) (2010) 139-154.
- [4] Dhakal, B. P.: Summability of a Jacobi series by lower triangular matrix method. *Amer. J. Math. Anal.* 1 (3) (2013) 42-47.
- [5] Liu Z. and Liu, S.: Randomization of the Fourier transform. *Opt. Lett.* 32 (5) (2007) 478-480.
- [6] Liu, Z. and Liu, S.: Random fractional Fourier transform. *Opt. Lett.* 32 (15) (2007) 2088-2090.
- [7] Lubinsky, D. S. and Totik, V.: Best weighted polynomial approximation via Jacobi expansions. *SIAM . Math. Anal.* 25 (1994) 555-570.
- [8] Lukacs, E.: *Stochastic convergence*. Second Ed., Academic Press, London (1975).
- [9] Maharana, P. and Sahoo, S.: On the convergence of random Fourier–Jacobi series of continuous functions. *Comm. Math.* 32 (1) (2024) 49-61.
- [10] Maharana, P. and Sahoo, S.: On the Convergence of Random Fourier–Jacobi Series in $L^p_{[-1,1]}(\eta, \tau)$ space. Preprint, arXiv:2212.08592v1.
- [11] Nayak, C., Pattanayak, S. and Mishra, M. N.: Random Fourier-Stieltjes series associated with stable process. *Tôhoku Math. J.* 39 (1987) 1-15.
- [12] Raghuvanshi, A. K.: On $|N, q_n, r_n|$ -summability of Jacob series. *Int. J. Sci. Inno. Math. Res.* 2 (2) (2014) 139-148.
- [13] Töeplitz, O.: Über allgemeine lineare Mittelbildungen. *Prace matfz.* 22 (1913) 113-119.
- [14] Towghi, N.: Random Fourier series and absolute summability. *Rocky Moutain J. Math.* 28 (1998) 749-765.
- [15] Zygmund, A.: *Trigonometric series*, Third Ed., Cambridge Univ. Press (2002).