

AN ORDER OUT OF NOWHERE: A NEW ALGORITHM FOR INFINITE-DOMAIN CSPs

ANTOINE MOTTET, TOMÁŠ NAGY, AND MICHAEL PINSKER

ABSTRACT. We consider constraint satisfaction problems whose relations are defined in first-order logic over any uniform hypergraph satisfying certain weak abstract structural conditions. Our main result is a P/NP-complete complexity dichotomy for such CSPs. Surprisingly, the large class of structures under consideration falls into a mixed regime where neither the classical complexity reduction to finite-domain CSPs can be used as a black box, nor does the class exhibit order properties, known to prevent the application of this reduction. We introduce an algorithmic technique inspired by classical notions from the theory of finite-domain CSPs, and prove its correctness based on symmetries that depend on a linear order that is external to the structures under consideration.

1. INTRODUCTION

The *Constraint Satisfaction Problem* (CSP) over a relational structure \mathbb{A} in a finite signature, denoted by $\text{CSP}(\mathbb{A})$, is the problem of deciding satisfiability of a given conjunction of atomic formulas in \mathbb{A} . Any satisfying assignment of values to the variables is called a *solution*, and \mathbb{A} is called the *template* of $\text{CSP}(\mathbb{A})$.

Some well-known computational problems like SAT or graph colorability can be formulated as finite-domain CSPs, i.e., as CSPs with finite templates. It was conjectured by Feder and Vardi that a finite-domain CSP is always polynomial-time solvable or NP-complete [30], and this conjecture was confirmed independently by Bulatov and Zhuk recently [29, 45, 46]. Moreover, the border between tractability and NP-completeness for finite-domain CSPs is determined by the identities that are satisfied by the *polymorphisms* of the template. On the other hand, many computational problems, even as simple as the digraph acyclicity problem, can be formulated as CSPs only with an infinite template. Infinite-domain CSPs also play an important role in several branches of computer science, e.g., they are used for the study of finite-domain promise CSPs [2] and they find applications in artificial intelligence [12, 13, 10, 17, 7, 18, 19].

A full complexity classification of infinite-domain CSPs can never be achieved since every computational decision problem is polynomial-time Turing-equivalent to some infinite-domain CSP [11]. Both proofs of the finite-domain CSP dichotomy theorem rely on the fact that the complexity of CSP of a finite template depends only on its polymorphisms. This is known to be true also for infinite templates which are ω -categorical, i.e., every instance of the CSP of such instance has only finitely many solutions up to automorphisms [23]. The sole assumption of ω -categoricity is still insufficient to assess the complexity of the CSP [31, 32] since the set of possible solutions of any instance up to automorphisms does not need to be algorithmically enumerable. A natural way to achieve this is to additionally require every solution to be described by the relations holding on its image (*homogeneity*) and that every solution must only be verified locally on subsets of a fixed size (*finite boundedness*). For templates whose relations are first-order definable in a finitely bounded homogeneous structure (so-called *first-order reducts* of this structure), a generalization of the original dichotomy conjecture for finite-domain CSPs was formulated by Bodirsky and Pinsker

This research was funded in whole or in part by the Austrian Science Fund (FWF) [P 32337, I 5948]. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

in 2011 [26]. The modern formulation of the conjecture based on recent progress [4, 3, 5] is the following:

Conjecture 1. *Let \mathbb{A} be a CSP template which is a first-order reduct of a finitely bounded homogeneous structure. Then one of the following applies.*

- *The clone of polymorphisms of \mathbb{A} has a uniformly continuous minion homomorphism to the clone of projections \mathcal{P} , and $\text{CSP}(\mathbb{A})$ is NP-complete.*
- *The clone of polymorphisms of \mathbb{A} has no uniformly continuous minion homomorphism to the clone of projections \mathcal{P} , and $\text{CSP}(\mathbb{A})$ is in P.*

The first item of [Conjecture 1](#) corresponds to the clone of polymorphisms being essentially structureless, while in the second item, the clone is supposed to have a rich algebraic structure witnessing the tractability of the CSP. Since, within the range of [Conjecture 1](#), a solution to $\text{CSP}(\mathbb{A})$ can be described by specifying coherent local solutions to instances of bounded size, CSP of any template within the range of the conjecture is in NP. Moreover, if the clone of polymorphisms of such a template has a uniformly continuous minion homomorphism to the clone of projections, then the CSP is in NP-hard [5]. It is also known that in order to prove [Conjecture 1](#), one can replace the assumption of the template \mathbb{A} being a first-order reduct of a finitely bounded homogeneous structure by \mathbb{A} being a *model-complete core* of such template.

[Conjecture 1](#) has been confirmed for many subclasses: for example for CSPs of all structures first-order definable in finitely bounded homogeneous graphs [24, 20], in $(\mathbb{Q}, <)$ [16], in any unary structure [22], in the random poset [34], in the random tournament [38], or in the homogeneous branching C-relation [14], as well as for all CSPs in the class MMSNP [19], for CSPs of representations of some relational algebras [17] and for CSPs of ω -categorical monadically stable structures [28].

There are two regimes in which the above-mentioned complexity classifications were proven. In the first regime, the aim is to show that already the clone of *canonical* polymorphisms admits a rich algebraic structure, which yields an efficient many-one reduction to a tractable finite-domain CSP [21, 22].

In order to prove that the structures under consideration have canonical polymorphisms enabling an efficient reduction of their CSPs to tractable finite-domain CSPs, a demanding case distinction had to be done until recently. Today, at least two general approaches that avoid this case distinction are available: the theory of *smooth approximations* of Mottet and Pinsker [38] and the work of Bodirsky and Bodor on the *unique interpolation property* [10].

In the second regime, the standard action of the canonical polymorphisms of the structure on orbits of tuples is trivial, and therefore ad hoc algorithms that do not use the standard reduction to the finite-domain CSP mentioned above need to be given to confirm [Conjecture 1](#). In [38], a non-standard action of the canonical polymorphisms is used to explain the known algorithms in the case of temporal CSPs, but the method does not seem to generalize. It was remarked in [10] that the boundary between the two regimes is roughly drawn by whether or not the template has the *strict order property* (SOP) (for details on the SOP, see [33]). Indeed, in all of the above-mentioned complexity classifications, it was possible to confirm [Conjecture 1](#) for first-order reducts of structures without the SOP within the first regime.

However, in [41, Example 1], a worrying example of a first-order reduct \mathbb{A} of the universal homogeneous 3-uniform hypergraph \mathbb{H} is given. Even though \mathbb{H} does not have the SOP, the structure \mathbb{A} should have a polynomial-time solvable CSP according to [Conjecture 1](#) but it does not possess canonical polymorphisms that would allow us to use the standard reduction to a tractable finite-domain CSP. The polymorphisms of \mathbb{A} depend on a linear order on the domain of \mathbb{H} even though the hypergraph \mathbb{H} is not ordered. This surprising discovery implies that new algorithmic techniques are needed to solve CSPs of first-order reducts of finitely bounded homogeneous hypergraphs.

1.1. Results. In this article, we will use the above-mentioned theory of smooth approximations to confirm [Conjecture 1](#) for first-order reducts of certain finitely bounded homogeneous ℓ -uniform hypergraphs (where $\ell \geq 3$). In particular, we show that the templates from [\[41, Example 1\]](#) have polynomial-time solvable CSPs. For the precise definitions of all the mentioned concepts, see [Section 2](#).

Theorem 2. *Let $\ell \geq 3$, let \mathbb{H} be a finitely bounded homogeneous ℓ -uniform hypergraph whose expansion with a freely added linear order $<$ is a Ramsey structure and whose automorphism group is n -“primitive” for every $n \geq 1$ and let \mathbb{A} be a first-order reduct of \mathbb{H} . Then precisely one of the following applies.*

- (1) *The clone of polymorphisms of \mathbb{A} has a uniformly continuous minion homomorphism to the clone of projections \mathcal{P} , and $\text{CSP}(\mathbb{A})$ is NP-complete.*
- (2) *The clone of polymorphisms of \mathbb{A} has no uniformly continuous minion homomorphism to the clone of projections \mathcal{P} , and $\text{CSP}(\mathbb{A})$ is in P.*

The complexity classification from [Theorem 2](#) is of particular interest for the following reasons:

- New algorithms are needed to prove [Theorem 2](#). Since proving [Conjecture 1](#) would in particular give an algorithm solving all tractable finite-domain CSPs, it seems likely that the methods existing in the finite either have to be used as a black box or have to be adapted in the infinite setting. The black box method from [\[22\]](#) does not work for the class of structures considered in this paper, so we resort to the second option and introduce algorithmic techniques inspired by Zhuk’s algorithm for finite-domain CSPs. These techniques are then coupled with the classical reduction to finite-domain CSPs, resulting in an intriguing interplay between infinitary and finitary methods.
- The result depends only on some general properties of the automorphism group of the base structure (n -“primitivity”) and on the fact that the base structure has a particular Ramsey expansion. Moreover, the base structures, i.e., ℓ -uniform hypergraphs satisfying the properties from [Theorem 2](#), are not classified and a complete classification seems to be demanding if not hopeless (some 3-uniform hypergraphs satisfying our assumptions were classified in [\[1\]](#)). This is the first classification where no structural results about the base structures are known.
- In [\[38\]](#), the scalability of the theory of smooth approximations, i.e., the fact that this theory does not require us to analyze all first-order reducts of the particular structure, was claimed to be one of the main contributions of this theory compared to the archaic case-distinction method from the early literature on the subject. [Theorem 2](#) provides us with the first complexity classification using smooth approximations that truly stands by this promise. By [\[44\]](#), even for the universal homogeneous ℓ -uniform hypergraph, the number of first-order reducts of this hypergraph grows with ℓ , putting an exhaustive case analysis out of reach.

Finally, in [Section 8](#), we obtain as an easy consequence of the proof of [Theorem 2](#) a classification of first-order expansions of the finitely bounded homogeneous hypergraphs under consideration whose CSP is solvable by local consistency methods.

2. PRELIMINARIES

For $k \geq 1$, we write $[k]$ for the set $\{1, \dots, k\}$. A tuple is called *injective* if its entries are pairwise distinct.

2.1. CSPs and Relational Width. A *CSP instance* over a set A is a pair $\mathcal{I} = (\mathcal{V}, \mathcal{C})$, where \mathcal{V} is a finite set of variables, and \mathcal{C} is a set of *constraints*; each constraint $C \in \mathcal{C}$ is a subset of A^U for some non-empty $U \subseteq \mathcal{V}$ (U is called the *scope* of C). For a relational structure \mathbb{A} , we say that \mathcal{I} is an *instance of $\text{CSP}(\mathbb{A})$* if for every $C \in \mathcal{C}$ with scope U , there exists an enumeration u_1, \dots, u_k of the elements of U and a k -ary relation R of \mathbb{A} such that for all $f: U \rightarrow A$ we have

$f \in C \Leftrightarrow (f(u_1), \dots, f(u_k)) \in R$. A mapping $s: \mathcal{V} \rightarrow A$ is a *solution* of the instance \mathcal{I} if we have $s|_U \in C$ for every $C \in \mathcal{C}$ with scope U . Given a constraint $C \subseteq A^U$ and a tuple $\mathbf{v} \in U^k$ for some $k \geq 1$, the *projection of C onto \mathbf{v}* is defined by $\text{proj}_{\mathbf{v}}(C) := \{f(\mathbf{v}) : f \in C\}$. Let $U \subseteq \mathcal{V}$. We define the *restriction of \mathcal{I} to U* to be an instance $\mathcal{I}|_U = (U, \mathcal{C}|_U)$ where the set of constraints $\mathcal{C}|_U$ contains for every $C \in \mathcal{C}$ the constraint $C|_U = \{g|_U \mid g \in C\}$.

We denote by $\text{CSP}_{\text{Inj}}(\mathbb{A})$ the restriction of $\text{CSP}(\mathbb{A})$ to those instances of $\text{CSP}(\mathbb{A})$ where for every constraint C and for every pair of distinct variables u, v in its scope, $\text{proj}_{(u,v)}(C) \subseteq \{(a, b) \in A^2 \mid a \neq b\}$.

Definition 3. Let $1 \leq m \leq n$. We say that an instance $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ is (m, n) -*minimal* if both of the following hold:

- every non-empty subset of at most m variables in \mathcal{V} is contained in the scope of some constraint in \mathcal{I} ;
- for every at most n -element tuple of variables \mathbf{v} and any two constraints $C_1, C_2 \in \mathcal{C}$ whose scopes contain all variables of \mathbf{v} , the projections of C_1 and C_2 onto \mathbf{v} coincide.

For $m \geq 1$, we say that an instance is m -*minimal* if it is (m, m) -minimal. We say that an instance \mathcal{I} of the CSP is *non-trivial* if it does not contain any empty constraint. Otherwise, \mathcal{I} is *trivial*.

For all $1 \leq m \leq n$ and for every instance \mathcal{I} of a $\text{CSP}(\mathbb{A})$ for some finite structure \mathbb{A} , an (m, n) -minimal instance with the same solution set as \mathcal{I} can be computed from \mathcal{I} in polynomial time. The same holds for any ω -categorical structure \mathbb{A} (see the next section for the definition of ω -categoricity, and see e.g., Section 2.3 in [40] for a description of the (m, n) -minimality algorithm in this setting). The resulting instance \mathcal{I}' is called the (m, n) -*minimal instance equivalent to \mathcal{I}* and the algorithm that computes this instance is called the (m, n) -*minimality algorithm*. Note that the instance \mathcal{I}' is not necessarily an instance of $\text{CSP}(\mathbb{A})$. However, \mathcal{I}' is an instance of $\text{CSP}(\mathbb{A}')$ where \mathbb{A}' is the expansion of \mathbb{A} by all at most n -ary relations pp-definable in \mathbb{A} . Moreover, $\text{CSP}(\mathbb{A}')$ has the same complexity as $\text{CSP}(\mathbb{A})$.

If \mathcal{I} is m -minimal and \mathbf{v} is a tuple of variables of length at most m , then by definition there exists a constraint of \mathcal{I} whose scope contains \mathbf{v} , and all the constraints who do have the same projection on \mathbf{v} . We write $\text{proj}_{\mathbf{v}}(\mathcal{I})$ for this projection, and call it the *projection of \mathcal{I} onto \mathbf{v}* .

Definition 4. Let $1 \leq m \leq n$ and let \mathbb{A} be a relational structure. We say that $\text{CSP}(\mathbb{A})$ has relational width (m, n) if every non-trivial (m, n) -minimal instance of $\text{CSP}(\mathbb{A})$ has a solution. $\text{CSP}(\mathbb{A})$ has *bounded width* if it has relational width (m, n) for some natural numbers $m \leq n$. We say that CSP_{Inj} has relational width (m, n) if every non-trivial (m, n) -minimal instance of CSP_{Inj} has a solution.

2.2. Hypergraphs and model-theoretic notions. A relational structure \mathbb{B} is *homogeneous* if every partial isomorphism between finite induced substructures of \mathbb{B} extends to an automorphism of \mathbb{B} . It is *finitely bounded* if there exists a finite set \mathcal{F} of forbidden finite substructures such that the *age* of \mathbb{B} , i.e., the set of its finite substructures up to isomorphism, consists precisely of those structures in the signature of \mathbb{B} which do not embed any member of \mathcal{F} . For a finitely bounded structure \mathbb{B} , we will denote by $b_{\mathbb{B}}$ the cardinality of the biggest forbidden substructure of \mathbb{B} .

Let $\ell \geq 2$. A structure $\mathbb{H} = (H; E)$ is an ℓ -uniform hypergraph if the relation E (called a *hyperedge*) is of arity ℓ , contains only injective tuples and is *fully symmetric*, i.e., for any tuple in E , all tuples obtained by permuting its components are in E as well.

Let $\ell \geq 2$. The *universal homogeneous ℓ -uniform hypergraph* is the up to isomorphism unique countably infinite homogeneous structure that is *universal* for the class of all finite ℓ -uniform hypergraphs, i.e., it contains every such hypergraph as an induced substructure.

A homogeneous structure \mathbb{B} is *Ramsey* if its age satisfies a certain structural analogue of Ramsey's theorem – the concrete definition will not be needed, we only need its consequence from [27] which is stated later in this section. The universal homogeneous ℓ -uniform hypergraph is not Ramsey for

any ℓ but it has a finitely bounded homogeneous *Ramsey expansion* – if we add a linear order *freely* (i.e., so that the new age consists of the structures from the old age ordered in all possible ways and the resulting structure is homogeneous), the resulting structure will be homogeneous, finitely bounded and Ramsey – this follows, e.g., from the Nešetřil–Rödl theorem [43].

Definition 5 (“Primitivity”). Let A be a set and $n \geq 1$. A permutation group \mathcal{G} acting on A is n -“*primitive*” if for every orbit $O \subseteq A^n$ of \mathcal{G} , every \mathcal{G} -invariant equivalence relation on O containing some pair (a, b) with a, b disjoint is full.

Example 6. The automorphism group of the universal homogeneous ℓ -uniform hypergraph \mathbb{H} is n -“*primitive*” for any $n \geq 1$. Indeed, let $n \geq 1$, let O be an orbit of n -tuples under $\text{Aut}(\mathbb{H})$ and let \sim be an equivalence relation on O containing (a, b) such that a, b are disjoint. Let $c, d \in O$ be arbitrary. We define \mathbb{X} to be an ℓ -uniform hypergraph over $3n$ elements $\{x_i^j \mid i \in [n], j \in [3]\}$ such that the following holds. The hypergraph induced by $(x_1^j, \dots, x_n^j, x_1^{j+1}, \dots, x_n^{j+1})$ is isomorphic to the structure induced by (a, b) in \mathbb{H} for every $j \in [2]$ and the hypergraph induced by $(x_1^1, \dots, x_n^1, x_1^3, \dots, x_n^3)$ is isomorphic to the structure induced by (c, d) in \mathbb{H} . By the universality of \mathbb{H} , \mathbb{X} embeds to \mathbb{H} . By the homogeneity of \mathbb{H} , we can assume that the embedding maps $(x_1^1, \dots, x_n^1, x_1^3, \dots, x_n^3)$ to (c, d) . By the transitivity of \sim , $c \sim d$.

In the whole paper, we fix $\ell \geq 3$ and a finitely bounded homogeneous ℓ -uniform hypergraph \mathbb{H} whose expansion with a freely added linear order $<$ is a Ramsey structure and whose automorphism group is n -“*primitive*” for every $n \geq 1$.

Note that not every finitely bounded homogeneous ℓ -uniform hypergraph satisfies our assumptions. However, the universal homogeneous ℓ -uniform hypergraph does satisfy the assumptions for every $\ell \geq 3$. Additionally, for every fixed $\ell \geq 3$ and $n > \ell$, there exists a homogeneous ℓ -uniform hypergraph that is universal for the class of K_n^ℓ -free hypergraphs, where K_n^ℓ is the ℓ -hypergraph on n vertices whose every ℓ -element subset forms a hyperedge; these hypergraphs also satisfy our assumptions.

For $n \geq 1$, we denote by I_n the set of injective n -tuples in \mathbb{H} and we write $I := I_\ell$. We write N for the complement of the hyperedge relation E in I and we call it the *non-hyperedge* relation.

A permutation group is *oligomorphic* if it has only finitely many orbits in its componentwise action on n -tuples of elements for all $n \geq 1$. A countable relational structure is ω -*categorical* if its automorphism group is oligomorphic. A *first-order reduct* of a structure \mathbb{B} is a structure \mathbb{A} on the same domain whose relations are first-order definable without parameters from \mathbb{B} . Every first-order reduct of a finitely bounded homogeneous structure is ω -categorical. A first-order reduct of a relational structure \mathbb{B} is a *first-order expansion* of \mathbb{B} if it has among its relations all relations from the signature of \mathbb{B} .

A *primitive-positive* (pp-)formula is a first-order formula built only from atomic formulae, existential quantifiers, and conjunction. A relation is *pp-definable* in a relational structure \mathbb{A} if it is first-order definable by a pp-formula in \mathbb{A} .

We say that an ω -categorical relational structure has *no algebraicity* if none of its elements is first-order definable using other elements as parameters. Note that \mathbb{H} has no algebraicity. To see this, suppose that there exists $a_0 \in H$ that is first-order definable using elements $a_1, \dots, a_n \in H$ as parameters. Let us define ordered ℓ -uniform hypergraphs $\mathbb{X} = (\{x_0, \dots, x_n\}, E^{\mathbb{X}}, <^{\mathbb{X}})$ and $\mathbb{Y} = (\{y_0, \dots, y_n\}, E^{\mathbb{Y}}, <^{\mathbb{Y}})$ such that both $(x_{i_1}, \dots, x_{i_\ell}) \in E^{\mathbb{X}}$ and $(y_{i_1}, \dots, y_{i_\ell}) \in E^{\mathbb{Y}}$ if, and only if, $(a_{i_1}, \dots, a_{i_\ell}) \in E$ for every $(i_1, \dots, i_\ell) \in \{0, \dots, n\}^\ell$. Let moreover both $x_i <^{\mathbb{X}} x_j$ and $y_i <^{\mathbb{Y}} y_j$ if, and only if $a_i < a_j$ for every $i, j \in [n]$ and let $x_0 < x_i$ and $y_0 > y_i$ for every $i \in [n]$. It follows that \mathbb{X} and \mathbb{Y} embeds to $(\mathbb{H}, <)$ by embeddings e_X and e_Y and by the homogeneity, we may suppose that $e_X(a_i) = e_Y(b_i) = x_i$ for every $i \in [n]$. Hence, $e_X(x_0), e_Y(y_0)$ and a_0 satisfy the same first-order formulas over $(\mathbb{H}, \{\{a_i\} \mid i \in [n]\})$ but $e_X(x_0) \neq e_Y(y_0)$, a contradiction.

2.3. Universal algebra. A *polymorphism* of a relational structure \mathbb{A} is a homomorphism from some finite power of \mathbb{A} to \mathbb{A} . The set of all polymorphisms of a structure \mathbb{A} , denoted by $\text{Pol}(\mathbb{A})$, is a *function clone*, i.e., a set of finitary operations on a fixed set which contains all projections and which is closed under arbitrary compositions. A relation is pp-definable in an ω -categorical relational structure \mathbb{A} if, and only if, it is invariant under $\text{Pol}(\mathbb{A})$.

For a function clone \mathcal{C} , we denote the domain of its functions by C ; we say that \mathcal{C} *acts on* C . The clone \mathcal{C} also naturally acts (componentwise) on C^l for any $l \geq 1$, on any invariant subset S of C (by restriction), and on the classes of any invariant equivalence relation \sim on an invariant subset S of C (by its action on representatives of the classes). We write $\mathcal{C} \curvearrowright C^l$, $\mathcal{C} \curvearrowright S$ and $\mathcal{C} \curvearrowright S/\sim$ for these actions. Any action $\mathcal{C} \curvearrowright S/\sim$ is called a *subfactor* of \mathcal{C} , and we also call the pair (S, \sim) a subfactor. A subfactor (S, \sim) is *minimal* if the equivalence relation \sim has at least two classes and no proper subset of S that intersects at least two \sim -classes is invariant under \mathcal{C} .

For $n \geq 1$, a k -ary operation f defined on the domain C of a permutation group \mathcal{G} is *n-canonical* with respect to \mathcal{G} if for all $a_1, \dots, a_k \in C^n$ and all $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ there exists $\beta \in \mathcal{G}$ such that $f(a_1, \dots, a_k) = \beta \circ f(\alpha_1(a_1), \dots, \alpha_k(a_k))$. In particular, f induces an operation on the set C^n/\mathcal{G} of orbits of n -tuples under \mathcal{G} . If all functions of a function clone \mathcal{C} are n -canonical with respect to \mathcal{G} , then \mathcal{C} acts on C^n/\mathcal{G} and we write $\mathcal{C}^n/\mathcal{G}$ for this action; if \mathcal{G} is oligomorphic then $\mathcal{C}^n/\mathcal{G}$ is a function clone on a finite set. A function that is n -canonical with respect to \mathcal{G} for all $n \geq 1$ is called *canonical* with respect to \mathcal{G} . We say that a function is *diagonally canonical* if it satisfies the definition of n -canonicity in the case $\alpha_1 = \dots = \alpha_k$ for every $n \geq 1$.

We write $\mathcal{G}_{\mathcal{C}}$ to denote the largest permutation group contained in a function clone \mathcal{C} , and say that \mathcal{C} is oligomorphic if $\mathcal{G}_{\mathcal{C}}$ is oligomorphic.

For a set of functions \mathcal{F} over the same fixed set C we write $\overline{\mathcal{F}}$ for the set of those functions g such that for all finite subsets $F \subseteq C$, there exists a function in \mathcal{F} which agrees with g on F . For $k \geq 1$, for k -ary functions f, g and for a permutation group \mathcal{G} such that f, g and \mathcal{G} act on the same domain, we say that f *locally interpolates* g modulo \mathcal{G} if $g \in \overline{\{\beta \circ f(\alpha_1, \dots, \alpha_k) \mid \beta, \alpha_1, \dots, \alpha_k \in \mathcal{G}\}}$. Similarly, we say that f *diagonally interpolates* g modulo \mathcal{G} if f locally interpolates g with $\alpha_1 = \dots = \alpha_k$. If \mathcal{G} is the automorphism group of a *Ramsey structure* in the sense of [9], then every function on its domain locally (diagonally) interpolates a canonical (diagonally canonical) function modulo \mathcal{G} [27, 25]. We say that a clone \mathcal{D} locally interpolates a clone \mathcal{C} modulo a permutation group \mathcal{G} if for every $g \in \mathcal{D}$ there exists $f \in \mathcal{C}$ such that g locally interpolates f modulo \mathcal{G} . A clone \mathcal{C} is a *model-complete core* if its unary functions are equal to $\overline{\mathcal{G}_{\mathcal{C}}}$. Hence, for every $n \geq 1$, every orbit of n -tuples under $\mathcal{G}_{\mathcal{C}}$ is preserved by \mathcal{C} .

A structure \mathbb{A} is called a *model-complete core* if its polymorphism clone is a model-complete core. By [8], for any ω -categorical structure \mathbb{A} , there exists an up to isomorphism unique ω -categorical model-complete core \mathbb{A}' and the CSPs of \mathbb{A} and \mathbb{A}' are the same computational problem. In any ω -categorical model-complete core \mathbb{A} , all orbits of n -tuples with respect to the automorphism group $\text{Aut}(\mathbb{A})$ are pp-definable, for all $n \geq 1$.

A *weak near-unanimity (WNU) operation of arity $n \geq 2$* is an operation w satisfying the equation $w(x, \dots, x, y) = \dots = w(y, x, \dots, x)$ for all x, y from its domain. A ternary operation m is a *minority operation* if it satisfies $m(x, x, y) = m(x, y, x) = m(y, x, x) = y$ for all x, y from its domain. A binary operation f on a two element domain is a *semilattice operation* if $f(x, y) = f(y, x) = f(x, x) = x$ and $f(y, y) = y$ for some enumeration $\{x, y\}$ of its domain.

A function is *idempotent* if it satisfies $f(x, \dots, x) = x$ for every x from its domain. A function clone is idempotent if all of its functions are. For a function f of arity $n \geq 1$ and for $i \in [n]$, we say that the i -th variable of f is *essential* if there exist a_1, \dots, a_n, a'_i from the domain of f such that $f(a_1, \dots, a_n) \neq f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_n)$. A function is called *essentially unary* if at most one of its variables is essential, otherwise is the function *essential*.

An arity-preserving map $\xi: \mathcal{C} \rightarrow \mathcal{D}$ between function clones is called a *minion homomorphism* if it preserves compositions with projections, i.e., it satisfies $\xi(f \circ (\pi_1, \dots, \pi_n)) = \xi(f) \circ (\pi_1, \dots, \pi_n)$

for all $n, m \geq 1$ and all n -ary $f \in \mathcal{C}$ and m -ary projections π_1, \dots, π_n . An arity-preserving map ξ is called a *clone homomorphism* if it preserves projections, i.e., maps every projection in \mathcal{C} to the corresponding projection in \mathcal{D} , and compositions, i.e., it satisfies $\xi(f \circ (g_1, \dots, g_n)) = \xi(f) \circ (\xi(g_1), \dots, \xi(g_n))$ for all $n, m \geq 1$ and all n -ary $f \in \mathcal{C}$ and m -ary $g_1, \dots, g_n \in \mathcal{C}$. We say that a function clone \mathcal{C} is *equationally trivial* if it has a clone homomorphism to the clone \mathcal{P} , and *equationally non-trivial* otherwise. We say that \mathcal{C} is *equationally affine* if it has a clone homomorphism to an *affine clone*, i.e., a clone of affine maps over a finite module. It is known that a finite idempotent clone is either equationally affine or it contains WNU operations of all arities $n \geq 3$ ([36], this stronger version is attributed to E. Kiss in [35, Theorem 2.8], a different proof can be found in [47]).

If \mathcal{C}, \mathcal{D} are function clones and \mathcal{D} has a finite domain, then a clone (or minion) homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{D}$ is *uniformly continuous* if for all $n \geq 1$ there exists a finite subset F of C^n such that $\xi(f) = \xi(g)$ for all n -ary $f, g \in \mathcal{C}$ which agree on F .

For a first-order reduct \mathbb{A} of \mathbb{H} , we will consider the following two subclones of $\text{Pol}(\mathbb{A})$:

- $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ is the clone of those polymorphisms of \mathbb{A} which preserve the equivalence of orbits of injective tuples under $\text{Aut}(\mathbb{H})$.
- $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ is the clone of those polymorphisms of \mathbb{A} which preserve the equivalence of orbits of injective tuples under $\text{Aut}(\mathbb{A})$.

3. OVERVIEW OF THE PROOF OF THEOREM 2

Let \mathbb{A} be a model-complete core of a first-order reduct \mathbb{A}' of \mathbb{H} . By [5], it is enough to prove that Theorem 2 holds for \mathbb{A} since there exists a uniformly continuous minion homomorphism from $\text{Pol}(\mathbb{A}')$ to $\text{Pol}(\mathbb{A})$ and from $\text{Pol}(\mathbb{A})$ to $\text{Pol}(\mathbb{A}')$. By Proposition 10, we can assume that \mathbb{A} is itself a first-order reduct of \mathbb{H} .

By applying some folklore results, as well as a compactness argument, we finally obtain in Proposition 13 that $\text{Pol}(\mathbb{A})$ contains an injective operation with certain properties unless $\text{Pol}(\mathbb{A})$ admits a uniformly continuous clone homomorphism to \mathcal{P} . This polymorphism witnesses that I is a *binary absorbing subuniverse* of H^ℓ . In the rest of the paper, we will prove that assuming that I is absorbing, then $\text{CSP}(\mathbb{A})$ is tractable if, and only if, $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial.

Let us therefore first suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial. In this case, $\text{CSP}_{\text{Inj}}(\mathbb{A})$ can be solved by the reduction to a tractable finite domain CSP from [21, 22]. We will show that $\text{CSP}(\mathbb{A})$ can be reduced to $\text{CSP}_{\text{Inj}}(\mathbb{A})$.

By the classification of clones on a two-element domain from [42], $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is either equationally non-affine, or it consists of affine maps over \mathbb{Z}_2 . In the first case, $\text{CSP}_{\text{Inj}}(\mathbb{A})$ has relational width $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ by an easy modification of the proof of Theorem 2 in [40]. Now, Theorem 2 follows from Corollary 15 (details in Section 5.1). In the second case, $\text{CSP}_{\text{Inj}}(\mathbb{A})$ amounts to solving linear equations over \mathbb{Z}_2 . In this situation, we can apply the following algorithm – for more details, see Section 5.2.

Let \mathcal{I} be an instance of $\text{CSP}(\mathbb{A})$. Our algorithm transforms \mathcal{I} into an equi-satisfiable instance \mathcal{I}' that is sufficiently minimal, such that the solution set of a certain relaxation of \mathcal{I}' is subdirect on all projections to an ℓ -tuple of pairwise distinct variables, and that additionally satisfies a condition which we call *inj-irreducibility*, inspired by Zhuk's notion of irreducibility [45, 46]. We then prove that any satisfiable instance satisfying those properties has an injective solution. This step is to be compared with the case of *absorbing reductions* in Zhuk's algorithm, and in particular with Theorem 5.5 in [46], in which it is proved that any sufficiently minimal and irreducible instance that has a solution also has a solution where an arbitrary variable is constrained to belong an absorbing subuniverse. Since in our setting I is an absorbing subuniverse of H^ℓ , this fully establishes a parallel between the present work and [46].

If $\mathcal{C}_{\mathbb{A}}^{\mathbb{H},\text{inj}} \curvearrowright \{E, N\}$ is equationally trivial, our goal is to prove that $\text{CSP}(\mathbb{A})$ is NP-hard. The first step is to establish that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}}$ is equationally trivial as well and that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \subseteq \mathcal{C}_{\mathbb{A}}^{\mathbb{H},\text{inj}}$ (Lemma 24). Moreover, Proposition 36 in [38] implies that there exists $k \geq \ell$ such that the action of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}}$ on orbits of injective k -tuples under $\text{Aut}(\mathbb{A})$ is equationally trivial. Therefore, there exists a *naked set* (S, \sim) for this action. A naked set of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ consists of an invariant subset $S \subseteq I_k/\text{Aut}(\mathbb{A})$ and an invariant equivalence relation \sim on S such that \sim has at least two equivalence classes and such that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ acts on S/\sim by projections. By classical results in finite clone theory, the existence of such a naked set is equivalent to the existence of a clone homomorphism from $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ to \mathcal{P} , which extends to a uniformly continuous clone homomorphism $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \rightarrow \mathcal{P}$.

Now, we can employ the theory of smooth approximations to extend this homomorphism further and obtain a uniformly continuous clone homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \mathcal{P}$. We recall below the relevant definitions from the theory of smooth approximations.

Definition 7 (Smooth approximations). Let A be a set and let \sim be an equivalence relation on $S \subseteq A$. We say that an equivalence relation η on a set S' with $S \subseteq S' \subseteq A$ *approximates* \sim if the restriction of η to S is a refinement of \sim . η is called an *approximation* of \sim .

For a permutation group \mathcal{G} acting on A and leaving the \sim -classes invariant as well as η , we say that the approximation η is *smooth* if every equivalence class C of \sim intersects some equivalence class C' of η such that $C \cap C'$ contains a \mathcal{G} -orbit. η is *very smooth* if orbit-equivalence with respect to \mathcal{G} is a refinement of η on S .

Recall that \mathbb{H} has no algebraicity and hence, the hypotheses of the loop lemma of smooth approximations [38, Theorem 10] are met and we may use the following reformulation of the lemma to our situation.

Theorem 8. *Let $k \geq 1$ and suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ is equationally trivial. Then there exists a naked set (S, \sim) of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ with $\text{Aut}(\mathbb{A})$ -invariant \sim -classes such that one of the following holds:*

- \sim is approximated by a $\text{Pol}(\mathbb{A})$ -invariant equivalence relation that is very smooth with respect to $\text{Aut}(\mathbb{A})$;
- every $\text{Pol}(\mathbb{A})$ -invariant binary symmetric relation $R \subseteq (I_k)^2$ that contains a pair $(a, b) \in S^2$ such that $a \neq b$ and such that $a \not\sim b$ contains a pseudo-loop modulo $\text{Aut}(\mathbb{A})$, i.e., a pair (c, c') where c, c' belong to the same orbit under $\text{Aut}(\mathbb{A})$.

In [38, Theorem 10], the first item of the statement gives an approximation \sim that is not very smooth, but *presmooth*. By [38, Lemma 8], under the assumption that $\text{Pol}(\mathbb{A})$ preserves I_2 and that $\text{Aut}(\mathbb{A})$ is n -“primitive”, every presmooth approximation is very smooth; this justifies our reformulation above.

Suppose that the first case of Theorem 8 applies. By a minor modification of the smooth approximation toolbox (Lemma 25), this implies that $\text{Pol}(\mathbb{A})$ has a uniformly continuous clone homomorphism to $\mathcal{C}_{\mathbb{A}}^{\mathbb{A},\text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ and hence to the clone of projection.

Suppose now that the second case of Theorem 8 applies. By [38, Lemma 13], $\text{Pol}(\mathbb{A})$ contains a *weakly commutative function*, i.e., a binary operation f with the property that $f(a, b) \sim f(b, a)$ holds for all $a, b \in I_k$ such that $f(a, b)$ and $f(b, a)$ are in S and disjoint. It follows from a fairly involved compactness argument in Lemma 27 that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H},\text{inj}} \curvearrowright \{E, N\}$ contains a semilattice operation and in particular, is equationally non-trivial, which is a contradiction.

4. MODEL-COMPLETE CORES AND INJECTIVE POLYMORPHISMS

In this section, we first prove some basic facts about model-complete cores of first-order reducts of \mathbb{H} . In the second part, we prove that such first-order reducts that are model-complete cores and

that are equationally non-trivial have binary injective polymorphisms acting as a projection or as a semilattice operation on $\{E, N\}$. These binary injections will play an important role in the algorithm for tractable CSPs in [Section 5](#).

4.1. Model-complete cores. Let \mathcal{G} be a permutation group and let $g: G \rightarrow G$ be a function. We say that g is *range-rigid* with respect to \mathcal{G} if all orbits of tuples under \mathcal{G} that intersect the range of g are invariant under g . We will use the following theorem to understand the model-complete cores of first-order reducts of \mathbb{A} .

Theorem 9 ([\[37\]](#)). *Let \mathbb{A} be a first-order reduct of a homogeneous Ramsey structure \mathbb{B} and let \mathbb{A}' be its model-complete core. Then \mathbb{A}' is a first-order reduct of a homogeneous Ramsey substructure \mathbb{B}' of \mathbb{B} .*

Moreover, there exists $g \in \text{End}(\mathbb{A})$ that is range-rigid with respect to $\text{Aut}(\mathbb{B})$ and such that the age of \mathbb{B}' is equal to the age of the structure induced by the range of g in \mathbb{B} .

Proposition 10. *Let \mathbb{A} be a first-order reduct of \mathbb{H} . Then the model-complete core of \mathbb{A} is a one-element structure or a first-order reduct of \mathbb{H} . Moreover, if \mathbb{A} is a model-complete core that is a first-order reduct of \mathbb{H} and not of $(H, =)$, then the range of every $f \in \text{End}(\mathbb{A})$ intersects every orbit under $\text{Aut}(\mathbb{H}, <)$.*

Proof. Using [Theorem 9](#), we obtain that the model-complete core \mathbb{A}' of \mathbb{A} is a first-order reduct of a homogeneous Ramsey substructure \mathbb{B}' of $(\mathbb{H}, <)$. Moreover, there exists $g \in \text{End}(\mathbb{A})$ which is range-rigid with respect to $\text{Aut}(\mathbb{H}, <)$ and such that the age of the structure induced by the range of g in $(\mathbb{H}, <)$ is equal to the age of \mathbb{B}' .

If the range of g contains just one element, \mathbb{B}' is a one-element structure. Otherwise, the range contains at least two elements $a < b$ and by the range-rigidity, it then contains infinitely many elements. In particular, it contains a hyperedge or a non-hyperedge.

If the range contains only hyperedges, then g sends every ℓ -tuple in N to an ℓ -tuple ordered in the same way as the original tuple that lies in the hyperedge relation E . It follows that for all injective tuples a, b of the same length, there exists an automorphism α of \mathbb{H} and an embedding from the range of g into \mathbb{B}' such that $e \circ g \circ \alpha(a) = b$. It follows that $\text{Aut}(\mathbb{A}')$ is the full symmetric group on the domain of \mathbb{A}' and hence, \mathbb{A}' is a first-order reduct of $(A', =)$ which is isomorphic to $(H, =)$. If the range of g contains only ℓ -tuples in N , \mathbb{A}' is a first-order reduct of $(A', =)$ by the same argument where the roles of E and N are switched. Finally, if the range of g contains both a hyperedge as well as a non-hyperedge, it follows from the range-rigidity of g that \mathbb{B}' is isomorphic to $(\mathbb{H}, <)$ and \mathbb{A}' is isomorphic to \mathbb{A} . In particular, \mathbb{A}' is a first-order reduct of \mathbb{H} .

Suppose now that \mathbb{A} is a model-complete core that is a first-order reduct of \mathbb{H} but not of $(H, =)$ and let $f \in \text{End}(\mathbb{A})$. Suppose that the range of f does not intersect every orbit under $\text{Aut}(\mathbb{H}, <)$. By [Lemma 15](#) and [Lemma 11](#) in [\[37\]](#) applied to $\text{End}(\mathbb{A})$, the range-rigid function g does not intersect every orbit under $\text{Aut}(\mathbb{H}, <)$ either and we obtain a contradiction with the previous paragraph. \square

4.2. Injective binary polymorphisms.

Lemma 11. *Let \mathbb{A} be a first-order reduct of a finitely bounded homogeneous hypergraph \mathbb{H} that is a model-complete core. If $\text{Pol}(\mathbb{A})$ does not have a uniformly continuous clone homomorphism to \mathcal{P} , then it contains a binary essential operation.*

Proof. It follows from [Corollary 6.9](#) in [\[6\]](#) that $\text{Pol}(\mathbb{A})$ contains a ternary essential operation. Moreover, the binary relation $O := \{(a, b) \mid a \neq b \in H\}$ is an orbit under $\text{Aut}(\mathbb{H})$ that is *free*, i.e., for every $(c, d) \in H^2$ there exists $a \in H$ such that $(a, c), (a, d) \in O$. Now, the lemma follows directly from [Proposition 22](#) in [\[38\]](#). \square

Note that for every binary injective operation f on H , there exists an embedding e of the substructure induced by the range of f into \mathbb{H} such that $f' := e \circ f$ acts lexicographically on the

order, i.e., $f'(x, y) < f'(x', y')$ if $x < x'$ or $x = x', y < y'$. To see this, let us define an ordered hypergraph $(\mathbb{Y}, E^{\mathbb{Y}}, <^{\mathbb{Y}})$ on H^2 such that $((x_1, y_1), \dots, (x_\ell, y_\ell)) \in E^{\mathbb{Y}}$ if $(f(x_1, y_1), \dots, f(x_n, y_n)) \in E$ and such that $(x, y) <^{\mathbb{Y}} (x', y')$ if $x < x'$ or $x = x', y < y'$. It is easy to see that there exists an isomorphism i from the substructure of \mathbb{H} induced by the range of f in \mathbb{H} to \mathbb{Y} . Since $(\mathbb{H}, <)$ is an expansion of \mathbb{H} by a linear order that is added freely, it is universal for the class of all l -uniform linearly ordered hypergraphs $(\mathbb{X}, <^{\mathbb{X}})$ such that \mathbb{X} embeds to \mathbb{H} . It follows that \mathbb{Y} embeds to $(\mathbb{H}, <)$ by an embedding e' . Finally, by setting $e := e' \circ i$, we get the desired embedding e .

We say that a relational structure \mathbb{B} has *finite duality* if it is finitely bounded and the set \mathcal{F} from the definition of finite boundedness is closed under homomorphic images.

Lemma 12. *Let \mathbb{A} be a first-order reduct of \mathbb{H} that is a model-complete core. Suppose that $\text{Pol}(\mathbb{A})$ contains a binary essential operation. Then $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a binary injection f such that one of the following holds:*

- f acts like a semilattice operation on $\{E, N\}$, or
- f acts like a projection on $\{E, N\}$ and is canonical with respect to $(\mathbb{H}, <)$.

Proof. The binary disequality relation \neq is clearly an orbit under $\text{Aut}(\mathbb{A})$ and since \mathbb{A} is a model-complete core, \mathbb{A} pp-defines \neq . \mathbb{H} is clearly transitive, i.e., $\text{Aut}(\mathbb{H})$ has only one orbit in its action on H . Moreover, $\text{Aut}(\mathbb{H})$ has only one orbit of injective pairs in its action on H^2 , namely the orbit of $\{(a, b) \mid a \neq b \in H\}$ and the structure (H, \neq) clearly has finite duality. Proposition 25 in [38] then implies that $\text{Pol}(\mathbb{A})$ contains a binary injection.

Note that if \mathbb{A} is a reduct of $(H, =)$, then this binary injection can be composed with a unary injection so that the result acts like a semilattice operation on $\{E, N\}$, so that we can assume below that \mathbb{A} is not a first-order reduct of $(H, =)$.

Since $(\mathbb{H}, <)$ is a homogeneous Ramsey structure and since any function interpolated by an injective function is injective, $\text{Pol}(\mathbb{A})$ contains a binary injection f which is canonical with respect to $(\mathbb{H}, <)$. Since \mathbb{A} is a model-complete core and not a first-order reduct of $(H, =)$, we can assume that the action of f on orbits of ℓ -tuples under $\text{Aut}(\mathbb{H}, <)$ is idempotent by Proposition 10 – the range of the endomorphism e defined by $e(x) := f(x, x)$ intersects every orbit under $\text{Aut}(\mathbb{H}, <)$ and hence, if f does not act idempotently on the orbits of ℓ -tuples under $\text{Aut}(\mathbb{H}, <)$, we may compose f finitely many times with itself until it does. Here, by composition of a binary function g with f we mean the function $g(f(x, y), f(x, y))$. Moreover, by the remark above, we may assume that f acts lexicographically on the order $<$.

Let α be a permutation of $\{1, \dots, \ell\}$. If O is an orbit of injective ℓ -tuples under $\text{Aut}(\mathbb{H}, <)$, then we write $\alpha(O)$ for the orbit obtained by changing the order of the tuples in O according to α (not by permuting the tuples in O). That way, α acts naturally on orbits of injective ℓ -tuples. We also apply this notation to unions of orbits of injective tuples under $\text{Aut}(\mathbb{H}, <)$. Let J be the set of strictly increasing ℓ -tuples with respect to $<$. For a fixed permutation α , the restriction of the natural action of f on ℓ -tuples to $J \times \alpha(J)$ acts idempotently on $\{E, N\}$; depending on what this action is, we say that f behaves like p_1 on input α if it behaves like the first projection on $\{E, N\}$ and we say that it behaves like p_2 if it behaves like the second projection in this action on input α ; in both cases, we say that f behaves “like a projection” on input α . Finally, we say that f behaves like \vee on input α if $f(E, N) = f(N, E) = f(E, E) = E, f(N, N) = N$ and that f behaves like \wedge on input α if $f(E, N) = f(N, E) = f(N, N) = N, f(E, E) = E$; in both cases, we say that f behaves “like a semilattice” on input α . Since the action of f on orbits is idempotent, it cannot behave like a constant in the above situation.

We write id for the identity on $\{1, \dots, \ell\}$. If f behaves like p_2 on input id , then $f(f(x, y), f(y, x))$ behaves like p_1 on id while still satisfying the behaviour with respect to the order we assumed in the beginning, so we assume that this is not the case.

If f behaves like p_1 on id , then $f(x, f(x, y))$ behaves like p_1 on $\{E, N\}$ on all inputs, and hence it acts on $\{E, N\}$ and is canonical on injective tuples with respect to \mathbb{H} .

Suppose that f behaves like a semilattice on $\{E, N\}$ on input id ; without loss of generality it behaves like \vee . If f behaves differently on some other input α , then $f(f(x, y), f(x, \alpha^{-1}(y)))$ behaves like \vee on input α as well as on all other inputs where f behaves like \vee . Hence, repeating this argument for all α , we can assume that f behaves like \vee on $\{E, N\}$ on any input. Note, however, that the expression $f(f(x, y), f(x, \alpha^{-1}(y)))$ only makes sense for the action on orbits unless α is the identity or reverses the order; that is, only in those cases there exists a canonical function on $(\mathbb{H}, <)$ whose action on orbits is that of α . The global argument has to be made local in the other cases, as follows. Let $(s, t) \in J \times \alpha(J)$. Let $\overline{\alpha^{-1}}$ be an automorphism of \mathbb{H} which changes the order on t as would α^{-1} . Then $f(f(x, y), f(x, \overline{\alpha^{-1}}(y)))$ acts like \vee on (s, t) , as well on all pairs of injective ℓ -tuples on which it previously acted like \vee . Repeating this procedure for all pairs of ℓ -tuples and applying a standard compactness argument, we obtain a function which acts like \vee on all pairs of injective tuples, and hence on all inputs. \square

A combination of [Lemma 11](#) and [Lemma 12](#) immediately yields the following proposition.

Proposition 13. *Let \mathbb{A} be a first-order reduct of \mathbb{H} that is a model-complete core. If $\text{Pol}(\mathbb{A})$ does not have a uniformly continuous clone homomorphism to \mathcal{P} , then $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a binary injection f such that one of the following holds:*

- f acts like a semilattice operation on $\{E, N\}$, or
- f acts like a projection on $\{E, N\}$ and is canonical with respect to $(\mathbb{H}, <)$.

5. THE TRACTABLE CASE

In this section, we will prove that if $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial, then $\text{CSP}(\mathbb{A})$ is in P. We distinguish two cases based on whether $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally affine.

5.1. $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine. Let \mathbb{A} be a relational structure, let $k \geq 1$, let S, R with $S \subseteq R$ be n -ary relations pp-definable in \mathbb{A} and let $f \in \text{Pol}(\mathbb{A})$ be binary. We say that S is a *binary absorbing subuniverse of R* if for every $s \in S, r \in R$, we have that $f(s, r), f(r, s) \in S$. In this case, we write $S \trianglelefteq_2 R$ and we say that f *witnesses* the binary absorption.

Lemma 14. *Let \mathbb{A} be an ω -categorical structure, let $1 \leq k \leq m \leq n$, let S, R with $S \subseteq R \subseteq A^k$ be relations pp-definable in \mathbb{A} and suppose that $S \trianglelefteq_2 R$. Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be a non-trivial, (m, n) -minimal instance of $\text{CSP}(\mathbb{A})$. Let \mathcal{I}' be an instance obtained from \mathcal{I} by adding for every $\mathbf{v} = (v_1, \dots, v_k) \in \mathcal{V}^k$ with $\text{proj}_{\mathbf{v}}(\mathcal{I}) \subseteq R$, $S \cap \text{proj}_{\mathbf{v}}(\mathcal{I}) \neq \emptyset$ and such that $S \cap \text{proj}_{\mathbf{v}}(\mathcal{I}) \trianglelefteq_2 R$ a constraint $\{c: \{v_1, \dots, v_k\} \rightarrow A \mid c(\mathbf{v}) \in S\}$. Then the (m, n) -minimal instance equivalent to \mathcal{I}' is non-trivial.*

Proof. Let \mathcal{I}'' be the (m, n) -minimal instance equivalent to \mathcal{I}' . We will find a non-trivial, (m, n) -minimal instance \mathcal{J} of CSP over the set A with the same set of variables as \mathcal{I}'' and such that every constraint of \mathcal{J} is a subset of a constraint of \mathcal{I}'' . Then it follows that \mathcal{I}'' is non-trivial.

Let f be a binary polymorphism of \mathbb{A} witnessing $S \trianglelefteq_2 R$, let $F := \{\alpha f(\beta, \gamma) \mid \alpha, \beta, \gamma \in \text{Aut}(\mathbb{A})\}$ and let $T := \{t \in \text{Pol}(\mathbb{A}) \mid t \text{ is a term in } F\}$. Let C be constraint of \mathcal{I} with scope $\{u_1, \dots, u_q\}$ and let $c_1, \dots, c_p \in C$ be such that $(c_1(u_1), \dots, c_1(u_q)), \dots, (c_p(u_1), \dots, c_p(u_q))$ are in pairwise different orbits of q -tuples under $\text{Aut}(\mathbb{A})$ and such that for every $d \in C$ there exists $i \in [p]$ such that $(d(u_1), \dots, d(u_q))$ is in the same orbit under $\text{Aut}(\mathbb{A})$ as $(c_i(u_1), \dots, c_i(u_q))$.

Set for every $C \in \mathcal{C}$, $C' := \{t(c_1, \dots, c_p, a_1, \dots, a_r) \mid t \in T, a_i \in C, \text{ all variables of } t \text{ essential}\}$. Set now \mathcal{J} to be the instance obtained from \mathcal{I} by replacing every constraint C by C' . Clearly, \mathcal{J} is non-trivial, for every k -tuple of variables \mathbf{v} it holds that if $\text{proj}_{\mathbf{v}}(\mathcal{I}) \subseteq R$ and $S \cap \text{proj}_{\mathbf{v}}(\mathcal{I}) \neq \emptyset$ then $S \cap \text{proj}_{\mathbf{v}}(\mathcal{J}) \trianglelefteq_2 R$ and $C' \subseteq C$ for every $C \in \mathcal{C}$. It remains to show that \mathcal{J} is (m, n) -minimal.

To this end, let $1 \leq m' \leq m$ and let $\mathbf{v} \in \mathcal{V}^{m'}$, let $C, D \in \mathcal{C}$ be such that $\text{proj}_{\mathbf{v}}(C) = \text{proj}_{\mathbf{v}}(D)$ and let $c \in C'$. We need to find $d \in D'$ such that $c(\mathbf{v}) = d(\mathbf{v})$. We may suppose that there exist

$d_1, d_2 \in D$ such that $d_1(\mathbf{v}), d_2(\mathbf{v})$ lie in different orbits under $\text{Aut}(\mathbb{A})$ since otherwise, $d(\mathbf{v})$ is in the same orbit of m' -tuples under $\text{Aut}(\mathbb{A})$ as $c(\mathbf{v})$ for any $d \in D$ by the m' -minimality of \mathcal{I} , and hence, for any $d \in D'$ we may find $\alpha \in \text{Aut}(\mathbb{A})$ such that $\alpha(d(\mathbf{v})) = c(\mathbf{v})$. By the definition of D' , it follows that $\alpha d \in D'$.

We have $c = t(c_1, \dots, c_p, a_1, \dots, a_r)$ for some $a_1, \dots, a_r \in C, t \in T$. Suppose that D' is defined using $d_1, \dots, d_{p'} \in D$ and let $s \in T$ be of arity $p' \geq 2$ (by our assumption from the previous paragraph). Then $\alpha_j \circ s(d_1, \dots, d_{p'}) (\mathbf{v}) = c_j(\mathbf{v})$ for some $j \in [p]$ and some $\alpha_j \in \text{Aut}(\mathbb{A})$. Find $x_i \in D$ with $c_i(\mathbf{v}) = x_i(\mathbf{v})$ for every $i \in [p]$ and $y_i \in D$ with $a_i(\mathbf{v}) = y_i(\mathbf{v})$ for every $i \in [r]$. Set now $d := t(x_1, \dots, x_{j-1}, \alpha_j(s(d_1, \dots, d_{p'})), x_{j+1}, \dots, x_p, y_1, \dots, y_r)$. It is easy to see that $d \in D'$.

It follows that $c(\mathbf{v}) = t(c_1, \dots, c_p, a_1, \dots, a_r)(\mathbf{v}) = d(\mathbf{v})$ as desired. \square

[Lemma 14](#) together with the results from [\[39, 40\]](#) give the following corollary.

Corollary 15. *Let \mathbb{A} be a first-order reduct of \mathbb{H} which is a model-complete core. Suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine. Then every non-trivial $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal instance of $\text{CSP}(\mathbb{A})$ has a solution. Moreover, such a solution s exists where $s(x) \neq s(y)$ for all variables x, y such that $\text{proj}_{(x,y)}(\mathcal{I}) \cap I_2 \neq \emptyset$.*

Proof. It follows from [\[36, 35\]](#) that for every $k \geq 3$, $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains an operation of arity k that acts as a WNU operation on $\{E, N\}$. An easy modification of the proof of Theorem 2 in [\[40\]](#) yields that $\text{CSP}_{\text{Inj}}(\mathbb{A})$ has relational width $(2\ell, \max(3\ell, b_{\mathbb{H}}))$. Moreover, by [Proposition 13](#), there exists a binary injection $f \in \text{Pol}(\mathbb{A})$. Therefore, for all $a, b \in A^2$ such that b is injective, $f(a, b), f(b, a) \in I_2$ and hence, $I_2 \triangleleft_2 A^2$. Finally, [Lemma 14](#) yields the result. \square

5.2. $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally affine. Let \mathbb{A} be a first-order reduct of \mathbb{H} which is a model-complete core. If $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally affine but equationally non-trivial, then \mathbb{A} is not a first-order reduct of $(\mathbb{H}, =)$ and it follows from [Proposition 13](#) that $\text{Pol}(\mathbb{A})$ contains a binary injection p_1 canonical with respect to $(\mathbb{H}, <)$ that acts as the first projection on $\{E, N\}$.

Moreover, we can suppose that the function $f(x, y) := p_1(y, x)$ acts lexicographically on the order by the remark above [Lemma 12](#). It follows from [\[42\]](#) and from [\[21\]](#) that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a ternary function m' that acts idempotently and as a minority on $\{E, N\}$. We may moreover suppose that m' is canonical with respect to $(\mathbb{H}, <)$ since any function locally interpolated by a function that acts idempotently and as a minority on $\{E, N\}$ acts idempotently and as a minority on $\{E, N\}$. It can be easily seen that the function $m(x, y, z) = m'(p_1(x, p_1(y, z)), p_1(y, p_1(z, x)), p_1(z, p_1(x, y)))$ is a ternary injection canonical with respect to $(\mathbb{H}, <)$ that acts like a minority on $\{E, N\}$.

For $\mathbf{a} \in H^\ell$, we write $O(\mathbf{a})$ for the orbit of \mathbf{a} under $\text{Aut}(\mathbb{H})$ and $O_{<}(\mathbf{a})$ for the orbit of \mathbf{a} under $\text{Aut}(\mathbb{H}, <)$. We say that a non-injective orbit O of ℓ -tuples under $\text{Aut}(\mathbb{H})$ is:

- *deterministic* if for every $\mathbf{a} \in O$, there exists $\alpha \in \text{Aut}(\mathbb{H})$ such that $p_1(O_{<}(\alpha(\mathbf{a})), E) = p_1(O_{<}(\alpha(\mathbf{a})), N)$, where the ordering on the second coordinate is strictly increasing,
- *non-deterministic* otherwise.

For a tuple \mathbf{a} contained in a deterministic orbit O , we call any $\alpha \in \text{Aut}(\mathbb{H})$ with the property that $p_1(O_{<}(\alpha(\mathbf{a})), E) = p_1(O_{<}(\alpha(\mathbf{a})), N)$ for the strictly increasing ordering in the second coordinate *deterministic for \mathbf{a}* . Note that for any $\beta \in \text{Aut}(\mathbb{H}, <)$, $\beta\alpha \in \text{Aut}(\mathbb{H})$ is deterministic for \mathbf{a} as well since p_1 is canonical with respect to $(\mathbb{H}, <)$.

5.2.1. Injectivisation of instances. Let \mathbb{A} be a first-order reduct of \mathbb{H} . Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be an instance of the $\text{CSP}(\mathbb{A})$. In this section, we always assume that the variable set \mathcal{V} is equipped with an arbitrary linear order; this assumption is inessential and is used to formulate the statements and proofs in a more concise way. We denote by $[\mathcal{V}]^\ell$ the set of injective increasing ℓ -tuples of variables from \mathcal{V} . Given any instance \mathcal{I} of $\text{CSP}(\mathbb{A})$, consider the following CSP instance \mathcal{I}_{fin} over the set \mathcal{O} of orbits of ℓ -tuples under $\text{Aut}(\mathbb{H})$:

- The variable set of \mathcal{I}_{fin} is the set $[\mathcal{V}]^\ell$.
- For every constraint $C \subseteq A^U$ in \mathcal{I} , \mathcal{I}_{fin} contains the constraint C' containing the maps $g: [U]^\ell \rightarrow \mathcal{O}$ such that there exists $f \in C$ satisfying $f(\mathbf{v}) \in g(\mathbf{v})$ for every $\mathbf{v} \in [U]^\ell$.

This instance was introduced in [40, Definition 3.1] in a more general setting, where following the notation from [40] it is denoted by $\mathcal{I}_{\text{Aut}(\mathbb{H}), \ell}$. In the original definition, ℓ -element subsets of \mathcal{V} were used as variables and the domain consisted of orbits of maps. However, the translation between the two definitions is straightforward.

Let $\mathcal{J} = (S, \mathcal{C})$ be an instance over the set \mathcal{O} of orbit of ℓ -tuples under $\text{Aut}(\mathbb{H})$, e.g., $\mathcal{J} = \mathcal{I}_{\text{fin}}$ for some \mathcal{I} . The *injectivisation* of \mathcal{J} , denoted by $\mathcal{J}^{(\text{inj})}$, is the instance obtained by replacing every constraint $C \in \mathcal{C}$ with scope $U \subseteq S$ by $\{g \in C \mid g(s) \in \{E, N\} \text{ for every } s \in U\}$.

Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be an instance of $\text{CSP}(\mathbb{A})$ and let $S \subseteq [\mathcal{V}]^\ell$. The *finite injectivisation of \mathcal{I} on S* is the instance $\mathcal{I}_{\text{fin}}^{(\text{inj})} \upharpoonright_S$. If $S = [\mathcal{V}]^\ell$, we call the finite injectivisation of \mathcal{I} on S just the finite injectivisation of \mathcal{I} . For any constraint $C \in \mathcal{C}$, the corresponding constraint in the finite injectivisation of \mathcal{I} is called the finite injectivisation of C . Note that if $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial, $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ is an instance of $\text{CSP}(\mathbb{A})$ and $S \subseteq [\mathcal{V}]^\ell$, the finite injectivisation of \mathcal{I} on S is solvable in polynomial time by Lemma 3.4 in [40] and by the dichotomy theorem for finite-domain CSPs [45, 46, 29].

Let \mathbb{A} be a first-order reduct of \mathbb{H} , let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be an instance of $\text{CSP}(\mathbb{A})$ and let $W \subseteq \mathcal{V}$. Let $g: W \rightarrow A$ and let $h: \mathcal{V} \rightarrow A$, we say that g is *ordered by h* if for all $u, v \in W$, we have $g(u) < g(v)$ if, and only if, $h(u) < h(v)$.

Let \mathbb{A} be a first-order reduct of \mathbb{H} preserved by m and by p_1 . We can assume that \mathbb{A} has among its relations all unions of orbits of ℓ -tuples under $\text{Aut}(\mathbb{H})$ that are preserved by p_1 and by the ternary injection m . Otherwise, we expand \mathbb{A} by these finitely many relations and we prove that the CSP of this expanded structure is solvable in polynomial time. Note that in particular, every orbit of ℓ -tuples under $\text{Aut}(\mathbb{H})$ is a relation of \mathbb{A} . Moreover, we will suppose that \mathbb{A} has among its relations all relations of arity at most $\max(3\ell, b_{\mathbb{H}})$ that are pp-definable in \mathbb{A} . Hence, for every instance \mathcal{I} of $\text{CSP}(\mathbb{A})$, the $(2\ell, \max(3, \ell, b_{\mathbb{H}}))$ -minimal instance equivalent to \mathcal{I} is again an instance of $\text{CSP}(\mathbb{A})$.

Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be an instance of $\text{CSP}(\mathbb{A})$ and let $C \in \mathcal{C}$. Since \mathbb{A} is preserved by m , there exists a set of linear equations over \mathbb{Z}_2 associated with the finite injectivisation of C . By abuse of notation, we write every linear equation as $\sum_{\mathbf{v} \in S} X_{\mathbf{v}} = P$, where $P \in \{E, N\}$ and S is a set of injective ℓ -tuples of variables from the scope of C . In these linear equations, we identify E with 1 and N with 0, so that e.g. $E + E = N$ and $N + E = E$.

We may assume that no equation $\sum_{\mathbf{v} \in S} X_{\mathbf{v}} = P$ associated with the finite injectivisation of any constraint $C \in \mathcal{C}$ splits into two equations $\sum_{\mathbf{v} \in S_1} X_{\mathbf{v}} = P_1$ and $\sum_{\mathbf{v} \in S_2} X_{\mathbf{v}} = P_2$ where $S_1 \cap S_2 = \emptyset$, $S_1 \cup S_2 = S$, $P_1 + P_2 = P$ and such that every element of the finite injectivisation of C satisfies both these equations. If this is not the case, let us consider only equations satisfying this assumption – it is clear that this new set of equations is satisfiable if, and only if, the original set of equations is satisfiable. We will call every equation satisfying this assumption *unsplittable*.

For an instance $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ of $\text{CSP}(\mathbb{A})$, we define an instance $\mathcal{I}_{\text{eq}} = (\mathcal{V}, \mathcal{C}_{\text{eq}})$ of the equality-CSP over the same base set H corresponding to the closure of the constraints under the full symmetric group on H . Formally, for every constraint $C \in \mathcal{C}$, the corresponding constraint $C_{\text{eq}} \in \mathcal{C}_{\text{eq}}$ contains all functions αh for all $h \in C$ and $\alpha \in \text{Sym}(H)$. Since \mathbb{A} is preserved by a binary injection, \mathcal{I}_{eq} is preserved by the same or indeed any binary injection and hence, its CSP has relational width $(2, 3)$ by the classification of equality CSPs [15].

```

INPUT:  $\ell$ -minimal instance  $\mathcal{I} = (\mathcal{V}, \mathcal{C})$  of  $\text{CSP}(\mathbb{A})$ ;
OUTPUT: eq-subdirect instance  $\mathcal{I}'$ ;
repeat
  changed:=false;
  for  $\mathbf{v} \in [\mathcal{V}]^\ell$  do
    if not changed then
       $P := \text{proj}_{\mathbf{v}}(\mathcal{I}) \cap I$ ;
      for  $O \subseteq \text{proj}_{\mathbf{v}}(\mathcal{I})$  non-injective orbit do
        if  $(\mathcal{I}^{\mathbf{v} \in O})_{\text{eq}}$  has a solution then
           $P := P \cup O$ ;
        end if
      end for
      for  $C \in \mathcal{C}$  containing all variables of  $\mathbf{v}$  in its scope do
        replace  $C$  by  $\{g \in C \mid g(\mathbf{v}) \in P\}$ ;
      end for
      if  $\mathcal{I}$  is not  $\ell$ -minimal then
        make  $\mathcal{I}$   $\ell$ -minimal;
        changed:=true;
      end if
    end if
  end for
until not changed
return  $\mathcal{I}$ 

```

FIGURE 1. Procedure EQ-SUBDIRECT

Let \mathcal{I} be an ℓ -minimal instance of $\text{CSP}(\mathbb{A})$, let $\mathbf{v} \in [\mathcal{V}]^\ell$ and let $R \subseteq \text{proj}_{\mathbf{v}}(\mathcal{I})$ be an ℓ -ary relation from the signature of \mathbb{A} . Let $\mathcal{I}^{\mathbf{v} \in R}$ be the instance obtained from \mathcal{I} by replacing every constraint C containing all variables from \mathbf{v} by $\{g \in C \mid g(\mathbf{v}) \in R\}$.

We call an ℓ -minimal instance of $\text{CSP}(\mathbb{A})$ *eq-subdirect* if for every $\mathbf{v} \in [\mathcal{V}]^\ell$ and for every non-injective orbit $O \subseteq \text{proj}_{\mathbf{v}}(\mathcal{I})$ under $\text{Aut}(\mathbb{H})$, the instance $(\mathcal{I}^{\mathbf{v} \in O})_{\text{eq}}$ has a solution. Note that by ℓ -minimality and since the instance is preserved by a binary injection, the instance $(\mathcal{I}^{\mathbf{v} \in O})_{\text{eq}}$ has a solution for every injective orbit $O \in \text{proj}_{\mathbf{v}}(\mathcal{I})$ under $\text{Aut}(\mathbb{H})$.

It is clear that we can obtain an eq-subdirect instance out of an ℓ -minimal instance in polynomial time by the algorithm in [Figure 1](#).

Lemma 16. *The instance \mathcal{I}' outputted by the algorithm in [Figure 1](#) is ℓ -minimal, eq-subdirect, and it is an instance of $\text{CSP}(\mathbb{A})$.*

Proof. It is easy to see that \mathcal{I}' is ℓ -minimal since in every run of the repeat... until not changed-loop, we check ℓ -minimality of the produced instance.

To prove that \mathcal{I}' is an instance of $\text{CSP}(\mathbb{A})$, it is sufficient to show that for every $\mathbf{v} \in [\mathcal{V}]^\ell$ the last P in the first for-loop is preserved by the binary injection p_1 as well as by the ternary injection m . To this end, let $\mathbf{v} \in [\mathcal{V}]^\ell$ be injective, let P be as in the algorithm, let O_1, O_2, O_3 be orbits of ℓ -tuples from P and let $g_i, i \in \{1, 2, 3\}$ be solutions to $(\mathcal{I}^{\mathbf{v} \in O_i})_{\text{eq}}$, $i \in \{1, 2, 3\}$.

Let $C \in \mathcal{C}$ be arbitrary and let U be its scope. By the definition of \mathcal{I}_{eq} , there exist $h_1, h_2, h_3 \in C$ and $\alpha_1, \alpha_2, \alpha_3 \in \text{Sym}(H)$ such that $g_i|_U = \alpha_i h_i$ for every $i \in \{1, 2, 3\}$. Since every binary injection is canonical in its action on $\{=, \neq\}$, it follows that $p_1(g_1, g_2)|_U = p_1(\alpha_1 h_1, \alpha_2 h_2) = \beta_1 p_1(h_1, h_2) \in C$ and $m(g_1, g_2, g_3)|_U = m(\alpha_1 h_1, \alpha_2 h_2, \alpha_3 h_3) = \beta_2 m(h_1, h_2, h_3) \in C$ for some $\beta_1, \beta_2 \in \text{Sym}(H)$. Hence, $p_1(g_1, g_2)|_U, m(g_1, g_2, g_3)|_U \in C$ and since C was chosen arbitrarily, both $p_1(g_1, g_2)$ and $m(g_1, g_2, g_3)$ are solutions to $(\mathcal{I}^{\mathbf{v} \in O_i})_{\text{eq}}$.

Let $\mathbf{v} \in [\mathcal{V}]^\ell$. If C contains all variables from \mathbf{v} in its scope, then either $p_1(g_1, g_2)(\mathbf{v})$ is injective and $p_1(g_1, g_2)(\mathbf{v}) \in P$ or $p_1(g_1, g_2)(\mathbf{v})$ and $p_1(h_1, h_2)(\mathbf{v})$ are non-injective and contained in the same orbit under $\text{Aut}(\mathbb{H})$ and it follows that $p_1(h_1, h_2)(\mathbf{v}) \in P$. Similar argument holds for $m(h_1, h_2, h_3)$ as well. Therefore, P is preserved by p_1 and m as desired. \square

Note that the eq-subdirect instance \mathcal{I}' outputted by the algorithm has the same solution set as \mathcal{I} . Moreover, for any $1 \leq m \leq n$ and any instance \mathcal{I} of $\text{CSP}(\mathbb{A})$, we can compute an instance that is both eq-subdirect and (m, n) -minimal and that has the same solution set as \mathcal{I} in polynomial time. Indeed, it is enough to repeat the above-mentioned algorithm and the (m, n) -minimality algorithm until no orbits under $\text{Aut}(\mathbb{H})$ are removed from any constraint.

5.2.2. Inj-irreducibility. Let $\mathcal{J} = (\mathcal{V}, \mathcal{C})$ be a CSP instance over a set B . A *path* in \mathcal{J} is a sequence $v_1, C_1, v_2, \dots, C_k, v_{k+1}$ where $k \geq 1$, $v_i \in \mathcal{V}$ for every $i \in [k+1]$, $C_i \in \mathcal{C}$ for every $i \in [k]$ and v_i, v_{i+1} are contained in the scope of C_i for every $i \in [k]$. We say that two elements $a, b \in B$ are *connected* by a path $v_1, C_1, v_2, \dots, C_k, v_{k+1}$ if there exists a tuple $(c_1, \dots, c_{k+1}) \in B^{k+1}$ such that $c_1 = a, c_{k+1} = b$ and such that $(c_i, c_{i+1}) \in \text{proj}_{(v_i, v_{i+1})}(C_i)$ for every $i \in [k]$. Let $\mathcal{J} = (\mathcal{V}, \mathcal{C})$ be a 1-minimal instance of $\text{CSP}(\mathbb{B})$ and let $v \in \mathcal{V}$. The *linkedness congruence* on $\text{proj}_v(\mathcal{J})$ is the equivalence relation λ on $\text{proj}_v(\mathcal{J})$ defined by $(a, b) \in \lambda$ if there exists a path from a to b in \mathcal{J} . Note that for a finite relational structure \mathbb{B} , for a $(2, 3)$ -minimal instance $\mathcal{J} = (\mathcal{V}, \mathcal{C})$ of $\text{CSP}(\mathbb{B})$ and for any $v \in \mathcal{V}$, the linkedness congruence λ on $\text{proj}_v(\mathcal{J})$ is a relation pp-definable in \mathbb{B} . Indeed, it is easy to see that for every $k \geq 1$, the binary relation containing precisely the pairs $(a, b) \in B^2$ that are connected by a particular path in \mathcal{I} is pp-definable in \mathbb{B} . If we concatenate all paths that connect two elements $(a, b) \in \lambda$, the resulting path will connect every pair $(a, b) \in \lambda$ since by the $(2, 3)$ -minimality of \mathcal{J} , every path from v to v connects c to c for every $c \in \text{proj}_v(\mathcal{J})$. It follows that λ is pp-definable.

Definition 17. Let \mathbb{A} be a first-order reduct of \mathbb{H} and let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be a non-trivial ℓ -minimal instance of $\text{CSP}(\mathbb{A})$. We call \mathcal{I} *inj-irreducible* if for every set $S \subseteq [\mathcal{V}]^\ell$, one of the following holds for the instance $\mathcal{J} = \mathcal{I}_{\text{fin}}|_S$:

- $\mathcal{J}^{(\text{inj})}$ has a solution,
- for some $\mathbf{v} \in S$, $\text{proj}_{\mathbf{v}}(\mathcal{J})$ contains the two injective orbits and the linkedness congruence on $\text{proj}_{\mathbf{v}}(\mathcal{J})$ does not connect them,
- for some $\mathbf{v} \in S$, the linkedness congruence on $\text{proj}_{\mathbf{v}}(\mathcal{J})$ links an injective orbit to a non-deterministic orbit.

Lemma 18. *Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be a non-trivial 2-minimal instance of $\text{CSP}(\mathbb{A})$ such that $\text{proj}_{(u,v)}(\mathcal{I}) \cap I_2 \neq \emptyset$ for every $u \neq v \in \mathcal{V}$. Let $S \subseteq [\mathcal{V}]^\ell$ be a set of variables appearing together in an unsplittable linear equation associated with the finite injectivisation of some constraint $C \in \mathcal{C}$. Then for every $g \in C$ either $g(\mathbf{s})$ is in an injective or deterministic orbit for all $\mathbf{s} \in S$ or $g(\mathbf{s})$ is in a non-deterministic orbit for all $\mathbf{s} \in S$.*

Proof. Suppose not and let $g \in C$ be a counterexample. Let $\emptyset \neq S' \subsetneq S$ be the set of all $\mathbf{s} \in S$ such that $g(\mathbf{s})$ is in an injective or deterministic orbit. Let $\sum_{\mathbf{s} \in S} X_{\mathbf{s}} = P$ be an unsplittable linear equation associated with the finite injectivisation of C and containing all the variables from S .

Let $\{\mathbf{s}_1, \dots, \mathbf{s}_n\}$ be the set of all $\mathbf{s}_i \in S'$ such that $g(\mathbf{s}_i)$ is in a deterministic orbit. For every $i \in [n]$, let $\alpha_i \in \text{Aut}(\mathbb{H})$ be deterministic for $g(\mathbf{s}_i)$. Let $p'_1(x, y) := p_1(\alpha_1 x, p_1(\alpha_2 x, p_1(\dots, p_1(\alpha_n x, y) \dots)))$. It is easy to see that for all $\mathbf{s} \in S$, we have that $p'_1(O_{<}(g(\mathbf{s})), E)$ and $p'_1(O_{<}(g(\mathbf{s})), N)$ are subsets of the same injective orbit $O_{\mathbf{s}}$ under $\text{Aut}(\mathbb{H})$ for the increasing order in the second coordinate. For all $\mathbf{s} \in S \setminus S'$, let $P_{\mathbf{s}}$ be N if $p'_1(O_{<}(g(\mathbf{s})), E) \subseteq E$ for the increasing order in the second coordinate and let $P_{\mathbf{s}}$ be E otherwise.

Since every binary projection of \mathcal{I} (and hence of C) has a non-empty intersection with I_2 , it follows that there exists a monotone injective $g' \in C$. Moreover, since $S' \subsetneq S$ and the equation

under consideration is unsplitable, one can assume that $\sum_{\mathbf{s} \in S'} O(g'(\mathbf{s})) = \sum_{\mathbf{s} \in S'} O_{\mathbf{s}} + \sum_{\mathbf{s} \in S \setminus S'} P_{\mathbf{s}} + E$.

Let us consider $h = p'_1(g, g') \in C$. h is clearly injective and we obtain:

$$\begin{aligned} \sum_{\mathbf{s} \in S} O(h(\mathbf{s})) &= \sum_{\mathbf{s} \in S'} O(h(\mathbf{s})) + \sum_{\mathbf{s} \in S \setminus S'} O(h(\mathbf{s})) = \sum_{\mathbf{s} \in S'} O_{\mathbf{s}} + \sum_{\mathbf{s} \in S \setminus S'} O(g'(\mathbf{s})) + \sum_{\mathbf{s} \in S \setminus S'} P_{\mathbf{s}} \\ &= \sum_{\mathbf{s} \in S'} O(g'(\mathbf{s})) + E + \sum_{\mathbf{s} \in S \setminus S'} O(g'(\mathbf{s})) = \sum_{\mathbf{s} \in S} O(g'(\mathbf{s})) + E \\ &= P + E \end{aligned}$$

Hence, the mapping $h': S \rightarrow \{E, N\}$ defined by $h'(\mathbf{s}) := O(h(\mathbf{s}))$ is a solution to the finite injectivisation of C on S but it does not satisfy the unsplitable linear equation, a contradiction. \square

Theorem 19. *Let \mathbb{A} be a first-order reduct of \mathbb{H} which is a model-complete core and suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial. Let \mathcal{I} be a $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal, eq-subdirect, inj-irreducible instance of $\text{CSP}(\mathbb{A})$ with variables \mathcal{V} such that for every distinct $u, v \in \mathcal{V}$, $\text{proj}_{(u,v)}(\mathcal{I}) \cap I_2 \neq \emptyset$. Then \mathcal{I} has a solution if, and only if, it has an injective one.*

Proof. In the case when $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine, the statement follows directly from Lemma 14 and Corollary 15, where only the $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimality of \mathcal{I} and the fact that for every distinct $u, v \in \mathcal{V}$, $\text{proj}_{(u,v)}(\mathcal{I}) \cap I_2 \neq \emptyset$ are needed.

Suppose henceforth that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally affine (and equationally non-trivial by assumption). Note that if \mathcal{I} has less than ℓ variables, it has an injective solution by the assumption on binary projections of \mathcal{I} and since it is preserved by a binary injection by Proposition 13. Let us therefore suppose that \mathcal{I} has at least ℓ variables. Let us assume for the sake of contradiction that \mathcal{I} does not have an injective solution. Let \mathcal{J} be \mathcal{I}_{fin} and let \mathcal{C} be the set of its constraints. By assumption, $\mathcal{J}^{(\text{inj})}$ does not have a solution. Note that $\mathcal{J}^{(\text{inj})}$ corresponds to a system of linear equations over \mathbb{Z}_2 , which is therefore unsatisfiable. In case this system can be written as a block matrix, there exists a set $S \subseteq [\mathcal{V}]^{\ell}$ of variables such that the system of equations associated with the injectivisation of $\mathcal{K} := \mathcal{J}|_S = (S, \mathcal{C}')$ corresponds to a minimal block, and is therefore unsatisfiable. By definition, this means that $\mathcal{K}^{(\text{inj})}$ is unsatisfiable. The instance \mathcal{K} has the property that for every constraint $C \in \mathcal{C}'$, and every non-trivial partition of the scope of C into parts S_1, S_2 , some unsplitable equation associated with C contains a variable from S_1 and a variable from S_2 .

Since \mathcal{I} is inj-irreducible, there exists $\mathbf{v} \in S$ such that the two injective orbits are elements of $\text{proj}_{\mathbf{v}}(\mathcal{K})$ and are not linked, or some injective orbit in $\text{proj}_{\mathbf{v}}(\mathcal{K})$ is linked to a non-deterministic orbit in $\text{proj}_{\mathbf{v}}(\mathcal{K})$.

In the first case, we note that for *all* $\mathbf{w} \in S$ such that $\text{proj}_{\mathbf{w}}(\mathcal{K})$ contains the two injective orbits, the two injective orbits are not linked. Indeed, suppose that there exists $\mathbf{w} \in S$ such that $E, N \in \text{proj}_{\mathbf{w}}(\mathcal{K})$ are linked, i.e., there exists a path $\mathbf{v}_1 = \mathbf{w}, C_1, \dots, C_k, \mathbf{v}_{k+1} = \mathbf{w}$ in \mathcal{K} connecting E and N . Since \mathcal{I} is $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal, Lemma 3.2 in [40] yields that \mathcal{J} and hence also \mathcal{K} is $(2, 3)$ -minimal. In particular, there exists $C \in \mathcal{C}'$ containing in its scope both \mathbf{v} and \mathbf{w} . Let $O_1, O_2 \in \{E, N\}$ be disjoint such that there exist $g_1, g_2 \in C$ with $g_1(\mathbf{v}) \in O_1, g_1(\mathbf{w}) \in E, g_2(\mathbf{v}) \in O_2, g_2(\mathbf{w}) \in N$. It follows that the path $\mathbf{v}, C, \mathbf{w} = \mathbf{v}_1, C_1, \dots, C_k, \mathbf{v}_{k+1} = \mathbf{w}, C, \mathbf{v}$ connects O_1 with O_2 in $\text{proj}_{\mathbf{v}}(\mathcal{J})$, a contradiction. Let $g: S \rightarrow \{E, N\}$ be defined as follows. For a fixed $\mathbf{v} \in S$, let $g(\mathbf{v})$ be an arbitrary element of $\text{proj}_{\mathbf{v}}(\mathcal{K}^{(\text{inj})})$. Next, for $\mathbf{w} \in S$, define $g(\mathbf{w})$ to be the unique injective orbit O such that there exists a path in \mathcal{K} from \mathbf{v} to \mathbf{w} connecting $g(\mathbf{v})$ to O . This g is a solution to $\mathcal{K}^{(\text{inj})}$, a contradiction.

Thus, it must be that a non-deterministic orbit in $\text{proj}_{\mathbf{v}}(\mathcal{K})$ is linked to an injective orbit in $\text{proj}_{\mathbf{v}}(\mathcal{K})$. Hence, there exists a path in \mathcal{K} from \mathbf{v} to \mathbf{v} and connecting an injective orbit to a non-deterministic one. Moreover, up to composing this path with additional constraints, one can assume that this path goes through all the variables in S . This follows by the $(2, 3)$ -minimality of

INPUT: instance $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ of $\text{CSP}(\mathbb{A})$;
OUTPUT: instance \mathcal{I}' that is either inj-irreducible or trivial and that has a solution if, and only, if \mathcal{I} has a solution;
repeat
 changed := false;
 $\mathcal{J} := (2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal eq-subdirect instance with the same solution set as \mathcal{I} ;
 $\mathcal{I} := \mathcal{J}$;
 $\mathcal{I} := \text{IDENTIFYALLEQUAL}(\mathcal{I})$;
 for $\mathbf{u} \in [\mathcal{V}]^\ell$ **do**
 for $\{E_{\mathbf{u}}^1, \dots, E_{\mathbf{u}}^m\}$ partition on $\text{proj}_{\mathbf{u}}(\mathcal{I})$ with pp-definable classes such that $\text{proj}_{\mathbf{u}}(\mathcal{I}) \cap I \subseteq E_{\mathbf{u}}^1$ and $E_{\mathbf{u}}^1$ contains no non-deterministic orbit **do**
 $S, \{E_{\mathbf{w}}^i \mid i \in [m], \mathbf{w} \in S\} := \text{EXTENDPARTITION}(\mathcal{I}, \{\mathbf{u}\}, \{E_{\mathbf{u}}^i \mid i \in [m]\})$;
 solve the finite injectivisation of \mathcal{I} on S ;
 if it does not have a solution and not changed **then**
 changed := true;
 for $\mathbf{v} \in S$ **do**
 for $C \in \mathcal{C}$ containing all variables of \mathbf{v} in its scope **do**
 replace C by $\{f \in C \mid f(\mathbf{v}) \notin E_{\mathbf{v}}^1\}$;
 end for
 end for
 end if
 end for
 end for
until not changed;
return \mathcal{I}

FIGURE 2. Procedure INJIRREDUCIBILITY

\mathcal{J} . Define a partition of S where $\mathbf{w} \in S_1$ if the first time that \mathbf{w} appears in the path, the element associated with \mathbf{w} is in an injective orbit, and $\mathbf{w} \in S_2$ otherwise. Since the system of unsplitable equations associated with $\mathcal{K}^{(\text{inj})}$ cannot be decomposed as a block matrix, some constraint $C \in \mathcal{C}'$ gives an equation in that system containing $\mathbf{u}_1 \in S_1$ and $\mathbf{u}_2 \in S_2$. Thus, there exists $g \in C$ with $g(\mathbf{u}_1)$ injective, and $g(\mathbf{u}_2)$ non-deterministic. This contradicts Lemma 18. \square

5.2.3. *Establishing inj-irreducibility.* We show that the algorithm in Figure 2 produces, given an instance \mathcal{I} of $\text{CSP}(\mathbb{A})$, an instance \mathcal{I}' of $\text{CSP}(\mathbb{A})$ that is either inj-irreducible or trivial and that has a solution if, and only if, \mathcal{I} has a solution. It uses the fact that the finite injectivisation of an instance \mathcal{I} of $\text{CSP}(\mathbb{A})$ on S is solvable in polynomial time for any set $S \subseteq [\mathcal{V}]^\ell$. This follows from the fact that the finite injectivisation of \mathcal{I} is preserved by a ternary minority by Lemma 3.4 from [40].

The algorithm uses the subroutines IDENTIFYALLEQUAL and EXTENDPARTITION. The subroutine IDENTIFYALLEQUAL takes as an input a 2-minimal instance $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ and returns an instance where all variables $u, v \in \mathcal{V}$ with $\text{proj}_{(u,v)}(\mathcal{I}) = \{(a, a) \mid a \in H\}$ are identified. It is clear that this can be implemented in polynomial time. Note moreover that for a $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal instance \mathcal{I} , the resulting instance IDENTIFYALLEQUAL(\mathcal{I}) is again $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal. The subroutine EXTENDPARTITION is given in Figure 3. We note that if $(S, \{E_{\mathbf{w}}^i\})$ is a set of partitions returned by EXTENDPARTITION, then for every $\mathbf{w} \in S$, the linkedness congruence on $\text{proj}_{\mathbf{w}}(\mathcal{I}_{\text{fin}})$ defined by the instance $\mathcal{I}_{\text{fin}}|_S$ is a refinement of the partition $\{E_{\mathbf{w}}^i\}$.

First of all, note that the instance \mathcal{I}' outputted by this algorithm is $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal and for every $u \neq v \in \mathcal{V}$, $\mathcal{I}'_{(u,v)} \cap I_2 \neq \emptyset$. We will show in Lemma 21 that \mathcal{I}' is an instance of $\text{CSP}(\mathbb{A})$.

INPUT: a tuple $(\mathcal{I}, S, \{E_{\mathbf{w}}^i \mid i \in [m], \mathbf{w} \in S\})$ where

- $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ is a $((2\ell, \max(3\ell, b_{\mathbb{H}})))$ -minimal instance of $\text{CSP}(\mathbb{A})$,
- $S \subseteq [\mathcal{V}]^\ell$,
- $\{E_{\mathbf{w}}^1, \dots, E_{\mathbf{w}}^m\}$ is a partition on $\text{proj}_{\mathbf{w}}(\mathcal{I})$ with pp-definable classes for every $\mathbf{w} \in S$ such that $E_{\mathbf{w}}^1$ contains no non-deterministic orbit and such that $E_{\mathbf{w}}^i := \{\mathbf{a} \in H^\ell \mid \exists \mathbf{b} \in H^\ell: (\mathbf{b}, \mathbf{a}) \in \text{proj}_{(\mathbf{v}, \mathbf{w})}(\mathcal{I})\}$ for every $\mathbf{v}, \mathbf{w} \in S$ and every $i \in [m]$;

OUTPUT: $S, \{E_{\mathbf{w}}^i : i \in [m], \mathbf{w} \in S\}$ as above such that no tuple \mathbf{w} can be added to S where a partition on $\text{proj}_{\mathbf{w}}(\mathcal{I})$ as above exists;

repeat
 added := false;
for $\mathbf{v} \in S, \mathbf{w} \in [\mathcal{V}]^\ell$ with $\mathbf{w} \notin S$ **do**
 $D := \text{proj}_{(\mathbf{v}, \mathbf{w})}(\mathcal{I})$;
 for $t = 1, \dots, m$ **do**
 $E_{\mathbf{w}}^t := \{\mathbf{a} \in H^\ell \mid \exists \mathbf{b} \in E_{\mathbf{v}}^t: (\mathbf{b}, \mathbf{a}) \in D\}$;
 end for
 if $E_{\mathbf{w}}^1, \dots, E_{\mathbf{w}}^m$ are disjoint and $E_{\mathbf{w}}^1$ contains no non-deterministic orbit **then**
 $S := S \cup \{\mathbf{w}\}$;
 added := true;
 end if
end for
until not added
return $S, \{E_{\mathbf{w}}^i : i \in [m], \mathbf{w} \in S\}$

FIGURE 3. Procedure EXTENDPARTITION

Then it follows that if \mathcal{I}' is non-trivial, then every constraint C' of \mathcal{I}' contains an injective tuple since \mathcal{I}' is preserved by the binary injection p_1 .

Let $S \subseteq [\mathcal{V}]^\ell$ be the set outputted by $\text{EXTENDPARTITION}(\mathcal{I}, \{\mathbf{u}\}, \{E_{\mathbf{u}}^i \mid i \in [m]\})$ for some $\mathbf{u} \in [\mathcal{V}]^\ell$ as in the algorithm for an instance \mathcal{I} with $\mathcal{I} = \text{IDENTIFYALLEQUAL}(\mathcal{I})$ – note that this subroutine is called in the algorithm only for instances satisfying this assumption. Note that for every $\mathbf{w} \in S$, $E_{\mathbf{w}}^1$ contains all injective orbits under $\text{Aut}(\mathbb{H})$ contained in $\text{proj}_{\mathbf{w}}(\mathcal{I})$. This can be easily shown by induction of the size of S . For S containing just the tuple \mathbf{u} , there is nothing to prove. Suppose that it holds for S , let $\mathbf{w} \in \mathcal{V}^\ell$ be a tuple added to S and let $\mathbf{v} \in S$ be such that $E_{\mathbf{w}}^1 = \{\mathbf{a} \in H^\ell \mid \exists \mathbf{b} \in H^\ell: (\mathbf{b}, \mathbf{a}) \in \text{proj}_{(\mathbf{v}, \mathbf{w})}(\mathcal{I})\}$. Let $O \subseteq \text{proj}_{\mathbf{w}}(\mathcal{I})$ be an injective orbit under $\text{Aut}(\mathbb{H})$ and let $C \in \mathcal{C}$ be a constraint containing all variables from the tuples \mathbf{v} and \mathbf{w} in its scope. Let $g_1 \in C$ be such that $g_1(\mathbf{w}) \in O$ and let $g_2 \in C$ be injective. Then $p_1(g_1, g_2) \in C$ is injective and hence, $p_1(g_1(\mathbf{v}), g_2(\mathbf{v})) \in E_{\mathbf{v}}^1$ by the induction assumption. Moreover, $p_1(g_1(\mathbf{w}), g_2(\mathbf{w})) \in O$ and $p_1(g_1(\mathbf{w}), g_2(\mathbf{w})) \in E_{\mathbf{w}}^1$ by the definition of $E_{\mathbf{w}}^1$.

It is easy to see that the algorithm runs in polynomial time: the algorithm of checking eq-subdirectness as well as the $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimality algorithm both run in polynomial time. Moreover, the number of runs of every for- or repeat...until-loop in the main algorithm as well as in the subroutine EXTENDPARTITION is bounded by $c^c(d + \ell^{\max(3\ell, b_{\mathbb{H}})} \binom{|\mathcal{V}|}{\ell})^2$, where c is the number of orbits of ℓ -tuples under $\text{Aut}(\mathbb{H})$ and d is the number of constraints of the original instance \mathcal{I} .

Before proving that \mathcal{I}' is an inj-irreducible instance of $\text{CSP}(\mathbb{A})$ and that it has a solution if, and only if, \mathcal{I} has a solution, we will show a few auxiliary statements.

Lemma 20. *Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be an ℓ -minimal non-trivial instance of $\text{CSP}(\mathbb{A})$ such that $\text{proj}_{(u,v)}(\mathcal{I}) \cap I_2 \neq \emptyset$ for every $u \neq v \in \mathcal{V}$. Let $S \subseteq [\mathcal{V}]^\ell$ be such that $\text{proj}_{\mathbf{v}}(\mathcal{I})$ contains some injective orbit and no non-deterministic orbit for every $\mathbf{v} \in S$. Suppose that \mathcal{I} has a solution. Then the finite injectivisation of \mathcal{I} on S has a solution as well.*

Proof. Let $g: \mathcal{V} \rightarrow H$ be a solution of \mathcal{I} such that $g(\mathbf{s})$ is not injective for some $\mathbf{s} \in S$. Let $\{\mathbf{s}_1, \dots, \mathbf{s}_n\}$ be the set of all $\mathbf{s}_i \in S'$ such that $g(\mathbf{s}_i)$ is in a non-injective orbit under $\text{Aut}(\mathbb{H})$. For every $i \in [n]$, let $\alpha_i \in \text{Aut}(\mathbb{H})$ be deterministic for $g(\mathbf{s}_i)$. Let $p'_1(x, y) := p_1(\alpha_1 x, p_1(\alpha_2 x, p_1(\dots, p_1(\alpha_n x, y))))$.

It is easy to see that for all $\mathbf{s} \in S$, we have that $p'_1(O_{<}(g(\mathbf{s})), E)$ and $p'_1(O_{<}(g(\mathbf{s})), N)$ are subsets of the same injective orbit $O_{\mathbf{s}}$ under $\text{Aut}(\mathbb{H})$ for the increasing order in the second coordinate.

We claim that $f: S \rightarrow \{E, N\}$ defined by $f(\mathbf{s}) = O_{\mathbf{s}}$ is a solution to the finite injectivisation of \mathcal{I} on S . To this end, let $C \in \mathcal{C}$ and let $g' \in C$ be monotone and injective. It follows that $p'_1(g, g') \in C$ and moreover, $O(p'_1(g(\mathbf{s}), g'(\mathbf{s}))) = O_{\mathbf{s}}$ for all $\mathbf{s} \in S$ such that all variables from \mathbf{s} are in the scope of C and the lemma follows. \square

Lemma 21. *Let $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ be a $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal, non-trivial, eq-subdirect instance of $\text{CSP}(\mathbb{A})$ such that $\text{proj}_{(u,v)}(\mathcal{I}) \cap I_2 \neq \emptyset$ for every $u \neq v \in \mathcal{V}$. Let $S \subseteq [\mathcal{V}]^\ell$ and let $\{E_{\mathbf{s}}^i \mid \mathbf{s} \in S, i \in [m]\}$ be sets of classes of the partitions from the algorithm. Suppose that there exists $\mathbf{s}' \in S$ such that the relation $\bigcup_{i \in \{2, \dots, m\}} E_{\mathbf{s}'}^i$ is not preserved by p_1 or by m . Then the finite injectivisation of \mathcal{I} on S has a solution.*

Proof. Suppose that the relation $R := \bigcup_{i \in \{2, \dots, m\}} E_{\mathbf{s}'}^i$ is not preserved by p_1 . Let $\mathbf{a}_1, \mathbf{a}_2 \in R$ be such that $p_1(\mathbf{a}_1, \mathbf{a}_2) \in E_{\mathbf{s}'}^1$.

For $j \in [2]$, let O^j be the orbit of \mathbf{a}_j under $\text{Aut}(\mathbb{H})$. Let $h^j: \mathcal{V} \rightarrow H$ be a solution to $(\mathcal{I}^{\mathbf{s}' \in O^j})_{\text{eq}}$ such that $h^j(\mathbf{s}') = \mathbf{a}_j$. Let $h := p_1(h^1, h^2)$. It follows that $h(\mathbf{s}') \in E_{\mathbf{s}'}^1$ and, by the way how the partitions are obtained, $h(\mathbf{s}) \in E_{\mathbf{s}}^1$ for every $\mathbf{s} \in S$. In particular, $h(\mathbf{s})$ lies in an injective or deterministic under $\text{Aut}(\mathbb{H})$ for every $\mathbf{s} \in S$.

Let $\mathbf{s}_1, \dots, \mathbf{s}_n$ be all tuples in S such that $h(\mathbf{s}_i)$ is not injective. For every $i \in [n]$, let $\beta_i \in \text{Aut}(\mathbb{H})$ be such that β_i is deterministic for $h(\mathbf{s}_i)$. Let now $p''_1(x, y) := p_1(\beta_1 x, p_1(\beta_2 x, p_1(\dots, p_1(\beta_n x, y))))$. Observe that for all $\mathbf{s} \in S$, we have that $p''_1(O_{<}(h(\mathbf{s})), E)$ and $p''_1(O_{<}(h(\mathbf{s})), N)$ are subsets of the same orbit $O_{\mathbf{s}} \in \{E, N\}$ under $\text{Aut}(\mathbb{H})$. Let $f: S \rightarrow \{E, N\}$ be defined as $f(\mathbf{s}) = O_{\mathbf{s}}$. We claim that f is a solution of the finite injectivisation of \mathcal{I} on S .

To this end, let $C \in \mathcal{C}$. By the choice of h^1, h^2 , there exist $g^1, g^2 \in C$ ordered by h^1 and h^2 , respectively. It follows that $g := p_1(g^1, g^2)$ is ordered by h and moreover, $g \in C$. For every $i \in [n]$, let $\alpha_i \in \text{Aut}(\mathbb{H})$ be such that $\alpha_i g$ is ordered by $\beta_i h$. Let $p'_1(x, y) := p_1(\alpha_1 x, p_1(\alpha_2 x, p_1(\dots, p_1(\alpha_n x, y))))$ and let $g' \in C$ be monotone and injective. It follows that for every $\mathbf{s} \in S$ such that all variables from \mathbf{s} are contained in the scope of C and for every $i \in [n]$, $O_{<}(\alpha_i g(\mathbf{s})) = O_{<}(\beta_i h(\mathbf{s}))$. In particular, $p'_1(g(\mathbf{s}), g'(\mathbf{s})) \in f(\mathbf{s})$ as desired.

The case when R is not preserved by m is similar. \square

Using the auxiliary statements above, we are able to prove the correctness of the main algorithm:

Theorem 22. *The instance \mathcal{I}' produced by the algorithm in [Figure 2](#) is an instance of $\text{CSP}(\mathbb{A})$ and it has a solution if, and only if, the original instance has a solution. Moreover, \mathcal{I}' is either trivial or inj-irreducible.*

Proof. It is easy to see that \mathcal{I}' is an instance of $\text{CSP}(\mathbb{A})$ by the algorithm – the only thing that needs to be proven is that if the algorithm constrains an injective tuple $\mathbf{v} \in \mathcal{V}^\ell$ by $\text{proj}_{\mathbf{v}}(\mathcal{I}) \setminus E_{\mathbf{v}}^1$, where $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ is an eq-subdirect, $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal instance obtained during the run of the algorithm, then $\text{proj}_{\mathbf{v}}(\mathcal{I}) \setminus E_{\mathbf{v}}^1$ is preserved by p_1 and by m . But this follows directly from [Lemma 21](#).

We will now show that if the original instance has a solution, the instance \mathcal{I}' has a solution as well. Suppose that in the repeat... until not changed loop, we get an instance $\mathcal{I} = (\mathcal{V}, \mathcal{C})$ and we removed a solution of \mathcal{I} during one run of this loop. Clearly, we didn't remove any solutions by making the instance eq-subdirect, $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal and by identifying variables in the IDENTIFYALLEQUAL subroutine. Hence, the only problem could occur when removing the classes $E_{\mathbf{w}}^1$.

Suppose therefore that there is a set $S \subseteq [\mathcal{V}]^\ell$ and a partition $\{E_{\mathbf{v}}^j \mid j \in \{1, \dots, m\}, \mathbf{v} \in S\}$ on the projections of \mathcal{I} to the tuples from S . Suppose moreover that \mathcal{I} had a solution $s: \mathcal{V} \rightarrow H$ such that $s(\mathbf{w}) \in E_{\mathbf{w}}^1$ for some $\mathbf{w} \in S$. Then $s(\mathbf{v}) \in E_{\mathbf{v}}^1$ for every $\mathbf{v} \in S$ by the definition of the partition.

Recall that $E_{\mathbf{v}}^1$ contains all injective orbits under $\text{Aut}(\mathbb{H})$ that are contained in $\text{proj}_{\mathbf{v}}(\mathcal{I})$ for all tuples $\mathbf{v} \in S$ by the reasoning below the algorithm. Let \mathcal{I}'' be an ℓ -minimal instance equivalent to the instance obtained from \mathcal{I} by adding for every $\mathbf{v} = (v_1, \dots, v_\ell) \in S$ the constraint $\{g: \{v_1, \dots, v_\ell\} \rightarrow H \mid g(\mathbf{v}) \in E_{\mathbf{v}}^1\}$. Since $E_{\mathbf{v}}^1$ is pp-definable in \mathbb{A} , it is by assumption one of the relations of \mathbb{A} and therefore \mathcal{I}'' is an instance of $\text{CSP}(\mathbb{A})$. Since s is a solution to \mathcal{I} and hence also to \mathcal{I}'' , \mathcal{I}'' is non-trivial. Since the classes $\{E_{\mathbf{v}}^1 \mid \mathbf{v} \in S\}$ were removed, the finite injectivisation of \mathcal{I} on S does not have a solution and it follows that the finite injectivisation of \mathcal{I}'' on S does not have a solution as well. Hence, s is not a solution to \mathcal{I}'' by [Lemma 20](#), a contradiction.

We finally prove that if \mathcal{I}' is non-trivial, it is inj-irreducible. Note that by the algorithm, \mathcal{I}' is $(2\ell, \max(3\ell, b_{\mathbb{H}}))$ -minimal and if it is non-trivial, then any projection of \mathcal{I}' to a pair of disjoint variables has a non-empty intersection with I_2 . In particular, for every injective tuple \mathbf{v} of variables of \mathcal{I}' , $\text{proj}_{\mathbf{v}}(\mathcal{I}')$ contains an injective orbit under $\text{Aut}(\mathbb{H})$. Let S be a subset of variables of $\mathcal{I}'_{\text{fin}}$.

Suppose that for some variable \mathbf{w} in S , the linkedness congruence on $\text{proj}_{\mathbf{w}}(\mathcal{I}'_{\text{fin}} \upharpoonright_S)$ links the injective orbits within this set, and separates the injective orbits from the non-deterministic ones.

Each block $B_i^{\mathbf{w}}$ of the linkedness congruence on $\text{proj}_{\mathbf{w}}(\mathcal{I}'_{\text{fin}} \upharpoonright_S)$ defines a subset of $\text{proj}_{\mathbf{w}}(\mathcal{I}')$ by taking $E_i^{\mathbf{w}} := \bigcup B_i^{\mathbf{w}}$. By assumption, this partition of $\text{proj}_{\mathbf{w}}(\mathcal{I}')$ extends to a partition $\{E_{\mathbf{v}}^1, \dots, E_{\mathbf{v}}^m\}$ of $\text{proj}_{\mathbf{v}}(\mathcal{I}')$ for $\mathbf{v} \in S$. Each $E_i^{\mathbf{w}}$ is pp-definable: for an arbitrary orbit O in $B_i^{\mathbf{w}}$, $E_i^{\mathbf{w}}$ consists of the tuples that are reachable from some $\mathbf{a} \in O$ by a suitable path of constraints. Since \mathbb{A} contains all orbits of ℓ -tuples under $\text{Aut}(\mathbb{H})$ by our assumptions, O is a relation of \mathbb{A} . Thus, the algorithm has checked that the finite injectivisation of \mathcal{I}' on S has a solution, showing the inj-irreducibility of \mathcal{I}' . \square

By combining [Theorems 19](#) and [22](#), we obtain the desired result.

Corollary 23. *Let \mathbb{A} be a first-order reduct of \mathbb{H} that is a model-complete core and such that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial. Then $\text{CSP}(\mathbb{A})$ is in P.*

6. THE NP-HARD CASE

In this section, we prove that if the first case of [Theorem 8](#) applies, $\text{CSP}(\mathbb{A})$ is NP-hard.

Lemma 24. *Let \mathbb{A} be a first-order reduct of \mathbb{H} that is a model-complete core. Suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally trivial. Then so is $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$. Moreover, $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \subseteq \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$.*

Proof. Suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ is equationally non-trivial. Then $\text{Pol}(\mathbb{A})$ contains a binary injection by [Lemma 12](#).

We claim that $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ has an operation s which satisfies the pseudo-Siggers identity modulo $\text{Aut}(\mathbb{A})$ on injective tuples. To see this, consider the actions of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ on $I_n/\text{Aut}(\mathbb{A})$ for all $n \geq 1$. Each of these actions has a Siggers operation s_n , and we may assume that the sequence $(s_n)_{n \geq 1}$ converges in $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ to a function s , which has the desired property.

Let s' be a polymorphism canonical with respect to $(\mathbb{H}, <)$ that is locally interpolated by s modulo $\text{Aut}(\mathbb{H}, <)$. Then on all injective tuples x, y, z of the same length, s' satisfies the pseudo-Siggers equation on x, y, z modulo $\text{Aut}(\mathbb{A})$, by a standard computation (see e.g. the proof of [Lemma 35](#) in [\[38\]](#)). Now for all $1 \leq i \leq 6$ let p_i be a 6-ary function which is canonical with respect to $(\mathbb{H}, <)$, behaves like the i -th projection on $\{E, N\}$, and which acts lexicographically on the order – such functions can easily be obtained from the binary injections from [Proposition 13](#) and by the remark above [Lemma 12](#). Set $s'' := s'(p_1(x_1, \dots, x_6), \dots, p_6(x_1, \dots, x_6))$. Then $s'' \in \mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ since application of p_1, \dots, p_6 makes forget the order. Say without loss of generality that s'' behaves like the first projection on $\{E, N\}$; the case where it flips E and N of the first coordinate is similar.

Let a, b be injective increasing tuples in distinct orbits under $\text{Aut}(\mathbb{A})$ (such tuples only exist if \mathbb{A} is not reduct of $(H; =)$, but we already know from [Lemmas 11](#) and [12](#) that in those cases, $\text{Pol}(\mathbb{A})$ consists of essentially unary operations only and thus the result is also true in this case). Then

$$a \sim_{\text{Aut}(\mathbb{H})} s''(a, b, \dots) \sim_{\text{Aut}(\mathbb{A})} s''(b, a, \dots) \sim_{\text{Aut}(\mathbb{H})} b,$$

a contradiction.

The proof that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \subseteq \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ is almost identical as the proof of [Lemma 32](#) in [\[38\]](#) but we provide it for the convenience of the reader. Observe that the action $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is by essentially unary functions by [\[42\]](#). Moreover, the action $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ determines the action of $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ on orbits of injective tuples under $\text{Aut}(\mathbb{H})$ since \mathbb{H} is homogeneous in an ℓ -ary language. Hence, for every $f \in \mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ of arity $n \geq 1$, there exists $1 \leq i \leq n$ such that the orbit of $f(a_1, \dots, a_n)$ under $\text{Aut}(\mathbb{H})$ is either equal to the orbit of a_i under $\text{Aut}(\mathbb{H})$ for all injective tuples a_1, \dots, a_n of the same length, or the orbit of $f(a_1, \dots, a_n)$ under $\text{Aut}(\mathbb{H})$ is equal to the orbit under $\text{Aut}(\mathbb{H})$ of a tuple obtained from a_i by changing hyperedges to non-hyperedges and vice versa. In the first case, $f \in \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ since the orbits under $\text{Aut}(\mathbb{H})$ are refinements of the orbits under $\text{Aut}(\mathbb{A})$. In the second case, $\text{Aut}(\mathbb{A})$ contains a function α changing hyperedges into non-hyperedges and vice versa and hence, $\alpha \circ f \in \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ and $f \in \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ follows. \square

The following is an adjusted version of the fundamental theorem of smooth approximations [\[38, Theorem 12\]](#). The original version of this theorem does not apply to our case since for a first-order reduct \mathbb{A} of \mathbb{H} , we do not have that $\text{Pol}(\mathbb{A})$ locally interpolates $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ modulo $\text{Aut}(\mathbb{H})$.

Lemma 25. *Let \mathbb{A} be a first-order reduct of \mathbb{H} that is a model-complete core. Assume that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \subseteq \mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ and that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ does not contain a binary function that acts as a semilattice operation on $\{E, N\}$. Let $k \geq 1$ be so that $I_k/\text{Aut}(\mathbb{A})$ has at least two elements and let (S, \sim) be a subfactor of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}} \curvearrowright I_k/\text{Aut}(\mathbb{A})$ with $\text{Aut}(\mathbb{A})$ -invariant equivalence classes. Assume moreover that η is a very smooth approximation of \sim which is invariant under $\text{Pol}(\mathbb{A})$. Then $\text{Pol}(\mathbb{A})$ has a uniformly continuous clone homomorphism to $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}} \curvearrowright S/\sim$.*

Proof. We may assume that $\text{Pol}(\mathbb{A})$ contains an essential operation – otherwise, $\text{Pol}(\mathbb{A})$ has a uniformly continuous clone homomorphism to the clone of projections and hence, to any clone. For all $n \geq 1$ and all $1 \leq i \leq n$, let $p_i^n \in \text{Pol}(\mathbb{A})$ be an n -ary injection which is canonical with respect to $\text{Aut}(\mathbb{H}, <)$, acts like the i -th projection on $\{E, N\}$, and follows the first coordinate for the order. These functions are easily obtained from the binary injection acting as a projection on $\{E, N\}$ from [Lemma 12](#). We define a map ϕ from $\text{Pol}(\mathbb{A})$ to $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}} \curvearrowright S/\sim$ as follows. For every $n \geq 1$ and every n -ary $f \in \text{Pol}(\mathbb{A})$, we take any function f' canonical on $(\mathbb{H}, <)$ which is locally interpolated by f modulo $\text{Aut}(\mathbb{H}, <)$, and then set $f'' := f'(p_1^n, \dots, p_n^n)$; this function is an element of $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$, and hence also of $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}}$ by assumption. We set $\phi(f)$ to be f'' acting on S/\sim .

Our first claim is that ϕ is well-defined, i.e., it does not depend on the choice of f' . Let $u_1, \dots, u_n \in S$. Then

$$f(u_1, \dots, u_n) (\eta \circ \sim) f'(u_1, \dots, u_n) \sim f''(u_1, \dots, u_n).$$

Hence, the action of f'' on S/\sim is completely determined by f , and consequently independent of f' .

It remains to show that ϕ is a uniformly continuous clone homomorphism. ϕ clearly preserves arities, let us show that it preserves compositions as well. Let $n, m \geq 1$, let $f \in \text{Pol}(\mathbb{A})$ be m -ary, let g_1, \dots, g_m be n -ary, and let $u_1, \dots, u_n \in S$. Pick, for all $1 \leq i \leq m$, any $v_i \in S$ such that $v_i \eta g_i(u_1, \dots, u_n)$. Then

$$f(g_1, \dots, g_m)(u_1, \dots, u_n) \eta f(v_1, \dots, v_m) (\eta \circ \sim) f''(v_1, \dots, v_m) (\eta \circ \sim) f''(g_1'', \dots, g_m'')(u_1, \dots, u_n),$$

the last equivalence holding since $g_i(u_1, \dots, u_n)$ ($\eta \circ \sim$) $g''(u_1, \dots, u_n)$ for all $i \in [n]$. Hence, $\phi(f(g_1, \dots, g_m)) = \phi(f)(\phi(g_1), \dots, \phi(g_m))$. It follows that ϕ is a clone homomorphism. Moreover, ϕ is uniformly continuous since for any fixed set of representatives of the classes of \sim and for any $f \in \text{Pol}(\mathbb{A})$, the action of f on these representatives determines $\phi(f)$ by the above proof showing that ϕ is well-defined. \square

7. THE IMPOSSIBLE CASE: FEW CANONICAL FUNCTIONS AND A WEAKLY COMMUTATIVE FUNCTION

In this section, we prove that if \mathbb{A} is a first-order reduct of \mathbb{H} which has among its polymorphisms a weakly commutative function then $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a binary function that acts as a semilattice operation on $\{E, N\}$. This will allow us to derive a contradiction if the second case of [Theorem 8](#) applies. To prove this, we will need the following consequence from [\[10\]](#) of the fact that $(\mathbb{H}, <)$ satisfies the Ramsey property.

Let \mathbb{C} be a relational structure and let $A, B \subseteq C$. We say that A is *independent from B* in \mathbb{C} if for every $n \geq 1$ and for all $a_1, a_2 \in A^n$ such that a_1, a_2 satisfy the same first-order formulas over \mathbb{C} , a_1 and a_2 satisfy the same first-order formulas over \mathbb{C} with parameters from B . We say that two substructures \mathbb{A} and \mathbb{B} of \mathbb{C} are *independent* if A is independent from B in \mathbb{C} and B is independent from A in \mathbb{C} . Finally, we say that a substructure \mathbb{A} of \mathbb{C} is *elementary* if the identity mapping from A to C preserves the truth of all first-order formulas in the signature of \mathbb{C} .

Theorem 26 (Theorem 4.4. in [\[10\]](#)). *Let \mathbb{C} be a countable homogeneous ω -categorical Ramsey structure. Then \mathbb{C} contains two independent elementary substructures.*

Let \mathbb{A} be a first-order reduct of \mathbb{H} which is a model-complete core and so that $\text{Pol}(\mathbb{A})$ is equationally non-trivial and let us assume that the second case of [Proposition 13](#) applies, i.e., $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a binary injection g_1 which acts as a projection on $\{E, N\}$ and which is canonical with respect to $(\mathbb{H}, <)$. We will show that in this case, there exist $p_1, p_2 \in \mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ with the following properties.

- p_1, p_2 are both canonical with respect to $(\mathbb{H}, <)$,
- p_i acts like the i -th projection on $\{E, N\}$ for all $i \in \{1, 2\}$,
- p_i acts lexicographically on the order for all $i \in \{1, 2\}$,
- for all $x, y \in H$, if $x < y$, then $p_1(x, y) < p_2(x, y)$, and if $y < x$, then $p_1(x, y) > p_2(x, y)$,
- the ranges of p_1 and p_2 are disjoint and independent as substructures of \mathbb{H} .

To show the existence of p_1, p_2 , assume without loss of generality that g_1 acts as the first projection on $\{E, N\}$. Then the function $g_2(x, y) := g_1(y, x)$ acts as the second projection on $\{E, N\}$. By the remark above [Lemma 12](#), we may assume that both g_1 and g_2 act lexicographically on the order $<$. Finally, by [Theorem 26](#), we may compose g_1 and g_2 with automorphisms of $(\mathbb{H}, <)$ and obtain binary injections g'_1, g'_2 that still satisfy all the assumptions above and whose ranges are disjoint and induce in $(\mathbb{H}, <)$ substructures that are independent.

Let us define a linear order $<^*$ on $U := \text{im}(g'_1) \cup \text{im}(g'_2)$ as follows. We set $u <^* v$ if one of the following holds.

- $u < v$ and $u, v \in \text{im}(g'_i)$ for some $i \in [2]$, or
- $u = g'_i(x_1, y_1), v = g'_j(x_2, y_2)$ for some $i \neq j \in [2]$, $x_1, x_2, y_1, y_2 \in H$ and one of the following holds
 - $i = 1, j = 2, x_2 \leq y_2$ and $u \leq g'_1(x_2, y_2)$, or
 - $i = 1, j = 2, x_1 \leq y_1$ and $g'_2(x_1, y_1) \leq v$, or
 - $i = 2, j = 1, x_2 > y_2$ and $u \leq g'_2(x_2, y_2)$, or
 - $i = 2, j = 1, x_1 > y_1$ and $g'_1(x_1, y_1) \leq v$.

It is easy to verify that $<^*$ is a linear order on U . Let us define a hyperedge relation E^* on U as the restriction of the relation E to U . It follows that (U, E^*) is isomorphic to the structure

induced by the union of the ranges of g'_1 and g'_2 in \mathbb{H} and hence, $(U, E^*, <^*)$ embeds to $(\mathbb{H}, <)$ by an embedding e since $(\mathbb{H}, <)$ is universal for the class of all ℓ -uniform linearly ordered hypergraphs $(\mathbb{X}, <^{\mathbb{X}})$ such that \mathbb{X} embeds to \mathbb{H} . Finally, we obtain the desired functions p_1 and p_2 by setting $p_1 := e \circ g'_1$ and $p_2 := e \circ g'_2$.

Lemma 27. *Let \mathbb{A} be a first-order reduct of \mathbb{H} which is a model-complete core. Suppose that $\text{Pol}(\mathbb{A})$ does not have a uniformly continuous clone homomorphism to \mathcal{P} and contains a binary function f such that there exist $k \geq 1$, $S \subseteq I_k$, and an equivalence relation \sim on S with at least two $\text{Aut}(\mathbb{A})$ -invariant classes such that $f(a, b) \sim f(b, a)$ for all disjoint injective tuples $a, b \in H^k$ with $f(a, b), f(b, a) \in S$. Then $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ contains a semilattice operation.*

Proof. Since \mathbb{A} is a model-complete core, we have $f(a, a) \in S$ for all $a \in S$, and hence picking any $b \in S$ which is not \sim -equivalent to a , we see on the values $f(a, a), f(b, b), f(a, b)$, and $f(b, a)$ that f must be essential. It follows from Proposition 13, from Theorem 26 and from the discussion above that $\text{Pol}(\mathbb{A})$ contains functions p_1, p_2 as above, or else the conclusion of the lemma follows immediately. Moreover, since its distinctive property is stable under diagonal interpolation modulo $\text{Aut}(\mathbb{H})$, we can assume that f is diagonally canonical with respect to $\text{Aut}(\mathbb{H}, <)$.

It is easy to see that for every $U', V' \subseteq H$ such that $u < v$ for every $u \in U', v \in V'$, f is canonical with respect to $(\mathbb{H}, <)$ on $U' \times V'$. Let $c_1, c_2 \in H$ be arbitrary, let $U := \{p_1(d, c_2) \mid d \in H, d < c_1\}$ and let $V := \{p_2(c_1, d) \mid d \in H, c_2 < d\}$. Hence, for every $u \in U, v \in V$, it holds that $u < v$. Note that for every $m \geq 1$ and for every $a \in H^m$, there exist $b \in U^m, c \in V^m$ such that a, b, c are in the same orbit under $\text{Aut}(\mathbb{H}, <)$. Note moreover that if $m \geq 1$ and (a, b) is any pair of m -tuples such that $(a_i, b_j) \in U \times V$ for all $i, j \in [m]$, then $p_1(a_i, b_j) < p_2(a_i, b_j)$ for every $i, j \in [m]$. Setting $g(x, y) := f(p_1(x, y), p_2(x, y))$, we obtain a function which is canonical with respect to \mathbb{H} on $U \times V$. Hence, the restriction of g to $U \times V$ naturally acts on $\{E, N\}$, and we may assume it does so as an essentially unary function as otherwise we are done. A similar statement holds for its restriction to $V \times U$. If one of the two mentioned essentially unary functions on $\{E, N\}$ is not a projection, then $\text{Aut}(\mathbb{A})$ contains a function flipping E and N . It follows that whenever $a, b \in S$ are so that $(a_i, b_j) \in U \times V$ for all $i, j \in \{1, \dots, k\}$, then $g(a, b)$ and $g(b, a)$ are elements of S . If a, b are moreover both increasing, then $g(a, b)$ and $f(a, b)$, as well as $g(b, a)$ and $f(b, a)$, belong to the same orbit under $\text{Aut}(\mathbb{A})$. Since $f(a, b) \sim f(b, a)$, the two essentially unary functions mentioned above cannot depend on the same argument, as witnessed by choosing $a, b \in S$ from distinct \sim -classes. Hence, we may assume that the one from $U \times V$ depends on the first argument and the other one on the second.

More generally, since $g(x, y) = f(p_1(x, y), p_2(x, y))$, the function g has the property that for all injective ℓ -tuples a, b where a is increasing, the type of their image under this function in \mathbb{H} only depends on the relations of \mathbb{H} on each of a and on b , respectively, and on the order relation on pairs (a_i, b_j) , where $i, j \in \{1, \dots, \ell\}$; this type is precisely the same type obtained when applying the function to a, b' , where b' is obtained from b by changing its order to be increasing.

In the following, for any pair (a, b) of injective increasing ℓ -tuples, we shall consider the set B of all pairs (a', b') of injective increasing ℓ -tuples such that the order relation on (a_i, b_j) and that on (a'_i, b'_j) agree for all $i, j \in \{1, \dots, \ell\}$; we call this set an *increasing diagonal order type*. We then have by the above that g acts naturally on $\{E, N\}$ within each such set B .

Set $h(x, y) := g(e_1 \circ g(x, y), e_2 \circ g(x, y))$, where e_1, e_2 are self-embeddings of $(\mathbb{H}, <)$ which ensure that for all injective increasing a, b , the diagonal order type of (a, b) is equal to that of $(e_1 \circ g(a, b), e_2 \circ g(a, b))$. Then h acts idempotently or as a constant function in its action on $\{E, N\}$ within each increasing diagonal order type B as above. In particular, h then acts like the first projection on $\{E, N\}$ when restricted to $U \times V$, and like the second when restricted to $V \times U$.

We will now rule out the possibility that h acts like a constant function on $\{E, N\}$ within some increasing diagonal order type of injective ℓ -tuples. In the following, we assign to every increasing diagonal order type B of pairs of increasing injective ℓ -tuples a number n_B as follows:

if $((x_1, \dots, x_\ell), (y_1, \dots, y_\ell)) \in B$, then n_B is the minimal number in $\{0, \dots, \ell\}$ such that for all $n_B < i \leq \ell$ it holds that $x_i < y_i$, and if $i > 1$, then moreover $y_{i-1} < x_i$.

Suppose that within some increasing diagonal order type B , we have that h acts as a constant function on $\{E, N\}$. Pick such B such that n_B is minimal. Assume without loss of generality that the constant value of h on B is E . We claim that \mathbb{A} has an endomorphism onto a clique, which contradicts the conjunction of the assumptions that \mathbb{A} is a model-complete core and not a reduct of $(H, =)$. We prove this claim by showing that any finite injective tuple can be mapped to a clique by an endomorphism of \mathbb{A} . Suppose that $m \geq 1$ and that (a_1, \dots, a_m) is an injective m -tuple of elements of H which does not induce a clique, i.e., there is a subtuple of length ℓ which is not an element of E . Then $m \geq \ell$, and we may assume without loss of generality that $(a_1, \dots, a_\ell) \notin E$. By applying a self-embedding e_1 of \mathbb{H} , we obtain an increasing tuple (x_1, \dots, x_m) such that $(x_1, \dots, x_\ell) \in N$. Applying an appropriate self-embedding e_2 of \mathbb{H} to (a_1, \dots, a_m) , we moreover obtain an increasing tuple (y_1, \dots, y_m) such that $((x_1, \dots, x_\ell), (y_1, \dots, y_\ell)) \in B$ and such that $y_{i-1} < x_i < y_i$ for all $i > \ell$. Then the increasing diagonal order type C for any pair $((x_{i_1}, \dots, x_{i_\ell}), (y_{i_1}, \dots, y_{i_\ell}))$ of subtuples, both increasing, has the property that $n_C \leq n_B$ (note that in order to compute n_C , the entries of both tuples receive the indices 1 to ℓ). If $n_C < n_B$, then h acts idempotently on $\{E, N\}$ within C , by the minimality of n_B ; if on the other hand $n_C = n_B$, then $B = C$ and h acts as a constant with value E . It follows that applying the endomorphism $h(e_1(x), e_2(x))$ to (a_1, \dots, a_m) , one obtains a tuple which has strictly more subtuples in E than (a_1, \dots, a_m) . The claim follows.

We may thus henceforth assume that within each increasing diagonal order type B , we have that h acts in an idempotent fashion on $\{E, N\}$. Thus, within each such type, h acts as a semilattice operation or as a projection on $\{E, N\}$.

Let $0 \leq j \leq \ell$, and consider the diagonal order type T given by a pair (c, d) of increasing injective j -tuples of elements of H ; we call j the *length* of T . In the following, we shall say that T is *categorical* if for all pairs (a, b) of increasing injective ℓ -tuples which extend (c, d) (we mean any extension, not just end-extension) the corresponding diagonal order type is one where h behaves like the first projection on, or if a similar statement holds for the second projection, or for the semilattice behaviour (both semilattice behaviours are considered the same here). Note that for length $j = 0$ every T is non-categorical, by the behaviours on $U \times V$ and $V \times U$. Note also that for length $j = \ell$ every T is trivially categorical. We claim that there exists T of length $j = \ell - 1$ which is not categorical. Suppose otherwise, and take any T which is categorical and implies the behaviour as the second projection; this exists by the behaviour on $V \times U$. Let (c, d) be a pair of injective j -tuples which are ordered such that they represent the diagonal order type T . Let (a_1, b_1) be obtained from (c, d) by extending both increasing tuples by a single element c' and d' at the end, respectively, in such a way that $c' < d'$. Let (c_1, d_1) be tuple obtained from (a_1, b_1) by taking away the first components. Then the order type represented by (c_1, d_1) is categorical but not for the first projection. We continue in this fashion until we arrive at a pair (a_ℓ, b_ℓ) in $U \times V$, a contradiction.

In the following, we assume that there exists T of length $j = \ell - 1$ which extends to diagonal order types where h behaves like different projections; the other case (projection + semilattice) is handled similarly. We show by induction on $m \geq 1$ that for all tuples $a, b \in I^m$ such that for every $i \in \{1, \dots, m\}$ at most one of the tuples a_i, b_i is in N there exists $u \in \text{Pol}(\mathbb{A})$ such that $u(a, b) \in E^m$. A standard compactness argument then implies that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ contains a function such that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is a semilattice operation.

The base case $m = 1$ is clearly achieved by applying an appropriate projection. For the induction step, let $a, b \in I^m$ for some $m \geq 2$. Since $\text{Pol}(\mathbb{A})$ contains p_1 , we may assume that the kernels of a and b are identical. We may then also assume that a_i, a_j induce distinct sets whenever $1 \leq i, j \leq m$ and $i \neq j$, for otherwise we are done by the induction hypothesis. By the induction hypothesis, we

may assume that all components of a are in E except for the second, and all components of b are in E except for the first. It is sufficient to show that there exists an increasing diagonal order type on the pair (a, b) (i.e., increasing diagonal order types for each of the pairs $(a_1, b_1), \dots, (a_m, b_m)$ which are consistent with the kernels of a and b) such that h behaves like the first projection within the order type of (a_1, b_1) and like the second within the order type of (a_2, b_2) . This, however, is obvious by our assumption. \square

8. BOUNDED WIDTH

In this section, we prove a characterization of first-order expansions of \mathbb{H} whose CSPs have bounded width.

Theorem 28. *Let \mathbb{A} be a first-order expansion of \mathbb{H} . Then precisely one of the following applies.*

- (1) *The clone $\text{Pol}(\mathbb{A})$ has a uniformly continuous minion homomorphism to the clone of affine maps over a finite module.*
- (2) *The clone $\text{Pol}(\mathbb{A})$ has no uniformly continuous minion homomorphism to the clone of affine maps over a finite module, and $\text{CSP}(\mathbb{A})$ has relational width $(2\ell, \max(3\ell, b_{\mathbb{H}}))$.*

We prove [Theorem 28](#) in a similar way as [Theorem 2](#). We prove that if \mathbb{A} is a first-order expansion of \mathbb{H} which is a model-complete core then $\text{CSP}(\mathbb{A})$ has bounded width if, and only if, $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine. If $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine, the result follows from [Corollary 15](#).

Let us therefore suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally affine. Moreover, we may assume that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial as otherwise, $\text{Pol}(\mathbb{A})$ has a uniformly continuous homomorphism to the clone of projections by the proof of [Theorem 2](#). We apply the second loop lemma of smooth approximations [[38](#), [Theorem 11](#)].

Theorem 29. *Let $k \geq 1$ and suppose that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-trivial. Let (S, \sim) be a minimal subfactor of $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ with $\text{Aut}(\mathbb{H})$ -invariant \sim -classes. Then one of the following holds:*

- *\sim is approximated by a $\text{Pol}(\mathbb{A})$ -invariant equivalence relation that is very smooth with respect to $\text{Aut}(\mathbb{H})$;*
- *every $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ -invariant binary symmetric relation $R \subseteq I^2$ that contains a pair $(a, b) \in S^2$ such that $a \neq b$ and such that $a \not\sim b$ contains a pseudo-loop modulo $\text{Aut}(\mathbb{H})$, i.e., a pair (c, c') where c, c' belong to the same orbit under $\text{Aut}(\mathbb{H})$.*

In the formulation of the first item of [Theorem 29](#), we are using [[38](#), [Lemma 8](#)]. If the first case of the theorem applies, i.e., the equivalence relation (S, \sim) on whose classes $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$ acts by a function from a clone \mathcal{M} of affine maps over a finite module is approximated by a $\text{Pol}(\mathbb{A})$ -invariant equivalence relation that is very smooth with respect to $\text{Aut}(\mathbb{H})$, [Lemma 24](#) (with $k = \ell$, $\mathcal{C}_{\mathbb{A}}^{\mathbb{A}, \text{inj}} = \mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}}$) implies that $\text{Pol}(\mathbb{A})$ has a uniformly continuous clone homomorphism to $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright (S, \sim)$ and hence to \mathcal{M} .

If the second case of [Theorem 29](#) applies, we get a weakly commutative function by [[38](#), [Lemma 13](#)], and the same argument mentioned at the end of [Section 3](#) gives that $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ contains a semilattice operation. In particular, $\mathcal{C}_{\mathbb{A}}^{\mathbb{H}, \text{inj}} \curvearrowright \{E, N\}$ is equationally non-affine, which is a contradiction.

REFERENCES

- [1] Reza Akhtar and Alistair H. Lachlan. On countable homogeneous 3-hypergraphs. *Archive for Mathematical Logic*, 34:331–344, 1995.

- [2] Libor Barto. Promises make finite (constraint satisfaction) problems infinitary. In *Proceedings of the 34th Annual ACM/IEEE Symposium on Logic in Computer Science – LICS’19*, 2019.
- [3] Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker. Equations in oligomorphic clones and the constraint satisfaction problem for ω -categorical structures. *Journal of Mathematical Logic*, 19(02):1950010, 2019.
- [4] Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker. The equivalence of two dichotomy conjectures for infinite domain constraint satisfaction problems. In *Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer Science – LICS’17*, 2017.
- [5] Libor Barto, Jakub Opršal, and Michael Pinsker. The wonderland of reflections. *Israel Journal of Mathematics*, 223(1):363–398, 2018.
- [6] Libor Barto and Michael Pinsker. Topology is irrelevant. *SIAM Journal on Computing*, 49(2):365–393, 2020.
- [7] Meghyn Bienvenu, Balder Ten Cate, Carsten Lutz, and Frank Wolter. Ontology-based data access: A study through disjunctive Datalog, CSP, and MMSNP. *ACM Transactions on Database Systems*, 39(4), 2015.
- [8] Manuel Bodirsky. Cores of countably categorical structures. *Logical Methods in Computer Science*, 3(1):1–16, 2007.
- [9] Manuel Bodirsky. Ramsey classes: Examples and constructions. In *Surveys in Combinatorics. London Mathematical Society Lecture Note Series 424*. Cambridge University Press, 2015. Invited survey article for the British Combinatorial Conference; preprint arXiv:1502.05146.
- [10] Manuel Bodirsky and Bertalan Bodor. Canonical polymorphisms of Ramsey structures and the unique interpolation property. In *Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science – LICS’21*, 2021.
- [11] Manuel Bodirsky and Martin Grohe. Non-dichotomies in constraint satisfaction complexity. In Luca Aceto, Ivan Damgård, Leslie Ann Goldberg, Magnús M. Halldórsson, Anna Ingólfssdóttir, and Igor Walukiewicz, editors, *Proceedings of the 35th International Colloquium on Automata, Languages and Programming, ICALP 2008*, pages 184–196, Berlin, Heidelberg, 2008. Springer-Verlag.
- [12] Manuel Bodirsky and Peter Jonsson. A model-theoretic view on qualitative constraint reasoning. *Journal of Artificial Intelligence Research*, 58(1):339–385, 2017.
- [13] Manuel Bodirsky, Peter Jonsson, Barnaby Martin, and Antoine Mottet. Classification transfer for qualitative reasoning problems. In *Proceedings of the Twenty-Seventh International Joint Conference on Artificial Intelligence, IJCAI-18*, pages 1256–1262. International Joint Conferences on Artificial Intelligence Organization, 2018.
- [14] Manuel Bodirsky, Peter Jonsson, and Trung Van Pham. The complexity of phylogeny constraint satisfaction problems. *ACM Transactions on Computational Logic*, 18(3), 2017. An extended abstract appeared in the conference STACS 2016.
- [15] Manuel Bodirsky and Jan Kára. The complexity of equality constraint languages. *Theory of Computing Systems*, 3(2):136–158, 2008. A conference version appeared in the proceedings of Computer Science Russia (CSR’06).
- [16] Manuel Bodirsky and Jan Kára. The complexity of temporal constraint satisfaction problems. *Journal of the ACM*, 57(2), 2010. An extended abstract appeared in the Proceedings of the Symposium on Theory of Computing (STOC).
- [17] Manuel Bodirsky and Simon Knäuer. Network satisfaction for symmetric relation algebras with a flexible atom. *Proceedings of the 35th AAAI Conference on Artificial Intelligence*, 35(7):6218–6226, 2021.
- [18] Manuel Bodirsky, Florent Madelaine, and Antoine Mottet. A proof of the algebraic tractability conjecture for monotone monadic SNP. *SIAM Journal on Computing*, 50(4):1359–1409, 2021.
- [19] Manuel Bodirsky, Florent R. Madelaine, and Antoine Mottet. A universal-algebraic proof of the complexity dichotomy for monotone monadic SNP. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science – LICS’18*, 2018.
- [20] Manuel Bodirsky, Barnaby Martin, Michael Pinsker, and András Pongrácz. Constraint satisfaction problems for reducts of homogeneous graphs. *SIAM Journal on Computing*, 48(4):1224–1264, 2019. A conference version appeared in the Proceedings of the 43rd International Colloquium on Automata, Languages, and Programming, ICALP 2016, pages 119:1–119:14.
- [21] Manuel Bodirsky and Antoine Mottet. Reducts of finitely bounded homogeneous structures, and lifting tractability from finite-domain constraint satisfaction. In *Proceedings of the 31th Annual ACM/IEEE Symposium on Logic in Computer Science – LICS’16*, 2016.
- [22] Manuel Bodirsky and Antoine Mottet. A dichotomy for first-order reducts of unary structures. *Logical Methods in Computer Science*, 14(2), 2018.
- [23] Manuel Bodirsky and Jaroslav Nešetřil. Constraint satisfaction with countable homogeneous templates. *Journal of Logic and Computation*, 16(3):359–373, 2006. A conference version appeared in the proceedings of Computer Science Logic (CSL 2003).
- [24] Manuel Bodirsky and Michael Pinsker. Schaefer’s theorem for graphs. *Journal of the ACM*, 62(3):19:1–19:52, 2015.

- [25] Manuel Bodirsky and Michael Pinsker. Canonical functions: a proof via topological dynamics. *Homogeneous Structures, A Workshop in Honour of Norbert Sauer's 70th Birthday, Contributions to Discrete Mathematics*, 16(2):36–45, 2021.
- [26] Manuel Bodirsky, Michael Pinsker, and András Pongrácz. Projective clone homomorphisms. *Journal of Symbolic Logic*, 86(1):148–161, 2021.
- [27] Manuel Bodirsky, Michael Pinsker, and Todor Tsankov. Decidability of definability. *Journal of Symbolic Logic*, 78(4):1036–1054, 2013. A conference version appeared in the Proceedings of the 26th Annual ACM/IEEE Symposium on Logic in Computer Science – LICS'11.
- [28] Bertalan Bodor. CSP dichotomy for ω -categorical monadically stable structures. PhD dissertation, Institute of Algebra, Technische Universität Dresden, 2021.
- [29] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In Chris Umans, editor, *58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017*, pages 319–330. IEEE Computer Society, 2017.
- [30] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: A study through Datalog and group theory. *SIAM Journal on Computing*, 28(1):57–104, 1998.
- [31] Pierre Gillibert, Julius Jonušas, Michael Kompatscher, Antoine Mottet, and Michael Pinsker. Hrushovski's encoding and ω -categorical CSP monsters. In Artur Czumaj, Anuj Dawar, and Emanuela Merelli, editors, *47th International Colloquium on Automata, Languages, and Programming, ICALP 2020, July 8-11, 2020, Saarbrücken, Germany (Virtual Conference)*, volume 168 of *LIPICs*, pages 131:1–131:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [32] Pierre Gillibert, Julius Jonušas, Michael Kompatscher, Antoine Mottet, and Michael Pinsker. When symmetries are not enough: a hierarchy of hard constraint satisfaction problems. *SIAM Journal on Computing*, 51(2):175–213, 2022.
- [33] Karim Khanaki. Dividing lines in unstable theories and subclasses of baire 1 functions. *Archive for Mathematical Logic*, 61(7-8):977–993, 2022.
- [34] Michael Kompatscher and Trung Van Pham. A complexity dichotomy for poset constraint satisfaction. *IfCoLog Journal of Logics and their Applications (FLAP)*, 5(8):1663–1696, 2018. A conference version appeared in the Proceedings of the 34th Symposium on Theoretical Aspects of Computer Science (STACS 2017), pages 47:1–47:12.
- [35] Marcin Kozik, Andrei Krokhin, Matt Valeriote, and Ross Willard. Characterizations of several Maltsev conditions. *Algebra universalis*, 73(3-4):205–224, 2015.
- [36] Miklós Maróti and Ralph McKenzie. Existence theorems for weakly symmetric operations. *Algebra Universalis*, 59(3):463–489, 2008.
- [37] Antoine Mottet and Michael Pinsker. Cores over Ramsey structures. *The Journal of Symbolic Logic*, 86(1):352–361, 2021.
- [38] Antoine Mottet and Michael Pinsker. Smooth approximations and CSPs over finitely bounded homogeneous structures. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science – LICS'22*, 2022.
- [39] Antoine Mottet, Tomáš Nagy, Michael Pinsker, and Michał Wrona. Smooth approximations and relational width collapses. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference)*, volume 198 of *LIPICs*, pages 138:1–138:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [40] Antoine Mottet, Tomáš Nagy, Michael Pinsker, and Michał Wrona. When symmetries are enough: collapsing the bounded width hierarchy for infinite-domain CSPs. arxiv:2102.07531, 2022.
- [41] Michael Pinsker. Current challenges in infinite-domain constraint satisfaction: Dilemmas of the infinite sheep. In *2022 IEEE 52nd International Symposium on Multiple-Valued Logic (ISMVL)*, pages 80–87, Los Alamitos, CA, USA, 2022. IEEE Computer Society.
- [42] Emil L. Post. The two-valued iterative systems of mathematical logic. *Annals of Mathematics Studies*, 5, 1941.
- [43] Jaroslav Nešetřil and Vojtěch Rödl. Ramsey classes of set systems. *Journal of Combinatorial Theory, Series A*, 34(2):183–201, 1983.
- [44] Simon Thomas. Reducts of random hypergraphs. *Annals of Pure and Applied Logic*, 80(2):165–193, 1996.
- [45] Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In Chris Umans, editor, *58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017*, pages 331–342. IEEE Computer Society, 2017.
- [46] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. *Journal of the ACM*, 67(5):30:1–30:78, 2020.
- [47] Dmitriy Zhuk. Strong subalgebras and the constraint satisfaction problem. *Journal of Multiple-Valued Logic and Soft Computing*, 36(4-5):455–504, 2021.

HAMBURG UNIVERSITY OF TECHNOLOGY, RESEARCH GROUP ON THEORETICAL COMPUTER SCIENCE, GERMANY

Email address: antoine.mottet@tuhh.de

URL: <http://amottet.github.io/~mottet/>

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, FG ALGEBRA, TU WIEN

Email address: tomas.nagy@email.com

URL: <http://dmg.tuwien.ac.at/nagy/>

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, FG ALGEBRA, TU WIEN

Email address: marula@gmx.at

URL: <http://dmg.tuwien.ac.at/pinsker/>