

REPRESENTATIONS AND IDENTITIES OF BAXTER MONOIDS WITH INVOLUTION

BIN BIN HAN¹, WEN TING ZHANG^{1,*}, YAN FENG LUO¹, AND JIN XING ZHAO²

ABSTRACT. Let $(\mathbf{baxt}_n, \sharp)$ be the Baxter monoid of finite rank n with Schützenberger’s involution \sharp . In this paper, it is shown that $(\mathbf{baxt}_n, \sharp)$ admits a faithful representation by an involution monoid of upper triangular matrices over any semiring from a large class including the tropical semiring under the skew transposition. Then a transparent combinatorial characterization of the word identities satisfied by $(\mathbf{baxt}_n, \sharp)$ is given. Further, it is proved that $(\mathbf{baxt}_n, \sharp)$ is finitely based if and only if $n \neq 3$, and shown that the identity checking problem for $(\mathbf{baxt}_n, \sharp)$ can be done in polynomial time.

1. INTRODUCTION

Identities and varieties of semigroups have long been studied, and several important questions arise in this area, such as the *finite basis problem*, that is the problem of classifying semigroups according to the finite basis property. The first example of non-finitely based finite semigroup was discovered by Perkins [49] in the 1960s; since then, the finite basis problem for semigroups has attracted much attention. Refer to Volkov [51] for a survey of work performed in this direction and for more information on the finite basis problem for semigroups in general. Other questions regarding the variety generated by a semigroup are those of whether it contains only finitely generated subvarieties (see, for example, [51]), or countably infinite subvarieties [28]. Another widely studied question is the *identity checking problem* $\text{CHECK-ID}(S)$ for a semigroup S , that is the decision problem whose instance is an arbitrary identity $\mathbf{u} \approx \mathbf{v}$, and the answer to such an instance is ‘YES’ if S satisfies $\mathbf{u} \approx \mathbf{v}$, and ‘NO’ if it does not. For a finite semigroup S , the identity checking problem $\text{CHECK-ID}(S)$ is always decidable but this is not necessarily true for an infinite semigroup [46]. Studying the computational complexity of identity checking in semigroups was proposed by Sapir [31, Problem 2.4], and up to now, there are many results in this study [6, 10, 16, 29, 33].

Recall that a unary operation $*$ on a semigroup S is an *involution* if S satisfies the identities

$$(x^*)^* \approx x \quad \text{and} \quad (xy)^* \approx y^*x^*. \quad (1.1)$$

An *involution semigroup* is a pair $(S, *)$ where S is a semigroup with involution $*$, and S is called the *semigroup reduct* of $(S, *)$. Common examples of involution semigroups include groups $(G, {}^{-1})$ with inversion ${}^{-1}$, multiplicative matrix semigroups $(M_n, {}^T)$ and $(M_n, {}^D)$ over any field with transposition T and skew transposition D respectively. Over the years, the identities and varieties of involution semigroups have received less attention than that for semigroups. Since the turn of the millennium, interest in involution semigroups has significantly increased. For example, many counterintuitive results were established and examples have been found

2010 *Mathematics Subject Classification.* 20M07, 20M30, 05E99, 12K10, 16Y60.

Key words and phrases. Baxter monoid, Schützenberger’s involution, representation, identity, finite basis problem.

The authors are partially supported by the National Natural Science Foundation of China (Nos. 12271224, 12171213, 12161062).

* Corresponding author.

to demonstrate that an involution semigroup $(S, *)$ and its semigroup reduct S need not be simultaneously finitely based [19, 20, 30, 35–39]. Refer to [1–3, 21, 40, 41] for more information on the identities and varieties of involution semigroups.

The plactic monoid, whose elements can be identified with Young tableaux, is famous for its interesting connection with combinatorics [34] and applications in symmetric functions [47], representation theory [17], Kostka-Foulkes polynomials [42, 43] and Schubert polynomials [44, 45]. Its finite-rank versions were shown to have faithful tropical representations by Johnson and Kambites [27, Theorem 2.8]. An important consequence of these representations, which are specifically representations using upper triangular tropical matrices, is that each finite-rank plactic monoid satisfies a non-trivial semigroup identity [27, Theorem 3.1]. In particular, the plactic monoids of rank equal to 2 and 3 are non-finitely based by [9, 16] and [24, 27] respectively, and their involution cases are also non-finitely based by using the result of [37, Theorem 4]. The finite basis problem for the plactic monoids of rank greater than 3 and their involution cases are still open. The plactic monoid of infinite rank is finitely based since it only satisfies trivial identity [12, Proposition 3.1].

Plactic monoids belong to a family of ‘plactic-like’ monoids which are connected with combinatorics and whose elements can be identified with combinatorial objects. Others in the family include the hypoplactic monoids, whose elements are quasi-ribbon tableaux and which play a role in the theory of quasi-symmetric functions analogous to that of the plactic monoid for symmetric functions [32, 48]; the sylvester and \sharp -sylvester monoids, whose elements are respectively right strict and left strict binary search trees [22]; the Baxter monoids, whose elements are pairs of twin binary search trees [18] and which are linked to the theory of Baxter permutations; the taiga monoids, whose elements are binary search trees with multiplicities [50]; the stalactic monoids, whose elements are stalactic tableaux [23, 50]; and the stylic monoids, whose elements are N -tableaux [4].

The hypoplactic, stalactic, taiga, sylvester and Baxter monoids of each finite rank were shown to have faithful tropical representations [11], and each of these monoids satisfy non-trivial identities [13]. It is shown that all Baxter [resp. hypoplactic, sylvester, stalactic and taiga] monoids of rank greater than or equal to 2 generate the same variety and are finitely based [11, 14, 15, 25]. Aird and Ribeiro have given a faithful representation of the stylic monoid and its involution case of each finite rank, and then solved the finite basis problems for them [5]. Also, Volkov has solved the finite basis problem for the stylic monoid by different mean [52]. The identity checking problems for the Baxter, sylvester and stylic monoids have been considered [5, 14], which can be done in polynomial time. By the characterization of the identities satisfied by the hypoplactic, stalactic and taiga monoids [11, 14, 25], it is easy to see that the identity checking problem for each of them is in the complexity class P.

The Baxter monoid \mathbf{baxt}_n of finite rank n , first studied by Giraudo [18], can be obtained by factoring the free monoid $\mathcal{A}_n^* = \mathcal{A}_n^+ \cup \{\varepsilon\}$ over the finite ordered alphabet $\mathcal{A}_n = \{1 < 2 < \dots < n\}$ by a congruence that can be defined by the insertion algorithm that computes a combinatorial object from a word. Each element of the Baxter monoid can be uniquely identified with a pair of twin binary trees, and it is presented by the relations defined for $\mathbf{u}, \mathbf{v} \in \mathcal{A}_n^*$ and $a, b, c, d \in \mathcal{A}_n$ by $\mathbf{cua}d\mathbf{v}b \equiv \mathbf{cud}a\mathbf{v}b$ with $a \leq b < c \leq d$ and $\mathbf{bud}a\mathbf{v}c \equiv \mathbf{buad}v\mathbf{c}$ with $a < b \leq c < d$. The Baxter monoid \mathbf{baxt}_n forms the involution monoid $(\mathbf{baxt}_n, \sharp)$ under the order-reversing permutation on \mathcal{A}_n .

In this paper, we investigate the representations and identities of finite-rank Baxter monoid with involution $(\mathbf{baxt}_n, \sharp)$. This paper is organized as follows.

Notation and background information of the paper are given in Section 2. In Section 3, we exhibit a faithful representation of $(\text{baxt}_n, \#)$ as an involution monoid of upper triangular matrices over any semiring from a large class including the tropical semiring under the skew transposition. In Section 4, a characterization of the word identities satisfied by $(\text{baxt}_n, \#)$ is given. In Section 5, we prove that $(\text{baxt}_n, \#)$ is finitely based if and only if $n \neq 3$ and that each variety generated by $(\text{baxt}_n, \#)$ with $n \geq 2$ contains continuum many subvarieties. And it is shown that the identity checking problem for $(\text{baxt}_n, \#)$ can be done in polynomial time in Section 6.

2. PRELIMINARIES

Most of the notation and definition of this article are given in this section. Refer to the monograph of Burris and Sankappanavar [7] for any undefined notation and terminology of universal algebra in general.

2.1. Words and content. Let \mathcal{X} be a countably infinite alphabet and $\mathcal{X}^* = \{x^* \mid x \in \mathcal{X}\}$ be a disjoint copy of \mathcal{X} . Elements of $\mathcal{X} \cup \mathcal{X}^*$ are called *variables*, elements of the free involution monoid $(\mathcal{X} \cup \mathcal{X}^*)^\times = (\mathcal{X} \cup \mathcal{X}^*)^+ \cup \{\emptyset\}$ are called *words*, and elements of $\mathcal{X}^+ \cup \{\emptyset\}$ are called *plain words*. A word \mathbf{u} is a *factor* of a word \mathbf{w} if $\mathbf{w} = \mathbf{aub}$ for some $\mathbf{a}, \mathbf{b} \in (\mathcal{X} \cup \mathcal{X}^*)^\times$.

Let $\mathbf{u} \in (\mathcal{X} \cup \mathcal{X}^*)^+$ be a word and $x \in \mathcal{X} \cup \mathcal{X}^*$ be a variable. The *content* $\text{con}(\mathbf{u})$ of a word \mathbf{u} is the set of variables that occur in \mathbf{u} , and $\text{occ}(x, \mathbf{u})$ is the number of occurrences of the letter x in \mathbf{u} . Let $\bar{\mathbf{u}}$ be the plain word obtained from \mathbf{u} by removing all occurrences of the symbol $*$. For any $x_1, x_2, \dots, x_n \in \mathcal{X} \cup \mathcal{X}^*$ such that $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_n \in \mathcal{X}$ are distinct variables, let $\mathbf{u}[x_1, x_2, \dots, x_n]$ denote the word obtained from \mathbf{u} by retaining only the variables $x_1, x_1^*, x_2, x_2^*, \dots, x_n, x_n^*$. Denote by $\overrightarrow{\text{occ}}_y(x, \mathbf{u})$ [resp. $\overleftarrow{\text{occ}}_y(x, \mathbf{u})$] the number of occurrences of x appearing before [resp. after] the first [resp. last] occurrence of y in \mathbf{u} . The *initial part* of \mathbf{u} , denoted by $\text{ip}(\mathbf{u})$, is the word obtained from \mathbf{u} by retaining the occurrence of each variable $x \in \text{con}(\mathbf{u})$ satisfying $x, x^* \notin \text{con}(\mathbf{u}_1)$ where $\mathbf{u} = \mathbf{u}_1 x \mathbf{u}_2$; the *final part* of \mathbf{u} , denoted by $\text{fp}(\mathbf{u})$, is the word obtained from \mathbf{u} by retaining the occurrence of each variable $x \in \text{con}(\mathbf{u})$ satisfying $x, x^* \notin \text{con}(\mathbf{u}_2)$ where $\mathbf{u} = \mathbf{u}_1 x \mathbf{u}_2$.

Example 2.1. Let $\mathbf{u} = x^* z x y^* x y z^2 x$ and $x, y, z \in \mathcal{X}$. Then

- $\text{con}(\mathbf{u}) = \{x, y, z, x^*, y^*\}$, $\bar{\mathbf{u}} = x z x y x y z^2 x$,
- $\text{occ}(x, \mathbf{u}) = 3$, $\text{occ}(x^*, \mathbf{u}) = \text{occ}(y, \mathbf{u}) = \text{occ}(y^*, \mathbf{u}) = 1$,
- $\mathbf{u}[x] = x^* x^3$, $\mathbf{u}[x, y] = x^* x y^* x y x$,
- $\overrightarrow{\text{occ}}_{y^*}(x, \mathbf{u}) = 1$, $\overleftarrow{\text{occ}}_{y^*}(x, \mathbf{u}) = 2$,
- $\text{ip}(\mathbf{u}) = x^* z y^*$, $\text{fp}(\mathbf{u}) = y z x$.

2.2. Terms and identities. The set $\text{T}(\mathcal{X})$ of *terms* over \mathcal{X} is the smallest set containing \mathcal{X} that is closed under concatenation and $*$. The proper inclusion $(\mathcal{X} \cup \mathcal{X}^*)^\times \subset \text{T}(\mathcal{X})$ holds and the identities (1.1) can be used to convert any nonempty term $\mathbf{t} \in \text{T}(\mathcal{X})$ into some unique word $[\mathbf{t}] \in (\mathcal{X} \cup \mathcal{X}^*)^+$. For instance, $[x(x^2(yx^*)^*)^*zy^*] = xy(x^*)^3zy^*$.

Remark 2.2. For any subterm \mathbf{s} of a term \mathbf{t} , either $[\mathbf{s}]$ or $[\mathbf{s}^*]$ is a factor of $[\mathbf{t}]$.

An *identity* is an expression $\mathbf{s} \approx \mathbf{t}$ formed by nonempty terms $\mathbf{s}, \mathbf{t} \in \text{T}(\mathcal{X})$, a *word identity* is an identity $\mathbf{u} \approx \mathbf{v}$ formed by words $\mathbf{u}, \mathbf{v} \in (\mathcal{X} \cup \mathcal{X}^*)^+$. We write $\mathbf{u} = \mathbf{v}$ if \mathbf{u} and \mathbf{v} are identical. An identity $\mathbf{u} \approx \mathbf{v}$ is *non-trivial* if $\mathbf{u} \neq \mathbf{v}$. An identity $\mathbf{s} \approx \mathbf{t}$ is directly deducible from an identity $\mathbf{p} \approx \mathbf{q}$ if there exists some substitution $\varphi : \mathcal{X} \rightarrow \text{T}(\mathcal{X})$ such that $\varphi(\mathbf{p})$ is a subterm of \mathbf{s} , and replacing this particular subterm $\varphi(\mathbf{p})$ of \mathbf{s} with $\varphi(\mathbf{q})$ results in the term \mathbf{t} . An identity $\mathbf{s} \approx \mathbf{t}$ is deducible from some set Σ of identities if there exists a sequence $\mathbf{s} = \mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_r = \mathbf{t}$ of

terms such that each identity $\mathbf{s}_i \approx \mathbf{s}_{i+1}$ is directly deducible from some identity in Σ .

Remark 2.3 ([8, Sublemma 2.2]). An identity $\mathbf{s} \approx \mathbf{t}$ is deducible from (1.1) if and only if $\lfloor \mathbf{s} \rfloor = \lfloor \mathbf{t} \rfloor$.

An involution semigroup $(S, *)$ satisfies an identity $\mathbf{s} \approx \mathbf{t}$, if for any substitution $\varphi : \mathcal{X} \rightarrow S$, the elements $\varphi(\mathbf{s})$ and $\varphi(\mathbf{t})$ of S coincide; in this case, $\mathbf{s} \approx \mathbf{t}$ is also said to be an *identity of* $(S, *)$.

Remark 2.4. Note that assigning the unit element to a variable x in a word identity is effectively the same as removing all occurrences of x and x^* . Therefore any involution monoid that satisfies a word identity $\mathbf{s} \approx \mathbf{t}$ also satisfies the word identity $\mathbf{s}[x_1, x_2, \dots, x_n] \approx \mathbf{t}[x_1, x_2, \dots, x_n]$ for any distinct variables $\overline{x_1}, \overline{x_2}, \dots, \overline{x_n} \in \mathcal{X}$.

For any involution semigroup $(S, *)$, a set Σ of identities of $(S, *)$ is an *identity basis* for $(S, *)$ if every identity satisfied by $(S, *)$ is deducible from Σ . An involution semigroup is *finitely based* if it has some finite identity basis; otherwise, it is *non-finitely based*.


The variety generated by semigroup S [resp. involution semigroup $(S, *)$] is denoted by $\text{Var}S$ [resp. $\text{Var}(S, *)$]. For any set Σ of identities, denote by $\text{Var}\Sigma$ the variety determined by Σ .

2.3. The Baxter monoid and its involution. Let $\mathcal{A} = \{1 < 2 < 3 < \dots\}$ denote the set of positive integers, viewed as an infinite ordered alphabet. The combinatorial objects and insertion algorithm related to the Baxter monoid are given in the following.

A *right strict binary search tree* is a labelled rooted binary tree where the label of each node is greater than or equal to the label of every node in its left subtree, and strictly less than every node in its right subtree. The associated insertion algorithm is as follows:

Algorithm 1. Input a right strict binary search tree T and a symbol $a \in \mathcal{A}$. If T is empty, create a node and label it a . If T is non-empty, examine the label x of the root node: if $a > x$, recursively insert a into the right subtree of the root node; otherwise recursively insert a into the left subtree of the root node. Output the resulting tree.

Let $w_1, \dots, w_k \in \mathcal{A}$ and $\mathbf{w} = w_1 \dots w_k \in \mathcal{A}^*$. Then the combinatorial object $P_{\text{sylv}_\infty}(\mathbf{w})$ of \mathbf{w} is obtained as follows: reading \mathbf{w} from right-to-left, one starts with an empty tree and inserts each symbol in \mathbf{w} into a right strict binary search tree according to Algorithm 1. For example, $P_{\text{sylv}_\infty}(36131512665)$ is given as follows:




A *left strict binary search tree* is a labelled rooted binary tree where the label of each node is strictly greater than the label of every node in its left subtree, and less than or equal to every node in its right subtree. The associated insertion algorithm is as follows:

Algorithm 2. Input a left strict binary search tree T and a symbol $a \in \mathcal{A}$. If T is empty, create a node and label it a . If T is non-empty, examine the label x of the root node: if $a < x$, recursively insert a into the left subtree of the root node;

otherwise recursively insert a into the right subtree of the root node. Output the resulting tree.

Let $w_1, \dots, w_k \in \mathcal{A}$ and $\mathbf{w} = w_1 \cdots w_k \in \mathcal{A}^*$. Then the combinatorial object $P_{\text{sylv}_\infty^\#}(\mathbf{w})$ of \mathbf{w} is obtained as follows: reading \mathbf{w} from left-to-right, one starts with an empty tree and inserts each symbol in \mathbf{w} into a left strict binary search tree according to Algorithm 2. For example, $P_{\text{sylv}_\infty^\#}(36131512665)$ is given as follows:



Let $w_1, \dots, w_k \in \mathcal{A}$ and $\mathbf{w} = w_1 \cdots w_k \in \mathcal{A}^*$. Then the combinatorial object $P_{\text{baxt}_\infty}(\mathbf{w})$ of \mathbf{w} is obtained by Algorithms 1 and 2, that is, $P_{\text{baxt}_\infty}(\mathbf{w}) = (P_{\text{sylv}_\infty^\#}(\mathbf{w}), P_{\text{sylv}_\infty}(\mathbf{w}))$.

Define the relation $\equiv_{\text{baxt}_\infty}$ by

$$\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v} \quad \text{if and only if} \quad P_{\text{baxt}_\infty}(\mathbf{u}) = P_{\text{baxt}_\infty}(\mathbf{v})$$

for any $\mathbf{u}, \mathbf{v} \in \mathcal{A}^*$. The relation $\equiv_{\text{baxt}_\infty}$ is a congruence on \mathcal{A}^* . The Baxter monoid baxt_∞ is the factor monoid $\mathcal{A}^*/\equiv_{\text{baxt}_\infty}$. The rank- n analogue baxt_n is the factor monoid $\mathcal{A}_n^*/\equiv_{\text{baxt}_\infty}$, where the relation $\equiv_{\text{baxt}_\infty}$ is naturally restricted to $\mathcal{A}_n^* \times \mathcal{A}_n^*$ and $\mathcal{A}_n = \{1 < 2 < \dots < n\}$ is the set of the first n natural numbers viewed as a finite ordered alphabet. It follows from the definition of $\equiv_{\text{baxt}_\infty}$ that each element $[\mathbf{u}]_{\equiv_{\text{baxt}_\infty}}$ of the factor monoid baxt_∞ can be identified with the combinatorial object $P_{\text{baxt}_\infty}(\mathbf{u})$. Clearly baxt_1 is a free monogenic monoid $\langle 1 \rangle = \{\varepsilon, 1, 1^2, 1^3, \dots\}$ and so it is commutative. Note that

$$\text{baxt}_1 \subset \text{baxt}_2 \subset \dots \subset \text{baxt}_i \subset \text{baxt}_{i+1} \subset \dots \subset \text{baxt}_\infty. \quad (2.1)$$

For any word $\mathbf{u} \in \mathcal{A}^*$, the *length* $|\mathbf{u}|$ of \mathbf{u} is the number of symbols occurring in \mathbf{u} , and $|\mathbf{u}|_a$ is the number of times the symbol a appearing in \mathbf{u} ; the *evaluation* of \mathbf{u} , denoted by $\text{ev}(\mathbf{u})$, is the infinite tuple of non-negative integers, indexed by \mathcal{A} , whose a -th element is $|\mathbf{u}|_a$, thus this tuple describes the number of each symbol in \mathcal{A} that appears in \mathbf{u} . It is immediate from the definition of the Baxter monoid that if $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$, then $\text{ev}(\mathbf{u}) = \text{ev}(\mathbf{v})$, and hence it makes sense to define the evaluation of each element of Baxter monoid to be the evaluation of any word representing it. The *support* of a word $\mathbf{u} \in \mathcal{A}^*$, denoted by $\text{sup}(\mathbf{u})$, is the set of letters that occur in \mathbf{u} . Note that $\text{ev}(\mathbf{u}) = \text{ev}(\mathbf{v})$ implies that $\text{sup}(\mathbf{u}) = \text{sup}(\mathbf{v})$.

Let $\mathbf{u} \in \mathcal{A}^*$ and $a, b \in \text{sup}(\mathbf{u})$ with $a < b$. We say that \mathbf{u} has a *b - a right precedence of index r* for some $r \geq 1$ if, when reading \mathbf{u} from right to left, b occurs r times before the first occurrence of a and, for any $c \in \text{sup}(\mathbf{u})$ such that $a < c < b$, c does not occur before the first occurrence of a . We say that \mathbf{u} has a *a - b left precedence of index ℓ* if, when reading \mathbf{u} from left to right, a occurs ℓ times before the first occurrence of b and, for any $c \in \text{sup}(\mathbf{u})$ such that $a < c < b$, c does not occur before the first occurrence of b . Note that, for any given $a \in \text{sup}(\mathbf{u})$, there is at most one $b \in \text{sup}(\mathbf{u})$ such that \mathbf{u} has a b - a right precedence of index r ; on the other hand, \mathbf{u} can have several right precedences of the form b - x for a fixed b ; and that, for any given $b \in \text{sup}(\mathbf{u})$, there is at most one $a \in \text{sup}(\mathbf{u})$ such that \mathbf{u} has a a - b left precedence of index ℓ ; on the other hand, \mathbf{u} can have several left precedences of the form a - x for a fixed a . Denote by

$$\text{rpi}(\mathbf{u}) = \{(b, r) : \mathbf{u} \text{ has a } b\text{-}a \text{ right precedence of index } r\}$$

and

$$\text{lpi}(\mathbf{u}) = \{(a-b, \ell) : \mathbf{u} \text{ has a } a\text{-}b \text{ left precedence of index } \ell\}.$$

For example, $\text{rpi}(36131512665) = \{(2-1, 1), (5-2, 1), (5-3, 2)\}$ and $\text{lpi}(36131512665) = \{(1-2, 3), (3-5, 2), (3-6, 1)\}$.

Proposition 2.5 ([14, Corollary 2.11]). *For any $\mathbf{u}, \mathbf{v} \in \mathcal{A}^*$, $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$ if and only if $\text{ev}(\mathbf{u}) = \text{ev}(\mathbf{v})$, $\text{rpi}(\mathbf{u}) = \text{rpi}(\mathbf{v})$ and $\text{lpi}(\mathbf{u}) = \text{lpi}(\mathbf{v})$.*

Note that if the word \mathbf{u} has a b - a right precedence or a a - b left precedence for $a < b$, then each word in $[\mathbf{u}]_{\text{baxt}_\infty}$ has a b - a right precedence or a a - b left precedence.

The Baxter monoid can also be defined by the presentation $\langle \mathcal{A} \mid \mathcal{R}_{\text{baxt}_\infty} \rangle$, where

$$\begin{aligned} \mathcal{R}_{\text{baxt}_\infty} = & \{(cuadvb, cudavb) : a \leq b < c \leq d\} \\ & \cup \{(budavc, buadvc) : a < b \leq c < d\}. \end{aligned}$$

For each $n \in \mathbb{N}$, a presentation for the Baxter monoid of rank n can be obtained by restricting generators and relations of the above presentation to generators in \mathcal{A}_n . Note that these relations are length-preserving.

If the Baxter monoid baxt_n under the unary operation $*$ is an involution monoid, then the relation $\equiv_{\text{baxt}_\infty}$ must be compatible with the involution operation, that is, if $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$, then $\mathbf{u}^* \equiv_{\text{baxt}_\infty} \mathbf{v}^*$.

Let $\mathcal{A}_n^\sharp := \{1^\sharp > 2^\sharp > \dots > n^\sharp\}$ be the alphabet \mathcal{A}_n on which the order relation has been reversed and $(\mathcal{A}_n^\sharp)^\sharp := \mathcal{A}_n$. For $w_1 w_2 \dots w_n \in \mathcal{A}^*$, $(w_1 w_2 \dots w_n)^\sharp := w_n^\sharp \dots w_2^\sharp w_1^\sharp$.

Proposition 2.6 ([18, Proposition 3.4]). *Let \mathbf{w} and \mathbf{w}' be two words in \mathcal{A}_n^* . Then $\mathbf{w} \equiv_{\text{baxt}_\infty} \mathbf{w}'$ if and only if $\mathbf{w}^\sharp \equiv_{\text{baxt}_\infty} (\mathbf{w}')^\sharp$.*

The relation $\equiv_{\text{baxt}_\infty}$ is compatible with the unary operation $^\sharp$. Thus $(\text{baxt}_n, ^\sharp)$ is an involution monoid, and the involution $^\sharp$ is always called *Schützenberger's involution*.

Proposition 2.7. *Schützenberger's involution is the unique involution on the Baxter monoid baxt_n for each $n \geq 1$.*

Proof. Suppose that $*$ is an involution operation on baxt_n . Note that the relations $\mathcal{R}_{\text{hypo}_\infty}$ are length-preserving. Then the involution of a generator in \mathcal{A}_n is still a generator in \mathcal{A}_n . Since baxt_1 has only one generator 1, we have $1^* = 1$. Thus the involution on baxt_1 is trivial. For baxt_n with $n \geq 2$, let $a < b \leq n$. Then $(buabva)^* = a^* \mathbf{v}^* b^* a^* \mathbf{u}^* b^* \equiv_{\text{baxt}_\infty} a^* \mathbf{v}^* a^* b^* \mathbf{u}^* b^* = (bubava)^*$ by $buabva \equiv_{\text{baxt}_\infty} bubava \in \mathcal{R}_{\text{hypo}_\infty}$. This implies $a^* \mathbf{v}^* b^* a^* \mathbf{u}^* b^* \equiv_{\text{baxt}_\infty} a^* \mathbf{v}^* a^* b^* \mathbf{u}^* b^* \in \mathcal{R}_{\text{baxt}_\infty}$, and so $b^* < a^*$. Hence for any $a < b$, we must have $b^* < a^*$, whence $*$ must be the order-reversing permutation on \mathcal{A}_n . Therefore $*$ is Schützenberger's involution, and so Schützenberger's involution is the unique involution on the Baxter monoid baxt_n . \square

2.4. Matrix representations over semirings. Recall that $\mathbb{S} = (S, +, \cdot)$ is a *commutative semiring* with additive identity $\mathbf{0}$ and multiplicative identity $\mathbf{1}$ if S is a set equipped with two binary operations $+$ and \cdot such that $(S, +)$ and (S, \cdot) are commutative monoids satisfying

$$a(b+c) = a \cdot b + a \cdot c \quad \text{and} \quad \mathbf{0} \cdot a = \mathbf{0}$$

for all $a, b, c \in \mathbb{S}$. Semiring \mathbb{S} is *idempotent* if $a + a = a$ for all $a \in \mathbb{S}$. An element $a \in \mathbb{S}$ is *infinite multiplicative order* if for any non-negative integers i, j , $a^i = a^j$ if and only if $i = j$. In this paper, we always assume that \mathbb{S} is a commutative and idempotent semiring with $\mathbf{0}, \mathbf{1}$ containing an element of infinite multiplicative order. A common example of such semiring is the tropical semiring $\mathbb{T} = (\mathbb{R} \cup \{-\infty\}, \oplus, \otimes)$,

which is the set \mathbb{R} of real numbers together with minus infinity $-\infty$, with the addition and multiplication defined as follows

$$a \oplus b = \max\{a, b\} \quad \text{and} \quad a \otimes b = a + b.$$

In other words, the tropical sum of two numbers is their maximum and the tropical product of two numbers is their sum, and $-\infty$ is the additive identity and 0 is the multiplicative identity. Note that except $-\infty$ and 0, all other elements in \mathbb{T} have infinite multiplicative order.

Note that the set of all $n \times n$ matrices with entries in \mathbb{S} forms a semigroup under the matrix multiplication induced from the operations in \mathbb{S} . We denote this semigroup by $M_n(\mathbb{S})$. Let $UT_n(\mathbb{S})$ be the subsemigroup of $M_n(\mathbb{S})$ of all upper triangular $n \times n$ matrices. For any matrix $A \in M_n(\mathbb{S})$, denote by A^D the matrix obtained by reflecting A with respect to the secondary diagonal (from the top right to the bottom left corner), that is, $(A^D)_{ij} = A_{(n+1-j)(n+1-i)}$. It is easy to verify that this unary operation D is an involution operation of $M_n(\mathbb{S})$. A (linear) representation of a semigroup S [resp. involution semigroup $(S, *)$] is a homomorphism $\rho : S \rightarrow M_n(\mathbb{S})$ [resp. $\rho : (S, *) \rightarrow (M_n(\mathbb{S}), ^D)$]. The homomorphism ρ is said to be *faithful* if it is injective. Note that an involution semigroup representation $\rho : (S, *) \rightarrow (M_n(\mathbb{S}), ^D)$ deduces a semigroup representation $\rho : S \rightarrow M_n(\mathbb{S})$, but a semigroup representation $\rho : S \rightarrow M_n(\mathbb{S})$ can not be an involution semigroup representation $\rho : (S, *) \rightarrow (M_n(\mathbb{S}), ^D)$. The tropical semiring is of interest as a natural carrier for representations of semigroups. For example, the bicyclic monoid $\mathcal{B} := \langle a, b \mid ba = 1 \rangle$, which is ubiquitous in infinite semigroup theory, admits no faithful finite dimensional representations over any field; however it has a number of natural representations over the tropical semiring [16, 26].

3. MATRIX REPRESENTATIONS OF $(\text{baxt}_n, \#)$

In this section, we exhibit a faithful representation of $(\text{baxt}_n, \#)$ for each finite n as an involution monoid of upper triangular matrices over \mathbb{S} under the skew transposition, and we prove that all involution semigroups $(\text{baxt}_n, \#)$ with $n \geq 4$ generate the same variety.

For convenience, denote by

$$P = \begin{pmatrix} s & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}, Q = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & s \end{pmatrix}, J = \begin{pmatrix} \mathbf{1} & \mathbf{1} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}, K = \begin{pmatrix} \mathbf{0} & \mathbf{1} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}$$

where $s \in \mathbb{S}$ is an element of infinite multiplicative order. Denote by $\text{diag}\{\Lambda_1, \Lambda_2, \dots, \Lambda_n\}$ the block diagonal matrix

$$\begin{pmatrix} \Lambda_1 & & & \\ & \Lambda_2 & & \\ & & \ddots & \\ & & & \Lambda_n \end{pmatrix}$$

where $\Lambda_1, \Lambda_2, \dots, \Lambda_n$ are square matrices. And let E_n be the $n \times n$ matrix with $\mathbf{1}$ s on the main diagonal and $\mathbf{0}$ s elsewhere.

First we give a matrix representation of $(\text{baxt}_1, \#)$. Define a map $\varphi_1 : \mathcal{A}_1 \cup \{\varepsilon\} \rightarrow UT_2(\mathbb{S})$ given by $\varepsilon \mapsto E_2$ and $1 \mapsto PQ$. Clearly, the map φ_1 induces a faithful representation of $(\text{baxt}_1, \#)$.

Next we consider the matrix representation of $(\text{baxt}_2, \#)$. Define a map $\varphi_2 : \mathcal{A}_2 \cup \{\varepsilon\} \rightarrow UT_6(\mathbb{S})$ given by $\varepsilon \mapsto E_6$,

$$1 \mapsto \text{diag}\{s, P, J, \mathbf{1}\}, \quad 2 \mapsto \text{diag}\{\mathbf{1}, K, Q, s\}.$$

Clearly, φ_2 can be extended to a homomorphism from \mathcal{A}_2^* to $UT_6(\mathbb{S})$. Note that $\varphi_2(1^\#) = \varphi_2(2) = \text{diag}\{\mathbf{1}, K, Q, s\} = (\varphi_2(1))^D$ and $\varphi_2(2^\#) = \varphi_2(1) =$

$\text{diag}\{s, P, J, \mathbf{1}\} = (\varphi_2(2))^D$. Thus φ_2 can be extended to a homomorphism from $(\mathcal{A}_2^*, \#)$ to $(UT_6(\mathbb{S}), {}^D)$. In fact, φ_2 induces a faithful representation of $(\text{baxt}_2, \#)$.

Theorem 3.1. *The map $\varphi_2 : (\text{baxt}_2, \#) \rightarrow (UT_6(\mathbb{S}), {}^D)$ is a faithful representation of $(\text{baxt}_2, \#)$.*

Proof. Note that φ_2 is a homomorphism from $(\mathcal{A}_2^*, \#)$ to $(UT_6(\mathbb{S}), {}^D)$. Then to show that the map φ_2 induces a homomorphism from $(\text{baxt}_2, \#)$ to $(UT_6(\mathbb{S}), {}^D)$, we only need to show that for any $\mathbf{u}, \mathbf{v} \in \mathcal{A}_2^*$, if $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$, then $\varphi_2(\mathbf{u}) = \varphi_2(\mathbf{v})$. By the definition of φ_2 , it is easy to verify that for any $\mathbf{w} \in \mathcal{A}_2^*$,

$$\varphi_2(\mathbf{w}) = \text{diag}\{\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4\}$$

where

$$\Lambda_1 = \begin{cases} s^{|\mathbf{w}|_1}, & \text{if } \{1\} \in \text{sup}(\mathbf{w}), \\ \mathbf{1}, & \text{if } \{1\} \notin \text{sup}(\mathbf{w}), \end{cases}$$

$$\Lambda_2 = \begin{cases} P^{|\mathbf{w}|_1}, & \text{if } \text{sup}(\mathbf{w}) = \{1\}, \\ K, & \text{if } \{2\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (1-2, \ell) \notin \text{lpi}(\mathbf{w}) \text{ for any } \ell > 0, \\ P^{\ell_1} K, & \text{if } \text{sup}(\mathbf{w}) = \{1, 2\} \text{ and } (1-2, \ell_1) \in \text{lpi}(\mathbf{w}), \end{cases}$$

$$\Lambda_3 = \begin{cases} Q^{|\mathbf{w}|_2}, & \text{if } \text{sup}(\mathbf{w}) = \{2\}, \\ J, & \text{if } \{1\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (2-1, r) \notin \text{rpi}(\mathbf{w}) \text{ for any } r > 0, \\ JQ^{r_1}, & \text{if } \text{sup}(\mathbf{w}) = \{1, 2\} \text{ and } (2-1, r_1) \in \text{rpi}(\mathbf{w}), \end{cases}$$

$$\Lambda_4 = \begin{cases} s^{|\mathbf{w}|_2}, & \text{if } \{2\} \in \text{sup}(\mathbf{w}), \\ \mathbf{1}, & \text{if } \{2\} \notin \text{sup}(\mathbf{w}). \end{cases}$$

Since $\text{ev}(\mathbf{u}) = \text{ev}(\mathbf{v})$, $\text{lpi}(\mathbf{u}) = \text{lpi}(\mathbf{v})$, $\text{rpi}(\mathbf{u}) = \text{rpi}(\mathbf{v})$ by Proposition 2.5, it is routine to show that $\varphi_2(\mathbf{u}) = \varphi_2(\mathbf{v})$.

Suppose that $\mathbf{u} \not\equiv_{\text{baxt}_\infty} \mathbf{v}$. Then $\text{ev}(\mathbf{u}) \neq \text{ev}(\mathbf{v})$, or $\text{lpi}(\mathbf{u}) \neq \text{lpi}(\mathbf{v})$, or $\text{rpi}(\mathbf{u}) \neq \text{rpi}(\mathbf{v})$ by Proposition 2.5. By the definition of φ_2 , it is routine to show that $\varphi_2(\mathbf{u}) \neq \varphi_2(\mathbf{v})$. Hence φ_2 is injective. Therefore the map $\varphi_2 : (\text{baxt}_2, \#) \rightarrow (UT_6(\mathbb{S}), {}^D)$ is a faithful representation of $(\text{baxt}_2, \#)$. \square

Next we consider the matrix representation of $(\text{baxt}_3, \#)$. Define a map $\varphi_3 : \mathcal{A}_3 \cup \{\varepsilon\} \rightarrow UT_{15}(\mathbb{S})$ given by $\varepsilon \mapsto E_{15}$,

$$1 \mapsto \text{diag}\{s, P, P, E_2, \mathbf{1}, J, E_2, J, \mathbf{1}\},$$

$$2 \mapsto \text{diag}\{\mathbf{1}, K, K, P, s, Q, J, J, \mathbf{1}\},$$

$$3 \mapsto \text{diag}\{\mathbf{1}, K, E_2, K, \mathbf{1}, E_2, Q, Q, s\}.$$

Clearly, φ_3 can be extended to a homomorphism from \mathcal{A}_3^* to $UT_{15}(\mathbb{S})$. Note that $\varphi_3(1^\#) = \varphi_3(3) = (\varphi_3(1))^D$, $\varphi_3(2^\#) = \varphi_3(2) = (\varphi_3(2))^D$ and $\varphi_3(3^\#) = \varphi_3(1) = (\varphi_3(1))^D$. Thus φ_3 can be extended to a homomorphism from $(\mathcal{A}_3^*, \#)$ to $(UT_{15}(\mathbb{S}), {}^D)$. In fact, φ_3 induces a faithful representation of $(\text{baxt}_3, \#)$.

Theorem 3.2. *The map $\varphi_3 : (\text{baxt}_3, \#) \rightarrow (UT_{15}(\mathbb{S}), {}^D)$ is a faithful representation of $(\text{baxt}_3, \#)$.*

Proof. Note that φ_3 is a homomorphism from $(\mathcal{A}_3^*, \#)$ to $(UT_{15}(\mathbb{S}), {}^D)$. Then to show that the map φ_3 induces a homomorphism from $(\text{baxt}_3, \#)$ to $(UT_{15}(\mathbb{S}), {}^D)$, we only need to show that for any $\mathbf{u}, \mathbf{v} \in \mathcal{A}_3^*$, if $\mathbf{u} \equiv_{\text{hyp}_\infty} \mathbf{v}$, then $\varphi_3(\mathbf{u}) = \varphi_3(\mathbf{v})$. By the definition of φ_3 , it is easy to verify that for any $\mathbf{w} \in \mathcal{A}_3^*$,

$$\varphi_3(\mathbf{w}) = \text{diag}\{\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5, \Lambda_6, \Lambda_7, \Lambda_8, \Lambda_9\}$$

where

$$\begin{aligned}
 \Lambda_1 &= \begin{cases} s^{|\mathbf{w}|_1}, & \text{if } \{1\} \in \text{sup}(\mathbf{w}), \\ \mathbf{1}, & \text{if } \{1\} \notin \text{sup}(\mathbf{w}), \end{cases} \\
 \Lambda_2 &= \begin{cases} P^{|\mathbf{w}|_1}, & \text{if } \text{sup}(\mathbf{w}) = \{1\}, \\ P^{\ell_1}K, & \text{if } \{1, 2\} = \text{sup}(\mathbf{w}) \text{ and } (1-2, \ell_1) \in \text{lpi}(\mathbf{w}), \\ P^{\ell_2}K, & \text{if } \{1, 3\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (1-3, \ell_2) \in \text{lpi}(\mathbf{w}), \\ P^{\ell_1}K, & \text{if } \{1, 2, 3\} = \text{sup}(\mathbf{w}) \text{ and } (1-2, \ell_1) \in \text{lpi}(\mathbf{w}), (2-3, \ell_3) \in \text{lpi}(\mathbf{w}), \\ K, & \text{otherwise,} \end{cases} \\
 \Lambda_3 &= \begin{cases} P^{|\mathbf{w}|_1}, & \text{if } \text{sup}(\mathbf{w}) = \{1\} \text{ or } \{1, 3\}, \\ E_2, & \text{if } \text{sup}(\mathbf{w}) = \{3\}, \\ P^{\ell_1}K, & \text{if } \{1, 2\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (1-2, \ell_1) \in \text{lpi}(\mathbf{w}), \\ K, & \text{otherwise,} \end{cases} \\
 \Lambda_4 &= \begin{cases} E_2, & \text{if } \text{sup}(\mathbf{w}) = \{1\}, \\ P^{|\mathbf{w}|_2}, & \text{if } \text{sup}(\mathbf{w}) = \{2\} \text{ or } \{1, 2\}, \\ P^{\ell_3}K, & \text{if } \{2, 3\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (2-3, \ell_3) \in \text{lpi}(\mathbf{w}), \\ K, & \text{otherwise,} \end{cases} \\
 \Lambda_5 &= \begin{cases} s^{|\mathbf{w}|_2}, & \text{if } \{2\} \in \text{sup}(\mathbf{w}), \\ \mathbf{1}, & \text{if } \{2\} \notin \text{sup}(\mathbf{w}), \end{cases} \\
 \Lambda_6 &= \begin{cases} Q^{|\mathbf{w}|_2}, & \text{if } \text{sup}(\mathbf{w}) = \{2\} \text{ or } \{2, 3\}, \\ E_2, & \text{if } \text{sup}(\mathbf{w}) = \{3\}, \\ JQ^{r_1}, & \text{if } \{1, 2\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (2-1, r_1) \in \text{rpi}(\mathbf{w}), \\ J, & \text{otherwise,} \end{cases} \\
 \Lambda_7 &= \begin{cases} E_2, & \text{if } \text{sup}(\mathbf{w}) = \{1\}, \\ Q^{|\mathbf{w}|_3}, & \text{if } \text{sup}(\mathbf{w}) = \{3\} \text{ or } \{1, 3\}, \\ JQ^{r_3}, & \text{if } \{2, 3\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (3-2, r_3) \in \text{rpi}(\mathbf{w}), \\ J, & \text{otherwise,} \end{cases} \\
 \Lambda_8 &= \begin{cases} Q^{|\mathbf{w}|_3}, & \text{if } \text{sup}(\mathbf{w}) = \{3\}, \\ JQ^{r_2}, & \text{if } \{1, 3\} \subseteq \text{sup}(\mathbf{w}) \text{ and } (3-1, r_2) \in \text{rpi}(\mathbf{w}), \\ JQ^{r_3}, & \text{if } \{2, 3\} = \text{sup}(\mathbf{w}) \text{ and } (3-2, r_3) \in \text{rpi}(\mathbf{w}), \\ JQ^{r_3}, & \text{if } \{1, 2, 3\} = \text{sup}(\mathbf{w}) \text{ and } (2-1, r_1) \in \text{rpi}(\mathbf{w}), (3-2, r_3) \in \text{rpi}(\mathbf{w}), \\ J, & \text{otherwise,} \end{cases} \\
 \Lambda_9 &= \begin{cases} s^{|\mathbf{w}|_3}, & \text{if } \{3\} \in \text{sup}(\mathbf{w}), \\ \mathbf{1}, & \text{if } \{3\} \notin \text{sup}(\mathbf{w}). \end{cases}
 \end{aligned}$$

Since $\text{ev}(\mathbf{u}) = \text{ev}(\mathbf{v})$, $\text{lpi}(\mathbf{u}) = \text{lpi}(\mathbf{v})$, $\text{rpi}(\mathbf{u}) = \text{rpi}(\mathbf{v})$ by Proposition 2.5, it is routine to show that $\varphi_3(\mathbf{u}) = \varphi_3(\mathbf{v})$.

Suppose that $\mathbf{u} \not\equiv_{\text{baxt}_\infty} \mathbf{v}$. Then $\text{ev}(\mathbf{u}) \neq \text{ev}(\mathbf{v})$, or $\text{rpi}(\mathbf{u}) \neq \text{rpi}(\mathbf{v})$, or $\text{lpi}(\mathbf{u}) \neq \text{lpi}(\mathbf{v})$ by Proposition 2.5. By the definition of φ_3 , it is routine to show that $\varphi_3(\mathbf{u}) \neq \varphi_3(\mathbf{v})$. Hence φ_3 is injective. Therefore the map $\varphi_3 : (\text{baxt}_3, \#) \rightarrow (UT_{15}(\mathbb{S}), \#)$ is a faithful representation of $(\text{baxt}_3, \#)$. \square

Now we consider the matrix representation of $(\text{baxt}_n, \#)$ for $n \geq 4$. For any $([\mathbf{u}]_{\text{baxt}_3}, [\mathbf{v}]_{\text{baxt}_3}) \in \text{baxt}_3 \times \text{baxt}_3$, define an involution operation $\#$ on $\text{baxt}_3 \times \text{baxt}_3$ by $([\mathbf{u}]_{\text{baxt}_3}, [\mathbf{v}]_{\text{baxt}_3})^\# = ([\mathbf{v}]_{\text{baxt}_3}^\#, [\mathbf{u}]_{\text{baxt}_3}^\#)$. For any $i < j \in \mathcal{A}_n$ with $n \geq 4$, by the

definition of \sharp , we have that $j^\sharp < i^\sharp$ and there is at most one $k \in \mathcal{A}_n$ satisfying $k = k^\sharp$ and $j - i = i^\sharp - j^\sharp$ when $i \neq i^\sharp, j \neq j^\sharp$. So there are seven cases about the order of i, j, i^\sharp, j^\sharp in \mathcal{A}_n : $i^\sharp = j, i < j = j^\sharp < i^\sharp, j^\sharp < i = i^\sharp < j, i < j < j^\sharp < i^\sharp, j^\sharp < i^\sharp < i < j, i < j^\sharp < j < i^\sharp, j^\sharp < i < i^\sharp < j$. For any $i < j \in \mathcal{A}_n$ with $n \geq 4$, define a map φ_{ij} from \mathcal{A}_n^* to $\mathbf{baxt}_3 \times \mathbf{baxt}_3$ which can be determined by the following four cases according to the order of i, i^\sharp, j, j^\sharp in \mathcal{A}_n .

Case 1. $i^\sharp = j$. Define a map $\lambda : \mathcal{A}_n \rightarrow \mathbf{baxt}_3$ given by

$$k \mapsto \begin{cases} [1]_{\mathbf{baxt}_3} & \text{if } k = i, \\ [3]_{\mathbf{baxt}_3} & \text{if } k = j, \\ [31]_{\mathbf{baxt}_3} & \text{if } i < k < j, \\ [\varepsilon]_{\mathbf{baxt}_3} & \text{otherwise.} \end{cases}$$

Clearly, this map can be extended to a homomorphism $\lambda : \mathcal{A}_n^* \rightarrow \mathbf{baxt}_3$. Define a map $\lambda_{ij} : \mathcal{A}_n \rightarrow \mathbf{baxt}_3 \times \mathbf{baxt}_3$ given by

$$k \mapsto (\lambda(k), \lambda(k)).$$

This map can be extended to a homomorphism $\lambda_{ij} : \mathcal{A}_n^* \rightarrow \mathbf{baxt}_3 \times \mathbf{baxt}_3$. Further, λ_{ij} is also a homomorphism from $(\mathcal{A}_n^*, \sharp)$ to $(\mathbf{baxt}_3 \times \mathbf{baxt}_3, \sharp)$. This is because for any $k \in \mathcal{A}_n$, $\lambda(k^\sharp) = (\lambda(k))^\sharp$ which follows from

$$\begin{cases} \lambda(k^\sharp) = (\lambda(k))^\sharp = [3]_{\mathbf{baxt}_3} & \text{if } k = i, \\ \lambda(k^\sharp) = (\lambda(k))^\sharp = [31]_{\mathbf{baxt}_3} & \text{if } i < k < j, \\ \lambda(k^\sharp) = (\lambda(k))^\sharp = [1]_{\mathbf{baxt}_3} & \text{if } k = j, \\ \lambda(k^\sharp) = (\lambda(k))^\sharp = [\varepsilon]_{\mathbf{baxt}_3} & \text{otherwise.} \end{cases}$$

Therefore for any $\mathbf{w} = k_1 k_2 \cdots k_n$,

$$\begin{aligned} \lambda_{ij}(\mathbf{w}^\sharp) &= (\lambda(\mathbf{w}^\sharp), \lambda(\mathbf{w}^\sharp)) \\ &= (\lambda(k_n^\sharp) \cdots \lambda(k_1^\sharp), \lambda(k_n^\sharp) \cdots \lambda(k_1^\sharp)) \\ &= ((\lambda(k_n))^\sharp \cdots (\lambda(k_1))^\sharp, (\lambda(k_n))^\sharp \cdots (\lambda(k_1))^\sharp) \\ &= ((\lambda(\mathbf{w}))^\sharp, (\lambda(\mathbf{w}))^\sharp) \\ &= (\lambda_{ij}(\mathbf{w}))^\sharp. \end{aligned}$$

Case 2. $i < j = j^\sharp < i^\sharp$ or $j^\sharp < i = i^\sharp < j$. For convenience, let $i_1 = i, i_2 = j$ and $i_3 = i^\sharp$ when $i < j = j^\sharp < i^\sharp$ and $i_1 = j^\sharp, i_2 = i$ and $i_3 = j$ when $j^\sharp < i = i^\sharp < j$. Define maps $\theta_1 : \mathcal{A}_n \rightarrow \mathbf{baxt}_3$ and $\theta_2 : \mathcal{A}_n \rightarrow \mathbf{baxt}_3$ by

$$k \mapsto \begin{cases} [1]_{\mathbf{baxt}_3} & \text{if } k = i_1, \\ [2]_{\mathbf{baxt}_3} & \text{if } k = i_2, \\ [21]_{\mathbf{baxt}_3} & \text{if } i_1 < k < i_2, \\ [\varepsilon]_{\mathbf{baxt}_3} & \text{otherwise,} \end{cases} \quad \text{and} \quad k \mapsto \begin{cases} [2]_{\mathbf{baxt}_3} & \text{if } k = i_2, \\ [3]_{\mathbf{baxt}_3} & \text{if } k = i_3, \\ [32]_{\mathbf{baxt}_3} & \text{if } i_2 < k < i_3, \\ [\varepsilon]_{\mathbf{baxt}_3} & \text{otherwise,} \end{cases}$$

respectively. Clearly, θ_1, θ_2 can be extended to homomorphisms from \mathcal{A}_n^* to \mathbf{baxt}_3 respectively. Define a map $\theta_{ij} : \mathcal{A}_n \rightarrow \mathbf{baxt}_3 \times \mathbf{baxt}_3$ by

$$k \mapsto (\theta_1(k), \theta_2(k)).$$

This map can be extended to a homomorphism $\theta_{ij} : \mathcal{A}_n^* \rightarrow \mathbf{baxt}_3 \times \mathbf{baxt}_3$. Further, θ_{ij} is also a homomorphism from $(\mathcal{A}_n^*, \sharp)$ to $(\mathbf{baxt}_3 \times \mathbf{baxt}_3, \sharp)$. This is because for

any $k \in \mathcal{A}_n$, $\theta_1(k^\sharp) = (\theta_2(k))^\sharp$, $(\theta_1(k))^\sharp = \theta_2(k^\sharp)$ which follows from

$$\begin{cases} \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [3]_{\text{baxt}_3} & \text{if } k = i_1, \\ \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [32]_{\text{baxt}_3} & \text{if } i_1 < k < i_2, \\ \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [2]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [2]_{\text{baxt}_3} & \text{if } k = i_2, \\ \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [21]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } i_2 < k < i_3, \\ \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [1]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } k = i_3, \\ \theta_1(k^\sharp) = (\theta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\theta_1(k))^\sharp = \theta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } k < i_1 \text{ or } k > i_3. \end{cases}$$

Therefore for any $\mathbf{w} = k_1 k_2 \cdots k_n$,

$$\begin{aligned} \theta_{ij}(\mathbf{w}^\sharp) &= (\theta_1(\mathbf{w}^\sharp), \theta_2(\mathbf{w}^\sharp)) \\ &= (\theta_1(k_n^\sharp) \cdots \theta_1(k_1^\sharp), \theta_2(k_n^\sharp) \cdots \theta_2(k_1^\sharp)) \\ &= ((\theta_2(k_n))^\sharp \cdots (\theta_2(k_1))^\sharp, (\theta_1(k_n))^\sharp \cdots (\theta_1(k_1))^\sharp) \\ &= ((\theta_2(\mathbf{w}))^\sharp, (\theta_1(\mathbf{w}))^\sharp) \\ &= (\theta_{ij}(\mathbf{w}))^\sharp. \end{aligned}$$

Case 3. $i < j < j^\sharp < i^\sharp$ or $j^\sharp < i^\sharp < i < j$. For convenience, let $i_1 = i, i_2 = j, i_3 = j^\sharp$ and $i_4 = i^\sharp$ when $i < j < j^\sharp < i^\sharp$ and $i_1 = j^\sharp, i_2 = i^\sharp, i_3 = i$ and $i_4 = j$ when $j^\sharp < i^\sharp < i < j$. Define maps $\eta_1 : \mathcal{A}_n \rightarrow \text{baxt}_3$ and $\eta_2 : \mathcal{A}_n \rightarrow \text{baxt}_3$ by

$$k \mapsto \begin{cases} [1]_{\text{baxt}_3} & \text{if } k = i_1, \\ [2]_{\text{baxt}_3} & \text{if } k = i_2, \\ [21]_{\text{baxt}_3} & \text{if } i_1 < k < i_2, \\ [\varepsilon]_{\text{baxt}_3} & \text{otherwise,} \end{cases} \quad \text{and} \quad k \mapsto \begin{cases} [2]_{\text{baxt}_3} & \text{if } k = i_3, \\ [3]_{\text{baxt}_3} & \text{if } k = i_4, \\ [32]_{\text{baxt}_3} & \text{if } i_3 < k < i_4, \\ [\varepsilon]_{\text{baxt}_3} & \text{otherwise,} \end{cases}$$

respectively. Clearly, η_1, η_2 can be extended to homomorphisms from \mathcal{A}_n^* to baxt_3 respectively. Define a map $\eta_{ij} : \mathcal{A}_n \rightarrow \text{baxt}_3 \times \text{baxt}_3$ by

$$k \mapsto (\eta_1(k), \eta_2(k)).$$

This map can be extended to a homomorphism from \mathcal{A}_n^* to $\text{baxt}_3 \times \text{baxt}_3$. Further, η_{ij} is also a homomorphism from $(\mathcal{A}_n^*, \sharp)$ to $(\text{baxt}_3 \times \text{baxt}_3, \sharp)$. This is because for any $k \in \mathcal{A}_n$, $\eta_1(k^\sharp) = (\eta_2(k))^\sharp$, $(\eta_1(k))^\sharp = \eta_2(k^\sharp)$ which follows from

$$\begin{cases} \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [3]_{\text{baxt}_3} & \text{if } k = i_1, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [32]_{\text{baxt}_3} & \text{if } i_1 < k < i_2, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [2]_{\text{baxt}_3} & \text{if } k = i_2, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [2]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } k = i_3, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [21]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } i_3 < k < i_4, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [1]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{if } k = i_4, \\ \eta_1(k^\sharp) = (\eta_2(k))^\sharp = [\varepsilon]_{\text{baxt}_3}, & (\eta_1(k))^\sharp = \eta_2(k^\sharp) = [\varepsilon]_{\text{baxt}_3} & \text{otherwise.} \end{cases}$$

Therefore it is routine to verify that $\eta_{ij}(\mathbf{w}^\sharp) = (\eta_{ij}(\mathbf{w}))^\sharp$ for any $\mathbf{w} \in \mathcal{A}_n^*$.

Case 4. $i < j^\sharp < j < i^\sharp$ or $j^\sharp < i^\sharp < i < j$. For convenience, let $i_1 = i, i_2 = j^\sharp, i_3 = j$ and $i_4 = i^\sharp$ when $i < j^\sharp < j < i^\sharp$ and $i_1 = j^\sharp, i_2 = i, i_3 = i^\sharp$ and $i_4 = j$ when $j^\sharp < i^\sharp < i < j$. Define maps $\kappa_1 : \mathcal{A}_n \rightarrow \text{baxt}_3$ and $\kappa_2 : \mathcal{A}_n \rightarrow \text{baxt}_3$ by

$$k \mapsto \begin{cases} [1]_{\text{baxt}_3} & \text{if } k = i_1, \\ [21]_{\text{baxt}_3} & \text{if } i_1 < k < i_3, \\ [2]_{\text{baxt}_3} & \text{if } k = i_3, \\ [\varepsilon]_{\text{baxt}_3} & \text{otherwise,} \end{cases} \quad k \mapsto \begin{cases} [2]_{\text{baxt}_3} & \text{if } k = i_2, \\ [3]_{\text{baxt}_3} & \text{if } k = i_4, \\ [32]_{\text{baxt}_3} & \text{if } i_2 < k < i_4, \\ [\varepsilon]_{\text{baxt}_3} & \text{otherwise} \end{cases}$$

respectively. Clearly, κ_1, κ_2 can be extended to homomorphisms from \mathcal{A}_n^* to baxt_3 respectively. Define a map $\kappa_{ij} : \mathcal{A}_n \rightarrow \text{baxt}_3 \times \text{baxt}_3$,

$$k \mapsto (\kappa_1(k), \kappa_2(k)).$$

Clearly, this map can be extended to a homomorphism from \mathcal{A}_n^* to $\text{baxt}_3 \times \text{baxt}_3$. Further, κ_{ij} is a homomorphism from $(\mathcal{A}_n^*, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$. This is because for any $k \in \mathcal{A}_n$, $\kappa_1(k^\#) = (\kappa_2(k))^\#, (\kappa_1(k))^\# = \kappa_2(k^\#)$ which follows from

$$\begin{cases} \kappa_1(k^\#) = (\kappa_2(k))^\# = [\varepsilon]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [3]_{\text{baxt}_3} & \text{if } k = i_1, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [\varepsilon]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [32]_{\text{baxt}_3} & \text{if } i_1 < k < i_2, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [2]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [32]_{\text{baxt}_3} & \text{if } k = i_2, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [21]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [32]_{\text{baxt}_3} & \text{if } i_2 < k < i_3, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [21]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [2]_{\text{baxt}_3} & \text{if } k = i_3, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [21]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [\varepsilon]_{\text{baxt}_3} & \text{if } i_3 < k < i_4, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [1]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [\varepsilon]_{\text{baxt}_3} & \text{if } k = i_4, \\ \kappa_1(k^\#) = (\kappa_2(k))^\# = [\varepsilon]_{\text{baxt}_3}, & (\kappa_1(k))^\# = \kappa_2(k^\#) = [\varepsilon]_{\text{baxt}_3} & \text{otherwise.} \end{cases}$$

Therefore it is routine to verify that $\kappa_{ij}(\mathbf{w}^\#) = (\kappa_{ij}(\mathbf{w}))^\#$ for any $\mathbf{w} \in \mathcal{A}_n^*$.

Now we can define the map $\varphi_{ij} : \mathcal{A}_n^* \rightarrow \text{baxt}_3 \times \text{baxt}_3$ by

$$\varphi_{ij} = \begin{cases} \lambda_{ij} & \text{if } i^\# = j, \\ \theta_{ij} & \text{if } i < j = j^\# < i^\# \text{ or } j^\# < i = i^\# < j, \\ \eta_{ij} & \text{if } i < j < j^\# < i^\# \text{ or } j^\# < i^\# < i < j, \\ \kappa_{ij} & \text{if } i < j^\# < j < i^\# \text{ or } j^\# < i < i^\# < j \end{cases}$$

where $\lambda_{ij}, \theta_{ij}, \eta_{ij}, \kappa_{ij}$ are defined as above. Since each of maps $\lambda_{ij}, \theta_{ij}, \eta_{ij}, \kappa_{ij}$ is a homomorphism from $(\mathcal{A}_n^*, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$. Therefore φ_{ij} is a homomorphism from $(\mathcal{A}_n^*, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$.

Lemma 3.3. *The homomorphism φ_{ij} induces a homomorphism $\varphi_{ij} : (\text{baxt}_n, \#) \rightarrow (\text{baxt}_3 \times \text{baxt}_3, \#)$ for any $n \geq 4$.*

Proof. Note that φ_{ij} is a homomorphism from $(\mathcal{A}_n^*, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$. Then to show that the homomorphism φ_{ij} induces a homomorphism from $(\text{baxt}_n, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$, we only need to show that for any $\mathbf{u}, \mathbf{v} \in \mathcal{A}_n^*$, if $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$, then $\varphi_{ij}(\mathbf{u}) = \varphi_{ij}(\mathbf{v})$. It follows from Proposition 2.5 and the definition of φ_{ij} that the evaluations of the first and the second components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ are the same respectively. In the following, we prove that the left and right precedences of the first and the second components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ are the same respectively.

Case 1. $\varphi_{ij} = \lambda_{ij}$. If there exists $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, since the 1-3 left precedence in the first component of $\lambda_{ij}(\mathbf{u})$ [resp. $\lambda_{ij}(\mathbf{v})$] corresponds to some i - h left precedence in \mathbf{u} [resp. \mathbf{v}] with $i < h \leq j$ and the 3-1 right precedence in the first component of $\lambda_{ij}(\mathbf{u})$ [resp. $\lambda_{ij}(\mathbf{v})$] corresponds to some h - i right precedence in \mathbf{u} [resp. \mathbf{v}] with $i < h \leq j$, it follows from Proposition 2.5 that the first components of $\lambda_{ij}(\mathbf{u})$ and $\lambda_{ij}(\mathbf{v})$ have the same 1-3 left precedence and the same 3-1 right precedence; if there is no $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, then since the 1-3 left precedence in the first component of $\lambda_{ij}(\mathbf{u})$ [resp. $\lambda_{ij}(\mathbf{v})$] corresponds to the i - j left precedence in \mathbf{u} [resp. \mathbf{v}] and the 3-1 right precedence in the first component of $\lambda_{ij}(\mathbf{u})$ [resp. $\lambda_{ij}(\mathbf{v})$] corresponds to the j - i right precedence in \mathbf{u} [resp. \mathbf{v}], it follows from Proposition 2.5 that the first component of $\lambda_{ij}(\mathbf{u})$ and the first component of $\lambda_{ij}(\mathbf{v})$ have the same 1-3 left precedence and the same 3-1 right precedence. A similar argument can show that the left and right precedences of the second components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ are the same.

Case 2. $\varphi_{ij} = \theta_{ij}, \eta_{ij}$ or κ_{ij} .

2.1. $i < j = j^\sharp < i^\sharp$, $i < j < j^\sharp < i^\sharp$ or $i < j^\sharp < j < i^\sharp$. If there exists $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, since the 1-2 left precedence in the first component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to some i - h left precedence in \mathbf{u} [resp. \mathbf{v}] with $i < h \leq j$ and the 2-1 right precedence in the first component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to some h - i right precedence in \mathbf{u} [resp. \mathbf{v}] for $i < h \leq j$, it follows from Proposition 2.5 that the first components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ have the same 1-2 left precedence and the same 2-1 right precedence; if there is no $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, then since the 1-2 left precedence in the first component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to the i - j left precedence in \mathbf{u} [resp. \mathbf{v}] and the 2-1 right precedence in the first component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to the j - i right precedence in \mathbf{u} [resp. \mathbf{v}], it follows from Proposition 2.5 that the first component of $\varphi_{ij}(\mathbf{u})$ and the first component of $\varphi_{ij}(\mathbf{v})$ have the same 1-2 left precedence and the same 2-1 right precedence. A similar argument can show that the left and right precedences of the second components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ are the same.

2.2. $j^\sharp < i = i^\sharp < j$, $j^\sharp < i^\sharp < i < j$, or $j^\sharp < i < i^\sharp < j$. If there exists $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, since the 2-3 left precedence in the second component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to some i - h left precedence in \mathbf{u} [resp. \mathbf{v}] with $i < h \leq j$ and the 3-2 right precedence in the second component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to some h - i right precedence in \mathbf{u} [resp. \mathbf{v}] with $i < h \leq j$, it follows from Proposition 2.5 that the second components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ have the same 2-3 left precedence and the same 3-2 right precedence; if there is no $k \in \text{sup}(\mathbf{u})$ satisfying $i < k < j$, then since the 2-3 left precedence in the second component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to some i - j left precedence in \mathbf{u} [resp. \mathbf{v}] and the 3-2 right precedence in the second component of $\varphi_{ij}(\mathbf{u})$ [resp. $\varphi_{ij}(\mathbf{v})$] corresponds to the j - i right precedence in \mathbf{u} [resp. \mathbf{v}], it follows from Proposition 2.5 that the second component of $\varphi_{ij}(\mathbf{u})$ and the second component of $\varphi_{ij}(\mathbf{v})$ have the same 2-3 left precedence and the same 3-2 right precedence. A similar argument can show that the left and right precedences of the first components of $\varphi_{ij}(\mathbf{u})$ and $\varphi_{ij}(\mathbf{v})$ are the same. \square

Lemma 3.4. *Let $\mathbf{u}, \mathbf{v} \in \mathcal{A}_n^*$ for any $n \geq 4$. Then $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$ if and only if $\varphi_{ij}(\mathbf{u}) = \varphi_{ij}(\mathbf{v})$ for all $1 \leq i < j \leq n$.*

Proof. The necessity follows from the proof of Lemma 3.3. Let $\mathbf{w} \in \mathcal{A}_n^*$ for some $n \geq 4$. Suppose $\text{sup}(\mathbf{w}) = \{a_1 < \dots < a_\ell\}$ for some $\ell \in \mathbb{N}$. We can obtain the evaluation of \mathbf{w} from $\varphi_{a_i, a_j}(\mathbf{w})$. For any i with $1 \leq i < \ell$, if $a_i^\sharp = a_{i+1}$, then since there is no $k \in \text{sup}(\mathbf{w})$ satisfying $a_i < k < a_{i+1}$, it follows from the definition of φ_{ij} that the number of occurrences of 1 in the first [resp. second] component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_i}$ and the number of occurrences of 3 in the first [resp. second] component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_{i+1}}$; if $a_i < a_{i+1} = a_{i+1}^\sharp < a_i^\sharp$, $a_i < a_{i+1} < a_{i+1}^\sharp < a_i^\sharp$ or $a_i < a_{i+1}^\sharp < a_{i+1} < a_i^\sharp$, then since there is no $k \in \text{sup}(\mathbf{w})$ satisfying $a_i < k < a_{i+1}$, it follows from the definition of φ_{ij} that the the number of occurrences of 1 in the first component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_i}$ and the number of occurrences of 2 in the first component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_{i+1}}$; if $a_{i+1}^\sharp < a_i = a_i^\sharp < a_{i+1}$, $a_{i+1}^\sharp < a_i^\sharp < a_i < a_{i+1}$, or $a_{i+1}^\sharp < a_i < a_i^\sharp < a_{i+1}$, then since there is no $k \in \text{sup}(\mathbf{w})$ satisfying $a_i < k < a_{i+1}$, it follows from the definition of φ_{ij} that the number of occurrences of 2 in the second component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_i}$ and the number of occurrences of 3 in the second component of $\varphi_{ij}(\mathbf{w})$ equals to $|\mathbf{w}|_{a_{i+1}}$. Therefore both the number of occurrences of a_i and a_{i+1} in \mathbf{w} can be derived from the maps $\varphi_{a_i a_{i+1}}$.

We can obtain the left and right precedences of \mathbf{w} from $\varphi_{a_i, a_j}(\mathbf{w})$. If a_i, a_j satisfy $a_i^\# = a_j$, $a_i < a_j = a_j^\# < a_i^\#$, $a_i < a_j < a_j^\# < a_i^\#$ or $a_i < a_j^\# < a_j < a_i^\#$, then we check the the first component of $\varphi_{a_i, a_j}(\mathbf{w})$; if a_i, a_j satisfy $a_j^\# < a_i = a_i^\# < a_j$, $a_j^\# < a_i^\# < a_i < a_j$, or $a_j^\# < a_i < a_i^\# < a_j$, then we check the the second component of $\varphi_{a_i, a_j}(\mathbf{w})$.

For each a_j satisfying $a_j^\# < a_j$, if a_i satisfying $a_j^\# < a_i < a_j$ is the largest number such that the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ starts with 2, then, when reading the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ from left-to-right, the first occurrence of 3 corresponds to the first occurrence of a_j , and all occurrences of 2 before the first occurrence of 3 correspond to all occurrences of a_i before the first occurrence of a_j . Hence \mathbf{w} has a a_i - a_j left precedence. Otherwise, if a_i satisfying $a_i^\# = a_j$ or $a_i < a_j^\# < a_j < a_i^\#$ is the largest number such that the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ starts with 1, then, when reading the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ from left-to-right, the first occurrence of 2 or 3 corresponds to the first occurrence of a_j , and all occurrences of 1 before the first occurrence of 2 or 3 correspond to all occurrences of a_i before the first occurrence of a_j , and so \mathbf{w} has a a_i - a_j left precedence; otherwise \mathbf{w} does not have a a_i - a_j left precedence. For each a_j satisfying $a_j \leq a_j^\#$, if $a_i < a_j$ is the largest number such that the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ starts with 1, then, when reading the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ from left-to-right, the first occurrence of 2 corresponds to the first occurrence of a_j , and all occurrences of 1 before the first occurrence of 2 correspond to all occurrences of a_i before the first occurrence of a_j , and so \mathbf{w} has a a_i - a_j left precedence; otherwise \mathbf{w} does not have a a_i - a_j left precedence.

For each a_i satisfying $a_i < a_i^\#$, if a_j satisfying $a_i < a_j \leq a_i^\#$ is the smallest number such that the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ ends with 2 or 3, then, when reading the first component of $\varphi_{a_i, a_j}(\mathbf{w})$ from right-to-left, the first occurrence of 1 corresponds to the first occurrence of a_i , and all occurrences of 2 or 3 before the first occurrence of 1 correspond to all occurrences of a_j before the first occurrence of a_i . Hence \mathbf{w} has a a_j - a_i right precedence. Otherwise, if a_j satisfying $a_j^\# < a_i < a_i^\# < a_j$ is the smallest number such that the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ ends with 3, then, when reading the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ from right-to-left, the first occurrence of 2 corresponds to the first occurrence of a_i , and all occurrences of 3 before the first occurrence of 2 correspond to all occurrences of a_j before the first occurrence of a_i , and so \mathbf{w} has a a_j - a_i right precedence; otherwise \mathbf{w} does not have a a_j - a_i right precedence. For each a_i satisfying $a_i \geq a_i^\#$, if a_j is the smallest number such that the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ ends with 3, then, when reading the second component of $\varphi_{a_i, a_j}(\mathbf{w})$ from right-to-left, the first occurrence of 2 corresponds to the first occurrence of a_i , and all occurrences of 3 before the first occurrence of 2 correspond to all occurrences of a_j before the first occurrence of a_i , and so \mathbf{w} has a a_j - a_i right precedence; otherwise \mathbf{w} does not have a a_j - a_i right precedence.

Therefore, if $\varphi_{ij}(\mathbf{u}) = \varphi_{ij}(\mathbf{v})$ for all $1 \leq i < j \leq n$, then by the above arguments, \mathbf{u} and \mathbf{v} have the same evaluations and left and right precedences. Consequently, $\mathbf{u} \equiv_{\text{baxt}_\infty} \mathbf{v}$. \square

For each $n \in \mathbb{N}$, with $n \geq 4$, let I_n be the index set

$$\{(i, j) : 1 \leq i < j \leq n\}.$$

Now, consider the map

$$\phi_n : (\text{baxt}_n, \#) \rightarrow \prod_{I_n} (\text{baxt}_3 \times \text{baxt}_3, \#),$$

whose (i, j) -th component is given by $\varphi_{ij}([\mathbf{w}]_{\text{baxt}_n})$ for $\mathbf{w} \in \mathcal{A}_n^*$ and $(i, j) \in I_n$.

Proposition 3.5. *The map ϕ_n is an embedding from $(\text{baxt}_n, \#)$ to $(\text{baxt}_3 \times \text{baxt}_3, \#)$.*

Proof. The map ϕ_n is a homomorphism by Lemma 3.3. It follows from the definition of ϕ_n and Lemma 3.4 that $[\mathbf{u}]_{\text{baxt}_n} = [\mathbf{v}]_{\text{baxt}_n}$ if and only if $\varphi_n([\mathbf{u}]_{\text{baxt}_n}) = \varphi_n([\mathbf{v}]_{\text{baxt}_n})$ for any $\mathbf{u}, \mathbf{v} \in \mathcal{A}_n^*$. Hence ϕ_n is an embedding. \square

For any $([\mathbf{u}_1]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}, \dots, [\mathbf{u}_{2|I_n|}]_{\text{baxt}_3}) \in \prod_{2I_n} \text{baxt}_3$, define an involution operation $\#$ on $\prod_{2I_n} \text{baxt}_3$ by

$$([\mathbf{u}_1]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}, \dots, [\mathbf{u}_{2|I_n|}]_{\text{baxt}_3})^\# = ([\mathbf{u}_{2|I_n|}]_{\text{baxt}_3}^\#, \dots, [\mathbf{u}_2]_{\text{baxt}_3}^\#, [\mathbf{u}_1]_{\text{baxt}_3}^\#).$$

Define a map $\psi : \prod_{I_n} (\text{baxt}_3 \times \text{baxt}_3, \#) \rightarrow (\prod_{2I_n} \text{baxt}_3, \#)$ given by

$$\begin{aligned} & (([\mathbf{u}_1]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}), ([\mathbf{u}_3]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}), \dots, ([\mathbf{u}_{2|I_n|-1}]_{\text{baxt}_3}, [\mathbf{u}_{2|I_n|}]_{\text{baxt}_3})) \\ & \mapsto ([\mathbf{u}_1]_{\text{baxt}_3}, [\mathbf{u}_3]_{\text{baxt}_3}, \dots, [\mathbf{u}_{2|I_n|-1}]_{\text{baxt}_3}, [\mathbf{u}_{2|I_n|}]_{\text{baxt}_3}, \dots, [\mathbf{u}_4]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}). \end{aligned}$$

It is routine to verify that the map ψ is an isomorphism. By Theorem 3.2, each element in $(\text{baxt}_3, \#)$ corresponds to a matrix in $UT_{15}(\mathbb{S})$ and the involution $\#$ on $(\text{baxt}_3, \#)$ corresponds to the skew transposition on $UT_{15}(\mathbb{S})$. It follows that there is an embedding, denoted by φ , from $(\prod_{2I_n} \text{baxt}_3, \#)$ to $(UT_{30|I_n|}(\mathbb{S}), {}^D)$. Let

$$\varphi_n = \varphi \circ \psi \circ \phi_n.$$

Then the following result hold.

Theorem 3.6. *For each $n \geq 4$, the map $\varphi_n : (\text{baxt}_n, \#) \rightarrow (UT_{30|I_n|}(\mathbb{S}), {}^D)$ is a faithful representation of $(\text{baxt}_n, \#)$.*

Let $a < b < c < d$ be a 4-element ordered alphabet and

$$B = \langle a, b, c, d \mid \mathcal{R}_{\text{baxt}_\infty}, ac = ca, ad = da, bc = cb, bd = db \rangle \cup \{1\}$$

be a monoid. The involution operation $\#$ on B can be defined by $a \mapsto d, b \mapsto c$.

Theorem 3.7. *For any $m, n \geq 4$, the involution monoids $(\text{baxt}_m, \#)$ and $(\text{baxt}_n, \#)$ generate the same variety.*

Proof. It follows from Proposition 3.5 that

$$\text{Var}(\text{baxt}_4, \#) \subseteq \text{Var}(\text{baxt}_5, \#) \subseteq \dots \subseteq \text{Var}(\text{baxt}_3 \times \text{baxt}_3, \#).$$

It suffices to show that $\text{Var}(\text{baxt}_3 \times \text{baxt}_3, \#) \subseteq \text{Var}(\text{baxt}_4, \#)$. Clearly $(B, \#)$ is a homomorphism image of $(\text{baxt}_4, \#)$, and so $\text{Var}(B, \#) \subseteq \text{Var}(\text{baxt}_4, \#)$. In the following, we show that $\text{Var}(\text{baxt}_3 \times \text{baxt}_3, \#) \subseteq \text{Var}(B, \#)$.

Let φ be a map from $(\text{baxt}_3 \times \text{baxt}_3, \#)$ to $(B, \#) \times (B, \#) \times (B, \#)$ given by

$$\begin{aligned} & ([\varepsilon]_{\text{baxt}_3}, [\varepsilon]_{\text{baxt}_3}) \mapsto (1, 1, 1), \\ & ([1]_{\text{baxt}_3}, [\varepsilon]_{\text{baxt}_3}) \mapsto (a, a, 1), \quad ([2]_{\text{baxt}_3}, [\varepsilon]_{\text{baxt}_3}) \mapsto (b, ba, a), \\ & ([3]_{\text{baxt}_3}, [\varepsilon]_{\text{baxt}_3}) \mapsto (1, b, b), \quad ([\varepsilon]_{\text{baxt}_3}, [1]_{\text{baxt}_3}) \mapsto (1, c, c), \\ & ([\varepsilon]_{\text{baxt}_3}, [2]_{\text{baxt}_3}) \mapsto (c, dc, d), \quad ([\varepsilon]_{\text{baxt}_3}, [3]_{\text{baxt}_3}) \mapsto (d, d, 1). \end{aligned}$$

First to show that φ is well-defined, that is, if $\mathbf{u}_1 \equiv_{\text{baxt}_\infty} \mathbf{v}_1, \mathbf{u}_2 \equiv_{\text{baxt}_\infty} \mathbf{v}_2$, then

$$(U_1, U_2, U_3) = \varphi([\mathbf{u}_1]_{\text{baxt}_3}, [\mathbf{u}_2]_{\text{baxt}_3}) = \varphi([\mathbf{v}_1]_{\text{baxt}_3}, [\mathbf{v}_2]_{\text{baxt}_3}) = (V_1, V_2, V_3).$$

Since $\mathbf{u}_1 \equiv_{\text{baxt}_\infty} \mathbf{v}_1, \mathbf{u}_2 \equiv_{\text{baxt}_\infty} \mathbf{v}_2$, it is easy to see that $\text{ev}(U_i) = \text{ev}(V_i)$ for $i = 1, 2, 3$. It follows from Proposition 2.5 and the definition of $(B, \#)$ that, for any

$([\mathbf{w}_1]_{\text{bax}t_3}, [\mathbf{w}_2]_{\text{bax}t_3}) \in \text{bax}t_3 \times \text{bax}t_3$, the left and right precedences of each component of $\varphi([\mathbf{w}_1]_{\text{bax}t_3}, [\mathbf{w}_2]_{\text{bax}t_3}) = (W_1, W_2, W_3)$ can be characterized as follows:

$$\begin{aligned} (a-b, \ell_1) \in \text{lpi}(W_1) \text{ [resp. } (c-d, \ell_1) \in \text{lpi}(W_3)] &\Leftrightarrow (1-2, \ell_1) \in \text{lpi}(\mathbf{w}_1) \text{ [resp. } \text{lpi}(\mathbf{w}_2)], \\ (a-b, \ell_2) \in \text{lpi}(W_3) \text{ [resp. } (c-d, \ell_2) \in \text{lpi}(W_1)] &\Leftrightarrow (2-3, \ell_2) \in \text{lpi}(\mathbf{w}_1) \text{ [resp. } \text{lpi}(\mathbf{w}_2)], \\ (a-b, \ell_3) \in \text{lpi}(W_2) \text{ [resp. } (c-d, \ell_3) \in \text{lpi}(W_2)] &\Leftrightarrow (1-3, \ell_3) \in \text{lpi}(\mathbf{w}_1) \text{ [resp. } \text{lpi}(\mathbf{w}_2)], \end{aligned}$$

where $\ell_3 \neq \ell_1$ and

$$\begin{aligned} (b-a, r_1) \in \text{rpi}(W_1) \text{ [resp. } (d-c, r_1) \in \text{rpi}(W_3)] &\Leftrightarrow (2-1, r_1) \in \text{rpi}(\mathbf{w}_1) \text{ [resp. } \text{rpi}(\mathbf{w}_2)], \\ (b-a, r_2) \in \text{rpi}(W_3) \text{ [resp. } (d-c, r_2) \in \text{rpi}(W_1)] &\Leftrightarrow (3-2, r_2) \in \text{rpi}(\mathbf{w}_1) \text{ [resp. } \text{rpi}(\mathbf{w}_2)], \\ (b-a, r_3) \in \text{rpi}(W_2) \text{ [resp. } (d-c, r_3) \in \text{rpi}(W_2)] &\Leftrightarrow (3-1, r_3) \in \text{rpi}(\mathbf{w}_1) \text{ [resp. } \text{rpi}(\mathbf{w}_2)]. \end{aligned}$$

where $r_3 \neq r_1$. Since $\mathbf{u}_1 \equiv_{\text{bax}t_\infty} \mathbf{v}_1, \mathbf{u}_2 \equiv_{\text{bax}t_\infty} \mathbf{v}_2$, it is routine to show that $\text{lpi}(U_i) = \text{lpi}(V_i)$ and $\text{rpi}(U_i) = \text{rpi}(V_i)$ for $i = 1, 2, 3$. Thus φ is well-defined, and so φ is a homomorphism.

Next to show that the homomorphism φ is injective. Suppose that $([\mathbf{u}_1]_{\text{bax}t_3}, [\mathbf{u}_2]_{\text{bax}t_3}) \neq ([\mathbf{v}_1]_{\text{bax}t_3}, [\mathbf{v}_2]_{\text{bax}t_3})$. Then $\text{ev}(\mathbf{u}_1) \neq \text{ev}(\mathbf{u}_2)$, or $\text{ev}(\mathbf{v}_1) \neq \text{ev}(\mathbf{v}_2)$, or $\text{lpi}(\mathbf{u}_1) \neq \text{lpi}(\mathbf{u}_2)$, or $\text{rpi}(\mathbf{u}_1) \neq \text{rpi}(\mathbf{u}_2)$, or $\text{lpi}(\mathbf{v}_1) \neq \text{lpi}(\mathbf{v}_2)$ or $\text{rpi}(\mathbf{v}_1) \neq \text{rpi}(\mathbf{v}_2)$. Hence, by the definition of φ , it is routine to show that $\varphi([\mathbf{u}_1]_{\text{bax}t_3}, [\mathbf{u}_2]_{\text{bax}t_3}) \neq \varphi([\mathbf{v}_1]_{\text{bax}t_3}, [\mathbf{v}_2]_{\text{bax}t_3})$. Therefore the homomorphism φ is injective. \square

4. THE IDENTITIES SATISFIED BY $(\text{bax}t_n, \sharp)$

In this section, a complete characterization of the word identities satisfied by $(\text{bax}t_n, \sharp)$ for each finite n is given. Clearly, a word identity $\mathbf{u} \approx \mathbf{v}$ holds in $(\text{bax}t_1, \sharp)$ if and only if $\text{occ}(x, \bar{\mathbf{u}}) = \text{occ}(x, \bar{\mathbf{v}})$ for any $x \in \text{con}(\bar{\mathbf{u}\bar{\mathbf{v}}})$.

Let

$$A = \langle a, b \mid ab = ba \rangle = \{a^m b^n \mid m, n \geq 0\}$$

be a monoid. The monoid A is an involution monoid $(A, *)$ under the unary operation $*$: $a^m b^n \mapsto a^n b^m$. A word identity $\mathbf{u} \approx \mathbf{v}$ is *balanced* if $\text{occ}(x, \mathbf{u}) = \text{occ}(x, \mathbf{v})$ for any $x \in \text{con}(\mathbf{u}\mathbf{v})$.

Lemma 4.1. *A word identity $\mathbf{u} \approx \mathbf{v}$ holds in $(A, *)$ if and only if $\mathbf{u} \approx \mathbf{v}$ is balanced.*

Proof. Suppose that $\mathbf{u} \approx \mathbf{v}$ is a word identity satisfied by $(A, *)$ such that either $\text{occ}(x, \mathbf{u}) \neq \text{occ}(x, \mathbf{v})$ or $\text{occ}(x^*, \mathbf{u}) \neq \text{occ}(x^*, \mathbf{v})$ for some $x \in \mathcal{X}$. Let φ be a homomorphism from $(\mathcal{X} \cup \mathcal{X}^*)^+$ to $(A, *)$ that maps x to a and any other variable to 1. Then $\varphi(\mathbf{u}) = a^{\text{occ}(x, \mathbf{u})} b^{\text{occ}(x^*, \mathbf{u})} \neq a^{\text{occ}(x, \mathbf{v})} b^{\text{occ}(x^*, \mathbf{v})} = \varphi(\mathbf{v})$, a contradiction.

Conversely, if $\mathbf{u} \approx \mathbf{v}$ is balanced, then $\varphi(\mathbf{u}) = \varphi(\mathbf{v})$ for any homomorphism φ from $(\mathcal{X} \cup \mathcal{X}^*)^+$ to $(A, *)$ since $(A, *)$ is commutative. Therefore $\mathbf{u} \approx \mathbf{v}$ holds in $(A, *)$. \square

If \mathbf{u} and \mathbf{v} are words such that $\text{occ}(x, \mathbf{u}) = \text{occ}(x, \mathbf{v})$ for each variable x , then we say that \mathbf{v} is a *permutation* of \mathbf{u} .

Theorem 4.2. *A word identity $\mathbf{u} \approx \mathbf{v}$ holds in $(\text{bax}t_2, \sharp)$ if and only if $\mathbf{u} \approx \mathbf{v}$ is balanced and satisfies that for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$,*

- (I) (a) if $\mathbf{u}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x^* x^\alpha$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x^* x^\alpha$.
- (II) (a) if $\mathbf{u}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x y^\alpha$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x y^\alpha$.

- (III) (a) if $\mathbf{u}[x, y] \in \mathbf{ax}\{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$, then $\mathbf{v}[x, y] \in \mathbf{a}'x\{x, x^*, y, y^*\}^\times$ or $\mathbf{v}[x, y] \in \mathbf{a}'y\{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} ,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x\mathbf{a}$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x\mathbf{a}'$ or $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times y\mathbf{a}'$ where \mathbf{a}' is a permutation of \mathbf{a} .

Proof. Let $\mathbf{u} \approx \mathbf{v}$ be any word identity satisfied by (baxt_2, \sharp) . It is routine to verify that $(A, *)$ is a homomorphic image of (baxt_2, \sharp) under the map given by $[1]_{\text{baxt}_2} \mapsto a$ and $[2]_{\text{baxt}_2} \mapsto b$. Then it follows from Lemma 4.1 that $\mathbf{u} \approx \mathbf{v}$ is balanced. Now to show that (Ia)–(IIIa) hold, and then (Ib)–(IIIb) hold by symmetry. Suppose that \mathbf{u} starts with x but \mathbf{v} starts with $y \neq x$. If $y = x^*$, then letting ϕ_1 be the substitution such that $x \mapsto [2]_{\text{baxt}_2}, z \mapsto [\varepsilon]_{\text{baxt}_2}$ for any $z \neq x, x^*$, we obtain

$$[2]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_1(\mathbf{u}) \neq \phi_1(\mathbf{v}) = [1]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction; if $y \neq x^*$, then letting ϕ_2 be the substitution such that $x \mapsto [2]_{\text{baxt}_2}, y \mapsto [1]_{\text{baxt}_2}, z \mapsto [\varepsilon]_{\text{baxt}_2}$ for any $z \neq x, y$, we obtain

$$[2]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_2(\mathbf{u}) \neq \phi_2(\mathbf{v}) = [1]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Thus \mathbf{u} and \mathbf{v} start with the same variable.

If $\mathbf{u}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$, then $\mathbf{v}[x, y] \in x^\beta z \{x, x^*, y, y^*\}^\times$ with $z \neq x$. Suppose that $z \neq x^*$. Let ϕ_3 be the substitution such that $x \mapsto [2]_{\text{baxt}_2}, z \mapsto [1^{\alpha 2}]_{\text{baxt}_2}$. Then

$$[1^{\alpha 2}]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_3(\mathbf{u}[x, y]) \neq \phi_3(\mathbf{v}[x, y]) = [1^{\alpha+\beta} 2]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Thus $z = x^*$. Suppose that $\alpha \neq \beta$. Let ϕ_4 be the substitution such that $x \mapsto [2]_{\text{baxt}_2}$. Then

$$[1^{\alpha 2}]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_4(\mathbf{u}[x, y]) \neq \phi_4(\mathbf{v}[x, y]) = [1^\beta 2]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Thus $\alpha = \beta$, and so $\mathbf{v}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$. Therefore (Ia) holds.

If $\mathbf{u}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$, then $\mathbf{v}[x, y] \in y^\beta z \{x, x^*, y, y^*\}^\times$ with $z \neq y$. Clearly $z \neq y^*$ by condition (Ia). If $z = x^*$, then $\mathbf{u}[x]$ starts with x but $\mathbf{v}[x]$ starts with x^* , which is impossible. Therefore $z = x$. Suppose that $\alpha \neq \beta$. Let ϕ_5 be the substitution such that $x \mapsto [2]_{\text{baxt}_2}, y \mapsto [1]_{\text{baxt}_2}$. Then

$$[1^{\alpha 2}]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_5(\mathbf{u}[x, y]) \neq \phi_5(\mathbf{v}[x, y]) = [1^\beta 2]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Thus $\alpha = \beta$, and so $\mathbf{v}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$. Therefore (IIa) holds.

If $\mathbf{u}[x, y] \in \mathbf{ax}\{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$, then by condition (IIa) $\mathbf{v}[x, y] \in \mathbf{a}'y\{x, x^*, y, y^*\}^\times$ or $\mathbf{v}[x, y] \in \mathbf{a}'x\{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}') = \{x^*, y^*\}$. Suppose that $|\mathbf{a}'| \neq |\mathbf{a}|$. Let ϕ_6 be the substitution from \mathcal{X} to baxt_2 such that $x \mapsto [2]_{\text{baxt}_2}, y \mapsto [2]_{\text{baxt}_2}$. Then

$$[1^{|\mathbf{a}'|} 2]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_6(\mathbf{u}[x, y]) \neq \phi_6(\mathbf{v}[x, y]) = [1^{|\mathbf{a}'|} 2]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Therefore $|\mathbf{a}'| = |\mathbf{a}|$. Suppose that $\text{occ}(x^*, \mathbf{a}') \neq \text{occ}(x^*, \mathbf{a})$. Let ϕ_7 be the substitution from \mathcal{X} to baxt_2 such that $x \mapsto [22]_{\text{baxt}_2}, y \mapsto [2]_{\text{baxt}_2}$. Then

$$[1^{|\mathbf{a}|+\text{occ}(x^*, \mathbf{u})} 2]_{\text{baxt}_2} \cdot [\mathbf{s}]_{\text{baxt}_2} = \phi_7(\mathbf{u}[x, y]) \neq \phi_7(\mathbf{v}[x, y]) = [1^{|\mathbf{a}'|+\text{occ}(x^*, \mathbf{v})} 2]_{\text{baxt}_2} \cdot [\mathbf{t}]_{\text{baxt}_2}$$

where $\mathbf{s}, \mathbf{t} \in \mathcal{A}_2^*$, a contradiction. Therefore $\text{occ}(x^*, \mathbf{a}') = \text{occ}(x^*, \mathbf{a}), \text{occ}(y^*, \mathbf{a}') = \text{occ}(y^*, \mathbf{a})$, and so condition (IIIa) holds.

Conversely, let $\mathbf{u} \approx \mathbf{v}$ be any balanced word identity satisfying (I)–(III) and ϕ be any substitution from \mathcal{X} to baxt_2 . Note that $\text{ev}(\phi(\mathbf{u})) = \text{ev}(\phi(\mathbf{v}))$ since $\mathbf{u} \approx \mathbf{v}$

is balanced. If $\text{sup}(\phi(\mathbf{u})) = \{1\}$ or $\{2\}$, then $\phi(\mathbf{u}) = \phi(\mathbf{v})$. If $\text{sup}(\phi(\mathbf{u})) = \{1, 2\}$, to show $\phi(\mathbf{u}) = \phi(\mathbf{v})$, it suffices to show $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$ and $\text{lpi}(\phi(\mathbf{u})) = \text{lpi}(\phi(\mathbf{v}))$. By symmetry, we only need to show that $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$. If $\phi(\mathbf{u})$ does not have a 2-1 right precedence, then $\phi(\mathbf{v})$ does not have a 2-1 right precedence by conditions (I) and (II). If $\phi(\mathbf{u})$ has a 2-1 right precedence of index r , then we may assume that $\phi(x) \neq [\varepsilon]_{\text{bax}t_2}$ for any $x \in \text{con}(\mathbf{u})$ by Remark 2.4, whence \mathbf{u} can be written into the form $\mathbf{u} = \mathbf{u}_1 z \mathbf{u}_2$ for some possibly empty words $\mathbf{u}_1, \mathbf{u}_2$ satisfying $\text{sup}(\phi(\mathbf{u}_2)) = \{2\}$ and $\text{sup}(\phi(z)) = \{1, 2\}$ or $\{1\}$. Note that $z \notin \text{con}(\mathbf{u}_2)$ and $z^* \in \text{con}(\mathbf{u}_2)$ if $\text{sup}(\phi(z)) = \{1\}$ and that for any $x \in \text{con}(\mathbf{u}_2), x^* \notin \text{con}(\mathbf{u}_2)$. As such, $\phi(z \mathbf{u}_2)$ has the same 2-1 right precedence of index r as $\phi(\mathbf{u})$. Now we consider the form of \mathbf{v} . There are two cases.

Case 1. $z^* \notin \text{con}(\mathbf{u}_2)$. It follows from (IIb) that $\mathbf{v} = \mathbf{v}_1 z \mathbf{v}_2$ where \mathbf{v}_2 is a permutation of \mathbf{u}_2 . This implies that $\text{sup}(\phi(\mathbf{v}_2)) = \{2\}$, hence $\phi(z \mathbf{v}_2)$ also has a 2-1 right precedence of index r . Since $\phi(z \mathbf{v}_2)$ has the same right precedence as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 2-1 right precedence of index r . Therefore $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$.

Case 2. $z^* \in \text{con}(\mathbf{u}_2)$. Let $\text{con}(\mathbf{u}_2) = \{z^*, x_1, x_2, \dots, x_n\}$. Without loss of generality, we may assume that $\text{fp}(\mathbf{u}_2) = z^* x_1 x_2 \dots x_n$. Note that $x_1^*, x_2^*, \dots, x_n^* \notin \text{con}(\mathbf{u}_2)$. Then $\mathbf{v} = \mathbf{v}_1 \mathbf{y} z^* \mathbf{x}_1 x_1 \mathbf{x}_2 x_2 \dots \mathbf{x}_n x_n$ by (IIb) where $\mathbf{y} \in \{x_1, x_1^*, \dots, x_n, x_n^*, z, z^*\}^+$ and $\mathbf{x}_i \in \{x_i, x_{i+1}, \dots, x_n\}^+$ for $i = 1, 2, \dots, n$. Therefore either $\mathbf{v} = \mathbf{v}_1 z \mathbf{v}_2$ or $\mathbf{v} = \mathbf{v}_1 x_i^* \mathbf{v}_2$ with $\text{con}(\mathbf{v}_2) = \text{con}(\mathbf{u}_2)$.

2.1. $\mathbf{v} = \mathbf{v}_1 z \mathbf{v}_2$. It follows from (IIIb) that \mathbf{v}_2 is a permutation of \mathbf{u}_2 . This implies that $\text{sup}(\phi(\mathbf{v}_2)) = \{2\}$, hence $\phi(z \mathbf{v}_2)$ also has a 2-1 right precedence of index r . Since $\phi(z \mathbf{v}_2)$ has the same right precedence as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 2-1 right precedence of index r . Therefore $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$.

2.2. $\mathbf{v} = \mathbf{v}_1 x_i^* \mathbf{v}_2$. It follows from (IIIb) that $\text{occ}(z^*, \mathbf{u}_2) = \text{occ}(z^*, \mathbf{v}_2)$ and $\text{occ}(x_i, \mathbf{u}_2) = \text{occ}(x_i, \mathbf{v}_2)$. Suppose that there exists some $x_j \in \text{con}(\mathbf{u}_2)$ such that $\text{occ}(x_j, \mathbf{u}_2) \neq \text{occ}(x_j, \mathbf{v}_2)$. Then $x_j \neq x_i, z^*$. If $x_j^* \notin \text{con}(\mathbf{u}_1)$, then it follows from (IIIb) that $\overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v})$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_z(x_j, \mathbf{v})$. It follows from the forms of \mathbf{u} and \mathbf{v} that $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) \geq 0$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) \leq 0$. Suppose that $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) > 0$. Then $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) > 0$, which is impossible. If $x_j^* \in \text{con}(\mathbf{u}_1)$, then there are four cases about the order of $\infty x_j^*, \infty x_i^*, \infty z$ in \mathbf{u}, \mathbf{v} : $\infty x_j^* \prec_{\mathbf{u}} \infty x_i^* \prec_{\mathbf{u}} \infty z$ and $\infty x_j^* \prec_{\mathbf{v}} \infty z \prec_{\mathbf{v}} \infty x_i^*$; $\infty x_j^* \prec_{\mathbf{u}} \infty x_i^* \prec_{\mathbf{u}} \infty z$ and $\infty z \prec_{\mathbf{v}} \infty x_j^* \prec_{\mathbf{v}} \infty x_i^*$; $\infty x_i^* \prec_{\mathbf{u}} \infty x_j^* \prec_{\mathbf{u}} \infty z$ and $\infty x_j^* \prec_{\mathbf{v}} \infty z \prec_{\mathbf{v}} \infty x_i^*$; and $\infty x_i^* \prec_{\mathbf{u}} \infty x_j^* \prec_{\mathbf{u}} \infty z$ and $\infty z \prec_{\mathbf{v}} \infty x_j^* \prec_{\mathbf{v}} \infty x_i^*$.

If $\infty x_j^* \prec_{\mathbf{u}} \infty x_i^* \prec_{\mathbf{u}} \infty z$ and $\infty x_j^* \prec_{\mathbf{v}} \infty z \prec_{\mathbf{v}} \infty x_i^*$, then it follows from (IIIb) that $\overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v})$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_z(x_j, \mathbf{v})$. By the forms of \mathbf{u} and \mathbf{v} , $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) \geq 0$, $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) \leq 0$. Suppose $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) > 0$. Then $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) > 0$, which is impossible.

If $\infty x_j^* \prec_{\mathbf{u}} \infty x_i^* \prec_{\mathbf{u}} \infty z$ and $\infty z \prec_{\mathbf{v}} \infty x_j^* \prec_{\mathbf{v}} \infty x_i^*$, then it follows from (IIIb) that $\overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v})$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v})$. By the forms of \mathbf{u} and \mathbf{v} , $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) \geq 0$, $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) \leq 0$. Suppose $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) > 0$. Then $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) > 0$, which is impossible.

If $\infty x_i^* \prec_{\mathbf{u}} \infty x_j^* \prec_{\mathbf{u}} \infty z$ and $\infty x_j^* \prec_{\mathbf{v}} \infty z \prec_{\mathbf{v}} \infty x_i^*$, then it follows from (IIIb) that $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v})$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_z(x_j, \mathbf{v})$. By the forms of \mathbf{u} and \mathbf{v} , $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) \geq 0$, $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) \leq 0$. Suppose $\overrightarrow{\text{occ}}_z(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) > 0$. Then $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{u}) > 0$, which is impossible.

If $\infty x_i^* \prec_{\mathbf{u}} \infty x_j^* \prec_{\mathbf{u}} \infty z$ and $\infty z \prec_{\mathbf{v}} \infty x_j^* \prec_{\mathbf{v}} \infty x_i^*$, then it follows from (IIIb) that $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v})$ and $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) = \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v})$. By the forms of

\mathbf{u} and \mathbf{v} , $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) \geq 0$, $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{u}) \leq 0$. Suppose $\overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{v}) - \overrightarrow{\text{occ}}_{x_i^*}(x_j, \mathbf{v}) > 0$. Then $\overrightarrow{\text{occ}}_z(x_j, \mathbf{u}) - \overrightarrow{\text{occ}}_{x_j^*}(x_j, \mathbf{u}) > 0$, which is impossible.

Thus $\mathbf{v} = \mathbf{v}_1 x_i^* \mathbf{v}_2$ where \mathbf{v}_2 is a permutation of \mathbf{u}_2 . This implies that $\phi(\mathbf{v}_2)$ has support $\{2\}$, hence $\phi(x_i^* \mathbf{v}_2)$ also has a 2-1 right precedence of index r . Since $\phi(x_i^* \mathbf{v}_2)$ has the same right precedence as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 2-1 right precedence of index r . Therefore $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$. \square

Theorem 4.3. *A word identity $\mathbf{u} \approx \mathbf{v}$ holds in (baxt_3, \sharp) if and only if $\mathbf{u} \approx \mathbf{v}$ is balanced and satisfies that for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$,*

- (I) (a) if $\mathbf{u}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in x^\alpha x^* \{x, x^*, y, y^*\}^\times$,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x^* x^\alpha$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x^* x^\alpha$.
- (II) (a) if $\mathbf{u}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in y^\alpha x \{x, x^*, y, y^*\}^\times$,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x y^\alpha$ for some $\alpha \geq 1$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x y^\alpha$.
- (III) (a) if $\mathbf{u}[x, y] \in \mathbf{a}x \{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$, then $\mathbf{v}[x, y] \in \mathbf{a}'x \{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} ,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times x \mathbf{a}$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times x \mathbf{a}'$ where \mathbf{a}' is a permutation of \mathbf{a} .
- (IV) (a) $\overrightarrow{\text{occ}}_y(x, \mathbf{u}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{u}) = \overrightarrow{\text{occ}}_y(x, \mathbf{v}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{v})$,
- (b) $\overrightarrow{\text{occ}}_y(x, \mathbf{u}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{u}) = \overrightarrow{\text{occ}}_y(x, \mathbf{v}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{v})$.
- (V) (a) if $\mathbf{u}[x, y] \in \mathbf{a}y \{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x, x^*\}$, then $\mathbf{v}[x, y] \in \mathbf{a}'y \{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} ,
- (b) if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times y \mathbf{a}$ with $\text{con}(\mathbf{a}) = \{x, x^*\}$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times y \mathbf{a}'$ where \mathbf{a}' is a permutation of \mathbf{a} .

Proof. Suppose that $\mathbf{u} \approx \mathbf{v}$ is a word identity satisfied by (baxt_3, \sharp) . It is routine to verify that the involution submonoid of (baxt_3, \sharp) generated by $[1]_{\text{baxt}_3}$ and $[3]_{\text{baxt}_3}$ is isomorphic to (baxt_2, \sharp) . It follows from Theorem 4.2 that $\mathbf{u} \approx \mathbf{v}$ is balanced and satisfies conditions (I)–(II). Now to show that (IIIa)–(Va) hold, and then (IIIb)–(Vb) hold by symmetry.

Suppose that $\mathbf{u}[x, y] \in \mathbf{a}x \{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x^*, y^*\}$. Then it follows from condition (IIIa) of Theorem 4.2 that $\mathbf{v}[x, y] \in \mathbf{a}'x \{x, x^*, y, y^*\}^\times$ or $\mathbf{v}[x, y] \in \mathbf{a}'y \{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} . Suppose that $\mathbf{v}[x, y] \in \mathbf{a}'y \{x, x^*, y, y^*\}^\times$. Let ϕ_1 be a substitution such that $x \mapsto [3]_{\text{baxt}_3}$, $y \mapsto [2]_{\text{baxt}_3}$ and any other variable to $[\varepsilon]_{\text{baxt}_3}$. Then $\phi_1(\mathbf{u})$ have a 2-3 left precedence of index $\text{occ}(y^*, \mathbf{a})$ but the index of 2-3 left precedence of $\phi_1(\mathbf{v})$ is greater than $\text{occ}(y^*, \mathbf{a})$, a contradiction. Therefore $\mathbf{v}[x, y] \in \mathbf{a}'x \{x, x^*, y, y^*\}^\times$, and so (IIIa) holds.

Suppose that $\overrightarrow{\text{occ}}_y(x, \mathbf{u}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{u}) \neq \overrightarrow{\text{occ}}_y(x, \mathbf{v}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{v})$ for some $x, x^* \neq y \in \text{con}(\mathbf{u})$. Let ϕ_2 be a substitution such that $x \mapsto [2]_{\text{baxt}_3}$, $y \mapsto [3]_{\text{baxt}_3}$ and any other variable to $[\varepsilon]_{\text{baxt}_3}$. Then $\phi_2(\mathbf{u})$ has a 2-3 left precedence of index $\overrightarrow{\text{occ}}_y(x, \mathbf{u}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{u})$ but $\phi_2(\mathbf{v})$ has a 2-3 left precedence of index $\overrightarrow{\text{occ}}_y(x, \mathbf{v}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{v})$, a contradiction. Therefore $\overrightarrow{\text{occ}}_y(x, \mathbf{u}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{u}) = \overrightarrow{\text{occ}}_y(x, \mathbf{v}) + \overrightarrow{\text{occ}}_y(x^*, \mathbf{v})$, and so (IVa) holds.

If $\mathbf{u}[x, y] \in \mathbf{a}y \{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x, x^*\}$, then it follows from (I) that $\mathbf{v}[x, y] \in \mathbf{a}'y \{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}') = \{x, x^*\}$. By (IVb), $\text{occ}(x, \mathbf{a}) + \text{occ}(x^*, \mathbf{a}) = \text{occ}(x, \mathbf{a}') + \text{occ}(x^*, \mathbf{a}')$. Suppose that $\text{occ}(x, \mathbf{a}) \neq \text{occ}(x, \mathbf{a}')$. Let ϕ_3 be a substitution such that $x \mapsto [1]_{\text{baxt}_3}$, $y \mapsto [2]_{\text{baxt}_3}$ and any other variable to $[\varepsilon]_{\text{baxt}_3}$.

Then $\phi_3(\mathbf{u})$ has a 1-2 left precedence of index $\text{occ}(x, \mathbf{a})$ but $\phi_3(\mathbf{v})$ has a 1-2 left precedence of index $\text{occ}(x, \mathbf{a}')$, a contradiction. Therefore $\mathbf{v}[x, y] \in \mathbf{a}'y\{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} , and so (Va) holds.

Conversely, let $\mathbf{u} \approx \mathbf{v}$ be any balanced word identity satisfying (I)–(V) and ϕ be any substitution from \mathcal{X} to baxt_3 . Note that $\text{ev}(\phi(\mathbf{u})) = \text{ev}(\phi(\mathbf{v}))$ since $\mathbf{u} \approx \mathbf{v}$ is balanced. Now, to show $\phi(\mathbf{u}) = \phi(\mathbf{v})$, it suffices to show $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$ and $\text{lpi}(\phi(\mathbf{u})) = \text{lpi}(\phi(\mathbf{v}))$. By symmetry, we only need to show that $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$. If $\phi(\mathbf{u})$ does not have a 2-1 [resp. 3-2, 3-1] right precedence, then $\phi(\mathbf{v})$ does not have a 2-1 [resp. 3-2, 3-1] right precedence by (I) and (II). In the following, we show that if $\phi(\mathbf{u})$ has a 2-1 [resp. 3-2, 3-1] right precedence of index r , then $\phi(\mathbf{v})$ has a 2-1 [resp. 3-2, 3-1] right precedence of index r .

If $\phi(\mathbf{u})$ has a 3-1 right precedence of index r , then we may assume that $\phi(x) \neq [\varepsilon]_{\text{baxt}_3}$ for any $x \in \text{con}(\mathbf{u})$ by Remark 2.4, whence \mathbf{u} can be written into the form $\mathbf{u} = \mathbf{u}_1z\mathbf{u}_2$ for some possibly empty words $\mathbf{u}_1, \mathbf{u}_2$ satisfying $\text{sup}(\phi(\mathbf{u}_2)) = \{3\}$ and $\text{sup}(\phi(z)) = \{1\}$ or $\{1, 3\}$. Note that $z \notin \text{con}(\mathbf{u}_2)$ and z^* may occur in \mathbf{u}_2 if $\text{sup}(\phi(z)) = \{1\}$ and that for any $x \in \text{con}(\mathbf{u}_2)$, $x^* \notin \text{con}(\mathbf{u}_2)$. As such, $\phi(z\mathbf{u}_2)$ has the same 3-1 right precedence of index r as $\phi(\mathbf{u})$. Now we consider the form of \mathbf{v} . If $z^* \notin \text{con}(\mathbf{u}_2)$, then it follows from (IIb) that $\mathbf{v} = \mathbf{v}_1z\mathbf{v}_2$ where \mathbf{v}_2 is a permutation of \mathbf{u}_2 . This implies that $\text{sup}(\phi(\mathbf{v}_2)) = \{3\}$, hence $\phi(z\mathbf{v}_2)$ also has a 3-1 right precedence of index r . Since $\phi(z\mathbf{v}_2)$ has the same 3-1 right precedence of index r as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 3-1 right precedence of index r . If $z^* \in \text{con}(\mathbf{u}_2)$. Let $\text{con}(\mathbf{u}_2) = \{z^*, x_1, x_2, \dots, x_n\}$. Without loss of generality, we may assume that $\text{fp}(\mathbf{u}_2) = z^*x_1x_2 \cdots x_n$. Note that $x_1^*, x_2^*, \dots, x_n^* \notin \text{con}(\mathbf{u}_2)$. Then it follows from (IIb) that $\mathbf{v} = \mathbf{v}_1\mathbf{y}z^*\mathbf{x}_1x_1\mathbf{x}_2x_2 \cdots \mathbf{x}_nx_n$ where $\mathbf{y} \in \{x_1, x_1^*, \dots, x_n, x_n^*, z, z^*\}^+$ and $\mathbf{x}_i \in \{x_i, x_{i+1}, \dots, x_n\}^+$ for $i = 1, 2, \dots, n$. Therefore $\mathbf{v} = \mathbf{v}_1z\mathbf{v}_2$ where \mathbf{v}_2 is a permutation of \mathbf{u}_2 by (IIIb). This implies that $\phi(\mathbf{v}_2)$ has support $\{3\}$, hence $\phi(z\mathbf{v}_2)$ also has a 3-1 right precedence of index r . Since $\phi(z\mathbf{v}_2)$ has the 3-1 right precedence of index r as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 3-1 right precedence of index r .

If $\phi(\mathbf{u})$ has a 2-1 right precedence of index r , then we may assume that $\phi(x) \neq [\varepsilon]_{\text{baxt}_3}$ for any $x \in \text{con}(\mathbf{u})$ by Remark 2.4, whence \mathbf{u} can be written into the form $\mathbf{u} = \mathbf{u}_1z\mathbf{u}_2$ for some possibly empty words $\mathbf{u}_1, \mathbf{u}_2$ satisfying $\text{sup}(\phi(\mathbf{u}_2)) \subseteq \{2, 3\}$ and $\text{sup}(\phi(z)) = \{1\}, \{1, 2\}, \{1, 3\}$ or $\{1, 2, 3\}$. Note that $z \notin \text{con}(\mathbf{u}_2)$ and z^* may occur in \mathbf{u}_2 if $\text{sup}(\phi(z)) = \{1\}$ or $\{1, 2\}$ and that for any $x \in \text{con}(\mathbf{u}_2)$, x^* may occur in $\text{con}(\mathbf{u}_2)$ if $\text{sup}(\phi(x)) = \{2\}$. As such, $\phi(z\mathbf{u}_2)$ has the same 2-1 right precedence of index r as $\phi(\mathbf{u})$. It follows from (IIb)–(Vb) that $\mathbf{v} = \mathbf{v}_1z\mathbf{v}_2$ satisfying that $z \notin \text{con}(\mathbf{v}_2)$, $\text{con}(\mathbf{u}_2) = \text{con}(\mathbf{v}_2)$, and $\text{occ}(z^*, \mathbf{u}_2) = \text{occ}(z^*, \mathbf{v}_2)$, and $\text{occ}(x, \mathbf{u}_2) = \text{occ}(x, \mathbf{v}_2)$ when $x \in \text{con}(\mathbf{u}_2)$, $x^* \notin \text{con}(\mathbf{u}_2)$, and $\text{occ}(x, \mathbf{u}_2) + \text{occ}(x^*, \mathbf{u}_2) = \text{occ}(x, \mathbf{v}_2) + \text{occ}(x^*, \mathbf{v}_2)$ when $x, x^* \in \text{con}(\mathbf{u}_2)$. This implies that $\phi(z\mathbf{v}_2)$ also has a 2-1 right precedence of index r . Since $\phi(z\mathbf{v}_2)$ has the same 2-1 right precedence of index r as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 2-1 right precedence of index r .

If $\phi(\mathbf{u})$ has a 3-2 right precedence of index r , then we may assume that $\phi(x) \neq [\varepsilon]_{\text{baxt}_3}$ for any $x \in \text{con}(\mathbf{u})$ by Remark 2.4, whence \mathbf{u} can be written into the form $\mathbf{u} = \mathbf{u}_1z\mathbf{u}_2$ satisfying $\text{sup}(\phi(\mathbf{u}_2)) \subseteq \{1, 3\}$ and $\text{sup}(\phi(z)) = \{2\}, \{1, 2\}, \{2, 3\}$ or $\{1, 2, 3\}$. Note that z, z^* cannot occur in \mathbf{u}_2 and that for any $x \in \text{con}(\mathbf{u}_2)$, x^* may occur in $\text{con}(\mathbf{u}_2)$. As such, $\phi(z\mathbf{u}_2)$ has the same 3-2 right precedence of index r as $\phi(\mathbf{u})$. It follows from (IIb)–(Vb) that $\mathbf{v} = \mathbf{v}_1z\mathbf{v}_2$ such that $z \notin \text{con}(\mathbf{v}_2)$, $\text{con}(\mathbf{u}_2) = \text{con}(\mathbf{v}_2)$, $\text{occ}(x, \mathbf{u}_2) = \text{occ}(x, \mathbf{v}_2)$ and $\text{occ}(x^*, \mathbf{u}_2) = \text{occ}(x^*, \mathbf{v}_2)$. This implies that $\phi(z\mathbf{v}_2)$ also has a 3-2 right precedence of index r . Since $\phi(z\mathbf{v}_2)$ has the same 3-2 right precedence of index r as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a 3-2 right precedence of index r . Therefore $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$. \square

Theorem 4.4. *A word identity $\mathbf{u} \approx \mathbf{v}$ holds in $(\mathbf{baxt}_4, \#)$ if and only if $\mathbf{u} \approx \mathbf{v}$ is balanced and $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$, $\overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ for any $x, y \in \text{con}(\mathbf{u})$.*

Proof. Suppose that $\mathbf{u} \approx \mathbf{v}$ is a word identity satisfied by $(\mathbf{baxt}_4, \#)$. It is routine to verify that the involution submonoid of $(\mathbf{baxt}_4, \#)$ generated by $[1]_{\mathbf{baxt}_4}$ and $[4]_{\mathbf{baxt}_3}$ is isomorphic to $(\mathbf{baxt}_2, \#)$. Then it follows from Theorem 4.2 that $\mathbf{u} \approx \mathbf{v}$ is balanced. Suppose that $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) \neq \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$. Let ϕ be a substitution such that $x \mapsto [1]_{\mathbf{baxt}_4}$, $y \mapsto [2]_{\mathbf{baxt}_4}$ and any other variable to $[\varepsilon]_{\mathbf{baxt}_4}$. Then $\phi(\mathbf{u})$ has a 2-1 right precedence of index $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u})$ but $\phi(\mathbf{v})$ has a 2-1 left precedence of index $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$, a contradiction. Therefore $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ for any $x, y \in \text{con}(\mathbf{u})$. Symmetrically, $\overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ for any $x, y \in \text{con}(\mathbf{u})$.

Conversely, let $\mathbf{u} \approx \mathbf{v}$ be any balanced word identity satisfying $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$, $\overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ for any $x, y \in \text{con}(\mathbf{u})$ and ϕ be any substitution from \mathcal{X} to \mathbf{baxt}_4 . Note that $\text{ev}(\phi(\mathbf{u})) = \text{ev}(\phi(\mathbf{v}))$ since $\mathbf{u} \approx \mathbf{v}$ is balanced. Now, to show $\phi(\mathbf{u}) = \phi(\mathbf{v})$, it suffices to show that $\text{rpi}(\phi(\mathbf{u})) = \text{rpi}(\phi(\mathbf{v}))$, and then $\text{lpi}(\phi(\mathbf{u})) = \text{lpi}(\phi(\mathbf{v}))$ by symmetry. If $\phi(\mathbf{u})$ does not have a j - i right precedence with $i < j$, then $\phi(\mathbf{v})$ does not have a j - i right precedence by $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$. If $\phi(\mathbf{u})$ has a j - i right precedence of index r , then \mathbf{u} can be written into the form $\mathbf{u} = \mathbf{u}_1 z \mathbf{u}_2$ satisfying $i \notin \text{sup}(\phi(\mathbf{u}_2))$ and $i \in \text{sup}(\phi(z))$. Note that z cannot occur in \mathbf{u}_2 . As such, $\phi(z \mathbf{u}_2)$ has the same j - i right precedence of index r as $\phi(\mathbf{u})$. It follows from $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ that $\mathbf{v} = \mathbf{v}_1 z \mathbf{v}_2$ such that $z \notin \text{con}(\mathbf{v}_2)$ and \mathbf{v}_2 is a permutation of \mathbf{u}_2 . This implies that $\phi(z \mathbf{v}_2)$ also has a j - i right precedence of index r . Since $\phi(z \mathbf{v}_2)$ has the same j - i right precedence of index r as $\phi(\mathbf{v})$, $\phi(\mathbf{v})$ has a j - i right precedence of index r . Therefore $\text{rpi}(\phi(\mathbf{u}_2)) = \text{rpi}(\phi(\mathbf{v}))$. \square

Remark 4.5. It follows from Theorems 4.2–4.4 that the involution monoids $(\mathbf{baxt}_2, \#)$, $(\mathbf{baxt}_3, \#)$ and $(\mathbf{baxt}_n, \#)$ with $n \geq 4$ generate different varieties. However, the monoids \mathbf{baxt}_n with $n \geq 2$ generate the same variety by [14, Theorem 3.12] or [11, Proposition 6.10]. And by Theorems 4.2–4.4, an identity $\mathbf{u} \approx \mathbf{v}$ holds in \mathbf{baxt}_n with $n \geq 2$ if and only if

- (i) $\mathbf{u} \approx \mathbf{v}$ is balanced;
- (ii) $\overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overrightarrow{\text{oc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ and $\overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{u}) = \overleftarrow{\text{dc}\check{c}_x}(\mathbf{u}, \mathbf{v})$ for any $x, y \in \text{con}(\mathbf{u})$.

This fact also has been shown in [14, Theorem 4.3].

5. FINITE BASIS PROBLEM FOR $(\mathbf{baxt}_n, \#)$

In this section, the finite basis problem for involution monoid $(\mathbf{baxt}_n, \#)$ for each finite n is solved. Clearly, $(\mathbf{baxt}_1, \#)$ is finitely based since its involution is trivial and it is commutative.

For any word $\mathbf{u} \in (\mathcal{A} \cup \mathcal{A}^*)^+$, denote by $\overleftarrow{\mathbf{u}}$ the word obtained from \mathbf{u} by writing \mathbf{u} in reverse. For example, if $\mathbf{u} = x^5 z y^* (z^*)^3 x^*$, then $\overleftarrow{\mathbf{u}} = x^* (z^*)^3 y^* z x^5$. The identity $\overleftarrow{\mathbf{u}} \approx \overleftarrow{\mathbf{v}}$ is called the *reverse* of $\mathbf{u} \approx \mathbf{v}$. It follows from the proof of [37, Lemma 7(i)] that if an involution semigroup $(S, *)$ satisfies the word identity $\mathbf{u} \approx \mathbf{v}$, then $(S, *)$ also satisfies $\overleftarrow{\mathbf{u}} \approx \overleftarrow{\mathbf{v}}$.

First, we show that $(\mathbf{baxt}_2, \#)$ is finitely based.

Theorem 5.1. *The identities (1.1) and*

$$\left. \begin{aligned} x^* h x k x y s x^* t x &\approx x^* h x k y x s x^* t x, & x^* h x k x y s x t x^* &\approx x^* h x k y x s x t x^*, \\ x h x^* k x y s x^* t x &\approx x h x^* k y x s x^* t x, & x h x^* k x y s x t x^* &\approx x h x^* k y x s x t x^*, \end{aligned} \right\} (5.1)$$

$$\left. \begin{aligned} x^* h x k x y s y^* t y &\approx x^* h x k y x s y^* t y, & x^* h x k x y s y t y^* &\approx x^* h x k y x s y t y^*, \\ x h x^* k x y s y^* t y &\approx x h x^* k y x s y^* t y, & x h x^* k x y s y t y^* &\approx x h x^* k y x s y t y^*, \end{aligned} \right\} (5.2)$$

$$x h y k x y s x t y \approx x h y k y x s x t y, \quad x h y k x y s y t x \approx x h y k y x s y t x, \quad (5.3)$$

$$x h y k x y s x^* t y^* \approx x h y k y x s x^* t y^*, \quad x h y k x y s y^* t x^* \approx x h y k y x s y^* t x^*, \quad (5.4)$$

$$x^*hy^*kxysx^*ty^* \approx x^*hy^*kyxsx^*ty^*, \quad x^*hy^*kxysy^*tx^* \approx x^*hy^*kyxsy^*tx^*, \quad (5.5)$$

$$\left. \begin{aligned} x^*h\dot{x}kxysxty &\approx x^*h\dot{x}kyxsxty, & x^*h\dot{x}kxysytx &\approx x^*h\dot{x}kyxsytx, \\ xh\dot{x}^*kxysxty &\approx xh\dot{x}^*kyxsxty, & xh\dot{x}^*kxysytx &\approx xh\dot{x}^*kyxsytx, \end{aligned} \right\} (5.6)$$

$$\left. \begin{aligned} x^*h\dot{x}kxysx^*ty^* &\approx x^*h\dot{x}kyxsx^*ty^*, & x^*h\dot{x}kxysy^*tx^* &\approx x^*h\dot{x}kyxsy^*tx^*, \\ xh\dot{x}^*kxysx^*ty^* &\approx xh\dot{x}^*kyxsx^*ty^*, & xh\dot{x}^*kxysy^*tx^* &\approx xh\dot{x}^*kyxsy^*tx^*, \end{aligned} \right\} (5.7)$$

and their reverses constitute an identity basis for $(\mathbf{baxt}_2, \#)$.

Proof. Clearly, $(\mathbf{baxt}_2, \#)$ satisfies (1.1), and the identities (5.1)–(5.7) and their reverses by Theorem 4.2.

Note that the identities (1.1) can be used to convert any non-empty term into some unique word. It suffices to show that each non-trivial word identity satisfied by $(\mathbf{baxt}_2, \#)$ can be deduced from (5.1)–(5.7) and their reverses. Note that any identity satisfied by $(\mathbf{baxt}_2, \#)$ is balanced. Let Σ be the set of all non-trivial balanced word identities satisfied by $(\mathbf{baxt}_2, \#)$ but can not be deducible from (5.1)–(5.7) and their reverses. Suppose that $\Sigma \neq \emptyset$. Note that any balanced identity $\mathbf{u} \approx \mathbf{v}$ in Σ can be written uniquely into the form $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ where a, b are distinct variables and $|\mathbf{u}'| = |\mathbf{v}'|$. Choose an identity, say $\mathbf{u} \approx \mathbf{v}$, from Σ such that when it is written as $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$, the lengths of the words \mathbf{u}' and \mathbf{v}' are as short as possible. Since $\mathbf{u} \approx \mathbf{v}$ is balanced, we have $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$ and $\text{occ}(x, \mathbf{u}'a) = \text{occ}(x, \mathbf{v}'b)$ for any $x \in \text{con}(\mathbf{u}'a)$.

Since $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$, we have $b \in \text{con}(\mathbf{u}')$, and so $\mathbf{u}'a = \mathbf{u}_1bc\mathbf{u}_2$ with $b \neq c$ and $b \notin \text{con}(\mathbf{u}_2)$. Since $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$, we have $c \in \text{con}(\mathbf{v}')$, and so $\mathbf{v}' = \mathbf{v}_1c\mathbf{v}_2$ with $c \notin \text{con}(\mathbf{u}_2)$. Thus $\mathbf{u} = \mathbf{u}_1bc\mathbf{u}_2\mathbf{w}$ and $\mathbf{v} = \mathbf{v}_1c\mathbf{v}_2b\mathbf{w}$. On the one hand, we claim that at least one of four cases $b, b^* \in \text{con}(\mathbf{u}_1)$, $c, c^* \in \text{con}(\mathbf{u}_1)$, $b^*, c^* \in \text{con}(\mathbf{u}_1)$ or $b, c \in \text{con}(\mathbf{u}_1)$ is true. If $b, b^*, c, c^* \notin \text{con}(\mathbf{u}_1)$, then $\text{occ}(b, \mathbf{u}'a) = 1$, and so $\mathbf{u}[b, c]$ starts with b but $\mathbf{v}[b, c]$ does not, which contradicts with (I) of Theorem 4.2. If only b occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with $b^{\text{occ}(b, \mathbf{u}'a)}c$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIa) of Theorem 4.2. If only b^* occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with $(b^*)^{\text{occ}(b^*, \mathbf{u}_1)}b$ but $\mathbf{v}[b, c]$ does not, which contradicts with (Ia) of Theorem 4.2. If only c occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with $c^{\text{occ}(c, \mathbf{u}_1)}b$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIa) of Theorem 4.2. If only c^* occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with $(c^*)^{\text{occ}(c^*, \mathbf{u}_1)}b$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIa) of Theorem 4.2. If only c, b^* occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with \mathbf{u}'_1b with $\text{con}(\mathbf{u}'_1) = \{c, b^*\}$ but $\mathbf{v}[b, c]$ does not start with \mathbf{v}'_1b or \mathbf{v}'_1c^* with $\text{con}(\mathbf{v}'_1) = \{c, b^*\}$ and $\text{occ}(c, \mathbf{v}'_1) = \text{occ}(c, \mathbf{u}'_1)$, which contradicts with (IIIa) of Theorem 4.2. If only c^*, b occurs in \mathbf{u}_1 , then $\mathbf{u}[b, c]$ starts with \mathbf{u}'_1c with $\text{con}(\mathbf{u}'_1) = \{c^*, b\}$ but $\mathbf{v}[b, c]$ does not start with \mathbf{v}'_1b^* or \mathbf{v}'_1c with $\text{con}(\mathbf{v}'_1) = \{c^*, b\}$ and $\text{occ}(b, \mathbf{v}'_1) = \text{occ}(b, \mathbf{u}'_1)$, which contradicts with (IIIa) of Theorem 4.2.

On the other hand, we claim that at least one of four cases $b, b^* \in \text{con}(\mathbf{u}_2\mathbf{w})$, $c, c^* \in \text{con}(\mathbf{u}_2\mathbf{w})$, $b^*, c^* \in \text{con}(\mathbf{u}_2\mathbf{w})$ or $b, c \in \text{con}(\mathbf{u}_2\mathbf{w})$ is true. If $b, b^*, c, c^* \notin \text{con}(\mathbf{u}_2\mathbf{w})$, then $\mathbf{u}[b, c]$ ends with c but $\mathbf{v}[b, c]$ ends with b , which contradicts with (I) of Theorem 4.2. If only b occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with $cb^{\text{occ}(b, \mathbf{w})}$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIb) of Theorem 4.2. If only b^* occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with $c(b^*)^{\text{occ}(b^*, \mathbf{u}_2\mathbf{w})}$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIb) of Theorem 4.2. If only c occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with $bc^{1+\text{occ}(c, \mathbf{u}_2\mathbf{w})}$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIb) of Theorem 4.2. If only c^* occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with $c(c^*)^{\text{occ}(c^*, \mathbf{u}_2\mathbf{w})}$ but $\mathbf{v}[b, c]$ does not, which contradicts with (IIb) of Theorem 4.2. If only c, b^* occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with \mathbf{u}'_2 with $\text{con}(\mathbf{u}'_2) = \{c, b^*\}$ but $\mathbf{v}[b, c]$ does not end with \mathbf{v}'_2 or $c^*\mathbf{v}'_2$ with $\text{con}(\mathbf{v}'_2) = \{c, b^*\}$ and $\text{occ}(c, \mathbf{v}'_2) = \text{occ}(c, \mathbf{u}'_2)$, which contradicts with (IIIb) of Theorem 4.2. If only c^*, b occurs in $\mathbf{u}_2\mathbf{w}$, then $\mathbf{u}[b, c]$ ends with \mathbf{u}'_2 with $\text{con}(\mathbf{u}'_2) = \{c^*, b\}$ but $\mathbf{v}[b, c]$ does not end with \mathbf{v}'_2 or $b^*\mathbf{v}'_2$ with $\text{con}(\mathbf{v}'_2) = \{c^*, b\}$ and $\text{occ}(b, \mathbf{v}'_2) = \text{occ}(b, \mathbf{u}'_2)$,

which contradicts with (IIIb) of Theorem 4.2. Therefore, the identities (5.1)–(5.7) and their reverses can be used to convert $\mathbf{u}_1bc\mathbf{u}_2\mathbf{w}$ into $\mathbf{u}_1cb\mathbf{u}_2\mathbf{w}$. By repeating this process, the word $\mathbf{u} = \mathbf{u}'a\mathbf{w} = \mathbf{u}_1bc\mathbf{u}_2\mathbf{w}$ can be converted into the word $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w}$ by the identities (5.1)–(5.7) and their reverses.

Clearly the identities $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ holds in (baxt_2, \sharp) . Note that $|\mathbf{u}_1c\mathbf{u}_2b\mathbf{w}| = |\mathbf{u}'a\mathbf{w}| = |\mathbf{v}'b\mathbf{w}|$ and words in the identity $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ have a longer common suffix than words in the identity $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$. Hence $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w} \notin \Sigma$ by the minimality assumption on the lengths \mathbf{u}' and \mathbf{v}' , that is, $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ can be deducible from (5.1)–(5.7) and their reverses. We have shown that $\mathbf{u}'a\mathbf{w} \approx \mathbf{u}_1c\mathbf{u}_2b\mathbf{w}$ can be deducible from (5.1)–(5.7) and their reverses. Hence $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ can be deducible from (5.1)–(5.7) and their reverses, which contradicts with $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w} \in \Sigma$. Therefore $\Sigma = \emptyset$. \square

Next, we show that (baxt_3, \sharp) is non-finitely based.

Theorem 5.2. *Suppose that $(M, *)$ is an involution monoid satisfying the following conditions:*

(I) *for each $k \geq 2$, $(M, *)$ satisfies the identity $\mathbf{p}_k \approx \mathbf{q}_k$ where*

$$\begin{aligned} \mathbf{p}_k &= x_1^*x_2^*\cdots x_{2k}^* \cdot xx^* \cdot x^*x_1 \cdots x_{2k}x \cdot x^*x \cdot x_1^*x_3^* \cdots x_{2k-1}^*x_2^*x_4^* \cdots x_{2k}^*, \\ \mathbf{q}_k &= x_1^*x_2^*\cdots x_{2k}^* \cdot xx^* \cdot xx_1 \cdots x_{2k}x^* \cdot x^*x \cdot x_1^*x_3^* \cdots x_{2k-1}^*x_2^*x_4^* \cdots x_{2k}^*; \end{aligned}$$

(II) *$\text{ip}(\mathbf{u}) = \text{ip}(\mathbf{v})$, $\text{fp}(\mathbf{u}) = \text{fp}(\mathbf{v})$ and $\mathbf{u} \approx \mathbf{v}$ is balanced for any $\mathbf{u} \approx \mathbf{v}$ satisfied by $(M, *)$;*

(III) *if $(M, *)$ satisfies a non-trivial identity $y^*xx^* \cdot x^*yx \cdot x^*xy^* \approx \mathbf{w}$, then $\mathbf{w} = y^*xx^* \cdot xyx^* \cdot x^*xy^*$;*

(IV) (a) *if $\mathbf{u}[x, y] \in \mathbf{a}y\{x, x^*, y, y^*\}^\times$ with $\text{con}(\mathbf{a}) = \{x, x^*\}$, then $\mathbf{v}[x, y] \in \mathbf{a}'y\{x, x^*, y, y^*\}^\times$ where \mathbf{a}' is a permutation of \mathbf{a} ,*
 (b) *if $\mathbf{u}[x, y] \in \{x, x^*, y, y^*\}^\times y\mathbf{a}$ with $\text{con}(\mathbf{a}) = \{x, x^*\}$, then $\mathbf{v}[x, y] \in \{x, x^*, y, y^*\}^\times y\mathbf{a}'$ where \mathbf{a}' is a permutation of \mathbf{a} .*

*Then $(M, *)$ is non-finitely based.*

The proof of Theorem 5.2 is given at the end of Lemma 5.4.

For each $k \geq 2$, define

$$\begin{aligned} \mathbf{P}_k &= x_1^*x_2^*\cdots x_{2k}^* \cdot xx^* \cdot x^*x_{1\pi} \cdots x_{2k\pi}x \cdot x^*x \cdot x_1^*x_3^* \cdots x_{2k-1}^*x_2^*x_4^* \cdots x_{2k}^*, \\ \mathbf{Q}_k &= x_1^*x_2^*\cdots x_{2k}^* \cdot xx^* \cdot xx_{1\sigma} \cdots x_{2k\sigma}x^* \cdot x^*x \cdot x_1^*x_3^* \cdots x_{2k-1}^*x_2^*x_4^* \cdots x_{2k}^*, \end{aligned}$$

where π, σ are any permutations on $\{1, 2, \dots, 2k\}$.

Lemma 5.3. *Let $(M, *)$ be an involution monoid satisfying conditions (II) and (III) in Theorem 5.2. Suppose that $\mathbf{p}_k \approx \mathbf{w}$ is any word identity satisfied by $(M, *)$ such that $\mathbf{p}_k \in \mathbf{P}_k$. Then $\mathbf{w} \in \mathbf{P}_k \cup \mathbf{Q}_k$.*

Proof. Note that $\mathbf{p}_k[x_i, x] = x_i^*xx^* \cdot x^*x_ix \cdot x^*xx_i^*$ for $i = 1, 2, \dots, 2k$. It follows from the condition (II) of Theorem 5.2 that $\mathbf{w}[x_i, x] = x_i^*xx^* \cdot x^*x_ix \cdot x^*xx_i^*$ or $\mathbf{w}[x_i, x] = x_i^*xx^* \cdot xx_ix^* \cdot x^*xx_i^*$. Then it follows from (III) Theorem 5.2 that $\mathbf{w} \in \mathbf{P}_k \cup \mathbf{Q}_k$. \square

A word identity $\mathbf{u} \approx \mathbf{v}$ is *k-limited* if $\text{con}(\overline{\mathbf{u}\mathbf{v}}) \leq k$. For any involution semigroup $(S, *)$, let $\text{id}_k(S, *)$ denote the set of all *k-limited* word identities of $(S, *)$.

Lemma 5.4. *Suppose that $(M, *)$ is an involution monoid satisfying conditions (II)–(IV) in Theorem 5.2. Let $\mathbf{s} \approx \mathbf{t}$ be an identity which is directly deducible from some identity in $\text{id}_{2k}(M, *)$ with $[\mathbf{s}] \in \mathbf{P}_k$. Then $[\mathbf{t}] \in \mathbf{Q}_k$.*

Proof. Let $\mathbf{u} \approx \mathbf{v}$ be a word identity in $\text{id}_{2k}(S, *)$ from which the identity $\mathbf{s} \approx \mathbf{t}$ is directly deducible. There is a substitution $\varphi : \mathcal{X} \rightarrow \mathbb{T}(\mathcal{X})$ such that $\varphi(\mathbf{u})$ is a subterm of \mathbf{s} , and replacing this particular subterm $\varphi(\mathbf{u})$ of \mathbf{s} with $\varphi(\mathbf{v})$ results in \mathbf{t} . By Remark 2.2, either $[\varphi(\mathbf{u})]$ or $[(\varphi(\mathbf{u}))^*]$ is a factor of $[\mathbf{s}]$. It suffices to consider the former case since the latter is similar. Hence there exist words $\mathbf{a}, \mathbf{b} \in (\mathcal{X} \cup \mathcal{X}^*)^+$ such that $[\mathbf{s}] = \mathbf{a}[\varphi(\mathbf{u})]\mathbf{b}$. Since \mathbf{t} is obtained by replacing $\varphi(\mathbf{u})$ in \mathbf{s} with $\varphi(\mathbf{v})$, it follows that $[\mathbf{t}] = \mathbf{a}[\varphi(\mathbf{v})]\mathbf{b}$. Since $[\mathbf{s}] \in \mathbb{P}_k$, it follows from Lemma 5.3 that $[\mathbf{t}] \in \mathbb{P}_k \cup \mathbb{Q}_k$. Working toward a contradiction, suppose that $[\mathbf{t}] \in \mathbb{Q}_k$. Since $[\mathbf{s}]$ and $[\mathbf{t}]$ share the same prefix \mathbf{a} and the same suffix \mathbf{b} and $\text{ip}([\varphi(\mathbf{u})]) = \text{ip}([\varphi(\mathbf{v})])$ and $\text{fp}([\varphi(\mathbf{u})]) = \text{fp}([\varphi(\mathbf{v})])$, $[\varphi(\mathbf{u})]$ contains the factor

$$x^* \cdot x^* x_{1\pi} \cdots x_{2k\pi} x \cdot x^*$$

and $[\varphi(\mathbf{v})]$ contains the factor

$$x^* \cdot x x_{1\pi} \cdots x_{2k\pi} x^* \cdot x^*.$$

It follows from (IV) of Theorem 5.2 that $\overleftarrow{\text{occ}}_{x_{i\pi}}(x_{i\pi}^*, \mathbf{u}\varphi) \neq 0$ and $\overrightarrow{\text{occ}}_{x_{i\pi}}(x_{i\pi}^*, \varphi(\mathbf{u})) \neq 0$ for each $i \in \{1, 2, \dots, 2k\}$, and so $\mathbf{a}, \mathbf{b} = \emptyset$. Since $\mathbf{u} \approx \mathbf{v}$ is $2k$ -limited and $[\varphi(\mathbf{u})] \approx [\varphi(\mathbf{v})]$ is $2k+1$ -limited, there exists a variable $s \in \text{con}(\mathbf{u})$ such that $[\varphi(s)]$ contains one of the following factors:

$$x^* x_{1\pi}, x_{1\pi} x_{2\pi}, x_{2\pi} x_{3\pi}, \dots, x_{2k-1\pi} x_{2k\pi}, x_{2k\pi} x.$$

Suppose that there exists a variable $s \in \text{con}(\mathbf{u})$ such that $[\varphi(s)]$ contains $x_{2k\pi} x$ as a factor. Then $[\mathbf{t}]$ also contains $x_{2k\pi} x$ as a factor, which contradicts with $[\mathbf{t}] \in \mathbb{Q}_k$. Similarly, there does not exist $s \in \text{con}(\mathbf{u})$ such that $[\varphi(s)]$ contains $x^* x_{1\pi}$ as a factor. Suppose that there exists a variable $s \in \text{con}(\mathbf{u})$ such that $[\varphi(s)]$ contains $x_{i\pi} x_{(i+1)\pi}$ as a factor for some $i = 1, 2, \dots, 2k-1$. Clearly, $\text{occ}(s, \mathbf{u}) = 1$. Then $x_{i\pi} x_{(i+1)\pi}$ is a factor of $[\varphi(\mathbf{v})]$ since $\mathbf{u} \approx \mathbf{v}$ is balanced. It follows from the definition of \mathbb{P}_k that $\mathbf{u} = \mathbf{u}_1 s \mathbf{u}_2$, and s^* can not occur in $\text{con}(\mathbf{u}_1)$ and $\text{con}(\mathbf{u}_2)$ simultaneously. Suppose that $s^* \notin \text{con}(\mathbf{u}_2)$. Note that $x x^* x$ is a factor of $[\mathbf{s}]$ occurring after $[\varphi(s)]$. Then there exist at most three variables in $\text{con}(\mathbf{u}_2)$, say s_1, s_2, s_3 , such that $[\varphi(s_1 s_2 s_3)]$ contains $x x^* x$ as a factor in $[\mathbf{s}]$. It follows from conditions (II) and (IV) of Theorem 5.2 that $\mathbf{v} = \mathbf{v}_1 s \mathbf{v}_2$ satisfying $s, s^* \notin \text{con}(\mathbf{v}_2)$, $\text{occ}(s_1, \mathbf{v}_2) = \text{occ}(s_1, \mathbf{u}_2)$, $\text{occ}(s_2, \mathbf{v}_2) = \text{occ}(s_2, \mathbf{u}_2)$ and $\text{occ}(s_3, \mathbf{v}_2) = \text{occ}(s_3, \mathbf{u}_2)$, and so $x x^* x$ occurs after $x_{i\pi} x_{i+1\pi}$ in $[\mathbf{t}]$, but this contradicts with $\mathbf{a}[\varphi(\mathbf{v})]\mathbf{b} \in \mathbb{Q}_k$. Consequently, $\mathbf{a}[\varphi(\mathbf{v})]\mathbf{b} \in \mathbb{P}_k$. \square

Proof of Theorem 5.2. Let $(M, *)$ be any involution monoid that satisfies conditions (I)–(IV) in Theorem 5.2. Then there exists some set Σ of word identities such that $(1.1) \cup \Sigma$ is an identity basis for $(M, *)$. Working toward a contradiction, suppose that $(M, *)$ is finitely based. Then there exists a finite subset Σ_{fin} of Σ such that all identities of $(M, *)$ are deducible from $(1.1) \cup \Sigma_{\text{fin}}$. Hence there exists some fixed integer k such that $\Sigma_{\text{fin}} \subseteq \Sigma \cap \text{id}_{2k}(M, *)$. By (I), the involution monoid $(M, *)$ satisfies some word identity $\mathbf{p}_k \approx \mathbf{q}_k$ with $\mathbf{p}_k \in \mathbb{P}_k$ and $\mathbf{q}_k \in \mathbb{Q}_k$. Therefore there exists some sequence

$$\mathbf{p}_k = \mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_m = \mathbf{q}_k$$

of terms such that each identity $\mathbf{s}_i \approx \mathbf{s}_{i+1}$ is directly deducible from some identity $\mathbf{u}_i \approx \mathbf{v}_i \in (1.1) \cup \Sigma_{\text{fin}}$. The equality $\mathbf{s}_1 = \mathbf{p}_k \in \mathbb{P}_k$ holds. If $\mathbf{s}_i \in \mathbb{P}_k$ for some $i \geq 1$, then there are two cases depending on whether the identity $\mathbf{u}_i \approx \mathbf{v}_i$ is from (1.1) or Σ_{fin} . If $\mathbf{u}_i \approx \mathbf{v}_i$ is from (1.1), then $\mathbf{s}_i = \mathbf{s}_{i+1}$ by Remark 2.3, whence $\mathbf{s}_{i+1} \in \mathbb{P}_k$. If $\mathbf{u}_i \approx \mathbf{v}_i$ is from Σ_{fin} , then $\mathbf{s}_{i+1} \in \mathbb{P}_k$ by Lemma 5.4. Therefore $\mathbf{s}_{i+1} \in \mathbb{P}_k$ in any case, whence by induction, $\mathbf{s}_i \in \mathbb{P}_k$ for all i . But this implies the contradiction $\mathbf{q}_k = \mathbf{s}_m \in \mathbb{P}_k$. Consequently, the involution monoid $(M, *)$ is non-finitely based. \square

Theorem 5.5. *The involution monoid $(\mathbf{baxt}_3, \sharp)$ is non-finitely based.*

Proof. It follows from Theorem 4.3 that $(\mathbf{baxt}_3, \sharp)$ satisfies all of conditions of Theorem 5.2. Therefore $(\mathbf{baxt}_3, \sharp)$ is non-finitely based. \square

Now, we consider the finite basis problem for $(\mathbf{baxt}_n, \sharp)$ with $n \geq 4$.

Theorem 5.6. *The identities (1.1) and*

$$xhykxy\,sxt\,y \approx xhyk\,yx\,sxt\,y, \quad xhykxy\,sytx \approx xhyk\,yx\,sytx, \quad (5.3)$$

constitute an identity basis for $(\mathbf{baxt}_n, \sharp)$ for $n \geq 4$.

Proof. By Theorem 3.7, we only need to show that the result holds for $(\mathbf{baxt}_4, \sharp)$. Clearly, $(\mathbf{baxt}_4, \sharp)$ satisfies the identities (1.1), and (5.3) by Theorem 4.4.

Note that the identities (1.1) can be used to convert any non-empty term into some unique word. It suffices to show that each non-trivial word identity satisfied by $(\mathbf{baxt}_4, \sharp)$ can be derived from (5.3). Note that any identity satisfied by $(\mathbf{baxt}_4, \sharp)$ is balanced. Let Σ be the set of all non-trivial balanced word identities satisfied by $(\mathbf{baxt}_4, \sharp)$ but can not be deducible from (5.3). Suppose that $\Sigma \neq \emptyset$. Note that any balanced identity $\mathbf{u} \approx \mathbf{v}$ in Σ can be written uniquely into the form $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ where a, b are distinct variables and $|\mathbf{u}'| = |\mathbf{v}'|$. Choose an identity, say $\mathbf{u} \approx \mathbf{v}$, from Σ such that when it is written as $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$, the lengths of the words \mathbf{u}' and \mathbf{v}' are as short as possible. Since $\mathbf{u} \approx \mathbf{v}$ is balanced, we have $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$ and $\text{occ}(x, \mathbf{u}'a) = \text{occ}(x, \mathbf{v}'b)$ for any $x \in \text{con}(\mathbf{u}'a)$.

Since $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$, we have $b \in \text{con}(\mathbf{u}')$, and so $\mathbf{u}'a = \mathbf{u}_1bc\mathbf{u}_2$ with $b \neq c, b \notin \text{con}(\mathbf{u}_2)$. Since $\text{con}(\mathbf{u}'a) = \text{con}(\mathbf{v}'b)$, we have $c \in \text{con}(\mathbf{v}')$, and so $\mathbf{v}' = \mathbf{v}_1c\mathbf{v}_2$ with $c \notin \text{con}(\mathbf{u}_2)$. Thus $\mathbf{u} = \mathbf{u}_1bc\mathbf{u}_2\mathbf{w}$ and $\mathbf{v} = \mathbf{v}_1c\mathbf{v}_2b\mathbf{w}$. We claim that $b, c \in \text{con}(\mathbf{u}_1)$. If $b, c \notin \text{con}(\mathbf{u}_1)$, then $\text{occ}(b, \mathbf{u}'a) = 1$, and so $\overline{\delta\text{cc}}_c(b, \mathbf{u}) = 1$ but $\overline{\delta\text{cc}}_c(b, \mathbf{v}) = 0$, which contradicts with Theorem 4.4. If only b occurs in \mathbf{u}_1 , then $\overline{\delta\text{cc}}_c(b, \mathbf{u}) > \overline{\delta\text{cc}}_c(b, \mathbf{v})$, which contradicts with Theorem 4.4. If only c occurs in \mathbf{u}_1 , then $\overline{\delta\text{cc}}_c(c, \mathbf{u}) < \overline{\delta\text{cc}}_c(c, \mathbf{v})$, which contradicts with Theorem 4.4. We also claim that $b, c \in \text{con}(\mathbf{u}_2\mathbf{w})$. If $b, c \notin \text{con}(\mathbf{u}_2\mathbf{w})$, then $\overline{\text{occ}}_b(c, \mathbf{u}) = 1$ but $\overline{\text{occ}}_b(c, \mathbf{v}) = 0$, which contradicts with Theorem 4.4. If only b occurs in $\mathbf{u}_2\mathbf{w}$, then $\overline{\text{occ}}_c(b, \mathbf{u}) < \overline{\text{occ}}_c(b, \mathbf{v})$, which contradicts with Theorem 4.4. If only c occurs in $\mathbf{u}_2\mathbf{w}$, then $\overline{\text{occ}}_b(c, \mathbf{u}) > \overline{\text{occ}}_b(c, \mathbf{v})$, which contradicts with Theorem 4.4. As such, we can deduce the word $\mathbf{u}_1cb\mathbf{u}_2\mathbf{w}$, by applying the identities (5.3). By repeating this process, the word $\mathbf{u}_1bc\mathbf{u}_2\mathbf{w}$ can be converted into the word $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w}$.

Clearly the identities $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ holds in $(\mathbf{baxt}_4, \sharp)$. Note that $|\mathbf{u}_1c\mathbf{u}_2b\mathbf{w}| = |\mathbf{u}'a\mathbf{w}| = |\mathbf{v}'b\mathbf{w}|$ and words in the identity $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ have a longer common suffix than words in the identity $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$. Hence $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w} \notin \Sigma$ by the minimality assumption on the lengths of \mathbf{u}' and \mathbf{v}' , that is, $\mathbf{u}_1c\mathbf{u}_2b\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ can be deducible from (5.3). We have shown that $\mathbf{u}'a\mathbf{w} \approx \mathbf{u}_1c\mathbf{u}_2b\mathbf{w}$ can be deducible from (5.3). Hence $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w}$ can be deducible from (5.3), which contradicts with $\mathbf{u}'a\mathbf{w} \approx \mathbf{v}'b\mathbf{w} \in \Sigma$. Therefore $\Sigma = \emptyset$. \square

In the end, we consider the number of subvarieties of $\text{Var}(\mathbf{baxt}_n, \sharp)$ for each $n \geq 2$. Recall that a word \mathbf{u} is an *isoterm* for an involution monoid if it does not satisfy any non-trivial word identity of the form $\mathbf{u} \approx \mathbf{v}$.

Lemma 5.7 ([21, Theorem 3.6]). *Let $(M, *)$ be any involution monoid with isoterms xx^*yy^* and xyy^*x^* . Then the variety $\text{Var}(M, *)$ contains continuum many subvarieties.*

Theorem 5.8. *The variety $\text{Var}(\mathbf{baxt}_n, \sharp)$ for each finite $n \geq 2$ contains continuum many subvarieties.*

Proof. By Theorems 4.2–4.4, it is routine to show that both the words xx^*yy^* and xyy^*x^* are isoterm for (baxt_n, \sharp) for each finite $n \geq 2$. Now the results follows from Lemma 5.7. \square

6. RECOGNIZING IDENTITIES OF (baxt_n, \sharp) IN POLYNOMIAL TIME

In this section, it is shown that the identity checking problem of (baxt_n, \sharp) for each finite n belongs to the complexity class P.

Let $\mathbf{u} \in (\mathcal{X} \cup \mathcal{X}^*)^+$. If $x, x^* \in \text{con}(\mathbf{u})$, then $\{x, x^*\}$ is called a *mixed pair* of \mathbf{u} . Denote by $\text{pre}(\mathbf{u})$ the longest prefix of \mathbf{u} containing only one variable, that is, $|\text{con}(\text{pre}(\mathbf{u}))| = 1$, and $\text{suf}(\mathbf{u})$ the longest suffix of \mathbf{u} containing only one variable, that is, $|\text{con}(\text{suf}(\mathbf{u}))| = 1$; and denote by $\text{pren}(\mathbf{u})$ the longest prefix of \mathbf{u} which does not contain any mixed pair, and $\text{sufn}(\mathbf{u})$ the longest suffix of \mathbf{u} which does not contain any mixed pair.

Theorem 6.1. *The decision problem $\text{CHECK-ID}(\text{baxt}_n, \sharp)$ for each finite n belongs to the complexity class P.*

Proof. For decision problem $\text{CHECK-ID}(\text{baxt}_1, \sharp)$, given any word identity $\mathbf{u} \approx \mathbf{v}$, it suffices to show that whether $\text{con}(\bar{\mathbf{u}}) = \text{con}(\bar{\mathbf{v}})$ and $\text{occ}(x, \mathbf{u}) + \text{occ}(x^*, \mathbf{u}) = \text{occ}(x, \mathbf{v}) + \text{occ}(x^*, \mathbf{v})$ for any $x \in \text{con}(\bar{\mathbf{u}})$. Clearly, these can be completed in polynomial time.

For decision problem $\text{CHECK-ID}(\text{baxt}_2, \sharp)$, given any word identity $\mathbf{u} \approx \mathbf{v}$, it suffices to show that one can check whether the identity $\mathbf{u} \approx \mathbf{v}$ is balanced and the words \mathbf{u} and \mathbf{v} satisfy conditions of Theorem 4.2 in polynomial time.

To check whether $\mathbf{u} \approx \mathbf{v}$ is balanced, it suffices to check whether $\text{con}(\mathbf{u}) = \text{con}(\mathbf{v})$ and $\text{occ}(x, \mathbf{u}) = \text{occ}(x, \mathbf{v})$ for any $x \in \text{con}(\mathbf{u})$. Clearly, these can be completed in polynomial time.

To check conditions (I)–(II) of Theorem 4.2, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$, whether $\text{pre}(\mathbf{u}[x, y]) = \text{pre}(\mathbf{v}[x, y])$, and the variable following immediately $\text{pre}(\mathbf{u}[x, y])$ is the same as the variable following immediately $\text{pre}(\mathbf{v}[x, y])$; and to check whether $\text{suf}(\mathbf{u}[x, y]) = \text{suf}(\mathbf{v}[x, y])$, and the variable preceding immediately $\text{suf}(\mathbf{u}[x, y])$ is the same as the variable preceding immediately $\text{suf}(\mathbf{v}[x, y])$. There are at most $\binom{|\text{con}(\mathbf{u})|}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time.

To check condition (III) of Theorem 4.2, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$, whether $\text{con}(\text{pren}(\mathbf{u}[x, y])) = \text{con}(\text{pren}(\mathbf{v}[x, y]))$ and $\text{occ}(z, \text{pren}(\mathbf{u}[x, y])) = \text{occ}(z, \text{pren}(\mathbf{v}[x, y]))$ for any $z \in \text{con}(\text{pren}(\mathbf{u}[x, y]))$; and whether $\text{con}(\text{sufn}(\mathbf{u}[x, y])) = \text{con}(\text{sufn}(\mathbf{v}[x, y]))$ and $\text{occ}(z, \text{sufn}(\mathbf{u}[x, y])) = \text{occ}(z, \text{sufn}(\mathbf{v}[x, y]))$ for any $z \in \text{con}(\text{sufn}(\mathbf{u}[x, y]))$. There are at most $\binom{|\text{con}(\mathbf{u})|}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time.

Therefore the decision problem $\text{CHECK-ID}(\text{baxt}_2, \sharp)$ belongs to the complexity class P.

For decision problem $\text{CHECK-ID}(\text{baxt}_3, \sharp)$, given any word identity $\mathbf{u} \approx \mathbf{v}$, by Theorem 4.2 and the above arguments, we only need to show that one can check whether the words \mathbf{u} and \mathbf{v} satisfy conditions (III)–(V) of Theorem 4.3 in polynomial time.

To check condition (III) of Theorem 4.3, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$, whether $\text{con}(\text{pren}(\mathbf{u}[x, y])) = \text{con}(\text{pren}(\mathbf{v}[x, y]))$, $\text{occ}(z, \text{pren}(\mathbf{u}[x, y])) = \text{occ}(z, \text{pren}(\mathbf{v}[x, y]))$ for any $z \in \text{con}(\text{pren}(\mathbf{u}[x, y]))$ and the variable following immediately $\text{pren}(\mathbf{u}[x, y])$ is the same as the variable following immediately $\text{pren}(\mathbf{v}[x, y])$; and to check whether $\text{con}(\text{sufn}(\mathbf{u}[x, y])) = \text{con}(\text{sufn}(\mathbf{v}[x, y]))$, $\text{occ}(z, \text{sufn}(\mathbf{u}[x, y])) = \text{occ}(z, \text{sufn}(\mathbf{v}[x, y]))$ for any $z \in \text{con}(\text{sufn}(\mathbf{u}[x, y]))$ and the variable preceding immediately $\text{pren}(\mathbf{u}[x, y])$ is the same

as the variable preceding immediately $\text{pren}(\mathbf{v}[x, y])$. There are at most $\binom{\text{con}(\mathbf{u})}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time.

To check condition (IV) of Theorem 4.3, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$, whether $\overrightarrow{\delta\overline{cc}}_y(x, \mathbf{u}) + \overrightarrow{\delta\overline{cc}}_y(x^*, \mathbf{u}) = \overrightarrow{\delta\overline{cc}}_y(x, \mathbf{v}) + \overrightarrow{\delta\overline{cc}}_y(x^*, \mathbf{v})$ and $\overrightarrow{\overline{cc}}_y(x, \mathbf{u}) + \overrightarrow{\overline{cc}}_y(x^*, \mathbf{u}) = \overrightarrow{\overline{cc}}_y(x, \mathbf{v}) + \overrightarrow{\overline{cc}}_y(x^*, \mathbf{v})$. There are at most $\binom{\text{con}(\mathbf{u})}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time.

To check condition (V) of Theorem 4.3, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$ with $x, x^* \neq y$ satisfying $\overrightarrow{\delta\overline{cc}}_y(y^*, \mathbf{u}) = 0$ whether $\overrightarrow{\delta\overline{cc}}_y(x, \mathbf{u}) = \overrightarrow{\delta\overline{cc}}_y(x, \mathbf{v})$ and $\overrightarrow{\delta\overline{cc}}_y(x^*, \mathbf{v}) = \overrightarrow{\delta\overline{cc}}_y(x^*, \mathbf{v})$; and for any $x, x^* \neq y \in \text{con}(\mathbf{u})$ satisfying $\overrightarrow{\overline{cc}}_y(y^*, \mathbf{u}) = 0$, whether $\overrightarrow{\overline{cc}}_y(x, \mathbf{u}) = \overrightarrow{\overline{cc}}_y(x, \mathbf{v})$ and $\overrightarrow{\overline{cc}}_y(x^*, \mathbf{v}) = \overrightarrow{\overline{cc}}_y(x^*, \mathbf{v})$. There are at most $\binom{\text{con}(\mathbf{u})}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time.

Therefore the decision problem $\text{CHECK-ID}(\text{baxt}_3, \#)$ belongs to the complexity class P.

For decision problem $\text{CHECK-ID}(\text{baxt}_n, \#)$ with $n \geq 4$, it suffices to check that, for any $x, y \in \text{con}(\mathbf{u})$, whether $\overrightarrow{\delta\overline{cc}}_x(y, \mathbf{u}) = \overrightarrow{\delta\overline{cc}}_x(y, \mathbf{v})$ and $\overrightarrow{\overline{cc}}_x(y, \mathbf{u}) = \overrightarrow{\overline{cc}}_x(y, \mathbf{v})$. There are at most $\binom{\text{con}(\mathbf{u})}{2}$ pairs of such x, y . Clearly, these can be completed in polynomial time. Therefore the decision problem $\text{CHECK-ID}(\text{baxt}_n, \#)$ with $n \geq 4$ belongs to the complexity class P. \square

REFERENCES

- [1] Auinger, K., Dolinka, I., Pervukhina, T.V., Volkov, M.V.: Unary enhancements of inherently non-finitely based semigroups. *Semigroup Forum* **89**, 41–51 (2014)
- [2] Auinger, K., Dolinka, I., Volkov, M.V.: Matrix identities involving multiplication and transposition. *J. Europ. Math. Soc.* **14**(3), 937–969 (2012)
- [3] Auinger, K., Dolinka, I., Volkov, M.V.: Equational theories of semigroups with involution. *J. Algebra* **369**, 203–225 (2012)
- [4] Abram, A., Reutenauer, C.: The stylic monoid. *Semigroup Forum* **105**(1), 1–45 (2022)
- [5] Aird, T., Ribeiro, D.: Tropical representations and identities of the stylic monoid. *Semigroup Forum* **106**, 1–23 (2023)
- [6] Almeida, J., Volkov, M.V., Goldberg, S.V.: Complexity of the identity checking problem for finite semigroups. *J. Math. Sci.* **158**(5), 605–614 (2009)
- [7] Burris, S., Sankappanavar, H.P.: *A Course in Universal Algebra*. Springer, New York, (1981)
- [8] Crvenković, S., Dolinka, I., Ésik, Z.: The variety of Kleene algebras with conversion is not finitely based. *Theor. Comput. Sci.* **230**, 235–245 (2000)
- [9] Chen, Y.Z., Hu, X., Kitov, N.V., Luo, Y.F., Sapir, O.: The finite basis problem for the monoid of two-by-two upper triangular tropical matrices. *Bull. Aust. Math. Soc.* **94**(1), 54–64 (2016)
- [10] Chen, Y.Z., Hu, X., Kitov, N.V., Luo, Y.F., Volkov, M.V.: Identities of the Kauffman monoid \mathcal{K}_3 . *Commun. Algebra* **48**(5), 1956–1968 (2020)
- [11] Cain, A.J., Johnson, M., Kambites, M., Malheiro, A.: Representations and identities of plactic-like monoids. *J. Algebra* **606**, 819–850 (2022)
- [12] Cain, A.J., Klein, G., Kubat, L., Malheiro, A., Okniński, J.: A note on identities in plactic monoids and monoids of upper-triangular tropical matrices. *ArXiv preprint* (2017) arXiv:1705.04596
- [13] Cain, A.J., Malheiro, A.: Identities in plactic, hypoplactic, sylvester, Baxter, and related monoids. *Electron. J. Combin.* **25**(3), Paper No.3.30 (2018)
- [14] Cain, A.J., Malheiro, A., Ribeiro, D.: Identities and bases in the sylvester and Baxter monoids. *J. Algebr. Comb.* (2023). <https://doi.org/10.1007/s10801-022-01202-6>
- [15] Cain, A.J., Malheiro, A., Ribeiro, D.: Identities and bases in the hypoplactic monoid. *Commun. Algebra* **50**(1), 146–162 (2022)
- [16] Daviaud, L., Johnson, M., Kambites, M.: Identities in upper triangular tropical matrix semigroups and the bicyclic monoid. *J. Algebra* **501**, 503–525 (2018)
- [17] Fulton, W.: *Young Tableaux: With Applications to Representation Theory and Geometry*, LMS Student Texts, vol. 35, Cambridge University Press, (1997)
- [18] Giraud, S.: Algebraic and combinatorial structures on pairs of twin binary trees. *J. Algebra* **360**, 115–157 (2012)
- [19] Gao, M., Zhang, W.T., Luo, Y.F.: The monoid of 2×2 triangular boolean matrices under skew transposition is non-finitely based. *Semigroup Forum* **100**, 153–168 (2020)

- [20] Gao, M., Zhang, W.T., Luo, Y.F.: A non-finitely based involution semigroup of order five. *Algebra Universalis* **81**(3), Paper No. 31 (2020)
- [21] Gao, M., Zhang, W.T., Luo, Y.F.: Varieties of involution J-trivial monoids with continuum many subvarieties. *Algebra Colloquium* (2022) (accepted)
- [22] Hivert, F., Novelli, J.C., Thibon, J.Y.: The algebra of binary search trees. *Theoret. Comput. Sci.* **339**(1), 129–165 (2005)
- [23] Hivert, F., Novelli, J.C., Thibon, J.Y.: Commutative combinatorial Hopf algebras. *J. Algebraic Combin.* **28**(1), 65–95 (2008)
- [24] Han, B.B., Zhang, W.T., Luo, Y.F.: Equational theories of upper triangular tropical matrix semigroups. *Algebra Universalis* **82**(3), Paper No. 44 (2021)
- [25] Han, B.B., Zhang, W.T.: Finite basis problems for stalactic, taiga, sylvester and baxter monoids. *J. Algebra Appl.* (2022). <https://doi.org/10.1142/S0219498823502043>
- [26] Izhakian, Z., Margolis, S.W.: Semigroup identities in the monoid of two-by-two tropical matrices. *Semigroup Forum* **80**(2), 191–218 (2010)
- [27] Johnson, M., Kambites, M.: Tropical representations and identities of plactic monoids. *Trans. Amer. Math. Soc.* **374**(6), 4423–4447 (2021)
- [28] Jackson, M.: Finite semigroups whose varieties have uncountably many subvarieties. *J. Algebra* **228**(2), 512–535 (2000)
- [29] Jackson, M., McKenzie, R.: Interpreting graph colorability in finite semigroups. *Internat. J. Algebra Comput.* **16**(1), 119–140 (2006)
- [30] Jackson, M., Volkov, M.V.: The algebra of adjacency patterns: Rees matrix semigroups with reversion. in: Blass A., Dershowitz N., Reisig W. (eds.), *Fields of Logic and Computation*, Lecture Notes in Computer Science, Springer, Berlin, pp. 414–443 (2010)
- [31] Kharlampovich, O.G., Sapir, M.V.: Algorithmic problems in varieties. *Internat. J. Algebra Comput.* **5**(4–5), 379–602 (1995)
- [32] Krob, D., Thibon, J. Y.: Noncommutative Symmetric Functions IV: Quantum Linear Groups and Hecke Algebras at $q = 0$. *J. Algebr. Comb.* **6**(4), 339–376 (1997)
- [33] Kitov, N.V., Volkov, M.V.: Identities of the Kauffman monoid \mathcal{K}_4 and of the Jones monoid \mathcal{J}_4 . In: Blass, A., Cegielski, P., Dershowitz, N., Droste, M., Finkbeiner, B. (eds) *Fields of Logic and Computation III*. Lecture Notes in Computer Science, vol 12180. Springer, Cham. (2020)
- [34] Lothaire, M.: *Algebraic Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, vol. 90, Cambridge University Press, 2002
- [35] Lee, E.W.H.: Finite involution semigroups with infinite irredundant bases of identities. *Forum Math.* **28**, 587–607 (2016)
- [36] Lee, E.W.H.: Finitely based finite involution semigroups with non-finitely based reducts. *Quaest. Math.* **39**(2), 217–243 (2016)
- [37] Lee, E.W.H.: Equational theories of unstable involution semigroups. *Electron. Res. Announc. Math. Sci.* **24**, 10–20 (2017)
- [38] Lee, E.W.H.: A sufficient condition for the absence of irredundant bases. *Houston J. Math.* **44**, 399–411 (2018)
- [39] Lee, E.W.H.: Non-finitely based finite involution semigroups with finitely based semigroup reducts. *Korean J. Math.* **27**(1), 53–62 (2019)
- [40] Lee, E.W.H.: Varieties of involution monoids with extreme properties. *Quart. J. Math.* **70**, 1157–1180 (2019)
- [41] Lee, E.W.H.: *Contributions to the Theory of Varieties of Semigroups*. Dr. Sci. thesis, National Research University Higher School of Economics, Moscow (2020)
- [42] Lascoux, A., Schützenberger, M.P.: Sur une conjecture de H. O. Foulkes. *C. R. Acad. Sci. Paris Sér. A–B* **286**(7), A323–A324 (1978)
- [43] Lascoux, A., Schützenberger, M.P.: Le monoïde plaxique. In: A. de Luca (ed.), *Noncommutative Structures in Algebra and Geometric Combinatorics*, Proceedings of the Colloquium Held at Arco Felice, Naples, July 24–26, 1978. *Quaderni della Ricerca Scientifica*, vol. 109, pp.129–156. Consiglio Nazionale delle Ricerche, Roma (1981)
- [44] Lascoux, A., Schützenberger, M.P.: Schubert polynomials and the Littlewood-Richardson rule. *Lett. Math. Phys.* **10**(2–3), 111–124 (1985)
- [45] Lascoux, A., Schützenberger, M. P.: Tableaux and noncommutative Schubert polynomials. *Funct. Anal. Its. Appl.* **23**(3), 223–225 (1990)
- [46] Murskiĭ, V.L.: Examples of varieties of semigroups, *Math. Notes* **3**(6), 423–427 (1968)
- [47] Macdonald, I.: *Symmetric Functions and Hall Polynomials*. Clarendon Press, Oxford, (2008)
- [48] Novelli, J.C.: On the hypoplactic monoid. *Discrete Math.* **217**(1–3), 315–336 (2000)
- [49] Perkins, P.: Bases for equational theories of semigroups. *J. Algebra* **11**, 298–314 (1969)
- [50] Priez, J.B.: Lattice of combinatorial Hopf algebras: binary trees with multiplicities. In: Goupil, A., Schaeffer, G. (eds.) *25th International Conference on Formal Power Series and*

Algebraic Combinatorics (FPSAC 2013), DMTCS Proceedings, vol. AS, pp.1137–1148. Discrete Mathematics and Theoretical Computer Science, Paris (2013)

- [51] Volkov, M.V.: The finite basis problem for finite semigroups. *Sci. Math. Jpn.* **53**(1), 171–199 (2001)
- [52] Volkov, M.V.: Identities of the stylic monoid. *Semigroup Forum* **105**(1), 345–349 (2022)

¹ SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, PR CHINA

Email address: hanbb19@lzu.edu.cn, zhangwt@lzu.edu.cn, luoyf@lzu.edu.cn

² SCHOOL OF MATHEMATICAL SCIENCES, INNER MONGOLIA UNIVERSITY, HOHHOT, INNER MONGOLIA 010021, PR CHINA

Email address: zhjxing@imu.edu.cn