

# Heat currents in qubit systems

Hans C. Fogedby\*

*Department of Physics and Astronomy,  
University of Aarhus, Ny Munkegade*

## Abstract

There is a current interest in quantum thermodynamics in the context of open quantum systems. An important issue is the consistency of quantum thermodynamics, in particular the second law of thermodynamics, i.e., the flow of heat from a hot reservoir to a cold reservoir. Recent emphasis has been on composite system. Here we discuss two cases, namely as an example a single qubit and as a simple composite system two coupled qubits driven by two heat reservoirs at different temperatures, respectively. We present explicit expressions for the heat currents in agreement with the second law of thermodynamics. The analysis is carried out in the Born-Markov approximation.

arXiv:2301.13544v1 [quant-ph] 31 Jan 2023

---

\*Electronic address: fogedby@phys.au.dk

## I. INTRODUCTION

There is a current interest in quantum thermodynamics [1, 1–9]. An important issue here is the emergence of thermodynamics from a microscopic point of view and in particular the validity of the second law of thermodynamics. In the Clausius formulation the second law implies that heat flows from a hot reservoir to a cold reservoir and this essential property should also hold in the quantum regime. In non equilibrium quantum statistical mechanics the instantaneous heat current is subject to both statistical and quantum fluctuations, however, the mean heat current should obey the second law of thermodynamics.

The natural framework for discussing these issues is the theory of open quantum systems [10, 11]. Here the focus is on a small controllable quantum system interacting with an environment which can be either radiation fields or heat baths. In order to obtain a steady heat current we must drive the quantum system by several heat reservoirs, typically two reservoirs, maintained at different temperatures.

In the context of energy transport in mesoscopic and atomic structures one typically deals with composite systems [1, 12, 13] and the issue of the validity of quantum thermodynamics and the appropriate master equation description has been discussed extensively, see e.g. [14, 15] and references cited therein. In [16] this issue was discussed and resolved for a composite system of two coupled oscillators.

The present work is to a large extent motivated by the study in [16]. We discuss two driven open quantum systems and derive explicit expressions for the heat currents. The first system serving as an illustration is the simple case of a single qubit driven by two heat reservoirs. The second composite system is the more complex case of two coupled qubits driven by two reservoirs. Within a Born-Markov approximation we derive explicit expressions for the heat currents. The heat currents have a similar structure and are both in agreement with the second law of thermodynamics, i.e., heat flows from a hot reservoir to a cold reservoir.

The paper is organised as follows. In Sec. II we set up the general scheme. In Sec. III A we derive the heat currents for a single qubit. In Sec. III B we derive the heat currents for two coupled qubits. In Sec. IV A we discuss the single qubit case, in Sec. IV B the coupled qubits and in Sec. IV C the off diagonal density matrix elements and the local master equation issue. Details are deferred to Appendix V.

## II. GENERAL ANALYSIS

The density operator for an open quantum system  $S$  characterised by the Hamiltonian  $H_S$  interacting with several heat reservoirs is in the Markov approximation governed by a master equation of the form [10, 11, 17]

$$\frac{d\rho(t)}{dt} = -i[H_S, \rho(t)] + L\rho(t). \quad (2.1)$$

Here the super operator  $L$  has the structure  $L = K + i\Delta$ , where  $\Delta$  is a Lamb type correction to the energy levels and the dissipative kernel  $K$  characterises the relaxation of the system. In the following we assume that the shift  $\Delta$  is incorporated in  $H$  and focus on the dissipative kernel  $K$ . In the energy basis of  $H_S$ ,  $H_S|n\rangle = E_n|n\rangle$ , the master equation in (2.1) takes the form

$$\frac{d\rho(t)_{pp'}}{dt} = -iE_{pp'}\rho(t)_{pp'} + \sum_{qq'} K_{pp',qq'}\rho(t)_{qq'}, \quad (2.2)$$

where  $E_{pp'} = E_p - E_{p'}$ ,  $\rho(t)_{pp'} = \langle p|\rho(t)|p'\rangle$  and  $K_{pp',qq'} = \langle p|\langle q'|K|q\rangle|p'\rangle$ .

By inspection of (2.2) it follows that the steady state of the system  $S$  is determined by the diagonal elements  $\rho_{pp}$  of the density matrix  $\rho_{pp'}$ , yielding the populations of the energy levels  $E_n$ . The off-diagonal terms vanish in the long time limit; this issue is discussed in more detail in Sec. IV C. We thus infer, including the trace condition  $\sum_n \rho_n = 1$ , the scheme

$$\sum_p K_{np}\rho_p = 0, \quad (2.3)$$

$$\sum_p \rho_p = 1; \quad (2.4)$$

we have used the notation  $K_{pp,qq} \equiv K_{pq}$  and  $\rho_{pp} \equiv \rho_p$ , where in the following  $\rho_p$  denotes the stationary diagonal density matrix elements.

The mean energy of the system is given by  $E(t) = \sum_n E_n\rho(t)_{nn}$  and we obtain by insertion of (2.2) for the mean heat current  $q(t) = dE(t)/dt$  in the steady state

$$q = \sum_{np} E_n K_{np}\rho_p. \quad (2.5)$$

For a single heat reservoir the system equilibrates and it follows directly from (2.3) and (2.5) that the heat current  $q$  vanishes. However, in the presence of several heat reservoirs maintained at different temperatures a steady state heat current will be generated.

In the case of two uncorrelated heat reservoirs  $A$  and  $B$  the kernel  $K$  in (2.2) is composed of two parts and we have

$$K_{pq} = K_{pq}^A + K_{pq}^B. \quad (2.6)$$

In this case the steady state density operator and total current are determined by the conditions

$$\sum_p (K_{np}^A + K_{np}^B) \rho_p = 0, \quad (2.7)$$

$$\sum_p \rho_p = 1, \quad (2.8)$$

$$q = \sum_{np} E_n (K_{np}^A + K_{np}^B) \rho_p. \quad (2.9)$$

The vanishing total current has components associated with each bath, i.e.,

$$q = q^A + q^B = 0, \quad (2.10)$$

$$q^A = \sum_{np} E_n K_{np}^A \rho_p, \quad (2.11)$$

$$q^B = \sum_{np} E_n K_{np}^B \rho_p. \quad (2.12)$$

We infer that the heat flux from the reservoir  $A$  to the system  $S$  given by  $q^A$  equals the heat flux  $q^A = -q^B$  flowing from the system  $S$  into the reservoir  $B$ , expressing energy conservation, i.e. the first law of thermodynamics. Regarding the second law of thermodynamics the issue is to show that a positive heat current flows from the hot heat reservoir to the cold heat reservoir.

From the above it follows that in order to evaluate the heat currents for a given open quantum system driven by two reservoirs we need three components: The energy spectrum of the system  $E_n$ , the diagonal kernel elements  $K_{np}^A$  and  $K_{np}^B$ , and the density operator  $\rho_p$ .

### III. QUBITS DRIVEN BY TWO HEAT RESERVOIRS

#### A. Single qubit

Referring for details to Appendix V B, we consider first the simple case of a single qubit driven by two heat reservoirs  $A$  and  $B$ . Here the energy or heat flows between the reservoirs via the qubit; the configuration is depicted in Fig. 1.

In a Pauli matrix basis [18] the Hamiltonian and coupling to the reservoirs are given by

$$H_S = \frac{\omega_0}{2} \sigma^z \quad (3.1)$$

$$H_{SA} = \sigma^+ A + \sigma^- A^\dagger, \quad (3.2)$$

$$H_{SB} = \sigma^+ B + \sigma^- B^\dagger, \quad (3.3)$$

$$A = \sum_k \lambda_k^A a_k^\dagger a_k, \quad (3.4)$$

$$B = \sum_k \lambda_k^B b_k^\dagger b_k, \quad (3.5)$$

where the levels  $|+\rangle$  and  $|-\rangle$  have energies  $E_+ = \omega_0/2$  and  $E_- = -\omega_0/2$ , corresponding to the energy splitting  $\omega_0$ . The qubit is driven by two heat reservoirs in the rotating wave approximation (RWA). The bath operators  $A$  and  $B$  sample the bath modes with wavenumber  $k$ , frequencies  $\omega_k^{A,B}$ , and strength  $\lambda_k^{A,B}$ ;  $a_k$  and  $b_k$  are the Bose operators. The heat reservoirs are characterised by the spectral functions and Planck distributions

$$g_{A,B}(\omega) = 2\pi \sum_k (\lambda_k^{A,B})^2 \delta(\omega - \omega_k^{A,B}), \quad (3.6)$$

$$n_{A,B}(\omega) = \frac{1}{\exp(\omega/T_{A,B}) - 1}. \quad (3.7)$$

With the abbreviations  $g_{A,B} = g_{A,B}(\omega_0)$  and  $n_{A,B} = n_{A,B}(\omega_0)$  we derive the dissipative kernel elements

$$K_{+,+}^{A,B} = -g_{A,B}(1 + n_{A,B}), \quad (3.8)$$

$$K_{-,+}^{A,B} = +g_{A,B}(1 + n_{A,B}), \quad (3.9)$$

$$K_{+,-}^{A,B} = +g_{A,B}n_{A,B}, \quad (3.10)$$

$$K_{-,-}^{A,B} = -g_{A,B}n_{A,B}, \quad (3.11)$$

yielding according to (2.3) and (2.4) in Sec. II the diagonal density matrix elements. For the populations of the energy levels we have

$$\rho_+ = \frac{g_A n_A + g_B n_B}{g_A(1 + 2n_A) + g_B(1 + 2n_B)}, \quad (3.12)$$

$$\rho_- = \frac{g_A(1 + n_A) + g_B(1 + n_B)}{g_A(1 + 2n_A) + g_B(1 + 2n_B)}. \quad (3.13)$$

Using (2.11) and (2.12) in Sec. II we readily find the steady heat currents from reservoir  $A$

and from reservoir  $B$  to the qubit

$$q^A = \frac{g_A g_B \omega_0}{g_A(1 + 2n_A) + g_B(1 + 2n_B)} [n_A - n_B], \quad (3.14)$$

$$q^B = \frac{g_A g_B \omega_0}{g_A(1 + 2n_A) + g_B(1 + 2n_B)} [n_B - n_A]. \quad (3.15)$$

Further discussions of these results are deferred to Sec. IV A.

## B. Coupled qubits

Referring for details to Appendix V C, we consider the case of two coupled qubits driven by two heat reservoirs as a model of a simple composite system. Heat reservoir  $A$  drives qubit 1 and heat reservoir  $B$  drives qubit 2. Here the heat flows between the reservoirs and the qubits transmitted across the system by the qubit coupling; the configuration is depicted in Fig. 2.

In a Pauli matrix basis [18] the Hamiltonian and coupling to the reservoirs are given by

$$H_S = \frac{\omega_1}{2} \sigma_1^z + \frac{\omega_2}{2} \sigma_2^z + \lambda(\sigma_1^+ \sigma_2^- + \sigma_1^- \sigma_2^+). \quad (3.16)$$

$$H_{SA} = \sigma_1^+ A + \sigma_1^- A^\dagger, \quad (3.17)$$

$$H_{SB} = \sigma_2^+ B + \sigma_2^- B^\dagger, \quad (3.18)$$

where  $\omega_1$  and  $\omega_2$  are the energy splittings of the unperturbed qubits and  $\lambda$  the coupling strength of the flip-flop interaction, i.e., the dipole coupling in the RWA. For the composite system we have the unperturbed states  $|--\rangle$ ,  $|+-\rangle$ ,  $|+ -\rangle$  and  $|++\rangle$  with energies  $-\omega_m$ ,  $-\Delta\omega$ ,  $\Delta\omega$ , and  $\omega_m$ , respectively, where

$$\omega_m = \frac{\omega_1 + \omega_2}{2}, \quad (3.19)$$

$$\Delta\omega = \frac{\omega_1 - \omega_2}{2}; \quad (3.20)$$

we assume  $\Delta\omega > 0$ .

Diagonalising  $H_S$  yields the four states

$$|1\rangle = |--\rangle, \quad (3.21)$$

$$|2\rangle = -\beta|+ -\rangle + \alpha| - +\rangle, \quad (3.22)$$

$$|3\rangle = \alpha|+ -\rangle + \beta| - +\rangle, \quad (3.23)$$

$$|4\rangle = |++\rangle, \quad (3.24)$$

with energies

$$E_1 = -\omega_m, \quad (3.25)$$

$$E_2 = -\sqrt{(\Delta\omega)^2 + \lambda^2}, \quad (3.26)$$

$$E_3 = \sqrt{(\Delta\omega)^2 + \lambda^2}, \quad (3.27)$$

$$E_4 = \omega_m. \quad (3.28)$$

The expansion coefficients and relevant energy shifts  $\omega_+ = E_{42} = E_{31}$  and  $\omega_- = E_{43} = E_{21}$  are given by

$$\alpha = \cos \theta/2, \quad (3.29)$$

$$\beta = \sin \theta/2, \quad (3.30)$$

$$\tan \theta = 2\lambda/(\omega_1 - \omega_2), \quad 0 < \theta < \pi/2, \quad (3.31)$$

$$\omega_+ = \omega_m + \sqrt{(\Delta\omega)^2 + \lambda^2}, \quad (3.32)$$

$$\omega_- = \omega_m - \sqrt{(\Delta\omega)^2 + \lambda^2}. \quad (3.33)$$

For vanishing coupling between the qubits,  $\lambda \rightarrow 0$ , we have  $\alpha \rightarrow 1$ ,  $\beta \rightarrow 0$ ,  $\omega_+ \rightarrow \omega_1$ , and  $\omega_- \rightarrow \omega_2$ ; moreover, assuming  $\omega_{\pm} > 0$  we require  $\lambda < \sqrt{\omega_1\omega_2}$ .

Introducing the notation

$$A_{++} = \alpha^2 g_A(\omega_+)(1 + n_A(\omega_+)), \quad (3.34)$$

$$A_{+-} = \beta^2 g_A(\omega_-)(1 + n_A(\omega_-)), \quad (3.35)$$

$$A_{-+} = \alpha^2 g_A(\omega_+)n_A(\omega_+), \quad (3.36)$$

$$A_{--} = \beta^2 g_A(\omega_-)n_A(\omega_-), \quad (3.37)$$

$$B_{++} = \beta^2 g_B(\omega_+)(1 + n_B(\omega_+)), \quad (3.38)$$

$$B_{+-} = \alpha^2 g_B(\omega_-)(1 + n_B(\omega_-)), \quad (3.39)$$

$$B_{-+} = \beta^2 g_B(\omega_+)n_B(\omega_+), \quad (3.40)$$

$$B_{--} = \alpha^2 g_B(\omega_-)n_B(\omega_-), \quad (3.41)$$

we find the kernel elements

$$K^A = \begin{pmatrix} -A_{-+} - A_{--} & A_{+-} & A_{++} & 0 \\ A_{--} & -A_{-+} - A_{+-} & 0 & A_{++} \\ A_{-+} & 0 & -A_{++} - A_{--} & A_{+-} \\ 0 & A_{-+} & A_{--} & -A_{++} - A_{+-} \end{pmatrix}, \quad (3.42)$$

$$K^B = \begin{pmatrix} -B_{-+} - B_{--} & B_{+-} & B_{++} & 0 \\ B_{--} & -B_{-+} - B_{+-} & 0 & B_{++} \\ B_{-+} & 0 & -B_{++} - B_{--} & B_{+-} \\ 0 & B_{-+} & B_{--} & -B_{++} - B_{+-} \end{pmatrix}. \quad (3.43)$$

Introducing the matrices

$$\rho_{\pm}^C \propto \begin{pmatrix} C_{+\pm} & 0 \\ 0 & C_{-\pm} \end{pmatrix}, \quad C = A, B \quad (3.44)$$

and using (2.3) and (2.4) in Sec. II we find the diagonal density matrix

$$\rho = \frac{(\rho_+^A + \rho_+^B) \otimes (\rho_-^A + \rho_-^B)}{\sum_n \rho_n}, \quad (3.45)$$

where  $\otimes$  denotes a direct matrix product. Explicitly the elements are given by

$$\rho_1 = \frac{(A_{++} + B_{++})(A_{+-} + B_{+-})}{(A_{++} + B_{++} + A_{-+} + B_{-+})(A_{--} + B_{--} + A_{+-} + B_{+-})}, \quad (3.46)$$

$$\rho_2 = \frac{(A_{++} + B_{++})(A_{--} + B_{--})}{(A_{++} + B_{++} + A_{-+} + B_{-+})(A_{--} + B_{--} + A_{+-} + B_{+-})}, \quad (3.47)$$

$$\rho_3 = \frac{(A_{-+} + B_{-+})(A_{+-} + B_{+-})}{(A_{++} + B_{++} + A_{-+} + B_{-+})(A_{--} + B_{--} + A_{+-} + B_{+-})}, \quad (3.48)$$

$$\rho_4 = \frac{(A_{-+} + B_{-+})(A_{--} + B_{--})}{(A_{++} + B_{++} + A_{-+} + B_{-+})(A_{--} + B_{--} + A_{+-} + B_{+-})}; \quad (3.49)$$

note that  $\det(K^{A,B}) = 0$ , however, imposing the trace constraint  $\sum_n \rho_n = 1$  and eliminating a component we readily obtain the solution above, satisfying the trace condition. In a similar manner we obtain using (2.11) and (2.12) in Sec. II the heat currents

$$q^A = \frac{B_{++}A_{-+} - B_{-+}A_{++}}{A_{++} + B_{++} + A_{-+} + B_{-+}}\omega_+ + \frac{B_{+-}A_{--} - B_{--}A_{+-}}{A_{--} + B_{--} + A_{+-} + B_{+-}}\omega_-, \quad (3.50)$$

$$q^B = \frac{A_{++}B_{-+} - A_{-+}B_{++}}{A_{++} + B_{++} + A_{-+} + B_{-+}}\omega_+ + \frac{A_{+-}B_{--} - A_{--}B_{+-}}{A_{--} + B_{--} + A_{+-} + B_{+-}}\omega_-, \quad (3.51)$$

and by insertion of  $A_{\pm\pm}$  and  $B_{\pm\pm}$

$$q^A = q_+^A + q_-^A, \quad (3.52)$$

$$q_+^A = \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) (1 + 2n_A(\omega_+)) + \beta^2 g_B(\omega_+) (1 + 2n_B(\omega_+))} [n_A(\omega_+) - n_B(\omega_+)], \quad (3.53)$$

$$q_-^A = \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) (1 + 2n_A(\omega_-)) + \alpha^2 g_B(\omega_-) (1 + 2n_B(\omega_-))} [n_A(\omega_-) - n_B(\omega_-)]. \quad (3.54)$$



and

$$q^B = q_+^B + q_-^B, \quad (3.55)$$

$$q_+^B = \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) (1 + 2n_A(\omega_+)) + \beta^2 g_B(\omega_+) (1 + 2n_B(\omega_+))} [n_B(\omega_+) - n_A(\omega_+)], \quad (3.56)$$

$$q_-^B = \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) (1 + 2n_A(\omega_-)) + \alpha^2 g_B(\omega_-) (1 + 2n_B(\omega_-))} [n_B(\omega_-) - n_A(\omega_-)]. \quad (3.57)$$

Further discussions of these results are deferred Sec. IV B.

## IV. DISCUSSION

Here we discuss the above results for the driven single qubit and the driven coupled qubits. First we note that the expressions for the heat currents in (3.14 - 3.15) and (3.52 - 3.57) exhibit a similar structure. Since  $q^A + q^B = 0$  energy is conserved consistent with the general discussion in Sec. II. Expressing the currents in the form  $q^A = F(n^A - n^B)$  and  $q^B = F(n^B - n^A)$  and noting that the prefactor  $F$  is positive and that  $n^A > n^B$  for  $T^A > T^B$  we infer that a positive heat current flows from the hot reservoir  $A$  to the system and back to reservoir  $B$  in accordance with the second law of thermodynamics, as illustrated in Figs. 1 and 2.

### A. Single qubit

#### 1. Density matrix

A few comments regarding the density matrix. Introducing the ratio  $r = \rho_+/\rho_-$  we have from (3.12) and (3.13)

$$r = \frac{g_A n_A + g_B n_B}{g_A (1 + n_A) + g_B (1 + n_B)}. \quad (4.1)$$

In general the ratio  $r$  depends on both the spectral densities  $g_{A,B}$ , i.e., the structure of the reservoirs, and the temperatures  $T_{A,B}$ . Turning off for example reservoir  $B$  by setting  $g_B = 0$  we obtain the well-studied case of a single qubit driven by a single reservoir, see e.g. [10]. The ratio  $r$  does not depend on the spectral strength and we obtain  $r = \exp(-\omega_0/T_A)$ , i.e., the Boltzmann distribution. For identical reservoirs with the same spectral densities, i.e.,  $g_A = g_B = g$ , we obtain  $r = (n_A + n_B)/(2 + n_A + n_B)$ . At high

temperatures  $r \rightarrow 1 - \omega_0/T$ , where  $T = (T_A + T_B)/2$  is the mean temperature of the combined reservoirs and the energy levels become equally populated. In the low temperature limit  $r \rightarrow (\exp(-\omega_0/T_A) + \exp(-\omega_0/T_B))/2$ . Introducing the temperature bias  $\Delta = T_A - T_B$  we have  $r \rightarrow \exp(-\omega_0/T) \cosh(2\omega_0/\Delta)$ , i.e., a correction to the Boltzmann factor. At high temperature,  $T \gg \omega_0$ , we have  $\rho_+ \rightarrow 1/2$  and  $\rho_- \rightarrow 1/2$ , i.e., equal population; at low temperature,  $T \ll \omega_0$ , we have  $\rho_+ \rightarrow 0$  and  $\rho_- \rightarrow 1$ , i.e., population of the lowest level (ground state).

## 2. Currents

Regarding the currents given by (3.14) and (3.15) we note that they depend on both  $g^A$  and  $g^B$ . Removing for example reservoir  $B$  by setting  $g^B = 0$  the currents vanish and the qubit coupled only to reservoir  $A$  equilibrates with temperature  $T_A$ . In the high temperature limit for  $T_{A,B} \gg \omega_0$  we obtain expanding (3.14) and (3.15) the classical results

$$q^A \sim \frac{1}{2} \frac{g_A g_B \omega_0}{g_A T_A + g_B T_B} [T_A - T_B], \quad (4.2)$$

$$q^B \sim \frac{1}{2} \frac{g_A g_B \omega_0}{g_A T_A + g_B T_B} [T_B - T_A], \quad (4.3)$$

at low temperatures for  $T_{A,B} \ll \omega_0$  the vanishing currents

$$q^A \sim \frac{g_A g_B \omega_0}{g_A + g_B} [e^{-\omega_0/T_A} - e^{-\omega_0/T_B}], \quad (4.4)$$

$$q^B \sim \frac{g_A g_B \omega_0}{g_A + g_B} [e^{-\omega_0/T_B} - e^{-\omega_0/T_A}]. \quad (4.5)$$

## B. Coupled qubits

### 1. Density matrix

Here we discuss first the expressions for the density matrix or level populations in (3.46 - 3.49). In the high temperature limit  $T_A, T_B \gg \omega_+, \omega_-$  we have  $A_{\pm\pm} \rightarrow g_A(\omega_{\pm})T_A/\omega_{\pm}$  and  $B_{\pm\pm} \rightarrow g_B(\omega_{\pm})T_B/\omega_{\pm}$  implying  $\rho_n \rightarrow 1/4$ , i.e., equal population of the four levels. At low temperature  $T_A, T_B \ll \omega_+, \omega_-$  we have  $\rho_1 \rightarrow 1$  and  $\rho_2, \rho_3, \rho_4 \rightarrow 0$ , i.e., occupation of the

lowest level. Finally, for  $\lambda = 0$  we have  $A_{+-} = A_{--} = 0$  and  $B_{++} = B_{-+} = 0$ , i.e.,

$$\rho_1 = \frac{A_{++}B_{+-}}{(A_{++} + A_{-+})(B_{--} + B_{+-})}, \quad (4.6)$$

$$\rho_2 = \frac{A_{++}B_{--}}{(A_{++} + A_{-+})(B_{--} + B_{+-})}, \quad (4.7)$$

$$\rho_3 = \frac{A_{-+}B_{+-}}{(A_{++} + A_{-+})(B_{--} + B_{+-})}, \quad (4.8)$$

$$\rho_4 = \frac{A_{-+}B_{--}}{(A_{++} + A_{-+})(B_{--} + B_{+-})}. \quad (4.9)$$

By inspection we conclude that the qubits decouple and we obtain for the density operator the direct product  $\rho = \rho^A \otimes \rho^B$ , where

$$\rho^A = \frac{1}{g_A(\omega_1)(1 + 2n_A(\omega_1))} \begin{pmatrix} g_A(\omega_1)(1 + n_A(\omega_1)) & 0 \\ 0 & g_A(\omega_1)n_A(\omega_1) \end{pmatrix}, \quad (4.10)$$

and

$$\rho^B = \frac{1}{g_B(\omega_2)(1 + 2n_B(\omega_2))} \begin{pmatrix} g_B(\omega_2)(1 + n_B(\omega_2)) & 0 \\ 0 & g_B(\omega_2)n_B(\omega_2) \end{pmatrix}, \quad (4.11)$$

are the equilibrium density operators for qubit 1 coupled to reservoir  $A$  and qubit 2 coupled to reservoir  $B$ , respectively, see e.g. [10].

## 2. Currents

The heat currents  $q^A$  and  $q^B$  in (3.52 - 3.57) fall in two parts associated with the energy shifts  $\omega_+$  and  $\omega_-$ . For vanishing coupling,  $\lambda = 0$ , we have  $\alpha = 1$  and  $\beta = 1$ , the qubits are uncoupled and the currents vanish. Likewise, quenching e.g. reservoir  $B$  by setting  $g_B = 0$  we obtain a vanishing current. In the high temperature limit for  $T_A, T_B \gg \omega_+, \omega_-$  we have

$$q_+^A = \frac{1}{2} \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) T_A + \beta^2 g_B(\omega_+) T_B} [T_A - T_B], \quad (4.12)$$

$$q_-^A = \frac{1}{2} \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) T_A + \alpha^2 g_B(\omega_-) T_B} [T_A - T_B]. \quad (4.13)$$

and

$$q_+^B = \frac{1}{2} \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) T_A + \beta^2 g_B(\omega_+) T_B} [T_B - T_A], \quad (4.14)$$

$$q_-^B = \frac{1}{2} \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) T_A + \alpha^2 g_B(\omega_-) T_B} [T_B - T_A], \quad (4.15)$$

at low temperatures  $T_A, T_B \ll \omega_+, \omega_-$ ,

$$q_+^A = \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) + \beta^2 g_B(\omega_+)} [e^{-\omega_+/T_A} - e^{-\omega_+/T_B}], \quad (4.16)$$

$$q_-^A = \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) + \alpha^2 g_B(\omega_-)} [e^{-\omega_-/T_A} - e^{-\omega_-/T_B}]. \quad (4.17)$$

and

$$q_+^B = \frac{(\alpha\beta)^2 g_A(\omega_+) g_B(\omega_+) \omega_+}{\alpha^2 g_A(\omega_+) + \beta^2 g_B(\omega_+)} [e^{-\omega_+/T_B} - e^{-\omega_+/T_A}], \quad (4.18)$$

$$q_-^B = \frac{(\alpha\beta)^2 g_A(\omega_-) g_B(\omega_-) \omega_-}{\beta^2 g_A(\omega_-) + \alpha^2 g_B(\omega_-)} [e^{-\omega_-/T_B} - e^{-\omega_-/T_A}]. \quad (4.19)$$

### C. Off diagonal terms - Local master equation

#### 1. Off diagonal density matrix elements

In Sec. II we argued heuristically that the steady state is determined by the diagonal density matrix elements  $\rho_p \equiv \rho_{pp}$ . Here we demonstrate that this assumption holds in the two-qubit case. The master equation in the basis of the system Hamiltonian  $H_S$  has the form given in (2.2). In the two-qubit case we have using the assignments in (5.19) and (5.20) and inserting in (5.2) the non vanishing kernel elements

$$K_{11,qq'} = (K_{11,11}, K_{11,22}, K_{11,33}, K_{11,23}, K_{11,32}), \quad (4.20)$$

$$K_{22,qq'} = (K_{22,11}, K_{22,22}, K_{22,44}), \quad (4.21)$$

$$K_{33,qq'} = (K_{33,11}, K_{33,33}, K_{33,44}), \quad (4.22)$$

$$K_{44,qq'} = (K_{44,22}, K_{44,33}, K_{44,44}, K_{44,23}, K_{44,32}), \quad (4.23)$$

$$K_{23,qq'} = (K_{23,11}, K_{23,44}, K_{23,23}), \quad (4.24)$$

$$K_{32,qq'} = (K_{32,11}, K_{32,44}, K_{32,32}). \quad (4.25)$$

Using the symmetries  $K_{pp,23} = K_{pp,32} = A_p$ ,  $K_{23,pp} = K_{32,pp} = B_p$  for  $p = 1, 4$ ,  $K_{23,23} = K_{32,32} = C$ , and  $E_{23} = -E_{32} = E$  we obtain the coupled equations of motion

$$\frac{d\rho_{11}}{dt} = K_{11,11}\rho_{11} + K_{11,22}\rho_{22} + K_{11,33}\rho_{33} + A_1(\rho_{23} + \rho_{32}), \quad (4.26)$$

$$\frac{d\rho_{22}}{dt} = K_{22,11}\rho_{11} + K_{22,22}\rho_{22} + K_{22,44}\rho_{44}, \quad (4.27)$$

$$\frac{d\rho_{33}}{dt} = K_{33,11}\rho_{11} + K_{33,33}\rho_{33} + K_{33,44}\rho_{44}, \quad (4.28)$$

$$\frac{d\rho_{44}}{dt} = K_{44,22}\rho_{22} + K_{44,33}\rho_{33} + K_{44,44}\rho_{44} + A_4(\rho_{23} + \rho_{32}), \quad (4.29)$$

$$\left(\frac{d}{dt} + iE - C\right)\rho_{23} = B_1\rho_{11} + B_4\rho_{44}, \quad (4.30)$$

$$\left(\frac{d}{dt} - iE - C\right)\rho_{32} = B_1\rho_{11} + B_4\rho_{44}. \quad (4.31)$$

Since  $C = -(1/2) \sum_{p=\pm, q=\pm} (A_{pq} + B_{pq})$  is negative it follows from inspection of (4.30) and (4.31) that  $\rho_{23}$  and  $\rho_{32}$  fall off exponentially with a time constant given by  $1/C$  and we conclude that the steady state is determined by the diagonal kernel elements  $K_{nn,pp}$ , i.e., the analysis in Sec. II.

## 2. Local master equation

In the context of open composite quantum systems there has been a recent discussion regarding the connection between a global master equation approach as applied in the present paper and a local master equation approach [14, 16] in which the interactions in the composite system is ignored in the dissipative kernels. This issue was discussed and resolved in [16] for the case of coupled quantum oscillators driven by two heat reservoirs.

In a Bose operator description two coupled oscillators with frequencies  $\omega_1$  and  $\omega_2$  and coupling strength  $\lambda$  driven by reservoir  $A$  and  $B$  the Hamiltonian and standard Lindblad master equation are given by

$$H_S = \omega_1 a_1^\dagger a_1 + \omega_2 a_2^\dagger a_2 + \lambda(a_1 a_2^\dagger + a_1^\dagger a_2), \quad (4.32)$$

$$\frac{d\rho}{dt} = -i[H_S, \rho] + (K^A + K^B)\rho, \quad (4.33)$$

$$\begin{aligned} K^A \rho &= g_A(\omega_1)(1 + n_A(\omega_1))(a_1 \rho a_1^\dagger - (1/2)\{a_1^\dagger a_1, \rho\}) \\ &\quad + g_A(\omega_1)n_A(\omega_1)(a_1^\dagger \rho a_1 - (1/2)\{a_1 a_1^\dagger, \rho\}), \end{aligned} \quad (4.34)$$

$$\begin{aligned} K^B \rho &= g_B(\omega_2)(1 + n_B(\omega_2))(a_2 \rho a_2^\dagger - (1/2)\{a_2^\dagger a_2, \rho\}) \\ &\quad + g_B(\omega_2)n_B(\omega_2)(a_2^\dagger \rho a_2 - (1/2)\{a_2 a_2^\dagger, \rho\}). \end{aligned} \quad (4.35)$$

In this model the interaction between the oscillators is ignored in the dissipative kernels  $K^A$  and  $K^B$ , an assumption often made in the analysis of composite systems and expected to hold for weakly coupled systems. This model has been studied in [16] and shown to be inconsistent with the laws of quantum thermodynamics. The current  $q^A$  turns out to be proportional to  $n_A(\omega_1) - n_B(\omega_2)$  which violates the second law of thermodynamics. The issue was resolved in [16] by expanding the dissipative kernels in the eigenbasis of  $H_S$ , yielding a current in compliance with the second law.

We have not carried out a similar analysis in the qubit case, but note that the field theoretical analysis in Appendix V A, corroborated in Sec. II, implies an analysis in the eigenspace of the full system Hamiltonian  $H_S$ .

## V. APPENDIX

### A. Field theoretical approach

The standard approach to open quantum systems in the Markov approximation is based on the Lindblad equation [10, 19, 20]. In recent work [17] we presented a detailed discussion of the application of field theoretical methods to open quantum systems. Including the rotating wave approximation (RWA) the field theoretical method is equivalent to the Lindblad approach. The field theoretical approach is based on a systematic expansion in terms of diagrams and applies a condensed matter quasi-particle approximation in order to implement the Markov limit. The field theoretical method is the basis for the present analysis.

In the Markov approximation the master equation in the energy representation and in Fourier space has the form

$$-i\omega\rho(\omega)_{pp'} = -i(E_p - E_{p'})\rho(\omega)_{pp'} + \sum_{qq'} L_{pp',qq'}\rho(\omega)_{pp'}, \quad (5.1)$$

where the constant kernel  $L = K + i\Delta$  is composed of a shift  $\Delta$  and a dissipative kernel  $K$ . Assuming that the shift is incorporated in a renormalisation of the energy levels, we have for the dissipative kernel in the Born approximation

$$\begin{aligned} K_{pp',qq'} = & -\delta_{p'q'}\frac{1}{2}\sum_{\alpha\beta,l} S_{pl}^\alpha S_{lq}^\beta D^{\alpha\beta}(E_{pl}) - \delta_{pq}\frac{1}{2}\sum_{\alpha\beta,l} S_{q'l}^\alpha S_{lp'}^\beta D^{\alpha\beta}(E_{p'l}) \\ & + \frac{1}{2}\sum_{\alpha\beta} S_{pq}^\beta S_{q'p'}^\alpha D^{\alpha\beta}(E_{q'p'}) + \frac{1}{2}\sum_{\alpha\beta} S_{pq}^\beta S_{q'p'}^\alpha D^{\alpha\beta}(E_{qp}); \end{aligned} \quad (5.2)$$

where we have introduced the energy shift  $E_{pq} = E_p - E_q$ . Referring to Sec. II we only need the kernel elements  $K_{pq} = K_{pp,qq}$  and (5.2) implies

$$K_{pq} = -\delta_{pq}\frac{1}{2}\sum_{\alpha\beta,l} (S_{pl}^\alpha S_{lq}^\beta + S_{ql}^\alpha S_{lp}^\beta) D^{\alpha\beta}(E_{pl}) + \sum_{\alpha\beta} S_{pq}^\beta S_{qp}^\alpha D^{\alpha\beta}(E_{qp}) \quad (5.3)$$

Above the system operators and bath operators in vector form are denoted by  $\mathbf{A}$  and  $\mathbf{S}$  and we choose a system - bath coupling of the form  $\mathbf{S}\mathbf{A}$ , denoting a scalar product. The bath is of the Caldeira-Leggett form [21], i.e., a collection of quantum oscillators characterised by the bath correlation functions

$$D^{\alpha\beta}(\omega) = \int dt \exp(i\omega(t-t')) \text{Tr}_B[\rho_B B^\alpha(t) B^\beta(t)], \quad (5.4)$$

$$\rho_B = \exp(-H/T) / \text{Tr}_B[\exp(-H_B/T)], \quad (5.5)$$

where  $H_B = \sum_k \omega_k b_k^\dagger b_k$  is the bath Hamiltonian and  $T$  the associated temperature.

For a single bath with assignment  $\mathbf{B} = (B^1, B^2)$ , where

$$B^1(t) = B(t) = \sum_k \lambda_k b_k(t), \quad (5.6)$$

$$B^2(t) = B^\dagger(t) = \sum_k \lambda_k b_k^\dagger(t); \quad (5.7)$$

we have the bath correlations

$$D^{12}(\omega) = g(\omega)(1 + n(\omega)), \quad \omega > 0, \quad (5.8)$$

$$D^{21}(\omega) = g(-\omega)n(-\omega), \quad \omega < 0, \quad (5.9)$$

where the spectral function and Planck distribution are given by

$$g(\omega) = 2\pi \sum_k \lambda_k^2 \delta(\omega - \omega_k), \quad \omega > 0, \quad (5.10)$$

$$n(\omega) = \frac{1}{\exp(\omega/T) - 1}, \quad \omega > 0; \quad (5.11)$$

here  $b_k$  is the Bose field,  $\omega_k$  the frequency associated with the wavenumber  $k$ , and  $\lambda_k$  the system - bath coupling strength. The above scheme is easily generalised to several independent heat baths by using the system - bath coupling  $\mathbf{S}\mathbf{A} + \mathbf{S}\mathbf{B}$  and correlations  $D(\omega) = D_A(\omega) + D_B(\omega)$ , yielding  $K = K^A + K^B$ .

## B. Single qubit

In the single qubit case we have the assignment  $\mathbf{S} = (S^1, S^2) = (\sigma^+, \sigma^-)$  and for the bath operators  $\mathbf{A} = (A^1, A^2) = (A, A^\dagger)$  and  $\mathbf{B} = (B^1, B^2) = (B, B^\dagger)$ . For the coupling to baths  $A$  and  $B$  we employ the dipole coupling in the RWA

$$H_{SA} + H_{SB} = \sigma^+(A + B) + \sigma^-(A^\dagger + B^\dagger). \quad (5.12)$$

For the bath correlations we have

$$D_{A,B}^{12}(\omega) = g_{A,B}(\omega)(1 + n_{A,B}(\omega)), \quad \omega > 0 \quad (5.13)$$

$$D_{A,B}^{21}(\omega) = g_{A,B}(-\omega)n_{A,B}(-\omega), \quad \omega < 0. \quad (5.14)$$



Inserting the nonvanishing matrix elements  $\langle +|\sigma^+|-\rangle = \langle -|\sigma^-|+\rangle = 1$  we obtain from (5.3) the diagonal kernel elements

$$K_{++,++}^{A,B} = -K_{--,++}^{A,B} = -g_{A,B}(\omega_0)(1 + n_{A,B}(\omega_0)), \quad (5.15)$$

$$K_{+,-,-}^{A,B} = -K_{-,-,-}^{A,B} = +g_{A,B}(\omega_0)n_{A,B}(\omega_0), \quad (5.16)$$

with spectral functions and Planck distributions

$$g_{A,B} = g_{A,B}(\omega_0) = 2\pi \sum_k (\lambda_k^{A,B})^2 \delta(\omega_0 - \omega_k^{A,B}), \quad (5.17)$$

$$n_{A,B} = n_{A,B}(\omega_0) = \frac{1}{\exp(\omega_0/T_{A,B}) - 1}. \quad (5.18)$$

### C. Coupled qubits driven by two reservoirs

In the case of interacting qubits driven by reservoirs  $A$  and  $B$  according to (3.17) and (3.18) we have  $K_{pq} = K_{pq}^A + K_{pq}^B$ , where in  $K^A$  we have the assignment  $(S^1, S^2) = (\sigma_1^+, \sigma_1^-)$  and in  $K^B$  the assignment  $(S^1, S^2) = (\sigma_2^+, \sigma_2^-)$  with associated bath correlations (5.13) and (5.14).

In order to evaluate the density operator and *a fortiori* the current according to the scheme in Sec. II we need the matrix elements of  $\sigma_{1,2}^\pm$  in the basis  $|n\rangle$ ,  $n = 1 \cdots 4$  of the system Hamiltonian  $H_S$ , given by (3.21) and (3.24) in Sec. IIIB. With the notation  $(\sigma_{1,2}^\pm)_{pq} = \langle p|\sigma_{1,2}^\pm|q\rangle$  we find the non-vanishing matrix elements

$$(\sigma_1^+)_{31} = (\sigma_1^+)_{42} = (\sigma_2^+)_{21} = (\sigma_2^+)_{43} = \alpha, \quad (5.19)$$

$$(\sigma_1^+)_{43} = -(\sigma_1^+)_{21} = (\sigma_2^+)_{31} = -(\sigma_2^+)_{42} = \beta. \quad (5.20)$$

Using  $(\sigma_{1,2}^+)_{pq} = (\sigma_{1,2}^-)_{qp}$  and inserting in (5.3) we obtain expressions for  $K_{p,q}^{A,B}$  presented as 4 by 4 matrices in (3.42) and (3.43) in Sec. IIIB.




FIG. 1: We depict the drive of a single qubit by a hot heat reservoir  $A$  and a cold heat reservoir  $B$ . The heat current  $q^A$  flows from  $A$  to the qubit and  $-q^B$  from qubit to  $B$ ;  $q^A + q^B = 0$  (energy conservation).




FIG. 2: We depict the drive of a composite two-qubit system by a hot heat reservoir  $A$  coupled to qubit 1 and a cold heat reservoir  $B$  coupled to qubit 2. Owing to the interaction  $\lambda$  the heat current is transmitted from qubit 1 to qubit 2. The heat current  $q^A$  flows from  $A$  to qubit 1, from qubit 1 to qubit 2 via the interaction and  $-q^B$  from qubit 2 to  $B$ ;  $q^A + q^B = 0$  (energy conservation).

- 
- [1] R. Kosloff and A. Levy, *Annu. Rev. Phys. Chem.* **65**, 365 (2014).
- [2] R. Alicki, *J. Phys. A* **12**, L103 (1979).
- [3] F. Barra, *Sci. Rep.* **5**, 14873 (2015).
- [4] R. Kosloff, *Entropy* **15**, 2100 (2013).
- [5] A. Mari and J. Eisert, *Phys. Rev. Lett.* **108**, 120602 (2012).
- [6] R. Kosloff and A. Levy, *J. Chem. Phys.* **150**, 204105 (2019).
- [7] A. Levy, M. Goeb, B. Dong, K. Singer, K. Torrontegui, and D. Wang, *New J. Phys.* **22**, 093020 (2020).
- [8] A. Colla and H.-P. Breuer, *Phys. Rev. A* **105**, 052216 (2021).
- [9] N. Linden, S. Popescu, and P. Skrzypczyk, *Phys. Rev. Lett.* **105**, 130401 (2010).
- [10] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2006).
- [11] A. Rivas and S. F. Huelga, *Open Quantum Systems* (Springer-Verlag, Berlin, 2012).
- [12] W. G. van der Wiel, S. D. Franceschi, J. M. Elzerman, T. Fujisawa, S. Tarucha, and L. P. Kouwenhoven, *Rev. Mod. Phys.* **75**, 1 (2002).
- [13] S. Kohler, J. Lehmann, and P. Haenggi, *Phys. Rep.* **406**, 379 (2005).
- [14] G. D. Chiara, G. Landi, A. Hewgill, B. Reid, A. Ferraro, A. Roncaglia, and M. Antezza, *New Journal of Physics* **20**, 113024 (2018).
- [15] A. Hewgill, G. D. Chiara, and A. Imparato, *Phys. Rev. Research* **3**, 013165 (2021).
- [16] A. Levy and R. Kosloff, *EPL* **107**, 20004 (2014).
- [17] H. C. Fogedby, *Phys. Rev. A* **106**, 022205 (2022).
- [18] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford University Press, Oxford, 1989).
- [19] G. Lindblad, *Commun. Math. Phys.* **48**, 119 (1976).
- [20] D. Manzano, *AIP Advances* 10 p. 025106 (2020).
- [21] A. O. Caldeira and A. J. Leggett, *Ann. Phys.* **149**, 374 (1983).