

W -representations of two-matrix models with infinite set of variables

Lu-Yao Wang^{a,*}, Yu-Sen Zhu^{a,†}, Ying Chen^{b,‡}, Bei Kang^{c§}

^a School of Mathematical Sciences, Capital Normal University, Beijing 100048, China

^bSchool of Mathematics and Statistics, Jiangsu Normal University, Xuzhou 221116, Jiangsu, China

^c School of Mathematics and Statistics, North China University of Water Resources and Electric Power, Zhengzhou 450046, Henan, China

Abstract

The Hermitian, complex and fermionic two-matrix models with infinite set of variables are constructed. We show that these two-matrix models can be realized by the W -representations. In terms of the W -representations, we derive the compact expressions of correlators for these two-matrix models.

Keywords: Two-matrix models, Conformal and W Symmetry

1 Introduction

Matrix models have been developed to solve non-perturbative two-dimensional gravity and provide a rich set of approaches to physical systems. For two-matrix model, there is the interaction between the two matrices. Hence it possesses a richer mathematical structure than single matrix models, and thus produces more applications in physics and mathematics. The two-matrix models have been studied as an important solvable example of statistical mechanical systems, i.e., Ising spins [1–3]. For fermionic two-matrix model, the complete sets of loop equations can be derived [4]. The Ward identities in Kontsevich-like one-matrix models are used to relate the degree of potential in Kontsevich-like two-matrix model to the W -constraints [5]. The spectral curves, loop equations and topological expansion for Hermitian two-matrix models were presented in Refs.[6–8].

For W -representation of matrix model, it realizes partition function by acting on elementary functions with exponents of the given W -operator [9]. Since W -representation plays an important role in understanding the structures of matrix models, much interest has been attributed to this direction. A variety of matrix models have been realized by W -representations and their correlators can be exactly calculated. Recently the (super) partition function hierarchies with W -representations were constructed [10, 11]. Some well known superintegrable matrix models were contained in these superintegrable hierarchies. In addition, the progress of W -representation has been made on tensor models [12–15] and super-eigenvalue models [16, 17].

Recently, the two-matrix models with multi-set of variables were proposed [18–21], which are the superintegrable matrix models. Their W -representations and character expansions were well investigated. In this paper, we'll construct the new two-matrix models with infinite set of variables and derive their W -representations.

*wangly100@outlook.com

†zhuyusen@cnu.edu.cn

‡chenying_math@jsnu.edu.cn

§Corresponding author:kangbei@ncwu.edu.cn

2 W -representation of new Hermitian two-matrix model

Let us construct the Hermitian two-matrix model

$$\begin{aligned} Z_{2H} &= \int dA dB \exp\left(-\frac{1}{2}\text{tr}A^2 - \frac{1}{2}\text{tr}B^2 + \sum_{k=0}^{\infty} t_k \text{tr}A^k + \sum_{k=0}^{\infty} g_k \text{tr}B^k\right. \\ &\quad \left. + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} \text{tr}A^{k_1} B^{k_2} A^{k_3} B^{k_4} \dots A^{k_{2l-1}} B^{k_{2l}}\right), \end{aligned} \quad (1)$$

where A and B are $N \times N$ matrices. When $B = 0$ in (1), it reduces to the well known Gaussian Hermitian matrix model.

By requiring the invariance of the integral (1) under the infinitesimal transformation $A \rightarrow A + \epsilon A^n$ ($n \geq 0$) or $B \rightarrow B + \epsilon B^n$ ($n \geq 0$), we obtain the Virasoro constraints

$$\mathcal{L}_{n-1} Z_{2H} = 0, \quad (2)$$

where the constraint operators are given by

$$\begin{aligned} \mathcal{L}_{n-1} &= \sum_{n=0}^{\infty} 2N \frac{\partial}{\partial t_{n-1}} + \sum_{n=0}^{\infty} \sum_{s=1}^{n-1} \frac{\partial^2}{\partial t_s \partial t_{n-1-s}} + \delta_{n,1} N^2 + \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k-1}} + \sum_{k_2=1}^{\infty} t_{1,k_2} \frac{\partial}{\partial g_{k_2}} \\ &\quad + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} \left[\sum_{a=1}^l \sum_{n=0}^{\infty} k_{2a-1} \frac{\partial}{\partial t_{k_1, \dots, n+k_{2a-1}, k_{2a}, \dots, k_{2l}}} + \delta_{k_1,1} \frac{\partial}{\partial t_{k_3, \dots, k_{2l-1}, k_{2l}+k_2}} \right. \\ &\quad \left. + \sum_{a=2}^l \delta_{k_{2a-1},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}+k_{2a}, k_{2a+1}, \dots, k_{2l}}} \right], \end{aligned} \quad (3)$$

which obey the Virasoro algebra

$$[\mathcal{L}_{n-1}, \mathcal{L}_{m-1}] = (n-m) \mathcal{L}_{n+m-2}. \quad (4)$$

Let us now consider the following five infinitesimal transformations, respectively,

$$(i) A \longrightarrow A + \epsilon \sum_{n=0}^{\infty} (n+1) t_{n+1} A^n, \quad (ii) A \longrightarrow A + \epsilon \sum_{n=1}^{\infty} (n+1) t_{1,n} B^n,$$

$$(iii) A \longrightarrow A + \epsilon \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r}=1}^{\infty} \mathcal{N}_1 t_{n_1+1, n_2, \dots, n_{2r}} A^{n_1} B^{n_2} \dots A^{n_{2r-1}} B^{n_{2r}},$$

$$(iv) A \longrightarrow A + \epsilon \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r}=1}^{\infty} \mathcal{N}_2 t_{1, n_2, \dots, n_{2r}} B^{n_2} A^{n_3} \dots A^{n_{2r-1}} B^{n_{2r}},$$

$$(v) B \longrightarrow B + \epsilon \sum_{n=0}^{\infty} (n+1) g_{n+1} B^n,$$

where $\mathcal{N}_1 = n_1 + 1 + n_2 + \dots + n_{2r}$ and $\mathcal{N}_2 = 1 + n_2 + \dots + n_{2r}$.

From the invariance of the integral (1), it gives

$$\hat{D}_i Z_{2H} = \hat{W}_i Z_{2H}, i = 1, 2, \dots, 5, \quad (5)$$

where the operators \hat{W}_i are listed in (A.1) and \hat{D}_i are

$$\begin{aligned}\hat{D}_1 &= \sum_{i=1}^{\infty} it_i \frac{\partial}{\partial t_i}, \quad \hat{D}_2 = \sum_{n=1}^{\infty} (n+1)t_{1,n} \frac{\partial}{\partial t_{1,n}}, \\ \hat{D}_3 &= \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r}=1}^{\infty} \mathcal{N}_1 t_{n_1+1, n_2, \dots, n_{2r}} \frac{\partial}{\partial t_{n_1+1, n_2, \dots, n_{2r}}}, \\ \hat{D}_4 &= \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r}=1}^{\infty} \mathcal{N}_2 t_{1, n_2, \dots, n_{2r}} \frac{\partial}{\partial t_{1, n_2, \dots, n_{2r}}}, \quad \hat{D}_5 = \sum_{i=1}^{\infty} i g_i \frac{\partial}{\partial g_i}.\end{aligned}\tag{6}$$

In the following, we'll focus on the sum of (5)

$$\hat{D}Z_{2H} = \hat{W}Z_{2H},\tag{7}$$

where $\hat{D} = \sum_{i=1}^5 D_i$ and $\hat{W} = \sum_{i=1}^5 W_i$.

Let us write the partition function (1) as the grading form $Z_{2H} = \sum_{d=0}^{\infty} Z_{2H}^{(d)}$ and

$$\begin{aligned}Z_{2H}^{(d)} &= e^{N(t_0+g_0)} \sum_{l=0}^{\infty} \frac{1}{l!} \sum_{\substack{l_1+l_2+l_3=l \\ \rho_1+\rho_2+\rho_3=d}} \langle \prod_{i=1}^{l_1} \text{tr} A^{k_i} \prod_{j=1}^{l_2} \text{tr} B^{r_j} \prod_{n=1}^{l_3} \text{tr} A^{S_{n,1}} B^{S_{n,2}} \dots A^{S_{n,2p_n-1}} B^{S_{n,2p_n}} \rangle \\ &\cdot \prod_{i=1}^{l_1} t_{k_i} \prod_{j=1}^{l_2} g_{r_j} \prod_{n=1}^{l_3} t_{S_{n,1}, \dots, S_{n,2p_n}} \cdot \int dA dB \exp(-\frac{1}{2} \text{tr} A^2 - \frac{1}{2} \text{tr} B^2),\end{aligned}\tag{8}$$

where $\rho_1 = \sum_{i=1}^{l_1} k_i$, $\rho_2 = \sum_{i=1}^{l_2} r_i$, $\rho_3 = \sum_{i=1}^{l_3} (S_{i,1} + \dots + S_{i,2p_i})$ correlators $\langle \dots \rangle$ are defined as

$$\langle \dots \rangle = \frac{\int dA dB \dots \exp(-\frac{1}{2} \text{tr} A^2 - \frac{1}{2} \text{tr} B^2)}{\int dA dB \exp(-\frac{1}{2} \text{tr} A^2 - \frac{1}{2} \text{tr} B^2)}.\tag{9}$$

We denote the degrees of operators as $\deg(t_k) = \deg(g_k) = k$, $\deg(\frac{\partial}{\partial t_k}) = \deg(\frac{\partial}{\partial g_k}) = -k$, $\deg(\frac{\partial}{\partial t_{k_1, k_2, \dots, k_{2l-1}, k_{2l}}}) = -(k_1 + \dots + k_{2l})$. Then it is easy to see that $\deg(\hat{D}) = 0$ and $\deg(\hat{W}) = 2$.

Due to the operators \hat{D} and $\hat{D} - \hat{W}$ being invertible and $\hat{D}e^{N(t_0+g_0)} = 0$, from (7), we have

$$\sum_{s=1}^{\infty} Z_{2H}^{(s)} = (\hat{D} - \hat{W})^{-1} \hat{W} e^{N(t_0+g_0)} = \sum_{k=1}^{\infty} (\hat{D}^{-1} \hat{W})^k e^{N(t_0+g_0)}.\tag{10}$$

Note that \hat{W} is an homogeneous operator with degree 2, and $\hat{D}f = \deg(f) \cdot f$ for any homogeneous function f . We may give the W -representation of the Hermitian two-matrix model (1)

$$Z_{2H} = e^{\frac{1}{2} \hat{W}} e^{N(t_0+g_0)}.\tag{11}$$

Let us formally write the $(m+1)$ -th power of the operator \hat{W} as

$$\begin{aligned}\hat{W}^{(m+1)} &= \sum_{l_1+l_2+l_3=1}^{2(m+1)} \sum_{\rho_1+\rho_2+\rho_3=2(m+1)} P_{(k_1, \dots, k_{l_1}); (r_1, \dots, r_{l_2})}^{(S_{1,1}, \dots, S_{1,2p_1}; \dots; S_{l_3,1}, \dots, S_{l_3,2p_{l_3}})} t_{k_1} \dots t_{k_{l_1}} g_{r_1} \dots g_{r_{l_2}} \\ &\cdot t_{S_{1,1}, \dots, S_{1,2p_1}} \dots t_{S_{l_3,1}, \dots, S_{l_3,2p_{l_3}}} + \dots.\end{aligned}\tag{12}$$

By means of the W -representation of (1), we derive the compact expression of correlators

$$\begin{aligned} & \left\langle \prod_{i=1}^{l_1} \text{tr} A^{k_i} \prod_{j=1}^{l_2} \text{tr} B^{r_j} \prod_{n=1}^{l_3} \text{tr} A^{S_{n,1}} B^{S_{n,2}} \cdots A^{S_{n,2p_n-1}} B^{S_{n,2p_n}} \right\rangle \\ &= \frac{l_1! l_2! l_3! \sum_{\rho_1+\rho_2+\rho_3=1}^{2(m+1)} \sum_{\sigma} P_{(\sigma(k_1), \dots, \sigma(k_{l_1})); (\sigma(r_1), \dots, \sigma(r_{l_2}))}^{(\sigma(S_{1,1}), \dots, \sigma(S_{1,2p_1}); \dots; \sigma(S_{l_3,1}), \dots, \sigma(S_{l_3,2p_{l_3}}))}}{2^{m+1}(m+1)! \lambda_{(k_1, \dots, k_{l_1})} \lambda_{(r_1, \dots, r_{l_2})} \lambda_{(S_{1,1}, \dots, S_{1,2p_1}; \dots; S_{l_3,1}, \dots, S_{l_3,2p_{l_3}})}}, \end{aligned} \quad (13)$$

where $(\sigma(k_1), \dots, \sigma(k_{l_1}))$ denotes all distinct permutations of $(k_{l_1}, \dots, k_{l_1})$, and $\lambda_{(k_1, \dots, k_{l_1})}$ is the number of distinct permutations (k_1, \dots, k_{l_1}) .

For example, let us consider the cases

$$\begin{aligned} \hat{W} &= t_1^2 N + 2t_2 N^2 + g_1^2 N + 2g_2 N^2 + \cdots, \\ \hat{W}^2 &= 8t_1^2 t_2 N + 24t_1 t_3 N^2 + 12t_4 N^3 + 8t_2^2 N^2 + 8t_1 t_{1,2} N^2 + 8t_1 g_1 t_{1,1} N + 8g_1 t_{2,1} N^2 \\ &\quad + 4t_{1,1}^2 N^2 + 8t_{2,2} N^3 + 8g_1^2 g_2 N + 24g_1 g_3 N^2 + 12g_4 N^3 + 8g_2^2 N^2 + \cdots. \end{aligned} \quad (14)$$

We may give some correlators in (13) as follows

$$\begin{aligned} \langle \text{tr} A \text{tr} A \rangle &= \langle \text{tr} B \text{tr} B \rangle = N, & \langle \text{tr} A^2 \rangle &= \langle \text{tr} B^2 \rangle = N^2, \\ \langle \text{tr} A \text{tr} B \text{tr} A^2 \rangle &= 2N + N^3, & \langle \text{tr} A \text{tr} A^3 \rangle &= \langle \text{tr} B \text{tr} B^3 \rangle = 3N^2, \\ \langle \text{tr} A^4 \rangle &= \langle \text{tr} B^4 \rangle = 3N^2, & \langle \text{tr} A^2 B^2 \rangle &= N^2, \\ \langle \text{tr} A B \text{tr} A B \rangle &= N^2, & \langle \text{tr} A \text{tr} B \text{tr} A B \rangle &= N, \\ \langle \text{tr} A \text{tr} A B^2 \rangle &= \langle \text{tr} A \text{tr} A^2 B \rangle = N^2, & \langle \text{tr} A^2 \text{tr} A^2 \rangle &= \langle \text{tr} B^2 \text{tr} B^2 \rangle = 2N^2 + N^4. \end{aligned} \quad (15)$$

3 W -representation of complex two-matrix model

Let us construct the complex two-matrix model

$$\begin{aligned} Z_{2C} &= \int d^2 M_1 d^2 M_2 \exp[-\mu \text{tr} M_1 M_1^\dagger - \mu \text{tr} M_2 M_2^\dagger + \sum_{k=0}^{\infty} t_k \text{tr} (M_1 M_1^\dagger)^k + \sum_{k=0}^{\infty} g_k \text{tr} (M_2 M_2^\dagger)^k \\ &\quad + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} \text{tr} (M_1 M_1^\dagger)^{k_1} (M_2 M_2^\dagger)^{k_2} \cdots (M_1 M_1^\dagger)^{k_{2l-1}} (M_2 M_2^\dagger)^{k_{2l}}], \end{aligned} \quad (16)$$

where M_1 and M_2 are $N \times N$ complex matrices.

By requiring the invariance of the integral (16) under the infinitesimal transformation $M_1 \rightarrow M_1 + \epsilon(M_1 M_1^\dagger)^n M_1$ ($n \geq 0$) or $M_2 \rightarrow M_2 + \epsilon(M_2 M_2^\dagger)^n M_2$ ($n \geq 0$), it gives the Virasoro constraints

$$\bar{\mathcal{L}}_n Z_{2C} = 0, \quad (17)$$

where

$$\begin{aligned} \bar{\mathcal{L}}_n &= \sum_{n=0}^{\infty} 2N \frac{\partial}{\partial t_n} + \sum_{n=0}^{\infty} \sum_{s=1}^{n-1} \frac{\partial^2}{\partial t_s \partial t_{n-s}} + \delta_{n,0} N^2 - \mu \sum_{n=0}^{\infty} \frac{\partial}{\partial t_{n+1}} + \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k}} \\ &\quad + \sum_{n=0}^{\infty} \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} \sum_{a=1}^l k_{2a-1} t_{k_1, \dots, k_{2l}} \frac{\partial}{\partial t_{k_1, k_2, \dots, n+k_{2a-1}, k_{2a}, \dots, k_{2l}}}. \end{aligned} \quad (18)$$

Similarly, the four constraints of (16) can be derived from the invariance of the integral under the following four infinitesimal transformations, respectively,

- (i) $M_1 \longrightarrow M_1 + \epsilon \sum_{n=0}^{\infty} (n+1)t_{n+1}(M_1 M_1^\dagger)^n M_1$,
- (ii) $M_1 \longrightarrow M_1 + \epsilon \sum_{n,m=0}^{\infty} [(n+1)+(m+1)]t_{n+1,m+1}(M_2 M_2^\dagger)^{m+1}(M_1 M_1^\dagger)^n M_1$,
- (iii) $M_1 \longrightarrow M_1 + \epsilon \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \bar{\mathcal{N}} t_{n_{2r+1}, n_2+1, \dots, n_{2r+1}} (M_2 M_2^\dagger)^{n_2+1} (M_1 M_1^\dagger)^{n_3+1} \dots (M_2 M_2^\dagger)^{n_{2r+1}} (M_1 M_1^\dagger)^{n_{2r+1}} M_1$,
- (iv) $M_2 \longrightarrow M_2 + \epsilon \sum_{m=0}^{\infty} (m+1)g_{m+1}(M_2 M_2^\dagger)^m M_2$,

where $\bar{\mathcal{N}} = (n_2+1) + (n_3+1) + \dots + (n_{2r+1}+1)$. The sum of these constraints are

$$\mu \bar{D} Z_{2C} = \bar{W} Z_{2C}, \quad (19)$$

where $\bar{D} = \sum_{i=1}^4 \bar{D}_i$, $\bar{W} = \sum_{i=1}^4 \bar{W}_i$, the operators \bar{D}_i are

$$\begin{aligned} \bar{D}_1 &= \sum_{n=0}^{\infty} (n+1)t_{n+1} \frac{\partial}{\partial t_{n+1}}, & \bar{D}_2 &= \sum_{n,m=0}^{\infty} \bar{T}_1 \frac{\partial}{\partial t_{n+1,m+1}}, \\ \bar{D}_2 &= \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \bar{T}_2 \frac{\partial}{\partial t_{n_1+1, \dots, n_{2r+1}+1}}, & \bar{D}_4 &= \sum_{m=0}^{\infty} (m+1)g_{m+1} \frac{\partial}{\partial g_{m+1}}, \end{aligned} \quad (20)$$

and the operators \bar{W}_i are

$$\begin{aligned} \bar{W}_1 &= \sum_{n=0}^{\infty} (n+1)t_{n+1} [2N \frac{\partial}{\partial t_n} (1 - \delta_{n,0}) + \sum_{a=1}^{n-1} \frac{\partial}{\partial t_a} \frac{\partial}{\partial t_{n-a}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k}}] + N^2 t_1 \\ &\quad + \sum_{n=0}^{\infty} \sum_{l=1}^{\infty} \sum_{a=1}^l \sum_{k_1, \dots, k_{2l}=1}^{\infty} (n+1)t_{n+1} k_{2a-1} t_{k_1, \dots, k_{2l}} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-1}+n, k_{2a}, \dots, k_{2l}}}, \\ \bar{W}_2 &= \sum_{n,m=0}^{\infty} \bar{T}_1 \{ (N \frac{\partial}{\partial t_{n,m+1}} + \frac{\partial^2}{\partial g_{m+1} \partial t_n}) (1 - \delta_{n,0}) + \sum_{a=1}^{n-1} \frac{\partial^2}{\partial t_{a,m+1} \partial t_{n-a}} + N \delta_{n,0} \frac{\partial}{\partial g_{m+1}} \\ &\quad + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k, m+1}} + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} [k_1 \frac{\partial}{\partial t_{n+k_1, \dots, k_{2l-1}, k_{2l}+m+1}} \\ &\quad + k_{2l-1} \sum_{a=2}^l \frac{\partial}{\partial t_{k_1, \dots, k_{2a-1}+n, k_{2a}, \dots, k_{2l}}} + \sum_{a=1}^l \sum_{s=1}^{k_{2a-1}-1} \frac{\partial}{\partial t_{k_1, \dots, s, m+1, n+k_{2a-1}-s, k_{2a}, \dots, k_{2l}}}] \} \} \\ \bar{W}_3 &= \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \bar{T}_2 \{ [N \frac{\partial}{\partial g_{n_2+1}} \frac{\partial}{\partial t_{n_3+n_{2r+1}+1, n_4+1, \dots, n_{2r+1}}} + N \frac{\partial}{\partial t_{n_3+1, \dots, n_{2r}+n_2+2}} \frac{\partial}{\partial t_{n_{2r+1}}} \\ &\quad + N \sum_{b=2}^{r-1} \frac{\partial}{\partial t_{n_3+1, \dots, n_{2b}+n_2+2}} \frac{\partial}{\partial t_{n_{2b+1}+1, \dots, n_{2r+1}}} + \sum_{s=1}^{n_{2r+1}-1} \frac{\partial}{\partial t_{s, n_2+1, \dots, n_{2r+1}}} \frac{\partial}{\partial t_{n_{2r+1}-s}} \\ &\quad + \sum_{b=1}^{r-1} \sum_{s=1}^{n_{2b+1}} \frac{\partial}{\partial t_{s, n_2+1, \dots, n_{2b}+1}} \frac{\partial}{\partial t_{\xi_1, n_{2b+2}+1, \dots, n_{2r+1}}} + N \frac{\partial}{\partial t_{n_{2r+1}, n_2+1, \dots, n_{2r+1}}}] \cdot \\ &\quad \cdot (1 - \delta_{n_{2r+1}, 0}) + \delta_{n_{2r+1}, 0} \frac{\partial}{\partial t_{n_3+1, \dots, n_{2r-1}+1, n_{2r}+n_2+2}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n_{2r+1}+k, n_2+1, \dots, n_{2r+1}}} \} \} \end{aligned}$$

$$\begin{aligned}
& + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} [k_1 \frac{\partial}{\partial t_{n_3+1, \dots, \bar{\xi}_2, k_2, \dots, \bar{\xi}_3}} + \sum_{a=2}^l (k_{2a-1} \frac{\partial}{\partial t_{k_1, \dots, \bar{\xi}_4, k_{2a}, \dots, k_{2l}}} \\
& + \sum_{s=1}^{k_{2a-1}-1} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}, s, n_2+1, \dots, \bar{\xi}_5, k_{2a}, \dots, k_{2l}})])], \\
\bar{W}_4 & = N^2 g_1 + \sum_{m=0}^{\infty} (m+1) g_{m+1} [2N \frac{\partial}{\partial g_m} (1 - \delta_{m,0}) + \sum_{a=1}^{m-1} \frac{\partial}{\partial g_a} \frac{\partial}{\partial g_{m-a}} + \sum_{k=0}^{\infty} k g_k \frac{\partial}{\partial g_{m+k}}] \\
& + \sum_{m=0}^{\infty} \sum_{l=1}^{\infty} \sum_{a=1}^l \sum_{k_1, \dots, k_{2l}=1}^{\infty} (m+1) g_{m+1} k_{2a} t_{k_1, \dots, k_{2l}} \frac{\partial}{\partial t_{k_1, \dots, k_{2a+m}, k_{2a+1}, \dots, k_{2l}}}, \tag{21}
\end{aligned}$$

where $\bar{T}_1 = (n+m+2)t_{n+1,m+1}$, $\bar{T}_2 = \bar{\mathcal{N}}t_{n_{2r+1}+1,n_2+1, \dots, n_{2r+1}}$ and $\bar{\xi}_1 = n_{2r+1} + n_{2b+1} + 1 - s$, $\bar{\xi}_2 = n_{2r+1} + k_1$, $\bar{\xi}_3 = k_{2l} + n_2 + 1$, $\bar{\xi}_4 = (k_{2a-2} + n_2 + 1, n_3 + 1, \dots, n_{2r} + 1, n_{2r+1} + k_{2a-1})$, $\bar{\xi}_5 = n_{2r+1} + k_{2a-1} - s$.

Similar to the case of the Hermitian two-matrix model (11), the complex two-matrix model (16) can be realized by the W -representation

$$Z_{2C} = e^{\frac{1}{\mu} \bar{W}} e^{N(t_0 + g_0)}. \tag{22}$$

There is also the compact expression of correlators

$$\begin{aligned}
& \langle \prod_{i=1}^{l_1} \text{tr}(M_1 M_1^\dagger)^{k_i} \prod_{j=1}^{l_2} \text{tr}(M_2 M_2^\dagger)^{r_j} \prod_{n=1}^{l_3} \text{tr}(M_1 M_1^\dagger)^{S_{n,1}} (M_2 M_2^\dagger)^{S_{n,2}} \cdots (M_2 M_2^\dagger)^{S_{n,2p_n}} \rangle \\
& = \frac{l_1! l_2! l_3! \sum_{\rho_1+\rho_2+\rho_3=1}^{m+1} \sum_{\sigma} \bar{P}_{(\sigma(k_1), \dots, \sigma(k_{l_1})); (\sigma(r_1), \dots, \sigma(r_{l_2}))}^{(\sigma(S_{1,1}), \dots, \sigma(S_{1,2p_1}); \dots; \sigma(S_{l_3,1}), \dots, \sigma(S_{l_3,2p_{l_3}}))}}{\mu^{m+1} (m+1)! \lambda_{(k_1, \dots, k_{l_1})} \lambda_{(r_1, \dots, r_{l_2})} \lambda_{(S_{1,1}, \dots, S_{1,2p_1}; \dots; S_{l_3,1}, \dots, S_{l_3,2p_{l_3}})}}, \tag{23}
\end{aligned}$$

where $\rho_1 = \sum_{i=1}^{l_1} k_i$, $\rho_2 = \sum_{i=1}^{l_2} r_i$, $\rho_3 = \sum_{i=1}^{l_3} (S_{i,1} + \dots + S_{i,2p_i})$, and $\bar{P}_{(\sigma(k_1), \dots, \sigma(k_{l_1})); (\sigma(r_1), \dots, \sigma(r_{l_2}))}^{(\sigma(S_{1,1}), \dots, \dots; \sigma(S_{l_3,2p_{l_3}}))}$ is

the coefficient of $t_{k_1} \cdots t_{k_{l_1}} g_{r_1} \cdots g_{r_{l_2}} t_{S_{1,1}, \dots, S_{1,2p_1}} \cdots t_{S_{l_3,1}, \dots, S_{l_3,2p_{l_3}}}$ in \bar{W}^{m+1} .

For example, we list some correlators

$$\begin{aligned}
\langle \text{tr} M_1 M_1^\dagger \rangle & = \langle \text{tr} M_2 M_2^\dagger \rangle = \frac{1}{\mu} N^2, \quad \langle \text{tr} M_1 M_1^\dagger \text{tr} M_2 M_2^\dagger \rangle = \frac{2}{\mu^2} N^4, \\
\langle \text{tr} M_1 M_1^\dagger \text{tr} M_1 M_1^\dagger \rangle & = \langle \text{tr} M_2 M_2^\dagger \text{tr} M_2 M_2^\dagger \rangle = \frac{1}{\mu^2} (N^2 + 1) N^2, \\
\langle \text{tr} M_1 M_1^\dagger M_2 M_2^\dagger \rangle & = \frac{2}{\mu^2} N^3, \quad \langle \text{tr}(M_1 M_1^\dagger)^3 \rangle = \langle \text{tr}(M_2 M_2^\dagger)^3 \rangle = \frac{6}{\mu^3} (N^2 + N^4), \\
\langle \text{tr} M_1 M_1^\dagger \text{tr} M_1 M_1^\dagger \text{tr} M_1 M_1^\dagger \rangle & = \langle \text{tr} M_2 M_2^\dagger \text{tr} M_2 M_2^\dagger \text{tr} M_2 M_2^\dagger \rangle = \frac{1}{\mu^3} (N^2 + 2)(N^2 + 1) N^2, \\
\langle \text{tr} M_1 M_1^\dagger \text{tr} M_1 M_1^\dagger \text{tr} M_2 M_2^\dagger \rangle & = \langle \text{tr} M_1 M_1^\dagger \text{tr} M_2 M_2^\dagger \text{tr} M_2 M_2^\dagger \rangle = \frac{3}{\mu^3} (N^4 + N^6), \\
\langle \text{tr} M_1 M_1^\dagger \text{tr} M_1 M_1^\dagger M_2 M_2^\dagger \rangle & = \langle \text{tr} M_2 M_2^\dagger \text{tr} M_1 M_1^\dagger M_2 M_2^\dagger \rangle = \frac{6}{\mu^3} (N^3 + N^5), \\
\langle \text{tr} M_1 M_1^\dagger \text{tr}(M_2 M_2^\dagger)^2 \rangle & = \langle \text{tr}(M_1 M_1^\dagger)^2 \text{tr} M_2 M_2^\dagger \rangle = \frac{8}{\mu^3} N^5, \\
\langle \text{tr} M_1 M_1^\dagger \text{tr}(M_1 M_1^\dagger)^2 \rangle & = \langle \text{tr} M_2 M_2^\dagger \text{tr}(M_2 M_2^\dagger)^2 \rangle = \frac{8}{\mu^3} (N^3 + N^5). \tag{24}
\end{aligned}$$

4 W -representation of fermionic two-matrix model

The fermionic matrix model Z_F with the super integrability is given by [14]

$$Z_F = \frac{\int d\psi d\bar{\psi} \exp[N \sum_{k>0} \frac{p_k}{k} \text{tr}(\bar{\psi}\psi)^k + N^2 \text{tr}(\bar{\psi}\psi)]}{\int d\psi d\bar{\psi} \exp(N^2 \text{tr}\bar{\psi}\psi)}$$

$$= \sum_R \left(\frac{-1}{N}\right)^{|R|} \frac{D_R(N)D_R(-N)}{d_R} S_R, \quad (25)$$

where ψ and $\bar{\psi}$ are independent complex Grassmann-valued $N \times N$ matrices, and $D_R(N) = S_R\{p_k = N\}$, $d_R = S_R\{p_k = \delta_{k,1}\}$ are respectively the dimension of representation R for the linear group $GL(N)$.

Let us extend (25) to the fermionic two-matrix model,

$$\begin{aligned} Z_{2F} &= \int d\psi d\bar{\psi} d\chi d\bar{\chi} \exp[-\mu \text{tr} \bar{\psi} \psi - \mu \text{tr} \bar{\chi} \chi + \sum_{k=0}^{\infty} t_k \text{tr}(\bar{\psi} \psi)^k + \sum_{k=0}^{\infty} g_k \text{tr}(\bar{\chi} \chi)^k \\ &\quad + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} t_{k_1, \dots, k_{2l}} \text{tr}(\bar{\psi} \psi)^{k_1} (\bar{\chi} \chi)^{k_2} (\bar{\psi} \psi)^{k_3} \dots (\bar{\psi} \psi)^{k_{2l-1}} (\bar{\chi} \chi)^{k_{2l}}], \end{aligned} \quad (26)$$

where χ and $\bar{\chi}$ are independent complex Grassmann-valued $N \times N$ matrices.

There are the Virasoro constraints

$$\check{\mathcal{L}}_n Z_{2F} = 0, \quad (27)$$

where

$$\begin{aligned} \check{\mathcal{L}}_n &= \sum_{n=0}^{\infty} \sum_{s=1}^{n-1} \frac{\partial^2}{\partial t_s \partial t_{n-s}} - \delta_{n,0} N^2 - \mu \sum_{n=0}^{\infty} \frac{\partial}{\partial t_{n+1}} + \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k}} \\ &\quad + \sum_{n=0}^{\infty} \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} \sum_{a=1}^l k_{2a-1} t_{k_1, \dots, k_{2l}} \frac{\partial}{\partial t_{k_1, \dots, n+k_{2a-1}, k_{2a}, \dots, k_{2l}}}. \end{aligned} \quad (28)$$

Similar to the complex two-matrix case, by considering the following infinitesimal transformations in the integral (26), respectively,

- (i) $\psi \rightarrow \psi + \epsilon \sum_{n=0}^{\infty} (n+1) t_{n+1} \psi (\bar{\psi} \psi)^n$,
- (ii) $\psi \rightarrow \psi + \epsilon \sum_{n,m=0}^{\infty} [(n+1) + (m+1)] t_{n+1, m+1} \psi (\bar{\chi} \chi)^{m+1} (\bar{\psi} \psi)^n$,
- (iii) $\psi \rightarrow \psi + \epsilon \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \check{\mathcal{N}} \check{T} \psi (\bar{\chi} \chi)^{n_2+1} (\bar{\psi} \psi)^{n_3+1} \dots (\bar{\chi} \chi)^{n_{2r}+1} (\bar{\psi} \psi)^{n_{2r+1}}$,
- (iv) $\chi \rightarrow \chi + \epsilon \sum_{m=0}^{\infty} (m+1) g_{m+1} \chi (\bar{\chi} \chi)^m$,

where $\check{\mathcal{N}} \check{T} = (n_2+1) + (n_3+1) + \dots + (n_{2r+1}+1) t_{n_{2r+1}, n_2+1, \dots, n_{2r+1}}$, we finally obtain

$$\mu \check{D} Z_{2F} = \check{W} Z_{2F}, \quad (29)$$

where $\check{D} = \sum_{i=1}^4 \check{D}_i$, $\check{W} = \sum_{i=1}^4 \check{W}_i$, the operators \check{D}_i and \check{W}_i are

$$\begin{aligned} \check{D}_1 &= \sum_{n=0}^{\infty} (n+1) t_{n+1} \frac{\partial}{\partial t_{n+1}}, & \check{D}_2 &= \sum_{n,m=0}^{\infty} \check{T}_1 \frac{\partial}{\partial t_{n+1, m+1}}, \\ \check{D}_3 &= \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \check{T}_2 \frac{\partial}{\partial t_{n_1+1, \dots, n_{2r+1}+1}}, & \check{D}_4 &= \sum_{m=0}^{\infty} (m+1) g_{m+1} \frac{\partial}{\partial g_{m+1}}, \end{aligned} \quad (30)$$

$$\begin{aligned}
\check{W}_1 &= \sum_{n=0}^{\infty} (n+1) t_{n+1} \left[\sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k}} + \sum_{a=1}^{n-1} \frac{\partial}{\partial t_a} \frac{\partial}{\partial t_{n-a}} + \sum_{l=1}^{\infty} \sum_{a=1}^l \sum_{k_1, \dots, k_{2l}=1}^{\infty} k_{2a-1} \check{T}_0 \right. \\
&\quad \left. \cdot \frac{\partial}{\partial t_{k_1, \dots, k_{2a-1}+n, k_{2a}, \dots, k_{2l}}} \right] - N^2 t_1, \\
\check{W}_2 &= \sum_{n,m=0}^{\infty} \sum_{a=1}^{n-1} \check{T}_1 \left\{ \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} \check{T}_0 \left[\sum_{a=2}^l \sum_{s=0}^{k_{2a-1}} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}, \xi_0, k_{2a}, \dots, k_{2l}}} + \sum_{s=0}^{k_1} \frac{\partial}{\partial t_{\xi_1, k_2, \dots, k_{2l}}} \right] \right. \\
&\quad \left. + \frac{\partial}{\partial t_{a,m+1}} \frac{\partial}{\partial t_{n-a}} - N \delta_{n,0} \frac{\partial}{\partial g_{m+1}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k, m+1}} \right\}, \\
\check{W}_3 &= \sum_{n_1, \dots, n_{2r+1}=0}^{\infty} \check{T}_2 \left\{ \left[\sum_{s=1}^{n_{2r+1}-1} \frac{\partial}{\partial t_{s, n_2+1, \dots, n_{2r+1}}} \frac{\partial}{\partial t_{n_{2r+1}-s}} + \sum_{b=1}^{r-1} \sum_{s=1}^{n_{2b+1}} \frac{\partial}{\partial t_{s, n_2+1, \dots, n_{2b+1}}} \right. \right. \\
&\quad \left. \left. \cdot \frac{\partial}{\partial t_{\xi_2, n_{2b+2}+1, \dots, n_{2r+1}}} \right] - N \delta_{n_{2r+1}, 0} \frac{\partial}{\partial t_{n_3+1, \dots, n_{2r-1}+1, \xi_3}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n_{2r+1}+k, n_2+1, \dots, n_{2r+1}}} \right. \\
&\quad \left. + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} \check{T}_0 \left[\sum_{s=0}^{k_1} \frac{\partial}{\partial t_{s+1, n_2+1, \dots, \xi_4, k_2, \dots, k_{2l}}} + \sum_{a=2}^l \sum_{s=0}^{k_{2a-1}} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}, s+1, \xi_5, k_{2a}, \dots, k_{2l}}} \right] \right\}, \\
\check{W}_4 &= \sum_{m=0}^{\infty} (m+1) g_{m+1} \left[\sum_{a=1}^{m-1} \frac{\partial}{\partial g_a} \frac{\partial}{\partial g_{m-a}} + \sum_{k=0}^{\infty} k g_k \frac{\partial}{\partial g_{m+k}} + \sum_{l=1}^{\infty} \sum_{a=1}^l \sum_{k_1, \dots, k_{2l}=1}^{\infty} k_{2a} \check{T}_0 \right. \\
&\quad \left. \cdot \frac{\partial}{\partial t_{k_1, \dots, k_{2a+m}, k_{2a+1}, \dots, k_{2l}}} \right] - N^2 g_1,
\end{aligned} \tag{31}$$

where $\check{T}_0 = t_{k_1, \dots, k_{2l}}$, $\check{T}_1 = (n+m+2)t_{n+1, m+1}$, $\check{T}_2 = \bar{N}t_{n_{2r+1}+1, n_2+1, \dots, n_{2r+1}}$ and $\xi_0 = (a+1, m+1, n+k_{2a-1}-k-1)$, $\xi_1 = (s+1, m+1, n+k_1-s-1)$, $\xi_2 = n_{2r+1}+n_{2b+1}+1-s$, $\xi_3 = n_{2r}+n_2+2$, $\xi_4 = n_{2r+1}+k_1-1-s$, $\xi_5 = (n_2+1, \dots, n_{2r}+1, n_{2r+1}+k_{2a-1}-s-1)$.

We find that the fermionic two-matrix model (26) can be realized by the W -representation

$$Z_{2F} = e^{\frac{1}{\mu} \check{W}} e^{N(t_0+g_0)}. \tag{32}$$

The compact expression of correlators is

$$\begin{aligned}
&\left\langle \prod_{i=1}^{l_1} \text{tr}(\bar{\psi}\psi)^{k_i} \prod_{j=1}^{l_2} \text{tr}(\bar{\chi}\chi)^{r_j} \prod_{n=1}^{l_3} \text{tr}(\bar{\psi}\psi)^{S_{n,1}} (\bar{\chi}\chi)^{S_{n,2}} \dots (\bar{\psi}\psi)^{S_{n,2p_n-1}} (\bar{\chi}\chi)^{S_{n,2p_n}} \right\rangle \\
&= \frac{l_1! l_2! l_3! \sum_{\sigma}^{\rho_1+\rho_2+\rho_3=1} \sum_{\sigma}^{m+1} \check{P}_{(\sigma(k_1), \dots, \sigma(k_{l_1})); (\sigma(r_1), \dots, \sigma(r_{l_2}))}^{(\sigma(S_{1,1}), \dots, \sigma(S_{1,2p_1}); \dots; \sigma(S_{l_3,1}), \dots, \sigma(S_{l_3,2p_{l_3}}))}}{\mu^{m+1} (m+1)! \lambda_{(k_1, \dots, k_{l_1})} \lambda_{(r_1, \dots, r_{l_2})} \lambda_{(S_{1,1}, \dots, S_{1,2p_1}; \dots; S_{l_3,1}, \dots, S_{l_3,2p_{l_3}})}},
\end{aligned} \tag{33}$$

where $\rho_1 = \sum_{i=1}^{l_1} k_i$, $\rho_2 = \sum_{i=1}^{l_2} r_i$, $\rho_3 = \sum_{i=1}^{l_3} (S_{i,1} + \dots + S_{i,2p_i})$, and $\check{P}_{(\sigma(k_1), \dots, \sigma(k_{l_1})); (\sigma(r_1), \dots, \sigma(r_{l_2}))}^{(\sigma(S_{1,1}), \dots, \dots, \sigma(S_{l_3,2p_{l_3}}))}$ is the coefficient of $t_{k_1} \dots t_{k_{l_1}} g_{r_1} \dots g_{r_{l_2}} t_{S_{1,1}, \dots, S_{1,2p_1}} \dots t_{S_{l_3,1}, \dots, S_{l_3,2p_{l_3}}}$ in \check{W}^{m+1} .

Here we list some correlators

$$\begin{aligned}
\langle \text{tr} \bar{\psi} \psi \rangle &= \langle \text{tr} \bar{\chi} \chi \rangle = -\frac{1}{\mu} N^2, \quad \langle \text{tr} \bar{\psi} \psi \text{tr} \bar{\chi} \chi \rangle = \frac{2}{\mu^2} N^4, \\
\langle \text{tr} \bar{\psi} \psi \text{tr} \bar{\psi} \psi \rangle &= \langle \text{tr} \bar{\chi} \chi \text{tr} \bar{\chi} \chi \rangle = \frac{1}{\mu^2} (N^2 - 1) N^2, \quad \langle \text{tr} \bar{\psi} \psi \bar{\chi} \chi \rangle = -\frac{2}{\mu^2} N^3, \\
\langle \text{tr} (\bar{\psi} \psi)^3 \rangle &= \langle \text{tr} (\bar{\chi} \chi)^3 \rangle = \frac{6}{\mu^3} (-N^2 + N^4), \\
\langle \text{tr} \bar{\psi} \psi \text{tr} \bar{\psi} \psi \text{tr} \bar{\psi} \psi \rangle &= \langle \text{tr} \bar{\chi} \chi \text{tr} \bar{\chi} \chi \text{tr} \bar{\chi} \chi \rangle = \frac{1}{\mu^3} (N^2 + 2)(N^2 - 1) N^2, \\
\langle \text{tr} \bar{\psi} \psi \text{tr} \bar{\psi} \psi \text{tr} \bar{\chi} \chi \rangle &= \langle \text{tr} \bar{\chi} \chi \text{tr} \bar{\psi} \psi \text{tr} \bar{\chi} \chi \rangle = \frac{3}{\mu^3} (N^4 - N^6), \\
\langle \text{tr} \bar{\psi} \psi \text{tr} \bar{\psi} \psi \bar{\chi} \chi \rangle &= \langle \text{tr} \bar{\chi} \chi \text{tr} \bar{\psi} \psi \bar{\chi} \chi \rangle = -\frac{1}{\mu^3} (6N^3 + 2N^5).
\end{aligned} \tag{34}$$

5 Conclusion

We have constructed the Hermitian, complex and fermionic two-matrix models with infinite set of variables and presented their Virasoro constraints. W -representation is important for understanding matrix model, since it provides a dual formula for partition function through differentiation. By considering the particular infinitesimal transformations of integration variables in the partition functions, we finally derived the desired operators preserving and increasing the grading. Thus it can be shown that the two-matrix models constructed in this paper can be realized by the W -representations. Moreover, by means of the W -representations, we derived the compact expressions of correlators for these two-matrix models. It should be noted that there are the infinite set of variables in these two-matrix models. It leads to that we can not give their character expansions. For further research, it would be interesting to study the case of β -deformed two-matrix models.

Appendix A The operators \hat{W}_i in (5)

$$\begin{aligned}
\hat{W}_1 &= \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} \left\{ t_1 T_1 [\delta_{k_1,1} \frac{\partial}{\partial t_{k_3, \dots, k_{2l-1}, k_{2l}+k_2}} + \sum_{a=2}^l \delta_{k_{2a-1},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}+k_{2a}, \dots, k_{2l}}} \right. \\
&\quad \left. + \sum_{n=0}^{\infty} \sum_{a=1}^l k_{2a-1} (n+1) t_{n+1} \frac{\partial}{\partial t_{k_1, \dots, n+k_{2a-1}-1, \dots, k_{2l}}} \right\} + \sum_{k_2=1}^{\infty} t_1 t_{1,k_2} \frac{\partial}{\partial g_{k_2}} + t_1^2 N + 2t_2 N^2 \\
&\quad + \sum_{n=0}^{\infty} (n+1) t_{n+1} \left[\sum_{b=1}^{n-2} \frac{\partial^2}{\partial t_b \partial t_{n-1-b}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n+k-1}} \right] + \sum_{n=2}^{\infty} 2N(n+1) t_{n+1} \frac{\partial}{\partial t_{n-1}}, \\
\hat{W}_2 &= \sum_{l,n=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} T_2 \left\{ (1 - \delta_{k_1,1}) \left(\frac{\partial}{\partial t_{k_1-1, k_2, \dots, k_{2l}+n}} + \frac{\partial}{\partial t_{k_1-1, k_2+n, \dots, k_{2l}}} \right) \right. \\
&\quad \left. + \delta_{k_1,1} \frac{\partial}{\partial t_{k_3, \dots, k_{2l}+n+k_2}} + \sum_{a=2}^l (1 - \delta_{k_{2a-1},1}) \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}+n, k_{2a-1}-1, k_{2a}, \dots, k_{2l}}} \right. \\
&\quad \left. + \sum_{a=2}^{l-1} \delta_{k_{2a-1},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}+k_{2a}+n, \dots, k_{2l}}} \right\} + \sum_{l=2}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} T_2 \delta_{k_{2l-1},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2l-2}+n+k_{2l}}} \\
&\quad + \sum_{l,n=1}^{\infty} \sum_{a=1}^l \sum_{k_{2a-1}=3}^{\infty} \sum_{b=1}^{k_{2a-1}-2} \sum_{\substack{k_1, \dots, k_{2a-2}, \\ k_{2a}, \dots, k_{2l}=1}}^{\infty} T_2 [(1 - \delta_{a,1}) \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}, b, n, k_{2a-1}-1-b, \dots, k_{2l}}} \\
&\quad + \delta_{a,1} \frac{\partial}{\partial t_{b, n, k_1-1-b, \dots, k_{2l}}}] + \sum_{n=1}^{\infty} (n+1) t_{1,n} \left[\sum_{k=2}^{\infty} k t_k \frac{\partial}{\partial t_{k-1,n}} + \sum_{k_2=1}^{\infty} t_{1,k_2} \frac{\partial}{\partial g_{k_2+n}} + t_1 \frac{\partial}{\partial g_n} \right],
\end{aligned}$$

$$\begin{aligned}
\hat{W}_3 &= \sum_{r=1}^{\infty} \sum_{n_1, \dots, n_{2r}=1}^{\infty} T_3 \left\{ N \delta_{n_1,1} \frac{\partial}{\partial t_{n_3, \dots, n_{2r}+n_2}} + (1 - \delta_{n_1,1}) \left(\sum_{a=1}^{n_1-2} \frac{\partial}{\partial t_a} \frac{\partial}{\partial t_{n_1-1-a, \dots, n_{2r}}} \right. \right. \\
&\quad + N \frac{\partial}{\partial t_{n_1-1, n_2, \dots, n_{2r}}} + \frac{\partial}{\partial t_{n_1-1}} \frac{\partial}{\partial t_{n_3, \dots, n_{2r}}} \left. \right) + \sum_{s=2}^r \sum_{a=0}^{n_{2s-1}-2} (1 - \delta_{n_{2s-1},1}) \frac{\partial}{\partial t_{n_1+a, n_2, \dots, n_{2s-2}}} \\
&\quad \cdot \frac{\partial}{\partial t_{n_{2s-1}-1-a, \dots, n_{2r}}} + \sum_{s=2}^{r-1} [\delta_{n_{2s-1},1} \frac{\partial}{\partial t_{n_1, \dots, n_{2s-2}}} + (1 - \delta_{n_1,1}) \frac{\partial}{\partial t_{n_1+n_{2s-1}-1, n_2, \dots, n_{2s-2}}} \\
&\quad \cdot \frac{\partial}{\partial t_{n_{2s+1}, \dots, n_{2r}+n_{2s}}} + (1 - \delta_{n_{2r-1},1}) \frac{\partial}{\partial t_{n_1+n_{2r-1}-1, \dots, n_{2r-2}}} \frac{\partial}{\partial g_{n_{2r}}} + \sum_{k=0}^{\infty} k t_k \frac{\partial}{\partial t_{n_1+k-1, \dots, n_{2r}}} \\
&\quad + \delta_{n_{2r-1},1} \frac{\partial^2}{\partial t_{n_1, \dots, n_{2r-2}} \partial g_{n_{2r}}} + \sum_{k_1, \dots, k_{2l}=1}^{\infty} T_1 \left[\sum_{i=1}^l \sum_{s=0}^{k_{2i-1}-2} \frac{\partial}{\partial t_{k_1, \dots, k_{2i-2}, s+n_1, \dots, n_{2r}, \xi_1, \dots, k_{2l}}} \right. \\
&\quad \left. + \frac{\partial}{\partial t_{n_1+k_1-1, n_2, \dots, n_{2r}+k_2, \dots, k_{2l}}} + \frac{\partial}{\partial t_{k_1, \dots, n_1+k_{2i-1}-1, \dots, n_{2r}+k_{2i}, \dots, k_{2l}}} \right] \\
&\quad + \sum_{n_1=2}^{\infty} \sum_{n_2=1}^{\infty} (n_1 + 1 + n_2) t_{n_1+1, n_2} \frac{\partial^2}{\partial t_{n_1-1} \partial g_{n_2}}, \\
\hat{W}_4 &= \sum_{r=1}^{\infty} \sum_{n_2, \dots, n_{2r}=1}^{\infty} T_4 \left\{ \sum_{b=1}^{n_3-2} \frac{\partial^2}{\partial t_{b, n_2} \partial t_{n_3-1-b, \dots, n_{2r}}} + (1 - \delta_{n_3,1}) \left(\frac{\partial^2}{\partial g_{n_2} \partial t_{n_3-1, \dots, n_{2r}}} \right. \right. \\
&\quad + \frac{\partial^2}{\partial t_{n_3-1, n_2, \partial t_{n_5, \dots, n_{2r}+n_4}}} + \delta_{n_3,1} \frac{\partial^2}{\partial g_{n_2} \partial t_{n_5, \dots, n_{2r}+n_4}} + \sum_{a=3}^{r-1} [\delta_{n_{2a-1},1} \frac{\partial}{\partial t_{n_3, \dots, n_{2a-2}+n_2}} \\
&\quad \cdot \frac{\partial}{\partial t_{n_{2a+1}, \dots, n_{2r}}} + (1 - \delta_{n_{2a-1},1}) \left(\frac{\partial}{\partial t_{n_3, \dots, n_{2a-1}-1, n_{2a}}} \frac{\partial}{\partial t_{n_{2a+1}, \dots, n_{2r}+n_{2a}}} + \frac{\partial}{\partial t_{n_3, \dots, n_{2a-2}+n_2}} \right. \\
&\quad \left. \left. \cdot \frac{\partial}{\partial t_{n_{2a-1}-1, n_{2a}, \dots, n_{2r}}} \right)] + [\delta_{n_{2r-1},1} \frac{\partial}{\partial t_{n_3, \dots, n_{2r-2}+n_2}} + (1 - \delta_{n_{2r-1},1}) \frac{\partial}{\partial t_{n_3, \dots, n_{2r-1}-1, n_2}}] \frac{\partial}{\partial g_{n_{2r}}} \right. \\
&\quad + \sum_{a=3}^r \sum_{b=1}^{n_{2a-1}-2} \frac{\partial}{\partial t_{n_3, \dots, n_{2a-2}, b, n_2}} \frac{\partial}{\partial t_{n_{2a-1}-1-b, \dots, n_{2r}}} + t_1 \frac{\partial}{\partial t_{n_3, \dots, n_{2r}+n_2}} + \sum_{k=2}^{\infty} k t_k \frac{\partial}{\partial t_{k-1, n_2, \dots, n_{2r}}} \\
&\quad + \sum_{l=1}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} T_1 [\delta_{k_1,1} \frac{\partial}{\partial t_{n_3, \dots, n_{2r}+k_2, \dots, k_{2l}+n_2}} + (1 - \delta_{k_1,1}) \left(\frac{\partial}{\partial t_{k_1-1, n_2, \dots, n_{2r}+k_2, \dots, k_{2l}}} \right. \\
&\quad \left. + \frac{\partial}{\partial t_{n_3, \dots, n_{2r}, k_1-1, \dots, k_{2l}+n_2}} \right) + \sum_{b=2}^l (1 - \delta_{k_{2b-1},1}) \left(\frac{\partial}{\partial t_{k_1, \dots, k_{2b-3}, \xi_2, \dots, k_{2l}}} + \frac{\partial}{\partial t_{k_1, \dots, \xi_3, \dots, n_{2r}+k_{2l}}} \right) \\
&\quad + \sum_{b=2}^l \delta_{k_{2b-1},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2b-1}+n_2, \xi_4, \dots, k_{2l}}} + \sum_{l=1}^{\infty} \sum_{a=1}^l \sum_{k_{2a-1}=3}^{\infty} \sum_{\substack{k_1, \dots, k_{2a-2}, \\ k_{2a}, \dots, k_{2l}=1}}^{\infty} \sum_{s=1}^{k_{2a-1}-2} T_1 \frac{\partial}{\partial t_{k_1, \dots, k_{2a-2}, s, \xi_5, \dots, k_{2l}}}], \\
\hat{W}_5 &= g_1^2 N + \sum_{n,k=0}^{\infty} (n+1) g_{n+1} k g_k \frac{\partial}{\partial g_{n+k-1}} + \sum_{k_l, k_2, k_3=1}^{\infty} g_1 t_{k_1, k_2, k_3, 1} \frac{\partial}{\partial t_{k_1+k_3, k_2}} + \sum_{k_1=1}^{\infty} t_{k_1, 1} g_1 \frac{\partial}{\partial t_{k_1}} \\
&\quad + \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} \sum_{k_1, \dots, k_{2l}=1}^{\infty} T_5 [\delta_{k_{2l},1} \sum_{a=1}^{l-1} \frac{\partial}{\partial t_{k_1+k_{2l-1}, \dots, k_{2l-2}}} k_{2a} \frac{\partial}{\partial t_{k_1, \dots, \xi_6, \dots, k_{2l}}} + \frac{\partial}{\partial t_{k_1+k_{2l-1}, \dots, k_{2l-2}}} \\
&\quad \cdot \frac{\partial}{\partial t_{k_1, \dots, k_{2l-1}, \xi_7}} + \delta_{n,0} (1 - \delta_{k_{2a},1}) \frac{\partial}{\partial t_{k_1, k_2-1, \dots, k_{2l}}} + \delta_{n,0} \sum_{a=1}^{l-1} \delta_{k_{2a},1} \frac{\partial}{\partial t_{k_1, \dots, k_{2a-1}+k_{2a+1}, \dots, k_{2l}}}]
\end{aligned}$$

$$+ \sum_{n=1}^{\infty} \sum_{s=1}^{n-2} (n+1)g_{n+1} \frac{\partial}{\partial g_s} \frac{\partial}{\partial g_{n-1-s}} + \sum_{n=2}^{\infty} 2N(n+1)g_{n+1} \frac{\partial}{\partial g_{n-1}} + 2g_2 N^2, \quad (\text{A.1})$$

where $T_1 = t_{k_1, \dots, k_{2l}}$, $T_2 = (n+1)t_{1,n}t_{k_1, \dots, k_{2l}}$, $T_3 = \mathcal{N}_1 t_{n_1+1, n_2, \dots, n_{2r}}$, $T_4 = \mathcal{N}_2 t_{1, n_2, \dots, n_{2r}}$, $T_5 = (n+1)g_{n+1}t_{k_1, \dots, k_{2l}}$ and $\xi_1 = k_{2i-1} - 1 - s$, $\xi_2 = (k_{2b-2} + n_2, \dots, n_{2r}, k_{2b-1} - 1)$, $\xi_3 = (k_{2b-1} - 1, n_2)$, $\xi_4 = (n_3, \dots, n_{2r} + k_{2b})$, $\xi_5 = (n_2, \dots, n_{2r}, k_{2a-1} - 1 - s)$, $\xi_6 = k_{2a} + n - 1$, $\xi_7 = k_{2l} + n - 1$.

Acknowledgment

This work is supported by the National Natural Science Foundation of China (No. 11875194).

References

- [1] F. David, Planar diagrams, two-dimensional lattice gravity and surface models, *Nucl. Phys. B* **45** (1985) 257.
- [2] S. Chadha, G. Mahoux and M.L. Mehta, A method of integration over matrix variables, *J. Phys. A: Math. Gen.* **14** (1981) 579.
- [3] V.A. Kazakov, Ising model on dynamical planar random lattice: exact solution, *Phys. Lett. A* **119** (1986) 140.
- [4] G.W. Semenoff and R.J. Szabo, Fermionic matrix models, *Int. J. Mod. Phys. A* **12** (1997) 2135 [arXiv:9605140].
- [5] A. Marshakov, A. Mironov and A. Morozov, From Virasoro constraints in Kontsevich's model to W -constraints in two-matrix models, *Mod. Phys. A* **07** (1992) 1345-1359 [arXiv:9201010].
- [6] V.A. Kazakov and A. Marshakov, Complex curve of the two matrix model and its tau-function, *J. Phys. A: Math. Gen.* **36** (2003) 3107 [arXiv:0211236].
- [7] B. Eynard and N. Orantin, Topological expansion of the 2-matrix model correlation functions: diagrammatic rules for a residue formula, *J. High Energy Phys.* **12** (2005) 034 [arXiv:math-ph/0504058].
- [8] M. Bergère, B. Eynard, O. Marchal and A. Prats-Ferrer, Loop equations and topological recursion for the arbitrary- β two-matrix model, *J. High Energy Phys.* **03** (2012) 098 [arXiv:1106.0332].
- [9] A. Morozov and Sh. Shakirov, Generation of matrix models by \hat{W} -operators, *J. High Energy Phys.* **04** (2009) 064 [arXiv:0902.2627].
- [10] R. Wang, F. Liu, C.H. Zhang and W.Z. Zhao, Superintegrability for (β -deformed) partition function hierarchies with W -representations, *Eur. Phys. J. C* **82** (2022) 902 [arXiv:2206.13038].
- [11] R. Wang, F. Liu, M.L. Li and W.Z. Zhao, Superintegrability for super partition function hierarchies with W -representations, arXiv:2208.03671.

- [12] H. Itoyama, A. Mironov and A. Morozov, Complete solution to Gaussian tensor model and its integrable properties, *Phys. Lett. B* **802** (2020) 135237 [arXiv:1910.03261].
- [13] B. Kang, L.Y. Wang, K. Wu, J. Yang and W.Z. Zhao, W -representation of rainbow tensor model, *J. High Energy Phys.* **05** (2021) 228 [arXiv:2104.01332].
- [14] L.Y. Wang, R. Wang, K. Wu and W.Z. Zhao, W -representations of the fermionic matrix and Aristotelian tensor models, *Nucl. Phys. B* **973** (2021) 115612 [arXiv:2110.14269].
- [15] B. Kang, L.Y. Wang, K. Wu and W.Z. Zhao, Rainbow tensor model with two tensors of rank three, arXiv:2301.06046.
- [16] Y. Chen, R. Wang, K. Wu and W.Z. Zhao, Correlators in the supereigenvalue model in the Ramond sector, *Phys. Lett. B* **807** (2020) 135563 [arXiv:2006.11013].
- [17] R. Wang, S.K. Wang, K. Wu and W.Z. Zhao, Correlators in the Gaussian and chiral supereigenvalue models in the Neveu-Schwarz sector, *J. High Energy Phys.* **11** (2020) 119 [arXiv:2009.02929].
- [18] A. Alexandrov, On W -operators and superintegrability for dessins d'enfant, arXiv:2212.10952.
- [19] A. Mironov, V. Mishnyakov, A. Morozov, A. Popolitov, R. Wang and W.Z. Zhao, Interpolating matrix models for WLZZ series, arXiv:2301.04107.
- [20] A. Mironov, V. Mishnyakov, A. Morozov, A. Popolitov and W.Z. Zhao, On KP-integrable skew Hurwitz τ -functions and their β -deformations, arXiv:2301.11877.
- [21] L.Y. Wang, V. Mishnyakov, A. Popolitov, F. Liu and R. Wang, W -representations for multi-character partition functions and their β -deformations, arXiv:2301.12763.