

Wiener densities for the Airy line ensemble

Duncan Dauvergne

February 2, 2023

Abstract

The parabolic Airy line ensemble \mathfrak{A} is a central limit object in the KPZ universality class and related areas. On any compact set $K = \{1, \dots, k\} \times [a, a + t]$, the law of the recentered ensemble $\mathfrak{A} - \mathfrak{A}(a)$ has a density X_K with respect to the law of k independent Brownian motions. We show that

$$X_K(f) = \exp(-S(f) + o(S(f)))$$

where S is an explicit, tractable, non-negative function of f . We use this formula to show that X_K is bounded above by a K -dependent constant, give a sharp estimate on the size of the set where $X_K < \epsilon$ as $\epsilon \rightarrow 0$, and prove a large deviation principle for \mathfrak{A} . We also give density estimates that take into account the relative positions of the Airy lines, and prove sharp two-point tail bounds that are stronger than those for Brownian motion. The paper is essentially self-contained, requiring only tail bounds on the Airy point process and the Brownian Gibbs property as inputs.

1 Introduction

The Airy line ensemble is a stationary random sequence of functions $\mathcal{A} = \{\mathcal{A}_i : \mathbb{R} \rightarrow \mathbb{R}, i \in \mathbb{N}\}$ satisfying $\mathcal{A}_1 > \mathcal{A}_2 > \dots$. It was first introduced by Prähofer and Spohn [38] via a determinantal formula, see (7) below. Prähofer and Spohn showed that the top line \mathcal{A}_1 —known as the Airy (or Airy₂) process—describes the scaling limit at a fixed time for a certain one-dimensional random growth model in the KPZ (Kardar-Parisi-Zhang) universality class started from a point. The remaining lines describe the scaling limit of a richer multi-layer random growth model. Since this work, the Airy process and the Airy line ensemble have been established to be universal limit objects in the KPZ universality class and for other closely related branches of probability. Among other results, we now know that:

- (i) The Airy process and the Airy line ensemble appear as one-parameter scaling limits of classical solvable random metric and random growth models in the KPZ universality class (e.g. zero temperature models including exponential/geometric/Poisson/Brownian last passage percolation and tasep, see [14, 16, 34, 38]; positive temperature models including the KPZ equation and the KPZ line ensemble, see [15, 40, 43, 44]). See also [10, 13, 24, 39, 41] and references therein for general background on the KPZ universality class and related areas.

- (ii) The richest scaling limit in the KPZ universality class—the directed landscape—can be described by a certain metric-like last passage structure on the Airy line ensemble, see [17]. Moreover, at least for last passage models, convergence to the Airy line ensemble is the only integrable input required for convergence to the directed landscape, see [20].
- (iii) The Airy line ensemble is the limit as $n \rightarrow \infty$ at the edge of a systems of n non-intersecting Brownian motions, see [2, 14]. In particular, this implies that it is limit at the edge for the eigenvalue process for Hermitian-matrix valued Brownian motion (Dyson’s Brownian motion with $\beta = 2$).
- (iv) The Airy line ensemble appears as the limit at the boundary between the frozen and liquid regions for a very general class of random lozenge tilings, see [4], also [22, 35–37] for earlier convergence results and work on related models. The Airy kernel has also been found at the more delicate liquid-gas boundary in the Aztec diamond [7, 8].

While the determinantal representation of the Airy line ensemble is useful for the definition, it can be difficult to work with in practice and does not give a good intuition for what the Airy line ensemble looks like. A better intuition is to think of \mathcal{A} as a system of infinitely many non-intersecting Brownian motions (i.e. a limit of (iii) above).

This idea was made rigorous by Corwin and Hammond [14] by identifying a **Brownian Gibbs property** for the Airy line ensemble. To describe this property, it is more natural to work with the **parabolic Airy line ensemble** $\mathfrak{A}_1 > \mathfrak{A}_2 > \dots$, where $\mathfrak{A}_i(t) = \mathcal{A}_i(t) - t^2$. The Brownian Gibbs property states that inside any region $K = \{1, \dots, k\} \times [a, b]$, conditionally on all values $\mathfrak{A}_i(t)$ for $(i, t) \notin K$, the parabolic Airy line ensemble on K is simply given by a sequence of k independent Brownian bridges (of variance 2) from $(a, \mathfrak{A}_i(a))$ to $(b, \mathfrak{A}_i(b))$ conditioned so that the entire ensemble remains non-intersecting ¹

This property turns out to be an extremely useful tool for understanding the Airy line ensemble. In fact, the Brownian Gibbs property is necessary for even rigorously showing that the determinantal formula of Prähofer and Spohn defines a system of non-intersecting lines. The Brownian Gibbs property also implies that that recentered Airy lines $\mathfrak{A}_i - \mathfrak{A}_i(0)$ are locally absolutely continuous with respect to Wiener measure (of variance 2), which gives information about the sample paths of \mathfrak{A} that seem to be extremely difficult to access with the determinantal formula alone.

Given that recentered Airy lines are locally absolutely continuous with respect to Wiener measure, the next natural question is to as exactly how this occurs. More precisely, we would like to find the density (i.e. the Radon-Nikodym derivative) of the Airy lines against Wiener measure. This is the goal of the present paper.

To state our main result, for $k \in \mathbb{N}$ and $t > 0$, let $\mathcal{C}_0^k([0, t])$ denote the space of k -tuples of continuous functions $f = (f_1, \dots, f_k), f_i : [0, t] \rightarrow \mathbb{R}$ with $f(0) = 0$, equipped with the topology of uniform convergence. Let $\nu_{k,t}$ be the law on $\mathcal{C}_0^k([0, t])$ of

$$\mathfrak{A}_i(r) - \mathfrak{A}_i(0), \quad r \in [0, t], i \in \llbracket 1, k \rrbracket, \tag{1}$$

¹Here we say that a Brownian bridge has variance v if its quadratic variation over any interval $[c, d]$ equals $v(d - c)$.

and let $\mu_{k,t}$ be the law on $\mathcal{C}_0^k([0,t])$ of a k -tuple of independent Brownian motions of variance 2. In (1) and throughout the paper we write $[[\ell,k]] := [\ell,k] \cap \mathbb{Z}$. Let $X_{k,t}$ be the density of $\nu_{k,t}$ against $\nu_{k,t}$.

Theorem 1.1. *Fix $k \in \mathbb{N}, t \geq 1$. Then there exists a function $S: \mathcal{C}_0^k([0,t]) \rightarrow [0, \infty)$ such that for $\mu_{k,t}$ -a.e. $f \in \mathcal{C}_0^k([0,t])$ we have*

$$X_{k,t}(f) = \exp(-S(f) + O_k(t^3 + \sqrt{t}S(f)^{5/6})). \quad (2)$$

In Theorem 1.1 and throughout the paper, we write $O_k(g)$ for a term that is bounded in absolute value by g times a k -dependent constant.

The function S is given as follows. First, let $\mathcal{C}^k([0,t])$ denote the space of k -tuples of continuous functions from $[0,t] \rightarrow \mathbb{R}$ with the topology of uniform convergence (without a condition on $f(0)$). For $f \in \mathcal{C}^k([0,t])$, let $\mathbb{T}f \in \mathcal{C}^k([0,t])$ be the minimal k -tuple of functions such that

$$\mathbb{T}f_1(r) \geq \mathbb{T}f_2(r) \geq \dots \geq \mathbb{T}f_k(r) \geq 0$$

for all $r \in [0,t]$ and such that $\mathbb{T}f - \mathbb{T}f(0) = f - f(0)$. Here and throughout we write $\mathbb{T}f_i = (\mathbb{T}f)_i, \mathbb{T}f(0) = (\mathbb{T}f)(0)$, etc. We can think of $\mathbb{T}f$ as the ‘‘Tetris of f ’’: the result of consecutively dropping the graphs of the functions f_k, f_{k-1}, \dots, f_1 onto a flat line, and letting them stack up on top of each other as in the game of Tetris, see Figure 1. Then

$$S(f) = \frac{2}{3} \left(\sum_{i=1}^k [\mathbb{T}f_i(0)]^{3/2} + [\mathbb{T}f_i(t)]^{3/2} \right).$$

Remark 1.2 (*t -dependence in Theorem 1.1*). We impose the restriction that $t \geq 1$ throughout the paper in order to simplify our expression of the error term. Naturally, Theorem 1.1 immediately implies that similar bounds hold for $t < 1$ by pushing forward the measures $\mu_{k,1}, \nu_{k,1}$ from $\mathcal{C}_0^k([0,1])$ to $\mathcal{C}_0^k([0,t])$ under the projection $f \mapsto f|_{[0,t]}$. Note that (2) also implies estimates on intervals of the form $[a, a+t]$ by the stationarity of $\mathcal{A}(t) = \mathfrak{A}(t) + t^2$.

The t^3 factor in the error is optimal as $t \rightarrow \infty$. Because of the stationarity of the Airy line ensemble $\mathcal{A}(r) = \mathfrak{A}(r) + r^2$ the law $\nu_{k,t}$ concentrates around functions that are relatively close to the parabola $-r^2$. On the other hand, the probability that a Brownian motion follows this parabola up to time t is easily checked to be $e^{-\Theta(t^3)}$.




Figure 1: A function $f \in \mathcal{C}_0^3([0,t])$ and its image under the map \mathbb{T} .

1.1 Motivation and some applications

Since the work [14], the Brownian Gibbs property and the local Brownian structure of the Airy line ensemble has been repeatedly exploited to prove results about the Airy process, the Airy line ensemble, and more general limit objects in the KPZ universality class, e.g. see [14, 31] for some early example and [27] for a survey with more recent applications. In particular, the construction of the directed landscape [17] relies heavily on Brownian Gibbs analysis from [19].

While it is quite useful to know that the Airy line ensemble is locally absolutely continuous with respect to Brownian motion, there are many situations when a more quantitative comparison is required. This approach was first taken up by Hammond [32] and then continued by Calvert, Hegde, and Hammond [11]. The culmination of these two papers is the following estimate from [11]. For any $k \in \mathbb{N}$, $t > 0$, and any Borel set $A \subset \mathcal{C}_0^1([0, t])$ we have

$$\mathbb{P}(\mathfrak{A}_k(\cdot) - \mathfrak{A}_k(0) \in A) \leq 2\mu_t(A) \exp(d_{k,t} \log^{5/6}[\mu_t(A)^{-1}]), \quad (3)$$

where $d_{k,t}$ is a k, t -dependent constant and $\mu_t = \mu_{1,t}$ is Wiener measure of variance 2 on $\mathcal{C}_0^1([0, t])$.

While this seems like a fairly technical statement, over the past five years it has proven to be extremely useful. Indeed, since the decay in (3) is faster than any polynomial, it shows that events for Brownian motion that occur with probability ϵ occur only with probability $\epsilon^{1-o(1)}$ for the k th Airy line. This is a key estimate for ruling out pathological behaviour of the Airy line ensemble, and has been used repeatedly for understanding limiting random geometry in the KPZ universality class, e.g. see [5, 6, 9, 18, 21, 25, 26, 29–31, 33].

Theorem 1.1 strengthens the estimate (3) In fact, it implies something more surprising: namely, that the Wiener density $X_{k,t}$ is bounded!

Corollary 1.3. *For every $k \in \mathbb{N}, t \geq 1$, we have that $X_{k,t}(f) \leq e^{O_k(t^3)}$ for $\mu_{k,t}$ -a.e. $f \in \mathcal{C}_0^k([0, t])$. In other words, $\nu_{k,t}(A) \leq e^{O_k(t^3)} \mu_{k,t}(A)$ for any Borel set $A \subset \mathcal{C}_0^k([0, t])$.*

It is worth noting that our proof of Theorem 1.1 and Corollary 1.3 is much shorter than the proof of (3) from [11, 32], and hence gives an easier route to this cornerstone result and its consequences.

In a similar vein, we can also identify how small $\nu_{k,t}(A)$ can be given $\mu_{k,t}(A)$.

Corollary 1.4. *Define the function $\Theta : \mathbb{R}^k \rightarrow \mathbb{R}$ by*

$$\Theta(\mathbf{x}) = \frac{2}{3} \left(\sum_{i=1}^k |x_i| \right)^{3/2} + \frac{4}{3} \sum_{j=2}^k \left(\sum_{i=j}^k |x_i| \right)^{3/2},$$

and let α_k be the maximum of Θ on the sphere $\|\mathbf{x}\|_2 = 1$. Then

$$\inf\{\nu_{k,t}(A) : \mu_{k,t}(A) = \epsilon\} = \epsilon \exp\left(-\alpha_k [4t \log(\epsilon^{-1})]^{3/4} + O_k(t^3 + t \log^{5/8}(\epsilon^{-1}))\right).$$

For general k , the minimum value of Θ does not have a nice closed form. However, when $k = 1$, then $\Theta(x) = 2/3$ when $x = \pm 1$. The proof of Corollary 1.4 also gives a sense of what types of sets will come close to achieving the supremum above.

Remark 1.5. To give some sense of the fragility and strength of the estimate in Theorem 1.1, it is interesting to note that Corollary 1.3 fails if we replace \mathfrak{A} by the stationary Airy line ensemble $\mathcal{A}(t) = \mathfrak{A}(t) + t^2$ (though Corollary 1.4 will still hold). See Proposition 7.3 for details.

As another application, we can quickly deduce a large deviation principle for \mathfrak{A} by using Corollaries 1.3 and 1.4 and Schilder's theorem. Note that this does not require the full strength of the above results, only that $X_{k,t}, X_{k,t}^{-1} \in L^p(\mu_{k,t})$ for all $p \in (1, \infty)$.

Corollary 1.6. *For every $\epsilon > 0$, let $\nu_{k,t}^\epsilon$ denote the pushforward on $\mathcal{C}_0^k([0, t])$ of $\nu_{k,t}$ under the map $f \mapsto \epsilon f$. Also, for $f \in \mathcal{C}_0^k([0, t])$, define the rate function*

$$I(f) = \frac{1}{4} \int_0^t \sum_{i=1}^k |f'_i(s)|^2 ds,$$

if f is absolutely continuous, and ∞ otherwise. Then for every open set $G \subset \mathcal{C}_0^k([0, t])$ we have

$$\liminf_{\epsilon \downarrow 0} \epsilon^2 \log \nu_{k,t}^\epsilon(G) \geq - \inf_{f \in G} I(f)$$

and for every closed set $F \subset \mathcal{C}_0^k([0, t])$ we have

$$\limsup_{\epsilon \downarrow 0} \epsilon^2 \log \nu_{k,t}^\epsilon(F) \leq - \inf_{f \in F} I(f).$$

1.2 A tool for Airy analysis

While the Brownian Gibbs property is a powerful tool for analyzing the Airy line ensemble, it can nonetheless be incredibly difficult to analyze a collection of non-intersecting Brownian bridges on $\llbracket 1, k \rrbracket \times [0, t]$ conditioned to stay above a possibly arbitrary and unknown boundary. Indeed, at this point there is a fairly substantial bag of tricks designed precisely to handle these issues (e.g. monotonicity ideas [14], the bridge representation [19], the Wiener candidate and jump ensemble methods [11, 32], tangent methods [28]).

One of the main contributions of this paper is to construct a line ensemble $\mathfrak{L}^{t,k}$ that is equal to the Airy line ensemble under a certain non-intersection conditioning, but is given by genuine *independent Brownian bridges* on $\llbracket 1, k \rrbracket \times [0, t]$. The existence of the line ensemble $\mathfrak{L}^{t,k}$ implies Wiener density bounds that are similar to (3) and (1.3), but crucially *take into account the locations of the Airy endpoints*.

Theorem 1.7. *Fix $t \geq 1, k \in \mathbb{N}$. Then there exists a random sequence of continuous functions $\mathfrak{L} = \mathfrak{L}^{t,k} = \{\mathfrak{L}_i^{t,k} : \mathbb{R} \rightarrow \mathbb{R}, i \in \mathbb{N}\}$ such that the following points hold:*

1. *Almost surely, \mathfrak{L} satisfies $\mathfrak{L}_i(r) > \mathfrak{L}_{i+1}(r)$ for all pairs $(i, r) \notin \llbracket 1, k \rrbracket \times (0, t)$.*

2. The ensemble \mathfrak{L} has the following Gibbs property. For any set $S = \llbracket 1, \ell \rrbracket \times [a, b]$, conditional on the values of $\mathfrak{L}_i(r)$ for $(i, r) \notin S$, the distribution of $\mathfrak{L}_i(r)$, $(i, r) \in S$ is given by ℓ independent Brownian bridges B_1, \dots, B_k from $(a, \mathfrak{L}_i(a))$ to $(b, \mathfrak{L}_i(b))$ for $i \in \llbracket 1, \ell \rrbracket$, conditioned on the event $B_i(r) > B_{i+1}(r)$ whenever $(i, r) \notin \llbracket 1, k \rrbracket \times (0, t)$.

In particular, the law of \mathfrak{L} on $\llbracket 1, k \rrbracket \times [0, t]$ is simply given by k independent Brownian bridges connecting $(0, \mathfrak{L}_i(0))$ to $(t, \mathfrak{L}_i(t))$.

3. We have

$$\mathbb{P}(\mathfrak{L}_1 > \mathfrak{L}_2 > \dots) \geq e^{-O_k(t^3)},$$

and conditional on the event $\{\mathfrak{L}_1 > \mathfrak{L}_2 > \dots\}$, the ensemble \mathfrak{L} is equal in law to the parabolic Airy line ensemble \mathfrak{A} .

We also have strong one- and two-point tail bounds on $\mathfrak{L}^{t,k}$ that make it useful for analysis in practice, see Sections 4 and 5 for details. In particular, the tail bounds on $\mathfrak{L}_i(0) - \mathfrak{L}_i(t)$ are stronger than for (variance 2) Brownian motion! We have

$$\mathbb{P}(|\mathfrak{L}_i(0) - \mathfrak{L}_i(t)| > m) \leq \exp\left(-\frac{m^2}{4t} - \frac{4i-2}{3}m^{3/2} + O_k(\sqrt{t}m^{5/4})\right) \quad (4)$$

for all $m > 0$. Note that we can deduce the same tail bounds for \mathfrak{A} by a careful examination of Theorem 1.1.

1.3 A heuristic for (2): Brownian motion avoiding a parabola

We start by giving a heuristic for where formula (2) comes from. For simplicity, we will only consider the case when $k = 1$. A good starting point for trying to understand the behaviour of the top curve \mathfrak{A}_1 is the well-known pair of tail estimates

$$\mathbb{P}(\mathfrak{A}_1(s) + s^2 > m) \sim e^{-(\frac{4}{3}+o(1))m^{3/2}} \quad \text{and} \quad \mathbb{P}(\mathfrak{A}_2(s) + s^2 < -m) \sim e^{-cm^3}. \quad (5)$$

Similar estimates hold for all Airy lines, see Lemmas 2.1, 2.2. The intuition from these estimates suggests first that \mathfrak{A}_1 typically adheres to the parabola $-s^2$, and secondly that it is much easier for \mathfrak{A}_1 to move up away from this parabola than it is for \mathfrak{A}_1 to push \mathfrak{A}_2 down below the parabola. Because of this, a good heuristic for \mathfrak{A}_1 is the following model. Consider a sequence of Brownian bridges B_n from $(-n, -n^2)$ to $(n, -n^2)$ conditioned to stay above the parabola $f(s) = -s^2$, and take n to ∞ to get a distributional limit \mathcal{B} . The process \mathcal{B} has been studied in [23] and can actually be described as a diffusion with an explicit drift involving Airy functions. However, for this discussion we will stick to softer intuition regarding \mathcal{B} .

One way to check the reasonability of this heuristic is to check the probability that $\mathcal{B}(0) \sim m$. The best way for $\mathcal{B}(0)$ to equal m should be if B only deviates from the parabola by following following straight lines from $(-\lambda, -\lambda^2)$ to $(0, m)$ and back to $(\lambda, -\lambda^2)$. Computing the Dirichlet energy of this path against the Dirichlet energy of the parabola, and optimizing over λ yields that our best guess for $\mathbb{P}(\mathcal{B}(0) \sim m)$ should be $e^{-(\frac{4}{3}+o(1))m^{3/2}}$, which agrees




Figure 2: The easiest way for $\mathcal{B}(r) - \mathcal{B}(0), r \in [0, t]$ to equal a function $f \in \mathcal{C}_0^1([0, t])$. The cost to lift \mathcal{B} far enough above of the parabola to achieve the function f is the difference between the Dirichlet energies of the solid blue and purple curves. This energy difference is $S(f)$ plus a lower order term in t .

with the known bound for $\mathfrak{A}_1(0)$. The optimal value is $\lambda = \sqrt{m}$; here the straight lines from $(-\lambda, -\lambda^2)$ and $(\lambda, -\lambda^2)$ to $(0, m)$ are *tangent* to the parabola.

Next, for $\mathcal{B}(r) - \mathcal{B}(0), r \in [0, t]$ to equal a function $f \in \mathcal{C}_0^1([0, t])$, then we must have $\mathcal{B}(0) - f(r) \geq -t^2$ for all r . In other words, we require $\mathcal{B}(0) \geq \mathbb{T}f(0) - t^2$ (or equivalently $\mathcal{B}(t) \geq \mathbb{T}f(t) - t^2$). Since it is costlier to raise \mathcal{B} up further away from the parabola, the optimal way to have $\mathcal{B}(r) - \mathcal{B}(0), r \in [0, t]$ equal to f is to have $\mathcal{B}(0) \sim \mathbb{T}f(0) - t^2, \mathcal{B}(t) \sim \mathbb{T}f(t) - t^2$. The cost of this, relative to the cost of simply having a Brownian motion equal to f can again be computed by comparing the Dirichlet energy of straight lines against a parabola. Assuming t is very small relative to the size of $\mathbb{T}f$, this amounts to computing the Dirichlet energy of the straight line from $(-\sqrt{\mathbb{T}f(0)}, -\mathbb{T}f(0))$ to $(0, \mathbb{T}f(0))$ plus the Dirichlet energy of the straight line from $(0, \mathbb{T}f(t))$ to $(\sqrt{\mathbb{T}f(t)}, -\mathbb{T}f(t))$ minus the Dirichlet energy of the parabola from $(-\sqrt{\mathbb{T}f(0)}, -\mathbb{T}f(0))$ to $(\sqrt{\mathbb{T}f(t)}, -\mathbb{T}f(t))$. This is

$$\frac{2}{3}([\mathbb{T}f(0)]^{3/2} + [\mathbb{T}f(t)]^{3/2}),$$

which is $S(f)$. See Figure 2 for an illustration. The intuition behind the expression (2) for more than one line is similar, except now we need to stack the k functions f_1, \dots, f_k on top of each other, *and* on top of the parabola in order to achieve the ordering $\mathfrak{A}_1 > \mathfrak{A}_2 > \dots$.

Understanding probabilistic objects built from non-crossing paths via a tangent picture is not a new idea. Indeed, tangent methods were proposed by Colomo and Sportiello [12] for studying limit shapes of the six-vertex model, and made rigorous by Aggarwal [3]. In the context of the KPZ and Airy line ensembles, tangent methods were used by Ganguly and Hegde [28] to pinpoint the upper tail one-point asymptotics in (5) and related upper tail two-point asymptotics using only minimal assumptions.

1.4 Outline of the proof and the paper

A priori, it is not clear why the hydrodynamic tangent heuristic should give fine control over the Wiener density. As a result, our proof of the upper bound in Theorem 1.1 does not proceed by directly trying to prove this heuristic. Rather, we first prove Theorem 1.7, which implies that the only contribution to the Wiener density $X_{k,t}(f)$ comes from the joint distribution of the endpoints $\mathfrak{L}(0), \mathfrak{L}(t)$ (here \mathfrak{L} is as in Theorem 1.7). To prove Theorem 1.7, by the Brownian Gibbs property we can abstractly define a σ -finite measure $\mu_{\mathfrak{L}}$ on the space $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$ of sequences of continuous functions $f = \{f_i : \mathbb{R} \rightarrow \mathbb{R}, i \in \mathbb{N}\}$ such that if $\mu_{\mathfrak{L}}(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))$ were finite, then after scaling it to a probability measure, it would be the law of a line ensemble \mathfrak{L} satisfying Theorem 1.7. The key difficulty is in showing that $\mu_{\mathfrak{L}}(\mathcal{C}^{\mathbb{N}}(\mathbb{R})) < \infty$: here we use a refinement of some ideas from [19] and [32], along with new insights about how the Brownian Gibbs property on different domains controls $\mu_{\mathfrak{L}}(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))$. The proof of Theorem 1.7 is contained in Section 3.

In Section 4, we analyze the joint distribution of $\mathfrak{L}(0), \mathfrak{L}(t)$ by again appealing to one-point bounds, using monotonicity arguments, and applying an induction argument to pass from control of $(\mathfrak{L}_i(0), \mathfrak{L}_i(t))$ for any fixed i to joint control of $(\mathfrak{L}_i(0), \mathfrak{L}_i(t)), i \in \llbracket 1, k \rrbracket$. It is only because the upper tail one-point bound in (5) sharply matches the bound coming from the tangent heuristic that the tangent picture emerges. In Section 5, we combine these tail bounds with Theorem 1.7 establishes that $\text{LHS} \leq \text{RHS}$ in (2).

The lower bound $\text{LHS} \geq \text{RHS}$ is easier, since we only need to work on a part of the probability space where the $(k+1)$ st line does not deviate too far from the parabola. It closely resembles a rigorous version of the heuristic discussed above, and is shown in Section 6. A final section (Section 7) proves the corollaries.

We have strived to keep the paper self-contained, relying only on standard tail bounds on the Airy points $\mathfrak{A}_1(0), \mathfrak{A}_2(0), \dots$, the Brownian Gibbs property, and basic monotonicity properties for non-intersecting Brownian bridges as inputs.

Conventions throughout the paper

In all sections of the paper after Section 2, $t \geq 1$ and $k \in \mathbb{N}$ are fixed. We will often suppress dependence of objects on k and t for notational convenience (e.g. we write \mathfrak{L} instead of $\mathfrak{L}^{k,t}$). We write $c, c', c'' > 0$ for large constants and $d, d' > 0$ for small constants. Our constants may change from line to line within proofs and we use a subscript k (i.e. c_k) if they depend on k . Constants will not depend on any other parameters. All Brownian objects in the paper (e.g. Brownian bridge, Wiener measure) will have variance 2, as is the convention for the Airy line ensemble. For a set $I \subset \mathbb{R}$, we write $I_{>}^k$ (respectively I_{\geq}^k) for all $\mathbf{x} \in I^k$ with $x_1 > \dots > x_k$ (respectively $x_1 \geq \dots \geq x_k$). Finally, we use the shorthand $\mathbb{E}_{\mathcal{F}}(\cdot) = \mathbb{E}(\cdot \mid \mathcal{F})$ for conditional expectation with respect to a σ -algebra \mathcal{F} .

2 Preliminaries

In this section, we introduce a few basic preliminaries about the Airy line ensemble and other non-intersecting collections of Brownian motions.

Let I be a closed interval in \mathbb{R} , let $k \in \mathbb{N}$, and define $\mathcal{C}^k(I)$ denote the space of k -tuples of continuous functions $f = (f_1, \dots, f_k)$ from I to \mathbb{R} with the **compact topology**

$$f_n \rightarrow f \quad \text{if} \quad \sup_{x \in K} |f_{n,i}(x) - f_i(x)| \rightarrow 0 \quad \text{for all compact sets } K \subset I, i \in \llbracket 1, k \rrbracket.$$

We also let $\mathcal{C}^{\mathbb{N}}(I)$ be the space of all sequences of continuous functions $f = (f_1, f_2, \dots)$ from I to \mathbb{R} , again equipped the with compact topology (we replace $\llbracket 1, k \rrbracket$ with \mathbb{N} in the definition above), and we write $\mathcal{C}(I) = \mathcal{C}^1(I)$.

We can also think of $f \in \mathcal{C}^k(I)$ as a function $f : \llbracket 1, k \rrbracket \times I \rightarrow \mathbb{R}$ with the topology of uniform convergence on compact subsets of $\llbracket 1, k \rrbracket \times I$, and with this in mind, we will often write $f|_U$ for the restriction of f to a subset of $\llbracket 1, k \rrbracket \times I$. For $f \in \mathcal{C}^k(I)$ with $\ell \leq k$ we write $f^\ell = (f_1, \dots, f_\ell)$ for the first ℓ coordinates of f .

The **parabolic Airy line ensemble** \mathfrak{A} is a random element of $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$ (a.k.a. a line ensemble) satisfying

$$\mathfrak{A}_1 > \mathfrak{A}_2 > \dots$$

almost surely. The process $\mathcal{A}_i(t) = \mathfrak{A}_i(t) + t^2$ is the usual **Airy line ensemble** and is stationary under shifts and flips. For any $r \in \mathbb{R}$ we have

$$\mathcal{A}(s) \stackrel{d}{=} \mathcal{A}(s) \stackrel{d}{=} \mathcal{A}(s+r), \quad \mathcal{A}(\cdot) \stackrel{d}{=} \mathcal{A}(\cdot+r), \quad (6)$$

where the equality in distribution is as random element of $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$. Throughout the paper we will typically use the parabolic version \mathfrak{A} , which has a more natural Gibbs resampling property. The Airy line ensemble \mathcal{A} is uniquely characterized by a determinantal formula. Indeed, for every finite set of times $\{t_1, \dots, t_k\} \subset \mathbb{R}$, the set of locations set $\{(\mathcal{A}_i(t_j), t_j) : i \in \mathbb{N}, j \in \llbracket 1, k \rrbracket\}$ is a determinantal point process on $\mathbb{R} \times \{t_1, \dots, t_k\}$ with with kernel

$$K((x, s); (y, r)) = \begin{cases} \int_0^\infty d\lambda e^{-\lambda(s-r)} \text{Ai}(x+\lambda) \text{Ai}(y+\lambda), & \text{if } s \geq t, \\ -\int_{-\infty}^0 d\lambda e^{-\lambda(s-r)} \text{Ai}(x+\lambda) \text{Ai}(y+\lambda), & \text{if } s < r, \end{cases} \quad (7)$$

where $\text{Ai}(\cdot)$ is the Airy function.

2.1 Tail bounds

The Airy points $\mathfrak{A}_i(0), i \in \mathbb{N}$ satisfy well-known tail bounds. A fairly sharp upper tail bound is a quick consequence of the determinantal formula.

Lemma 2.1. *For all $k \in \mathbb{N}$ there exists $c_k > 0$ such that for $\mathbf{m} \in [0, \infty)_{\geq}^k$ we have*

$$\mathbb{P}(\mathfrak{A}_i(0) > m_i \quad \text{for all } i \in \llbracket 1, k \rrbracket) \leq c_k \exp\left(-\frac{4}{3} \sum_{i=1}^k m_i^{3/2}\right).$$

Proof. This is a simple calculation involving the Airy kernel at a single time. Let $\Lambda = \{\mathfrak{A}_i(0), i \in \mathbb{N}\}$. Then for $\mathbf{m} \in \mathbb{R}_{>}^k$ and $\epsilon > 0$ we have

$$\mathbb{P}(\mathfrak{A}_i(0) \in [m_i - \epsilon, m_i + \epsilon] \text{ for all } i \in \llbracket 1, k \rrbracket) \leq \mathbb{E} \left(\prod_{i=1}^k \#(\Lambda \cap [m_i - \epsilon, m_i + \epsilon]) \right).$$

As long as $\epsilon > 0$ is small enough, the right hand side above equals

$$\int_{m_k - \epsilon}^{m_k + \epsilon} \cdots \int_{m_1 - \epsilon}^{m_1 + \epsilon} \det_{1 \leq i, j \leq k} K(x_i, 0; x_j, 0) dx_1 \dots dx_k,$$

where K is the Airy kernel. Taking $\epsilon \rightarrow 0$, we see that $(\mathfrak{A}_1(0), \dots, \mathfrak{A}_k(0))$ has a Lebesgue density that at $\mathbf{x} \in \mathbb{R}_{\geq}^k$ is bounded above by

$$\det_{1 \leq i, j \leq k} K(x_i, 0; x_j, 0).$$

Using the formula (7) and the standard bound $\text{Ai}(x) \leq c \exp(-\frac{2}{3}x^{3/2})$ for a universal constant c (see formula 10.4.59 in [1]), we have that

$$K((x, 0), (y, 0)) \leq c' \exp(-\frac{2}{3}(x^{3/2} + y^{3/2})).$$

and so

$$\det_{1 \leq i, j \leq k} K(x_i, 0; x_j, 0) \leq k! c' \exp(-\frac{4}{3} \sum_{i=1}^k x_i^{3/2}).$$

Integrating this over the set of all $\mathbf{x} \in \mathbb{R}_{\geq}^k$ with $x_i \geq m_i$ for all i yields the result. \square

Lemma 2.2. *For every $k \in \mathbb{N}$ there exists $d_k > 0$ such that for all $m > 0$ we have*

$$\mathbb{P}(\mathcal{A}_k(0) < -m) \leq 2 \exp(-d_k m^3). \quad (8)$$

Proof. For $k = 1$ this is well-known tail bound goes back to [42]. For $k > 1$, this is a limiting version of Theorem 3.1 of [19], equation (5). More precisely, that theorem states that for a sequence of random variables $\mathcal{A}_k^n(0) \xrightarrow{d} \mathcal{A}_k(0)$ as $n \rightarrow \infty$, that (8) holds for all large enough n . \square

Later on in the paper, we will need a quick way to move between two-point tail bounds and a maximum bound on a stochastic process. This is accomplished through the following lemma.

Lemma 2.3. *Let $t > 0$, $W \in \mathcal{C}([0, t])$ and suppose that there exist constants $\alpha, \beta > 0$ such that for all $m > 0$ and all $s, s+r \in [0, t]$, we have the estimate*

$$\mathbb{P}(|W(s) - W(s+r)| > m\sqrt{r}) \leq \alpha \exp(-\beta m^2).$$

Then for $m > 20\sqrt{\beta}$ we have

$$\mathbb{P} \left(\sup_{s, s+r \in [0, t]} |W(s) - W(s+r)| \geq m\sqrt{t} \right) \leq 2\alpha \exp(-\beta m^2/64).$$

Proof of Lemma 2.3. The proof is a standard chaining argument. We may assume $W(0) = 0$, and focus on bounding $2\|W\|_\infty$. For every $k = 0, 1, \dots$, let G_k be the function with $G_k(s) = W(s)$ for $s = it/(2^k)$ for some $i \in \llbracket 0, 2^k \rrbracket$, and such that G_k is linear on each of the intervals $[it/(2^k), (i+1)t/(2^k)]$. For $k = 0, 1, \dots$, let $H_k = G_k - G_{k-1}$, where $G_{-1} := 0$. We can write $W = \sum_{k=0}^\infty H_k$. Also,

$$\|H_k\|_\infty \leq \sup_{i \in \llbracket 1, 2^k \rrbracket} |W(it2^{-k}) - W((i-1)t2^{-k})|.$$

and hence we have the union bound

$$\mathbb{P}(\|H_k\|_\infty > m\sqrt{t}2^{-k/2}) \leq \sum_{i=1}^{2^k} \mathbb{P}(|W(it2^{-k}) - W((i-1)t2^{-k})| > m\sqrt{t}2^{-k/2}) \leq 2^k \alpha \exp(-\beta m^2).$$

Therefore letting

$$M = \max_{k \geq 0} (t^{-1/2} 2^{k/2} \|H_k\|_\infty - \sqrt{k/\beta}),$$

we have $\mathbb{P}(M' > m) \leq 2\alpha \exp(-\beta m^2)$. Next,

$$2\|W\|_\infty \leq 2 \sum_{k=0}^\infty \|H_k\|_\infty \leq 2\sqrt{t} \sum_{k=0}^\infty (M + \sqrt{k/\beta}) 2^{-k/2} \leq \sqrt{t}(4M + 10/\sqrt{\beta}),$$

which yields the result after converting the tail bound on M . \square

2.2 Gibbs resampling

As discussed in the introduction, the Airy line ensemble satisfies a resampling property called the Brownian Gibbs property. Here we make introduce notation that will help us work with the Brownian Gibbs property, and state a few useful lemmas.

First, for an interval $I \subset \mathbb{R}$, an interval $J \subset I$ and an $\mathbb{R} \cup \{-\infty\}$ -valued function g whose domain contains J , let

$$\text{NI}(g, J) = \{f \in \mathcal{C}^k(I) : f_1(s) > f_2(s) > \dots > f_k(s) > g(s) \quad \text{for all } s \in J\}.$$

Note that the value of k and the interval I is not explicit in the notation $\text{NI}(g, J)$. These will always be clear from context (i.e. by specifying the initial space $\mathcal{C}^k(I)$). We will use the shorthand $\text{NI}(g) = \text{NI}(g, I)$. Next, we say that a k -tuple of **independent Brownian bridges** $B = (B_1, \dots, B_k) \in \mathcal{C}^k([s, t])$ goes from (s, \mathbf{x}) to (t, \mathbf{y}) for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^k$ if each of the B_i is a Brownian bridge with $B_i(s) = x_i, B_i(t) = y_i$. Note that our bridges will always have variance 2. We write

$$\mathbb{P}_{s,t}(\mathbf{x}, \mathbf{y}, f, J)$$

for the probability that a k -tuple of independent Brownian bridges from (s, \mathbf{x}) to (t, \mathbf{y}) is in the non-intersection set $\text{NI}(f, J)$. Again, we omit J from all notation if $J = [s, t]$. We say that the k -tuple of Brownian bridges is **non-intersecting** if it is conditioned on the event $\text{NI}(-\infty)$.

The Brownian Gibbs property for \mathfrak{A} can now be more formally stated as follows:

For a set $S = \llbracket 1, k \rrbracket \times [a, b]$, let $\mathcal{F}_{k,a,b}$ denote the σ -algebra generated by $\{\mathfrak{A}_i(x) : (i, x) \notin S\}$. Then the conditional distribution of $\mathfrak{A}|_S$ given $\mathcal{F}_{k,a,b}$ is equal to the law of k independent Brownian bridges from $(a, \mathfrak{A}^k(a))$ to $(b, \mathfrak{A}^k(b))$ conditioned on the event $\text{NI}(\mathfrak{A}_{k+1})$.

In the statement above, recall that $\mathfrak{A}^k(a) = (\mathfrak{A}_1(a), \dots, \mathfrak{A}_k(a))$ and note that when applying the Gibbs property, the bridges are always drawn independently of the endpoint vectors $\mathfrak{A}^k(a), \mathfrak{A}^k(b)$ and of the lower boundary \mathfrak{A}_{k+1} .

We record two key lemmas that will allow us to work with the Brownian Gibbs property. The first is a crucial monotonicity property.

Lemma 2.4. *Let $[s, t], J$ be closed intervals in \mathbb{R} with $J \subset [s, t]$, let $\mathbf{x}^1 \leq \mathbf{x}^2, \mathbf{y}^1 \leq \mathbf{y}^2 \in \mathbb{R}_{>}^k$ where \leq is the coordinate-wise partial order, and let $g_1 \leq g_2$ be bounded Borel measurable functions from $[s, t] \rightarrow \mathbb{R} \cup \{-\infty\}$. For $i = 1, 2$ be B^i be a k -tuple of Brownian bridges from (s, \mathbf{x}^i) to (t, \mathbf{y}^i) , conditioned on the event $\text{NI}(g_i, J)$. Then there exists a coupling such that $B_j^1(r) \leq B_j^2(r)$ for all $r \in [s, t], j \in \llbracket 1, k \rrbracket$.*

Lemma 2.4 is proven as Lemmas 2.6 and 2.7 in [14] in the case when $[s, t] = J$. The proof goes through verbatim in the slight extension when J is a strict subset of $[s, t]$.

The next lemma is a lower bound on the probability of Brownian bridges being non-intersecting. This bound will allow us to compare regular Brownian bridge probabilities with non-intersecting bridge probabilities.

Lemma 2.5. *Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}_{>}^k$ be vectors with*

$$\min_{i \in \llbracket 1, k-1 \rrbracket} (x_i - x_{i+1}) \wedge (y_i - y_{i+1}) \geq \alpha$$

for some $\alpha > 0$. Let $t > 0$. Then setting $\epsilon = \alpha^2/t$ we have

$$\mathbb{P}_{0,t}(\mathbf{x}, \mathbf{y}, -\infty) \geq e^{-k^3\epsilon/6} (\epsilon/2)^{k(k-1)/2}.$$

This lemma is fairly standard and follows from the Karlin-McGregor formula. For example, it is very similar to Proposition 3.1.1 in [32], and we follow the same steps used in the proof of that proposition.

Proof. First, $\mathbb{P}_{0,t}(\mathbf{x}, \mathbf{y}, -\infty) \geq \mathbb{P}_{0,t}(\alpha\iota, \alpha\iota, -\infty)$ where $\iota = (k, k-1, \dots, 1)$. Therefore letting $\mathbf{x} = \alpha\iota$, by the Karlin-McGregor formula, after simplification we can write

$$P_{0,t}(\alpha\iota, \alpha\iota, -\infty) = \exp\left(-\frac{1}{2t} \sum_{i=1}^k x_i^2\right) \det(e^{x_i x_j / (2t)})_{1 \leq i, j \leq k}.$$

See the middle of p. 48 in [32] for this version of the Karlin-McGregor formula. The determinant is a Vandermonde, and satisfies

$$\det(e^{x_i x_j t^{-1}})_{1 \leq i, j \leq k} = \prod_{1 \leq i < j \leq k} (e^{\alpha^2 j (2t)^{-1}} - e^{\alpha^2 i (2t)^{-1}}) \geq \prod_{i=1}^k \alpha^2 (2t)^{-1} (j-i) \geq (\alpha^2 (2t)^{-1})^{k(k-1)/2}.$$

Finally

$$\exp(-(2t)^{-1} \sum_{i=1}^k x_i^2) \geq \exp(-k^3 \alpha^2 t^{-1}/6),$$

yielding the result. \square

3 Constructing $\mathfrak{L}^{t,k}$

In this section, we construct the line ensemble $\mathfrak{L} = \mathfrak{L}^{t,k}$ and prove Theorem 1.7. Throughout the section, we fix $t \geq 1, k \in \mathbb{N}$.

Let \mathfrak{A} be the parabolic Airy line ensemble. We will define a σ -finite measure $\eta = \eta^{t,k}$ on $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$ according to the following procedure.

1. First, define a line ensemble \mathfrak{B} by letting $\mathfrak{B}_i(r) = \mathfrak{A}_i(r)$ for $(i, r) \notin \llbracket 1, k \rrbracket \times [0, t]$, and let $\mathfrak{B}^k|_{[0,t]}$ be given by k independent Brownian bridges from $(0, \mathfrak{A}^k(0))$ to $(t, \mathfrak{A}^k(t))$. These bridges are independent of each other and of \mathfrak{A} . Let $\tilde{\eta}$ denote the law of \mathfrak{B} .
2. Let η be the measure which is absolutely continuous with respect to $\tilde{\eta}$ with density given by

$$\frac{d\eta}{d\tilde{\eta}}(f) = \frac{1}{\mathbb{P}_{0,t}(f^k(0), f^k(t), f_{k+1})} \quad (9)$$

for $f \in \mathcal{C}^{\mathbb{N}}(\mathbb{R})$. This density exists for $\tilde{\eta}$ -a.e. f since $\mathfrak{B}^k(0) = \mathfrak{A}^k(0)$, $\mathfrak{B}^k(t) = \mathfrak{A}^k(t)$, $\mathfrak{B}_{k+1} = \mathfrak{A}_{k+1}$ and almost surely, $\mathfrak{A}_1(0) > \dots > \mathfrak{A}_{k+1}(0)$ and $\mathfrak{A}_1(t) > \dots > \mathfrak{A}_{k+1}(t)$.

We would like to say that η is a finite measure, and hence can be normalized to a probability measure. This normalized measure will be the law of the line ensemble $\mathfrak{L} = \mathfrak{L}^{t,k}$. Once we know this, Theorem 1.7 will follow easily. However, it is highly non-trivial to show that η is finite. The bulk of this section is devoted to proving this. Indeed, by (9) we can observe that

$$\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R})) = \mathbb{E} \left(\frac{1}{\mathbb{P}_{0,t}(\mathfrak{A}^k(0), \mathfrak{A}^k(t), \mathfrak{A}_{k+1})} \right) \quad (10)$$

and so concretely we just need to show that the **inverse acceptance probability**

$$[\mathbb{P}_{0,t}(\mathfrak{A}^k(0), \mathfrak{A}^k(t), \mathfrak{A}_{k+1})]^{-1}$$

has finite expectation. In order to do this, we will study alternate constructions of the measure η . Fix $s > t$, and define a measure $\eta_s = \eta_s^{t,k}$ by the following procedure:

1. Define a line ensemble \mathfrak{B}_s by letting $\mathfrak{B}_{s,i}(r) = \mathfrak{A}_i(r)$ for $(i, r) \notin \llbracket 1, k \rrbracket \times [-s, s]$, and let $\mathfrak{B}^{k,s}|_{[-s,s]}$ have the conditional law given \mathfrak{A} outside of $\llbracket 1, k \rrbracket \times [-s, s]$ of k independent Brownian bridges from $(-s, \mathfrak{A}^k(-s))$ to $(s, \mathfrak{A}^k(s))$ conditioned on the event

$$\text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]).$$

Let $\tilde{\eta}_s$ denote the law of \mathfrak{B}^s .

2. Let η_s be the measure which is absolutely continuous with respect to $\tilde{\eta}$ with density given by

$$\frac{d\eta_s}{d\tilde{\eta}_s}(f) = \frac{\mathbb{P}_{-s,s}(f^k(-s), f^k(s), f_{k+1}, [-s, 0] \cup [t, s])}{\mathbb{P}_{-s,s}(f^k(-s), f^k(s), f_{k+1})} \quad (11)$$

for $f \in \mathcal{C}^{\mathbb{N}}(\mathbb{R})$.

Lemma 3.1. *For every $s > t$, we have $\eta = \eta_s$, and letting \mathcal{F}_s be the σ -algebra generated by the Airy line ensemble \mathfrak{A} outside of the set $\llbracket 1, k \rrbracket \times [-s, s]$, we have*

$$\mathbb{E}_{\mathcal{F}_s} \left(\frac{1}{\mathbb{P}_{0,t}(\mathfrak{A}^k(0), \mathfrak{A}^k(t), \mathfrak{A}_{k+1})} \right) = \left(\frac{1}{\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1})]} \right). \quad (12)$$

Equation (12) exchanges the expectation of an inverse for the inverse of an expectation, which is much easier to bound.

The idea that the inverse acceptance probability $[\mathbb{P}_{0,t}(\mathfrak{A}^k(0), \mathfrak{A}^k(t), \mathfrak{A}_{k+1})]^{-1}$ can be controlled by an inverse of a conditional expectation by ‘stepping back’ onto the interval $[-s, s]$ and applying the Brownian Gibbs property on $[-s, s]$ is inspired by the Wiener candidate method of Brownian Gibbs resampling introduced by Hammond [32], see also [11].

Proof. We first compute the density of \mathfrak{B} against \mathfrak{B}^s (more precisely, the density of $\tilde{\eta}_s$ against $\tilde{\eta}$). First, letting $U = \llbracket 1, k \rrbracket \times [-s, s]$ we have that $\mathfrak{B} = \mathfrak{B}_s = \mathfrak{A}$ outside of U . Therefore the density of \mathfrak{B} against \mathfrak{B}^s is the same as the density of $\mathfrak{B}|_U$ against $\mathfrak{B}^s|_U$.

Conditional on \mathcal{F}_s , all of $\mathfrak{B}^s|_U$, $\mathfrak{B}|_U$, and $\mathfrak{A}|_U$ have densities with respect to the law of k independent Brownian bridges from $(-s, \mathfrak{A}^k(-s))$ to $(s, \mathfrak{A}^k(s))$. The density of $\mathfrak{B}^s|_U$ is given by

$$\frac{\mathbf{1}(f \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]))}{\mathbb{P}_{-s,s}(f(-s), f(s), \mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])} \quad (13)$$

for $f \in \mathcal{C}^k([-s, s])$. The density of $\mathfrak{A}|_U$ is

$$\frac{\mathbf{1}(f \in \text{NI}(\mathfrak{A}_{k+1}, [-s, s]))}{\mathbb{P}_{-s,s}(f(-s), f(s), \mathfrak{A}_{k+1})} \quad (14)$$

for $f \in \mathcal{C}^k([-s, s])$. We obtain \mathfrak{B} from \mathfrak{A} by replacing \mathfrak{A} with independent Brownian bridges on $[0, t]$ from $(0, \mathfrak{A}^k(0))$ to $(t, \mathfrak{A}^k(t))$. The density of \mathfrak{A} on $\llbracket 1, k \rrbracket \times [0, t]$ with respect to independent Brownian bridges between these points is given (again by the Brownian Gibbs property) by

$$\frac{\mathbf{1}(g \in \text{NI}(\mathfrak{A}_{k+1}, [0, t]))}{\mathbb{P}_{0,t}(g(0), g(t), \mathfrak{A}_{k+1})} \quad (15)$$

for $g \in \mathcal{C}^k([0, t])$. Therefore the density of $\mathfrak{B}|_U$ with respect to the law of k independent Brownian bridges from $(-s, \mathfrak{A}^k(-s))$ to $(s, \mathfrak{A}^k(s))$ is simply the product of (14) with the inverse of (15) at the function $g = f|_{[0,t]}$. This is

$$\frac{\mathbf{1}(f \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])) \mathbb{P}_{0,t}(f(0), f(t), \mathfrak{A}_{k+1})}{\mathbb{P}_{-s,s}(f(-s), f(s), \mathfrak{A}_{k+1})}. \quad (16)$$

The ratio of (16) and (13) gives the density of $\mathfrak{B}|_U$ against $\mathfrak{B}^s|_U$, conditional on \mathcal{F}_s :

$$\left(\frac{\mathbb{P}_{-s,s}(\mathfrak{A}^k(-s), \mathfrak{A}^k(s), \mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])}{\mathbb{P}_{-s,s}(\mathfrak{A}^k(-s), \mathfrak{A}^k(s), \mathfrak{A}_{k+1})} \right) \times \mathbb{P}_{0,t}(f(0), f(t), \mathfrak{A}_{k+1}), \quad (17)$$

for $f \in \mathcal{C}^k([-s, s])$. Using this, and the definitions (9), (11) of η, η_s from $\tilde{\eta}, \tilde{\eta}_s$ gives that $\eta = \eta_s$. For the equality (12), using that the first factor in (17) is \mathcal{F}_s -measurable, we have that

$$\begin{aligned} \mathbb{E}_{\mathcal{F}_s} \left(\frac{1}{\mathbb{P}_{0,t}(\mathfrak{A}^k(0), \mathfrak{A}^k(t), \mathfrak{A}_{k+1})} \right) &= \mathbb{E}_{\mathcal{F}_s} \left(\frac{1}{\mathbb{P}_{0,t}(\mathfrak{B}^k(0), \mathfrak{B}^k(t), \mathfrak{A}_{k+1})} \right) \\ &= \frac{\mathbb{P}_{-s,s}(\mathfrak{A}^k(-s), \mathfrak{A}^k(s), \mathfrak{A}_{k+1})}{\mathbb{P}_{-s,s}(\mathfrak{A}^k(-s), \mathfrak{A}^k(s), \mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])}, \end{aligned}$$

which equals $1/\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1})]$. \square

The goal of the next few lemmas is to estimate the right-hand side of (12). The first lemma gives a bound on the right-hand side of (12) in terms of a few simple \mathcal{F}_s -measurable random variables.

Lemma 3.2. *Fix $s \geq 2t$ and define \mathcal{F}_s -measurable random variables*

$$\begin{aligned} D &= D(k, t) = \sqrt{t} + \max_{r, r' \in [0, t]} |\mathfrak{A}_{k+1}(r) - \mathfrak{A}_{k+1}(r')|, \\ M &= M(k, s) = \sqrt{s} + \max_{r, r' \in [-s, s]} |\mathfrak{A}_{k+1}(r) - \mathfrak{A}_{k+1}(r')| + \max_{i \in [1, k]} |\mathfrak{A}_i(s) - \mathfrak{A}_i(-s)|. \end{aligned}$$

Then with \mathfrak{B}_s as in the previous lemma, we have

$$\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1})] \geq \exp\left(-\frac{ck^3(D^2 + MD)}{s} - cks\right).$$

We separate out one of the main calculations for Lemma 3.2, as it will also be required later in the paper.

Lemma 3.3. *Let $s \geq 2t$, and $\mathbf{x}, \mathbf{y} \in \mathbb{R}_{>}^k$. Let $g \in \mathcal{C}([-s, s])$ be such that $g(-s) < x_k, g(s) < y_k$. Let B be a k -tuple on independent Brownian bridges from $(-s, \mathbf{x})$ to (s, \mathbf{y}) , conditioned on the event*

$$\text{NI}(g, [-s, 0] \cup [t, s]).$$

Define $\iota = (k, \dots, 1), \mathbf{1} = (1, 1, \dots, 1) \in \mathbb{R}^k$, and for $\alpha, \beta \geq 0$ define $f^{\alpha, \beta} \in \mathcal{C}^k([-s, s])$ by letting

$$f^{\alpha, \beta}(-s) = 0, \quad f^{\alpha, \beta}(0) = f^{\alpha, \beta}(t) = \alpha\iota + \beta\mathbf{1}, \quad f^{\alpha, \beta}(s) = 0,$$

and so that $f^{\alpha, \beta}$ is linear on each of the pieces $[-s, 0], [0, t], [t, s]$.

Then for $f \in \text{NI}(g, [-s, 0] \cup [t, s])$ we have that

$$\frac{\mathbb{P}(B = f + f^{\alpha, \beta})}{\mathbb{P}(B = f)} \geq \exp\left(-\frac{k^3(\alpha + \beta)^2}{s} - \frac{k(\alpha + \beta) \sum_{i=1}^k [(f_i(0) - x_i)^+ + (f_i(t) - y_i)^+]}{s}\right). \quad (18)$$

The more formal way to interpret (18) is as a bound on the density of the law μ'_B of $B - f^{\alpha,\beta}$ against the law μ_B of B , on the set $\text{NI}(g, [-s, 0] \cup [t, s])$ where μ'_B is absolutely continuous with respect to μ_B .

Proof. Let ν be the law of k independent Brownian bridges from $(-s, \mathbf{x})$ to (s, \mathbf{y}) . Observe that μ_B, μ'_B are absolutely continuous with respect to ν with densities

$$\begin{aligned} \frac{d\mu_B}{d\nu}(f) &= \frac{1}{Z} \mathbf{1}(f \in \text{NI}(g, [-s, 0] \cup [t, s])), \\ \frac{d\mu'_B}{d\nu}(f) &= \frac{1}{Z} \mathbf{1}(f + f^{\alpha,\beta} \in \text{NI}(g, [-s, 0] \cup [t, s])) \\ &\times \exp\left(-\frac{2(f(0) - \mathbf{x}) \cdot f^{\alpha,\beta}(0) + \|f^{\alpha,\beta}(0)\|^2}{4s} - \frac{2(f(t) - \mathbf{y}) \cdot f^{\alpha,\beta}(t) + \|f^{\alpha,\beta}(t)\|^2}{4(s-t)}\right). \end{aligned} \quad (19)$$

where $Z = \mathbb{P}_{-s,s}(\mathbf{x}, \mathbf{y}, g, [-s, 0] \cup [t, s])$ is a normalizing factor. Now, if f is in the set $\text{NI}(g, [-s, 0] \cup [t, s])$, then so is $f + f^{\alpha,\beta}$. Hence the right-hand side of (18) is bounded below by the exponential factor (19). We can bound (19) below by using that $s - t \geq s/2$, and that $0 \leq f^{\alpha,\beta} \leq k(\alpha + \beta)\mathbf{1}$, which yields the desired bound. \square

Proof of Lemma 3.2. All statements in the proof are conditional on \mathcal{F}_s . Define the \mathcal{F}_s -measurable vector

$$\mathbf{z} = (D + k, D + (k - 1), \dots, D + 1)$$

and the \mathcal{F}_s -measurable set

$$O = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}_>^k \times \mathbb{R}_>^k : x_k > \mathfrak{A}_{k+1}(0), y_k > \mathfrak{A}_{k+1}(t)\}. \quad (20)$$

By the definition of D , for $(\mathbf{x}, \mathbf{y}) \in O$ we have

$$\mathbb{P}_{0,t}(\mathbf{x} + \mathbf{z}, \mathbf{y} + \mathbf{z}, \mathfrak{A}_{k+1}) \geq \mathbb{P}(|B(s)| \geq 1/2)^k, \quad (21)$$

where B is a Brownian bridge from $(0, 0)$ to $(0, t)$. By a standard bound, the right hand side above is bounded below by e^{-ckt} . Therefore letting $\mu_{\mathfrak{B}}$ denote the conditional law of $(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t))$ given \mathcal{F}_s , to complete the proof it suffices to find a set $A \subset O$ such that $\mu_{\mathfrak{B}}(A + (\mathbf{z}, \mathbf{z}))$ is large. Fix $\Delta > 0$ and let A_Δ be the \mathcal{F}_s -measurable subset of $(\mathbf{x}, \mathbf{y}) \in O$ where

$$x_i \leq \mathfrak{A}_i(-s) + \Delta, \quad y_i \leq \mathfrak{A}_i(s) + \Delta$$

for all $i \in \llbracket 1, k \rrbracket$. Then by Lemma 3.3 with $\alpha = 1, \beta = D$, we have

$$\begin{aligned} &\mu_{\mathfrak{B}}(A_\Delta + (\mathbf{z}, \mathbf{z})) \\ &\geq \mu_{\mathfrak{B}}(A_\Delta) \sup_{(\mathbf{x}, \mathbf{y}) \in A} \exp\left(-\frac{k^3(\sqrt{t} + D)^2}{s} - \frac{k(\sqrt{t} + D) \sum_{i=1}^k ((x_i - \mathfrak{A}_i(-s))^+ + (y_i - \mathfrak{A}_i(-s))^+)}{s}\right) \\ &\geq \mu_{\mathfrak{B}}(A_\Delta) \exp\left(-\frac{4k^3 D^2}{s} - \frac{4k^2 D \Delta}{s}\right). \end{aligned} \quad (22)$$

It remains to find Δ where $\mu_{\mathfrak{B}}(A_\Delta)$ is large. Define vectors $\mathbf{w}^s, \mathbf{w}^{-s}$ where

$$w_i^{\pm s} = \mathfrak{A}_i(\pm s) + M + k + 1 - i.$$

By Lemma 2.4, on the interval $[-s, s]$, the k -tuple $(\mathfrak{B}_1^s, \dots, \mathfrak{B}_k^s)$ is stochastically dominated by k independent Brownian bridges $B = (B_1, \dots, B_k)$ from $(-s, \mathbf{w}^{-s})$ to (s, \mathbf{w}^s) conditioned on the event

$$\text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]).$$

Now, let $L \in \mathcal{C}^k([-s, s])$ be the function whose i th coordinate L_i is the linear function satisfying $L_i(\pm s) = w_i^{\pm s}$. Observe that by the construction of $\mathbf{w}^{\pm s}$, we have $f \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])$ for any sequence of bridges f from $(-s, \mathbf{w}^{-s})$ to (s, \mathbf{w}^s) with $\|f - L\|_\infty < 1/2$. The probability that this condition holds for a sequence of independent Brownian bridges is bounded below by $\exp(-cks)$ (similarly to the bound on (21)). Therefore

$$\begin{aligned} \mathbb{P}(B_i(r) \leq M + k - i + 3/2 + \mathfrak{A}_i(s) \vee \mathfrak{A}_i(-s) \quad \forall i \in \llbracket 1, k \rrbracket, r = 0, t) \\ \geq \mathbb{P}(\|B - L\|_\infty < 1) \geq \exp(-cks). \end{aligned}$$

Observing that

$$M + k - i + 3/2 + \mathfrak{A}_i(s) \vee \mathfrak{A}_i(-s) - \mathfrak{A}_i(\pm s) \leq 2M + 3k$$

for all i , we can conclude that

$$\mu_{\mathfrak{B}}(A_{2M+3k}) \geq \mathbb{P}((B(0), B(t)) \in A_{2M+3k}) \geq \exp(-cks).$$

Combining this with the bound on (21) and (22) and simplifying yields the result. \square

It remains to prove tails bounds on M, D . The key is a two-point estimate for the Airy line ensemble. In Section 4.2 will prove a highly refined two-point estimate directly for \mathfrak{L} , whose proof goes through verbatim for \mathfrak{A} . However, the full power of such an estimate is not necessary and we make do with the following lemma, whose proof is much less involved.

Lemma 3.4. *For every $k \in \mathbb{N}$ there exists $c_k > 0$ such that for all $s \in \mathbb{R}, r \in (0, \infty), a > k^2 r^2$ we have*

$$\begin{aligned} \mathbb{P}(|\mathfrak{A}_k(s) + s^2 - \mathfrak{A}_k(s+r) - (s+r)^2| > a) \\ \leq c_k \exp\left(-\frac{a^2}{4r} - \frac{ka^{3/2}}{8} + 3a^{3/2} + \frac{a}{\sqrt{r}} + 2k^2 |\log(\sqrt{a}/r)|\right). \end{aligned} \quad (23)$$

The proof for Lemma 3.4 is similar to the proof of Lemma 6.1 in [19], except here we keep closer track of the error terms. Somewhat remarkably, this already gives a stronger tail bound than for Brownian motion when k is large enough. We will need the more nuanced method in Section 4.2 to give a stronger tail bound than Brownian motion for all k .

Proof. By stationarity, it suffices to prove the lemma when $s = 0$. Moreover, since $\mathfrak{A}(\cdot) = \mathfrak{A}(-\cdot)$, it suffices to prove the bound with the absolute value removed. For $r > 0$, define

$$L_r = \mathfrak{A}_k(0) - \mathfrak{A}_k(r) - r^2.$$

Let $\lambda > r$ be a parameter that we will optimize over. Let \mathcal{F}_λ be the σ -algebra generated by \mathfrak{A} outside of the set $\llbracket 1, k \rrbracket \times [0, \lambda]$. Let $\mathbf{v} = \sqrt{r}(1/k, 2/k \dots, 1)$, and let B be a k -tuple of non-intersecting Brownian bridges from $(0, \mathfrak{A}_k(0) - \mathbf{v})$ to $(\lambda, \mathfrak{A}_k(\lambda) - \mathbf{v})$. By Lemma 2.4, given \mathcal{F}_λ , the Airy lines $\mathfrak{A}_{\llbracket 1, k \rrbracket \times [0, \lambda]}$ stochastically dominate B , so

$$\mathbb{P}(L_r > a \mid \mathcal{F}_\lambda) \leq \mathbb{P}(\mathfrak{A}_k(0) - B_k(r) > r^2 + a \mid \mathcal{F}_\lambda) = \mathbb{P}(B_k(0) - B_k(r) > r^2 - \sqrt{r} + a \mid \mathcal{F}_\lambda).$$

Now let \tilde{B} be a collection of k -tuple of independent Brownian bridges from $(0, \mathfrak{A}_k(0) - \mathbf{v})$ to $(\lambda, \mathfrak{A}_k(\lambda) - \mathbf{v})$ without a non-intersection condition. By Lemma 2.5 with $t = \lambda, \alpha = \sqrt{r}/k$,

$$\begin{aligned} & \mathbb{P}(B_k(0) - B_k(r) > r^2 - \sqrt{r} + a \mid \mathcal{F}_\lambda) \\ & \leq \exp\left(-\frac{kr}{6\lambda} + \frac{k(k-1)\log(k^2\lambda/(2r))}{2}\right) \mathbb{P}(\tilde{B}_k(0) - \tilde{B}_k(r) > r^2 - \sqrt{r} + a \mid \mathcal{F}_\lambda) \\ & \leq \exp(k^2 \log(k\lambda/r)) \mathbb{P}(\tilde{B}_k(0) - \tilde{B}_k(r) > a - \sqrt{r} \mid \mathcal{F}_\lambda). \end{aligned} \quad (24)$$

Given \mathcal{F}_λ , the increment $\tilde{B}_k(0) - \tilde{B}_k(r)$ is normal with variance $2r(1 - r/\lambda)$ and mean $rL_\lambda/\lambda + r\lambda$.

Therefore we have

$$\begin{aligned} \mathbb{P}(\tilde{B}_k(0) - \tilde{B}_k(r) > a - \sqrt{r} \mid \mathcal{F}_\lambda) & \leq c \exp\left(-\frac{(a - \sqrt{r} - rL_\lambda/\lambda - r\lambda)^2}{4(1 - r/\lambda)r}\right) \\ & \leq \exp\left(-\frac{(a^2 - 2a(\sqrt{r} + rL_\lambda/\lambda + r\lambda))(1 + r/\lambda)}{4r}\right) \\ & \leq \exp\left(-\frac{a^2}{4r} - \frac{a^2}{4\lambda} + \frac{a}{\sqrt{r}} + a(L_\lambda)^+/\lambda + a\lambda\right). \end{aligned} \quad (25)$$

In the final inequality, we have used that $1 + r/\lambda \leq 2$. Now, by Lemmas 2.1 and 2.2, there exists $c_k > 0$ such that for all $m > 0$,

$$\mathbb{P}(L_\lambda > m) \leq c_k e^{-\frac{4k-1}{3}m^{3/2}}.$$

Therefore L_λ is stochastically dominated by a random variable with Lebesgue density bounded above by $c'_k e^{-\frac{4k-1}{3}m^{3/2}}$ and so

$$\mathbb{E}e^{aL_\lambda/\lambda} \leq c'_k \int_0^\infty e^{am/\lambda - \frac{4k-1}{3}m^{3/2}} dm.$$

For any $\alpha, \beta > 0$, if $f(x) = \alpha x - \beta x^{3/2}$ and $u = 4\alpha^2/(9\beta^2)$ is the argmax of f , then

$$\int_0^\infty e^{f(x)} dx \leq e^{f(u)} \left(2 + \int_{-\infty}^\infty e^{-|f'(u+1) \wedge f'(u-1)|x} dx\right) \leq \frac{4e^{f(u)}}{|f'(u+1) \wedge f'(u-1)|} \leq \frac{4e^{f(u)}}{\alpha}.$$

Using this inequality and simplifying the result gives that $\mathbb{E}e^{aL_\lambda/\lambda} \leq c''_k(\lambda/a)e^{4a^3\lambda^{-3}/(3(4k-1)^2)}$, and so the expectation of (25) is bounded above by

$$c''_k \exp\left(-\frac{a^2}{4r} - \frac{a^2}{4\lambda} + \frac{a}{\sqrt{r}} + \frac{4a^3}{3\lambda^3(4k-1)^2} + a\lambda + \log(\lambda/a)\right). \quad (26)$$

A quick inspection reveals that the minimum value of (26) is taken when $\lambda = O(k^{-1}\sqrt{a})$. Plugging in $\lambda = \sqrt{ak}^{-1}$ (which is bounded below by r since $a > k^2r^2$) and simplifying gives that (26) is bounded above by

$$c_k'' \exp\left(-\frac{a^2}{4r} - \frac{ka^{3/2}}{8} + 3a^{3/2} + \frac{a}{\sqrt{r}}\right).$$

Combining this with (24) yields the result. \square

We can deduce tail bounds on M and D immediately from Lemma 3.4.

Lemma 3.5. *For every $k \in \mathbb{N}$, there exists $c_k > 0$ such that the following bounds hold for all $t \geq 1, s \geq 2t$. Letting $D = D(k, t)$ and $M = M(k, s)$ we have*

$$\mathbb{P}(D > a) \leq \exp\left(c_k t^3 - \frac{a^2}{320t}\right), \quad \mathbb{P}(M > a) \leq \exp\left(c_k s^3 - \frac{a^2}{320s}\right).$$

Proof. By Lemma 3.4, the process $\mathfrak{A}_k(s), s \in [0, t]$ satisfies the conditions of Lemma 2.3 with $\beta = 1/5$ and $\alpha = \exp(c_k t^3)$. Appealing to that lemma yields the result. The proof of the bound on M is essentially the same. \square

We can finally give a concrete bound on $\mathbb{E}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1} \mid \mathcal{F}_s)]$, which in turn shows that the measure η is finite.

Corollary 3.6. *There exists $d_k > 0$ such that for all $s \geq 2t$ we have*

$$\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1})] \leq \epsilon \tag{27}$$

with probability at most $e^{O_k(s^{5/2}\sqrt{t})} \epsilon^{d_k\sqrt{s/t}}$. In particular, $\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R})) \leq e^{O_k(t^3)}$.

Proof. Let $\alpha = \log(\epsilon^{-1})$. Without loss of generality, we may assume $\alpha \geq c_k s^2 t$ for some large constant c_k , since otherwise the bound holds trivially. Next, let $D = D(k, t), M = M(k, s)$ and write $D' = t^{-1/2}D - c'_k t^{3/2}, M' = s^{-1/2}M - c'_k s^{3/2}$. Then by Lemma 3.5, for $X = D' \vee M'$ and $a > 0$ we have $\mathbb{P}(X > a) \leq e^{-d'_k a^2}$. Moreover, by Lemma 3.2, we have

$$\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}_s^k(0), \mathfrak{B}_s^k(t), \mathfrak{A}_{k+1})] \geq \exp\left(-c_k(X^2\sqrt{t/s} + s\sqrt{t}X + st^2)\right).$$

Therefore the probability of the event (27) is at most

$$\begin{aligned} \mathbb{P}(c_k(X^2\sqrt{t/s} + s\sqrt{t}X + st^2) \geq \alpha) &\leq \mathbb{P}(X(X + s^{3/2}) \geq c_k^{-1}\sqrt{s/t}\alpha/2) \\ &\leq \mathbb{P}(X^2 \geq c_k^{-1}\sqrt{s/t}\alpha/4) + \mathbb{P}(X \geq c_k^{-1}\alpha/(4s\sqrt{t})) \\ &\leq 2 \exp\left(-d_k(\alpha\sqrt{s/t}) \wedge (\alpha^2/(s^2t))\right) \\ &= 2 \exp\left(-d_k\alpha\sqrt{s/t}\right) = 2\epsilon^{d_k\sqrt{s/t}}, \end{aligned}$$

where we have used the lower bound on α and the tail bound on X to achieve the bound. \square

By Corollary 3.6, η is a finite measure, and so the normalized measure $\eta' = \frac{1}{\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))}\eta$ is a probability measure. Let $\mathfrak{L} = \mathfrak{L}^{t,k}$ denote a line ensemble whose law is equal to η' . It is easy to check that \mathfrak{L} satisfies Theorem 1.7.

Proof of Theorem 1.7. Part 1 is immediate from the definition of η . Next, by the definition (9), restricting η to the set

$$\mathcal{C}_{>}^{\mathbb{N}}(\mathbb{R}) = \{f \in \mathcal{C}^{\mathbb{N}}(\mathbb{R}) : f_1 > f_2 > \dots\}$$

gives a probability measure. By the Brownian Gibbs property, this measure is the law of \mathfrak{A} . Therefore conditioning \mathfrak{L} to lie in $\mathcal{C}_{>}^{\mathbb{N}}(\mathbb{R})$ gives a parabolic Airy line ensemble. The bound $\mathbb{P}(\mathfrak{L} \in \mathcal{C}_{>}^{\mathbb{N}}(\mathbb{R})) \geq e^{-O_k(t^3)}$ follows from the bound on $\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))$ in Corollary 3.6. This gives Theorem 1.7.3.

For part 2, first observe that if \mathfrak{L} satisfies the desired Gibbs property on all sets $S = \llbracket 1, \ell \rrbracket \times [a, b]$ where $a < 0 < t < b$ and $\ell > k$, then it satisfies the Gibbs property everywhere. Fix such a set S . Recalling the definition of \mathfrak{B} from the construction of η , and let \mathfrak{A} be coupled to \mathfrak{B} as in that construction. We first compute the conditional density of $\mathbb{P}_{\sigma(\mathfrak{B}|_{S^c})}(\mathfrak{B}|_S \in \cdot)$ against the law of ℓ independent Brownian bridges B from $(a, \mathfrak{B}^\ell(a))$ to $(b, \mathfrak{B}^\ell(b))$. Let $S_1 = \llbracket 1, k \rrbracket \times [0, t]$ and $S_2 = S \setminus S_1$. First, $\mathfrak{B}|_{S_2} = \mathfrak{A}|_{S_2}$ so by the Brownian Gibbs property for \mathfrak{A} we have

$$\frac{\mathbb{P}_{\sigma(\mathfrak{B}|_{S^c})}(\mathfrak{B}|_{S_2} = f)}{\mathbb{P}_{\sigma(\mathfrak{B}|_{S^c})}(B|_{S_2} = f)} = \frac{\mathbf{1}(f_i(r) > f_{i+1}(r) \text{ for all } (i, r) \in S_2) \mathbb{P}_{0,t}(f^k(0), f^k(t), f_{k+1})}{Z}. \quad (28)$$

where in the indicator above we say that $f_{\ell+1} = \mathfrak{B}_{\ell+1}$, and Z is the $\sigma(\mathfrak{B}|_{S^c})$ -measurable normalization factor $\mathbb{P}_{a,b}(\mathfrak{B}^\ell(a), \mathfrak{B}^\ell(b), \mathfrak{B}_{k+1})$. Next, given $\mathfrak{B}|_{S_2 \cup S^c}$, the distribution of $\mathfrak{B}|_{S_1}$ is that of k independent Brownian bridges from $(0, \mathfrak{B}^k(0))$ to $(t, \mathfrak{B}^k(t))$. As the same is true of the unconditioned bridges B , the density

$$\frac{\mathbb{P}_{\sigma(\mathfrak{B}|_{S^c})}(\mathfrak{B}|_S = f)}{\mathbb{P}_{\sigma(\mathfrak{B}|_{S^c})}(B|_S = f)}$$

is the same as the right-hand side of (28). Therefore by (9), if B' consists of ℓ independent Brownian bridges B from $(a, \mathfrak{L}^\ell(a))$ to $(b, \mathfrak{L}^\ell(b))$ then

$$\frac{\mathbb{P}_{\sigma(\mathfrak{L}|_{S^c})}(\mathfrak{L}|_S = f)}{\mathbb{P}_{\sigma(\mathfrak{L}|_{S^c})}(B|_S = f)} = \frac{\mathbf{1}(f_i(r) > f_{i+1}(r) \text{ for all } (i, r) \in S_2)}{Z'},$$

where Z' is a new $\sigma(\mathfrak{L}|_{S^c})$ -measurable normalization factor. This is the desired Gibbs property. \square

Remark 3.7. It is interesting to note that Theorem 1.7 already implies that on the set $\llbracket 1, k \rrbracket \times [0, t]$, if $L \in \mathcal{C}^k([0, t])$ is the function whose coordinates L_i are linear functions given by the conditions $L_i(0) = \mathfrak{A}_i(0)$, $L_i(t) = \mathfrak{A}_i(t)$, then $[\mathfrak{A} - L]_{\llbracket 1, k \rrbracket \times [0, t]}$ has a uniformly bounded density against the law of k independent Brownian bridges.

4 Tail bounds for $\mathfrak{L}^{t,k}$

In this section, we prove one- and two-time tail bounds on $\mathfrak{L} = \mathfrak{L}^{t,k}$ (including the bound (4)). As in the previous section, we fix $k \in \mathbb{N}$ and $t \geq 1$ throughout the section. These bounds will be used in Section 5 to move from Theorem 1.7 to the upper bound in Theorem 1.1.

4.1 The one-time bound

The single-time bound is easier, so we start there. The idea is to first prove tail bounds on the line ensembles \mathfrak{B}_s introduced for Lemma 3.1, and then deduce tail bounds on \mathfrak{L} through the definition (11) and the bounds in Corollary 3.6.

Lemma 4.1. *Let $\mathbf{m} \in [0, \infty)_{\geq}^k$ and define the set*

$$A(\mathbf{m}, r) = \{f \in \mathcal{C}^{\mathbb{N}}(\mathbb{R}) : f_i(r) > m_i + r^2 \quad \text{for all } i \in \llbracket 1, k \rrbracket\}.$$

Then for $r \notin (0, t)$ we have

$$\mathbb{P}(\mathfrak{B}_s \in A(\mathbf{m}, r)) \leq \exp\left(-\frac{4}{3} \sum_{i=1}^k m_i^{3/2} + c_k \log(\|\mathbf{m}\|_2 + s)\right).$$

Proof. First, the bound is immediate from Lemma 2.1 when $|r| \geq s$ since $\mathfrak{A}(r) = \mathfrak{B}_s(r)$ when $r \geq s$. The case when $|r| < s$ is more involved. Let

$$S = \llbracket 1, k \rrbracket \times [-s, s], \quad S_1 = \llbracket 1, k \rrbracket \times ([-s, 0] \cup [t, s]), \quad S_2 = \llbracket 1, k \rrbracket \times [0, t].$$

Define a line ensemble \mathfrak{M}_s by letting $\mathfrak{M}_{s,i}(r) = \mathfrak{A}_i(r)$ for $(i, r) \notin S$, and let $\mathfrak{M}_s^s|_S$ have the conditional law given $\mathfrak{A}|_{S^c}$ of k independent Brownian bridges from $(-s, \mathfrak{A}^k(-s))$ to $(s, \mathfrak{A}^k(s))$ conditioned on the event

$$\text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]) \cap \text{NI}(-\infty, [0, t]).$$

In words, \mathfrak{M}_s is defined similarly to \mathfrak{B}_s , except with the extra condition that $\mathfrak{M}_{s,1} > \dots > \mathfrak{M}_{s,k}$ on the interval $[0, t]$. Then by Lemma 2.4, \mathfrak{M}_s is stochastically dominated by \mathfrak{A} , so by Lemma 2.1, we have

$$\mathbb{P}(\mathfrak{M}_i(r) + r^2 > m_i \quad \text{for all } i \in \llbracket 1, k \rrbracket) \leq c_k \exp\left(-\frac{4}{3} \sum_{i=1}^k m_i^{3/2}\right) \quad (29)$$

for all $r \in \mathbb{R}$, $\mathbf{m} \in [0, \infty)_{\geq}^k$. Our goal is to effectively compare $\mathfrak{M}_s^k|_{S_2}$ to $\mathfrak{B}_s^k|_{S_2}$ conditionally on the σ -algebra $\mathcal{F}_s := \sigma(\mathfrak{A}|_{\llbracket 1, k \rrbracket \times [-s, s]^c})$ in order to apply this bound. By Lemma 3.3 (and with notation as in that lemma), for any $g \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])$ we have

$$\frac{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{B}_s|_{S_2} = [g + f^{\epsilon, 0}]|_{S_2})}{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{B}_s|_{S_2} = g|_{S_2})} \geq \exp\left(-\frac{k^3 \epsilon^2}{s} - \frac{k^2 \epsilon (g_1(0) \vee g_1(t) - \mathfrak{A}_{k+1}(-s) \wedge \mathfrak{A}_{k+1}(s))^+}{s}\right).$$

On the other hand, the conditional laws of $\mathfrak{B}_s|_{S_2}$ and $\mathfrak{M}_s|_{S_2}$ given \mathcal{F}_s are mutually absolutely continuous with density

$$\begin{aligned} \frac{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{M}_s|_{S_2} = g)}{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{B}_s|_{S_2} = g)} &= \frac{\mathbb{P}_{0,t}(g(0), g(t), -\infty) \mathbb{P}_{\mathcal{F}_s}(\mathfrak{A}^k|_{[-s,s]} \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]))}{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{A}^k|_{[-s,s]} \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]) \cap \text{NI}(-\infty, [0, t]))} \\ &\geq \mathbb{P}_{0,t}(g(0), g(t), -\infty). \end{aligned}$$

In particular, if $g_i(r) \geq g_{i+1}(r) + \epsilon$ for all $i \in \llbracket 1, k-1 \rrbracket, r \in \{0, 2t\}$ for some $\epsilon < 1/2$, then by Lemma 2.5, this is bounded below by $e^{-c_k \log(t/\epsilon)}$. Putting this together with the previous bound implies that for $g \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])$, setting

$$Y_g = 2s + (g_1(0) \vee g_1(t) - \mathfrak{A}_{k+1}(-s) \wedge \mathfrak{A}_{k+1}(s))^+,$$

and $\epsilon = s/Y_g$, we have

$$\frac{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{M}_s|_{S_2} = [g + f^{\epsilon, 0}]|_{S_2})}{\mathbb{P}_{\mathcal{F}_s}(\mathfrak{B}_s|_{S_2} = g|_{S_2})} \geq \exp(-c_k \log Y_g). \quad (30)$$

Therefore for every $\mathbf{m} \in [0, \infty)_{\geq}^k, r \in [-s, 0] \cup [t, s]$, and $\lambda > 0$ we have

$$\mathbb{P}_{\mathcal{F}_s}(\mathfrak{B}_s \in A(\mathbf{m}, r)) \leq e^{c_k \log \lambda} \mathbb{P}_{\mathcal{F}_s}(\mathfrak{M}_s \in A(\mathbf{m}, r)) + \mathbb{P}_{\mathcal{F}_s}(Y_{\mathfrak{B}} > \lambda).$$

Upon taking expectations of both sides, we get that

$$\mathbb{P}(\mathfrak{B} \in A(\mathbf{m}, r)) \leq e^{c_k \log \lambda} \mathbb{P}(\mathfrak{M} \in A(\mathbf{m}, r)) + \mathbb{P}(Y_{\mathfrak{B}} > \lambda).$$

It remains to bound $\mathbb{P}(Y_{\mathfrak{B}} > \lambda)$ and then optimize over λ . Define vectors $\mathbf{w}^s, \mathbf{w}^{-s}$ where

$$w_i^{\pm s} = \mathfrak{A}_i(\pm s) + \sup_{r \in [-s, s]} |\mathfrak{A}_{k+1}(r) - \mathfrak{A}_{k+1}(s)| + (k+1-i)s.$$

By Lemma 2.4, on the interval $[-s, s]$ \mathfrak{B}_s^k is stochastically dominated by k independent Brownian bridges $B = (B_1, \dots, B_k)$ from $(-s, \mathbf{w}^{-s})$ to (s, \mathbf{w}^s) conditioned on the event

$$\text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s]).$$

Now, by the construction of $\mathbf{w}^{\pm s}$, the probability that k independent Brownian bridges $B = (B_1, \dots, B_k)$ from $(-s, \mathbf{w}^{-s})$ to (s, \mathbf{w}^s) satisfy $B \in \text{NI}(\mathfrak{A}_{k+1}, [-s, 0] \cup [t, s])$ is bounded below by e^{-c_k} . Therefore

$$\begin{aligned} \mathbb{P}(Y_{\mathfrak{B}} > \lambda) &\leq \mathbb{P}(Y_B > \lambda) \\ &\leq \mathbb{P}(\mathfrak{A}_1(s) \vee \mathfrak{A}_1(-s) - \mathfrak{A}_{k+1}(-s) \wedge \mathfrak{A}_{k+1}(s) + \sup_{r \in [-s, s]} |\mathfrak{A}_{k+1}(r) - \mathfrak{A}_{k+1}(s)| \geq \lambda/2 - 3ks) \\ &\quad + e^{c_k} \mathbb{P}(\sup_{r \in [-s, s]} W(s) > \lambda/2), \end{aligned}$$

where W is an independent Brownian bridge from $(-s, 0)$ to $(-s, 0)$. We can bound the first of these terms with Lemma 2.1, Lemma 2.2, and Lemma 3.5. The second of these terms can be bounded with a standard Brownian bridge bound. Altogether, we get that

both of terms are of size $e^{-d_k \lambda^{3/2}}$ as long as $\lambda > c_k s^3$. Therefore setting $\lambda = c_k (s^3 + \|\mathbf{m}\|_2)$ and applying the bound (29), we get

$$e^{c_k \log \lambda} \mathbb{P}(\mathfrak{M} \in A(\mathbf{m}, r)) + \mathbb{P}(M_{\mathfrak{B}} > \lambda) \leq \exp\left(c_k \log(\|\mathbf{m}\|_2 + s) - \frac{4}{3} \sum_{i=1}^k m_i^{3/2}\right).$$

which gives the desired result. \square

We can translate Lemma 4.1 into a one-time bound on \mathfrak{L} .

Lemma 4.2. *For any $r \notin [0, t]$ and $\mathbf{m} \in [0, \infty)_{\geq}^k$ we have*

$$\mathbb{P}(\mathfrak{L} \in A(\mathbf{m}, r)) \leq \exp\left(-\frac{4}{3} \sum_{i=1}^k m_i^{3/2} + c_k \sqrt{t} \left(\sum_{i=1}^k m_i^{3/2}\right)^{5/6}\right).$$

Proof. First, we may assume $\sum_{i=1}^k m_i^{3/2} > t^3$ since otherwise the bound is trivial. We use the notation introduced prior to Lemma 3.1. For every $s > t$, we have that

$$\mathbb{P}(\mathfrak{L} \in A(\mathbf{m}, r)) = \frac{\eta_s(A(\mathbf{m}, r))}{\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))} = \frac{1}{\eta(\mathcal{C}^{\mathbb{N}}(\mathbb{R}))} \int_{A(\mathbf{m}, r)} \frac{d\eta_s}{d\tilde{\eta}_s}(f) d\tilde{\eta}_s(f).$$

Now, the density $\frac{d\eta_s}{d\tilde{\eta}_s}$ can be thought of as an \mathbb{R} -valued random variable whose domain is the Borel space $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$ with probability measure η_s . When thought of this way, it is equal in distribution to

$$\frac{1}{\mathbb{E}_{\mathcal{F}_s}[\mathbb{P}_{0,t}(\mathfrak{B}^{s,k}(0), \mathfrak{B}^{s,k}(t), \mathfrak{A}_{k+1})]}.$$

By the tail bounds in Corollary 3.6 on this random variable, for $s > c_k t$ we then have

$$\int_{A(\mathbf{m}, r)} \frac{d\eta_s}{d\tilde{\eta}_s}(f) d\eta_s(f) \leq [\tilde{\eta}_s(A(\mathbf{m}, r))]^{1-d_k \sqrt{s/t}} e^{c_k s^{5/2} \sqrt{t}}.$$

Therefore by Lemma 4.1, we have that

$$\mathbb{P}(\mathfrak{L} \in A(\mathbf{m}, r)) \leq \exp\left(c_k s^{5/2} \sqrt{t} - (1 - d_k \sqrt{t/s}) \left[\frac{4}{3} \sum_{i=1}^k m_i^{3/2} + c_k \log(\|\mathbf{m}\|_2 + s)\right]\right).$$

We can now minimize over s . This occurs roughly when $s = c_k (\sum_{i=1}^k m_i^{3/2})^{1/3}$ (which is greater than t by our assumption), which gives the desired bound after simplification. \square

4.2 The two-time bound

The proof of a sharp two-time bound is a much more refined version of the argument used Lemma 3.4 that takes into account the parabola. The argument is also applied to the ensemble \mathfrak{L} rather than directly to \mathfrak{A} . The first step is to achieve a lower bound on the probability that \mathfrak{L}_{k+1} is small. This is achieved via the following lemma.

Lemma 4.3. *Let $S = \llbracket k + 1, \infty \rrbracket \times \mathbb{R}$. Then the law of $\mathfrak{L}|_S$ is absolutely continuous with respect to the law of $\mathfrak{A}|_S$ and for any Borel set A with $\mathbb{P}(\mathfrak{A}|_S \in A) = \epsilon > 0$ have*

$$\mathbb{P}(\mathfrak{L}|_S \in A) \leq \epsilon e^{c_k \sqrt{t} (\log^{5/6}(\epsilon^{-1}))}$$

Proof. First, we may assume $\log^{1/3}(\epsilon^{-1}) > c_k t$ for a large constant c_k , since otherwise the bound is trivial. Letting $\mathfrak{B} = \mathfrak{B}_s$ for some $s > c_k t$ we have $\mathfrak{B}_s|_S = \mathfrak{A}|_S$. Moreover, as in the proof of Lemma 4.2 we have

$$\mathbb{P}(\mathfrak{L}|_S \in A) \leq [\mathbb{P}(\mathfrak{B}_s \in A)]^{1-c'_k \sqrt{t/s}} e^{c_k s^{5/2} \sqrt{t}},$$

and minimizing at $s = c_k (\log \mathbb{P}(\mathfrak{B}^s \in A))^{-1/3}$ (which is greater than $c_k t$ by our lower bound on ϵ), gives the result. \square

Corollary 4.4. *Fix an interval $[-\lambda, \lambda]$ for $\lambda \geq 2$. Then for $m > c_k t$ we have*

$$\mathbb{P}\left(\min_{r \in [-\lambda, \lambda]} \mathfrak{L}_{k+1}(r) + r^2 \leq -m\right) \leq \lambda e^{-d_k m^3}.$$

Proof. Let $\tilde{\mathfrak{A}}(r) = \mathfrak{A}(r) + r^2$. First, by Lemma 2.2 and a union bound we have that

$$\mathbb{P}\left(\min_{r \in [-\lambda - \epsilon, \lambda], r \in \mathbb{Z}} \tilde{\mathfrak{A}}_{k+1}(r) \leq -m + 1\right) \leq 2\lambda \epsilon^{-1} e^{-d_k m^3},$$

and by Lemma 3.4, Lemma 2.3, and a union bound

$$\mathbb{P}\left(\min_{r \in [-\lambda - \epsilon, \lambda], r \in \mathbb{Z}, \delta \in [0, \epsilon]} |\tilde{\mathfrak{A}}_{k+1}(r + \delta) - \tilde{\mathfrak{A}}_{k+1}(r)| \leq 1\right) \leq c_k \lambda \epsilon^{-1} e^{-d_k / \epsilon}.$$

Combining these bounds with $\epsilon = m^{-3}$ gives that

$$\mathbb{P}\left(\min_{r \in [-\lambda, \lambda]} \mathfrak{A}_{k+1}(r) + r^2 \leq -m\right) \leq c_k \lambda m^6 e^{-d_k m^3} \leq \lambda e^{-d'_k m^3}.$$

Applying Lemma 4.3 with $\epsilon = \lambda e^{-d'_k m^3}$ and simplifying yields the result. \square

One of the main technical ingredients we need to achieve a two-point bound is a probability estimate for a Brownian bridge avoiding a parabola.

Lemma 4.5. *Fix $\lambda \geq 1$, and for $x, y \geq 0$ with $\sqrt{x} < \lambda/2, \sqrt{y} < \lambda/2$, define*

$$E(x, y, \lambda) = \frac{2}{3} x^{3/2} + (\lambda - \sqrt{y})^2 \sqrt{y} + \frac{1}{3} (\lambda - \sqrt{y})^3, \quad \text{and}$$

$$J(x, y, \lambda) = E(x, y, \lambda) - \frac{(x - y + \lambda^2)^2}{4\lambda}.$$

Next let $\alpha \in \mathbb{R}$, and let $f(s) = -s^2 + \alpha s$. Then we have the following bounds.

1. *For $\lambda > 1$, any $x, y > 0$ with $\sqrt{x} < \lambda/2, \sqrt{y} < \lambda/2$, and any $t \in [0, \sqrt{x} + 1]$, we have*

$$\mathbb{P}_{0, \lambda}(x, y + f(\lambda), f, [t, \lambda]) \leq \exp(-J(x, y, \lambda) + c\lambda \log(1 + \lambda)).$$

2. Let $\lambda > 1, x, y > 0$ with $\sqrt{x} < \lambda/2, \sqrt{y} < \lambda/2$, and let $t \in (0, \sqrt{x} + 1], a \in (0, x]$. If B is a Brownian bridge from $(0, x)$ to $(\lambda, f(\lambda) + y)$, then

$$\mathbb{P}(B(s) > f(s) \text{ for all } s \in [t, \lambda], B(t) \leq x - a) \leq \exp\left(-J(x, y, \lambda) - \frac{a^2}{4t} + \frac{2}{3}x^{3/2} - \frac{2}{3}(x - a)^{3/2} + c(t^3 + tx + \lambda \log(1 + \lambda))\right).$$

3. Let $\mathbf{x}, \mathbf{y} \in [1, \infty)_{\geq}^k$ with $x_i - x_{i+1} \geq 1$ and $y_i - y_{i+1} \geq 1$ for all $i \in \llbracket 1, k-1 \rrbracket$, and suppose that $\sqrt{x_1} < \lambda/2, \sqrt{y_1} < \lambda/2$. Then

$$\mathbb{P}_{0, \lambda}(\mathbf{x}, \mathbf{y} + f(\lambda), f) \geq \exp\left(-\sum_{i=1}^k J(x_i, y_i, \lambda) - ck^2\lambda\right).$$

The proof of Lemma 4.5 is essentially a standard Dirichlet energy computation. We sketch the idea here, and leave the details to the appendix.

First, we can set $\alpha = 0$, as Brownian bridge law commutes with affine shifts. This simplifies the computations. Next, the cost for a Brownian bridge B from $(0, x)$ to $(\lambda, -\lambda^2 + y)$ to roughly follow a path $g : [0, \lambda] \rightarrow \mathbb{R}$ with $g(0) = x, g(\lambda) = -\lambda^2 + y$ is the exponentiated Dirichlet energy

$$\exp\left(-E(g) + \frac{(x - y + \lambda^2)^2}{4\lambda}\right), \quad \text{where} \quad E(g) = \frac{1}{4} \int_0^\lambda |g'(s)|^2 dt.$$

This follows, for example, from the Cameron-Martin theorem. The factor of $1/4$ here (as opposed to the usual $1/2$) is due to the fact that we are working with variance 2 bridges, and the added factor of $(x - y + \lambda^2)^2/(4\lambda)$ comes from conditioning a Brownian motion to end at the point $y - \lambda^2$ to give rise to a Brownian bridge. Hence we should look for a path g of minimal Dirichlet energy that stays above the function $f(s) = -s^2$. This is the minimal concave function $g = g_{x,y}$ from $(0, x)$ to $(\lambda, -\lambda^2 + y)$ that dominates f . Since f is just a parabola, the function g is easily computed, and is determined by the following constraints:

- $g(0) = 0, g(\lambda) = -\lambda^2 + y$, and there are values $0 \leq a < b \leq \lambda$ such that g is linear on the intervals $[0, a]$ and $[b, \lambda]$.
- $g(z) = f(z)$ for all $z \in (a, b)$.
- g is differentiable at a and b . This follows from forcing g to be both concave and to dominate f .

We can compute that $a = \sqrt{x}, b = \lambda - \sqrt{y}$, and that $E(g) = E(x, y, \lambda)$. This gives rise to the bound in Lemma 4.5.1, and the bound in part 3; the cost of having the k bridges stay ordered is a lower order term. The bound in part 2 follows similarly to part 1, by computing the path of minimal Dirichlet energy from $(0, x)$ to $(\lambda, -\lambda^2 + y)$ that dips below $x - a$ at time t .

We can now state the first version of the two-time bound for \mathfrak{L} .

Proposition 4.6. Let $\mathbf{m} \in [0, \infty)_{\geq}^k$, let $\mathbf{a} \in [0, \infty)^k$ and let $\sigma \in \{0, t\}^k$. Define $\hat{\sigma} \in \{0, t\}^k$ to be the unique vector with $\hat{\sigma}_i \neq \sigma_i$ for all i . Next, define

$$A(\mathbf{m}, \mathbf{a}, \sigma) = \{f \in \mathcal{C}^k(\mathbb{R}) : f(\sigma_i) \geq m_i, \quad f_i(\sigma_i) - f_i(\hat{\sigma}_i) \geq a_i \text{ for all } i \in \llbracket 1, k \rrbracket\}.$$

Then

$$\mathbb{P}(\mathfrak{L}^k \in A(\mathbf{m}, \mathbf{a}, \sigma)) \leq \exp\left(-\sum_{i=1}^k \left(\frac{a_i^2}{4t} + \frac{2}{3}m_i^{3/2} + \frac{2}{3}(m_i - a_i)^{3/2}\right) + E(\mathbf{a}, \mathbf{m}, t)\right), \quad (31)$$

where

$$E(\mathbf{a}, \mathbf{m}, t) = O_k(t^3 + \sqrt{t}\|\mathbf{m}\|_2^{5/4} + t\|\mathbf{a}\|_2 + t^{-1/6}\|\mathbf{a}\|_2^{2/3}\|\mathbf{m}\|_2^{1/2}).$$

The proof of Proposition 4.6 is the most technical proof in the paper, and it is based on a backwards induction on $\ell \in \llbracket 1, k \rrbracket$. The next lemma gives the key inductive step.

Lemma 4.7. *In the setup of Proposition 4.6, for $\ell \in \llbracket 1, k+1 \rrbracket$ define the sets*

$$A_\ell(\mathbf{m}, \mathbf{a}, \sigma) = \{f \in \mathcal{C}^k(\mathbb{R}) : f(\sigma_i) \geq m_i, \quad f_i(\sigma_i) - f_i(\hat{\sigma}_i) \geq a_i \text{ for all } i \in \llbracket \ell, k \rrbracket\}.$$

and for $r = 0, t$ define

$$D_\ell^r(\mathbf{m}) = \{f \in \mathcal{C}^k(\mathbb{R}) : f_i(r) \geq m_i \text{ for all } i \in \llbracket 1, \ell \rrbracket\}.$$

Note that with these definitions, $A_{k+1}(\mathbf{m}, \mathbf{a}, \sigma)$, $D_0^0(\mathbf{m})$, and $D_0^t(\mathbf{m})$ are the whole space. Then for all $\ell \in \llbracket 1, k+1 \rrbracket$ and $r = 0, t$ we have

$$\begin{aligned} & \mathbb{P}(\mathfrak{L}^k \in A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^r(\mathbf{m})) \\ & \leq \exp\left(-\sum_{i=\ell}^k \left(\frac{a_i^2}{4t} + \frac{2}{3}m_i^{3/2} + \frac{2}{3}(m_i - a_i)^{3/2}\right) - \sum_{i=1}^{\ell-1} \frac{4}{3}m_i^{3/2} + E(\mathbf{a}, \mathbf{m}, t)\right). \end{aligned} \quad (32)$$

Proposition 4.6 is immediate from Lemma 4.7 since

$$A_1(\mathbf{m}, \mathbf{a}, \sigma) \cap D_0^r(\mathbf{m}) = A(\mathbf{m}, \mathbf{a}, \sigma).$$

Proof. The proof is based on a downwards induction on ℓ . In the base case when $\ell = k+1$, since $A_{k+1}(\mathbf{m}, \mathbf{a}, \sigma)$ is the whole space, the bound is equivalent to the claim that

$$\mathbb{P}(D_k^r(\mathbf{m})) \leq \exp\left(-\sum_{i=1}^k \frac{4}{3}m_i^{3/2} + E(\mathbf{a}, \mathbf{m}, t)\right)$$

for $r = 0, t$. This follows from Lemma 4.2. Now assume $\ell \leq k$, and that the lemma holds for $\ell+1$ for any choice of $\mathbf{a}, \mathbf{m}, \sigma, r$. Without loss of generality, we assume that $\sigma_\ell = 0$; the case when $\sigma_\ell = t$ follows from the distributional equality

$$\mathfrak{L}(r) + r^2 \stackrel{d}{=} \mathfrak{L}(t-r) + (t-r)^2$$

as functions in $\mathcal{C}^{\mathbb{N}}(\mathbb{R})$. This equality follows from the same result for \mathfrak{A} . There are two cases here, depending on whether $r = 0$ or $r = t$. We start with the simpler case when $r = 0$.

Step 1: Setting up a Gibbs resampling. Let $\beta_k > 0$ be a large k -dependent constant and set

$$\gamma = \gamma(\mathbf{a}, \mathbf{m}, t) = \beta_k(t^{3/2} + t^{-1/2}\|\mathbf{a}\|_2 + \|\mathbf{m}\|_2^{3/4}).$$

How large we need to take β_k will be made clear in the proof. Let \mathcal{F} denote the σ -algebra generated by \mathfrak{L} outside of the set $S = \llbracket 1, \ell \rrbracket \times [0, \gamma^{2/3}]$. The basic idea of the induction is a Gibbs resampling on the set S . To see how the structure will work, observe that

$$A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^0(\mathbf{m}) = A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_\ell^0(\mathbf{m}) \cap \{\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell\}.$$

The set $A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_\ell^0(\mathbf{m})$ is \mathcal{F} -measurable, so for any set E we have

$$\begin{aligned} & \mathbb{P}_{\mathcal{F}}(A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^0(\mathbf{m})) \\ &= \mathbf{1}(A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_\ell^0(\mathbf{m})) \mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) \\ &\leq \mathbf{1}(A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_\ell^0(\mathbf{m}) \cap E) \mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) + \mathbf{1}(E^c). \end{aligned} \quad (33)$$

Now, for every $n = 0, 1, \dots$, let

$$\mathbf{m}_n = (m_1 \vee (m_\ell + n), \dots, m_{\ell-1} \vee (m_\ell + n), m_\ell + n, m_{\ell+1}, \dots, m_k),$$

and let $D_n = D_\ell^0(\mathbf{m}_{n-1}) \setminus D_\ell^0(\mathbf{m}_n)$ so that $D_\ell^0(\mathbf{m}) = \bigcup_{n=1}^\infty D_n$. We can write

$$\begin{aligned} & \mathbf{1}(A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_\ell^0(\mathbf{m}) \cap E) \mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) \\ &\leq \sum_{n=1}^\infty \mathbf{1}(A_{\ell+1}(\mathbf{m}, \mathbf{a}, \sigma) \cap D_n) \|\mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) \mathbf{1}(E \cap D_n)\|_\infty, \end{aligned}$$

Taking expectations in (33) and applying the inductive hypothesis we get

$$\begin{aligned} & \mathbb{P}(A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^0(\mathbf{m})) \leq \mathbb{P}(E^c) + \sum_{n=1}^\infty \|\mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) \mathbf{1}(E \cap D_n)\|_\infty \times \\ & \times \exp\left(-\sum_{i=\ell+1}^k \left(\frac{a_i^2}{4t} + \frac{2}{3}m_i^{3/2} + \frac{2}{3}(m_i - a_i)^{3/2}\right) - \frac{4}{3}(m_\ell + n - 1)^{3/2} - \sum_{i=1}^{\ell-1} \frac{4}{3}m_i^{3/2} + E(\mathbf{a}, \mathbf{m}, t)\right). \end{aligned}$$

Concretely then, to complete the proof it is enough to find a set E such that $\mathbb{P}(E^c)$ is bounded above by the right-hand side of (32) and

$$\begin{aligned} & \|\mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) \mathbf{1}(E \cap D_n)\|_\infty \\ &\leq \exp\left(-\frac{a_\ell^2}{4t} + \frac{2}{3}m_\ell^{3/2} - \frac{2}{3}(m_\ell - a_\ell)^{3/2} + E(\mathbf{a}, \mathbf{m}, t) - n\right). \end{aligned} \quad (34)$$

Step 2: Defining E and comparing to a simpler ensemble.

Let E_r be the set

where:

- For all $s \in [-\gamma^{2/3}, \gamma^{2/3}]$, we have $\mathfrak{L}_{\ell+1}(s) + s^2 > -\beta_k \gamma^{2/3}$.
- We have the inequalities

$$\begin{aligned} & -0.9\sqrt{\beta_k} \gamma^{2/3} \leq \mathfrak{L}_\ell(r) + r^2 \leq \mathfrak{L}_1(r) + r^2 \leq \frac{1}{10} \gamma^{4/3} \quad \text{and} \\ & -0.9\sqrt{\beta_k} \gamma^{2/3} \leq \mathfrak{L}_\ell(\pm \gamma^{2/3}) + \gamma^{4/3} \leq \mathfrak{L}_1(\pm \gamma^{-2/3}) + \gamma^{4/3} \leq \frac{1}{10} \gamma^{4/3}, \end{aligned}$$

and set $E = E_0$ (in the final step of the proof we will also use the set E_t). By Lemma 4.2, Corollary 4.4 and a union bound, as long as the constant β_k is large enough we have that

$$\mathbb{P}(E^c) \leq e^{-\gamma^2}, \quad (35)$$

which is bounded above by the right-hand side of (32) for β_k large enough.

Now let $\bar{t} = (1, \dots, \ell)$, and let $B = (B_1, \dots, B_\ell)$ be a k -tuple of independent Brownian bridges from $(0, \mathfrak{L}^\ell(0) - \bar{t})$ to $(\gamma^{2/3}, \mathfrak{L}^\ell(\gamma^{2/3}) - \bar{t})$. Let $f : [0, \gamma^{2/3}] \rightarrow \mathbb{R}$ be given by $f(s) = -s^2 - \sqrt{\beta_k} \gamma^{2/3}$, and let \tilde{B} denote B conditioned on the non-intersection event $\text{NI}(f; [t, \gamma^{2/3}])$.

By Lemma 2.4 and the Gibbs property for \mathfrak{L} on the set S , on the event E the bottom line $\mathfrak{L}_\ell|_{[0, \gamma^{2/3}]}$ stochastically dominates \tilde{B}_ℓ . Therefore on E :

$$\begin{aligned} \mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \mathfrak{L}_\ell(t) > a_\ell) &\leq \mathbb{P}_{\mathcal{F}}(\mathfrak{L}_\ell(0) - \tilde{B}_\ell(t) > a_\ell) \\ &= \mathbb{P}_{\mathcal{F}}(\tilde{B}_\ell(0) - \tilde{B}_\ell(t) > a_\ell - (\ell - 1)). \end{aligned}$$

Now,

$$\mathbb{P}_{\mathcal{F}}(\tilde{B}_\ell(0) - \tilde{B}_\ell(t) > a_\ell - (\ell - 1)) = \frac{\mathbb{P}_{\mathcal{F}}(B \in \text{NI}(f; [t, \gamma^{2/3}]), B_\ell(t) < B_\ell(0) - a + (\ell - 1))}{\mathbb{P}_{0, \gamma^{2/3}}(B(0), B(\gamma^{2/3}), f; [t, \gamma^{2/3}])}, \quad (36)$$

and we can bound the numerator above and denominator below on the event E_0 using Lemma 4.5.

Step 3: Bounding (36). Starting with the denominator in (36), we have that

$$\begin{aligned} \mathbb{P}_{0, \gamma^{2/3}}(B(0), B(\gamma^{2/3}), f; [t, \gamma^{2/3}]) &\geq \mathbb{P}_{0, \gamma^{2/3}}(B(0), B(\gamma^{2/3}), f) \\ &= \mathbb{P}_{0, \gamma^{2/3}}(B(0) + \sqrt{\beta_k} \gamma^{2/3}, B(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3}, g), \end{aligned}$$

where $g(s) = -s^2$. Moreover, on E we have that

$$\max(\mathfrak{L}_1(0) + \sqrt{\beta_k} \gamma^{2/3}, \mathfrak{L}_1(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3} - g(\gamma^{2/3})) \leq \frac{1}{5} \gamma^{4/3}$$

as long as β_k was chosen large enough. Similarly,

$$\mathfrak{L}_\ell(0) + \sqrt{\beta_k} \gamma^{2/3} \geq 2k, \quad \mathfrak{L}_\ell(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3} - g(\gamma^{2/3}) \geq 2k.$$

Therefore we are in the setting of Lemma 4.5.3, and so

$$\begin{aligned} &\mathbb{P}_{0, \gamma^{2/3}}(B(0) + \sqrt{\beta_k} \gamma^{2/3}, B(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3}, g) \\ &\geq \exp\left(-\sum_{i=1}^{\ell} J(B_i(0) + \sqrt{\beta_k} \gamma^{2/3}, B_i(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3} + \gamma^{4/3}, \gamma^{2/3}) - c\ell^2 \gamma^{2/3}\right). \end{aligned}$$

Turning to the numerator, we have that

$$\begin{aligned} &\mathbb{P}_{\mathcal{F}}(B \in \text{NI}(f; [t, \gamma^{2/3}]), B_\ell(t) < B_\ell(0) - a_\ell + (\ell - 1)) \\ &\leq \mathbb{P}_{\mathcal{F}}(B_\ell \in \text{NI}(f; [t, \gamma^{2/3}]), B_\ell(t) < B_\ell(0) - a_\ell + (\ell - 1)) \\ &\quad \times \prod_{i=1}^{\ell-1} \mathbb{P}_{0, \gamma^{2/3}}(B_i(0) + \sqrt{\beta_k} \gamma^{2/3}, B_i(\gamma^{2/3}) + \sqrt{\beta_k} \gamma^{2/3}, g; [t, \gamma^{2/3}]) \end{aligned}$$

The product of $\ell - 1$ terms above can be bounded above by Lemma 4.5.1. Note that $t \leq 0.1\sqrt{\beta_k}\gamma^{2/3} \leq B_i(0) + \sqrt{\beta_k}\gamma^{2/3}$ by the definition of γ as long as β is large enough. The remaining ℓ -term can be bounded using Lemma 4.5.2. Together we get that the right-hand side above is bounded above by

$$\exp\left(-\sum_{i=1}^{\ell} J(B_i(0) + \sqrt{\beta_k}\gamma^{2/3}, B_i(\gamma^{2/3}) + \sqrt{\beta_k}\gamma^{2/3} + \gamma^{4/3}, \gamma^{2/3}) + c_k(t\gamma^{2/3} \log(\gamma) + t^3 + tB_\ell(0))\right. \\ \left. - \frac{(a_\ell - \ell + 1)^2}{4t} + \frac{2}{3}(B_\ell(0) + \sqrt{\beta_k}\gamma^{2/3})^{3/2} - \frac{2}{3}(B_\ell(0) + \sqrt{\beta_k}\gamma^{2/3} - a_\ell + \ell - 1)^{3/2}\right).$$

Putting together the two bounds in this section, and using that $B_\ell(0) = \mathfrak{L}_\ell(0) - (\ell - 1)$ and that $\gamma \gg k$ gives that (36) is bounded above by

$$\exp\left(-\frac{a_\ell^2}{4t} + \frac{2}{3}[\mathfrak{L}_\ell(0)]^{3/2} - \frac{2}{3}[\mathfrak{L}_\ell(0) - a_\ell]^{3/2}\right. \\ \left.+ O_k(t\gamma^{2/3} \log(\gamma) + t^3 + t\mathfrak{L}_\ell(0) + \frac{a_\ell}{t} + \gamma^{2/3}\sqrt{\mathfrak{L}_\ell(0) + \gamma})\right). \quad (37)$$

Now, on the set $E \cap D_n$, we know that $\mathfrak{L}_\ell(0) \in [m_\ell + n - 1, m_\ell + n]$ and as long as β_k was chosen large enough we have $|\mathfrak{L}_\ell(0)| \leq \gamma^{4/3}$. Applying this, and simplifying the error, we get that (37) is bounded above by

$$\exp\left(-\frac{a_\ell^2}{4t} + \frac{2}{3}[m_\ell + n - 1]^{3/2} - \frac{2}{3}[m_\ell - a_\ell]^{3/2} + O_k(t^{3/2}\gamma + t\gamma^{2/3}\sqrt{m_\ell + n})\right) \\ \leq \exp\left(-\frac{a_\ell^2}{4t} + \frac{2}{3}m_\ell^{3/2} - \frac{2}{3}[m_\ell - a_\ell]^{3/2} + O_k(t^{3/2}\gamma) - n\right)$$

which implies the desired bound in (34).

Step 4: Completing the proof when $\sigma_\ell = 0, r = t$. First, with E_t as defined in Step 2 above, by noting that $\mathbb{P}(E_t^c)$ satisfies the same bound as $\mathbb{P}(E^c)$ in (35), it is enough to bound

$$\mathbb{P}(A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^t(\mathbf{m}) \cap E_t).$$

Our strategy here is to bootstrap from the previous case, by using that the Gibbs property implies that it is unlikely for the lines in \mathfrak{L} to be large at time t without being large at time 0. Let \mathcal{G} be the σ -algebra generated by \mathfrak{L} outside of the set $\llbracket 1, \ell - 1 \rrbracket \times [-\gamma^{2/3}, t]$. Noting that the sets $A_\ell(\mathbf{m}', \mathbf{a}, \sigma), D_{\ell-1}^t(\mathbf{m}), E_t$ are \mathcal{G} -measurable for any $\mathbf{m}', \mathbf{a}, \sigma$, for any \mathbf{m}' which equals \mathbf{m} on coordinates ℓ, \dots, k we can write

$$\mathbb{P}(D_{\ell-1}^0(\mathbf{m}') \cap A_\ell(\mathbf{m}, \mathbf{a}, \sigma)) \geq \mathbb{E}[\mathbb{P}_{\mathcal{G}}(D_{\ell-1}^0(\mathbf{m}'))\mathbf{1}(A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^t(\mathbf{m}) \cap E_t)] \\ \geq \frac{1}{2}\mathbb{P}(A_\ell(\mathbf{m}, \mathbf{a}, \sigma) \cap D_{\ell-1}^t(\mathbf{m}) \cap E_t)$$

as long as $\mathbb{P}_{\mathcal{G}}(D_{\ell-1}^0(\mathbf{m}')) \geq 1/2$ almost surely on $D_{\ell-1}^t(\mathbf{m}) \cap E_t$. Now, on $D_{\ell-1}^t(\mathbf{m}) \cap E$ and conditional on \mathcal{G} , by Lemma 2.4 the vector $\mathfrak{L}^k(0)$ stochastically dominates an $(\ell - 1)$ -tuple of Brownian bridges B from $(-2\gamma^{4/3} - 1, \dots, -2\gamma^{4/3} - \ell)$ at time $-\gamma^{2/3}$ to $(m_1 - 1, \dots, m_\ell - \ell)$ at time t , conditioned not to intersect each other. The probability of non-intersection is

bounded below by $\exp(-ck \log \gamma)$ by Lemma 2.5, so $\mathbb{P}(\mathfrak{L}^k(0) \geq \mathbf{m}') \geq 1/2$ for any \mathbf{m}' for which $\mathbb{P}(B(0) \geq \mathbf{m}') \geq 1 - e^{-ck \log \gamma}/2$. This holds with $\mathbf{m}' = \mathbf{m} - c_k \sqrt{t} \gamma^{2/3}$. Therefore

$$\mathbb{P}(D_{\ell-1}^t(t) \cap A_\ell(\mathbf{m}, \mathbf{a}, \sigma)) \leq 2\mathbb{P}(D_{\ell-1}^0(t) \cap A_\ell(\mathbf{m} - c_k \sqrt{t} \gamma^{2/3}, \mathbf{a}, \sigma)),$$

which satisfies the same upper bound as $\mathbb{P}(D_{\ell-1}^0(t) \cap A_\ell(\mathbf{m}, \mathbf{a}, \sigma))$, up to a term that gets folded into the error. \square

Proposition 4.6 implies the following simpler two-point bound, of which inequality (4) is a special case.

Corollary 4.8. *Let $\mathbf{a} \in [0, \infty)^k$ with $\|\mathbf{a}\|_2 \geq c_k t^2$ and let $\sigma \in \{0, t\}^k$. Let $\hat{\sigma} \in \{0, t\}^k$ be the unique vector with $\hat{\sigma}_i \neq \sigma_i$ for all i . Next, let $f(\mathbf{a}, \sigma) \in \mathcal{C}_0^k([0, t])$ be the unique k -tuple of linear functions satisfying $f_i(\sigma_i) - f_i(\hat{\sigma}_i) = a_i$, and recall the definition of the map \mathcal{S} introduced for Theorem 1.1. Then*

$$\begin{aligned} & \mathbb{P}(\mathfrak{L}_i(\sigma_i) - \mathfrak{L}_i(\hat{\sigma}_i) \geq a_i \text{ for all } i \in \llbracket 1, k \rrbracket) \\ & \leq \exp\left(-\mathcal{S}(f(\mathbf{a}, \sigma)) - \sum_{i=1}^k \frac{a_i^2}{4t} + O_k(\sqrt{t} \|\mathbf{a}\|_2^{5/4})\right) \end{aligned} \quad (38)$$

Proof. Without loss of generality, assume $\sigma_k = 0$. Then using the notation \mathbb{T} for the tetris map introduced after Theorem 1.1 we have that

$$\mathfrak{L}_k(t) \geq -\alpha \quad \implies \quad \mathfrak{L}(0) \geq \mathbb{T}f(0) - \alpha, \quad \mathfrak{L}(t) \geq \mathbb{T}f(t) - \alpha,$$

since the vectors $\mathfrak{L}^k(0), \mathfrak{L}^k(t)$ are ordered. Therefore for any $\alpha > 0$, with notation as in Proposition 4.6 we have

$$\mathbb{P}(\mathfrak{L}_i(\sigma_i) - \mathfrak{L}_i(\hat{\sigma}_i) \geq a_i \forall i \in \llbracket 1, k \rrbracket) \leq \mathbb{P}(\mathfrak{L} \in A(\mathbb{T}f(0) \vee \mathbb{T}f(t) - \alpha, \mathbf{a}, \sigma)) + \mathbb{P}(\mathfrak{L}_k(t) \leq -\alpha).$$

Setting $\alpha = c_k(t^2 + t^{-1/3} \|\mathbf{a}\|_2^{2/3})$, by Corollary 4.4, we have that $\mathbb{P}(\mathfrak{L}_k(t) \leq -\alpha)$ is bounded above by the right-hand side of (38) for large enough c_k . This uses that $Tf_1 \leq c_k \|\mathbf{a}\|_2$ and that $\|\mathbf{a}\|_2 \geq t^2$. For this choice of α , the left-hand side is similarly bounded via Proposition 4.6. We have simplified the error by using that $Tf_1 \leq c_k \|\mathbf{a}\|_2$ and that $\|\mathbf{a}\|_2 \geq t^2$. \square

5 The upper bound in Theorem 1.1

In this section we prove the upper bound in Theorem 1.1. Throughout the section we fix $k \in \mathbb{N}, t \geq 1$. For this, it is enough to prove that the upper bound holds for $\mathfrak{A}^k|_{[1, t+1]} - \mathfrak{A}^k(1)$, since in law $\mathfrak{A}^k|_{[1, t+1]} - \mathfrak{A}^k(1)$ and $\mathfrak{A}^k|_{[0, t]} - \mathfrak{A}^k(0)$ only differ by a linear shift which can be wrapped into the error. Moreover, we can work with $\mathfrak{L} = \mathfrak{L}^{t+2, k}$ rather than \mathfrak{A} , since the conditional probability that \mathfrak{L} is non-intersecting $-e^{-O_k(t^3)}$ -can again be wrapped into the error term in Theorem 1.1. The key step is to convert Proposition 4.6 into a density estimate.

For this lemma, we define a σ -finite measure μ on $X = \mathbb{R}^k \times \mathbb{R}^k \times \mathcal{C}^k([1, t+1])$ as follows. First let μ' denote the product measure on X given by Lebesgue measure on $\mathbb{R}^k \times \mathbb{R}^k$ and the law of k independent Brownian bridges from $(1, 0)$ to $(t+1, 0)$ on $\mathcal{C}^k([1, t+1])$. Let μ denote the pushforward of μ' under the map

$$(\mathbf{x}, \mathbf{y}, f) \mapsto (\mathbf{x}, \mathbf{y}, f + L_{\mathbf{x}, \mathbf{y}}),$$

where $L_{\mathbf{x}, \mathbf{y}}$ is the function which is linear in each coordinate and satisfies $L(1) = \mathbf{x}, L(t+1) = \mathbf{y}$. Then the law of $(\mathfrak{A}^k(1), \mathfrak{A}^k(t+1), \mathfrak{A}^k|_{[1, t+1]})$ is absolutely continuous with respect to μ . The next lemma bounds its density.

Lemma 5.1. *Let $Y = Y(\mathbf{x}, \mathbf{y}, g)$ denote the density on $\mathbb{R}^k \times \mathbb{R}^k \times \mathcal{C}^k([t, t+1])$ of $(\mathfrak{A}^k(1), \mathfrak{A}^k(t+1), \mathfrak{A}^k|_{[1, t+1]})$ against the measure μ . Then we have the following three bounds:*

1. For all $(\mathbf{x}, \mathbf{y}, g) \in \mathbb{R}^k \times \mathbb{R}^k \times \mathcal{C}^k([t, t+1])$ we have

$$Y(\mathbf{x}, \mathbf{y}, g) \leq \exp(-d_k |\inf g_k|^3 \mathbf{1}(\inf g_k \leq -c_k t^2) + c_k t^3).$$

2. For all $(\mathbf{x}, \mathbf{y}, g) \in \mathbb{R}^k \times \mathbb{R}^k \times \mathcal{C}^k([t, t+1])$ we have

$$Y(\mathbf{x}, \mathbf{y}, g) \leq \exp\left(-\left(\frac{\|\mathbf{x} - \mathbf{y}\|_2^2}{4t} + \sum_{i=1}^k \frac{2}{3} [(x_i^+)^{3/2} + (y_i^+)^{3/2}]\right) + O_k(t^3 + \sqrt{t}(\|\mathbf{x}\|_2^{5/4} + \|\mathbf{y}\|_2^{5/4}))\right).$$

Proof. Throughout, we work with \mathfrak{L} conditional on the event $\mathfrak{L}_1 > \dots > \mathfrak{L}_k > \mathfrak{L}_{k+1}$, which is equal in law to \mathfrak{A} . Note that this event has probability $e^{-O_k(t^3)}$.

Bound 1: First, since $\mathfrak{L}^k|_{[0, t+2]}$ given $\mathfrak{L}^k(0), \mathfrak{L}^k(t+2)$ is simply given by k independent Brownian bridges, the Lebesgue density of $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1))$ is bounded above by a constant $c_k > 0$. Moreover, $\mathfrak{L}^k|_{[1, t+1]}$ has the law of k independent Brownian bridges from $(1, \mathfrak{L}^k(1))$ to $(t+1, \mathfrak{L}^k(t+1))$ given $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1))$. Therefore the density of $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1), \mathfrak{L}^k|_{[1, t+1]})$ against μ is uniformly bounded by a constant c_k , and so Y is a.s. by $e^{O_k(t^3)}$.

Note that this density bound also holds a.s. if we condition on any information about \mathfrak{L} outside of $[[1, k] \times [0, t+2]]$. In particular, if we let Y_α be the conditional density of $(\mathfrak{A}^k(1), \mathfrak{A}^k(t+1), \mathfrak{A}^k|_{[1, t+1]})$ against μ on the event $A_\alpha = \{\inf \mathfrak{L}_{k+1}^c|_{[0, t+2]} \leq -\alpha\}$ and Z_α the conditional density on A_α^c we have

$$\begin{aligned} Y(\mathbf{x}, \mathbf{y}, g) &= Y_\alpha(\mathbf{x}, \mathbf{y}, g) \mathbb{P}(A_\alpha) + Z_\alpha(\mathbf{x}, \mathbf{y}, g) \mathbb{P}(A_\alpha^c) \\ &\leq e^{c_k t^3} \mathbb{P}(A_\alpha) + Z_\alpha(\mathbf{x}, \mathbf{y}, g). \end{aligned}$$

If $-\alpha > \inf g_k$, then $Z_\alpha(\mathbf{x}, \mathbf{y}, g) = 0$ by the ordering of the lines in \mathfrak{A} . Moreover, $\mathbb{P}(A_\alpha) \leq (t+2)e^{-d_k \alpha^3}$ by Corollary 4.4 for $\alpha \geq c_k t^2$, which yields the desired bound when $\inf g_k \leq -c_k t^2$.

Bound 2: Let f be the Lebesgue density of $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1))$. It is enough to prove the second bound with $Y(\mathbf{x}, \mathbf{y}, g)$ replaced by $f(\mathbf{x}, \mathbf{y})$, since the $e^{O(t^3)}$ cost of conditioning \mathfrak{L} to be non-intersecting folds into the error, and $\mathfrak{L}^k|_{[1, t+1]}$ has the law of k independent Brownian bridges from $(1, \mathfrak{L}^k(1))$ to $(t+1, \mathfrak{L}^k(t+1))$ given $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1))$.

For $\mathbf{m}, \mathbf{n} \in \mathbb{Z}_{\geq}^k$, let $f(\mathbf{x}, \mathbf{y}, \mathbf{m}, \mathbf{n})$ be the conditional Lebesgue density of $(\mathfrak{L}^k(1), \mathfrak{L}^k(t+1))$ at (\mathbf{x}, \mathbf{y}) on the event

$$C(\mathbf{m}, \mathbf{n}) = \{[\mathfrak{L}^k(0)] = \mathbf{m}, [\mathfrak{L}^k(t+2)] = \mathbf{n}\}.$$

Since $\mathfrak{L}^k|_{[0, t+2]}$ is simply a k -tuple of Brownian bridges from $\mathfrak{L}^k(0)$ to $\mathfrak{L}^k(t+2)$, we have that $f(\mathbf{x}, \mathbf{y}, \mathbf{m}, \mathbf{n})$ is bounded above by

$$c_k \exp\left(-\frac{\|\mathbf{x} - \mathbf{y}\|_2^2}{4t} - \frac{\|\mathbf{x} - \mathbf{m}\|_2^2}{4} - \frac{\|\mathbf{y} - \mathbf{n}\|_2^2}{4} + \frac{\|\mathbf{m} - \mathbf{n}\|_2^2}{4(t+2)} + \frac{\sum_{i=1}^k |x_i - m_i| + |y_i - n_i| + |n_i - m_i|}{4}\right).$$

The final error term comes from the fact that $\mathfrak{L}^k(0) \in [\mathbf{m}, \mathbf{m} + 1]$, and $\mathfrak{L}^k(t) \in [\mathbf{n}, \mathbf{n} + 1]$. Next, from Proposition 4.6, we have the bound

$$\mathbb{P}(C(\mathbf{m}, \mathbf{n})) \leq \exp\left(-\frac{\|\mathbf{m} - \mathbf{n}\|_2^2}{4(t+2)} - \sum_{i=1}^k \left(\frac{2}{3}[(m_i^+)^{3/2} + (n_i^+)^{3/2}]\right) + O_k(\sqrt{t}(\|\mathbf{m}\|_2^{5/4} + \|\mathbf{n}\|_2^{5/4}) + t^3)\right)$$

and by Corollary 4.4 we have $\mathbb{P}(\cup_{m_k \wedge n_k \leq -\alpha} C(\mathbf{m}, \mathbf{n})) \leq 2 \exp(-d_k \alpha^3)$ for $\alpha > c'_k t^2$. Therefore for $\alpha > c'_k t^2$, we have

$$f(\mathbf{x}, \mathbf{y}) \leq 2 \exp(-d_k \alpha^3) + \sum_{\substack{\mathbf{m}, \mathbf{n} \in \mathbb{Z}_{\geq}^k \\ n_k \wedge m_k \geq \alpha}} \exp(-D(\mathbf{n}, \mathbf{m})) \quad (39)$$

where

$$D(\mathbf{n}, \mathbf{m}) = \frac{\|\mathbf{x} - \mathbf{y}\|_2^2}{4t} + O_k(t^3 + \|\mathbf{x}\|_2 + \|\mathbf{y}\|_2) + \sum_{i=1}^k D_{x_i}(m_i) + D_{y_i}(n_i)$$

and

$$D_z(\ell) = \frac{(z - \ell)^2}{4} + \frac{2}{3}(\ell^+)^{3/2} - c_k \sqrt{t} |\ell|^{5/4}.$$

We set $\alpha = c'_k(t^2 + \|\mathbf{x}\|_2^{3/4} + \|\mathbf{y}\|_2^{3/4})$ so that

$$2 \exp(-d_k \alpha^3) \leq \exp(-100D(\mathbf{x}, \mathbf{y})). \quad (40)$$

Now,

$$-D(\mathbf{n}, \mathbf{m}) \leq -\frac{1}{3}(m_1 \vee n_1)^{3/2} - 100D(\mathbf{x}, \mathbf{y}) \quad (41)$$

whenever $m_1 \vee n_1 > (\|\mathbf{x}\|_2 + \|\mathbf{y}\|_2 + 2)^{c'_k}$. All but $(2t^2 + \|\mathbf{x}\|_2 + \|\mathbf{y}\|_2)^{c'_k}$ terms in the sum in (39) satisfy this bound. Therefore by (39), (40), and (41) we have

$$\begin{aligned} f(\mathbf{x}, \mathbf{y}) &\leq \exp(-100D(\mathbf{x}, \mathbf{y}) + c'_k \log(\|\mathbf{x}\|_2 + \|\mathbf{y}\|_2 + 2)) \\ &\quad + \max_{\mathbf{m}, \mathbf{n} \in \mathbb{Z}_{\geq}^k, n_k \wedge m_k \geq \alpha} \exp(-D(\mathbf{n}, \mathbf{m}) + c'_k \log(\|\mathbf{x}\|_2 + \|\mathbf{y}\|_2 + 2)) \\ &\leq \max_{\mathbf{m}, \mathbf{n} \in \mathbb{Z}_{\geq}^k, n_k \wedge m_k \geq \alpha} \exp(-D(\mathbf{n}, \mathbf{m})), \end{aligned}$$

since the log factor is lower order compared with the O_k -term in D . It just remains to bound $D(\mathbf{m}, \mathbf{n})$ below. Indeed, the bound in the lemma will follow if we can show that for every $z \in \mathbb{R}$ we have

$$D_z(\ell) \geq \frac{2}{3}(z^+)^{3/2} + O_k(t^3 + \|\mathbf{x}\|_2 + \|\mathbf{y}\|_2 + \sqrt{t}|z|^{5/4}) \quad (42)$$

whenever $\ell \geq -\alpha$. First, for $\ell \in [-\alpha, 0]$ we have

$$D_z(\ell) \geq \frac{(z^+)^2}{4} + c_k \sqrt{t} |\alpha|^{5/4} = \frac{(z^+)^2}{4} + O_k(t^3 + \|\mathbf{x}\|_2 + \|\mathbf{y}\|_2), \quad (43)$$

which is bounded below by the right-hand side of (42) for any z . Next, for $\ell \geq 0$ we always have the bound

$$D_z(\ell) \geq \frac{2}{3}\ell^{3/2} - c_k \sqrt{t} \ell^{5/4} = O_k(t^3)$$

which is bounded below by the right-hand side of (42) when $z \leq c_k'' t^2$. Finally, for $z \geq c_k'' t^2$ and $\ell \geq 0$, we can differentiate D_z to get

$$D'_z(\ell) = \frac{2(z - \ell)}{t} - \sqrt{\ell} + \frac{5}{4} c_k \sqrt{t} \ell^{1/4}$$

As long as c_k'' was chosen large enough relative to c_k , D'_z has all its roots in the region where $|\ell - z| \leq 20t\sqrt{z}$, and hence the minimum value of D_z over $\ell \geq 0$ is taken in this interval. Finally, for $|\ell - z| \leq 20t\sqrt{z}$ and $z \geq c_k'' t^2$ we have the estimate

$$D_z(\ell) \geq \frac{2}{3}z^{3/2} - 20tz - 2c_k \sqrt{t}|z|^{5/4},$$

which completes the proof of (42) when $z \geq c_k'' t^2$. \square

We are now ready to prove the upper bound in Theorem 1.1.

Corollary 5.2. *With notation as in Theorem 1.1, for $\mu_{k,t}$ -a.e. $f \in \mathcal{C}_0^k([0, t])$ we have that*

$$X_{k,t}(f) = \exp(-S(f) + O_k(t^3 + \sqrt{t}S(f)^{5/6})). \quad (44)$$

Proof. With notation as in Lemma 5.1 we can write

$$X_{k,t}(f) = \int Y(\mathbf{x}, \mathbf{x} + f(t), f) \sqrt{4\pi t} \exp\left(\frac{\|f(t)\|_2^2}{4t}\right) d\mathbf{x}. \quad (45)$$

Observe that $Y(\mathbf{x}, \mathbf{x} + f(t), f) = 0$ unless $f_1 + x_1 > f_2 + x_2 > \dots > f_k + x_k$. Equivalently, we require

$$\mathbf{x} - x_k \mathbf{1} \geq \mathbb{T}f(0) - \mathbb{T}f_k(0) \mathbf{1} \quad (46)$$

where $\mathbf{1} = (1, \dots, 1)$. here the functional \mathbb{T} is the Tetris map from Theorem 1.1. Next, by Lemma 5.1.2 we can see that the integrand in (45) is bounded above by

$$\exp\left(-\frac{2}{3} \sum_{i=1}^k [(x_i^+)^{3/2} + ([x_i + f_i(t)]^+)^{3/2}] + O_k(t^3 + \sqrt{t}(\|\mathbf{x}\|_2^{5/4} + \|f(t)\|_2^{5/4}))\right). \quad (47)$$

Also, by Lemma 5.1.1, for $x_k - \mathbb{T}f_k(0) \leq -c_k t^2$, the integrand is bounded above by

$$\exp\left(-d_k |x_k - \mathbb{T}f_k(0)|^3 + \|f(t)\|_2^2 + c_k t^3\right),$$

Together with (47), this bounds implies that the contribution to the integral from the region where either $x_1 \geq c_k(t^2 + S^{2/3}(f))$ or $x_k \leq \mathbb{T}f_k(0) - c'_k(t^2 + S(f)^{1/3})$ is at most $e^{-S(f)}$ which is bounded above by the right-hand side of (44).

The size of the remaining region is $\exp(O_k(\log(S(f) + 2)))$, which folds in to the error, so it enough to simply maximize (47) subject to the constraint (46) on the set where $x_1 \geq c_k(t^2 + S(f)^{2/3})$ or $x_k \leq \mathbb{T}f_k(0) - c'_k(t^2 + S(f)^{1/3})$. Observe that on this set, the error term in (47) is at most $O_k(t^3 + \sqrt{t}S(f)^{5/6})$ so it suffices to understand the main term. As the main term $\frac{2}{3} \sum_{i=1}^k [(x_i^+)^{3/2} + ([x_i + f_i(t)]^+)^{3/2}]$ is decreasing in each x_i , the minimal value occurs at $\mathbf{x} = \mathbb{T}f(0) - c'_k(t^2 + S(f)^{1/3})$. Plugging in this value for \mathbf{x} yields

$$S(f) - O_k((t^2 + S(f)^{1/3})\sqrt{t^2 + S(f)^{2/3}}) = S(f) - O_k(t^3 + \sqrt{t}S(f)^{5/6}),$$

as desired. \square

6 The lower bound in Theorem 1.1

In this section we complete the proof of Theorem 1.1. For this section $k \in \mathbb{N}, t \geq 1$, and $f \in \mathcal{C}^k([0, t])$ are fixed throughout and we use the notation \mathbb{S}, \mathbb{T} from Theorem 1.1. The proof of the lower bound for Theorem 1.1 involves a Brownian Gibbs resampling on a staircase-shaped region of the form

$$U = \bigcup_{i=1}^k \{i\} \times [x_i^-, x_i^+],$$

where $\mathbf{x}^-, \mathbf{x}^+ \in \mathbb{R}^k$, and as in Figure 3 we roughly take $x_i^- \sim -\sqrt{\mathbb{T}f_i(0)}$ and $x_i^+ \sim t + \sqrt{\mathbb{T}f_i(t) + t^2}$. For technical reasons, we need to space the values x_i^\pm out slightly. Define

$$x_i^- = -\sqrt{\mathbb{T}f_i(0)} - (k + 2 - i), \quad x_i^+ = \sqrt{\mathbb{T}f_i(t) + t^2} + t + (k + 2 - i), \quad (48)$$

With these definitions, we have the separation

$$x_i^- + 1 \leq x_{i+1}^-, \quad x_{i+1}^+ + 1 \leq x_i^+,$$

and $x_i^- \leq -2, x_i^+ \geq 2t$ for all i .

One major simplification when applying a Gibbs resampling argument for the lower bound is that we can work only with a part of space where \mathfrak{A} behaves nicely on the outer boundary of U . The next lemma sets up this nice boundary behaviour for \mathfrak{A} . For this lemma and throughout this section, let

$$E_{t,f} = 2t^2 + \mathbb{T}f_1(0) + \mathbb{T}f_1(t).$$

The errors in the section will be expressed in terms of $E_{t,f}$.




Figure 3: The setup for the lower bound in Theorem 1.1. The goal is to apply a Brownian Gibbs resampling to create the above configuration, where the red lines are tangent to the parabola. The start and end locations of the red lines for the top curve are indicated. The cost $S(f)$ is (up to a lower order term) the Dirichlet energy difference between the red lines and the corresponding parabolic segments.

Lemma 6.1. *Let A_m be the event where*

$$|\mathfrak{A}_i(y) + y^2| \leq m \log^{2/3}(3kE_{t,f}) \quad \text{for all } y \in [-3kE_{t,f}, 3kE_{t,f}], i \in \llbracket 1, k \rrbracket.$$

Also, for each $i \in \llbracket 1, k \rrbracket$, let $C_\epsilon^{i,-}$ be the set where

$$\mathbb{P}_{x_i^-, x_{i+1}^-}(\mathfrak{A}_i(x_i^-), \mathfrak{A}_i(x_i^- + 1), \mathfrak{A}_{i+1}) \geq \epsilon.$$

In words, $C_\epsilon^{i,-}$ is the set where the probability that an independent Brownian bridge B from $(x_i^-, \mathfrak{A}_i(x_i^-))$ to $(x_i^- + 1, \mathfrak{A}_i(x_i^- + 1))$ stays above \mathfrak{A}_{i+1} is at least ϵ . Similarly let $C_\epsilon^{i,+}$ be the set where

$$\mathbb{P}_{x_i^+ - 1, x_i^+}(\mathfrak{A}_i(x_i^+ - 1), \mathfrak{A}_i(x_i^+), \mathfrak{A}_{i+1}) \geq \epsilon.$$

Then there exists a k -dependent constant $m_k > 0$ such that for $m \geq m_k$ we have

$$\mathbb{P}(D_m) \geq 1/2, \quad \text{where} \quad D_m = A_m \cap \bigcap_{i=1}^k C_{1/m}^{i,-} \cap C_{1/m}^{i,+}$$

Note the implicit dependence on t, f in the events in the lemma; the claim is that we can take m_k independent of t, f .

Proof. By a union bound, it is enough to show that

$$\lim_{m \rightarrow \infty} \sup_{t \geq 1, f} \mathbb{P}A_m = 1, \quad \lim_{\epsilon \rightarrow 0} \sup_{t \geq 1, f} \mathbb{P}C_\epsilon^{i,*} = 1 \quad \text{for all } i \in \llbracket 1, k \rrbracket, * \in \{+, -\}.$$

By Lemmas 2.1 and 2.2 and a union bound, we have

$$|\mathfrak{A}_j(i) + i^2| \leq m \log^{2/3}(3kE_{t,f})$$

for all integers $i \in \mathbb{Z} \cap [-3kE_{t,f} - 1, 3kE_{t,f} + 1]$ and all $j \in \llbracket 1, k \rrbracket$ with probability tending to 1 with m , uniformly in t and f . Next, for all $i \in \mathbb{Z} \cap [-3kE_{t,f} - 1, 3kE_{t,f} + 1]$ and $j \in \llbracket 1, k \rrbracket$, by Lemma 3.4 each of the processes $\mathfrak{A}_j(s) + s^2$, $s \in [i, i + 1]$ satisfies the conditions of the modulus of continuity Lemma 2.3 with common $\beta = 1/5$ and $\alpha = c_k$. Therefore by another union bound, $\mathbb{P}A_m \rightarrow 1$ as $m \rightarrow \infty$ uniformly in t and f .

Next, since $t \mapsto \mathfrak{A}(t) + t^2$ is stationary, the probabilities $\mathbb{P}C_\epsilon^{i,\pm}$ do not depend on the exact values of x_i^\pm , and hence they do not depend on f, t . Moreover, the strict ordering of the Airy line ensemble implies that for every i, ϵ , we have $\mathbb{P}C_\epsilon^{i,*} \rightarrow 1$ as $\epsilon \rightarrow 0$, as desired. \square

One of the main difficulties in applying a Gibbs resampling to make rigorous the picture in Figure xxx is that dealing with non-intersection costs that come from an $O(1)$ region near the boundary of U . We can handle this on the event D_m with the following lemma.

Setting some notation, let $\mathbf{x} \in \mathbb{R}_{\geq}^k$, $\mathbf{y} \in \mathbb{R}_{\leq}^k$ and define $I_k = [x_k, y_k]$, $I_j = [x_j, y_j] \setminus I_{j+1}$ for $j < k$. Consider a k -tuple of functions $h = (h_2, \dots, h_{k+1})$ such that the domain of f_i contains I_{i-1} for all i . Let

$$\text{NI}(\mathbf{x}, \mathbf{y}, h) = \left\{ g \in \prod_{j=1}^k \mathcal{C}([x_j, y_j]) : g_1(r) > \dots > g_j(r) > h_{j+1}(r) \text{ for all } r \in I_j, j \in \llbracket 1, k \rrbracket \right\},$$

and let $\mathbb{P}_{\mathbf{x}, \mathbf{y}}(\mathbf{u}, \mathbf{v}, h)$ be the probability that k independent Brownian bridges from (x_i, u_i) to (y_i, v_i) lie in the set $\text{NI}(\mathbf{x}, \mathbf{y}, f)$. Finally we set

$$\zeta = \zeta_m = m \log^{2/3}(3kE_{t,f}).$$

Lemma 6.2. *Define $\mathfrak{A}^k(\mathbf{x}^\pm) \in \mathbb{R}_{>}^k$ by*

$$\mathfrak{A}^k(\mathbf{x}^\pm) = (\mathfrak{A}_1(x_1^\pm), \dots, \mathfrak{A}_k(x_k^\pm)),$$

and let $\iota = (k, k-1, \dots, 1)$. Then there exists a k -dependent constant $m'_k \geq 0$ such that for $m \geq m'_k$, on D_m we have the following almost sure lower bounds on non-intersection probabilities:

$$\mathbb{P}_{\mathbf{x}^-, 0}(\mathfrak{A}^k(\mathbf{x}^-), Tf(0) + 2\zeta_m \iota, \mathfrak{A}|_{U^c}) \geq e^{-c_k m^2 \sqrt{E_{t,f}}}. \quad (49)$$

and

$$\mathbb{P}_{t, \mathbf{x}^+}(Tf(t) + 2\zeta_m \iota, \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}) \geq e^{-c_k m^2 \sqrt{E_{t,f}}}. \quad (50)$$

Proof. Step 1: Defining events for the proof of (49). Define

$$\ell_i(s) = 2\sqrt{\Gamma f_i(0)}s + \Gamma f_i(0) + 2(k-i+1)\zeta_m.$$

Observe that for all $i \in \llbracket 1, k \rrbracket$, $s \in \mathbb{R}$ we have

$$\ell_i(s) \geq -s^2 + 2\zeta_m. \quad (51)$$

Moreover, for $i \in \llbracket 1, k-1 \rrbracket$ we have

$$\ell_i(s) \geq \ell_{i+1}(s) + 2\zeta_m, \quad s \in [x_{i+1}^- + 1, 0]. \quad (52)$$

The event D_m implies that for all $(i, s) \in \llbracket 1, k+1 \rrbracket \times [x_1^-, x_1^+]$ we have $\mathfrak{A}_i(s) \leq -s^2 + \zeta_m$. Therefore if B is a k -tuple independent Brownian bridges from $(x_i^-, \mathfrak{A}^k(x_i^-))$ to $(0, \mathbb{T}f(0) + 2\zeta_m\iota)$, then $B \in \text{NI}(\mathbf{x}^-, 0, \mathfrak{A}|_{U^c})$ as long as the following three events hold for all $i \in \llbracket 1, k \rrbracket$ and m is sufficiently large.

1. $G_{1,i}$: We have $B_i(x_i^- + 1) \in [\ell_i(x_i^- + 1) - 1, \ell_i(x_i^- + 1)]$.
2. $G_{2,i}$: For all $s \in [x_i^- + 1, 0]$ we have $|B_i(s) - \ell_i(s)| \leq 2$.
3. $G_{3,i}$: For all $s \in [x_i^-, x_i^- + 1]$ we have $\mathfrak{A}_{i+1}(s) < B_i(s) < \ell_{i-1}(s) - \zeta_m$. Here the upper inequality is dropped if $i = 1$.

Therefore conditional on $\mathfrak{A}|_U$, we have that

$$\begin{aligned} \mathbb{P}_{\mathbf{x}^-, 0}(\mathfrak{A}^k(\mathbf{x}^-), \mathbb{T}f(0) + 2\zeta_m\iota, \mathfrak{A}|_{U^c}) &\geq \mathbb{P}\left(\bigcap_{i=1}^k G_{1,i} \cap G_{2,i} \cap G_{3,i}\right) \\ &\geq \prod_{i=1}^k \mathbb{P}(G_{1,i})\mathbb{P}(G_{2,i} | G_{1,i})\mathbb{P}(G_{3,i} | G_{1,i}). \end{aligned}$$

Here the final inequality uses independence properties of the Brownian bridges B_i .

Step 2: Bounding $\mathbb{P}(G_{1,i})$, $\mathbb{P}(G_{2,i} | G_{1,i})$, and $\mathbb{P}(G_{3,i} | G_{1,i})$ on D_m . We have

$$\mathbb{P}(G_{1,i}) = \frac{1}{\sqrt{4\pi(|x_i^-| - 1)/|x_i^-|}} \int_{\ell_i(x_i^- + 1) - 1}^{\ell_i(x_i^- + 1)} \exp\left(-\frac{(u - \mu_i)^2}{4(|x_i^-| - 1)/|x_i^-|}\right) du,$$

where

$$\mu_i = \frac{|x_i^-| - 1}{|x_i^-|} \mathfrak{A}_i(x_i^-) + \frac{1}{|x_i^-|} [\mathbb{T}f_i(0) + 2(k - i + 1)\zeta_m].$$

Now, on D_m we have $|\mathfrak{A}_i(x_i^-) + (x_i^-)^2| \leq \zeta_m$, so using that $x_i^- = -\sqrt{\mathbb{T}f_i(0)} - (k + 2 - i)$ and $x_i^- \leq -2$ we can check that

$$|\mu_i + (x_i^- + 1)^2| \leq c_k \zeta_m$$

for a k -dependent constant $c_k > 0$. The same bound holds with $\ell_i(x_i^- + 1)$ in place of μ_i . Therefore

$$\mathbb{P}(G_{1,i}) \geq e^{-c_k \zeta_m^2} = e^{-c_k m^2 \log^{4/3}(3kE_{t,f})}. \quad (53)$$

Next, $\mathbb{P}(G_{2,i} | G_{1,i})$ is bounded below by the probability that a Brownian bridge from $(x_i^- + 1, 0)$ to $(0, 0)$ has maximum absolute value at most 1. Therefore

$$\mathbb{P}(G_{2,i} | G_{1,i}) \geq e^{-cx_i^-} \quad (54)$$

for an absolute constant $c > 0$. We turn our attention to $G_{3,i}$. We will condition on the value of $u = B_i(x_i^- + 1)$ given $u \in [\ell_i(x_i^- + 1) - 1, \ell_i(x_i^- + 1)]$; the bounds on u are implied

by the event $G_{1,i}$. Let L_u be the linear function with $L_u(x_i^-) = \mathfrak{A}_i(x_i^-)$ and $L_u(x_i^- + 1) = u$. Since $\mathfrak{A}_i(x_i^-) \leq -(x_i^-)^2 + \zeta_m$, by (51) and (52) we have $\ell_{i-1}(s) - \zeta_m \geq L_u(s)$. Therefore for all large enough m , we have

$$\mathbb{P}(B_i(s) < \ell_{i-1}(s) - \zeta_m \text{ for all } s \in [x_i^-, x_{i+1}^-] \mid u = B_i(x_i^- + 1)) \geq 1 - (2m)^{-1}$$

for all $u \in [\ell_i(x_i^- + 1) - 1, \ell_i(x_i^- + 1)]$. Moreover, $\ell_i(x_i^- + 1) - 1 \geq \mathfrak{A}_i(x_i^- + 1)$ by (51) on the event D_m . Therefore on D_m ,

$$\begin{aligned} \mathbb{P}(B_i(s) > \mathfrak{A}_{i+1}(s) \text{ for all } s \in [x_i^-, x_{i+1}^-] \mid u = B_i(x_i^- + 1)) \\ \geq \mathbb{P}_{x_i^-, x_{i+1}^-}(\mathfrak{A}_i(x_i^-), \mathfrak{A}_i(x_i^- + 1), \mathfrak{A}_{i+1}) \geq 1/m, \end{aligned}$$

where the final inequality follows since D_m implies the event $C_{1/m}^{i,-}$. Combining these two bounds by a union bound gives that

$$\mathbb{P}(G_{3,i} \mid G_{1,i}) \geq 1/(2m). \quad (55)$$

The bounds (53), (54), and (55) then yield (49).

Step 3: The proof of (50). Define

$$\tilde{\mathfrak{A}}_i(s) = \mathfrak{A}_i(t - s) + t^2 - 2st, \quad \tilde{U} = \bigcup_{i=1}^k \{i\} \times [-x_i^+ + t, -x_i^- + t],$$

and note that $\tilde{\mathfrak{A}}$ is another parabolic Airy line ensemble. Now, since Brownian bridge law commutes with affine shifts,

$$\mathbb{P}_{t, \mathbf{x}^+}(\Upsilon f(t) + 2\zeta_m t, \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}) = \mathbb{P}_{-\mathbf{x}^+ + t, 0}(\tilde{\mathfrak{A}}^k(-\mathbf{x}^+ + t), Z(f) + 2\zeta_m t, \tilde{\mathfrak{A}}^k|_{\tilde{U}^c}),$$

where $Z(f) = \Upsilon f(t) + t^2$. Via this transformation, the proof of (50) follows in the same way as the proof of (49) with Z in place of Y , and $-x_i^+ + t$ in place of x_i^- . Indeed, the only inputs we required from \mathfrak{A} also hold for $\tilde{\mathfrak{A}}$ on D . These are parabolic shape bounds implied by the event A_m , and the bound $\mathbb{P}_{x_i^-, x_{i+1}^-}(\mathfrak{A}_i(x_i^-), \mathfrak{A}_i(x_i^- + 1), \mathfrak{A}_{i+1}) \geq 1/m$; the analogue

$$\mathbb{P}_{-x_i^+ + t, -x_{i+1}^- + t}(\tilde{\mathfrak{A}}_i(-x_i^+ + t), \tilde{\mathfrak{A}}_i(-x_i^+ + t + 1), \tilde{\mathfrak{A}}_{i+1}) \geq 1/m$$

is implied by the event $C_{1/m}^{i,+}$. □

We can now prove the upper bound in Theorem 1.1. In fact, we get a slightly stronger estimate on the error than what is claimed in the introduction.

Proposition 6.3. *We have*

$$X_{k,t}(f) \geq \exp(-S(f) + O_k(tE_{t,f})) = \exp(-S(f) + O_k(t^3 + tS(f)^{2/3})).$$

Proof. Step 1: A conditional density formula. Let \mathcal{F} be the σ -algebra generated by \mathfrak{A} on the set U^c . We first use the Brownian Gibbs property to give a formula for the conditional expectation

$$\mathbb{E}_{\mathcal{F}}(X_{k,t}(f)). \quad (56)$$

Let $\nu_{\mathbf{u}^-, \mathbf{u}^+}$ be the law of k independent Brownian bridges B_1, \dots, B_k , where B_i goes from (x_i^-, u_i^-) to (x_i^+, u_i^+) , and let $\tilde{\nu}_{\mathbf{u}^-, \mathbf{u}^+}$ be the pushforward of $\nu_{\mathbf{u}^-, \mathbf{u}^+}$ onto $\mathbb{R}^k \times \mathcal{C}_0^k([0, t])$ under the map

$$(g_1, \dots, g_k) \mapsto [(g_1(0), \dots, g_k(0)); (g_1|_{[0, t]} - g_1(0), \dots, g_k|_{[0, t]} - g_k(0))]. \quad (57)$$

Then

$$\begin{aligned} \frac{d\tilde{\nu}_{\mathbf{u}^-, \mathbf{u}^+}}{d\mathbf{y}d\mu_{k, t}}(\mathbf{y}, f) &= \left(\frac{(x_i^+ + |x_i^-|)}{(x_i^+ - t)|x_i^-|} \right)^{k/2} \\ &\times \prod_{i=1}^k \exp \left(-\frac{(y_i - u_i^-)^2}{4|x_i^-|} - \frac{(u_i^+ - y_i - f_i(t))^2}{4(x_i^+ - t)} + \frac{(u_i^+ - u_i^-)^2}{4(x_i^+ + |x_i^-|)} \right). \end{aligned} \quad (58)$$

Now, for a function $h : \llbracket 1, k+1 \rrbracket \times [x_1^-, x_1^+] \setminus U \rightarrow \mathbb{R}$, define the conditional measure $\eta_{\mathbf{u}^-, \mathbf{u}^+, h}$ on $\mathcal{C}(U)$ by

$$\eta_{\mathbf{u}^-, \mathbf{u}^+, h}(A) = \frac{\nu_{\mathbf{u}^-, \mathbf{u}^+}(A \cap \text{NI}(\mathbf{x}^-, \mathbf{x}^+, h))}{\mathbb{P}_{\mathbf{x}^-, \mathbf{x}^+}(\mathbf{u}^-, \mathbf{u}^+, h)},$$

and let $\tilde{\eta}_{\mathbf{u}^-, \mathbf{u}^+, h}$ be the pushforward of $\eta_{\mathbf{u}^-, \mathbf{u}^+, h}$ under the map (57). Then

$$\frac{d\tilde{\eta}_{\mathbf{u}^-, \mathbf{u}^+, h}}{d\tilde{\nu}_{\mathbf{u}^-, \mathbf{u}^+}}(\mathbf{y}, f) = \frac{\mathbb{P}_{\mathbf{x}^-, 0}(\mathbf{u}^-, \mathbf{y}, h) \mathbb{P}_{t, \mathbf{x}^+}(\mathbf{y} + f(t), \mathbf{u}^+, h) \mathbf{1}(f_1 + y_1 > \dots > f_k + y_k > h_{k+1})}{\mathbb{P}_{\mathbf{x}^-, \mathbf{x}^+}(\mathbf{u}^-, \mathbf{u}^+, h)}. \quad (59)$$

Now, the conditional expectation (56) equals

$$\int_{\mathbb{R}^k} \frac{d\tilde{\eta}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}}}{d\mathbf{y}d\mu^k}(\mathbf{y}, f) d\mathbf{y} = \int_{\mathbb{R}^k} \frac{d\tilde{\eta}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}}}{d\tilde{\nu}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+)}} \frac{d\tilde{\nu}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+)}}{d\mathbf{y}d\mu^k}(\mathbf{y}, f) d\mathbf{y} \quad (60)$$

Let $m > 0$ be large enough so that Lemmas 6.1 and 6.2 hold, and let \tilde{D}_m be the \mathcal{F} -measurable event where $\mathbb{P}_{\mathcal{F}}(D_m) > 0$. We will use (59) and (58) to bound (60) below on the event \tilde{D}_m .

Step 2: Bounding (59) on \tilde{D}_m . Let $\zeta_m = m \log^{2/3}(3kE_{t, f})$ be as in Lemma 6.2, and let W be the set of all $\mathbf{y} \in \mathbb{R}^k$ satisfying

$$y_i \in (\mathbb{T}f_i(0) + (k-i+1)\zeta_m, \mathbb{T}f_i(0) + (k-i+1)\zeta_m + 1). \quad (61)$$

On the set \tilde{D}_m (which implies $\mathfrak{A}_{k+1} \leq \zeta_m$) if $\mathbf{y} \in W$ we have

$$\mathbf{1}(f_1 + y_1 > f_2 + y_2 > \dots > f_k + y_k > \mathfrak{A}_{k+1}) = 1. \quad (62)$$

Next, for any $\mathbf{u}^-, \mathbf{u}^+, h$, the avoidance probabilities $\mathbb{P}_{\mathbf{x}^-, 0}(\mathbf{u}^-, \mathbf{y}, h)$ and $\mathbb{P}_{t, \mathbf{x}^+}(\mathbf{y} + f(t), \mathbf{u}^+, h)$ are monotone increasing if we add $\mathbf{z} \in [0, \infty)_>$ to the coordinate \mathbf{y} . Therefore if we let $\mathbf{y} \in W$, and set $\iota = (k, k-1, \dots, 1)$, then

$$\begin{aligned} &\mathbb{P}_{\mathbf{x}^-, 0}(\mathfrak{A}^k(\mathbf{x}^-), \mathbf{y}, \mathfrak{A}|_{U^c}) \mathbb{P}_{t, \mathbf{x}^+}(\mathbf{y} + f(t), \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}) \\ &\geq \mathbb{P}_{\mathbf{x}^-, 0}(\mathfrak{A}^k(\mathbf{x}^-), \mathbb{T}f(0) + 2\zeta_m \iota, \mathfrak{A}|_{U^c}) \mathbb{P}_{y, \mathbf{x}^+}(\mathbb{T}f(0) + 2\zeta_m \iota, \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}) \\ &\geq e^{-c_k m^2 \sqrt{E_{t, f}}}. \end{aligned} \quad (63)$$

where the second inequality holds on \tilde{D}_m by Lemma 6.2.

Moreover, $\mathfrak{A}_{k+1}(x) \geq -x^2 - \zeta_m$ for $x \in [x_1^-, x_1^+]$ on \tilde{D}_m . Therefore for any \mathbf{u}^\pm , letting $p(x) = -x^2$ we have

$$\mathbb{P}_{\mathbf{x}^-, \mathbf{x}^+}(\mathbf{u}^-, \mathbf{u}^+, \mathfrak{A}|_{U^c}) \leq \prod_{i=1}^k \mathbb{P}_{x_i^-, x_i^+}(u_i^-, u_i^+, \mathfrak{A}_{k+1}) \leq \prod_{i=1}^k \mathbb{P}_{x_i^-, x_i^+}(u_i^-, u_i^+, -p - \zeta_m).$$

We bound each of the terms in this product when $|u_i^\pm + (x_i^\pm)^2| \leq \zeta_m$. First suppose that $x_i^+ - x_i^- \geq 2\sqrt{2\zeta_m}$. Then we can apply Lemma 4.5.1 to get that the i th term in the product is bounded above by

$$\exp\left(c(x_i^+ - x_i^-) \log(x_i^+ - x_i^-) - J(u_i^- + (x_i^-)^2 + \zeta_m, u_i^+ + (x_i^+)^2 + \zeta_m, x_i^+ - x_i^-)\right).$$

Since $|u_i^\pm - (x_i^\pm)^2| \leq \zeta_m$, simple algebra shows that this is bounded above by

$$\begin{aligned} & \exp\left(c(x_i^+ - x_i^-) \log(x_i^+ - x_i^-) + \frac{(u_i^- - u_i^+)^2}{4(x_i^+ - x_i^-)} - \frac{(x_i^+)^3 + |x_i^-|^3}{3} - 2\zeta_m(x_i^+ - x_i^-)\right) \\ & \leq \exp\left(\frac{(u_i^- - u_i^+)^2}{4(x_i^+ - x_i^-)} - \frac{1}{3}\left([\Upsilon f_i(0)]^{3/2} + [\Upsilon f_i(t)]^{3/2}\right) + c_{k,m}E_{t,f}\right) \end{aligned}$$

where $c_{k,m}$ is a k, m -dependent constant. On the other hand, if $x_i^+ - x_i^- \leq 2\sqrt{2\zeta_m}$, then the same bound holds trivially as long as $c_{k,m}$ is large enough, since $\Upsilon f_i(0) \leq |x_i^-|^2$ and $\Upsilon f_i(t) \leq |x_i^+|^2$.

Putting this computation together with (59), (62), and (63) implies that for $\mathbf{y} \in W$ we have

$$\begin{aligned} & \frac{d\tilde{\eta}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+), \mathfrak{A}|_{U^c}}}{d\tilde{\nu}_{\mathfrak{A}^k(\mathbf{x}^-), \mathfrak{A}^k(\mathbf{x}^+)}} \geq \\ & \exp\left(\sum_{i=1}^k \left[\frac{1}{3}\left([\Upsilon f_i(0)]^{3/2} + [\Upsilon f_i(t)]^{3/2}\right) - \frac{(\mathfrak{A}_i(x_i^+) - \mathfrak{A}_i(x_i^-))^2}{4(x_i^+ + |x_i^-|)}\right] - c_{k,m}E_{t,f}\right). \end{aligned} \quad (64)$$

Step 3: Bounding (58) below on \tilde{D}_m . We turn our attention to bounding (58) below on \tilde{D}_m for $\mathbf{y} \in W$, and $\mathbf{u}^\pm = \mathfrak{A}^k(\mathbf{x}^\pm)$.

First, noting that $|x_i^-|, x_i^+ \leq \sqrt{E_{t,f}}$ and $x_i^+ \geq 2t$, the factor prior to the exponential in (58) is bounded below by

$$\exp(-ck \log(E_{t,f}))$$

for an absolute constant $c > 0$. Next, on \tilde{D}_m we have that $u_i^\pm + (x_i^\pm)^2 \in [-\zeta_m, \zeta_m]$ and for $\mathbf{y} \in W$ we have $y_i \geq \zeta_m$ for all i , so $y_i - u_i^+ \geq 0$. Moreover, using that $x_i^+ = \sqrt{\Upsilon f_i(t) + t^2} + t + (k+2-i)$ and that $y_i + f_i(t) - \Upsilon f_i(t) \in [2\zeta_m, 3k\zeta_m]$ for $i \in \llbracket 1, k \rrbracket$, $\mathbf{y} \in W$ we have

$$\begin{aligned} \frac{(u_i^+ - y_i - f_i(t))^2}{4(x_i^+ - t)} & \leq \frac{(2\Upsilon f_i(t) + c_k t \sqrt{E_{t,f}})^2}{4\sqrt{\Upsilon f_i(t) + t^2}} \\ & \leq [\Upsilon f_i(t)]^{3/2} + c_k t E_{t,f}. \end{aligned}$$

Similarly,

$$\frac{(y_i - u_i^-)^2}{4|x_i^-|} \leq [\mathbb{T}f_i(0)]^{3/2} + c_k t E_{t,f}$$

Combining all three of these bounds, on \tilde{D}_m for $\mathbf{y} \in W$, and $\mathbf{u}^\pm = \mathfrak{A}^k(\mathbf{x}^\pm)$, (58) is bounded below by

$$\prod_{i=1}^k \exp\left(-[\mathbb{T}f_i(0)]^{3/2} - [\mathbb{T}f_i(t)]^{3/2} + \frac{(\mathfrak{A}_i(x_i^+) - \mathfrak{A}_i(x_i^-))^2}{4(x_i^+ + |x_i^-|)} - c_k t E_{t,f}\right).$$

Putting this together with (64) and using that the Lebesgue measure of W is 1 gives that on \tilde{D}_m , (56) is bounded below by

$$\exp\left(-\sum_{i=1}^k \frac{2}{3} \left([\mathbb{T}f_i(0)]^{3/2} + [\mathbb{T}f_i(t)]^{3/2}\right) - c_k t E_{t,f}\right).$$

The proposition follows since $\mathbb{P}(\tilde{D}_m) \geq \mathbb{P}(D_m) \geq 1/2$ by Lemma 6.1. \square

7 Proofs of Corollaries and Remark 1.5

Corollaries 1.3 and 1.6 have easy proofs.

Proof of Corollary 1.3. By Theorem 1.1, letting $u_f = \mathbb{T}f_1(0) + \mathbb{T}f_1(t)$ we have

$$X_{k,t}(f) \leq \exp\left(c_k t^3 + c_k \sqrt{t} u_f^{5/4} - \frac{1}{3} u_f^{3/2}\right),$$

which is bounded above by $e^{O_k(t^3)}$ for all f . \square

Proof of Corollary 1.6. The usual Schilder's theorem is Corollary 1.6 with $\nu_{k,t}$ replaced by the Brownian measure $\mu_{k,t}$. The version for $\nu_{k,t}$ follows since for any Borel set A and any $\epsilon > 0$, we have

$$\log \mu_{k,t}^\epsilon(A) - c_k t^3 - d_k \log^{3/4}[\mu_{k,t}^\epsilon(A)^{-1}] \leq \log \nu_{k,t}^\epsilon(A) \leq \log \mu_{k,t}^\epsilon(A) + c_k t^3$$

by Corollaries 1.3 and 1.4. \square

We need to work harder to prove Corollary 1.4. We will need the following elementary lemma about Brownian motion.

Lemma 7.1. *Let $B \in \mathcal{C}_0([0, t])$ be a Brownian motion, and let*

$$S = S(B) = \sup_{s \in [0, t]} B(s), \quad I = I(B) = - \inf_{s \in [0, t]} B(s).$$

Then $(S, I, B(t))$ has a Lebesgue density on the set $\{(a, b, x) \in [0, \infty)^2 \times \mathbb{R} : -b \leq x \leq a\}$ given by $h(a, b, x) + h(a, b, -x)$, where

$$h(a, b, x) = \left[\frac{(2a + 2b - x)^2}{8\sqrt{\pi}t^{5/2}} - \frac{1}{2\sqrt{\pi}t^{3/2}} \right] \exp\left(-\frac{(2a + 2b - x)^2}{4t}\right)$$

Proof. This follows from a reflection principle argument. Fix $a \geq 0, b \geq 0$ and let

$$\tau_a = \inf\{s \in [0, t] : B(s) = a\}, \quad \tau_b = \inf\{s \in [0, t] : B(s) = -b\},$$

and let $\tau_1 = \tau_a \wedge \tau_b, \tau_2 = \tau_a \vee \tau_b$. Let \tilde{B} be the unique continuous process satisfying

$$\tilde{B}|_{[0, \tau_1]} = B|_{[0, \tau_1]}, \quad \tilde{B}|_{[\tau_1, \tau_2]} - \tilde{B}(\tau_1) = -B|_{[\tau_1, \tau_2]} + B(\tau_1), \quad B|_{[\tau_2, t]} - \tilde{B}(\tau_2) = B|_{[\tau_2, t]} - B(\tau_2).$$

Then by the strong Markov property, \tilde{B} is another Brownian motion, and for $x \in [-b, a]$ we have

$$\{S \geq a, I \geq b, B(t) = x\} = \{\tilde{B}(t) = 2a + 2b + x\} \cup \{-\tilde{B}(t) = 2a + 2b - x\}.$$

Therefore

$$\mathbb{P}(S \geq a, I \geq b, B(t) \in dx) = \frac{1}{\sqrt{4\pi t}} \left[\exp\left(-\frac{(2a + 2b - x)^2}{4t}\right) + \exp\left(-\frac{(2a + 2b + x)^2}{4t}\right) \right] dx.$$

Differentiating in both a and b gives the joint density. \square

We also require a straightforward optimization lemma.

Lemma 7.2. Define $W^k = \{\mathbf{u} = (\mathbf{a}, \mathbf{b}, \mathbf{x}) \in [0, \infty)^{2k} \times \mathbb{R}^k : -b_i \leq x_i \leq a_i \text{ for all } i\}$, and define $g : W^k \rightarrow \mathbb{R}$ by

$$g(\mathbf{u}) = \frac{2}{3} \left(\sum_{i=1}^k (Y_i(\mathbf{a}, \mathbf{b}))^{3/2} + (Y_i(\mathbf{a}, \mathbf{b}) + x_i)^{3/2} \right), \quad Y_i(\mathbf{a}, \mathbf{b}) = \sum_{j=i+1}^k a_j + \sum_{j=i}^k b_j.$$

Now, let

$$D_\beta = \{\mathbf{u} = (\mathbf{a}, \mathbf{b}, \mathbf{x}) \in W^k : \sum_{i=1}^k (2a_i + 2b_i - |x_i|)^2 \leq \beta^2\}.$$

Then

$$\max\{g(\mathbf{u}) : \mathbf{u} \in D_\beta\} = \beta^{3/2} \max\{\Theta(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^k, \|\mathbf{x}\|_2^2 = 1\} = \beta^{3/2} \alpha_k$$

where $\Theta : [0, \infty)^k \rightarrow \mathbb{R}, \alpha_k$ are as in Corollary 1.4.

Proof. First, observe that

$$\begin{aligned} \beta^{3/2} \max\{\Theta(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^k, \|\mathbf{x}\|_2^2 = 1\} &= \max\{\Theta(\mathbf{x}) : \mathbf{x} \in [0, \infty)^k, \|\mathbf{x}\|_2^2 = \beta\} \\ &= \max\{g(\mathbf{x}, 0, \mathbf{x}) : \mathbf{x} \in [0, \infty)^k, \|\mathbf{x}\|_2^2 = \beta\} \\ &= \max\{g(\mathbf{x}, 0, \mathbf{x}) : (\mathbf{x}, 0, \mathbf{x}) \in D_\beta\}, \end{aligned}$$

so it suffices to prove that the maximum of g is obtained at a point of the form $(\mathbf{x}, 0, \mathbf{x})$. For $i \in \llbracket 1, k \rrbracket$, observe that the map from $W^k \rightarrow W^k$ sending

$$(a_i, b_i, x_i) \mapsto (a_i - x_i, b_i + x_i, -x_i)$$

and fixing all other coordinates of $\mathbf{u} = (\mathbf{a}, \mathbf{b}, \mathbf{x})$ is an involution on D_β that fixes g . Therefore the maximum of g on D_β is attained at a point where all $x_i \geq 0$. Next, when $x_i \geq 0$, we have

$$[2\partial_{a_i} - \partial_{b_i} + 2\partial_{x_i}]g \geq 0 \quad \text{and} \quad [\partial_{b_i} - \partial_{a_i}]g \geq 0.$$

Therefore letting $\mathbf{e}_i = (0, \dots, 1, \dots, 0) \in \mathbb{R}^k$ be the i th coordinate vector, the maximum of g on D_β is attained at a point $(\mathbf{a}, \mathbf{b}, \mathbf{u})$ where all $x_i \geq 0$ and

$$(\mathbf{a} + 2\epsilon\mathbf{e}_i, \mathbf{b} - \epsilon\mathbf{e}_i, \mathbf{x} + 2\epsilon\mathbf{e}_i) \notin D_\beta, \quad (\mathbf{a} - \epsilon\mathbf{e}_i, \mathbf{b} + \epsilon\mathbf{e}_i, \mathbf{x}) \notin D_\beta$$

for all $\epsilon > 0, i \in \llbracket 1, k \rrbracket$. Since $[2\partial_{a_i} - \partial_{b_i} + 2\partial_{x_i}](2a_i + 2b_i - |x_i|) = 0$ and $[\partial_{b_i} - \partial_{a_i}](2a_i + 2b_i - |x_i|) = 0$ for $x_i \geq 0$, the first of these constraints implies that $b_i = 0$. The second constraint then implies that $a_i = x_i$, as desired. \square

Proof of Corollary 1.4. See Figure 4 for a sketch of the types of functions that maximize $S(f)$ relative to the probability that Brownian motion is close to f (which is controlled by the Dirichlet energy of f). Throughout we will use the notation from the statement of Lemma 7.2. Let B be a k -dimensional Brownian motion in $\mathcal{C}_0^k([0, t])$, and define $S = (S(B_1), \dots, S(B_k)), I = (I(B_1), \dots, I(B_k))$ with $S(B_i), I(B_i)$ as in Lemma 7.1. By Lemma 7.1, $(S, I, B(t))$ has a joint density $f(\mathbf{a}, \mathbf{b}, \mathbf{x})$ on W^k which is bounded above by

$$u^2 t^{-5/2} e^{-u^2/4t}, \quad \text{where } u = u(\mathbf{a}, \mathbf{b}, \mathbf{x}) = \sqrt{\sum_{i=1}^k (2a_i + 2b_i - |x_i|)^2}.$$

Now, shells in W^k of the form $D_{v+\epsilon} \setminus D_v$ with $\epsilon \leq 1$ have volume bounded above by $c_k \epsilon v^{k-1}$, so

$$\int_{W^k \setminus D_v} f(\mathbf{a}, \mathbf{b}, \mathbf{x}) d\mathbf{a} d\mathbf{b} d\mathbf{x} \leq t^{-5/2} \int_v^\infty u^{k+1} e^{-u^2/4t} du \leq c_k v^k e^{-v^2/4t}.$$

In particular, if A is any set with $\mathbb{P}(B \in A) \geq \epsilon$, then on a set $A' \subset A$ with $\mathbb{P}(B \in A') \geq \epsilon/2$, we have

$$(S, I, B(t)) \in D_{h(\epsilon)}, \quad \text{where } h(\epsilon) = [4t \log(\epsilon^{-1})]^{1/2} + c_k \sqrt{t}.$$

To lower bound $\nu_{k,t}(A)$, using the notation of Lemma 7.2 we have

$$\top B_i(0) \leq Y_i(S, I), \quad \top B_i(t) \leq Y_i(S, I) + B_i(t).$$

Therefore by Theorem 1.1, we have

$$\nu_k(A) \geq \epsilon \exp(-\sup\{g(\mathbf{a}, \mathbf{b}, \mathbf{x}) + e(\mathbf{a}, \mathbf{b}, \mathbf{x}) : (\mathbf{a}, \mathbf{b}, \mathbf{x}) \in D_{h(\epsilon)}\}),$$

where g is as in Lemma 7.2 and

$$e(\mathbf{a}, \mathbf{b}, \mathbf{x}) = c_k (t^3 + \sqrt{t} (Y_1(\mathbf{a}, \mathbf{b}) + |\mathbf{x}_1(t)|)^{5/4}).$$

First, since $|x_i| \leq a_i \vee b_i$ for all i , we have

$$\sqrt{\sum_{i=1}^k (2a_i + 2b_i - |x_i|)^2} \geq \sqrt{\sum_{i=1}^k (a_i + b_i)^2} \geq \frac{1}{k} \sum_{i=1}^k (a_i + b_i).$$

Therefore on $D_{h(\epsilon)}$,

$$e(\mathbf{a}, \mathbf{b}, \mathbf{x}) \leq c_k (t^3 + \sqrt{t} [h(\epsilon)]^{5/4}).$$

Now, we maximized g over $D_{h(\epsilon)}$ in Lemma 7.2. This gives that

$$\nu_k(A) \geq \epsilon \exp\left(-[h(\epsilon)]^{3/2} \alpha_k - c_k(t^3 + \sqrt{t}[h(\epsilon)]^{5/4})\right),$$

which gives the desired lower bound in the corollary after simplification.

For the upper bound, let $\mathbf{y} \in [0, \infty)^k$ be any argmax of Θ on the unit sphere. For $\beta > 0$, define let $A_\beta \subset \mathcal{C}_0^k([0, t])$ be the set of functions where

$$\{(-1)^i f_i(t) \geq \beta y_i, i \in \llbracket 1, k \rrbracket\}.$$

We can find $\beta(\epsilon)$ such that $\mathbb{P}(B \in A_{\beta(\epsilon)}) = \epsilon$. To bound the value of $\beta(\epsilon)$ observe that for $\beta > 2\sqrt{t}$ we have

$$\mathbb{P}(B \in A_\beta) \geq \exp\left(-c_k \log(\beta^2/(4t)) - \frac{\beta^2}{4t}\right).$$

Hence $\beta(\epsilon) \geq [4t \log(\epsilon^{-1})]^{1/2} - c_k \sqrt{t}$. Now, for $f \in A_\beta$ we have

$$\Upsilon f(0) \geq \beta \left(y_i \mathbf{1}(i \notin 2\mathbb{Z}) + \sum_{j=i+1}^k y_j \right), \quad \Upsilon f(t) \geq \beta \left(y_i \mathbf{1}(i \in 2\mathbb{Z}) + \sum_{j=i+1}^k y_j \right).$$

Hence for $f \in A_\beta$, by Theorem 1.1 we have

$$X_{t,k}(f) \leq \exp\left(-\beta^{3/2} \Theta(\mathbf{y}) - O_k(t^3 + \sqrt{t}\beta^{5/4})\right),$$

and so using that $\Theta(\mathbf{y}) = \alpha_k$ we get

$$\nu_{k,t}(A_{\beta(\epsilon)}) \leq \epsilon \exp\left(-\beta(\epsilon)^{3/2} \alpha_k - O_k(t^3 + \sqrt{t}\beta(\epsilon)^{5/4})\right),$$

which gives the desired upper bound after simplifying the expression and grouping errors. \square

Finally, we prove Remark 1.5.

Proposition 7.3. *Let $\hat{\mathcal{A}}_1 = \mathcal{A}_1 - \mathcal{A}_1(0)$, the recentered Airy line ensemble. Then we can find a sequence of sets $D_n \subset \mathcal{C}_0([0, 1])$ with $\mu_{1,1}(D_n) \rightarrow 0$ such that*

$$\mathbb{P}(\hat{\mathcal{A}}_1 \in D_n) \geq \mu_{1,1}(D_n) e^{c \log^{1/2}(\mu_{1,1}(D_n)^{-1})}$$

for all $n \in \mathbb{N}$.

It is not difficult to use Theorem 1.1 to find the optimal value of c and show that the $\log^{1/2}$ -decay in the exponent is sharp, but we do not pursue this as it is a rather peripheral point.

Proof. Let $\hat{\mathfrak{A}}_1(r) = \mathfrak{A}_1(r) - \mathfrak{A}_1(0) = \mathfrak{A}_1(r) - r^2 - \mathcal{A}_1(0)$ and let $B : [0, t] \rightarrow \mathbb{R}$ be a Brownian motion. Fix $m > 0$ and consider the class of functions $A_{m,s} \subset \mathcal{C}_0([0, 1])$ such that

$$f(0) = 0, \quad f(1) \in [-1, s], \quad \inf f \geq -1, \quad f(1/2) \geq m.$$




Figure 4: A sketch of $\mathbb{T}f$ for a function $f \in \mathcal{C}_0^k([0, t])$ that maximizes $\mathbb{S}(f)$ relative to the probability that Brownian motion is close to f . Here (y_1, y_2, y_3) is a nonnegative maximizer of the function Θ on the sphere and $\beta > 0$.

Equation (2) implies that $X_{1,1}(f)$ is bounded below by an absolute constant on $A_{m,1}$, and $\hat{\mathcal{A}}_1 \in A_{m,1}$ implies $\hat{\mathcal{A}}_1 \in A_{m+1/4,2}$. Combining this observation with standard estimates on Brownian motion, we get that

$$\mathbb{P}(\hat{\mathcal{A}}_1 \in A_{m+1/4,2}) \geq c\mathbb{P}(B \in A_{m,1}) \geq c'e^{m/4}\mathbb{P}(B \in A_{m+1/4,2}).$$

where c, c' , are absolute constants. Since $\mathbb{P}(B \in A_{m+1/4,2}) \geq de^{-c''m^2}$, taking $D_n = A_{n+1/4,2}$ gives the result. \square

8 Appendix: Proof of Lemma 4.5

As discussed after the statement of Lemma 4.5, we may assume that $\alpha = 0$, since Brownian bridges commute with linear shifts.

Proof of Lemma 4.5.1. First, the bound is trivial for small λ if $c > 0$ is large enough. Therefore throughout the proof we may assume $\lambda > 1000$.

Step 1: A Brownian bridge estimate. Let B be a Brownian bridge from $(0, x)$ to $(\lambda, y - \lambda^2)$. Now, let $\zeta = \lambda/|\lambda|$ and define the mesh

$$\Pi = \{\pi_0, \dots, \pi_m\} := \{0, i\zeta, \dots, [\lambda]\zeta\}.$$

Since $t \leq \sqrt{x} + 1$ we have

$$\mathbb{P}_{0,\lambda}(x, y + f(\lambda), f, [t, \lambda]) \leq \mathbb{P}(B(n) > f(n) \text{ for all } n \in \Pi, n \geq \sqrt{x} + 1). \quad (65)$$

Now, letting π_j be the smallest element of $\Pi \cap [\sqrt{x} + 1, \lambda]$ we have that

$$\mathbb{P}(B(n) > f(n) \text{ for all } n \in \Pi, n < \sqrt{x} + 1 \mid B(\pi_j) > f(\pi_j)) \geq e^{-c\lambda}$$

for an absolute constant $c > 0$. To see this, observe that if $L : \mathbb{R} \rightarrow \mathbb{R}$ is the linear function satisfying $L(0) = x$ and $L(\pi_j) = B(\pi_j)$ for some $B(\pi_j) > f(\pi_j)$ then $L(s) + c' > -s^2$ for an absolute constant $c' > 0$, and the probability that $B(n) \geq L(n) + c'$ for $n \in \Pi, n < \sqrt{x} + 1$ is easily bounded below by $e^{-c\lambda}$. Therefore (65) is bounded above by

$$e^{c\lambda} \mathbb{P}(B(n) > f(n) \text{ for all } n \in \Pi). \quad (66)$$

Step 2: Converting to a Dirichlet energy bound. Consider the vector $\Delta B \in \mathbb{R}^m$ given by $\Delta B_i = B(\pi_i) - B(\pi_{i-1})$, and let E be the minimal value of $\frac{1}{4\zeta} \|\Delta B\|_2^2$ among all possible choices of B with $B(n) > f(n)$ for all $n \in \Pi$. The vector $\Delta B = \sqrt{2\zeta} N$, where N is a standard normal vector on \mathbb{R}^m conditioned on the event $\sum_{i=1}^m N_i = (y - \lambda^2 - x)/\sqrt{2\zeta}$, and so (66) is bounded above by $e^{c\lambda} \mathbb{P}(\|N\|_2^2 \geq 2E)$. Next, by rotational invariance of the standard normal distribution, $\|N\|_2^2$ has the same distribution as if we conditioned N to lie in the hyperplane $\{\mathbf{x} \in \mathbb{R}^m : x_1 = (y - \lambda^2 - x)/\sqrt{2m\zeta}\}$. Therefore

$$\|N\|_2^2 = (y - \lambda^2 - x)^2 / (2m\zeta) + \chi_{m-1},$$

where χ_{m-1} is a χ -squared random variable with $m-1$ degrees of freedom. Therefore letting $I = 2E - (y - \lambda^2 - x)^2 / (2m\zeta)$, we have

$$\mathbb{P}(\|N\|_2^2 > 2E) = \mathbb{P}(\chi_{m-1} \geq I) = \frac{1}{2^{(m-1)/2} \Gamma((m-1)/2)} \int_I^\infty x^{(m-1)/2-1} e^{-x/2} dx.$$

Making the substitution $y = x/2$, the right hand side equals

$$\frac{1}{\Gamma((m-1)/2)} \int_{I/2}^\infty y^{(m-1)/2-1} e^{-y} dy.$$

If $I \geq 2$, then this is bounded above by

$$\begin{aligned} \frac{1}{\Gamma((m-1)/2)} \int_{I/2}^\infty y^{[(m-1)/2]-1} e^{-y} dy &= \frac{([[(m-1)/2]-1]!}{\Gamma((m-1)/2)} e^{-I/2} \sum_{k=0}^{[(m-1)/2]-1} \frac{(I/2)^k}{k!} \\ &\leq e^{-I/2} (I/2 + 9)^{m/2}. \end{aligned}$$

If $I \leq 2$, then $\mathbb{P}(\chi_{m-1} \geq I) \leq e^{-I/2} (I/2 + 9)^{m/2}$ trivially. Putting everything together and using that $m\zeta = \lambda$ we get that

$$\mathbb{P}_{0,\lambda}(x, y + f(\lambda), f, [t, \lambda]) \leq \exp\left(c\lambda - E - \frac{(y - \lambda^2 - x)^2}{4\lambda} + c\lambda \log(E + 9)\right). \quad (67)$$

Step 2: Estimating the Dirichlet energy E . At this point, the problem is purely deterministic. Let $f_\Pi : [0, \lambda] \rightarrow \mathbb{R}$ be the function given by linearly interpolating between the values of f on the mesh Π . We can then equivalently write E as the minimum value of the Dirichlet energy

$$E(g) = \frac{1}{4} \int_0^\lambda |g'(s)|^2 ds$$

among all functions $g : [0, \lambda] \rightarrow \mathbb{R}$ such that $g \geq f_\Pi$ and $g(0) = x, g(\lambda) = y - \lambda^2$. The Dirichlet energy minimizer g_Π is the smallest concave function with these properties. Since f_Π itself is concave by the concavity of f , g_Π has the following form:

- There are values $a \leq b \in \Pi$ such that g_Π is linear on $[0, a]$ and $[b, \lambda]$.
- $g_\Pi(z) = f(z)$ for all $z \in (a, b)$.

The fact that g_Π dominates f_Π forces $a \leq \sqrt{x}$. Also, the fact that g_Π is concave forces

$$\frac{(-a^2 - x)}{a} \geq \frac{-(a + \zeta)^2 + a^2}{\zeta},$$

which implies that $a > \sqrt{x} - \zeta$. Hence $a = \lfloor \sqrt{x} \rfloor_\Pi$, the largest element of Π that is less than or equal to \sqrt{x} . Similarly $b = \lceil \lambda - \sqrt{y} \rceil_\Pi$, the smallest element of Π that is greater than or equal to $\lambda - \sqrt{y}$. The function g_Π can be thought of as a discretized version of the function g introduced after the statement of Lemma 4.5 whose energy equals $E(x, y, \lambda)$. Moreover, a computation shows that $|E(x, y, \lambda) - E(g_\Pi)| \leq c\lambda$. Finally, since $\sqrt{x} < \lambda/2$, $\sqrt{y} < \lambda/2$, we have that $E(x, y, \lambda) \leq 2\lambda^3$, so (67) is bounded above by

$$\exp\left(c\lambda \log(1 + \lambda) - E(x, y, \lambda) - \frac{(y - \lambda^2 - x)^2}{4\lambda}\right),$$

as desired. \square

Proof of Lemma 4.5.2. First, the bound holds trivially for small λ , so we may assume $\lambda > 1000$. Also, to simplify notation we write $u = x - a$ in the proof. We have

$$\begin{aligned} & \mathbb{P}(B(s) > f(s) \text{ for all } s \in [t, \lambda], B(t) \leq -a) \\ &= \mathbb{P}(B(s) > f(s) \text{ for all } s \in [t, \lambda] \mid B(t) \leq -a) \mathbb{P}(B(t) \leq -a) \\ &\leq \mathbb{P}_{t, \lambda}(u, f(\lambda) + y, f) \mathbb{P}(B(t) \leq -a) \\ &\leq c \mathbb{P}_{t, \lambda}(u, f(\lambda) + y, f) \exp\left(-\frac{a^2}{4t} - \frac{(x - a - y - \lambda^2)^2}{4(\lambda - t)} + \frac{(x - y - \lambda^2)^2}{4\lambda}\right). \end{aligned} \quad (68)$$

Now, since Brownian bridge law commutes with affine shifts,

$$\mathbb{P}_{t, \lambda}(u, f(\lambda) + y, f) = \mathbb{P}_{0, \lambda - t}(u + t^2, f(\lambda - t) + y, f).$$

Bounding this with Lemma 4.5.1, we get that (68) is bounded above by

$$\exp\left(-\frac{a^2}{4t} - E(u + t^2, y, \lambda - t) + \frac{(u + t^2 - y - (\lambda - t)^2)^2}{4(\lambda - t)} - \frac{(u - y - \lambda^2)^2}{4(\lambda - t)} + \frac{(x - y - \lambda^2)^2}{4\lambda} + c\lambda \log(1 + \lambda)\right).$$

Now,

$$\frac{(u + t^2 - y - (\lambda - t)^2)^2}{4(\lambda - t)} - \frac{(u - y - \lambda^2)^2}{4(\lambda - t)} \leq -\lambda^2 t + O(t^3 + tx),$$

where we have bounded the error term by using that $t \leq \sqrt{x} + 1 \leq 2\lambda/3$ and $\sqrt{x}, \sqrt{y} \leq \lambda/2$. We also have that

$$E(x, y, \lambda) - E(u + t^2, y, \lambda - t) \leq \frac{2}{3}x^{3/2} - \frac{2}{3}u^{3/2} + \lambda^2 t + \frac{t^3}{3}$$

which combined with the previous computations gives the desired result. \square

Proof of Lemma 4.5.2. For $x, y > 0$ with $\sqrt{x} < \lambda/2, \sqrt{y} < \lambda/2$, let $g_{x,y}$ be the function defined immediately after Lemma 4.5, and for every i , define $h_i : [0, \lambda] \rightarrow \mathbb{R}$ by

$$h_i(s) = g_{x_i - (k+1-i)/2, y_i - (k+1-i)/2}(s) + (k+1-i)/2.$$

Then $h_i(0) = x_i, h_i(\lambda) = y_i - \lambda^2$. Moreover, since $g_{x',y'} \geq g_{x,y} \geq f$ for any $x' \geq x$ and $y' \geq y$ and

$$x_k, y_k \geq 1, \quad x_i - x_{i+1} \geq 1, \quad y_i - y_{i+1} \leq 1,$$

we have that $h_k - f \geq 1/2$ and $h_i - h_{i+1} \geq 1/2$ for all $i \leq k-1$. Therefore letting B denote a k -tuple of independent Brownian bridges from $(0, \mathbf{x})$ to $(\lambda, \mathbf{y} - \lambda^2)$, we have that

$$\begin{aligned} \mathbb{P}_{0,\lambda}(\mathbf{x}, \mathbf{y} + f(\lambda), f) &\geq \mathbb{P}(\|B_i - h_i\|_\infty < 1/4 \text{ for all } i \in \llbracket 1, k \rrbracket) \\ &= \prod_{i=1}^k \mathbb{P}(\|B_i - h_i\|_\infty < 1/4). \end{aligned} \tag{69}$$

It just remains to bound $\mathbb{P}(\|B_i - h_i\|_\infty < 1/4)$ below for each i . We use the Cameron-Martin theorem for Brownian bridges. Indeed, letting μ_i denote the law of B_i and ν_i denote the law of $B_i + h_i$, the Cameron-Martin theorem yields that

$$\frac{d\mu_i}{d\nu_i}(W) = \exp\left(\frac{1}{4} \int_0^\lambda [h'_i(s)]^2 ds - \frac{1}{2} \int_0^\lambda h'_i dW + \frac{(x_i + \lambda^2 - y_i)^2}{4\lambda}\right).$$

Note that the first term in the exponent above equals $E(x_i - (k+1-i)/2, y_i - (k+1-i)/2, \lambda)$. Now, standard estimates on the maximum absolute value of a Brownian bridge imply that

$$\nu_i(w \in \mathcal{C}([0, \lambda]) : \|w - h_i\|_\infty < 1/4) \geq e^{-c\lambda}.$$

On the other hand, for a Brownian sample path W with $W(0) = h_i(0), W(\lambda) = h_i(\lambda)$, by integration by parts and stochastic integration by parts we have that

$$\begin{aligned} \left| \int_0^\lambda h'_i dW - \int_0^\lambda (h'_i(s))^2 ds \right| &= \left| \int_0^\lambda h''_i(s) W(s) ds - \int_0^\lambda h''_i(s) h_i(s) ds \right| \\ &\leq \int_0^\lambda |h''_i(s)| |W(s) - h_i(s)| ds \end{aligned}$$

This is bounded above by $\lambda/2$ when $\|W - h_i\|_\infty < 1/4$ since $|h''_i| \leq 2$. Therefore $\mathbb{P}(\|B_i - h_i\|_\infty < 1/4)$ is bounded below by

$$\exp\left(-c\lambda - E(x_i - (k+1-i)/2, y_i - (k+1-i)/2, \lambda) + \frac{(x_i + \lambda^2 - y_i)^2}{4\lambda}\right).$$

We can complete the proof by appealing to (69) and recognizing that

$$E(x_i - (k+1-i)/2, y_i - (k+1-i)/2, \lambda) \leq E(x_i, y_i, \lambda) + ck(\sqrt{x} + \sqrt{y}) \leq E(x_i, y_i, \lambda) + ck\lambda. \quad \square$$

References

- [1] Abramowitz, M. and Stegun, I. A. [1972]. *Handbook of mathematical functions: with formulas, graphs, and mathematical tables*, Vol. 55, Dover publications New York.
- [2] Adler, M. and Van Moerbeke, P. [2005]. PDEs for the joint distributions of the Dyson, Airy and sine processes, *Ann. Probab.* **33**(4): 1326–1361.
- [3] Aggarwal, A. [2020]. Arctic boundaries of the ice model on three-bundle domains, *Inventiones mathematicae* **220**(2): 611–671.
- [4] Aggarwal, A. and Huang, J. [2021]. Edge statistics for lozenge tilings of polygons, II: Airy line ensemble, *arXiv preprint arXiv:2108.12874* .
- [5] Basu, R., Ganguly, S. and Zhang, L. [2021]. Temporal correlation in last passage percolation with flat initial condition via brownian comparison, *Communications in Mathematical Physics* **383**: 1805–1888.
- [6] Bates, E., Ganguly, S. and Hammond, A. [2022]. Hausdorff dimensions for shared endpoints of disjoint geodesics in the directed landscape, *Electron. J. Probab.* **27**: 1–44.
- [7] Beffara, V., Chhita, S. and Johansson, K. [2018]. Airy point process at the liquid-gas boundary, *The Annals of Probability* **46**(5): 2973–3013.
- [8] Beffara, V., Chhita, S. and Johansson, K. [2022]. Local geometry of the rough-smooth interface in the two-periodic aztec diamond, *The Annals of Applied Probability* **32**(2): 974–1017.
- [9] Bhatia, M. [2022]. Atypical stars on a directed landscape geodesic, *arXiv preprint arXiv:2211.05734* .
- [10] Borodin, A. and Gorin, V. [2016]. Lectures on integrable probability, *Probability and statistical physics in St. Petersburg* **91**(155-214).
- [11] Calvert, J., Hammond, A. and Hegde, M. [2019]. Brownian structure in the KPZ fixed point, *arXiv preprint arXiv:1912.00992* .
- [12] Colomo, F. and Sportiello, A. [2016]. Arctic curves of the six-vertex model on generic domains: the tangent method, *Journal of Statistical Physics* **164**(6): 1488–1523.
- [13] Corwin, I. [2012]. The Kardar–Parisi–Zhang equation and universality class, *Random matrices: Theory and applications* **1**(01): 1130001.
- [14] Corwin, I. and Hammond, A. [2014]. Brownian Gibbs property for Airy line ensembles, *Invent. Math.* **195**(2): 441–508.
- [15] Corwin, I. and Hammond, A. [2016]. KPZ line ensemble, *Probab. Theory Related Fields* **166**(1-2): 67–185.

- [16] Dauvergne, D., Nica, M. and Virág, B. [2019]. Uniform convergence to the Airy line ensemble, *arXiv:1907.10160*.
- [17] Dauvergne, D., Ortmann, J. and Virág, B. [2018]. The directed landscape, *arXiv preprint arXiv:1812.00309*.
- [18] Dauvergne, D., Sarkar, S. and Virág, B. [2020]. Three-halves variation of geodesics in the directed landscape. *arXiv preprint arXiv:2010.12994*.
- [19] Dauvergne, D. and Virág, B. [2021a]. Bulk properties of the Airy line ensemble, *Ann. Probab.* **49**(4): 1738–1777.
- [20] Dauvergne, D. and Virág, B. [2021b]. The scaling limit of the longest increasing subsequence, *arXiv preprint arXiv:2104.08210*.
- [21] Dauvergne, D. and Zhang, L. [2021]. Disjoint optimizers and the directed landscape, *arXiv preprint arXiv:2102.00954*.
- [22] Ferrari, P. L. and Spohn, H. [2003]. Step fluctuations for a faceted crystal, *Journal of statistical physics* **113**(1): 1–46.
- [23] Ferrari, P. L. and Spohn, H. [2005]. Constrained Brownian motion: fluctuations away from circular and parabolic barriers, *The Annals of Probability* **33**(4): 1302–1325.
- [24] Ferrari, P. L. and Spohn, H. [2010]. Random growth models, *arXiv preprint arXiv:1003.0881*.
- [25] Ganguly, S. and Hammond, A. [2020a]. The geometry of near ground states in gaussian polymer models. *arXiv preprint arXiv:2010.05836*.
- [26] Ganguly, S. and Hammond, A. [2020b]. Stability and chaos in dynamical last passage percolation. *arXiv preprint arXiv:2010.05837*.
- [27] Ganguly, S. and Hegde, M. [2022a]. Fractal structure in the directed landscape, *epprint: <http://math.columbia.edu/~milind>*.
- [28] Ganguly, S. and Hegde, M. [2022b]. Sharp upper tail estimates and limit shapes for the KPZ equation via the tangent method, *arXiv preprint arXiv:2208.08922*.
- [29] Ganguly, S. and Zhang, L. [2022]. Fractal geometry of the space-time difference profile in the directed landscape via construction of geodesic local times, *arXiv preprint arXiv:2204.01674*.
- [30] Hammond, A. [2019]. A patchwork quilt sewn from Brownian fabric: Regularity of polymer weight profiles in Brownian last passage percolation, *Forum of Mathematics, Pi* **7**.
- [31] Hammond, A. [2020]. Exponents governing the rarity of disjoint polymers in Brownian last passage percolation, *Proceedings of the London Mathematical Society* **120**(3): 370–433.

- [32] Hammond, A. [2022]. Brownian regularity for the Airy line ensemble, and multi-polymer watermelons in Brownian last passage percolation, *Mem. Amer. Math. Soc.* **277**: 1363.
- [33] Hammond, A. and Sarkar, S. [2018]. Modulus of continuity for polymer fluctuations and weight profiles in Poissonian last passage percolation, arXiv:1804.07843.
- [34] Johansson, K. [2003]. Discrete polynuclear growth and determinantal processes, *Communications in Mathematical Physics* **242**(1-2): 277–329.
- [35] Johansson, K. [2005]. The arctic circle boundary and the Airy process, *Ann. Probab.* **33**(1): 1–30.
- [36] Okounkov, A. and Reshetikhin, N. [2007]. Random skew plane partitions and the Pearcey process, *Communications in mathematical physics* **269**(3): 571–609.
- [37] Petrov, L. [2014]. Asymptotics of random lozenge tilings via gelfand–tsetlin schemes, *Probability theory and related fields* **160**(3): 429–487.
- [38] Prähofer, M. and Spohn, H. [2002]. Scale invariance of the PNG droplet and the Airy process, *J. Stat. Phys.* **108**(5-6): 1071–1106.
- [39] Quastel, J. [2011]. Introduction to KPZ, *Current Developments in Mathematics* **2011**(1).
- [40] Quastel, J. and Sarkar, S. [2023]. Convergence of exclusion processes and the KPZ equation to the KPZ fixed point, *Journal of the American Mathematical Society* **36**(1): 251–289.
- [41] Romik, D. [2015]. *The surprising mathematics of longest increasing subsequences*, Vol. 4, Cambridge University Press.
- [42] Tracy, C. A. and Widom, H. [1994]. Level-spacing distributions and the Airy kernel, *Communications in Mathematical Physics* **159**(1): 151–174.
- [43] Virág, B. [2020]. The heat and the landscape I, *arXiv preprint arXiv:2008.07241* .
- [44] Wu, X. [2021]. Tightness and local fluctuation estimates for the KPZ line ensemble, *arXiv preprint arXiv:2106.08051* .