

Limits of graded Gorenstein algebras of Hilbert function $(1, 3^k, 1)$

Nancy Abdallah

University of Borås, Borås, Sweden.

Jacques Emsalem*

Paris, France.

Anthony Iarrobino

Department of Mathematics, Northeastern University, Boston, MA 02115, USA.

J. Yaméogo

Université Côte d'Azur, CNRS, LJAD, France

February 10, 2023

Abstract

Let $R = k[x, y, z]$, the polynomial ring over a field k . Several of the authors previously classified nets of ternary conics and their specializations over an algebraically closed field [AEI]. Building on this work, we here show that when k is algebraically closed, and the

*Our friend and colleague Jacques Emsalem passed on 12 July, 2022, after an illness; he had participated actively and enthusiastically in our work.

Keywords: Artinian Gorenstein algebra, closure, deformation, Hilbert function, irreducible component, isomorphism class, limits, nets of conics, normal form, parametrization.

2021 Mathematics Subject Classification: Primary: 13E10; Secondary: 14A05.

Hilbert function sequence $T = (1, 3^k, 1), k \geq 2$, then the family G_T parametrizing graded Artinian algebra quotients $A = R/I$ of R having Hilbert function T is irreducible, and G_T is the closure of the family $\text{Gor}(T)$ of Artinian Gorenstein algebras of Hilbert function T . We then classify up to isomorphism the elements of these families $\text{Gor}(T)$ and of G_T . Finally, we give examples of codimension three Gorenstein sequences, such as $(1, 3, 5, 3, 1)$, for which G_T has several irreducible components, one being the Zariski closure of $\text{Gor}(T)$.

1 Introduction.

Let $R = \mathbb{k}[x_1, \dots, x_r]$ be a standard-graded polynomial ring over a field \mathbb{k} , and denote by $S = \mathbb{k}_{DP}[X_1, \dots, X_r]$ the corresponding ring of divided powers, upon which R acts by contraction: that is

$$x_i^k \circ X_j^u = \delta_{i,j} X^{u-k} \text{ if } u \geq k \text{ or } 0 \text{ if } u < k \quad (1)$$

We have $R = \bigoplus_{i=0}^{\infty} R_i$ the sum of its homogeneous degree- i components.

Let T be a *Gorenstein sequence*, a sequence that may occur as the Hilbert function $H(A)$ of a graded Artinian Gorenstein quotient $A = R/I$. The family of graded Gorenstein algebras $\text{Gor}(T)$ of Hilbert function T is an open subfamily of G_T , the family of all graded algebra quotients of R having Hilbert function T .

For codimension $r = 2$ the Gorenstein algebras A are complete intersections. Also, in codimension 2 for \mathbb{k} algebraically closed, we have that $\text{Gor}(T)$ is irreducible of known dimension and is open dense in G_T [I1]. For codimension $r = 3$, the Gorenstein sequences T are known as a consequence of the D. Buchsbaum-D. Eisenbud Pfaffian resolution theorem [BuEi, St]; S.J. Diesel showed also for $r = 3$ that over an algebraically closed field the variety $\text{Gor}(T)$ is irreducible; and she determined a poset of minimal resolution strata, and their associated dimensions [Di]. A consequence for $T = (1, 3, 3, 1)$ is that $\text{Gor}(T) = \overline{CI}_T$ the closure of the family of complete intersections of Hilbert function T . For $T = (1, 3^k, 1), k \geq 3$ every $A \in \text{Gor}(T)$ has five generators. We consider the

Question. Given T a Gorenstein sequence of codimension three, is $\text{Gor}(T)$ open dense in G_T ?

We will show that for $T = (1, 3^k, 1)$ the answer is “Yes” (Theorem 1) but in general is “No” (Theorem 3).

For codimension $r \geq 4$, the family $\text{Gor}(T)$ itself may have several irreducible components [IaSr].

A *net of conics* in the projective plane \mathbb{P}^2 is a 3-dimensional vector space V of conics, elements of R_2 for $r = 3$. These nets are parametrized by the Grassmanian $\text{Grass}(R_2, 3)$. The projective linear group $\text{PGL}(3)$ acts on $R_1 = \langle x, y, z \rangle$, hence on $R = \mathbb{k}[x, y, z]$. In [AEI] the first three authors determined the isomorphism classes under $\text{PGL}(3)$ of nets of conics over an algebraically closed field \mathbb{k} not of characteristic 2 or 3, and as well their specializations (see our Figure 1 below, based on [AEI, Tables 1,5]).¹ We will assume henceforth throughout this paper that \mathbb{k} is an algebraically closed field of arbitrary characteristic (unless we otherwise specify).

We will denote by $\text{Gor}_{\#6a}(T)$ the family of Gorenstein algebras $A = R/I$ of Hilbert function T , such that the net $V = I_2$ is in the isomorphism class #6a of Table 1, that is $I_2 \cong \langle xy, xz, yz \rangle$ under $\text{PGL}(3)$ -action. Equivalently, $\text{Gor}_{\#6a}(T)$ parametrizes those $A = R/I$ for which $I \cong \text{Ann}F, F = X^j + \alpha_1 Y^j + \alpha_2 Z^j, \alpha_1, \alpha_2 \in \mathbb{k}$.²

We recall from [AEI, Section 6.3], the *scheme length* $\ell(V)$ of a net of conics V is the length of the scheme $\text{Proj}(R/(V))$; this is just the integer a for which the Hilbert function

$$H(R/(V)) = (1, 3, 3, \dots, \bar{a})$$

has an infinite tail $\bar{a} = (a, a, \dots, a, \dots)$ of a 's. The scheme \mathfrak{Z}_V determined by V is empty if $\ell(V) = 0$; it is a finite set of $\ell(V)$ points if $\ell(V) \neq 0$; and is a line in the case $V \cong \langle x^2, xy, xz \rangle$ (#2a in Table 1).

Our first main result is

Theorem 1. Let $T = (1, 3^k, 1), k \geq 2$. For $k = 2$, the family G_T of graded quotients of $R = \mathbb{k}[x, y, z]$ having Hilbert function T , is the Zariski closure of the family CI_T of graded complete intersection quotients of R having Hilbert function T ; and G_T has dimension 9.

¹These classes had been determined also by C.T.C. Wall - who considered as fields both the complexes and the reals, and also gave an account of specializations [Wall]; A. Conca gives further information about the nets [Co]. N. Onda - apparently unaware of previous results - gave recently an independent determination of the isomorphism classes over an algebraically closed field of characteristic not 2 or 3, without discussion of specialization [Onda]. See [AEI, Appendices A,B] for some historical account, and discussion of related problems, in particular [AEI, Table 10, Appendix B] for a concordance of the results of [Co, Wall, AEI] classifying nets of conics.

²When \mathbb{k} is closed under j -th roots, we may assume that $F \cong X^j + Y^j + Z^j$.

When $k \geq 3$ then G_T is in the closure of $\text{Gor}_{\#6a}(T)$, and G_T has dimension eight.

We show this in Theorem 3.3 (scheme length of $V = I_2$ is three), Theorems 3.6 (case $k = 2$) and 3.8 (case $k \geq 3$).

In our second main result we show that the pattern of specializations of algebras $A = R/I$ having Hilbert function T corresponds to that for the corresponding nets of conics $V = I_2$, provided we stay within constant scheme-length families of conics.

Theorem 2. Assume that the net $V \in \text{Grass}(R_2, 3)$ is in the Zariski closure of nets isomorphic to V' , and that the scheme length $\ell(V) = \ell(V')$ and that V, V' is not $\#2a$. Then any algebra $A = R/I$ of Hilbert function $T = (1, 3^k, 1)$ such that $I_2 \cong V$ is in the closure of a family $A(x) = R/I(x)$ of algebras having $I_2(x) \cong V'$, that is $A = \lim_{x \rightarrow x_0} A(x), x \in X \setminus x_0$. We may take X to be an irreducible variety or curve.

We show Theorem 2 in Theorem 3.3 (case scheme length of V is three - including the case $k \geq 3$), and in Theorem 3.4 (case $k = 2$). Remark 3.10 shows that an analogous statement to Theorem 2 does not hold when the scheme lengths of V, V' are not the same.

We show in Theorem 3.16,

Theorem 3 (Smallest G_T not in $\overline{\text{Gor}(T)}$). Let $5 \leq a \leq 6$ and $T = (1, 3, a, 3, 1)$, or let $T = (1, 3, 6, 6, 3, 1)$. Then G_T has two irreducible components, $\overline{\text{Gor}(T)}$, and a second, larger component U_T (Definition 3.13), whose dimensions are given in Figure 3.

In Section 4 we study G_T : we classify graded algebras of Hilbert function $T = (1, 3^k, 1)$ up to isomorphism, over an algebraically closed field k . We discuss briefly their specializations. Given a net $V \subset R_2$ we denote by $\text{Gor}_V(T)$ the family of Gorenstein algebra quotients $A = R/I$ with $I_2 = V$, of Hilbert function T ; and by $G_V(T)$ the family of Artinian algebra quotients $A = R/I$ with $I_2 = V$, of Hilbert function T . We denote by $\text{Gor}_{\{V\}}(T)$ and $G_{\{V\}}(T)$ the corresponding families where $I_2 \cong V$ under $\text{PGL}(3)$ action.

In Section 4, we show first the following result (Theorem 4.1).

Theorem 4 (Dimension theorem). Let $T = (1, 3^k, 1), k \geq 2$. Then $G_V(T)$ and $G_{\{V\}}(T)$ are irreducible. We let $d = d(V)$ be the dimension of the family of nets of conics that are $\text{PGL}(3)$ equivalent to V .

- (i). Let $T = (1, 3, 3, 1)$. Then the family of Artinian algebras $G_{\{V\}}(T)$ has dimension d if $\ell = 0$, dimension $d + \ell - 1$ if $1 \leq \ell \leq 3$; and has dimension 5 if $\ell = \infty$ (the case $V = \#2a$).
- (ii). Let $T = (1, 3^k, 1)$ with $k \geq 3$. Then $G_{\{V\}}(T)$ has dimension $d+2$ if $\ell = 3$ and dimension 7 if $\ell = \infty$ (case $V = \#2a$), and is empty otherwise.

We then show our main classification result for Artinian quotients of R . Recall that the nets of scheme length zero determine CI algebras: these are the nets of $\#8b$, and the three nets $\#8c, \#7c$, and $\#6d$.

Theorem 5. Let $T = (1, 3^k, 1)$. We determine the isomorphism classes of algebras in $G_V(T)$, the family of Artinian quotients $A = R/I$ in G_T for which $I_2 = V$.

- (i). The number of classes for V of scheme lengths $\ell = 2$ and 3 satisfy

	Cases $\ell = 2$.		Cases $\ell = 3$.					
Class of I_2	$\#7a$	$\#6b$	$\#6c$	$\#5b$	$\#6a$	$\#5$	$\#4$	$\#2b$
# Isom classes	6	∞	3	3	3	7	3	3.

There are among the nets V of scheme length 3, exactly four isomorphism classes of Gorenstein algebras in $G_V(T)$, namely $\#6a.i, \#5a.i, \#5a.ii$, and $\#4.iii.$, (see the Betti C table in Figure 8). There are no further non-CI Gorenstein classes of algebras in G_T .

- (ii). Classes for the $V = xR_1$ ($\#2a$) of scheme length ∞ :
When $T = (1, 3, 3, 1)$: there are 3 classes;

$$T = (1, 3^k, 1), k \geq 3: I = (V, f, f_5, f_6), f \in I_3; f_5, f_6 \in I_j.$$

$$f = yz(y - z) : \infty^2, \infty; f = yz^2: 5 \text{ classes}; f = z^3: 4 \text{ classes.}$$

Note. Here ∞^2 means a two-parameter family of classes, see Corollary 6. Theorem 5 summarizes the results of Theorems 4.6, 4.8 and 4.11 for V of scheme length two, three and ∞ , respectively. We also specify the Betti tables for the classes for V of scheme lengths two and three.

Recall that the family of nets having the form $\#8b$ in Table 1 is a continuous family of complete intersections, whose isomorphism classes are determined by their j -invariant $j(\lambda)$ (see [AEI, Footnote 9]).

Question. Fix any net $V \subset R_3$. Do we have continuous families of Artinian algebras $A = R/I$ in G_T with $I_2 = V$?

We answer this:

Corollary 6. Aside of the complete intersection class of nets #8b, there are three continuous families of isomorphism classes in these G_T . First, when $T = (1, 3, 3, 1)$ and $V \cong \langle x^2, xy, yz + z^2 \rangle$ (#6b), Second, when $T = (1, 3^k, 1), k \geq 3$ and $V \cong xR_1$ (#2a) and $f = yz(y - z)$ we have the families $I(a, b), I(a)$ with $I(a, b) = I(a^{-1}, a^{-1}b)$ if $a \neq 0$ (Equations (39), (40),(42)). For all other nets V there are only a finite number of non-isomorphic Artin algebras having $I_2 = V$.

Proof. See Conclusion 4.7 for # 6b and Theorem 4.11 for #2a. \square

In Section 2 we recall some definitions and results about nets of conics. Section 3 is devoted to the deformations of graded algebras to graded Gorenstein algebras; Section 4 is devoted to the classification up to isomorphism of graded algebra quotients of R having Hilbert function $T = (1, 3^k, 1), k \geq 2$. In Section 4.5.1 we consider some questions about limits of Betti tables.

Contents

1	Introduction.	2
2	Preliminaries.	7
2.1	Nets of conics.	7
2.1.1	Hilbert functions $H(R/(V))$ for V a net of conics in \mathbb{P}^2	8
3	Deformation of algebras in G_T to Gorenstein algebras.	12
3.0.1	The families $\text{Gor}_V(T), \text{Gor}_V(T)$ and $G_V(T), G_{\{V\}}(T)$.	13
3.1	When $A = R/I$ and the scheme length of $V = I_2$ is three.	13
3.2	The case $T = (1, 3, 3, 1)$, G_T is in the closure of CI_T .	14
3.3	Explicit deformations to AG algebras, $T = (1, 3, 3, 1)$.	17
3.3.1	Scheme length of $V = I_2$ is two, $T = (1, 3, 3, 1)$.	17
3.3.2	Scheme length of $V = I_2$ is three, $T = (1, 3, 3, 1)$.	20
3.4	The case $T = (1, 3^k, 1), k \geq 3$: G_T is in the closure of $\text{Gor}_{\#6a}(T)$.	21
3.4.1	Scheme \mathfrak{Z}_V a line, $\ell(V) = \infty$.	22
3.5	Graded algebras G_T in codimension three not in the closure of $\text{Gor}(T)$.	23

4	Isomorphism classes of graded Artinian algebras of Hilbert function $T = (1, 3^k, 1)$.	28
4.1	Parametrizing Artinian algebras in $G_{\{V\}}(T)$	29
4.1.1	Betti tables and cancellation.	31
4.2	Classification of isomorphism classes of Artinian algebras of Hilbert function T by the net of conics $V = I_2$: the family $G_V(T)$	32
4.2.1	Classification of algebras in $\text{Gor}(T)$, $T = (1, 3^k, 1)$	33
4.2.2	Isomorphism classes in $G_{\{V\}}T$ when $V = I_2$ defines a length 0 or 1 scheme.	34
4.2.3	Isomorphism classes in $G_{\{V\}}T$ when $V = I_2$ defines a length two scheme.	34
4.2.4	Betti diagrams for $A \in G_T$, $T = (1, 3, 3, 1)$	37
4.3	Isomorphism classes in G_T when I_2 defines a length 3 scheme, $T = (1, 3^k, 1)$	40
4.4	Isomorphism classes in G_T , $k \geq 2$ when $V = I_2$ defines a line (case #2a).	45
4.5	Questions and Problems	50
4.5.1	Limits of Betti tables in $\text{Gor}(T)$ and G_T	50
4.5.2	Specialization of G_T isomorphism classes.	52
4.5.3	Possible future directions involving Jordan type or applications.	53
	List of Figures	53
	References	54

2 Preliminaries.

2.1 Nets of conics.

We adapt these notions from [AEI]. Figure 1 below combines information from [AEI, Tables 1,5]. Recall that $\text{PGL}(3)$ acts on the \mathbf{k} -vector space $\langle x, y, z \rangle$, hence on $R = \mathbf{k}[x, y, z]$.

Definition 2.1. Let $V \subset R_2$ be a subvector space of \mathbf{k} -dimension three. The *scheme length* $\ell(V)$ of V is the length of the scheme $\mathfrak{Z}_V = \text{Proj}(R/(V))$, or zero if $R/(V)$ is a complete intersection (CI), or ∞ if $V \cong xR_1$, in which case the associated scheme \mathfrak{Z}_V is a line.

The *associated cubic* $\Gamma(V)$ of the net of conics $V = \langle v_1, v_2, v_3 \rangle$ is the cubic $f(a_1, a_2, a_3)$ such that $f = 0$ is the locus of elements

$a_1v_1 + a_2v_2 + a_3v_3$ that are decomposable - that is - may be factored into a product of linear forms. Here $a_1, a_2, a_3 \in R_1$ - are linear.

The *double line* subset $D_2(V) \subset \Gamma(V)$ denotes the set of perfect squares, that is the elements of V having the form L^2 for some $L \in R_1$.

Organization of Figure 1:

In referencing the isomorphism classes of nets in Figure 1, we will use #7a, #7b, #7c to denote respectively the nets of dimension 7, left-most, middle, and right - so #7a is the net $\langle x^2 + yz, xy, xz \rangle$, or, used loosely, any net that is $\text{PGL}(3)$ -equivalent to it.³

In Figure 1 we give for each $\text{PGL}(3)$ orbit of nets of conics

- i. a basis for a net belonging to it,
- ii. a diagram indicating the isomorphism type of its associated cubic $\Gamma(V)$, where each point of $\Gamma(V) \cap D_2$ is marked by a circle around it,
- iii. the scheme length $\ell(V)$ for V in the orbit, as an italicized integer $0, 1, 2$ or 3 to the right of the diagram, or as ∞ for the net V of #2a.
- iv. the dimension of the orbit is indicated by the integer in the left column.

The arrows oriented downward $V \rightarrow V'$ indicate that the orbit of V' is included in the closure of the orbit of V . The arrows downward from the one-parameter family $V_j \rightarrow V'$ indicate that the orbit of V' is in the closure of each orbit V_j , with j , the invariant of the cubic, held constant.

2.1.1 Hilbert functions $H(R/(V))$ for V a net of conics in \mathbb{P}^2 .

The Lemma 2.6 below is a consequence of the G. Gotzmann theorems on the Hilbert scheme - see [Gotz] and the exposition in [IaKa, Appendix C]. We explain the concept briefly.

³The nets in #8b correspond to $J\phi$ where ϕ is a smooth cubic not isomorphic to the harmonic cubic $x^3 + y^3 + z^3$. Here ω is a primitive cube root of one. The omitted values $-1, -\omega, -\omega^2$ correspond to singular cubics ϕ giving rise to the net in #6a. The omitted values $0, 2, 2\omega, 2\omega^2$ give a cubic isomorphic to the harmonic cubic. For further details see [AEI].




Figure 1: Specializations of nets of conics and scheme length.

Lemma 2.2. [Macaulay inequality] Let $V \subset R_j$ be a d -dimensional vector space of forms in $R = \mathbb{k}[x_1, \dots, x_r]$. Let $V_0(r, j, d)$ be the span of the first d monomials of R_j in lexicographic order, and let $f(r, j, d) = \dim_{\mathbb{k}} R_1 V_0(r, j, d)$.

$$V \subset R_j, \text{ the } \dim_{\mathbb{k}} R_1 V \geq f(r, j, d). \quad (2)$$

To state the Gotzmann theorem, in the form that we will use to delimit the possible Hilbert functions of $R/(V)$ and of related algebras, we need some definitions. Recall that we denote by \bar{a} the infinite sequence (a, a, \dots) of a 's.

The *Castelnuovo-Mumford regularity degree* of a finitely-generated R -module M is the smallest degree m such the M is m -regular; M is m -regular if $\text{Ext}^j(M, R)_n = 0$ for all j and all $n \leq -m - j - 1$ ([Ei, Proposition 20.16]). These definitions become simpler when A is Artinian or has dimension one.

Definition 2.3. The *regularity degree* of an Artinian algebra A is the socle degree of A . A graded algebra A has dimension one when $H(A) = (1, \dots, s_u, s, \bar{s})$: here, the regularity degree of A is the smallest degree u for which $H(A)_u = s$. More generally the

The *Hilbert polynomial* $\mathfrak{h}_A(t)$ of an Artinian algebra is its length $|A|$; that of a dimension one algebra is s above.

The *Gotzmann regularity degree* of a polynomial $h(t)$ is the smallest degree $\tau(h)$ such that all algebras of Hilbert polynomial $h(t)$ are regular in that degree.

We now define the ancestor ideal \bar{V} of a vector space of forms $V \subset R_j$.

Definition 2.4. [I3] Let $R = \mathbb{k}[x_1, \dots, x_r]$ and $V \subset R_j$ be a subvector space. Then the *ancestor ideal* \bar{V} of V in R satisfies

$$\begin{aligned} \bar{V} &= \bigoplus_{i=1}^j \bar{V}_i + (V) \text{ where for } 1 \leq i \leq j \\ \bar{V}_i &= V : R_{j-i} = \{h \in R_i \mid R_{j-i}h \subset V\}. \end{aligned} \quad (3)$$

The Gotzmann theorem ([Gotz],[IaKa, Appendix C]) states

Proposition 2.5.

(i). *Persistence:* When $V \subset R_j$ satisfies $\dim_{\mathbb{k}} R_1 V = f(r, j, d)$ then $\dim R_2 V$ satisfies

$$\dim R_2 V = f(r, j, d_1), \text{ where } d_1 = f(r, j, d).$$

(ii). Scheme \mathfrak{Z}_V : The ancestor ideal $J = \overline{V}$ (Definition 2.4) - which agrees with the ideal (V) in degree at least j - defines a subscheme $\mathfrak{Z}_V = \text{Proj}(R/J) \subset \mathbb{P}^{r-1}$ whose Hilbert polynomial p_3 is that giving $\dim_{\mathbb{k}} R_i/R_{i-j}V$ for i at least the Gotzmann regularity degree $\mathfrak{r}(p_3)$ of p_3 .

For the next Lemma see [IaKa, Proposition C.32].

Lemma 2.6 (The Gotzmann regularity degree of the constant Hilbert polynomial). *Let $R = \mathbb{k}[x_1, \dots, x_r]$ be a polynomial ring, let s be a positive integer. Assume that the ideal $I \subset R$, satisfies*

$$H(R/I)_i = H(R/I)_{i+1} = s \text{ for some integer } i \geq s.$$

Then $H(R/(I_i))_k = s$ for $k \geq i$, where (I_i) denotes the ideal generated by I_i .

Equivalently, the Gotzmann regularity degree of the constant Hilbert polynomial $p_3 = s$ is s .

This result assures that the sequence $H(R/(V)) = (1, 3, 3, 2, 2, \overline{1})$ cannot occur as the Hilbert function for $A = R/(V)$ for a net V of ternary conics, as $(V)_3$ would be defined by a length 2 scheme so $(V)_{\geq 3} = (J_3)_{\geq 3}$ and we would have $H(R/(V)) = (1, 3, 3, \overline{2})$.

We next recall the following semicontinuity result.

Lemma 2.7. *Let $V(w), w \in \mathcal{W}$ be a family of constant dimension d subspaces of R_j . Then the condition $\dim_{\mathbb{k}} R_1V > k$ is an open condition on the family $V(w)$. Also, the Hilbert function of $H(R/(V))$ is semicontinuous.*

Proof. That $\dim_{\mathbb{k}} R_1V > k$ is the condition that the $rd \times r_{j+1}$ matrix defining the span of $R_1V = \langle x_1V, \dots, x_rV \rangle$ in R_{j+1} have rank greater than k , an open condition. The same can be said for $R_iV(w)$, so $H(R/(V(w))), w \in \mathcal{W}$ is semicontinuous. \square

The following result is well-known.

Lemma 2.8. *A complete intersection of generator degrees $(2, 2, 2)$ has Hilbert function $(1, 3, 3, 1, 0)$. No ideal $I = (V)$ generated by a net of conics can have a different Hilbert function ending in zero.*

Thus, for example, $H(R/(V)) = (1, 3, 3, 2, 1, 0)$ cannot occur.

Lemma 2.9 (Hilbert functions possible for a net of ternary cubics). *Let V be a net of conics in $R = \mathbb{k}[x, y, z]$. Then the Hilbert function $H(R/(V))$ determined by the ideal generated by V must be one of the following:*

$$(1, 3, 3, 1, 0), (1, 3, 3, \bar{1}), (1, 3, 3, \bar{2}), (1, 3, 3, \bar{3}) \text{ or } (1, 3, 3, 4, 5, \dots), \quad (4)$$

the last only occurring for a net isomorphic to $V = xR_1 = \langle x^2, xy, xz \rangle$ (#2a). The sequences $(1, 3, 3, 2, \bar{1}), (1, 3, 3, 3, 2, \bar{1}), (1, 3, 3, 3, \bar{2})$ do not occur as Hilbert functions $H(R/(V))$. The sequence ending in $\bar{1}$ occurs only for the $\text{PGL}(3)$ isomorphism classes #8a, #7b in Figure 1; that ending in $\bar{2}$ occurs only for #7a, 6b, 6c, 5d in Figure 1; and that ending in $\bar{3}$ occurs only for the isomorphism classes #6a, 5a, 4, 2b.

Proof. We can eliminate $H(R/(V)) = (1, 3, 3, a, \bar{1}), a > 1$, by considering Figure 1 #7b, the most special with tail $\bar{1}$ whose Hilbert function is $H(R/(V)) = (1, 3, 3, \bar{1})$. Any family specializing to it must have $a = 1$, as $\dim R_1V = 10 - H(R/(V))_3$ can only decrease under specialization, as a consequence of Lemma 2.7.

We may eliminate $(1, 3, 3, 3, \bar{2})$ by considering #5b of Figure 1, $V = (y^2, z^2, xy)$, which is the most special for which $H(R/(V))$ ends in $\bar{2}$, so it will have the termwise highest Hilbert function: and here $H(R/(V))_3 = 2$, not 3.

The last statement can be read from the scheme lengths in Figure 1. □

Question. Do we have all but the last statement of Lemma 2.9 over an arbitrary field? The Table 1 requires $\text{char } \mathbb{k} \neq 2, 3$ and k algebraically closed (see remark before Theorem A1 in [AEI]) but the G. Gotzmann theorem does not have a restriction on $\text{char } \mathbb{k}$, and holds more generally for B -flat graded modules over a ring $B[x_1, \dots, x_r]$ (see [IaKa, Theorem C.17 and C.29]).

3 Deformation of algebras in G_T to Gorenstein algebras.

We recall our Theorem 1:

Theorem 1. Let $T = (1, 3^k, 1), k \geq 2$. For $k = 2$, the family G_T of graded quotients of $R = \mathbb{k}[x, y, z]$ having Hilbert function T , is

the Zariski closure of the family CI_T of graded complete intersection quotients of R having Hilbert function T ; and G_T has dimension 9.

When $k \geq 3$ then G_T is in the closure of $\text{Gor}_{\#6a}(T)$, and G_T has dimension eight.

We show this in Theorem 3.3 (scheme length of $V = I_2$ is three), Theorem 3.6 (case $k = 2$) and Theorem 3.8 (case $k \geq 3$).

3.0.1 The families $\text{Gor}_V(T)$, $\text{Gor}_V(T)$ and $G_V(T)$, $G_{\{V\}}(T)$.

Definition 3.1. Let V be a net of planar conics. We denote by $\text{Gor}_V(T)$ the family of Artinian Gorenstein quotients $A = R/I$ of R having Hilbert function $H(A) = T$, and satisfying $I_2 = V$. We denote by $\text{Gor}_{\{V\}}(T)$ the family of Artinian Gorenstein quotients $A = R/I$ of R for which $I_2 \cong V$ - is equivalent to V under $\text{PGL}(3)$ isomorphism. We define $G_V(T)$ and $G_{\{V\}}(T)$ similarly.

Lemma 3.2. *Let k be algebraically closed. For $T = (1, 3^k, 1)$ and V a net of conics. Then $\text{Gor}_V(T)$, $\text{Gor}_{\{V\}}(T)$, $G_V(T)$ and $G_{\{V\}}(T)$ are irreducible varieties.*

Proof. The irreducibility of $G_V(T)$ and $G_{\{V\}}(T)$ is shown in Theorem 4.1, page 29 below: the result depends on the irreducibility of each of the families of nets in Figure 1 and as well a review of the fibre of $G_V(T)$ over V . The irreducibility of the $\text{Gor}_V(T)$ follows from the classification of Gorenstein algebras $\text{Gor}(T)$ in Theorem 5. \square

3.1 When $A = R/I$ and the scheme length of $V = I_2$ is three.

We here show our

Theorem 2 Assume that the net $V \in \text{Grass}(R_2, 3)$ is in the Zariski closure of nets isomorphic to V' , and that the scheme length $\ell(V) = \ell(V')$ and that V, V' is not $\#2a$. Then any algebra $A = R/I$ of Hilbert function $T = (1, 3^k, 1)$ such that $I_2 \cong V$ is in the closure $A = \lim_{x \rightarrow x_0} A(x)$, $x \in X$ of a family $A(x) = R/I(x)$ of algebras having $I_2(x) \cong V'$. We may take X to be an irreducible variety or curve.

We show Theorem 2 in Theorem 3.3 (case scheme length of V is three - including the case $k \geq 3$), and in Theorem 3.4 (case $k = 2$).

Our proof of the following result uses the classification of nets of conics of [AEI].

Theorem 3.3. *Let $T = (1, 3^k, 1)$, $k \geq 2$ and fix a net of conics V of scheme length three, and assume the algebra $A = R/I$ satisfies $A \in G_T$, and $A = R/I$ where $I_2 = V$.*

(i). *Assume further V is not isomorphic to the class #2b of Table 1. Then $A \in \overline{\text{Gor}_V(T)}$.*

(ii). *Assume V is isomorphic to #2b. Then $\text{Gor}_V(T)$ is empty but $A \in \overline{\text{Gor}_W(T)}$ where W is the class of #4 in Table 1.*

Proof of (i). The Table 2 below gives the dual vector spaces $((V)_j)^\perp$ for V of scheme length three. A check shows that a generic F in each except #2b, satisfies $\dim_{\mathbb{k}} R_1 \circ F = 3$. We have in general for a vector space $K \subset R_{j-1}$ that $R_1 \circ \langle R_1 K \rangle^\perp \subset K^\perp \subset S_{j-1}$: thus we have for these V that $R_1 \circ F = ((V)_{j-1})^\vee$. It is easy to see then that $\dim_{\mathbb{k}} R^u \circ F = 3$ for $1 \leq u \leq j - 2$, hence that $A_F = R/\text{Ann}F$ is Gorenstein of Hilbert function T .

Now let the algebra $A = R/I$ of Hilbert function T satisfy $I_2 = V$ of scheme length 3, and V not $\text{PGL}(3)$ -congruent to #2b. Then

$$I_k = (V)_k \text{ for } 2 \leq k \leq j - 1 \quad (5)$$

as $(V)_k \subset I_k$ and $\text{cod}I_k = \text{cod}(V)_k = 3$ in R_k .

Also, $I_j = \text{Ann}G$ for some $G \in ((V)_j)^\perp$: since $G = \lim_{t \rightarrow t_0} F(t)$ where $F(t)$ is generic for $t \neq t_0$, we have $A_G = R/I_G = \lim_{t \rightarrow t_0} A_{F(t)}$, the limit of elements in $\text{Gor}_V(T)$.

The proof of (ii) is similar, we use that for $A = R/I \in G_{\{V\}}(T)$, V isomorphic to #2b, that $I_j = \text{Ann}G$, $G = \lim_{t \rightarrow t_0} F(t)$ where $F(t) \in (W_j)^\perp$ - using the specialization from Table 1 (letting $X \rightarrow 0$ in $W_{\#4}$ of Table 2). And $A_{F(t)} \in \text{Gor}_W(T)$. \square

Recall that the nets V of scheme length 3 are #6a, #5a, #4, and #2b, from Table 1; their perpendicular spaces for $j \geq 2$ are given in Figure 2 below, where $W_{\#4} = \langle (X^2 - YZ)Z^{j-2}, XZ^{j-1}, Z^j \rangle$.

3.2 The case $T = (1, 3, 3, 1)$, G_T is in the closure of CI_T .

Fix a net V of conics of $R = \mathbb{k}[x, y, z]$, fix $T = (1, 3, 3, 1)$. Let $A \in \text{Gor}_V(T)$; When $\ell = 2$, we have by Lemma 2.9 that

#	6a	5a	4	2b
V	xy, xz, yz	xy, xz, z^2	$x^2 + yz, xy, y^2$	$x^2, y^2, (x+y)^2$
$((V)_j)^\perp$	X^j, Y^j, Z^j	$X^j, Y^j, Y^{j-1}Z$	$W_{\#4}$	XZ^{j-1}, YZ^{j-1}, Z^j

Figure 2: Nets of scheme length three and their duals.

$H(R/(V))_3 = 2$, therefore there exists a non zero $f \in R_3/R_1V$ such that $I = (V, f, \mathfrak{m}^4)$. When $\ell = 3$, again by Lemma 2.9, we have $H(R/(V))_3 = 3$ and there exist non zero polynomials $f, g \in R_3/R_1V$ such that $I = (V, f, g, \mathfrak{m}^4)$. The following Theorem shows that the specialization of such algebras follows the specialization pattern of their corresponding nets, as in Table 1.

We take two approaches here to parametrization. Often w is an informal parameter in the field \mathbf{k} . Or, more precisely, we may take w to be a new variable and write $A(w) = R[w]/I(w)$ for an element of the family $A_{\mathcal{W}} \rightarrow \mathcal{W}$, with $w \in \mathcal{W} \cong \mathbb{A}^1$.

Theorem 3.4. *Let $T = (1, 3, 3, 1)$ and let $V_w, w \in \mathcal{W}$ be a family of nets of conics of finite scheme length ℓ and let $V = V_{w_0}$ be a specific net. Assume further that for $w \neq w_0, V_w$ is isomorphic to a fixed net V' . Now let the algebra $A(w_0) = R/I(w_0) \in G_T$ satisfy $I(w_0)_2 = V_{w_0}$. Then $A(w_0)$ is a specialization of a family $A(w) = R/I(w)$ of Artinian algebras, also of Hilbert function T , for which $I(w)_2$ for $w \neq w_0$ is isomorphic to V' .*

Proof. For $\ell(V) = 0$, then $A = R/(V)$ so specializing the algebra $A(w)$ is the same as specializing the corresponding net $V_w = I_2$.

For $\ell(V) = 1$, $A = R/(I_2, \mathfrak{m}^4)$ is simply obtained by quotienting $R/(I_2)$ of Hilbert function $(1, 3, 3, \bar{1})$ by \mathfrak{m}^4 . Hence, again, specializing such algebras $A(w)$ is the same as specializing the corresponding nets of conics.

Suppose that $\ell(V) = 2$. We have $A(0) = R/(V, f, \mathfrak{m}^4)$ with $f \notin (V)_3$, and let $A(w), w \in \mathcal{W}$ be the family of algebras over $V(w), w \neq 0, V(w) \cong V'$ fixed. The quotient vector space $R_3/R_1V_w \cong \mathbf{k}^2$, so $R_3/R_1V_w, w \in \mathcal{W}$ is an affine \mathbb{A}^2 bundle over \mathcal{W} . Given $f \in R_3/R_1V_{w_0}$ there is a Zariski open subset $\mathcal{U} \subset \mathcal{W}$ parametrizing nets such that for $V_w \in \mathcal{U}$ the same $f \neq R_1V_w$. Thus

(V_w, f, \mathfrak{m}^4) for $w \in \mathcal{U}$ is an ideal defining an Artinian algebra $A(w)$ of constant Hilbert function T , whose limit as $w \rightarrow w_0$ is $A = A(0)$.

A similar proof holds for $\ell(V) = 3$, where f above is replaced by (f, g) and $R_3/R_1V_w, w \in \mathcal{W}$ is an affine \mathbb{A}^3 bundle over \mathcal{W} . An alternative proof for $\ell(V) = 3$ is in Theorem 3.3. \square

Constancy of the dimension $\ell(V)$ of the space $R_3/R_1V_w, w \in \mathcal{W}$ is key to this proof.

Remark 3.5. These results are valid over an algebraically closed field \mathbf{k} of characteristic zero or characteristic $p > 3$ - as then the classification of nets in Figure 1 is valid ([AEI, Appendix B, p. 80] or [Onda]). They are of course also valid over any field for which the classification of isomorphism classes of nets in Figure 1 is valid, but in general the classification of nets over non-algebraically-closed fields is more complicated (see [AEI, Appendix B]⁴

The deformation of algebras of Hilbert function $T = (1, 3, 3, 1)$ to Gorenstein algebras are treated in 5 different cases, depending on the length of the algebra. For the cases scheme length $\ell(V) = 0, 1$, the conclusion is easy. For the other cases scheme length $\ell(V) = 2, 3$ or ∞ we give explicit deformations. The proof of the following main result (first half of Theorem 1), is given below and in the following subsections.

Theorem 3.6. *Let $T = (1, 3, 3, 1)$. The family G_T of graded quotients of $R = \mathbf{k}[x, y, z]$ having Hilbert function T , is the Zariski closure of the family CI_T of graded complete intersection quotients of R having Hilbert function T ; and G_T has dimension 9.*

Proof outline. We have $I_2 = V$ where $\ell(V) \in \{0, 1, 2, 3, \infty\}$. Scheme length $\ell(V) = 0$ implies that $R/(V)$ is a complete intersection (CI), of Hilbert function $H(R/(V)) = (1, 3, 3, 1, 0)$, hence that the algebra A is a CI and there is nothing to show. These algebras are classified up to isomorphism in [AEI, Table 1]. These

⁴[AEI, Classification of nets of conics over algebraically closed fields of small characteristic, or over finite fields \mathbf{k} , Appendix B].

are our Table 1 cases #8b (continuous family of isomorphisms, parametrized by a function of λ), #8c, #7c, and #6d.

We suppose next that the scheme length $\ell(V) = 1$. These are the cases #8a or #7b of Figure 1. We have that the nets of #8a are in the closure of #8b since the latter parametrize a dense open in $\text{Grass}(R_2, 3)$ that defines CI algebras $A = R/(V)$; thus schemes with $I_2 \in \#8a$ are in the closure of CI_T . Hence the algebras A with I_2 a net of #7b are also in the closure of CI_T by Theorem 3.4.

For the cases $\ell(V) = 2$ or 3 , again by Theorem 3.4 it is enough to prove that the classes #7a and #6a are in the closure of CI_T . We show this by explicit deformations of A to a complete intersection (Sections 3.3.1 and 3.3.2). The case $\ell(V) = 3$ is also shown in Theorem 3.3.

When $\ell(V), V = I_2$ is infinite, the scheme $\mathfrak{Z}(V)$ (Proposition 2.5) may be chosen to be the line $x = 0$. This corresponds to the case #2a which is the special case of Section 3.4.1, Proposition 3.9 for $k = 2$.

□

3.3 Explicit deformations to AG algebras, $T = (1, 3, 3, 1)$.

By Theorem 3.4 the only explicit deformations needed to prove our main result that algebras $A \in G_T$ are in the closure of $\text{Gor}(T)$, when the scheme length $\ell(V), V = I_2$ is two or three, are those for the cases #7a and #6a. We nevertheless give here explicit deformations to Gorenstein algebras for all $A \in G_T$ such that the net $V = I_2$ has scheme length 2 or 3. We explain the cases #7a and #6a in detail.

3.3.1 Scheme length of $V = I_2$ is two, $T = (1, 3, 3, 1)$.

We consider first the nets $V = I_2$ of scheme length is equal to 2, i.e. the cases #7a, #6b, #6c and #5c. Any algebra $A = R/I$ in G_T having $V = I_2$ satisfies

$$I = (V, f, m^4) \text{ where } f \in R_3 \quad f \notin (V). \quad (6)$$

We give below the deformations needed on I so that the deformed algebra $A(t), t \neq 0$ is a complete intersection (CI).

1. **Case #7a.** Let the Artinian algebra $A = R/I$ of Hilbert function $H(A) = (1, 3, 3, 1)$ satisfy $I_2 = V = (zy+x^2, zx, yx)$, #7a in Table 1. Here V is symmetric in y, z and $R_3/R_1V \cong \langle z^3, y^3 \rangle$. We take $f = z^3 + ay^3$ with $a \neq 0$.

(When $f = y^3$ or z^3 we argue that the algebra A deforms first to a case where $f = z^3 + ay^3, a \neq 0$.)

Claim: A is in the closure of the complete intersections of the form $A = R/I(t)$ such that $V(t) = I(t)_2$ is in #8b for $t \neq 0$.

Proof of Claim. (i). Consider $f_1 = zy + x^2, f_2 = zx + t_1y^2, f_3 = t_2z^2 + xy$, and let $I(t) = (f_1, f_2, f_3, f)$, where t denotes (t_1, t_2) .

We assume henceforth that

$$at_1t_2 \neq 0 \text{ and } t_1t_2 \neq -1. \quad (7)$$

Then $x^4 \in I(t)$, since for $at_1t_2 \neq 0$ and $t_1t_2 \neq -1$,

$$(t_1t_2 + 1)x^4 = ((t_1t_2 + 1)^2 - t_1t_2yz)f_1 + t_2z^2f_2 - xzf_3.$$

Also, since $(1 + t_1t_2)x^2y = xf_3 - t_2zf_2 + t_1t_2yf_1$, then

$$R_1x^2y = (x^3y, x^2y^2, x^2yz) \subset I(t).$$

Similarly, since $(1 + t_1t_2)xy^2 = yf_3 + t_2xf_2 - t_2zf_1$, then

$$R_1xy^2 = (x^2y^2, xy^3, xy^2x) \subset I(t);$$

and since $(1 + t_1t_2)yz^2 = t_1yf_3 - xf_2 + zf_1$, then

$$R_1yz^2 = (xyz^2, y^2z^2, yz^3) \subset I(t) \quad \text{for } t \neq 0.$$

Also,

$$\begin{aligned} x^3z &= x^2f_2 - t_1x^2y^2, \quad t_2xz^3 = xzf_3 - x^2yz, \quad y^3z = y^2f_1 - x^2y^2 \\ z^4 &= zf - ay^3z, \quad x^2z^2 = xzf_2 - t_1xy^2z, \quad \text{and } ay^4 = fy - z^3y. \end{aligned}$$

Hence, for $a \neq 0$ and t satisfying Equation (7) (that is, $t_1 t_2 \neq 0, -1$) the ideal $I(t)$ contains \mathfrak{m}^4 and $HF(R/I(t)) = (1, 3, 3, 1)$.

(ii.) We now want to find a relation between t_1 and t_2 so that $I(t) = (f_1, f_2, f_3)$. A calculation by Macaulay2 shows that

$$t_1 t_2 f - t_1 z f_3 + (t_1 + at_2) x f_1 - at_2 y f_2 = (t_1 + at_2) x^3$$

Therefore, $f \in (f_1, f_2, f_3)$ if and only if $t_1 = -at_2 \neq 0$. Write $t_2 = t$ and $t_1 = -at$ and then we have that

$$I(t) = (zy + x^2, zx - aty^2, tz^2 + xy) \quad (8)$$

is a complete intersection, as desired, provided $t_1 = -at, t_2 = t \neq 0$, and $t_1 t_2 = -at^2 \neq -1$. This is Figure 1, entry #8b. \square

2. **Case #6b.** Let $V = (x^2, xy, z(z + y))$. Since $R_3/R_1 V \cong \langle y^2 z, y^3 \rangle$ we take $f = zy^2 + ay^3$ in Equation (6); and consider the deformed algebra $A(t) = R(t)/I(t)$ where

$$I(t) = (z^2 + yz, at^2 yz + x^2, aty^2 + xy).$$

Then $A(t)$ is a CI, provided that $at(at - 1) \neq 0$.

3. **Case #6c.** $V = (z^2 + x^2, yz, xz)$ and $f = y^3 + axy^2$. The deformation $A(t) = R(t)/I(t)$ with

$$I(t) = ((1 + at)x^2 + t(1 + at)xy + t^2 y^2 + z^2, yz + tx^2, xz),$$

is a CI if $at(1 + at) \neq 0$.

4. **Case #5b.** $V = (z^2, y^2, xy)$ and $f = x^2 z + ax^3$. Then, with $at \neq 0$ we have that $A(t) = R(t)/I(t)$ is a CI for

$$I(t) = (z^2, y^2 + at^2 xz, xy + atx^2, x^2 z + ax^3).$$

3.3.2 Scheme length of $V = I_2$ is three, $T = (1, 3, 3, 1)$.

The cases #6a, #5a, #4 and #2b of Table 1 correspond to nets V having scheme length 3. An algebra $A = R/I$ in G_T having $V = I_2$, with $\ell(V) = 3$ satisfies

$$I = (V, f, g, \mathfrak{m}^4) \text{ where } f, g \in R^3 \text{ and } \langle f, g \rangle \cap R_1V = 0. \quad (9)$$

1. **Case #6a**, Let $V = (xy, yz, xz)$ of scheme length 3. The ideal $I \in G_{\{V\}}(T)$ with $T = (1, 3, 3, 1)$ is up to isomorphism of the form (9) where $f = z^3 + ax^3$ and $g = y^3 + bx^3$, with $ab \neq 0$. Consider the following deformation:

Let $I(\mathbf{t}) = I(t_1, t_2, t_3) = (f_1, f_2, f_3, f, g)$ where $f_1 = yz + t_1x^2$, $f_2 = xz + t_2y^2$, $f_3 = xy + t_3z^2$. A computation by Macaulay2 shows that the Hilbert function of $A/I(\mathbf{t})$ is $(1, 3, 3, 1)$ with the conditions $abt_1t_2t_3 \neq 0$ and $1 + t_1t_2t_3 \neq 0$. We want to find a relation between the parameters so that f and g are in (f_1, f_2, f_3) . Using Macaulay2 again we get

$$(t_1 + at_3)x^3 = xf_1 - zf_3 + t_3f$$

and

$$(bt_2 - at_3)x^3 = -yf_2 + zf_3 - t_3f + t_2g.$$

Assuming furthermore that $t_1 + at_3 = 0$ and $bt_2 - at_3 = 0$, and with a change of variable $t_1 = -abt$, $t_2 = at$ and $t_3 = bt$, we set

$$I(t) = (yz - abtx^2, xz + aty^2, xy + btz^2) \supset (f, g)$$

with $abt \neq 0$ and $1 - a^2b^2t^3 \neq 0$. Then $A(t) = R(t)/I(t)$ is a subfamily of CI's in the isomorphism classes #8b.

2. **Case #5a**. $V = (z^2, xz, xy)$. We can take - possibly after an initial deformation - $f = yz^2 + ax^3$ and $g = yz^2 + by^3$. The deformed algebra $A(t) = R/I(t)$ where for $abt \neq 0$,

$$I(t) = (z^2 - a^2tx^2, xz + aty^2 - btyz, xy),$$

defining a complete intersection $A(t) = R(t)/I(t)$.

3. **Case #4.** $V = (x^2 + yz, xy, y^2)$, $f = xz^2 + az^3$ and $g = yz^2 + bz^3$. The desired deformation is

$$I(t) = (x^2 + yz + t_1z^2, xy + t_2z^2, t_3xz + y^2 + t_4z^2)$$

where $t_1 = at$, $t_2 = \frac{b}{a^2-b}t(b-at)$, $t_3 = bt$ and $t_4 = \frac{ab}{a^2-b}t(b-at)$ for $atb(a^2-b)(b-at) \neq 0$ and $t_2 + ab \neq 0$. Then $A(t) = R(t)/I(t)$ is a CI.

4. **Case #2b.** After a change of variables and an initial deformation we have $V = (z^2, yz, y^2)$, $f = zx^2 + ax^3$ and $g = yx^2 + bx^3$. We consider the following deformation

$$I(t) = (z^2 + (at^2 - bt)zy + atzx + a^2t^2x^2, yz, btxz + y^2)$$

which defines a complete intersection $A(t) = R(t)/I(t)$ when $abt \neq 0$.

3.4 The case $T = (1, 3^k, 1)$, $k \geq 3$: G_T is in the closure of $\text{Gor}_{\#6a}(T)$.

We begin with a simple observation.

Remark 3.7. Since $H(R/(V))_i = \ell(V)$ for $i \geq 3$, the only $V = I_2$ that are possible for a graded algebra A of Hilbert function $T = (1, 3^k, 1)$, $k \geq 3$ are

- (i). those of scheme length $\ell(V) = 3$: #6a, #5a, #4, #2b; OR
- (ii). V from #2a from Table 1 where the scheme $\mathfrak{Z}(V)$ is a line.

Theorem 3.8.

Let $T = (1, 3^k, 1)$ with $k \geq 3$. Then G_T is in the closure of $\text{Gor}_{\#6a}(T)$, the Artinian Gorenstein algebras $A = R/I$ where $I_2 \cong (xy, xz, yz)$, or, in other words those for which $I = \text{Ann}F$, $F \cong X^j + \alpha_1Y^j + \alpha_2Z^j$; and G_T has dimension eight.

Assume further that \mathbf{k} is closed under taking j -th roots. Then G_T is in the closure of the family $A_{\mathcal{L}} = R/I_{\mathcal{L}}$ where $I_{\mathcal{L}} = \text{Ann}F$ for $F = L_1^j + L_2^j + L_3^j$ with $\mathcal{L} = \{L_1, L_2, L_3\}$ linearly independent in S_1 .

This is the second part of Theorem 1. It follows from Remark 3.7, from Theorem 3.3 above and Proposition 3.9 below.

3.4.1 Scheme \mathfrak{Z}_V a line, $\ell(V) = \infty$.

The case V in #2a is the only case where the length of the scheme is infinite, with Hilbert function $(1, 3, 4, 5, \dots)$: the scheme \mathfrak{Z}_V is the line $x = 0$. We next show that the algebra $A = R/I$, with $V = I_2$ in #2a, deforms to an algebra $A(w)$ where $V(w), w \neq w_o$ is in #6a. Then Theorem 3.3 provides a deformation to a Gorenstein algebra. Note that the socle degree $j_A = k + 1$.

Proposition 3.9. *Let $T = (1, 3^k, 1), k \geq 2$. Suppose $A = R/I$ where $V = I_2 = \langle x^2, xy, xz \rangle$ (the class #2a) and $H(A) = T$. Then $A = A(w_o)$ is a specialization of a family $A(w), w \in W$ where $A(w) = R(w)/I(w), w \neq w_o$ satisfies $V(w) = I(w)_2$ is in class #6a, for $w \neq w_o$, and $H(A(w)) = T$.*

Proof. Since $H(R/I_2) = (1, 3, 3, 4, 5, \dots)$, in order to achieve the Hilbert function $H(A) = T$ we need that I has a generator $f \in \mathbf{k}[y, z]_3$, and further generators $g, h \in \mathbf{k}[y, z]_j$. The ideal (V, f) satisfies $H(R/(V, f)) = (1, 3, \bar{3})$ (if only because it is essentially generated in $\mathbf{k}[y, z]$ by a degree three form), so (V, f) defines a scheme $\mathfrak{Z}(w_0)$ of length 3 concentrated on the line $x = 0$ of $\mathbb{P}^2 = \text{Proj } R$. We have $I = (V, f, g, h, \mathbf{m}^{j+1})$ (we might need a generator for I in degree $j + 1$). There is a parameter variety $W \subset \text{Hilb}^3(\mathbb{P}^2)$ (we can take W to be a curve) such that the scheme $\mathfrak{Z} = \mathfrak{Z}(w_0)$ defined by (V, f) deforms to a scheme $\mathfrak{Z}(w) \subset \mathbb{P}^2$, defined by the ideal $(V(w)) \subset R$ such that for $w \neq w_0$ the scheme $\mathfrak{Z}(w)$ is smooth, comprised of three non-collinear points. Thus, for $w \neq w_0$, we have that $V(w) = I(w)_2$ defines $\mathfrak{Z}(w)$, and lies in the $\text{PGL}(3)$ equivalence class of #6a (here #6a: $V = (xy, xz, yz)$ corresponds to the scheme defined by the three points $(1, 0, 0), (0, 1, 0)$ and $(0, 0, 1)$). Asking that the ideal $(I(w)_2)$ have intersection 0 with $\langle g, h \rangle$ is an open condition on w , satisfied for $w = w_0$ by (V, f) .

Thus, we may deform $A = A(w_0)$ to the algebra $A(w) = R/I(w)$ where $I(w) = (V(w), g, h, \mathbf{m}^4)$ $H(A(w)) = T$ and $I(w)_2 = V(w)$ is in the isomorphism class #6a of nets of conics for $w \neq w_0$.

Remark 3.10 (Counterexample). We will show that for $V = (xy, xz, x^2)$ (# 2a) the family $G_{\{V\}}(T)$ of graded algebras $A = R/I$ of Hilbert function $T = (1, 3^k, 1), k \geq 3$ having $I_2 \cong V$,

form an irreducible variety of dimension seven, of known structure (Theorem 4.1(iii)). We will show also that the analogous family of quotients $A = R/I$ for #4, #5b or #6d, has dimension 6, and for #5a, #6b, #6c dimension 7 (Theorem 4.1(i),(ii)). It follows that when $\ell(V) \neq \ell(V')$ and V is a limit of nets in the class of V' , it may occur that not all algebras in $G_{\{V\}}(T)$ are in the closure of $G_{\{V'\}}(T)$.

Although we cannot make an analogous counter example involving only V', V of finite lengths for $T = (1, 3^k, 1), k \geq 3$, as the only occurring finite length is three, we can readily make an analogous example for $k = 2$, relying on Theorem 4.1 and Corollary 4.2 (see Remark 4.3). Namely, choose any pair (V', V) from Table 1 with dimension

$$d(V') = d(V) + 1 \text{ and } \ell(V) \geq \min\{2, \ell(V') + 1\};$$

then $G_{\{V\}}(T) \not\subseteq \overline{G_{\{V'\}}(T)}$, for dimension reasons. Examples of such pairs are (#7a, #6a), (#6b, #5a), and (#5b, #4).

□

3.5 Graded algebras G_T in codimension three not in the closure of $\text{Gor}(T)$.

We have proven in Theorem 1 that for $T = (1, 3^k, 1)$ $\overline{\text{Gor}(T)} = G_T$. To show that this result is sharp we determine the smallest-length codimension three Gorenstein sequences $T = (1, 3, 5, 3, 1)$, $(1, 3, 6, 3, 1)$ and $(1, 3, 6, 6, 3, 1)$ such that G_T has two irreducible components, $\text{Gor}(T)$ and another component U_T that we will describe.

Notation and preliminary facts. Below we set $R = \mathbf{k}[x, y, z]$ and $S = \mathbf{k}_{DP}[X, Y, Z]$, over an infinite field \mathbf{k} . Recall that R acts on S by contraction (Equation (1)). Recall

Lemma 3.11. *[ACGH, Prop. p.67 §4.2]. The rank r locus constructed from a generic $m \times n$ matrix M_{gen} satisfies*

$$\text{codimension of rank } r \text{ locus} = (m - r)(n - r). \quad (10)$$

This is also the maximum codimension for the rank r locus of any $m \times n$ matrix M .

We denote by $r_i = \dim_k R_i = \binom{i+2}{2}$. We recall

Definition 3.12. A *compressed* Gorenstein sequence H of codimension three and socle degree j satisfies

$$H = H(j) = (1, 3, \dots, h_u, \dots, 1_j) \text{ where } h_u = \min(r_u, r_{j-u}). \quad (11)$$

When j is even these have been called *extremal* Gorenstein sequences [Sch]. We will term *almost compressed* the Hilbert function $H(j) - 1_{j/2}$ when j is even. For example, $T = (1, 3, 6, 3, 1)$ is compressed, and $T = (1, 3, 5, 3, 1)$ is almost compressed.

	$a = 6$	$a = 5$	$a = 4$	$b = 6$	$b = 5$
$\dim \text{Gor}(T)$	14	13	11	20	15
$\dim U_T$	≥ 16	≥ 14	$\geq 10^*$	≥ 22	$\geq 14^*$
	$T = (1, 3, a, 3, 1)$			$T = (1, 3, b, b, 3, 1)$	

Figure 3: Dimensions of multiple components of $\text{Gor}(T), T = (1, 3, a, 3, 1), a = 5, 6$, and $T = (1, 3, 6, 6, 3, 1)$.

- $\geq 10^*$: $U_T, T = (1, 3, 4, 3, 1)$ is likely in the closure of $\text{Gor}(T)$.
- $\geq 14^*$: $U_T, T = (1, 3, 5, 5, 3, 1)$ may be in the closure of $\text{Gor}(T)$.

Definition 3.13. Given a Hilbert function $T = (1, 3, \dots, 1_j, 0)$ we will denote by $U_T \subset G_T$ the family of graded Artinian quotients $A = R/I$ of R such that $I_j^\perp = L^{[j]}$ the (divided) power of a linear form $L \in S_1$. We denote by $U_{T,s}, 1 \leq s \leq 3$ the subfamily of G_T for which the socle-degree j dual generator F_j satisfies $\dim_k R_1 \circ F = s$. Equivalently $H(R/\text{Ann}F_j) = (1, s, \dots, s, 1_j)$, satisfying $h_1 = h_{j-1} = s$. This is just the subfamily of G_T where F_j , after a suitable coordinate change, involves a minimum of s variables. Here $U_T = U_{T,1}$.

Example 3.14 (Algebras in $U_{T,i}$). Let $T = (1, 3, 3, 1)$. Then the Artinian algebra $A(1) = R/I(1), I(1) = \text{Ann}(X^3, Y^2, Z^2) =$

$(xz, yz, xy, y^3, z^3, x^4)$ is in $U_{T,1}$. The algebra $A(2) = R/I(2)$, $I(2) = \text{Ann}(X^2Y, Z^2) = (xz, yz, y^2, x^3, z^3)$ is in $U_{T,2}$; and the algebra $A(3) = R/I(3)$, $I(3) = \text{Ann}(XYZ) = (x^2, y^2, z^2)$ is in $U_{T,3}$.

By *closure* of an algebraic set we will mean its Zariski closure.

Lemma 3.15. *For $T = (1, 3, \dots, 3_{j-1}, 1_j, 0)$ the closure $\overline{U_T} \subset G_T$ is just $\overline{U_T} = U_T$. The closure of $U_{T,2}$ can contain no elements of $U_{T,3}$. When $T = (1, 3^k, 1)$ we have $U_{T,3} = \text{Gor}(T)$.*

For any Gorenstein sequence $T = (1, r, \dots, r, 1)$ with $r > 1$ we have $\overline{U_T} \cap \text{Gor}(T) = \emptyset$.

Proof. Let $A(w)$, $w \in W$ be a subfamily of G_T , and let $F(w) \in S_j$ (up to non-constant multiple) be the degree- j dual generator of $A(w)$. The condition $\dim_{\mathbb{k}} R_1 \circ F(w) \leq s$ asks that the module $R_1 \circ F \subset S_{j-1}$ have rank no greater than s , which defines a closed condition on W . This also implies the last statement of the Lemma.

When $T = (1, 3^k, 1)$ and $F \in S_j$ is a dual generator satisfying $\dim R_1 \circ F = 3$ and $\dim R_i \circ F \leq 3$ for $1 \leq i \leq j-1$, we have by symmetry that the Hilbert function of $A_F = R/\text{Ann}F$ satisfies $H_F = (1, 3, \dots, 3, 1)$; but the Macaulay equation (2) implies that for $i \geq 2$,

$$H(R/\text{Ann}F)_i = 2 \Rightarrow H(R/\text{Ann}F)_{i+1} \leq 2.$$

Since $(H_F)_{j-1} = 3$ this implies that $H_F = (1, 3^k, 1)$ so $A = R/\text{Ann}F$ and is an element of $U_{T,3} = \text{Gor}(T)$. Evidently, if $A_F \in \text{Gor}(T)$ then $\dim R_1 \circ F = 3$: we have shown $U_{T,3} = \text{Gor}(T)$. \square

Consider the family $\text{CI}(\mathcal{D})$ of graded complete intersection quotients of $R = \mathbb{k}[x, y, z]$ having generator degrees $\mathcal{D} = (d_1, d_2, d_3)$, and denote by $T(\mathcal{D})$ their Hilbert function, so $\text{CI}(\mathcal{D}) = CI_{T(\mathcal{D})}$. It is well-known that with $T = T(\mathcal{D}) = (t_0, t_1, \dots)$,

$$\dim \text{CI}(\mathcal{D}) = t_{d_1} + t_{d_2} + t_{d_3}. \quad (12)$$

Theorem 3.16 (Smallest G_T not in $\overline{\text{Gor}(T)}$). *Let $5 \leq a \leq 6$ and $T = (1, 3, a, 3, 1)$, or let $T = (1, 3, 6, 6, 3, 1)$. Then G_T has two irreducible components, $\overline{\text{Gor}(T)}$, and a second, larger component U_T of Definition 3.13, whose dimensions are given in Figure 3.*

Proof. For each of the cases $T = (1, 3, a, 3, 1)$ and $T = (1, 3, b, b, 3, 1)$ we show the step-by-step construction of U_T .

i. First consider $T = (1, 3, a, 3, 1)$. Choose an element $L \in S_1$: this is a choice parametrized by $\mathbb{P}(S_1) = \mathbb{P}^2$. We will take $F = L^4$ as our first dual generator, so $A^\vee \supset \langle 1, L, L^2, L^3, L^4 \rangle$.

i.a. Now let $a = 6$. Let W_L parametrize the three-dimensional sub-vector spaces of S_3 that contain $\langle L^3 \rangle$. Here $W_L \cong \text{Grass}(9, 2)$, in effect parametrizing two dimension subspaces of $S_3/\langle L^3 \rangle$.

Let $U_L \subset G_T$ be the subset comprised of algebra quotients $A_{\mathcal{B}}$ of R having dual generators $\langle L^4, \mathcal{B} \rangle$ where $\mathcal{B} \in W_L$ is general enough so that $R_1 \circ \mathcal{B} = S_2$. That this occurs for \mathcal{B} in a Zariski open dense subset of W_L is a consequence of the theory of relatively compressed algebras [I2, Theorem III]. Thus, each Artin algebra $A_{\mathcal{B}}, \mathcal{B} \in U_L$ satisfies $A_{\mathcal{B}} = R/\text{Ann} \langle \mathcal{B}, L^4 \rangle$, and has Hilbert function $H(A_{\mathcal{B}}) = T = (1, 3, 6, 3, 1)$.

Evidently, U_T of Definition 3.13 satisfies

$$\pi : U_T \rightarrow \mathbb{P}^2 : \pi(A_{\mathcal{B}}) = L, \quad (13)$$

is the bundle whose fibre over $L \in S_1$ is U_L . We have $U_T \subset G_T$, and

$$\dim U_T = \dim \mathbb{P}^2 + \dim \text{Grass}(9, 2) = 16.$$

But here the algebras $A \in \text{Gor}(T)$ are parametrized by their dual generators $F_A \in S_4 \bmod \mathfrak{k}^*$ -multiple: this is an open dense subvariety of $\mathbb{P}(S_4) \cong \mathbb{P}^{14}$ so $\text{Gor}(T)$ has dimension 14, smaller than that of U_T . It follows that U_T and $\overline{\text{Gor}(T)}$ are two irreducible components of G_T , when $T = (1, 3, 6, 3, 1)$.

i.b. Now let $a = 5$. Again we fix $L \in S_1$ and take the dual generator $F = L^4$. Then to determine an element of G_T , we must consider the elements \mathcal{B} of W_L (as above, three-dimensional subspaces of S_3 containing L^3) such that

$$\dim R_1 \circ \mathcal{B} = 5. \quad (14)$$

Recall that we have $\mathcal{B} = \langle L^3, v_1, v_2 \rangle$, so

$$R_1 \circ \mathcal{B} = \langle L^2, R_1 \circ v_1, R_1 \circ v_2 \rangle \subset S_3. \quad (15)$$

The condition (14) here asks that an ostensibly 6×7 matrix of Equation (15) have rank 5. The codimension of this condition, by Lemma 3.11 and Equation (10), is no greater than $(7-5)(6-5) = 2$. We let U_L parametrize those elements (3-dimensional vector spaces) $\mathcal{B} \in W_L$ satisfying the condition (14), so $A_{\mathcal{B}}$ has Hilbert function $H(\mathcal{B}) = T(a) = (1, 3, 5, 3, 1)$. Thus, U_T is a bundle over \mathbb{P}^2 (parametrizing the L) whose fibre over a linear form L is U_L .

From the above codimension count and Equation (13) we have that $\dim U_T(5) = 16 - 2 = 14$; however, $A \in \text{Gor } T, T = (1, 3, 5, 3, 1)$, is defined by the vanishing of $\text{Cat}_{2,2}(F_A)$, the catalecticant of the dual generator $F_A \in S_4$, so $\text{Gor}(T)$ has dimension $14 - 1 = 13$, respectively, of dimensions 14 and 13, respectively.

Thus, when $a = 5$, G_T has (at least) the two irreducible components U_T and $\overline{\text{Gor}(T)}$.

ii. Now consider $T = (1, 3, 6, 6, 3, 1)$. The dimension of $\text{Gor}(T) = \dim \mathbb{P}(S_5) = 20$, as $A \in \text{Gor}(T)$ is parametrized by the dual generator $F_A \in S_5$, up to non-constant scalar multiple. We next construct U_T similarly to the previous case (i).

As before, choose $L \in S_1$, a choice parametrized by $\mathbb{P}(S_1) = \mathbb{P}^2$, and now take $F = L^5$, so $A^\vee \supset \langle 1, L, L^2, L^3, L^4, L^5 \rangle$. Let W_L parametrize the three-dimensional sub-vector spaces $\mathcal{B} \subset S_4$ containing L^4 . Here

$$W_L \cong \text{Grass}(S_4/\langle L^4 \rangle, 2) = \text{Grass}(14, 2) \text{ of dimension } 24. \quad (16)$$

By the theory of relatively compressed algebras, for a dense open set of $\mathcal{B} \in W_L$ we have $\dim R_1 \circ \mathcal{B} = 7$ [I2, Theorem III]. That is, $\mathcal{B} = \langle L^4, v_1, v_2 \rangle$ and the 6 partials of general enough elements v_1, v_2 are linearly independent of L^3 . But we need $\dim_{\mathbb{k}} R_1 \circ \mathcal{B} = 6$. The matrix for $R_1 \circ \mathcal{B} \subset S_3$ is a 7×10 matrix: by Lemma 3.11 and Equation (10) the rank six locus has codimension at most $(7-6)(10-6) = 4$. Then U_T is the family of algebras $A_{\mathcal{B}} = R/\text{Ann}\langle L^5, \mathcal{B} \rangle$ such that $\dim R_1 \circ \mathcal{B} = 6$ and $R_2 \circ \mathcal{B} = S_2$ (the latter is again true generically for an open dense family of \mathcal{B} by [I2, Theorem III]).

We now will show that in general $R_2 \circ \mathcal{B} = S_2$, even on the locus where $\dim R_1 \circ \mathcal{B} = 6$. Suppose $R_2 \circ \mathcal{B} \neq S_2$. Choose $f \in R_2$, a choice of $f \in \mathbb{P}(R_2) = \mathbb{P}^5$. then $H(R/(f)) = (1, 3, 5, 7, 9, \dots)$

and \mathcal{B} must be a 3-dimensional space inside a 9 dimensional space $f^\perp \cap S_4$, or in effect, since L is chosen, a 2-dimensional space inside an 8 dimensional space. It follows that such \mathcal{B} are parametrized by a $\text{Grass}(8, 2)$, a 12 dimensional family: adding the choice of f and of L gives a 19-dimensional family. But the family U_L of $\mathcal{B} \supset L^4$ and satisfying $\dim R_1 \circ B = 6$ has by Lemma 3.11 and Equation (16) dimension at least

$$\dim U_T \geq 2 + 24 - 4 = 22,$$

so generically \mathcal{B} satisfying the rank condition $\dim_k R_1 \circ \mathcal{B} = 6$ and $R_2 \circ \mathcal{B} = S_2$ has dimension at least 22, compared to 20 for $\text{Gor}(\overline{T})$ so G_T has (at least) the two irreducible components U_T and $\text{Gor}(T)$ when $T = (1, 3, 6, 6, 3, 1)$.

This completes the proof of the Theorem. \square

We include an example due to the fourth author.

Example 3.17 (J. Yaméogo). [IaKa, Lemma 8.3, p. 251]. When $T = (1, 3, 4, 3, 1)$ then

- (i). The ideal $I = \text{Ann}W, W = \langle L^4, X^3, Y^3, XY \rangle$ which satisfies $I = (xz, yz, x^2y, xy^2, x^4, y^4, z^5)$, is a singular point of G_T .
- (ii). Let $f \in S_4$ be in the closure $\overline{\text{Gor}(T)}_4^\perp$ of the dual generators of ideals in $\text{Gor}(T)$. Then the fibre of G_T over f (those algebras $A = R/I \in G_T$ such that $I_4 = (\text{Ann}f)_4$ consists of those ideals I “in” G_T with $I \subset J = \text{Ann}f$.

Question 3.18. For $T = (1, 3, 4, 3, 1)$ do we have $\overline{\text{Gor}(T)} = G_T$?

The answer “Yes” would imply that the cases $T = (1, 3, 6, 3, 1)$ and $T = (1, 3, 5, 3, 1)$ are the only socle-degree 4, embedding dimension three, where $\overline{\text{Gor}(T)} \neq G_T$.

4 Isomorphism classes of graded Artinian algebras of Hilbert function $T = (1, 3^k, 1)$.

Recall that $R = \mathbb{k}[x, y, z], S = \mathbb{k}_{DP}[X, Y, Z]$ the action of R on S is contraction. Here $\{V\}$ denotes the $\text{PGL}(3)$ isomorphism class of

nets equivalent to the vector space V , $\text{Grass}_{\{V\}} = \text{Grass}_{\{V\}}(R_2, 3)$ parametrizes the family of nets of conics $\text{PGL}(3)$ isomorphic to V , and $G_{\{V\}}(T)$ parametrizes the family of all graded algebras having $I_2 \cong V$. The dimension $d(V)$ of the family $\text{Grass}_{\{V\}}$ is indicated at the left of Table 1. We will denote by $\text{Gor}_V(T)$ and $G_V(T)$ respectively the family of Artinian Gorenstein quotients $A = R/I$ having $I_2 = V$, and all such Artinian quotients; and by $\text{Gor}_{\{V\}}(T)$ and $G_{\{V\}}(T)$, respectively, those quotients for which $I_2 \cong V$, that is, is congruent to V under $\text{PGL}(3)$. We will also denote the later by $\mathcal{G}_{\#6b}(T)$ - for $I_2 \cong V$ of $\#6b/$

4.1 Parametrizing Artinian algebras in $G_{\{V\}}(T)$.

When $T = (1, 3^k, 1)$ with $k \geq 3$ and $A = R/I \in G_T$, then the vector space $V = I_2$ must have scheme length three, or be in case $\#2a$, as otherwise $H(R/(I_2)) = (1, 3, 3, \ell, \dots)$, and $H(R/I)$ would be smaller than T .

We will first show

Theorem 4.1. *We let $d = d(V)$ be the dimension of the family of nets of conics that are $\text{PGL}(3)$ equivalent to V . Let $T = (1, 3^k, 1)$. Then $G_V(T)$ and $G_{\{V\}}(T)$ are irreducible.*

- (i). *Let $T = (1, 3, 3, 1)$. Then the family of Artinian algebras $G_{\{V\}}(T)$ has dimension d if $\ell = 0$, dimension $d + \ell - 1$ if $1 \leq \ell \leq 3$; and has dimension 5 if $\ell = \infty$ (the case $V = \#2a$).*
- (ii). *Let $T = (1, 3^k, 1)$ with $k \geq 3$. Then $G_{\{V\}}(T)$ has dimension $d + 2$ if $\ell = 3$ and dimension 7 if $\ell = \infty$ (case $V = \#2a$), and is empty otherwise.*

Proof of (i). For V not in the class $\#2a$, the Hilbert function $H(R/(V)) = (1, 3, 3, \bar{\ell}), \ell = \ell(V)$. Then the family $G_{\{V\}}(T)$ is fibred over $\text{Grass}_{\{V\}}$ parametrizing the nets, by a fibre that is the projectivization of $R_3/\langle(V) \cap R_3\rangle$, a vector space of k -dimension ℓ : hence the fibre has dimension $\ell - 1$ if $1 \leq \ell \leq 3$. Otherwise, the fiber comprises a single point. For V in the case $\#2a$, the Hilbert function $H(R/(V)) = (1, 3, 3, 4, \dots)$ and I_3 is determined by a 3-dimensional quotient of R_3/R_1V , hence by an element

of a projective 3-space $\mathbb{P}(R_3/R_1V) \cong \mathbb{P}^3$, so the dimension of $G_{\#2a}(T) = 5$.

Proof of (ii). When $k \geq 3$, the case $\ell(V) = 3$ is treated in Theorem 3.3, in particular Equation (5). When V is from #2a in Table 1 the Hilbert function of $H(R/(V)) = (1, 3, 3, 4, \dots)$, and $\dim \text{Grass}_{\{V\}} = 2$. Given V we first choose $f \in \mathbb{k}[y, z]_3$, so that $H(R/(V, f)) = (1, 3, 3, \overline{4})$, parametrized by \mathbb{P}^3 , then we choose a two dimensional subspace of the 3-dimensional space $R_j/\langle (V, f) \cap R_j \rangle$, so an element of \mathbb{P}^2 . This shows $G_{\{V\}}(T)$ is an irreducible family of dimension 7 when $k \geq 3$. □

We may also use that when $T = (1, 3, 3, 1)$ the fibre of $G_{\{V\}}(T)$ over $\text{Grass}_{\{V\}}$ is isomorphic to $\mathbb{P}(S_3/((V)_3^\perp)$, the dual of R_3/R_1V . Recall that specialization of nets V' to V must satisfy $\ell(V') \leq \ell(V)$. We see from Table 1 that the net $V = (xy, xz, x^2)$ (#2a) is in the closure of all families of nets in Figure 1 except #2b.

Corollary 4.2 (Specialization). *(i). Let $T = (1, 3, 3, 1)$. Suppose the net V' has dimension $d(V') = d(V) + 1$ and scheme length $\ell(V') = \ell(V) - 1$, and assume V is not in #7b. Then the Zariski closure of $G_{\{V'\}}(T)$ cannot include $G_{\{V\}}(T)$.*

(ii). Let $T = (1, 3^k, 1), k \geq 3$. Let $V = (xy, xz, x^2)$ (Case #2a). Then $G_{\{V\}}(T), T = (1, 3^k, 1), k \geq 3$ is not in the closure of any $G_{\{V'\}}(T)$ for which $d(V') \leq 5$.

Proof. Corollary 4.2 (i) follows from Theorem 4.1(i) and the scheme lengths in Table 1. This is so since the dimension of both $G_{\{V'\}}(T)$ and $G_{\{V\}}(T)$ are the same under the hypotheses, and each is irreducible. Corollary 4.2 (ii.) follows from Theorem 4.1(ii). □

Recall that Proposition 3.9 shows, pertaining to (ii) above, that if the net V is in #2a, then $G_{\{V\}}(T)$ is in the closure of $G_{\#6a}(T)$.

Remark 4.3. Part (i) of the Corollary applies, for example, to the pairs (#5b, #4), (#6b, #5a), (#6c, #5a), (#6d, #5b), and (#7b,c, #6b,c). Since $G_{\{V'\}}(T)$ is $\text{PGL}(3)$ invariant, so are their closures. Since all classes but #8b of nets of conics are “discrete” (do not involve parameters), the closure of an isomorphism stratum $\text{Grass}_{\{V'\}}$ of nets over an algebraically closed field of characteristic zero, or of characteristic $p \neq 2, 3$ is the union of certain

lower strata, specified in [AEI] and in our Figure 1. In contrast, Theorem 4.1 shows that - provided certain expected intersections are non-empty - that certain $G_{\{V\}}(T)$ for a fixed isomorphism class $\{V\}$ of nets of conics must contain several isomorphism classes of algebras A , of different dimensions! However, each $G_{\{V\}}(T)$ is itself irreducible.

Question 4.4. The above results then lead to a typical question in parametrization, analogue of one that baffled the Nice school of Hilbert scheme specialists concerning closures of families Z_T of non-graded Artinian quotients of $\mathbf{k}[y, z]$ having a given Hilbert function T .⁵ Our problem here is to determine the intersections $\overline{G_{\{V'\}}(T)} \cap G_{\{V\}}(T)$, when the scheme length of V' is less than that of V , in cases as in Remark 4.3 when for dimension reasons there cannot be an inclusion $\overline{G_{\{V'\}}(T)} \supset G_{\{V\}}(T)$. Are these intersections irreducible?

4.1.1 Betti tables and cancellation.

We recall that given any finitely generated graded Artinian algebra A of codimension three, quotient of $R = \mathbf{k}[x_1, x_2, x_3]$, has a minimal graded finite free resolution

$$\mathcal{F} : 0 \rightarrow F_3 \rightarrow F_2 \rightarrow F_1 \rightarrow R \rightarrow A \rightarrow 0, \quad (17)$$

where $F_i = \bigoplus_j R(-j)^{\beta_{i,j}}$, and $\beta_{i,j} = \text{Tor}_i^R(A, \mathbf{k})_j$. We represent this in a *Betti table*, whose (i, j) entry in column i , row j is $\beta_{i,i+j}$. For example, the minimal resolution for $A = R/I, I = (xy, xz, yz, x^3 - y^3, z^3)$, the ideal in #6a.ii. below, whose terms F_3, F_2, F_1, F_0 are

$$R(-6) \oplus R(-5) \rightarrow R(-5) \oplus R(-4)^3 \oplus R(-3)^2 \rightarrow R(-3)^2 \oplus R(-2)^3 \rightarrow R, \quad (18)$$

which is encoded in the Betti diagram C, Figure 8.

We will in Section 4.5.1 discuss problems involving limits of Betti tables. A *consecutive cancellation* from a resolution \mathcal{F} to

⁵The fundamental result of the irreducibility of $\text{Hilb}^n(\mathbf{k}[y, z])$: that the closure of the stratum $T = (1^n)$ is all length- n Artinian algebra quotients of $\mathbf{k}[y, z]$ had been shown by J. Briançon in [Bri], but the closures of other strata are even now only partially understood, see the last author's [Y1, Y2, Y3].

\mathcal{F}' is given by setting for some pair $(i, j), (i + 1, j)$ of adjacent indices where $\beta_{i,j}, \beta_{i,j+1} \neq 0$, the Betti indices $\beta'_{i,j} = \beta_{i,j} - 1$, $\beta'_{i+1,j} = \beta_{i+1,j} - 1$. In a Betti table, this will lead to two indices in adjacent columns and rows, one diagonally above the other, being each decreased by one. For example the Betti diagram C of Figure 8 arises by consecutive cancellation of $\beta_{3,5} = \beta_{2,5} = 1$ from Equation (18) above and Betti diagram D of Figure 9. Recall that the *lexicographic ideal* J of Hilbert function $H(R/J) = H$ is the unique monomial ideal J first in the lex-order. I. Peeva has shown [Pe, Theorem 1.1]

Lemma 4.5. (*I. Peeva*) *Let I be a graded ideal and L be the lexicographic ideal with the same Hilbert function. The graded Betti numbers $\beta_{i,j}(S/I)$ can be obtained from the graded Betti numbers $\beta_{i,j}(S/L)$ by a sequence of consecutive cancellations.*

This was extended to local algebras A , and a comparison between the graded Betti numbers for the associated graded algebra and A by M.E. Rossi and L. Sharifan [RoSh], using negative and consecutive cancellations. Maria Evelina Rossi pointed out to us that L above in Lemma 4.5 may be replaced by $Lt(I)$ the leading terms of I in any term order (see [Pe, Proof (2) of Thm. 1.1]).

4.2 Classification of isomorphism classes of Artinian algebras of Hilbert function T by the net of conics $V = I_2$: the family $G_V(T)$.

We here show

Theorem 5. Let $T = (1, 3^k, 1)$. We determine the isomorphism classes of algebras in $G_V(T)$, the family of Artinian quotients $A = R/I$ in G_T for which $I_2 = V$.

There are among the nets V of scheme length 3, exactly four isomorphism classes of Gorenstein algebras in $G_V(T)$, namely #6a.i, #5a.i, #5a.ii, and #4.iii., (see the Betti C table in Figure 8). There are no further non-CI Gorenstein classes of algebras in G_T .

This Theorem summarizes the results of Theorems 4.6, 4.8 and 4.11 for V of scheme length two, three and ∞ , respectively, where we also specify the Betti tables. A detailed statement of

Theorem 5 (minus the Betti tables) is in the Introduction. We assume throughout the section that \mathbf{k} is closed under taking j -th roots, or is algebraically closed, of characteristic not 2 or 3.

4.2.1 Classification of algebras in $\text{Gor}(T), T = (1, 3^k, 1)$.

Previous classification of isomorphism types of Artinian Gorenstein (AG) algebras having Hilbert function $T = (1, 3^k, 1), k \geq 2$.

Recall from Proposition 2.5(ii). and Lemma 2.6 that when $k \geq 4$, there is necessarily a length 3 punctual subscheme $\mathfrak{Z} \subset \mathbb{P}^2$, determined as $\mathfrak{Z} = \text{Spec} R/(\overline{I_3})$, where $\overline{I_3}$ is the ancestor ideal of I_3 . This occurs since $(3_3, 3_4)$ is a maximal growth Using the structure of nets of conics, we can show this for $k \geq 3$. Thus, determining G_T is closely related to the study of length 3 punctual subschemes of \mathbb{P}^2 . The article of A. Bernardi et al [BGI, §4.2, Theorem 37] classifies the isomorphism types of length 3 subschemes of \mathbb{P}^n in characteristic zero. When $n = 2$ we may derive this classification from the scheme-length 3 cases of Table 1, together with a study of length-3 schemes that are on a line: the latter is related to Section 4.3 below. P. LeBarz and G. Elencwajg classify triangles in \mathbb{P}^2 : they determine the cohomology ring structure of $\text{Hilb}^3(\mathbb{P}^2)$ [EL] - see also E. Vasserot's more general [Va].

G. Casnati, J. Jelisiejew, and R. Notari determine, for arbitrary characteristic, the three isomorphism classes of forms $F \in \mathbf{k}[X, Y, Z]_j$ whose apolar algebra $A = R/\text{Ann}F$ has Hilbert function $H(A) = (1, 3^k, 1)$ when $k \geq 3$ [CJN, Proposition 4.9]. These are

$$X^j + Y^j + Z^j, X^{j-1}Y + Z^j, X^{j-2}(XZ + Y^2). \quad (19)$$

The statements there are given for characteristic zero, but the arguments hold for \mathbf{k} algebraically closed of any characteristic $p \neq 2, 3$. J. Jelisiejew [Je] determines the isomorphism types of AG algebras A of Hilbert function $H(A) = (1, 3, 3, 3, 1)$ including those for non-graded A .

When $T = (1, 3^k, 1)$ with $k \geq 3$ then all Gorenstein algebras $A = R/I$ of Hilbert function T have five generators. The generators are comprised of the net, which must have scheme length

three, and two in the socle degree $j = k + 1$. When $k = 2$ there may also be complete intersections Gorenstein algebras whose ideal is generated by the net (scheme length zero), or those with five generators - two of degree four. We verify this in our classification, see the statement of Theorem 5 above.

We next classify all graded algebras $A \in G_T$, for these Hilbert functions, and specify the dimension of their isomorphism classes, as well as their Betti tables, dividing our classification according to the scheme length of $V = I_2$.

4.2.2 Isomorphism classes in $G_{\{V\}}T$ when $V = I_2$ defines a length 0 or 1 scheme.

Evidently, these ideals are just $I = (V, \mathfrak{m}^4)$ so their isomorphism classes and dimensions are those of the corresponding nets $V = I_2$. When the scheme length of V is 0 then I_2 defines a complete intersection, with CI betti table of Figure 4: these are #8b, #8c, #7c, #6d.

The Betti resolutions in both #8a, #7b, are the table SL1 of Figure 5.

4.2.3 Isomorphism classes in $G_{\{V\}}T$ when $V = I_2$ defines a length two scheme.

Here $T = (1, 3, 3, 1)$ is the only Hilbert function possible. We give isomorphism classes $G_{\{V\}}T$ of Artinian algebras where the net $V = I_2$ defines a length two scheme by cases, according to isomorphism class of the net V in Figure 1 (derived from [AEI, Tables 1 and 5]). These are the cases #7a, #6b, #6c, and #5b.

Theorem 4.6. *The isomorphism classes of graded Artinian algebras of Hilbert function $T = (1, 3, 3, 1)$ for which $V = I_2$ has scheme length two, and their Betti tables are listed as follows:*

- (i). *Case V in class #7a: two classes, Equation (20);*
- (ii). *Case V in class #6b. There is a one-parameter family of non-isomorphic Artinian algebras, parametrized by \mathbb{P}^1 , having $V = I_2$ in #6b (Conclusion 4.7);*

(iii). Case $V \in \#6c$: three classes, Equation (22);

(iv). Case $V \in \#5b$: three classes, Equation (23).

There are no Gorenstein algebras having such $V = I_2$, that is $\text{Gor}_V(T)$ is empty for each V of scheme length two.

Proof. We work out the proof by cases.

Case #7a: Here $V = \langle x^2 + yz, xy, xz \rangle$; any algebra in $G_{\{V\}}(T)$ has the form $A(\alpha, \beta) = R/I(\alpha, \beta)$, $I(\alpha, \beta) = (V, f(\alpha, \beta), \mathfrak{m}^4)$ where $f = \alpha y^3 + \beta z^3$. $R_1 V^\perp = \langle Z^3, Y^3 \rangle$. Here V is invariant under scaling $\omega_a x \rightarrow x', y \rightarrow ay', z \rightarrow a^{-1}z'$, and also under $\theta_b : x \rightarrow bx', y \rightarrow b^2 y', z \rightarrow z'$, and under $\sigma\theta : x \rightarrow bx', y \rightarrow y', z \rightarrow b^2 z'$.

When both $\alpha, \beta \neq 0$, we may scale z by a cube root of β/α , to obtain $A(\alpha, \beta) \cong A(1, 1)$ giving # 7a.i, of Betti table A from Figure 6. When one of $\alpha, \beta = 0$, then, by symmetry $y \rightarrow z, z \rightarrow y$, there is a single isomorphism class #7a.ii. of Betti table B from Figure 7. Also, there are no Gorenstein elements of $G_V(T)$.

$$\begin{aligned} \#7a.i : I &= (z^2 + xy, yz, xz, y^3 + z^3, \mathfrak{m}^4), & \dim 8; \\ \#7a.ii : I &= (z^2 + xy, yz, xz, x^3, \mathfrak{m}^4). & \dim 7. \end{aligned} \quad (20)$$

Case #6b: Here any algebra in $G_{\{V\}}(T)$ has the form $A = R/I, I = (V, f, \mathfrak{m}^4)$ where $V = \langle x^2, (y+z)z, xy \rangle$ and $f \in R_3/R_1 V \cong \langle y^3, z^3 \rangle$, note $R_1 V^\perp = \langle Y^3, Y^2 Z - Y Z^2 + Z^3 \rangle$. Notice that the stabilizer $G = \text{Stab}_{\text{PGL}(3)} V$ in $\text{PGL}(3)$ is two dimensional, as the dimension of the orbit of #6b is six, and the dimension $\text{PGL}(3)$ is 8. The stabilizer G is generated by the two scalings: $x \rightarrow ax$ and $y, z \rightarrow by, bz, a, b \neq 0$. These each act on $f(\alpha, \beta) = \alpha z^3 + \beta y^3$ by scaling it, by 1, or b^3 , respectively. Thus, when $\alpha\beta \neq 0$ we may take $\alpha = 1, f(1, \beta) = z^3 + \beta y^3$, and we have a one-dimensional parametrized family of non-isomorphic algebras

$$A_V(1, \beta) = R/(V, z^3 + \beta y^3, \mathfrak{m}^4) \text{ with } I_2 = V.$$

For $\beta \neq 0$, $A_V(1, \beta)$ has the Betti diagram⁶ A of Figure 6.

The Betti diagram A of Figure 6 occurs also for $A_V(0, 1)$.

For $\beta = 0$ the algebra $A_V(1, 0) = R/(V, z^3, \mathfrak{m}^4)$ has the Betti diagram B of Figure 7. Note, there are no Gorenstein Artin

⁶tested by Macaulay2 for various β .

algebras in $\text{Gor}_V(T)$ - this can be seen from the Betti diagrams, or from $F \in (V)_3^\perp \in \langle Y^3, Z^3 \rangle$, so $R_1 \circ F \subset \langle Y^2, Z^2 \rangle$ has dimension two not three.

Conclusion 4.7. [Projective line of isomorphism classes] For $V = I_2$ isomorphic to #6b, there is a projective line $\beta \in \mathbb{P}^1$ of non-isomorphic algebras $A(1, \beta)$, elements of $G_V(T)$, $T = (1, 3, 3, 1)$, including $A_V(1, 0)$ and $A_V(0, 1) \equiv A_V(1, \infty)$. The Betti diagram A occurs for all classes except $A_V(1, 0)$, which has Betti diagram B. There are no Gorenstein elements of $G_V(T)$.

total	1	3	3	1
0	1	-	-	-
1	-	3	0	-
2	-		3	0
3	-			1

Figure 4: Betti diagram CI, #8b,#8c,#7c,#6d

total	1	4	6	3
0	1	-	-	-
1	-	3		-
2	-		4	2
3	-	1	2	1

Figure 5: Betti diagram SL1 (Scheme length 1) #8a,#7b.

4.2.4 Betti diagrams for $A \in G_T, T = (1, 3, 3, 1)$

total	1	4	5	2
0	1	-	-	-
1	-	3	1	-
2	-	1	3	1
3	-	-	1	1

Figure 6: Betti diagram A, #7a.i.,#6b(1, β), #5b i,ii

total	1	5	7	3
0	1	-	-	-
1	-	3	1	-
2	-	1	4	2
3	-	1	2	1

Figure 7: Betti diagram B, #7a.ii, #6b $A_V(0, 1)$, #5b.iii.

total	1	5	5	1
0	1	-	-	-
1	-	3	2	-
2	-	2	3	
3	-			1

Figure 8: Gorenstein Betti diagram C #6a.i, #5a.i, #5a.ii. For Betti C_j move rows $(2, 3)$ to rows $\{j - 1, j\}$.

total	1	5	6	2
0	1	-	-	-
1	-	3	2	-
2	-	2	3	1
3	-		1	1

Figure 9: Betti diagram D, #6a.ii; #5a.iii; #5a.iv; #5a.vii; #4.ii; #2b.i, #2b.ii; for Betti D_j move rows $(2, 3)$ to $\{j - 1, j\}$.

total	1	6	8	3
0	1	-	-	-
1	-	3	2	-
2	-	2	4	2
3	-	1	2	1

Figure 10: Betti diagram E, #6a.iii., #5a.v, #5a.vi, 4.iii., #2b.iii. Betti E_j move rows $(2, 3)$ to rows $\{j - 1, j\}$

total	1	6	8	3
0	1	-	-	-
1	-	3	3	1
2	-	3	4	1
3	-		1	1

Figure 11: Betti diagram G #2a.i,ii Equation (35)

total	1	7	10	4
0	1	-	-	-
1	-	3	3	1
2	-	3	5	2
3	-	1	2	1

Figure 12: Betti diagram H #2a.iii (Lex ideal) Equation (35).

Case #6c: We have $V = \langle xz, yz, x^2 + z^2 \rangle$ and $R_1V^\perp = \langle XY^2, Y^3 \rangle$. Any element $A = R/I$ of $G_V(T)$ satisfies $I = (V, f(\alpha, \beta), \mathfrak{m}^4)$ where $f(\alpha, \beta) = \alpha y^3 - \beta xy^2$.

Case (i). When both $\alpha, \beta \neq 0$, we have

$$\begin{aligned} I(\alpha, \beta) &= (xz, yz, x^2 + z^2, \alpha y^3 - \beta xy^2, \mathfrak{m}^4) = I(1, \lambda) \\ &= (xz, yz, x^2 + z^2, f(1, \lambda) = y^3 - \lambda xy^2, \mathfrak{m}^4), \end{aligned} \quad (21)$$

where $\lambda = \beta/\alpha$. By scaling $(x, z) \rightarrow \lambda(x', z')$ and $y \rightarrow y'$ we have $I(1, \lambda) = I(1, 1)$, and its Betti table is that of Figure 6. We have three isomorphism classes:

$$\begin{aligned} I(1, 1) &= (xz, yz, x^2 + z^2, y^3 - xy^2, \mathfrak{m}^4), \text{ dimension 7, Betti table A} \\ I(1, 0) &= (xz, yz, x^2 + z^2, y^3, \mathfrak{m}^4), \text{ dimension 6, Betti table A} \\ I(0, 1) &= (xz, yz, x^2 + z^2, x^2y, \mathfrak{m}^4), \text{ dimension 6. Betti table B.} \end{aligned} \quad (22)$$

There are no Gorenstein elements of $G_{\#6c}(T)$.

Case #5b: We have $V = \langle xy, y^2, z^2 \rangle$ and $R_1V^\perp = \langle X^3, X^2Z \rangle$, so $A = R/I$ with $I(\alpha, \beta) = (V, f(\alpha, \beta), \mathfrak{m}^4)$ with $f(\alpha, \beta) = \alpha x^3 + \beta x^2z$. Note that there can be no Gorensteins: $\text{Gor}_V(T)$ is empty, as $R_1 \circ (R_1V)^\perp \subset \langle X^2, XZ \rangle$.

When $\alpha, \neq 0$ we have $I(\alpha, \beta) = I(1, \lambda), \lambda = \beta/\alpha \in \mathbf{k}$. Here $f = x^3 + \lambda x^2 z$ for $\lambda \neq 0$ can be scaled to $\lambda = 1$ by scaling x . Thus, we have three isomorphism classes. Note that for 5b.ii the ideal $I_{\leq 3}$ contains powers of all the variables, so defines an Artinian quotient of R , in contrast to 5b.iii: thus these cases are non-isomorphic. By dimension they cannot be isomorphic to #5b.i . \square

Thus we have - here $V = \langle xy, y^2, z^2 \rangle$ -

$$\begin{aligned}
\#5b.i : I(1, 1) &= (V, x^3 + x^2 z, \mathfrak{m}^4), \quad \dim = 6, \text{ Betti table A;} \\
\#5b.ii : I(1, 0) &= (V, x^3, \mathfrak{m}^4), \quad \dim = 5, \text{ Betti table A;} \\
\#5b.iii : I(0, 1) &= (V, x^2 z, \mathfrak{m}^4), \quad \dim = 5, \text{ Betti table B.}
\end{aligned}
\tag{23}$$

Question. We have by Theorem 4.1 that the dimension of #5b.i is 6; and we can check that #5b.ii and #5b.iii both have dimension 5. Thus, $A \in G_{\{\#6d\}}(T)$ of scheme length zero and dimension 6 must specialize to either or both of #5b.ii or #5b.iii. Which?

4.3 Isomorphism classes in G_T when I_2 defines a length 3 scheme, $T = (1, 3^k, 1)$.

From Figure 1 these are the cases $V = I_2$ from #6a, #5a, #4, and #2b. Recall from Theorem 3.3, Equation (5) that $I_k = (V)_k$ for $2 \leq k \leq j - 1$. Thus, the fiber of $G_{\{V\}}(T)$ over a fixed $V = I_2$ parametrizes algebras A entirely determined by the pair $(V, \langle F \rangle)$, where $\langle F \rangle \in (V_j)^\perp \subset S_j$, as $I_j = (\text{Ann} F)_j$. The three-dimensional vector spaces $(V_j)^\perp$ are given in Table 2. We now just need to

- (i). parametrize the choices of $\langle F \rangle$ from these.
- (ii). Determine their isomorphism classes.

Theorem 4.8. *The isomorphism classes of Artinian algebras of Hilbert function $H = (1, 3^k, 1), k \geq 2$, for which $V = I_2$ has scheme length three are given as follows:*

- (i). Case $V = I_2 = (xy, xz, yz)$ (#6a), 3 classes, Equation (25).

- (ii). Case $V = I_2 = (xy, xz, z^2)$ (#5a), 7 classes, Equation (27).
(iii). Case $V = I_2 = (x^2 + yz, xy, y^2)$ (#4), 3 classes, Equation (31).
(iv). Case $V = I_2 = (x^2, xy, y^2)$ (#2b), 3 classes, Equation (32).

Proof and discussion of cases:

#6a Here $V = (xy, xz, yz)$ defining the scheme $\mathfrak{Z} = (1, 0, 0) \cup (0, 1, 0) \cup (0, 0, 1)$, and $R_{j-1}V^\perp = \langle X^j, Y^j, Z^j \rangle$. Assuming that \mathbf{k} is closed under cube roots, we have up to isomorphism, three classes determined by choosing $I_j^\perp = X^j + \alpha Y^j + \beta Z^j$ (case 6a.i where A is Gorenstein), or $I_j^\perp = X^j + \alpha Y^j$ (case 6a.ii.) or $I_3^\perp = X^j$ (case 6a.iii.). These lead to the families,

$$\begin{aligned}
\#6a.i &: (V, \alpha x^j - y^j, \beta x^j - z^j). \text{ Gorenstein family, dim 8,} \\
\#6a.ii &: (V, \alpha x^j - y^j, z^j). \quad \text{dim 7,} \\
\#6a.iii &: (V, y^j, z^j, x^{j+1}). \quad \text{dim 6.} \tag{24}
\end{aligned}$$

whose isomorphism classes, assuming \mathbf{k} is closed under j -th roots, are

$$\begin{aligned}
6a.i &: (V, x^j - y^j, x^j - z^j); \quad 6a.ii : (V, x^j - y^j, z^j) \text{ and} \\
6a.iii &: (V, y^j, z^j, x^{j+1}). \tag{25}
\end{aligned}$$

When $k = 2$ #6a.i has the Betti diagram C of Figure 8, 6a ii has the Betti diagram D of Figure 9, while #6a.iii. has the Betti diagram E of Figure 10.

#5a: $V = (xy, xz, z^2)$ so $(R_{j-1}V)^\perp = \langle X^j, Y^j, Y^{j-1}Z \rangle$. So $I(\alpha, \beta, \gamma; j) = (V, W(\alpha, \beta, \gamma; j), \mathfrak{m}^{j+1})$ for some $\alpha, \beta, \lambda \in \mathbf{k}$, where

$$\begin{aligned}
W(\alpha, \beta, \gamma; j) &= (\text{Ann}F) \cap R_j, F = \alpha X^j + \beta Y^j + \lambda Y^{j-1}Z \\
&= \langle \beta x^j - \alpha y^j, \gamma x^j - \alpha y^{j-1}z, \gamma y^j - \beta y^{j-1}z \rangle. \tag{26}
\end{aligned}$$

If \mathbf{k} is closed under j -th roots, we may, up to isomorphism, scale x, y, z by j -th roots, keeping V fixed. It follows that we have the following isomorphism classes, where each named parameter

among α, β, γ is non-zero and where $j \geq 3$.

$$\begin{aligned}
\#5a.i : I(\alpha, \beta, \gamma; j) &\equiv I(1, 1, 1; j), \text{ for } \alpha, \beta, \gamma \neq 0, \\
I(j) &= (V, x^j - y^j, y^{j-1}z - y^j), \text{ Betti } C_j, \text{ dim } 7; \\
\#5a.ii : I(\alpha, 0, \gamma; j) &\equiv I(1, 0, 1; j), \\
I &= (V, x^j - y^{j-1}z, y^j), \text{ Betti } C_j, \text{ dim } 6; \\
\#5a.iii : I(\alpha, \beta, 0; j) &\equiv I(1, 1, 0; j), \\
I &= (V, x^j - y^j, y^{j-1}z), \text{ Betti } D_j, \text{ dim } 6; \\
\#5a.iv : I(0, \beta, \gamma; j) &\equiv I(0, 1, 1; j), \\
I &= (V, y^j - y^{j-1}z, x^j), \text{ Betti } D_j, \text{ dim } 6; \\
\#5a.v : I(\alpha, 0, 0) &= I(1, 0, 0), \\
I &= (V, y^3, y^{j-1}z, x^{j+1}), \text{ Betti } E_j, \text{ dim } 5; \\
\#5a.vi : I(0, \beta, 0; j) &= I(0, 1, 0; j), \\
I &= (V, x^j, y^{j-1}z, y^{j+1}), \text{ Betti } E_j, \text{ dim } 5; \\
\#5a.vii : I(0, 0, \gamma; j) &= I(0, 0, 1; j), \\
I &= (V, x^j, y^j) \text{ Betti } D_j, \text{ dim } 5. \tag{27}
\end{aligned}$$

#4: $V = \langle x^2 + yz, xy, y^2 \rangle$, then $R_1V^\perp = \langle Z^3, XZ^2, X^2Z - YZ^2 \rangle$. We know that V defines a length three scheme in \mathbb{P}^2 , supported on $(0, 0, 1)$; the ideal of the scheme \mathfrak{Z} in $\mathbf{k}\{x, y\}$ is $(y + x^2, yx, y^2) = (y + x^2, x^3)$, which is curvilinear. The following Lemma arose from considering the automorphisms of the scheme \mathfrak{Z} : the group $\Phi_{\underline{t}}$ describes all the automorphisms of \mathbb{P}^2 that fix the curvilinear length three scheme defined by (V) .

Lemma 4.9. *Let $\underline{t} = (\alpha, \beta, \gamma, \lambda)$ with $\alpha \neq 0$. Then the following four-dimensional parameter space $\Phi_{\underline{t}} = \{\phi_{\underline{t}}\}$ of algebraic automorphisms of $\mathbf{k}[x, y, z]$ fixes V : we define $\phi_{\underline{t}}$ by*

$$\begin{aligned} \phi_{\underline{t}}(x) &= x + \lambda y \\ \phi_{\underline{t}}(y) &= (1/\alpha)y \\ \phi_{\underline{t}}(z) &= \gamma x + \beta y + \alpha z \end{aligned} = \begin{pmatrix} 1 & \lambda & 0 \\ 0 & (1/\alpha) & 0 \\ \gamma & \beta & \alpha \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

Proof of Lemma. Just check that $\phi_{\underline{t}}(V) = V$. □

Let $I \in G_V(T)$. Then $I = (V, f_4, f_5, \mathbf{m}^4)$ where $\mathbf{m} = (x, y, z)$ and

$$\begin{aligned} f_4 &= a_1xz^2 + a_2yz^2 + a_3z^3 \\ f_5 &= b_1xz^2 + b_2yz^2 + b_3z^3, \end{aligned} \quad (28)$$

with $\dim_{\mathbf{k}}\langle f_4, f_5 \rangle = 2$. It is clear that a minimal set of generators for $I \in G_V(T)$ will have five or six elements.

We will use the following in studying the orbit of $\Phi_{\underline{t}}$ acting on an ideal I . Note that modulo R_1V we have

$$\begin{aligned} \phi_{\underline{t}}(xz^2) &= \alpha^2xz^2 + (\lambda\alpha^2 - 2\alpha\gamma)yz^2 \\ \phi_{\underline{t}}(yz^2) &= \alpha yz^2 \\ \phi_{\underline{t}}(z^3) &= 3\alpha^2\gamma xz^2 + 3(\alpha^2\beta - \alpha\gamma^2)yz^2 + \alpha^3z^3. \end{aligned} \quad (29)$$

- (1). If $a_1 = b_1 = 0$, then $I = (V, yz^2, z^3)$, is not Gorenstein: $A = R/I$ has a two dimensional socle $\langle xz^2, x^2 - yz \rangle$, and A has the Betti tabel D of Figure 9. One can see that

$$\begin{aligned} \phi_{\underline{t}}(I) &= (V, yz^2, 3\gamma xz^2 + \alpha z^3) \\ &= (V, yz^2, xz^2 + \nu z^3) \end{aligned} \quad (30)$$

so the orbit $\Phi_{\underline{t}}(V)$ has dimension one, so we have a five-dimensional isomorphism class.

- (2). Suppose $a_1 \neq 0$ in Equation (28). Then we may set $f_4 = xz^2 + a_2yz^2 + a_3z^3$; $f_5 = b_2yz^2 + b_3z^3$.
- (a). if $b_3 = 0$ and $a_3 \neq 0$, then $I = (V, yz^2, xz^2 + a_3z^3)$. One can see that setting $\alpha = a_3, 3\gamma = 1$ this ideal belongs to $\Phi_{\underline{t}}(V, yz^2, z^3)$, which is (1) above.
- (b). if $b_3 = 0$ and $a_3 = 0$, then $I = (V, xz^2, yz^2, z^4)$. Here $A = R/I$ is not Gorenstein: its socle $\langle z^3, x^2 - yz, xz \rangle$ has dimension three, and it has Betti diagram E of Figure 10. We have $\Phi_{\underline{t}}(I) = I$, so the fibre of the orbit over V has dimension zero, so the orbit of A has dimension four.
- (c). if $b_3 \neq 0$, then we may set $f_4 = xz^2 + a_2yz^2$, $f_5 = b_2yz^2 + z^3$. In this case the ideal $I = (V, f_4, f_5)$ is Gorenstein with Betti diagram C of Figure 8, and Macaulay dual generator $F = a_2XZ^2 + X^2Z - YZ^2 + b_2Z^3$. Such an ideal is in the orbit of (V, xz^2, z^3) under $\Phi_{\underline{t}}$, the fibre of the orbit has dimension two, parametrized by a_2, b_2 ; or also because this is a generic choice of $\langle f_4, f_5 \rangle$ from a three-dimensional space R_3/R_1V . The orbit of A has dimension six.

So we have three isomorphism classes in $G_V(T)$, $V = (x^2 + yz, xy, y^2)$ (take $a_2, b_2, \nu \neq 0$)

- 4.i. $I = (V, xz^2 + a_2yz^2, b_2yz^2 + z^3, \dim. 6, \text{Betti C (Gorenstein)})$;
4.ii. $I = (V, yz^2, xz^2 + \nu z^3), \dim 5, \text{Betti D}$;
4.iii. $I = (V, xz^2, yz^2, z^4), \dim 4, \text{Betti E}.$ (31)

The analysis of $G_V(T)$ for $V = I_2 = \#4$ is similar for $T = (1, 3^k, 1)$ and leads to analogous families, of Betti tables C_j, D_j , and E_j , respectively.

#2b. $V = (x^2, xy, y^2) = \text{Ann}\langle XZ, YZ, Z^2 \rangle$, and $(R_1V)^\perp = \langle XZ^2, YZ^2, Z^3 \rangle$. So $V = \text{Ann}\langle ZS_1 \rangle \cap R_2$, and $R_1V = \text{Ann}Z^2S_1 \cap R_3$, each depending on a 2-dimensional choice of $Z \in \mathbb{P}(S_1)$.

Case $T = (1, 3, 3, 1)$. Then $I_3^\perp = \langle F \rangle$ and $I = (V, \text{Ann}F \cap R_3, \mathfrak{m}^4)$. We may take $F = LZ^2 + \gamma Z^3$, where $L = \alpha X + \beta Y$. But up to isomorphism $L = X$ or $L = 0$ and we can scale Z , so up to

isomorphism we have $F = XZ^2 + Z^3$, or $F = XZ^2$, or $F = Z^3$, and we have, respectively, up to isomorphism,

$$\begin{aligned} & \#2b.i.(V, xz^2 - z^3, yz^2), \text{ Betti D, dim 4; ,} \\ & \#2b.ii.V, xz^2, z^3), \text{ Betti D, dim 3;} \\ & \#2b.iii.(V, xz^2, yz^2, z^4), \text{ Betti E, dim 2.} \end{aligned} \quad (32)$$

For the dimensions it suffices to consider group action on the dual generator $F = R_3^\perp$.

Case $T = (1, 3^k, 1)$, $k \geq 3$ Here $(V^\perp)_j = \langle XZ^{j-1}, YZ^{j-1}, Z^j \rangle$ and the characterization of isomorphism types is a straightforward generalization of the case $k = 2$. Here the diagram Betti D_j adjusts the degree of the later generators from three, four to $j, j + 1$. We have

$$\begin{aligned} & \#2b.i.(V, xz^{j-1} - z^j, yz^{j-1}), \text{ Betti } D_j, \text{ dim 4; ,} \\ & \#2b.ii.V, xz^{j-1}, z^j), \text{ Betti } D_j, \text{ dim 3;} \\ & \#2b.iii.(V, xz^{j-1}, yz^{j-1}, z^j), \text{ Betti } E_j, \text{ dim 2.} \end{aligned} \quad (33)$$

4.4 Isomorphism classes in G_T , $k \geq 2$ when $V = I_2$ defines a line (case #2a).

Here, up to isomorphism we may take $V = xR_1 = \langle x^2, xy, xz \rangle$. The stabilizer of V in $\text{PGL}(3)$ is

$$\begin{aligned} x' &= x \\ y' &= ux + ay + bz \\ z' &= vx + cy + dz, \end{aligned} \quad (34)$$

where $(u, v, a, b, c, d) \in \mathbf{k}^6$ and $ad - cd \neq 0$.

We have shown in Section 3.4.1, Proposition 3.9 that each algebra $A \in G_V(T)$ is in the closure of $\text{Gor}_{V'}(T)$ where V' is from #6a. We now determine their isomorphism classes. We let $R' = \mathbf{k}[y, z]$ and $S' = \mathbf{k}_{DP}[Y, Z]$.

Consider first $T = (1, 3, 3, 1)$. Then $I_3 = \text{Ann}F \cap R_3$, with $F \in S'_3 = \langle Y^3, Y^2Z, YZ^2, Z^3 \rangle$: it is well known that up to

isomorphism, for k algebraically closed, then $F \in S'_3$ satisfies $F \cong ZY(Z + Y), YZ^2$ or Z^3 , whose isomorphism classes are of dimension three (open in $\mathbb{P}^3 = \mathbb{P}(S'_3)$), two (choose $Y, Z \in S'_1$ up to non-constant k -multiple), or one (choose Z up to non-zero constant multiple), respectively. We have the following result, where the fibre dimension is that of the family $G_V(T)$ over a fixed V ,

Lemma 4.10. *Let $T = (1, 3, 3, 1)$ and suppose $V = \langle x^2, xy, xz \rangle$. Then there are three isomorphism classes of algebra $A = R/I$ in $G_V(T)$, namely*

$$\begin{aligned}
2a.i. & I = (x^2, xy, xz, yz^2 - y^2z, y^3, z^3), \text{ fibre dim } 3, \text{ Betti } G; \\
2a.ii. & I = (x^2, xy, xz, y^2z, y^3, z^3), \text{ fibre dim } 2, \text{ Betti } G; \\
2a.iii. & I = (x^2, xy, xz, y^3, y^2z, yz^2, z^4), \text{ (lex ideal) fibre dim } 1, \text{ Betti } H.
\end{aligned}
\tag{35}$$

Proof. The given ideals satisfy $I = (V, W, \mathfrak{m}^4)$, where $W = \text{Ann}F \cap R'_3$; this also gives the dimensions of the isomorphism classes. \square

Consider $T = (1, 3^k, 1), k \geq 3$ with $I_2 = V = \langle x^2, xy, xz \rangle$. Then, since $\text{cod}R_1V = 4$ in R_3 , the ideal $I \in G_V(T)$ must have a degree-three generator, which we may consider to be $f \in R_3/R_1V \cong \langle y^3, y^2z, yz^2, z^3 \rangle = R'_3$. As before for $F \in S'_3$, up to isomorphism of R' we may assume that $f = yz(y-z), yz^2$, or z^3 , choices of fiber dimension 3, 2 and 1 respectively, given V . The Hilbert function $H(R/(V, f)) = (1, 3, 3, \bar{3})$ as $(1, 3, 3, 3, 4, \dots)$ is not possible by Macaulay's bound Equation (2) and Lemma 2.6. We then must choose generators $g, h \in R_j/(V, f)_j$, so $I = (V, f, g, h, \mathfrak{m}^{j+1})$; since $(V) \cap R_j \supset xR_{j-1}$, we may assume that $g, h \in R'_j$ and are determined by a Macaulay dual generator $F \in S'_j$ that is annihilated by f . In the three isomorphism classes for f we have, respectively, as isomorphism classes for F , determining those for I

Theorem 4.11. *[$I_2 = \#2a$] Let $T = (1, 3^k, 1), k \geq 3$ and assume that $\text{char } k \neq 2, 3$. There are the following isomorphism classes and dimensions for algebras $A = R/I \in G_V(T)$ where $V = I_2 \cong \langle x^2, xy, xz \rangle$ ($\#2a$) is fixed and $k \geq 3$, as in Figures 13, 14, 15.*

f	$F \in ((V, f)^\perp)_j$	F up to isomorphism
$yz(y-z)$	$\langle Y^j, (Y+Z)^{[j]}, Z^j \rangle$	$Y^j, Y^j + aZ^j, Y^j + aZ^j + b(Y+Z)^{[j]}$
y^2z	$\langle Z^j, YZ^{j-1}, Y^j \rangle$	$Z^j, YZ^{j-1}, Y^j, Y^j + Z^j, Y^j + YZ^{j-1}$
y^3	$\langle Z^j, YZ^{j-1}, Y^2Z^{j-2} \rangle$	$Z^j, YZ^{j-1}, Y^2Z^{j-2}, Y^2Z^{j-2} + Z^j$

Figure 13: Isomorphism classes of (f, F) for $A \in G_V(T)$, $V = \langle x^2, xy, xz \rangle$ (Theorem 4.11, Equation (36)).

Each ideal I defining $A \in G_V(T)$ satisfies

$$I = (V, f, \text{Ann}F \cap R_j, \mathfrak{m}^{j+1})$$

where $f \in R_3/xR_2 \cong R'_3$, and $F \in (V, f)_j^\perp \subset S'_j$, (36)

where $f \cong yz(y-z)$ or y^2z or z^3 . Here the fibre $G_V(T)$ over V is irreducible of dimension five: it is fibred over V by a projective plane \mathbb{P}^2 parametrizing the element F , over the 3-space \mathbb{P}^3 parametrizing $f \in R'_3$.

There are over $f = yz(y-z)$ a two-parameter family $I(a, b)$ of isomorphism classes (Equations (39) and (41)), and a one-parameter family $I(a)$ (Equation (42)) as well as the specific class $I(0, 0)$. Over each of y^2z, z^3 there are a finite set of isomorphism classes, as listed in Figure 15.

Proof. Fix $V = I_2 = (x^2, xy, xz)$.

For $i \geq 2$ we have $R/(I_2)_i \cong R'_i$ and $(I_2^\perp)_i = S'_i$. Thus, an ideal I defining $A \in G_V(T)$ satisfies

$$I = (I_2, f, W, \mathfrak{m}^{j+1}), \quad (37)$$

where we may regard $f \in R_3/xR_2$, parametrized by a projective space $P(R'_3) \cong \mathbb{P}^3$, the two-dimensional $W \text{ Grass}(R_j/(I_2, f)_j, 2)$, parametrized by a projective plane \mathbb{P}^2 , so a total fibre dimension of five; here also $W = \text{Ann}F \cap R'_j / (V, f)_j$, where $F \in S'_j \cap ((V, f)_j)^\perp$. The dimension of $G_{\{V\}}(T) = G_{\#2a}(T)$ is thus 7.

Consider $f \in R_3/xR_2 \cong R'_3$: up to isomorphism f is one of $yz(y-z), y^2z$ or y^3 , families of dimension, respectively, 3, 2, 1. For $f = y^2z$ or y^3 we will use F to parametrize $I_j = \text{Ann}(F)_j$. In Figure 13 we write the space $\langle (f)_j^\perp \rangle$ and then the isomorphism

classes of elements F from that space. However, for $f = yz(y - z)$ we prefer the direct calculation of isomorphism classes of W , just below.

Calculating W when $f = yz(y - z)$. Consider the vector subspace $\mathcal{W} = \langle y^j, yz^{j-1}, z^j \rangle \subset R_j$, which induces a basis for $R_j/((V, f) \cap R_j)$.

We may write $W = \langle f_5, f_6 \rangle$ Then $I = (V, yz(y - z), W, \mathfrak{m}^4)$ where $\mathfrak{m} = (x, y, z)$ and $W = \langle f_5, f_6 \rangle$, where

$$\begin{aligned} f_5 &= a_1 y^j + a_2 y z^{j-1} + a_3 z^j \\ f_6 &= b_1 y^j + b_2 y z^{j-1} + b_3 z^j \end{aligned} \quad (38)$$

- (i). If $a_1 \neq 0$ we may write $f_6 = b_2 y z^{j-1} + b_3 z^j$; if now also $b_2 \neq 0$, we may, after scaling, write the two parameter family of isomorphism classes $I(a, b) = (xR_1, yz(y - z), W(a, b))$ where

$$W(a, b) = \langle f_5, f_6 \rangle \text{ where } f_5 = y^j - a z^j, f_6 = y z^{j-1} + b z^j. \quad (39)$$

However, we may also apply the involution $\sigma : y \rightarrow z, z \rightarrow y$. Then if $a \neq 0$ we have $\sigma(f_5) = a y^j - z^j$ which we may scale to $y^j - a^{-1} z^j$ and $\sigma(f_6) = z y^{j-1} + b y^j = y z^{j-1} + a^{-1} b z^j$, so we have

$$W(a, b) = W(a^{-1}, a^{-1}b). \quad (40)$$

Using a dual generator $F(a, b) = aY^j + Z^j - bYZ^{j-1}$, we may write

$$I(a, b) = (xR_1, f, \langle \text{Ann } F(a, b) \cap R_j', \mathfrak{m}^{j+1} \rangle). \quad (41)$$

- (ii). If $a_1 \neq 0$ but $b_2 = 0$, we may reduce to

$$W(a) = \langle f_5, f_6 \rangle, f_5 = y^j + a y z^{j-1}, f_6 = z^j, \quad (42)$$

a one-parameter family of isomorphism classes with degree- j dual generator $F = aY^j - YZ^{j-1}$.

- (iii). If $a_1 = b_1 = 0$, then, since W is two-dimensional, we must have $W = \langle yz^{j-1}, z^j \rangle$ isomorphic to $\sigma(W) = \langle y^{j-1}z, y^j \rangle \cong \langle y^j, yz^{j-1} \rangle = W(0, 0)$.

In Figure 14, we give the dimension of these isomorphism classes, and in Figure 15 we give the isomorphism classes in terms of their defining ideals. For example, the isomorphism class of the pair $(f = y^3, F = Z^j)$ is determined by a choice of $y \in R_1/\langle x \rangle$ then $z + ay$ (in place of $Z \in \langle Y, Z \rangle$, each a one-dimensional choice. The base dimension of the isomorphism class $y^2z \in R_3/\langle xR_2 \rangle$ is 2, giving the dimensions $(3, 3, 3, 4)$ for the respective classes of F ; likewise, the base dimension of $yz(y - z)$ is 3 (parametrized by an element of $R_3/xR_2 \cong \mathbf{k}^4$ up to \mathbf{k}^* action, so by \mathbb{P}^3); there are just three perfect powers in the space $(V, yz(y - z))_j^\perp$, yielding dimension 3 - the base dimension - when $F = Z^3$. The generic element of $G_V(T)$ is, evidently the family with two-dimensional fibre $I(a, b) = (V, yz(y - z), W(a, b))$ of Equation (41), where $I(a, b)_j = \langle \text{Ann } F(a, b) \cap R_j \rangle$, with $F(a, b) = aY^j + Z^j - bYZ^{j-1}$. This family has total dimension 7.

f	F	dim. of fibre $G_V(T)$ over V
$yz\ell$	$Z^j, Y^j + aZ^j, aY^j + Z^j - bYZ^{j-1}$	3, 4, 5;
y^2z	$Z^j, YZ^{j-1}, Y^j, Y^j + Z^j, YZ^{j-1} + Y^j$	3, 3, 3, 4, 4;
y^3	$Z^j, YZ^{j-1}, Y^2Z^{j-2}, Y^2Z^{j-2} + Z^j$	2, 2, 2, 3.

where $T = (1, 3^k, 1)$, $k \geq 3$ and $\ell = y - z$.

Figure 14: Dimension of isomorphism classes in $G_V(T)$, $T = (1, 3^k, 1)$, $k \geq 3$, $V = \langle x^2, xy, xz \rangle$ (Theorem 4.11)

□

Note. The Betti tables J and K for these classes of Figure 15 were calculated using Macaulay 2 for $j = 4$. Evidently Betti table K of Figure 17 is the table for the lexicographic ideal (having highest column sums). Note that the lex ideal of socle degree 4, so $T = (1, 3^3, 1)$ is that corresponding to (y^3, Z^4) in Figure 14.

f	F_1	F_2	F_3
$yz(y-z)$	Z^j	$Y^j + aZ^j,$	$aY^j + Z^j - bYZ^{j-1}$
	$W(0,0)$	$W(a)$	$W(a,b)$ of Equation (39)

f	F_1	F_2	F_3	F_4	F_5
y^2z	Z^j	YZ^{j-1}	Y^j	$Y^j + Z^j$	$Y^j + YZ^{j-1}$
	(y^j, yz^{j-1})	(y^j, z^j)	(y^j, yz^{j-1})	$(y^j - z^j, yz^{j-1})$	$(y^j - yz^{j-1}, z^j)$

f	F_1	F_2	F_3	F_4
y^3	Z^j	YZ^{j-1}	Y^2Z^{j-2}	$Y^2Z^{j-2} + Z^j$
	(y^2z^{j-2}, yz^{j-1})	(y^2z^{j-2}, z^j)	(yz^{j-1}, z^j)	$(y^2z^{j-2} - z^j, yz^{j-1})$

where $T = (1, 3^k, 1), k \geq 3$ and $\ell = y + z$.

Figure 15: Isomorphism classes of ideals in $G_V(T), T = (1, 3^k, 1), k \geq 3, V = \langle x^2, xy, xz \rangle$ (Theorem 4.11).

total	1	6	8	3
0	1	-	-	-
1	-	3	3	1
2	-	1	1	
3	-	2	3	1
4	-		1	1

Figure 16: Betti diagram $J_4 \#2a.(f, F) = (y^3, YZ^3)$, also $(f = yz(y-z), all F)$, and $(f, F) = (y^2z, F \neq Z^4)$. See Figure 15.

For J_j move the last two rows to indices $(j-1, j)$.

4.5 Questions and Problems

4.5.1 Limits of Betti tables in $\text{Gor}(T)$ and G_T .

Let T be an arbitrary Gorenstein sequence - one possible for a graded Artinian Gorenstein algebra. For $\text{Gor}(T)$ in codimension three, S.J. Diesel and D. Buchsbaum-D. Eisenbud determined that, given T the minimal Betti resolutions compatible with T have a poset structure, and satisfy a frontier property: the Zariski closure of a Betti stratum of $\text{Gor}(T)$ is the union of Betti strata ([Di], see also ([IaKa, Theorem 5.25] for a statement). For our special $T = (1, 3^k, 1)$ the Gorenstein minimal resolution table is unique if $k > 2$ (Betti table C_j , Figure 8); there are two Goren-

total	1	7	10	4
0	1	-	-	-
1	-	3	3	1
2	-	1	1	
3	-	2	4	2
4	-	1	2	1

Figure 17: Betti diagram K_4 #2a., for $(f, F) = (y^3, Z^4)$ (the lex ideal for $T = (1, 3, 3, 3, 1)$) and (y^2z, Z^4) ; for the lex table K_j of socle degree $j \geq 4$ move the last two rows to indices $(j - 1, j)$.

stein minimal resolutions when $k = 2$ (Betti table CI - Figure 4, and Betti table C). H. Schenck et al in [KKRSSY, p. 65-66] show that in an analogous codimension four case, the Betti tables do *not* satisfy a frontier property. I. Peeva has shown that all Betti tables for a given Hilbert function may be obtained by consecutive cancellation from that of a lex-segment ideal (Lemma 4.5). She also showed [Pe, Remark after Theorem 1.1]:

Lemma 4.12. *Consider a flat family $\mathcal{F}(t)$ over $\mathbf{k}[t]$ of graded Artinian quotients of R whose fiber $\mathcal{F}(0)$ over 0 is S/J and whose fiber $\mathcal{F}(t)$ over $t \in \mathbf{k} \setminus 0$ is S/J' . Then the graded Betti numbers $\beta'_{i,j} = \beta_{i,j}(S/J')$ of the general fiber may be obtained from the graded Betti numbers $\beta_{i,j}(S/J)$ of the special fiber by a sequence of consecutive cancellations.*

An example is the C,D,E Betti diagrams (Figures 8, 9 and 10, respectively) for the three isomorphism classes #6a.i., #6a.ii., and #6a.iii., respectively where $I_2 = (xy, xz, yz)$ (#6a), for $G_T, T = (1, 3, 3, 1)$: each is obtained by consecutive cancellation from the next.⁷

Question. Lemma 4.12 assumes a “jump specialization” - there is a single isomorphism class among the more general algebras. Is there a similar statement when we allow the more general family to not be in a single isomorphism class of algebras: that is, a family $\mathcal{F}(t)$ whose fiber over $t \in \mathbf{k}$ is $\mathcal{F}(t) = S/J'(t)$, but which

⁷We thank Alexandra Seceleanu who pointed out an error in a previous writing of the Betti diagram for #6a.ii.; we have made consequent changes here.

have a common Betti table β' for $t \neq t_0$? Then is β' obtained by consecutive cancellation from the Betti table $\beta(S/J(t_0))$?

Question 4.13. Do the Betti table strata for $T = (1, 3^k, 1)$ satisfy the frontier property? If so, is this a more general result for T that are codimension three Gorenstein sequences?

Since we have shown that $\overline{Gor(T)}$ need not be all of G_T for some codimension three Gorenstein sequences, this question appears to be open for such T in general.

4.5.2 Specialization of G_T isomorphism classes.

Note: As the dimension of the classes $G_V(T)$ of Artin algebras corresponding to the nets #7b,#7c are seven, we can expect that they each specialize to a dimension six class,#6c.ii, or #6c.iii? (latter, when char $k = 3$). Which specializations occur?

Question 4.14. We found in specializing nets of conics, that because each family but #8b was a complete isomorphism class, there were no jump specializations, that is if we have specializations $U \rightarrow Z$ and the pair $U \rightarrow Y, Y \rightarrow Z$ then the specialization $U \rightarrow Z$ comes from first specializing $U \rightarrow Y$, then $Y \rightarrow Z$. Now we would need to consider - and perhaps rule out - that the algebra class say in #8a, might jump, that is, specialize to a class in #6a, but not specialize there through a class in #7a. Can we rule such out?

Goal. Make tables of specializations of Artinian algebras of Hilbert function $T = (1, 3^k, 1)$, extending Table 1 for nets of conics. The first table would be for isomorphism classes of $T = (1, 3, 3, 1)$ algebras, and the second for isomorphism classes of $T = (1, 3^k, 1), k \geq 3$ algebras.

Remark 4.15. The only direct specializations to #6a algebras can be from #7a algebras: here #7a has two classes, one of dimension 8, and the other of dimension 7. In specializing we must drop dimension of a family by at least one, so we have that #6a.i. cannot be the specialization of any family in #7a; #6a.ii could only be a specialization from #7a.i of dimension 8; and #6a.iii might be a specialization of either class in #7a.

Question. According to Remark 4.3 we expect that $\overline{G_{\#8a}(T)} \cap G_{\#7a}(T) = G_{\#7a.ii}(T)$, the dimension seven component. What is the intersection of the closure of a 8-dimensional cell of $\#8a$ with the 8-dimensional cell here?

4.5.3 Possible future directions involving Jordan type or applications.

The Jordan degree type of multiplication m_ℓ on a graded Artinian algebra A , adds to the information of the lengths of Jordan strings, their initial degrees. Note that [AAIY] has a table with the classification of Jordan degree types for graded AG algebras of Hilbert function $(1, 3^k, 1)$: these are all symmetric, by a general result of T. Harima and J. Watanabe [HW] (see also [IMM, Prop. 2.38]). But their limit algebras may have JDT that are not symmetric. The notation $1_{2\uparrow 3}$ means two strings of length 1, beginning in degree 2 and 3, respectively.

Example 4.16 (Limit algebra has non-symmetric JDT, using non-generic ℓ). Take $I = (x^2, xy, xz, y^3, z^3, \mathfrak{m}^4)$ of Hilbert function $H(R/I) = (1, 3, 3, 1)$ and $\ell = x$, then $x^2 = 0$ and the Jordan degree type $S_{x,A} = (2_0, (1_{1\uparrow 2})^2, 1_{2\uparrow 3})$ which is not symmetric.

A. What are the possible generic JDT for these algebras in G_T that are not Gorenstein?

B. Can we specify how Jordan types correspond to the Betti tables, as in a recent paper of N. Abdallah and H. Schenck [AbSc]?

C. Some people have applied nets of conics, Can we apply these algebras? Mapping germs?

See Appendix B, p. 82 of [AEI] for some applications others have made.

D. The classification of pencils of conics up to isomorphism is well known. Describe the related isomorphism classes in the family G_T of graded Artinian algebras of Hilbert function $T = (1, 3, 4^k, 3, 1)$?

List of Figures

1	Specializations of nets of conics and scheme length.	9
2	Nets of scheme length three and their duals. . . .	15
3	Dimensions of multiple components of $\text{Gor}(T)$, $T = (1, 3, a, 3, 1)$, $a = 5, 6$, and $T = (1, 3, 6, 6, 3, 1)$	24
4	Betti diagram CI, #8b, #8c, #7c, #6d	37
5	Betti diagram SL1 (Scheme length 1) #8a, #7b.	37
6	Betti diagram A, #7a.i., #6b(1, β), #5b i,ii	37
7	Betti diagram B, #7a.ii, #6b $A_V(0, 1)$, #5b.iii.	38
8	Gorenstein Betti diagram C #6a.i, #5a.i, #5a.ii. For Betti C_j move rows $(2, 3)$ to rows $\{j - 1, j\}$	38
9	Betti diagram D, #6a.ii; # 5a iii; #5a iv; #5a vii; #4.ii; #2b i, #2bii; for Betti D_j move rows $(2, 3)$ to $\{j - 1, j\}$	38
10	Betti diagram E, #6a.iii., #5a.v, #5a.vi, 4.iii., #2b.iii. Betti E_j move rows $(2, 3)$ to rows $\{j - 1, j\}$	38
11	Betti diagram G #2a.i,ii Equation (35)	39
12	Betti diagram H #2a.iii (Lex ideal) Equation (35).	39
13	Isomorphism classes of (f, F) for $A \in G_V(T)$, $V = \langle x^2, xy, xz \rangle$ (Theorem 4.11, Equation (36)).	47
14	Dimension of isomorphism classes in $G_V(T)$, $T = (1, 3^k, 1)$, $k \geq 3$, $V = \langle x^2, xy, xz \rangle$ (Theorem 4.11)	49
15	Isomorphism classes of ideals in $G_V(T)$, $T = (1, 3^k, 1)$, $k \geq 3$, $V = \langle x^2, xy, xz \rangle$ (Theorem 4.11).	50
16	Betti diagram J_4 #2a. $(f, F) = (y^3, YZ^3)$, also $(f = yz(y - z), \text{all } F)$, and $(f, F) = (y^2z, F \neq Z^4)$. See Figure 15.	50
17	Betti diagram K_4 #2a., for $(f, F) = (y^3, Z^4)$ (the lex ideal for $T = (1, 3, 3, 3, 1)$) and (y^2z, Z^4) ; for the lex table K_j of socle degree $j \geq 4$ move the last two rows to indices $(j - 1, j)$	51

Acknowledgment. *We appreciate comments of Pedro Macias Marques, Hal Schenck, Alexandra Secealanu, and Maria Evelina Rossi. The first author would like to thank GS Magnusons Fond for financing research trips to University of Côte-D’Azur in Nice and to Northeastern University in Boston.*

References

- [AEI] N. Abdallah, J. Emsalem and A. Iarrobino: *Nets of Conics and associated Artinian algebras of length 7*. Translation and update of J. Emsalem and A. Iarrobino: *Réseaux de Coniques et algèbres de longueur sept associées*, (1977), preprint, 69p., Univ. Paris VII. To appear, European J. of Math., 2023. arXiv:math.AG/2110.04436.
- [AAIY] N. Abdallah, N. Altafi, A. Iarrobino, and J. Yaméogo: *Jordan type for codimension three Gorenstein algebras*, preprint, 2023.
- [AbSc] N. Abdallah and H. Schenck: *Free resolutions and Lefschetz properties of some Artin Gorenstein rings of codimension four*, arXiv:math.AC/2208.01536.
- [ACGH] E. Arborello, M. Cornalba, P.A. Griffiths, J. Harris: *Geometry of Algebraic Curves*, Vol I, Grundlehren der math. Wissenschaften, #267 (1985), Springer-Verlag, New York.
- [BGI] A. Bernardi, A. Gimigliano, M. Idà: *Computing symmetric rank for symmetric tensors*, J. Symbolic Comput. 46 (2011), no. 1, 34-53.
- [Bo] M. Boij: *Components of the space parametrizing graded Gorenstein Artin algebras with a given Hilbert function*, Pacific J. Math. 187 (1999), no. 1, 1-11.
- [Bri] J. Briançon: *Description de $\text{Hilb}^n C[x, y]$* , [Preprint, Univ. Nice, 1972], Invent. Math. 41 (1977), 45-89.
- [BuEi] D. Buchsbaum and D. Eisenbud: *Algebra structures for finite free resolutions, and some structure theorems for codimension three*, Amer. J. Math. **99** (1977), 447-485.
- [CJN] G. Casnati, J. Jelisiejew, and R. Notari: *Irreducibility of the Gorenstein loci of Hilbert schemes via ray families*, Algebra Number Theory 9 (2015), no. 7, 1525-1570.
- [Co] A. Conca: *Gröbner bases for spaces of quadrics of codimension 3*, J. Pure Appl. Algebra 213 (2009), no. 8, 1564-1568.

- [Di] S. J. Diesel: *Some irreducibility and dimension theorems for families of height 3 Gorenstein algebras*, Pacific J. Math. **172** (1996), 365–397.
- [Ei] D. Eisenbud: *Commutative Algebra, with a view toward Algebraic Geometry*, Springer GTM #150, (1995), Springer-Verlag, New York.
- [EL] G. Elencwajg and P. LeBarz: *L’anneau de Chow des triangles du plan*, Compositio Math. 71 (1989), no. 1, 85-119.
- [Gotz] G. Gotzmann: *Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes*, Math. Z. 158 (1978), no. 1, 61-70.
- [HW] T. Harima and J. Watanabe: *The central simple modules of Artinian Gorenstein algebras*, J. Pure Appl. Algebra 210 (2007), no. 2, 447–463.
- [I1] A. Iarrobino: *Punctual Hilbert schemes*, 111 p., Mem. Amer. Math. Soc., Vol. 10 (1977), #188, American Mathematical Society, Providence, (1977).
- [I2] A. Iarrobino: *Compressed algebras: Artin algebras having given socle degrees and maximal length*, Trans. AMS 285 (1984), 337–378.
- [I3] A. Iarrobino, *Ancestor ideals of vector spaces of forms and level algebras*, J. Algebra **272** (2004), 530–580.
- [IaKa] A. Iarrobino and V. Kanev: *Power Sums, Gorenstein Algebras, and Determinantal Varieties*, Appendix C by Iarrobino and Steven L. Kleiman *The Gotzmann Theorems and the Hilbert scheme*, Lecture Notes in Mathematics, 1721. Springer-Verlag, Berlin, 1999. xxxii+345 pp. ISBN: 3-540-66766-0 (1999), 345+xxvii p.
- [IMM] A. Iarrobino, P. Macias Marques, and C. McDaniel: *Artinian algebras and Jordan type*, J. Commut. Algebra 14 (3), 365-414, (2022)
- [IaSr] A. Iarrobino and H. Srinivasan: *Some Gorenstein Artin algebras of embedding dimension four*, components of $PGOR(H)$ for $H = (1, 4, 7, \dots, 1)$, J. of Pure and Applied Algebra **201** (2005) no. 1-3. 62–96.

- [Je] J. Jelisiejew *Classifying local Artinian Gorenstein algebras*, Collect. Math. 68 (2017), no. 1, 101-127.
- [KKRSSY] G. Kapustka, M. Kapustka, K. Ranestad, H. Schenck, M. Stillman, and B. Yuan: *Quarternary quartic forms and Gorenstein rings*, arXiv:math.AC/2111.05817.
- [Onda] N. Onda: *Isomorphism types of commutative algebras of rank 7 over an arbitrary algebraically closed field of characteristic not 2 or 3*, arXiv:math.AC/2107.04959.
- [Pe] I. Peeva: *Consecutive cancellations in Betti numbers*, Proc. Amer. Math. Soc. 132 (2004), no. 12, 3503-3507.
- [RoSh] M.E. Rossi and L. Sharifan: *Consecutive cancellations in Betti numbers of local rings* Proc. Amer. Math. Soc. 138 (2010), no. 1, 61-73.
- [Sch] P. Schenzel: *Über die freien Auflösungen extremaler Cohen-Macaulay Ringe*, J. Algebra 64 (1980), no. 1, 93-101.
- [St] R. Stanley: *Hilbert functions of graded algebras*, Advances in Math. 28 (1978), 58-73.
- [Va] E. Vasserot: *Sur l'anneau de cohomologie du schéma de Hilbert de \mathbb{C}^2* , C. R. Acad. Sci. Paris Sér. I Math. 332 (2001), no. 1, 7-12. MathSciNet review #1805619 (2001k:14012).
- [Wall] C.T.C. Wall: *Nets of conics*, Math. Proc. Cambridge Philos. Soc. 81 (1977), no. 3, 351-364.
- [Y1] J. Yaméogo: *Sur l'alignement dans les schémas de Hilbert ponctuels du plan*, Math Ann. 285, (1989), 511-525 .
- [Y2] J. Yaméogo: *Décomposition cellulaire de variétés paramétrant des idéaux homogènes de $C[[x,y]]$. Incidence des cellules. I*, Compositio Math. 90 (1) (1994), 81-98.
- [Y3] J. Yaméogo: *Décomposition cellulaire de variétés paramétrant des idéaux homogènes de $C[[x,y]]$. Incidence des cellules. II*, J. Reine Angew. Math. 450 (1994), 123-137.