

# UNIQUENESS AND HOMOGENEITY OF NON-SEPARABLE URYSOHN UNIVERSAL ULTRAMETRIC SPACES

YOSHITO ISHIKI

ABSTRACT. Urysohn constructed a separable complete universal metric space homogeneous for all finite subspaces, which is today called the Urysohn universal metric space. Some authors have recently investigated an ultrametric analogue of this space. The isometry problem of such ultrametric spaces is our main subject in this paper. We first introduce the new notion of petaloid ultrametric spaces, which is intended to be a standard class of non-separable Urysohn universal ultrametric spaces. Next we prove that all petaloid spaces are isometric to each other and homogeneous for all finite subspaces (and compact subspaces). Moreover, we show that the following spaces are petaloid, and hence they are isometric to each other and homogeneous: (1) The space of all continuous functions, whose images contain the zero, from the Cantor set into the space of non-negative real numbers equipped with the nearly discrete topology, (2) the space of all continuous ultrametrics on a zero-dimensional infinite compact metrizable space, (3) the non-Archimedean Gromov–Hausdorff space, and (4) the space of all maps from the set of non-negative real numbers into the set of natural numbers whose supports are finite or decreasing sequences convergent to the zero.

## 1. INTRODUCTION

For a class  $\mathcal{C}$  of metric spaces, a metric space  $(X, d)$  is said to be  $\mathcal{C}$ -*injective* if for all  $(A, a)$  and  $(B, b)$  in  $\mathcal{C}$  and for all isometric embeddings  $\phi: (A, a) \rightarrow (B, b)$  and  $\psi: (A, a) \rightarrow (X, d)$ , there exists an isometric embedding  $\theta: (B, b) \rightarrow (X, d)$  such that  $\theta \circ \phi = \psi$ . Let  $\mathcal{F}$  denote the class of all finite metric spaces. Urysohn [11] constructed a separable complete  $\mathcal{F}$ -injective metric space, which is today called the *Urysohn universal (metric) space*. A remarkable fact is that all separable complete  $\mathcal{F}$ -injective metric spaces are isometric to each other. This isometry theorem is proven by the so-called back-and-forth argument, which is a variant of the mathematical induction. Some authors have

---

*Date:* February 22, 2023.

*2020 Mathematics Subject Classification.* Primary 54E35, Secondary 51F99.

*Key words and phrases.* Urysohn universal ultrametric space.

recently investigated a non-Archimedean analogue of the Urysohn universal space (see [5], [12], [2], and [13]), which is also our main subject of the paper.

A metric  $d$  on a set  $X$  is said to be an *ultrametric* if for all  $x, y, z \in X$ , it satisfies the *strong triangle inequality*  $d(x, y) \leq d(x, z) \vee d(z, y)$ , where  $\vee$  stands for the maximum operator on  $\mathbb{R}$ . If a pseudo-metric  $d$  satisfies the strong triangle inequality, then  $d$  is called a *pseudo-ultrametric*.

A set  $R$  is said to be a *range set* if it is a subset of  $[0, \infty)$  and  $0 \in R$ . If an ultrametric  $d$  (resp. pseudo-ultrametric) on a set  $X$  satisfies  $d(x, y) \in R$  for all  $x, y \in X$ , then we call  $d$  an  *$R$ -ultrametric* or  *$R$ -valued ultrametrics* (resp.  *$R$ -pseudo-ultrametric* or  *$R$ -valued pseudo-ultrametrics*).

For a range set  $R$ , there are several constructions of complete  $\mathcal{N}(R)$ -injective  $R$ -valued ultrametric spaces. For example, in [5], the author provided some new constructions of complete  $\mathcal{N}(R)$ -injective  $R$ -ultrametric space (see also [12], [2], and [13]). If  $R$  is countable, then, by the back-and-forth argument, a separable complete  $\mathcal{N}(R)$ -injective  $R$ -ultrametric space is unique up to isometry whichever construction we choose. However, when  $R$  is uncountable, it was not known whether  $\mathcal{N}(R)$ -injective spaces are isometric to each other. In this paper, we solve the isometry problem of Urysohn universal ultrametric spaces in the non-separable case.

By observing and comparing constructions of  $\mathcal{N}(R)$ -injective spaces, we reach the new notion of  *$R$ -petaloid spaces*, which is intended to be a standard class of non-separable Urysohn universal spaces.

Before explaining petaloid spaces, we prepare some notations and notions. A subset  $E$  of  $[0, \infty)$  is said to be *semi-sporadic* if there exists a strictly decreasing sequence  $\{a_i\}_{i \in \mathbb{Z}_{\geq 0}}$  in  $(0, \infty)$  such that  $\lim_{i \rightarrow \infty} a_i = 0$  and  $E = \{0\} \cup \{a_i \mid i \in \mathbb{Z}_{\geq 0}\}$ . A subset of  $[0, \infty)$  is said to be *tenuous* if it is finite or semi-sporadic (see [5]). For a range set  $R$ , we denote by  $\mathbf{TEN}(R)$  the set of all tenuous range subsets of  $R$ . In this paper, we often represent a restricted metric as the same symbol to the ambient one. For a metric space  $(X, d)$ , and for a subset  $A$  of  $X$ , we write  $d(x, A) = \inf_{a \in A} d(x, a)$ .

**Definition 1.1.** Let  $R$  be an uncountable range set. We say that a metric space  $(X, d)$  is  *$R$ -petaloid* if it is an  $R$ -valued ultrametric space and there exists a family  $\{\Pi(X, S)\}_{S \in \mathbf{TEN}(R)}$  of subspaces of  $X$  satisfying the following properties:

- (P1) The set  $\Pi(X, \{0\})$  is a singleton.
- (P2) If  $S \in \mathbf{TEN}(R)$  satisfies  $S \neq \{0\}$ , then  $(\Pi(X, S), d)$  is a separable complete  $\mathcal{N}(S)$ -injective  $S$ -ultrametric space.
- (P3) We have  $\bigcup_{S \in \mathbf{TEN}(R)} \Pi(X, S) = X$ .
- (P4) If  $S, T \in \mathbf{TEN}(R)$ , then  $\Pi(X, S) \cap \Pi(X, T) = \Pi(X, S \cap T)$ .

(P5) If  $S, T \in \mathbf{TEN}(R)$  and  $x \in \Pi(X, T)$ , then  $d(x, \Pi(X, S))$  belongs to  $(T \setminus S) \cup \{0\}$ .

(P6) For all  $S \in \mathbf{TEN}(R)$ , and for all  $x \in X$ , there exists  $p \in \Pi(X, S)$  such that  $d(x, \Pi(X, S)) = d(x, p)$ .

We call the family  $\{\Pi(X, S)\}_{S \in \mathbf{TEN}(R)}$  an  $R$ -petal of  $X$ , and call  $\Pi(X, S)$  the  $S$ -piece of the  $R$ -petal  $\{\Pi(X, S)\}_{S \in \mathbf{TEN}(R)}$ . We simply write  $\Pi(S) = \Pi(X, S)$  when the whole space is clear by the context.

Section 2 presents some basic statements on petaloid spaces. In particular, it is shown that all petaloid spaces are complete.

The next is our first main result:

**Theorem 1.1.** *Let  $R$  be an uncountable range set. If  $(X, d)$  and  $(Y, e)$  are  $R$ -petaloid ultrametric spaces, then  $(X, d)$  and  $(Y, e)$  are isometric to each other. Namely, an  $R$ -petaloid ultrametric space is unique up to isometry.*

A metric space  $(X, d)$  is said to be *ultrahomogeneous* (resp. *compactly ultrahomogeneous*) if for every finite subset (resp. compact subset)  $A$  of  $X$ , and for every isometric embedding  $\phi: A \rightarrow X$ , there exists an isometric bijection  $F: X \rightarrow X$  such that  $F|_A = \phi$ . Remark that the usual Urysohn universal metric space and the separable  $\mathcal{N}(R)$ -injective complete  $R$ -ultrametric spaces are ultrahomogeneous and compactly ultrahomogeneous, which are also proven by the back-and-forth argument. For the ultrahomogeneity, see [6, Theorem 3.2] and [2, Proposition 2.7]. For the compact ultrahomogeneity, see [2, Corollary 6.18], [4], [1], and [6, Subsection 4.5].

Our second result states that all petaloid spaces satisfy the compact ultrahomogeneity despite their non-separability.

**Theorem 1.2.** *For every uncountable range set  $R$ , every  $R$ -petaloid ultrametric space is compactly ultrahomogeneous. Consequently, it is ultrahomogeneous.*

The proofs of Theorems 1.1 and 1.2 will be presented in Section 3, and both of them are reduced to Theorem 3.3 asserting that for all uncountable range set  $R$ ,  $S \in \mathbf{TEN}(R)$ , and  $R$ -petaloid spaces  $(X, d)$  and  $(Y, e)$ , every isometric bijection  $f: \Pi(X, S) \rightarrow \Pi(Y, S)$  can be extended to an isometric bijection  $F: X \rightarrow Y$ . In the proof of Theorem 3.3, to obtain an isometric bijection  $F: X \rightarrow Y$ , we construct isometric bijections between  $\Pi(X, T)$  and  $\Pi(Y, T)$  for all  $T \in \mathbf{TEN}(R)$  such that  $T \setminus S$  is finite using back-and-forth argument, and we glue them together by the transfinite induction.

In the next theorem, we show that injective spaces constructed in [5], [13], and [2] become naturally petaloid. The proof and the precise definition of the spaces will be given in Section 4.

**Theorem 1.3.** *For every uncountable range set  $R$ , all the following spaces are  $R$ -petaloid.*

- (1) The space  $(C_0(\Gamma, R), \nabla)$  of all continuous functions  $f$  from the Cantor set  $\Gamma$  into the space  $R$  equipped with the nearly discrete topology such that  $0 \in f(X)$ .
- (2) The ultrametric space  $(Cpu(X, R), \mathcal{UD}_X^R)$  of all  $R$ -valued continuous pseudo-ultrametrics on a compact ultrametrizable space  $X$  with an accumulation point.
- (3) The non-Archimedean Gromov–Hausdorff space  $(\mathcal{U}_R, \mathcal{NA})$  associated with  $R$ .
- (4) The space  $(G(R, \omega_0), \Delta)$  of all maps  $f: R \rightarrow \omega_0$  for which  $f(0) = 0$  and the support of  $f$  are tenuous, where  $\omega_0$  is the set of all natural numbers.

*Remark 1.1.* Intriguingly, it is unknown whether petaloid ultrametric spaces can be obtained by the classical constructions, i.e., the method using the Katětov function spaces, and the Urysohn-type amalgamation (the way of the Fraïssé limit).

## 2. PRELIMINARIES

From the properties (P4) or (P5), we deduce the following:

**Lemma 2.1.** *Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid space. If  $S, T \in \mathbf{TEN}(R)$  satisfy  $S \subseteq T$ , then  $\Pi(X, S) \subseteq \Pi(X, T)$ .*

The proof of the next lemma is presented in [5, Lemma 2.12].

**Lemma 2.2.** *Let  $K$  be a subset of  $[0, \infty)$ . Then  $K$  is tenuous if and only if  $K$  is a closed subset of  $[0, \infty)$  with respect to the Euclidean topology and satisfies that  $K \cap [r, \infty)$  is finite for all  $r \in (0, \infty)$ .*

By the definition of the tenuous sets, we obtain:

**Lemma 2.3.** *If  $\{S_i\}_{i=0}^k$  is a finite family of tenuous subset of  $[0, \infty)$ , then the unison  $\bigcup_{i=0}^k S_i$  is also tenuous.*

In what follows, without referring to Lemma 2.3, we use the property that the union of finite many tenuous sets is tenuous.

We now discuss the compactness of petaloid spaces. Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid ultrametric space. For all  $x \in X$ , we define the *trace*  $\text{Tr}(x)$  of  $x$  by  $\text{Tr}(x) = \bigcap \{S \in \mathbf{TEN}(R) \mid x \in \Pi(S)\}$ . Note that  $\text{Tr}(x) \in \mathbf{TEN}(R)$  and  $\text{Tr}(x) \subseteq S$  whenever  $x \in \Pi(S)$ .

**Lemma 2.4.** *Let  $R$  be an uncountable range set, and  $X$  be an  $R$ -petaloid ultrametric space. Then the following are true:*

- (1) *For all  $x \in X$ , and for all  $r \in (0, \infty)$ , there exists  $S \in \mathbf{TEN}(R)$  such that  $\text{Tr}(x) \cap (r, \infty) = S \cap (r, \infty)$  and  $x \in \Pi(S)$ .*
- (2) *For all  $x \in X$ , we have  $x \in \Pi(\text{Tr}(x))$ .*

*Proof.* First we show (1). By Lemma 2.2, the set  $\text{Tr}(x) \cap (r, \infty)$  is finite. From the definition of  $\text{Tr}(x)$ , there exists finite sets  $S_0, \dots, S_k \in \mathbf{TEN}(R)$  such that  $\text{Tr}(x) \cap (r, \infty) = \left(\bigcap_{i=0}^k S_i\right) \cap (r, \infty)$ . Put  $S = \bigcap_{i=0}^k S_i$ . According to the property (P4), the set  $S$  satisfies  $x \in \Pi(S)$ . Thus  $S$  is a desired set.

Second we verify (2). For all  $i \in \mathbb{Z}_{>0}$ , put  $t_i = 2^{-i}$ . According to the statement (1), we can find a family  $\{S_i\}_{i \in \mathbb{Z}_{>0}}$  of tenuous range subsets of  $R$  such that  $\bigcap_{i \in \mathbb{Z}_{>0}} S_i = \text{Tr}(x)$ , and for each  $i \in \mathbb{Z}_{>0}$ , we have  $x \in \Pi(S_i)$  and  $S_i \cap (t_i, \infty) = \text{Tr}(x) \cap (t_i, \infty)$ . Combining the fact that  $x \in \Pi(S_i)$  and the property (P5), we obtain  $d(x, \Pi(\text{Tr}(x))) \leq t_i$ . Since  $t_i \rightarrow 0$  as  $i \rightarrow \infty$ , we conclude that  $x \in \Pi(\text{Tr}(x))$ .  $\square$

We see the relationship between the traces and the distances.

**Lemma 2.5.** *Let  $R$  be an uncountable range set, and  $X$  be an  $R$ -petaloid ultrametric space. If  $S \in \mathbf{TEN}(R)$  and  $x \notin \Pi(S)$ , then we obtain  $d(x, \Pi(S)) = \min\{t \in \text{Tr}(x) \mid \text{Tr}(x) \cap (t, \infty) \subseteq S \cap (t, \infty)\}$ .*

*Proof.* Put  $u = d(x, \Pi(S))$ . By  $x \notin \Pi(S)$ , it is true that  $u \in \text{Tr}(x) \setminus S$ . For the sake of contradiction, suppose that  $\text{Tr}(x) \cap (u, \infty) \not\subseteq S \cap (u, \infty)$ . Then there exists  $s \in \text{Tr}(x) \cap (u, \infty)$  such that  $s \notin S$ . In particular, we have  $u < s$ . Put  $A = (\text{Tr}(x) \setminus \{s\}) \cup S$ . From the definition of  $\text{Tr}(x)$  and  $s \in \text{Tr}(x)$ , it follows that  $x \notin \Pi(A)$ . Put  $v = d(x, \Pi(A))$ . Since  $\Pi(S) \subseteq \Pi(A)$  (see Lemma 2.1), we obtain  $v \leq u$ . The property (P5) yields  $v \in \text{Tr}(x) \setminus A = \{s\}$ , and hence  $s \leq v$ . This is a contradiction to  $u < s$ . Thus we have  $\text{Tr}(x) \cap (u, \infty) \subseteq S \cap (u, \infty)$ . The minimality of  $u$  follows from  $u \in \text{Tr}(x) \setminus S$  and  $\text{Tr}(x) \cap (u, \infty) \subseteq S \cap (u, \infty)$ .  $\square$

**Corollary 2.6.** *Let  $R$  be an uncountable range set, and  $X$  be an  $R$ -petaloid ultrametric space. If  $x, y \in X$  and  $w \in R$  satisfies  $x \neq y$  and  $d(x, y) \leq w$ , then we have  $\text{Tr}(x) \cap (w, \infty) = \text{Tr}(y) \cap (w, \infty)$ .*

**Lemma 2.7.** *Let  $\{S_i\}_{i \in \mathbb{Z}_{>0}}$  be a family of tenuous range sets. If there exists a sequence  $\{t_i\}_{i \in \mathbb{Z}_{>0}}$  in  $(0, \infty)$  such that for all  $i \in \mathbb{Z}_{>0}$ , we have  $t_{i+1} \leq t_i$  and  $S_i \cap (t_i, \infty) = S_{i+1} \cap (t_i, \infty)$ , and  $t_i \rightarrow 0$  as  $i \rightarrow \infty$ , then the union  $\bigcup_{i \in \mathbb{Z}_{>0}} S_i$  is also a tenuous range set.*

*Proof.* Put  $S = \bigcup_{i \in \mathbb{Z}_{>0}} S_i$ . For all  $r \in (0, \infty)$ , take  $N \in \mathbb{Z}_{>0}$  with  $t_N < r$  and  $t_{N+1} < t_N$ . In this setting, by induction, we notice that if  $n \in \mathbb{Z}_{>0}$  satisfies  $n \geq N$ , then  $S_n \cap [r, \infty) = S_N \cap [r, \infty)$ . Thus we obtain  $S \cap [r, \infty) = \bigcup_{i=0}^N (S_i \cap [r, \infty))$ . Due to Lemma 2.2, each  $S_i \cap [r, \infty)$  is finite, and hence so is  $S \cap [r, \infty)$ . By the properties that  $0 \in S$  and  $S_i \cap [r, \infty)$  is finite for all  $r \in (0, \infty)$ , we see that the set  $S$  is closed in  $[0, \infty)$  with respect to the Euclidean topology. Using Lemma 2.2 again, the set  $S$  is tenuous.  $\square$

For a metric space  $(X, d)$ , for  $\epsilon \in (0, \infty)$  a subset  $A$  of  $X$  is said to be an  $\epsilon$ -net of  $X$  if  $A$  is finite and satisfies that for all  $x \in X$ , there

exists  $a \in A$  with  $d(x, a) \leq \epsilon$ . We say that a metric space  $(X, d)$  is *totally bounded* if for all  $\epsilon \in (0, \infty)$ , there exists an  $\epsilon$ -net of  $X$ .

**Proposition 2.8.** *Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid ultrametric space. If  $K$  is a totally bounded subset of  $X$ , then there exists  $S \in \mathbf{TEN}(R)$  such that  $K \subseteq \Pi(S)$ .*

*Proof.* For each  $n \in \mathbb{Z}_{\geq 0}$ , let  $L_n$  be a  $(2^{-n})$ -net of  $K$ . We may assume that  $L_n \subseteq L_{n+1}$  for all  $n \in \mathbb{Z}_{\geq 0}$ . Put  $T_n = \bigcup_{x \in L_n} \text{Tr}(x)$ . Then  $T_n \in \mathbf{TEN}(R)$  and  $T_n \subseteq T_{n+1}$  for all  $n \in \mathbb{Z}_{\geq 0}$ . Since  $L_n$  is a  $(2^{-n})$ -net of  $L_{n+1}$ , for each  $x \in L_{n+1}$ , there exists  $a \in L_n$  with  $d(x, a) \leq 2^{-n}$ . According to Corollary 2.6, we have  $\text{Tr}(x) \cap (2^{-n}, \infty) = \text{Tr}(a) \cap (2^{-n}, \infty)$ . Thus  $T_n \cap (2^{-n}, \infty) = T_{n+1} \cap (2^{-n}, \infty)$  for all  $n \in \mathbb{Z}_{\geq 0}$ . Put  $S = \bigcup_{n \in \mathbb{Z}_{\geq 0}} T_n$ . Therefore Lemma 2.2 shows that the set  $S$  belongs to  $\mathbf{TEN}(R)$ . From the definition of the traces, it follows that  $\bigcup_{n \in \mathbb{Z}_{\geq 0}} L_n \subseteq \Pi(S)$ . Since  $\bigcup_{n \in \mathbb{Z}_{\geq 0}} L_n$  is dense in  $K$  and  $\Pi(X, S)$  is complete (see (P2)), we can conclude that  $K \subseteq \Pi(S)$ .  $\square$

We now prove that all petaloid spaces are complete.

**Proposition 2.9.** *Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid ultrametric space. Then  $(X, d)$  is complete.*

*Proof.* Let  $\{x_i\}_{i \in \mathbb{Z}_{\geq 0}}$  be a Cauchy sequence in  $(X, d)$ . Then the set  $K = \{x_i \mid i \in \mathbb{Z}_{\geq 0}\}$  is totally bounded. Then Proposition 2.8 enables us to take  $S \in \mathbf{TEN}(R)$  such that  $K \subseteq \Pi(S)$ . Since  $\Pi(S)$  is complete, then the sequence  $\{x_i\}_{i \in \mathbb{Z}_{\geq 0}}$  has a limit. Thus  $(X, d)$  itself is complete.  $\square$

To confirm the naturality of petaloid spaces, we observe their injectivity for all finite  $R$ -ultrametric spaces.

**Proposition 2.10.** *Let  $R$  be an uncountable range set. Then all  $R$ -petaloid spaces are  $\mathcal{N}(R)$ -injective.*

*Proof.* Let  $(X, d)$  be an arbitrary  $R$ -petaloid space. Take  $(A, e)$  and  $(B, e)$  be finite metric spaces in  $\mathcal{N}(R)$  with  $A \subseteq B$ . Take an isometric embedding  $\phi: A \rightarrow X$ . Put  $S = e(B \times B) \cup \bigcup_{x \in A} \text{Tr}(\phi(x))$ . Since  $B$  is finite, we notice that  $S \in \mathbf{TEN}(R)$  and  $(B, e)$  belongs to  $\mathcal{N}(S)$ . By the definition of the trace, we also have  $\phi(A) \subseteq \Pi(S)$ . The property (P2) implies that  $\Pi(S)$  is  $\mathcal{N}(S)$ -injective, and hence there exists an isometric embedding  $F: B \rightarrow \Pi(S)$  such that  $F|_B = \phi$ . Therefore we conclude that  $(X, d)$  is  $\mathcal{N}(R)$ -injective.  $\square$

For range sets  $R$  and  $S$  with  $S \subseteq R$ , we denote by  $\mathbf{F}(R/S)$  the set of all  $T \in \mathbf{TEN}(R)$  such that  $S \subseteq T$  and  $T \setminus S$  is finite. The following lemma plays an important role in the proofs of our main theorems. The proof is deduced from Lemma 2.5 and the property (P3).

**Lemma 2.11.** *If  $R$  be an uncountable range set,  $S \in \mathbf{TEN}(R)$ , and  $(X, d)$  be an  $R$ -petaloid ultrametric space, then  $\bigcup_{T \in \mathbf{F}(R/S)} \Pi(T)$  is dense in  $X$ .*

*Proof.* Take  $x \in X$ . For all  $n \in \mathbb{Z}_{\geq 0}$ , we put  $T_n = S \cup (\text{Tr}(x) \cap [2^{-n}, \infty))$ . Then  $T_n \in \mathbf{F}(R/S)$ . By the property (P5), we have  $d(x, \Pi(X, T_n)) \leq 2^{-n}$  for all  $n \in \mathbb{Z}_{\geq 0}$ . Thus, the set  $\bigcup_{T \in \mathbf{F}(R/S)} \Pi(T)$  is dense in  $X$ .  $\square$

### 3. PROOFS OF UNIQUENESS AND HOMOGENEITY

The following lemma is a generalization of the injectivity.

**Lemma 3.1.** *Let  $R$  be an uncountable range set and  $X$  and  $Y$  be  $R$ -petaloid ultrametric spaces. Let  $k \in \mathbb{Z}_{\geq 0}$ ,  $T_0, \dots, T_k \in \mathbf{TEN}(R)$  and  $S \in \mathbf{TEN}(R)$  such that  $\bigcup_{i=0}^k T_i \subseteq S$ . Let  $A \sqcup \{\omega\}$  be a finite subset of  $\Pi(X, S)$  and  $B$  be a finite subset of  $\Pi(Y, S)$ . Put  $G = \bigcup_{i=0}^k \Pi(X, T_i)$  and  $H = \bigcup_{i=0}^k \Pi(Y, T_i)$ . If  $f: G \sqcup A \rightarrow H \cup B$  is an isometric bijection such that  $f(G) = H$  and  $f(A) = B$ , then there exists  $\theta \in \Pi(Y, S)$  for which  $d(f(x), \theta) = d(x, \omega)$  for all  $x \in G \cup A$ . Namely, we obtain an isometric bijection  $F: G \cup A \cup \{\omega\} \rightarrow H \cup B \cup \{\theta\}$  such that  $F|_{G \cup A} = f$ .*

*Proof.* If  $\omega \in G$ , then putting  $\theta = f(\omega)$  proves the lemma. We may assume that  $\omega \notin G$ . For each  $i \in \{0, \dots, k\}$ , the property (P6) enables us to take  $p_i \in \Pi(X, T_i)$  such that  $d(\omega, \Pi(X, T_i)) = d(\omega, p_i)$ . Put  $P = \{p_0, \dots, p_k\}$ ,  $Q = \{f(p_0), \dots, f(p_k)\}$ , and  $\phi = f|_{A \cup P}$ . Then  $\phi: A \cup P \rightarrow \Pi(Y, S)$  is an isometric embedding. Due to (P2), we can apply the  $\mathcal{N}(R)$ -injectivity of  $\Pi(Y, S)$  to  $\phi$ ,  $A \cup P$ , and  $(A \cup P) \cup \{\omega\}$ . Then we obtain  $\theta \in \Pi(Y, S)$  such that  $e(\phi(x), \theta) = d(x, \omega)$  for all  $x \in P \cup A$ .

We now prove that  $e(\phi(x), \theta) = d(x, \omega)$  for all  $x \in G \cup A$ . Take an arbitrary point  $x \in G \cup A$ . If  $x \in A \cup P$ , then we have  $e(f(x), \theta) = d(x, \omega)$ . If  $x \notin A \cup P$ , then we can take  $j \in \{0, \dots, k\}$  such that  $x \in \Pi(X, T_j)$ . Put  $c = d(\omega, \Pi(X, T_j)) (= d(\omega, p_j))$ . Then we have  $e(\theta, f(p_j)) = c$  and the assumption that  $\omega \notin G$  shows  $c > 0$ . The property (P5) implies that  $c \notin T_j$ . In particular, we obtain  $e(f(x), f(p_j)) \neq c$ . Recall that  $e(f(x), f(p_j)) = d(x, p_j)$ . We divide the proof into two parts.

Case 1. [ $e(f(x), f(p_j)) < c$ ]: In this case,  $e(f(x), f(p_j)) < e(\theta, f(p_j))$  and  $d(x, p_j) < d(\omega, p_j)$ . The strong triangle inequality shows that  $e(\theta, f(x)) = e(\theta, f(p_j))$  and  $d(\omega, x) = d(\omega, p_j)$ . Hence  $e(f(x), \theta) = d(x, \omega)$ .

Case 2. [ $c < e(f(x), f(p_j))$ ]: We have  $e(\theta, f(p_j)) < e(f(x), f(p_j))$  and  $d(\omega, p_j) < d(x, p_j)$ . Using the strong triangle inequality again, we also have  $e(f(x), f(p_j)) = e(f(x), \theta)$  and  $d(x, p_j) = d(x, \omega)$ . Hence  $e(f(x), \theta) = d(x, \omega)$ . This finishes the proof.  $\square$

**Lemma 3.2.** *Let  $R$  be an uncountable range set and  $X$  and  $Y$  be  $R$ -petaloid ultrametric spaces. Let  $k \in \mathbb{Z}_{\geq 0}$ ,  $T_0, \dots, T_k \in \mathbf{TEN}(R)$  and  $S \in \mathbf{TEN}(R)$  such that  $\bigcup_{i=0}^k T_i \subseteq S$ . For each  $i \in \{0, \dots, k\}$ , let  $g_i: \Pi(X, T_i) \rightarrow \Pi(Y, T_i)$  be an isometric bijection. Assume that the following condition is satisfied:*

(G) For all  $i, j \in \{0, \dots, k\}$ , we have  $g_i(x) = g_j(x)$  for all  $x \in \Pi(X, T_i \cap T_j)$ .

Then we can glue  $\{g_i\}_{i=0}^k$  together and extend it, i.e., there exists an isometric bijection  $g: \Pi(X, S) \rightarrow \Pi(Y, S)$  such that  $g|_{\Pi(X, T_i)} = g_i$  for all  $i \in \{0, \dots, k\}$ .

*Proof.* Put  $G = \bigcup_{i=0}^k \Pi(X, T_i)$  and  $H = \bigcup_{i=0}^k \Pi(Y, T_i)$ . We notice that  $\Pi(X, S) \setminus G$ , and  $\Pi(Y, S) \setminus H$  are separable and take a countable dense subsets  $A = \{a_i\}_{i \in \mathbb{Z}_{\geq 1}}$  and  $B = \{b_i\}_{i \in \mathbb{Z}_{\geq 1}}$  of  $\Pi(X, S) \setminus G$ , and  $\Pi(Y, S) \setminus H$ , respectively. We put  $A_i = \{a_1, \dots, a_i\}$  and  $B_i = \{b_1, \dots, b_i\}$  for all  $i \in \mathbb{Z}_{\geq 0}$ . We also put  $A_0 = B_0 = \emptyset$ . Define a map  $u: G \rightarrow H$  by  $u(x) = g_i(x)$  if  $x \in \Pi(X, T_i)$ . By the assumption (G), the map  $u$  is well-defined. To construct an isometric bijection  $g$ , we use the back-and-forth argument, in which we repeat the two types of operations alternately; in the first operation, we extend the domain of an isometric map, and in the second operation, we extend the codomain using the inverse map. Namely, we construct a sequence  $\{w_i: P_i \rightarrow Q_i\}_{i \in \mathbb{Z}_{\geq 0}}$  of maps such that

- (W1) for all  $i \in \mathbb{Z}_{\geq 0}$ ,  $P_i$  and  $Q_i$  are subsets of  $\Pi(X, S)$  and  $\Pi(Y, S)$ , respectively, satisfying that  $G \cup A_i \subseteq P_i$  and  $H \cup B_i \subseteq Q_i$ ;
- (W2) each  $w_i$  is an isometric bijection;
- (W3) for all  $i \in \mathbb{Z}_{\geq 0}$ , we have  $w_i|_G = u$ ;
- (W4) for all  $i \in \mathbb{Z}_{\geq 0}$ , we have  $P_i \subseteq P_{i+1}$ ,  $Q_i \subseteq Q_{i+1}$ , and  $w_{i+1}|_{P_i} = w_i$ .

First, we define  $P_0 = G$ ,  $Q_0 = H$ , and  $w_0 = u$ . Next fix  $k \in \mathbb{Z}_{\geq 0}$  and assume that we have already obtained  $P_k$ ,  $Q_k$ , and an isometric bijection  $w_k: P_k \rightarrow Q_k$  such that  $w_k|_G = u$  and  $G \cup A_i \subseteq P_i$  and  $H \cup B_i \subseteq Q_i$ . We now define  $v_{k+1}: P_k \cup \{a_{k+1}\} \rightarrow Q_k \cup \{v_{k+1}(a_{k+1})\}$ . Let  $v_{k+1}: P_k \cup \{a_{k+1}\} \rightarrow Q_k \cup \{v_{k+1}(a_{k+1})\}$  be an extended map of  $w_k$  stated in Lemma 3.1. Applying the same argument to  $v_{k+1}^{-1}$ , we obtain an isometric map  $m_{k+1}: Q_k \cup \{v_{k+1}(a_{k+1})\} \cup \{b_{k+1}\} \rightarrow P_k \cup \{a_{k+1}\} \cup \{m_{k+1}(b_{k+1})\}$ . Then we define  $w_{k+1} = m_{k+1}^{-1}$ ,  $P_{k+1} = P_k \cup \{a_{k+1}\} \cup \{m_{k+1}(b_{k+1})\}$  and  $Q_{k+1} = Q_k \cup \{v_{k+1}(a_{k+1})\} \cup \{b_{k+1}\}$ .

Therefore we obtain sequences  $\{w_i\}_{i \in \mathbb{Z}_{\geq 0}}$ ,  $\{P_i\}_{i \in \mathbb{Z}_{\geq 0}}$ , and  $\{Q_i\}_{i \in \mathbb{Z}_{\geq 0}}$  as required. Put  $K = \bigcup_{i \in \mathbb{Z}_{\geq 0}} P_i$  and  $L = \bigcup_{i \in \mathbb{Z}_{\geq 0}} Q_i$ . We define  $h: K \rightarrow L$  by  $h(x) = w_k(x)$ , where  $k \in \mathbb{Z}_{\geq 0}$  satisfies  $x \in P_k$ . Due to the conditions (W1)–(W4), the map  $h$  is well-defined. Using a canonical method by Cauchy sequences together with the completeness of  $\Pi(X, S)$  and  $\Pi(Y, S)$ , we obtain an isometric bijection  $g: \Pi(X, S) \rightarrow \Pi(Y, S)$  such that  $g|_G = u$ . Since  $h$  is an isometric bijection, and since  $K$  and  $L$  are dense in  $\Pi(X, S)$  and  $\Pi(Y, S)$ , respectively, the map  $g$  is an isometric bijection between  $\Pi(X, S)$  and  $\Pi(Y, S)$ . This proves the lemma.  $\square$

**Theorem 3.3.** *Let  $R$  be an uncountable range set, and let  $(X, d)$  and  $(Y, e)$  be  $R$ -petaloid ultrametric spaces. Let  $f: \Pi(X, S) \rightarrow \Pi(Y, S)$  be*



an isometric bijection, where  $S \in \mathbf{TEN}(R)$ . Then there exists an isometric bijection  $F: X \rightarrow Y$  such that  $F|_{\Pi(X,S)} = f$ .

*Proof.* Let  $\kappa$  be the cardinal of  $\mathbf{F}(R/S)$ . We represent  $\mathbf{F}(R/S) = \{S_\alpha\}_{\alpha < \kappa}$  such that  $R_0 = S$  and  $S_\alpha \neq S_\beta$  for all distinct  $\alpha, \beta < \kappa$ . We shall construct a family  $\{g_\alpha: \Pi(X, S_\alpha) \rightarrow \Pi(Y, S_\alpha)\}_{\alpha < \kappa}$  of isometric bijections such that

(C) If  $\alpha, \beta < \kappa$  satisfy  $S_\beta \subseteq S_\alpha$ , then  $g_\alpha|_{\Pi(X, S_\beta)} = g_\beta$ .

We use transfinite induction. If  $\alpha = 0$ , we define  $g_0 = f$ . We next consider general steps. At the  $\alpha$ -th step ( $\alpha < \kappa$ ), we will define not only  $g_\alpha$  but also  $g_\gamma$  for all  $\gamma < \kappa$  for which  $S_\gamma \subseteq S_\alpha$ . Fix  $\alpha < \kappa$  and assume that we have already passed through the  $\beta$ -th step for all  $\beta < \alpha$ . Then we walk up the  $\alpha$ -th step as follows:

Case 1. [if for all  $\beta < \alpha$ , we have  $\Pi(X, S_\alpha) \not\subseteq \Pi(X, S_\beta)$ ]: In this case, we denote by the  $\Lambda$  the set of all  $\beta < \kappa$  such that  $g_\beta$  is already defined and  $S_\beta \subseteq S_\alpha$ . Since  $S_\alpha \setminus S$  is a finite set, so is  $\Lambda$ . By the condition (C) with respect to all  $\gamma < \kappa$  for which  $g_\gamma$  is already defined, the set  $S_\alpha$ , the family  $\{S_\beta\}_{\beta \in \Lambda}$ , and the maps  $\{g_\beta\}_{\beta \in \Lambda}$  satisfy the assumptions of Lemma 3.2 (especially, the condition (G)). Then we obtain an isometric map  $g_\alpha: \Pi(X, S_\alpha) \rightarrow \Pi(Y, S_\alpha)$  such that  $g_\alpha|_{\Pi(X, S_\beta)} = g_\beta$  for all  $\beta \in \Lambda$ . Moreover, for all  $\gamma < \kappa$  with  $\alpha < \gamma$  such that  $S_\gamma \subseteq S_\alpha$ , we define  $g_\gamma = g_\beta|_{\Pi(X, S_\gamma)}$ .

Case 2. [there exists  $\beta < \kappa$  such that  $\Pi(X, S_\alpha) \subseteq \Pi(X, S_\beta)$ ]: Take a minimal ordinal  $\beta < \kappa$  such that  $S_\alpha \subseteq S_\beta$ . Then, at the  $\beta$ -th step, the map  $g_\alpha$  has been already defined by  $g_\alpha = g_\beta|_{\Pi(X, S_\alpha)}$ .

Therefore, we obtain a family  $\{g_\alpha: \Pi(X, S_\alpha) \rightarrow \Pi(Y, S_\alpha)\}_{\alpha < \kappa}$  of isometric bijections with the condition (C). Put  $K = \bigcup_{T \in \mathbf{F}(R/S)} \Pi(X, T)$  and  $L = \bigcup_{T \in \mathbf{F}(R/S)} \Pi(Y, T)$ . We define a map  $G: K \rightarrow L$  by  $G(x) = g_\alpha(x)$  if  $x \in \Pi(X, S_\alpha)$ . By the condition (C), the map  $G$  is well-defined. Lemma 2.11 shows that  $K$  and  $L$  are dense in  $X$  and  $Y$ , respectively. In a canonical way using Cauchy sequences, owing to Proposition 2.9, we can obtain  $F: X \rightarrow Y$  satisfying that  $F|_K = G$ . Then  $F$  is an isometric bijection as required.  $\square$

Using Theorem 3.3, we can prove our main results.

*Proof of Theorem 1.1.* Let  $R$  be an uncountable range set, and  $(X, d)$  and  $(Y, e)$  be  $R$ -petaloid  $R$ -ultrametric spaces. Put  $S = \{0\}$  and let  $f: \Pi(X, S) \rightarrow \Pi(X, S)$  be the trivial map (see (P1)). Applying Theorem 3.3 to  $S$  and  $f$ , we obtain an isometric bijection  $F: X \rightarrow Y$ .  $\square$

*Proof of Theorem 1.2.* Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid ultrametric space. Assume that  $A$  is a compact subset of  $X$  and  $f: A \rightarrow X$  is an isometric embedding. Using Proposition 2.8, we take  $S \in \mathbf{TEN}(R)$  with  $K \subseteq \Pi(X, S)$  and  $\phi(K) \subseteq \Pi(X, S)$ . Since  $\Pi(X, S)$  is compactly ultrahomogeneous (see [2, Corollary 6.18]), we obtain an isometric bijection  $g: \Pi(X, S) \rightarrow \Pi(X, S)$  with  $g|_K = f$ .

Theorem 3.3 implies that there exists an isometric bijection  $F: X \rightarrow X$  such that  $F|_{\Pi(S)} = g$ . The map  $F$  is a desired one, and hence the proof of Theorem 1.2 is finished.  $\square$

#### 4. EXAMPLES

We first briefly review the four constructions of injective spaces.

For a range set  $R$ , we define an ultrametric  $M_R$  on  $R$  by  $M_R(x, y) = x \vee y$  if  $x \neq y$ ; otherwise, 0. We call it the *nearly discrete (ultra)metric* and call the topology generated by  $M_R$  the *nearly discrete topology*. For a topological space  $X$  and a range set  $R$ , we denote by  $C_0(X, R)$  the set of all continuous maps  $f: X \rightarrow R$  from  $X$  to  $(R, M_R)$  such that  $0 \in f(X)$ . For  $f, g \in C(X, R)$ , we define  $\nabla(f, g)$  by the infimum of all  $\epsilon \in (0, \infty)$  such that  $f(x) \leq g(x) \vee \epsilon$  and  $g(x) \leq f(x) \vee \epsilon$  for all  $x \in X$ .

Let  $X$  be a topological space, and  $R$  be a range set, we denote by  $\text{Cpu}(X, R)$  the set of all continuous  $R$ -valued pseudo-ultrametrics  $d: X \times X \rightarrow R$  on  $X$ , where  $X \times X$  and  $R$  are equipped with the product topology and the Euclidean topology, respectively. For  $d, e \in \text{Cpu}(X, S)$ , we define  $\mathcal{UD}_X^S(d, e)$  the infimum of all  $\epsilon \in S$  such that  $d(x, y) \leq e(x, y) \vee \epsilon$  and  $e(x, y) \leq d(x, y) \vee \epsilon$  for all  $x, y \in X$ . For more details on  $C_0(X, R)$  and  $\text{Cpu}(X, R)$ , we refer the readers to [5].

For a range set  $R$ , we denote by  $\mathcal{U}_R$  the set of all isometry classes of compact  $R$ -ultrametric spaces and denote by  $\mathcal{NA}$  the non-Archimedean Gromov–Hausdorff distance on  $\mathcal{U}_R$ , i.e., the value  $\mathcal{NA}((X, d), (Y, e))$  is the infimum of all  $\mathcal{HD}(i(X), j(Y); Z, h)$ , where  $(Z, h)$  is an  $R$ -valued ultrametric space,  $\mathcal{HD}(*, *; Z, h)$  is the Hausdorff distance associated with  $(Z, h)$ , and  $i: X \rightarrow Z$  and  $j: Y \rightarrow Z$  are isometric embedding. We call  $(\mathcal{U}_R, \mathcal{NA})$  the Gromov–Hausdorff  $R$ -ultrametric space (For more details, see [14], [10] and [13]).

We denote by  $\omega_0$  the set of all non-negative integers. Let  $R$  be a range set. We also denote by  $G(R, \omega_0)$  the set of all function  $f: R \rightarrow \omega_0$  such that  $f(0) = 0$  and the set  $\{0\} \cup \{x \in R \mid f(x) \neq 0\}$  is tenuous. For  $f, g \in G(R, \omega_0)$ , we define an  $R$ -ultrametric  $\Delta$  on  $G(R, \omega_0)$  by  $\Delta(f, g) = \max\{r \in R \mid f(r) \neq g(r)\}$  if  $f \neq g$ ; otherwise,  $\Delta(f, g) = 0$ . For more information, we refer the readers to [2] and [9].

**Theorem 4.1.** *For every range set  $R$ , the following spaces are all  $\mathcal{N}(R)$  and complete. Moreover, if  $R$  is finite or countable, they are separable.*

- (1) *The space  $(C_0(\Gamma, R), \nabla)$  of all continuous functions from the Cantor set  $\Gamma$  into  $(R, M_R)$ .*
- (2) *The ultrametric space  $(\text{Cpu}(X, R), \mathcal{UD}_X^R)$  of all  $R$ -valued continuous pseudo-ultrametrics on a compact ultrametrizable space  $X$  with an accumulation point.*
- (3) *The Gromov–Hausdorff  $R$ -ultrametric space  $(\mathcal{U}_R, \mathcal{NA})$ .*
- (4) *The  $R$ -ultrametric space  $(G(R, \omega_0), \Delta)$ .*

*Proof.* The cases (1) and (2) are proven in [5, Theorem 3.2 and Theorem 4.2], respectively. For the separable part, see [5, Proposition 3.5 and Proposition 4.3]. The case (3) is proven by Wan [13]. The case (4) can be found in [2, Section 6], [9, Proposition 11] and [8, Subsection 1.3].  $\square$

We now prove that all the spaces described above are petaloid.

*Proof of Theorem 1.3.* Let  $R$  be an uncountable range set, and  $X$  be a compact ultrametrizable space with an accumulation point. Let  $\Gamma$  stands for the Cantor set. For every  $S \in \mathbf{TEN}(R)$ , we define the petals of the spaces stated in the theorem by  $\Pi(C_0(X, R), S) = C_0(X, S)$ ,  $\Pi(\text{Cpu}(X, R), S) = \text{Cpu}(X, S)$ ,  $\Pi(\mathcal{U}_R, S) = \mathcal{U}_S$ , and  $\Pi(G(R, \omega_0), S) = G(S, \omega_0)$ . Then the families satisfy the properties (P1), (P3), and (P4). Theorem 4.1 implies that they enjoy the property (P2).

We now consider the case of  $(G(R, \omega_0), \Delta)$ . To verify the properties (P5) and (P6), we need the following claim that is deduced from the definition of  $\Delta$ :

**Claim 4.2.** *Let  $R$  be a range set. Let  $r \in R \setminus \{0\}$ , and  $f, g \in G(R, \omega_0)$ . Then  $r = \Delta(f, g)$  if and only if  $f(r) \neq g(r)$  and  $f(x) = g(x)$  whenever  $r < x$ .*

Let  $S \in \mathbf{TEN}(R)$ . For every  $f \in G(R, \omega_0)$ , put  $T = \{0\} \cup \{x \in R \mid f(x) \neq 0\} \in \mathbf{TEN}(R)$ . If  $T \subseteq S$ , then  $\Delta(f, C_0(X, S)) = \Delta(f, f) = 0$ . If  $T \not\subseteq S$ , then let  $r$  be the maximum of  $T \setminus S$ . Thus we have  $T \cap (r, \infty) \subseteq S \cap (r, \infty)$ . We define  $g \in G(R, \omega_0)$  by

$$g(x) = \begin{cases} f(x) & \text{if } r < x; \\ f(r) + 1 & \text{if } x = r; \\ 0 & \text{otherwise.} \end{cases}$$

Then by Claim 4.2, we have  $r = \Delta(f, g)$  and  $\Delta(f, g) = \Delta(f, C_0(X, S))$ . This proves the property (P5), and also proves (P6).

Remark that using the statements corresponding to Claim 4.2 such as [5, Corollary 2.17], [5, Corollary 2.30], and [7, Theorem 5.1], in a similar way to the case of  $(G(R, \omega_0), \Delta)$ , we can prove that the spaces  $(C_0(X, R), \nabla)$ ,  $(\text{Cpu}(X, R), \mathcal{UD}_X^R)$ , and  $(\mathcal{U}_R, \mathcal{NA})$  satisfy (P5) and (P6), respectively, and hence they are  $R$ -petaloid. Since there is a small gap, we give a little explanation of the case of  $(G(R, \omega_0), \Delta)$ , for example. Let  $S \in \mathbf{TEN}(R)$ , and  $f \in G(R, \omega_0)$ , and put  $T = f(\Gamma) \in \mathbf{TEN}(R)$ . If  $T \subseteq S$ , then put  $g = f$ . If  $T \not\subseteq S$ , then we take non-empty clopen subsets  $A$  and  $B$  of  $X$  such that  $A \cap B = \emptyset$  and  $A \cup B = f^{-1}([0, r])$ . We then define  $g \in \Pi(G(R, \omega_0), S)$  by

$$g(x) = \begin{cases} f(x) & \text{if } r < f(x); \\ r & \text{if } x \in A; \\ 0 & \text{if } x \in B. \end{cases}$$

Thus, by [5, Corollary 2.17], we have  $\nabla(f, g) = \nabla(f, \Pi(G(R, \omega_0), S))$ .

This completes the proof of Theorem 1.3.  $\square$

**Question 4.3.** Let  $R$  be a range set, and  $\Gamma$  be the Cantor set. In this setting, we see that  $(\mathcal{U}_R, \mathcal{NA})$  and  $(\text{Cpu}(\Gamma, R), \mathcal{UD}_X^R)$  are isometric to each other by Theorem 1.1. Does there exist a natural isometric bijection between them?

**Question 4.4.** Let  $R$  be an uncountable range set, and  $(X, d)$  be an  $R$ -petaloid space. We denote by  $\text{Isom}(X, d)$  the group of isometric bijections from  $X$  into itself. What is the isometry group  $\text{Isom}(X, d)$ ? Namely, what interesting properties does  $\text{Isom}(X, d)$  satisfy?

**Question 4.5.** Let  $(X, d)$  and  $(Y, e)$  be complete  $\mathcal{N}(R)$ -injective  $R$ -valued ultrametric spaces with the same topological weight. In this setting, are  $(X, d)$  and  $(Y, e)$  isometric to each other? If not, is there a sufficient and necessary condition for isometry? The author thinks that we can find counterexamples since there exists a complete  $\mathcal{F}$ -injective metric space  $(X, d)$  such that  $\text{Isom}(X, d)$  is trivial (see [3]).

#### REFERENCES

1. S. A. Bogatyĭ, *Compact homogeneity of the universal metric Uryson space*, *Uspekhi Mat. Nauk* **55** (2000), no. 2(332), 131–132.
2. S. Gao and C. Shao, *Polish ultrametric Urysohn spaces and their isometry groups*, *Topology Appl.* **158** (2011), no. 3, 492–508.
3. J. Grebik, *A rigid Urysohn-like metric space*, *Proc. Amer. Math. Soc.* **145** (2017), no. 9, 4049–4060.
4. G. E. Huhunaišvili, *On a property of Uryson's universal metric space*, *Dokl. Akad. Nauk SSSR (N.S.)* **101** (1955), 607–610.
5. Y. Ishiki, *Constructions of Urysohn universal ultrametric spaces*, (2023), arXiv:2302.00305.
6. J. Melleray, *Some geometric and dynamical properties of the Urysohn space*, *Topology Appl.* **155** (2008), no. 14, 1531–1560.
7. F. Mémoli and Z. Wan, *On  $p$ -metric spaces and the  $p$ -Gromov-Hausdorff distance*, *p-Adic Numbers Ultrametric Anal. Appl.* **14** (2022), no. 3, 173–223.
8. Y. A. Neretin, *Groups of isometries of ultrametric urysohn space and their unitary representations*, (2022), preprint, arXiv:2212.02607.
9. L. Nguyen Van Thé, *Structural Ramsey theory of metric spaces and topological dynamics of isometry groups*, *Mem. Amer. Math. Soc.* **206** (2010), no. 968, 155 pages.
10. D. Qiu, *Geometry of non-Archimedean Gromov-Hausdorff distance*, *p-Adic Numbers Ultrametric Anal. Appl.* **1** (2009), no. 4, 317–337.
11. P. Urysohn, *Sur un espace métrique universel*, *Bull. Sci. Math.* **51** (1927), 43–64 and 74–90.
12. I. A. Vestfrid, *On a universal ultrametric space*, *Ukrain. Mat. Zh.* **46** (1994), no. 12, 1700–1706.
13. Z. Wan, *A novel construction of Urysohn universal ultrametric space via the Gromov-Hausdorff ultrametric*, *Topology Appl.* **300** (2021), Paper No. 107759.
14. I. Zarichnyi, *Gromov-Hausdorff ultrametric*, (2005), arXiv:math/0511437.

(Yoshito Ishiki)

PHOTONICS CONTROL TECHNOLOGY TEAM  
RIKEN CENTER FOR ADVANCED PHOTONICS  
2-1 HIRASAWA, WAKO, SAITAMA 351-0198, JAPAN  
*Email address:* [yoshito.ishiki@riken.jp](mailto:yoshito.ishiki@riken.jp)