

Structure theorem for mod p^m singular Siegel modular forms

Siegfried Böcherer and Toshiyuki Kikuta

February 2, 2023

2020 Mathematics subject classification: Primary 11F33 · Secondary 11F46

Key words: Siegel modular forms, Congruences for modular forms, Fourier coefficients, mod p^m singular.

Abstract

We prove that all mod p^m singular forms of level N , degree $n + r$, and p -rank r with $n \geq r$ are congruent mod p^m to linear combinations of theta series of degree r attached to quadratic forms of some level. Moreover, we prove that, the levels of theta series are of the form “ p -power $\times N$ ”. Additionally, in some cases of mod p singular forms with smallest possible weight, we prove that the levels of theta series should be p .

1 Introduction

Freitag [10] showed the following properties concerning singular (Siegel modular) forms.

- A Siegel modular form of weight k and degree n is singular if and only if $k < n/2$.
- The weight k of a singular form with singular rank r should be $k = r/2$.
- All singular modular forms of singular rank r are linear combinations of theta series attached to quadratic forms of matrix size r .

We should also mention here that Resnikoff [21] and much later Shimura [24] considered singular modular forms on more general tube domains.

In [4], the authors defined a notion of mod p^m singular form and proved that a congruence holds between the weight k and the mod p singular rank r (we call it p -rank). More precisely we proved that $2k - r \equiv 0 \pmod{(p-1)p^{m-1}}$. This can be regarded as a mod p^m analogue to Freitag’s result concerning weights. We also mentioned in the same paper that we can construct some examples from any true singular modular forms and also from some Siegel-Eisenstein series of level 1.

Another type of examples is provided by the work of Nagaoka [18] and Katsurada-Nagaoka [14], who studied p -adic limits of Siegel-Eisenstein series and showed in some cases that they are equal to certain genus theta series of singular rank. In particular, some of the Siegel-Eisenstein series studied by them are then congruent to linear combinations of singular theta series, i.e. true singular forms. Our conjecture is that, all mod p^m singular forms are congruent mod p^m to true singular forms with some level, i.e., linear combinations of theta series with singular weights ($< \frac{n}{2}$) and some level.

In this paper, we mainly discuss this conjecture and prove it for many cases, in particular for “strongly” mod p^m singular forms: More precisely, we prove the following properties.

- All mod p^m singular forms of level N , degree $n+r$, and p -rank r with $n \geq r$ are congruent mod p^m to linear combinations of theta series of degree r attached to quadratic forms of some level (Theorem 3.2 (1)).
- We show that the levels of the theta series are of the form $p^e N$ with some $e \in \mathbb{N}$ (Theorem 3.2 (2)).
- In the case where the weights are the smallest possible in some sense, then the levels of theta series should be p ; this is proved for the case of mod p singular forms of p -rank 2 (Theorem 3.6).

Of course, our results can be regarded as the congruence version of Freitag’s results. However, there are many new difficulties in mod p cases. For example, if a degree n theta series for a quadratic form S is congruent mod p to a level one modular form, then this does not specify the level of S . This is one of the main differences to the case over \mathbb{C} . We can use the filtration (weight) of the theta series as a kind of substitute, but this has rather weak properties. As new tools we use a strong version of the q -expansion principle and a degree n version of Kitaoka’s transformation formula for theta series. These are shown within appendices of this paper and may be of independent interest.

We remark that we chose to consider only modular forms for groups of type $\Gamma_0(N)$ and quadratic nebentypus character. This allows us to consider congruences modulo rational primes (not modulo prime ideals in suitable algebraic number fields); also we wanted to avoid some delicate issues concerning congruences for Siegel modular forms of nonquadratic nebentypus. Instead, we show at the end of the main text of this paper how some of our results can be extended to the setting of mod p^m singular forms for arbitrary m .

This paper is organized as follows. In Section 2, we fix notation and definitions, and review known facts. In Section 3, we explain our conjectures and main results of this paper. In Section 4, we provide one of our main tools: An expansion of the rank r -part of the Fourier expansion of any modular form and its formulation for mod p^m singular forms of p -rank r . Then in Section 5 we focus on the mod p case. We first give a kind of Sturm bound for mod p singular forms and use it to show that every strongly mod p singular modular form is represented by a linear combination

of theta series for some quadratic forms. In Section 6, we try to specify the levels of their quadratic forms. We discuss also mod p singular forms with p -rank 2 whose weights are the smallest possible. In Section 7, we extend the results for the mod p case (precisely, results for general strongly mod p singular forms) proved in Section 5 and 6 to the mod p^m case by induction on m . Section 8 and 9 are appendices: We provide two tools that are needed for specification of the levels that we do in Section 6, namely a modified version of the q -expansion principle, and an extension to degree n of Kitaoka's transformation formula for theta series.

2 Preliminaries

2.1 Siegel modular forms

Let \mathbb{H}_n be the Siegel upper half space of degree n . We put

$$\mathrm{GSp}_n^+(\mathbb{R}) := \{g \in \mathrm{GL}_{2n}(\mathbb{R}) \mid {}^t g J_n g = n(g) J_n \text{ for some } n(g) > 0\},$$

where $J_n := \begin{pmatrix} 0_n & 1_n \\ -1_n & 0_n \end{pmatrix}$. We define the action of $\mathrm{GSp}_n^+(\mathbb{R})$ on \mathbb{H}_n by $gZ = (AZ + B)(CZ + D)^{-1}$ for $Z \in \mathbb{H}_n$, $g \in \mathrm{GSp}_n^+(\mathbb{R})$. For a holomorphic function $F : \mathbb{H}_n \rightarrow \mathbb{C}$ and a matrix $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{GSp}_n^+(\mathbb{R})$, we define the slash operator in the usual way;

$$(F|_k g)(Z) := n(g)^{\frac{nk}{2}} \det(CZ + D)^{-k} F(gZ).$$

Let N be a natural number and $\Gamma_n := \mathrm{Sp}_n(\mathbb{Z})$ the Siegel modular group (symplectic group with components in \mathbb{Z}). In this paper, we deal mainly with the congruence subgroup $\Gamma_0^{(n)}(N)$ with level N of Γ_n defined as

$$\Gamma_0^{(n)}(N) := \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_n \mid C \equiv 0_n \pmod{N} \right\}.$$

We will also use the groups

$$\begin{aligned} \Gamma_1^{(n)}(N) &:= \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_n \mid C \equiv 0_n \pmod{N}, \det A \equiv \det D \equiv 1 \pmod{N} \right\}, \\ \Gamma^{(n)}(N) &:= \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_n \mid B \equiv C \equiv 0_n \pmod{N}, A \equiv D \equiv 1_n \pmod{N} \right\}, \end{aligned}$$

where $\Gamma^{(n)}(N)$ is the so-called principal congruence subgroup of level N . We remark that

$$\Gamma^{(n)}(N) \subset \Gamma_1^{(n)}(N) \subset \Gamma_0^{(n)}(N) \subset \Gamma_n.$$

For a natural number k and a Dirichlet character $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$, the space $M_k(\Gamma_0^{(n)}(N), \chi)$ of Siegel modular forms of weight k with character χ consists of all of holomorphic functions $F : \mathbb{H}_n \rightarrow \mathbb{C}$ satisfying

$$(F|_k g)(Z) = \chi(\det D) F(Z) \quad \text{for } g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_0^{(n)}(N).$$

If $n = 1$, the usual condition in the cusps should be added. When χ is a trivial character, we write simply $M_k(\Gamma_0^{(n)}(N))$ for $M_k(\Gamma_0^{(n)}(N), \chi)$.

Let $\Gamma \supset \Gamma^{(n)}(N)$. Similarly as above, we denote by $M_k(\Gamma)$ the space consists of all of holomorphic functions $F : \mathbb{H}_n \rightarrow \mathbb{C}$ satisfying

$$(F|_k g)(Z) = F(Z) \quad \text{for } g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma.$$

In this case also, we have to add the usual condition in the cusps when $n = 1$.

For a prime p , let Γ be $\Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N)$ or $\Gamma_0^{(n)}(p^m) \cap \Gamma_1^{(n)}(N)$, and χ a Dirichlet character mod p . In this paper, symbols $M_k(\Gamma, \chi)$ and $M_k(\Gamma)$ also for such Γ are sometimes used, but these are defined in the same way as above.

Note that, for any Dirichlet character χ mod N , we have

$$M_k(\Gamma^{(n)}(N)) \supset M_k(\Gamma_1^{(n)}(N)) \supset M_k(\Gamma_0^{(n)}(N), \chi).$$

When $F \in M_k(\Gamma)$ with $\Gamma \supset \Gamma^{(n)}(N)$, N is called the ‘‘level’’ of F . Sometimes Γ itself is also called the level of F .

Any $F \in M_k(\Gamma^{(n)}(N))$ has a Fourier expansion of the form

$$F(Z) = \sum_{0 \leq T \in \frac{1}{N}\Lambda_n} a_F(T) e(\text{tr}(TZ)), \quad Z \in \mathbb{H}_n,$$

where $e(x) := e^{2\pi i x}$,

$$\Lambda_n := \{T = (t_{ij}) \in \text{Sym}_n(\mathbb{Q}) \mid t_{ii}, 2t_{ij} \in \mathbb{Z}\},$$

and $\text{Sym}_n(R)$ is the set of symmetric matrices of size n with components in R . In particular, if $F \in M_k(\Gamma_1^{(n)}(N))$, the Fourier expansion of F is given in the form

$$F(Z) = \sum_{0 \leq T \in \Lambda_n} a_F(T) e(\text{tr}(TZ)).$$

It is known that

$$M_k(\Gamma_1^{(n)}(N)) = \bigoplus_{\chi: (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times} M_k(\Gamma_0^{(n)}(N), \chi),$$

where χ runs over all the Dirichlet characters mod N . We remark that, if $F \in M_k(\Gamma_0^{(n)}(N), \chi)$, then we have

$$a_F(T[U]) = (\det U)^k \chi(\det U) a_F(T)$$

for each $T \in \Lambda_n$ and $U \in \text{GL}_n(\mathbb{Z})$. Here we write as $T[U] := {}^t U T U$. In particular, if $\chi(-1) = (-1)^k$, we have $a_F(T[U]) = a_F(T)$ for each $T \in \Lambda_n$ and $U \in \text{GL}_n(\mathbb{Z})$.

Let Φ be the Siegel Φ -operator defined by

$$\Phi(F)(Z') := \lim_{t \rightarrow \infty} F \begin{pmatrix} Z' & 0 \\ 0 & it \end{pmatrix},$$

where $F \in M_k(\Gamma_0^{(n)}(N), \chi)$, $Z' \in \mathbb{H}_{n-1}$, and $t \in \mathbb{R}$. As is well-known, we have $\Phi(F) \in M_k(\Gamma_0^{(n-1)}(N), \chi)$ and the Fourier expansion of $\Phi(F)$ is described as

$$\Phi(F)(Z') = \sum_{0 \leq T \in \Lambda_{n-1}} a_F \begin{pmatrix} T & 0 \\ 0 & 0 \end{pmatrix} e(\text{tr}(TZ')).$$

Suppose that $F \in M_k(\Gamma_0^{(n)}(N), \chi)$ satisfies $\Phi(F) \neq 0$ (and then $F \neq 0$). Taking $g \in \Gamma_0^{(n)}(N)$ as $g = \begin{pmatrix} -1_n & 0_n \\ 0_n & -1_n \end{pmatrix}$, we have $F|_k g = (-1)^{nk} F = \chi(-1)^n F$ because of the transformation law of F . Therefore we have $\chi(-1)^n = (-1)^{nk}$. On the other hand, by the same property of $\Phi(F) \neq 0$, we have $\chi(-1)^{n-1} = (-1)^{(n-1)k}$. These imply that $\chi(-1) = (-1)^k$. In this case (of $\Phi(F) \neq 0$), we have automatically $a_F(T[U]) = a_F(T)$ for each $T \in \Lambda_n$ and $U \in \text{GL}_n(\mathbb{Z})$.

For a subring R of \mathbb{C} , let $M_k(\Gamma, \chi)_R$ (resp. $M_k(\Gamma)_R$) denote the R -module of all modular forms in $M_k(\Gamma, \chi)$ (resp. $M_k(\Gamma)$) whose Fourier coefficients are in R .

2.2 Congruences for modular forms

Let p be a prime and $\mathbb{Z}_{(p)}$ the set of p -integral rational numbers. Let F_i ($i = 1, 2$) be two formal power series of the form

$$F_i = \sum_{T \in \frac{1}{N}\Lambda_n} a_{F_i}(T) e(\text{tr}(TZ))$$

with $a_{F_i}(T) \in \mathbb{Z}_{(p)}$ for all $T \in \frac{1}{N}\Lambda_n$. We write $F_1 \equiv F_2 \pmod{p^m}$ if $a_{F_1}(T) \equiv a_{F_2}(T) \pmod{p^m}$ for all $T \in \frac{1}{N}\Lambda_n$.

Let $\widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$ be the set of $\widetilde{F} = \sum_T \widetilde{a_F}(T) e(\text{tr}(TZ))$ with $F \in M_k(\Gamma_0^{(n)}(N), \chi)_{\mathbb{Z}_{(p)}}$, where $\widetilde{a_F}(T) := a_F(T) \pmod{p^m}$. If $m = 1$, we write simply $\widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)$ for $\widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$. Note that, $\widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)$ is a vector space over \mathbb{F}_p .

We define the filtration weight as

$$\omega_{N, \chi, p^m}^n(F) := \min\{k \mid \widetilde{F} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}\}.$$

We write also $\omega_{N, \chi}^n(F) := \omega_{N, \chi, p}^n(F)$.

Definition 2.1. Let $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}_{(p)}}$. We say that F is “*mod p^m singular*” if

- we have $a_F(T) \equiv 0 \pmod{p^m}$ for any $T \in \Lambda_{n+r}$ with $\text{rank}(T) > r$,

- there exists $T \in \Lambda_{n+r}$ with $\text{rank}(T) = r$ satisfying $a_F(T) \not\equiv 0 \pmod{p}$.

We call such r “ p -rank” of F . Additionally if $n \geq r$, we say that such F is “strongly mod p^m singular”.

Remark 2.2. (1) Such a condition “strongly singular” also plays a crucial role in Freitag’s book [10] for the theory over \mathbb{C} for arbitrary congruence subgroups.

- (2) If F is nontrivial mod p^m singular, then F satisfies $\Phi(F) \neq 0$. This implies that $\chi(-1) = (-1)^k$ and $a_F(T[U]) = a_F(T)$ for each $T \in \Lambda_n$ and $U \in \text{GL}_n(\mathbb{Z})$ (see Page 4). In other words, when dealing with nontrivial mod p^m singular modular forms, we may assume that $\chi(-1) = (-1)^k$.

Theorem 2.3 (Böcherer-Kikuta [4]). *Let n, r, k, N be positive integers and p a prime with $p \geq 5$. Let χ be a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}(p)}$ is mod p^m singular of p -rank r . Then we have $2k - r \equiv 0 \pmod{(p-1)p^{m-1}}$. In particular, r should be even.*

It is a classical result by Serre [23] that a congruence mod p for two elliptic modular forms f and g for level one implies a congruence of their weights mod $p-1$. We need a version for degree n including levels and quadratic nebentypus:

Proposition 2.4. *Let p be a prime with $p \geq 5$ and N a positive integer with $p \nmid N$. Let χ and χ' be two quadratic Dirichlet characters mod p and $F \in M_k(\Gamma_0^{(n)}(p^m) \cap \Gamma_1^{(n)}(N), \chi)$, $F' \in M_{k'}(\Gamma_0^{(n)}(p^m) \cap \Gamma_1^{(n)}(N), \chi')$ be two modular forms satisfying $F \equiv F' \pmod{p}$. Then $k - k' = t \cdot \frac{p-1}{2}$ holds for some $t \in \mathbb{Z}$ and we have*

$$\chi = \chi' \iff t \text{ even.}$$

Proof. We want to apply the results from Böcherer-Nagaoka [5] to get the desired congruences for the weights. Note that in [5] only the case of level $\Gamma_1^{(n)}(N)$ with N coprime to p is covered. We may apply level change to F and F' to arrive at G and G' of level $\Gamma_1^{(n)}(N)$ with $F \equiv G \pmod{p}$ and $G' \equiv F' \pmod{p}$ with weights l and l' .

If $\chi = \chi'$, we have $k \equiv l \pmod{p-1}$ and $k' \equiv l' \pmod{p-1}$ and then, by [5] $k \equiv k' \pmod{p-1}$, i.e. t is even.

If $\chi \neq \chi'$, let us assume that χ is nontrivial. Then $l \equiv k + \frac{p-1}{2} \pmod{p-1}$ and the congruence $l \equiv l' \pmod{p-1}$ implies that t has to be odd. This completes the proof. \square

To formulate our results efficiently, we introduce the following (somewhat non-standard).

Definition 2.5. *Suppose that $k - k' = t \cdot \frac{p-1}{2}$ holds for some $t \in \mathbb{Z}$. For a prime p and a natural number N coprime to p let χ and χ' be two quadratic Dirichlet characters mod pN . We write*

$$\chi =^t \chi'$$

if $\chi_N = \chi'_N$, and χ_p and χ'_p are related as in Proposition 2.4; i.e.

$$\chi_p = \chi'_p \iff t \text{ even.}$$

Here χ_N and χ_p are the N -component and p -component of χ (and the same for χ'). In other words, we have

$$\chi = \chi' \iff \chi = \chi' \left(\frac{*}{p} \right)^t,$$

where $\left(\frac{*}{p} \right)$ is the unique nontrivial quadratic character mod p .

Note that this notation depends on k, k' but it will always be clear from the context, which weights are involved.

2.3 Theta series for quadratic forms

Let m be a positive integer. For $S, T \in \Lambda_m$, we write $S \sim T \pmod{\mathrm{GL}_m(\mathbb{Z})}$ if there exists $U \in \mathrm{GL}_m(\mathbb{Z})$ such that $S[U] = T$. Here we put $S[U] := {}^tUSU$. We say that S and T are “ $\mathrm{GL}_m(\mathbb{Z})$ -equivalent” if $S \sim T \pmod{\mathrm{GL}_m(\mathbb{Z})}$. We denote by Λ_m^+ the set of all positive definite elements of Λ_m . We put $L := \Lambda_m$ or Λ_m^+ . We write $L/\mathrm{GL}_m(\mathbb{Z})$ for L/\sim the set of representatives of $\mathrm{GL}_m(\mathbb{Z})$ -inequivalence classes in L .

Let m be even. For $S \in \Lambda_m^+$, we define the theta series of degree n in the usual way:

$$\theta_S^{(n)}(Z) := \sum_{X \in \mathbb{Z}^{m,n}} e(\mathrm{tr}(S[X]Z)) \quad (Z \in \mathbb{H}_n),$$

where $\mathbb{Z}^{m,n}$ is the set of $m \times n$ matrices with integral components and $S[X] := {}^tXSX$ and we write $e(x) := e^{2\pi ix}$. We define the level of S as

$$\mathrm{level}(S) := \min\{N \in \mathbb{Z}_{\geq 1} \mid N(2S)^{-1} \in 2\Lambda_m\}.$$

Then $\theta_S^{(n)}$ defines an element of $M_{\frac{m}{2}}(\Gamma_0^{(n)}(N), \chi_S)$, where $N = \mathrm{level}(S)$, χ_S is a Dirichlet character mod N defined by

$$\chi_S(d) = \mathrm{sign}(d)^{\frac{m}{2}} \left(\frac{(-1)^{\frac{m}{2}} \det 2S}{|d|} \right).$$

We denote by $\mathrm{cont}(S)$ the content of S defined as

$$\mathrm{cont}(S) := \max\{C \in \mathbb{Z}_{\geq 1} \mid C^{-1}S \in \Lambda_n\}.$$

For fixed $S \in \Lambda_m^+$ and $T \in \Lambda_n$, we put

$$A(S, T) := \#\{X \in \mathbb{Z}^{m,n} \mid S[X] = T\}.$$

Using this notation, we can write the Fourier expansion of the theta series in the form

$$\theta_S^{(n)}(Z) = \sum_{T \in \Lambda_n} A(S, T) e(\mathrm{tr}(TZ)).$$

In the above definitions, S and T are restricted to symmetric half integral matrices, but the same symbols (such as $\theta_S^{(n)}(Z)$ and $A(S, T)$) are used for symmetric matrices with rational components. Then $A(S, S)$ is the order of automorphism group of S ;

$$A(S, S) = \#\{U \in \mathrm{GL}_m(\mathbb{Z}) \mid S[U] = S\}.$$

By looking at minimal polynomials, one sees that $\mathrm{GL}_m(\mathbb{Z})$ cannot contain elements of order p if $p > m + 1$. From this we obtain the very useful statement that

$$A(S, S) \not\equiv 0 \pmod{p} \quad \text{if } p > m + 1 \tag{2.1}$$

for any rational positive definite symmetric matrix S of size m .

For later use, we introduce some results on the representation numbers for binary quadratic forms:

Theorem 2.6 (Dirichlet, Weber (see Kani [13], Lemma 8, page 4)). *Let $T, T_i \in \Lambda_2$ ($i = 1, 2$) be primitive forms, i.e. $\mathrm{cont}(T) = \mathrm{cont}(T_i) = 1$. Assume that T_1 and T_2 have a same discriminant $D < 0$. Then we have the following statements.*

- (1) *There are infinitely many primes l such that $A(T, l) > 0$.*
- (2) *If there exists a prime l with $l \nmid D$ such that $A(T_1, l) > 0$ and $A(T_2, l) > 0$, then $T_1 \sim T_2 \pmod{\mathrm{GL}_2(\mathbb{Z})}$.*

Theorem 2.7 (Böcherer-Nagaoka [7]). *Let $S \in \Lambda_2^+$ be of level p . Let n be a positive integer. Then there exists $F \in M_{\frac{p+1}{2}}(\Gamma_n)_{\mathbb{Z}(p)}$ such that $F \equiv \theta_S^{(n)} \pmod{p}$.*

For a quadratic form $S \in \Lambda_m^+$, we put $\omega_{N, \chi, p^m}^n(S) := \omega_{N, \chi, p^m}^n(\theta_S^{(n)})$, $\omega_{N, \chi}^n(S) := \omega_{N, \chi, p}^n(\theta_S^{(n)})$.

3 Conjectures and Results

We begin by stating our conjecture in the most general situation which does not specify the weight and level.

Conjecture 3.1. *Any mod p^m singular form is congruent mod p^m to some true singular form.*

Our first result states that this conjecture is true in the strongly mod p^m singular case (if p is not small).

Theorem 3.2. *Let n, k, N be positive integers, r an even integer with $n \geq r$. Let p be a prime with $p > r + 1$ and χ a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}(p)}$ is mod p^m singular of p -rank r . Then we have the following statements.*

(1) There are finitely many $S \in \Lambda_r^+$ such that

$$F \equiv \sum_S c_S \theta_S^{(n+r)} \pmod{p^m} \quad (c_S \in \mathbb{Z}_{(p)})$$

and $\widetilde{\theta_S^{(n)}} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^{m-\nu}}$ (and hence $\omega_{N, \chi, p^{m-\nu}}^n(S) \leq k$). Here $\nu := \nu_p(c_S)$ and ν_p is the additive valuation on \mathbb{Q} normalized so that $\nu_p(p) = 1$. Moreover, all S involved satisfy $\chi = ' \chi_S$.

(2) For a suitable $e \in \mathbb{N}$, all of $S \in \Lambda_r^+$ appearing in (1) satisfy that $\text{level}(S) \mid p^e N$.

Remark 3.3. (1) Actually, each c_S is described in terms of the primitive Fourier coefficient for S of F . For details, see the proof in Section 7.

(2) The statement on $\widetilde{\theta_S^{(n)}} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^{m-\nu}}$ can be rephrased by $c_S \theta_S^{(n)} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$. Note also that $\nu = 0$ if $m = 1$.

(3) We emphasize that we do not know anything about e .

We expect that the theorem above holds in the most general case:

Conjecture 3.4. Theorem 3.2 should hold for any prime (not only for $p > r + 1$) and without the assumption $n \geq r$.

The smallest weights (see Theorem 2.3) where we can expect mod p singular forms, which are not true singular forms, is of special interest. In these cases, we expect also the power e of p in the levels of theta series to be the smallest possible:

Conjecture 3.5. Let n be a positive integer, r an even integer, and p a prime with $p \geq n + r + 3$. We put

$$k = k(p, r) := \begin{cases} r/2 + (p-1)/2 & \text{if } r \equiv 2 \pmod{4}, p \equiv -1 \pmod{4} \\ r/2 + p - 1 & \text{if } r \equiv 0 \pmod{4}. \end{cases}$$

Let N be a positive integer and χ a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}_{(p)}}$ is mod p singular of p -rank r . Then we have

$$F \equiv \sum_{\substack{S \in \Lambda_r / \text{GL}_r(\mathbb{Z}) \\ \text{level}(S) \mid pN}} c_S \theta_S^{(n+r)} \pmod{p} \quad (c_S \in \mathbb{Z}_{(p)}).$$

In the special case of $N = 1$, $r = 2$, we can prove this conjecture:

Theorem 3.6. Let n be a positive integer and p a prime with $p \geq n + 5$. Suppose that $F \in M_{\frac{p+1}{2}}(\Gamma_{n+2})_{\mathbb{Z}_{(p)}}$ is mod p singular of p -rank 2. Then we have

$$F \equiv \sum_{\substack{S \in \Lambda_2 / \text{GL}_2(\mathbb{Z}) \\ \text{level}(S) = p}} c_S \theta_S^{(n+2)} \pmod{p} \quad (c_S \in \mathbb{Z}_{(p)}).$$

Moreover for any such F the degree one form $f := \Phi^{n+1}(F)$ is nonzero mod p .

Remark 3.7. We remark that Theorem 3.6 has some potential for showing the nonvanishing of Fourier coefficients mod p in some interesting cases, e.g. a degree 3 modular form satisfying $\Phi^2(F) \equiv 0 \pmod{p}$ cannot be mod p singular. This implies the existence of some nonvanishing rank 3-Fourier coefficients $a_F(T) \pmod{p}$ of Klingen-Eisenstein series $F := E_k^{3,2}$ attached to a cusp form h of degree 2, provided that its Fourier coefficients are in $\mathbb{Z}_{(p)}$. Note that such Fourier coefficients are quite delicate, given by some critical values of L -series attached to h and to $T \in \Lambda_3^+$, if h is a Hecke eigenform (see [3, 17] for more details on Klingen-Eisenstein series). To cover more general cases, versions of this for congruences modulo prime ideals in the field generated by the Hecke eigenvalues of h will be necessary.

4 Refinement of Freitag’s expansion

In this section, inspired by Freitag [9, 10], we give a formal expansion of the “singular part” of the Fourier expansion of any modular form and apply it to mod p^m singular forms.

We fix some notation. Let $M_n(R)$ be the set of all $n \times n$ matrices whose components are in R . We put $M_n^{(r)}(\mathbb{Z}) := \{M \in M_n(\mathbb{Z}) \mid \text{rank}(M) = r\}$ and $M_n^*(\mathbb{Z}) := M_n^{(n)}(\mathbb{Z})$. Similarly we write $\Lambda_n^{(r)} := \{T \in \Lambda_n \mid \text{rank}(T) = r\}$ ($\Lambda_n^+ = \Lambda_n^{(n)}$). Let F be a modular form of degree $n + r$ with Fourier expansion

$$F(Z) = \sum_{T \in \Lambda_{n+r}} a(T) e(\text{tr}(TZ)).$$

We write $a(S) := a\left(\begin{smallmatrix} 0 & 0 \\ 0 & S \end{smallmatrix}\right)$ when $S \in \Lambda_r$. We define a subseries $F_{[r]}$ of F as

$$F_{[r]}(Z) := \sum_{T \in \Lambda_{n+r}^{(r)}} a(T) e(\text{tr}(TZ)).$$

In Böcherer-Raghavan [8] (see page 82 and 83), the notion of “primitive Fourier coefficient” was introduced; we denote it by $a^*(S)$ for S positive definite. Namely, $a^*(S)$ is defined by the formula

$$a(S) = \sum_{\substack{G \in \text{GL}_r(\mathbb{Z}) \setminus M_r^*(\mathbb{Z}) \\ S[G^{-1}] \in \Lambda_r}} a^*(S[G^{-1}]).$$

We recall that by this formula, we can define a new $\text{GL}_r(\mathbb{Z})$ -invariant function $a^*(S)$ starting from the $\text{GL}_r(\mathbb{Z})$ -invariant function $T \mapsto a(T)$ on Λ_r .

As explained in [8], this can also be written as

$$a(T) = \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{1}{\epsilon(S)} \sum_{\substack{W \in M_r^*(\mathbb{Z}) \\ S[W]=T}} a^*(S), \quad (4.1)$$

where $\epsilon(S) := A(S, S)$. We use this version.

Using the Fourier coefficients $a(S)$ with $S \in \Lambda_r^+$ and their modification, a slight refinement of Freitag's argument gives by formal rearrangement of the Fourier expansion our crucial identity. Note that the statements are only claiming the equality of the Fourier coefficients on both sides (ignoring questions concerning convergence!).

Lemma 4.1. *Let χ be a Dirichlet character mod N such that $\chi(-1) = (-1)^k$. Let $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)$. Then we have*

$$F_{[r]}(Z) = \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} \sum_{\substack{(X_1, X_2) \in \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \text{rank}(X_1, X_2) = r}} e(\text{tr}(S[(X_1, X_2)]Z)). \quad (4.2)$$

Note that

$$\sum_{(X_1, X_2) \in \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n}} e(\text{tr}(S[(X_1, X_2)]Z)) = \sum_{X \in \mathbb{Z}^{r, n+r}} e(\text{tr}(S[X]Z)) = \theta_S^{(n+r)}(Z).$$

Hence if we can prove this lemma, then $F_{[r]}$ can be expressed by a infinite linear combination of (subseries of) theta series;

$$F_{[r]} = \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} (\theta_S^{(n+r)})_{[r]}. \quad (4.3)$$

Proof of Lemma 4.1. For $X_1 \in \mathbb{Z}^{r,r}$, $X_2 \in \mathbb{Z}^{r,n}$, we can take $W \in M_r^*(\mathbb{Z})$ such that $(X_1, X_2) = W(G_1, G_2)$ and $\begin{pmatrix} * & * \\ G_1 & G_2 \end{pmatrix} \in \text{GL}_{n+r}(\mathbb{Z})$. This W can be regarded as ‘‘gcd’’ of X_1 and X_2 . We observe that such W is unique up to a factor in $\text{GL}_r(\mathbb{Z})$ from the right. We switch this action of $\text{GL}_r(\mathbb{Z})$ to (G_1, G_2) . Then we can write the right hand side of (4.2) as

$$\sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} \sum_{W \in M_r^*(\mathbb{Z})} \sum_{\substack{(G_1, G_2) \in \text{GL}_r(\mathbb{Z}) \backslash \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \begin{pmatrix} * & * \\ G_1 & G_2 \end{pmatrix} \in \text{GL}_{n+r}(\mathbb{Z})}} e(\text{tr}(S[W][(G_1, G_2)]Z)) \quad (4.4)$$

where $\text{GL}_r(\mathbb{Z}) \backslash \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} = \sim \backslash \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n}$ and $(X_1, X_2) \sim (Y_1, Y_2)$ means that $(X_1, X_2) = G(Y_1, Y_2)$ for some $G \in \text{GL}_r(\mathbb{Z})$.

We put $S[W] = T$ and we rewrite the summation over S as over T . Then (4.4) becomes

$$\sum_{T \in \Lambda_r^+} a(T) \sum_{\substack{(G_1, G_2) \in \text{GL}_r(\mathbb{Z}) \backslash \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \begin{pmatrix} * & * \\ G_1 & G_2 \end{pmatrix} \in \text{GL}_{n+r}(\mathbb{Z})}} e(\text{tr}(T[(G_1, G_2)]Z)) \quad (4.5)$$

because of (4.1).

If we put $U := \begin{pmatrix} * & * \\ G_1 & G_2 \end{pmatrix}$, then we have $T[(G_1, G_2)] = \begin{pmatrix} 0 & 0 \\ 0 & T \end{pmatrix} [U]$. Therefore we have $a(T) = a\left(\begin{pmatrix} 0 & 0 \\ 0 & T \end{pmatrix}\right) = a(T[(G_1, G_2)])$. Then (4.5) can be written as

$$\begin{aligned} & \sum_{T \in \Lambda_r^+} \sum_{\substack{(G_1, G_2) \in \mathrm{GL}_r(\mathbb{Z}) \setminus \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \begin{pmatrix} * & * \\ G_1 & G_2 \end{pmatrix} \in \mathrm{GL}_{n+r}(\mathbb{Z})}} a(T[(G_1, G_2)]) \mathbf{e}(\mathrm{tr}(T[(G_1, G_2)]Z)) \\ &= \sum_{T \in \Lambda_{n+r}^{(r)}} a(T) \mathbf{e}(\mathrm{tr}(TZ)) = F_{[r]}(Z). \end{aligned}$$

Here the first equality in this formula follows from the fact that, if $T \in \Lambda_r^+$ and (G_1, G_2) run as in the subscript, then $T[(G_1, G_2)]$ runs over all elements of $\Lambda_{n+r}^{(r)}$. \square

Note that $F_{[r]}$ is not a modular form because some part of the Fourier expansion is missing.

Let $Z_1 \in \mathbb{H}_r$, $Z_2 \in \mathbb{H}_n$. Consider the restriction of F to $Z = \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix}$;

$$F \left(\begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \right) = \sum_{\substack{\begin{pmatrix} T & * \\ * & * \end{pmatrix} \in \Lambda_{n+r} \\ T \in \Lambda_r}} \phi_T(Z_2) \mathbf{e}(\mathrm{tr}(TZ_1)).$$

Then we have $\phi_T(Z_2) \in M_k(\Gamma_0^{(n)}(N), \chi)$ for any $T \in \Lambda_r$. For the proof of this fact, we refer to Andrianov [1] (page 83 and 84).

On the other hand, we denote by F^\sharp the subseries of $F = \sum_{\mathfrak{T} \in \Lambda_{n+r}} a_F(\mathfrak{T}) \mathbf{e}(\mathrm{tr}(\mathfrak{T}Z))$, characterized by

$$\mathfrak{T} = \begin{pmatrix} T & * \\ * & * \end{pmatrix} \quad \text{with } T \in \Lambda_r^+.$$

Namely we put

$$F^\sharp(Z) := \sum_{\substack{\mathfrak{T} = \begin{pmatrix} T & * \\ * & * \end{pmatrix} \in \Lambda_{n+r} \\ T \in \Lambda_r^+}} a_F(\mathfrak{T}) \mathbf{e}(\mathrm{tr}(\mathfrak{T}Z)).$$

Then we have

$$F^\sharp \left(\begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \right) = \sum_{\substack{\begin{pmatrix} T & * \\ * & * \end{pmatrix} \in \Lambda_{n+r} \\ T \in \Lambda_r^+}} \phi_T(Z_2) \mathbf{e}(\mathrm{tr}(TZ_1)).$$

Note that still we have $\phi_T(Z_2) \in M_k(\Gamma_0^{(n)}(N), \chi)$ for any $T \in \Lambda_r^+$.

Now assume that F is mod p^m singular of p -rank r . Then from F being mod p^m

singular we obtain $F^\sharp \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \equiv (F_{[r]})^\sharp \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \pmod{p^m}$. By Lemma 4.1, we have

$$\begin{aligned}
(F_{[r]})^\sharp \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} &\equiv \left(\sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} \sum_{\substack{(X_1, X_2) \in \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \text{rank}(X_1, X_2) = r}} \mathbf{e}(\text{tr}(S[X_1]Z_1)) \mathbf{e}(\text{tr}(S[X_2]Z_2)) \right)^\sharp \\
&\equiv \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} \sum_{\substack{(X_1, X_2) \in \mathbb{Z}^{r,r} \times \mathbb{Z}^{r,n} \\ \text{rank}(X_1) = r}} \mathbf{e}(\text{tr}(S[X_1]Z_1)) \mathbf{e}(\text{tr}(S[X_2]Z_2)) \\
&\equiv \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} \sum_{\substack{X_1 \in \mathbb{Z}^{r,r} \\ \text{rank}(X_1) = r}} \mathbf{e}(\text{tr}(S[X_1]Z_1)) \sum_{X_2 \in \mathbb{Z}^{r,n}} \mathbf{e}(\text{tr}(S[X_2]Z_2)) \\
&\equiv \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} \frac{a^*(S)}{\epsilon(S)} (\theta_S^{(r)}(Z_1))_{[r]} \theta_S^{(n)}(Z_2) \pmod{p^m}.
\end{aligned}$$

This implies that

$$\phi_T(Z_2) \equiv \sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} A(S, T) \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n)}(Z_2) \pmod{p^m}$$

for any $T \in \Lambda_r^+$. Hence we obtain that

$$\sum_{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z})} A(S, T) \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n)}(Z_2) \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}.$$

In the ordinary case (over \mathbb{C}), the key in Freitag's setting would be that $a(S)$ can be different from zero only if $\text{level}(S) \mid N$. In our mod p^m setting, we get a condition on the filtration of $\theta_S^{(n)}$ for $S \in \Lambda_r^+$ with $a^*(S) \not\equiv 0 \pmod{p^m}$. More precisely, we get the following property.

Proposition 4.2. *Let n, k, N be positive integers and r an even integer. Let p be a prime with $p > r + 1$ and χ a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}_{(p)}}$ is mod p^m singular of p -rank r . Then we have $a^*(S) \theta_S^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$ for any $S \in \Lambda_r^+$. In particular, if $S \in \Lambda_r^+$ satisfies $a^*(S) \not\equiv 0 \pmod{p^m}$, then we have $\theta_S^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^{m-\nu}}$ (and hence $\omega_{N, \chi, p^{m-\nu}}^n(S) \leq k$) with $\nu := \nu_p(a^*(S))$. Moreover $\chi = \chi_S$ holds.*

Proof. Seeking a contradiction, we suppose that there exists S such that the claim is not true. Let S_0 be one of S such that $\det S_0$ is minimal among such S . Then we

consider

$$\begin{aligned}
\phi_{S_0} &= \sum_{\substack{S \in \Lambda_r^+ / \mathrm{GL}_r(\mathbb{Z}) \\ \det S < \det S_0}} A(S, S_0) \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n)} \\
&\equiv \sum_{\substack{S \in \Lambda_r^+ / \mathrm{GL}_r(\mathbb{Z}) \\ \det S \geq \det S_0}} A(S, S_0) \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n)} \\
&\equiv A(S_0, S_0) \frac{a^*(S_0)}{\epsilon(S_0)} \theta_{S_0}^{(n)} \pmod{p^m}.
\end{aligned}$$

Here the last congruence follows from the facts that $\det S > \det S_0$ implies $A(S, S_0) = 0$ and $\det S = \det S_0$, $S \not\sim S_0 \pmod{\mathrm{GL}_r(\mathbb{Z})}$ implies $A(S, S_0) = 0$. By the assumption, we have

$$\phi_{S_0} = \sum_{\substack{S \in \mathrm{GL}_r(\mathbb{Z}) \setminus \Lambda_r^+ \\ \det S < \det S_0}} A(S, S_0) \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}.$$

This implies $A(S_0, S_0) \frac{a^*(S_0)}{\epsilon(S_0)} \theta_{S_0}^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$. By the fact (2.1), we have $p \nmid A(S_0, S_0)$. This shows $a^*(S_0) \theta_{S_0}^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$. This is a contradiction. Hence we have $a^*(S) \theta_S^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^m}$ for any $S \in \Lambda_r^+$. In particular if $a^*(S) \not\equiv 0 \pmod{p^m}$, then we have $\theta_S^{(n)} \pmod{p^m} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^{m-\nu}}$ with $\nu := \nu_p(a^*(S))$, and therefore $\omega_{N, \chi, p^{m-\nu}}^n(S) \leq k$. The statement $\chi = \chi_S$ follows from Proposition 2.4. \square

For later use, we mention a simple consequence of Proposition 4.2 for the mod p case.

Corollary 4.3. *Let n, k, N be positive integers, r an even integer with $n \geq r$. Let p be a prime with $p > r + 1$ and χ a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}_{(p)}}$ is mod p singular of p -rank r .*

- (1) *For any $S \in \Lambda_r^+$ with $a^*(S) \not\equiv 0 \pmod{p}$, we have $\widetilde{\theta}_S^{(r)} \in \widetilde{M}_k(\Gamma_0^{(r)}(N), \chi)$.*
- (2) *For $S \in \Lambda_r^+$ with $a(S) \not\equiv 0 \pmod{p}$ such that $\det S$ is minimal in such S , we have $\widetilde{\theta}_S^{(n)} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)$ and therefore $\omega_{N, \chi}^n(S) \leq k$.*

Remark 4.4. *The statement (2) can be proved also by the mod p version of Freitag's original arguments in [10]. Our strategy can be viewed as a refinement of his method.*

Proof of Corollary 4.3. (1) By Proposition 4.2, we have $\widetilde{\theta}_S^{(n)} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)$. Hence the claim follows from $\Phi^{n-r}(\theta_S^{(n)}) = \theta_S^{(r)}$ immediately.

(2) If $S \in \Lambda_r^+$ satisfies such the minimality condition, we have $a^*(S) = a(S)$ by the definition of $a^*(S)$. The claim follows from this fact. \square

5 Description by theta series for mod p case

In this section, we prove Theorem 3.2 (1) for $m = 1$. Therefore, we treat with the case of mod p in this section. The goal is to show that all strongly mod p singular modular forms are represented by a linear combination of finitely many theta series. In our method, two new tools are important: the existence of an “abstract Sturm bound” for detecting mod p singular forms and the refinement of Freitag’s expansion as exposed in the previous section. The nice thing is that we do not need to consider the exact level of the theta series.

5.1 Abstract Sturm bounds

In general, “Sturm bounds” give an explicit finite set of $T \in \Lambda_n$ such that a modular form of degree n with Fourier coefficients in $\mathbb{Z}_{(p)}$ must be congruent mod p to zero if the Fourier coefficients for all T in that finite set are divisible by p , see e.g. [20, 25]. Such an explicit finite set would usually contain quadratic forms of all ranks. We need a version, which involves only quadratic forms of rank $> r$ (finitely many) to detect mod p singular forms. We only need the existence of such a set, but we do not discuss explicit bounds for it.

Let χ be a quadratic Dirichlet character mod N . Let $M_{k,r}^{p\text{-sing}}(\Gamma_0^{(n+r)}(N), \chi)$ be the submodule of $M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}_{(p)}}$ consisting of all mod p singular modular forms with p -rank $\leq r$. We denote by $\widetilde{M}_{k,r}^{p\text{-sing}}(\Gamma_0^{(n+r)}(N), \chi)$ the set of reduction mod p of elements of $M_{k,r}^{p\text{-sing}}(\Gamma_0^{(n+r)}(N), \chi)$. This is a subspace over \mathbb{F}_p of the vector space $\widetilde{M}_k(\Gamma_0^{(n+r)}(N), \chi)$. We consider the quotient space

$$V = V_{k,r} := \widetilde{M}_k(\Gamma_0^{(n+r)}(N), \chi) / \widetilde{M}_{k,r}^{p\text{-sing}}(\Gamma_0^{(n+r)}(N), \chi).$$

Then we have $\dim V < \infty$. For fixed $T \in \Lambda_{n+r}$ with $\text{rank}(T) > r$ we define a linear map $\ell_T : V \rightarrow \mathbb{F}_p$ by

$$\ell_T(\widetilde{F} + \widetilde{M}_{k,r}^{p\text{-sing}}(\Gamma_0^{(n+r)}(N), \chi)) := \widetilde{a_F(T)}.$$

Clearly, the set L of all such ℓ_T is “total” for V , i.e. the intersection of the kernels of all ℓ_T is trivial. By linear algebra (V is of finite dimension!) we can choose a finite subset $\mathcal{T}_{n+r,r} = \{\ell_{T_1}, \dots, \ell_{T_d}\}$ of L which is still total for V .

Conclusion: *In the situation above, there exist finitely many T_1, \dots, T_d with all T_j of rank larger than r such that for all $F \in M_k(\Gamma_0^{(n)}(N), \chi)_{\mathbb{Z}_{(p)}}$ the vanishing of $\widetilde{a_F(T_j)}$ for all T_j implies that F is mod p singular of p -rank $\leq r$.*

We call such $\mathcal{T}_{n+r,r}$ a “Sturm set” for mod p singular forms.

5.2 Proof of Theorem 3.2 (1) for $m = 1$

For the Sturm set $\mathcal{T}_{r,r-1}$ ($\subset \Lambda_r^+$) corresponding to $M_{k,r-1}^{p\text{-sing}}(\Gamma_0^{(r)}(N))$, we take a natural number M such that

$$M > \max\{\det T \mid T \in \mathcal{T}_{r,r-1}\}.$$

We put

$$G := F - \sum_{\substack{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z}) \\ \det S < M}} \frac{a^*(S)}{\epsilon(S)} \theta_S^{(n+r)}$$

and consider $g := \Phi^n(G)$. Note here that the summation over S is finite. Then we have

$$g = \Phi^n(F) - \sum_{\substack{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z}) \\ \det S < M}} \frac{a^*(S)}{\epsilon(S)} \theta_S^{(r)}$$

and $\tilde{g} \in \widetilde{M}_k(\Gamma_0^{(r)}(N), \chi)$ by Corollary 4.3 (1).

We prove now that $\tilde{g} \in \widetilde{M}_{k,r-1}^{p\text{-sing}}(\Gamma_0^{(r)}(N), \chi)$. It suffices to check the Fourier coefficients $a_g(T)$ for all $T \in \mathcal{T}_{r,r-1}$. It follows from (4.3) that

$$a_g(T) = \sum_{\substack{S \in \Lambda_r^+ / \text{GL}_r(\mathbb{Z}) \\ \det S \geq M}} \frac{a^*(S)}{\epsilon(S)} A(S, T)$$

for any $T \in \Lambda_r^+$. If $S[X] = T$ with $X \in \mathbb{Z}^{r,r}$ then $\det S[X] = \det S(\det X)^2 = \det T$ and hence $\det S \leq \det T$. This implies that $A(S, T) = 0$ for T with $\det T < \det S$. Therefore we have $a_g(T) = 0$ for any $T \in \mathcal{T}_{r,r-1}$. This implies $g \in M_{k,r-1}^{p\text{-sing}}(\Gamma_0^{(r)}(N), \chi)$.

Suppose that $g \not\equiv 0 \pmod{p}$. Then g is mod p singular of some p -rank $r' < r$. By Theorem 2.3, we have $2k - r \equiv 2k - r' \equiv 0 \pmod{p-1}$ and hence $r' \equiv r \pmod{p-1}$. Since $r' < r < p-1$, this is impossible. This shows $g \equiv 0 \pmod{p}$. Taking into account that F and G are mod p singular, we obtain that $G \equiv 0 \pmod{p}$. This completes the proof of (1) in Theorem 3.2. \square

Remark 5.1. *The condition “ $n \geq r$ ” was necessary to assure $\tilde{g} \in \widetilde{M}_k(\Gamma_0^{(r)}(N), \chi)$.*

6 Specification of levels

6.1 Proof of Theorem 3.2 (2) for $m = 1$

In the situation of Theorem 3.2 we have to specify the levels of the $S \in \Lambda_r^+$ involved: To do so, we use Corollary 9.3, which follows from the modified q -expansion principle and Kitaoka’s formula. We observe that $\theta_S^{(r)} \in \widetilde{M}_k(\Gamma_0^{(r)}(N), \chi)$; here we need the stronger condition $p \geq r + 3$ (and also $n \geq r$). Then Corollary 9.3 shows that the

level of S is of the form requested and also the statement about the nebentypus follows. \square

Now we prove Theorem 3.6 in the remainder of this section. Namely, we discuss the case where the p -rank is 2 and the weight is the minimal one allowing mod p singular modular forms, which are not truly singular, i.e. $1 + \frac{p-1}{2}$.

6.2 Proof of Theorem 3.6

In Theorem 3.2, we assumed $n \geq r$. In this subsection, we discuss the special case of $n = 1$ and $r = 2$; this is the simplest case where the condition “strongly mod p singular” is violated. We use instead very special properties of binary quadratic forms.

Proposition 6.1. *Let p be an odd prime. Let $S \in \Lambda_2^+$ be of level $N = p^j N'$ with N' coprime to p . Suppose that $\theta_S^{(1)} \equiv \phi \pmod{p}$ for some $\phi \in M_k(\Gamma_1)_{\mathbb{Z}(p)}$. Then $N' = 1$, i.e. $\text{level}(S) = p^j$.*

Remark 6.2. *To include the case of $n = 1$, we should look at the degree 1 theta series $\theta_S^{(1)}$ and $\phi \in M_k(\Gamma_1)$.*

Proof. Seeking a contradiction we suppose $N' > 1$. We apply Theorem 9.1 of Kitaoka’s original result for $n = 1$ to $\theta_S^{(1)}$ with $M = \begin{pmatrix} * & * \\ p^j & N' \end{pmatrix}$. Then we have

$$\theta_S^{(1)}|M = \kappa \cdot \theta_{S'}^{(1)},$$

where S' is the same as in the proof of Proposition 9.2. Then $N'S'$ is half integral, $(N', \text{cont}(N'S')) = 1$, and $\text{cont}(N'S') = p^\alpha$ when $p^\alpha \parallel \text{cont}(S)$. Then $\frac{N'}{p^\alpha} S' \in \Lambda_2^+$ is primitive. Therefore we can take a prime l with $l \nmid N'$ such that

$$A\left(S', \frac{l \cdot p^\alpha}{N'}\right) = A\left(\frac{N'S'}{p^\alpha}, l\right) \not\equiv 0 \pmod{p}.$$

However, since ϕ is of level 1, we have also

$$A\left(S', \frac{l \cdot p^\alpha}{N'}\right) \equiv a_\phi\left(\frac{l \cdot p^\alpha}{N'}\right) = 0 \pmod{p}.$$

This contradicts and we get $N' = 1$. \square

Hence $\text{level}(S)$ is a power of p in the situation of Proposition 6.1. Then $\det(2S)$ also should be a power of p . By the elementary divisor theorem, we can find $U, V \in \text{GL}_2(\mathbb{Z})$ such that

$$U(2S)V = \begin{pmatrix} p^s & 0 \\ 0 & p^{s+t} \end{pmatrix}.$$

Then we have $U(2S) = \begin{pmatrix} p^s & 0 \\ 0 & p^{s+t} \end{pmatrix} V^{-1}$ and hence

$$U(2S)^t U = p^s \begin{pmatrix} 1 & 0 \\ 0 & p^t \end{pmatrix} V^{-1t} U.$$

If we put $W := V^{-1t} U = \begin{pmatrix} w_1 & w_2 \\ w_3 & w_4 \end{pmatrix}$, then we can write

$$U(2S)^t U = p^s \begin{pmatrix} w_1 & p^t w_3 \\ w_2 & p^t w_4 \end{pmatrix}.$$

Since $U(2S)^t U$ is symmetric, we have $w_2 = p^t w_3$ and

$$U(2S)^t U = p^s \begin{pmatrix} w_1 & p^t w_3 \\ p^t w_3 & p^t w_4 \end{pmatrix}.$$

From these argument, we may assume that S is of this form. In particular, if $\det(2S)$ is an odd power of p , by putting $b := p^j w_3$, we may assume that S is of the form

$$S = S(i, j) = p^i \begin{pmatrix} a & bp^{j+1} \\ bp^{j+1} & dp^{2j+1} \end{pmatrix}$$

with $adp - b^2 p^2 = p$.

Now we try to show that high powers of p in the level of S imply high filtration weight of $\theta_S^{(1)}$:

Proposition 6.3. *Let p be an odd prime and $S \in \Lambda_2^+$. Suppose that $\det(2S)$ is an odd power of p . Then we have*

$$\omega(\theta_S^{(1)}) \geq p^{i+j} \cdot \frac{p+1}{2},$$

where i, j are determined by

$$S = S(i, j) = p^i \begin{pmatrix} a & bp^{j+1} \\ bp^{j+1} & dp^{2j+1} \end{pmatrix}$$

with $adp - b^2 p^2 = p$.

Before proving this proposition, we confirm some notation and facts on filtration of modular forms mod p for degree 1 given by Serre [23] and Swwinerton-Dyer [26]. Let $f = \sum_{n=0}^{\infty} a_f(n) \mathbf{e}(nz)$ be a formal power series with $a_f(n) \in \mathbb{Z}_{(p)}$. We write $\omega(f)$ for $\omega_{1,1}^1(f)$. Let $V(p), U(p)$ be the operators defined by

$$f|V(p) = \sum_{n=0}^{\infty} a_f(n) \mathbf{e}(pnz),$$

$$f|U(p) = \sum_{n=0}^{\infty} a_f(pn) \mathbf{e}(nz).$$

Then we have $\omega(f|V(p)) = p\omega(f)$, $\omega(f|U(p)) \leq \omega(f)$.

We put $\omega(S(i, j)) := \omega(\theta_{S(i,j)}^{(1)})$.

Proof of Proposition 6.3. Using the facts that $\theta_{S(i,j)}^{(1)} = \theta_{S(i-1,j)}^{(1)}|V(p)$ and $\omega(f|V(p)) = p\omega(f)$, we have

$$\omega(S(i, j)) = p^i \omega(S(0, j)).$$

Furthermore, by comparing the representation numbers of $A(S(0, j), pn)$ and $A(S(1, j-1), n)$, we can easily confirm that

$$\theta_{S(0,j)}^{(1)}|U(p) = \theta_{S(1,j-1)}^{(1)}.$$

Since $\omega(f) \geq \omega(f|U(p))$ in general, we have

$$\omega(S(0, j)) \geq \omega(S(1, j-1)) = p\omega(S(0, j-1)).$$

This implies

$$\omega(S(i, j)) \geq p^{i+j} \omega(S(0, 0)) = p^{i+j} \omega(S(0, 0)) = p^{i+j} \cdot \frac{p+1}{2}.$$

Here the last equality follows from $\omega(S(0, 0)) = \frac{p+1}{2}$ and this is due to Theorem 2.7 (or Serre's result in [23]). This completes the proof. \square

Corollary 6.4. *Let p be a prime and $S \in \Lambda_2^+$. Suppose that $\theta_S^{(1)} \equiv \phi \pmod{p}$ for some $\phi \in M_{\frac{p+1}{2}}(\Gamma_1)_{\mathbb{Z}(p)}$. Then we have $\text{level}(S) = p$.*

Proof. The assumption $\theta_S^{(1)} \equiv \phi \pmod{p}$ implies

$$\omega(S) = \omega(S(i, j)) \leq \frac{p+1}{2}. \quad (6.1)$$

On the other hand, the level of S is a power of p because of Proposition 6.1. Then $\det(2S)$ is a power of p . Note here that, actually in this case, the power of p is odd. Because, the quadratic character χ_S should have nontrivial p -component, otherwise $\frac{p+1}{2} \equiv 1 \pmod{p-1}$ should hold and this is impossible. Hence we can assume that $S = S(i, j)$ as in Corollary 6.3. Then we have

$$\omega(S) = \omega(S(i, j)) \geq p^{i+j} \cdot \frac{p+1}{2}.$$

Combing this with (6.1), we have $i = j = 0$. Namely, we obtain $\text{level}(S) = p$. \square

We can now prove our result for the case of p -rank 2.

Proof of Theorem 3.6. Let $\{T_1, \dots, T_{h_p}\} \subset \Lambda_2/\text{GL}_2(\mathbb{Z})$ be a set of the representatives of $\text{GL}_2(\mathbb{Z})$ -inequivalence classes of binary quadratic forms with level p . Note that $\theta_{T_i}^{(n)} \in M_1(\Gamma_0^{(n)}(p), (\frac{*}{p}))$, and there exists $G_i \in M_{\frac{p+1}{2}}(\Gamma_n)$ such that $G_i \equiv \theta_{T_i}^{(n)}$

mod p by Theorem 2.7. Then G_i is mod p singular, because $\theta_{T_i}^{(n)}$ is true singular. Then we consider

$$H := F - \sum_{i=1}^{h_p} \frac{1}{2} a_F \begin{pmatrix} 0 & 0 \\ 0 & T_i \end{pmatrix} G_i \in M_{\frac{p+1}{2}}(\Gamma_{n+2})$$

This H is a mod p singular of some p -rank $r' \leq 2$.

Now we suppose that still $r' = 2$. Then we can take $S \in \Lambda_2^+$ with $a_H \begin{pmatrix} 0 & 0 \\ 0 & S \end{pmatrix} \not\equiv 0 \pmod{p}$ such that $\det S$ is minimal. By Corollary 4.3, we have $\theta_S^{(1)} \in \widetilde{M}_{\frac{p+1}{2}}(\Gamma_1)$. By Corollary 6.4, we have $\text{level}(S) = p$ and hence $S \sim T_j \pmod{\text{GL}_2(\mathbb{Z})}$ for some j . However, from $a_H \begin{pmatrix} 0 & 0 \\ 0 & T_i \end{pmatrix} \equiv 0 \pmod{p}$ for any i , we have $a_H \begin{pmatrix} 0 & 0 \\ 0 & S \end{pmatrix} \equiv 0 \pmod{p}$. This is a contradiction.

Therefore we have $r' \leq 1$ or $H \equiv 0 \pmod{p}$. If $H \not\equiv 0 \pmod{p}$, then we have $p+1-r' \equiv 0 \pmod{p-1}$ because of Theorem 2.3. This is impossible. This implies $H \equiv 0 \pmod{p}$. This shows the main statement of Theorem 3.6.

To prove the final statement concerning $\Phi^{n+1}(F) \not\equiv 0 \pmod{p}$ it is sufficient to assure the mod p linear independence of the binary theta series in question: Let $\{T_1, \dots, T_{h_p}\}$ be as above. As stated in Kani [13], $\theta_{T_j}^{(1)}$ ($j = 1, \dots, h_p$) are linearly independent over \mathbb{C} . Now we can prove that $\theta_{T_j}^{(1)}$ ($j = 1, \dots, h_p$) are linearly independent over \mathbb{F}_p : Let $\sum_{i=1}^{h_p} c_i \theta_{T_j}^{(1)} \equiv 0 \pmod{p}$. Then, for any $n \geq 0$, we have

$$\sum_{i=1}^{h_p} c_i A(T_i, n) \equiv 0 \pmod{p}.$$

For each T_i , we can find infinitely many primes l such that $A(T_i, l) > 0$. Then we have $A(T_j, l) = 0$ for any $j \neq i$. Since $A(T_i, l) = 2, 4, 6 \not\equiv 0 \pmod{p}$, we have $c_j \equiv 0 \pmod{p}$. Hence we have $c_i \equiv 0 \pmod{p}$. This shows $c_i \equiv 0 \pmod{p}$ for any i with $1 \leq i \leq h_p$. Therefore $\theta_{T_j}^{(1)}$ ($j = 1, \dots, h_p$) are linearly independent over \mathbb{F}_p . This completes the proof of Theorem 3.6. \square

7 From mod p to mod p^m

In this section, using induction, we extend Theorem 3.2 (proved for $m = 1$ in the previous sections) to the case of general m . The main reason why we preferred to give a mod p version first and then extend to arbitrary powers is the technical difficulty which one encounters, when one tries to give a mod p^m analogue of our abstract Sturm bound. We avoid such a problem in this way.

We start with proving the following weaker version of Theorem 3.2.

Theorem 7.1 (Weaker version of Theorem 3.2). *Let n, k, N be positive integers, r an even integer with $n \geq r$. Let p be a prime with $p > r+1$ and χ a quadratic Dirichlet character mod N with $\chi(-1) = (-1)^k$. Suppose that $F \in M_k(\Gamma_0^{(n+r)}(N), \chi)_{\mathbb{Z}(p)}$ is*

mod p^m singular of p -rank r . Then there are finitely many $S \in \Lambda_r^+$ of level dividing $p^e N$ with $\widetilde{\theta}_S^{(n)} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)$ (only mod p) such that

$$F \equiv \sum_S c_S \theta_S^{(n+r)} \pmod{p^m} \quad (c_S \in \mathbb{Z}_{(p)}). \quad (7.1)$$

Moreover, all S involved satisfy $\chi =' \chi_S$.

Proof of Theorem 7.1. We prove the statement by induction on m . We have already proved the statement for $m = 1$. Suppose that the statements (1), (2) in Theorem 3.2 are true for any m with $m < m_0$. We consider the case $m = m_0$. Note that $2k - r \equiv 0 \pmod{(p-1)p^{m_0-1}}$ by Theorem 2.3. Therefore we can write $k = \frac{r}{2} + t \cdot \frac{p-1}{2} p^{m_0-1}$.

By the assumption, F is mod p^{m_0-1} singular (because of mod p^{m_0} singular) and therefore by induction hypothesis, we have

$$F \equiv \sum_{\substack{S \in \Lambda_r^+ \\ \text{level}(S) | p^{e_1} N}} c_S \theta_S^{(n+r)} \pmod{p^{m_0-1}} \quad \text{with } c_S \in \mathbb{Z}_{(p)},$$

for some e_1 , and all S involved satisfy $\chi =' \chi_S$.

We take $\mathcal{E} \in M_{\frac{p-1}{2}}(\Gamma_0^{(n+r)}(p), (\frac{*}{p})_{\mathbb{Z}_{(p)}})$ from [6] such that $\mathcal{E} \equiv 1 \pmod{p}$ and put $G := \sum_S c_S \theta_S^{(n+r)} \cdot \mathcal{E}^{tp^{m_0-1}}$. Note here that $F, G \in M_k(\Gamma_0^{(n+r)}(p^{e_1} N), \chi)_{\mathbb{Z}_{(p)}}$ because of $\chi =' \chi_S$. Consider $H := \frac{1}{p^{m_0-1}}(F - G) \in M_k(\Gamma_0^{(n+r)}(p^{e_1} N), \chi)_{\mathbb{Z}_{(p)}}$. Then for any T , we have

$$a_F(T) = a_G(T) + p^{m_0-1} a_H(T).$$

Since $a_F(T) \equiv a_G(T) \equiv 0 \pmod{p^{m_0}}$ for any T with $\text{rank}(T) > r$, we have $a_H(T) \equiv 0 \pmod{p}$ for any T with $\text{rank}(T) > r$. This means that H is mod p singular of some p -rank $r' \leq r$. By $2k - r' \equiv 2k - r \equiv 0 \pmod{p-1}$ and $0 \leq r' \leq r \leq p-1$, we have $r' = r$. Therefore $a_H(T) \not\equiv 0 \pmod{p}$ for some T with $\text{rank}(T) = r$. Again by our theorem for $m = 1$, we have

$$H \equiv \sum_{\substack{R \in \Lambda_r^+ \\ \text{level}(R) | p^{e_2} N}} d_R \theta_R^{(n+r)} \pmod{p} \quad \text{with } d_R \in \mathbb{Z}_{(p)},$$

and all R involved satisfy $\chi =' \chi_R$. Noting $G := \sum_S c_S \theta_S^{(n+r)} \cdot \mathcal{E}^{tp^{m_0-1}} \equiv \sum_S c_S \theta_S^{(n+r)} \pmod{p^{m_0}}$, we have

$$\begin{aligned} F &\equiv G + p^{m_0-1} H \\ &\equiv \sum_{\substack{S \in \Lambda_r^+ \\ \text{level}(S) | p^{e_1} N}} c_S \theta_S^{(n+r)} + p^{m_0-1} \sum_{\substack{R \in \Lambda_r^+ \\ \text{level}(R) | p^{e_2} N}} d_R \theta_R^{(n+r)} \pmod{p^{m_0}}. \end{aligned}$$

This completes the proof of Theorem 7.1. \square

Completion of the proof of Theorem 3.2. We first claim that the coefficients c_S in (7.1) coincide mod p^m with $\frac{a^*(S)}{\epsilon(S)}$ provided that the summation in (7.1) is restricted to pairwise $\mathrm{GL}_r(\mathbb{Z})$ -inequivalent elements of Λ_r^+ . To see this, we take an arbitrary $S_0 \in \Lambda_r^+$. Then (using freely the notation from Section 4)

$$a(S_0) \equiv \sum_S c_S A(S, S_0) \pmod{p^m}$$

holds and passing to primitive Fourier coefficients gives

$$a^*(S_0) \equiv \sum_S c_S A^*(S, S_0) \pmod{p^m}.$$

Now $A^*(S, S_0) = \epsilon(S_0)$ if S and S_0 are $\mathrm{GL}_r(\mathbb{Z})$ -equivalent and zero otherwise; this proves the claim from above.

To complete the proof of Theorem 3.2, we have to prove $\widetilde{\theta}_S^{(n)} \in \widetilde{M}_k(\Gamma_0^{(n)}(N), \chi)_{p^{m-\nu}}$; this follows now directly from Proposition 4.2. This completes the proof of Theorem 3.2. \square

8 Appendix A: Modified q -expansion principle

To specify the level of theta series in our theorems, we need some control over the behaviour of congruences of modular forms, when we switch to other cusps. The “ q -expansion principles” available in the literature do not exactly provide the information necessary for our purpose.

We fix a prime p and a natural number N coprime to p . For $m \geq 0$ let R_m be a subring of \mathbb{C} containing $\mathbb{Z}[e^{\frac{2\pi i}{N \cdot p^m}}, e^{\frac{2\pi i}{3}}, \frac{1}{6}, \frac{1}{N}]$.

Theorem 8.1. *Let p be a prime with $p \geq n + 3$ and N a positive integer with $p \nmid N$. Let $f \in M_k(\Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N))$. Then for all $\gamma \in \Gamma_0^{(n)}(p^m)$ we have*

$$f \in M_k(\Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N))_{R_m} \iff f|_k \gamma \in M_k(\Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N))_{R_m}.$$

The case $m = 0$ is the statement of the usual q -expansion principle in the formulation of Pitale et al. [19] proved by Katz [15] for $n = 1$ and Ichikawa [12] for $n > 1$. The main point in our version is that f and $f|_k \gamma$ share the same p -integrality property. Note that one only needs to prove one direction of the theorem. The aim of Appendix A is to prove this property.

8.1 Proof for $m = 1$

We will do induction on m . We need the existence of a modular form $\mathcal{E} \in M_l(\Gamma_0^{(n)}(p))_{\mathbb{Z}(p)}$ such that

$$\begin{aligned} \mathcal{E} &\equiv 1 \pmod{p} \\ \mathcal{E}|_l \omega_j &\equiv 0 \pmod{p} \quad (1 \leq j \leq n). \end{aligned}$$

Here ω_j can be any element $\begin{pmatrix} * & * \\ C & D \end{pmatrix} \in \Gamma_n$ with C being of rank i in $M_n(\mathbb{F}_p)$. The existence of such \mathcal{E} was explained in [2], under the condition $p \geq n + 3$. We mention that the congruence condition of \mathcal{E} can be rephrased as saying that for any $\gamma \in \Gamma_n$, we have

$$\mathcal{E}|\gamma \equiv 1 \pmod{p} \iff \gamma \in \Gamma_0^{(n)}(p)$$

and $\mathcal{E}|\gamma \equiv 0 \pmod{p}$ for $\gamma \notin \Gamma_0^{(n)}(p)$.

We prove the statement for $m = 1$. We chose an appropriate power α of \mathcal{E} and consider a trace from level $\Gamma_0^{(n)}(p) \cap \Gamma^{(n)}(N)$ to level $\Gamma^{(n)}(N)$:

$$F = f\mathcal{E}^\alpha + \sum_{j \pmod{p}} (f\mathcal{E}^\alpha)|\delta_j$$

where δ_j are representatives of the left cosets not in $\Gamma_0^{(n)}(p) \cap \Gamma^{(n)}(N)$, in particular, $\mathcal{E}|\delta_j \equiv 0 \pmod{p}$. Applying γ , we see that $\delta_j \cdot \gamma \notin \Gamma_0^{(n)}(p)$ and hence $(f\mathcal{E}^\alpha)|\delta_j \cdot \gamma \equiv 0 \pmod{p}$. We observe that F is of level $\Gamma^{(n)}(N)$ and hence we may apply the ordinary q -expansion principle: Then we have

$$F|\gamma \equiv (f|\gamma)\mathcal{E}^\alpha \equiv f|\gamma \pmod{p}.$$

In particular $F|\gamma$ is p -integral and hence $f|\gamma$ is also p -integral. This completes the proof for $m = 1$. \square

8.2 Proof for $m \geq 2$

We prove it for $m \geq 2$ by induction on the power m . We suppose that the statement is true for the level $\Gamma_0^{(n)}(p^{m-1})$. We put

$$\mathcal{F} := \mathcal{E}(p^{m-1}z) = p^{-(m-1)\frac{nl}{2}} \mathcal{E} \left| \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \right. \in M_l(\Gamma_0^{(n)}(p^m))_{\mathbb{Z}(p)}.$$

Here l is the weight of \mathcal{E} . Taking some positive integer α , we consider $f\mathcal{F}^\alpha$ and the trace of this from $\Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N)$ to $\Gamma_0^{(n)}(p^{m-1}) \cap \Gamma^{(n)}(N)$;

$$\begin{aligned} F = \text{tr}(f\mathcal{F}^\alpha) &= f\mathcal{F}^\alpha + \sum_{w \pmod{p}}^{p-1} (f\mathcal{F}^\alpha)|\gamma_w \\ &= f\mathcal{F}^\alpha + \sum_{j=1}^{p-1} (f|\gamma_w)(\mathcal{F}|\gamma_w)^\alpha, \end{aligned} \tag{8.1}$$

where $\gamma_w = \begin{pmatrix} 1_n & 0_n \\ p^{m-1}N_w & 1_n \end{pmatrix}$ and w runs over a complete set of representatives of integral symmetric matrices of size $n \pmod{p}$ except the trivial coset $p \cdot \text{Sym}_n(\mathbb{Z})$. Now we prove that $(f|\gamma_w)(\mathcal{F}|\gamma_w)^\alpha \equiv 0 \pmod{p}$. By a direct calculation, we have

$$\begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \begin{pmatrix} 1_n & 0_n \\ p^{m-1}N_w & 1_n \end{pmatrix} = \begin{pmatrix} 1_n & 0_n \\ N_w & 1_n \end{pmatrix} \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix}. \tag{8.2}$$

This implies

$$\begin{aligned}\mathcal{F} \left| \begin{pmatrix} 1_n & 0_n \\ p^{m-1}Nw & 1_n \end{pmatrix} \right. &= p^{-(m-1)\frac{nl}{2}} \mathcal{E} \left| \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \begin{pmatrix} 1_n & 0_n \\ p^{m-1}Nw & 1_n \end{pmatrix} \right. \\ &= p^{-(m-1)\frac{nl}{2}} \mathcal{E} \left| \begin{pmatrix} 1_n & 0_n \\ Nw & 1_n \end{pmatrix} \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \right.\end{aligned}$$

Since $\mathcal{E} \left| \begin{pmatrix} 1_n & 0_n \\ Nw & 1_n \end{pmatrix} \right. \equiv 0 \pmod{p}$ and we can put $\mathcal{E} \left| \begin{pmatrix} 1_n & 0_n \\ Nw & 1_n \end{pmatrix} \right. = pX$ for some p -integral modular form $X = X_w$. Then we have

$$\begin{aligned}\mathcal{F} \left| \begin{pmatrix} 1_n & 0_n \\ p^{m-1}Nw & 1_n \end{pmatrix} \right. &= p^{-(m-1)\frac{nl}{2}} \mathcal{E} \left| \begin{pmatrix} 1_n & 0_n \\ Nw & 1_n \end{pmatrix} \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \right. \\ &= p^{-(m-1)\frac{nl}{2}} (pX) \left| \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \right. \\ &= pX(p^{m-1}z) \equiv 0 \pmod{p}.\end{aligned}$$

Therefore $(f|\gamma_w)(\mathcal{F}|\gamma_w)^\alpha \equiv 0 \pmod{p}$ follows if α is large. By the assumption that f is p -integral, F is also p -integral.

Now let $\delta \in \Gamma_0^{(n)}(p^m)$, we may apply induction because F is of level $\Gamma_0^{(n)}(p^{m-1}N)$. We start from (8.1) for F and apply δ to it: If f is p -integral, then F is also p -integral, and then $F|\delta$ is also p -integral. By (8.1), we have

$$F|\delta = (f|\delta)\mathcal{F}^\alpha + \sum_{j=1}^{p-1} (f\mathcal{F}^j) |\gamma_w \delta$$

The first term on the right hand side of (8.1) is congruent mod p to $f|\delta$. We prove $(f\mathcal{F}^\alpha) |\gamma_w \delta \equiv 0 \pmod{p}$. To prove this, we must multiply (8.2) from the right by $\delta = \begin{pmatrix} a & b \\ p^m c & d \end{pmatrix}$. We get

$$\begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix} \gamma_w \delta = \begin{pmatrix} a & p^{m-1}b \\ wNa + pc & wNp^{m-1}b + d \end{pmatrix} \begin{pmatrix} p^{m-1} \cdot 1_n & 0_n \\ 0_n & 1_n \end{pmatrix}.$$

Since $p \nmid wNa$ and by a similar argument as above, we obtain $\mathcal{F}|\gamma_w \delta \equiv 0 \pmod{p}$. This shows $(f\mathcal{F}^\alpha) |\gamma_w \delta \equiv 0 \pmod{p}$ for large α . This implies that $F|\delta \equiv f|\delta \pmod{p}$ and hence $f|\delta$ is p -integral. \square

Remark 8.2. A minor variant of the proof above applies if f is of “quadratic nebentypus mod p ”, i.e. f satisfies (under the same conditions as in the theorem above) $f|_k \gamma = \left(\frac{\det(d)}{p}\right) \cdot f$ for all $\gamma \in \Gamma_0^{(n)}(p^m) \cap \Gamma^{(n)}(N)$ with the character $\left(\frac{*}{p}\right)$. The proof above needs to be modified only for the step “ $m = 1$ ”, where one has to use a function \mathcal{E} with nebentypus $\left(\frac{*}{p}\right)$.

9 Appendix B: Kitaoka’s formula

In this Appendix, we generalize Kitaoka’s theorem to higher degree to specify the level of theta series in our theorem. We then apply this in practice and discuss levels.

9.1 Generalization of Kitaoka's formula

We freely switch here between the language of (positive integral) quadratic forms S (or integral symmetric matrices) and the language of lattices L in an euclidean space $(\mathbb{R}^m, \langle \cdot, \cdot \rangle)$. The aim is to get a degree n version of a result of Kitaoka; we state it here only in the version sufficient for our application and allowing a rather short proof. A more sophisticated version generalizing the computation of Kitaoka would lead to a more general statement.

Theorem 9.1. *Let L be an even integral positive definite lattice of rank $m = 2k$ and level N . Let d be a positive divisor of N with $(d, \frac{N}{d}) = 1$, and $c := \frac{N}{d}$. Choose integers a, b such that $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and denote by \mathfrak{M} the n -fold diagonal embedding of M into $\mathrm{Sp}_n(\mathbb{Z})$, namely, $\mathfrak{M} := \begin{pmatrix} a \cdot 1_n & b \cdot 1_n \\ c \cdot 1_n & d \cdot 1_n \end{pmatrix}$.*

Then, with a suitable constant κ

$$\theta_L^{(n)}|_k \mathfrak{M} = \kappa \cdot \theta_{L'}^{(n)}, \quad (9.1)$$

where L' is a lattice characterized by

$$L' \otimes \mathbb{Z}_p = \begin{cases} L \otimes \mathbb{Z}_p & \text{if } p \nmid d, \\ (L \otimes \mathbb{Z}_p)^* & \text{if } p \mid d, \end{cases}$$

and $*$ denotes the dual.

Before we begin the proof, we need to prepare some more notation and recall the facts: Kitaoka [16] proved the degree one version of this statement for a more general type of theta series

$$\theta_L^{(1)}(q)(z) = \sum_{\mathfrak{x} \in L} q(\mathfrak{x}) e^{2\pi i \langle \mathfrak{x}, \mathfrak{x} \rangle \cdot z}$$

allowing homogeneous harmonic polynomials q on \mathbb{C}^m as coefficients in the theta series. An inspection of his proof shows that the constants of proportionality do NOT depend on those harmonic polynomials.

Ibukiyama [11] investigated holomorphic differential operators \mathcal{D} on \mathbb{H}_n , which are polynomials in the partial holomorphic derivatives, evaluated in $\mathrm{diag}(z_1, \dots, z_n)$; Such a \mathcal{D} maps holomorphic functions F on \mathbb{H}_n to holomorphic functions $\mathcal{D}(F)$ defined on \mathbb{H}_1^n such that, with the notation above

$$\mathcal{D}(F)|_k \mathfrak{M} = \mathcal{D}(F)|_{k_1}^{(z_1)} M \cdots |_{k_n}^{(z_n)} M$$

holds for arbitrary $M \in \mathrm{SL}_2(\mathbb{R})$ with certain weights $k_i \geq k$. The upper index z_i at the slash-operator indicates that $|_{k_i} M$ should be applied w.r.t. the variable z_i .

He also showed that from all the $\mathcal{D}(F)$ one can recover the Taylor expansion of F with respect to the non-diagonal coefficients of $Z = (z_{ij}) \in \mathbb{H}_n$. To show the

theorem, it is then enough, to show the equality of both sides after applying all such differential operators.

To do this we need some more notation: For a fixed differential operator \mathcal{D} of Ibukiyama type and $w_1, \dots, w_n \in \mathbb{C}^m$ changing the automorphy factor from \det^k to k_1, \dots, k_n in the variables z_1, \dots, z_n we define

$$\mathcal{D}e^{2\pi i \sum \langle w_i, w_j \rangle z_{ij}} = P(w_1, \dots, w_n) e^{2\pi i \sum \langle w_i, w_i \rangle z_i}$$

with $z_i := z_{ii}$. Then P is a harmonic polynomial in each of the variables w_i and we may write it as a finite sum of pure tensors:

$$P(w_1, \dots, w_n) = \sum_t q_{1,t}(w_1) \cdots q_{n,t}(w_n),$$

where the $q_{i,t}$ are harmonic polynomials in the variables w_i .

Proof of Theorem 9.1. Applying \mathcal{D} to the right hand side of (9.1) gives

$$\kappa \cdot \sum_t \theta_{L'}^{(1)}(q_{1,t})(z_1) \cdots \theta_{L'}^{(1)}(q_{n,t})(z_n).$$

Applying \mathcal{D} to the left hand side gives

$$\sum_t \theta_L^{(1)}(q_{1,t})(z_1) \cdots \theta_L^{(1)}(q_{n,t})(z_n) |_{k_1} M \cdots |_{k_n} M.$$

Now one has to use Kitaoka's result in degree 1 to get our assertion. \square

9.2 Application

Kitaoka's formula allows us to exclude certain congruences between modular forms and theta series of higher level (as far as the prime to p components of the levels are concerned):

Proposition 9.2. *Let n be an even positive integer and p a prime with $p \geq n + 1$. Let N, N' be coprime to p and $S \in \Lambda_n^+$ with level $p^a N$. Assume that*

$$p^b N' \mid p^a N \quad \text{and} \quad \phi \equiv \theta_S^{(n)} \pmod{p}$$

for some $\phi \in M_k(\Gamma_0^{(n)}(p^b N'), \chi)_{\mathbb{Z}(p)}$ with $\chi^2 = 1$. Then $N = N'$ and $\chi_N = (\chi_S)_N$.

Proof. We want to apply the modified q -expansion principle. To do so, we must modify ϕ and $\theta_S^{(n)}$ to arrive at the same weights and same (quadratic) p -component of nebentypus: We recall from Böcherer-Nagaoka [6] that there exist degree n modular forms \mathcal{E} for $\Gamma_0^{(n)}(p)$ of weight $\frac{p-1}{2}$ of nontrivial quadratic nebentypus. Then Proposition 2.4 allows us to choose $t \in \mathbb{N}$ such that $\theta_S^{(n)} \cdot \mathcal{E}^t$ and ϕ have the same

weight and the same p -component of nebentypus (this holds if $k \geq \frac{n}{2}$; the modification for the case of the opposite inequality is obvious). Then we can indeed apply Theorem 8.1 to $\frac{1}{p} \left(\phi - \theta_S^{(n)} \cdot \mathcal{E}^t \right)$.

We may enlarge b (if necessary) to get $a = b$. Assume that there is a prime q different from p with $q^r \parallel N$ and $q^s \parallel N'$ with $s < r$. We apply Kitaoka's theorem to $\theta_S^{(n)}$ with

$$M = \begin{pmatrix} * & * \\ p^a \cdot \frac{N}{q^r} & q^r \end{pmatrix}.$$

By the modified q -expansion principle Theorem 8.1, we have $\theta_S^{(n)} | \mathfrak{M} \equiv \pm \phi | \mathfrak{M} \pmod{p}$, where the \pm comes in from applying M to \mathcal{E}^t .

Let L correspond to S and correspond L' to S' . Then S' is a rational symmetric matrix with q^r appearing in its denominator. The Fourier expansion of $\theta_S^{(n)} | \mathfrak{M} = \kappa \cdot \theta_{S'}^{(n)}$ has a S' th Fourier coefficient which is

$$\kappa \cdot A(S', S').$$

By the condition $p > n + 1$, we see that $A(S', S') \not\equiv 0 \pmod{p}$. On the other hand, $\phi | \mathfrak{M}$ does not have a nonzero Fourier coefficient at S' , because the q -part of the width of the cusp \mathfrak{M} for $\Gamma_0^{(n)}(p^a N')$ is q^s with $s < r$. Therefore, $\kappa \cdot A(S', S')$ and hence κ must be divisible by p , i.e. $\kappa \in p \cdot R_{p^a}$. But the modified q -expansion principle tells us that $\theta_{S'}^{(n)} | \mathfrak{M}^{-1} = \frac{1}{\kappa} \cdot \theta_S^{(n)}$ is p -integral, in particular $\frac{1}{\kappa} \in R_{p^a}$. This is a contradiction, and we obtain $N' = N$.

The statement about the N -components of nebentypus characters follows by applying the modified q -expansion principle Theorem 8.1 with any integral matrix $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ satisfying $C \equiv 0_n \pmod{p^r N}$ and $\det(D) \equiv 1 \pmod{p}$. \square

Corollary 9.3. *Let n be a positive even integer and p a prime with $p > n + 3$. Let $S \in \Lambda_n^+$. Assume that $\theta_S^{(n)} \equiv \phi \pmod{p}$ for some $\phi \in M_k(\Gamma_0^{(n)}(N), \chi)$ with $\chi^2 = 1$. Then $\text{level}(S)$ is of the form “ p -power $\times N'$ ” with some $N' \mid N$ and $\chi = ' \chi_S$.*

Acknowledgment

The authors would like to thank Professors G. Nebe and R. Schulze-Pillot for helpful advices, especially on the order of automorphism groups for quadratic forms and on the generalization of Kitaoka's transformation formula. They would also like to thank Professor T. Yamauchi for giving them information on q -expansion principles. This work was supported by JSPS KAKENHI Grant Number 22K03259.

References

- [1] A. N. Andrianov, Quadratic forms and Hecke operators. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 286. Springer-Verlag, Berlin, 1987. xii+374 pp.
- [2] S. Böcherer, Elementary introduction to p -adic Siegel modular forms. L -functions and automorphic forms, 317-332, Contrib. Math. Comput. Sci., 10, Springer, Cham, 2017.
- [3] S. Böcherer, Über die Fourier-Jacobi-Entwicklung Siegelscher Eisensteinreihen. Math. Z. 183 (1983), no. 1, 21-46.
- [4] S. Böcherer, T. Kikuta, On mod p singular modular forms. Forum Math. 28 (2016), no. 6, 1051-1065.
- [5] S. Böcherer, S. Nagaoka, Congruences for Siegel modular forms and their weights. Abh. Math. Semin. Univ. Hambg. 80 (2010), no. 2, 227-231.
- [6] S. Böcherer, S. Nagaoka, On mod p properties of Siegel modular forms. Math. Ann. 338 (2007), no. 2, 421-433.
- [7] S. Böcherer, S. Nagaoka, On Siegel modular forms of level p and their properties mod p . Manuscripta Math. 132 (2010), no. 3-4, 501-515.
- [8] S. Böcherer, S. Raghavan, On Fourier coefficients of Siegel modular forms. J. Reine Angew. Math. 384 (1988), 80-101.
- [9] E. Freitag Stabile Modulformen. Math. Ann. 230 (1977), no. 3, 197-211.
- [10] E. Freitag, Singular modular forms and theta relations. Lecture Notes in Mathematics, 1487. Springer-Verlag, Berlin, 1991. iv+172 pp.
- [11] T. Ibukiyama, Generic differential operators on Siegel modular forms and special polynomials. Selecta Math. (N.S.) 26 (2020), no. 5, Paper No. 66, 50 pp.
- [12] T. Ichikawa, Congruences between Siegel modular forms. Math. Ann. 342 (2008), no. 3, 527-532.
- [13] E. Kani, The space of binary theta series. Ann. Sci. Math. Québec 36 (2012), no. 2, 501-534 (2013).
- [14] H. Katsurada, S. Nagaoka, A remark on p -adic Siegel Eisenstein series, arXiv:2204.01261v2 [math.NT].
- [15] N. M. Katz, A result on modular forms in characteristic p . Modular functions of one variable, V (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), pp. 53-61. Lecture Notes in Math., Vol. 601, Springer, Berlin, 1977.

- [16] Y. Kitaoka, A remark on the transformation formula of theta functions associated to positive definite quadratic forms. *J. Number Theory* 12 (1980), no. 2, 224-229.
- [17] H. Klingen, Zum Darstellungssatz für Siegelische Modulformen. *Math. Z.* 102 (1967), 30-43.
- [18] S. Nagaoka, A remark on Serre's example of p -adic Eisenstein series. *Math. Z.* 235 (2000), no. 2, 227-250.
- [19] A. Pitale, R. Schmit, A. Saha, Integrality and cuspidality of pullbacks of nearly holomorphic Siegel Eisenstein series. *Publ. Mat.* 66 (2022), no. 1, 405-434.
- [20] M. Raum, O. Richter, Sturm bounds for Siegel modular forms. *Res. Number Theory* 1 (2015), Paper No. 5, 8 pp.
- [21] H. L. Resnikoff, Automorphic forms of singular weight are singular forms. *Math. Ann.* 215 (1975), 173-193.
- [22] J.-P. Serre, Divisibilité de certaines fonctions arithmétiques. *Enseign. Math.* (2) 22 (1976), no. 3-4, 227-260.
- [23] J.-P. Serre, Formes modulaires et fonctions zêta p -adiques. Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 191-268. *Lecture Notes in Math.*, Vol. 350, Springer, Berlin, 1973.
- [24] G. Shimura, Differential operators, holomorphic projection, and singular forms. *Duke Math. J.* 76 (1994), no. 1, 141-173.
- [25] J. Sturm, On the congruence of modular forms. *Number theory* (New York, 1984-1985), 275-280, *Lecture Notes in Math.*, 1240, Springer, Berlin, 1987.
- [26] H. P. F. Swinnerton-Dyer, On l -adic representations and congruences for coefficients of modular forms. *Modular functions of one variable, III* (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 1-55. *Lecture Notes in Math.*, Vol. 350, Springer, Berlin, 1973.

Siegfried Böcherer
 Kunzenhof 4B
 79177 Freiburg, Germany
 Email: siegfried.boecherer@uni-mannheim.de

Toshiyuki Kikuta
 Faculty of Information Engineering
 Department of Information and Systems Engineering
 Fukuoka Institute of Technology
 3-30-1 Wajiro-higashi, Higashi-ku, Fukuoka 811-0295, Japan
 E-mail: kikuta@fit.ac.jp