

EFFECTIVE DIVISORS ON PROJECTIVIZED HODGE BUNDLES AND MODULAR FORMS

GERARD VAN DER GEER AND ALEXIS KOUVIDAKIS

ABSTRACT. We construct vector-valued modular forms on moduli spaces of curves and abelian varieties using effective divisors in projectivized Hodge bundles over moduli of curves. Cycle relations tell us the weight of these modular forms. In particular we construct basic modular forms for genus 2 and 3. We also discuss modular forms on the moduli of hyperelliptic curves. In that case the relative canonical bundle is a pull back of a line bundle on a \mathbb{P}^1 -bundle over the moduli of hyperelliptic curves and we extend that line bundle to a compactification so that its push down is (close to) the Hodge bundle and use this to construct modular forms.

1. INTRODUCTION

Moduli spaces of curves and of abelian varieties come with a natural vector bundle, the Hodge bundle \mathbb{E} . Starting from this vector bundle one can construct other natural vector bundles by applying Schur functors, like $\mathrm{Sym}^n(\mathbb{E})$ or $\det(\mathbb{E})^{\otimes m}$. Sections of such bundles are called modular forms. For example, for the moduli space \mathcal{A}_g of principally polarized abelian varieties of dimension g these are Siegel modular forms, and for the moduli space \mathcal{M}_g of curves of genus g these are Teichmüller modular forms. The Hodge bundle extends to appropriate compactifications of such moduli spaces and in many cases the sections also extend automatically to the compactifications, e.g. for \mathcal{A}_g with $g \geq 2$ by the so-called Koecher principle.

In this paper we try to construct modular forms in a geometric way. It is well-known that an effective divisor on \mathcal{A}_g or on $\overline{\mathcal{M}}_g$ with $g \geq 2$ representing the cycle class $m\lambda$ with $\lambda = c_1(\det(\mathbb{E}))$ and $m \in \mathbb{Z}_{>0}$ yields a scalar-valued modular form of weight m . We will exploit explicit effective divisors on projectivized vector bundles to construct vector-valued modular forms. In particular, we will construct in this way certain modular forms that play a pivotal role in low genera.

For example, in the case of $g = 2$ there is the modular form $\chi_{6,8}$, a section of $\mathrm{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^8$, that appeared in [4] as follows. Recall that the Torelli morphism $\mathcal{M}_2 \hookrightarrow \mathcal{A}_2$ has dense image and we have an equality of standard compactifications $\overline{\mathcal{M}}_2 = \tilde{\mathcal{A}}_2$. The moduli space \mathcal{M}_2 has another description as a stack quotient. This derives from the fact that a smooth complete curve of genus 2 over a field k of characteristic not 2 is a double cover of \mathbb{P}^1 ramified at six points, so can be given as $y^2 = f$ with f a polynomial of degree 6 with non-vanishing discriminant. Writing f as a homogeneous polynomial in two variables, say $f \in \mathrm{Sym}^6(W)$ with W the k -vector space generated by x_1, x_2 , and observing that we may change the base of W , we find a presentation of \mathcal{M}_2 as a stack

quotient

$$\mathcal{M}_2 \sim [W_{6,-2}^0/\mathrm{GL}(W)],$$

where we write $W_{a,b}$ for the $\mathrm{GL}(W)$ -representation $\mathrm{Sym}^a(W) \otimes \det(W)^b$. Here the space $W_{6,-2}$ can be seen as the vector space of binary sextics f with an action of $\mathrm{GL}(W)$ by

$$f(x_1, x_2) \mapsto (ad - bc)^{-2} f(ax_1 + bx_2, cx_1 + dx_2).$$

The subspace $W_{6,-2}^0$ of $W_{6,-2}$ is the space of f with non-vanishing discriminant. The twisting by $\det(W)^{-2}$ is required to get the right stabilizer for the generic f , namely $\pm \mathrm{Id}_W$.

This interpretation of \mathcal{M}_2 was used in [4] to construct vector-valued Siegel modular forms of degree 2 by using invariant theory of binary sextics, thus extending and simplifying the description of scalar-valued Siegel modular forms by invariants by Igusa [15, 16]. Covariants define vector-valued modular forms and all Siegel modular forms of degree 2 on \mathcal{A}_2 can be constructed this way. In [4] it was shown that the most basic covariant, the universal binary sextic, defines a meromorphic Siegel modular form $\chi_{6,-2}$ of weight $(6, -2)$, that is, it defines a meromorphic section of $\mathrm{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^{-2}$ on \mathcal{A}_2 . After multiplying $\chi_{6,-2}$ by Igusa's cusp form χ_{10} one obtains the holomorphic modular form $\chi_{6,8}$, the 'first' vector-valued Siegel modular cusp form of degree 2.

In the case of $g = 3$, there is an analogous form $\chi_{4,0,8}$, a section of $\mathrm{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$. Here it derives from the description of the moduli space \mathcal{M}_3^{nh} of non-hyperelliptic curves of genus three as a stack quotient

$$\mathcal{M}_3^{nh} \sim [W_{4,0,-1}^0/\mathrm{GL}(W)],$$

where W is now of dimension 3 and $W_{4,0,-1}^0 \subset \mathrm{Sym}^4(W) \otimes \det(W)^{-1}$ represents ternary quartics defining smooth curves. In [5] this description led to the construction of a meromorphic Teichmüller modular form $\chi_{4,0,-1}$ of weight $(4, 0, -1)$ and a (holomorphic) Siegel modular form $\chi_{4,0,8}$ of degree 3 and weight $(4, 0, 8)$. Also in this case all Teichmüller and Siegel modular forms of genus 3 on $\overline{\mathcal{M}}_3$ and \mathcal{A}_3 can be constructed from these forms by invariant theory.

This paper arises from the desire to construct these basic forms and similar forms in a geometric way. We use cycle relations for effective divisors (or almost effective divisors) on the projectivized Hodge bundle to construct our forms. It is based on the observation that an effective divisor D on the projectivized Hodge bundle $\mathbb{P}(\mathbb{E})$ with cycle class

$$[D] = [O(j)] + k\lambda - \Delta$$

with positive integers j, k and Δ an effective boundary class gives rise to a section of $\mathrm{Sym}^j(\mathbb{E}) \otimes \det(\mathbb{E})^k$ vanishing on boundary divisors, that is, a modular form. This method produces the basic modular forms $\chi_{6,8}$ and $\chi_{4,0,8}$ of degree 2 and 3 in an efficient way.

This connection between divisors and modular forms can also be used in the other direction, obtaining cycle classes for divisors on projectivized Hodge bundles. We give some examples of this.

Another objective of this paper is to construct modular forms on moduli spaces of hyperelliptic curves of genus g . For this we work with two descriptions of the moduli, a description as a stack quotient and a description as a Hurwitz space. The latter space

$\mathcal{H}_{g,2}$ has as compactification the space $\overline{\mathcal{H}}_{g,2}$ of admissible degree 2 covers of genus g . In the stack description modular forms pull back to covariants for the action of $\mathrm{GL}(2)$ on the space of binary forms of degree $2g + 2$.

In the Hurwitz space description the relative canonical bundle of the universal curve over $\mathcal{H}_{g,2}$ can be viewed as the pull back of $O(g - 1)$ from the trivial \mathbb{P}^1 -bundle P over $\mathcal{H}_{g,2}$ equipped with $2g + 2$ non-intersecting sections. Using the theory of admissible covers, P is compactified to a space \overline{P} , a fibration of rational stable curves with $2g + 2$ marked points over $\overline{\mathcal{H}}_{g,2}$, and we show that the line bundle $O(g - 1)$ on P extends to a line bundle on \overline{P} with the property that its push down to $\overline{\mathcal{H}}_{g,2}$ is close to the Hodge bundle. This allows us to construct modular forms on $\overline{\mathcal{H}}_{g,2}$.

ACKNOWLEDGEMENTS

We thank Fabien Cléry, Carel Faber and Gavril Farkas for useful remarks.

2. THE CASE OF GENUS TWO

Let k be a field of characteristic not 2. We consider the moduli space \mathcal{M}_2 of curves of genus 2 over k . This is a Deligne-Mumford stack and it carries a universal curve $\pi : \mathcal{C} \rightarrow \mathcal{M}_2$ of genus 2. The relative dualizing sheaf ω_π is base point free and thus defines a morphism $\varphi : \mathcal{C} \rightarrow \mathbb{P}(\mathbb{E})$. Projectivization is in the Grothendieck sense, so that for a vector space V the projective space $\mathbb{P}(V)$ parametrizes hyperplanes in V . For a curve C the map $\varphi : C \rightarrow \mathbb{P}(\mathbb{E}_C)$ associates to a point the space of differentials vanishing in that point. We have a commutative diagram with u the natural morphism

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\varphi} & \mathbb{P}(\mathbb{E}) \\ & \searrow \pi & \downarrow u \\ & & \mathcal{M}_2 \end{array}$$

We let $\overline{\mathcal{M}}_2$ be the Deligne-Mumford compactification and $\overline{\pi} : \overline{\mathcal{C}} \rightarrow \overline{\mathcal{M}}_2$ the corresponding universal curve. However, the extension $\omega_{\overline{\pi}}$ of ω_π does not define an extension of the morphism φ to $\mathbb{P}(\mathbb{E})$ over the boundary component Δ_1 that parametrizes reducible curves.

We consider the branch divisor $D \subset \mathbb{P}(\mathbb{E})$ of the morphism φ . The divisor D is of relative degree 6 in the \mathbb{P}^1 -bundle $\mathbb{P}(\mathbb{E})$ over the base \mathcal{M}_2 . We define \overline{D} to be the closure of D in $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_2$. In the rational Picard group of $\mathbb{P}(\mathbb{E})$ we can write

$$[\overline{D}] = [O(6)] + u^*(A)$$

with A a class in the rational Picard group of $\overline{\mathcal{M}}_2$ and u denotes the canonical projection $\mathbb{P}(\mathbb{E}) \rightarrow \overline{\mathcal{M}}_2$.

We want to determine A in terms of the generators λ, δ_0 of the Picard group of $\overline{\mathcal{M}}_2$. We write λ for the first Chern class of \mathbb{E} and δ_1 (resp. δ_0) for the class of Δ_1 (resp. Δ_0) in the Picard group of the stack $\overline{\mathcal{M}}_2$.

In order to do this we extend the morphism φ . First we extend it over a Zariski open part of Δ_0 . Let $f : C \rightarrow B$ be a 1-dimensional family of stable curves of genus 2 over a

discrete valuation ring (or disk) with smooth generic fibre and special fibre of type Δ_0 . A curve C of genus 2 of type Δ_0 is obtained by identifying two distinct points p and q on a curve E of genus 1. The fibre \mathbb{E}_C can be identified with $H^0(C, \omega) = H^0(E, O(p+q))$. The corresponding linear system is base point free, hence we get a surjective morphism $f^*(\mathbb{E}_B) \rightarrow \omega_f$ which extends our map φ .

For a similar family $f : C \rightarrow B$ over a discrete valuation ring with smooth generic fibre and special fibre of type Δ_1 , we cannot argue like that, since the corresponding linear system does not define such a map. In this case the special fibre is a union $C' \cup C''$ with C' and C'' of genus 1 intersecting in one nodal point p . We blow up this nodal point obtaining as special fibre a chain $C' + 2T + C''$ with T an exceptional curve and do a quadratic base change after which we have a family $\tilde{f} : \tilde{C} \rightarrow \tilde{B}$ with special fibre a chain of curves $C' + R + C''$, where now R is a (-2) -curve lying over T with degree 2. Instead of $\omega_{\tilde{f}}$ we now consider $\omega_{\tilde{f}}(-R)$ on our family. We claim that $\tilde{f}_*\omega_{\tilde{f}}(-R) = \mathbb{E}_{\tilde{B}}$. Indeed, the restriction of $\omega_{\tilde{f}}$ to $C' + R + C''$ is $(O_{C'}(p'), O_R, O_{C''}(p''))$, but since $H^0(C', O_{C'}(p')) = H^0(C', O_{C'})$ and $\omega_{\tilde{f}}|_R \cong O_R$, a section of the restriction of $\omega_{\tilde{f}}$ to the central fibre necessarily vanishes on R . By the exact sequence $0 \rightarrow \omega_{\tilde{f}}(-R) \rightarrow \omega_{\tilde{f}} \rightarrow \omega_{\tilde{f}}|_R \rightarrow 0$ and $\omega_{\tilde{f}}|_R = O_R$ we see that $\tilde{f}_*\omega_{\tilde{f}}(-R) \cong \tilde{f}_*(\omega_{\tilde{f}}) = \mathbb{E}_{\tilde{B}}$. The special fibre of $\tilde{f}_*(\omega_{\tilde{f}}(-R))$ is of codimension 1 in $H^0(C, \omega_{\tilde{f}}(-R))$ and defines a base point free linear system. For this we refer to Proposition 17.1 in the appendix Section 17. In this way we get a map $\varphi' : \tilde{C} \rightarrow \mathbb{P}(\mathbb{E}_{\tilde{B}})$ that is of degree 2 on R and contracts C' and C'' . It has the property that

$$\varphi'^*(O(1)) = \omega_{\tilde{f}}(-R),$$

with $O(1)$ the hyperplane bundle on $\mathbb{P}(\mathbb{E}_{\tilde{B}})$.

Proposition 2.1. *We have $[\overline{D}] = 6 [O(1)] + u^*(8\lambda - \delta_0 - \delta_1)$.*

Proof. We write $[\overline{D}] = 6 [O(1)] + u^*(A)$. We work with the above two types of 1-dimensional families $f : C \rightarrow B$. The morphism φ is ramified over D , and thus φ' is ramified over \overline{D} and contracts C' and C'' . We denote the ramification divisor by S . We thus get (writing abusively line bundles and divisors for the corresponding divisor classes)

$$\omega_{\tilde{f}} = \varphi'^*\omega_u + S + 2(C' + C''), \quad \varphi'^*\overline{D}/2 = S + 3(C' + C''),$$

where the first equation comes from adjunction $\omega_{\tilde{f}} + C'_{|C'} = O_{C'}$ for C' and similarly for C'' , and the second one from $C' \cdot \varphi'^*\overline{D} = 0 = C'' \cdot \varphi'^*\overline{D}$. This gives

$$\begin{aligned} \omega_{\tilde{f}} &= \varphi'^*(\omega_u + \overline{D}/2) - (C' + C'') \\ &= \varphi'^*(O(-2) + u^*(\lambda) + O(3) + u^*(A/2)) - (C' + C'') \\ &= \varphi'^*(O(1) + u^*(\lambda + A/2)) - (C' + C'') \\ &= \omega_{\tilde{f}} - R + \tilde{f}^*(\lambda + A/2) - (C' + C'') \\ &= \omega_{\tilde{f}} + \tilde{f}^*(\lambda + A/2 - b_1) \end{aligned}$$

with b_1 the special point of \tilde{B} . This shows that $A = -2\lambda + 2b_1$. Because of the base change that we executed, we have $2b_1 = \delta_1$ and we obtain $A = -2\lambda + \delta_1$. Now we use the well-known relation $10\lambda = \delta_0 + 2\delta_1$ (see [21]) and thus get $A = -2\lambda + \delta_1 = 8\lambda - \delta_0 - \delta_1$. \square

Remark 2.2. We give an alternative proof of this result in Remark 14.2.

We remark that $u_*(O(1)) = \mathbb{E}$ and $u_*(O(m)) = \text{Sym}^m(\mathbb{E})$ for $m \geq 1$. The divisor \overline{D} with

$$[\overline{D}] = [O(6)] + u^*(8\lambda - \delta_0 - \delta_1)$$

is an effective divisor on $\mathbb{P}(\mathbb{E})$. We apply u_* to the corresponding section 1 of $O(\overline{D})$. By Proposition 2.1 we see that we get a regular section $\chi_{6,8}$ of the vector bundle $\text{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^8$ over $\overline{\mathcal{M}}_2$. Moreover, this section vanishes on the divisors Δ_0 and Δ_1 . Note that the Torelli map extends to an isomorphism $\overline{\mathcal{M}}_2 \cong \tilde{\mathcal{A}}_2$ with $\tilde{\mathcal{A}}_2$ the standard smooth compactification of \mathcal{A}_2 . Therefore our section defines a Siegel modular form $\chi_{6,8}$ of weight $(6, 8)$ that is a cusp form.

The relation $10\lambda = \delta_0 + 2\delta_1$ quoted above implies that there exists a Siegel modular cusp form χ_{10} of degree 2 and of weight 10 with divisor $\delta_0 + 2\delta_1$. The quotient $\chi_{6,-2} := \chi_{6,8}/\chi_{10}$ defines a meromorphic section of $\text{Sym}^6(\mathbb{E}) \otimes \det \mathbb{E}^{-2}$ that is regular outside Δ_1 .

Corollary 2.3. *Let \overline{D} be the closure in $\mathbb{P}(\mathbb{E})$ of the branch divisor of the canonical map for the universal curve over \mathcal{M}_2 . The push forward $u_*(s)$, with s the natural section 1 of $O(\overline{D})$ on $\mathbb{P}(\mathbb{E})$, defines a Siegel modular cusp form $\chi_{6,8}$ of degree 2 and weight $(6, 8)$.*

We now analyze the orders of vanishing along δ_1 of $\chi_{6,8}$ and $\chi_{6,-2}$. When identifying $\overline{\mathcal{M}}_2$ with $\tilde{\mathcal{A}}_2$ we also write $\mathcal{A}_{1,1}$ for δ_1 ; it is the locus of products of elliptic curves.

We analyze the orders by working locally on a family over a local base B with central fibre a general point b_1 of the boundary divisor Δ_1 . As we mentioned before, the map $\varphi : \mathcal{C} \rightarrow \mathbb{P}(\mathbb{E})$ defined over \mathcal{M}_2 does not extend to the whole $\overline{\mathcal{C}}$ over $\overline{\mathcal{M}}_2$ due to the fact that the canonical system has base points at the nodes of the curves over the boundary divisor Δ_1 . On the other hand, by the theory of admissible covers, the ramification divisor of the above map φ extends to a divisor S on $\overline{\mathcal{C}}$ in a way that avoids the above nodal locus. Namely, over $b_1 \in \Delta_1$ the fibre is a nodal curve C which is the union of two elliptic curves C_1 and C_2 meeting at a point p . The restriction of the ramification divisor on each component is the union of the three -additional to p - ramification points of the system $|O(2p)|$. Therefore the extension of the map φ is defined on the ramification divisor S . The map φ maps $C_1 \setminus \{p\}$ and $C_2 \setminus \{p\}$ to two distinct points p_1 and p_2 respectively which are defined as follows. The fibre of $\mathbb{P}(\mathbb{E})$ over b_1 can be identified with $\mathbb{P}(H^0(C, \omega_C))$. The points of $H^0(C, \omega_C)$ have the form (s_1, s_2) , with s_i a section of $H^0(C_i, O(p))$. Then the point p_1 corresponds to the hyperplane $\{(0, s_2), s_2 \in H^0(C_2, O(p))\}$ and the point p_2 corresponds to the hyperplane $\{(s_1, 0), s_1 \in H^0(C_1, O(p))\}$.

The divisor \overline{D} , the image of the ramification divisor under the extended map φ , splits then into six irreducible components denoted by D_1, \dots, D_6 . Over our local base B we thus have the six local sections D_i ($i = 1, \dots, 6$) of the family $\mathbb{P}(\mathbb{E}) \rightarrow B$. By the above

description of the extension of the map φ , we may conclude that D_1, D_2, D_3 pass through p_1 and D_4, D_5, D_6 through p_2 .

Lifting the sections D_i locally to sections σ_i of \mathbb{E} and choosing a basis e_1, e_2 of \mathbb{E} over B such that e_1 and e_2 determine p_1 and p_2 in the fibre of $\mathbb{P}(\mathbb{E})$ over $z = 0$, we can write $\sigma_i = a_i e_1 + b_i e_2$ for $i = 1, \dots, 6$. Then at $z = 0$ the functions b_1, b_2, b_3 and a_4, a_5, a_6 vanish, while $a_1, a_2, a_3, b_4, b_5, b_6$ do not vanish. Since by blowing up once we can separate, we may assume that these sections vanish with order 1 at $z = 0$. By construction the section χ of $\text{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^{-2}$ is locally given by

$$\frac{\sigma_1 \cdots \sigma_6}{z}.$$

We may write $\sigma_1 \cdots \sigma_6$ as

$$\begin{aligned} & a_1 \cdots a_6 e_1^6 + (a_1 a_2 a_3 a_4 a_5 b_6 + \dots + b_1 a_2 a_3 a_4 a_5 a_6) e_1^5 e_2 + \\ & (a_1 a_2 a_3 a_4 b_5 b_6 + \dots + b_1 b_2 a_3 a_4 a_5 a_6) e_1^4 e_2^2 + \\ & (a_1 a_2 a_3 b_4 b_5 b_6 + \dots + b_1 b_2 b_3 a_4 a_5 a_6) e_1^3 e_2^3 + \dots + b_1 \cdots b_6 e_2^6. \end{aligned}$$

The order at $z = 0$ of the coefficient of $e_1^i e_2^{6-i}$ equals

$$\min_{\#\Lambda=i} (\#\Lambda^c \cap \{1, 2, 3\} + \#\Lambda \cap \{4, 5, 6\})$$

with Λ running through the subsets of $\{1, \dots, 6\}$ of cardinality i and Λ^c denoting the complement. We find for these orders $(3, 2, 1, 0, 1, 2, 3)$ for $i = 0, \dots, 6$, hence for the section χ given by $\sigma_1 \cdots \sigma_6 / z$ we find the orders $(2, 1, 0, -1, 0, 1, 2)$.

Corollary 2.4. *The section 1 of the line bundle $O(\overline{D})$ on $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_2$ pushes down via $\mathbb{P}(\mathbb{E}) \rightarrow \overline{\mathcal{M}}_2$ to the meromorphic modular form $\chi_{6,-2}$ on $\overline{\mathcal{M}}_2 = \tilde{\mathcal{A}}_2$. The orders of the seven coordinates of $\chi_{6,-2}$ along $\mathcal{A}_{1,1}$ in \mathcal{A}_2 are $(2, 1, 0, -1, 0, 1, 2)$.*

These orders are in agreement with the result of [4] where $\chi_{6,-2}$ was constructed by invariant theory and properties of modular forms were used to determine these orders.

A different way to construct the form $\chi_{6,8}$ uses the so-called Weierstrass divisor W in the dual bundle:

$$W := \{(C, \eta) \in \mathbb{P}(\mathbb{E}^\vee) : \text{div}(\eta) \text{ contains a Weierstrass point}\}$$

over \mathcal{M}_2 . Here C denotes a curve of genus 2 and η a differential form on C . We let \overline{W} be the closure of W over $\overline{\mathcal{M}}_2$. We then have an identity due to Gheorghita [11, Thm 1]

$$[\overline{W}] = 6 [O_{\mathbb{P}(\mathbb{E}^\vee)}(1)] + 34 \lambda - 3 \delta_0 - 5 \delta_1,$$

where we write λ and δ_i for the pullback of λ and δ_i to $\mathbb{P}(\mathbb{E}^\vee)$. Now \overline{W} is an effective divisor and the push forward of the section 1 of $O(\overline{W})$ is a section of $\text{Sym}^6(\mathbb{E}^\vee) \otimes \det(\mathbb{E})^{34} \otimes O(-3\delta_0 - 5\delta_1)$. For $g = 2$ we have $\mathbb{E}^\vee \cong \mathbb{E} \otimes \det(\mathbb{E})^{-1}$, hence $\text{Sym}^6(\mathbb{E}^\vee) \cong \text{Sym}^6(\mathbb{E}) \otimes \det(\mathbb{E})^{-6}$. This implies that under the isomorphism of \mathbb{P}^1 -bundles $\mathbb{P}(\mathbb{E}) \cong \mathbb{P}(\mathbb{E}^\vee)$ the isomorphism identifies $[\overline{W}]$ with $[\overline{D}]$, and we get in the dual bundle

$$[\overline{W}] = 6 [O(1)] + 28 \lambda - 3 \delta_0 - 5 \delta_1 = 6 [O(1)] + 8 \lambda - \delta_0 - \delta_1.$$

Using push forward we find again a form of weight $(6, 8)$ vanishing on δ_1 and δ_0 . Up to a multiplicative non-zero constant this is $\chi_{6,8}$.

Remark 2.5. The identity $[\overline{W}] = 6 [O(1)] + 28 \lambda - 3 \delta_0 - 5 \delta_1$ implies that there exists a modular form of weight $(6, 28)$ vanishing with multiplicity 3 on Δ_0 and multiplicity 5 on Δ_1 , but this is (up to a multiplicative constant) the form $\chi_{10}^2 \chi_{6,8}$ with χ_{10} the form of weight 10 with divisor $\delta_0 + 2\delta_1$ that displays the relation $10 \lambda = \delta_0 + 2\delta_1$.

3. THE CASE OF GENUS THREE

Here there is no restriction on the characteristic. We consider the moduli stack \mathcal{M}_3 of curves of genus 3 over our field k and the universal curve $\pi : \mathcal{C} \rightarrow \mathcal{M}_3$. The canonical map defines a morphism $\varphi : \mathcal{C} \rightarrow \mathbb{P}(\mathbb{E})$ and we thus obtain the image divisor D in $\mathbb{P}(\mathbb{E})$ over \mathcal{M}_3 . We have a diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\varphi} & \mathbb{P}(\mathbb{E}) \\ & \searrow \pi & \downarrow u \\ & & \mathcal{M}_3 \end{array}$$

We consider the closure \overline{D} of D in $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_3$, the Deligne-Mumford compactification of \mathcal{M}_3 . The canonical image of the generic curve is a quartic curve. Thus we have a relation in the rational Picard group of $\mathbb{P}(\mathbb{E})$

$$[\overline{D}] = [O(4)] + u^*(A),$$

with A a divisor class on $\overline{\mathcal{M}}_3$. We now determine A by using test families.

Proposition 3.1. *We have $A = 8\lambda - \delta_0 + c\delta_1$ for some integer c .*

Remark 3.2. The constant c equals -2 as we shall see later, see Lemma 14.1.

Proof. We write the class A as $A = a\lambda + b\delta_0 + c\delta_1$. The map φ can be extended over $\overline{\mathcal{M}}_3 - \Delta_1$, but the relative dualizing sheaf $\omega_{\overline{\mathcal{C}}}$ restricted to a fibre of type Δ_1 has base points. Thus we consider the morphism $\varphi' : \overline{\mathcal{C}} \rightarrow \mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_3 - \Delta_1$. We use test families without fibres of type Δ_1 . The first type of test family is a pencil of quartic curves defined by $sF + tG$ with F and G ternary quartics defining generic smooth quartic curves in \mathbb{P}^2 . This is a family $\mathcal{X} \rightarrow B$ with $B = \mathbb{P}^1$. By standard theory (see [8, 14.5.1] or [13, Table 3.167, p. 189]) the degrees of the classes λ, δ_0 and δ_1 on B are given by

$$\deg(\lambda, \delta_0, \delta_1)|_B = (3, 27, 0).$$

In particular this family does not have hyperelliptic fibres since the class \mathfrak{h} of the hyperelliptic locus $\overline{\mathcal{H}}_3$ in $\overline{\mathcal{M}}_3$ satisfies (cf. [12, p. 140])

$$\mathfrak{h} = 9\lambda - \delta_0 - 3\delta_1. \tag{1}$$

In view of the relation (where again we abusively write line bundles for their corresponding divisor classes)

$$\begin{aligned}\omega_\pi &= \varphi'^*(\overline{D} + \omega_u) \\ &= \varphi'^*(O(4) + u^*(A) + O(-3) + u^*(\lambda)) \\ &= \varphi'^*(O(1) + u^*(A + \lambda))\end{aligned}$$

and the fact that $\omega_\pi = \varphi'^*(O(1))$ we find $A|_B = -\lambda|_B$. This implies $3a + 27b = -3$, so

$$a + 9b = -1. \quad (2)$$

The second test family is a family $sF + tQ^2$ with F a generic smooth quartic and Q a smooth conic. For such a family we have

$$\deg(\lambda, \delta_0, \delta_1)|_B = (3/2, 13, 0). \quad (3)$$

see [13, Table 3.167, p. 189]. We now have

$$\begin{aligned}\omega_f &= \varphi'^*((\overline{D} + \omega_u)|_{\overline{D}} - \mathfrak{h}) \\ &= \varphi'(O(4) + u^*(A) + O(-3) + u^*(\lambda) - u^*(\mathfrak{h})) \\ &= \varphi'(O(1) + u^*(A + \lambda - \mathfrak{h})),\end{aligned}$$

where we used the fact that φ' restricted to a generic fibre is of degree 1, but of degree 2 on a hyperelliptic fibre leading to the correction term $-\mathfrak{h}$. On the other hand $\omega_\pi = \varphi'^*(O(1))$, hence we find $A + \lambda - \mathfrak{h} = 0$. Using (1) and (3) this gives

$$(a + 1 - 9)\frac{3}{2} + (b + 1)13 + (c - 3)0 = 0, \quad \text{that is, } 3a + 26b = -2. \quad (4)$$

The relations (2) and (4) determine a and b . \square

The divisor \overline{D} is effective over $\overline{\mathcal{M}}_3 - \Delta_1$. Because of the relation

$$[\overline{D}] = [O(4)] + u^*(8\lambda - \delta_0 - c\delta_1)$$

the corresponding section 1 of $O(\overline{D})$ maps under u_* to a section ψ of $\text{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$ that is regular outside Δ_1 and vanishes on Δ_0 . This section ψ is invariant under the action of -1 on the fibres of \mathbb{E} . Therefore it descends to a section $\chi_{4,0,8}$ of $\text{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$ on the image of $\overline{\mathcal{M}}_3 - \Delta_1$ under the Torelli morphism $\overline{\mathcal{M}}_3 \rightarrow \tilde{\mathcal{A}}_3$, with $\tilde{\mathcal{A}}_3$ the standard second Voronoi compactification of \mathcal{A}_3 . Since the image of Δ_1 in $\tilde{\mathcal{A}}_3$ is of codimension 2, the section $\chi_{4,0,8}$ extends to a regular section of $\text{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$ on all of \mathcal{A}_3 , and then by the Koecher Principle it extends to $\tilde{\mathcal{A}}_3$. Thus it defines a regular Siegel modular cusp form $\chi_{4,0,8}$ of degree 3 and weight $(4, 0, 8)$.

Corollary 3.3. *Let \overline{D} be the closure of the canonical curve over \mathcal{M}_3 in $\mathbb{P}(\mathbb{E})$ and s the natural section 1 of $O(\overline{D})$. Then $\chi_{4,0,8} = u_*(s)$ is a Teichmüller modular form and it descends to a Siegel modular cusp form of degree 3 and weight $(4, 0, 8)$.*

The relation (1) shows that there exists a scalar-valued Teichmüller modular form χ_9 of weight 9 on $\overline{\mathcal{M}}_3$. Its square is invariant under the action of -1 on the fibres of \mathbb{E} ,

hence descends to a Siegel modular form of weight 18. Up to a multiplicative scalar this is Igusa's modular form χ_{18} .

If we divide $\chi_{4,0,8}$ by χ_9 we obtain a meromorphic section of $\text{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^{-1}$ on $\overline{\mathcal{M}}_3$ that is regular on \mathcal{M}_3 outside the hyperelliptic locus. This form was used in [5] to construct Teichmüller modular forms and Siegel modular forms by invariant theory.

4. MODULI OF HYPERELLIPTIC CURVES AS A STACK QUOTIENT

In this section we discuss the stack quotient description of the moduli of hyperelliptic curves. We consider hyperelliptic curves in characteristic not 2. A hyperelliptic curve of genus g is a morphism $\alpha : C \rightarrow \mathbb{P}^1$ of degree 2 where C is a smooth curve of genus g . A morphism $a : \alpha \rightarrow \alpha'$ between two hyperelliptic curves is a commutative diagram

$$\begin{array}{ccc} C & \longrightarrow & C' \\ \alpha \downarrow & & \downarrow \alpha' \\ \mathbb{P}^1 & \longrightarrow & \mathbb{P}^1 \end{array}$$

A hyperelliptic curve C of genus g can be written as $y^2 = f(x)$ with $f \in \mathbb{C}[x]$ of degree $2g + 2$. In fact, choosing a basis (x_1, x_2) of the g_2^1 defines the morphism α . Let $W = \langle x_1, x_2 \rangle$, a vector space (over our algebraically closed base field) of dimension 2, and $L = \alpha^*(\mathcal{O}_{\mathbb{P}^1}(1))$. By Riemann-Roch we have $\dim H^0(C, L^{g+1}) = g + 3$, while $\dim \text{Sym}^{g+1}(W) = g + 2$, so we have a non-zero element $y \in H^0(C, L^{g+1})$ which is anti-invariant under the involution corresponding to α . The anti-invariant subspace of $H^0(C, L^{g+1})$ has dimension 1. Then y^2 is invariant and lies in $\text{Sym}^{2g+2}(W)$. Thus we find the equation $y^2 = f(x_1, x_2)$ with f homogeneous of degree $2g + 2$ and with non-zero discriminant.

We have made two choices here: a generator y of $H^0(C, L^{g+1})^{(-1)}$, a space of dimension 1, and a basis of W . We can change the choice of y (by a non-zero scalar) and the choice of a basis of W by $\gamma = (a, b, c, d) \in \text{GL}(W)$. The action of $\text{GL}(W)$ is on the right via

$$f(x_1, x_2) \mapsto f(ax_1 + bx_2, cx_1 + dx_2).$$

If we let $\text{GL}(W)$ act on y by a power of the determinant, then this action preserves the type of equation. In inhomogeneous form the action by $\text{GL}(W)$ is by

$$f \mapsto f\left(\frac{ax + b}{cx + d}\right), \quad y \mapsto y/(cx + d)^{g+1},$$

with the following effect on the equation:

$$y^2 = f(x) \mapsto y^2 = (cx + d)^{2g+2} f\left(\frac{ax + b}{cx + d}\right).$$

The last expression on the right-hand-side can be written as binary form of degree $2g + 2$.

The stabilizer of a generic $f \in \text{Sym}^{2g+2}(W)$ is μ_{2g+2} , the roots of unity of order dividing $2g + 2$. Since we want a stabilizer of order 2 for the generic element, we consider a twisted action: define the $\text{GL}(W)$ -representation

$$W_{a,b} = \text{Sym}^a(W) \otimes \det(W)^{\otimes b}.$$

This can be identified with $\text{Sym}^a(W)$ as a vector space, but the action by $\text{GL}(W)$ is different. Inside this space $W_{a,b}$ we have the open subspace $W_{a,b}^0$ of homogeneous polynomials of degree a with non-zero discriminant. We now distinguish two cases.

Case 1. g even. Here we consider the stack quotient

$$[W_{2g+2,-g}^0/\text{GL}(W)].$$

This stack quotient can be identified with the moduli stack \mathcal{H}_g of hyperelliptic curves of genus g for g even. Indeed, the action of $t \cdot \text{Id}_W$ is (on inhomogeneous equations) by

$$f \mapsto t^{-2g} f, \quad y \mapsto y/t^{g+1},$$

hence $y^2 = f$ maps to $y^2 = t^2 f$, so that the stabilizer is μ_2 , as required. Note also that the action of $-1 \in \text{GL}(W)$ is by $y \mapsto -y$, so y is an odd element. A basis of $H^0(C, \Omega_C^1)$ is given by

$$x^i dx/y, \quad (i = 0, \dots, g-1).$$

The action on dx is by $(ad - bc)dx/(cx + d)^2$ resulting in the action on the space of differentials by

$$x^i dx/y \mapsto (ad - bc)(cx + d)^{g-1-i}(ax + b)^i dx/y,$$

If we forget the twisted action on y , we can identify $H^0(C, \Omega_C^1)$ with $W_{g-1,1}$. But y^2 must be viewed as an element of $W_{2g+2,-g}$, so the action of $t 1_W$ on y should be twisted by $t^{-g} = \det^{-g/2}$. We get

$$H^0(C, \Omega_C^1) \cong W_{g-1, (2-g)/2} \quad \text{for } g \text{ even.}$$

We see that under the identification $h : [W_{2g+2,-g}^0/\text{GL}(W)] \xrightarrow{\sim} \mathcal{H}_g$ the pullback $h^*(\mathbb{E})$ of the Hodge bundle \mathbb{E} is the equivariant bundle $W_{g-1, (2-g)/2}$. The action of -1_W is by -1 on $W_{g-1, (2-g)/2}$. We also observe $h^*(\det(\mathbb{E})) = \det(W)^{g/2}$.

Case 2. g odd. Here we take $W_{2g+2,-g+1}$.

Remark 4.1. If we consider $W_{2g+2,r}$ then r has to be even, since as above we later view y^2 as an element of $W_{2g+2,r}$ and we need an action by $\det^{r/2}$ on y .

Here the stabilizer of a generic element is μ_4 . Now on inhomogeneous equations the action is by

$$f \mapsto t^{-2g+2} f, \quad y \mapsto y/t^{g+1},$$

hence $y^2 = f$ maps to $y^2 = t^4 f$. Note that here -1_W acts by $f \mapsto f$ and $y \mapsto y$. But $\sqrt{-1}_W$ acts by $f \mapsto f$ and $y \mapsto -y$. To get the right stack quotient with stabilizer of the generic element of order 2, we take

$$[W_{2g+2,1-g}^0/(\text{GL}(W)/(\pm 1_W))].$$

The action on the differentials $x^i dx/y$ with $i = 0, \dots, g-1$ is by

$$x^i dx/y \mapsto (ad - bc)^{(1-g)/2}(cx + d)^{g-1-i}(ax + b)^i,$$

hence without twisting we get $H^0(C, \Omega_C^1) = W_{g,1}$. Since we view y^2 as element of $W_{g+3,1-g}$ we find under $h : [W_{2g+2,1-g}^0/(\text{GL}(W)/(\pm 1_W))]$ that $h^*(\mathbb{E}) = W_{g-1, (3-g)/2}$. The element $\sqrt{-1} 1_W$ acts on the differentials as $(-1)^{(3-g)/2}(\sqrt{-1})^{g-1} = -1$.

We summarize.

Proposition 4.2. *Writing $W_{a,b} = \text{Sym}^a(W) \otimes \det(W)^b$ we have the identification of stacks*

$$h^{-1} : \mathcal{H}_g \xrightarrow{\sim} \begin{cases} [W_{2g+2,-g}^0/\text{GL}(W)] & g \text{ even} \\ [W_{2g+2,1-g}^0/(\text{GL}(W)/(\pm 1_W))] & g \text{ odd,} \end{cases}$$

and

$$h^*(\mathbb{E}) \cong \begin{cases} W_{g-1,(2-g)/2} & g \text{ even} \\ W_{g-1,(3-g)/2} & g \text{ odd,} \end{cases} \quad h^*(\det(\mathbb{E})) = \begin{cases} \det(W)^{g/2} & g \text{ even} \\ \det(W)^g & g \text{ odd.} \end{cases}$$

For a somewhat different description see [2, Cor. 4.7, p. 654].

Recall that the moduli stack \mathcal{H}_g has as compactification the closure $\overline{\mathcal{H}}_g$ of \mathcal{H}_g inside the moduli stack $\overline{\mathcal{M}}_g$. The Picard group of \mathcal{H}_g is known by [2] to be finite cyclic for $g \geq 2$ of order $4g+2$ if g is even and $8g+4$ else. The rational Picard group of $\overline{\mathcal{H}}_g$ is known (see [7]) to be free abelian of rank g generated by classes δ_i and ξ_j for $i = 0, \dots, \lfloor g/2 \rfloor$ and $j = 1, \dots, \lfloor (g-1)/2 \rfloor$. Cornalba gives also the first Chern class λ of the Hodge bundle \mathbb{E} on $\overline{\mathcal{H}}_g$

$$(8g+4)\lambda = g\delta_0 + 4 \sum_{i=1}^{\lfloor g/2 \rfloor} i(g-i)\delta_i + 2 \sum_{i=1}^{\lfloor (g-1)/2 \rfloor} (i+1)(g-i)\xi_i,$$

where the generic point of the divisor ξ_i has an admissible model $C' \cup C''$ with two nodes $C' \cap C'' = \{p, q\}$ mapping to a union of two \mathbb{P}^1 , with $2i+2$ marked points on C' , see Figure 1 in Section 6.

5. MODULAR FORMS ON THE HYPERELLIPTIC LOCUS OF GENUS THREE

Let \mathbb{E} be the Hodge bundle on $\overline{\mathcal{H}}_3$. By a modular form of weight k on $\overline{\mathcal{H}}_3$ we mean a section of $\det(\mathbb{E})^{\otimes k}$. The construction in the preceding section shows that a modular form of weight k on $\overline{\mathcal{H}}_3$ when pulled back to the stack $[W_{2,0}^0/(\text{GL}(W)/\pm \text{id}_W)]$ gives rise to an invariant of degree $3k/2$. Indeed, it defines a section of the equivariant bundle $\det(W)^{3k}$ invariant under $\text{SL}(W)$, but in view of the fact that we divide by the action of $\text{GL}(W)/(\pm \text{id}_W)$ this yields an invariant of degree $3k/2$.

Let $M_k(\Gamma_3) = H^0(\mathcal{A}_3, \det(\mathbb{E})^k)$ be the space of Siegel modular forms of degree 3 on $\Gamma_3 = \text{Sp}(6, \mathbb{Z})$. In [17] Igusa considered an exact sequence

$$0 \rightarrow M_{k-18}(\Gamma_3) \xrightarrow{\cdot \chi_{18}} M_k(\Gamma_3) \rightarrow I_{3k/2}(2, 8)$$

with $I_d(2, 8)$ the vector space of invariants of degree d of binary octics. We can interpret Igusa's sequence in the following way. A Siegel modular form of weight k defines by restriction to the hyperelliptic locus a modular form of weight k on $\overline{\mathcal{H}}_3$ and it thus defines an invariant of degree $3k/2$.

For each irreducible representation ρ of $\text{GL}(3)$ we have a vector bundle \mathbb{E}_ρ made from \mathbb{E} by a Schur functor. By a modular form of weight ρ on $\overline{\mathcal{H}}_3$ we mean a section of a line bundle \mathbb{E}_ρ . We can pull back to the stack $[W_{8,-2}^0/(\text{GL}(W)/\pm \text{id}_W)]$, but the situation

is more involved as $\mathrm{Sym}^n(\mathrm{Sym}^2(W))$ decomposes as a representation of $\mathrm{GL}(W)$. For example, we have

$$h^*(\mathrm{Sym}^4(\mathbb{E})) = \mathrm{Sym}^4(\mathrm{Sym}^2(W)) = W_{8,0} \oplus W_{4,2} \oplus W_{0,4}.$$

Here and in the rest of this section we assume that the characteristic is 0, or not 2, and high enough for the representation theory (plethysm) to work¹. In this case we can consider the restriction of the Siegel modular form $\chi_{4,0,8}$ to the hyperelliptic locus and we know that it does not vanish identically by [5, Lemma 7.7]. On the other hand we have the basic covariant $f_{8,-2}$, the diagonal section of $W_{8,-2}$ over the stack $[W_{8,-2}^0/(\mathrm{GL}(W)/\pm \mathrm{id}_W)]$.

The discriminant form \mathfrak{d} of binary octics, an invariant of degree 14, does not define a modular form, but its third power \mathfrak{d}^3 does. It defines a modular form of weight 28, see [24, p. 811].

Via the projection $p_{8,0} : \mathrm{Sym}^4(\mathrm{Sym}^2(W)) \rightarrow W_{8,0}$ a section of $\mathrm{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$ defines a covariant of bi-degree $(8, 24/2) = (8, 12)$ for the action of $\mathrm{GL}(W)$.

Proposition 5.1. *The restriction to the hyperelliptic locus of the section $\chi_{4,0,8}$ corresponds via the projection $p_{8,0}$ to the covariant $f_{8,-2} \cdot \mathfrak{d}$ with \mathfrak{d} the discriminant of binary octics.*

Proof. By restricting and projecting we obtain a covariant of bi-degree $(8, 12)$. This covariant is divisible by the discriminant and does not vanish on the locus of smooth hyperelliptic curves. Therefore, division by \mathfrak{d} gives a non-vanishing covariant of bidegree $(8, -2)$. Taking into account the ‘twisting’ by $\det(W)^{-2}$ this must be a multiple of the universal binary octic. \square

We will discuss the other two projections later. Note that the divisor D , the canonical image of the universal curve in $\mathbb{P}(\mathbb{E})$ that defines $\chi_{4,0,8}$, has a restriction to the locus of smooth hyperelliptic curves which is divisible by 2. Indeed, the canonical image of a hyperelliptic curve is a double conic. This suggests that we can take the ‘square root’ of the restriction of $\chi_{4,0,8}$ to the hyperelliptic locus. However, the boundary divisors prevent this. If we take a level cover of the moduli space we can construct a modular form of weight $(2, 0, 4)$. We will carry this out later (in Corollary 12.4), working on a Hurwitz space that we shall introduce in the next section.

6. THE HURWITZ SPACE OF ADMISSIBLE COVERS OF DEGREE TWO

In this and the following sections will use the other description of the moduli of hyperelliptic curves, namely moduli space $\overline{\mathcal{H}}_{g,2}$ of admissible covers of degree 2 and genus g in the sense of [14], see [13]. Thus we are looking at covers $f : C \rightarrow P$ of degree 2 with C nodal of genus g and P a stable b -pointed curve of genus 0. Here the $b = 2g + 2$ branch points are ordered and $\mathcal{H}_{g,2} \rightarrow \mathcal{H}_g$ is a Galois cover with Galois group the symmetric group \mathfrak{S}_{2g+2} .

¹Alternatively one could use divided powers as in [1, 3.1]

The boundary $\overline{\mathcal{H}}_{g,2} - \mathcal{H}_{g,2}$ consists of finitely many divisors that we shall denote by $\Delta_b^\Lambda = \Delta^\Lambda$, where we omit the index b if g is clear. Here the index Λ defines a partition

$$\{1, 2, \dots, b\} = \Lambda \sqcup \Lambda^c,$$

and the generic point of Δ^Λ corresponds to an admissible cover that maps to a stable curve of genus 0 that is the union of two copies of \mathbb{P}^1 , one containing the points with mark in Λ , the other one those with mark in Λ^c . Here we will assume that $\#\Lambda = j$ with $2 \leq j \leq g+1$.

The parity of $\#\Lambda$ plays an important role here. If $\#\Lambda = 2i+2$ is even, then the generic admissible cover corresponding to a point of Δ^Λ is a union $C_i \cup C_{g-i-1}$ that is a double cover of a union of two rational curves \mathbb{P}_1 and \mathbb{P}_2 with C_i lying over \mathbb{P}_1 and C_{g-i-1} over \mathbb{P}_2 . Here C_i (resp. C_{g-i-1}) has genus i (resp. $g-i-1$) with $0 \leq i \leq (g-1)/$ and is ramified over the points of Λ (resp. Λ^c).




Fig. 1: Λ even

If $\#\Lambda = 2i+1$ is odd with $1 \leq i \leq g/2$, then we have a union $C_i \cup C_{g-i}$ lying over $\mathbb{P}_1 \cup \mathbb{P}_2$, where C_i (resp. C_{g-i}) of genus i (resp. $g-i$) is ramified over Λ and in p (resp. over Λ^c and in p). Note that p is a simple node.




Fig. 2: Λ odd

Assuming that g and $b = 2g+2$ are fixed we will write

$$\Delta_j = \sum_{\#\Lambda=j} \Delta^\Lambda \quad \text{with } 2 \leq j < b/2$$

and provide the symmetric case with a factor $1/2$, that is, $\Delta_{b/2} = \frac{1}{2} \sum_{\#\Lambda=b/2} \Delta^\Lambda$.

7. DIVISORS ON THE MODULI OF STABLE CURVES OF GENUS ZERO

For later use we recall some notation and facts concerning divisors on the moduli spaces $\overline{\mathcal{M}}_{0,n}$. We refer to [18]. The boundary divisors on $\overline{\mathcal{M}}_{0,n}$ are denoted by S_n^Λ and are indexed by partitions $\{1, \dots, n\} = \Lambda \sqcup \Lambda^c$ into two disjoint sets with $2 \leq \#\Lambda \leq n-2$ and we have $S_n^\Lambda = S_n^{\Lambda^c}$. Via the natural map $\pi_{n+1} : \overline{\mathcal{M}}_{0,n+1} \rightarrow \overline{\mathcal{M}}_{0,n}$ we may view $\overline{\mathcal{M}}_{0,n+1}$ as the universal curve and π_{n+1} has n sections. The generic point of S_n^Λ corresponds to a stable curve with two rational components, one of which contains the points marked by Λ . For pullback by π_{n+1} we have the relation

$$\pi_{n+1}^*(S_n^\Lambda) = S_{n+1}^{\{\Lambda, n+1\}} \cup S_{n+1}^{\{\Lambda^c, n+1\}}.$$

The n sections of π_{n+1} have images $S_{n+1}^{\{i, n+1\}}$ with $i = 1, \dots, n$.

We can collect these boundary divisors on $\overline{\mathcal{M}}_{0,n+1}$ via

$$T_{n+1,j} = \sum_{\#\Lambda=j} S_{n+1}^{\{\Lambda, n+1\}}, \quad T_{n+1,j}^c = \sum_{\#\Lambda=j} S_{n+1}^{\{\Lambda^c, n+1\}},$$

with the convention that in view of the symmetry we add a factor $1/2$ for even n and $j = n/2$

$$T_{n+1,n/2} = \frac{1}{2} \sum_{\#\Lambda=n/2} S_{n+1}^{\{\Lambda, n+1\}}, \quad T_{n+1,n/2}^c = \frac{1}{2} \sum_{\#\Lambda=n/2} S_{n+1}^{\{\Lambda^c, n+1\}}.$$

Later, when a fixed index k is given we will split these divisors as $T = T(k^+) + T(k^-)$ where (k^+) (resp. (k^-)) indicates that the sum is taken over Λ containing k (resp. not containing k). So $T_{n+1,j}(k^+) = \sum_{\#\Lambda=j, k \in \Lambda} S_{n+1}^{\{\Lambda, n+1\}}$ (and with a factor $1/2$ if $j = n/2$).

8. A GOOD MODEL

We now will work with a ‘good model’ of the universal admissible cover over $\overline{\mathcal{H}}_{g,2}$. Such a model was constructed in [10, Section 4]. We start with the observation that the space $\overline{\mathcal{H}}_{g,2}$ is not normal, and we therefore normalize it. The result $\tilde{\mathcal{H}}_{g,2}$ is now a smooth stack over which we have a universal curve $\tilde{\mathcal{C}} \rightarrow \tilde{\mathcal{H}}_{g,2}$.

We have a natural map $h : \tilde{\mathcal{H}}_{g,2} \rightarrow \overline{\mathcal{M}}_{0,b}$ with $b = 2g + 2$ and the universal curve now fits into a commutative diagram

$$\begin{array}{ccc} \tilde{\mathcal{C}} & \xrightarrow{c} & \overline{\mathcal{M}}_{0,b+1} \\ \varpi \downarrow & & \downarrow \pi_{b+1} \\ \overline{\mathcal{H}}_{g,2} & \xleftarrow{\nu} \tilde{\mathcal{H}}_{g,2} \xrightarrow{h} & \overline{\mathcal{M}}_{0,b} \end{array}$$

We can construct a proper flat map that extends the relative canonical morphism $\mathcal{C} \rightarrow \mathbb{P}_{\overline{\mathcal{H}}_{g,2}}^1$ by taking the fibre product \mathbb{P} of $\overline{\mathcal{M}}_{0,b+1}$ and $\tilde{\mathcal{H}}_{g,2}$ over $\overline{\mathcal{M}}_{0,b}$ and thus obtain a

commutative diagram

$$\begin{array}{ccccc}
 \tilde{\mathcal{C}} & \xrightarrow{\alpha} & \mathbb{P} & \xrightarrow{c'} & \overline{\mathcal{M}}_{0,b+1} \\
 & \searrow \varpi & \downarrow \varpi' & & \downarrow \pi_{b+1} \\
 & & \tilde{\mathcal{H}}_{g,2} & \xrightarrow{h} & \overline{\mathcal{M}}_{0,b}
 \end{array}$$

The resulting space \mathbb{P} is not smooth, but has rational singularities. Resolving these in a minimal way gives a model $\tilde{\mathbb{P}}$; taking the resolution \tilde{Y} of the normalization Y of the fibre product of $\tilde{\mathbb{P}}$ and $\tilde{\mathcal{C}}$ over \mathbb{P} gives us finally a commutative diagram

$$\begin{array}{ccc}
 \tilde{Y} & \xrightarrow{f} & \tilde{\mathbb{P}} \\
 & \searrow t & \downarrow \pi \\
 & & B \xrightarrow{\varphi} \tilde{\mathcal{H}}_{g,2}
 \end{array}$$

where B is our base $\tilde{\mathcal{H}}_{g,2}$ or any other base mapping to it. We write π for the resulting morphism $\tilde{\mathbb{P}} \rightarrow B$, h for the natural map $B \rightarrow \overline{\mathcal{M}}_{0,b}$ and ν for $B \rightarrow \overline{\mathcal{H}}_{g,2}$. We refer to [10, Section 4] for additional details.

In the following we will assume that we have a physical family over a base B . We will abuse the notation Δ^Λ for the pull back of the divisor Δ^Λ under $\nu : B \rightarrow \overline{\mathcal{H}}_{g,2}$.

In the case that $\#\Lambda$ is even, say $\#\Lambda = 2i + 2$ with $0 \leq i \leq (g-1)/2$, the pull back of Δ^Λ decomposes as

$$\pi^*(\Delta^\Lambda) = \Pi^\Lambda + \Pi^{\Lambda^c},$$

corresponding to the two components of a general fibre of π , with Π^Λ mapping to $S_{b+1}^{\{\Lambda, b+1\}}$ under $\tilde{\mathbb{P}} \rightarrow \overline{\mathcal{M}}_{0,b+1}$, and similarly Π^{Λ^c} mapping to $S_{b+1}^{\{\Lambda^c, b+1\}}$. Note that we restrict $\#\Lambda$ by $\leq g+1$, hence the notation Π^Λ should not lead to confusion.

In the case $\#\Lambda$ is odd we find a similar decomposition

$$\pi^*(\Delta^\Lambda) = \Pi^\Lambda + R^\Lambda + \Pi^{\Lambda^c},$$

corresponding now to the fact that the general fibre of π has three components, one coming from the blowing up.

We notice

$$h^*(S_b^\Lambda) = \begin{cases} \Delta^\Lambda & \#\Lambda \equiv 0 \pmod{2} \\ 2\Delta^\Lambda & \#\Lambda \equiv 1 \pmod{2} \end{cases}$$

If we use the notation $\Delta_j = \sum_{\#\Lambda=j} \Delta^\Lambda$, we find for the tautological classes $\lambda = c_1(\mathbb{E})$ and $h^*(\psi_k)$, simply denoted by ψ_k , ($k = 1, \dots, b$) on our base B , the following formulas (see [9])

$$\lambda = \sum_{i=0}^{(g-1)/2} \frac{(i+1)(g-i)}{2(2g+1)} \Delta_{2i+2} + \sum_{i=1}^{g/2} \frac{i(g-i)}{2g+1} \Delta_{2i+1} \quad (5)$$

and

$$\begin{aligned} \psi_k = & \sum_{i=0}^{(g-1)/2} \left(\frac{(g-i)(2g-2i-1)}{g(2g+1)} \Delta_{2i+2}(k^+) + \frac{(i+1)(2i+1)}{g(2g+1)} \Delta_{2i+2}(k^-) \right) \\ & + 2 \sum_{i=1}^{g/2} \left(\frac{(g-i)(2g-2i+1)}{g(2g+1)} \Delta_{2i+1}(k^+) + \frac{i(2i+1)}{g(2g+1)} \Delta_{2i+1}(k^-) \right) \end{aligned} \quad (6)$$

where we use the notation (k^+) (resp. (k^-)) to denote the condition $k \in \Lambda$ (resp. $k \notin \Lambda$) as above. The relation (5) implies the following.

Corollary 8.1. *There exists a scalar-valued modular form of weight $2(2g+1)$ on the moduli space $\widetilde{\mathcal{H}}_{g,2}$ whose divisor is a union of boundary divisors. It descends to the hyperelliptic locus $\overline{\mathcal{H}}_g$ and corresponds to a power of the discriminant of the binary form of degree $2g+2$.*

9. EXTENDING THE LINEAR SYSTEM

The canonical system on a hyperelliptic curve is defined by the pull back of the sections of the line bundle of $O(g-1)$ of degree $g-1$ on the projective line. We now try to extend this line bundle over our compactification.

A first attempt would be to consider the divisor $(g-1)\tilde{S}_k$ with \tilde{S}_k the pullback to $\widetilde{\mathbb{P}}$ of the section S_k of $\pi_{b+1} : \overline{\mathcal{M}}_{0,b+1} \rightarrow \overline{\mathcal{M}}_{0,b}$. We can add a boundary divisor Ξ_k to it such that $f^*O_{\widetilde{\mathbb{P}}}(D_k)$ with $D_k = (g-1)\tilde{S}_k + \Xi_k$ coincides with ω_t on the fibres of t , namely take Ξ_k equal to

$$\begin{aligned} & \sum_{i=0}^{(g-1)/2} \left((g-1-i) \Pi_{2i+2}(k^+) + i \Pi_{2i+2}^c(k^-) \right) + \\ & \sum_{i=1}^{g/2} \left((g-i-1) \Pi_{2i+1}(k^+) - (g-i) \Pi_{2i+1}^c(k^+) \right) + \sum_{i=1}^{g/2} \left((i-1) \Pi_{2i+1}^c(k^-) - i \Pi_{2i+1}(k^-) \right) \end{aligned}$$

Here $\Pi_j = \sum_{\#\Lambda=j} \Pi^\Lambda$ and $\Pi_j^c = \sum_{\#\Lambda=j} \Pi^{\Lambda^c}$ and (k^+) (resp. (k^-)) indicates the condition that $k \in \Lambda$ (resp. $k \notin \Lambda$); moreover, we add a factor $1/2$ in case $j = b/2$.

Now $f^*O(D_k)$ and ω_t agree on the fibres of t , so they differ by a pull back under $t = \pi \circ f$, see diagram (1). We therefore will change D_k by a pull back under π . Define a divisor class on B by

$$\begin{aligned} E_k = & \frac{2g-1}{2} \psi_k - \sum_{i=0}^{(g-1)/2} \left((g-i-1) \Delta_{2i+2}(k^+) + i \Delta_{2i+2}(k^-) \right) \\ & - \sum_{i=1}^{g/2} \left((g-i-1) \Delta_{2i+1}(k^+) + (i-1) \Delta_{2i+1}(k^-) \right) \end{aligned}$$

and define a line bundle on $\widetilde{\mathbb{P}}$ by

$$M = O(D_k + \pi^* E_k). \quad (7)$$

Lemma 9.1. *The line bundle M does not depend on k , satisfies $f^*(M) = \omega_t$ and restricts to the general fibre \mathbb{P}^1 of π as $O(g-1)$. For $\#\Lambda = 2i+2$ its restriction to the general fibre $\mathbb{P}_1 \cup \mathbb{P}_2$ over Δ_b^Λ is of degree $(i, g-i-1)$, while for $\#\Lambda = 2i+1$ its restriction to the general fibre $\mathbb{P}_1 \cup R \cup \mathbb{P}_2$ it is of degrees $(i, -1, g-i)$.*

Proof. We use the section τ_k of $t : \tilde{Y} \rightarrow B$ with $f\tau_k = \tilde{s}_k$ with \tilde{s}_k the natural sections of the map π with image \tilde{S}_k . Then we have $\tau_k^*\omega_t = \psi_k/2$ and $\tau_k^*f^*D_k = \tilde{s}_k^*D_k$ for which we have

$$\begin{aligned} \tilde{s}_k^*D_k &= -(g-1)\psi_k + \sum_{i=0}^{(g-1)/2} ((g-i-1)\Delta_{2i+2}(k^+) + i\Delta_{2i+2}(k^-)) \\ &\quad + \sum_{i=1}^{g/2} ((g-1-i)\Delta_{2i+1}(k^+) + (i-1)\Delta_{2i+1}(k^-)). \end{aligned}$$

From this we obtain $\tau_k^*(\omega_t) - \tau_k^*f^*D_k = \pi^*(E_k)$, so that $\omega_t = f^*(O(D_k + E_k))$. We also see that the restriction of M on the fibres of π does not depend on k . Moreover, we have

$$\tilde{s}_j^*(D_k + \pi^*E_k) = \tau_j^*f^*(D_k + E_k) = \tau_j^*(\omega_t) = \psi_j/2 = \tilde{s}_j^*(D_j + \pi^*E_j),$$

showing that the restrictions of $O(D_k + \pi^*E_k)$ and $O(D_j + \pi^*E_j)$ agree on \tilde{S}_j . The restrictions of the fibres of π over the general points of Δ_b^Λ are easily checked. \square

We now want to compare $\pi_*(M)$ with the Hodge bundle $\mathbb{E} = t_*(\omega_t)$ on B . The next proposition shows that these agree up to codimension 2.

Proposition 9.2. *We have an exact sequence $0 \rightarrow \pi_*(M) \rightarrow \mathbb{E} \rightarrow \mathcal{T} \rightarrow 0$, where \mathcal{T} is a coherent sheaf that is a torsion sheaf supported on the boundary. Moreover, we have $c_1(\pi_*(M)) = \lambda$.*

Proof. By Lemma 9.1 we have $\omega_t = f^*(M)$. But $R^1\pi_*(M) = (0)$, so we have

$$\pi_*(M \otimes f_*O_{\tilde{Y}}) = \pi_*f_*(f^*(M)) = \pi_*f_*(\omega_t) = t_*(\omega_t).$$

We have an exact sequence $0 \rightarrow O_{\tilde{\mathbb{P}}} \rightarrow f_*O_{\tilde{Y}} \rightarrow \mathcal{F} \rightarrow 0$ with \mathcal{F} a coherent sheaf of rank 1 that restricted to the smooth fibers of π has degree $-(g+1)$, as one sees by applying Riemann-Roch to f and $O_{\tilde{Y}}$. Tensoring the sequence with M and applying π_* gives the exact sequence

$$0 \rightarrow \pi_*(M) \rightarrow \pi_*(M \otimes f_*O_{\tilde{Y}}) \rightarrow \pi_*(M \otimes \mathcal{F}) \rightarrow 0.$$

On the smooth fibers of π the sheaf $M \otimes \mathcal{F}$ restricts to a line bundle of degree $(g-1) - (g+1) = -2$, hence $\pi_*(M \otimes \mathcal{F})$ is a torsion sheaf.

We now calculate $c_1(\pi_*(M))$. We apply Grothendieck-Riemann-Roch to π and $O(D_k)$. It says

$$\text{ch}(\pi_!(O(D_k))) = \pi_*(\text{ch}(O(D_k)))\text{Td}^\vee(\omega_\pi),$$

which by $\pi_*(\text{Td}_2^\vee(\omega_\pi)) = 0$ gives

$$c_1(\pi_*(O(D_k))) = \frac{1}{2}\pi_*(-D_k\omega_\pi + D_k^2).$$

We calculate

$$\pi_*(D_k \omega_\pi) = (g-1)\psi_k - \sum_{i=0}^{(g-1)/2} ((g-i-1)\Delta_{2i+2}(k^+) + i\Delta_{2i+2}(k^-)) + \sum_{i=1}^{g/2} \Delta_{2i+1}, \quad (8)$$

and

$$\begin{aligned} \pi_*(D_k^2) &= -(g-1)^2\psi_k + \sum_{i=0}^{(g-1)/2} (g-i-1)(g+i-1)\Delta_{2i+2}(k^+) \\ &\quad + \sum_{i=0}^{(g-1)/2} (2g-2-i)i\Delta_{2i+2}(k^-) + \sum_{i=1}^{g/2} ((2g-i-1)(i-1) - i^2)\Delta_{2i+1}. \end{aligned} \quad (9)$$

Adding $\pi_*(\pi^*E_k)$ gives

$$\begin{aligned} c_1(\pi_*(M)) &= \frac{g^2}{2}\psi_k - \frac{1}{2} \sum_{i=0}^{(g-1)/2} ((g-i-1)(g-i)\Delta_{2i+2}(k^+) + i(i+1)\Delta_{2i+2}(k^-)) \\ &\quad - \sum_{i=1}^{g/2} ((g-i)^2\Delta_{2i+1}(k^+) + i^2\Delta_{2i+1}(k^-)). \end{aligned}$$

Substituting the formula for ψ_k we find

$$c_1(\pi_*(M)) = \sum_{i=0}^{(g-1)/2} \frac{(g-i)(i+1)}{2(2g+1)}\Delta_{2i+2} + \sum_{i=1}^{g/2} \frac{i(g-i)}{2g+1}\Delta_{2i+1} = \lambda.$$

□

The line bundle M on $\tilde{\mathbb{P}}$ is not base point free as Proposition 9.2 shows; the restriction to the R -part has negative degree. We can make it base point free by defining

$$N = M(-R) = O(D_k + \pi^*E_k - R). \quad (10)$$

Now the restriction of N to a general fibre over Δ_{2i+1} , which is a chain of three rational curves $\mathbb{P}_1, R, \mathbb{P}_2$, has degrees $(i-1, 1, g-i-1)$ and one checks that N is base point free.

Lemma 9.3. *Up to codimension 2 we have on B that $\pi_*(N) = \mathbb{E}$.*

Proof. We have $R^1\pi_*(N) = 0$. Therefore the exact sequence $0 \rightarrow N \rightarrow M \rightarrow M|_R \rightarrow 0$ yields the exact sequence

$$0 \rightarrow \pi_*(N) \rightarrow \pi_*(M) \rightarrow \pi_*(M|_R) \rightarrow 0.$$

We now show that $c_1(\pi_*(N)) = c_1(\pi_*(M))$. Since $R^1\pi_*(M) = 0 = R^1\pi_*(N)$ we find by Grothendieck-Riemann-Roch that

$$c_1(\pi_*(M)) = \frac{1}{2}\pi_*(c_1(M)^2 - c_1(M)\omega_\pi), \quad c_1(\pi_*(N)) = \frac{1}{2}\pi_*(c_1(N)^2 - c_1(N)\omega_\pi).$$

By the definition of N and the fact that R is a (-2) -curve if we take a base B of dimension 1, and thus has intersection number 0 with a fibre, we have

$$\pi_*(c_1(N))^2 = \pi_*(c_1(M))^2$$

and $c_1(N)\omega_\pi = c_1(M)\omega_\pi$ since the restriction of ω_π to R is trivial. \square

10. THE RATIONAL NORMAL CURVE

The image of a hyperelliptic curve by the canonical map is a rational normal curve, that is, \mathbb{P}^1 embedded in \mathbb{P}^{g-1} via the linear system of degree $g-1$.

In our setting we can see the rational normal curve and its degenerations using the extension N of the line bundle of degree $g-1$, as defined in (10), to the compactification as constructed in the preceding section.

We let $u : \mathbb{P}(\mathbb{E}) \rightarrow B$ be natural projection. Now N is base point free and up to codimension 2 we have $\pi_*(N) = \mathbb{E}$, so the global-to-local map $\pi^*\pi_*(N) \rightarrow N$ induces a surjective map $\nu : \pi^*(\mathbb{E}) \rightarrow N$ over $\tilde{\mathbb{P}}$. This induces a morphism $\varphi : \tilde{\mathbb{P}} \rightarrow \mathbb{P}(\mathbb{E})$ by associating to a point of $\tilde{\mathbb{P}}$ the kernel of ν . It fits into a diagram

$$\begin{array}{ccc} \tilde{\mathbb{P}} & \xrightarrow{\varphi} & \mathbb{P}(\mathbb{E}) \\ & \searrow \pi & \downarrow u \\ & & B \end{array}$$

Proposition 10.1. *For a point of B with smooth fibre under π the image of φ is a rational normal curve of degree $g-1$. For a general point $\beta \in \Delta_{2i+2}$ with fibre $\mathbb{P}_1 \cup \mathbb{P}_2$ the image is a union of two rational normal curves of degree i and $g-i-1$. For a general point $\beta \in \Delta_{2i+1}$ with fibre $\mathbb{P}_1, R, \mathbb{P}_2$ the image is a union of three rational normal curves of degree $i-1, 1$ and $g-i-1$. Here we interpret the case of degree 0 as a contracted curve.*

Proof. The proposition follows almost immediately from Lemma 9.1. \square

Remark 10.2. If $i = 1$ then \mathbb{P}_1 is contracted. If also $g = 2$ then both \mathbb{P}_1 and \mathbb{P}_2 are contracted and the image of R coincides with the fibre of $\mathbb{P}(\mathbb{E})$.

Remark 10.3. The sections $\tilde{s}_i : B \rightarrow \tilde{\mathbb{P}}$ for $i = 1, \dots, b$ induce sections $\sigma_i = \varphi \circ s_i : B \rightarrow \mathbb{P}(\mathbb{E})$ by sending β to the kernel of $\mathbb{E} = \tilde{s}_i^*\pi^*(\mathbb{E}) \rightarrow s_i^*(N)$.

Remark 10.4. In the case $g = 2$ the map φ is a birational map $\tilde{\mathbb{P}} \rightarrow \mathbb{P}(\mathbb{E})$ that blows down boundary components. More precisely, over Δ_2 it blows down Π_2 and over Δ_3 the components supported at $\Pi_3 = \Pi_3^c$.

11. SYMMETRIZATION

We have been working with the moduli space $\mathcal{H}_{g,2}$ and $\mathcal{M}_{0,b}$ and their compactifications. Here the symmetric group \mathfrak{S}_b acts. We therefore make our construction symmetric.

We put $D = \sum_{k=1}^b D_k$ and $E = \sum_{k=1}^b E_k$ and set

$$\tilde{M} = O(D + E), \quad \psi = \sum_{k=1}^b \psi_k, \quad \text{and} \quad \tilde{S} = \sum_{k=1}^b \tilde{S}_k.$$

We find

$$\psi = 4 \sum_{i=0}^{(g-1)/2} \frac{(g-i)(i+1)}{2g+1} \Delta_{2i+2} + 2 \sum_{i=1}^{g/2} \frac{(2g-2i+1)(2i+1)}{2g+1} \Delta_{2i+1}$$

and

$$D = (g-1)\tilde{S} + 2 \sum_{i=0}^{(g-1)/2} ((g-i-1)(i+1)\Pi_{2i+2} + i(g-i)\Pi_{2i+2}^c) + \\ \sum_{i=1}^{g/2} ((g-4i-1)\Pi_{2i+1} - (3g-4i+1)\Pi_{2i+1}^c)$$

and

$$E = \frac{2g-1}{2}\psi - 2 \sum_{i=0}^{(g-1)/2} ((g-i)(2i+1) - (i+1)) \Delta_{2i+2} - \sum_{i=1}^{g/2} (4i(g-i) - (g+2)) \Delta_{2i+1}$$

12. THE CASE OF HYPERELLIPTIC GENUS THREE

The Hurwitz space $\mathcal{H}_{3,2}$ admits a compactification $\overline{\mathcal{H}}_{3,2}$ with boundary components Δ^Λ with $\#\Lambda \in \{2, 3, 4\}$. Taking the components with Λ of fixed cardinality together gives boundary components Δ_2, Δ_3 and Δ_4 . Under the morphism $\overline{\mathcal{H}}_{3,2} \rightarrow \overline{\mathcal{M}}_3$ the components Δ_2 and Δ_4 are mapped to δ_0 , while Δ_3 goes to δ_1 . The formulas (5) and (6) specialize to

$$\lambda = \frac{3}{14}\Delta_2 + \frac{2}{7}\Delta_3 + \frac{2}{7}\Delta_4. \quad (11)$$

and

$$\psi_k = \frac{5}{7}\Delta_2(k^+) + \frac{1}{21}\Delta_2(k^-) + \frac{20}{21}\Delta_3(k^+) + \frac{2}{7}\Delta_3(k^-) + \frac{2}{7}\Delta_4.$$

Remark 12.1. The equation (11) shows that on $\overline{\mathcal{H}}_{3,2}$ there exists a scalar-valued modular form of weight 14 whose square equals χ_{28} , a form mentioned in Section 5. Since on $\overline{\mathcal{H}}_3$ we have $28\lambda = 3\delta_0 + 8\Delta_1 + 8\xi_1$, an integral class not divisible by 2, there is not a modular form of weight 14 on $\overline{\mathcal{H}}_3$ with square χ_{28} .

We have the line bundle M on $\tilde{\mathbb{P}}$ defined in (7) corresponding to the divisor class $D_k + E_k$ given by

$$D_k = 2\tilde{S}_k + 2\Pi_2(k^+) + 2\Pi_4(k^+) + \Pi_3(k^+) - 2\Pi_3^c(k^+) - \Pi_3(k^-)$$

and

$$E_k = \frac{5}{2}\psi_k - (2\Delta_2(k^+) + \Delta_3(k^+) + \Delta_4),$$

where ψ_k is given in (6). Define the rational divisor class

$$U := \frac{1}{14}\Delta_2 + \frac{3}{7}(\Delta_3 + \Delta_4) = \frac{3}{2}\psi_k - (\Delta_2(k^+) + \Delta_3(k^+)).$$

The divisor class of $D_k + E_k$ is independent of k as observed in Lemma 9.1, but this can be seen also directly from the next lemma.

Lemma 12.2. *We have the linear equivalence $D_k + E_k \sim -\omega_\pi + \Pi_2 + \Pi_3 + \pi^*(U)$.*

Proof. One checks that $-\omega_\pi + \Pi_2 + \Pi_3$ and $D_k + E_k$ have the same restriction to fibres of π . We have $s_k^*(-\omega_\pi + \Pi_2 + \Pi_3) = -\psi_k + \Delta_2(k^+) + \Delta_3(k^+)$ and $s_k^*(D_k + E_k) = \psi_k/2$. \square

Let Q be the image of $\varphi : \tilde{\mathbb{P}} \rightarrow \mathbb{P}(\mathbb{E})$, see Proposition 10.1. The map φ is the composition of a map $\varphi' : \tilde{\mathbb{P}} \rightarrow Q$ with the inclusion map $\iota : Q \hookrightarrow \mathbb{P}(\mathbb{E})$. The generic fibre of $Q \rightarrow B$ is a conic, hence $O(Q) = O(2) \otimes O(u^*A)$ for some divisor A on B . We determine A .

Lemma 12.3. *On $\mathbb{P}(\mathbb{E})$ we have the linear equivalence*

$$[Q] \sim [O(2)] + u^*(4\lambda - (\Delta_2 + \Delta_3 + \Delta_4)).$$

Proof. We have $\omega_{\mathbb{P}(\mathbb{E})} \otimes u^*(\omega_B^{-1}) = O(-3) \otimes u^*(\det \mathbb{E})$ and by adjunction $\omega_Q = \iota^*(O(Q) \otimes \omega_{\mathbb{P}(\mathbb{E})})$. Since φ' is a blow down we have $\omega_{\tilde{\mathbb{P}}} = (\varphi')^*\omega_Q \otimes O(\Pi_2 + \Pi_3)$. We get

$$\begin{aligned} \varphi^*(O(Q)) &= \varphi'^*\omega_Q \otimes \varphi'^*\omega_{\tilde{\mathbb{P}}}^{-1} \\ &= \omega_{\tilde{\mathbb{P}}} \otimes O(-\Pi_2 - \Pi_3) \otimes \varphi'^*O(3) \otimes \pi^*\det(\mathbb{E})^{-1} \otimes \pi^*\omega_B^{-1} \\ &= \omega_\pi \otimes O(-\Pi_2 - \Pi_3) \otimes \varphi'^*O(3) \otimes \pi^*\det(\mathbb{E})^{-1}. \end{aligned}$$

On the other hand we have $\varphi^*O(Q) = \varphi^*O(2) \otimes O(\pi^*A)$ and $\varphi^*O(1) = N$, hence we get

$$O(u^*A) = N \otimes \omega_\pi \otimes O(-\Pi_2 - \Pi_3) \otimes \pi^*\det(\mathbb{E})^{-1}. \quad (12)$$

By Lemma 12.2 we have $N = \omega_\pi^{-1} \otimes O(\Pi_2 + \Pi_3) \otimes O(U)$. Substituting this in (12) we get the desired result. \square

The effective divisor Q yields a modular form and Lemma 12.3 gives its weight.

Corollary 12.4. *The effective divisor Q on $\mathbb{P}(\mathbb{E})$ defines a modular form $\chi_{2,0,4}$ on $\tilde{\mathcal{H}}_{3,2}$ of weight $(2, 0, 4)$, that is, a non-zero section of $\text{Sym}^2(\mathbb{E}) \otimes \det(\mathbb{E})^4$.*

Since the divisor $\Delta_2 + \Delta_3 + \Delta_4$ is not a pull back from the moduli space $\overline{\mathcal{H}}_3$, the modular form does not descend to $\overline{\mathcal{H}}_3$. Recall that the modular form $\chi_{4,0,8}$ restricted to the hyperelliptic locus was associated to a divisor D that equals $2Q$.

13. COMPARISON WITH THE HODGE BUNDLE

We know by Lemma 9.3 that the line bundle $N = O_{\tilde{\mathbb{P}}}(D_k + E_k - R)$ on $\tilde{\mathbb{P}}$ over $\tilde{\mathcal{H}}_{3,2}$ has the property that $\pi_*(N) \cong \mathbb{E}$ up to codimension 2. We now deal with the push forward of the tensor powers of N .

Lemma 13.1. *We have for $m \in \mathbb{Z}_{\geq 1}$*

$$c_1(\pi_*(N^{\otimes m})) = \frac{2m^2 + m}{14} \Delta_2 + \frac{5m^2 - m}{14} (\Delta_3 + \Delta_4).$$

Proof. We apply Grothendieck-Riemann-Roch to π and $N^{\otimes m}$ as in (9) in the proof of Proposition 9.2. Recall that N corresponds to the divisor(class) $D_k + E_k - R$. We use that $R^1\pi_*N^{\otimes m} = 0$ for all m and find

$$c_1(\pi_*(N^{\otimes m})) = \frac{1}{2}\pi_*(m^2(D_k + E_k - R)^2 - m\omega_\pi \cdot (D_k + E_k - R))$$

and using the relations (8) and (9) of the proof of Proposition 9.2 we get

$$c_1(\pi_*(N^{\otimes m})) = \frac{2m^2 + m}{14} \Delta_2 + \frac{5m^2 - m}{14} (\Delta_3 + \Delta_4)$$

as required. \square

Proposition 13.2. *On B we have the exact sequence*

$$0 \rightarrow \mathrm{Sym}^{m-2}(\mathbb{E}) \otimes O(-A) \rightarrow \mathrm{Sym}^m(\mathbb{E}) \rightarrow \pi_*(N^{\otimes m}) \rightarrow 0,$$

with $A = 4\lambda - (\Delta_2 + \Delta_3 + \Delta_4)$.

Proof. By Lemma 12.3 we have on $\mathbb{P}(\mathbb{E})$ the exact sequence

$$0 \rightarrow O(m-2) \otimes u^*O(-A) \rightarrow O(m) \rightarrow O(m)|_Q \rightarrow 0$$

Applying u_* and observing that $R^1u_*O(m-2)$ vanishes gives the result. \square

A section of $\mathrm{Sym}^j(\mathbb{E}) \otimes \det(\mathbb{E})^k$ over \mathcal{H}_3 pulls back to the stack $[W_{8,-2}^0/(\mathrm{GL}(W)/(\pm 1_W))]$ as a section of $\mathrm{Sym}^j(\mathrm{Sym}^2(W)) \otimes \det(W)^{k/2}$ for even k . We have an isotypical decomposition

$$\mathrm{Sym}^j(\mathrm{Sym}^2(W)) = \bigoplus_{n=0}^{\lfloor j/2 \rfloor} \mathrm{Sym}^{2j-4n}(W) \otimes \det(W)^{2n},$$

where we assume here and in the rest of this section that the characteristic is 0 or not 2 and high enough for this identity to hold (or use divided powers as in [1, 3.1]). A section of $\mathrm{Sym}^j(\mathbb{E}) \otimes \det(\mathbb{E})^k$ over \mathcal{M}_3^{nh} pulls back to $[V_{4,0,-1}/\mathrm{GL}(V)]$, where we now write V for the standard space of dimension 3. An identification $V \cong \mathrm{Sym}^2(W)$ corresponds to an embedding $\mathbb{P}^1 \hookrightarrow \mathbb{P}^2$ with image a smooth quadric. If we view V with basis x, y, z , the kernel of the projection

$$\mathrm{Sym}^j(V) = \mathrm{Sym}^j(\mathrm{Sym}^2(W)) \rightarrow \mathrm{Sym}^{2j}(W)$$

consists of the polynomials of degree j in x, y, z that vanish on the quadric. Thus in view of the isotypical decomposition above the exact sequence

$$0 \rightarrow \mathrm{Sym}^{m-2}(\mathbb{E}) \otimes O(-A) \rightarrow \mathrm{Sym}^m(\mathbb{E}) \rightarrow \pi_*(N^{\otimes m}) \rightarrow 0.$$

corresponds to (the pullback to $W_{8,-2}^0$ of) an exact sequence

$$0 \rightarrow (\mathrm{Sym}^{m-2}(\mathrm{Sym}^2W)) \otimes \det(W)^2 \rightarrow \mathrm{Sym}^m(\mathrm{Sym}^2W) \rightarrow \mathrm{Sym}^{2m}(W) \rightarrow 0.$$

The section $\chi_{4,0,8}$ of $\text{Sym}^4(\mathbb{E}) \otimes \det(\mathbb{E})^8$ restricted to the hyperelliptic locus allows three projections according to the decomposition

$$\text{Sym}^4(\text{Sym}^2 W) \otimes \det(W)^{24} = W_{8,24} \oplus W_{4,26} \oplus W_{0,28}. \quad (13)$$

Lemma 13.3. *The projections to the three summands in (13) of the pull back of $\chi_{4,0,8}$ to $\mathcal{H}_{3,2}$ define modular forms on $\overline{\mathcal{H}}_{3,2}$ of weights $(4, 0, 8)$, $(2, 0, 4)$ and $(0, 0, 14)$ and these are given by the covariants $f_{8,-2} \mathfrak{d}$, $f_{4,-1} \mathfrak{d}$ and the discriminant \mathfrak{d} .*

Proof. The identification of \mathbb{E} with $\text{Sym}^2(W)$ corresponds to the embedding of \mathbb{P}^1 as a conic C in \mathbb{P}^2 . A ternary quartic Q contains C either 0, 1 or 2 times, say $Q = mC + R$ with $0 \leq m \leq 2$. The three projections correspond to $R \cap C$ and give the universal binary octic, the universal binary quartic and 1 up to twisting. The first projection was identified in Proposition 5.1. The argument for the second is similar, while the third descends to $\overline{\mathcal{H}}_3$ and does not vanish on \mathcal{H}_3 . Therefore it must be a multiple of the discriminant. Taking into account the action of $\text{GL}_2/\pm 1_W$ we get the indicated weights (namely $2(14 + \epsilon)$ with $\epsilon = -2, -1, 0$). \square

14. MORE MODULAR FORMS FOR GENUS THREE

We will use more effective divisors on projectivized Hodge bundles to produce more modular forms. Here pull back to the Hurwitz space can be useful. We start by calculating the coefficient of δ_1 in the cycle relation for the canonical curve of genus 3 given in Proposition 3.1. Let \overline{D} be the closure in $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_3$ of the canonical image D of the universal curve of genus 3.

Lemma 14.1. *We have $[\overline{D}] = [O(4)] + u^*(8\lambda - \delta_0 - 2\delta_1)$.*

Proof. We will prove this by pulling back the divisor \overline{D} to the Hurwitz space $\mathcal{H}_{3,2}$ and take its closure there in $\mathbb{P}(\mathbb{E})$ over the space of admissible covers $\overline{\mathcal{H}}_{3,2}$. Since \overline{D} is defined as the closure of the canonical image D of the universal curve over \mathcal{M}_3 , it does not contain the whole fibre of $\mathbb{P}(\mathbb{E})$ over a boundary component of $\overline{\mathcal{M}}_3$. But in order to be sure that the pull back of \overline{D} on $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{H}}_{3,2}$ coincides with that of the closure of the pull back of D over $\mathcal{H}_{3,2}$, we have to check that \overline{D} does not contain the whole fibre of $\mathbb{P}(\mathbb{E})$ over boundary components of $\overline{\mathcal{H}}_{3,2}$.

For boundary components of type Δ_0 this follows from the fact that in a 1-dimensional family $X \rightarrow B$ with central fibre C of type Δ_0 the canonical image in the fibre of $\mathbb{P}(\mathbb{E})$ is well-defined and depends only on C . For a 1-dimensional family with central fibre of type Δ_1 we blow up the node of the central fibre, do a quadratic base change and obtain a family $\tilde{X} \rightarrow \tilde{B}$ with central fibre of type $C = C_1 \cup R \cup C_2$ with R a (-2) -curve. We show in the appendix Section 17 that also in this case the image of C in the central fibre of $\mathbb{P}(\mathbb{E})$ is well-defined and depends only on C . In fact, the \mathbb{P}^2 that is the projectivized central fibre of \mathbb{E} , contains a line L and a point P corresponding to the cotangent spaces of $\text{Jac}(C_i)$ in that of $\text{Jac}(C)$. The image of the central fibre is $2L + 2L'$ with L' the line connecting P with the image of a Weierstrass point on L . Thus we may pull back to $\overline{\mathcal{H}}_{3,2}$ and do the calculation of the class there.

We have $[\overline{D}] = O(4) + u^*(a\lambda + b\delta_0 + c\delta_1)$. The pull back of the class to the space of admissible covers $\overline{\mathcal{H}}_{3,2}$ is $O(4) + a\lambda + 2b(\Delta_2 + \Delta_4) + c\Delta_3$. Using the class of Q given in Lemma 12.3 we see that this equals $O(4) + 8\lambda - 2(\Delta_2 + \Delta_3 + \Delta_4)$. By the relation $14\lambda = 3\Delta_2 + 4\Delta_3 + 4\Delta_4$ this gives

$$(3a + 28b + 4)\Delta_2 + (4a + 14c - 4)\Delta_3 + (4a + 28b - 4)\Delta_4 = 0$$

and by the linear independence of the boundary classes this implies $a = 8$, $b = -1$ and $c = -2$. \square

Remark 14.2. Since the case $g = 2$ is hyperelliptic the method just used in the proof of Lemma 14.1 can be applied as well to determine in an alternative way the class of the closure \overline{D} of the ramification divisor D of the universal genus 2 curve in Proposition 2.1.

By the theory of admissible covers there is a natural map $\overline{\mathcal{H}}_{2,2} \rightarrow \overline{\mathcal{M}}_2$ with the property that the pull back of the Hodge bundle on $\overline{\mathcal{M}}_2$ is the Hodge bundle on $\overline{\mathcal{H}}_{2,2}$ associated to the corresponding family of admissible covers. Hence the pull back of the $O(1)$ of the bundle $\mathbb{P}(\mathbb{E}) \rightarrow \overline{\mathcal{M}}_2$ equals the $O(1)$ of the bundle $\mathbb{P}(\mathbb{E}) \rightarrow \overline{\mathcal{H}}_{2,2}$. Let $\Sigma = \varphi_*(\sum_{k=1}^6 \tilde{S}_k)$, with $\varphi : \mathbb{P} \rightarrow \mathbb{P}(\mathbb{E})$ the map defined in Section 10. By geometry, the pull back of \overline{D} to the bundle $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{H}}_{2,2}$ equals Σ . By Remark 10.4 we have $\varphi^*\Sigma = \tilde{S} + 2\Pi_2 + 6\Pi_3$. By using the formulas of Section 11 11, we have for $g = 2$:

$$\varphi^*O(6) = \tilde{M} - 6R = \tilde{S} + \frac{12}{5}\Pi_2 + \frac{2}{5}\Pi_2^c + \frac{24}{5}\Pi_3 - \frac{3}{5}R.$$

We now write $[\overline{D}] = O(6) + u^*(a\delta_0 + b\delta_1)$. By pulling back to $\tilde{\mathbb{P}}$ and using the above formulas, we get (we refer to the diagram in Section 10 for notation)

$$\tilde{S} + \frac{12}{5}\Pi_2 + \frac{2}{5}\Pi_2^c + \frac{24}{5}\Pi_3 - \frac{3}{5}R + \pi^*(2a\Delta_0 + b\Delta_3) = \tilde{S} + 2\Pi_2 + 6\Pi_3.$$

This implies

$$\pi^*(2a\Delta_0 + b\Delta_3) = -\frac{2}{5}(\Pi_2 + \Pi_2^c) + \frac{3}{5}(2\Pi_3 + R) = \pi^*(-\frac{2}{5}\Delta_2 + \frac{3}{5}\Delta_3),$$

hence $a = -1/5$ and $b = 3/5$ and the result follows by using the formula $10\lambda = \delta_1 + 2\delta_2$.

The connection between divisors on projectivized Hodge bundles and modular forms can also be used in the other direction: obtaining results on cycle classes using modular forms. We give a few examples. To a canonical quartic plane curve C we can associate a curve \check{S} in the dual plane of lines intersecting C equianharmonically. It corresponds to a contravariant (concomitant) σ of the ternary quartic given by Salmon in [22, p. 264] and it is defined by an equivariant $\mathrm{GL}(3)$ embedding $W[4, 4, 0] \hookrightarrow \mathrm{Sym}^2(\mathrm{Sym}^4(W))$. It gives rise to a divisor in $\mathbb{P}(\mathbb{E}^\vee)$ and a modular form $\chi_{0,4,16}$ of weight $(0, 4, 16)$. We refer to [5, p. 54] for the relation between invariant theory of ternary quartics and modular forms. The Siegel modular form $\chi_{0,4,16}$ vanishes with order 2 at infinity and order 4 along the locus $\mathcal{A}_{2,1}$ of decomposable abelian threefolds. With $\check{u} : \mathbb{P}(\mathbb{E}^\vee) \rightarrow \overline{\mathcal{M}}_3$ the projection we have $\check{u}_*(O_{\mathbb{P}(\mathbb{E}^\vee)}(1)) = \mathbb{E}^\vee \cong \wedge^2 \mathbb{E} \otimes \det(\mathbb{E})^{-1}$ and we thus find an effective divisor on

$\mathbb{P}(\mathbb{E}^\vee)$ over $\tilde{\mathcal{A}}_3$ with class $[\check{S}] = [O_{\mathbb{P}(\mathbb{E}^\vee)}(4)] + 20\lambda - 2\delta$ and it vanishes with multiplicity 4 along $\mathcal{A}_{2,1}$. We thus find on $\mathbb{P}(\mathbb{E}^\vee)$ over $\overline{\mathcal{M}}_3$ a relation

$$[\check{S}] = [O_{\mathbb{P}(\mathbb{E}^\vee)}(4)] + 20\lambda - 2\delta_0 - 4\delta_1,$$

where we identify λ and δ_i with their pullbacks to $\mathbb{P}(\mathbb{E}^\vee)$. Similarly, in the dual plane we have the sextic \check{T} of lines intersecting the quartic curve in a quadruple of points with j -invariant 1728. The corresponding concomitant τ corresponds to $W[6, 6, 0] \hookrightarrow \text{Sym}^3(\text{Sym}^4(W))$ and defines a modular form of weight $(0, 6, 24)$ vanishing with multiplicity 3 at infinity and multiplicity 6 along $\mathcal{A}_{2,1}$. We thus get a cycle relation

$$[\check{T}] = [O_{\mathbb{P}(\mathbb{E}^\vee)}(6)] + 30\lambda - 3\delta_0 - 6\delta_1.$$

The concomitant $\sigma^3 - 27\tau^2$ vanishes on the locus of double conics and the corresponding modular form of weight $(0, 12, 48)$ is divisible by χ_{18}^2 as can be checked using the methods of [5]. Dividing by χ_{18}^2 gives a cusp form of weight $(0, 12, 12)$ vanishing with multiplicity 2 at infinity and multiplicity 3 along $\mathcal{A}_{2,1}$. It is classically known (see e.g. [3, p. 43]) that this concomitant defines the dual curve \check{C} to the canonical image C in $\mathbb{P}(\mathbb{E})$. We thus find an effective divisor in $\mathbb{P}(\mathbb{E}^\vee)$ containing the closure of the dual curve with class

$$12[O_{\mathbb{P}(\mathbb{E}^\vee)}(1)] + 24\lambda - 2\delta_0 - 3\delta_1.$$

This effective divisor class can also be given by the cycle

$$B = \{(C, \eta) \in \mathbb{P}(\mathbb{E}^\vee) : \text{div}(\eta) \text{ has a point of multiplicity } 2\}$$

over \mathcal{M}_3 and Korotkin and Zograf in [19] determined the class of its closure \overline{B}

$$[\overline{B}] = 12[O_{\mathbb{P}(\mathbb{E}^\vee)}(1)] + 24\lambda - 2\delta_0 - 3\delta_1.$$

Another example of an effective divisor for genus 3 is provided by the Weierstrass divisor W with class

$$[\overline{W}] = 24[O_{\mathbb{P}(\mathbb{E}^\vee)}(1)] + 68\lambda - 6\delta_0 - 12\delta_1$$

as given by Gheorghita in [11]. Here we get a section of

$$\text{Sym}^{24}(\wedge^2 \mathbb{E}) \otimes \det(\mathbb{E})^{44}(-6\delta_0 - 12\delta_1)$$

This gives a Teichmüller modular form of weight $(0, 24, 44)$ vanishing with multiplicity 6 at the cusp. It descends to a Siegel modular form.

Corollary 14.3. *The dual of the canonical curve defines a Siegel modular cusp form of degree 3 of weight $(0, 12, 12)$ vanishing with multiplicity 2 at infinity. The Weierstrass divisor defines a cusp form of weight $(0, 24, 44)$ vanishing with multiplicity 6 at infinity.*

15. THE HYPERTANGENT DIVISOR

A generic canonically embedded curve C of genus 3 has 24 (Weierstrass) points where the tangent line intersects C with multiplicity 3. The union of these 24 lines forms a divisor in \mathbb{P}^2 . Taking the closure of this divisor for the universal curve over \mathcal{M}_3 defines a divisor H in $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{M}}_3$ which we call the hypertangent line divisor. We calculate

the class of this divisor over $\overline{\mathcal{M}}_3$ and also calculate the class of a corresponding divisor over $\overline{\mathcal{H}}_{3,2}$.

The calculation over $\overline{\mathcal{M}}_3$ uses the divisors \check{S} and \check{T} in $\mathbb{P}(\mathbb{E}^\vee)$ over $\overline{\mathcal{M}}_3$ as defined in the preceding section. It is a classical result that the intersection $\check{S} \cdot \check{T}$ in the generic fibre is the 0-cycle consisting of the 24 points defining the 24 hyperflexes of the generic curve C , see [22]. We consider the incidence variety

$$I = \{(p, \ell) \in \mathbb{P}(\mathbb{E}) \times_{\overline{\mathcal{M}}_3} \mathbb{P}(\mathbb{E}^\vee) : p \in \ell\}$$

Let $\pi : I \rightarrow \mathbb{P}(\mathbb{E})$ and $\check{\pi} : I \rightarrow \mathbb{P}(\mathbb{E}^\vee)$ be the two projections fitting in the commutative diagram

$$\begin{array}{ccc} I & \xrightarrow{\check{\pi}} & \mathbb{P}(\mathbb{E}^\vee) \\ \downarrow \pi & & \downarrow \check{u} \\ \mathbb{P}(\mathbb{E}) & \xrightarrow{u} & \overline{\mathcal{M}}_3 \end{array}$$

We have the tautological sequence on $\mathbb{P}(\mathbb{E})$

$$0 \rightarrow F \rightarrow u^*(\mathbb{E}) \rightarrow \mathcal{O}_{\mathbb{P}(\mathbb{E})}(1) \rightarrow 0$$

and a similar one on $\mathbb{P}(\mathbb{E}^\vee)$

$$0 \rightarrow \check{F} \rightarrow \check{u}^*(\mathbb{E}^\vee) \rightarrow \mathcal{O}_{\mathbb{P}(\mathbb{E}^\vee)}(1) \rightarrow 0.$$

Now note that I can be identified with the \mathbb{P}^1 -bundle $\mathbb{P}(F^\vee)$ on $\mathbb{P}(\mathbb{E})$, but also with the \mathbb{P}^1 -bundle $\mathbb{P}(\check{F}^\vee)$ on $\mathbb{P}(\mathbb{E}^\vee)$.

The tautological inclusion $F \rightarrow u^*\mathbb{E}$ induces a surjection $u^*\mathbb{E}^\vee \rightarrow F^\vee$ and this gives an inclusion $\mathbb{P}(F^\vee) \rightarrow \mathbb{P}(u^*\mathbb{E}^\vee)$ of projective bundles over $\mathbb{P}(\mathbb{E})$ which composed with natural map $\mathbb{P}(u^*\mathbb{E}^\vee) \rightarrow \mathbb{P}(\mathbb{E}^\vee)$ gives the map $\check{\pi} : I = \mathbb{P}(F^\vee) \rightarrow \mathbb{P}(\mathbb{E}^\vee)$. This implies

$$\mathcal{O}_{\mathbb{P}(F^\vee)}(1) = \check{\pi}^*\mathcal{O}_{\mathbb{P}(\mathbb{E}^\vee)}(1) \quad \text{and similarly} \quad \mathcal{O}_{\mathbb{P}(\check{F}^\vee)}(1) = \pi^*\mathcal{O}_{\mathbb{P}(\mathbb{E})}(1). \quad (14)$$

With

$$f = c_1(F), \quad \check{f} = c_1(\check{F}), \quad h = c_1(\mathcal{O}_{\mathbb{P}(\mathbb{E})}(1)), \quad \check{h} = c_1(\mathcal{O}_{\mathbb{P}(\mathbb{E}^\vee)}(1)),$$

this gives the identities of pullbacks of the first Chern classes $c_1(\mathbb{E}) = -c_1(\mathbb{E}^\vee) = \lambda$

$$\pi^*(f) + \pi^*(h) = \pi^*u^*(\lambda) = \check{\pi}^*\check{u}^*(\lambda) = -\check{\pi}^*(\check{f}) - \check{\pi}^*(\check{h}),$$

Since $I = \mathbb{P}(F^\vee)$ over $\mathbb{P}(\mathbb{E})$ and $\check{\pi}^*\check{h} = c_1(\mathcal{O}_{\mathbb{P}(F^\vee)}(1))$, the Chern classes of F^\vee and the first Chern class of the tautological line bundle satisfy the relation

$$\check{\pi}^*\check{h}^2 + \pi^*(f)\check{\pi}^*(\check{h}) + \pi^*(c_2(F)) = 0. \quad (15)$$

Corollary 15.1. *Under the map $\pi_*\check{\pi}^*$ we have*

$$\check{h}^2 \mapsto h - u^*(\lambda), \quad \check{h}\check{u}^*(\xi) \mapsto u^*(\xi), \quad \check{u}^*(\eta) \mapsto 0$$

for $\xi \in CH^1(\overline{\mathcal{M}}_3)$ and $\eta \in CH^2(\overline{\mathcal{M}}_3)$.

Proof. Using relation (15) gives

$$\pi_*(\check{\pi}^*(\check{h})^2) = -\pi_*(\pi^*(f)\check{\pi}^*(\check{h}) + \pi^*(c_2(F))) = -f = h - u^*(\lambda).$$

The other properties follow from general intersection theory. \square

Let now ψ be the class of the codimension 2 cocycle $\check{S} \cdot \check{T}$.

Lemma 15.2. *We have $\pi_* \check{\pi}^* \psi = 24h + 216\lambda - 24\delta_0 - 48\delta_1$.*

Proof. By the results of the preceding section we have

$$\psi = 24\check{h}^2 + 240\check{h}\lambda - 24\check{h}\delta_0 - 48\check{h}\delta_1 + r$$

with $r \in \check{\pi}^* CH^2(\overline{\mathcal{M}}_3)$. Corollary 15.1 implies the result. \square

We now claim that the codimension 2 cycle $\check{S} \cdot \check{T}$ when restricted to the hyperelliptic locus is of the form $12\check{h}$, in other words, by (1) it contains an effective codimension 2 cycle with class

$$12(9\lambda - \delta_0 - 3\delta_1)\check{h} + \check{u}^*(\xi)$$

with ξ a codimension 2 class on $\overline{\mathcal{M}}_3$. We check this using the explicit form of the two concomitants σ and τ defining \check{S} and \check{T} . Here σ is a polynomial of degree 4 in a_0, \dots, a_{14} and degree 4 in the coordinates u_0, u_1, u_2 where a_0, \dots, a_{14} are the coefficients of the general ternary quartic. A calculation shows that σ restricted to the locus of double conics becomes a square q^2 with q of degree 2 in the u_i , while τ becomes a cube q^3 . Hence the cycle $S^\vee \cdot T^\vee$ restricted to the hyperelliptic locus is represented by an effective cycle representing $6q \sim 12\check{h}$. By Corollary 15.1 under $\pi_* \hat{\pi}^*$ this is sent to an effective cycle with class $12(9\lambda - \delta_0 - 3\delta_1)$. Since H is defined as the closure of the hypertangent divisor in the generic fibre, the class of H equals $\pi_* \check{\pi}^* \psi$ minus 12 times the class of the hyperelliptic locus; by Lemma 15.2 we get

$$24h + 216\lambda - 24\delta_0 - 48\delta_1 - 12(9\lambda - \delta_0 - 3\delta_1) = 24h + 108\lambda - 12\delta_0 - 12\delta_1.$$

We summarize.

Proposition 15.3. *The class $[H]$ of the hypertangent divisor H in $\mathbb{P}_{\overline{\mathcal{M}}_3}(\mathbb{E})$ equals $[O_{\mathbb{P}(\mathbb{E})}(24)] + 108\lambda - 12\delta_0 - 12\delta_1$. It gives rise to a Siegel modular form of degree 3 and weight $(24, 0, 108)$ vanishing with multiplicity 12 along the boundary.*

We now work on the Hurwitz space and define and calculate the class of a hypertangent H_h divisor there. It is defined by taking the eight tangent lines at the ramification points of the canonical image. More precisely, on $\tilde{\mathbb{P}}$ we have the line bundle N defined in (10). Recall that \tilde{S}_k for $1 \leq k \leq 8$ is the pullback of the section S_k of $\pi_9 : \overline{\mathcal{M}}_{0,9} \rightarrow \overline{\mathcal{M}}_{0,8}$. Under restriction to the hyperelliptic locus the Weierstrass points degenerate to the ramification points. We define the corresponding hypertangent divisor H_h in $\mathbb{P}(\mathbb{E})$ over $\overline{\mathcal{H}}_{3,2}$ by taking the tangents to the canonical image of the generic curve at the points of the sections \tilde{S}_k , $k = 1, \dots, 8$ over $\mathcal{H}_{3,2}$ and then taking the closure over $\overline{\mathcal{H}}_{3,2}$.

We now consider the bundle $N(-2\tilde{S}_k)$ on $\tilde{\mathbb{P}}$. This line bundle is trivial on the generic fibre of $\pi : \tilde{\mathbb{P}} \rightarrow B$, so $\pi_*(N(-2\tilde{S}_k))$ is a line bundle on B .

Lemma 15.4. *We have*

$$c_1(R^1 \pi_* N(-2\tilde{S}_k)) = \Delta_2(k^+) + \Delta_3(k^+), \quad \text{and} \quad c_1(\pi_* N(-2\tilde{S}_k)) = -\Delta_3(k^+) - \Delta_3 + E_k.$$

Proof. Recall that $N = O(D_k + \pi^*(E_k) - R)$. The first statement follows by analyzing the restrictions over the boundary components. For the second we apply Grothendieck-Riemann-Roch as in the proof of Proposition 9.2. By (8) and (9) we have

$$\begin{aligned} c_1(\pi_*N(-2\tilde{S}_k)) &= -\Delta_2(k^+) - 2\Delta_3(k^+) - \Delta_3 + c_1(R^1\pi_*N(-2\tilde{S}_k)) \\ &= -\Delta_3(k^+) - \Delta_3 + E_k. \end{aligned}$$

□

Put $\mathcal{F}_k = \pi_*(N(-2\tilde{S}_k))$. The injection $N(-2\tilde{S}_k) \hookrightarrow N$ induces an injection $\mathcal{F}_k \rightarrow \mathbb{E}$. Pulling back to $\mathbb{P}(\mathbb{E})$ via u^* and composing with the canonical surjection $u^*(\mathbb{E}) \rightarrow O_{\mathbb{P}(\mathbb{E})}(1)$ we get an induced map

$$q : u^*\mathcal{F}_k \rightarrow O_{\mathbb{P}(\mathbb{E})}(1).$$

The degeneracy locus of q is an effective divisor F_k that is the vanishing divisor of a section of $O_{\mathbb{P}(\mathbb{E})}(1) \otimes u^*\mathcal{F}_k^{-1}$. The interpretation is as follows. The map φ defines an embedding of the generic fibre of $\tilde{\mathbb{P}}$ into the generic fibre $\mathbb{P}(\mathbb{E})$. If we identify $H^0(\mathbb{P}^1, O(2))$ with the fibre of \mathbb{E} and projectivize, the divisor $p_1 + p_2 \in |O(2)|$ is mapped to the line through the points $\varphi(p_1), \varphi(p_2)$. We now sum these divisors F_k and get an effective divisor H_h with class

$$\begin{aligned} [H_h] &= 8[O(1)] - \sum_{k=1}^8 [u^*\mathcal{F}_k] = [O(8)] + u^*(3\Delta_3 + 8\Delta_3 - \sum_{k=1}^8 E_k) \\ &= [O(8)] + u^*(8\lambda - 2\Delta_2 + \Delta_3), \end{aligned}$$

where we use the formulas of Section 9 and Section 11.

We can now compare the class of the hyperelliptic hypertangent divisor H_h with that of the pull back of the hypertangent divisor H to the Hurwitz space. By Proposition 15.3 the pull back of H has class $[O(24)] + 108\lambda - 24(\Delta_2 + \Delta_4) - 12\Delta_1$. Since the 24 Weierstrass points collapse with multiplicity 3 to the 8 ramification points we compare the class (of the pull back of) $[H]$ with that of $3[H_h]$. Substituting the formula for λ we get

$$[H] - 3[H_h] = 9\Delta_1,$$

which means that the pull back of H vanishes with multiplicity 9 at the hyperelliptic boundary component Δ_1 .

16. GENUS FOUR

For a smooth curve C of genus 4 the natural map $\text{Sym}^2(H^0(C, \omega_C)) \rightarrow H^0(C, \omega_C^{\otimes 2})$ is surjective and the kernel has dimension 1. It determines a quadric in $\mathbb{P}^3 = \mathbb{P}(H^0(C, \omega_C))$ containing the canonical curve. Over $\overline{\mathcal{M}}_4$ we find a corresponding exact sequence

$$0 \rightarrow U \rightarrow \text{Sym}^2(\mathbb{E}) \rightarrow \pi_*\omega_C^{\otimes 2} \rightarrow 0.$$

The line bundle U has first Chern class $5\lambda - (13\lambda - \delta) = -8\lambda + \delta$ by Mumford's calculation of $c_1(\pi_*\omega_C^{\otimes 2})$ [20, Thm. 5.10]. In the bundle $\mathbb{P}(\mathbb{E})$ the quadric containing the canonical curve determines a divisor Q . Let $u : \mathbb{P}(\mathbb{E}) \rightarrow \overline{\mathcal{M}}_4$ be the projection.

Lemma 16.1. *The divisor class of Q satisfies: $[Q] = [O(2)] + u^*(8\lambda - \delta)$.*

Proof. Observe that $u^*u_*O(2) = u^*(\text{Sym}^2(\mathbb{E}))$. The natural morphism $u^*u_*O(2) \rightarrow O(2)$ induces $u^*U \rightarrow O(2)$. The divisor Q is the vanishing locus of this morphism, hence has class $[O(2)] + u^*(8\lambda - \delta)$. \square

Corollary 16.2. *The effective divisor Q defines a Teichmüller modular cusp form χ of genus 4 and weight $(2, 0, 0, 8)$.*

If we view a section of $\text{Sym}^2(\mathbb{E})$ as a quadratic form on \mathbb{E}^\vee we can take the discriminant, cf. [6]. Doing this with the form χ of weight $(2, 0, 0, 8)$ just constructed we get a scalar-valued modular form $D(\chi)$ of weight 34. This modular form vanishes on the closure of the locus of curves whose canonical model lies on a quadric cone. This locus has class $34\lambda - 4\delta_0 - 14\delta_1 - 18\delta_2$ by Teixidor i Bigas [23, Prop. 3.1] and equals the divisor of curves with a vanishing theta null. The modular form $D(\chi)$ is the square root of the restriction to \mathcal{M}_4 of the product of the even theta characteristics on \mathcal{A}_4 .

17. APPENDIX ON BASE-POINT FREENESS

We consider a family $\pi : X \rightarrow B$ of semistable curves of genus g with B a disc or the spectrum of a discrete valuation ring. We assume that the central fibre C is a nodal curve which is a chain of three curves $C = C_1 \cup R \cup C_2$ with C_1, C_2 smooth curves of genus g_1 and g_2 and R a rational (-2) -curve. We let $q = C_1 \cap R$ and $p = C_2 \cap R$.

Proposition 17.1. *Let ω the relative dualizing sheaf of $\pi : X \rightarrow B$. Then $\pi_*(\omega(-R)) \cong \pi_*(\omega)$ and the central fibre of $\pi_*(\omega(-R))$ is of codimension 1 in $H^0(C, \omega(-R))$ and defines a base point free linear system on C .*

Proof. The exact sequence $0 \rightarrow \omega(-R) \rightarrow \omega \rightarrow \omega|_R \rightarrow 0$ induces a sequence

$$0 \rightarrow \pi_*(\omega(-R)) \rightarrow \pi_*(\omega) \xrightarrow{r} \pi_*(\omega)|_R$$

and the map r is zero because $\omega|_C = (\omega_{C_1}(q), O_R, \omega_{C_2}(p))$, therefore the restrictions to C_1 (resp. C_2) must vanish at q (resp. p), hence extend by 0 on R . We thus see by the exactness that $\pi_*(\omega(-R)) \cong \pi_*(\omega)$.

Next we observe that $\dim H^0(C, \omega(-R)) = g_1 + g_2 + 1$ with g_i the genus of C_i . This follows directly from $\omega(-R)|_C = (\omega_{C_1}, O_R(2), \omega_{C_2})$.

We have the exact sequence

$$0 \rightarrow \omega(-R - C_1) \rightarrow \omega(-R) \rightarrow \omega(-R)|_{C_1} \rightarrow 0, \quad (22)$$

where $\omega(-R)|_{C_1} \cong \omega_{C_1}$ and $\omega(-R - C_1)|_C = (\omega_{C_1}(q), O_R(1), \omega_{C_2})$. For a section $(s_1, s, s_2) \in H^0(C, \omega(-R - C_1))$ the section s is the unique section of O_R that vanishes at q and with $s(p) = s_2(p)$. We thus see $\dim H^0(C, \omega(-R - C_1)) = g_1 + g_2$. Therefore $h^0(\omega(-R - C_1))$ has constant rank $g_1 + g_2$ on the fibres of π , hence $R^1\pi_*(\omega(-R - C_1))$ is a line bundle. We conclude that the special fibre of $\pi_*(\omega(-R - C_1))$ equals $H^0(C, \omega(-R - C_1))$. But $\pi_*(\omega(-R)|_{C_1})$ is a torsion sheaf, hence the connecting homomorphism $\pi_*(\omega(-R)|_{C_1}) \rightarrow R^1\pi_*(\omega(-R - C_1))$ of (22) must be zero and we get an induced exact sequence

$$0 \rightarrow \pi_*(\omega(-R - C_1)) \xrightarrow{l} \pi_*(\omega(-R)) \xrightarrow{j} \pi_*(\omega(-R)|_{C_1}) \rightarrow 0.$$

Consider now a section σ of $\pi_*(\omega(-R - C_1))$ with restriction (s_1, s, s_2) to C . Suppose that $s \neq 0$. If we multiply σ with a local section τ of O_{C_1} on X with divisor C_1 then $\iota(\sigma) = \sigma \cdot \tau|_C$ has as restriction to R a section of $O_R(2)$ vanishing with multiplicity 2 at q and therefore it does not vanish anywhere else. Hence the subspace of the special fibre V of $\pi_*(\omega(-R))$ of sections vanishing on C_1 has q as only base point on R . Furthermore, the map j is surjective, and choosing a section $s_1 \in H^0(C_1, \omega_{C_1})$ with $s_1(q) \neq 0$ we see that q is not a base point. Therefore there are no base points on R . By the surjectivity of j the restriction of V to C_1 is $H^0(C_1, \omega_{C_1})$ and therefore there are no base points on C_1 . By symmetry the same holds for C_2 . \square

Similarly to (22) we have exact sequence

$$0 \rightarrow \omega(-R - C_1 - C_2) \rightarrow \omega(-R) \rightarrow \omega(-R)|_{C_1+C_2} \rightarrow 0$$

and by a similar reasoning we see that we get an exact sequence

$$0 \rightarrow \pi_*(\omega(-R - C_1 - C_2)) \xrightarrow{\iota} \pi_*(\omega(-R)) \xrightarrow{j} \pi_*(\omega(-R))|_{C_1+C_2} \rightarrow 0.$$

This implies that given $s_1 \in H^0(C_1, \omega_{C_1})$ and $s_2 \in H^0(C_2, \omega_{C_2})$ there is a unique element (s_1, s, s_2) in the special fibre V of $\pi_*(\omega(-R))$ mapping to (s_1, s_2) under j . The morphism $X \rightarrow \mathbb{P}(\pi_*(\omega(-R)))$ is given by the surjection $\pi^*\pi_*(\omega(-R)) \rightarrow \omega(-R)$. The image of the curve C in the special fibre of $\mathbb{P}(\mathbb{E})$ consists of the canonical images of C_1 and C_2 , provided with with image of p and q and the image of R , that is, the line connecting the images of p and q . If the genus $g(C_i) = 1$ then the image of C_i is a point.

REFERENCES

- [1] M. Aprodu, G. Farkas, S. Papadima, C. Raicu, J. Weyman: *Koszul modules and Green's conjecture*. *Inventiones Math.* **218** (2019), 657–720.
- [2] A. Arsie, A. Vistoli: *Stacks of cyclic covers of projective spaces*. *Compositio Math.* **140** (2004) 647–666.
- [3] A. Clebsch: *Ueber symbolische Darstellung algebraischer Formen*. *Journal für die Reine und Angewandte Mathematik* **59**, 1–62 (1861).
- [4] F. Cléry, C. Faber, and G. van der Geer: *Covariants of binary sextics and vector-valued Siegel modular forms of genus 2*. *Mathematische Annalen* **369**(3–4), 1649–1669 (2017).
- [5] F. Cléry, C. Faber, and G. van der Geer: *Concomitants of ternary quartics and vector-valued Siegel modular and Teichmueller modular forms of genus three*. *Selecta Mathematica* (2020) 26:55.
- [6] F. Cléry and G. van der Geer: *Modular forms via invariant theory*. [arXiv: 2211.05611](https://arxiv.org/abs/2211.05611)
- [7] M. Cornalba: *The Picard group of the moduli stack of stable hyperelliptic curves*. *Rend. Lincei Mat. Appl.* **18** (2007), 109–115.
- [8] D. Eisenbud, J. Harris: *3264 and all that: Intersection theory in algebraic geometry*. Cambridge, UK. Cambridge University Press, 2016.
- [9] G. van der Geer, A. Kouvidakis: *The Hodge bundle on Hurwitz Spaces*. *Pure Appl. Math. Quarterly* **7** (2011), No. 4, Special Issue: In memory of Eckart Viehweg, 1297–1307.
- [10] G. van der Geer, A. Kouvidakis: *The cycle classes of divisorial Maroni loci*. *International Mathematics Research Notices*, Vol. 2016, No. 00, pp. 1–47 doi: 10.1093/imrn/rnw133
- [11] I. Gheorghita: *Effective divisors in the projectivized Hodge bundle*. *International Mathematics Research Notices*, Vol. 2021, No. 1, 303–325
- [12] J. Harris: *Curves and their moduli*. In: *Algebraic Geometry Bowdoin 1985*. *Proc. Symp. Pure Math.* **48** (1987), 99–143.

- [13] J. Harris, I. Morrison: *Moduli of Curves*. Graduate Texts in Mathematics **187**, 1998, Springer Verlag.
- [14] J. Harris, D. Mumford: *On the Kodaira dimension of the moduli space of curves*. *Inventiones Mathematicae* **67**, (1982), 23–86.
- [15] J.-I. Igusa: *Arithmetic moduli for genus two*. *Annals of Math.*, **72**, (1960), pp. 612–649.
- [16] J.-I. Igusa: *On Siegel modular forms of genus two*. *Amer. J. of Math.*, **84**, (1962), pp. 175–200.
- [17] J.-I. Igusa: *Modular forms and projective invariants*. *Amer. J. Math.* **89** (1967), 817–855.
- [18] S. Keel: *Intersection theory of moduli space of stable N -pointed curves of genus zero*. *Transactions A.M.S.* **330**, 1992, 545–574.
- [19] D. Korotkin, P. Zograf. *Tau function and moduli of differentials*. *Math. Research Letters* **18**, no. 3 (2011): 447–58.
- [20] D. Mumford: *Stability of projective varieties*. *L'Enseignement Mathématique XXIII* (1977), 39–110.
- [21] D. Mumford: *Towards an enumerative geometry of the moduli space of curves*. *Arithmetic and Geometry*. Papers dedicated to I. R. Shafarevich, Vol. II: *Geometry*. *Progress in Math.* 36, pp. 271–328.
- [22] G. Salmon: *A treatise on the higher plane curves*. Dublin, Hodges, Foster and Figgis, 1879.
- [23] M. Teixidor i Bigas: *The divisor of curves with a vanishing theta-null*. *Compositio Math.* **66**, (1988), 15–22.
- [24] S. Tsuyumine: *On Siegel modular forms of degree three*. *American Journal of Mathematics* **108**, (1986), pp. 755–862.