

Quasicontral modulus and self-similar sets: a supplementary result to Voiculescu's work

Kozo Ikeda and Masaki Izumi
Graduate School of Science
Kyoto University
Sakyo-ku, Kyoto 606-8502, Japan

March 14, 2023

Abstract

In his recent work, Voiculescu generalized his remarkable formula for the quasicontral modulus of a commuting n -tuple of hermitian operators with respect to the $(n, 1)$ -Lorentz ideal to the case where its spectrum is contained in a Cantor-like self-similar set in a certain class. In this note, we treat general self-similar sets satisfying the open set condition, and obtain lower and upper bounds of the quasicontral modulus. Our proof shows that Voiculescu's formula holds for a class of self-similar sets including the Sierpinski gasket and the Sierpinski carpet.

1 Introduction

Pursuing his idea in the proof of a non-commutative Weyl-von Neumann theorem for C^* -algebras [8], D. Voiculescu introduced in [9] the quasicontral modulus $k_\Phi(\tau)$ for an n -tuple τ of Hilbert space operators with respect to a symmetric normed ideal \mathfrak{S}_Φ . With this quantity, he established a far reaching generalization of the classical Weyl-von Neumann theorem in perturbation theory: an n -tuple of commuting hermitian operators τ is a $\mathfrak{S}_\Phi^{(0)}$ -perturbation of a commuting tuple of diagonal hermitian operators if and only if $k_\Phi(\tau)$ vanishes. Moreover he showed that $k_n(\tau) := k_{\mathcal{C}_n}(\tau)$ always vanishes for the Schatten-von Neumann ideal \mathcal{C}_n (see also [1], [2], [10], [11], [12] for later development).

Besides the significance of whether it vanishes or not, the quantity $k_n^-(\tau) := k_{\mathcal{C}_n^-}(\tau)$ itself is of particular interest for the $(n, 1)$ -Lorentz ideal \mathcal{C}_n^- because Voiculescu obtained the following remarkable formula:

Theorem 1.1 ([9, Theorem 4.5]). *Let τ be an n -tuple of commuting hermitian operators, and let \mathbf{m} be the multiplicity function of the Lebesgue absolutely continuous part of τ . Then we have*

$$k_n^-(\tau)^n = \gamma_n \int_{\mathbb{R}^n} \mathbf{m}(s) d\lambda(s),$$

where $0 < \gamma_n < \infty$ is a constant independent of τ and λ is the Lebesgue measure.

In his recent work [13], Voiculescu generalized the above result to the case where the joint spectrum $\sigma(\tau)$ is a subset of a Cantor-like self-similar set K in a certain class and the Lebesgue measure is replaced by the Hausdorff measure (see Theorem 3.1 below). The purpose of this note is to treat a larger class of self-similar sets than Voiculescu did. Although we cannot show the precise equality as above, we obtain upper and lower bounds establishing that $k_p^-(\tau)^p$ behaves like the Hausdorff measure \mathcal{H}_p with p the Hausdorff dimension of K for general self-similar sets satisfying the open set condition (Theorem 3.2). Our proof shows that Voiculescu's formula holds for a class of self-similar sets including the Sierpinski gasket and the Sierpinski carpet (Remark 1).

The authors would like to thank Dan Voiculescu for useful discussion, and Kenneth Falconer and Jun Kigami for having informed the authors of the reference [6].

2 Preliminaries

Our basic references for normed ideals are [4] and [7], while we adopt the notation in [9] and [13].

Throughout the paper H denotes a separable Hilbert space. The symbols $\mathcal{L}(H)$, $\mathcal{K}(H)$, $\mathcal{P}(H)$, $\mathcal{F}(H)$, $\mathcal{R}(H)_1^+$ will respectively denote the bounded operators, the compact operators, the finite rank orthogonal projections, the finite rank operators, and the finite rank positive contractions on H . $\mathcal{P}(H)$ and $\mathcal{R}_1^+(H)$ are both directed sets by natural order of self-adjoint operators.

For $T \in \mathcal{K}(H)$, we denote by $\{\mu_j(T)\}_{j=1}^\infty$ the singular value sequence of T , that is, the set of eigenvalues of $|T|$ in decreasing order. For a symmetric function Φ , we denote by $|T|_\Phi$ the norm of the operator ideal corresponding to Φ . For $1 \leq p < \infty$, let

$$|T|_p = \left(\sum_{j=1}^{\infty} \mu_j(T)^p \right)^{1/p},$$

and for $1 < p \leq \infty$, let

$$|T|_p^- = \sum_{j=1}^{\infty} \frac{\mu_j(T)}{j^{1-\frac{1}{p}}}.$$

The Schatten-von Neumann ideal \mathcal{C}_p is the set of compact operators T with finite $|T|_p$, and the $(p, 1)$ -Lorentz ideal \mathcal{C}_p^- is the set of compact operators T with finite $|T|_p^-$. Then we have

$$\bigcup_{r < p} \mathcal{C}_r \subset \mathcal{C}_p^- \subset \mathcal{C}_p.$$

Let $\tau^{(j)} = (T_1^{(j)}, T_2^{(j)}, \dots, T_n^{(j)}) \in (\mathcal{L}(H))^n$, $\sigma = (S_1, S_2, \dots, S_n) \in (\mathcal{L}(H))^m$, $X, Y \in \mathcal{L}(H)$, and $Z \in \mathcal{L}(K)$. We will write

$$\tau^{(1)} \oplus \tau^{(2)} \equiv (T_1^{(1)} \oplus T_1^{(2)}, T_2^{(1)} \oplus T_2^{(2)}, \dots, T_n^{(1)} \oplus T_n^{(2)}) \in \mathcal{L}(H \oplus H)^n$$

$$\tau^{(1)} + \tau^{(2)} \equiv (T_1^{(1)} + T_1^{(2)}, T_2^{(1)} + T_2^{(2)}, \dots, T_n^{(1)} + T_n^{(2)}) \in \mathcal{L}(H)^n$$

$$X\tau Y = (XT_1Y, XT_2Y, \dots, XT_nY) \in \mathcal{L}(H)^n$$

$$\tau \otimes Z = (T_1 \otimes Z, T_2 \otimes Z, \dots, T_n \otimes Z) \in \mathcal{L}(H \otimes K)^n$$

$$[X, \tau] = ([X, T_1], [X, T_2], \dots, [X, T_n]) \in \mathcal{L}(H)^n$$

$$\|\tau\| = \max_{1 \leq j \leq n} \|T_j\|$$

$$|\tau|_\Phi = \max_{1 \leq j \leq n} |T_j|_\Phi, \quad |\tau|_p = \max_{1 \leq j \leq n} |T_j|_p, \quad |\tau|_p^- = \max_{1 \leq j \leq n} |T_j|_p^-.$$

Definition 2.1 ([9, Section 1]). For $\tau \in \mathcal{L}(H)^n$, we define

$$k_\Phi(\tau) = \liminf_{A \in \mathcal{R}_1^+(H)} |[A, \tau]|_\Phi.$$

When $|T|_\Phi = |T|_p$, we simply denote it by $k_p(\tau)$, and when $|T|_\Phi = |T|_p^-$, we denote it by $k_p^-(\tau)$.

We collect basic facts frequently used in this paper (see [9, Section 1]).

Theorem 2.1. *Let the notation be as above. The following hold:*

- (1) *If $\{A_i\}$ is a sequence in $\mathcal{R}_1^+(H)$ converging to I in the weak operator topology, then*

$$k_\Phi(\tau) \leq \liminf_{i \rightarrow \infty} |[A_i, \tau]|_\Phi.$$

- (2) *If $k_\Phi(\tau)$ is finite, there exists an increasing sequence $\{A_i\}$ in $\mathcal{R}_1^+(H)$ converging to I satisfying*

$$\lim_{i \rightarrow \infty} |[A_i, \tau]|_\Phi = k_\Phi(\tau).$$

- (3) *Let $\tau^{(j)} \in \mathcal{L}(H)^n$ for $j \in \mathbb{N}$. We have*

$$\max_{j=1,2} k_\Phi(\tau^{(j)}) \leq k_\Phi(\tau^{(1)} \oplus \tau^{(2)}) \leq k_\Phi(\tau^{(1)}) + k_\Phi(\tau^{(2)})$$

and

$$k_\Phi \left(\bigoplus_{j=1}^{\infty} \tau^{(j)} \right) = \lim_{m \rightarrow \infty} k_\Phi \left(\bigoplus_{j=1}^m \tau^{(j)} \right).$$

The following properties can be shown by the averaging trick (see [9, Proof of Proposition 1.6]),

Theorem 2.2. *Let the notation be as above. Then the following hold:*

- (1) *For $m \in \mathbb{N}$, there exists an increasing sequence $\{A_i\}$ in $\mathcal{R}_1^+(H)$ converging to I and satisfying*

$$\lim_{i \rightarrow \infty} |[A_i \otimes I_m, \tau \otimes I_m]|_{\Phi} = k_{\Phi}(\tau \otimes I_m),$$

where I_m is the identity operator of \mathbb{C}^m .

- (2) *Let $\tau^{(j)} \in \mathcal{L}(H)^n$ for $j = 1, 2, \dots, m$. Then there exists an increasing sequence $\{A_i\}_{i=1}^{\infty}$ in $\mathcal{R}_1^+(H)^{\oplus m} \subset \mathcal{R}_1^+(H^{\oplus m})$ converging to I and satisfying*

$$\lim_{i \rightarrow \infty} |[A_i, \tau^{(1)} \oplus \tau^{(2)} \oplus \dots \oplus \tau^{(m)}]|_{\Phi} = k_{\Phi}(\tau^{(1)} \oplus \tau^{(2)} \oplus \dots \oplus \tau^{(m)}).$$

- (3) *Let $\tau^{(j)} \in \mathcal{L}(H)^n$ and $\lambda^{(j)} \in (\mathbb{C}I)^n$ for $j = 1, 2, \dots, m$. Then we have*

$$k_{\Phi}(\tau^{(1)} \oplus \dots \oplus \tau^{(m)}) = k_{\Phi}((\tau^{(1)} - \lambda^{(1)}) \oplus \dots \oplus (\tau^{(m)} - \lambda^{(m)})).$$

3 Main result

Our basic references for self-similar sets are [3] and [5]. Throughout this section, we treat \mathbb{R}^n as a metric space equipped with the Euclidean metric. For $X \subset \mathbb{R}^n$, we denote by $\dim_H X$ the Hausdorff dimension of X . We denote by $O(n)$ the real orthogonal group.

We say that a map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a contraction if there exists a constant $0 < \lambda < 1$ such that for all $x, y \in \mathbb{R}^n$ the inequality

$$|F(x) - F(y)| \leq \lambda|x - y|$$

holds. If moreover equality holds for all $x, y \in \mathbb{R}^n$, we say that F is a *similitude* and λ is the *contraction ratio* of F .

Let F_1, F_2, \dots, F_m be contractions of \mathbb{R}^n . Then it is known that there exists a unique compact set $K \subset \mathbb{R}^n$ satisfying

$$K = \bigcup_{j=1}^m F_j(K),$$

which we call the self-similar set for $\{F_j\}_{j=1}^m$ ([3, Theorem 8.3], [5, Theorem 1.1.4]).

We say that the open set condition holds for $\{F_j\}_{j=1}^m$ if there exists a bounded non-empty open set $V \subset \mathbb{R}^n$ such that

$$\prod_{j=1}^m F_j(V) \subset V$$

with this union disjoint.

Assume that similitudes $\{F_j\}_{j=1}^m$ satisfy the open set condition, and let $0 < \lambda_j < 1$ be the contraction ratio of F_j for $j = 1, 2, \dots, m$. It follows from [3, Theorem 8.6],[5, Corollary 1.5.9] that the Hausdorff dimension of K is the number $p > 0$ determined by

$$\sum_{j=1}^m \lambda_j^p = 1,$$

and the Hausdorff measure $\mathcal{H}_p(K)$ of K is a non-zero finite number. Let $\mathcal{W}_l = \{1, 2, \dots, m\}^l$, and let $\mathcal{W} = \bigcup_{l=0}^{\infty} \mathcal{W}_l$. For $w \in \mathcal{W}$, we denote its word length by $|w|$. For $w \in \mathcal{W}_l$, we define $F_w = F_{w_1} \circ F_{w_2} \circ \dots \circ F_{w_l}$, $K_w = F_w(K)$, and

$$\lambda_w = \lambda_{w_1} \lambda_{w_2} \dots \lambda_{w_l}.$$

Then we have

$$\mathcal{H}_p(K_w) = \lambda_w^p \mathcal{H}_p(K).$$

Now we consider the following 4 conditions for self-similar sets.

(A1) Let F_1, F_2, \dots, F_m be similitudes in \mathbb{R}^n , and let K be the corresponding self-similar set. For $j = 1, 2, \dots, m$, we can express F_j as

$$F_j(x) = \lambda_j U_j(x) + b_j,$$

where $0 < \lambda_j < 1$, U_j is an orthogonal matrix in $O(n)$, and $b_j \in \mathbb{R}^n$. We assume that the similitudes F_1, F_2, \dots, F_m satisfy the open set condition, and the Hausdorff dimension p of K is strictly larger than 1.

(A2) In addition to (A1), we assume $\lambda_1 = \lambda_2 = \dots = \lambda_m$.

(A3) In addition to (A2), we assume $U_1 = U_2 = \dots = U_m = I_m$.

(A4) In addition to (A3), we assume $F_i(K) \cap F_j(K) = \emptyset$ for any $i \neq j$.

Voiculescu showed the following theorem in his recent work:

Theorem 3.1 ([13, Theorem 5.1]). *Let τ be an n -tuple of commuting Hermitian operators whose spectrum $\sigma(\tau)$ is a subset of a self-similar set K satisfying (A4). Then there exists a constant $0 < \gamma_K < \infty$, depending only on K , satisfying*

$$k_p^-(\tau)^p = \gamma_K \int_K \mathbf{m}(x) d\mathcal{H}_p(x),$$

where \mathbf{m} is the multiplicity function of the \mathcal{H}_p -absolutely continuous part of τ .

Remark 1. As we can see from our arguments below, the same statement holds even if the assumption (A4) is relaxed to (A3). For example, the statement is true for the Sierpinski gasket and the Sierpinski carpet.

The following is our main theorem, where we assume only (A1), but our conclusion requires two constants.

Theorem 3.2. *Let $\tau \in \mathcal{L}(H)^n$ be an n -tuple of commuting hermitian operators whose spectrum $\sigma(\tau)$ is a subset of a self-similar set K satisfying (A1). Then there exist two positive constants C_1, C_2 determined by only F_1, F_2, \dots, F_m such that*

$$C_1 \int_K \mathbf{m} d\mathcal{H}_p \leq k_p^-(\tau)^p \leq C_2 \int_K \mathbf{m} d\mathcal{H}_p,$$

where \mathbf{m} is the multiplicity function of the \mathcal{H}_p -absolutely continuous part of τ

We prove Theorem 3.2 in several steps following Voiculescu's strategy in [13] with necessary modifications.

In the sequel, we assume that K and τ satisfy the assumption of Theorem 3.2. We denote by $E(\tau; \cdot)$ the spectral measure of τ . We denote by $\text{diam}(K_w)$ the diameter of K_w . We set $\lambda_* = \min\{\lambda_j\}_{j=1}^m$. We say that two Borel subsets A and B of K are *virtually disjoint* if $\dim_H A \cap B < p$. For virtually disjoint A and B , we have $\mathcal{H}_p(A \cap B) = 0$.

The first step immediately follows from [2, Corollary 4.7].

Lemma 3.3. *We denote by τ_K the n -tuple of multiplication operators of coordinate functions in \mathbb{R}^n acting on $L^2(K, \mathcal{H}_p)$. Then we have $k_p^-(\tau_K) > 0$.*

Next we prove [13, Lemma 5.1] under the assumption of (A1).

Lemma 3.4. *Assume that τ has a cyclic vector ξ . Then for some constant $C > 0$ depending only on K ,*

$$k_p^-(\tau) \leq C \mathcal{H}_p(\sigma(\tau))^{1/p}.$$

Proof. We enumerate the family of sets $F_j(K) \cap F_k(K)$ for $j \neq k$, and denote them by L_1, L_2, \dots, L_N (some of them can be empty). We further enumerate the family of the sets $F_w(L_i)$ for $|w| \geq 1$ and $1 \leq i \leq N$, and denote them by L_{N+1}, L_{N+2}, \dots . Note that we have $\dim_H L_i < p$ thanks to [6, Theorem 3.3].

We set $H_1 = E(\tau; L_1)H$, and inductively define

$$H_i = E(\tau; L_i)H \cap \left(\bigcup_{j=1}^{i-1} H_j \right)^\perp$$

for $i \geq 2$, and

$$H_0 = \left(\bigcup_{i=1}^{\infty} H_i \right)^\perp.$$

Then τ is decomposed as

$$(\tau, H) = \bigoplus_{i=0}^{\infty} (\tau_i, H_i).$$

We claim $k_p^-(\tau_i) = 0$ for all $i \geq 1$. Indeed, note that we have $\sigma(\tau_i) \subset L_i$ and $\dim_H L_i < p$. We choose $r_i > 1$ satisfying $\dim_H L_i < r_i < p$. Then $\mathcal{H}_{r_i}(L_i) = 0$. Since τ_i is decomposed into its cyclic components, we get $k_{r_i}(\tau) = 0$ thanks to [14, Theorem 6.1] and Theorem 2.1,(3). Since $\mathcal{C}_{r_j} \subset C_p^-$, the claim follows from [9, Corollary 2.6].

By Theorem 2.1,(3), the claim implies $k_p^-(\tau) = k_p^-(\tau_0)$. Therefore to show Lemma 3.4, we may and do assume $\tau = \tau_0$ as we have $\mathcal{H}_p(\sigma(\tau_0)) \leq \mathcal{H}_p(\sigma(\tau))$.

For $r \in \mathbb{N}$, let $\Omega(r)$ be the set of all $w = (w_1, w_2, \dots, w_l) \in \mathcal{W}$ satisfying

$$\lambda_w \leq \frac{1}{r} < \lambda_{(w_1, w_2, \dots, w_{l-1})}$$

and $K_w \cap \sigma(\tau) \neq \emptyset$, and let

$$G(r) = \bigcup_{w \in \Omega(r)} K_w.$$

Then we have $\sigma(\tau) \subset G(r)$, and

$$\frac{\lambda_*}{r} \text{diam}(K) \leq \text{diam}(K_w) < \frac{1}{r} \text{diam}(K),$$

for $w \in \Omega(r)$ by definition. Note that K_{w_1} and K_{w_2} are virtually disjoint for all $w_1 \neq w_2 \in \Omega(r)$, and $E(\tau; K_{w_1}) \perp E(\tau; K_{w_2})$. For a given $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ so that we have

$$n_0 \leq r \Rightarrow \mathcal{H}_p(G(r)) \leq \mathcal{H}_p(\sigma(\tau)) + \varepsilon$$

because $\sigma(\tau) = \bigcap_r G(r)$. We denote $E_w = E(\tau; K_w)$. Then we have

$$\sum_{w \in \Omega(r)} E_w = I.$$

We also have

$$|\Omega(r)| \leq \frac{r^p}{\lambda_*^p \mathcal{H}_p(K)} (\mathcal{H}_p(\sigma(\tau)) + \varepsilon)$$

for $r \geq n_0$ as $\sum_{w \in \Omega(r)} \mathcal{H}_p(K_w) = \mathcal{H}_p(G(r))$ and if $w \in \Omega(r)$ we have

$$\left(\frac{\lambda_*}{r}\right)^p \mathcal{H}_p(K) \leq \mathcal{H}_p(K_w).$$

Let P_r be the orthogonal projection onto the linear span of $\{E_w\xi \mid w \in \Omega(r)\}$. We can express P_r as

$$P_r = \sum_{w \in \Omega(r)} e_w \otimes e_w^*,$$

where $e_w = \frac{1}{\|E_w\xi\|} E_w\xi$ if $E_w\xi \neq 0$ and $e_w = 0$ if $E_w\xi = 0$. Since E_w commutes with τ and P_r , we have

$$\begin{aligned} \|[P_r, \tau]\| &= \max_{w \in \Omega(r)} \|[P_r, \tau] E_w\| \\ &= \max_{1 \leq i \leq n} \max_{w \in G(r)} \|e_w \otimes (T_i e_w)^* - (T_i e_w) \otimes e_w^*\|. \end{aligned}$$

Since $\text{diam}(K_w) \leq \text{diam}(K)/r$, we get $\|[P_r, \tau]\| \leq 2 \text{diam}(K)/r$.

Since $\{P_r\}_{r=1}^\infty$ is an increasing sequence of projections, it strongly converges to a projection, say P . Since $P_r\xi = \xi$ for $r \in \mathbb{N}$ and $\lim_{r \rightarrow \infty} \|[P_r, \tau]\| = 0$, we have $P\xi = \xi$ and $[P, \tau] = 0$. Thus we get $P = I$ as ξ is cyclic for τ .

Since $\text{rank}(P_r) \leq |\Omega(r)|$,

$$\begin{aligned} |[P_r, \tau]^-|_p &\leq \sum_{k=1}^{2|\Omega(r)|} k^{-1+1/p} \|[P_r, \tau]\| \leq \sum_{k=1}^{2|\Omega(r)|} k^{-1+1/p} \frac{2 \text{diam}(K)}{r} \\ &\leq p(2|\Omega(r)|)^{1/p} \frac{2 \text{diam}(K)}{r} \leq C(\mathcal{H}_p(\sigma(\tau)) + \varepsilon)^{1/p}, \end{aligned}$$

where $C = p \cdot 2^{1+1/p} \lambda_*^{-1} \cdot \mathcal{H}_p(K)^{-1/p}$. Using Theorem 2.1,(1), we get the statement. \square

We can prove [13, Lemma 5.2] under the assumption (A1) now.

Lemma 3.5. *Assume that $\sigma(\tau) \subset K$ and the spectral measure of τ is singular with respect to the Hausdorff measure \mathcal{H}_p . Then $k_p^-(\tau) = 0$.*

Proof. As in the proof of the previous lemma, we may assume that $\tau = \tau_0$. Then we can prove the statement exactly in the same way as in the proof of [13, Lemma 5.2] by using Lemma 3.4. \square

The main technical improvement in [13] is the ampliation homogeneity [13, Theorem 3.1] showing $k_p^-(\tau \otimes I_m) = m^{\frac{1}{p}} k_p^-(\tau)$. To deal with the contraction rates λ_j depending on j , we modify it as follows.

Lemma 3.6. *Let Φ be a symmetric function, let $0 \leq c_1 \leq a_1, a_2, \dots, a_m \leq c_2$, and let $\tau \in \mathcal{L}(H)^n$. Then we have*

$$c_1 k_\Phi(\tau \otimes I_m) \leq k_\Phi(\tau \otimes \text{diag}(a_i)_{i=1}^m) \leq c_2 k_\Phi(\tau \otimes I_m).$$

In particular,

$$c_1 m^{\frac{1}{p}} k_p^-(\tau) \leq k_p^-(\tau \otimes \text{diag}(a_i)_{i=1}^m) \leq c_2 m^{\frac{1}{p}} k_p^-(\tau).$$

Proof. For $X_1, X_2, \dots, X_m \in \mathcal{L}(H)$ and $a_1, a_2, \dots, a_m, b_1, b_2, \dots, b_m \in \mathbb{R}$ satisfying $0 \leq a_j \leq b_j$, we denote the singular values of X_j by $\{s_i^{(j)}\}_{i=1}^\infty$. If we define $A = \bigoplus_{j=1}^m a_j X_j$ and $B = \bigoplus_{j=1}^m b_j X_j$, then the singular values of A and B are respectively $\{a_j s_i^{(j)}\}_{i,j}$ and $\{b_j s_i^{(j)}\}_{i,j}$. So for all $n \in \mathbb{N}$ we have

$$\sum_{k=1}^n \mu(A)_k = \sum_{k=1}^n a_{j_k} s_{i_k}^{(j_k)} \leq \sum_{k=1}^n b_{j_k} s_{i_k}^{(j_k)} \leq \sum_{k=1}^n \mu(B)_k.$$

Then it follows from [7, Theorem 1.16 (b)] that $|A|_\Phi \leq |B|_\Phi$.

Thanks to Theorem 2.2,(2), there exists an increasing sequence $\{A_i\}_{i=1}^\infty$ in $\mathcal{R}_1^+(H)^{\oplus m} \subset \mathcal{R}_1^+(H^{\oplus m})$ such that $\{A_i\}$ strongly converges to I and

$$\lim_{i \rightarrow \infty} |[\tau \otimes \text{diag}(a_j), A_i]|_\Phi = k_\Phi(\tau \otimes \text{diag}(a_j)).$$

Expressing A_i as

$$A_i = A_1^{(i)} \oplus A_2^{(i)} \oplus \dots \oplus A_m^{(i)} \in \mathcal{R}_1^+(H)^{\oplus m},$$

we get

$$\begin{aligned} c_1 |[\tau \otimes I_m, A_i]|_\Phi &= \left| \left[\bigoplus_{j=1}^m c_1 \tau, \bigoplus_{j=1}^m A_j^{(i)} \right] \right|_\Phi = \left| \bigoplus_{j=1}^m c_1 [\tau, A_j^{(i)}] \right|_\Phi \\ &\leq \left| \bigoplus_{j=1}^m a_j [\tau, A_j^{(i)}] \right|_\Phi = |[\tau \otimes \text{diag}(a_j), A_i]|_\Phi. \end{aligned}$$

Then by Theorem 2.1,(1), we get

$$c_1 k_\Phi(\tau \otimes I_m) \leq k_\Phi(\tau \otimes \text{diag}(a_j)).$$

By Theorem 2.2,(1), we can take an increasing sequence $\{B_i\} \in \mathcal{R}_1^+(H)$ strongly converges to I and satisfying

$$\lim_{i \rightarrow \infty} |[\tau \otimes I_m, B_i \otimes I_m]|_\Phi = k_\Phi(\tau \otimes I_m).$$

Then we have

$$\begin{aligned} |[\tau \otimes \text{diag}(a_j), B_i \otimes I_m]|_\Phi &= \left| \bigoplus_{j=1}^m a_j [\tau, B_i] \right|_\Phi \\ &\leq \left| \bigoplus_{j=1}^m c_2 [\tau, B_i] \right|_\Phi = c_2 |[\tau \otimes I_m, B_i \otimes I_m]|_\Phi. \end{aligned}$$

Using Theorem 2.1,(1), we obtain

$$k_\Phi(\tau \otimes \text{diag}(a_j)) \leq c_2 k_\Phi(\tau \otimes I_m). \quad \square$$

To deal with the rotation part in F_j , we use a modified quasicentral modulus, as suggested in [13, Remark 6.2].

Definition 3.1 ([9, Theorem 1.7]). For a symmetric function Φ and $\tau \in \mathcal{L}(H)^n$, we define $\widetilde{k}_\Phi(\tau)$ by

$$\widetilde{k}_\Phi(\tau) = \liminf_{A \in \mathcal{R}_1^+} |[A, \tau]|_{\widetilde{\Phi}},$$

where $|\tau|_{\widetilde{\Phi}} = \left| \left(\sum_{i=1}^n T_i^* T_i \right)^{\frac{1}{2}} \right|_{\Phi}$.

Let e_1, e_2, \dots, e_n be the orthonormal basis of \mathbb{C}^n and define $d_\tau : H \rightarrow H \otimes \mathbb{C}^n$ by

$$d_\tau(\xi) = \sum_{i=1}^n T_i \xi \otimes e_i$$

Then we have $d_\tau^* d_\tau = \sum_{i=1}^n T_i^* T_i$ and $|\tau|_{\widetilde{\Phi}} = |d_\tau|_{\Phi}$. Since

$$|\tau|_{\Phi} \leq |d_\tau|_{\Phi} \leq n|\tau|_{\Phi},$$

we have

$$k_\Phi(\tau) \leq \widetilde{k}_\Phi(\tau) \leq nk_\Phi(\tau).$$

\widetilde{k}_Φ and k_Φ share similar properties such as Theorem 2.1, Theorem 2.2, Lemma 3.5, and Lemma 3.6 etc. However, \widetilde{k}_Φ is distinguished from k_Φ in that \widetilde{k}_Φ has the following property.

For $U = (u_{i,j})_{i,j} \in O(n)$ and $\tau = (T_i)_i \in \mathcal{L}(H)^n$, we denote

$$U\tau = \left(\sum_{i=1}^n u_{1,i} T_i, \sum_{i=1}^n u_{2,i} T_i, \dots, \sum_{i=1}^n u_{n,i} T_i \right) \in \mathcal{L}(H)^n.$$

Lemma 3.7. *Let Φ be a symmetric function, let $\tau^{(1)}, \tau^{(2)}, \dots, \tau^{(m)} \in \mathcal{L}(H)^n$, and let $U_1, U_2, \dots, U_m \in O(n)$. Then we have*

$$\widetilde{k}_\Phi \left(\bigoplus_{j=1}^m \tau^{(j)} \right) = \widetilde{k}_\Phi \left(\bigoplus_{j=1}^m U_j \tau^{(j)} \right).$$

Proof. By symmetry, it suffices to show

$$\widetilde{k}_\Phi \left(\bigoplus_{j=1}^m U_j \tau^{(j)} \right) \leq \widetilde{k}_\Phi \left(\bigoplus_{j=1}^m \tau^{(j)} \right).$$

From Theorem 2.2,(2), there exists an increasing sequence $\{A_i\}$ in the set $\mathcal{R}_1^+(H)^{(m)}$ converging to I and

$$\lim_{i \rightarrow \infty} \left\| \left[A_i, \bigoplus_{j=1}^m \tau^{(j)} \right] \right\|_{\tilde{\Phi}} = \widetilde{k}_{\Phi} \left(\bigoplus_{j=1}^m \tau^{(j)} \right).$$

Denoting

$$A_j = A_j^{(1)} \oplus A_j^{(2)} \oplus \cdots \oplus A_j^{(n)},$$

we have

$$\begin{aligned} \left\| \left[A_i, \bigoplus_{j=1}^m \tau^{(j)} \right] \right\|_{\tilde{\Phi}} &= \left\| \bigoplus_{j=1}^m \left[A_i^{(j)}, \tau^{(j)} \right] \right\|_{\tilde{\Phi}} = \left\| \bigoplus_{j=1}^m U_j \left[A_i^{(j)}, \tau^{(j)} \right] \right\|_{\tilde{\Phi}} \\ &= \left\| \bigoplus_{j=1}^m \left[A_i^{(j)}, U_j \tau^{(j)} \right] \right\|_{\tilde{\Phi}} = \left\| \left[A_i, \bigoplus_{j=1}^m U_j \tau^{(j)} \right] \right\|_{\tilde{\Phi}}. \end{aligned}$$

Using Theorem 2.1,(1), we get

$$\widetilde{k}_{\Phi} \left(\bigoplus_{j=1}^m \tau^{(j)} \right) = \lim_{i \rightarrow \infty} \left\| \left[A_i, \bigoplus_{j=1}^m U_j \tau^{(j)} \right] \right\|_{\tilde{\Phi}} \geq \widetilde{k}_{\Phi} \left(\bigoplus_{j=1}^m U_j \tau_j \right). \quad \square$$

Note that Lemma 3.3 and Lemma 3.4 show $0 < \widetilde{k}_p^-(\tau_K) < \infty$. We set

$$\widetilde{\gamma}_K = \frac{\widetilde{k}_p^-(\tau_K)^p}{\mathcal{H}_p(K)}.$$

Now Theorem 3.2 follows from the following theorem.

Theorem 3.8. *Under the assumption of Theorem 3.2, the following holds:*

$$\lambda_*^p \widetilde{\gamma}_K \int_K \mathbf{m} \, d\mathcal{H}_p \leq \widetilde{k}_p^-(\tau)^p \leq \lambda_*^{-p} \widetilde{\gamma}_K \int_K \mathbf{m} \, d\mathcal{H}_p.$$

Proof. We denote by $\tau_{p,a}$ and $\tau_{p,s}$ respectively the \mathcal{H}_p -absolutely continuous part of τ and the \mathcal{H}_p -singular part of τ . Thanks to Lemma 3.5, we have $\widetilde{k}_p^-(\tau_{p,s}) = 0$. Thus Theorem 2.1,(3) shows that to prove the statement, we may and do assume $\tau = \tau_{p,a}$, that is, the spectral measure of τ is absolutely continuous with respect to \mathcal{H}_p .

Up to unitary equivalence, we may assume

$$\tau = \bigoplus_{k=1,2,\dots,\infty} \tau_{X_k}^{\oplus k},$$

where $\{X_k\}_{k=1}^\infty$ are mutually disjoint Borel subsets of K , and τ_{X_k} is the n -tuple of multiplication operators by coordinate functions in \mathbb{R}^n acting on $L^2(X_k, \mathcal{H}_p)$. We have $\mathbf{m}(x) = k$ if and only if $x \in X_k$.

First, we prove the statement in the case of multiplicity 1, that is, $\tau = \tau_X$ for a Borel subset $X \subset K$. In the same way as in the proof of [13, Theorem 5.1], this case is reduced to the case where X is a finite union of K_{w_j} . Thus we assume

$$X = \bigcup_{j=1}^l K_{w_j} \text{ with}$$

$$F_{w_j}(x) = r_j V_j(x) + a_j,$$

and that $\{K_{w_j}\}_{j=1}^l$ are mutually virtually disjoint. If necessary, we decompose each K_{w_j} and assume that

$$\lambda_* r_1 \leq r_j \leq r_1$$

for $j = 1, 2, \dots, m$. We have a unitary φ from $L^2(F_{w_j}(X), \mathcal{H}_p)$ to $L^2(X, \mathcal{H}_p)$ defined by

$$\varphi(f) = r_j^{\frac{p}{2}} f \circ F_{w_j},$$

and $\tau_{K_{w_j}} = \tau_{F_{w_j}(K)}$ is unitarily equivalent to $F_{w_j}(\tau_K)$, where for

$$F = (F_1, F_2, \dots, F_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

and $\tau = (T_1, T_2, \dots, T_n) \in \mathcal{L}(H)^n$,

$$F(\tau) = (F_1(T_1, T_2, \dots, T_n), F_2(T_1, T_2, \dots, T_n), \dots, F_n(T_1, T_2, \dots, T_n)).$$

Then we can see that

$$\begin{aligned} \widetilde{k}_p^-(\tau_X) &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l \tau_{K_{w_j}} \right) = \widetilde{k}_p^- \left(\bigoplus_{j=1}^l F_{w_j}(\tau_K) \right) \\ &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l (r_j V_j(\tau_K) + a_j) \right) \\ &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l r_j V_j(\tau_K) \right) \text{ (by Theorem 2.2, (3))} \\ &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l r_j \tau_K \right) \text{ (by Theorem 3.7)} \\ &= \widetilde{k}_p^- (\tau_K \otimes \text{diag}(r_j)_{j=1}^l). \end{aligned}$$

By Lemma 3.6, we get,

$$\lambda_* r_1 l^{1/p} \widetilde{k}_p^- (\tau_K) \leq \widetilde{k}_p^- (\tau_K \otimes \text{diag}(r_j)_{j=1}^l) \leq r_1 l^{1/p} \widetilde{k}_p^- (\tau_K),$$

and on the other hand,

$$\lambda_*^p r_1^p l \mathcal{H}_p(K) \leq \mathcal{H}_p(X) \leq r_1^p l \mathcal{H}_p(K).$$

Therefore we obtain the statement.

Next we show the statement in the case of finite multiplicity. In this case, we may assume $\tau = \bigoplus_{j=1}^l \tau_{Y_j}$ with $Y_j \subset K$. We choose a sufficiently large number e and compose F_1 and F_2 e times each to make similitudes $F_{w_1}, F_{w_2}, \dots, F_{w_m}$ so that $\{F_{w_j}(X_j)\}_{j=1}^l$ are virtually disjoint. Now we can express

$$F_{w_j}^{-1}(x) = rW_j(x) + c_j$$

with $r = \lambda_1^{-e} \lambda_2^{-e} > 0$ and some $W_j \in O(n)$, $c_j \in \mathbb{R}^n$. Then we have

$$\begin{aligned} \widetilde{k}_p^-(\tau) &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l \tau_{Y_j} \right) = \widetilde{k}_p^- \left(\bigoplus_{j=1}^l F_{w_j}^{-1} F_{w_j}(\tau_{Y_j}) \right) \\ &= \widetilde{k}_p^- \left(\bigoplus_{j=1}^l (rW_j(F_{w_j}(\tau_{w_j})) + c_j) \right) = r \widetilde{k}_p^- \left(\bigoplus_{j=1}^l F_{w_j}(\tau_{w_j}) \right) \\ &= r \widetilde{k}_p^- \left(\bigoplus_{j=1}^l \tau_{F_{w_j}(Y_j)} \right) = r \widetilde{k}_p^-(\tau_Y), \end{aligned}$$

where $Y = \bigcup_{j=1}^l F_{w_j}(Y_j)$. On the other hand,

$$r^p \mathcal{H}_p(Y) = \sum_{j=1}^l r^p \mathcal{H}_p(F_{w_j}(Y_j)) = \sum_{j=1}^l \mathcal{H}_p(Y_j) = \int_K \mathfrak{m} d\mathcal{H}_p.$$

The general case is reduced to the finite multiplicity case thanks to Theorem 2.1,(3), and we are done. \square

Corollary 3.9. *Let τ and K be as above. Then the following conditions are equivalent:*

- (1) $k_p^-(\tau) = 0$.
- (2) *The spectral measure of τ is singular with respect to \mathcal{H}_p .*

An easy modification of the proof of Theorem 3.8 shows the following:

Corollary 3.10. *In addition to the assumption of Theorem 3.2, we assume (A2). Then the following holds:*

$$\widetilde{k}_p^-(\tau)^p = \widetilde{\gamma}_K \int_K \mathfrak{m} d\mathcal{H}_p.$$

Question 1. Let $\tau \in \mathcal{L}(H)^n$, let $1 < p < \infty$, and let $a = (a_j) \in \mathbb{R}^k$. In view of the proof of Theorem 3.8, we can see that if

$$\widetilde{k}_p^-(\tau \otimes \text{diag}(a_j)_{j=1}^k) = \widetilde{k}_p^-(\tau) \|a\|_p \quad (3.1)$$

is true, the formula in Corollary 3.10 would hold under the condition (A1). As was already asked in [13, Remark 6.1], a more fundamental question is: does the equality

$$\widetilde{k}_p^-(\tau_1 \oplus \tau_2)^p = \widetilde{k}_p^-(\tau_1)^p + \widetilde{k}_p^-(\tau_2)^p \quad (3.2)$$

hold in general?

Thanks to [13, Lemma 3.2], one might hope to prove (3.2) (or (3.1)) by reducing it to the following question. Let $\{S_l\}$ and $\{T_m\}$ be sequences of finite rank operators satisfying

$$\begin{aligned} \lim_{l \rightarrow \infty} \|S_l\| &= \lim_{m \rightarrow \infty} \|T_m\| = 0, \\ \lim_{l \rightarrow \infty} |S_l|_p^- &= \alpha, \quad \lim_{m \rightarrow \infty} |T_m|_p^- = \beta. \end{aligned}$$

Then does the equality

$$\inf_{l_r, m_r} \liminf_{r \rightarrow \infty} |S_{l_r} \oplus T_{m_r}|_p^- = (\alpha^p + \beta^p)^{1/p}, \quad (3.3)$$

where the infimum is taken over all subsequences, hold true? However, there exists an easy counter example to (3.3). Let P_l be a projection of rank l , and let $S_l = p^{-1} 2^{-l/p} P_{2^l}$ and $T_m = 3^{1/p} S_m$. Then it is straightforward to show that (3.3) does not hold for $\{S_l\}$ and $\{T_m\}$, and the left-hand side is strictly larger than the right-hand side. This means that we should use, not only subsequences, but also convex combinations of $\{S_l\}$ and $\{T_m\}$ in the left-hand side of (3.3) to raise a reasonable question.

Question 2. For our purpose, we can relax (3.1) as follows. Let $\{N_n\}_{n=1}^\infty$ be an increasing sequence of natural numbers converging to infinity, and let $a_{n,j}$, $n \in \mathbb{N}$, $1 \leq j \leq N_n$, be positive numbers such that there exists $c > 0$ satisfying

$$\sum_{j=1}^{N_n} |a_{n,j}|^p = 1,$$

$$\max\{a_{n,j}\}_{j=1}^{N_n} \leq c \min\{a_{n,j}\}_{j=1}^{N_n},$$

for all $n \in \mathbb{N}$. Then we can see that if

$$\lim_{n \rightarrow \infty} \widetilde{k}_p^-(\tau \otimes \text{diag}(a_{n,j})_{j=1}^{N_n}) = \widetilde{k}_p^-(\tau) \quad (3.4)$$

is true, the formula in Corollary 3.10 would hold under the condition (A1). It might be interesting to ask what kind of condition on $a_{n,j}$ assures (3.4).

Acknowledgement

The authors would like to thank Dan Voiculescu for useful discussion, and Kenneth Falconer and Jun Kigami for having informed the authors of the reference [6].

Declarations

M. I. is supported in part by JSPS KAKENHI Grant Number JP20H01805. The authors have no relevant financial or non-financial interests to disclose. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- [1] Bercovici, Hari and Voiculescu, Dan. *The analogue of Kuroda's theorem for n -tuples*. The Gohberg anniversary collection, Vol. II (Calgary, AB, 1988), 57–60, Oper. Theory Adv. Appl., 41, Birkhäuser, Basel, 1989.
- [2] David, Guy and Voiculescu, Dan. *s -Numbers of singular integrals for the invariance of absolutely continuous spectra in fractional dimensions*. Journal of Functional Analysis **94** (1990), no. 1, 14–26.
- [3] Falconer, Kenneth. *The geometry of fractal sets*. Cambridge Tracts in Mathematics, 85. Cambridge University Press, Cambridge, 1986.
- [4] Gohberg, Israel and Krein, Mark. *Introduction to the Theory of Linear non-selfadjoint operators*. Translated from the Russian by A. Feinstein Translations of Mathematical Monographs, Vol. 18 American Mathematical Society, Providence, R.I. 1969.
- [5] Kigami, Jun. *Analysis on fractals*. Cambridge Tracts in Mathematics, 143. Cambridge University Press, Cambridge, 2001.
- [6] Morán, Manuel. *Dynamical boundary of a self-similar set*. Fund. Math. **160** (1999), no. 1, 1–14.
- [7] Simon, Barry. *Trace ideals and their applications*. Mathematical Surveys and Monographs, 120. American Mathematical Society, Providence, RI, 2005.
- [8] Voiculescu, Dan. *A non-commutative Weyl-von Neumann theorem*. Rev. Roumaine Math. Pures Appl. **21** (1976), no. 1, 97–113.
- [9] Voiculescu, Dan. *Some results on norm-ideal perturbations of Hilbert space operators*. Journal of Operator Theory **2** (1979), no. 1, 3–37.

- [10] Voiculescu, Dan. *Some results on norm-ideal perturbations of Hilbert space operators. II.* Journal of Operator Theory **5** (1981), no. 1, 77–100.
- [11] Voiculescu, Dan. *Hilbert space operators modulo normed ideals.* Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), 1041–1047, PWN, Warsaw, 1984.
- [12] Voiculescu, Dan. *On the existence of quascentral approximate units relative to normed ideals. Part I.* Journal of Functional Analysis **91** (1990), no. 1, 1–36.
- [13] Voiculescu, Dan. *The formula for the quascentral modulus in the case of spectral measures on fractals.* Journal of Fractal Geometry **8** (2021), no. 4, 347–361.
- [14] Xia, Jingbo. *Diagonalization modulo norm ideals with Lipschitz estimates.* J. Funct. Anal. **145** (1997), no. 2, 491–526.