

Characterising Solutions of Anomalous Cancellation

Satvik Saha^{1*}, Sohom Gupta², Sayan Dutta¹ and Sourin Chatterjee¹

¹Department of Mathematics and Statistics, Indian Institute of Science Education and Research, Kolkata, India.

²Department of Physical Sciences, Indian Institute of Science Education and Research, Kolkata, India.

*Corresponding author(s). E-mail(s): ss19ms154@iiserkol.ac.in;
 Contributing authors: sg19ms141@iiserkol.ac.in;
sd19ms148@iiserkol.ac.in; sc19ms170@iiserkol.ac.in;

Abstract

Anomalous cancellation of fractions is a mathematically inaccurate method where cancelling the common digits of the numerator and denominator correctly reduces it. While it appears to be *accidentally* successful, the property of anomalous cancellation is intricately connected to the number of digits of the denominator as well as the base in which the fraction is represented. Previous work have been mostly surrounding three digit solutions or specific properties of the same. This paper seeks to get general results regarding the structure of numbers that follow the cancellation property (denoted by $P_{\ell; k}^*$) and an estimate of the total number of solutions possible in a given base representation. In particular, interesting properties regarding the *saturation* of the number of solutions in general and p^n bases (where p is a prime) have been studied in detail.

Keywords: Anomalous cancellation, Diophantine equation

1 Introduction

A zeal for some interesting mathematical problems brought us to a very peculiar problem, a quest to find all odd digit integers $[a_1 a_2 \dots a_{2k+1}]$ (all a_i 's are

2 *Characterising Solutions of Anomalous Cancellation*

digits, $a_1 \neq 0$) such that the following property holds.

$$[a_1 a_2 \dots a_k] \cdot [a_{k+1} a_{k+2} \dots a_{2k+1}] = [a_1 a_2 \dots a_{k+1}] \cdot [a_{k+2} a_{k+2} \dots a_{2k+1}]$$

An elementary example is the number 164, which has the property as shown $1 \times 64 = 16 \times 4$. Similarly $24 \times 996 = 249 \times 96$ implies that 24996 also fits our problem requirement. Now the question is, can one find a way to generate all such numbers? A brute force algorithm always works, but it is never satisfying to leave things at that – a proof of being a mathematics aspirant. This prompted us to go through existing literature which, though scarce, hide a gold mine of information. Adding on to previous work, we arrived at several interesting results that demand a place in this paper. The paper has been structured in a format that the general reader can be presented with all the beautiful results, while the seasoned readers can move to the appendix to get a flavour of the relevant proofs. All proofs use elementary number theory techniques and hence this paper is directed at undergraduate students and beyond.

The anomalous cancellation property for fractions of the form $[ab]_B/[bc]_B$, where the numerator and denominator are two digit integers in base B , has been studied extensively [2]. Finding such fractions amounts to solving the following Diophantine equation in a, b, c .

$$(aB + b)c = a(bB + c).$$

We shall quickly state some nice results in literature, to keep readers up to date.

Apart from trivial solutions such as $a = b = c$ or $b = c = 0$, it has been shown that b is the largest digit. With this, the transformation $(a, b, c) \mapsto (b - c, b, b - a)$ is an involution on the set of non-trivial solutions [5]. Ekhad [6] has examined a much more general anomalous cancellation property where a digit is allowed to be cancelled from anywhere in the numerator and denominator. Fixing the base, denominator, and the digit indices used during cancellation reduces the problem of finding all solutions to a linear Diophantine equation, which can be solved easily.

It should be good to go dive in our work now. For no future confusion, we shall take some time to sort out the notations used frequently. All other symbols have their usual meaning.

Notation.

Let $B \geq 2$ be an integer base, and let x_1, x_2, \dots, x_k be integer digits, with each $0 \leq x_i < B$. Then, we denote their concatenation as

$$x = [x_1 x_2 \dots x_k] \equiv \sum_{i=1}^k x_i B^{k-i}.$$

In our problem, we consider natural numbers $N_{\ell; k}$ in base B , where $\ell, k \geq 1$. Its digits are denoted as

$$N_{\ell; k} = [a_1 a_2 \dots a_\ell b c_1 c_2 \dots c_k] = [abc],$$

with blocks

$$a = [a_1 a_2 \dots a_\ell], \quad c = [c_1 c_2 \dots c_k].$$

Note that we may write

$$N_{\ell; k} = aB^{k+1} + bB^k + c.$$

Definition 1 We say that the number $N_{\ell; k}$ has property $P_{\ell; k}$, or is a solution of $P_{\ell; k}$, if

$$[a_1 a_2 \dots a_\ell b] \times [c_1 c_2 \dots c_k] = [a_1 a_2 \dots a_\ell] \times [b c_1 c_2 \dots c_k].$$

Equivalently,

$$\frac{[a_1 a_2 \dots a_\ell \cancel{b}]}{[\cancel{b} c_1 c_2 \dots c_k]} = \frac{[a_1 a_2 \dots a_\ell]}{[c_1 c_2 \dots c_k]}$$

which means that the digit 'b' can be anomalously cancelled in the LHS fraction to *correctly* reduce to the RHS equation. This can also be expressed as the homogeneous Diophantine equations

$$(aB + b)c = a(bB^k + c), \quad \frac{1}{a} + \frac{B-1}{b} = \frac{B^k}{c}.$$

In the special case $\ell = k$, we abbreviate $N_{k; k} \equiv N_k$ and $P_{k; k} \equiv P_k$.

Convention.

Certain types of solutions of $P_{\ell; k}$ exist given any base B . These are of the following forms.

- $[dd \dots dd]$.
- $[a_1 a_2 \dots a_k 00 \dots 0]$ (in particular, any two of $a, b, c = 0$.)

We term these as *trivial solutions*. In our discussion, we ignore such solutions, and focus on the case $\ell = k$, i.e. we are searching for non-trivial solutions to the P_k problem.

Definition 2 We say that $N_{\ell; k}$ has property $P_{\ell; k}^*$ if it has $P_{\ell; k}$ non-trivially, i.e. no two of $a, b, c = 0$ and not all digits $a_i = b = c_i$.

2 Characterizing how solutions look like.

The first important observation is that solutions of $P_{\ell; k}$ can be extended very simply to solutions of $P_{\ell+1; k+1}$, by replacing the central digit b with the digits $[bbb]$. This is a special case of the Pumping Lemma described in [7].

4 Characterising Solutions of Anomalous Cancellation

Proposition 1 *Let B be an arbitrary integer base, and let*

$$N = [a_1 a_2 \dots a_\ell b c_1 c_2 \dots c_k], \quad N^+ = [a_1 a_2 \dots a_\ell b b b c_1 c_2 \dots c_k].$$

Then, N has property $P_{\ell; k}$ if and only if N^+ has property $P_{\ell+1; k+1}$. We say that N^+ is an extension of N .

Proof The number N^+ has property $P_{\ell+1; k+1}$ precisely when

$$(aB^2 + bB + b)(bB^k + c) = (aB + b)(bB^{k+1} + bB^k + c),$$

which after expanding and cancelling reduces to

$$(aB + b)cB = aB(bB^k + c), \quad (aB + b)c = a(bB^k + c),$$

which is precisely the statement that N has property $P_{\ell; k}$. \square

This shows that the number of solutions of P_k^* in a particular base B cannot decrease with increasing k . A natural question is whether the number of such solutions gets arbitrarily large; can we keep producing solutions of the P_k problem that aren't merely extensions of old ones? To answer this, we show that the digits of any such solution must obey very rigid rules, culminating in the following.

Theorem 2 *Let B be an arbitrary integer base and let N_k have property P_k^* . Then, the digits must satisfy the following constraints.*

1. $a_1 < B/2$.
2. $b = c_1 = c_2 = \dots = c_{k-1} > c_k > 1$.
3. $\gcd(c_k, B) > 1$.
4. $\gcd(a_k - b, B) > 1$ if $a_k \neq b$.

The proof of Theorem 2 will follow by combining Corollary 12 and Lemmas 14, 15, 17, which we prove later.

An immediate consequence is that all solutions of P_k^* look like

$$[a_1 a_2 \dots a_k b b \dots b c_k].$$

This means that the blocks are of the form

$$a = \frac{bM}{B} + \frac{bc_k B^{k-1}}{bB - (B-1)c_k}, \quad c = bM + c_k, \quad M = \frac{B^k - B}{B-1}.$$

Theorem 2 guarantees that when the base B is even, plugging in $b = B-1$, $c_k = B-2$ corresponds to the largest solution of P_k^* ,

$$a = \frac{1}{2}B^k - 1, \quad b = B - 1, \quad c = B^k - 2.$$

For example, in base $B = 10$, the largest solution of P_3^* is 4999998.

Theorem 2 also gives us a nice criterion for determining whether solutions of P_k^* exist in the first place.

Theorem 3 *The P_k^* problem admits solutions if and only if the base B is composite.*

Proof If the base $B = mn$ for $m, n > 1$, then $N_k = [abc]$ with

$$(a, b, c) = (mB^{k-1} - 1, B - 1, B^k - n)$$

has property P_k^* . Conversely, if the base $B = p$ for prime p , then any solution N_k of P_k^* must satisfy $0 < c_k < p$ and $\gcd(c_k, p) > 1$ simultaneously by Proposition 2, a contradiction. \square

3 Saturation of solutions.

In Section 2, we have shown that every candidate solution of P_k^* can be completely written in terms of the two single digits b and c_k . Here, we say that (b, c_k) *generates* such a solution of P_k^* . Figure 1 visualizes the solution space of P_{101}^* in base $B = 126$ via such generating tuples. Thus, finding solutions amounts to testing pairs (b, c_k) , with $1 < c_k < b < B$. Importantly, this search space depends only on B , not k (!) Thus, for a given base B , the number of solutions of P_k^* *saturates*.

Proposition 4 *Let B be an arbitrary integer base. There are at most*

$$\frac{(B-2)(B-3)}{2}$$

solutions of P_k^ , for any $k \geq 1$.*

Proof For each b , we have $2 \leq c_k < b$, i.e. $b-2$ choices. Putting $3 \leq b < B$, we obtain a total of $1 + 2 + \dots + (B-3) = (B-2)(B-3)/2$ candidate tuples (b, c_k) . \square

The above estimate is clearly very crude: we have only used the relation $b = c_{i < k} > c_k$. Indeed, numerical evidence suggests that a much sharper bound can be established. To illustrate this, the only solutions of P_k^* for base $B = 10$ are extensions of the following (in order of appearance with increasing k).

$$164, 265, 195, 498, 21775, 24996, 1249992, 340277776.$$

We next describe when the full gamut of solutions for a base B is achieved.

Theorem 5 *Let B be an arbitrary integer base. Then, the number of solutions of P_k^* become constant beyond*

$$k = \max\{5, 2 \log_2(B-1) + 2\}.$$

6 Characterising Solutions of Anomalous Cancellation




Fig. 1 Generating tuples for $P_{\ell; k}^*$ with $B = 126$, $k = 101$, represented by blue dots. Points above the orange curve generate solutions of P_k^* with the candidate block $a \geq B^{k-1}$. Points underneath the orange curve generate solutions of $P_{\ell; k}^*$ with $\ell < k$.

This means that for k beyond the above *saturation point*, all solutions of P_k^* are mere extensions of old ones. A proof of Theorem 5 is supplied in Appendix D.

The bound on k described in Theorem 5 is especially bad for one family of bases in particular. When $B = p^n$ for prime p , $n \geq 2$, the solutions of P_k^* saturate in the very first step, $k = 1$.

Theorem 6 *Let $B = p^n$ for prime p . Then, all solutions of P_k^* are extensions of those of P_1^* .*

Proof By Lemma 18, any solution N_k of P_k^* looks like

$$N_k = [a_1 b \dots b b b \dots b c_k].$$

Proposition 1 can be used $k - 1$ times to reduce this to the solution $[a_1 b c_k]$ of P_1^* . \square

For example, the only solutions of P_k^* for base $B = 3^2$ are of the form

$$14\dots 43, \quad 28\dots 86.$$

4 Discussion.

Our main goal was to characterize all solutions of the anomalous cancellation problem P_k^* , which still seems to be a far way off. However, we have successfully characterized solutions in certain special bases, and given rudimentary bounds on their number. Our major findings are summarized below.

- There are no solutions of P_k^* for prime bases.
- The only solutions of P_k^* for prime-power bases p^n are extensions of solutions of P_1^* .
- There are at least as many solutions of P_k^* for composite bases B as there are non-trivial factors of B . These are of the form $[abc]$ with

$$(a, b, c) = (mB^{k-1} - 1, B - 1, B^k - n)$$

for $B = mn$, $m, n > 1$, $M = (B^k - B)/(B - 1)$.

Some very interesting and peculiar observations were made from numerical solutions; while these haven't been proved in this paper, there is a lot of scope for future expansion.

- Suppose in base B , we have no new non-trivial solutions (except extensions) in $(2k + 1)$ digits. We have observed that there would be no new solutions in $(2k + 3)$ digits. In other words, as long as the number of solutions do not saturate, they will keep on increasing.
- Currently we have a quadratic bound on the total number of solutions possible in a given base B , but observed solution counts are much smaller than that.
- An interesting result discussed is that composite bases are guaranteed to have solutions that are of the form $(a, B - 1, c)$. Therefore, finding out no solutions of this form guarantees primality. While the current search space is only as good as a brute force method, some better ideas relating b and c_k might lead to a much faster primality test.

Acknowledgments. The authors would like to thank Prof. Soumya Bhattacharya for his careful reading of the manuscript and many helpful discussions on the topic of this paper.

Appendix A Trivial solutions.

This section focuses on trivial solutions of $P_{\ell; k}$. We supply a few criteria which can be used to swiftly identify candidate solutions as trivial, based on a subset of their digits.

8 Characterising Solutions of Anomalous Cancellation

Lemma 7 Let $N_{\ell; k}$ have property $P_{\ell; k}$. Then,

1. $a \mid bc$.
2. $b \mid ac(B - 1)$.
3. $c \mid abB^k$.

Proof Using the fact that N has the $P_{\ell; k}$ property, we have

$$(aB + b)c = a(bB^k + c).$$

This can be rewritten by collecting each of a, b, c successively on one side, giving

$$a(bB^k - (B - 1)c) = bc, \quad b(aB^k - c) = ac(B - 1), \quad c(a(B - 1) + b) = abB^k,$$

from which the desired rules follow. \square

Lemma 8 Let $N_{\ell; k}$ have property $P_{\ell; k}$. If any one of $a, b, c = 0$, then at least one of the others is also 0, i.e. N is a trivial solution for the $P_{\ell; k}$ problem.

Proof This follows from the divisibility conditions in Lemma 7. \square

Lemma 9 Let N_k have property P_k and at least one of the following hold.

1. $a = c$.
2. $a_i = b$ for all $1 \leq i \leq k$.
3. $c_i = b$ for all $1 \leq i \leq k$.

Then, all the digits $a_i = c_i = b$, i.e. N is a trivial solution for the P_k problem.

Proof Let N have property P_k , whence $(aB + b)c = a(bB^k + c)$. Denote

$$I = \underbrace{[11 \dots 1]}_k = \frac{B^k - 1}{B - 1}.$$

1. Putting $a = c$,

$$a = \frac{B^k - 1}{B - 1} \cdot b, \quad \sum_{i=1}^k a_i B^{k-i} = \sum_{i=1}^k b B^{k-i}.$$

By the uniqueness of representation in the base B , each $a_i = b$.

2. Putting $a = bI$,

$$c = \frac{abB^k}{a(B - 1) + b} = \frac{b^2 IB^k}{b(I(B - 1) + 1)} = bI.$$

3. Putting $c = bI$,

$$a = \frac{bc}{bB^k - (B-1)c} = \frac{b^2I}{bB^k - b(B-1)I} = \frac{b^2I}{b(B^k - (B^k - 1))} = bI. \quad \square$$

Appendix B Uneven blocks.

Although we primarily deal with solutions of P_k^* in this paper, it is necessary to make a short detour and examine a few aspects of the more general $P_{\ell; k}^*$ problem in order to prove Theorem 2.

Lemma 10 *Let B be an arbitrary integer base and let $N_{\ell; k}$ have property $P_{\ell; k}$. If $c_k = 0$, then the number*

$$N^- = [a_1 a_2 \dots a_\ell b c_1 c_2 \dots c_{k-1}]$$

has property $P_{\ell; k-1}$

Proof Denote

$$c' = [c_1 c_2 \dots c_{k-1}], \quad c = c'B + c_k = c'B.$$

Since N has property $P_{\ell; k}$, we have $(aB + b)c = a(bB^k + c)$, hence

$$(aB + b)c'B = a(bB^k + c'B) \quad (aB + b)c' = a(bB^{k-1} + c'),$$

which is precisely the statement that N^- has property $P_{\ell; k-1}$. □

Lemma 11 *Let B be an arbitrary integer base and let $N_{\ell; k}$ have property $P_{\ell; k}^*$. Then, $\ell \leq k$. In other words, there are no solutions of $P_{\ell; k}^*$ when $\ell > k$.*

Proof If $N_{\ell; k}$ has property $P_{\ell; k}^*$, then $a, b, c > 0$,

$$(aB + b)c = a(bB^k + c), \quad a(bB^k + c - Bc) = bc.$$

Put $bB^k + c - Bc = bc/a = d$, which is a positive integer. Suppose that $1 \leq d < B$, i.e. d is a single digit in base B . Expanding $bB^k + c = cB + d$ gives us

$$bB^k + c_1 B^{k-1} + \dots + c_{k-1} B + c_k = c_1 B^k + c_2 B^{k-1} + \dots + c_k B + d.$$

By the uniqueness of representation of integers in the base B , we equate the coefficients $b = c_1$, $c_1 = c_2$, \dots , $c_{k-1} = c_k$, $c_k = d$; specifically, $b = d$. Thus, $a = bc/d = c$, hence the solution is trivial by Lemma 9.

This means that for N to be a non-trivial solution, we must have $d \geq B$. Now, $0 < b < B$ and $0 < c < B^k$, hence

$$a = \frac{bc}{d} < \frac{B \cdot B^k}{B} = B^k.$$

This shows that a can have at most k digits, hence $\ell \leq k$. □

Corollary 12 *If N_k has property P_k^* , then $c_k \neq 0$.*

Proof If $c_k = 0$, we see that

$$N^- = [a_1 a_2 \dots a_k b c_1 c_2 \dots c_{k-1}]$$

has property $P_{k; k-1}$, and hence must be a trivial solution by Lemma 11. Furthermore, it must be trivial in the sense that one of $a, b, c = 0$; if not, then $a < B^{k-1}$ from the lemma contradicts the fact that a is a k -digit number. Thus, the original number N_k is also a trivial solution. \square

Corollary 13 *If N_k has property $P_{\ell; k}^*$, then the integer $d = bc/a \geq B$.*

The technique used in Lemma 11 can be employed to obtain an even sharper bound on the leading block a of a solution N_k .

Lemma 14 *Let B be an arbitrary integer base and let N_k have property P_k^* . Then, $a < B^k/2$. As a result, the leading digit $a_1 < B/2$.*

Proof Continuing along the same lines as the proof of Lemma 11, suppose that N_k has property P_k^* . Then $a, b, c > 0$,

$$(aB + b)c = a(bB^k + c), \quad a(bB^k + c - Bc) = bc,$$

and $bB^k + c - Bc = bc/a = d$ is a positive integer. We saw that when $1 \leq d < B$, the solution N is trivial. Furthermore, when $d \geq 2B - 2$, observe that

$$a = \frac{bc}{d} \leq \frac{(B-1) \cdot (B^k - 1)}{2B - 2} = \frac{1}{2}(B^k - 1).$$

We now examine the remaining case $B \leq d < 2B - 2$. Setting $d' = d - B$, we have $0 \leq d' < B - 2$, i.e. d' is a single digit in base B . Expanding $bB^k + c = cB + d$ gives us

$$\begin{aligned} bB^k + c_1 B^{k-1} + \dots + c_{k-1} B + c_k \\ = c_1 B^k + c_2 B^{k-1} + \dots + c_k B + (B + d'). \end{aligned}$$

This implies $B \mid c_k - d'$; but $0 \leq |c_k - d'| < B$ forcing $c_k = d'$. Since $k \geq 1$, we can subtract $c_k = d'$ and divide B , yielding

$$\begin{aligned} bB^{k-1} + c_1 B^{k-2} + \dots + c_{k-1} \\ = c_1 B^{k-1} + c_2 B^{k-2} + \dots + c_{k-1} B + d' + 1. \end{aligned} \quad (\star)$$

Since $d' < B - 2$, the number $d' + 1 < B - 1$ is a single digit, so we can equate coefficients and see that $b = c_1$, $c_1 = c_2$, \dots , $c_{k-2} = c_{k-1}$, $c_{k-1} = d' + 1$, hence $b = c_1 = \dots = c_{k-1} = d' + 1$. In other words, all the digits of $c + 1$ are exactly $d' + 1$, so if we set

$$I = [1_1 1_2 \dots 1_k] = \frac{B^k - 1}{B - 1},$$

then $c + 1 = (d' + 1)I$. Then,

$$a = \frac{bc}{d} = \frac{(d' + 1)[(d' + 1)I - 1]}{B + d'} < \frac{d' \cdot [(B - 1)I - 1]}{2d'} = \frac{1}{2}[(B - 1)I - 1]$$

hence putting $(B - 1)I = B^k - 1$ gives

$$a < \frac{1}{2}(B^k - 2).$$

□

Appendix C The trailing block.

This section deals with the trailing block c of a solution N_k , which can be almost completely described in terms of the central block b .

Lemma 15 *Let B be an arbitrary integer base and let N_k have property P_k^* . Then, the digits in the last block satisfy $c_k < c_{i < k} = b$. In other words, all solutions of P_k^* look like*

$$N = [a_1 a_2 \dots a_k b b b \dots b c_k]$$

Proof As before, suppose that N_k has property P_k^* . Then $a, b, c > 0$,

$$(aB + b)c = a(bB^k + c), \quad a(bB^k + c - Bc) = bc,$$

and $bB^k + c - Bc = bc/a = d$ is a positive integer. By Corollary 13, we have $d \geq B$. Since

$$d = \frac{bc}{a} < \frac{(B - 1) \cdot B^k}{B^{k-1}} < (B - 1)B < B^2,$$

we have $d = qB + d'$, with $0 < q, d' < B$; the fact that $q > 0$ follows from $d \geq B$. Expanding $bB^k + c = cB + d$, we have

$$\begin{aligned} bB^k + c_1 B^{k-1} + \dots + c_{k-1} B + c_k \\ = c_1 B^k + c_2 B^{k-1} + \dots + c_k B + (qB + d'). \end{aligned} \quad (\star)$$

We have $B \mid c_k - d'$, forcing $c_k = d'$.

Consider the case $k = 1$, where our equation now reads $bB + c = cB + d$, hence $bB + d' = d'B + qB + d'$, so $bB = (d' + q)B$. Thus, $b = d' + q > d' = c$ as desired.

Now let $k \geq 2$. Subtracting $c_k = d'$ from both sides of (\star) and dividing by B gives

$$\begin{aligned} bB^{k-1} + c_1 B^{k-2} + \dots + c_{k-2} B + c_{k-1} \\ = c_1 B^{k-1} + c_2 B^{k-2} + \dots + c_{k-1} B + d' + q. \end{aligned}$$

Note that $2 \leq d' + q \leq 2B - 2 < 2B$, so expand $d' + q = q'B + r$ for some $0 \leq r < B$, and $q' = 0, 1$. If $q' = 0$, then

$$\begin{aligned} bB^{k-1} + c_1 B^{k-2} + \dots + c_{k-2} B + c_{k-1} \\ = c_1 B^{k-1} + c_2 B^{k-2} + \dots + c_{k-1} B + r, \end{aligned}$$

hence we can equate coefficients yielding $b = c_1$, $c_1 = c_2$, \dots , $c_{k-2} = c_{k-1}$, $c_{k-1} = r = d' + q = c_k + q > c_k$. In other words, all $b = c_{i < k} > c_k$.

12 *Characterising Solutions of Anomalous Cancellation*

Otherwise, $q' = 1$, and

$$\begin{aligned} bB^{k-1} + c_1B^{k-2} + \cdots + c_{k-2}B + c_{k-1} \\ = c_1B^{k-1} + c_2B^{k-2} + \cdots + (c_{k-1} + 1)B + r. \end{aligned}$$

This gives $B \mid c_{k-1} - r$, hence $c_{k-1} = r$ anyways. Now if $r = B - 1$, we would have $d' + q = q'B + r = 2B - 1$; this contradicts $q' + d \leq 2B - 2$. Thus, $r \leq B - 2$, so $c_{k-1} + 1 = r + 1 \leq B - 1$ is a single digit in base B . Equating coefficients, $b = c_1$, $c_1 = c_2, \dots, c_{k-2} = c_{k-1} = r$. Furthermore, $c_k = d' = (q'B + r) - q = (B - q) + r < r = c_{k-1}$. Thus, we again have $b = c_{i < k} > c_k$. \square

Corollary 16 *Let B be an arbitrary integer base and let N_k have property P_k^* . Then,*

$$a = \frac{bM}{B} + \frac{bc_kB^{k-1}}{bB - (B-1)c_k}, \quad c = bM + c_k, \quad M = \frac{B^k - B}{B - 1}.$$

Since $bM/B = b(B^{k-1} - 1)/(B - 1)$ is an integer for $k > 1$, so is $bc_kB^{k-1}/(bB - (B - 1)c_k)$.

Proof Recall that

$$(aB + b)c = a(bB^k + c), \quad a(bB^k + c - Bc) = bc.$$

Since $b = c_{i < k}$ by Lemma 15, we can write

$$c = [bb \dots bc_k] = bB^{k-1} + \cdots + bB + c_k = b \frac{B^k - B}{B - 1} + c_k = bM + c_k.$$

Thus,

$$a = \frac{bc}{bB^k - (B - 1)c} = \frac{b(bM + c_k)}{bB^k - (b(B^k - B) + (B - 1)c_k)} = \frac{b(bM + c_k)}{bB - (B - 1)c_k}.$$

Now, note that

$$(bB - (B - 1)c_k) \frac{M}{B} = bM - (B^{k-1} - 1)c_k = bM + c_k - B^{k-1}c_k.$$

Thus,

$$\frac{bM + c_k}{bB - (B - 1)c_k} = \frac{M}{B} + \frac{c_kB^{k-1}}{bB - (B - 1)c_k},$$

whence

$$a = \frac{b(bM + c_k)}{bB - (B - 1)c_k} = \frac{bM}{B} + \frac{bc_kB^{k-1}}{bB - (B - 1)c_k}.$$

\square

The following observation regarding the final digit c_k ties up the proof of Theorem 2. This requires diving into the prime factorisation of the base B .

Lemma 17 *Let B be an arbitrary integer base and let N_k have property P_k^* . Then, $\gcd(c_k, B) > 1$, i.e. the last digit c_k must share some factor $p > 1$ with B . Furthermore, if $a_k \neq b$, then $\gcd(a_k - b, B) > 1$.*

Proof Suppose that N_k has property P_k^* . Then $a, b, c > 0$, $c_k > 0$, and

$$(aB + b)c = a(bB^k + c), \quad (ac - abB^{k-1})B = (a - b)c.$$

This gives $B \mid (a - b)c$; writing

$$\begin{aligned} a' &= [a_1 a_2 \dots a_{k-1}], & a &= a'B + a_k, \\ c' &= [c_1 c_2 \dots c_{k-1}], & c &= c'B + c_k, \end{aligned}$$

we have $B \mid (a - b)(c'B + c_k)$, hence $B \mid (a - b)c_k = (a'B + a_k - b)c_k$, hence $B \mid (a_k - b)c_k$. Let

$$B = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_m^{\alpha_m}$$

be the prime factorization of B , with each $\alpha_i \geq 1$. Then, for each prime factor p_i of B , $p_i^{\alpha_i} \mid (a_k - b)c_k$. Let β_i be the greatest integer such that $p_i^{\beta_i} \mid c_k$. Then, we must have $p_i^{\alpha_i - \beta_i} \mid a_k - b$; if not, the greatest power of p_i dividing $(a_k - b)c_k$ would have been strictly less than $\beta_i + (\alpha_i - \beta_i) = \alpha_i$, a contradiction.

It is clear that we cannot have all $\beta_i \geq \alpha_i$; if so, we would have all $p_i^{\alpha_i} \mid c_k$, hence the product $p_1^{\alpha_1} \dots p_m^{\alpha_m} = B \mid c_k$, but $0 < c_k < B$, a contradiction. Thus, there must be some $\beta_j < \alpha_j$ corresponding to which $p_j^{\alpha_j - \beta_j} \mid a_k - b$, hence $\gcd(a_k - b, B) \geq p_j > 1$. This proves the second part of our lemma.

To prove the first part, we use induction on the block size k . Consider $k = 1$, where $a_k = a$, $c_k = c$, and suppose that all $\beta_i = 0$. This forces all $p_i^{\alpha_i - \beta_i} = p_i^{\alpha_i} \mid a - b$, hence their product $p_1^{\alpha_1} \dots p_m^{\alpha_m} = B \mid a - b$. But $0 \leq |a - b| < B$, forcing $a - b = 0$. By Lemma 9, this gives a trivial solution, a contradiction. Thus, there must be some $\beta_{j'} > 0$, hence $\gcd(c, B) \geq p_j^{\beta_j} > 1$.

Next, suppose that the statement holds for some $k \geq 1$, and let

$$N = [a_1 a_2 \dots a_k a_{k+1} b c_1 c_2 \dots c_{k+1}] = [a b c]$$

have property P_{k+1}^* . Then, $a, b, c, > 0$, $c_{k+1} > 0$, and $B \mid (a_{k+1} - b)c_{k+1}$. Again, if all $\beta_i = 0$, then we must have all $p_i^{\alpha_i - \beta_i} = p_i^{\alpha_i} \mid a_{k+1} - b$, hence their product $B \mid a_{k+1} - b$. Since $0 \leq |a_{k+1} - b| < B$, we have $a_{k+1} - b = 0$. But we also have $b = c_1 > c_{k+1}$ by Lemma 15. By Proposition 1, the number with the digits $a_{k+1}, c_1 = b$ removed, i.e.

$$N' = [a_1 a_2 \dots a_k b c_2 \dots c_{k+1}],$$

must have the P_k^* property (each block remains non-zero, and $b > c_{k+1}$ so not all digits are equal). Applying our induction hypothesis, we have $\gcd(c_{k+1}, B) > 1$.

Thus, by induction, our statement holds for all $k \geq 1$. \square

Appendix D Estimate of saturation points.

We are now ready to prove Theorem 5 using Corollary 16.

Proof of Theorem 5 Suppose that the digits b, c_k generate a solution of the P_k^* problem as per Lemma 15; further suppose that they do *not* generate a solution for the P_{k-1}^* problem. Then, Corollary 16 guarantees that

$$\frac{bc_k B^{k-1}}{bB - (B-1)c_k}$$

is an integer. Since b, c_k do not generate a solution for the P_{k-1}^* problem, either

$$\frac{bc_k B^{k-2}}{bB - (B-1)c_k}$$

is not an integer, or the first block a' is too small, i.e. the block

$$a' = \frac{b(B^{k-2} - 1)}{B - 1} + \frac{bc_k B^{k-2}}{bB - (B-1)c_k} < B^{k-2}.$$

The divisibility conditions in the first case are enough to obtain certain relations between k and the prime factors of B and bc_k . The bound on k in the second case requires only direct algebraic manipulation.

Consider the former case, and factorize

$$B = p_1^{\alpha_1} \cdots p_r^{\alpha_r}, \quad bc_k = p_1^{\beta_1} \cdots p_r^{\beta_r},$$

where p_1, \dots, p_r are primes, and each $\alpha_i, \beta_i \geq 0$. Then, the denominator

$$bB - (B-1)c_k \mid bc_k B^{k-1} = p_1^{\alpha_1(k-1)+\beta_1} \cdots p_r^{\alpha_r(k-1)+\beta_r},$$

hence its prime factorization cannot have any primes apart from p_1, \dots, p_r . Write

$$bB - (B-1)c_k = p_1^{\gamma_1} \cdots p_r^{\gamma_r}$$

for $\gamma_i \geq 0$. On the other hand, the denominator

$$bB - (B-1)c_k \nmid bc_k B^{k-2} = p_1^{\alpha_1(k-2)+\beta_1} \cdots p_r^{\alpha_r(k-2)+\beta_r}.$$

This means that we must have some j for which $\gamma_j > \alpha_j(k-2) + \beta_j$. Furthermore, $\gamma_j \leq \alpha_j(k-1) + \beta_j$ so as to satisfy the first divisibility, ensuring that $\alpha_j \neq 0$. Now,

$$(B-1)^2 \geq (B-1)(B-c_k) \geq bB - (B-1)c_k \geq p_j^{\gamma_j} > p_j^{\alpha_j(k-2)+\beta_j},$$

which gives

$$(B-1)^2 > p_j^{\alpha_j(k-2)+\beta_j} \geq 2^{\alpha_j(k-2)}, \quad 2 \log_2(B-1) > \alpha_j(k-2).$$

Since $\alpha_j \geq 1$,

$$k-2 < \frac{2 \log_2(B-1)}{\alpha_j} \leq 2 \log_2(B-1), \quad k < 2 \log_2(B-1) + 2.$$

Next, we consider the latter case where the block a' for the P_{k-1}^* problem is too small, hence

$$a' = \frac{b(B^{k-2} - 1)}{B - 1} + \frac{bc_k B^{k-2}}{bB - (B-1)c_k} < B^{k-2}.$$

However, the corresponding block a for the P_k^* problem is of the right size, hence

$$a = \frac{b(B^{k-1} - 1)}{B - 1} + \frac{bc_k B^{k-1}}{bB - (B-1)c_k} \geq B^{k-1}.$$

Collecting the powers of B , we have the equations

$$B^{k-2} \left[\frac{b^2 B}{(B-1)(bB - (B-1)c_k)} - 1 \right] < \frac{b}{B-1},$$

$$B^{k-1} \left[\frac{b^2 B}{(B-1)(bB - (B-1)c_k)} - 1 \right] \geq \frac{b}{B-1}.$$

Note that the term inside the square bracket must be positive by the second equation. Thus, we can divide the first equation by this and obtain the estimate

$$\begin{aligned} B^{k-2} &< \frac{b}{B-1} \cdot \frac{(B-1)(bB - (B-1)c_k)}{b^2B - (B-1)(bB - (B-1)c_k)} \\ &= \frac{b^2B - (B-1)bc_k}{b^2B - bB(B-1) + (B-1)^2c_k} \\ &\leq b^2B - (B-1)bc_k \\ &\leq (B-1)^3. \end{aligned}$$

Thus,

$$(B-1)^{k-2} < B^{k-2} \leq (B-1)^3, \quad k-2 < 3, \quad k < 5.$$

This means that new solutions cannot appear with increasing k , when

$$k \geq 2 \log_2(B-1) + 2, \quad \text{and} \quad k \geq 5,$$

that is,

$$k \geq \max\{5, 2 \log_2(B-1) + 2\}.$$

□

Appendix E Powers of primes.

In this section, we examine solutions of P_k^* where the base $B = p^n$ for prime p , $n \geq 2$.

Lemma 18 *Let $B = p^n$ where p is prime, $n > 1$. If N_k has property P_k^* for $k \geq 1$, then $a_{i \neq 1} = b = c_{i \neq k}$. Furthermore, $p \mid c_k$.*

Proof Suppose that N_k has property P_k^* for $k > 1$. Then, $a, b, c > 0$ via Lemma 8, the last digit $c_k > 0$, and $p \mid c_k$ using Lemma 17. By Corollary 16, write

$$(a, b, c) = \left(\frac{bM}{B} + \frac{bc_k B^{k-1}}{bB - (B-1)c_k}, b, bM + c_k \right), \quad M = \frac{B^k - B}{B-1}.$$

Specifically,

$$\frac{bc_k B^{k-1}}{bB - (B-1)c_k} = q$$

is an integer. Write $B = p^n$, and $c_k = p^r c'_k$, $q = p^s q'$, $b = p^t b'$ with $p \nmid c'_k, q', b'$. Since $c_k < B = p^n$ is a single digit, we must have $r < n$. Now, we have

$$bc_k p^{n(k-1)} = q(bp^n - (p^n - 1)c_k) = q((b - c_k)p^n + c_k),$$

hence

$$bc'_k p^r p^{n(k-1)} = q((b - c_k)p^n + p^r c'_k), \quad bc'_k p^{n(k-1)} = q((b - c_k)p^{n-r} + c'_k).$$

Now,

$$b' c'_k p^t p^{n(k-1)} = q' p^s ((b - c_k)p^{n-r} + c'_k),$$

hence

$$b' c'_k p^{n(k-1)+t-s} = q'(b - c_k)p^{n-r} + q' c'_k.$$

Note that we have integers on both sides. Since $r < n$, we have $p \mid q'(b - c_k)p^{n-r}$; but by construction, $p \nmid q'c'_k$. Thus, the left hand side $p \nmid b'c'_k p^{n(k-1)+t-s}$. Again, $p \nmid b'c'_k$, hence we have $s = n(k - 1) + t$. Thus,

$$\frac{bc_k}{bB - (B - 1)c_k} = \frac{q}{p^{n(k-1)}} = q'p^{s-n(k-1)} = q'p^t$$

is an integer.

Now, the number $N_* = [a_* b_* c_*]$ where

$$(a_*, b_*, c_*) = \left(\frac{bc_k}{bB - (B - 1)c_k}, b, c_k \right)$$

has property P_1^* . To see this, note that $c_k < b$ gives

$$a_* = \frac{bc_k}{bB - (B - 1)c_k} < \frac{bc_k}{bB - (B - 1)b} = c_k < B$$

ensuring that a_* is a single digit, and that

$$\frac{1}{a_*} + \frac{B - 1}{b_*} = \frac{bB - (B - 1)c_k}{bc_k} + \frac{B - 1}{b} = \frac{B}{c_k} = \frac{B^1}{c_*},$$

satisfying the P_k property.

By Proposition 1, its extension

$$N_*^+ = [a_* b_* \dots b_* b_* b_* \dots b_* c] = [a_* b \dots b b b \dots b c_k] = [a_*^+ b c]$$

has property P_k^+ . However, the digits b, c_k uniquely determine the first block a_*^+ , and we already have a solution $N = [a b c]$ generated by b, c_k . This forces $a = a_*^+$, hence all $a_{i \neq 1} = b$ as desired. In other words, our original solution N for P_k is an extension of the solution N_* for P_1 . \square

References

- [1] Sholander, M. (1961). Problems and Questions. *Mathematics Magazine*, 34(6): 363–373. doi:10.1080/0025570x.1961.11975268
- [2] Boas, R. P. (1972). Anomalous Cancellation. *The Two-Year College Mathematics Journal*, 3(2): 21–24. doi:10.2307/3026913
- [3] Ogilvy, C.S, Anderson J. T. (1996). Excursions in Number Theory. *Oxford University Press, New York*.
- [4] N. J. A. Sloane, The On Line Encyclopedia of Integer Sequences, <https://oeis.org>
- [5] Michael Glickman. Think Logically! Anomalous Cancellation in Depth, 07/23/2021, Challenging Math video, 41:02, https://youtu.be/0_PCqkbGs5E
- [6] Ekhad, Shalosh B. (2017). Automated Generation of Anomalous Cancellations. *arXiv*. doi:10.48550/ARXIV.1709.03379

- [7] Fried, M. N., & Goldberg, M. (2010). A Pumping Lemma for Invalid Reductions of Fractions. *The College Mathematics Journal*, 41(5): 357–364. doi:10.4169/074683410x521955
- [8] Stuffelbeam, R. (2013). How Weird Are Weird Fractions? *The College Mathematics Journal*, 44(3): 202—209. doi:10.4169/college.math.j.44.3.2