

# THE FULL SUPPORT PROPERTY OF PARABOLIC STOCHASTIC PDES WITH NONDEGENERATE HÖLDER DIFFUSION COEFFICIENTS

YI HAN

ABSTRACT. In this paper we work with parabolic SPDEs of the form

$$\partial_t u(t, x) = \partial_x^2 u(t, x) + g(t, x, u) + \sigma(t, x, u) dW(t, x)$$

where  $x \in [0, 1]$ ,  $W$  is the space-time white noise on  $[0, 1]$  and  $g$  is uniformly bounded. The diffusion coefficient  $\sigma$  is assumed to be uniformly elliptic but only Hölder continuous in  $u$ . Previously support theorems have only been established when  $\sigma$  is Lipschitz continuous in  $u$ . We obtain new support theorems and small ball probabilities in this  $\sigma$  Hölder continuous case via the recently established sharp two sided estimates of stochastic integrals.

## 1. INTRODUCTION

The support theorem for stochastic processes has a long history. One of its simplest forms can be phrased as follows: let  $B_t$  be a  $d$ -dimensional Brownian motion started from 0, then for any  $\epsilon, t > 0$  we have

$$\mathbb{P}(\sup_{s \leq t} |B_s| < \epsilon) > 0.$$

This follows from the reflection principle of Brownian motion. Via a Girsanov change of measure, we can deduce that for any continuous  $\psi : [0, t] \rightarrow \mathbb{R}^d$  with  $\psi(0) = 0$ , we have

$$\mathbb{P}(\sup_{s \leq t} |B_s - \psi(s)| < \epsilon) > 0.$$

See for example [3], page 59-60, (6.5) and (6.6). Both claims rely on Gaussian structure of the process  $B$ .

There is another notion, usually named Stroock–Varadhan support theorem for stochastic processes, that has a different flavour. Consider the parabolic SPDE

$$\partial_t u(t, x) = \partial_x^2 u(t, x) + g(t, x, u) + \sigma(t, x, u) dW(t, x), \quad x \in [0, 1]. \quad (1.1)$$

Denote by  $\mathcal{H} := \{h : [0, T] \times [0, 1] \rightarrow \mathbb{R}, h \text{ absolutely continuous}, \dot{h} \in L^2([0, T] \times [0, 1])\}$  and let  $S(h)$  denote the solution of (1.1) when we take  $\dot{h} \in \mathcal{H}$  in place of the white noise  $dW$ . If we assume  $\sigma$  is Lipschitz,  $u(0, \cdot)$  is Hölder continuous and  $g$  has sufficiently many derivatives, it is proved in [2] that the support of  $u$  on the Wiener space is the closure of  $S(h), h \in \mathcal{H}$ .

Several extensions of this support theorem have appeared, see for example [4], [5] and [7], with the same conclusion that the topological support of the solution  $u$  is the closure of  $S(h)$ , the solution of the SPDE driven by the control  $h$  in place of the noise. The support theorem to singular SPDEs has been obtained in [10], with the feature that the coset structure in the renormalization group plays a key role in characterizing topological

---

Supported by EPSRC grant EP/W524141/1.

supports of the solution. In all these works, the coefficients  $g$  and  $\sigma$  are nice enough so that the control problem  $S(h)$  can be properly solved, and the support of (1.1) can be characterized in terms of solutions to the control problem  $S(h)$ .

In this paper we consider support theorems of SPDEs in the lens of the regularization by noise phenomenon. In the setting of (1.1), this means that the coefficients  $g$  or  $\sigma$  (or both) are not necessarily locally Lipschitz continuous. One usually requires  $\sigma$  to be uniformly elliptic, so that roughness of the driving noise can restore well-posedness of the equation. We generally do not have a Stroock–Varadhan support theorem since the ODE or PDE for the control process  $S(h)$  is in general not well-posed. However it is still possible to prove the solution has full support or obtain small ball probability estimates

For additive noise, i.e.,  $\sigma = I_d$ , and assuming the drift  $g$  is not too singular, we can remove the drift via Girsanov transform and show the solution has full support because white noise has. A related example for finite dimensional SDEs with singular drifts can be found in [12]. The story is the same when  $\sigma$  is Lipschitz continuous and uniformly elliptic.

We are particularly interested in the *robustness* of the support theorems, in the remaining (hardest) case that  $\sigma$  is only  $\alpha$ -Hölder continuous in  $u$ ,  $\alpha \in (0, 1]$ . More precisely, we assume that for some  $\mathcal{C}_1, \mathcal{C}_2, \mathcal{D} > 0$  we have

$$|\sigma(t, x, u) - \sigma(t, x, v)| \leq \mathcal{D}|u - v|^\alpha, \quad \mathcal{C}_1 \leq |\sigma(t, x, u)| \leq \mathcal{C}_2.$$

for any  $t > 0$ ,  $x \in [0, 1]$  and  $u, v \in \mathbb{R}$ . In this case the Girsanov transform or the control processes  $S(h)$  do not tell us the answer in the same way as before.

Before stating our support theorems, it is crucial to discuss the well-posedness issues of (1.1). For  $\sigma$  to be  $\alpha$ -Hölder continuous in  $u$ ,  $\alpha > \frac{3}{4}$ , strong existence and uniqueness have been established in [14] (see also [11] for a different perspective, also with  $\alpha > \frac{3}{4}$ ). In general we may consider probabilistic weak solutions to (1.1) without the knowledge of uniqueness. So long as  $\alpha > 0$ , the lower bound in our main theorem still holds, and in particular every weak solution to (1.1) has full support.

Our strategy to characterize the support of (1.1) goes as follows. Instead of viewing (1.1) merely as an evolution equation, we regard it as a random field and probe into fine geometric properties of its fluctuations. Since  $\sigma$  is non-degenerate, one expects that (1.1) lies in the same universality class as the linear stochastic heat equation

$$\partial_t u(t, x) = \partial_x^2 u(t, x) + dW(t, x), \tag{1.2}$$

which is also named as the Edwards-Wilkinson universality class. In this universality class we observe nontrivial limiting behavior under the 1:2:4 scaling relation  $u(t, x) \mapsto \epsilon^{-1}u(\epsilon^{-4}t, \epsilon^{-2}x)$ . From this scaling relation we can expect fairly sharp two-sided probability estimates for the solution of (1.1) on small scales, even in the case that  $\sigma$  is not constant in  $u$ . Such computations have been carried out in [1], obtaining matching small ball probability estimates of solutions to (1.1) when  $\sigma$  is Lipschitz continuous in  $u$ , with a Lipschitz constant sufficiently small.

When  $\sigma$  is merely Hölder continuous, we expect that it induces a highly nonlinear stretching of space and time. To overcome this, we adjust the 1:2 space-time parabolic scaling through a reduction of the temporal length scale while keeping the spatial scale fixed. Consequently, we can still obtain nontrivial (upper and lower) probability estimates of fine scale properties of the solution, and this is already sufficient for us to prove the support theorem. The upper and lower bounds of small ball probabilities in this Hölder continuous case however do not have matching exponents in  $\epsilon$ , reflecting the fact that

irregularity of  $\sigma$  induces high order stretching in space and time. We note that for the linear stochastic heat equation (1.2), we can obtain small ball probabilities where the upper and lower bounds have matching exponents in  $\epsilon$ , see [6], page 171, Theorem 5.1.

We now state the main theorem.

Denote by  $\mathcal{P}$  the  $\sigma$ -field of the noise  $W(t, x)$ , generated by functions of the form  $f(x, t, \omega) = X(\omega) \cdot 1_A(x) \cdot 1_{(a,b]}(t)$ , with  $A \subset [0, 1]$  and  $X$  some  $\mathcal{F}_a$ -measurable random variable. We say  $h \in \mathcal{PC}_b^2$  if almost surely,  $h, \partial_t h, \partial_x^2 h$  are bounded by a fixed constant  $C$ .

**Theorem 1.1.** *Consider solutions  $u(t, x)$  to the stochastic heat equation*

$$\partial_t u(t, x) = \frac{1}{2} \partial_x^2 u(t, x) + g(t, x, u(t, x)) + \sigma(t, x, u(t, x)) dW(t, x), \quad u(0, x) = u_0(x). \quad (1.3)$$

Assume  $u_0, h \in \mathcal{PC}_b^2$  with

$$\sup_{x \in [0, 1]} |u_0(x) - h(0, x)| < \epsilon/2,$$

and that for some constants  $D, \mathcal{C}_1, \mathcal{C}_2 > 0$ ,  $\alpha \in (\frac{3}{4}, 1]$ , we have

$$|\sigma(t, x, u) - \sigma(t, x, v)| \leq \mathcal{D}|u - v|^\alpha,$$

$$\mathcal{C}_1 \leq |\sigma(t, x, u)| \leq \mathcal{C}_2$$

for all  $x \in [0, 1]$ ,  $u, v \in \mathbb{R}$  and  $t \geq 0$ .

Then for any  $\beta > 2 - \alpha$  we may find positive constants  $C_0, C_1, C_2, C_3$  and  $\epsilon_0$  depending on  $\beta, \mathcal{C}_1, \mathcal{C}_2$  and  $\sup_{t, x, u} |b(t, x, u)|$ , such that for any  $0 < \epsilon < \epsilon_0$ , we have

$$C_0 \exp\left(-\frac{C_1 T}{\epsilon^{2+4\beta}}\right) \leq P\left(\sup_{0 \leq t \leq T, x \in [0, 1]} |u(t, x) - h(t, x)| \leq \epsilon\right) \leq C_2 \exp\left(-\frac{C_3 T}{(1 + \mathcal{D}^2)\epsilon^{4+2\alpha}}\right). \quad (1.4)$$

If we only assume  $\alpha > 0$ , then the lower bound in (1.4) holds, that is,

$$C_0 \exp\left(-\frac{C_1 T}{\epsilon^{2+4\beta}}\right) \leq P\left(\sup_{0 \leq t \leq T, x \in [0, 1]} |u(t, x) - h(t, x)| \leq \epsilon\right).$$

Since  $h$  is arbitrary, this in particular implies that the solution  $u$  has full support on Wiener space with respect to the supremum norm.

*Remark 1.2.* The technical assumption  $\alpha \in (\frac{3}{4}, 1]$  is only used to match with the well-posedness results in [14] (see also [11]). Via a compactness argument it is easy to show existence of a (probabilistic) weak solution to (1.3) for any  $\alpha > 0$  (see for example [9]).

A natural question is whether every weak solution to (1.3) has the full support property. For the upper bound to be proved in Section 2.3, we are not sure if the proof carries over for any  $\alpha > 0$  or not because we have to solve another SPDE (2.3)<sup>1</sup> on the same probability space. This procedure requires strong well-posedness for every diffusion coefficient with the given Hölder continuity.

For the lower bound which is more relevant to the full support property, a careful check of the proof in Section 2.4 shows that the lower bound in (1.4) holds for any weak solution of (1.3) whenever  $\alpha > 0$ . Therefore, every weak solution to (1.3) must have full support.

*Remark 1.3.* We have for simplicity worked on the unit interval  $[0, 1]$ , but everything carries over to finite intervals  $[0, J]$  for any  $J > 0$ . In this paper we assume the solution  $u$  is real-valued, but everything carries over to the vector valued case  $\mathbf{u}(t, x) \in \mathbb{R}^d$ ,  $d \in \mathbb{N}_+$  without change. These extensions can be found in [1].

<sup>1</sup>with the diffusion coefficient  $\sigma$  only  $\alpha$ -Hölder continuous in the solution argument

There are a few remaining questions. One of them is to obtain support theorems in Hölder semi-norm rather than the supremum norm. Fairly sharp results have been obtained when  $\sigma$  is Lipschitz continuous in  $u$  (see [8] for recent progress), but adapting estimates in the existing literature to our Hölder continuous  $\sigma$  seems a bit out of reach. Another more fundamental question is if we assume  $\sigma$  is uniformly elliptic, it is not clear whether the assumption that the Hölder index  $\alpha > \frac{3}{4}$  is necessary for (strong or weak) well-posedness of (1.3). We believe that  $\alpha > 0$  is enough for weak well-posedness but could not give a proof. Note that when  $\sigma$  is not assumed to be uniformly elliptic, then  $\alpha > \frac{3}{4}$  is a sharp condition, see [13] for the case  $\alpha < \frac{3}{4}$ .

## 2. PROOF OF MAIN THEOREM

**2.1. Reduction to simple cases.** We first show that after some simple reductions we can assume  $g = 0$  and  $h = 0$ . These reductions follow from Girsanov theorem and the non-degeneracy of  $\sigma$ .

The solution to (1.3) can be reformulated as

$$\partial_t u(t, x) = \frac{1}{2} \partial_x^2 u(t, x) + \sigma(t, x, u(t, x)) [dW(t, x) + \sigma^{-1} g(t, x, u(t, x))],$$

we consider the probability law  $Q_t$  defined as

$$\frac{dQ_t}{dP_t} = \int_0^t \int_0^1 \sigma^{-1} g(t, x, u(t, x)) dW(t, x) - \frac{1}{2} \int_0^t \int_0^1 (\sigma^{-1} g(t, x, u(t, x)))^2 dx dt. \quad (2.1)$$

By Girsanov theorem,  $dW(t, x) + \sigma^{-1} g(t, x, u(t, x))$  is a space-time white noise with respect to  $Q_t$ . Denote by  $A$  the event that

$$A = \left\{ \sup_{s \in [0, T], y \in [0, 1]} |u(s, y) - h(s, y)| < \epsilon \right\},$$

then by Cauchy-Schwartz inequality

$$Q_T(A) \leq \sqrt{P_T(A)} \sqrt{E\left(\frac{dQ_T}{dP_T}\right)^2} \leq \sqrt{P_T(A)} M, \quad (2.2)$$

where  $M$  depends only on  $T, \mathcal{C}_1$  and  $\sup_{t, x, y} |g(t, x, y)|$ . This implies the lower bound in (1.4) for general  $g$  can be deduced from the lower bound in the case  $g = 0$ .

For the upper bound, a similar argument holds: one only needs to swap  $P$  and  $Q$  in (2.2) and replace  $g$  by  $-g$  in (2.1).

Now we show why we can take  $h = 0$ . This is outlined in page 6 of [1] but we reproduce here for completeness. Let  $H := \partial_t - \frac{1}{2} \partial_x^2$  and consider the process

$$w(t, x) = u(t, x) - u_0(x) - h(t, x) + h_0(x),$$

so that  $w(0, x) = 0$ . If we set  $\sigma_1(t, x, w) = \sigma(t, x, u)$  and

$$g_1(t, x, w) = g(t, x, u) - H u_0(x) - H h(t, x) + H h_0(x),$$

we have

$$\partial_t w(t, x) = \frac{1}{2} \partial_x^2 w(t, x) + g_1(t, x, w) + \sigma_1(t, x, w) dW(t, x).$$

Since  $u_0, h \in \mathcal{P}\mathcal{C}_b^2$ ,  $\sup_{t, x, \omega} |g_1(t, x, \omega)| < \infty$ , so we are reduced to the case  $h = 0$  and  $u(0, x) \equiv 0, x \in [0, 1]$ .

**2.2. Sharp two-sided estimates.** Recall the heat kernel on  $[0, 1]$  is given by

$$G(t, x) = \sum_{n \in \mathbb{Z}} (2\pi t)^{-1/2} \exp\left(-\frac{(x+n)^2}{2t}\right).$$

Consider the noise term  $\mathbf{N}$  defined as

$$\mathbf{N}(t, x) := \int_0^t \int_0^1 G(t-s, x-y) \sigma(s, y, u(s, y)) W(dy ds).$$

We quote the following large deviations estimate from [1], Proposition 3.4 and Remark 3.1, which is a very precise formulation of the 1:2:4 scaling of Gaussian processes:

**Proposition 2.1.** *Assume that  $\sup_{s,y} |\sigma(s, y, u(s, y))| \leq C < \infty$ . Then we can find universal constants  $K_1$  and  $K_2$  such that, for any  $\alpha, \lambda, \epsilon > 0$ ,*

$$\mathbb{P} \left( \sup_{0 \leq t \leq \alpha \epsilon^4, x \in [0, \epsilon^2]} |\mathbf{N}(t, x)| > \lambda \epsilon \right) \leq \frac{K_1}{1 \wedge \sqrt{\alpha}} \exp\left(-K_2 \frac{\lambda^2}{C^2 \sqrt{\alpha}}\right).$$

We fix a sufficiently small  $c_0 > 0$  such that  $0 < c_0 < \max\left\{\left(\frac{K_2}{36 \log K_1 C_2^2}\right)^2, 1\right\}$ , and define the discretized mesh of time as:<sup>2</sup>

$$t_n = n c_0 \epsilon^4, \quad n \geq 0,$$

and denote by  $I_n := [t_n, t_{n+1}]$  the time interval with numbering  $n$ . Choose some  $\theta = \theta(C_1, C_2) > 0$  sufficiently large (with the precise condition given in [1], (2.11)) and fix  $c_1^2 = \theta c_0$ , the spatial mesh points are chosen as

$$x_n = n c_1 \epsilon^2, \quad n \geq 0.$$

This time-space mesh respects the parabolic 1:2 scaling. Fix a terminal time  $T > 0$  and define the terminal index

$$n_1 := \min\{n \geq 1 : t_n > T\}, \quad n_2 := \min\{n \geq 1 : x_n > 1\}.$$

Write  $p_{i,j} = (t_i, x_j)$ , and consider the following two series of events

$$A_n = \{|u(t_{n+1}, x)| \leq \frac{\epsilon}{3}, \quad x \in [0, 1], \text{ and } |u(t, x)| \leq \epsilon, t \in I_n, x \in [0, 1]\},$$

and

$$F_n = \{|u(p_{n,j})| < \epsilon \text{ for all } j \leq n_2 - 2\}.$$

The strategy of proof is first fix the  $u$  component of  $\sigma$  and obtain an estimate in the Gaussian case, then deduce the general case via an interpolation argument. For the Gaussian case (when  $\sigma$  does not depend on  $u$ ), we quote the following result from [1], Proposition 2.1:

**Proposition 2.2.** *Under the assumptions of Theorem 1.1, assume further that  $g = 0$ ,  $u_0(x) \equiv 0$  and  $\sigma(t, x, u)$  does not depend on  $u$ .*

*Then there exists constants  $\epsilon_0, C_4, C_5 > 0$  which depend only on  $C_1$  and  $C_2$  such that for any  $0 < \epsilon < \epsilon_0$ ,*

$$P(F_n \mid \cap_{k=0}^{n-1} F_k) \leq C_4 \exp(-C_5 \epsilon^{-2}),$$

---

<sup>2</sup>Later we will introduce a different scheme to divide time intervals, which doesn't follow the 1:2 parabolic scaling.

and we can find constants  $C_6, C_7 > 0$  which depend only on  $\mathcal{C}_1, \mathcal{C}_2$  such that for any  $0 < \epsilon < \epsilon_0$ ,

$$P(A_n \mid \cap_{k=0}^{n-1} A_k) \geq C_6 \exp(-C_7 \epsilon^{-2}).$$

Now we prove the general case (i.e.,  $\sigma$  depends on  $u$ ).

**2.3. Upper bound, general case.** Define a function

$$f_\epsilon(z) = \begin{cases} z, & |z| < \epsilon, \\ \frac{\epsilon}{|z|}z, & |z| > \epsilon, \end{cases}$$

so that  $|f_\epsilon(z)| \leq \epsilon$  and  $f_\epsilon$  is Lipschitz continuous. We solve the following SPDE

$$\partial_t v(t, x) = \frac{1}{2} \partial_x^2 v(t, x) + \sigma(t, x, f_\epsilon(v(t, x))) \cdot dW(t, x) \quad (2.3)$$

with  $v(0, x) = u_0(x)$ , which is well posed because  $\sigma(t, x, f_\epsilon(u))$  is  $\alpha$ -Hölder continuous in  $u$ , for  $\alpha > \frac{3}{4}$ .

As long as  $|u(t, x)| \leq \epsilon$  for all  $x \in [0, 1]$  and  $t \in [0, t_1]$ , we have  $v(t, x) = u(t, x)$ , so we proceed with the proof for  $v$ .

The point is to compare  $v$  with an auxiliary process  $v_g$  defined by

$$\partial_t v_g(t, x) = \frac{1}{2} \partial_x^2 v_g(t, x) + \sigma(t, x, f_\epsilon(u_0(x))) \cdot dW(t, x),$$

with  $v_g(0, x) = u_0(x)$ , where the diffusion coefficient is independent of  $v_g$ .

The difference process  $D(t, x) := V(t, x) - V_g(t, x)$  is a stochastic integral satisfying

$$D(t, x) = \int_0^t \int_0^1 G(t-s, x-y) [\sigma(s, y, f_\epsilon(v(s, y))) - \sigma(s, y, f_\epsilon(u_0(y)))] \cdot W(dy ds).$$

Define

$$H_j = \{v(p_{1j}) \mid \leq \epsilon\}$$

and consider the events

$$A_{1,j} = \{|v_g(p_{1j}) \mid \leq 2\epsilon\},$$

$$A_{2,j} = \{|D(p_{1j}) \mid \geq \epsilon\}.$$

It is clear that  $H_j \subset A_{1,j} \cup A_{2,j}$ . Define another sequence of events

$$B_n = \{|u(t, x)| \leq \epsilon, t \in I_{n-1}, x \in [0, 1]\}, \quad n \geq 1.$$

Then clearly  $B_n \subset F_n$  and on  $B_n$ , we have  $u(t_n, x) = v(t_n, x)$ . Therefore

$$P(B_1) \leq P(\cap_{j=1}^{n_2-2} H_j) \leq P(\cap_{j=1}^{n_2-2} (A_{1,j} \cup A_{2,j})).$$

An elementary set-inclusion argument implies

$$P(B_1) \leq P(\cap_{j=1}^{n_2-2} A_{1,j}) + \sum_{j=1}^{n_2-2} P(A_{2,j}).$$

We now apply Proposition 2.2 to the process  $v_g$  to deduce

$$P(\cap_{j=1}^{n_2-2} A_{1,j}) = P(|v_g(p_{1,j}) \mid \leq 2\epsilon, \quad j = 1, \dots, n_2 - 2) \leq C_2 \exp(-C_3 \epsilon^{-2}).$$

By Hölder continuity of  $\sigma$  in  $u$ , we deduce that

$$|\sigma(s, y, f_\epsilon(v(s, y))) - \sigma(s, y, f_\epsilon(v_0(y)))| \leq \mathcal{D}(2\epsilon)^\alpha,$$

so that by Proposition 2.1, we have for  $j = 1, \dots, n_2 - 2$ ,

$$P(A_{2,j}) \leq K_1 \exp\left(-\frac{K_2}{4\epsilon^{2\alpha}\mathcal{D}^2\sqrt{c_0}}\right).$$

Therefore

$$\begin{aligned} P(B_1) &\leq C_2 \exp(-C_3\epsilon^{-2}) + \sum_{j=1}^{n_2-2} K_1 \exp\left(-\frac{K_2}{4\epsilon^{2\alpha}\mathcal{D}^2\sqrt{c_0}}\right) \\ &\leq C_2 \exp(-C_3\epsilon^{-2}) + \frac{1}{c_1\epsilon^2} K_1 \exp\left(-\frac{K_2}{4\epsilon^{2\alpha}\mathcal{D}^2\sqrt{c_0}}\right) \\ &\leq C_4 \exp\left(-\frac{C_5}{8(1+C_2)^2(1+\mathcal{D}^2)\epsilon^{2\alpha}\sqrt{c_0}}\right), \end{aligned}$$

whenever  $\epsilon$  is small enough, the  $\epsilon^{-2\alpha}$  term in the exponent wins over the  $\epsilon^{-2}$  term, so we keep the former.<sup>3</sup>  $C_4$  and  $C_5$  are universal constants that depend only on  $C_2$ .

The expression shows that when  $\sigma$  is merely Hölder continuous, i.e.  $\alpha < 1$ , the  $\epsilon^{2\alpha}$  term dominates in the upper bound. The upper and lower bounds we obtain will not have matching exponents of  $\epsilon$  (they do if  $\alpha = 1$ ), but both bounds are nontrivial and in particular they lead to the desired support theorem.

By the Markov property, for each  $n \leq n_1$ ,

$$P(B_n \mid \cap_{j=1}^{n-1} B_j) \leq \exp\left(-\frac{C_7}{(1+\mathcal{D}^2)\epsilon^{2\alpha}}\right)$$

where  $C_7$  depends only on  $C_1, C_2$  and  $\mathcal{D}$ .

Therefore

$$\begin{aligned} P(|u(t, x)| \leq \epsilon, t \in [0, T], x \in [0, 1]) &= P(\cap_{j=1}^{n_1-1} B_j) \\ &\leq \exp\left(-\frac{C_7}{(1+\mathcal{D}^2)\epsilon^{2\alpha}}\right)^{\frac{T}{\epsilon^4}} \\ &\leq \exp\left(-\frac{C_7 T}{(1+\mathcal{D}^2)\epsilon^{2\alpha+4}}\right). \end{aligned}$$

This establishes the upper bound in (1.4).

**2.4. Lower bound, general case.** We now proceed to prove the corresponding lower bound. The argument roughly follows that in [1], while the last key estimates are different.

Fix some  $\beta > 1$  to be determined later, consider a new time mesh as follows:

$$\hat{t}_n := nc_0\epsilon^{4\beta}, n \geq 0,$$

and the corresponding time intervals  $\hat{I}_n := [\hat{t}_n, \hat{t}_{n+1}]$ . This introduces a finer grid of time when  $\epsilon$  is sufficiently small. We analogously define the events

$$\hat{A}_n := \{|u(\hat{t}_{n+1}, x)| \leq \frac{\epsilon}{3}, \quad x \in [0, 1], \text{ and } |u(t, x)| \leq \epsilon, t \in \hat{I}_n, x \in [0, 1]\},$$

---

<sup>3</sup>We have used another approximation which follows from the elementary inequality

$$\sup_{x \geq 0} x^n e^{-Cx} < \infty$$

for any  $C > 0$  and  $n > 0$ .

Assuming that  $|u_0(x)| \leq \frac{\epsilon}{3}$ ,  $x \in [0, 1]$ . Define the stopping time

$$\tau = \inf\{t \geq 0 : \sup_{x \in [0,1]} |u(t, x) - u_0(x)| > 2\epsilon\},$$

such that on the event  $\hat{A}_0$  we have  $\tau \geq \hat{t}_1$ . Consider the process

$$\tilde{D}(t, x) = \int_0^t \int_0^1 G(t-s, x-y) [\sigma(s, y, u(s \wedge \tau, y)) - \sigma(s, y, u_0(y))] W(dy ds),$$

and the auxiliary comparison process  $u_g$  solving

$$\partial_t u_g = \frac{1}{2} \partial_x^2 u_g + \sigma(t, x, u_0(x)) dW(t, x), \quad u_g(0, x) = u_0(x),$$

and we write  $u(t, x) = u_g(t, x) + D(t, x)$  as before, so that

$$D(t, x) = \int_0^t \int_0^1 G(t-s, x-y) [\sigma(s, y, u(s, y)) - \sigma(s, y, u_0(y))] W(dy ds).$$

It is clear that  $D(t, x) = \tilde{D}(t, x)$  whenever  $\tau \geq \hat{t}_1$ . Consider the event

$$\tilde{B}_0 := \{|u_g(\hat{t}_1, x)| \leq \frac{\epsilon}{6}, x \in [0, 1], \quad |u_g(t, x)| \leq \frac{2\epsilon}{3} \forall t \in \hat{I}_0, x \in [0, 1]\}.$$

Then we have the following sequence of set inclusions

$$\begin{aligned} P(\hat{A}_0) &\geq P\left(\tilde{B}_0 \cap \left\{ \sup_{0 \leq t \leq \hat{t}_1, x \in [0,1]} |D(t, x)| \leq \frac{\epsilon}{6} \right\}\right) \\ &= P\left(\tilde{B}_0 \cap \left\{ \sup_{0 \leq t \leq \hat{t}_1, x \in [0,1]} |\tilde{D}(t, x)| \leq \frac{\epsilon}{6} \right\}\right) \\ &\geq P(\tilde{B}_0) - P\left(\sup_{0 \leq t \leq \hat{t}_1, x \in [0,1]} |\tilde{D}(t, x)| > \frac{\epsilon}{6}\right). \end{aligned} \tag{2.4}$$

The equality in the second line needs some explanation. If  $\tau \geq \hat{t}_1$ , then  $D = \tilde{D}$  on  $[0, \hat{t}_1]$ . On  $\tilde{B}_0 \cap \{\tau < \hat{t}_1\}$ , one must have  $\sup_x |u_g(\tau, x) - u_0(x)| > \epsilon$ , so that  $\sup_x |\tilde{D}(\tau, x)| > \epsilon$ .

Since  $\beta > 1$  and  $\epsilon < 1$ , one must have  $\hat{t}_1 < t_1$ , so that by Proposition 2.2,

$$P(\tilde{B}_0) \geq P(\bar{B}_0) \geq C_1 \exp\left(-\frac{C_2}{\epsilon^2}\right),$$

where  $C_1, C_2$  depend only on  $\mathcal{C}_1, \mathcal{C}_2$ , and where we define

$$\bar{B}_0 := \{|u_g(t_1, x)| \leq \frac{\epsilon}{6}, x \in [0, 1], \quad |u_g(t, x)| \leq \frac{2\epsilon}{3} \forall t \in I_0, x \in [0, 1]\}.$$

It remains to estimate the probability of the last event in (2.4).

$$\begin{aligned} P\left(\sup_{0 \leq t \leq \hat{t}_1, x \in [0,1]} |\tilde{D}(t, x)| > \frac{\epsilon}{6}\right) &\leq \frac{1}{\sqrt{c_0} \epsilon^2} P\left(\sup_{0 \leq t \leq \hat{t}_1, x \in [0, \sqrt{c_0} \epsilon^2]} |\tilde{D}(t, x)| > \frac{\epsilon}{6}\right) \\ &\leq \frac{1}{\sqrt{c_0} \epsilon^2} \frac{1}{\epsilon^{2\beta-2}} K_1 \exp\left(-\frac{K_2}{\epsilon^{2\alpha} \mathcal{D}^2 \mathcal{C}_2^2 \sqrt{c_0} \epsilon^{2\beta-2}}\right). \end{aligned}$$

To be a bit more precise about the exponent of  $\epsilon$ , we take  $\alpha = \epsilon^{4\beta-4}$  and the constant  $\mathcal{C} := \mathcal{D}(2\epsilon)^\alpha$  in the setting of Proposition 2.1. Comparing the exponents, we see that as



long as we take  $\alpha + \beta > 2$ , we can find some  $C_8, C_9$  depending only on  $C_1, C_2$  and  $\mathcal{D}$  such that

$$P\left(\sup_{0 \leq t \leq \hat{t}_1, x \in [0,1]} |\tilde{D}(t, x)| > \frac{\epsilon}{6}\right) \leq C_8 \exp\left(-\frac{C_9}{\epsilon^{2(\alpha+\beta-1)}}\right)$$

and finally

$$\mathbb{P}(\hat{A}_0) \geq C_1 \exp\left(-\frac{C_2}{\epsilon^2}\right) - C_8 \exp\left(-\frac{C_9}{\epsilon^{2(\alpha+\beta-1)}}\right),$$

with the first term dominating. So we conclude that when  $\epsilon > 0$  is sufficiently small, we can find  $C_3, C_4$  depending only on  $C_1, C_2, \mathcal{D}$  such that

$$\mathbb{P}(\hat{A}_0) \geq C_3 \exp\left(-\frac{C_4}{\epsilon^2}\right).$$

By the Markov property, for each  $n \leq \hat{n}_1 := \lfloor \frac{T}{\epsilon^{4\beta}} \rfloor$  we have

$$\mathbb{P}(\hat{A}_n \mid \cap_{j=0}^{n-1} \hat{A}_j) \geq C_3 \exp\left(-\frac{C_4}{\epsilon^2}\right).$$

Thus for some constant  $C_1 > 0$  depending on  $C_1, C_2$ ,

$$P(|u(t, x)| \leq \epsilon, t \in [0, T], x \in [0, 1]) \geq \exp\left(-\frac{C_1 T}{\epsilon^2 \epsilon^{4\beta}}\right) \geq \exp\left(-\frac{C_1 T}{\epsilon^{4\beta+2}}\right).$$

The various constants  $C_1, C_2, \dots, C_8, C_9$  depend only on  $C_1, C_2$  and may change from line to line. This establishes the lower bound in (1.4).

#### REFERENCES

- [1] Siva Athreya, Mathew Joseph, and Carl Mueller. “Small ball probabilities and a support theorem for the stochastic heat equation”. In: *The Annals of Probability* 49.5 (2021), pp. 2548–2572.
- [2] Vlad Bally, Annie Millet, and Marta Sanz-Solé. “Approximation and support theorem in Holder norm for parabolic stochastic partial differential equations”. In: *The Annals of Probability* (1995), pp. 178–222.
- [3] Richard F Bass. *Probabilistic techniques in analysis*. Springer Science & Business Media, 1994.
- [4] Caroline Cardon-Weber and Annie Millet. “A support theorem for a generalized Burgers SPDE”. In: *Potential Analysis* 15.4 (2001), pp. 361–408.
- [5] Igor Chueshov and Annie Millet. “Stochastic two-dimensional hydrodynamical systems: Wong-Zakai approximation and support theorem”. In: *Stochastic analysis and applications* 29.4 (2011), pp. 570–611.
- [6] Robert C Dalang et al. *A minicourse on stochastic partial differential equations*. Vol. 1962. Springer, 2009.
- [7] Francisco J Delgado-Vences and Marta Sanz-Solé. “Approximation of a stochastic wave equation in dimension three, with application to a support theorem in Hölder norm”. In: *Bernoulli* 20.4 (2014), pp. 2169–2216.
- [8] Mohammud Foondun, Mathew Joseph, and Kunwoo Kim. “Small ball probability estimates for the Hölder semi-norm of the stochastic heat equation”. In: *Probability Theory and Related Fields* (2022), pp. 1–61.
- [9] Dariusz Gatarek and Benjamin Goldys. “On weak solutions of stochastic equations in Hilbert spaces”. In: *Stochastics: An International Journal of Probability and Stochastic Processes* 46.1-2 (1994), pp. 41–51.

- [10] Martin Hairer and Philipp Schönbauer. “The support of singular stochastic partial differential equations”. In: *Forum of Mathematics, Pi*. Vol. 10. Cambridge University Press. 2022.
- [11] Yi Han. “Exponential ergodicity of stochastic heat equations with Hölder coefficients”. In: *arXiv preprint arXiv:2211.08242* (2022).
- [12] Chengcheng Ling, Sebastian Riedel, and Michael Scheutzow. “A Wong-Zakai theorem for SDEs with singular drift”. In: *Journal of Differential Equations* 326 (2022), pp. 344–363.
- [13] Carl Mueller, Leonid Mytnik, and Edwin Perkins. “Nonuniqueness for a parabolic SPDE with  $\frac{3}{4} - \epsilon$ -Hölder diffusion coefficients”. In: *The Annals of Probability* 42.5 (2014), pp. 2032–2112.
- [14] Leonid Mytnik and Edwin Perkins. “Pathwise uniqueness for stochastic heat equations with Hölder continuous coefficients: the white noise case”. In: *Probability theory and related fields* 149.1 (2011), pp. 1–96.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE.  
Email address: yh482@cam.ac.uk