

A note on the generalized maximal numerical range of operators

Abderrahim Baghdad ¹_{a,b}, El Hassan Benabdi ² and Kais Feki ³_{a,b}

ABSTRACT. The paper considers some new properties of the so-called A -maximal numerical range of operators, denoted by $W_{\max}^A(\cdot)$, where A is a positive bounded linear operator acting on a complex Hilbert space \mathcal{H} . Some characterizations of A -normaloid operators are also given. In particular, we extend a recent result by Spitkovsky in [Oper. Matrices, 13, 3(2019)]. Namely, it is shown that an A -bounded linear operator T acting on \mathcal{H} is A -normaloid if and only if $W_{\max}^A(T) \cap \partial W_A(T) \neq \emptyset$. Here $\partial W_A(T)$ stands for the boundary of A -numerical range of T . Some new A -numerical radius inequalities generalizing and improving earlier well-known results are also given.

1. INTRODUCTION AND PRELIMINARIES

Throughout this work \mathcal{H} stands for a non trivial complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and associated norm $\|\cdot\|$. By $\mathbb{B}(\mathcal{H})$, we denote the C^* -algebra of bounded linear operators acting on \mathcal{H} with the identity operator $I_{\mathcal{H}}$ (or simply I if no confusion arises). For simplicity, by an operator we mean an operator in $\mathbb{B}(\mathcal{H})$. For every operator T , its adjoint is denoted by T^* , its range by $\mathcal{R}(T)$ and its null space by $\mathcal{N}(T)$.

For the sequel, the following facts are useful. An operator T is said to be positive if $\langle Tx, x \rangle \geq 0$ for every $x \in \mathcal{H}$. By $\mathbb{B}(\mathcal{H})^+$, we denote the cone of

Date: February 2, 2023.

2020 Mathematics Subject Classification. 47B20, 47A12, 46C05, 47A10.

Key words and phrases. Positive operator, A -maximal numerical range, A -normaloid operator.

[¹] Department of Mathematics, Faculty of Sciences-Semlalia, University Cadi Ayyad, Marrakesh, Morocco.

Email address: bagabd66@gmail.com

[²] Department of Mathematics, Laboratory of Mathematics, Statistics and Applications, Faculty of Sciences, Mohammed V University in Rabat, Rabat, Morocco.

Email address: e.benabdi@um5r.ac.ma

[³_a] University of Monastir, Faculty of Economic Sciences and Management of Mahdia, Mahdia, Tunisia

[³_b] Laboratory Physics-Mathematics and Applications (LR/13/ES-22), Faculty of Sciences of Sfax, University of Sfax, Sfax, Tunisia

Email address: kais.feki@hotmail.com; kais.feki@fsegma.u-monastir.tn

positive (semi-definite) operators, i.e.

$$\mathbb{B}(\mathcal{H})^+ = \{T \in \mathbb{B}(\mathcal{H}) ; \langle Tx, x \rangle \geq 0, \forall x \in \mathcal{H}\}.$$

For the rest of this article, we suppose that $A \in \mathbb{B}(\mathcal{H})^+$ is a nonzero operator which clearly defines the following positive semidefinite sesquilinear form:

$$\langle \cdot, \cdot \rangle_A : \mathcal{H} \times \mathcal{H} \longrightarrow \mathbb{C}, (x, y) \longmapsto \langle x, y \rangle_A := \langle Ax, y \rangle = \langle A^{1/2}x, A^{1/2}y \rangle.$$

Here $A^{1/2}$ means the square root of A . We denote by $\|\cdot\|_A$ the seminorm induced by $\langle \cdot, \cdot \rangle_A$ which is given by $\|x\|_A = \sqrt{\langle x, x \rangle_A} = \sqrt{\|A^{1/2}x\|}$ for every $x \in \mathcal{H}$. It can be checked that $\|x\|_A = 0$ if and only if $x \in \mathcal{N}(A)$. So, $\|\cdot\|_A$ is a norm on \mathcal{H} if and only if A is one-to-one. Furthermore, one may verify that the semi-Hilbert space $(\mathcal{H}, \|\cdot\|_A)$ is complete if and only if $\mathcal{R}(A)$ is closed in $(\mathcal{H}, \|\cdot\|)$. For a given $T \in \mathbb{B}(\mathcal{H})$, if there exists $c > 0$ such that $\|Tx\|_A \leq c\|x\|_A$ for all $x \in \mathcal{R}(A)$, then it holds:

$$\|T\|_A := \sup_{\substack{x \in \mathcal{R}(A), \\ x \neq 0}} \frac{\|Tx\|_A}{\|x\|_A} = \sup_{\substack{x \in \mathcal{R}(A), \\ \|x\|_A=1}} \|Tx\|_A < \infty.$$

If $A = I$, we get the classical norm of an operator T which will be simple denoted by $\|T\|_A$. From now on, we denote $\mathbb{B}^A(\mathcal{H}) := \{T \in \mathbb{B}(\mathcal{H}) ; \|T\|_A < \infty\}$. It is important to note that $\mathbb{B}^A(\mathcal{H})$ is not generally a subalgebra of $\mathbb{B}(\mathcal{H})$ (see [13]). Further, it is difficult to check that $\|T\|_A = 0$ if and only if $ATA = 0$. Recently, there are many papers that study operators defined on a semi-Hilbert space $(\mathcal{H}, \|\cdot\|_A)$. One may see [5, 6, 7, ?, 8, 14, 18, 19] and their references.

Let $T \in \mathbb{B}(\mathcal{H})$. An operator $S \in \mathbb{B}(\mathcal{H})$ is called an A -adjoint operator of T if $\langle Tx, y \rangle_A = \langle x, Sy \rangle_A$ for all $x, y \in \mathcal{H}$ (see [1]). Clearly, S is an A -adjoint of T if and only if $AS = T^*A$, i.e., S is a solution in $\mathbb{B}(\mathcal{H})$ of the equation $AX = T^*A$. We mention here that this type of operator equations can be studied by using the following famous theorem due to Douglas (for its proof see [11]).

Theorem A. *If $T, U \in \mathbb{B}(\mathcal{H})$, then the following statements are equivalent:*

- (i) $\mathcal{R}(U) \subseteq \mathcal{R}(T)$,
- (ii) $TS = U$ for some $S \in \mathbb{B}(\mathcal{H})$,
- (iii) *There exists $\lambda > 0$ such that $\|U^*x\| \leq \lambda\|T^*x\|$ for all $x \in \mathcal{H}$.*

If one of these conditions holds, then there exists a unique solution of the operator equation $TX = U$, denoted by Q , such that $\mathcal{R}(Q) \subseteq \overline{\mathcal{R}(T^)}$. Such Q is called the reduced solution of $TX = U$.*

Let $\mathbb{B}_{A^{1/2}}(\mathcal{H})$ denote the set of all operators that admit $A^{1/2}$ -adjoints. An application of Theorem A shows that

$$\mathbb{B}_{A^{1/2}}(\mathcal{H}) = \{T \in \mathbb{B}(\mathcal{H}) ; \exists \lambda > 0 \text{ such that } \|Tx\|_A \leq \lambda\|x\|_A, \forall x \in \mathcal{H}\}.$$

If $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$, then T is said A -bounded. It can be observed that if $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$, then $T(\mathcal{N}(A)) \subseteq \mathcal{N}(A)$. Further, the following property $\|TS\|_A \leq \|T\|_A\|S\|_A$ holds for all $T, S \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Also, if $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$, then the authors

of [15] showed that

$$\begin{aligned} \|T\|_A &= \sup \{ \|Tx\|_A; x \in \mathcal{H}, \|x\|_A = 1 \} \\ &= \sup \{ |\langle Tx, y \rangle_A|; x, y \in \mathcal{H}, \|x\|_A = \|y\|_A = 1 \}. \end{aligned}$$

For more details regarding the class of A -bounded operators, we refer the reader to [3, 13, 19] and the references therein. Note that $\mathbb{B}_{A^{1/2}}(\mathcal{H})$ is a subalgebra of $\mathbb{B}(\mathcal{H})$ which is neither closed nor dense in $\mathbb{B}(\mathcal{H})$. Moreover, the following inclusions:

$$\mathbb{B}_{A^{1/2}}(\mathcal{H}) \subseteq \mathbb{B}^A(\mathcal{H}) \subseteq \mathbb{B}(\mathcal{H}) \quad (1.1)$$

hold. Note that in general the inclusions in (1.1) are proper. However, if A is an injective operator, then obviously $\mathbb{B}_{A^{1/2}}(\mathcal{H}) = \mathbb{B}^A(\mathcal{H})$. Further, if A has a closed range in \mathcal{H} , then it can be seen that $\mathbb{B}^A(\mathcal{H}) = \mathbb{B}(\mathcal{H})$. So, the inclusions in (1.1) remain equalities if A is injective and has a closed range. We refer to [1, 2, 3, 13] and the references therein for an account of results related the theory of semi-Hilbert spaces.

The notion of the maximal numerical range induced by a positive operator A has recently been introduced by Baklouti et al. in [5]. More precisely, we have the following definition.

Definition B. Let $T \in \mathbb{B}^A(\mathcal{H})$. The A -maximal numerical range of T , denoted by $W_{\max}^A(T)$, is defined as

$$\begin{aligned} W_{\max}^A(T) &= \{ \lambda \in \mathbb{C}; \exists (x_n) \subseteq \mathcal{H}; \|x_n\|_A = 1, \lim_n \langle Tx_n, x_n \rangle_A = \lambda, \\ &\quad \text{and } \lim_n \|Tx_n\|_A = \|T\|_A \}. \end{aligned}$$

For every $T \in \mathbb{B}(\mathcal{H})$, it was shown in [5] that $W_{\max}^A(T)$ is non-empty, convex and compact subset of \mathbb{C} .

Notice that the notion of the maximal numerical range of an operator $T \in \mathbb{B}(\mathcal{H})$, denoted by $W_{\max}(T)$ (that is when $A = I$; the identity operator), was first introduced by Stampfli in [21], in order to determine the norm of the inner derivation acting on $\mathbb{B}(\mathcal{H})$. Recall that the inner derivation δ_T associated with $T \in \mathbb{B}(\mathcal{H})$ is defined by

$$\delta_T : \mathbb{B}(\mathcal{H}) \longrightarrow \mathbb{B}(\mathcal{H}), \quad X \longmapsto TX - XT.$$

For this, in the same paper [21], the author first established the following.

Theorem C. Let $T \in \mathbb{B}(\mathcal{H})$. Then the following conditions are equivalent:

- (1) $0 \in W_{\max}(T)$,
- (2) $\|T\|^2 + |\lambda|^2 \leq \|T + \lambda\|^2$ for any $\lambda \in \mathbb{C}$,
- (3) $\|T\| \leq \|T + \lambda\|$ for any $\lambda \in \mathbb{C}$.

Here $T + \lambda$ is denoted to be $T + \lambda I$ for any $\lambda \in \mathbb{C}$.

Corollary 1.1. Let $T \in \mathbb{B}(\mathcal{H})$. Then, there is a unique scalar c_T such that

$$\|T - c_T\|^2 + |\lambda|^2 \leq \|(T - c_T) - \lambda\|^2, \quad \text{for all } \lambda \in \mathbb{C}.$$

Moreover, $0 \in W_{\max}(T)$ if and only if $c_T = 0$.

The scalar c_T is called the center of mass of T . Note that c_T is the unique scalar satisfying the following

$$\|T - c_T\| = \inf_{\lambda \in \mathbb{C}} \|T - \lambda\|.$$

The scalar $\|T - c(T)\|$ is denoted by $d_A(T)$ and is called the distance of T to scalars. The author [21] then proved that for any $T \in \mathbb{B}(\mathcal{H})$

$$\|\delta_T\| = 2d(T).$$

Recall that an operator $T \in \mathbb{B}(\mathcal{H})$ is said to be normaloid if $\omega(T) = \|T\|$, where $\omega(T)$ is denoted to be the numerical radius of T which is given by

$$\omega(T) = \sup\{|\lambda|; \lambda \in W(T)\}.$$

Here $W(T)$ is denoted to be the numerical range of T and it is defined by Toeplitz in [22] as

$$W(T) := \{\langle Tx, x \rangle; x \in \mathcal{H} \text{ with } \|x\| = 1\}.$$

Equivalent condition is $r(T) = \|T\|$, see, [17]. Here, $r(T)$ is the spectral radius of T . Recently, Spitkovsky in [20] gave the following characterization of a normaloid operator.

Theorem D. *Let $T \in \mathbb{B}(\mathcal{H})$. Then the following conditions are equivalent:*

- (1) T is a normaloid operator,
- (2) $W_{\max}(T) \cap \partial W(T) \neq \emptyset$.

Here ∂L stands for the boundary of a subset L in the complex plane.

Notions of the numerical range and numerical radius are generalized in [5] as follows.

Definition E. *Let $T \in \mathbb{B}(\mathcal{H})$. The A -numerical range and the A -numerical radius of T are respectively given by*

$$W_A(T) := \{\langle Tx, x \rangle_A; x \in \mathcal{H} \text{ with } \|x\|_A = 1\},$$

and

$$\omega_A(T) := \sup\{|\lambda|; \lambda \in W_A(T)\}.$$

It is important to mention that $\omega_A(T)$ may be equal to $+\infty$ for some $T \in \mathbb{B}(\mathcal{H})$ (see [13]). However, $\omega_A(\cdot)$ defines a seminorm on $\mathbb{B}_{A^{1/2}}(\mathcal{H})$ which is equivalent to $\|T\|_A$. More precisely, for any $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$, we have

$$\frac{1}{2} \|T\|_A \leq \omega_A(T) \leq \|T\|_A, \quad (1.2)$$

see [5].

Recently, the concept of A -normaloid operators is introduced by the third author in [13] as follows.

Definition F. *An operator $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$ is said to be A -normaloid if $r_A(T) = \|T\|_A$, where*

$$r_A(T) = \lim_n \|T^n\|_A^{\frac{1}{n}}.$$

Some characterizations of A -normaloid operators are proved in [13]. In particular, we have the following proposition.

Proposition G ([13]). *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then, the following assertions are equivalent:*

- (1) T is A -normaloid,
- (2) $\|T^n\|_A = \|T\|_A^n$ for all positive integer n ,
- (3) $\omega_A(T) = \|T\|_A$,
- (4) There exists a sequence $(x_n) \subseteq \mathcal{H}$ such that $\|x_n\|_A = 1$.

$$\lim_n \|Tx_n\|_A = \|T\|_A \quad \text{and} \quad \lim_n |\langle Tx_n, x_n \rangle_A| = \omega_A(T).$$

Our aim in this work is to give some new characterizations of A -normaloid operators. Mainly, by considering the operator range $\mathcal{R}(A^{1/2})$ endowed with its canonical Hilbertian structure, which will be denoted by $\mathbf{R}(A^{1/2})$, and then using the connection between A -bounded operators and operators acting on the Hilbert space $\mathbf{R}(A^{1/2})$, we extend Theorem D to the context of semi-Hilbert spaces. Moreover, several new properties concerning the A -maximal numerical range of A -bounded operators are established. One main target of this article is to generalize Theorem C for $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. In addition, we give a sufficient and necessary condition for which the A -center of mass of an operator $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$ belongs to $W_{\max}^A(T)$. Other properties are also studied.

In the sequel, if T is any operator in $\mathbb{B}_{A^{1/2}}(\mathcal{H})$, we define

$$\Gamma_A(T) := \{z \in \mathbb{C}; |z| = \|T\|_A\}.$$

2. MAIN RESULTS

We begin this section with the following theorem which gives another useful characterization of A -normaloid operators. We will denote by \overline{L} the closure of any subset L in the complex plane.

Theorem 2.1. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then,*

- (1) T is A -normaloid,
- (2) $\Gamma_A(T) \cap \overline{W_A(T)} \neq \emptyset$.

Proof. (1) \Rightarrow (2): Assume that T is A -normaloid. Then, by Proposition G we have $\omega_A(T) = \|T\|_A$. So, there exists a sequence $(z_n) \subseteq W_A(T)$ such that $\lim_n |z_n| = \|T\|_A$. By compactness of $\overline{W_A(T)}$ we can, taking a subsequence of (z_n) if needed, assume that (z_n) converges to some $z \in \overline{W_A(T)}$. Therefore, $|z| = \|T\|_A$, so $z \in \Gamma_A(T) \cap \overline{W_A(T)}$.

(2) \Rightarrow (1): Let $z \in \Gamma_A(T) \cap \overline{W_A(T)}$. We have $\omega_A(T) \geq |z| = \|T\|_A$. From Inequalities (1.2), we deduce that $\omega_A(T) = \|T\|_A$. That is, T is A -normaloid. \square

Now, we aim to generalize Theorems C and D for $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. To accomplish this goal, some facts from [3] should be recalled. Let $X = \mathcal{H}/\mathcal{N}(A)$ be the quotient space of \mathcal{H} by $\mathcal{N}(A)$. It can be observed that $\langle \cdot, \cdot \rangle_A$ induces on X the following inner product:

$$[\overline{x}, \overline{y}] = \langle x, y \rangle_A = \langle Ax, y \rangle,$$

for every $\bar{x}, \bar{y} \in X$. We note that $(X, [\cdot, \cdot])$ is not complete unless $\mathcal{R}(A)$ is a closed subspace in \mathcal{H} . However, de Branges et al. proved in [9] (see also [16]) that the completion of X under the inner product $[\cdot, \cdot]$ is isomorphic to the Hilbert space $\mathcal{R}(A^{1/2})$ endowed with the following inner product:

$$(A^{1/2}x, A^{1/2}y) := \langle Px, Py \rangle, \quad \forall x, y \in \mathcal{H},$$

where P stands for the orthogonal projection of \mathcal{H} onto the closure of $\mathcal{R}(A)$. From now on, the Hilbert space $(\mathcal{R}(A^{1/2}), (\cdot, \cdot))$ will be simply denoted by $\mathbf{R}(A^{1/2})$. Further, the symbol $\|\cdot\|_{\mathbf{R}(A^{1/2})}$ represents the norm induced by (\cdot, \cdot) . It is crucial to note that $\mathcal{R}(A)$ is dense in $\mathbf{R}(A^{1/2})$ (see [13]). Since $\mathcal{R}(A) \subseteq \mathcal{R}(A^{1/2})$, then we see that

$$(Ax, Ay) = (A^{1/2}A^{1/2}x, A^{1/2}A^{1/2}y) = \langle PA^{1/2}x, PA^{1/2}y \rangle = \langle x, y \rangle_A, \quad \forall x, y \in \mathcal{H}, \quad (2.1)$$

whence,

$$\|Ax\|_{\mathbf{R}(A^{1/2})} = \|x\|_A, \quad (2.2)$$

for any $x \in \mathcal{H}$. For more information concerning the Hilbert space $\mathbf{R}(A^{1/2})$, the interested reader is referred to [3].

Let us consider now the operator Z_A defined by:

$$Z_A : \mathcal{H} \longrightarrow \mathbf{R}(A^{1/2}), \quad x \longmapsto Z_A x = Ax.$$

Further, the following useful proposition is stated in [3].

Proposition H. *Let $T \in \mathbb{B}(\mathcal{H})$. Then $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$ if and only if there exists a unique $\widehat{T} \in \mathbb{B}(\mathbf{R}(A^{1/2}))$ such that $Z_A T = \widehat{T} Z_A$.*

Before we move on, it is important to state the following lemmas. The proof of the first one can be found in [13].

Lemma 2.1. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then*

- (i) $\|T\|_A = \|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}$.
- (ii) $\omega_A(T) = \omega(\widehat{T})$.

Lemma 2.2. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then*

$$W_{\max}^A(T) = W_{\max}(\widehat{T}),$$

where \widehat{T} is the operator given by Proposition H.

Proof. We have $Z_A T = \widehat{T} Z_A$, that is, $ATx = \widehat{T}Ax$ for all $x \in \mathcal{H}$. Now, let $\lambda \in W_{\max}^A(T)$, then there exists $(x_n) \subseteq \mathcal{H}$ such that $\|x_n\|_A = 1$,

$$\lim_{n \rightarrow +\infty} \langle Tx_n, x_n \rangle_A = \lambda, \quad \text{and} \quad \lim_{n \rightarrow +\infty} \|Tx_n\|_A = \|T\|_A.$$

Set $y_n = Ax_n \in \mathcal{R}(A^{1/2})$. By using (2.1) together with (2.2), we have $\|y_n\|_{\mathbf{R}(A^{1/2})} = \|x_n\|_A = 1$ and

$$\langle Tx_n, x_n \rangle_A = \langle ATx_n, Ax_n \rangle = \langle \widehat{T}y_n, y_n \rangle,$$

Again, by (2.2), we infer that

$$\|Tx_n\|_A = \|ATx_n\|_{\mathbf{R}(A^{1/2})} = \|\widehat{T}y_n\|_{\mathbf{R}(A^{1/2})}.$$

On the other hand, by Lemma 2.1 we have $\|T\|_A = \|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}$. This implies that $\lambda \in W_{\max}(\widehat{T})$ and so $W_{\max}^A(T) \subseteq W_{\max}(\widehat{T})$. Conversely, let $\lambda \in W_{\max}(\widehat{T})$, then there exists $(y_n) \subseteq \mathcal{R}(A^{1/2})$ such that $\|y_n\|_{\mathbf{R}(A^{1/2})} = 1$,

$$\lim_{n \rightarrow +\infty} (\widehat{T}y_n, y_n) = \lambda, \text{ and } \lim_{n \rightarrow +\infty} \|\widehat{T}y_n\|_{\mathbf{R}(A^{1/2})} = \|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))} = \|T\|_A.$$

Since $(y_n) \subseteq \mathcal{R}(A^{1/2})$ for all n , then there exists $(x_n) \subseteq \mathcal{H}$ such that $y_n = A^{1/2}x_n$. So, $\|A^{1/2}x_n\|_{\mathbf{R}(A^{1/2})} = 1$,

$$\lim_{n \rightarrow +\infty} (\widehat{T}A^{1/2}x_n, A^{1/2}x_n) = \lambda \text{ and } \lim_{n \rightarrow +\infty} \|\widehat{T}A^{1/2}x_n\|_{\mathbf{R}(A^{1/2})} = \|T\|_A. \quad (2.3)$$

On the other hand, since $\mathcal{R}(A)$ is dense in $\mathbf{R}(A^{1/2})$, then for any $n \in \mathbb{N}$, there exists $(x_{n,k}) \subseteq \mathcal{H}$ such that

$$\lim_{k \rightarrow +\infty} \|Ax_{n,k} - A^{1/2}x_n\|_{\mathbf{R}(A^{1/2})} = 0.$$

This gives

$$\lim_{k \rightarrow +\infty} \|Ax_{n,k}\|_{\mathbf{R}(A^{1/2})} = 1. \quad (2.4)$$

Moreover, by (2.3) we have

$$\lim_{n,k \rightarrow +\infty} (\widehat{T}Ax_{n,k}, Ax_{n,k}) = \lambda \text{ and } \lim_{n,k \rightarrow +\infty} \|\widehat{T}Ax_{n,k}\|_{\mathbf{R}(A^{1/2})} = \|T\|_A.$$

Let $z_k = \frac{x_{n,k}}{\|Ax_{n,k}\|_{\mathbf{R}(A^{1/2})}}$. So, by using (2.4), we obtain

$$\lim_{k \rightarrow +\infty} (\widehat{T}Az_k, Az_k) = \lambda \text{ and } \lim_{k \rightarrow +\infty} \|\widehat{T}Az_k\|_{\mathbf{R}(A^{1/2})} = \|T\|_A.$$

On the other hand, we have

$$(\widehat{T}Az_k, Az_k) = (ATz_k, Az_k) \text{ and } \|\widehat{T}Az_k\|_{\mathbf{R}(A^{1/2})} = \|ATz_k\|_{\mathbf{R}(A^{1/2})}.$$

So, by applying (2.1) together with (2.2), we infer that

$$\lim_{k \rightarrow +\infty} \langle Tz_k, z_k \rangle_A = \lambda \text{ and } \lim_{k \rightarrow +\infty} \|Tz_k\|_A = \|T\|_A.$$

Furthermore, $\|Az_k\|_{\mathbf{R}(A^{1/2})} = \|z_k\|_A = 1$. So, we deduce that $\lambda \in W_{\max}^A(T)$. Hence, the proof is complete. \square

Now, we are in a position to prove the following three theorems. The first one has been proved in [5], however we can obtain the same result as an immediate consequence of Lemma 2.2 and [21, Lemma 2].

Theorem 2.2. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then $W_{\max}^A(T)$ is convex.*

Theorem 2.3. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then the following conditions are equivalent:*

- (1) $0 \in W_{\max}^A(T)$.
- (2) $\|T\|_A^2 + |\lambda|^2 \leq \|T + \lambda\|_A^2$ for any $\lambda \in \mathbb{C}$.
- (3) $\|T\|_A \leq \|T + \lambda\|_A$ for any $\lambda \in \mathbb{C}$.

Proof. Note first that by using Theorem C, we obtain the equivalence between the following assertions:

- (i) $0 \in W_{\max}(\widehat{T})$.

- (ii) $\|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}^2 + |\lambda|^2 \leq \|\widehat{T} + \lambda I_{\mathbf{R}(A^{1/2})}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}^2$ for any $\lambda \in \mathbb{C}$.
- (iii) $\|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))} \leq \|\widehat{T} + \lambda I_{\mathbf{R}(A^{1/2})}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}$ for any $\lambda \in \mathbb{C}$.

On the other hand, by Lemma 2.2, we have $W_{\max}^A(T) = W_{\max}(\widehat{T})$. Moreover, by Lemma 2.1, we have $\|T\|_A = \|\widehat{T}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}$. Also, notice that $T + \lambda \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$ for any $\lambda \in \mathbb{C}$ since $\mathbb{B}_{A^{1/2}}(\mathcal{H})$ is a subalgebra of $\mathbb{B}(\mathcal{H})$. Then, from Proposition H, for any $\lambda \in \mathbb{C}$ there exists a unique $\widehat{T + \lambda} \in \mathbb{B}(\mathbf{R}(A^{1/2}))$ such that $Z_A(T + \lambda) = \widehat{T + \lambda} Z_A$. So, all what remains to prove is that $\|T + \lambda\|_A = \|\widehat{T} + \lambda I_{\mathbf{R}(A^{1/2})}\|_{\mathbb{B}(\mathbf{R}(A^{1/2}))}$ for any $\lambda \in \mathbb{C}$. But the above equality follows by applying Lemma 2.1 (i) together with the fact that $\widehat{T + \lambda} = \widehat{T} + \lambda I_{\mathbf{R}(A^{1/2})}$ (see [16]). \square

Now, we state the third theorem which generalizes Theorem D for A -bounded operators. We need the following lemma.

Lemma 2.3. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then,*

$$\Gamma_A(T) \cap W_{\max}^A(T) = \Gamma_A(T) \cap \overline{W_A(T)}.$$

Proof. Since $W_{\max}^A(T) \subseteq \overline{W_A(T)}$ then the first inclusion holds. Now, let $\lambda \in \Gamma_A(T) \cap \overline{W_A(T)}$. Then, $\lambda = \|T\|_A$ and there exists a sequence $(\lambda_n) \subseteq W_A(T)$ such that $\lambda = \lim_n \lambda_n$. So, there is a sequence $(x_n) \subseteq \mathcal{H}$ such that $\|x_n\|_A = 1$ and $\lambda_n = \langle Tx_n, x_n \rangle_A$ for all n . By applying the Cauchy-Schwarz inequality we get

$$\begin{aligned} |\langle Tx_n, x_n \rangle_A| &= |\langle A^{1/2}Tx_n, A^{1/2}x_n \rangle| \\ &\leq \|Tx_n\|_A \|x_n\|_A \\ &= \|Tx_n\|_A \\ &\leq \|T\|_A. \end{aligned}$$

So, $\lim_n \|Tx_n\|_A = \|T\|_A$. Hence, $\lambda \in \Gamma_A(T) \cap W_{\max}^A(T)$. \square

Now, we are able to prove one of our main results of this article. We will denote by $\overset{\circ}{L}$ the interior of any subset L in the complex plane.

Theorem 2.4. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then, the following statements are equivalent*

- (1) T is an A -normaloid operator,
- (2) $W_{\max}^A(T) \cap \partial W_A(T) \neq \emptyset$.

Proof. (1) \Rightarrow (2): Assume that T is an A -normaloid operator. Then, by applying Theorem 2.1 together with Lemma 2.3, we get

$$\Gamma_A(T) \cap \overline{W_A(T)} = \Gamma_A(T) \cap W_{\max}^A(T) \neq \emptyset.$$

So, there exist $z \in \Gamma_A(T) \cap \overline{W_A(T)}$. Thus, z must lie on the boundary of $W_A(T)$. Since z is also in $W_{\max}^A(T)$, then $W_{\max}^A(T) \cap \partial W_A(T) \neq \emptyset$ as required.

(2) \Rightarrow (1): Assume that $W_{\max}^A(T) \cap \partial W_A(T) \neq \emptyset$. Notice that in view of Lemma 2.1 we have T is A -normaloid if and only if \widehat{T} is a normaloid operator on the Hilbert space $\mathbf{R}(A^{1/2})$. So, in order to prove (1), it suffices to show that

$$W_{\max}(\widehat{T}) \cap \partial W(\widehat{T}) \neq \emptyset.$$

It was shown in [13] that $\overline{W(\widehat{T})} = \overline{W_A(T)}$. Hence $\overline{\partial W(\widehat{T})} = \overline{\partial W_A(T)}$. It is well known that if C is a convex subset in the complex plane, then $\overset{\circ}{C} = \overset{\circ}{\overline{C}}$. Thus $\partial C = \overline{C} \setminus \overset{\circ}{C} = \overline{C} \setminus \overset{\circ}{\overline{C}} = \overline{\partial C}$. Therefore, since both of $W(\widehat{T})$ and $W_A(T)$ are convex, the equality $\overline{\partial W(\widehat{T})} = \overline{\partial W_A(T)}$ implies $\partial W(\widehat{T}) = \partial W_A(T)$. Moreover, $W_{\max}^A(T) = W_{\max}(\widehat{T})$ by Lemma 2.2. We deduce that $W_{\max}(\widehat{T}) \cap \partial W(\widehat{T}) \neq \emptyset$. This completes the proof. \square

Remark 2.1. In [10], the authors gave the following characterization in terms of the numerical radius of a normaloid operator. An operator $T \in \mathbb{B}(\mathcal{H})$ is normaloid if and only if $\omega(T) = \omega_{\max}(T)$. Here, $\omega_{\max}(T)$ is the maximal numerical radius defined by

$$\omega_{\max}(T) := \sup\{|\lambda|; \lambda \in W_{\max}(T)\}.$$

Therefore, by Lemmas 2.1 and 2.2, we can easily obtain the following analogous characterization of A -normaloid operators as follows. Notice that this characterization has been also proved by the third author in [13]. However, our approach here is different from that used in [13].

Theorem 2.5. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then, the following statements are equivalent*

- (1) T is an A -normaloid operator,
- (2) $\omega_A(T) = \omega_{\max}^A(T)$,

where $\omega_{\max}^A(T)$ is the A -maximal numerical radius defined by

$$\omega_{\max}^A(T) := \sup\{|\lambda|; \lambda \in W_{\max}^A(T)\}.$$

On the other hand, by a similar argument as in the proof of Theorem 2.3 and using Corollary 1.1, we obtain the following corollary.

Corollary 2.1. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then, there is a unique scalar $c_A(T)$ such that*

$$\|T - c_A(T)\|_A^2 + |\lambda|^2 \leq \|(T - c_A(T)) - \lambda\|_A^2, \text{ for all } \lambda \in \mathbb{C}. \quad (2.5)$$

Moreover, $0 \in W_{\max}^A(T)$ if and only if $c_A(T) = 0$.

Note that $c_A(T) = c_{\widehat{T}}$; center of mass of \widehat{T} . We call $c_A(T)$ the A -center of mass of T and we denote $d_A(T) = \|T - c_A(T)\|_A$ that we call the A -distance of T to scalars. Clearly, $c_A(T)$ is the unique scalar satisfying

$$d_A(T) = \inf_{\lambda \in \mathbb{C}} \|T - \lambda\|_A.$$

In the following, we give a formula for $d_A(T)$, where $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$.

Theorem 2.6. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then,*

$$d_A^2(T) = \sup_{\|x\|_A=1} \{ \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2 \}.$$

Proof. For any $x \in \mathcal{H}$ with $\|x\|_A = 1$, we have

$$\begin{aligned} d_A^2(T) &= \|T - c_A(T)\|_A^2 \geq \|(T - c_A(T))x\|_A^2 \\ &= \|Tx\|_A^2 + |c_A(T)|^2 - 2\operatorname{Re}(\overline{c_A(T)}\langle Tx, x \rangle_A) \\ &\geq \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2 + |c_A(T) - \langle Tx, x \rangle_A|^2 \\ &\geq \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2. \end{aligned}$$

Whence,

$$d_A^2(T) \geq \sup_{\|x\|_A=1} \{ \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2 \}.$$

Conversely,

$$\|T - c_A(T)\|_A = \inf_{\lambda \in \mathbb{C}} \|T - \lambda\|_A = \inf_{\lambda \in \mathbb{C}} \|(T - c_A(T)) - \lambda\|_A.$$

Then, $\|T - c_A(T)\|_A \leq \|(T - c_A(T)) - \lambda\|_A$ for any $\lambda \in \mathbb{C}$. Since $T - c_A(T) \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$, from Theorem 2.3 we get $0 \in W_{\max}^A(T - c_A(T))$. So, there exists a sequence $(x_n) \subseteq \mathcal{H}$ with $\|x_n\|_A = 1$ such that

$$\lim_n \langle (T - c_A(T))x_n, x_n \rangle_A = 0 \quad \text{and} \quad \lim_n \|(T - c_A(T))x_n\|_A = \|T - c_A(T)\|_A.$$

Then, $\lim_n \langle Tx_n, x_n \rangle_A = c_A(T)$ and

$$\begin{aligned} \|T - c_A(T)\|_A^2 &= \lim_n \|(T - c_A(T))x_n\|_A^2 \\ &= \lim_n \{ \|Tx_n\|_A^2 - |\langle Tx_n, x_n \rangle_A|^2 + |c_A(T) - \langle Tx_n, x_n \rangle_A|^2 \} \\ &= \lim_n \{ \|Tx_n\|_A^2 - |\langle Tx_n, x_n \rangle_A|^2 \} \\ &\leq \sup_{\|x\|_A=1} \{ \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2 \}. \end{aligned}$$

Consequently,

$$d_A^2(T) = \sup_{\|x\|_A=1} \{ \|Tx\|_A^2 - |\langle Tx, x \rangle_A|^2 \}.$$

The proof is complete. \square

Remark 2.2. Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. There is a sequence $(x_n) \subseteq \mathcal{H}$ with $\|x_n\|_A = 1$ such that $\lim_n \langle Tx_n, x_n \rangle_A = c_A(T)$. We derive that $c_A(T) \in \overline{W_A(T)}$. However, $c_A(T)$ need not be contained in $W_{\max}^A(T)$. Indeed, the following corollary gives sufficient and necessary conditions to have $c_A(T) \in W_{\max}^A(T)$.

Corollary 2.2 (Pythagorean Relation). *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. Then, the following statements are equivalent:*

- (1) $c_A(T) \in W_{\max}^A(T)$,
- (2) $d_A^2(T) + |c_A(T)|^2 = \|T\|_A^2$.

Proof. (1) \Rightarrow (2): Assume that $c_A(T) \in W_{\max}^A(T)$. There is a sequence $(x_n) \subseteq \mathcal{H}$ with $\|x_n\|_A = 1$ such that

$$\lim_n \langle Tx_n, x_n \rangle_A = c_A(T) \quad \text{and} \quad \lim_n \|Tx_n\|_A = \|T\|_A.$$

As above, we have

$$\begin{aligned} \|T - c_A(T)\|_A^2 &\geq \lim_n \|(T - c_A(T))x_n\|_A^2 \\ &= \lim_n \{ \|Tx_n\|_A^2 - |\langle Tx_n, x_n \rangle_A|^2 \} \\ &= \|T\|_A^2 - |c_A(T)|^2. \end{aligned}$$

Hence,

$$\|T - c_A(T)\|_A^2 + |c_A(T)|^2 \geq \|T\|_A^2.$$

Taking $\lambda = -c_A(T)$ in Inequality (2.5), we obtain

$$\|T - c_A(T)\|_A^2 + |c_A(T)|^2 \leq \|T\|_A^2. \quad (2.6)$$

Hence,

$$\|T - c_A(T)\|_A^2 + |c_A(T)|^2 = \|T\|_A^2.$$

(2) \Rightarrow (1): Assume that $d_A^2(T) + |c_A(T)|^2 = \|T\|_A^2$. From the proof of Theorem 2.6, there is a sequence $(x_n) \subseteq \mathcal{H}$ with $\|x_n\|_A = 1$ such that $\lim_n \langle Tx_n, x_n \rangle_A = c_A(T)$ and

$$\begin{aligned} d_A^2(T) &= \|T - c_A(T)\|_A^2 = \lim_n \{ \|Tx_n\|_A^2 - |\langle Tx_n, x_n \rangle_A|^2 \} \\ &= \lim_n \|Tx_n\|_A^2 - |c_A(T)|^2. \end{aligned}$$

Remembering the hypothesis, we infer that $\lim_n \|Tx_n\|_A = \|T\|_A$. Consequently, $c_A(T) \in W_{\max}^A(T)$. \square

Remark 2.3. Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. From Remark 2.2, $c_A(T) \in \overline{W_A(T)}$. So, $|c_A(T)| \leq \omega_A(T)$. We know that $W_{\max}^A(T) \subseteq \overline{W_A(T)}$, the following question arises: what about $|c_A(T)|$ and $\omega_{\max}^A(T)$?

Define

$$m_{\max}^A(T) := \inf\{|\lambda|; \lambda \in W_{\max}^A(T)\},$$

for any $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$. The following answers this question.

Theorem 2.7. *Let $T \in \mathbb{B}_{A^{1/2}}(\mathcal{H})$ Then,*

$$|c_A(T)| \leq m_{\max}^A(T).$$

In particular,

$$|c_A(T)| \leq \omega_{\max}^A(T).$$

Proof. By an argument of compactness, there exists $\alpha \in W_{\max}^A(T)$ such that $|\alpha| = m_{\max}^A(T)$. Hence, there is a sequence $(x_n) \subseteq \mathcal{H}$ with $\|x_n\|_A = 1$ satisfying

$$\alpha = \lim_n \langle Tx_n, x_n \rangle_A \quad \text{and} \quad \lim_n \|Tx_n\|_A = \|T\|_A.$$

Therefore, we have

$$\begin{aligned} \|T - c_A(T)\|_A^2 &\geq \|(T - c_A(T))x_n\|_A^2 \\ &= \|Tx_n\|_A^2 + |c_A|^2 - 2\operatorname{Re}(\overline{c_A(T)}\langle Tx_n, x_n \rangle_A) \\ &\geq \|Tx_n\|_A^2 + |c_A(T)|^2 - 2|c_A| |\langle Tx_n, x_n \rangle_A|. \end{aligned}$$

It results that

$$\begin{aligned} \|T - c_A(T)\|_A^2 &\geq \|T\|_A^2 + |c_A(T)|^2 - 2|c_A(T)|m_{\max}^A(T) \\ &= \|T\|_A^2 - (m_{\max}^A(T))^2 + (m_{\max}^A(T) - |c_A(T)|)^2. \end{aligned} \quad (2.7)$$

Thus,

$$\|T - c_A(T)\|_A^2 + (m_{\max}^A(T))^2 \geq \|T\|_A^2 + (m_{\max}^A(T) - |c_A(T)|)^2.$$

We see that

$$\|T - c_A(T)\|_A^2 + (m_{\max}^A(T))^2 \geq \|T\|_A^2 \quad (2.8)$$

and from Inequality (2.6), we get $m_{\max}^A(T) \geq |c_A(T)|$. \square

Remark 2.4. In [12], it is proved that

$$\|T\|^2 \leq d^2(T) + \omega^2(T) \quad (2.9)$$

for any $T \in \mathbb{B}(\mathcal{H})$. From Inequality (2.8), we have

$$\|T\|_A^2 \leq d_A^2(T)^2 + (m_{\max}^A(T))^2 \leq d_A^2(T) + \omega_A^2(T). \quad (2.10)$$

Note that, taking $A = I$, Inequality (2.10) is a refinement of Inequality (2.9). On the other hand, from Inequality (2.7), we have

$$\|T\|_A^2 + |c_A(T)|^2 \leq d_A^2(T) + 2|c_A(T)|m_{\max}^A(T).$$

Then

$$2\|T\|_A|c_A(T)| \leq d_A^2(T) + 2|c_A(T)|m_{\max}^A(T).$$

Consequently, if $c_A(T) \neq 0$ (i.e., $0 \notin W_{\max}^A(T)$), then

$$\|T\|_A \leq m_{\max}^A(T) + \frac{1}{2} \frac{d_A^2(T)}{|c_A(T)|}.$$

Therefore, if $c_A(T) \neq 0$, we get

$$\|T\|_A \leq \inf \left\{ (d_A^2(T)^2 + (m_{\max}^A(T))^2)^{1/2}, m_{\max}^A(T) + \frac{1}{2} \frac{d_A^2(T)}{|c_A(T)|} \right\}.$$

Note that if $c_A(T) = 0$, then $\|T\|_A = d_A(T)$.

Conflict of interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data availability: Data sharing not applicable to the present paper as no data sets were generated or analyzed during the current study.

REFERENCES

- [1] M. L. Arias, G. Corach, M. C. Gonzalez, Partial isometries in semi-Hilbertian spaces, *Linear Algebra Appl.* 428 (7) (2008), 1460-1475.
- [2] M. L. Arias, G. Corach, M. C. Gonzalez, Metric properties of projections in semi-Hilbertian spaces, *Integral Equations and Operator Theory*, 62 (2008), pp.11-28.
- [3] M. L. Arias, G. Corach, M. C. Gonzalez, Lifting properties in operator ranges, *Acta Sci. Math. (Szeged)* 75:3-4 (2009), 635-653.
- [4] N. Altwaijry, K. Feki, N. Minculete, Further inequalities for the weighted numerical radius of operators, *Mathematics* (2022) to appear.
- [5] H. Baklouti, K. Feki, O. A. M. Sid Ahmed, Joint numerical ranges of operators in semi-Hilbertian spaces, *Linear Algebra Appl.* 555 (2018), 266-284.

- [6] H. Baklouti, S. Namouri, Closed operators in semi-Hilbertian spaces, *Linear Multilinear Algebra* (2021) <https://doi.org/10.1080/03081087.2021.1932709>.
- [7] H. Baklouti, S. Namouri, Spectral analysis of bounded operators on semi-Hilbertian spaces, *Banach J. Math. Anal.* **16**, 12 (2022).
- [8] P. Bhunia, S. S. Dragomir, M. S. Moslehian and K. Paul, *Lectures on numerical radius inequalities*, Infosys Science Foundation Series in Mathematical Sciences. Springer, 2022.
- [9] L. de Branges and J. Rovnyak, *Square Summable Power Series*, Holt, Rinehart and Winston, New York, 1966.
- [10] J-T. Chan, K. Chan, An observation about normaloid operators, operators and matrices, Volume 11, Number 3 (2017), 885-890.
- [11] R. G. Douglas, On majorization, factorization and range inclusion of operators in Hilbert space, *Proc. Amer. Math. Soc.* **17** (1966), 413–416.
- [12] S. S. Dragomir, Inequalities for the norm and the numerical radius of linear operators in Hilbert spaces, *Demonstratio Math.* **40**(2) (2007), 411–417.
- [13] K. Feki, Spectral radius of semi-Hilbertian space operators and its applications, *Ann. Funct. Anal.* **11** (2020), 929-946. <https://doi.org/10.1007/s43034-020-00064-y>.
- [14] K. Feki, Some A -spectral radius inequalities for A -bounded Hilbert space operators, *Banach J. Math. Anal.* **16**, 31 (2022)
- [15] M. Faghieh-Ahmadi, F. Gorjizadeh, A -numerical radius of A -normal operators in semi-Hilbertian spaces, *Ital. J. Pure Appl. Math.* **36**, 73–78 (2016)
- [16] K. Feki, On tuples of commuting operators in positive semidefinite inner product spaces, *Linear Algebra Appl.* **603**, 313-328 (2020)
- [17] K. E. Gustafson, D. K. M. Rao, *Numerical range: The Field of Values of Linear Operators and Matrices*, New York, NY, USA, (1997).
- [18] F. Kittaneh and A. Zamani, *Bounds for A -numerical radius based on an extension of A -Buzano inequality*, *Journal of Computational and Applied Mathematics*, 2023. <https://doi.org/10.1016/j.cam.2023.115070>
- [19] W. Majdak , N. A. Secelean, L. Suciu, Ergodic properties of operators in some semi-Hilbertian spaces, *Linear and Multilinear Algebra*, **61**:2 (2013), 139–159.
- [20] I. Spitkovsky, A note on the maximal numerical range, operators and matrices, Volume 13, Number 3 (2019), 601–605
- [21] J. G. Stampfli, The norm of derivation, *Pacific J. Math.* **33** (1970), 737–747.
- [22] O. Toeplitz, Das algebraische Analogou zu einem satze von fejer, *Math. Zeit.* **2** (1918), 187–197.