

ON TORUS QUOTIENTS OF SCHUBERT VARIETIES IN ORTHOGONAL GRASSMANNIAN-II

ARPITA NAYEK AND PINAKINATH SAHA

ABSTRACT. Let $G = SO(8n + 4, \mathbb{C})$ ($n \geq 1$). Let B be a Borel subgroup of G containing a maximal torus T of G . Let $P(\supset B)$ denote the maximal parabolic subgroup of G corresponding to the simple root α_{4n+2} . In this article, we prove projective normality of the GIT quotients of certain Schubert varieties in the orthogonal Grassmannian G/P with respect to the descent of a suitable T -linearized very ample line bundle.

1. INTRODUCTION

Let $G = SO(8n + 4, \mathbb{C})$. Let B be a Borel subgroup of G containing a maximal torus T of G . Let W be the Weyl group of G with respect to T . Let $P(\supset B)$ denote the maximal parabolic subgroup of G corresponding to the simple root α_{4n+2} . Let ϖ_{4n+2} be the fundamental weight corresponding to the simple root α_{4n+2} . Let W_P be the Weyl group of P . Let W^P denote the set of minimal coset representatives of W/W_P . Let “ \leq ” denote the Bruhat-Chevalley order on W^P . For $w \in W^P$, let $X(w) := \overline{BwP/P}$ denote the Schubert variety in G/P corresponding to w .

To state the main result of our article we introduce some notations.

For $1 \leq i \leq 2n$, let $\tau_{2i-1} := \begin{cases} s_{2i-1}s_{2i} \cdots s_{4n-1}s_{4n}s_{4n+1} & \text{for } i \text{ even} \\ s_{2i-1}s_{2i} \cdots s_{4n-1}s_{4n}s_{4n+2} & \text{for } i \text{ odd} \end{cases}$ where s_i denotes the simple reflection corresponding to the simple root α_i .

Let $\lambda = \varpi_{4n+2}$. Let $v_1 = s_{4n+2} \prod_{i=2n}^1 \tau_{2i-1}$. Then v_1 is a unique minimal element in W^P such that $v_1(\lambda) \leq 0$. Consider the following v_i 's such that $v_1 \leq v_i$ for all $2 \leq i \leq 6$:




Figure 1: Sublattice of the Bruhat lattice W^P

2010 *Mathematics Subject Classification.* 14M15.

Key words and phrases. orthogonal grassmannian, GIT-quotient, line bundle, semistable point.

$v_2 = s_{4n-2}v_1, v_3 = s_{4n}v_1, v_4 = s_{4n-2}s_{4n}v_1, v_5 = s_{4n+1}s_{4n}v_1, v_6 = s_{4n-2}s_{4n+1}s_{4n}v_1$. Since $v_1 \leq v_i$ and $v_1(\lambda) \leq 0$, we have $v_i(\lambda) \leq 0$ for all $2 \leq i \leq 6$. Thus, by [8, Lemma 2.1, p.470] $X(v_i)_{T}^{ss}(\mathcal{L}(2\lambda)) \neq \emptyset$ for all $1 \leq i \leq 6$.

In this article, our main results are the following.

Theorem 1.1. *The GIT quotient $T \backslash \backslash (X(v_6))_{T}^{ss}(\mathcal{L}(4\lambda))$ is projectively normal with respect to the descent of the T -linearized very ample line bundle $\mathcal{L}(4\lambda)$.*

Proposition 1.2. (i) *The GIT quotient $T \backslash \backslash (X(v_1))_{T}^{ss}(\mathcal{L}(4\lambda))$ is a point.*

(ii) *The GIT quotient $T \backslash \backslash (X(v_2))_{T}^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2))$ as a polarized variety.*

(iii) *The GIT quotient $T \backslash \backslash (X(v_3))_{T}^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2))$ as a polarized variety.*

(iv) *The GIT quotient $T \backslash \backslash (X(v_4))_{T}^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(2))$ as a polarized variety.*

(v) *The GIT quotient $T \backslash \backslash (X(v_5))_{T}^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$ as a polarized variety.*

When $G = SO(8n, \mathbb{C}) (n \geq 1)$, analogous statements of Theorem 1.1 and Proposition 1.2 are proved in [15].

The layout of the article is as follows. In Section 2, we introduce some notation and recall some preliminaries. In Section 3, we prove Theorem 1.1 and Proposition 1.2.

2. NOTATION AND PRELIMINARIES

We refer [5], [6], [7], [11] and [13] for preliminaries in algebraic groups, Lie algebras, Standard Monomial Theory and Geometric Invariant Theory.

2.1. Projective normality. A polarized variety (X, \mathcal{L}) , where \mathcal{L} is a very-ample line bundle on a projective variety X is said to be *projectively normal* if its homogeneous coordinate ring $\bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X, \mathcal{L}^{\otimes k})$ is integrally closed and is generated as a \mathbb{C} -algebra by $H^0(X, \mathcal{L})$ (see [4, Chapter II, Exercise 5.14, p.126]).

2.2. Geometric Invariant Theory. Let G, P and T be as in the previous section.

Let \mathcal{L} be a T -linearized very ample line bundle on G/P . We also denote the restriction of the line bundle \mathcal{L} on $X(w)$ by \mathcal{L} .

A point $p \in X(w)$ is said to be a *semi-stable* point with respect to the T -linearized line bundle \mathcal{L} if there is a T -invariant section $s \in H^0(X(w), \mathcal{L}^{\otimes m})$ for some positive integer m such that $s(p) \neq 0$. We denote the set of all semi-stable points of $X(w)$ with respect to \mathcal{L} by $X(w)_{T}^{ss}(\mathcal{L})$.

A point p in $X(w)_{T}^{ss}(\mathcal{L})$ is said to be a *stable* point if T -orbit of p is closed in $X(w)_{T}^{ss}(\mathcal{L})$ and stabilizer of p in T is finite. We denote the set of all stable points of $X(w)$ with respect to \mathcal{L} by $X(w)_{T}^s(\mathcal{L})$.

2.3. Notation and preliminaries for $SO(8n+4, \mathbb{C})$. Let us recall some well-known properties of $SO(8n+4, \mathbb{C})$ (see [11]). Let $V = \mathbb{C}^{8n+4}$ together with a non-degenerate symmetric bilinear form $(-, -)$. Let $\{v_1, \dots, v_{8n+4}\}$ be the standard basis of V . Let $E = \begin{pmatrix} 0 & J \\ J & 0 \end{pmatrix}$ be the matrix

of the form $(-, -)$ with respect to the standard basis $\{v_1, \dots, v_{8n+4}\}$, where J denotes the anti-diagonal matrix of size $4n+2 \times 4n+2$ with all the entries are 1.

Let $H = SL(V)$. Let $\sigma : H \rightarrow H$ be the involution defined by $\sigma(A) = E(A^t)^{-1}E$, where $A \in H$ and A^t denote transpose of A . Let $G = H^\sigma$. Then $G = SO(V)$. Let T_H (resp. B_H) be the maximal torus in H consisting of diagonal matrices (resp. the Borel subgroup in H consisting of the upper triangular matrices). Let $T = T_H^\sigma$ and $B = B_H^\sigma$. Then B is a Borel subgroup of G containing T as a maximal torus of G . Let $N_G(T)$ (resp. $N_H(T_H)$) be the normalizer in G (resp. H) of T (resp. T_H). Then we have $N_G(T) \subseteq N_H(T_H)$. Thus, we obtain a homomorphism

$$W \rightarrow W_H,$$

where W, W_H denote the Weyl groups of G, H with respect to T, T_H respectively.

2.3.1. Weyl group. The Weyl group W_H is identified with S_{8n+4} , the symmetric group on the $8n+4$ letters $\{1, 2, \dots, 8n+4\}$. Further, σ induces an involution $W_H \rightarrow W_H$ given by

$$w = (a_1, \dots, a_{8n+4}) \mapsto \sigma(w) = (c_1, \dots, c_{8n+4}),$$

where $w = (a_1, \dots, a_{8n+4})$ is written in one line notation of the permutation and $c_i = 8n+5 - a_{8n+5-i}$. Thus, we have $W = \{w \in W_H^\sigma : w \text{ is an even permutation in } W_H\}$. In one line notation, we have $W = \{(a_1, \dots, a_{8n+4}) \in S_{8n+4} : a_i = 8n+5 - a_{8n+5-i}, 1 \leq i \leq 8n+4 \text{ and } m_w \text{ is even}\}$, where $m_w = \#\{i \in \{1, \dots, 4n+2\} : a_i > 4n+2\}$. Thus, $w = (a_1, \dots, a_{8n+4}) \in W$ is known once (a_1, \dots, a_{4n+2}) is known.

2.3.2. Root System and some other properties. Let $X(T_H)$ (resp. $X(T)$) be the group of characters of T_H (resp. T). For $1 \leq i \leq 8n+4$, the elements $\epsilon_i \in X(T_H)$ are defined by

$$\epsilon_i(t) = t_i,$$

where $t = \text{diag}(t_1, t_2, \dots, t_{8n+4}) \in T_H$.

Then the set $R_H = \{\epsilon_i - \epsilon_j : i \neq j\}$ can be identified with the set of roots of H with respect to T_H and that $R_H^+ = \{\epsilon_i - \epsilon_j : 1 \leq i < j \leq 8n+4\}$ is the set of positive roots with respect to B_H . For $1 \leq i \leq 8n+3$, we denote $\beta_i = \epsilon_i - \epsilon_{i+1}$. Then $S_H = \{\beta_i : 1 \leq i \leq 8n+3\}$ is the set of simple roots of R_H .

Note that σ induces an involution $\sigma : X(T_H) \rightarrow X(T_H)$, on $X(T_H)$ defined by $\chi \mapsto \sigma(\chi)$, where $\sigma(\chi)(t) = \chi(\sigma(t))$ for $t \in T_H$. Then we have $\sigma(\epsilon_i) = -\epsilon_{8n+5-i}$ for $1 \leq i \leq 8n+4$. Since $T \subset T_H$, there is a surjective map $\varphi : X(T_H) \rightarrow X(T)$, defined by $\varphi(\epsilon_i) = -\varphi(\epsilon_{8n+5-i})$ for $1 \leq i \leq 8n+4$.

Then σ leaves R_H (resp. R_H^+) stable. Let R (resp. R^+) be the set of roots (resp. positive roots) of G with respect to T (resp. B). Then from the explicit nature of the adjoint representation of G on \mathfrak{g} (Lie algebra of G) it follows that $R = \varphi(R_H \setminus R_H^\sigma)$ and $R^+ = \varphi(R_H^+ \setminus R_H^{\sigma})$. In particular, we have R (resp. R^+) can be identified with the orbit space of R_H (resp. R_H^+) under the action of σ minus the fixed point set under σ . We see now that R^+ can be identified with the subset $\{\epsilon_i - \epsilon_j : 1 \leq i < j \leq 4n+2\} \cup \{\epsilon_i + \epsilon_j : 1 \leq i < j \leq 4n+2\}$ of $X(H)$ through φ . For $1 \leq i \leq 4n+1$, we denote $\alpha_i = \epsilon_i - \epsilon_{i+1}$ and $\alpha_{4n+2} = \epsilon_{4n+1} + \epsilon_{4n+2}$. Then $S = \{\alpha_i : 1 \leq i \leq 4n+2\}$ is the set of simple roots of R .

Let Q (resp. P) denote the maximal parabolic subgroup of H (resp. G) corresponding to the simple root β_{4n+2} (resp. α_{4n+2}). Let W_Q (resp. W_P) be the Weyl group of the Q (resp. P). Then the Schubert varieties in H/Q (resp. G/P) are parameterized by the minimal coset

representatives of W_H/W_Q (resp. W/W_P). We denote the set of all minimal coset representatives of W_H/W_Q (resp. W/W_P) by W^Q (resp. W^P).

Let ϖ'_{4n+2} (resp. ϖ_{4n+2}) be the fundamental weight corresponding to the simple root β_{4n+2} (resp. α_{4n+2}) in H (resp. in G). Note that the very ample generator of the Picard group of H/Q (resp. of G/P) is the line bundle $\mathcal{L}(\varpi'_{4n+2})$ (resp. $\mathcal{L}(\varpi_{4n+2})$). Further, the restriction of the very ample line bundle $\mathcal{L}(\varpi'_{4n+2})$ via the inclusion $G/P \subset H/Q$ is the very ample line bundle $\mathcal{L}(2\varpi_{4n+2})$.

Note that the Bruhat-Chevalley order “ \leq ” on W^P is induced by the Bruhat-Chevalley order “ \leq ” on W^Q . Define $I_{4n+2,8n+4} := \{\underline{i} = (i_1, \dots, i_{4n+2}) : 1 \leq i_1 < \dots < i_{4n+2} \leq 8n+4\}$. Note that there is a natural partial order “ \leq ” on $I_{4n+2,8n+4}$ defined by $\underline{i} \leq \underline{j}$ if and only if $i_t \leq j_t$ for all $1 \leq t \leq 4n+2$. Then there is a natural order preserving isomorphism between (W^Q, \leq) and $(I_{4n+2,8n+4}, \leq)$. Let $\underline{i} := (i_1, \dots, i_{4n+2}) \in I_{4n+2,8n+4}$, and $p_{\underline{i}}$ be the associated Plücker co-ordinate on $G_{4n+2,8n+4} (\simeq H/Q)$. Let $f_{\underline{i}}$ denote the restriction of $p_{\underline{i}}$ to the set M_{4n+2} of matrices of size $4n+2 \times 4n+2$ identified with the opposite big cell O_H^- of H/Q . Then $f_{\underline{i}}(Y) = p(\underline{i}(A), \underline{i}(B))$, where $\underline{i}(A), \underline{i}(B)$ are given as follows: Let r be such that $i_r \leq 4n+2, i_{r+1} > 4n+2$. Let $s = 4n+2 - r$. Then $\underline{i}(A)$ is the s -tuple given by $(8n+5 - i_{4n+2}, \dots, 8n+5 - i_{r+1})$, while $\underline{i}(B)$ is the s -tuple given by the complement of (i_1, \dots, i_r) in $(1, \dots, 4n+2)$. Then the pair $(\underline{i}(A), \underline{i}(B))$ is the canonical dual pair associated to \underline{i} .

Note that $\mathcal{L}(\varpi'_{4n+2})$ is the tautological line bundle on $H/Q (\simeq G_{4n+2,8n+4})$, the Grassmannian of $4n+2$ -dimensional subspaces of V . Then $H^0(H/Q, \mathcal{L}(\varpi'_{4n+2}))$ is the dual of the irreducible H -module with highest weight ϖ'_{4n+2} . Let \mathcal{L} be the restriction of $\mathcal{L}(\varpi'_{4n+2})$ to G/P . For $\underline{i} \in I_{4n+2,8n+4}$, let $p'_{\underline{i}}$ be the restriction of $p_{\underline{i}}$ to G/P . Note that $p'_{\underline{i}} \in H^0(G/P, \mathcal{L})$. Let $p(\underline{i}(A), \underline{i}(B))$ be the restriction of $p_{\underline{i}}$ to M_{4n+2} with the above identification. Let $p'(\underline{i}(A), \underline{i}(B))$ be the restriction of $p(\underline{i}(A), \underline{i}(B))$ to the set SkM_{4n+2} of skew-symmetric matrices of size $4n+2 \times 4n+2$ (identified with the opposite big cell O^- of G/P). Let $Y \in SkM_{4n+2}$. Consider $\underline{i} \in I_{4n+2,8n+4}$ such that $\underline{i}(A) = \underline{i}(B)$. Then $p'(\underline{i}(A), \underline{i}(A))(Y)$ is a principal minor of Y . Hence, $p'(\underline{i}(A), \underline{i}(A))(Y)$ is a square denoting $q_{\underline{i}}(Y)$ the corresponding Pfaffian (i.e., $q_{\underline{i}}^2(Y) = p'(\underline{i}(A), \underline{i}(A))(Y)$). Thus we obtain a regular function $q_{\underline{i}} : SkM_{4n+2} \rightarrow \mathbb{C}$. Let $r = \#\underline{i}(A)$. Then we have that $q_{\underline{i}}$ is non zero if only if r is even (since the determinant of a skew-symmetric $r \times r$ matrix is zero, if r is odd). Further, we have $\mathcal{L}^2 = \mathcal{L}(\varpi'_{4n+2})$.

2.4. Descends to a line bundle. Now, we recall the following theorem due to Shrawan Kumar.

Theorem 2.1 ([10], Theorem 3.10, p.764). *Let $G = Spin(8n+4, \mathbb{C})$. Then the line bundle $\mathcal{L}(\lambda)$ descends to a line bundle on the GIT quotient $T \setminus \setminus (G/P)_T^{ss}(\mathcal{L}(\lambda))$ if and only if $\lambda = 4m\varpi_{4n+2}$ for some positive integer m .*

2.5. Young tableau for $Spin(8n+4, \mathbb{C})$. Now, we present here the simplified version of the definition of Young tableau and standard Young tableau for $Spin(8n+4, \mathbb{C})$ associated to the weight $\lambda = 2m\varpi_{4n+2}$, where m is a positive integer (for more general see [12, Appendix p.363-365]).

2.5.1. Young tableau for $\lambda = 2m\varpi_{4n+2}$, where $m \in \mathbb{N}$. Define $p_i = 2m$ for $1 \leq i \leq 4n+2$. Associated to λ , we define a partition $p(\lambda) := (p_1, p_2, \dots, p_{4n+2})$. Then a Young diagram of shape $p(\lambda) = (p_1, \dots, p_{4n+2})$ associated to λ is denoted by Γ consists of p_1 boxes in the first column, p_2 boxes in the second column and so on.

A Young tableau of shape λ is a Young diagram filled with integers between 1 and $8n + 4$.

A Young tableau is said to be standard if the entries along any column is non-decreasing from top to bottom and along any row is strictly increasing from left to right.

A Young tableau Γ of shape $p(\lambda)$ is said to be $Spin(8n + 4, \mathbb{C})$ -standard if Γ is standard and if $r = (i_1, i_2, \dots, i_{4n+2})$ is a row of length $4n + 2$ such that if i_j is an entry of the row, then $8n + 5 - i_j$ is not an entry of this row. The number of integers greater than $4n + 2$ in a row of Γ is even.

Given a $Spin(8n + 4, \mathbb{C})$ -standard Young tableau Γ of shape λ , we associate a section $p_\Gamma \in H^0(G/P, \mathcal{L}^{\otimes m}(2\varpi_{4n+2}))^T$ which is called standard monomial of degree m . It is well known that standard monomials of shape λ forms a basis of $H^0(G/P, \mathcal{L}^{\otimes m}(2\varpi_{4n+2}))^T$

Remark 2.2 ([14] Lemma 3.2, p.5). *Fix an integer $1 \leq t \leq 8n + 4$. Let $c_\Gamma(t)$ denote the number of times t appears in Γ . A monomial $p_\Gamma \in H^0(G/P, \mathcal{L}(\lambda))$ is T -invariant if and only if $c_\Gamma(t) = c_\Gamma(8n + 5 - t)$ for all $1 \leq t \leq 8n + 4$.*

2.6. Properties of Pfaffian. To proceed further we recall some result on Pfaffian of a skew-symmetric matrix from [2] which we will use to determine the straightening law.

The Pfaffian $\text{Pf}(A)$ of a skew-symmetric $4n + 2 \times 4n + 2$ -matrix $A = (a_{ij})$ (i.e., $a_{ii} = 0$ and $a_{ij} = -a_{ji}$ for $i \neq j$) with coefficients in \mathbb{C} is defined to be $\text{Pf}(A)^2 = \det(A)$. One can describe the Pfaffian equivalently as a polynomial

$$\text{Pf}(X_{12}, X_{13}, \dots, X_{14n+2}, X_{23}, \dots, X_{24n+2}, \dots, X_{4n+14n+2}) \in \mathbb{Z}[X_{12}, X_{13}, \dots, X_{4n+14n+2}]$$

such that for each $A = (a_{ij}) \in SkM_{4n+2}$ one has

$$\text{Pf}(a_{12}, a_{13}, \dots, a_{14n+2}, a_{23}, \dots, a_{24n+2}, \dots, a_{4n+14n+2})^2 = \det(A),$$

normalized such that

$$\text{Pf}(a_{12}, a_{13}, \dots, a_{14n+2}, a_{23}, \dots, a_{24n+2}, \dots, a_{4n+14n+2}) := 1 \quad (2.1)$$

$$\text{for } a_{ij} := \begin{cases} 1 & \text{for } j - i = 2n + 1 \\ 0 & \text{otherwise.} \end{cases}$$

To be more explicit, consider as above a skew-symmetric $4n + 2 \times 4n + 2$ -matrix A and for each subset $I \subseteq \{1, 2, \dots, 4n + 2\}$ of cardinality, say, m . Let $A(I)$ denote the skew-symmetric $m \times m$ -matrix one gets from A by eliminating all rows and columns not indexed by indices from I . Moreover, put

$$P(I) := \text{Pf}(A(I)) \quad (2.2)$$

for every $I \subseteq \{1, 2, \dots, 4n + 2\}$, where as usual $\text{Pf}(A(\emptyset)) := 1$. Then the following result holds.

Theorem 2.3 ([2], Theorem 1, p.122). *For any two subsets $I_1, I_2 \subseteq \{1, 2, \dots, 4n + 2\}$ of odd cardinality and elements $i_1, i_2, \dots, i_t \in \{1, 2, \dots, 4n + 2\}$ with $i_1 < i_2 < \dots < i_t$ and $\{i_1, \dots, i_t\} = I_1 \Delta I_2 := (I_1 \setminus I_2) \cup (I_2 \setminus I_1)$ one has*

$$\sum_{\tau=1}^t (-1)^\tau \cdot P(I_1 \Delta \{i_\tau\}) \cdot P(I_2 \Delta \{i_\tau\}) = 0. \quad (2.3)$$

Remark 2.4. *For $\underline{i} = (i_1, i_2, \dots, i_{4n+2}) \in I_{4n+2, 8n+4}$, we have*

$$q_{\underline{i}} = P(\underline{i}(B)) = \begin{array}{|c|c|c|c|} \hline i_1 & i_2 & \cdots & i_{4n+2} \\ \hline \end{array}$$

Remark 2.5. Let $I = \{i_1, i_2, \dots, i_r\} \subset \{1, 2, \dots, 4n+2\}$ be such that $i_1 < i_2 < \dots < i_r$. Let $\{j_1, \dots, j_{4n+2-r}\}$ be the complements of $\{i_1, \dots, i_r\}$ in $\{1, 2, \dots, 4n+2\}$ such that $j_1 < j_2 < \dots < j_{4n+2-r}$. Then we have $P(I) = q_{\underline{i}}$, where $\underline{i} = (j_1, j_2, \dots, j_{4n+2-r}, 8n+5-i_r, \dots, 8n+5-i_1)$.

3. MAIN THEOREM

Recall that by [9, Corollary 1.9, p.85] there exists a unique minimal $v_1 \in W^P$ such that $v_1(\lambda) \leq 0$. An easy computation shows that

$$v_1 = s_{4n+2} \prod_{i=2n}^1 \tau_{2i-1},$$

where $\tau_{2i-1} = s_{2i-1}s_{2i} \cdots s_{4n-1}s_{4n}s_{4n+1}$ for i even and $\tau_{2i-1} = s_{2i-1}s_{2i} \cdots s_{4n-1}s_{4n}s_{4n+2}$ for i odd. In one line notation v_i 's ($1 \leq i \leq 6$) are the following:

- $v_1 = (2, 4, 6, \dots, 4n-4, 4n-2, 4n, 4n+3, 4n+4, 4n+6, 4n+8, \dots, 8n+4)$
- $v_2 = (2, 4, 6, \dots, 4n-4, 4n-1, 4n, 4n+3, 4n+4, 4n+7, 4n+8, \dots, 8n+4)$
- $v_3 = (2, 4, 6, \dots, 4n-4, 4n-2, 4n+1, 4n+3, 4n+5, 4n+6, 4n+8, \dots, 8n+4)$
- $v_4 = (2, 4, 6, \dots, 4n-4, 4n-1, 4n+1, 4n+3, 4n+5, 4n+7, 4n+8, \dots, 8n+4)$
- $v_5 = (2, 4, 6, \dots, 4n-4, 4n-2, 4n+2, 4n+4, 4n+5, 4n+6, 4n+8, \dots, 8n+4)$
- $v_6 = (2, 4, 6, \dots, 4n-4, 4n-1, 4n+2, 4n+4, 4n+5, 4n+7, 4n+8, \dots, 8n+4)$.

Since $v_1 \leq v_i$ and $v_1(\lambda) \leq 0$, we have $v_i(\lambda) \leq 0$ for all $2 \leq i \leq 6$. Thus, by [8, Lemma 2.1, p.470] $X(v_i)_{T^{\text{ss}}}^{\text{ss}}(\mathcal{L}(2\lambda))$ is non-empty for all $1 \leq i \leq 6$.

Let $X = T \setminus \setminus (X(v_6))_{T^{\text{ss}}}^{\text{ss}}(\mathcal{L}(2\lambda))$. Let $R = \bigoplus_{k \in \mathbb{Z}_{\geq 0}} R_k$, where $R_k = H^0(X(v_6), \mathcal{L}^{\otimes k}(2\lambda))^T$. Note that R_k 's are finite dimensional vector spaces, and $X = \text{Proj}(R)$.

Let p_{Γ} be the standard monomial in R_k associated to a $Spin(8n+4, \mathbb{C})$ -standard Young tableau Γ . Thus, Γ is of the shape $(2k, 2k, \dots, 2k)$ ($4n+2$ times). Further, the entries along any row of Γ are strictly increasing from left to right and entries along any column of Γ are non-decreasing from top to bottom (see Section 2.5.1). Since p_{Γ} is T -invariant, by Remark 2.2 we have $c_{\Gamma}(t) = c_{\Gamma}(8n+5-t)$ for all $1 \leq t \leq 8n+4$. Thus, we have

$$c_{\Gamma}(t) = k \tag{3.1}$$

for all $1 \leq t \leq 8n+4$.

Let $\underline{i} \in I_{4n+2, 8n+4}$ be such that both i_t and $8n+4-i_t$ simultaneously does not appear in \underline{i} and the number of integers greater than or equal to $4n+3$ are even. Note that such \underline{i} is uniquely determined by the tuple (i_1, i_2, \dots, i_r) of increasing sequence of integers less than or equal to $4n+2$ appear in \underline{i} , where r is even. We denote the Young tableau of shape $k\lambda$ ($k \geq 1$) with first row \underline{i}_1 , second row \underline{i}_2 , and so on the k -th row \underline{i}_k by $\text{YT}^{k\lambda}_{(i_1(1), i_1(2), \dots, i_1(r_1)) \ (i_2(1), i_2(2), \dots, i_2(r_2)) \ \cdots \ (i_k(1), i_k(2), \dots, i_k(r_k))}$, where the tuple $(i_j(1), i_j(2), \dots, i_j(r_j))$ denotes the increasing sequence of integers less than or equal to $4n+2$ appear in \underline{i}_j .

Consider the following tableaux

$$\begin{aligned} X_1 &= \text{YT}^{2\lambda}_{(1, 3, \dots, 4n-3, 4n-1, 4n+1, 4n+2) \ (2, 4, \dots, 4n-4, 4n-2, 4n)} \\ X_2 &= \text{YT}^{2\lambda}_{(1, 3, \dots, 4n-3, 4n-2, 4n+1, 4n+2) \ (2, 4, \dots, 4n-4, 4n-1, 4n)} \\ X_3 &= \text{YT}^{2\lambda}_{(1, 3, \dots, 4n-3, 4n-1, 4n, 4n+2) \ (2, 4, \dots, 4n-4, 4n-2, 4n+1)} \end{aligned}$$

$$\begin{aligned}
X_4 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2) \ (2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} \\
X_5 &= \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1) \ (2,4,\dots,4n-4,4n-2,4n+2)}^{2\lambda} \\
X_6 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda} \\
Y_1 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2) \ (1,3,\dots,4n-3,4n,4n+1,4n+2) \ (2,4,\dots,4n-4,4n-2,4n) \ (2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda} \\
Y_2 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1) \ (1,3,\dots,4n-3,4n,4n+1,4n+2) \ (2,4,\dots,4n-4,4n-2,4n) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\
Y_3 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2) \ (1,3,\dots,4n-3,4n) \ (2,4,\dots,4n-4,4n-2,4n+1) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\
Y_4 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n) \ (1,3,\dots,4n-3,4n,4n+1,4n+2) \ (2,4,\dots,4n-4,4n-2,4n+1) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\
Z_1 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n) \ (1,3,\dots,4n-3,4n-2,4n+1,4n+2) \ (1,3,\dots,4n-3,4n,4n+1,4n+2) \ (2,4,\dots,4n-4,4n-2,4n)}^{4\lambda} \\
&\quad \text{YT}_{(2,4,\dots,4n-4,4n-1,4n+1) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda} \\
Z_2 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2) \ (1,\dots,4n-3,4n-2,4n+1,4n+2) \ (1,3,\dots,4n-3,4n) \ (2,4,\dots,4n-4,4n-2,4n)}^{4\lambda} \\
&\quad \text{YT}_{(2,4,\dots,4n-4,4n-1,4n+1) \ (2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda}.
\end{aligned}$$

Lemma 3.1. *The homogeneous coordinate ring R of X is generated by X_i, Y_j, Z_1 and Z_2 .*

Proof. Let $p_\Gamma \in R_k$ be a standard monomial. We show that $p_\Gamma = p_{\Gamma_1} p_{\Gamma_2}$, where p_{Γ_1} is in R_1 or R_2 or R_3 .

For $1 \leq i \leq 2k$ and $1 \leq j \leq 4n+2$, let row_i denote the i -th row, $e_{i,j}$ denote the (i,j) -th entry of Γ . Let $n_{t,j}$ denote the number of times t appear in the j -th column of Γ .

Since we are considering $X(v_6)$, we have $\text{row}_{2k} \leq v_6$. Thus, for $1 \leq j \leq 2n-2$ or $2n+4 \leq j \leq 4n+2$, we have $e_{2k,j} \leq 2j$. Therefore, for $1 \leq j \leq 2n-2$ or $2n+5 \leq j \leq 4n+2$, we have

$$e_{i,j} = \begin{cases} 2j-1 & \text{for } 1 \leq i \leq k \\ 2j & \text{for } k+1 \leq i \leq 2k. \end{cases}$$

Further, since $e_{2k,2n-2} = 4n-4$ (resp. $e_{1,2n+5} = 4n+9$) we have $e_{i,2n-1} = 4n-3$ (resp. $e_{i,2n+4} = 4n+8$) for all $1 \leq i \leq k$ (resp. $k+1 \leq i \leq 2k$).

Note that $e_{2k,2n-1} \leq 4n-1$. Thus, $e_{2k,2n-1}$ is either $4n-2$ or $4n-1$. Also, since $e_{2k,2n} \leq 4n+2$, we have $e_{2k,2n}$ is either $4n, 4n+1$ or $4n+2$. Thus, the followings are the possibilities for row_{2k}

- (i) $\text{YT}_{(2,4,\dots,4n-4,4n-2,4n)}^\lambda$
- (ii) $\text{YT}_{(2,4,\dots,4n-4,4n-1,4n)}^\lambda$
- (iii) $\text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1)}^\lambda$
- (iv) $\text{YT}_{(2,4,\dots,4n-4,4n-1,4n+1)}^\lambda$
- (v) $\text{YT}_{(2,4,\dots,4n-4,4n-2,4n+2)}^\lambda$
- (vi) $\text{YT}_{(2,4,\dots,4n-4,4n-1,4n+2)}^\lambda$.

Case (i): Assume that $\text{row}_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n)}^\lambda$.

Then

$$e_{i,j} = \begin{cases} 4n-2 & \text{for } k+1 \leq i \leq 2k \text{ and } j = 2n-1 \\ 4n-1 & \text{for } 1 \leq i \leq k \text{ and } j = 2n \\ 4n & \text{for } k+1 \leq i \leq 2k \text{ and } j = 2n \\ 4n+1 & \text{for } 1 \leq i \leq k \text{ and } j = 2n+1. \end{cases}$$

Therefore, $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n+1,4n+2)}^\lambda$ and hence, row_1, row_{2k} together give a factor X_1 of p_Γ .

Case (ii): Assume that $row_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-1,4n)}^\lambda$.

Then $e_{i,2n} = 4n$ for all $k+1 \leq i \leq 2k$ and $e_{1,2n} = 4n-2$ or $4n-1$. If $e_{1,2n} = 4n-1$, then all $4n-2$ appear in the $(2n-1)$ -th column. Further, since $e_{2k,2n-1} = 4n-1$, $n_{4n-2,2n-1} \leq k-1$, which is a contradiction to Eq. (3.1). Thus, $e_{1,2n} = 4n-2$. Therefore, $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2)}^\lambda$ and hence, row_1, row_{2k} together give a factor X_2 of p_Γ .

Case (iii): Assume that $row_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1)}^\lambda$.

Then $e_{i,2n-1} = 4n-2$ for all $k+1 \leq i \leq 2k$ and $e_{i,2n} = 4n-1$ for all $1 \leq i \leq k$. Since $e_{2k,2n} = 4n+1$, we have $n_{4n,2n} \leq k-1$. Thus, $n_{4n,2n+1} \geq 1$. Thus, we have $e_{1,2n+1} = 4n$. Further, since Γ satisfies the property that number of entries greater than $4n+2$ in each row is even, we have $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$ or $\text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+2)}^\lambda$ (see Section 2.5.1).

Claim: $row_1 \neq \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$.

Assume that $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$. Let k' denote the number of rows in Γ those contain exactly $2n+2$ entries less than or equal to $4n+2$. Then remaining $2k-k'$ number of rows in Γ contains exactly $2n$ entries less than or equal to $4n+2$. Since by Eq. (3.1), $c_\Gamma(t) = k$ for all $1 \leq t \leq 4n+2$, we have $(2n+2)k' + 2n(2k-k') = (4n+2)k$. Thus, we have $k' = k$. Thus, row_i ($1 \leq i \leq k$) contains exactly $2n$ entries less than or equal to $4n+2$ and row_i ($k+1 \leq i \leq 2k$) contains exactly $2n+2$ entries less than or equal to $4n+2$. Further, since $e_{2k,2n} = 4n+1$, all $4n+2$ appear in the $(2n+2)$ -th column and hence, $n_{4n+2,2n+2} \leq k-1$, which is a contradiction to Eq. (3.1).

On the other hand, if $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+2)}^\lambda$, then row_1, row_{2k} together give a factor X_3 of p_Γ .

Case (iv): Assume that $row_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-1,4n+1)}^\lambda$.

Since $e_{2k,2n-1} = 4n-1$, we have $n_{4n-2,2n-1} \leq k-1$. Hence, $n_{4n-2,2n} \geq 1$. Therefore, $e_{1,2n} = 4n-2$. Now we claim that $e_{1,2n+1} \leq 4n$. If $e_{1,2n+1} \geq 4n+1$, then all integers less than or equal to $4n$ appear in the first $2n$ columns. Since $e_{2k,2n} = 4n+1$, we have $\sum_{t=1}^{4n} \sum_{j=1}^{2n} n_{t,j} \leq 4nk-1$, which is a contradiction to Eq. (3.1). Therefore, $e_{1,2n+1} \leq 4n$. Hence, row_1 is one of the following:

- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2)}^\lambda$.

If $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n)}^\lambda$, $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1)}^\lambda$, or $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$, then by using the argument as in the third paragraph (line no. 7–12) of case (iii), we see that row_i ($1 \leq i \leq k$) contains exactly $2n+2$ entries less than or equal to $4n+2$ and row_i ($k+1 \leq i \leq 2k$)

contains exactly $2n$ entries less than or equal to $4n + 2$. Further, since $e_{2k,2n} = 4n + 1$, all $4n + 2$ appear in the $(2n + 2)$ -th column. Since $e_{1,2n+2} = 4n$ or $4n + 1$, we have $n_{4n+2,2n+2} \leq k - 1$, which is a contradiction to Eq. (3.1).

If $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2)}^\lambda$, then for $k+1 \leq i \leq 2k-1$, we have $e_{i,2n-1} = 4n-2$ or $4n-1$. Further, since $e_{1,2n+1} = 4n-1$, we have $n_{4n-2,2n-1} + n_{4n-1,2n-1} + n_{4n-1,2n+1} \geq k+1$. Thus, $n_{4n-2,2n} + n_{4n-1,2n} \leq k-1$. Thus, $e_{k,2n} = 4n$. Hence, $row_k = \text{YT}_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$ or $\text{YT}_{(1,3,\dots,4n-3,4n)}^\lambda$. If $row_k = \text{YT}_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$, then $\sum_{j=2n+1}^{4n+2} \sum_{t=4n+3}^{8n+4} n_{t,j} \leq (4n+2)k-2$, which is a contradiction to Eq. (3.1). If $row_k = \text{YT}_{(1,3,\dots,4n-3,4n)}^\lambda$, then $e_{i,2n+1} = 4n+3$ for all $k \leq i \leq 2k$, which is a contradiction to Eq. (3.1).

If $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2)}^\lambda$, then hence, row_1, row_{2k} together give a factor X_4 of p_Γ .

If $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2)}^\lambda$, then by using the argument as in the third paragraph (line no. 7–12) of case (iii), we see that row_i ($1 \leq i \leq k$) contains exactly $2n+2$ entries less than or equal to $4n+2$ and row_i ($k+1 \leq i \leq 2k$) contains exactly $2n$ entries less than or equal to $4n+2$. Further, since $e_{1,2n+2} = 4n+2$, we have $e_{i,2n+2} = 4n+2$ for all $1 \leq i \leq k$. Since $n_{4n-2,2n-1} + n_{4n-1,2n-1} + n_{4n-1,2n+1} \geq k+1$, we have $n_{4n-2,2n} + n_{4n-1,2n} \leq k-1$. Hence, $e_{k,2n} \geq 4n$. Further, since $e_{2k,2n} = 4n+1$, we have $e_{k,2n} = 4n$. Thus, $row_k = \text{YT}_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$. Since $e_{k,2n+1} = 4n+1$ we have $n_{4n+1,2n} \leq k-1$. Thus, $e_{k+1,2n} = 4n$. Therefore, $row_{k+1} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n)}^\lambda$ and hence, $row_1, row_k, row_{k+1}, row_{2k}$ together give a factor Y_1 of p_Γ .

Case (v): Assume that $row_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+2)}^\lambda$. Then $e_{i,2n-1} = 4n-2$ for all $k+1 \leq i \leq 2k$ and $e_{i,2n} = 4n-1$ for all $1 \leq i \leq k$. Further, since $e_{2k,2n} = 4n+2$, we have $n_{4n,2n} \leq k-1$. Thus, $n_{4n,2n+1} \geq 1$. Hence, $e_{1,2n+1} = 4n$. Therefore, $e_{1,2n+2} = 4n+1$ or $4n+2$.

Claim: $e_{1,2n+2} \neq 4n+2$.

Assume that $e_{1,2n+2} = 4n+2$. By using the argument as in the third paragraph (line no. 7–12) of case (iii), we see that row_i ($1 \leq i \leq k$) contains exactly $2n+2$ entries less than or equal to $4n+2$ and row_i ($k+1 \leq i \leq 2k$) contains exactly $2n$ entries less than or equal to $4n+2$. Further, since $e_{2k,2n} = 4n+2$, we have $n_{4n+2,2n+2} + n_{4n+2,2n} \geq k+1$, which is a contradiction to Eq. (3.1).

On the other hand, if $e_{1,2n+2} = 4n+1$, then $row_1 = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$. Therefore, row_1, row_{2k} together give a factor X_5 of p_Γ .

Case (vi): Assume that $row_{2k} = \text{YT}_{(2,4,\dots,4n-4,4n-1,4n+2)}^\lambda$.

Then $e_{1,2n} = 4n-2$. Since $e_{2k,2n+1} = 4n+4$, we have $e_{1,2n+2} \leq 4n+1$. Indeed, if $e_{1,2n+2} > 4n+2$, then $\sum_{j=2n+1}^{4n+2} \sum_{t=4n+3}^{8n+4} n_{t,j} \geq (4n+2)k+1$, which is a contradiction to Eq. (3.1). If $e_{1,2n+2} = 4n+2$, then by using the argument as in the third paragraph (line no. 7–12) of case (iii), we see that row_i ($1 \leq i \leq k$) contains exactly $2n+2$ entries less than or equal to $4n+2$ and row_i ($k+1 \leq i \leq 2k$) contains exactly $2n$ entries less than or equal to $4n+2$. Thus, $e_{i,2n+2} = 4n+2$, for all $1 \leq i \leq k$. Further, since $e_{2k,2n} = 4n+2$, we have $n_{4n+2,2n} + n_{4n+2,2n+2} \geq k+1$, which is a contradiction to Eq. (3.1). Therefore, row_1 is one of the following:

- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$

- $YT_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2)}^\lambda$
- $YT_{(1,3,\dots,4n-3,4n-2,4n-1,4n)}^\lambda$

Subcase (i): Assume that $row_1 = YT_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1)}^\lambda$.

Since $n_{4n-1,2n+1} \geq 1$, we have $e_{k+1,2n-1} = 4n - 2$. Further, since $n_{4n-2,2n-1} + n_{4n-1,2n-1} + n_{4n-1,2n+1} \geq k + 1$, we have $n_{4n-2,2n} + n_{4n-1,2n} \leq k - 1$. Thus, $e_{k,2n} \geq 4n$. Also, by using the argument as in the third paragraph (line no. 7–12) of case (iii), we see that row_i ($1 \leq i \leq k$) contains exactly $2n + 2$ entries less than or equal to $4n + 2$ and row_i ($k + 1 \leq i \leq 2k$) contains exactly $2n$ entries less than or equal to $4n + 2$. Therefore, $row_k = YT_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$. Hence, $e_{k+1,2n} \geq 4n$. If $e_{k+1,2n} \geq 4n + 1$, then $\sum_{t=4n+1}^{4n+2} \sum_{j=2n}^{2n+2} n_{t,j} \geq 2k + 1$, which is a contradiction to Eq. (3.1). Hence, $e_{k+1,2n} = 4n$ and therefore, $row_1, row_k, row_{k+1}, row_{2k}$ together give a factor Y_2 of p_Γ .

Subcase (ii): If $row_1 = YT_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$, then row_1, row_{2k} together give a factor X_6 of p_Γ .

Subcase (iii): Assume that $row_1 = YT_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2)}^\lambda$.

Since $e_{1,2n+1} = 4n - 1$, we have $e_{k+1,2n-1} = 4n - 2$. Further, since $n_{4n-2,2n-1} + n_{4n-1,2n-1} + n_{4n-1,2n+1} \geq k + 1$, we have $n_{4n-2,2n} + n_{4n-1,2n} \leq k - 1$. Hence, $e_{k,2n} \geq 4n$. Next we show that $e_{k,2n} = 4n$. If $e_{k,2n} = 4n + 2$, then we have $n_{4n+2,2n} \geq k + 1$, which is a contradiction to Eq. (3.1). If $e_{k,2n} = 4n + 1$, then $row_k = YT_{(1,3,\dots,4n-3,4n+1)}^\lambda$. Thus, $e_{i,2n+2} = 4n + 5$ for all $k \leq i \leq 2k$, which is a contradiction to Eq. (3.1). Thus, $e_{k,2n} = 4n$. Therefore, row_k is either $YT_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$ or $YT_{(1,3,\dots,4n-3,4n)}^\lambda$. If $row_k = YT_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$, then $\sum_{j=2n+1}^{4n+2} \sum_{t=4n+3}^{8n+4} n_{t,j} \leq (4n + 2)k - 2$, which is a contradiction to Eq. (3.1).

Now assume that $row_k = YT_{(1,3,\dots,4n-3,4n)}^\lambda$.

Since $e_{1,2n+4} = 4n + 2$, we have $e_{k+1,2n} = 4n$ or $4n + 1$. If $e_{k+1,2n} = 4n + 1$, then $row_{k+1} = YT_{(2,4,\dots,4n-4,4n-2,4n+1)}^\lambda$. Therefore, $row_1, row_k, row_{k+1}, row_{2k}$ together give a factor Y_3 of p_Γ .

If $e_{k+1,2n} = 4n$, then $row_{k+1} = YT_{(2,4,\dots,4n-4,4n-2,4n)}^\lambda$. Since row_k does not contain $4n + 1$, there exists l for some $k + 2 \leq l \leq 2k - 1$ such that row_l contains $4n + 1$. Hence, row_l is either $YT_{(2,4,\dots,4n-4,4n-2,4n+1)}^\lambda$ or $YT_{(2,4,\dots,4n-4,4n-1,4n+1)}^\lambda$. If $row_l = YT_{(2,4,\dots,4n-4,4n-2,4n+1)}^\lambda$, then $row_1, row_k, row_l, row_{2k}$ together give a factor Y_3 of p_Γ .

Consider $row_l = YT_{(2,4,\dots,4n-4,4n-1,4n+1)}^\lambda$. Then we study $row_{\frac{k}{2}+1}$.

Assume that k is even.

If $e_{\frac{k}{2}+1,2n+1} = 4n - 1$, then $\sum_{t=4n-2}^{4n-1} (n_{t,2n-1} + n_{t,2n} + n_{t,2n+1}) \geq 2k + 2$, which is a contradiction to Eq. (3.1). If $e_{\frac{k}{2}+1,2n+1} \geq 4n + 3$, then $e_{\frac{k}{2}+1,2n+1} = 4n + 3$ and $e_{\frac{k}{2}+1,2n+2} = 4n + 4$. Thus, $\sum_{t=4n+3}^{4n+4} \sum_{j=2n+1}^{2n+2} n_{t,j} \geq 2k + 1$, which is a contradiction to Eq. (3.1). Hence, $row_{\frac{k}{2}+1}$ is in the following set of rows.

- $YT_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$
- $YT_{(1,3,\dots,4n-3,4n-2,4n,4n+2)}^\lambda$

- $\text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+2)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n-1,4n+1,4n+2)}^\lambda$
- $\text{YT}_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$.

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1)}^\lambda$, then $\text{row}_{\frac{k}{2}+1}, \text{row}_{2k}$ give a factor X_6 of p_Γ .

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2)}^\lambda$, then $\text{row}_{\frac{k}{2}+1}, \text{row}_l$ give a factor X_4 of p_Γ .

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2)}^\lambda$, then $\text{row}_i (i = 1, \frac{k}{2} + 1, k, k + 1, l, 2k)$ give a factor Z_2 of p_Γ .

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+1)}^\lambda$ or $\text{YT}_{(1,3,\dots,4n-3,4n-1,4n,4n+2)}^\lambda$, then $n_{4n-2,2n} \leq \frac{k}{2}$. Thus, $n_{4n-2,2n-1} \geq \frac{k}{2}$. Further, since $\text{row}_l = \text{YT}_{(2,4,\dots,4n-4,4n-1,4n+1)}^\lambda$, we have $\text{row}_{\frac{3k}{2}} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n)}^\lambda$. Therefore, for $1 \leq i \leq \frac{3k}{2}$, $e_{i,2n} = 4n - 2, 4n - 1$ or $4n$ and for $1 \leq i \leq \frac{k}{2} + 1$, $e_{i,2n+1} = 4n - 1$ or $4n$. Hence, $\sum_{j=2n-1}^{2n+1} \sum_{t=4n-2}^{4n} n_{t,j} \geq k + \frac{3k}{2} + \frac{k}{2} + 1 = 3k + 1$, which is a contradiction to Eq. (3.1).

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n-1,4n+1,4n+2)}^\lambda$, then $\text{row}_{\frac{k}{2}+1}, \text{row}_{k+1}$ give a factor X_1 of p_Γ .

If $\text{row}_{\frac{k}{2}+1} = \text{YT}_{(1,3,\dots,4n-3,4n,4n+1,4n+2)}^\lambda$, then for $1 \leq i \leq \frac{k}{2}$, $e_{i,2} = 4n - 2$ and $e_{i,3} = 4n - 1$. Thus, $\text{row}_i = \text{YT}_{(1,3,\dots,4n-2,4n)}^\lambda$ for all $k + 1 \leq i \leq \frac{3k}{2}$. Hence, $e_{i,2} = 4n$ for all $\frac{k}{2} + 1 \leq i \leq \frac{3k}{2}$. Since $e_{1,4} = 4n$, we have $n_{4n,2} + n_{4n,4} \geq k + 1$, which is a contradiction to Eq. (3.1).

Similarly, if $k + 1$ is even, then the proof is similar.

Subcase (iv): If $\text{row}_1 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n)}^\lambda$, then the proof is similar to subcase (iii). □

Lemma 3.2. *The following relations in X_i 's ($1 \leq i \leq 6$), Y_j 's ($1 \leq j \leq 4$) hold in R_2 .*

- (i) $X_2X_3 - X_1X_4 + Y_1 - Y_3 = 0$.
- (ii) $X_4X_5 - X_3X_6 + Y_4 - Y_3 = 0$.
- (iii) $X_2X_5 - X_1X_6 + Y_2 - Y_3 = 0$.

Proof. (i) : Note that

$$X_2X_3 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2), (1,3,\dots,4n-3,4n-1,4n,4n+2), (2,4,\dots,4n-4,4n-1,4n), (2,4,\dots,4n-4,4n-2,4n+1)}^{4\lambda}$$

Now we see that $\text{YT}_{(1,\dots,4n-3,4n-2,4n+1,4n+2), (1,\dots,4n-3,4n-1,4n,4n+2)}^{2\lambda}$ is a nonstandard monomial. We apply Theorem 2.3, Remark 2.4 and Remark 2.5 to express the above non-standard monomial as a linear combination of standard monomials.

Let $\underline{i}_1 = (1, 3, \dots, 4n - 3, 4n - 2, 4n + 1, 4n + 2, 4n + 5, 4n + 6, 4n + 9, \dots, 8n + 3)$ and $\underline{i}_2 = (1, 3, \dots, 4n - 3, 4n - 1, 4n, 4n + 2, 4n + 4, 4n + 7, 4n + 9, \dots, 8n + 3)$. Then $\underline{i}_1(B) = (2, 4, \dots, 4n - 4, 4n - 1, 4n)$ and $\underline{i}_2(B) = (2, 4, \dots, 4n - 4, 4n - 2, 4n + 1)$ are two subsets of $\{1, 2, \dots, 4n + 2\}$.

In order to apply Theorem 2.3 set $I_1 = \{2, 4, \dots, 4n-4, 4n-1\}$ and $I_2 = \{2, 4, \dots, 4n-4, 4n-2, 4n, 4n+1\}$. Note that $I_1 \Delta I_2 = \{4n-2, 4n-1, 4n, 4n+1\}$. Then we have

$$P(I_1 \Delta \{4n-2\}) \cdot P(I_2 \Delta \{4n-2\}) - P(I_1 \Delta \{4n-1\}) \cdot P(I_2 \Delta \{4n-1\}) + P(I_1 \Delta \{4n\}) \cdot P(I_2 \Delta \{4n\}) - P(I_1 \Delta \{4n+1\}) \cdot P(I_2 \Delta \{4n+1\}) = 0.$$

By using Remark 2.5 we have the following:

$$\begin{aligned} P(I_1 \Delta \{4n-2\}) &= q_{\{1,3,\dots,4n-3,4n,4n+1,4n+2,4n+6,4n+7,4n+9,\dots,8n+4\}} \\ P(I_2 \Delta \{4n-2\}) &= q_{\{1,3,\dots,4n-3,4n-2,4n-1,4n+2,4n+4,4n+5,4n+9,\dots,8n+4\}} \\ P(I_1 \Delta \{4n-1\}) &= q_{\{1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2,4n+9,\dots,8n+4\}} \\ P(I_2 \Delta \{4n-1\}) &= q_{\{1,3,\dots,4n-3,4n+2,4n+4,4n+5,4n+6,4n+7,4n+9,\dots,8n+4\}} \\ P(I_1 \Delta \{4n\}) &= q_{i_1} \\ P(I_2 \Delta \{4n\}) &= q_{i_2} \\ P(I_1 \Delta \{4n+1\}) &= q_{\{1,3,\dots,4n-3,4n-2,4n,4n+2,4n+4,4n+6,4n+9,\dots,8n+4\}} \\ P(I_2 \Delta \{4n+1\}) &= q_{\{1,3,\dots,4n-3,4n-1,4n+1,4n+2,4n+5,4n+7,4n+9,\dots,8n+4\}}. \end{aligned}$$

Therefore, we have the following straightening laws in $G/P^{\alpha_{4n+2}}$

$$\begin{aligned} & \text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2),(1,3,\dots,4n-3,4n-1,4n,4n+2)}^{2\lambda} \\ &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n+1,4n+2)}^{2\lambda} \\ & \quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2),(1,3,\dots,4n-3,4n,4n+1,4n+2)}^{2\lambda} \\ & \quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n+2)}^{2\lambda}. \end{aligned}$$

Similarly, by using Theorem 2.3, Remark 2.4 and Remark 2.5 we have

$$\begin{aligned} \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n)}^{2\lambda} &= \text{YT}_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} \\ & \quad - \text{YT}_{(2,4,\dots,4n-4,4n-2,4n-1),(2,4,\dots,4n-4,4n,4n+1)}^{2\lambda} \\ & \quad + \text{YT}_{(2,4,\dots,4n-4,4n-2,4n-1,4n,4n+1),(2,4,\dots,4n-4)}^{2\lambda}. \end{aligned}$$

Since we are working in the Schubert variety $X(v_6)$, above straightening law becomes

$$\text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n)}^{2\lambda} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda}. \quad (3.2)$$

Therefore, by using the above straightening laws we have

$$\begin{aligned} X_2 X_3 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda} \\ & \quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda} \\ & \quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda}. \end{aligned}$$

Since we are working in the Schubert variety $X(v_6)$, by using Theorem 2.3 and Remark 2.5 we have

$$\text{YT}_{(1,3,\dots,4n-3,4n+2),(2,4,\dots,4n-4,4n-2,4n)}^{2\lambda} = \text{YT}_{(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+2)}^{2\lambda}.$$

Therefore,

$$\begin{aligned} X_2 X_3 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda} \\ &\quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda} \\ &\quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda}. \end{aligned}$$

We have the following straightening law in $G/P^{\alpha_{4n+2}}$.

$$\begin{aligned} \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} &= \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda} \\ &\quad - \text{YT}_{(2,4,\dots,4n-4,4n-2,4n-1),(2,4,\dots,4n-4,4n+1,4n+2)}^{2\lambda} \\ &\quad + \text{YT}_{(2,4,\dots,4n-4,4n-2,4n-1,4n+1,4n+2),(2,4,\dots,4n-4)}^{2\lambda}. \end{aligned}$$

Since we are working in the Schubert variety $X(v_6)$, above straightening law becomes

$$\text{YT}_{(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} = \text{YT}_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda}. \quad (3.3)$$

Therefore, we have

$$\begin{aligned} X_2 X_3 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} \\ &\quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+2),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda} \\ &\quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{2\lambda} \\ &= X_1 X_4 - Y_1 + Y_3. \end{aligned}$$

(ii) : Note that

$$X_4 X_5 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n,4n+1),(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n+1)}^{4\lambda}.$$

By using Theorem 2.3 and Remark 2.5, the following straightening law holds in $G/P^{\alpha_{4n+2}}$.

$$\begin{aligned} \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+2),(1,3,\dots,4n-3,4n-1,4n,4n+1)}^{2\lambda} &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1),(1,3,\dots,4n-3,4n-1,4n,4n+2)}^{2\lambda} \\ &\quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n),(1,3,\dots,4n-3,4n,4n+1,4n+2)}^{2\lambda} \\ &\quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n)}^{2\lambda}. \end{aligned} \quad (3.4)$$

Therefore, by using Eq. (3.3) and Eq. (3.4) we have

$$\begin{aligned} X_4 X_5 &= \text{YT}_{(1,3,\dots,4n-3,4n-2,4n,4n+1),(1,3,\dots,4n-3,4n-1,4n,4n+2),(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\ &\quad - \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\ &\quad + \text{YT}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)}^{4\lambda} \\ &= X_3 X_6 - Y_4 + Y_3. \end{aligned}$$

(iii) : Note that

$$X_2 X_5 = \text{YT}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2),(1,3,\dots,4n-3,4n-1,4n,4n+1),(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n)}^{4\lambda}.$$

By using Theorem 2.3 and Remark 2.5, the following straightening law holds in $G/P^{\alpha_{4n+2}}$.

$$\begin{aligned}
& \text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n-2,4n+1,4n+2),(1,3,\dots,4n-3,4n-1,4n,4n+1)} \\
= & \text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n-2,4n,4n+1),(1,3,\dots,4n-3,4n-1,4n+1,4n+2)} \\
& - \text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1),(1,3,\dots,4n-3,4n,4n+1,4n+2)} \\
& + \text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n+1)}
\end{aligned}$$

and

$$\begin{aligned}
\text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n)} &= \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& - \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n-1),(2,4,\dots,4n-4,4n,4n+2)} \\
& + \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n-1,4n,4n+2),(2,4,\dots,4n-4)}
\end{aligned}$$

Since we are working in the Schubert variety $X(v_6)$, the above straightening law becomes

$$\text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n+2),(2,4,\dots,4n-4,4n-1,4n)} = \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)}$$

Therefore, by using the above straightening laws we have

$$\begin{aligned}
X_2 X_5 &= \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n,4n+1),(1,3,\dots,4n-3,4n-1,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& - \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& + \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n+1),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)}
\end{aligned}$$

Since we are working in the Schubert variety $X(v_6)$, by using Theorem 2.3 and Remark 2.5 we have

$$\text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n+1),(2,4,\dots,4n-4,4n-2,4n)} = \text{YT}^{2\lambda}_{(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+1)}$$

Therefore, we have

$$\begin{aligned}
X_2 X_5 &= \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n,4n+1),(1,3,\dots,4n-3,4n-1,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& - \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n+1),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& + \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n+2)} \\
& = X_1 X_6 - Y_2 + Y_3. \quad \square
\end{aligned}$$

Lemma 3.3. *The following relations in X_i 's ($1 \leq i \leq 6$), Y_j 's ($1 \leq j \leq 4$) and Z_1, Z_2 hold in R_3 .*

- (i) $Z_1 - X_2 Y_4 = 0$.
- (ii) $Z_2 - X_2 Y_3 = 0$.

Proof. (i): Note that

$$\begin{aligned}
X_2 Y_4 &= \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n),(1,3,\dots,4n-3,4n-2,4n+1,4n+2),(1,3,\dots,4n-3,4n,4n+1,4n+2),(2,4,\dots,4n-4,4n-2,4n+1)} \\
& \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-1,4n),(2,4,\dots,4n-4,4n-1,4n+2)}
\end{aligned}$$

Recall that $\text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n)} = \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}$ (see Eq. (3.2)). Therefore, we have $X_2 Y_4 = Z_1$.

(ii): Note that

$$\begin{aligned}
X_2 Y_3 &= \text{YT}^{4\lambda}_{(1,3,\dots,4n-3,4n-2,4n-1,4n,4n+1,4n+2),(1,3,\dots,4n-3,4n-2,4n+1,4n+2),(1,3,\dots,4n-3,4n),(2,4,\dots,4n-4,4n-2,4n+1)} \\
& \text{YT}^{2\lambda}_{(2,4,\dots,4n-4,4n-1,4n),(2,4,\dots,4n-4,4n-1,4n+2)}
\end{aligned}$$

Recall that $YT_{(2,4,\dots,4n-4,4n-2,4n+1),(2,4,\dots,4n-4,4n-1,4n)}^{2\lambda} = YT_{(2,4,\dots,4n-4,4n-2,4n),(2,4,\dots,4n-4,4n-1,4n+1)}^{2\lambda}$ (see Eq. (3.2)). Therefore, we have $X_2Y_3 = Z_2$. \square

Lemma 3.4. *R is generated by X_i , ($1 \leq i \leq 6$) and Y_3 as a \mathbb{C} -algebra.*

Proof. Follows from Lemma 3.1, Lemma 3.2, and Lemma 3.3. \square

Recall that by Theorem 2.1, the very ample line bundle $\mathcal{L}(4\lambda)$ descends to a line bundle on the GIT quotient $T \backslash \backslash (G/P^{\alpha_{4n+2}})^{ss}_T(\mathcal{L}(4\lambda))$. We prove that the GIT quotient $T \backslash \backslash (X(v_6))^{ss}_T(\mathcal{L}(4\lambda))$ is projectively normal with respect to the descent of the T -linearized very ample line bundle $\mathcal{L}(4\lambda)$.

Theorem 3.5. *The homogeneous coordinate ring of $T \backslash \backslash (X(v_6))^{ss}_T(\mathcal{L}(4\lambda))$ is generated by elements of degree one.*

Proof. Let $f \in H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T = H^0(X(v_6), \mathcal{L}^{\otimes 2k}(2\lambda))^T$. Then by Lemma 3.4, we have

$$f = \sum a_{\underline{m}, n_1} X^{\underline{m}} Y_3^{n_1},$$

where $\underline{m} = (m_1, m_2, m_3, m_4, m_5, m_6) \in \mathbb{Z}_{\geq 0}^6$ and $n_1 \in \mathbb{Z}_{\geq 0}$ such that $m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + 2n_1 = 2k$, $X^{\underline{m}}$ denotes $X_1^{m_1} X_2^{m_2} X_3^{m_3} X_4^{m_4} X_5^{m_5} X_6^{m_6}$ and $a_{\underline{m}, n_1}$'s are non-zero scalars.

Now to prove that the homogeneous coordinate ring of $T \backslash \backslash (X(v_6))^{ss}_T(\mathcal{L}(4\lambda))$ is generated by $H^0(X(v_6), \mathcal{L}(4\lambda))^T$ as a \mathbb{C} -algebra, it is enough to show that for each f as above and each monomial appearing in the expression of f is a product of k elements of $H^0(X(v_6), \mathcal{L}(4\lambda))^T$.

Consider the monomial $X^{\underline{m}} Y_3^{n_1}$ in the expression of f . Note that $m_1 + m_2 + m_3 + m_4 + m_5 + m_6$ is an even integer. Thus, $X^{\underline{m}}$ can be written as $\prod_{(i,j)} X_i X_j$, where the number of pairs (i, j) is $k - n_1$ and repetitions of X_i 's are allowed. Therefore, $X^{\underline{m}}$ can be written as product of $k - n_1$ number of monomials in $H^0(X(v_6), \mathcal{L}(4\lambda))^T$. On the other hand, since $Y_3 \in H^0(X(v_6), \mathcal{L}(4\lambda))^T$, $X^{\underline{m}} Y_3^{n_1}$ is a product of k elements of $H^0(X(v_6), \mathcal{L}(4\lambda))^T$. \square

Proof of Theorem 1.1. Since $X(v_6)$ is normal, the variety $T \backslash \backslash (X(v_6))^{ss}_T(\mathcal{L}(4\lambda))$ is normal. Therefore, by using Theorem 3.5, it follows that $T \backslash \backslash (X(v_6))^{ss}_T(\mathcal{L}(4\lambda))$ is projectively normal with respect to the descent of the T -linearized very ample line bundle $\mathcal{L}(4\lambda)$. \square

Corollary 3.6. *Let $v \in W^{P^{\alpha_{4n+2}}}$ be such that $v_1 \leq v \leq v_6$. Then the GIT quotient $T \backslash \backslash (X(v))^{ss}_T(\mathcal{L}(4\lambda))$ is projectively normal with respect to the descent of the T -linearized very ample line bundle $\mathcal{L}(4\lambda)$.*

Proof. Since T is linearly reductive, the restriction map

$$\phi : H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T \longrightarrow H^0(X(v), \mathcal{L}^{\otimes k}(4\lambda))^T$$

is surjective for all $k \geq 1$.

So, by Theorem 1.1, $T \backslash \backslash (X(v))^{ss}_T(\mathcal{L}(4\lambda))$ is projectively normal with respect to the descent of the T -linearized very ample line bundle $\mathcal{L}(4\lambda)$. \square

Proof of Proposition 1.2. Note that $Y_3 = 0$ on $X(v_i)$ for all $1 \leq i \leq 5$.

(i): Since T is linearly reductive, the restriction map

$$\phi : H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T \longrightarrow H^0(X(v_1), \mathcal{L}^{\otimes k}(4\lambda))^T$$

is surjective for all $k \geq 1$. Thus, by Lemma 3.4, any standard monomial in $H^0(X(v_1), \mathcal{L}^{\otimes k}(4\lambda))^T$ is of the form X_1^{2k} . Hence, $T \setminus \setminus (X(v_1))_T^{ss}(\mathcal{L}(4\lambda)) = Proj(\mathbb{C}[X_1^2])$.

(ii): Since T is linearly reductive, the restriction map

$$\phi : H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T \longrightarrow H^0(X(v_2), \mathcal{L}^{\otimes k}(4\lambda))^T$$

is surjective for all $k \geq 1$. Thus, by Lemma 3.4, X_1^2, X_1X_2, X_2^2 are standard monomials in $H^0(X(v_2), \mathcal{L}(4\lambda))^T$.

Note that by Corollary 3.6, $T \setminus \setminus (X(v_2))_T^{ss}(\mathcal{L}(4\lambda))$ is projectively normal. Therefore, there is a surjective homomorphism of \mathbb{C} -algebras

$$\phi : \mathbb{C}[z_0, z_1, z_2] \longrightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_2), \mathcal{L}^{\otimes k}(4\lambda))^T,$$

defined by

$$(z_0, z_1, z_2) \mapsto (X_1^2, X_1X_2, X_2^2).$$

Let I be the ideal of $\mathbb{C}[z_0, z_1, z_2]$ generated by the relation

$$z_0z_2 = z_1^2. \tag{3.5}$$

Clearly, $I \subseteq \ker \phi$ and ϕ induces a homomorphism of \mathbb{C} -algebras

$$\tilde{\phi} : \mathbb{C}[z_0, z_1, z_2]/I \longrightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_2), \mathcal{L}^{\otimes k}(4\lambda))^T.$$

Now, we show that $\tilde{\phi}$ is an isomorphism. To complete the proof we use (3.5) as a reduction system. The process is of replacing a monomial M in z_i 's which is divisible by a term L_i 's on the left hand side of the reduction rule $L_i = R_i$ by $(M/L_i)R_i$, where R_i is the right hand side of the reduction rule. We show that the diamond lemma of ring theory holds for this reduction system (see [1]). That is any monomial in z_i 's reduces after applying the reduction rule to a unique expression in z_i 's, in which no term is divisible by a term appearing on the left hand side of the above reduction system.

Since we have only one reduction rule, it is enough to check for $z_0z_1z_2$. Note that $z_0z_1z_2 = (z_0z_2)z_1 = z_1^3$, for which no further reduction is possible. Therefore, $\tilde{\phi}$ is an isomorphism. Hence, the GIT quotient $T \setminus \setminus (X(v_2))_T^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $Proj(\frac{\mathbb{C}[z_0, z_1, z_2]}{(z_1^2 - z_0z_2)})$. Thus, the GIT quotient $T \setminus \setminus (X(v_2))_T^{ss}(\mathcal{L}(\lambda))$ is isomorphic to $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2))$ as a polarized variety.

Proof of (iii) is similar to the proof of (ii).

(iv): Since T is linearly reductive, the restriction map

$$\phi : H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T \longrightarrow H^0(X(v_4), \mathcal{L}^{\otimes k}(4\lambda))^T$$

is surjective for all $k \geq 1$. Thus, by Lemma 3.1, $X_1^2, X_2^2, X_3^2, X_4^2, X_1X_2, X_1X_3, X_1X_4, X_2X_4, X_3X_4, Y_1$ are standard monomials in $H^0(X(v_4), \mathcal{L}(4\lambda))^T$. Further, by Lemma 3.2, $X_2X_3 = X_1X_4 - Y_1$. Note that by Corollary 3.6, $T \setminus \setminus (X(v_4))_T^{ss}(\mathcal{L}(4\lambda))$ is projectively normal. Let $A = \mathbb{C}[z_1, z_2, z_3, z_4, z_5, z_6, z_7, z_8, z_9, z_{10}]$ be the polynomial algebra with variables z_i 's ($1 \leq i \leq 10$). Then there is a surjective homomorphism

$$\phi : A \rightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_4), \mathcal{L}^{\otimes k}(4\lambda))^T,$$

defined by

$$(z_1, z_2, \dots, z_{10}) \mapsto (X_1^2, X_2^2, X_3^2, X_4^2, X_1X_2, X_1X_3, X_1X_4, X_2X_3, X_2X_4, X_3X_4).$$

Let I be the ideal of A generated by the following relations:

| | | | | |
|------------------------|------------------------|------------------------|------------------------|------------------------|
| $z_1 z_2 = z_5^2$ | $z_1 z_3 = z_6^2$ | $z_1 z_4 = z_7^2$ | $z_1 z_8 = z_5 z_6$ | $z_1 z_9 = z_5 z_7$ |
| $z_1 z_{10} = z_6 z_7$ | $z_2 z_3 = z_8^2$ | $z_2 z_4 = z_9^2$ | $z_2 z_6 = z_5 z_8$ | $z_2 z_7 = z_5 z_9$ |
| $z_2 z_{10} = z_8 z_9$ | $z_3 z_4 = z_{10}^2$ | $z_3 z_5 = z_6 z_8$ | $z_3 z_7 = z_6 z_{10}$ | $z_3 z_9 = z_8 z_{10}$ |
| $z_4 z_5 = z_7 z_9$ | $z_4 z_6 = z_7 z_{10}$ | $z_4 z_8 = z_9 z_{10}$ | $z_5 z_{10} = z_6 z_9$ | $z_6 z_9 = z_7 z_8$ |

Clearly, $I \subseteq \ker \phi$ and ϕ induces a homomorphism of \mathbb{C} -algebras

$$\tilde{\phi} : A/I \longrightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_4), \mathcal{L}^{\otimes k}(4\lambda))^T.$$

Now, we show that $\tilde{\phi}$ is an isomorphism.

To complete the proof we use the above relations as a reduction system. We show that diamond lemma (see [1]) holds for this reduction system by looking at the reduction of the minimal overlapping ambiguities.

Note that $z_1 z_2 z_3 = (z_1 z_2) z_3 = z_3 z_5^2$ (using $z_1 z_2 = z_5^2$) $= (z_3 z_5) z_5 = z_5 z_6 z_8$ (using $z_3 z_5 = z_6 z_8$).

Again, $z_1 z_2 z_3 = z_1 (z_2 z_3) = z_1 z_8^2$ (using $z_2 z_3 = z_8^2$) $= (z_1 z_8) z_8 = z_5 z_6 z_8$ (using $z_1 z_8 = z_5 z_6$).

Also, $z_1 z_2 z_3 = (z_1 z_3) z_2 = z_2 z_6^2$ (using $z_1 z_3 = z_6^2$) $= (z_2 z_6) z_6 = z_5 z_6 z_8$ (using $z_2 z_6 = z_5 z_8$).

Therefore, $z_1 z_2 z_3$ reduces to a unique expression.

Likewise, we can show that the remaining overlapping ambiguities reduce to a unique expression in z_i 's after applying different reduction rules. Therefore, $\tilde{\phi}$ is an isomorphism.

Consider the embedding

$$\psi : \mathbb{P}^3 \hookrightarrow \mathbb{P}^9$$

given by

$$(x_1, x_2, x_3, x_4) \mapsto (x_1^2, x_2^2, x_3^2, x_4^2, x_1 x_2, x_1 x_3, x_1 x_4, x_2 x_3, x_2 x_4, x_3 x_4).$$

Let J be the homogeneous ideal of A generated by 2×2 minors of the matrix

$$\begin{pmatrix} z_1 & z_5 & z_6 & z_7 \\ z_5 & z_2 & z_8 & z_9 \\ z_6 & z_8 & z_3 & z_{10} \\ z_7 & z_9 & z_{10} & z_4 \end{pmatrix},$$

where z_i 's are the homogeneous coordinates of \mathbb{P}^9 . Note that $\psi(\mathbb{P}^3)$ is given by J . Further, note that $I \subseteq J$ and $\dim(A/I) = \dim(A/J) = 4$ (computed using Macaulay2 [3]). Hence, the GIT quotient $T \setminus \setminus (X(v_4))_T^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(2))$ as a polarized variety.

(v): Since T is linearly reductive, the restriction map

$$\phi : H^0(X(v_6), \mathcal{L}^{\otimes k}(4\lambda))^T \longrightarrow H^0(X(v_5), \mathcal{L}^{\otimes k}(4\lambda))^T$$

is surjective for all $k \geq 1$. Thus, by Lemma 3.1, $X_1^2, X_3^2, X_5^2, X_1 X_3, X_1 X_5, X_3 X_5$ are standard monomials in $H^0(X(v_5), \mathcal{L}(4\lambda))^T$. Note that by Corollary 3.6, $T \setminus \setminus (X(v_5))_T^{ss}(\mathcal{L}(4\lambda))$ is projectively normal. Let $A = \mathbb{C}[z_1, z_2, z_3, z_4, z_5, z_6]$ be the polynomial algebra with variables z_i 's. Then there is a surjective homomorphism

$$\phi : A \longrightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_5), \mathcal{L}^{\otimes k}(4\lambda))^T,$$

defined by

$$(z_1, z_2, \dots, z_6) \mapsto (X_1^2, X_3^2, X_5^2, X_1 X_3, X_1 X_5, X_3 X_5).$$

Let I be the ideal of A generated by the following relations:

| | | |
|---------------------|---------------------|---------------------|
| $z_1 z_2 = z_4^2$ | $z_1 z_3 = z_5^2$ | $z_2 z_3 = z_6^2$ |
| $z_1 z_6 = z_4 z_5$ | $z_2 z_5 = z_4 z_6$ | $z_3 z_4 = z_5 z_6$ |

Clearly, $I \subseteq \ker \phi$ and ϕ induces a homomorphism

$$\tilde{\phi} : A/I \longrightarrow \bigoplus_{k \in \mathbb{Z}_{\geq 0}} H^0(X(v_5), \mathcal{L}^{\otimes k}(4\lambda))^T.$$

Now, we show that $\tilde{\phi}$ is an isomorphism.

To complete the proof we use the above relations as a reduction system. We show that diamond lemma (see [1]) holds for this reduction system by looking at the reduction of the minimal overlapping ambiguities: $z_1 z_2 z_3, z_1 z_2 z_5, z_1 z_2 z_6, z_1 z_3 z_4, z_1 z_3 z_6, z_2 z_3 z_4, z_2 z_3 z_5$.

Note that $z_1 z_2 z_3 = z_3(z_1 z_2) = z_3 z_4^2$ (using $z_1 z_2 = z_4^2$) $= (z_3 z_4) z_4 = z_4 z_5 z_6$ (using $z_3 z_4 = z_5 z_6$).

Again $z_1 z_2 z_3 = z_1(z_2 z_3) = z_1 z_6^2$ (using $z_2 z_3 = z_6^2$) $= (z_1 z_6) z_6 = z_4 z_5 z_6$ (using $z_1 z_6 = z_4 z_5$).

Also, $z_1 z_2 z_3 = z_2(z_1 z_3) = z_2 z_5^2$ (using $z_1 z_3 = z_5^2$) $= (z_2 z_5) z_5 = z_4 z_5 z_6$ (using $z_2 z_5 = z_4 z_6$).

Likewise, we can show that the remaining overlapping ambiguities:

$z_1 z_2 z_5, z_1 z_2 z_6, z_1 z_3 z_4, z_1 z_3 z_6, z_2 z_3 z_4, z_2 z_3 z_5$ reduce to unique expressions in z_i 's namely, $z_4^2 z_5, z_4^2 z_6, z_4 z_5^2, z_4 z_6^2, z_5 z_6^2$ after applying different reduction rules. Therefore, $\tilde{\phi}$ is an isomorphism.

Consider the embedding

$$\psi : \mathbb{P}^2 \hookrightarrow \mathbb{P}^5$$

given by

$$(x_1, x_2, x_3) \mapsto (x_1^2, x_2^2, x_3^2, x_1 x_2, x_1 x_3, x_2 x_3).$$

Let J be the homogeneous ideal of A generated by 2×2 minors of the matrix

$$\begin{pmatrix} z_1 & z_4 & z_5 \\ z_4 & z_2 & z_6 \\ z_5 & z_6 & z_3 \end{pmatrix},$$

where z_i 's are the homogeneous coordinates of \mathbb{P}^5 . Note that $\psi(\mathbb{P}^2)$ is given by J . Further, note that $I \subseteq J$ and $\dim(A/I) = \dim(A/J) = 3$ (computed using Macaulay2 [3]). Thus, the GIT quotient $T \setminus \setminus (X(v_5))_T^{ss}(\mathcal{L}(4\lambda))$ is isomorphic to $(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$ as a polarized variety. \square

Acknowledgement. Both the authors would like to thank the Indian Institute of Technology Bombay for the postdoctoral fellowships and the hospitality during their stay.

REFERENCES

- [1] G. M. Bergman, The diamond lemma for ring theory, *Adv. Math.* **29**(2)(1978), 178–218.
- [2] A. W. M. Dress, W. Wenzel, A Simple Proof of an Identity Concerning Pfaffians of Skew Symmetric Matrices, *Advances in Mathematics*, **112**(1995), 120–134.
- [3] D. R. Grayson, M. E. Stillman, Macaulay2, a software system for research in algebraic geometry, Available at <https://faculty.math.illinois.edu/Macaulay2/>.
- [4] R. Hartshorne, *Algebraic Geometry, Graduate Texts in Mathematics book series*, (GTM **52**, Springer, New York, 1977.
- [5] J. E. Humphreys, *Introduction to lie Algebras and Representation Theory*, Springer-Verlag, Berlin, Heidelberg, New York, 1972.
- [6] J. E. Humphreys, *Linear Algebraic Groups*, Springer-Verlag, Berlin, Heidelberg, New York, 1975.

- [7] J. C. Jantzen, *Representations of Algebraic Groups*, second edition, Mathematical Surveys and Monographs **107**, Amer. Math. Soc., Providence, RI, 2003.
- [8] S. S. Kannan, S. K. Pattanayak, Torus quotients of homogeneous spaces-minimal dimensional Schubert varieties admitting semi-stable points, *Proc. Indian Acad. Sci. (Math.Sci.)* **119**(4), 2009, 469–485.
- [9] S. S. Kannan, P. Sardar, Torus quotients of homogeneous spaces of the general linear group and the standard representation of certain symmetric groups, *Proc. Indian Acad. Sci.* **119** (2009), 81–100.
- [10] S. Kumar, Descent of line bundles to GIT quotients of flag varieties by maximal torus, *Transformation groups*, **13**(2008), 757–771.
- [11] V. Lakshmibai, K. N. Raghavan, *Standard Monomial Theory Invariant Theoretic Approach*, Encyclopaedia of Mathematical Sciences **137**, Springer-Verlag, Berlin, 2008.
- [12] P. Littelmann, A Generalization of the Littelwood-Richardson Rule, *Journal of Algebra*, **130**(1990), 328–368.
- [13] D. Mumford, J. Fogarty, F. Kirwan, *Geometric Invariant Theory*, Third Edition, Springer-Verlag, Berlin, Heidelberg, New York, 1994.
- [14] A. Nayek, S. K. Pattanayak, S. Jindal, Projective normality of torus quotients of flag varieties, *Journal of Pure and Applied Algebra*, **224**(2020).
- [15] A. Nayek, P. Saha, On torus quotients of Schubert varieties in Orthogonal Grassmannian, *Preprint*.

ARPITA NAYEK, DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI, 400076, INDIA.

Email address: arpitan@math.iitb.ac.in

PINAKINATH SAHA, DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI, 400076, INDIA.

Email address: psaha@math.iitb.ac.in