

ON THE AUERBACH BASES OF l_p^n SPACES

ARUN MAITI AND DEBMALYA SAIN

ABSTRACT. In this paper, we study Auerbach basis of the Banach spaces l_p^n . We provide a complete classification of the spaces in terms of the cardinality of their bases. We also give a complete description of these bases for l_p^3 (l_p^2 is easy).

1. INTRODUCTION

Birkhoff-James orthogonality is a generalization of the notion of orthogonality in Hilbert spaces to Banach spaces introduced by Birkhoff in [2]. Let \mathbb{X} be a n dimensional real Banach space and $S_{\mathbb{X}}$ denote its unit sphere. For $x, y \in \mathbb{X}$, x is said to be orthogonal to y in the sense of Birkhoff-James, denoted by $x \perp_B y$, if

$$\|x\| \leq \|x + \lambda y\| \quad \text{for all } \lambda \in \mathbb{R}.$$

The concept of Auerbach basis of a Banach space is a close analogue of the concept of orthonormal basis of a Hilbert space. A basis \mathcal{B} of a Banach space $(X, \|\cdot\|)$ is called an Auerbach basis if for all $x \in \mathcal{B}$,

$$\|x\| = 1 \quad \text{and } x \perp_B \text{span}\{\mathcal{B} \setminus x\}.$$

Equivalently, $\mathcal{B} = \{v_1, \dots, v_n\}$ is an Auerbach basis if

$$\|v_i\| = 1 \quad \text{and } \|v^i\| = 1 \quad \text{for } i = 1, 2, \dots, n, \quad (1.1)$$

where v^1, \dots, v^n is a basis of the dual space X^* of X dual to v_1, \dots, v_n , i.e., $v^i(v_j) = \delta_{ij}$.

A basis vector of an Auerbach basis will be referred to as a *Auerbach basis vector*.

For the space $l_p^n := (\mathbb{R}^n, \|\cdot\|_p)$, the norm $\|\cdot\|_p$ is invariant under signed permutations of co-ordinates and hence the Birkhoff-James orthogonality. We shall often express Auerbach bases of l_p^n by $n \times n$ matrices whose rows are basis vectors, and two bases are said to be *equivalent* if one can be obtained from the other by negating rows or columns, or by interchanging rows or columns.

One can also use semi-inner product $[\cdot, \cdot] : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$ to interpret the Birkhoff-James orthogonality. For the space l_p^n , $1 < p < \infty$, there is a unique semi-inner product $[\cdot, \cdot]$ corresponding to the norm $\|\cdot\|_p$ given by

$$[y, x] = \frac{\sum_{j=1}^n y_j x_j |x_j|^{p-2}}{\|x\|_p^{p-2}},$$

where $x = (x_1, x_2, \dots, x_n) \neq \mathbf{0}$ and $y = (y_1, y_2, \dots, y_n)$. We refer the readers to [7, 5, 10] for more information on semi-inner-products and their potential applications in studying geometry of Banach spaces, and [3] particularly for the l_p^n spaces.

Denoting by ${}^p x := (x_1|x_1|^{p-2}, x_2|x_2|^{p-2}, \dots, x_n|x_n|^{p-2})$, the gradient of the function $\|\cdot\|_p^p$, we have

$$x \perp_B y \iff [y, x] = 0 \iff y \cdot {}^p x = 0. \quad (1.2)$$

We see that Birkhoff-James orthogonality is not sensitive to scaling. Further, for vectors with components in $\{0, 1, -1\}$, Birkhoff-James orthogonality in l_p^n spaces is equivalent to the Euclidean orthogonality. So, up to scalar multiples of the rows, matrices with mutually orthogonal rows and entries in $\{0, 1, -1\}$ correspond to Auerbach bases of l_p^n . We refer to these bases as *stationary bases*.

2010 *Mathematics Subject Classification*. 47I05; 46B20; 52A21.

Key words and phrases. l_p^n -spaces; Birkhoff-James orthogonality; Auerbach bases.

This includes, for example, weighing matrices which are the matrices with entries in $\{0, 1, -1\}$ and satisfying $WW^T = mI_n$ for some integer $m \leq n$ with I_n denoting the identity matrix. We refer readers to [11] for more information on such matrices.

In §2, we give a brief account of some of the known results on the existence of Auerbach bases for general Banach spaces. In §3, we prove that there are only finitely many Auerbach bases of l_p^n for $1 < p < \infty$, $p \neq 2$ for all n , Theorem 3.1. In §3.1, we show, however, that there are infinitely many Auerbach bases of l_p^n for $p = 1, 2, \infty$, $n \geq 3$.

In §4, we use elementary analysis to characterize the Auerbach bases of l_p^3 . This produces concrete examples of non-stationary bases for all p .

A *strong Auerbach basis* of the Banach space l_p^n , $1 < p < \infty$ is defined to be an Auerbach basis such that the span of any subset of m basis vectors is isometrically isomorphic to the Banach space l_p^m for all $m \leq n$. In §5, we give a complete description of the strong Auerbach bases of l_p^n for all n, p .

2. PRELIMINARIES

Auerbach first established the existence of at least one Auerbach basis for any finite dimensional Banach space in his doctoral thesis [1]. Proofs were then published by Day in [4] and independently by Taylor in [12]. In their proofs the basis vectors are selected by maximizing the volume of the convex symmetric envelope of n -tuples of vectors from the unit sphere. In [9], Plichko proved the existence of two different Auerbach bases for infinite dimensional Banach space that are not a Hilbert space. Here bases are identified if they differ only by a permutation or multiplication by scalars of absolute value one.

We may view $S_{\mathbb{X}}^n$, the cartesian product of n copies of $S_{\mathbb{X}}$, as the set of $n \times n$ real matrices whose rows are unit vectors in \mathbb{X} . We denote by $S_{\mathbb{X}, > 0}^n$, the subspace of non singular matrices in $S_{\mathbb{X}}^n$. Recall that a Banach space \mathbb{X} is called *smooth* if at every point of the unit ball of \mathbb{X} there is only one supporting hyperplane. We have the following alternative description of the bases for smooth Banach spaces.

Proposition 2.1. *A set of vectors $\{v_1, \dots, v_n\}$ form an Auerbach basis of a smooth Banach space \mathbb{X} if and only if it is a critical point of the determinant function $\det : S_{\mathbb{X}, > 0}^n \rightarrow \mathbb{R}$.*

Proof. By Theorem 4.1 in [6], the smoothness of \mathbb{X} is equivalent to the Gateaux differentiability of the norm at all non-zero points of \mathbb{X} . The norm is also a Lipschitz function in the finite dimension, so Gateaux derivative is the usual derivative. Consequently, the unit sphere $S_{\mathbb{X}}^n$ is a differentiable manifold, as does the subspace $S_{\mathbb{X}, > 0}^n$, and the determinant $\det : S_{\mathbb{X}, > 0}^n \rightarrow \mathbb{R}$ is a differentiable function [8]. By definition, a set of vectors $\{v_1, \dots, v_n\} \in S_{\mathbb{X}, > 0}^n$ is a critical point of the determinant if for all vector $u_k \in T_{v_k} S_{\mathbb{X}}$, the tangent space to $S_{\mathbb{X}}$ at v_k , we have

$$\sum_{k=1}^n \det(v_1, \dots, v_{k-1}, u_k, v_{k+1}, \dots, v_n) = 0.$$

This happens if and only if $\det(v_1, \dots, v_{k-1}, u_k, v_{k+1}, \dots, v_n) = 0$ for all k and for all $u_k \in T_{v_k} S_{\mathbb{X}}$. This means

$$\text{span}(v_1, \dots, v_{k-1}, v_{k+1}, \dots, v_n) = T_{v_k} S_{\mathbb{X}}$$

for all k . Now by definition, the set of vectors (v_1, v_2, \dots, v_n) form an Auerbach basis. \square

In [13], using a similar interpretation of Auerbach bases for general Banach spaces and the Lusternik–Schnirelman theory of the critical points, Weber and Wojciechowski proved that there exist at least $(n - 1)n/2 + 1$ Auerbach bases of Banach spaces of dimension n .

$$\text{where } A_i = \begin{bmatrix} a_1^{(1)} | a_1^{(1)} |_{\bar{p}} & a_2^{(1)} | a_2^{(1)} |_{\bar{p}} & \cdots & a_n^{(1)} | a_n^{(1)} |_{\bar{p}} \\ a_1^{(2)} | a_1^{(2)} |_{\bar{p}} & a_2^{(2)} | a_2^{(2)} |_{\bar{p}} & \cdots & a_n^{(2)} | a_n^{(2)} |_{\bar{p}} \\ \vdots & \vdots & \ddots & \vdots \\ p\epsilon_i^p a_1^{(i)} | a_1^{(i)} |_{\bar{p}} & p\epsilon_i^p a_2^{(i)} | a_2^{(i)} |_{\bar{p}} & \cdots & p\epsilon_i^p a_n^{(i)} | a_n^{(i)} |_{\bar{p}} \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{(n)} | a_1^{(n)} |_{\bar{p}} & a_2^{(n)} | a_2^{(n)} |_{\bar{p}} & \cdots & a_n^{(n)} | a_n^{(n)} |_{\bar{p}} \end{bmatrix} \text{ and } \hat{p} = p - 1, \bar{p} = p - 2.$$

By definition, $a^{(i)} \cdot_p a^{(j)} = 0$ for $i \neq j$, and $a^{(i)} \cdot_p a^{(i)} \neq 0$ for all i . So, by induction, the row vectors $({}_p a^{(1)}, {}_p a^{(2)}, \dots, {}_p a^{(n)})$ are linearly independent, consequently, A_i 's are all non-singular matrices for all $\epsilon_i > 0$.

We can treat the determinant of the submatrix $M_{i,i}$, $i \leq n$, as a function of the variables ϵ_j , $j \leq i$. We will show by induction that $M_{i,i}$ is non-singular for some values of ϵ_j s. For $i = 1$, $M_{1,1} = A_1$ is clearly non-singular for all $\epsilon_1 > 0$. Let us assume that the submatrix $M_{i-1,i-1}$ is non singular for

$$(\epsilon_1, \epsilon_2, \dots, \epsilon_{i-1}) \in (0, \delta_1) \times (0, \delta_2) \times \cdots \times (0, \delta_{i-1}) \text{ for some } \delta_1, \delta_2, \dots, \delta_{i-1} > 0.$$

Note that each term in the last n rows and columns of $M_{i,i}$ outside the matrix A_i is a multiple of ϵ_i , and only the entries in the i -th column of A_i^\top are multiple of ϵ_i . So we can write

$$\det(M_{i,i}) = \epsilon_i^p \det(A_{i,i}) \det(M_{i-1,i-1}) + \epsilon_i^{p+1} P(\epsilon_i),$$

where P is a bounded function. So, for sufficiently small $\delta_i > 0$,

$$\det(M_{i,i}) \neq 0 \text{ for all } \epsilon_j \in (0, \delta_j) \text{ for } j = 1, 2, \dots, i.$$

This completes the induction, and thus $M_{n^2, n^2} = M$ is nonsingular. Using inverse function theorem, we deduce that the solutions of Equation 3.2 are isolated, and so do the solutions of Equation 3.1. The first part of the statement of the theorem now follows as the solutions lies in a compact space. For the second part, let $[b_j^{(i)}]$ be a solution of the equations $f_{ij}^p = \delta_{ij}$ for some t . Then by the first part of the theorem, there is a neighborhood B_ϵ of $[b_j^{(i)}]$ which contains no other solution of $f_{ij}^p = \delta_{ij}$, and the function

$$f : B_\epsilon \times (2, \infty) \rightarrow \mathbb{R}^{n^2}, \quad f([x_j^{(i)}], t) = f_{ij}^t([x_j^{(i)}])$$

is a submersion at $([b_j^{(i)}], p)$. By the local submersion theorem, there is an open interval $(p - \delta_1, p + \delta_2)$ and a differentiable function $g : (p - \delta_1, p + \delta_2) \rightarrow \mathbb{R}^n$ such that

$$f(g(t), t) = \delta_{ij} \text{ for all } t \in (p - \delta_1, p + \delta_2).$$

Let J be the maximal domain of definition of g . Since the above argument holds for any $p \in (2, \infty)$, J is open in $(2, \infty)$. By continuity, J is also closed in $(2, \infty)$, hence $J = (2, \infty)$. This completes the proof of the theorem. \square

Remark 3.2. *The above theorem does not hold true in general for Banach spaces that are not Hilbert spaces. For example, for $2 < p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$, the Banach space $(\mathbb{R}^2, \|\cdot\|)$ with the norm $\|\cdot\|$ given by the l_p^2 norm in the first quadrant and the l_q^2 norm in the second quadrant of \mathbb{R}^2 .*

3.1. Auerbach bases of l_2^n , l_∞^n , and l_1^n . The following example shows that there are infinitely many Auerbach bases of l_∞^3 ,

$$J_\infty = \begin{bmatrix} 1 & 1 & 1 \\ -1 & 1 & 1 \\ t & 1 & -1 \end{bmatrix}$$

for $-1 \leq t \leq 1$. It follows that there are infinitely many bases of l_∞^n , $n \geq 3$. The same can be said about the space l_1^n for $n \geq 3$, by 1.1 and the duality between l_∞^n and l_1^n .

For the Hilbert space l_2^n Birkhoff-James orthogonality is the same as the standard orthogonality, hence there are infinitely many bases.

4. AUERBACH BASES OF l_p^3 , $p \neq 1, 2, \infty$

Let P_3 denote solid cube which is the convex hull of the points $(\pm 1, \pm 1, \pm 1) \in \mathbb{R}^3$. We denote

$$\pm e_1 = (\pm 1, 0, 0), \pm e_2 = (0, \pm 1, 0), \pm e_3 = (0, 0, \pm 1).$$

Let $F_{\pm e_i}$ denote the facet of the hollow cube ∂P_3 containing $\pm e_i$, respectively for $i = 1, 2, 3$. We observe that

- 1) for $x = (x_1, x_2, x_3)$, x_i and ${}^p x_i$ have the same signs and $|{}^p x_i| \leq |x_i|$ for $i = 1, 2, 3$;
- 2) if $x \in F_e^\circ$ (interior of F_e) or $x \in \partial F_e$ (boundary of F_e) if and only if ${}^p x \in F_e^\circ$ or ${}^p x \in \partial F_e$, respectively.

The following lemma turns out to be useful in characterizing the Auerbach bases of l_p^3 .

Lemma 4.1. *For $a = (a_1, a_2, a_3) \in P_3$, the straight-line joining a and ${}^p a$ either intersect one of the faces F_{+e_i} , F_{-e_i} or the plane portion $\{x_i = 0\} \cap P_3$, for $i = 1, 2, 3$.*

Proof. Without loss of generality we may assume that $a_1, a_2, a_3 \geq 0$. Let L be the projection of the straight line joining a and ${}^p a$ on the plane (e_1, e_2) . The equation of L is given by

$$x_1 = a_1 + \frac{a_1 - (a_1)^p}{a_2 - (a_2)^p} (x_2 - a_2) = a_1 + a_1 \frac{1 - (a_1)^{p-1}}{1 - (a_2)^{p-1}} \cdot \frac{x_2 - a_2}{a_2}.$$

For $a_1 \geq a_2$, we have $1 - (a_1)^p \leq 1 - (a_2)^p$, therefore, if $x_2 = 0$ on this straight line, then $-1 \leq x_1 \leq 1$. This also means that if $x_1 = 1$ then $-1 \leq x_2 \leq 1$. Analogously, for $a_1 < a_2$, if $x_2 = 1$ then $-1 \leq x_1 \leq 1$.

If the statement of the lemma is false for $i = 1$, then it contradicts what we have just proved. A similar argument can be made for $i = 2, 3$. \square

In light of the above result, the following proposition holds true immediately.

Assertion 4.2. *Let Q denote an octant in \mathbb{R}^3 . Suppose H be a plane such that $\{x_k = 1\} \cap Q \cap H = \emptyset$ and $\{x_k = 0\} \cap Q \cap H = \emptyset$, then for $i \neq k$ and for all $y \in F_{e_i} \cap Q \cap H$, ${}^p y$ lies in the same connected component of $F_{e_i} \setminus H$ as do e_i .*

We note that for $p > 0$, the function $x + x^p$ is monotone on $[0, 1]$. This implies that there exist an unique $r_p \in [0, 1]$ depending on p such that

$$1 - r_p - r_p^p = 0. \tag{4.1}$$

Consider the function $f(r) = (1 - r)^p + r - 1$ and its derivative $f'(r) = r(1 - r^p)$. Clearly $f(0) = f(1) = 0$. The function $f'(r)$ can be shown to be concave by looking further at the second derivative $f''(r)$. This together with the fact that $f'(r)$ has only two zeros in $(0, 1)$ implies that f has only one zero in $(0, 1)$. Thus we obtain the following functional inequalities which will be crucial in the proof of Theorem 4.4. For $1 < p < \infty$,

$$(1 - x_p)^p > 1 - x \text{ for } x \in [0, r_p], (1 - x_p)^p > 1 - x \text{ for } x \in [r_p, 1], \tag{4.2}$$

and for $0 < p < 1$,

$$(1 - x_p)^p > 1 - x \text{ for } x \in [0, 1 - r_p], (1 - x_p)^p < 1 - x \text{ for } x \in [1 - r_p, 1]. \tag{4.3}$$

The following technical lemma will be useful in the proof of Theorem 4.4.

Lemma 4.3. For a fixed $p > 0$, let γ be the curve defined by $\gamma(x) = (t^p x^{\frac{1}{p}} + s^p)^p$, where t and s are the parameters.

1. For $t < 0$, $s > 0$ and $1 < p < \infty$ (respectively $0 < p < 1$), γ is concave (respectively convex) on $[0, 1]$.
2. For $0 < s \leq t \leq 1$ and $1 < p < \infty$ (respectively $0 < p < 1$), γ intersect the line $y = tx + s$ in the first quadrant $\{(x, y) : 0 \leq x, y \leq 1\}$ if and only if $(\frac{1-s^p}{t^p})^p \leq \frac{1-s}{t}$ (respectively $(\frac{1-s^p}{t^p})^p \geq \frac{1-s}{t}$).

Proof. The proof of the first assertion can be inferred from the derivative of γ .

The "if" part of the second assertion is also easy to deduce. For the "only if" part we first observe that γ is convex for $x \geq 0$. So, it is enough to show that the tangent to γ at $y = 1$ has slope greater than t . We compute the slope to be $t^p((1 - s^p t^p)^p)^{1-p}$, and it is clearly greater than t for $(1 - s^p t^p)^p < t$.

For $(1 - s^p t^p)^p > t$, we note that the coordinates of a point on the curve γ can be expressed as $(x^p, (t^p x + s^p)^p)$. In order to show that the curve does not intersect the line $y - tx = s$, we argue that it lies entirely in one component of $\{[0, 1] \times [0, 1]\} \setminus \gamma$. For that it is sufficient to show that the following holds:

$$s > (-t, 1) \cdot (x^p, (t^p x + s^p)^p) \quad \text{equivalently,} \quad s + tx^p > (t^p x + s^p)^p. \quad (4.4)$$

Since by our hypothesis $s^p + t^p > 1$ which again implies $\frac{1-s}{t} < t$, if the curve $s + tx^p$ intersect the line $t^p x + s^p$, then by the Rolle's theorem, the slope of the curve must be equal to t^p for some $x \in [0, t)$ (see Figure 4.1). Since $\frac{d}{dx} s + tx^p = t(p-1)x^{p-1}$,

$$t(p-1)x^{p-1} = t^p \implies x = (p-1)^{\frac{1}{p-1}} t \geq t \quad (\text{since } p-1 \geq 1).$$

Since $(t^p x + s^p)^p < t^p x + s^p$, the required inequality of Equation 4.4 holds.




Fig. 4.1

□

Theorem 4.4. For $1 < p < \infty$, $p \neq 2$, an Auerbach basis vector of l_p^3 must be one of the following up to a signed permutation of the components: $(0, 0, 1)$, $(0, \frac{1}{\sqrt[2]{2}}, \frac{1}{\sqrt[2]{2}})$ or $(\frac{r_p}{\sqrt[2]{2+r_p^p}}, \frac{1}{\sqrt[2]{2+r_p^p}}, \frac{1}{\sqrt[2]{2+r_p^p}})$.

Proof. We first consider the case for $2 < p < \infty$. It is easy to see that $(0, 0, 1)$ and $(0, \frac{1}{\sqrt[2]{2}}, \frac{1}{\sqrt[2]{2}})$ are Auerbach basis vectors.

For the other possibilities, we look for three vectors in ∂P_3 which can be rescaled to an Auerbach bases of l_p^3 . We divide P_3 into the following regions, see Figure 4.2.




Fig. 4.2. Case 1




Fig. 4.3. Case 2

$$\begin{aligned}
 R_0 &:= \{(x, y, 1) \mid 0 \leq x \leq 1, 0 \leq y \leq 1\}; & R_1 &:= \{(-1, y, z) \mid 0 \leq y \leq 1, -1 \leq z \leq 0\} \\
 R_2 &:= \{(-1, y, z) \mid -1 \leq y \leq 0, 0 \leq z \leq 1\}; & R_3 &:= \{(x, -1, z) \mid -1 \leq x \leq 0, 0 \leq z \leq 1\} \\
 R_4 &:= \{(x, -1, z) \mid 0 \leq x \leq 1, 0 \leq z \leq 1\}; & R_5 &:= \{(x, y, 1) \mid -1 \leq x \leq 0, -1 \leq y \leq 0, \} \\
 R_x &:= \{(-1, y, z) \mid 0 \leq y \leq 1, 0 \leq z \leq 1\}. & & (4.5)
 \end{aligned}$$



Further, for a region R , $-R$ denote the region consisting of all antipodal points of R .

Let \perp_x denote the plane perpendicular to the vector x . By Equation 1.2, x, y are mutually orthogonal in the sense of Birkhoff-James iff

$$y \in \perp_{p_x}, \quad \text{and} \quad {}^p y \in \perp_x. \quad (4.6)$$

Let $\{u, v, w\}$ form an Auerbach basis of l_p^3 . We may choose $u = (a, b, 1)$, $0 \leq a \leq b \leq 1$ up to equivalence. Note that $a = 0$ reduces the problem to l_p^2 , so as shown in §3, $u = (0, 0, 1)$ or $(0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$. So we may further choose $a > 0$.

Case 1: $a + b \leq 1$: In this case, both the planes \perp_u and \perp_{p_u} do not intersect the face $F_{\pm e_3}$. The equation of the straight lines $\perp_u \cap \{x = -1\}$ and $\perp_{p_u} \cap \{x = -1\}$ are given by $z = by + a$ and $z = b|b|^{p-2}y + a|a|^{p-2} = b^{p-1}y + a^{p-1}$, respectively. So, the intersection $\{^p \zeta \mid \zeta \in \perp_{p_u}\} \cap \{x = -1\}$ is given by the curve $\gamma : y \rightarrow (b^{p-1} \text{sgn}(y)|y|^{\frac{1}{p-1}} + a^{p-1})^{p-1}$. Then it follows from Lemma 4.3 and Assertion 4.2 that v and w can possibly lie only in the regions R_1 and R_4 . Since the region $-R_1$ and R_4 are adjacent, the plane \perp_{p_v} does not intersect R_4 for $v \in R_1$. Consequently, v and w cannot be mutually Birkhoff orthogonal.

Fig. 4.4. Region R_2 Fig. 4.5. Region R_0

Case 2: $a + b > 1$ but $a^p + b^p \leq 1$: The plane \perp_{p_u} does not intersect the interior of the facet $z = 1$ in this case as well. Thus we arrive at the same conclusion as in Case 1.

Case 3: $a^p + b^p > 1$. In this case, both the planes \perp_u and \perp_{p_u} intersect the interior of the facet $z = 1$. Suppose v lies in the R_2 region with $v = (-1, -c, d)$. By Lemma 4.3, if 4.6 holds for $x = u$ and $y = v$ then following inequality must be satisfied (see figure 4.4)

$$\left(\frac{1-a^p}{b^p}\right) \leq \frac{1-a}{b} \implies (1-a^p)^p \leq b^{p^2-1}(1-a) \quad (4.7)$$

Since $b \leq 1$, so by 4.2 we must have $a \geq r_p$ and for equality, i.e., for $a = r_p$, we must have $b = 1$. On the other hand, if 4.6 holds for $x = v$ and $y = u$, then the following inequality must be satisfied (see Fig. 4.5)

$$(d^p - c^p)^p \geq d - c \implies d^{p^2-1}(1-l^p)^p \geq 1-l \text{ with } l = c/d \quad (4.8)$$

By inequality 4.2, $l > r_p$ and for equality, i.e., for $l = r_p$, we must have $d = 1$. If u and v lies in the R_0 and R_2 , respectively, then it is easy to see that w can possibly lie only in the regions R_1 or R_4 . Suppose w lies in the region R_4 , then by Lemma 4.4, the following must hold, see figure 4.6.

$$(b^p - a^p)^p \geq b - a \implies b^{p^2-1}(1-k^p)^p \geq 1-k \text{ with } k = a/b. \quad (4.9)$$

Therefore, by inequality 4.2, we have $k \leq r_p$. However, this contradicts 4.7 unless $k = r_p$, in which case $a = r_p, b = k = 1$. This also implies $w = (r_p, -1, 1)$. On the other hand, if w lies in R_1 region, then ${}^p w \in \perp_u \cap \perp_v$. But $\perp_u \cap \perp_v = \emptyset$, unless the following holds (see figure 4.7)

$$\frac{1-c}{d} \leq b-a.$$

Therefore, $c \geq a$, which implies $l = \frac{c}{d} \geq \frac{a}{d} \geq r_p$. Since $l \leq r_p$ by 4.8, then we must have $l = r_p$. This means $c = a = r_p$ and $d = 1, b = 1$.

This produces an isometry.

For the converse it is sufficient to show that if $h : l_p^2 \rightarrow V_2 = \text{span}(w_1, w_2)$ is an isometry with $w_1, w_2 \in \mathbb{R}^n$, then $\text{span}(w_1, w_2)$ must be spanned by two vectors with disjoint support. Suppose we have an isometry defined by extending the following assignment linearly

$$e_1 \rightarrow \nu_1 \quad \text{and} \quad e_2 \rightarrow \nu_2,$$

where $\nu_1, \nu_2 \in \text{span}(w_1, w_2)$.

Let us call a point x on the unit sphere of l_p^n spherical if x and the normal vector on the unit sphere at x are linearly dependent. It is not difficult to see that a spherical point must be one of the eight points: $\pm e_1, \pm e_2, \pm(e_1 \pm e_2)$.

Recall that the normal vector at a point $x = (x_1, x_2, \dots, x_n)$ on the unit sphere in l_p^n is given by ${}^p x = (x_1|x_1|^{p-2}, x_2|x_2|^{p-2}, \dots, x_n|x_n|^{p-2})$. So, $x = (x_1, x_2, \dots, x_n)$ is a spherical point if and only if either $x_i = 0$ or $|x_i| = c$ for some constant c for $i = 1, 2, \dots, n$. Since $h(e_1 + e_2) = v_1 + v_2$, $v_1 + v_2$ is also a spherical point. v_1, v_2 and $v_1 + v_2$ are all spherical point, implies v_1 and v_2 have mutually disjoint support. \square

In light of the above proposition and Theorem 4.4, it is now easy to deduce the following about strong Auerbach bases (defined in the introduction) of l_p^n .

Theorem 5.2. *A strong Auerbach basis vector of l_p^n with $1 < p < \infty$ must be of the form $(1, 0, \dots, 0)$ or $(\frac{1}{\sqrt[p]{2}}, \frac{1}{\sqrt[p]{2}}, 0, \dots, 0)$ up to a signed permutation of the components.*

ACKNOWLEDGEMENT

The first author was partially supported by Department of Science and Technology grant EMR/2016/006624 and by the UGC Centre for Advanced Studies. The research of the second author was sponsored by SERB-NPDF under the mentorship of Professor Apoorva Khare.

REFERENCES

1. A. M. Auerbach, *O polu krzywych wypuklych o srednicach sprzezonych (on the area of convex curves with conjugate diameters)*, Ph.D. thesis, University of Lwow (1930).
2. G. Birkhoff, *Orthogonality in linear metric spaces*, Duke Math. J. **2** (1935), 169–172.
3. Arup Chattopadhyay, Debmalya Sain, and Tanusri Senapati, *characterization of symmetric points in l_{pn} -spaces*, Linear and Multilinear Algebra **0** (2019), no. 0, 1–12.
4. Mahlon M. Day, *Polygons circumscribed about closed convex curves*, Trans. Amer. Math. Soc. **62** (1947), 315–319. MR 22686
5. J. R. Giles, *Classes of semi-inner-product spaces*, Trans. Amer. Math. Soc. **129** (1967), 436–446. MR 217574
6. Robert C. James, *Orthogonality and linear functionals in normed linear spaces*, Trans. Amer. Math. Soc. **61** (1947), 265–292. MR 21241
7. G. Lumer, *Semi-inner-product spaces*, Trans. Amer. Math. Soc. **100** (1961), 29–43. MR 133024
8. I. Madsen and J. Tornehave, *From calculus to cohomology. de rham cohomology and characteristic classes*, Cambridge University Press, Cambridge, 1997.
9. A. M. Plichko, *On the volume method in the study of auerbach bases of finite dimensional banach spaces*, Colloq. Math **69** (1995), no. 2.
10. Debmalya Sain, *On the norm attainment set of a bounded linear operator and semi-inner-products in normed spaces*, Indian J. Pure Appl. Math. **51** (2020), no. 1, 179–186. MR 4076205
11. Jennifer Seberry, *Orthogonal designs*, Springer, Cham, 2017, Hadamard matrices, quadratic forms and algebras, Revised and updated edition of the 1979 original [MR0534614]. MR 3729402
12. Angus E. Taylor, *A geometric theorem and its application to biorthogonal systems*, Bull. Amer. Math. Soc. **53** (1947), 614–616. MR 20718
13. Andrzej Weber and Michał Wojciechowski, *On the Pełczyński conjecture on Auerbach bases*, Commun. Contemp. Math. **19** (2017), no. 6, 1750016, 19. MR 3691505