

Universal lower bound on topological entanglement entropy

Isaac H. Kim,¹ Michael Levin,² Ting-Chun Lin,^{3,4} Daniel Ranard,⁵ and Bowen Shi³

¹*Department of Computer Science, University of California, Davis, CA 95616, USA*

²*Kadanoff Center for Theoretical Physics, University of Chicago, Chicago, Illinois 60637, USA*

³*Department of Physics, University of California at San Diego, La Jolla, CA 92093, USA*

⁴*Hon Hai Research Institute, Taipei, Taiwan*

⁵*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139*

(Dated: February 3, 2023)

Entanglement entropies of two-dimensional gapped ground states are expected to satisfy an area law, with a constant correction term known as the topological entanglement entropy (TEE). In many models, the TEE takes a universal value that characterizes the underlying topological phase. However, the TEE is not truly universal: it can differ even for two states related by constant-depth circuits, which are necessarily in the same phase. The difference between the TEE and the value predicted by the anyon theory is often called the *spurious* topological entanglement entropy. We show that this spurious contribution is always nonnegative, thus the value predicted by the anyon theory provides a universal lower bound. This observation also leads to a definition of TEE which is invariant under constant-depth quantum circuits.

Ground states of 2D gapped Hamiltonians are believed to satisfy an area law: the entanglement entropy of a region cannot increase faster than its perimeter. In many examples, one finds the entropy of the reduced density matrix on a disk A takes the form

$$S(\sigma_A) = \alpha|\partial A| - \gamma + \dots \quad (1)$$

where $\alpha|\partial A|$ is the leading “area law” term proportional to the boundary length, γ is a constant term, and the ellipsis represents terms that vanish for large regions.

The constant term γ , under natural assumptions, was argued to be universal, i.e. the same for all gapped ground states in a given phase [1, 2]. In particular, γ takes a form determined solely by the underlying anyon theory of the phase, $\gamma = \log \mathcal{D}$, where $\mathcal{D} = \sqrt{\sum_a d_a^2}$ is the total quantum dimension of the anyons and d_a is the quantum dimension of the anyon a . Given its connection to anyons, the constant γ has been termed the “topological entanglement entropy” (TEE). The TEE can be computed in both non-solvable [3, 4] and solvable models [5–9], and it is often used as a smoking gun signature of topological order, or to distinguish two phases.

A common way to extract the TEE is to use a judicious linear combination of entropies of adjacent regions. We focus on the definition in Ref. [2], where the TEE γ is defined using the conditional mutual information $I(A : C|B)_\sigma := S(\sigma_{AB}) + S(\sigma_{BC}) - S(\sigma_B) - S(\sigma_{ABC})$ of regions A, B , and C forming an annulus as in Fig. 1(a):

$$I(A : C|B)_\sigma \equiv 2\gamma, \quad (2)$$

where σ is a ground state.

While the TEE is a useful diagnostic of topological order, it was soon observed [10] that it is not a genuine invariant of the topological phase. Two ground states are in the same phase if they are connected by a constant-depth circuit consisting of local gates. But γ as defined by Eq. (2) can change under such a circuit. In fact, a shallow circuit acting on a product state may achieve a

nonzero value of $I(A : C|B)$ for arbitrarily large regions [11–13]. Deviations of γ from the purportedly universal value $\log \mathcal{D}$ have been called “spurious” contributions, or the spurious TEE. There exist examples of states with spurious TEE in both trivial and non-trivial topological phases. While these examples are often associated with symmetry-protected topological phases (SPTs) [11, 13, 14], it appears that this is not always the case [15].

Our main result partially restores the universality of the TEE by showing that the spurious contribution is always nonnegative. Thus $\log \mathcal{D}$ provides a universal *lower bound* for the TEE γ :

$$\gamma \geq \log \mathcal{D}. \quad (3)$$

This observation leads directly to a definition of TEE on infinite systems which is invariant under constant-depth circuits, by minimizing the ordinary TEE over such circuits. More specifically, for a state ρ defined on the infinite 2D plane, the following quantity

$$\gamma_{\min} = \lim_{R \rightarrow \infty} \min_U \frac{1}{2} I(A_R : C_R|B_R)_{U\rho U^\dagger}, \quad (4)$$

yields $\log \mathcal{D}$ (for a class of states elaborated below), where the regions A_R, B_R, C_R have a radius and thickness of order R , and the minimum is taken over circuits U of depth $d < cR$ for some fixed constant $c \in (0, 1)$.

Our result helps restore the TEE as a rigorous diagnostic to distinguish topological phases, albeit with limitations. For instance, if a state has $I(A : C|B) = \log 2$ for some large regions, it may still be in the trivial phase (where $2 \log \mathcal{D} = 0$), and indeed such examples exist. But it *cannot* be in the same phase as the toric code, which has $2 \log \mathcal{D} = 2 \log 2$; the latter would require a negative spurious TEE, which we rule out.

Setup — We now explain our main result more precisely. We consider a special class of bosonic quantum many-body states, defined on infinite two-dimensional lattices, which we refer to as “reference states” and which are closely related to the states considered in Ref. [16].

Definition 1. A state σ is a reference state if (i) the TEE calculated as $\gamma_0 = \frac{1}{2}I(A : C|B)_\sigma$ is the same for any choice of regions topologically equivalent to Fig. 1(a) and (ii) the mutual information between two subsystems is zero for any two non-adjacent subsystems.

Our main technical result is the following inequality, which holds for any reference state σ , and for any circuit U whose depth is small compared to the radius and thickness of the annulus ABC :

$$I(A : C|B)_{U\sigma U^\dagger} \geq I(A : C|B)_\sigma \equiv 2\gamma_0. \quad (5)$$

In other words, we show that constant-depth circuits can never decrease the TEE, *when acting on a reference state*.

To understand the implications of this result, note that the set of reference states includes all ground states of string-net [7–9] and quantum double models [6], and more generally any state satisfying the entanglement bootstrap axioms [16]. Furthermore, for any of these examples, the RHS of Eq. (5) is known to be equal to $2 \log \mathcal{D}$ [1, 2, 16]. Therefore, (5) implies the claimed lower bound (3) for any state obtained by a constant-depth circuit acting on a string-net, quantum double, or entanglement bootstrap state. (In fact, with heuristic arguments, we can go further and derive (3) for general 2D gapped ground states, as we explain in the discussion section). Similarly, we deduce (4) with $\gamma_{\min} = \log \mathcal{D}$ for any state ρ given by a finite-depth circuit V applied to a reference state, where the minimum is achieved by the circuit $U = V^{-1}$.

As a final comment, while we work in the plane for concreteness, our proof also applies to the TEE defined on any disk-like region embedded in an arbitrary manifold, such as a torus.

Example: Toric code — To explain the key idea behind our proof, it is instructive to first focus on a concrete reference state σ , namely the toric code ground state [6] on a plane. For this state, $I(A : C|B)_\sigma = 2 \log 2$ so that $\gamma_0 = \log 2$ [5]. If we now apply a constant-depth quantum circuit U , defining $\tilde{\sigma} = U\sigma U^\dagger$, in general $I(A : C|B)_{\tilde{\sigma}} \neq 2 \log 2$. Nevertheless, we will show that for a sufficiently large annulus ABC , we still have the lower bound

$$I(A : C|B)_{\tilde{\sigma}} \geq 2 \log 2. \quad (6)$$

We first prove the bound (6) for a special class of constant-depth circuits U , namely those that are supported within a constant distance of BC [Fig. 1(b)]. Later we will extend this result to general constant-depth circuits.

Our basic strategy is to construct a state $\tilde{\lambda}$ that is “locally indistinguishable” from $\tilde{\sigma}$. More precisely, we will construct a state $\tilde{\lambda}$ that is indistinguishable from $\tilde{\sigma}$ over AB and BC : that is, $\tilde{\lambda}_{AB} = \tilde{\sigma}_{AB}$ and $\tilde{\lambda}_{BC} = \tilde{\sigma}_{BC}$. We can then express $I(A : C|B)_{\tilde{\sigma}}$ in terms of $I(A : C|B)_{\tilde{\lambda}}$ using the identity

$$I(A : C|B)_{\tilde{\sigma}} = I(A : C|B)_{\tilde{\lambda}} + S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC}). \quad (7)$$




FIG. 1. (a) The partition used to calculate the TEE. (b) Support of the unitary U considered. (c) String operator V_a creates an anyon a in the interior of the annulus and its antiparticle in the exterior.

Applying the strong subadditivity of the entropy (SSA) [17], we know that $I(A : C|B)_{\tilde{\lambda}} \geq 0$, and therefore

$$I(A : C|B)_{\tilde{\sigma}} \geq S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC}). \quad (8)$$

We will obtain the desired lower bound (6) from a judicious choice of $\tilde{\lambda}$.

The easiest way to construct an appropriate $\tilde{\lambda}$ is to first find a state λ that is locally indistinguishable from the toric code ground state σ . More precisely, we need a λ that is indistinguishable from σ over the past light cone of AB and BC (with respect to U). Once we find such a λ , we can then set $\tilde{\lambda} = U\lambda U^\dagger$.

How do we construct such a λ ? We use a more general approach later; in the present case of the toric code, our approach is to consider a probabilistic mixture of toric code *excited* states. More specifically, for each anyon type $a \in \mathcal{C} = \{1, e, m, \epsilon\}$, we define a corresponding excited state $\rho^{(a)}$ by $\rho^{(a)} = V_a \sigma V_a^\dagger$, where V_a is a unitary (open) string operator that places an anyon excitation a in the interior of the annulus and its antiparticle in the exterior [Fig. 1(c)]. We then define $\lambda = \sum_a p_a \rho^{(a)}$ for some probability distribution $\{p_a : a \in \mathcal{C}\}$. Note that λ has the requisite indistinguishability property as long as the endpoints of the string operators V_a (where the anyons are created) are sufficiently far from the annulus that they are outside the past light cones of AB and BC .

To proceed further, we need to evaluate the entropy difference $S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC})$. To do this, it is convenient to choose the path of the string operators V_a so that they avoid the region of support of the constant-depth circuit U [which by assumption is supported near BC]. Then V_a commutes with U so $\tilde{\lambda}$ can be written as a probabilistic mixture of the form

$$\tilde{\lambda} = \sum_a p_a \tilde{\rho}^{(a)}, \quad \tilde{\rho}^{(a)} = V_a \tilde{\sigma} V_a^\dagger. \quad (9)$$

Crucially, the $\tilde{\rho}^{(a)}$ states have two simplifying properties: (i) different $\tilde{\rho}_{ABC}^{(a)}$ are orthogonal to one another, and (ii) $S(\tilde{\rho}_{ABC}^{(a)}) = S(\tilde{\sigma}_{ABC})$. At an intuitive level, property (i) follows from the fact that each $\tilde{\rho}_{ABC}^{(a)}$ belongs to a different anyon sector on the annulus. More formally, (i) follows from the existence of a collection of (closed) string operators that are supported within ABC and that




FIG. 2. For any constant-depth circuit U , and for any subsystem S , we can obtain a circuit U' of same depth acting trivially on S , by removing from U the “light cone” (white gates) of S .

take on different eigenvalues in each state $\tilde{\rho}_{ABC}^{(a)}$. (These string operators are simply the closed versions of UV_aU^\dagger and they can be drawn within ABC as long as ABC is wider than twice the circuit depth of U). As for property (ii), this follows from the fact that the V_a are products of single-site unitaries; in particular, each V_a can be written as a product of a unitary acting entirely within ABC and a unitary acting entirely outside ABC , neither of which changes the entanglement entropy of ABC .

Given properties (i) and (ii) of $\tilde{\rho}^{(a)}$, the entropy difference can be computed as

$$S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC}) = H(\{p_a\}), \quad (10)$$

where $H(\{p_a\}) = -\sum_a p_a \log(p_a)$ is the Shannon entropy of the probability distribution $\{p_a\}$. Substituting (10) into (8), we obtain

$$I(A : C|B)_{\tilde{\sigma}} \geq H(\{p_a\}). \quad (11)$$

To get the best bound, we choose the probability distribution that maximizes $H(\{p_a\})$, namely $p_a = \frac{1}{4}$ for all $a \in \mathcal{C}$. For this choice, $H(\{p_a\}) = 2 \log 2$, yielding the desired bound (6).

To complete the argument, we now extend the bound (11) to general constant-depth circuits U . First, recall that the entanglement entropy of a subsystem is invariant under unitaries that act exclusively within the subsystem or its complement. Using this fact, we can make the replacement

$$I(A : C|B)_{U\sigma U^\dagger} = I(A : C|B)_{U'\sigma U'^\dagger}, \quad (12)$$

where U' is a constant-depth quantum circuit that acts trivially deep in the interior of A and also trivially far outside ABC [Fig. 3(a)]. Here, we are using the fact that U is a constant-depth quantum circuit, and therefore we can “cancel out” its action in a subsystem S by multiplying by an appropriate unitary supported in the light cone of S (Fig. 2).

Next, using SSA, we know that $I(A : C|B)$ is non-increasing when the region A is reduced in size. Therefore

$$I(A : C|B)_{U'\sigma U'^\dagger} \geq I(A' : C|B)_{U'\sigma U'^\dagger} \quad (13)$$

where $A' \subset A$ is shown in Fig. 3(b). Finally, applying the same reasoning as in (12), we can replace

$$I(A' : C|B)_{U'\sigma U'^\dagger} = I(A' : C|B)_{U''\sigma U''^\dagger} \quad (14)$$




FIG. 3. (a) We first remove gates from U on a “hole” within A ; we call the new circuit U' . (b) We then deform A to $A' \subset A$ such that the boundary of the annulus $A'BC$ is only partially covered. (c) We then further remove some of the gates in the vicinity of A' , obtaining U'' .

where U'' is a constant-depth circuit acting on the region shown in Fig. 3(c). Combining (12-14), we deduce that

$$I(A : C|B)_{U\sigma U^\dagger} \geq I(A' : C|B)_{U''\sigma U''^\dagger}. \quad (15)$$

The lower bound (15) is useful because it allows us to leverage our results from the first part of the proof. In particular, U'' is precisely the kind of constant-depth quantum circuit that we analyzed above, and therefore we know that $I(A' : C|B)_{U''\sigma U''^\dagger} \geq 2 \log 2$ for any sufficiently large annulus $A'BC$. Substituting this inequality into (15), we obtain the desired bound (6).

General case — Our proof for the toric code proceeded in three steps. First, we derived a lower bound (8) for $I(A : C|B)_{\tilde{\sigma}}$ in terms of the entropy difference $S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC})$ where $\tilde{\lambda}$ is any state that is indistinguishable from $\tilde{\sigma}$ over AB and BC . Second, we constructed an appropriate $\tilde{\lambda}$ and computed the desired entropy difference (10) in the special case where U is a constant-depth circuit supported within a constant distance of BC [Fig. 1(b)]. Combining these two results, we obtained the desired lower bound, but only for this special class of circuits U . In the third and final step, we extended this bound to arbitrary constant-depth circuits U using the inequality (15).

Conveniently, the first and third steps of our proof immediately generalize to any reference state σ since they do not use any properties of the toric code. On the other hand, in the second step, we used the specific structure of the toric code string operators¹, so we need a different argument for this step in the general case. In particular, instead of defining $\tilde{\lambda}$ in terms of a mixture of excited states, we will now define it in terms of the *maximum-entropy state*, or *max-entropy state* for short. Consider a larger annulus $Y = ABC \cup \text{Supp}(U)$, with U again as in [Fig. 1(b)]. Define a density matrix λ to be the maximum-entropy state consistent with the reduced

¹ In particular, we used the fact that the string operators of the toric code are depth-1 quantum circuits. In contrast, general anyon models can require higher depths, e.g., non-Abelian anyon strings need linear depth ([18], Chapter 7).




FIG. 4. Using the procedure in Fig. 2, we can remove the gates in U that act deep in the interior of the annulus without changing the entropy; starting from a U , whose support is depicted as the blue region in (a), we obtain a unitary \bar{U} , whose support is shown in (b). At this point, the support of \bar{U} is a union of two disks, which, under regrouping of sites, is topologically equivalent to (c).

density matrices of σ over the past light cones of AB and BC . We then define $\tilde{\lambda} = U\lambda U^\dagger$.

By construction, $\tilde{\lambda}$ is indistinguishable from $\tilde{\sigma}$ over AB and BC and therefore the lower bound (8) still holds. The only remaining question is the value of the entropy difference on the right-hand side. We claim that

$$S(\tilde{\lambda}_{ABC}) - S(\tilde{\sigma}_{ABC}) = S(\lambda_Y) - S(\sigma_Y) \quad (16)$$

and in turn

$$S(\lambda_Y) - S(\sigma_Y) = 2\gamma_0 \quad (17)$$

provided that each subsystem A, B , and C is sufficiently large compared to the circuit depth. See Eq. (E1) in the Supplemental Material (SM) for a self-contained derivation of Eq. (17) starting from Definition 1. We will then be finished once we prove Eq. (16).

We now present the proof of (16). Our main tool is the following lemma, which gives a condition under which the entropy difference of two bipartite states ρ and ρ' remains invariant under a quantum channel \mathcal{R} .

Lemma 1. *Let ρ and ρ' be density matrices over PQ such that $\rho'_Q = \rho_Q$ and $\rho'_P = \rho_P$. Let \mathcal{R}, \mathcal{T} be a pair of quantum channels $\mathcal{R} : Q \rightarrow \hat{Q}$ and $\mathcal{T} : \hat{Q} \rightarrow Q$. If*

$$\begin{aligned} \mathcal{T} \circ \mathcal{R}(\rho_{PQ}) &= \rho_{PQ} \\ \mathcal{T} \circ \mathcal{R}(\rho'_{PQ}) &= \rho'_{PQ} \end{aligned} \quad (18)$$

then

$$S(\rho_{PQ}) - S(\rho'_{PQ}) = S(\mathcal{R}(\rho_{PQ})) - S(\mathcal{R}(\rho'_{PQ})). \quad (19)$$

We present the proof of Lemma 1 in the SM.

To apply Lemma 1 to our setup, we let $\rho = \tilde{\lambda}$ and $\rho' = \tilde{\sigma}$ where $\tilde{\lambda} = \bar{U}\lambda\bar{U}^\dagger$ and $\tilde{\sigma} = \bar{U}\sigma\bar{U}^\dagger$ and where \bar{U} is a unitary obtained by removing the gates in U that are deep in the interior of the annulus ABC [Fig. 4(a-b)]. We then let P, Q be a partition of the annulus ABC of the form shown in Fig. 4, such that $P \subset A$ is sufficiently far away from the support of \bar{U} .² By construction, $\tilde{\lambda}$ and




FIG. 5. A schematic description of the construction of \mathcal{R} . The first step is the partial trace $\text{Tr}_{\bar{u}}$ over the support of \bar{U} , and the second step is an application of the Petz map $\Phi_{v \rightarrow v\bar{u}}^\sigma$ where $v = v_1 v_2$ and $\bar{u} = \bar{u}_1 \bar{u}_2$. The blue subsystem is \bar{u} , the support of \bar{U} .

$\tilde{\sigma}$ are indistinguishable on P . In Appendix E, we show that the two states are indistinguishable on Q as well [Eq. (E2)], thus fulfilling the premise of Lemma 1.

Below we will construct quantum channels $\mathcal{R} : Q \rightarrow \hat{Q}$ and $\mathcal{T} : \hat{Q} \rightarrow Q$, with $\hat{Q} \equiv Q \cup \bar{u}$ where \bar{u} is the support of \bar{U} . These will obey (18) with $\mathcal{R}(\tilde{\lambda}_{PQ}) = \lambda_{PQ\cup\bar{u}}$ and $\mathcal{R}(\tilde{\sigma}_{PQ}) = \sigma_{PQ\cup\bar{u}}$. Because $PQ\cup\bar{u} = Y$ and also $S(\tilde{\lambda}_{PQ}) = S(\tilde{\lambda}_{PQ})$ and $S(\tilde{\sigma}_{PQ}) = S(\tilde{\sigma}_{PQ})$, once we construct these channels, we can immediately deduce Eq. (16) from Lemma 1. This will then complete our proof of the bound (5), as explained earlier.

Now let us discuss our construction of \mathcal{R} and \mathcal{T} . These maps are constructed from compositions of \bar{U} , partial trace, and the Petz map [19].

In order to make the construction of \mathcal{R} and \mathcal{T} transparent, we will depict \bar{u} as two disks of smaller sizes, as shown in Fig. 4(c). Loosely speaking, \mathcal{R} removes the circuits in the disks and \mathcal{T} adds them back.

The map \mathcal{R} is constructed by applying a partial trace followed by a Petz map. The construction is best described by Fig. 5. In the first step, we trace out the region $Q \cap \bar{u}$. This step effectively removes the circuit \bar{U} , mapping $\tilde{\lambda}_{PQ}$ to $\lambda_{PQ\setminus\bar{u}}$ and $\tilde{\sigma}_{PQ}$ to $\sigma_{PQ\setminus\bar{u}}$. In the second step, we apply the Petz map $\Phi_{v \rightarrow v\bar{u}}^\sigma$. We show in the Supplemental Material, Eq. (E3), that this step extends $\lambda_{PQ\setminus\bar{u}}$ to $\lambda_{PQ\cup\bar{u}}$ and $\sigma_{PQ\setminus\bar{u}}$ to $\sigma_{PQ\cup\bar{u}}$:

$$\lambda_{PQ\cup\bar{u}} = \Phi_{v \rightarrow v\bar{u}}^\sigma(\lambda_{PQ\setminus\bar{u}}), \quad \sigma_{PQ\cup\bar{u}} = \Phi_{v \rightarrow v\bar{u}}^\sigma(\sigma_{PQ\setminus\bar{u}}). \quad (20)$$

Combining the two steps we see that \mathcal{R} maps $\tilde{\lambda}_{PQ}$ to $\lambda_{PQ\cup\bar{u}}$ and $\tilde{\sigma}_{PQ}$ to $\sigma_{PQ\cup\bar{u}}$, as required. As for the map \mathcal{T} , this can be constructed by simply applying \bar{U} and tracing out $\bar{u} \setminus Q$. Clearly these operations map $\lambda_{PQ\cup\bar{u}}$ to $\tilde{\lambda}_{PQ}$ and $\sigma_{PQ\cup\bar{u}}$ to $\tilde{\sigma}_{PQ}$, as required.

Discussion — We remark that our lower bound (5) can be easily generalized to the case where σ contains an anyon in the interior of the annulus. All that was needed

² More precisely, P must be at least two lattice spacings away from the support of \bar{U} ; see Fig. 14 in the Supplemental Material for details.

in our proof was the invariance of the TEE of σ under topology-preserving deformations of the annulus, a fact that remains true even in the presence of an anyon. In this case, we get a lower bound of $\gamma_0 = \log(\mathcal{D}/d_a)$ [16], assuming anyon a is placed in the interior.

We emphasize again that we have proven the lower bound (3) for any state obtained by a constant-depth circuit acting on a valid reference state, where in particular, the reference state could be any string-net ground state [7–9]. Using this result, we can argue heuristically that (3) holds for general 2D bosonic gapped ground states. The key point is that string-net ground states are believed to realize all “doubled” 2D topological phases obtained by stacking a bosonic topological phase onto its time-reversed partner. So one expects that for any 2D bosonic gapped ground state ρ , we should be able to write the doubled state $\rho \otimes \rho^*$ (realized in a bilayer geometry) as $\rho \otimes \rho^* = U\sigma U^\dagger$ for some valid reference state σ . Hence the lower bound (3) should apply to $\rho \otimes \rho^*$ and therefore also to ρ itself (since the TEE of $\rho \otimes \rho^*$ is exactly twice that of ρ , as is the total quantum dimension $\log \mathcal{D}$). However, a careful argument in this direction would require a treatment of *quasi-local* circuits U , i.e. circuits with decaying tails, which arise from the adiabatic flow of Hamiltonians [20, 21]. We leave this question to future work.

Our proof relies on Markov state techniques supported by strong subadditivity and the fact that constant-depth circuits can be truncated in certain ways without affecting the entropies of regions. Due to the generality of these techniques, we expect that similar lower bounds can be derived in a variety of other setups, including sys-

tems with defects or higher dimensional systems. Note, however, that the TEE can also be defined using the alternative partition in Ref. [1], for which it is unclear if our argument applies. It would be interesting to understand this partition better.

It is worth noting that our Definition 1 and bound (5) apply beyond the scope of area law states. As a concrete example, coupling an area law reference state to a hot surface (modeled by an identity density matrix) for a short period of time cannot decrease the TEE of the joint system. We also speculate that similar arguments may apply to show that the TEE of the 3D toric code at finite temperature [22] cannot decrease under a circuit.

An important question for future work is to understand how generically our bound (3) is *saturated*. That is, how much fine-tuning is required to obtain a spurious contribution to the TEE that does not decay for regions of increasing size? There are some hints in this direction [11, 13, 14, 23], but the general question remains open.

ACKNOWLEDGEMENTS

D.R. thanks Aram Harrow and Jonathan Sorce for discussion. D.R. acknowledges support from NTT (Grant AGMT DTD 9/24/20). This work was supported by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651442, M.L.; 652264, B.S.). T.C.L. was supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under the cooperative research agreement DE-SC0009919.

-
- [1] A. Kitaev and J. Preskill, Topological entanglement entropy, *Phys. Rev. Lett.* **96**, 110404 (2006).
 - [2] M. Levin and X.-G. Wen, Detecting topological order in a ground state wave function, *Phys. Rev. Lett.* **96**, 110405 (2006).
 - [3] S. V. Isakov, M. B. Hastings, and R. G. Melko, Topological entanglement entropy of a bose–hubbard spin liquid, *Nature Physics* **7**, 772 (2011).
 - [4] H.-C. Jiang, Z. Wang, and L. Balents, Identifying topological order by entanglement entropy, *Nature Physics* **8**, 902 (2012).
 - [5] A. Hamma, R. Ionicioiu, and P. Zanardi, Bipartite entanglement and entropic boundary law in lattice spin systems, *Physical Review A* **71**, 022315 (2005).
 - [6] A. Y. Kitaev, Fault-tolerant quantum computation by anyons, *Annals of Physics* **303**, 2 (2003), arXiv:quant-ph/9707021 [quant-ph].
 - [7] M. A. Levin and X.-G. Wen, String-net condensation: A physical mechanism for topological phases, *Phys. Rev. B* **71**, 045110 (2005).
 - [8] A. Hahn and R. Wolf, Generalized string-net model for unitary fusion categories without tetrahedral symmetry, *Phys. Rev. B* **102**, 115154 (2020).
 - [9] C.-H. Lin, M. Levin, and F. J. Burnell, Generalized string-net models: A thorough exposition, *Physical Review B* **103**, 195155 (2021).
 - [10] S. Bravyi, Unpublished, (2008).
 - [11] L. Zou and J. Haah, Spurious long-range entanglement and replica correlation length, *Phys. Rev. B* **94**, 075151 (2016).
 - [12] J. Cano, T. L. Hughes, and M. Mulligan, Interactions along an entanglement cut in 2+ 1 d abelian topological phases, *Physical Review B* **92**, 075104 (2015).
 - [13] D. J. Williamson, A. Dua, and M. Cheng, Spurious topological entanglement entropy from subsystem symmetries, *Phys. Rev. Lett.* **122**, 140506 (2019).
 - [14] D. T. Stephen, H. Dreyer, M. Iqbal, and N. Schuch, Detecting subsystem symmetry protected topological order via entanglement entropy, *Physical Review B* **100**, 115112 (2019).
 - [15] K. Kato and F. G. S. L. Brandão, Toy model of boundary states with spurious topological entanglement entropy, *Phys. Rev. Research* **2**, 032005 (2020).
 - [16] B. Shi, K. Kato, and I. H. Kim, Fusion rules from entanglement, *Annals of Physics* **418**, 168164 (2020).
 - [17] E. H. Lieb and M. B. Ruskai, Proof of the strong subadditivity of quantum-mechanical entropy,

Journal of Mathematical Physics **14**, 1938 (1973), <https://doi.org/10.1063/1.1666274>.

- [18] B. Shi, *Anyon theory in gapped many-body systems from entanglement*, Ph.D. thesis (2020).
- [19] D. Petz, Sufficiency of channels over von Neumann algebras, *Q. J. Math.* **39**, 97 (1988).
- [20] M. B. Hastings and X.-G. Wen, Quasiadiabatic continuation of quantum states: The stability of topological ground-state degeneracy and emergent gauge invariance, *Phys. Rev. B* **72**, 045141 (2005), [arXiv:cond-mat/0503554 \[cond-mat.str-el\]](https://arxiv.org/abs/cond-mat/0503554).
- [21] B. Nachtergaele, R. Sims, and A. Young, Quasi-locality bounds for quantum lattice systems. i. lieb-robinson bounds, quasi-local maps, and spectral flow automorphisms, *Journal of Mathematical Physics* **60**, 061101 (2019).
- [22] C. Castelnovo and C. Chamon, Topological order in a three-dimensional toric code at finite temperature, *Phys. Rev. B* **78**, 155120 (2008).
- [23] S. F. E. Oliviero, L. Leone, Y. Zhou, and A. Hamma, Stability of topological purity under random local unitaries, *SciPost Physics* **12**, 096 (2022).
- [24] F. Hiai, M. Mosonyu, D. Petz, and C. Bény, Quantum f-divergences and error correction, *Reviews in Mathematical Physics* **23**, 691 (2011), <https://doi.org/10.1142/S0129055X11004412>.
- [25] I. H. Kim, Entropy scaling law and the quantum marginal problem, *Physical Review X* **11**, 021039 (2021).

Appendix A: Proof of Lemma 1

Here we present the proof of Lemma 1.

Proof. Define $\tilde{\rho}_{P\hat{Q}} \equiv \mathcal{R}(\rho_{PQ})$ and $\tilde{\rho}'_{P\hat{Q}} \equiv \mathcal{R}(\rho'_{PQ})$. Then $\tilde{\rho}_{\hat{Q}} = \tilde{\rho}'_{\hat{Q}}$, because $\rho_Q = \rho'_Q$. Likewise, $\tilde{\rho}_P = \tilde{\rho}'_P$. The mutual information obeys

$$I(P : Q)_\rho = I(P : \hat{Q})_{\tilde{\rho}}. \quad (\text{A1})$$

To see this, note that mutual information is nonincreasing under quantum channels acting on one party, which we can use to establish two inequalities between $I(P : Q)_\rho$ and $I(P : \hat{Q})_{\tilde{\rho}}$ in two opposite directions (using channels \mathcal{R} and \mathcal{T} respectively), yielding the desired equality. Deriving the same equality for ρ' and taking the difference, we find

$$S(\rho_{PQ}) - S(\rho'_{PQ}) = S(\tilde{\rho}_{P\hat{Q}}) - S(\tilde{\rho}'_{P\hat{Q}}), \quad (\text{A2})$$

which completes the proof. \square

Appendix B: Quantum Markov chains

In this Section, we collect various known facts about quantum entropies and Markov states, used extensively in this paper. We begin with the strong subadditivity of entropy (SSA), which states that [17]

$$I(A : C|B)_\rho \geq 0 \quad (\text{B1})$$

for any tripartite density matrix ρ_{ABC} . A straightforward application of SSA leads to the following inequalities,

$$\begin{aligned} I(AA' : C|B)_\rho &\geq I(A' : C|B)_\rho, \\ I(AA' : CC'|B)_\rho &\geq I(A : C|A'BC')_\rho, \end{aligned} \quad (\text{B2})$$

for any density matrix ρ .

If a tripartite quantum state ρ_{ABC} satisfies the SSA with equality, i.e., $I(A : C|B)_\rho = 0$, such a state is called a quantum Markov chain [19]. An important property of quantum Markov chains is the existence of the *Petz map*. If ρ_{ABC} is a quantum Markov chain, there is a quantum channel $\Phi_{B \rightarrow BC}^\rho : \mathcal{B}(\mathcal{H}_B) \rightarrow \mathcal{B}(\mathcal{H}_{BC})$ such that

$$\mathcal{I}_A \otimes \Phi_{B \rightarrow BC}^\rho(\rho_{AB}) = \rho_{ABC}, \quad (\text{B3})$$

where $\mathcal{B}(\mathcal{H})$ is the space of bounded operators acting on \mathcal{H} and \mathcal{I}_A is the identity channel on A . We remark that it is customary to omit the superoperator \mathcal{I}_A if the domain of the Petz map is obvious from context, a convention that we shall adhere to, i.e.,

$$\Phi_{B \rightarrow BC}^\rho(\rho_{AB}) = \rho_{ABC}. \quad (\text{B4})$$

We also remark that (B4) holds *if and only if* $I(A : C|B)_\rho = 0$. In particular, (B4) implies that ρ_{ABC} is a quantum Markov chain.

The Petz map depends only on the reduced density matrix ρ_{BC} and exists independent of whether ρ_B is invertible or not [24]. If ρ_B is invertible, it admits a particularly simple form:

$$\Phi_{B \rightarrow BC}^\rho(\cdot) = \rho_{BC}^{\frac{1}{2}} \rho_B^{-\frac{1}{2}} (\cdot) \rho_B^{-\frac{1}{2}} \rho_{BC}^{\frac{1}{2}}. \quad (\text{B5})$$

Nevertheless, the precise form of the Petz map will not be important in our analysis.

We remark that the property (B4) of the Petz map directly implies that two locally indistinguishable quantum Markov chains are globally the same. Without loss of generality, let ρ and σ be quantum Markov chains, i.e., $I(A : C|B)_\rho = I(A : C|B)_\sigma = 0$. This implies

$$\begin{aligned} \rho_{ABC} &= \Phi_{B \rightarrow BC}^\rho(\rho_{AB}), \\ \sigma_{ABC} &= \Phi_{B \rightarrow BC}^\sigma(\sigma_{AB}). \end{aligned} \quad (\text{B6})$$

If $\rho_{AB} = \sigma_{AB}$ and $\rho_{BC} = \sigma_{BC}$, it follows from the property of the Petz map that $\rho_{ABC} = \sigma_{ABC}$. We therefore conclude the following.

Lemma 2. *Let ρ_{ABC} and σ_{ABC} be states such that $I(A : C|B)_\rho = I(A : C|B)_\sigma = 0$. If $\rho_{AB} = \sigma_{AB}$ and $\rho_{BC} = \sigma_{BC}$, then $\rho_{ABC} = \sigma_{ABC}$.*

The notion of a quantum Markov chain can be straightforwardly generalized to n -partite states as well, for $n > 3$. Let $\{X_1, \dots, X_n\}$ be an *ordered set* of subsystems that partition X . Intuitively, one can think of these subsystems as ‘‘spins’’ arranged along a one-dimensional chain, with the index representing the location of the spin.




FIG. 6. We call three non-empty subsystems A , B , and C “chain-like” when they form three contiguous subsets as in the example above.

For the remainder of the Appendix, we will frequently consider three non-empty subsystems A , B , and C given by contiguous subsets of this spin chain. In such cases, we say that A , B , and C together are *chain-like* subsystems [Fig. 6].

Using the notion of chain-like subsystems, we can now present a generalization of quantum Markov chains to n -partite systems.

Definition 2. A quantum state σ over ordered subsystems $\{X_1, \dots, X_n\}$ is a Markov chain if $I(A : C|B)_\sigma = 0$ for any chain-like subsystems A, B, C .

Appendix C: Locally Quantum Markov Chains

We can also define a *locally quantum Markov chain*, or locally Markov chain for short, by only demanding the conditional mutual information constraints on proper subsystems.

Definition 3. A quantum state σ over $\{X_1, \dots, X_n\}$ is a locally Markov chain if $I(A : C|B)_\sigma = 0$ for any chain-like subsystems A, B, C such that ABC is a proper subset of X .

A locally Markov chain need not satisfy the conditional mutual information constraint globally. That is, $I(A : C|B) \neq 0$ is allowed when $ABC = \cup_{i=1}^n X_i$.

However, this violation must be a constant for any partition of X into chain-like subsystems. In fact, we show below that having a constant (possibly zero) violation is a necessary and sufficient condition for a state to be locally Markov: that is, a state σ over $\cup_{i=1}^n X_i$ is locally Markov if and only if $I(A : C|B)_\sigma$ is a constant for any chain-like A, B , and C that partition X .

Proposition 1. Let σ be a state over $X = \cup_{i=1}^n X_i$, where $n \geq 4$. Then the following statements are equivalent:

1. The state σ is a locally Markov chain over X , i.e., $I(A : C|B)_\sigma = 0$ for every choice of chain-like subsystems A, B , and C that form a proper subset of X .
2. $I(A : C|B)_\sigma$ is a constant for any chain-like A, B , and C that partition X .

Proof. The equivalence of the two statements hinges on a simple identity. Consider contiguous non-empty subsystems A, B, C , and D that partition X ; see Fig. 7. Here




FIG. 7. Partition of X into contiguous non-empty subsystems A, B, C , and D .

is the key observation:

$$\begin{aligned} I(A : C|B)_\sigma = 0 &\Leftrightarrow I(A : CD|B)_\sigma = I(A : D|BC)_\sigma, \\ I(B : D|C)_\sigma = 0 &\Leftrightarrow I(AB : D|C)_\sigma = I(A : D|BC)_\sigma. \end{aligned} \quad (\text{C1})$$

These facts can be shown straightforwardly by expanding the conditional mutual information in terms of von Neumann entropies. As we explain below, the left and the right hand side of Eq. (C1) can be related to 1 and 2 respectively, leading to the main claim.

We first show 1 \implies 2. From 1, $I(A : C|B)_\sigma = 0$ and $I(B : D|C)_\sigma = 0$ for any non-empty and contiguous subsystems A, B, C , and D that partition X . By Eq. (C1), $I(A : CD|B)_\sigma = I(A : D|BC)_\sigma = I(AB : D|C)_\sigma$.

These identities can be understood better with the help of Fig. 7. Consider two different partitions of X , one in terms of chain-like subsystems A, B, CD , and the other in terms of chain-like subsystems A, BC, D . In both cases, X is partitioned into three parts, the subsystem on the left, middle, and right, which we denote as L, M , and R , respectively. Viewed this way, the identity $I(A : CD|B)_\sigma = I(A : D|BC)_\sigma$ means that the conditional mutual information $I(L : R|M)$ for $X = LMR$ is invariant under shifting the boundary between the middle and the right subsystem, as long as both subsystems remain non-empty.

Similarly, the identity $I(A : D|BC)_\sigma = I(AB : D|C)_\sigma$ means that the conditional mutual information $I(L : R|M)$ is invariant under shifting the boundary between the left and the middle subsystem, again assuming both subsystems remain non-empty. Together, we see that $I(L : R|M)_\sigma$ is invariant under arbitrary shifts of both boundaries, assuming the subsystems L, R , and M all remain non-empty. Since any partition of X into chain-like subsystems can be obtained from an appropriate shift of the two boundaries, 2 follows.

Now we show 2 \implies 1. Without loss of generality, consider chain-like subsystems A, B , and C , which together form a proper subset of X . Because ABC is a proper subset, either $X_1 \not\subset A$ or $X_n \not\subset C$.

Consider the latter case first. Without loss of generality, let $A = \cup_{k=i}^j X_k$. We can define $A' = \cup_{k=1}^j X_k \supset A$. Consider a partition of X into chain-like subsystems A', B, CD and also A', BC, D . From 2, we see that

$$I(A' : CD|B)_\sigma = I(A' : D|BC)_\sigma. \quad (\text{C2})$$

From (C1) we obtain

$$I(A' : C|B)_\sigma = 0. \quad (\text{C3})$$

Since $A \subset A'$, from Eq. (B2) we deduce that $I(A : C|B)_\sigma = 0$. The case in which $X_1 \not\subset A$ can be dealt with in a similar way, establishing 2 \implies 1.

□

We now show that, given a locally Markov chain σ over $n \geq 4$ subsystems, there is a unique Markov chain state (in the sense of Definition 2) that is locally indistinguishable from σ over every connected proper subsystem. By connected, we mean the subsystem is a union of contiguous subsystems, e.g., $\{X_i, X_{i+1}, \dots, X_{j-1}, X_j\}$. This state, which we denote as τ , is defined in the following recursive fashion

$$\tau_{k+1} = \Phi_{X_k \rightarrow X_k X_{k+1}}^\sigma(\tau_k), \quad \forall k \in \{2, \dots, n-1\} \quad (\text{C4})$$

where $\tau := \tau_n$, $\tau_2 := \sigma_{X_1 X_2}$, and $\Phi_{X_k \rightarrow X_k X_{k+1}}^\sigma$ is the Petz map defined in terms of σ .

Here are some useful properties of τ , which we collate for the readers' convenience.³

- $\tau_Y = \sigma_Y$ for any connected proper subsystem Y . (Proposition 2)
- τ is a Markov chain. (Proposition 3)
- Let A, B , and C be chain-like subsystems that partition $\{X_1, \dots, X_n\}$. Then τ is the maximum-entropy state indistinguishable from σ over AB and BC . (Proposition 4)

Because τ is the unique maximum-entropy state indistinguishable from σ over every connected proper subsystem, there is a sense in which τ is canonical. We thus refer to it as the *canonical Markov chain* from now on. A more formal definition is stated below.

Definition 4. Let σ be a locally Markov chain over an ordered set $\{X_1, \dots, X_n\}$, where $n \geq 4$. The canonical Markov chain τ associated to σ and $\{X_1, \dots, X_n\}$ is given by

$$\tau_{k+1} = \Phi_{X_k \rightarrow X_k X_{k+1}}^\sigma(\tau_k), \quad \forall k \in \{2, \dots, n-1\} \quad (\text{C5})$$

where $\tau := \tau_n$ and $\tau_2 := \sigma_{X_1 X_2}$.

We now prove properties of canonical Markov chains. Throughout these proofs, τ will represent the canonical Markov chain associated to a locally Markov chain σ , defined over $\{X_1, \dots, X_n\}$.

Proposition 2. For every connected proper subsystem $Y \subset X$, $\tau_Y = \sigma_Y$.

Note $\tau_Y = \text{Tr}_{X \setminus Y} \tau$ refers to the restriction of τ on X to $Y \subset X$, rather than the canonical Markov chain defined with respect to σ_Y .

³ Some of these properties follow from [25, Section 5.2], but we provide self-contained proofs here for completeness.

Proof. We first prove the claim for $Y = \cup_{i=2}^n X_i$. Because σ is a locally Markov chain, $I(X_k : X_2 \dots X_{k-2} | X_{k-1}) = 0$ for every $4 \leq k \leq n$. Therefore, one can obtain σ_Y by repeatedly applying the Petz map $\Phi_{X_k \rightarrow X_k X_{k+1}}^\sigma$, from $k = 3$ to $n-1$:

$$\sigma_Y = \Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma \circ \dots \circ \Phi_{X_3 \rightarrow X_3 X_4}^\sigma(\sigma_{X_2 X_3}). \quad (\text{C6})$$

Now let us find an expression for $\tau_Y = \text{Tr}_{X_1}(\tau)$. Because all the channels appearing in the definition of τ [Definition 4] act trivially on X_1 , the partial trace over X_1 can be applied before applying those channels. The resulting expression is

$$\tau_Y = \Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma \circ \dots \circ \Phi_{X_2 \rightarrow X_2 X_3}^\sigma(\sigma_{X_2}). \quad (\text{C7})$$

Since $\Phi_{X_2 \rightarrow X_2 X_3}^\sigma(\sigma_{X_2}) = \sigma_{X_2 X_3}$, we derive $\tau_Y = \sigma_Y$.

Now we prove the claim for $Y' = \cup_{i=1}^{n-1} X_i$. Note that, because σ is a locally Markov chain, we can relate τ and σ in the following way:

$$\tau = \Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma(\sigma_{X_1 \dots X_{n-1}}), \quad (\text{C8})$$

Next we convert $\sigma_{X_1 \dots X_{n-1}}$ into another form, using the fact that $I(X_1 : X_3 \dots X_n | X_2)_\sigma = 0$:

$$\sigma_{X_1 \dots X_{n-1}} = \Phi_{X_2 \rightarrow X_1 X_2}^\sigma(\sigma_{X_2 \dots X_{n-1}}). \quad (\text{C9})$$

Note that the channel appearing in (C9) and the channel appearing in (C8) act on disjoint subsystems, and as such, they can be exchanged. Therefore,

$$\begin{aligned} \tau &= \Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma(\Phi_{X_2 \rightarrow X_1 X_2}^\sigma(\sigma_{X_2 \dots X_{n-1}})) \\ &= \Phi_{X_2 \rightarrow X_1 X_2}^\sigma(\Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma(\sigma_{X_2 \dots X_{n-1}})) \\ &= \Phi_{X_2 \rightarrow X_1 X_2}^\sigma(\sigma_{X_2 \dots X_n}), \end{aligned} \quad (\text{C10})$$

where in the last line we used the fact that $I(X_n : X_2 \dots X_{n-2} | X_{n-1})_\sigma = 0$.

Tracing out X_n , we obtain

$$\begin{aligned} \tau_{Y'} &= \Phi_{X_2 \rightarrow X_1 X_2}^\sigma(\sigma_{X_2 \dots X_{n-1}}) \\ &= \sigma_{Y'}. \end{aligned} \quad (\text{C11})$$

Since any proper connected subsystem is included in Y or Y' , the main claim follows. □

Proposition 3. τ is a quantum Markov chain.

Proof. By Proposition 2, τ is indistinguishable from σ over any proper connected subsystem. Since σ is a locally Markov chain, it follows that $I(A : C | B)_\tau = 0$ for any chain-like subsystems A, B , and C forming a proper subset of X . Therefore, to prove the main claim, it suffices to prove $I(A : C | B)_\tau = 0$ for chain-like subsystems A, B , and C that partition X .

Choose $C = X_n$ and $B = X_{n-1}$. Because σ is a locally Markov chain, $\tau = \Phi_{X_{n-1} \rightarrow X_{n-1} X_n}^\sigma(\sigma_{X_1 \dots X_{n-1}})$. Therefore,

$$\begin{aligned} \tau_{ABC} &= \Phi_{B \rightarrow BC}^\sigma(\sigma_{AB}) \\ &= \Phi_{B \rightarrow BC}^\tau(\tau_{AB}), \end{aligned} \quad (\text{C12})$$




FIG. 8. A chain-like region ABC in 2D.

where we used the fact that $\sigma_{AB} = \tau_{AB}$ and $\sigma_{BC} = \tau_{BC}$, consequences of Proposition 2. Therefore, $I(A : C|B)_\tau = 0$ for this particular choice of ABC .

Note that τ is in fact also a locally Markov chain, because it is indistinguishable from σ over any proper connected subsystem and σ is a locally Markov chain. By Proposition 1, $I(A : C|B)_\tau$ is a constant for any chain-like A, B , and C that partition X . Moreover, it was shown above that this constant is 0. Therefore, $I(A : C|B)_\tau = 0$ for any chain-like subsystems A, B , and C . This completes the proof. \square

Proposition 4. *Let A, B , and C be chain-like subsystems that partition $\{X_1, \dots, X_n\}$. Then τ is the maximum-entropy state indistinguishable from σ over AB and BC .*

Proof. Suppose there exists another such state, say τ' . Partition X into ABC which are chain-like. $I(A : C|B)_\tau = 0$. By the maximum-entropy condition, $S(\tau') \geq S(\tau)$. Because τ and τ' are indistinguishable over AB, BC , and B , we conclude $I(A : C|B)_{\tau'} \leq 0$. However, by SSA $I(A : C|B)_{\tau'} \geq 0$. Hence, $I(A : C|B)_{\tau'} = 0$.

Therefore, we obtain the following expression for τ and τ' :

$$\begin{aligned} \tau &= \Phi_{B \rightarrow BC}^\tau(\tau_{AB}) \\ \tau' &= \Phi_{B \rightarrow BC}^{\tau'}(\tau'_{AB}). \end{aligned} \quad (\text{C13})$$

Because $\tau_{AB} = \tau'_{AB}$ and $\tau_{BC} = \tau'_{BC}$, we deduce $\tau = \tau'$. This completes the proof. \square

Appendix D: Chain-like subsystems in 2D

In this Appendix, we prove a certain fact about *chain-like* subsystems in 2D. Roughly speaking, if three subsystems A, B , and C are arranged in a form shown in Fig. 8, we say these subsystems are chain-like. Our main claim is that for such subsystems $I(A : C|B)_\sigma = 0$. For concreteness, we provide a precise definition and a statement below.

Definition 5 (Chain-like regions in 2D). *Subsystems A, B, C of a 2D disk (or plane) form a chain-like region if their relation is topologically equivalent to the regions A, B and C shown in Fig. 8.*

Proposition 5 (Chain-like regions in 2D are Markov). *Let σ be a reference state as in Definition 1 of the main text. For any chain-like region ABC in 2D, we have $I(A : C|B)_\sigma = 0$.*




FIG. 9. We can view an annulus in 2D as a 1D “spin chain.” The shown partition defines a spin chain over the ordered subsystems $\{X_1, X_2, X_3, X_4, X_5\}$.

Proof. To prove this claim, we shall identify the chain-like region ABC as a subset of an annulus. Specifically, consider an annulus shown in Fig. 9 such that A, B , and C are identified with the upper connected components of X_2, X_3 , and X_4 , respectively.

Note that, by the definition of the reference state [Definition 1], the state σ over $\{X_1, X_2, X_3, X_4, X_5\}$ has a constant value of $I(A : C|B)_\sigma$ for any chain-like A, B , and C that partition $X = \cup_{i=1}^5 X_i$. By Proposition 1, we thus conclude that σ is a locally Markov chain over $\{X_1, X_2, X_3, X_4, X_5\}$. In particular,

$$I(X_2 : X_4|X_3)_\sigma = 0. \quad (\text{D1})$$

Because the reference state has vanishing mutual information between any two subsystems that are not adjacent to each other, the state over the upper and the lower components factorize. Therefore,

$$I(A : C|B)_\sigma = 0. \quad (\text{D2})$$

\square

Appendix E: Max-entropy state on a 2D annulus

The argument presented in the main text was based on three facts. Specifically, we considered a partition of an annulus ABC into P and $Q = ABC \setminus P$ such that $P \subset A$ is sufficiently far from the support of the unitary U [Fig. 10].⁴ The density matrix λ was then defined as the maximum-entropy state consistent with σ over the past light cones of AB and BC [Fig. 11]. Denoting the support of U as u , we claimed that

$$S(\lambda_{PQ \cup u}) - S(\sigma_{PQ \cup u}) = 2\gamma_0, \quad (\text{E1})$$

$$\bar{\lambda}_P = \bar{\sigma}_P, \bar{\lambda}_Q = \bar{\sigma}_Q. \quad (\text{E2})$$

Moreover, we also claimed that

$$\begin{aligned} \lambda_{PQ \cup \bar{u}} &= \Phi_{v \rightarrow v\bar{u}}^\sigma(\lambda_{PQ \setminus \bar{u}}), \\ \sigma_{PQ \cup \bar{u}} &= \Phi_{v \rightarrow v\bar{u}}^\sigma(\sigma_{PQ \setminus \bar{u}}), \end{aligned} \quad (\text{E3})$$

⁴ More specifically, we assumed that P must be at least two lattice spacing away from the support of \bar{U} [Fig. 14]. Because the support of \bar{U} is included in the support of U , the requisite condition can be fulfilled by demanding P is at least two lattice spacing away from the support of U .




FIG. 10. Two different partitions of the annulus, one into ABC (a) and one into PQ (b). $P \subset A$ is chosen so that it is sufficiently far from the support of the unitary U (blue).




FIG. 11. (a) Past light cone of AB (red). (b) Past light cone of BC (green). (c) The enlarged annulus $PQ\cup u$ can be partitioned into the new subsystems \tilde{A}, \tilde{B} , and \tilde{C} . The past light cone of AB and BC is $\tilde{A}\tilde{B}$ and $\tilde{B}\tilde{C}$, respectively. The blue region is the support of U .

where \bar{U} is the unitary obtained by removing the gates in U that are deep inside the annulus, \bar{u} is its support, and v is the region in the annulus that surrounds \bar{u} [Fig. 5 of the main text]. Below, we prove these facts.

The first two facts Eq. (E1) and (E2) follow straightforwardly from an observation that the annulus $PQ\cup u$ can be partitioned into a set of subsystems, after which the reference state σ over these subsystems can be viewed as a locally Markov chain [Appendix C]. This implies that λ is the canonical Markov chain [Definition 4] associated with the reference state, with respect to this partition. Then from the properties of the canonical Markov chain, Eq. (E1) and (E2) both follow.

More specifically, consider the partition shown in Fig. 11(c). Here \tilde{B} is the past light cone of B . The remaining subsystems \tilde{A} and \tilde{C} are obtained by subtracting \tilde{B} from the past light cone of AB and BC , respectively. Together, \tilde{A}, \tilde{B} , and \tilde{C} partition the enlarged annulus $ABC\cup u = PQ\cup u$.

By our assumption, P is separated from the support of U . Since the support of U includes $\tilde{B}\tilde{C}$, $\tilde{B}\tilde{C}$ is separated from P . Therefore, the enlarged annulus can be further partitioned into $P, \tilde{A}\setminus P, \tilde{B}, \tilde{C}$ [Fig. 12]. With respect to this partition, we now invoke facts about locally Markov chain [Appendix C]. To draw a parallel with the discussion in Appendix C, let us momentarily denote $P, \tilde{A}\setminus P, \tilde{B}, \tilde{C}$ as X_1, X_2, X_3, X_4 , respectively. Because we assumed that the reference state σ has a constant TEE, it immediately follows that $\sigma_{X_1 X_2 X_3 X_4}$ is a locally Markov




FIG. 12. Partition of an enlarged annulus $PQ\cup u$ into $P, \tilde{A}\setminus P, \tilde{B}$, and \tilde{C} .




FIG. 13. Partition of an enlarged annulus $PQ\cup u$ into $P, Q_1, \tilde{A}\setminus(PQ_1), \tilde{B}$, and \tilde{C} .

chain with respect to the partition $\{X_1, X_2, X_3, X_4\}$.

Since σ is a locally Markov chain, there is also a canonical Markov chain associated to σ . Crucially, this canonical Markov chain is precisely λ . Recall that the state λ was defined as the maximum-entropy state consistent with the past light cone of AB and BC . Thus, by Proposition 4, we immediately see that λ is the canonical Markov chain associated with σ , with respect to the partition $\{X_1, X_2, X_3, X_4\}$.

Now that we established that λ is the canonical Markov chain, we can use our machinery to prove Eqs. (E1) and (E2). In particular, because the canonical Markov chain is a Markov chain [Proposition 3], we know $I(X_1 X_2 : X_4 | X_3)_\lambda = 0$. Then (E1) follows from the fact that $I(X_1 X_2 : X_4 | X_3)_\sigma = 2\gamma_0$, and that $\lambda_{X_1 X_2 X_3} = \sigma_{X_1 X_2 X_3}$ and $\lambda_{X_3 X_4} = \sigma_{X_3 X_4}$ [Proposition 2]. Similarly, to establish (E2), we use Proposition 2 to deduce $\lambda_{X_1} = \sigma_{X_1}$ and $\lambda_{X_2 X_3 X_4} = \sigma_{X_2 X_3 X_4}$. Then since X_1 is the past light cone of P and $X_2 X_3 X_4$ includes the past light cone of Q , we derive (E2).

The proof of (E3) involves two different partitions of the enlarged annulus $PQ\cup u$. The first partition is described in Fig. 13, which is obtained by dividing $\tilde{A}\setminus P$ further into Q_1 and $\tilde{A}\setminus(PQ_1)$. The second partition is based on the subsystems defined in terms of \bar{U} , which is the unitary obtained from U by removing the gates deep in the annulus. Provided that the annulus is sufficiently thick, the support of \bar{U} can be split into two non-overlapping disks, each of which can be furthermore surrounded by non-overlapping disks lying inside the annulus [Fig. 14(a-b)].

We use the first partition to establish the following fact:

$$\lambda_{PQ\cup\bar{u}} = \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q\cup\bar{u}}). \quad (\text{E4})$$




FIG. 14. (a) If the annulus is sufficiently thick, upon removing the gates deep inside the annulus, we obtain a unitary \bar{U} acting on two non-overlapping disks (blue). Furthermore, one can choose two disconnected disks in the annulus that surround each disk (orange). Figure (b) can be obtained from (a) by an appropriate regrouping of sites and topologically equivalent re-drawing. Here $Q = Q_1 Q_2$, $\bar{u} = \bar{u}_1 \bar{u}_2$, and $v = v_1 v_2$. To be able to partition the annulus this way, the support of \bar{U} must be at least two lattice spacing away from P .

We can view λ as the canonical Markov chain associated with the reference state σ with respect to the partition $\{X_1, X_2, X_3, X_4, X_5\} = \{P, Q_1, \tilde{A} \setminus (PQ_1), \tilde{B}, \tilde{C}\}$. This implies that λ is a Markov chain satisfying $I(X_1 : X_3 X_4 X_5 | X_2)_\lambda = 0$ [Proposition 3]. Furthermore, λ and σ are indistinguishable over $X_1 X_2 = PQ_1$ and $X_2 X_3 X_4 X_5 = \tilde{A} \tilde{B} \tilde{C} P = Q \cup \bar{u}$ [Proposition 2]. Eq. (E4) follows directly from these facts.

We use the second partition to prove the following fact:

$$I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma = 0. \quad (\text{E5})$$

To see why this is true, note that

$$\begin{aligned} I(\bar{u}_1 : Q \setminus (v_1 \bar{u}) | v_1)_\sigma &= 0, \\ I(\bar{u}_2 : (Q \cup \bar{u}_1) \setminus (v_2 \bar{u}_2) | v_2)_\sigma &= 0, \end{aligned} \quad (\text{E6})$$

because \bar{u}_1, v_1 , and $Q \setminus (v_1 \bar{u})$ is chain-like and so is \bar{u}_2, v_2 , and $(Q \cup \bar{u}_1) \setminus (v_2 \bar{u}_2)$ [Proposition 5]. More explicitly, these equations can be written as

$$\begin{aligned} I(\bar{u}_1 : Q \setminus (v_1 \bar{u}) | v_1)_\sigma &= \\ S(\sigma_{\bar{u}_1 v_1}) + S(\sigma_{Q \setminus \bar{u}}) - S(\sigma_{v_1}) - S(\sigma_{(Q \cup \bar{u}_1) \setminus \bar{u}_2}) &= 0 \end{aligned} \quad (\text{E7})$$

and

$$\begin{aligned} I(\bar{u}_2 : (Q \cup \bar{u}_1) \setminus (v_2 \bar{u}_2) | v_2)_\sigma &= \\ S(\sigma_{\bar{u}_2 v_2}) + S(\sigma_{(Q \cup \bar{u}_1) \setminus \bar{u}_2}) - S(\sigma_{v_2}) - S(\sigma_{(Q \cup \bar{u})}) &= 0. \end{aligned} \quad (\text{E8})$$

Summing the two, we obtain

$$I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma + I(v_1 \bar{u}_1 : v_2 \bar{u}_2) - I(v_1 : v_2) = 0. \quad (\text{E9})$$

Because the mutual information between two non-adjacent subsystems is zero, we conclude $I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma = 0$.

Combining these two facts, we now prove (E3), starting with the first identity. First note that, upon taking a partial trace on \bar{u} , we get

$$\begin{aligned} \text{Tr}_{\bar{u}}(\lambda_{PQ \cup \bar{u}}) &= \text{Tr}_{\bar{u}}(\Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q \cup \bar{u}})) \\ &= \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\text{Tr}_{\bar{u}}(\sigma_{Q \cup \bar{u}})) \\ &= \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q \setminus \bar{u}}). \end{aligned} \quad (\text{E10})$$

On the second line, the map $\text{Tr}_{\bar{u}}$ and $\Phi_{Q_1 \rightarrow PQ_1}^\sigma$ can be exchanged because they act on disjoint subsystems. Therefore, we obtain

$$\lambda_{PQ \setminus \bar{u}} = \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q \setminus \bar{u}}). \quad (\text{E11})$$

Next note that, by construction, the Petz maps $\Phi_{v \rightarrow v\bar{u}}^\sigma$ and $\Phi_{Q_1 \rightarrow PQ_1}^\sigma$ act on disjoint subsystems. As such, their order can be exchanged. We therefore obtain

$$\begin{aligned} \Phi_{v \rightarrow v\bar{u}}^\sigma(\lambda_{PQ \setminus \bar{u}}) &= \Phi_{v \rightarrow v\bar{u}}^\sigma(\Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q \setminus \bar{u}})) \\ &= \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\Phi_{v \rightarrow v\bar{u}}^\sigma(\sigma_{Q \setminus \bar{u}})) \\ &= \Phi_{Q_1 \rightarrow PQ_1}^\sigma(\sigma_{Q \cup \bar{u}}) \\ &= \lambda_{PQ \cup \bar{u}}. \end{aligned} \quad (\text{E12})$$

In the second line we exchanged the order of the channels. In the third and the fourth line, we used (E5) and (E4), respectively. This establishes the first identity in (E3).

Now we prove the second identity in (E3). The key point is that

$$I(\bar{u} : PQ \setminus (v\bar{u}) | v)_\sigma = 0, \quad (\text{E13})$$

which follows from (E5) and the invariance of TEE under topology-preserving deformations. Specifically, note the following identity

$$\begin{aligned} I(\bar{u} : PQ \setminus (v\bar{u}) | v)_\sigma - I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma &= \\ = I(P : Q_2 | Q_1)_\sigma - I(P : Q_2 \setminus \bar{u} | Q_1)_\sigma &= 0. \end{aligned} \quad (\text{E14})$$

The equivalence of the first and the second line follows from the definition of the conditional mutual information. More specifically, note the following more explicit expressions for the two conditional mutual information quantities in the first line:

$$\begin{aligned} I(\bar{u} : PQ \setminus (v\bar{u}) | v)_\sigma &= \\ = S(\sigma_{\bar{u}v}) + S(\sigma_{PQ \setminus \bar{u}}) - S(\sigma_v) - S(\sigma_{PQ}) &= 0 \end{aligned} \quad (\text{E15})$$

and

$$\begin{aligned} I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma &= \\ = S(\sigma_{\bar{u}v}) + S(\sigma_{Q \setminus \bar{u}}) - S(\sigma_v) - S(\sigma_Q). &= 0 \end{aligned} \quad (\text{E16})$$

Taking the difference between the two, we obtain

$$\begin{aligned} I(\bar{u} : PQ \setminus (v\bar{u}) | v)_\sigma - I(\bar{u} : Q \setminus (v\bar{u}) | v)_\sigma &= \\ = (S(\sigma_Q) - S(\sigma_{PQ})) - (S(\sigma_{Q \setminus \bar{u}}) - S(\sigma_{PQ \setminus \bar{u}})). &= 0 \end{aligned} \quad (\text{E17})$$

By adding $S(\sigma_{PQ_1}) - S(\sigma_{Q_1})$ for each paranthesis (so that they cancel out each other), we obtain the second line of (E14). The last line in (E14) follows from the fact that TEE is a constant. The second identity in (E3) is a direct consequence of (E13). This completes the proof.




FIG. 15. Topologically equivalent but geometrically distinct partitions of annulus X . (a) X is ABC . (b) X is $A'B'C'$.

Appendix F: Canonical nature of max-entropy state

This appendix establishes further properties of the max-entropy state, but we emphasize our results here are not necessary for the argument in the main text nor in the other appendices.

The motivating question is as follows. Consider two topologically equivalent partitions of a chosen annulus X as $X = ABC$ and $X = A'B'C'$ as in Fig. 15. Is it true that the max-entropy state (λ_X) consistent with the reference state σ on AB and BC must be the same as the max-entropy state (λ'_X) consistent with the reference state σ on $A'B'$ and $B'C'$? We argue that the answer is yes for an appropriate notion of topological equivalence that we explain below. In particular, $A'B'C'$ need *not* be obtainable from ABC by using a quasi-1D approach (that is, by refining the partition only in the “angular direction” and regrouping sites); see the partitions in Fig. 15 for an example of partitions that are not related in the quasi-1D way.

The key technical result is the following lemma about partitions of an annulus X into five ordered regions and two moves in Fig. 16. Note that the combination of these two moves can “deform” every region X_i ($i = 1, \dots, 5$) in the partition.

Lemma 3. *Let X be an annulus and σ be a reference state [Definition 1 of the main text]. Let $\{X_1, X_2, X_3, X_4, X_5\}$ be an ordered set of subsystems topologically equivalent to that shown in Fig. 16. If $\{X_1, X_2, X_3, X_4, X_5\}$ can be converted to a new partition $\{Y_1, Y_2, Y_3, Y_4, Y_5\}$ of X by a finite sequence of the two moves shown in Fig. 16, then $\tau_X(\{X_i\}) = \tau_X(\{Y_i\})$ where $\tau_X(\{X_i\})$ and $\tau_X(\{Y_i\})$ are the canonical Markov chains (Definition 4) with respect to the partitions $\{X_1, \dots, X_5\}$ and $\{Y_1, \dots, Y_5\}$ respectively.*

Proof. From the definition of σ [Definition 1], it follows that σ has a constant $I(A : C|B)_\sigma$ for any chain-like A, B , and C that partition $X = \cup_{i=1}^5 X_i$. By Proposition 1, σ_X is a locally Markov chain. Therefore, a canonical Markov chain associated with σ_X and $\{X_1, X_2, X_3, X_4, X_5\}$ exists, which we refer to as $\tau_X(\{X_i\})$.

Next, we show that the two moves in Fig. 16 do not change the canonical quantum Markov chain. The first move “deforms” $\{X_1, X_2, X_3, X_4, X_5\}$ to




FIG. 16. A partition of annulus X into ordered regions $\{X_1, X_2, X_3, X_4, X_5\}$ and two moves preserving the topology. The first move gives $\{X'_1, X'_2, X_3, X'_4, X'_5\}$, where $X'_1 X'_2 = X_1 X_2$ and $X'_4 X'_5 = X_4 X_5$. The second move gives $\{X_1, X'_2, X'_3, X'_4, X_5\}$ such that $X'_2 X'_3 X'_4 = X_2 X_3 X_4$.

$\{X'_1, X'_2, X_3, X'_4, X'_5\}$ such that $X_1 X_2 = X'_1 X'_2$ and $X_4 X_5 = X'_4 X'_5$ and such that the topology of the two sets is identical. (Note that X_3 is preserved.) We claim that $\tau_X(\{X_i\}) = \tau_X(\{X'_i\})$. This is because $\tau_X(\{X_i\})$ is the max-entropy state consistent with σ on $X_1 X_2 X_3$ and $X_3 X_4 X_5$ (by Proposition 4), and thus also the max-entropy state consistent with σ on $X'_1 X'_2 X_3 = X_1 X_2 X_3$ and $X_3 X'_4 X'_5 = X_3 X_4 X_5$.

The same idea works for the second move. The second move “deforms” $\{X_1, X_2, X_3, X_4, X_5\}$ to $\{X_1, X'_2, X'_3, X'_4, X_5\}$ such that $X_2 X_3 X_4 = X'_2 X'_3 X'_4$ and such that the topology of the two sets is identical. (Note that X_1 and X_5 are preserved.) The state τ_X is the max-entropy state consistent with $X_1 X_2 X_3 X_4$ and $X_2 X_3 X_4 X_5$ (by Proposition 4), and therefore it is also the max-entropy state consistent with $X_1 X'_2 X'_3 X'_4 = X_1 X_2 X_3 X_4$ and $X'_2 X'_3 X'_4 X_5 = X_2 X_3 X_4 X_5$.

Thus, as long as an ordered partition of X as $\{X_1, X_2, X_3, X_4, X_5\}$ can be converted to the ordered partition $\{Y_1, Y_2, Y_3, Y_4, Y_5\}$ by a finite sequence of the two moves depicted in Fig. 16, the canonical Markov chain associated with the former partition, $\tau_X(\{X_i\})$, must be identical with $\tau_X(\{Y_i\})$, the canonical Markov chain associated with the latter partition. This completes the proof. \square

We can now use Lemma 3 to argue that the max-entropy state (λ_X) consistent with the reference state on AB and BC in Fig. 15(a) is identical to the max-entropy state (λ'_X) consistent on the “spiral-shaped” subsystems $A'B'$ and $B'C'$, shown in Fig. 15(b). First, we partition B further as $B = X_2 X_3 X_4$ and similarly, $B' = Y_2 Y_3 Y_4$, and let $A = X_1, C = X_5, A' = Y_1, C' = Y_5$. Suppose that the size (and thickness) of each subsystem is large compared to the lattice spacing, so that it is possible to deform $\{X_1, X_2, X_3, X_4, X_5\}$ to $\{Y_1, Y_2, Y_3, Y_4, Y_5\}$ by a finite sequence of moves discussed in Lemma 3, and Fig. 16. Then $\tau_X(\{X_i\}) = \tau_X(\{Y_i\})$ according

to Lemma 3. By Proposition 4, $\lambda_X = \tau_X(\{X_i\})$ and $\lambda'_X = \tau_X(\{Y_i\})$. Therefore, $\lambda_X = \lambda'_X$. This completes the argument.⁵

This argument leads to the conclusion that the max-entropy state for a given annulus X is *canonical*. Insofar as two different partitions of an annulus are “topo-

logically equivalent” (in the sense that the partitions can be deformed using a sequence of moves described in Lemma 3), the max-entropy states associated with those partitions are identical. This generalizes the observation that one can associate a canonical Markov chain to a one-dimensional chain [Appendix C] to two dimensions.

⁵ In order for this argument to work, it is important to assume that the chosen subsystems are sufficiently large. Oth-

erwise, it may not be possible to deform $\{X_1, X_2, X_3, X_4, X_5\}$ to $\{Y_1, Y_2, Y_3, Y_4, Y_5\}$ using the moves discussed in Lemma 3.