

DENSITY OF RATIONAL POINTS ON DIAGONAL BIDEGREE (1, 2) HYPERSURFACE IN $\mathbb{P}^{s-1} \times \mathbb{P}^{s-1}$

XUN WANG

ABSTRACT. In this paper we establish an asymptotic formula for the number of rational points of bounded anticanonical height which lie on a certain Zariski dense subset of the biprojective hypersurface

$$x_1 y_1^2 + \dots + x_s y_s^2 = 0$$

in $\mathbb{P}^{s-1} \times \mathbb{P}^{s-1}$ with $s \geq 7$. This confirms the Manin conjecture for this variety.

CONTENTS

1.	Introduction	1
2.	Asymptotic formula using lattice counting	3
3.	Counting points using circle method	7
4.	Combination of results	12
5.	Finale	20
	References	32

1. INTRODUCTION

Consider the biprojective hypersurface V_s in $\mathbb{P}^{s-1} \times \mathbb{P}^{s-1}$ defined by the equation

$$x_1 y_1^2 + \dots + x_s y_s^2 = 0.$$

The purpose of this article is to establish Manin's conjecture for such varieties with $s \geq 7$. For a point $(x, y) \in V_s(\mathbb{Q})$ represented by a vector $(\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s$, the anticanonical height function is given by $H(x, y) = |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2}$, with $|\cdot| : \mathbb{R}^s \rightarrow \mathbb{R}$ the usual sup-norm.

Theorem 1.1. *If $s \geq 7$ then there exists an open subset $U \subset V_s$ such that*

$$\#\{(x, y) \in U(\mathbb{Q}) : H(x, y) \leq B\} \sim c_{\text{Peyre}} B \log B,$$

as $B \rightarrow \infty$.

Note that the leading constant arising in the theorem is as predicted by Peyre [12] and that, since $\text{Pic}(V) \cong \mathbb{Z}^2$, the order of magnitude agrees with the prediction of Manin [5].

In [14], Schindler proves Manin's conjecture for any biprojective hypersurface of bidegree (d_1, d_2) in $\mathbb{P}^{n_1-1} \times \mathbb{P}^{n_2-2}$ for $d_1, d_2 \geq 2$ and n_1, n_2 sufficiently large. Due to the assumption on the degrees her results do not cover the case in which we are interested. In general, the literature is lacking an equivalent result to Schindler's which will handle the bidegree $(1, d)$ setting for sufficiently many variables. Diagonal equations have long been an important testing ground in the study of rational points and there has been a great deal of study of Manin's conjecture for diagonal hypersurfaces of the form

$$(1.1) \quad x_1^{d_1} y_1^{d_2} + \dots + x_s^{d_1} y_s^{d_2} = 0.$$

If $d_1 = d_2 = 1$ then Manin's conjecture holds for $n \geq 2$ and this has been shown in multiple different ways: the work of Franke–Manin–Tschinkel [5] proves this using the theory of related Eisenstein series, Thunder [17] provides a proof using the geometry of numbers and Robbiani [13] and Spencer [16] prove Manin's conjecture via the circle method. In the case, $d_1 = 1, d_2 = 2$ (the case in which the present paper is interested) upper and lower bounds of the correct order of magnitude were established for $s = 3$ by Le Boudec [10]. Recently, the $s = 4$ case was studied by Browning–Heath-Brown [3]. They showed that the expected asymptotic growth holds under the condition that one removes a thin subset. In particular, one needs to avoid counting those points (\mathbf{x}, \mathbf{y}) such that $x_1 x_2 x_3 x_4 \in \mathbb{Q}^2$, which is an infinite union of closed subsets. This thin set phenomenon was in fact first observed, by Batyrev–Tschinkel [1], for varieties of the shape (1.1) with $d_1 = 1, d_2 = 3$. Our goal is to expand the literature by considering the $(1, 2)$ setting for higher values of s . We note that in our result, one does not need to worry about the removal of a thin set, as a closed subset will suffice. However, it is certainly necessary to remove a closed subset as, for instance, the points such that $x_1 = 0$ and $y_2 = \dots = y_s = 0$ will contribute $\gg B^s$ to the count.

Our approach is very similar to that of Browning–Heath-Brown. We proceed in two stages: for small \mathbf{y} we may fix the \mathbf{y} and consider the resulting linear equation in \mathbf{x} . We will study the number of solutions to this equation using techniques from the geometry of numbers. When \mathbf{x} is small, we fix these variables and consider the resulting quadratic equation. This we can attack using the circle method. It is in this step that we diverge from the work of Browning–Heath-Brown. In [3], the authors apply the δ -method in the form introduced by Heath-Brown [6] since they are dealing with a quadratic equation in 4 variables. For quadratic equations in at least 5 variables, the traditional form of the circle method is effective and so it is this form which

we have chosen to use. One may consider the equation which we aim to solve as being very similar to studying Waring's problem, except with coefficients, a problem which is well understood classically (see [4, Chapter 8]). However, we need all our results to be uniform in the coefficients as we need to sum over them after, and this addition to the classical problem is novel. As such we need to develop uniform versions of the key ingredients in the study of Waring's problem (such as Weyl's inequality, Hua's lemma, etc). Much of this uniform machinery could be used later to study varieties of a similar shape.

Acknowledgements. During the preparation of this paper the author is supported by NSF under Grant No. DMS-101601, while working at University of Michigan as part of the Math REU program. The author would like to thank Nicholas Rome and Will Dudarov for the numerous helpful comments and University of Michigan Math department for this opportunity.

Notation Throughout this paper η and ε will denote arbitrarily small constants, thus $c\varepsilon$ will thus be rewritten as ε and similar cases hold for other expressions depending on η and ε . The constants c_i , d and δ with $i \in \mathbb{N}$, whenever used, will denote certain small positive constants that's difficult to keep track of but easy to verify the value of.

2. ASYMPTOTIC FORMULA USING LATTICE COUNTING

This section is devoted to solving the equation when $|\mathbf{y}|$ is small. We follow closely with the work of Browning and Heath-Brown (See [8]) and arrive at various results using counting arguments from the geometry of numbers.

Denote

$$M_1(\mathbf{y}; R) = \#\{\mathbf{x} \in \mathbb{Z}^s : |\mathbf{x}| \leq R, F(\mathbf{x}; \mathbf{y}) = 0\}$$

for fixed $\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s$. Define

$$N_1(Y; B) = \#\left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; Y \leq |\mathbf{y}| < 2Y \end{array} \right\}.$$

Then from our construction

$$N_1(Y; B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} M_1(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)}).$$

The rest of this section is devoted to proving asymptotic formula for $N_1(Y; B)$. This was made explicit in the next theorem.

Theorem 2.1. *Let $Y \geq 1$. Then*

$$(2.1) \quad N_1(Y; B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O(Y^s) + O(B^{(s-2)/(s-1)} Y^{(s+2)/(s-1)}),$$

where

$$(2.2) \quad \varrho_\infty(\mathbf{y}) = \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(-\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{x} d\theta.$$

We note that if $\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s$ has at least 2 nonzero components then

$$\int_{[-1,1]^s} e(-\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{x} = \prod_{j \leq s} \frac{\sin(2\pi\theta y_j^2)}{\pi\theta y_j^2} \ll_{\mathbf{y}} (1 + |\theta|)^{-2},$$

and so the outer integral is absolutely convergent. If $\mathbf{y} = (1, 0, \dots)$ then

$$\int_{[-1,1]^s} e(-\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{x} = 2^{s-1} \frac{\sin(2\pi\theta)}{\pi\theta},$$

and the outside integral is conditionally convergent with value 2^{s-1} .

We start by estimating $M_1(\mathbf{y}; R)$ for individual $\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s$.

Lemma 2.2. *Let $\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s$ and let $d(\mathbf{y}) = \sqrt{y_1^2 + \dots + y_s^2}$. Let $V(\mathbf{y})$ be the volume of the intersection of unit cube $[-1, 1]^s$ and the hyperplane*

$$\{\mathbf{x} \in \mathbb{R}^s; F(\mathbf{x}; \mathbf{y}) = 0\}.$$

Then there exists constant $c > 0$, $\mathbf{x}_1 \in \mathbb{Z}_{\text{prim}}^s$ satisfying

$$0 < |\mathbf{x}_1| < c|\mathbf{y}|^{2/(s-1)} \text{ and } F(\mathbf{x}_1; \mathbf{y}) = 0,$$

such that

$$M_1(\mathbf{y}; R) = \frac{V(\mathbf{y})}{d(\mathbf{y})} R^{s-1} + O\left(\frac{R^{s-2}}{|\mathbf{x}_1|^{s-2}}\right) + O(1).$$

Proof. We first note that $M_1(\mathbf{y}; R)$ counts the number of vectors $\mathbf{x} \in \mathbb{Z}^s$ in an $s-1$ dimensional lattice of determinant $d(\mathbf{y})$ (c.f. [15] Lemma 1(i)). Lemma 2 of Schmidt [15] implies

$$M_1(\mathbf{y}; R) = \frac{V(\mathbf{y})}{d(\mathbf{y})} R^{s-1} + O\left(\sum_{i=1}^{s-2} \frac{R^i}{|\mathbf{x}_1| \dots |\mathbf{x}_i|}\right) + O(1),$$

after replacing the L^∞ norm with L^2 norm by applying Minkowski's second theorem. In particular, in Schmidt's notation the \mathbf{x}_i corresponds to the i -th successive minimas. It follows that $R^i / (|\mathbf{x}_1| \dots |\mathbf{x}_i|) \ll R^i / |\mathbf{x}_1|^i \ll \min(1, R^{s-2} / |\mathbf{x}_1|^{s-2})$ for all $i \leq s-2$. Finally we see that

$$|\mathbf{x}_1| \leq (|\mathbf{x}_1| \dots |\mathbf{x}_{s-1}|)^{1/(s-1)} \ll d(\mathbf{y})^{2/(s-1)} \ll |\mathbf{y}|^{2/(s-1)}.$$

Lemma 2.2 hence follows. \square

An immediate consequence of Lemma 2.2 is that

$$(2.3) \quad N_1(Y; B) = B \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{V(\mathbf{y})}{|\mathbf{y}|^{s-2}d(\mathbf{y})} + O(B^{(s-2)/(s-1)}|\mathbf{y}|^{-(s-2)^2/(s-1)}\Sigma_1) + O(Y^s),$$

where

$$\Sigma_1 = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{1}{|\mathbf{x}_1|^{s-2}}.$$

Consider the following sum

$$E(Y) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^s \\ 0 < |\mathbf{x}| \leq cY^{2/(s-1)}}} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y \\ F(\mathbf{x}; \mathbf{y}) = 0}} \frac{1}{|\mathbf{x}|^{s-2}}.$$

Clearly $\Sigma_1 \leq E(Y)$. We shall prove an upper bound for $E(Y)$, which shows that Σ_1 makes a satisfactory contribution to the error term appeared in Theorem 2.1.

Lemma 2.3. $E(Y) \ll Y^{4/(s-1)+s-2}$ for all $Y \geq 1$.

Proof. We divide the first sum in accordance with the ranges of $|\mathbf{x}|$ into dyadic intervals, and note that the constant c in this case can be ignored for our error term. It's immediate that

$$E(Y) \ll \sum_L L^{-s+2} \# \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0 \\ L/2 \leq |\mathbf{x}| \leq L, Y \leq |\mathbf{y}| < 2Y \end{array} \right\},$$

where summation is over $L \ll Y^{2/(s-1)}$ with L being powers of 2 only. From Browning and Hu [2], Theorem 2.1 we obtain that

$$\begin{aligned} E(Y) &\ll \sum_L L^{-s+2} \sum_{|\mathbf{x}| \sim L/2} (Y^{s-2} + L^{3s/4}Y^{(s-2)/2}) \\ &\ll \sum_L L^2 Y^{s-2} \ll Y^{4/(s-1)+s-2}. \end{aligned}$$

This proves Lemma 2.3. \square

In particular we see that the error term from (2.3) is satisfactory for Theorem 2.1 as a consequence of Lemma 2.3. It remains to show that

$$(2.4) \quad \frac{V(\mathbf{y})}{d(\mathbf{y})} = \varrho_\infty(\mathbf{y}).$$

Let $\mathbf{w} = (w_1, \dots, w_s)$ where $w_i = y_i^2 d(\mathbf{y})$ for $1 \leq i \leq s$, then one sees that $\|\mathbf{w}\|_2 = 1$ and

$$\varrho_\infty(\mathbf{y})d(\mathbf{y}) = \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(-\theta\mathbf{w} \cdot \mathbf{x})d\mathbf{x}d\theta.$$

In the case where only one of the components y_i of \mathbf{y} is nonzero, from remark at the beginning of this section we see that

$$\varrho_\infty(\mathbf{y})d(\mathbf{y}) = \int_{-\infty}^{\infty} 2^{s-1} \frac{\sin(2\pi\theta)}{\pi\theta} d\theta$$

converges conditionally to 2^{s-1} , which suffices for Theorem 2.1 since one easily sees that $V(\mathbf{y}) = 2^{s-1}$ in this case. In the case where \mathbf{y} has more than 1 nonzero components we can rewrite the integral, again by remark at the beginning of this section, as

$$\begin{aligned} & \lim_{\delta \downarrow 0} \int_{-\infty}^{\infty} \left(\frac{\sin(\pi\delta\theta)}{\pi\delta\theta} \right)^2 \int_{[-1,1]^s} e(-\theta\mathbf{w} \cdot \mathbf{x})d\mathbf{x}d\theta \\ &= \lim_{\delta \downarrow 0} \int_{-\infty}^{\infty} \int_{[-1,1]^s} \left(\frac{\sin(\pi\delta\theta)}{\pi\delta\theta} \right)^2 e(-\theta\mathbf{w} \cdot \mathbf{x})d\mathbf{x}d\theta. \end{aligned}$$

Using results from fourier analysis(c.f. [4], Lemma 20.1), one set

$$K(u; \delta) = \int_{-\infty}^{\infty} \left(\frac{\sin(\pi\delta\theta)}{\pi\delta\theta} \right)^2 e(-\theta u)d\theta,$$

where

$$K(u; \delta) = \begin{cases} \delta^{-2}(\delta - |u|), & \text{if } |u| \leq \delta, \\ 0, & \text{if } |u| \geq \delta. \end{cases}$$

We then see that

$$\varrho_\infty(\mathbf{y})d(\mathbf{y}) = \lim_{\delta \downarrow 0} \int_{[-1,1]^s} K(\mathbf{w} \cdot \mathbf{x}; \delta)d\mathbf{x}.$$

Since $\|\mathbf{w}\|_2 = 1$, there exists $M \in O_s(\mathbb{R})$ such that $M\mathbf{w} = (1, 0, \dots)$, then $\mathbf{w} \cdot \mathbf{x} = M\mathbf{w} \cdot M\mathbf{x}$. Hence, applying change of variables $\mathbf{z} = M\mathbf{x}$ and integrate over $Z = M[-1, 1]^s$, one has

$$\varrho_\infty(\mathbf{y})d(\mathbf{y}) = \lim_{\delta \downarrow 0} \int_Z K(z_1; \delta)d\mathbf{z} = \text{meas}\{\mathbf{z} \in Z; z_1 = 0\} = V(M\mathbf{w}) = V(\mathbf{w}) = V(\mathbf{y}).$$

This proves the claim and Theorem 2.1 follows as a consequence.

3. COUNTING POINTS USING CIRCLE METHOD

In this section we seek to prove an asymptotic formula for the counting function

$$M_3(\mathbf{x}; P) = \#\{\mathbf{y} \in \mathbb{Z}^s; |\mathbf{y}| \leq P; F(\mathbf{x}; \mathbf{y}) = 0\}$$

for small $|\mathbf{x}|$. We will use the classical Hardy-Littlewood circle method.(c.f. Davenport [4].) This contrasts the new δ -method developed by Heath-Brown and used in [3], which we have been following very closely to throughout this paper. (A reason for this comes from the fact that the classical Hardy-Littlewood circle method is applicable to any quadratic equations for $s \geq 5$, whereas Heath-Brown and Browning in [3] study the case for when $s = 4$ in [3]).

In correspondence with the classical Hardy-Littlewood circle method, we make the following definition for major and minor arcs: For arbitrarily small η , let

$$\mathfrak{M}_{a,q} = \left\{ \alpha \in [0, 1] : \left| \alpha - \frac{a}{q} \right| < (2q|\mathbf{x}|)^{-1} P^{-1-\eta} \right\},$$

for $a \leq q$, $(a, q) = 1$, $1 \leq q \leq P|\mathbf{x}|$ and η arbitrarily small. Define major arcs

$$\mathfrak{M} = \bigcup_{q \leq P|\mathbf{x}|} \bigcup_{\substack{a=1 \\ (a,q)=1}}^q \mathfrak{M}_{a,q},$$

and minor arcs

$$\mathfrak{m} = [0, 1] \setminus \mathfrak{M}.$$

Finally notice that

$$M_3(\mathbf{x}; P) = \left(\int_{\mathfrak{M}} + \int_{\mathfrak{m}} \right) \left(T_1(\alpha) \dots T_s(\alpha) d\alpha \right),$$

where

$$T_j(\alpha) = \sum_{|y_j| \leq P, y_j \in \mathbb{Z}} e(\alpha x_j y_j^2).$$

Our strategy will be as follows: along the major arcs we will be using the classical Hardy-Littlewood circle method to produce the truncated singular integral and singular series. This provides an expression immediately applicable in §4. For the integral along the minor arcs we will prove an upper bound, using Weyl estimate and Hua's lemma, which should suffice for small $|\mathbf{x}|$. The case when $|\mathbf{x}|$ is large will be dealt with in §4.

Our result is the following:

Theorem 3.1. *For $s \geq 7$ we have*

$$(3.1) \quad M_3(\mathbf{x}; P) = P^{s-2} \sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x}; P) + O(T) + O(E_2) + O(E_3),$$

where

$$\begin{aligned} T &= \int_{\mathfrak{m}} T_1(\alpha) \dots T_s(\alpha) d\alpha \ll E_1(\mathbf{x}; P), \\ \sigma_\infty(\mathbf{x}) &= \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{y} d\theta, \\ \mathfrak{S}(\mathbf{x}; P) &= \sum_{q \leq P^{|\mathbf{x}|}} q^{-s} S_q(\mathbf{x}), \quad S_q(\mathbf{x}) = \sum_{\substack{a \bmod q \\ (a,q)=1}} \sum_{\mathbf{b} \bmod q} e_q(aF(\mathbf{x}; \mathbf{b})), \end{aligned}$$

and

$$\begin{aligned} E_1(\mathbf{x}; P) &= P^{s/2+\varepsilon} |\Delta(\mathbf{x})|^{(s-4)/2s}, \\ E_2(\mathbf{x}; P) &= P^{s-2-\eta} |\mathbf{x}|^{-1} \sum_{q \leq P^{|\mathbf{x}|}} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s/2} \prod_{j \leq s} (q, x_j)^{1/2}, \\ E_3(\mathbf{x}; P) &= P^{s/2-1+c\eta} |\Delta(\mathbf{x})|^{-1/2} |\mathbf{x}|^{s/2-1} \sum_{q \leq P^{|\mathbf{x}|}} \prod_{j \leq s} (q, x_j)^{1/2}, \end{aligned}$$

for arbitrarily small ε and some constant c depending only on s where we denote

$$\Delta(\mathbf{x}) = \prod_{j \leq s} x_j.$$

We begin by evaluating the minor arcs.

Lemma 3.2. *(Uniform Weyl Estimate) Suppose that there exists natural numbers a, q such that*

$$(a, q) = 1, q > 0, \left| \alpha - \frac{a}{q} \right| \leq \frac{1}{q^2},$$

then for any $\varepsilon > 0$, we have

$$(3.2) \quad \left| \sum_{y_j = -P}^P e(\alpha x_j y_j^2) \right| \ll P^{1/2} + P^{1+\varepsilon} \left(\left(\frac{P}{|x_j|} \right)^{-1/2} + \left(\frac{q}{|x_j|} \right)^{-1/2} + \left(\frac{P^2}{q} \right)^{-1/2} \right).$$

Proof. This follows straight from the classical Weyl estimate (c.f. [4], Lemma 3.1). The only thing to notice is to replace the number of blocks $\frac{P}{q} + 1$ with $\frac{|x_j|P}{q} + 1$. \square

Lemma 3.3. *We have*

$$\int_0^1 |T_j(\alpha)|^4 d\alpha \ll P^{2+\varepsilon}.$$

where $T_j(\alpha)$ is as previously defined.

Proof. This is an immediate consequence of Hua's lemma. (c.f. [4], Lemma 3.2 and comments on page 40). \square

One thus comes to an estimate for the integral over the minor arc.

Lemma 3.4. *For $s \geq 7$*

$$T = \int_{\mathfrak{m}} |T_1(\alpha)T_2(\alpha)\dots T_s(\alpha)| d\alpha \ll E_1(\mathbf{x}; P),$$

where $T_j(\alpha)$ are defined as before.

Proof. By Holder's inequality, one has

$$\int_{\mathfrak{m}} |T_1(\alpha)T_2(\alpha)\dots T_s(\alpha)| d\alpha \ll \left(\int_{\mathfrak{m}} |T_1(\alpha)|^s d\alpha \right)^{\frac{1}{s}} \dots \left(\int_{\mathfrak{m}} |T_s(\alpha)|^s d\alpha \right)^{\frac{1}{s}},$$

so we simply need to evaluate one of these integrals. By Dirichlet approximation theorem one can find for every $\alpha \in [0, 1]$ a pair $(a, q) = 1$, $a \leq q$ satisfying for arbitrarily small η

$$(3.3) \quad 1 \leq q \leq 2P^{1+\eta}|\mathbf{x}|, \quad \left| \alpha - \frac{a}{q} \right| \leq (2q|\mathbf{x}|)^{-1}P^{-1-\eta}.$$

One sees immediately that the latter bound is exactly the one used for the major arcs. In particular if $\alpha \in \mathfrak{m}$, we must have $q \geq P|\mathbf{x}|$. Note that by our construction we have $P/|\mathbf{x}|, q/|\mathbf{x}|, P^2/q \gg P^{1-\eta}/|\mathbf{x}|$. Hence lemma 3.3 implies that after possibly refixing η and ε :

$$|T_j(\alpha)| \ll P^{s/2+\varepsilon}|x_j|^{1/2}.$$

Thus using lemma 3.5, we have

$$\begin{aligned} \left(\int_{\mathfrak{m}} |T_j(\alpha)|^s d\alpha \right)^{1/s} &= \max |T_j(\alpha)|^{(s-4)/s} \left(\int_{\mathfrak{m}} |T_1(\alpha)|^4 d\alpha \right)^{1/s} \\ &\ll \left(p^{(s-4)/2+\varepsilon}|x_j|^{1/2} \right)^{1/s}. \end{aligned}$$

Combining using Holder's inequality, we obtain the desired result, namely

$$\int_{\mathfrak{m}} |T_1(\alpha)T_2(\alpha)\dots T_s(\alpha)| d\alpha \ll P^{s/2+\varepsilon}|\Delta(\mathbf{x})|^{(s-4)/2s}.$$

\square

This provides an estimate for the integral along the minor arc. As indicated in the beginning of this section, we will treat the integral along the minor arcs more carefully.

We are now ready to move on to evaluating the exponentials along the major arcs. This will follow from the next few lemmas. In particular, we recall a classical estimate of quadratic Gauss sum.

Lemma 3.5. *For any natural numbers a, q such that $(a, q) = 1$, define*

$$S_{x_j a, q} = \sum_{b_j \bmod q} e_q(ax_j b_j^2),$$

then we have

$$|S_{x_j a, q}| = q^{1/2}(q, |x_j|)^{1/2}.$$

Proof. c.f. [9], Lemma 4.8 for coprime case. One can also deduce this from Lemma 3.2, setting $P = q$ and $\alpha = a/q$. \square

Lemma 3.6. *For α in $\mathfrak{M}_{a, q}$, let $\alpha = \beta + a/q$, then we have*

$$T_j(\alpha) = q^{-1} S_{x_j a, q} I_j(\beta) + O(q^{1/2}(q, x_j)^{1/2}),$$

where

$$I_j(\beta) = \int_{-P}^P e(x_j \beta u^2) du.$$

Proof. Note that one has

$$\begin{aligned} T_j(\alpha) &= \sum_{y=-P}^P e(\alpha x_j y^2) \\ &= \sum_{r \bmod q} e_q(ax_j r^2) \sum_b e(x_j \beta (bq + r)^2), \end{aligned}$$

where the summation for b takes place such that $bq + r$ runs over $\mathbb{N} \cap [-P, P]$. In particular, we now seek to replace the second sum by an integral as indicated, and then reevaluate the error terms. This follows immediately, or after taking complex conjugates, from van der Corput([4], Lemma 9.1); here by our set up we have $0 < |x_j| \beta q (bq + r) \leq 1/2$ and note that the first sum is bounded by $q^{1/2}(x_j, q)^{1/2}$ from Lemma 3.6. \square

This gives an estimate for the integral over the major arc.

Lemma 3.7. *For $s \geq 7$, we have*

$$(3.4) \quad \int_{\mathfrak{M}} T_1(\alpha) \dots T_s(\alpha) d\alpha = P^{s-2} \sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x}; P) + O(E_2) + O(E_3),$$

where

$$\begin{aligned}\sigma_\infty(\mathbf{x}) &= \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{y} d\theta, \\ \mathfrak{S}(\mathbf{x}; P) &= \sum_{q \leq P|\mathbf{x}|} q^{-s} S_q(\mathbf{x}), \quad S_q(\mathbf{x}) = \sum_{\substack{a \bmod q \\ (a,q)=1}} \sum_{\mathbf{b} \bmod q} e_q(aF(\mathbf{x}; \mathbf{b})),\end{aligned}$$

where we remind the readers from Theorem 3.1 that

$$\begin{aligned}E_2 &= P^{s-2-\eta} |\mathbf{x}|^{-1} \sum_{q \leq P|\mathbf{x}|} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s/2} \prod_{j \leq s} (q, x_j)^{1/2}, \\ E_3 &= P^{s/2-1+c\eta} |\Delta(\mathbf{x})|^{-1/2} |\mathbf{x}|^{s/2-1} \sum_{q \leq P|\mathbf{x}|} \prod_{j \leq s} (q, x_j)^{1/2},\end{aligned}$$

for some constant c not depending on η .

Proof. Using Lemma 3.5, we see that since

$$|q^{-1} S_{x_j a, q} I_j(\beta)| \ll P q^{-1/2} (q, x_j)^{1/2},$$

the error terms from multiplying $T_i(\alpha)$'s are bounded by $O(P^{s-1} T(q, \mathbf{x})^{1/2} q^{-(s-2)/2})$ where

$$T(q, \mathbf{x}) = \prod_{j \leq s} (q, x_j).$$

Integrating over major arcs and summing over respective ranges for q and a , one obtains

$$\begin{aligned}\int_{\mathfrak{M}} T_1(\alpha) \dots T_s(\alpha) d\alpha &= \sum_{q \leq P|\mathbf{x}|} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s} S_{x_1 a, q} \dots S_{x_s a, q} \int_{\mathfrak{M}_{a,q}} \prod_{i=1}^s I_i(\alpha - a/q) \\ &\quad + O(P^{s-2-\eta} |\mathbf{x}|^{-1} \sum_{q \leq P|\mathbf{x}|} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s/2} \prod_{j \leq s} (q, x_j)^{1/2}),\end{aligned}$$

applying change of variables with $\theta = P^2 \beta$ and $\mathbf{y} = P\mathbf{z}$, we have

$$\begin{aligned}\int_{\mathfrak{M}} T_1(\alpha) \dots T_s(\alpha) d\alpha &= P^{s-2} \sum_{q \leq P} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s} S_{x_1 a, q} \dots S_{x_s a, q} \int_{|\theta| \leq P^{1-\eta} (2q|\mathbf{x}|)^{-1}} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{y} d\theta \\ &\quad + O(E_2).\end{aligned}$$

To complete the singular integral, one begins by noticing that by Holder's inequality we have

$$\begin{aligned} \int_{|\theta| > P^{1-\eta}/(2q|\mathbf{x}|)^{-1}} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{y} d\theta &\ll \int_{|\theta| > P^{1-\eta}/(2q|\mathbf{x}|)^{-1}} \prod_{j \leq s} \min(1, |\theta|^{-1/2} |x_j|^{-1/2}) d\theta \\ &\ll P^{-s/2+1+c\eta} (q|\mathbf{x}|)^{s/2-1} |\Delta(\mathbf{x})|^{-1/2}. \end{aligned}$$

After summing this gives E_3 , noting again that

$$|q^{-s} S_{ax_1, q} \dots S_{ax_s, q}| \ll q^{-s/2} T(q, \mathbf{x})^{1/2}.$$

This allows us to replace the finite integral with singular integral $\sigma_\infty(\mathbf{x})$ and leads to E_3 , which gives (3.4). \square

We can then combine Lemma 3.8 and Lemma 3.5; Theorem 3.1 follows as a consequence.

4. COMBINATION OF RESULTS

This section treats the results in §2, 3, summing over respective ranges for \mathbf{x} and \mathbf{y} . To begin we remind readers that

$$M_1(\mathbf{y}; R) = \#\{\mathbf{x} \in \mathbb{Z}^s : |\mathbf{x}| \leq R, F(\mathbf{x}; \mathbf{y}) = 0\}$$

and

$$N_1(Y; B) = \#\left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; Y \leq |\mathbf{y}| < 2Y \end{array} \right\}.$$

By construction, $N_1(Y; B)$ can be rewritten as

$$N_1(Y; B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} M_1(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)}).$$

In particular define in analogy

$$M_2(\mathbf{y}; R) = \#\{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^s : |\mathbf{x}| \leq R, F(\mathbf{x}; \mathbf{y}) = 0\},$$

and let

$$N_2(Y; B) = \#\left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; Y \leq |\mathbf{y}| < 2Y \end{array} \right\}.$$

Again by construction we see that

$$N_2(Y; B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} M_2(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)}).$$

Notice that $N_2(Y; B)$ is the expression we seek to evaluate in Theorem 1.1, and the following lemma gives an asymptotic for this expression.

Lemma 4.1. *For $1 \leq Y \leq B^{1/(s+2)}$ we have that*

$$\begin{aligned} N_2(B; Y) &= \frac{B}{\zeta(s-1)} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O(B^{1/(s-1)} Y^{s-(s-2)/(s-1)}) \\ &\quad + O(B^{(s-2)/(s-1)} Y^{(s+2)/(s-1)}) + O(B^{1/(s-1)} Y^{(s-2)^2/(s-1)}). \end{aligned}$$

Proof. One immediately sees from the definition that

$$\begin{aligned} N_2(Y; B) &= \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} M_2(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)}) \\ &= \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \sum_{d \leq M_2(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)})} \mu(d) M_1(\mathbf{y}; (B/|\mathbf{y}|^{s-2})^{1/(s-1)}/d) \\ &= \sum_{d \leq (B/Y^{s-2})^{1/(s-1)}} \mu(d) N_1(Y; B/d^{s-1}) \\ &= B \sum_{d \leq (B/Y^{s-2})^{1/(s-1)}} \frac{\mu(d)}{d^{s-1}} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + \\ &\quad O(Y^s (B/Y^{s-2})^{1/(s-1)}) + O(B^{(s-2)/(s-1)} Y^{(s+2)/(s-1)}). \end{aligned}$$

Which follows from Theorem 2.1. We note that

$$\frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} = \frac{V(\mathbf{y})}{d(\mathbf{y})|\mathbf{y}|^{s-2}} \ll 1,$$

and therefore we can safely replace the first sum with a series, obtaining

$$\frac{B}{\zeta(s-1)} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ Y \leq |\mathbf{y}| < 2Y}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O(B^{1-(s-2)/(s-1)} Y^{(s-2)^2/(s-1)}).$$

which is satisfactory. \square

One notices that when $Y \leq B^{1/(s+2)}$, all 3 error terms contribute $O(B)$. In particular one sees that after dyadic summation this leads to the overall

contribution and the following asymptotics:

$$\begin{aligned} & \# \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; |\mathbf{y}| \leq B^{1/(s+2)} \end{array} \right\} \\ &= \frac{B}{\zeta(s-1)} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{y}| \leq B^{1/(s+2)}}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O(B). \end{aligned}$$

Next we turn to the circle method treatment in §3; we remind the readers that

$$M_3(\mathbf{x}; P) = \#\{\mathbf{y} \in \mathbb{Z}^s; |\mathbf{y}| \leq P; F(\mathbf{x}; \mathbf{y}) = 0\},$$

and we define

$$N_3(X; B) = \# \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; X \leq |\mathbf{x}| < 2X \end{array} \right\}.$$

Then it follows from the construction that

$$N_3(X; B) = \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} M_3(\mathbf{x}; (B/|\mathbf{x}|^{s-1})^{1/(s-2)}).$$

We make use of the following Lemma about minor arc contribution, which shall be proved at the end of this section.

Lemma 4.2. *For $1 \leq X \leq B^{4/(s+2)(s-1)}$ and some small enough c_5 , we see that*

$$\sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T \ll B^{1-c_5}$$

where T is as defined in Theorem 3.1.

Assuming Lemma 4.2, we provide an estimate for $N_3(X; B)$.

Lemma 4.3. *For $1 \leq X \leq B^{4/(s+2)(s-1)}$ and some small enough c_4 we have*

$$N_3(X; B) = B \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + O(B^{1-c_4}),$$

where

$$\mathfrak{S}(\mathbf{x}) = \sum_{q=1}^{\infty} q^{-s} S_q(\mathbf{x}), \quad S_q(\mathbf{x}) = \sum_{\substack{a \bmod q \\ (a,q)=1}} \sum_{\mathbf{b} \bmod q} e_q(aF(\mathbf{x}; \mathbf{b})).$$

Proof. We wish to sum $M_3(\mathbf{x}; P)$ over the range $|\mathbf{x}| \sim X$ and $\mathbf{x} \in \mathbb{Z}_{\text{prim}}^s$. It suffices to evaluate each of error terms in Theorem 3.1, namely E_i 's and to replace the finite sum by a singular series with bounded error term.

We begin by evaluating the error terms. Following from Lemma 4.2, it suffices to evaluate error terms E_2 and E_3 , namely,

$$E_2(\mathbf{x}; P) = P^{s-2-\eta} |\mathbf{x}|^{-1} \sum_{q \leq P|\mathbf{x}|} \sum_{\substack{a \bmod q \\ (a,q)=1}} q^{-s/2} \prod_{j \leq s} (q, x_j)^{1/2},$$

$$E_3(\mathbf{x}; P) = P^{s/2-1+c\eta} |\Delta(\mathbf{x})|^{-1/2} |\mathbf{x}|^{s/2-1} \sum_{q \leq P|\mathbf{x}|} \prod_{j \leq s} (q, x_j)^{1/2}.$$

We first seek to evaluate E_2 , noting that

$$\begin{aligned} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} E_2(\mathbf{x}; (B/|\mathbf{x}|^{s-1})^{1/(s-2)}) &\ll \frac{B^{1-c\eta}}{X^{s-1-d\eta}} X^{-1} \sum_{\mathbf{x}} \sum_{q \leq (B/|\mathbf{x}|^{s-1})^{1/(s-2)} |\mathbf{x}|} q^{-s/2+2} \prod_{j \leq s} (q, x_j)^{1/2}, \\ &\ll \frac{B^{1-c\eta}}{X^{s-d\eta}} \sum_{q \leq (B/|\mathbf{x}|^{s-1})^{1/(s-2)} |\mathbf{x}|} q^{-s/2+2} \prod_{j \leq s} \sum_{\substack{x_j \in \mathbb{Z} \\ |x_j| \leq X}} (q, x_j)^{1/2}, \end{aligned}$$

where c, d are some small fixed positive constants depending only on s . For each individual sum in the final product we see that

$$\begin{aligned} \sum_{\substack{x_j \in \mathbb{Z} \\ x_j \leq X}} (q, x_j)^{1/2} &\ll \sum_{\substack{u_1 \leq X \\ u_1 | q^\infty}} (q, u_1)^{1/2} \sum_{\substack{u_2 \leq X/u_1 \\ (u_2, q)=1}} 1, \\ &\ll X \sum_{\substack{u_1 \leq X \\ u_1 | q^\infty}} \frac{(q, u_1)^{1/2}}{u_1}, \\ &\ll_\varepsilon X^{1+\varepsilon} q^\varepsilon, \end{aligned}$$

since there are $O(X^\varepsilon q^\varepsilon)$ choices in the last sum, which one may obtain by applying Rankin's method to the sum. In particular we have

$$\begin{aligned} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} E_2(\mathbf{x}; (B/|\mathbf{x}|^{s-1})^{1/(s-2)}) &\ll_\varepsilon X^{s+\varepsilon} \frac{B^{1-c\eta}}{X^{s-d\eta}} \sum_q q^{-s/2+2+\varepsilon}, \\ &\ll_{\varepsilon, \eta} B^{1-c_2}, \end{aligned}$$

for small enough c_2 and $X \leq B^{4/(s+2)(s-1)}$. Finally we start by assuming that $|x_j| \sim X$ for all $j \leq s$. It's easy to see, following similar evaluation process:

$$\sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} E_3(\mathbf{x}; (B/|\mathbf{x}|^{s-1})^{1/(s-2)}) \ll_{\eta, \varepsilon} X^{s+\varepsilon} (B/X^{s-1})^{(s/2+c\eta)/(s-2)} \ll B^{1-c_3},$$

for small enough c_3, η and $X \leq B^{4/(s+2)(s-1)}$. We can then remove the condition $|x_j| \sim X$ by the following argument: if we assume that $X^{1-p_j} \gg |x_j| \gg X^{1-t_j}$ for some t_j then we would be replacing X^s with $X^{s-\sum p_j}$ and $|\Delta(\mathbf{x})|^{-1/2} |\mathbf{x}|^{s/2-1}$, originally treated as X^{-1} , with $X^{\sum t_j/2-1}$. Choosing appropriate and finite number of p_j 's and t_j 's, one can show that the contribution from this range won't exceed $B^{1-c_3/2}$, which is satisfactory.

At this point we note that after summing

$$N_3(X; B) = B \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x}; B^{1/(s-2)} |\mathbf{x}|^{1/(s-2)})}{|\mathbf{x}|^{s-1}} + O(B^{1-c_4}),$$

so it suffices to prove the validity of replacing the finite sum with the singular series. We begin by noting that

$$\sum_{q \leq C} q^{-s/2-1} S_q(\mathbf{x}) \ll \sum_{q \leq C} \frac{\prod_{j \leq s} (x_j, q)^{1/2}}{q} \ll C |\Delta_{\text{bad}}(\mathbf{x})|^{(s-1)/2s},$$

where

$$\Delta_{\text{bad}}(\mathbf{x}) = \prod_{\substack{p|\Delta(\mathbf{x}) \\ \nu_p(\Delta(\mathbf{x})) \geq 2}} p^{\nu_p(\Delta(\mathbf{x}))}.$$

Hence by partial summation we see that

$$\sum_{q \geq (B/|\mathbf{x}|^{s-1})^{1/(s-2)}} q^{-s} S_q(\mathbf{x}) \ll (B|\mathbf{x}|)^{(-s/2+2)/(s-2)} |\Delta_{\text{bad}}(\mathbf{x})|^{(s-1)/2s}.$$

On noting that $\sigma_\infty(\mathbf{x}) \ll |\Delta(\mathbf{x})|^{-1/2}$ (c.f. [4], Theorem 7.1) and $|\mathbf{x}|^{s-1} \geq |\Delta(\mathbf{x})|^{(s-1)/s}$ we see that the tail of the singular series contributes

$$\ll (BX)^{(-s/2+2)/(s-2)} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{|\Delta_{\text{bad}}(\mathbf{x})|^{(s-1)/2s}}{|\Delta(\mathbf{x})|^{1/2+(s-1)/s}}.$$

Rewriting $n = |\Delta_{\text{bad}}(\mathbf{x})|$ and $t = |\Delta(\mathbf{x})|$

$$\begin{aligned} &\ll (BX)^{(-s/2+2)/(s-2)} \sum_{\substack{n \ll X^s \\ n \text{ squarefull}}} n^{(s-1)/2s} \sum_{\substack{t \ll X^s \\ n|t}} \frac{\tau_s(t)}{t^{1/2+(s-1)/s}} \\ &\ll B^{-c} \sum_{n \text{ squarefull}} \frac{1}{n^{(s+1)/2s}}. \end{aligned}$$

for small positive constant c where $s \geq 7$ and $X \leq B^{4/(s+2)(s-1)}$. This concludes the proof, noting that the last sum converges. \square

Similar as before, define

$$M_4(\mathbf{x}; P) = \#\{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s; |\mathbf{y}| \leq P; F(\mathbf{x}; \mathbf{y}) = 0\}$$

and let

$$N_4(X; B) = \#\left\{(\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; X \leq |\mathbf{x}| < 2X \end{array} \right\}.$$

By construction we have

$$N_4(X; B) = \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} M_4(\mathbf{x}; (B/|\mathbf{x}|^{s-1})^{1/(s-2)}).$$

Notice that the expression $N_4(X; B)$ is exactly what we are looking for in Theorem 1.1 and thus we seek an asymptotic for it.

Lemma 4.4. *For $1 \leq X \leq B^{4/(s+2)(s-1)}$, we have*

$$N_4(X; B) = \frac{B}{\zeta(s-2)} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + O(B^{1-\eta})$$

for small enough η .

Proof. By previous lemma we have for suitably small η

$$\begin{aligned} N_4(X; B) &= \sum_{d \leq (B/X^{s-1})^{1/(s-2)}} \mu(d) N_3(X; B/d^{s-2}) \\ &= B \sum_{d \leq (B/X^{s-1})^{1/(s-2)}} \frac{\mu(d)}{d^{s-1}} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + O(B^{1-\eta}). \end{aligned}$$

Replacing the summation over mobius functions μ with a series, we conclude that

$$N_4(X; B) = \frac{B}{\zeta(s-2)} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{G}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + O(B^{1-\eta}).$$

□

In particular, Lemma 4.1 and 4.4 lead to the following asymptotics.

$$\begin{aligned} & \# \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{Z}_{\text{prim}}^s \times \mathbb{Z}_{\text{prim}}^s : \begin{array}{l} F(\mathbf{x}; \mathbf{y}) = 0; \\ |\mathbf{x}|^{s-1} |\mathbf{y}|^{s-2} \leq B; |\mathbf{x}| \leq B^{4/(s+2)(s-1)} \end{array} \right\} \\ &= \frac{B}{\zeta(s-2)} \sum_{\substack{|\mathbf{x}| \in \mathbb{Z}_{\text{prim}}^s \\ \mathbf{x} \leq B^{4/(s+2)(s-1)}}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{G}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + O(B^{1-\eta} \log B) + O(B), \end{aligned}$$

and in particular we see that

$$\begin{aligned} N(\Omega, B) &= \frac{1}{4} \left(\frac{B}{\zeta(s-2)} \sum_{\substack{|\mathbf{x}| \in \mathbb{Z}_{\text{prim}}^s \\ \mathbf{x} \leq B^{4/(s+2)(s-1)}}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{G}(\mathbf{x})}{|\mathbf{x}|^{s-1}} + \frac{B}{\zeta(s-1)} \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{y}| \leq B^{1/(s+2)}}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} \right) \\ &+ O(B) \\ &= \frac{B}{4\zeta(s-2)} M_1(B) + \frac{B}{4\zeta(s-1)} M_2(B) + O(B), \end{aligned}$$

with

$$M_1(B) = \sum_{\substack{|\mathbf{x}| \in \mathbb{Z}_{\text{prim}}^s \\ \mathbf{x} \leq B^{4/(s+2)(s-1)}}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{G}(\mathbf{x})}{|\mathbf{x}|^{s-1}} \quad \text{and} \quad M_2(B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{y}| \leq B^{1/(s+2)}}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}}.$$

It remains to prove Lemma 4.2.

Proof. We recall that it suffices to show

$$\sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T \ll B^{1-c_5},$$

for some small enough c_5 , where we remind the readers that

$$T = \int_{\mathfrak{m}(\mathbf{x})} T_1(\alpha) \dots T_s(\alpha) d\alpha$$

Our strategy will be as follows. When X is small, we will apply Lemma 3.4 to provide a satisfying upper bound; when X is large, we will treat the sum over

$|\mathbf{x}| \leq X$ and $|\mathbf{y}| \leq P$ as a cubic form in $2s$ variables, with $P = (B/X^{s-1})^{1/(s-2)}$. To begin, we recall from Lemma 3.4 that

$$\begin{aligned} \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T &\ll \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} E_1(\mathbf{x}; (B/X^{s-1})^{1/(s-2)}) \\ &\ll X^s \frac{B^{s/2+\varepsilon}}{X^{s(s-1)/2(s-2)}} X^{(s-4)/2} \ll B^{1-c_5} \end{aligned}$$

for $s \geq 8$, $X \leq B^{4/(s+2)(s-1)}$ or $s = 7$, $X \leq B^{3.46/(s+2)(s-1)}$ and $c_5 > 0$ small enough. We now make the following definition

$$C(\mathbf{u}) = \sum_{i=1}^s u_i u_{2s+1-i}^2$$

for C cubic form on \mathbb{Z}^{2s} . In particular, one notices that

$$(4.1) \quad \sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T \ll \sup_{\alpha} \left| \sum_{\substack{|\mathbf{x}| \sim X, \Delta(\mathbf{x}) \neq 0 \\ |\mathbf{y}| \leq P \\ \mathbf{x}, \mathbf{y} \in \mathbb{Z}^s}} e(\alpha C(\mathbf{x}, \mathbf{y})) \right|.$$

where the sup is taken over all α such that if there exists $(a, q) = 1$ such that $|\alpha - a/q| \leq 1/2(qXP^{1+\eta})$ then $q \geq PX$; one should think of this as taking supremum over unions of minor arcs for each $\mathbf{x} \in \mathbb{Z}^s$ (c.f. Lemma 3.4), and in particular, for all such α , $PX \leq q \leq 4PX$.

Denote the right hand sum by $S(\alpha)$, then tracing the arguments of [7], §2, one finds that

$$|S(\alpha)|^4 \ll P^{3s+\varepsilon} X^s \sum_{\substack{|u_1 \dots u_s| \neq 0, \max_{i \leq s} |u_i| \leq X \\ |u_{s+1}| \dots |u_{2s}| \leq P}} M(\alpha, P, X),$$

where

$$M(\alpha, P, X) := \# \left\{ \mathbf{v} \in \mathbb{Z}^{2s} : \begin{array}{l} 0 < |v_1| \dots |v_s| \leq X, |v_{s+1}| \dots |v_{2s}| \leq P, \\ \|6\alpha B_i(\mathbf{u}, \mathbf{v})\| \leq P^{-1} \forall i \leq 2s \end{array} \right\}$$

and where

$$B_i(\mathbf{u}, \mathbf{v}) = \begin{cases} \frac{1}{3} u_{2s+1-i} v_{2s+1-i} & \text{if } i \leq s, \\ \frac{1}{3} u_i v_{2s+1-i} + \frac{1}{3} v_i u_{2s+1-i} & \text{if } i > s. \end{cases}$$

Then trivially we have

$$M(\alpha, P, X) \ll \# \left\{ \mathbf{v} \in \mathbb{Z}^{2s} : \begin{array}{l} 0 < |v_1| \dots |v_s|, |v_1| \dots |v_{2s}| \leq P \\ \|6\alpha B_i(\mathbf{u}, \mathbf{v})\| \leq P^{-1} \forall i \leq 2s \end{array} \right\}.$$

It therefore follows from [7] Lemma 2.2, 2.3 and equation (2.7) that

$$|S(\alpha)|^4 \ll X^{3s} P^s Z^{-4s} \# \left\{ (\mathbf{u}, \mathbf{v}) \in \mathbb{Z}^{4s} : \begin{array}{l} |u_1 \dots u_s|, |v_1 \dots v_s| \neq 0 \\ |v_1| \dots |v_{2s}|, |u_1| \dots |u_{2s}| \leq ZP \\ \|6\alpha B_i(\mathbf{u}, \mathbf{v})\| = 0 \ \forall i \leq 2s \end{array} \right\},$$

with

$$0 < Z < 1, \quad Z^2 < (12cq|\beta|P^2)^{-1}, \quad Z^2 < P/2q,$$

and

$$Z^2 < \max\left(\frac{q}{6cP^2}, qP|\theta|\right),$$

where $\alpha = a/q + \beta$ is the rational approximation to α as indicated before (or c.f. Lemma 3.4 with $|\mathbf{x}|$ replaced by X). In particular, from our remark at the beginning of the proof of this lemma, $4P^{1+\eta}X \geq q \geq PX$. Finally notice that there is only 1 solution to the $\|6\alpha B_i(\mathbf{u}, \mathbf{v})\| = 0$ with $u_1 \dots u_s, v_1 \dots v_s \neq 0$, namely when $v_{s+1} = \dots = v_{2s} = u_{s+1} = \dots = u_{2s} = 0$ by definition of the bilinear form B_i 's. Using (4.1),

$$\left(\sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T \right)^4 \ll X^{3s} P^{s+\eta} (1 + (q|\theta|P^2)^{2s} + q^{2s} P^{-2s} + P^{2s} q^{-2s}),$$

Finally by our construction of minor arcs and α , we have that $\theta \ll 1/qXP^{1+\eta}$ and $PX \leq q \leq 2P^{1+\eta}X$, we obtain for $B^{3.46/(s+2)(s-1)} \leq X \leq B^{4/(s+2)(s-1)}$,

$$\sum_{\substack{X \leq |\mathbf{x}| < 2X \\ \mathbf{x} \in \mathbb{Z}_{\text{prim}}^s}} T \ll X^{3s/4} P^{3s/4+\eta} + P^{5s/4+\eta} X^{-s/4} \ll B^{1-c_5},$$

where $P = (B/X^{s-1})^{1/(s-2)}$, small enough η and $s \geq 7$. We see that this is satisfactory for Lemma 4.2. (When $s = 6$, the error term becomes $O(B^{81/80+\eta})$, which is not satisfactory.) \square

5. FINALE

In this section we produce asymptotic formulae for $M_1(B)$ and $M_2(B)$ as given in §4; we will then show that the overall contribution is satisfactory in conclusion for Theorem 1.1. Again we follow very closely with the work of Heath-Brown and Browning [3].

We will first treat $M_2(B)$ as it is slightly easier to handle. We remind the readers that

$$M_2(B) = \sum_{\substack{\mathbf{y} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{y}| \leq B^{1/(s+2)}}} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}},$$

where

$$\varrho_\infty(\mathbf{y}) = \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{x} d\theta.$$

Lemma 5.1. *For $s \geq 7$, we have*

$$M_2(B) = \frac{(s-2)}{(s+2)\zeta(s)} \tau_\infty \log B + O(1),$$

where τ_∞ is as given in Theorem 1.1.

To begin, note that we can add in a Mobius function to check primitivity condition

$$M_2(B) = \sum_{k \leq B^{1/(s+2)}} \frac{\mu(k)}{k^{s-2}} \sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_0} \frac{\varrho_\infty(k\mathbf{y})}{|\mathbf{y}|^{s-2}},$$

where

$$T_0 = T_0(k) = \{\mathbf{y} \in \mathbb{R}^s; 1 \leq |\mathbf{y}| \leq B^{1/(s+2)}/k\},$$

in particular noting that $\varrho_\infty(k\mathbf{y}) = k^{-2}\varrho_\infty(\mathbf{y})$, we have that

$$(5.1) \quad M_2(B) = \sum_{k \leq B^{1/(s+2)}} \frac{\mu(k)}{k^s} \sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}}.$$

The main idea is then to convert the second sum into an integral around $[-1,1]^s$ to produce τ_∞ from ϱ_∞ . Throughout this subsection we make use of the fact that

$$\varrho_\infty(\mathbf{y}) \ll |\mathbf{y}|^{-2}$$

which is clear from (2.4). We begin with an estimate.

Lemma 5.2. *For $\min_j(|y_j|) \geq 2$, then*

$$(5.2) \quad \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} = \int_{[0,1]^s} \frac{\varrho_\infty(\mathbf{y} + \mathbf{t})}{|\mathbf{y} + \mathbf{t}|^{s-2}} d\mathbf{t} + O(|\mathbf{y}|^{2-s} \min_j(|y_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{y})|^{-2/(s-1)}).$$

Proof. We begin by proving that

$$\nabla \varrho_\infty(\mathbf{u}) \ll \min_j(|u_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{u})|^{-2/(s-1)}.$$

Note that

$$\varrho_\infty(\mathbf{u}) = \int_{-\infty}^{\infty} \prod_{i=1}^s I(-\theta u_i^2) d\theta,$$

where

$$I(\psi) = \int_{-1}^1 e(\psi x) dx.$$

One can then try to compute the derivative explicitly, noting that $I(\psi) \ll \min(1, |\psi|^{-1})$,

$$\frac{\partial}{\partial u_j} I(-\theta u_i^2) = -4\pi i \theta u_j \int_{-1}^1 x e(-\theta x u_j^2) dx \ll |u_j|^{-1},$$

and therefore

$$\begin{aligned} \frac{\partial}{\partial u_j} \varrho_\infty(\mathbf{u}) &\ll |u_j|^{-1} \int_{-\infty}^{\infty} \min(1, |\theta|^{-(s-1)}) \prod_{i=1; i \neq j}^s |u_i|^{-2} d\theta \\ &\ll |u_j|^{-1} \prod_{i=1; i \neq j}^s |u_i|^{-2/(s-1)} \\ &\ll \min_j (|u_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{u})|^{-2/(s-1)}, \end{aligned}$$

as needed.

We then make the following observation

$$\frac{\varrho_\infty(\mathbf{y} + \mathbf{t})}{|\mathbf{y} + \mathbf{t}|^{s-2}} - \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} = \frac{\varrho_\infty(\mathbf{y} + \mathbf{t}) - \varrho_\infty(\mathbf{y})}{|\mathbf{y} + \mathbf{t}|^{s-2}} + \varrho_\infty(\mathbf{y}) (|\mathbf{y} + \mathbf{t}|^{2-s} - |\mathbf{y}|^{2-s}).$$

Then if $\min_j (|y_j|) \geq 2$ and $|t_i| \leq 1$, then clearly

$$|y_i + t_i| \geq |y_i| - |t_i| \geq \frac{1}{2} |y_i|,$$

and thus $|\mathbf{y} + \mathbf{t}|^{2-s} \ll |\mathbf{y}|^{2-s}$. In particular, noting that by mean value theorem

$$|\varrho_\infty(\mathbf{y} + \mathbf{t}) - \varrho_\infty(\mathbf{y})| \leq \sup_{0 \leq \xi \leq 1} \left| \frac{\partial}{\partial \xi} \varrho_\infty(\mathbf{y} + \xi \mathbf{t}) \right|,$$

and therefore it follows from previous discussion that

$$\frac{\varrho_\infty(\mathbf{y} + \mathbf{t}) - \varrho_\infty(\mathbf{y})}{|\mathbf{y} + \mathbf{t}|^{s-2}} \ll |\mathbf{y}|^{2-s} \min_j (|y_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{y})|^{-2/(s-1)}.$$

We also have since $\min_j (|y_j|) \geq 2$ and $|t_i| \leq 1$, that by mean value theorem

$$(|\mathbf{y} + \mathbf{t}|^{2-s} - |\mathbf{y}|^{2-s}) \ll |\mathbf{y}|^{1-s},$$

thus

$$\varrho_\infty(\mathbf{y})(|\mathbf{y} + \mathbf{t}|^{2-s} - |\mathbf{y}|^{2-s}) \ll |\mathbf{y}|^{1-s} \varrho_\infty(\mathbf{y}) \ll |\mathbf{y}|^{2-s} \min_j (|y_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{y})|^{-2/(s-1)},$$

and therefore

$$(5.3) \quad \frac{\varrho_\infty(\mathbf{y} + \mathbf{t})}{|\mathbf{y} + \mathbf{t}|^{s-2}} = \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O(|\mathbf{y}|^{2-s} \min_j (|y_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{y})|^{-2/(s-1)}).$$

Integrating over $[0, 1]^s$ gives the result. \square

It's now appropriate to estimate the second sum in (5.1) with an integral.

Lemma 5.3. *We have:*

$$\sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} = J_2(B; k) + O(1),$$

where

$$J_2(B; k) = \int_{T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} d\mathbf{y}.$$

Proof. We begin by defining

$$X = \{\mathbf{y} \in \mathbb{Z}^s : |\mathbf{y}| \leq B^{1/(s+2)}/k - 2; \min_j |y_j| \geq 2\},$$

and let

$$Y = \bigcup_{\mathbf{y} \in X} \mathbf{y} + (0, 1]^s.$$

One begins to note that $Y \subset T_0$ and $T_0 \setminus Y \subset T_1 \cup T_2$, where

$$T_1 = \{\mathbf{t} \in T_0 : B^{1/(s+2)}/k - 3 \leq |\mathbf{y}| \leq B^{1/(s+2)}/k\},$$

and

$$T_2 = \{\mathbf{t} \in T_0 : \min_j |y_j| \leq 3\}.$$

Thus using the previous lemma we see that

$$\begin{aligned} \sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} &= \sum_{\mathbf{y} \in X} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} + O\left(\sum_{\mathbf{y} \in T_1} + \sum_{\mathbf{y} \in T_2} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}}\right) \\ &= \int_{T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} d\mathbf{y} + O\left(\sum_{i=0}^2 E_i\right), \end{aligned}$$

where

$$E_0 = \sum_{\substack{\mathbf{y} \in \mathbb{Z}^s \cap T_0 \\ \min_j |y_j| \geq 2}} |\mathbf{y}|^{2-s} \min_j (|y_j|)^{-(s-3)/(s-1)} |\Delta(\mathbf{y})|^{-2/(s-1)},$$

and

$$E_i \ll \sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_i} |\mathbf{y}|^{-s} + \int_{T_i} |\mathbf{y}|^{-s} d\mathbf{y},$$

for $i = 1, 2$. In particular, one easily see that $E_i \ll 1$ for $i = 1, 2$, and a similar claim holds for E_0 , by using the same argument as in the proof of Lemma 4.3 in assuming that $\mathbf{y} \sim |\mathbf{y}|$. This concludes the proof. \square

Finally to conclude the proof of Lemma 5.1 it remains to investigate $J_2(B; k)$.

Lemma 5.4. *Let $k \leq B^{1/(s+2)}$. Then*

$$J_2(B; k) = (s - 2)\tau_\infty \log(B^{1/(s+2)}/k).$$

Proof. One immediately notice that $\varrho_\infty(\mathbf{y})$ is unchanged in response to permuting coordinates, and that $\varrho_\infty(\mathbf{y}) = |\mathbf{y}|^{-2} \varrho_\infty(t_1 \dots t_{s-1}, t_s)$, when $|y_s| = |\mathbf{y}|$ and $t_i = y_i/|\mathbf{y}|$ for $i \leq s - 1$. We then divide T_0 into subsections for which $|\mathbf{y}| = |y_j|$ and permute coordinates,

$$\begin{aligned} J_2(B; k) &= 2s \int_1^{B^{1/(s+2)}/k} \frac{1}{y_s^{s-2}} \int_{-y_s}^{y_s} \varrho_\infty(y_1, \dots, y_{s-1}, y_s) dy_1 \dots dy_{s-1} dy_s \\ &= 2s \int_1^{B^{1/(s+2)}/k} \frac{dy_s}{y_s} \int_{[-1,1]^{s-1}} \varrho_\infty(t_1 \dots t_s, 1) dt \\ &= 2s \log(B^{1/(s+2)}/k) \int_{[-1,1]^{s-1}} \varrho_\infty(t_1 \dots t_s, 1) dt, \end{aligned}$$

we also note that

$$\begin{aligned} \tau_\infty &= \int_{[-1,1]^s} \varrho_\infty(\mathbf{y}) d\mathbf{y} \\ &= 2s \int_0^1 y_s^{s-1-2} dy_s \int_{[-1,1]^{s-1}} \varrho_\infty(t_1 \dots t_{s-1}, 1) dt \\ &= 2s \frac{1}{s-2} \int_{[-1,1]^{s-1}} \varrho_\infty(t_1 \dots t_{s-1}, 1) dt, \end{aligned}$$

and therefore

$$\begin{aligned} J_2(B; k) &= 2s \log B^{1/(s+2)}/k \int_{[-1,1]^{s-1}} \varrho_\infty(t_1 \dots t_s, 1) dt \\ &= (s - 2)\tau_\infty \log(B^{1/(s+2)}/k). \end{aligned}$$

\square

Finally to finish the proof of lemma 5.1 we note that

$$\begin{aligned} M_2(B) &= \sum_{k \leq B^{1/(s+2)}} \frac{\mu(k)}{k^s} \sum_{\mathbf{y} \in \mathbb{Z}^s \cap T_0} \frac{\varrho_\infty(\mathbf{y})}{|\mathbf{y}|^{s-2}} \\ &= \sum_{k \leq B^{1/(s+2)}} \frac{\mu(k)}{k^s} J_2(B; k) + O(1), \end{aligned}$$

which follows from Lemma 5.3. From

$$\sum_{k \leq B^{1/(s+2)}} \frac{\mu(k)}{k^s} = \zeta(s)^{-1} + O(B^{-(s-1)/(s+2)}),$$

Lemma 5.4 implies that

$$M_2(B) = \frac{(s-2)}{(s+2)\zeta(s)} \log B + O(1).$$

Concluding the proof of lemma 5.1.

We then turn our attentions to $M_1(B)$. Again we remind the readers that

$$M_1(B) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{x}| \leq B^{4/(s+2)(s-1)}}} \frac{\sigma_\infty(\mathbf{x}) \mathfrak{S}(\mathbf{x})}{|\mathbf{x}|^{s-1}},$$

where

$$\begin{aligned} \sigma_\infty(\mathbf{x}) &= \int_{-\infty}^{\infty} \int_{[-1,1]^s} e(\theta F(\mathbf{x}; \mathbf{y})) d\mathbf{y} d\theta \\ \mathfrak{S}(\mathbf{x}) &= \sum_{q=1}^{\infty} q^{-s} S_q(\mathbf{x}), \quad S_q(\mathbf{x}) = \sum_{\substack{a \bmod q \\ (a,q)=1}} \sum_{\mathbf{b} \bmod q} e_q(aF(\mathbf{x}; \mathbf{b})). \end{aligned}$$

The remainder of this section is devoted to producing an estimate for $M_1(B)$.

Lemma 5.5. *For $s \geq 7$*

$$M_1(B) = \frac{4\zeta(s-2)}{(s+2)\zeta(s)\zeta(s-1)} \tau_\infty \log B + O(1).$$

We note that by reordering summation due to absolute convergence of the singular series $\mathfrak{S}(\mathbf{x})$ for $s \geq 7$, we have that

$$M_1(B) = \sum_{q=1}^{\infty} q^{-s} \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod q \\ (q, \mathbf{a})=1}} c_q(F(\mathbf{a}; \mathbf{b})) \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^s \\ |\mathbf{x}| \leq B^{4/(s+2)(s-1)} \\ \mathbf{x} \equiv \mathbf{a} \bmod q}} \frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}},$$

where $c_q(F(\mathbf{a}; \mathbf{b}))$ denotes the Ramanujan sum.

Again, one can use Mobius function to detect residual primitivity and then define, similar as to the previous subsection:

$$T_0 = T_0(k) = \{\mathbf{x} \in (\mathbb{R}_{\neq 0})^s : 1 \leq |\mathbf{x}| \leq B^{4/(s+2)(s-1)}/k\},$$

.Noting that since $\sigma_\infty(k\mathbf{x}) = k^{-1}\sigma_\infty(\mathbf{x})$, we have thus proved:

Lemma 5.6. *For $s \geq 7$,*

$$M_1(B) = \sum_{q=1}^{\infty} q^{-s} \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod q \\ (q, \mathbf{a})=1}} c_q(F(\mathbf{a}; \mathbf{b})) \sum_{\substack{k \leq B^{4/(s+2)(s-1)} \\ (k, q)=1}} \frac{\mu(k)}{k^s} \sum_{\substack{\mathbf{x} \in \mathbb{R}^s \cap T_0 \\ \mathbf{x} \equiv k^{-1}\mathbf{a} \bmod q}} \frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}}.$$

Similar as to the previous subsection, we try to compare \mathbf{x} sum into an integral.

Lemma 5.7. *If $\min_j |x_j| \geq 2q$ then we have*

$$\frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}} = q^{-s} \int_{[0, q]^s} \frac{\sigma_\infty(\mathbf{x} + \mathbf{t})}{|\mathbf{x} + \mathbf{t}|^{s-1}} d\mathbf{t} + O(q|\mathbf{x}|^{-(s-1)}(\min_j |x_j|)^{-1}|\Delta(\mathbf{x})|^{-1/s}).$$

Proof. Again we first try to prove an estimate for $\nabla\sigma_\infty(\mathbf{x})$. We will begin by showing that

$$\nabla\sigma_\infty(\mathbf{x}) \ll (\min_j |x_j|)^{-1}|\Delta(\mathbf{x})|^{-1/s}.$$

It suffices then to analyze the partial derivatives. One note that by definition,

$$\sigma_\infty(\mathbf{x}) = \int_{-\infty}^{\infty} \prod_{i=1}^s I(\theta x_i) d\theta,$$

where we write

$$I(\psi) = \int_{-1}^1 e(\psi y^2) dy,$$

One note that by [18], Lemma 4.4, we have that $I(\psi) \ll \min(1, |\psi|^{-1/2})$, and thus differentiating with respect to x_j gives

$$\frac{\partial}{\partial x_j} I(-\theta x_j) = \frac{1}{2x_j} \int_{-1}^1 y \frac{\partial}{\partial y} e(-\theta x_j y^2) dy.$$

One can see that the second integral is uniformly bounded through integration by parts, and therefore

$$\frac{\partial}{\partial x_j} I(-\theta x_j) \ll |x_j|^{-1}.$$

In particular we see that

$$\begin{aligned} \frac{\partial}{\partial x_j} \sigma_\infty(\mathbf{x}) &\ll |x_j|^{-1} \int_{-\infty}^{\infty} \min(1, |\theta|^{(s-1)/2} \prod_{i=1; i \neq j}^s |x_i|^{-1/2}) d\theta \\ &\ll |x_j|^{-1} \prod_{i=1; i \neq j}^s |x_i|^{-1/(s-1)} \\ &\ll (\min_j |x_j|)^{-1} |\Delta(\mathbf{x})|^{-1/s}. \end{aligned}$$

Similar as Lemma 5.2, we notice that

$$\begin{aligned} \frac{\sigma_\infty(\mathbf{x} + \mathbf{t})}{|\mathbf{x} + \mathbf{t}|^{s-1}} - \frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}} \\ = \frac{\sigma_\infty(\mathbf{x} + \mathbf{t}) - \sigma_\infty(\mathbf{x})}{|\mathbf{x} + \mathbf{t}|^{s-1}} + \sigma_\infty(\mathbf{x}) (|\mathbf{x} + \mathbf{t}|^{-(s-1)} - |\mathbf{x}|^{-(s-1)}). \end{aligned}$$

One immediately notice that if $|\mathbf{t}| \leq q$ and $\min_j |x_j| \geq 2q$, then by mean value theorem

$$\begin{aligned} \frac{\sigma_\infty(\mathbf{x} + \mathbf{t}) - \sigma_\infty(\mathbf{x})}{|\mathbf{x} + \mathbf{t}|^{s-1}} &\ll |\mathbf{x}|^{-(s-1)} q \sup_{0 \leq \xi \leq 1} \left| \frac{\partial}{\partial \xi} \sigma_\infty(\mathbf{y} + \xi \mathbf{t}) \right| \\ &\ll |\mathbf{x}|^{-(s-1)} q (\min_j |x_j|)^{-1} |\Delta(\mathbf{x})|^{-1/s}. \end{aligned}$$

We also note that again by mean value theorem,

$$|\mathbf{x} + \mathbf{t}|^{-(s-1)} - |\mathbf{x}|^{-(s-1)} \ll q |\mathbf{x}|^{-s},$$

and therefore by [7, chapter 8] we have that

$$\begin{aligned} \sigma_\infty(\mathbf{x}) (|\mathbf{x} + \mathbf{t}|^{-(s-1)} - |\mathbf{x}|^{-(s-1)}) &\ll q |\mathbf{x}|^{-s} \sigma_\infty(\mathbf{x}) \\ &\ll q |\mathbf{x}|^{-(s-1)} (\min_j |x_j|)^{-(s/2)} \ll q |\mathbf{x}|^{-(s-1)} (\min_j |x_j|)^{-1} |\Delta(\mathbf{x})|^{-1/s}, \end{aligned}$$

and the lemma follows by integrating and multiplying q^{-s} . \square

We can then convert the \mathbf{x} -sum into an integral.

Lemma 5.8. *For $s \geq 7$,*

$$\sum_{\substack{\mathbf{x} \in \mathbb{Z}^s \cap T_0 \\ \mathbf{x} \equiv k^{-1} \mathbf{a} \pmod{q}}} \frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}} = q^{-s} J_1(B; k) + O(q),$$

where

$$J_1(B; k) = \int_{T_0(k)} \frac{\sigma_\infty(\mathbf{y})}{|\mathbf{y}|^{s-1}} d\mathbf{y}.$$

Proof. Similar as before, we define

$$X = \{\mathbf{x} \in \mathbb{Z}^s : 2q \leq |\mathbf{x}| \leq B^{4/(s+2)(s-1)}/k - 2q; \mathbf{x} \equiv k^{-1}\mathbf{a} \pmod{q}\},$$

and let

$$Y = \bigcup_{\mathbf{x} \in X} \mathbf{x} + [0, q]^s,$$

then we note that $Y \in T_0$ and $T_0 \setminus Y \subset T_1 \cup T_2$, where

$$T_1 = \{\mathbf{t} \in T_0 : |\mathbf{t}| \leq 3q\}$$

$$T_2 = \{\mathbf{t} \in T_0 : B^{4/(s+2)(s-1)}/k - 3q \leq |\mathbf{t}| \leq B^{4/(s+2)(s-1)}/k\}.$$

Using previous lemma we obtain

$$\sum_{\substack{\mathbf{x} \in \mathbb{Z}^s \cap T_0 \\ \mathbf{x} \equiv k^{-1}\mathbf{a} \pmod{q}}} \frac{\sigma_\infty(\mathbf{x})}{|\mathbf{x}|^{s-1}} = q^{-s} J_1(B; k) + O\left(\sum_{i=1}^3 E_i\right),$$

with

$$E_0 = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^s \cap T_0 \\ \mathbf{x} \equiv k^{-1}\mathbf{a} \pmod{q} \\ \min |x_j| \geq 2q}} q |\mathbf{x}|^{-(s-1)} (\min_j |x_j|)^{-1} |\Delta(\mathbf{x})|^{-1/s}.$$

Using the estimate $|\sigma_\infty(\mathbf{x})| \ll |\mathbf{x}|^{-1/2}$, which holds for $s \geq 7$ (c.f. [4], chapter 8, page 46, where $\sigma_\infty(\mathbf{x})$ showed up as the constant term in $\mathcal{N}(P)$ for $s \geq 7$), we obtain

$$E_i = \sum_{\mathbf{x} \in \mathbb{Z}^s \cap T_i} |\mathbf{x}|^{-(s-1)} |\Delta(\mathbf{x})|^{-1/2} + q^{-s} \int_{T_i} |\mathbf{y}|^{-(s-1)} |\Delta(\mathbf{y})|^{-1/2} d\mathbf{y},$$

for $i = 1, 2$. Following the same procedure as in the proof of Lemma 4.3, we can assume $|x_j| \sim |\mathbf{x}|$ in evaluation of E_1 and E_2 , which, from the construction of T_i , implies that $E_i \ll q$ for $i = 1, 2$. Finally, notice that

$$\begin{aligned} E_0 &\ll q \sum_{\substack{\mathbf{x} \in \mathbb{Z}^s \cap T_0 \\ 2q \leq x_1 \leq x_2, \dots, x_{s-1} \leq x_s}} x_1^{-(s+1)/s} x_2^{-1/s} \dots x_{s-1}^{-1/s} x_s^{-(s-1)-1/s} \\ &\ll q^{(s-1)/s} \sum_{x_s \geq 2q} x_s^{-(s-1)-1/s+(s-1)(s-2)/s} \ll q^{(s-1)/s} \end{aligned}$$

Since the last sum converges. This is satisfactory for Lemma 5.8. \square

We can then combine everything to get

$$M_1(B) = \sum_{q=1}^{\infty} q^{-2s} \sum_{\substack{\mathbf{a}, \mathbf{b} \pmod{q} \\ (q, \mathbf{a})=1}} c_q(F(\mathbf{a}; \mathbf{b})) \sum_{\substack{k \leq B^{4/(s+2)(s-1)} \\ (k, q)=1}} \frac{\mu(k)}{k^s} J_1(B; k) + O(1).$$

Denote

$$\psi(q) = \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod q \\ (q, \mathbf{a})=1}} c_q(F(\mathbf{a}; \mathbf{b})),$$

we will determine how ψ behaves to give a final expression for $M_1(B)$.

Lemma 5.9. *ψ is multiplicative, and*

$$\psi(p^f) = \begin{cases} \varphi(p^f)p^{3sf/2}(1-p^{-s}) & \text{if } 2 \mid f, \\ 0 & \text{if } 2 \nmid f. \end{cases}$$

for positive integer f and prime p .

Proof. It's easy to see that, by construction and the Chinese remainder theorem, the function is multiplicative. Therefore, it suffices to investigate its behavior on prime powers. Specifically,

$$\begin{aligned} \psi(p^f) &= \sum_{\substack{c \bmod p^f \\ (c,p)=1}} \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod p^f \\ p \nmid \mathbf{a}}} e_{p^f}(cF(\mathbf{a}; \mathbf{b})) \\ &= \varphi(p^f) \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod p^f \\ p \nmid \mathbf{a}}} e_{p^f}(F(\mathbf{a}; \mathbf{b})) \\ &= \varphi(p^f) \sum_{\mathbf{a}, \mathbf{b} \bmod p^f} e_{p^f}(F(\mathbf{a}; \mathbf{b})) - \varphi(p^f) \sum_{\substack{\mathbf{a}, \mathbf{b} \bmod p^f \\ p \mid \mathbf{a}}} e_{p^f}(F(\mathbf{a}; \mathbf{b})). \end{aligned}$$

Denote $\psi_1(p^f) = \sum_{\mathbf{a}, \mathbf{b} \bmod p^f} e_{p^f}(F(\mathbf{a}; \mathbf{b}))$, we see that

$$\psi(p^f) = \varphi(p^f)\psi_1(p^f) - p^s\varphi(p^f)\psi_1(p^{f-1}),$$

in particular one should find that

$$\psi_1(p^f) = p^{sf} \#\{\mathbf{b} \bmod p^f : p^f \mid (b_1^2 \dots b_s^2)\} = p^{sf} p^{s[f/2]},$$

the lemma then follows through a simple computation. \square

We also get an estimate for $J_1(B; k)$, using similar techniques as Lemma 5.4.

Lemma 5.10. *For $s \geq 7$, we have*

$$J_1(B; k) = (s-1)\tau_\infty \log B^{4/(s+2)(s-1)}/k.$$

Proof. The proof is analogous to the one on $M_2(B)$. One begins by noticing that $\sigma_\infty(\mathbf{x}) = |\mathbf{x}|^{-1}\sigma_\infty(t_1 \dots t_{s-1}, 1)$, when $|x_s| = |\mathbf{x}|$ and $t_i = x_i/|\mathbf{x}|$. Again,

noticing that permuting coordinates doesn't change the value of the expression, we find that

$$\begin{aligned}
J_1(B; k) &= 2s \int_1^{B^{4/(s+2)(s-1)}/k} \int_{-x_s}^{x_s} \frac{\sigma_\infty(x_1 \dots x_{s-1}, x_s)}{|x_s|^{s-1}} dx_1 \dots dx_{s-1} dx_s \\
&= 2s \int_1^{B^{4/(s+2)(s-1)}/k} \frac{dx_s}{x_s} \int_{[-1,1]^{s-1}} \sigma_\infty(t_1 \dots t_{s-1}, 1) dt \\
&= 2s \log B^{4/(s+2)(s-1)}/k \int_{[-1,1]^{s-1}} \sigma_\infty(t_1 \dots t_{s-1}, 1) dt.
\end{aligned}$$

One also note that

$$\begin{aligned}
\tau_\infty &= \int_{[-1,1]^s} \sigma_\infty(\mathbf{x}) d\mathbf{x} \\
&= 2s \int_0^1 x_s^{s-2} \int_{[-1,1]^{s-1}} \sigma_\infty(t_1 \dots t_{s-1}, 1) dt \\
&= 2 \frac{s}{s-1} \int_{[-1,1]^{s-1}} \sigma_\infty(t_1 \dots t_{s-1}, 1) dt.
\end{aligned}$$

Equating the 2 expression gives the result

$$\begin{aligned}
J_1(B; k) &= 2s \log B^{4/(s+2)(s-1)}/k \int_{[-1,1]^{s-1}} \sigma_\infty(t_1 \dots t_{s-1}, 1) dt \\
&= (s-1) \tau_\infty \log B^{4/(s+2)(s-1)}/k.
\end{aligned}$$

□

The lemma shows that

$$\begin{aligned}
\sum_{\substack{k \leq B^{4/(s+2)(s-1)} \\ (k,q)=1}} \frac{\mu(k)}{k^s} J_1(B; k) &= (s-1) \tau_\infty \log B^{4/(s+2)(s-1)} \sum_{\substack{k \leq B^{4/(s+2)(s-1)} \\ (k,q)=1}} \frac{\mu(k)}{k^s} \\
&= (s-1) \tau_\infty \log B^{4/(s+2)(s-1)} \sum_{\substack{k=1 \\ (k,q)=1}}^{\infty} \frac{\mu(k)}{k^s} + O(B^{-4/(s+2)} \log B) \\
&= \frac{(s-1) \tau_\infty \log B^{4/(s+2)(s-1)}}{\zeta(s)} \prod_{p|q} (1 - p^{-s})^{-1} + O(B^{-4/(s+2)} \log B).
\end{aligned}$$

Finally we see that

$$M_1(B) = \frac{(s-1) \tau_\infty \log B^{4/(s+2)(s-1)}}{\zeta(s)} \sum_{q=1}^{\infty} q^{-2s} \psi(q) \prod_{p|q} (1 - p^{-s})^{-1} + O(1).$$

And using Lemma 5.9, one obtains

$$\begin{aligned}
 \sum_{q=1}^{\infty} q^{-2s} \psi(q) \prod_{p|q} (1 - p^{-s})^{-1} &= \prod_p \left(1 + \frac{p^{-4s} \psi(p^2) + p^{-8s} \psi(p^4) + \dots}{1 - p^{-s}} \right) \\
 &= \prod_p \left(1 + (1 - p^{-1}) \left(\sum_{f=1}^{\infty} p^{(2-s)f} \right) \right) \\
 &= \prod_p \left(1 + \frac{p-1}{p} \frac{p^{3-s} - p^{2-s}}{p - p^{3-s}} \right) \\
 &= \prod_p \left(\frac{1 - p^{1-s}}{1 - p^{2-s}} \right) \\
 &= \frac{\zeta(s-2)}{\zeta(s-1)}.
 \end{aligned}$$

We can then conclude from Lemma 5.6, 5.8, 5.9 and 5.10 that

$$M_1(B) = \frac{4\zeta(s-2)}{(s+2)\zeta(s)\zeta(s-1)} \tau_{\infty} \log B + O(1).$$

Combining with results from §4 we see that

$$\begin{aligned}
 N(\Omega, B) &= \frac{B}{4\zeta(s-2)} M_1(B) + \frac{B}{4\zeta(s-1)} M_2(B) + O(B^{1-\eta} \log B) \\
 &\sim \frac{1}{4\zeta(s)\zeta(s-1)} \tau_{\infty} B \log B,
 \end{aligned}$$

as $B \rightarrow \infty$.

To complete the proof of the theorem we finally need to show that the leading constant agrees with the constant c_{Peyre} predicted by Peyre in [12]. According to Schindler [14], §3,

$$c = \frac{1}{4\zeta(s-1)\zeta(s-2)} \tau_{\infty} \prod_p \sigma_p,$$

where τ_{∞} is as previously defined in Theorem 1.1 and

$$\sigma_p = \lim_{l \rightarrow \infty} p^{-(2s-1)l} \#\{(\mathbf{x}, \mathbf{y}) \bmod p^l : F(x; y) \equiv 0 \bmod p^l\}.$$

We note that

$$\begin{aligned} n(p^l) &= \sum_{j=0}^l \sum_{\substack{y \bmod p^l \\ (y, p^l) = p^j}} \#\{\mathbf{x} \in (\mathbb{Z}/p^l\mathbb{Z})^s : F(x; y) \equiv 0 \pmod{p^l}\} \\ &= \sum_{j=0}^{\lfloor l/2 \rfloor} \sum_{\substack{(u, p) = 1 \\ u \bmod p^{l-j}}} \#\{\mathbf{x} \in (\mathbb{Z}/p^l\mathbb{Z})^s : F(x; u) \equiv 0 \pmod{p^{l-2j}}\} + O(p^{3s/2l}). \end{aligned}$$

Here we note that since $(u, p^j) = 1$, the number of solutions for each $x_i \in \mathbb{Z}/p^{l-2j}\mathbb{Z}$ is $p^{(l-2j)(s-1)}$, and thus we have

$$\begin{aligned} &= \sum_{j=0}^{\lfloor l/2 \rfloor} \sum_{\substack{(u, p) = 1 \\ u \bmod p^{l-j}}} p^{(l-2j)(s-1)} p^{2sj} + O(p^{3s/2l}) \\ &= \sum_{j=0}^{\lfloor l/2 \rfloor} (p^{s(l-j)} - p^{s(l-j)-s}) p^{(l-2j)(s-1)} p^{2sj} + O(p^{3s/2l}) \\ &= (1 - p^{-s}) \sum_{j=0}^{\lfloor l/2 \rfloor} p^{2sl-l+(2-s)j} + O(p^{3s/2l}) \\ &= \frac{1 - p^{-s}}{1 - p^{s-2}} p^{(2s-1)l} + O(p^{3s/2l}). \end{aligned}$$

In particular, taking it into limit as $l \rightarrow \infty$, we see that for $s \geq 7$,

$$\begin{aligned} c &= \frac{1}{4\zeta(s-1)\zeta(s-2)} \tau_\infty \prod_p \lim_{l \rightarrow \infty} p^{-(2s-1)l} n(p^l) \\ &= \frac{1}{4\zeta(s-1)\zeta(s)} \tau_\infty. \end{aligned}$$

Thus our prediction agrees with the constant predicted by Peyre. This proves the claim and Theorem 1.1 follows.

REFERENCES

- [1] V.V. Batyrev and Y. Tschinkel, *Rational points on some Fano cubic bundles*, C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), 41–46.
- [2] T.D. Browning and L .Q. Hu, *Counting rational points on biquadratic hypersurfaces* (2018).
- [3] T.D. Browning and D.R. Heath-Brown, *Density of Rational Points on a Quadric Bundle in $\mathbb{P}^s \times \mathbb{P}^s$* (2020).

- [4] H. Davenport, *Analytic Methods for Diophantine Equations and Diophantine Inequalities*, Cambridge Mathematical Library, 2005.
- [5] J. Franke and Y. I. Manin and Y. Tschinkel, *Rational points of bounded height on Fano varieties*, *Inventiones Mathematicae* **95** (1989), 421–435.
- [6] D. R. Heath-Brown, *A new form of the circle method, and its application to quadratic forms*, *J. Reine Angew. Math.* **481** (1996), 149–206.
- [7] ———, *Cubic forms in 14 variables*, *Invent. math.* **170** (2007), 199–230.
- [8] D.R. Heath-Brown, *The density of rational points on curves and surfaces*, Vol. 155, *Annals of Mathematics*, 2002.
- [9] H. Iwaniek, *Topics in Classical Automorphic Forms*, Amer. Math. Soc., 1997.
- [10] P. Le Boudec, *Density of rational points on a certain smooth bihomogeneous threefold*, *Int. Math. Res. Not. IMRN* **21** (2015), 10703–10715.
- [11] H. Minkowski, *Geometrie der Zahlen*, Teubner, Leipzig-Berlin (1896), 199–218.
- [12] E. Peyre, *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, *Duke Math Journal* **79** (1995).
- [13] M. Robbiani, *On the number of rational points of bounded height on smooth bilinear hypersurfaces in biprojective space.*, *Journal of the London Mathematical Society* (2) **63** (2001), 33–51.
- [14] D. Schindler, *Manin's conjecture for certain biprojective hypersurfaces*, *J. reine angew* **714** (2016).
- [15] W.M. Schmidt, *Asymptotic formulae for point lattices of bounded determinant and subspaces of bounded height*, *Duke Math Journal* **35** (1968).
- [16] C. V. Spencer, *The Manin conjecture for $x_0y_0 + \dots + x_sy_s = 0$* , *Journal of Number Theory* **129** (2009), 1505–1521.
- [17] J. L. Thunder, *Asymptotic estimates for rational points of bounded height on flag varieties*, *Compositio Mathematica* **88** (1993), 155–186.
- [18] E.C. Titchmarsh and D.R.Heath-Brown, *The theory of the Riemann zeta-function. 2nd ed.*, Oxford University Press, 1986.