

Sparse High-Dimensional Vector Autoregressive Bootstrap*

Robert Adamek^a, Stephan Smeekes^b, and Ines Wilms^b

^aDepartment of Economics and Business Economics, Aarhus University

^bDepartment of Quantitative Economics, Maastricht University
r.adamek@econ.au.dk, {s.smeekes, i.wilms}@maastrichtuniversity.nl

February 3, 2023

Abstract

We introduce a high-dimensional multiplier bootstrap for time series data based capturing dependence through a sparsely estimated vector autoregressive model. We prove its consistency for inference on high-dimensional means under two different moment assumptions on the errors, namely sub-gaussian moments and a finite number of absolute moments. In establishing these results, we derive a Gaussian approximation for the maximum mean of a linear process, which may be of independent interest.

JEL codes: C15, C32, C55;

Keywords: High-dimensional data, Time series, Bootstrap, vector autoregression, linear process.

1 Introduction

We introduce theory for bootstrapping the distribution of high-dimensional means of sparse, finite order, stationary vector autoregressive (VAR) processes. For an N -dimensional vector of time series $\mathbf{x}_t = (x_{1,t}, \dots, x_{N,t})'$, we provide an approximation for the distribution of $\max_{1 \leq j \leq N} \left| \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t} \right|$, where the number of variables N is potentially much larger than the sample size T , and can asymptotically grow faster than T . This prototypical statistic is commonly considered in high-dimensional settings, see e.g. the closely related work of Chernozhukov et al. (2013), Chernozhukov et al. (2017), Zhang and Wu (2017), Chernozhukov et al. (2020), Giessing and Fan (2020), or the review by Chernozhukov et al. (2022), who investigate the properties of this estimator for independent data. In this paper, we extend these results to high-dimensional linear processes, including stationary VARs. Related work in time series settings include Zhang and Cheng (2018), who provide Gaussian approximations in the general framework of functional dependence of Wu (2005).

The VAR sieve bootstrap is well-known in the low-dimensional time series bootstrapping literature, see e.g. Paparoditis (1996), Park (2002), Chang and Park (2003), Meyer and Kreiss (2015),

*The first and second author were financially supported by the Dutch Research Council (NWO) under grant number 452-17-010.

and Section 12.2 of [Kilian and Lütkepohl \(2017\)](#). It fits a VAR to the time series data, resamples the residuals of the estimated VAR, and re-applies the VAR recursively to place the dependence back into the bootstrap sample. Under appropriate conditions, the VAR sieve bootstrap allows for valid inference. We extend this approach to high dimensions where the VAR is estimated by the lasso ([Tibshirani, 1996](#)) or another sparse estimation method, and use a multiplier (or wild) bootstrap to resample the residuals. Our work is related to that of [Trapani \(2013\)](#), [Bi et al. \(2021\)](#) and [Krampe et al. \(2021\)](#). The two former papers assume a dense structure on the data, and apply the VAR sieve bootstrap to a low-dimensional set of factors. The latter consider a sparse setting, providing bootstrap inference for desparsified estimators of VAR coefficients. We assume a data-generating process (DGP) similar to the one considered in [Krampe et al. \(2021\)](#).

All theoretical results in this paper are established under two different sets of assumptions on the errors. First, we assume the errors have sub-gaussian moments, which generally allows N to grow at an exponential rate of T . Second, we assume that the errors have some finite number of absolute moments, which effectively restricts the growth of N to some polynomial rate of T . In Section 2, we introduce the multiplier bootstrap for sparsely estimated high-dimensional VARs. In Section 3, we start by providing a high-dimensional central limit theorem (HDCLT) for linear processes in Theorem 1, which may be of independent interest. In Section 4, we introduce the stationary VAR model, and show that under consistent estimation, the long run covariance structure is recovered with high probability. Theorem 2 provides a consistency result for the covariance matrix. In Section 5, we show that the bootstrap's behaviour is asymptotically similar to that of the original sample. In particular, Theorem 3 provides a HDCLT for the bootstrap process which mirrors that of Theorem 1, and Theorem 4 shows consistency of the bootstrap. Section 6 then shows how these results can be used to establish validity of inference in VARs estimated by the lasso.

Notation. For a random variable x , $\|x\|_{L_p} = (\mathbb{E} |x|^p)^{1/p}$, $\|x\|_{\psi_2} = \inf \left\{ c > 0 : \mathbb{E} \exp(|x|^2/c^2) \leq 2 \right\}$ denote the L_p and Orlicz norms. For any N dimensional vector \mathbf{x} , $\|\mathbf{x}\|_p = \left(\sum_{j=1}^N |x_j|^p \right)^{1/p}$ denotes the p -norm, with the familiar convention that $\|\mathbf{x}\|_0 = \sum_i 1(|x_i| > 0)$ and $\|\mathbf{x}\|_\infty = \max_i |x_i|$. For a matrix \mathbf{A} , we let $\|\mathbf{A}\|_p = \max_{\|\mathbf{x}\|_p=1} \|\mathbf{A}\mathbf{x}\|_p$ for any $p \in [0, \infty]$ and $\|\mathbf{A}\|_{\max} = \max_{i,j} |a_{i,j}|$. $\Lambda_{\min}(\mathbf{A})$ and $\Lambda_{\max}(\mathbf{A})$ denote the smallest and largest eigenvalues of \mathbf{A} , and $\rho(\mathbf{A})$ the spectral radius of \mathbf{A} , i.e. the largest absolute eigenvalue of \mathbf{A} , or equivalently $\rho(\mathbf{A}) = \lim_{k \rightarrow \infty} \|\mathbf{A}^k\|^{1/k}$ for any induced norm $\|\cdot\|$. For \mathbf{A} a square matrix, we let its zero-th power $\mathbf{A}^0 = \mathbf{I}$. We use \xrightarrow{p} and \xrightarrow{d} to denote convergence in probability and distribution respectively. Depending on the context, \sim denotes equivalence in order of magnitude of sequences, or equivalence in distribution. We frequently make

use of arbitrary positive finite constants C (or its sub-indexed version C_i) whose values may change from line to line throughout the paper, but they are always independent of the time and cross-sectional dimension. Similarly, generic sequences converging to zero as $T \rightarrow \infty$ are denoted by η_T (or its sub-indexed version $\eta_{i,t}$). When they are used, it should be understood that there exists some constant C or sequence $\eta_T \rightarrow 0$ such that the given statement holds.

2 Vector Autoregressive Bootstrap

We introduce our proposed bootstrap procedure for sparsely estimated high-dimensional VARs and subsequently discuss how it can be used to perform inference on high-dimensional time series.

2.1 Bootstrap for High-Dimensional VARs

Let \mathbf{x}_t be an N -dimensional time series process. We assume the data is generated by a stationary, finite order, high-dimensional VAR(K) model

$$\mathbf{x}_t = \sum_{k=1}^K \mathbf{A}_k \mathbf{x}_{t-k} + \boldsymbol{\epsilon}_t, \quad t \in \mathbb{Z}, \quad (1)$$

with autoregressive parameter matrices \mathbf{A}_k ($k = 1, \dots, K$), and independent errors $\boldsymbol{\epsilon}_t$ with $\mathbb{E}\boldsymbol{\epsilon}_t = \mathbf{0}$ and covariance matrix $\boldsymbol{\Sigma}_\epsilon := \frac{1}{T} \sum_{t=1}^T \mathbb{E}\boldsymbol{\epsilon}_t \boldsymbol{\epsilon}_t'$. We can re-write eq. (1) as a collection of linear equations

$$x_{j,t} = \sum_{k=1}^K \mathbf{a}_{j,k} \mathbf{x}_{t-k} + \epsilon_{j,t} = \underset{1 \times KN}{\boldsymbol{\beta}'_j} \underset{KN \times 1}{\boldsymbol{\mathcal{X}}_t} + \epsilon_{j,t}, \quad j = 1, \dots, N, \quad t \in \mathbb{Z},$$

where $\mathbf{a}_{j,k}$ is the j th row of \mathbf{A}_k , $\boldsymbol{\beta}_j = (\mathbf{a}_{j,1}, \dots, \mathbf{a}_{j,K})'$, and $\boldsymbol{\mathcal{X}}_t = (\mathbf{x}'_{t-1}, \dots, \mathbf{x}'_{t-K})'$. We observe a sub-sequence of length $T + K$ from the process \mathbf{x}_t , indexed by $t = -K + 1, \dots, T$, which we can denote in a stacked matrix $\underset{(T+K) \times N}{\mathbf{X}} = (\mathbf{x}'_{-K+1}, \dots, \mathbf{x}'_T)'$.

The lasso estimator of equation j is defined as

$$\hat{\boldsymbol{\beta}}_j = \arg \min_{\boldsymbol{\beta}_j^* \in \mathbb{R}^{KN}} \frac{1}{T} \sum_{t=1}^T (x_{j,t} - \boldsymbol{\beta}_j^{*'} \boldsymbol{\mathcal{X}}_t)^2 + 2\lambda_j \|\boldsymbol{\beta}_j^*\|_1, \quad (2)$$

where λ_j is a tuning parameter that determines the degree of penalization in equation j , and can be selected independently in each equation. For tuning parameter selection, one could use the iterative plug-in procedure described in Section 5.1 of [Adamek et al. \(2022\)](#), information criteria or time series cross-validation.

Once all equations $j = 1, \dots, N$ are estimated by the lasso, we collect the VAR coefficient

Algorithm 1: VAR Multiplier Bootstrap

- 1 Let $\hat{\mathbf{A}}_1, \dots, \hat{\mathbf{A}}_K$ be the lasso estimates;
 - 2 Set $\hat{\boldsymbol{\epsilon}}_t = \mathbf{x}_t - \sum_{k=1}^K \hat{\mathbf{A}}_k \mathbf{x}_{t-k}$ for $t = 1, \dots, T$;
 - 3 **for** $b \in \{1, \dots, B\}$ **do**
 - 4 Generate $\gamma_1, \dots, \gamma_T$ from a $N(0, 1)$ distribution;
 - 5 Set $\boldsymbol{\epsilon}_t^* = \hat{\boldsymbol{\epsilon}}_t \gamma_t$ for $t = 1, \dots, T$;
 - 6 Let $\mathbf{x}_t^* = \mathbf{x}_t$ for $t = -K + 1, \dots, 0$;
 - 7 Build \mathbf{x}_t^* recursively from $\mathbf{x}_t^* = \sum_{k=1}^K \hat{\mathbf{A}}_k \mathbf{x}_{t-k}^* + \boldsymbol{\epsilon}_t^*$ for $t = 1, \dots, T$;
 - 8 Compute and store the statistic $Q^{*b} = \max_{1 \leq j \leq N} \left| \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t}^* \right|$
-

estimates as follows

$$\begin{bmatrix} \hat{\mathbf{A}}_1 & \dots & \hat{\mathbf{A}}_k \end{bmatrix} = \begin{bmatrix} \hat{\boldsymbol{\beta}}'_1 \\ \vdots \\ \hat{\boldsymbol{\beta}}'_N \end{bmatrix}.$$

Our object of interest is the scaled high-dimensional mean

$$Q = \max_{1 \leq j \leq N} \left| \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t} \right|$$

of the sparse VAR. To approximate its distribution, we apply the VAR multiplier bootstrap summarized in Algorithm 1. When B is sufficiently large, the CDF of Q can be approximated by the quantiles of the ordered statistics $Q^{*(1)}, \dots, Q^{*(B)}$. Note that while we derive results for the maximum absolute mean, this bootstrap procedure is equally valid for statistics such as $\max_{1 \leq j \leq N} \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t}$ or $\min_{1 \leq j \leq N} \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t}$, which would allow for one-sided tests, or tests with an asymmetric rejection region.

Remark 1. So far, we treated the number of lags K in the VAR as known, which is typically not the case in practice. Indeed, Algorithm 1 requires one to choose K . One of the lasso's advantages is that it performs well when the number of regressors is large, provided the parameters are sparse. This means it is less harmful to include many redundant lags, compared to low-dimensional estimation methods which suffer in terms of efficiency. Therefore, if the practitioner believes the true VAR order is some $K \leq K_{\max}$, one may simply take $K = K_{\max}$, and let the lasso penalize any redundant lags to 0. In the absence of such an upper bound, one may use autocorrelation tests on $\hat{\boldsymbol{\epsilon}}_t$ as a guide, since we assume that the errors $\boldsymbol{\epsilon}_t$ are independent. As the number of equations is large in this setting, one should use a multiple testing correction (e.g. [Holm, 1979](#)); the conservative nature

of such corrections is not a large issue for the lasso. Alternatively, one could use the hierarchical lag structure approach of [Nicholson et al. \(2020\)](#) that embeds lag selection into the estimation procedure.

Remark 2. It may happen that the estimated VAR is not stationary, even if the true underlying process is. Proper functioning of our method requires, however, that the bootstrap process is stationary. stationary estimates, such as Yule-Walker estimation. However, to our knowledge, a similar method has not yet been proposed for high-dimensional settings. We suggest, in case of non-stationarity, to manually correct the estimates by uniformly shrinking all entries of $\hat{\mathbf{A}}_1, \dots, \hat{\mathbf{A}}_K$ towards 0 to ensure stationarity of the bootstrap process. We elaborate on this correction in [Section 4](#), and justify that it is asymptotically negligible.

2.2 Bootstrap Inference on (Approximate) Means

Statistics such as the scaled mean Q are useful in high-dimensional settings, since they allow us to simultaneously test a high-dimensional set of hypotheses. For example, let $\mu_j = \mathbb{E}x_{j,t}$ be the means of a high-dimensional stationary autoregressive process, and assume we are interested in testing the hypothesis

$$H_0 : \mu_1 = \dots = \mu_N = 0 \text{ vs. } H_1 : \mu_j \neq 0 \text{ for at least one } j.$$

Under the null hypothesis, this process follows [eq. \(1\)](#), which allows us to directly test the null using the quantiles of $Q^{*(1)}, \dots, Q^{*(B)}$. Specifically, one would reject the null at significance level α if $Q > Q^{*(B[1-\alpha])}$ for any j . To know for *which* means the null can be rejected, one can use the stepdown procedure of [Romano and Wolf \(2005\)](#), as detailed in [Algorithm 2](#) in line with the description in [Section 5](#) of [Chernozhukov et al. \(2013\)](#). Importantly, this procedure is asymptotically exact – non-conservative – as it takes into account the possible correlations amongst statistics, instead of using the conservative worst case of independence.

More generally, this bootstrap procedure can be used to test any high-dimensional set of hypotheses, provided its test statistic can be expressed as an approximate mean, that is, $\frac{1}{\sqrt{T}} \sum_{t=1}^T x_{j,t} + o_p(1)$. While we do not formally consider this extension here, we can adapt the arguments in [Section 5](#) of [Chernozhukov et al. \(2013\)](#) (which do not rely on independent data) to establish this result in our context as well. This opens up the way for applications to statistics that are much more general than just sample means, as many statistics of practical interest, such as (high-dimensional) regression estimates, can be written in this form. Our results therefore form a first step towards a more general bootstrap theory for high-dimensional inference using VAR models on statistics that

Algorithm 2: Stepdown Procedure

- 1 Estimate a VAR model with the lasso in eq. (2) and obtain $\hat{\mathbf{A}}_1, \dots, \hat{\mathbf{A}}_K$;
 - 2 Use Algorithm 1 to compute the bootstrap means $\bar{x}_j^* = \frac{1}{T} \sum_{t=1}^T x_{j,t}^*$, and store $\bar{x}_j^{*(1)}, \dots, \bar{x}_j^{*(B)}$ for all $j = 1, \dots, N$;
 - 3 Let the index set $\mathcal{S} = \{1, \dots, N\}$;
 - 4 **while** $\mathcal{S} \neq \emptyset$ **do**
 - 5 Compute the bootstrap statistics $Q_{\mathcal{S}}^{*b} = \max_{j \in \mathcal{S}} \left| \sqrt{T} \bar{x}_j^* \right|$, and store $Q_{\mathcal{S}}^{*(1)}, \dots, Q_{\mathcal{S}}^{*(B)}$;
 - 6 Set $\mathcal{R} = \left\{ k \in \mathcal{S} : \left| \frac{1}{\sqrt{T}} \sum_{t=1}^T x_{k,t} \right| > Q_{\mathcal{S}}^{*(B[1-\alpha])} \right\}$ and $\mathcal{S} = \mathcal{S} \setminus \mathcal{R}$;
 - 7 **if** $\mathcal{R} = \emptyset$ **then**
 - 8 **break**;
 - 9 Reject $H_0 : \mu_j = 0$ for all $j \in \mathcal{S}^c$;
-

can be well-approximated by the mean of a linear process.

3 HDCLT for Linear Processes

In this section, we establish a high-dimensional CLT for linear processes, which is a useful result in its own right, but also a vital building block to establish theoretical results for the bootstrap. We therefore give it a self-contained treatment in this section, before applying it to the VAR process in eq. (1) and covering the theory for the bootstrap in the following sections.

Under appropriate invertibility conditions, it is well-known that the VAR process in eq. (1) can be written as the following infinite order vector moving average (VMA)

$$\mathbf{x}_t = \sum_{k=0}^{\infty} \mathbf{B}_k \boldsymbol{\epsilon}_{t-k} = \mathcal{B}(L) \boldsymbol{\epsilon}_t, \quad t \in \mathbb{Z}, \quad (3)$$

where $\mathcal{B}(z) = \sum_{k=0}^{\infty} \mathbf{B}_k z^k = \left(\mathbf{I} - \sum_{k=1}^K \mathbf{A}_k z^k \right)^{-1}$, and L is the lag operator. We derive a Gaussian approximation for linear processes of the form in eq. (3), which builds on and extends similar approximations for independent and identically distributed (i.i.d.) processes by Chernozhukov et al. (2020) and others (see Section 1).

Specifically, we show that the distribution of $\max_{1 \leq j \leq N} \left| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right| = \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty}$ can be asymptotically approximated by $\|z\|_{\infty}$, with $z \sim N(\mathbf{0}, \boldsymbol{\Sigma})$ and $\boldsymbol{\Sigma}$ an appropriate covariance matrix. This result parallels well-known results in low-dimensional settings, where scaled means of linear processes converge in distribution to a Gaussian random variable as $T \rightarrow \infty$. However, in our high-dimensional setting, we consider the case where N and T diverge simultaneously, and $\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty}$ does not converge to a well defined limit; the maximum over a growing number of elements generally also grows. As such, we instead show that their distributions grow closer together asymptotically,

in the sense that the Kolmogorov distance between $\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty}$ and $\|\mathbf{z}\|_{\infty}$ converges to 0. Even though to our knowledge, there does not exist a closed-form expression for the CDF of $\|\mathbf{z}\|_{\infty}$, it can be approximated for any N by Monte Carlo simulation, making it a useful asymptotic approximation in practice.

The broad sketch of our proof is as follows. We use the Beveridge-Nelson decomposition to write

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t = \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \boldsymbol{\epsilon}_t - \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) (\boldsymbol{\epsilon}_T - \boldsymbol{\epsilon}_0), \quad (4)$$

where $\tilde{\mathcal{B}}(z) = \sum_{j=0}^{\infty} \sum_{k=j+1}^{\infty} \mathbf{B}_k z^j$. The first term is a scaled sum of independent errors with covariance matrix $\boldsymbol{\Sigma} := \mathcal{B}(1) \boldsymbol{\Sigma}_{\boldsymbol{\epsilon}} \mathcal{B}(1)'$, $\sigma_j^2 := \boldsymbol{\Sigma}_{(j,j)}$, and can therefore be approximated by a Gaussian maximum thanks to Chernozhukov et al. (2020) when $\boldsymbol{\Sigma}$ is non-degenerate and the $\boldsymbol{\epsilon}_t$'s satisfy certain moment conditions (see Lemma A.2). The second term involves only two errors at $t = 0$ and $t = T$, and is an asymptotically negligible leftover under certain summability conditions on the VMA coefficient matrices \mathbf{B}_k (see Lemma A.3). We therefore make the following assumptions:

Assumption 1. Let $\Lambda_{\min}(\boldsymbol{\Sigma}) \geq 1/C$ and $\max_{1 \leq j \leq N} \sigma_j \leq C$.

Assumption 2. Let the vector $\boldsymbol{\epsilon}_t$ satisfy *one* of the following moment conditions

1. $\max_{j,t} \|\boldsymbol{\epsilon}_{j,t}\|_{\psi_2} \leq C$.
2. $\max_{j,t} \|\boldsymbol{\epsilon}_{j,t}\|_{L_m} \leq C$, for some constant $m \geq 4$.

We derive our results under two different moment assumptions. In Assumption 2.1 we require that the errors are uniformly sub-gaussian over j and t ; or in Assumption 2.2 that the moments possess some number (m) of finite absolute moments. By equation (2.15) in Vershynin (2019), Assumption 2.2 follows automatically for all m from Assumption 2.1, making the latter a considerably less stringent assumption. Under these assumptions, Theorem 1 provides an upper bound on the Kolmogorov distance between our statistic of interest and a Gaussian maximum:

Theorem 1 (Gaussian approximation for linear processes). *Consider a linear process \mathbf{x}_t as in eq. (3), let Assumption 1 hold, and define $\tilde{S} := \sum_{j=0}^{\infty} \|\mathbf{B}_j\|_{\infty}$, $S_m := \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \|\mathbf{B}_k\|_{\infty} \right)^m$, and*

$$J_{N,T} := \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} \leq y \right) - \mathbb{P} (\|\mathbf{z}\|_{\infty} \leq y) \right|,$$

where $\mathbf{z} \sim N(\mathbf{0}, \boldsymbol{\Sigma})$.

1. Under Assumption 2.1,

$$J_{N,T} \leq C \left(\frac{(\tilde{S}d_N)^2 \log(N)^{3/2} \log(T)}{\sqrt{T}} + \frac{(\tilde{S}d_N)^2 \log(N)^2}{\sqrt{T}} + \frac{\log(N)d_N\sqrt{S_2}}{\sqrt{T}} + \frac{1}{\log(N)} \right),$$

where $d_N = C\sqrt{\log(N)}$.

2. Under Assumption 2.2,

$$J_{N,T} \leq C \left(\frac{(\tilde{S}d_N)^2 (\log N)^{3/2} \log(T)}{\sqrt{T}} + \frac{(\tilde{S}d_N)^4 \log(N)^2 \log(T)}{T^{1-2/m}} + \left[\frac{(\tilde{S}d_N)^{2m} \log(N)^{3m/2-4} \log(T) \log(NT)}{T^{m/2-1}} \right]^{\frac{1}{m-2}} + (Nd_N^m S_m)^{\frac{1}{m+1}} \left[\frac{\sqrt{\log(N)}}{\sqrt{T}} \right]^{\frac{m}{m+1}} \right),$$

where $d_N = CN^{1/m}$.

Under Assumption 2.1, convergence of this upper bound to 0 depends on the size of the terms \tilde{S} and S_2 , and the relative growth rates of N and T . As N only enters in logs compared to \sqrt{T} in the denominator, it is possible to have N grow at some exponential rate of T . Under Assumption 2.2, N enters the numerator at a polynomial rate through the sequence d_N ; this effectively restricts the growth rate of N to some polynomial of T , though it can still grow faster than T when m is sufficiently large. Our results under these two sets of assumptions therefore mainly differ (apart from the different proof strategies required for each case), in this regard: if exponential growth of N is desirable, we need finite exponential moments of ϵ_t ; whereas if polynomial growth of N is sufficient, we only need finite polynomial moments of ϵ_t .

4 Application to VAR Models

Theorem 1 is a key building block in our derivations for the bootstrap, as it can be applied to our VAR in eq. (1) under appropriate conditions. In this section, we explain our assumptions on the VAR process, and on the consistency properties of lasso estimation. While the lasso is our running example, the following theoretical results do not rely on the lasso specifically, and are equally valid for any other estimation method which satisfies our consistency conditions. We return to the lasso in Section 6, where we show examples of it satisfying these conditions.

For the following exposition, it is useful to define the companion matrix

$$\mathbb{A} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \dots & \mathbf{A}_K \\ \mathbf{I} & \mathbf{0} & \dots & \mathbf{0} \\ \vdots & \ddots & & \vdots \\ \mathbf{0} & \dots & \mathbf{I} & \mathbf{0} \end{pmatrix}.$$

of the VAR in eq. (1). This matrix allows us to re-write the VAR(K) as a VAR(1) with

$$\mathcal{X}_t = \mathbb{A}\mathcal{X}_{t-1} + \begin{bmatrix} \boldsymbol{\epsilon}_t \\ \mathbf{0} \end{bmatrix},$$

and allows for a simple expression for the corresponding VMA coefficients in eq. (3): $\mathbf{B}_k = \mathbf{J}\mathbb{A}^k\mathbf{J}'$, where $\mathbf{J}_{N \times KN} = (\mathbf{I}, \mathbf{0}, \dots, \mathbf{0})$.¹ This inversion is only possible if the VAR is invertible.

Assumption 3. Let $\|\mathbb{A}^j\|_\infty \leq \psi_N \lambda^j$, for some $\lambda \leq C < 1$, all $j \in \mathbb{N}_0$, and $1 \leq \psi_N < \infty$ a sequence potentially growing as $N \rightarrow \infty$.

Assumption 3 is based on Assumption 1(ii) of [Krampe et al. \(2021\)](#), and its purpose is twofold. First, it implies that the VAR process in eq. (1) is stationary, since $\rho(\mathbb{A}) = \lim_{k \rightarrow \infty} \|\mathbb{A}^k\|_\infty^{1/k} \leq \lim_{k \rightarrow \infty} (\psi_N \lambda^k)^{1/k} = \lambda$, and it can therefore be inverted into a VMA. Second, it allows us to derive summability properties for the quantities \tilde{S} and S_m in Section 3, since $\|\mathbf{B}_j\|_\infty \leq \|\mathbb{A}^j\|_\infty \leq \psi_N \lambda^j$. The sequence ψ_N controls the growth rate of the $\|\cdot\|_\infty$ as the dimension of \mathbb{A} grows. For large j , this becomes a non-issue as \mathbb{A}^j approaches the zero matrix when $\rho(\mathbb{A}) < 1$. However, the power beyond which the norm becomes smaller than λ can generally grow with N , see Example 2.3 of [Liu and Zhang \(2021\)](#). As such, we allow for the possibility that this uniform bound on all powers of \mathbb{A} also potentially grows with N .

Next, we make the following assumptions about consistency of the estimators $\hat{\mathbb{A}}$, and the residuals $\hat{\boldsymbol{\epsilon}}_t$:

Assumption 4. For some sequences $\xi_{N,T}, \psi_N$, define the set $\mathcal{P} := \left\{ \left\| \hat{\mathbb{A}} - \mathbb{A} \right\|_\infty \leq \xi_{N,T} \psi_N \right\}$. Assume that $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{P}) = 1$.

Assumption 5. For some sequence $\phi_{N,T}$, define the set $\mathcal{Q} := \left\{ \max_{1 \leq j \leq N} \frac{1}{T} \|\hat{\boldsymbol{\epsilon}}_j - \boldsymbol{\epsilon}_j\|_2^2 \leq \phi_{N,T} \right\}$, where $\boldsymbol{\epsilon}_j = (\epsilon_{j,1}, \dots, \epsilon_{j,T})'$ and similarly for $\hat{\boldsymbol{\epsilon}}_j$. Assume that $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{Q}) = 1$.

While we leave the bounds $\psi_N \xi_{N,T}$ and $\phi_{N,T}$ unspecified and derive later results in terms of these sequences, the reader may think of them as $\psi_N \xi_{N,T}$ converging at a rate close to $\frac{1}{\sqrt{T}}$ and $\phi_{N,T}$ close to $\frac{1}{T}$, which will be shown later in Section 6.

¹See page 279 of [Papadoditis \(1996\)](#).

In our proof strategy, we make use of the probabilistic sets denoted by calligraphic letters \mathcal{P} to \mathcal{U} . They describe events involving functions of the random variables in \mathbf{X} and $\boldsymbol{\epsilon}_t$, and can therefore only hold with a certain probability. For the sets \mathcal{P} and \mathcal{Q} , we simply assume that they hold with probability converging to 1 as $N, T \rightarrow \infty$. For the other sets, they are chosen in such a way that we can show they hold with probability converging to 1 under our assumptions. For example, relevant to this section are the sets

$$\mathcal{R}_1 := \left\{ \max_{1 \leq j \leq N} \left| \frac{1}{T} \sum_{t=1}^T \epsilon_{j,t}^2 \right| \leq C \log(N) \right\}, \quad \mathcal{R}_2 := \left\{ \max_{1 \leq j \leq N} \left| \frac{1}{T} \sum_{t=1}^T \epsilon_{j,t}^2 \right| \leq CN^{2/m} \right\},$$

and

$$\mathcal{S}_1 := \left\{ \left\| \frac{1}{T} \sum_{t=1}^T \boldsymbol{\epsilon}_t \boldsymbol{\epsilon}_t' - \boldsymbol{\Sigma}_\epsilon \right\|_{\max} \leq C \frac{\sqrt{\log(N)}}{\sqrt{T}} \right\}, \quad \mathcal{S}_2 := \left\{ \left\| \frac{1}{T} \sum_{t=1}^T \boldsymbol{\epsilon}_t \boldsymbol{\epsilon}_t' - \boldsymbol{\Sigma}_\epsilon \right\|_{\max} \leq \frac{N^{4/m}}{T^{(m-2)/m}} \eta_T^{-1} \right\}.$$

The different subscripts of these sets indicate for which version of Assumption 2 they are intended. We show they hold with high probability in Lemmas A.9 and A.11. Note that many of our intermediate results are phrased as non-random bounds on random quantities, which hold on these sets, i.e., these bounds hold with probability 1 conditionally on these random events occurring. For the main result in Theorem 4, we then show that the probability of all these random events occurring jointly converges to 1, such that these non-random bounds hold asymptotically.

The main result of this section concerns the consistency of our estimate of $\boldsymbol{\Sigma}$, namely $\hat{\boldsymbol{\Sigma}} := \hat{\mathcal{B}}(1) \hat{\boldsymbol{\Sigma}}_\epsilon \hat{\mathcal{B}}(1)'$, with $\hat{\boldsymbol{\Sigma}}_\epsilon := \frac{1}{T} \sum_{t=1}^T \hat{\boldsymbol{\epsilon}}_t \hat{\boldsymbol{\epsilon}}_t'$, $\hat{\mathcal{B}}(z) = \mathbf{I} + \sum_{k=1}^{\infty} \hat{\mathbf{B}}_k z^k$, $\hat{\mathcal{B}}(z) = \mathbf{I} + \sum_{k=1}^{\infty} \hat{\mathbf{B}}_k z^k$. Unsurprisingly, the form of $\hat{\boldsymbol{\Sigma}}$ mirrors that of $\boldsymbol{\Sigma}$, since we apply the same Beveridge-Nelson decomposition in eq. (4) to the bootstrap process. To do so, the estimated VAR is required to be invertible, i.e. $\rho(\hat{\mathbf{A}}) < 1$, at least with probability converging to 1. We show that this is the case when $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$ in Lemma A.7. This justifies our suggested invertibility correction in Section 2, since it is asymptotically negligible. In finite samples, one can perform this correction by dividing each element of $\hat{\mathbf{A}}$ by $\rho(\hat{\mathbf{A}}) + \epsilon$ for some small $\epsilon > 0$. In Theorem 2 we establish a covariance closeness result which plays a crucial role in showing consistency of our proposed bootstrap method in the next section.

Theorem 2. *Assume that $\xi_{N,T} \psi_N^2 \rightarrow 0$, $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$, and let Assumption 3 hold. Define the set*

$$\mathcal{T}_1 := \left\{ \left\| \hat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma} \right\|_{\max} \leq C \psi_N^2 \left[\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{\sqrt{\log(N)}}{\sqrt{T}} + \xi_{N,T} \psi_N^2 \right] \right\}.$$

Under Assumption 2.1, on $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_1 \cap \mathcal{S}_1$, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{T}_1) = 1$.

Furthermore, define the set

$$\mathcal{T}_2 := \left\{ \left\| \hat{\Sigma} - \Sigma \right\|_{\max} \leq C\psi_N^2 \left[\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{(m-2)/m}} \eta_T^{-1} + \xi_{N,T} \psi_N^2 \right] \right\}.$$

Under Assumption 2.2, on $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_2 \cap \mathcal{S}_2$, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{T}_2) = 1$.

5 Bootstrap Consistency

In this section, we introduce some of the bootstrap-related notation, and flesh out the exact properties of the processes \mathbf{x}_t^* and $\boldsymbol{\epsilon}_t^*$. In Theorem 3, we then give a Gaussian approximation for the bootstrap process, mirroring Theorem 1. Finally, Theorem 4 provides the main result of bootstrap consistency.

As is customary in the bootstrap literature, we define the following bootstrap conditional notation: Let $\mathbb{P}^*(\cdot)$ denote the bootstrap probability conditional on the sample \mathbf{X} , and $\mathbb{E}^*(\cdot)$ the expectation with respect to \mathbb{P}^* , and similarly the conditional norms $\|x\|_{\psi_2}^* := \inf \left\{ c > 0 : \mathbb{E}^* \exp(|x|^2/c^2) \leq 2 \right\}$ and $\|x\|_{L_p}^* := (\mathbb{E}^* |x|^p)^{1/p}$. To apply the Beveridge-Nelson decomposition to the bootstrap series constructed as in Algorithm 1, we need to verify that \mathbf{x}_t^* and $\boldsymbol{\epsilon}_t^*$ follow a VAR process; in particular, we need to consider what the choice of initial values for $\mathbf{x}_{-K+1}^*, \dots, \mathbf{x}_0^*$ implies about the initial errors $\boldsymbol{\epsilon}_{-K+1}^*, \dots, \boldsymbol{\epsilon}_0^*$. While these errors are generally not important for the bootstrap statistics themselves, the error $\boldsymbol{\epsilon}_0^*$ does appear in one of the leftover terms of the decomposition. We let

$$\boldsymbol{\epsilon}_t^* := \begin{cases} \hat{\boldsymbol{\epsilon}}_t \gamma_t & t = 1, \dots, T \\ \mathbf{0} & t \leq -K \end{cases}, \quad \gamma_t \stackrel{iid}{\sim} N(0, 1),$$

and \mathbf{x}_t^* built from $\boldsymbol{\epsilon}_t^*$

$$\mathbf{x}_t^* := \begin{cases} \sum_{k=1}^K \mathbf{A}_k^* \mathbf{x}_{t-k}^* + \boldsymbol{\epsilon}_t^* & t = 1, \dots, T \\ \mathbf{x}_t & t = -K + 1, \dots, 0, \\ \mathbf{0} & t \leq -K \end{cases} \quad (5)$$

where $\mathbf{A}_k^* := \hat{\mathbf{A}}_k$. The bootstrap errors for $t = -k + 1, \dots, 0$ are then chosen such that

$$\boldsymbol{\epsilon}_t^* := \mathbf{x}_t^* - \sum_{k=1}^K \mathbf{A}_k^* \mathbf{x}_{t-k}^*,$$

which implies they are functions of $\hat{\mathbf{A}}$ and the original sample \mathbf{x}_t , $t = -K + 1, \dots, 0$.² We therefore

²Note that this implies that we observe our original process from time $t = -K + 1$ on. Alternatively, we can define $t = 1$ as the first observation, in which case we move all indices K periods forward. We do not consider this more conventional notation here as starting at $t = -K + 1$ lightens the notation in the proofs.

have $\boldsymbol{\epsilon}_0^* = \boldsymbol{x}_0 + \sum_{k=1}^{K-1} \mathbf{A}_k^* \boldsymbol{x}_t$, where the sum runs over one fewer elements since $\boldsymbol{x}_{-K} = \mathbf{0}$. The remaining bootstrap errors can be constructed similarly. By construction, the bootstrap processes \boldsymbol{x}_t^* and $\boldsymbol{\epsilon}_t^*$ then follow a VAR process mirroring eq. (1), and can be inverted under appropriate conditions to a VMA process mirroring eq. (3). This then also leads to the bootstrap versions of \tilde{S} and S_m , and the following bootstrap equivalent of Theorem 1.

Theorem 3 (Gaussian approximation for the bootstrap process). *Let \boldsymbol{x}_t^* be a linear process as in eq. (5), let Assumptions 1 and 3 hold, let $\xi_{N,T} \psi_N^2 \rightarrow 0$ and $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$. Define the sets*

$$\mathcal{U}_1 := \left\{ \max_{j,t} |\epsilon_{j,t}| \leq \sqrt{\log(N)} \log(T) \right\}, \quad \mathcal{U}_2 := \left\{ \max_{j,t} |\epsilon_{j,t}| \leq (NT)^{1/m} \eta_T^{-1} \right\},$$

the bootstrap VMA coefficient sums $\tilde{S}^* := \sum_{j=0}^{\infty} \|\hat{\mathbf{B}}_j\|_{\infty}$, $S_m^* := \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \|\hat{\mathbf{B}}_k\|_{\infty} \right)^m$, and

$$J_{N,T}^* := \sup_{y \in \mathbb{R}} \left| \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{x}_t^* \right\|_{\infty} \leq y \right) - \mathbb{P}^* (\|\boldsymbol{z}\|_{\infty} \leq y) \right|,$$

where $\boldsymbol{z} \sim N(\mathbf{0}, \boldsymbol{\Sigma})$.

1. Under Assumption 2.1, on $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{T}_1 \cap \mathcal{U}_1$, for sufficiently large N, T ,

$$J_{N,T}^* \leq C \left\{ \log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \sqrt{\frac{\log(N)}{T}} + \xi_{N,T} \psi_N^2 \right] + \frac{\sqrt{K} \log(N) d_N^* \psi_N^2}{\sqrt{T}} + \frac{1}{\log(N)} \right. \\ \left. + (\tilde{S}^* d_N^*)^2 \left[\frac{\log(N)^{3/2} \log(T)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)^2}{T} \right] + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \right\},$$

$$\text{where } d_N^* = C \left(\sqrt{T \phi_{N,T}} + \sqrt{\log(N)} \log(T) \right).$$

2. Under Assumption 2.2, on $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{T}_1 \cap \mathcal{U}_1$, for sufficiently large N, T ,

$$J_{N,T}^* \leq C \left\{ \log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{\frac{m-2}{m}}} + \xi_{N,T} \psi_N^2 \right] + (NK d_N^{*m} \psi_N^m)^{\frac{1}{m+1}} \left(\frac{\sqrt{\log(N)}}{\sqrt{T}} \right)^{\frac{m}{m+1}} \right. \\ \left. + (\tilde{S}^* d_N^*)^2 \left[\frac{\log(N)^{3/2} \left(\log(T) + (\tilde{S}^* d_N^*)^{\frac{1}{m-1}} \right)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)}{T^{\frac{m-2}{m}}} \right] + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \right\},$$

$$\text{where } d_N^* = C \left(\sqrt{T \phi_{N,T}} + (NT)^{1/m} \eta_T^{-1} \right).$$

Since \boldsymbol{z} in Theorem 3 is the same as in Theorem 1, we can combine both theorems and a telescopic sum argument to bound the Kolmogorov distance between $\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{x}_t \right\|_{\infty}$ and $\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{x}_t^* \right\|_{\infty}$, giving us bootstrap consistency in the following theorem.

Theorem 4. *Let Assumptions 1 to 5 hold, and define*

$$D_{N,T} = \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} \leq y \right) - \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* \right\|_{\infty} \leq y \right) \right|$$

When $\xi_{N,T} \psi_N^2 \rightarrow 0$ and $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$, the following hold with probability converging to 1 as $N, T \rightarrow \infty$.

Under Assumption 2.1,

$$D_{N,T} \leq C \left\{ \psi_N^2 \left[\frac{\ell_N^3}{\sqrt{T}} + \frac{\ell_N^{5/2} \ell_T^3}{\sqrt{T}} + \sqrt{\phi_{N,T}} \ell_N^{3/2} \ell_T + \phi_{N,T} \sqrt{T} \ell_N^{3/2} \ell_T + \psi_N^2 \xi_{N,T} \ell_N \ell_T \right. \right. \\ \left. \left. + \sqrt{K} \left(\sqrt{\phi_{N,T}} + \frac{\ell_N^{3/2} \ell_T}{\sqrt{T}} \right) \right] + \frac{1}{\ell_N} \right\},$$

where $\ell_T = \log(T)$, $\ell_N = \log(N)$.

Under Assumption 2.2,

$$D_{N,T} \leq C \eta_T^{-1} \left\{ \psi_N^2 \left[\ell_N \ell_T \sqrt{\phi_{N,T}} + \ell_N \ell_T \xi_{N,T} \psi_N^2 + \left(T \phi_{N,T} + (NT)^{2/m} \right) \left(\frac{\ell_N^{3/2} \ell_T}{\sqrt{T}} + \frac{\ell_N^2 \ell_T^2}{T} \right) \right] \right. \\ \left. + \frac{\psi_N N^{1/m} \ell_N^{3/2} \ell_T}{\sqrt{T}} + \frac{\psi_N^m N \ell_N^{3m/2-4} \ell_T \ell_{NT}}{T^{m/2-1}} + \left[\left(\frac{\sqrt{\ell_N}}{\sqrt{T}} \psi_N \right)^m NK \left(\sqrt{T \phi_{N,T}}^m + NT \right) \right]^{\frac{1}{m+1}} \right\}.$$

where $\ell_{NT} = \log(NT)$.

6 Bootstrap Consistency for VAR Estimation by the Lasso

The application of our proposed bootstrap method requires that the lasso satisfies Assumptions 4 and 5 with sequences $\psi_N, \xi_{N,T}$, and $\phi_{N,T}$ such that the bound in Theorem 4 converges to 0. In this section, we show that this is the under both options of Assumption 2, and under both weak and exact row-wise sparsity of the underlying VAR.

As described in Section 2 we propose to estimate the VAR equation-by-equation, using the lasso estimators in eq. (2). Our goal is therefore to find bounds on $\max_j \left\| \hat{\beta}_j - \beta_j \right\|_1$ and $\max_j \frac{1}{T} \|\hat{\epsilon}_j - \epsilon_j\|_2^2 = \max_j \frac{1}{T} \sum_{t=1}^T \left[(\hat{\beta}_j - \beta_j)' \mathcal{X}_t \right]^2$. For this purpose, we will be using error bounds in Corollary 1 of our previous work in Adamek et al. (2022), though similar error bounds have been derived in different contexts by other authors; see e.g. Bickel et al. (2009), Kock and Callot (2015), Medeiros and Mendes (2016), and Masini et al. (2021). Next, we will elaborate on the assumptions under which these error bounds hold.

For Assumption 1 of Adamek et al. (2022), we have $\mathbb{E} \mathbf{x}_t = 0 \implies \mathbb{E} \mathcal{X}_t = 0$ by the structure of eq. (1), and $\mathbb{E} \mathbf{x}_t \epsilon_{j,t} = 0$, $\forall j$, by independence of the errors. We then need to assume that

$\max_{j,t} \mathbb{E} |x_{j,t}|^m \leq C$ in addition to Assumption 2.2 in this paper to ensure the first part of the assumption is satisfied. This high-level assumption on moments of $x_{j,t}$ can also be shown to hold under more primitive conditions, such as moment condition on linear combinations of the errors,

$\max_{\|\mathbf{u}\|_2 \leq 1, t} \mathbb{E} |\mathbf{u}' \boldsymbol{\epsilon}_t|^m \leq C$, and a new summability condition on the rows of \mathbf{B}_k , $\max_j \sum_{k=0}^{\infty} \|\mathbf{b}_{j,k}\|_2^m \leq C$:

$$\max_{j,t} \|x_{j,t}\|_{L_m} \leq \sum_{k=0}^{\infty} \max_{j,t} \|\mathbf{b}_{j,k} \boldsymbol{\epsilon}_{t-k}\|_{L_m} = \sum_{k=0}^{\infty} \|\mathbf{b}_{j,k}\|_2 \left\| \frac{\mathbf{b}_{j,k}}{\|\mathbf{b}_{j,k}\|_2} \boldsymbol{\epsilon}_{t-k} \right\|_{L_m} = \|\mathbf{u}' \boldsymbol{\epsilon}_{t-k}\|_{L_m} \sum_{k=0}^{\infty} \|\mathbf{b}_{j,k}\|_2.$$

Note that m in this paper corresponds to $2\bar{m}$ in Adamek et al. (2022). The NED assumption is satisfied trivially, since Assumption 3 ensures that the VMA coefficients decay at an exponential rate. It therefore satisfies any polynomial decay rate on the NED sequence, and the assumption is satisfied for any arbitrarily large d . Assumption 2 of Adamek et al. (2022) requires that the rows of \mathbf{A} are weakly sparse, in the sense that $\|\boldsymbol{\beta}_j\|_r^r = \|[\mathbf{A}]_{j,\cdot}\|_r^r \leq s_{r,j}$ for some $0 \leq r < 1$. Assumption 3 of Adamek et al. (2022) requires that the covariance matrix of the regressors satisfies a form of compatibility condition; for simplicity, we can assume that $\Lambda_{\min} \left(\frac{1}{T} \sum_{t=1}^T \mathbb{E} \mathcal{X}_t \mathcal{X}_t' \right)$ is bounded away from zero, which is sufficient to satisfy the condition simultaneously for all equations. For an example of conditions when this is satisfied, see Equation 6 of Masini et al. (2021). Under these conditions, we have by Corollary 1 of Adamek et al. (2022) that

$$\frac{1}{T} \|\hat{\boldsymbol{\epsilon}}_j - \boldsymbol{\epsilon}_j\|_2^2 \leq C \lambda_j^{2-r} s_{r,j}, \quad \|\hat{\boldsymbol{\beta}}_j - \boldsymbol{\beta}_j\|_1 \leq C \lambda_j^{1-r} s_{r,j},$$

with probability converging to 1 under appropriate restrictions on the λ_j , detailed in Theorem 1 of Adamek et al. (2022). To further simplify this result, we can use the asymptotic setup of Example C.1 of Adamek et al. (2022) where N , λ_j , and $s_{r,j}$ grow at a polynomial rate of T . While this example provides the full details on the tradeoff between r , the number of moments, and the growth rates of $s_{r,j}$ and N relative to T , we fix $r = 1/2$ and $s_{r,j} \sim T^{1/8}$, $\forall j$ for illustrative purposes.

Corollary 1. *Let Assumptions 1, 2.2, and 3-5 hold. Furthermore, assume $\max_{j,t} \mathbb{E} |x_{j,t}|^m \leq C$, $\sum_{k=1}^{KN} \left| [\mathbf{A}]_{j,k} \right|^{1/2} \leq CT^{1/8}$, and $\Lambda_{\min} \left(\frac{1}{T} \sum_{t=1}^T \mathbb{E} \mathcal{X}_t \mathcal{X}_t' \right) \geq 1/C$. Let $K \leq C$, $N \sim T^a$ for $a > 0$, $\psi_N = \log(N) = C \log(T)$, and $\lambda_j \sim T^{-\ell}$ for all j and $\ell < \frac{m-1}{m} - \frac{4a}{m} - \frac{1}{4}$. When $52a + 12 < m$, $D_{N,T} \rightarrow 0$ with probability converging to 1 as $N, T \rightarrow \infty$.*

While Corollary 1 shows an example of conditions for bootstrap consistency using the finite absolute moments in Assumption 2.2, the stronger assumption of subgaussian moments in Assumption 2.1 allows for faster growth of N relative to T . In this scenario, we can consider the error

bounds in Theorem 2 of [Kock and Callot \(2015\)](#),

$$\frac{1}{T} \|\hat{\epsilon}_j - \epsilon_j\|_2^2 \leq C \lambda_j^2 s_{0,j} / \kappa_j, \quad \|\hat{\beta}_j - \beta_j\|_1 \leq C \lambda_j s_{0,j} / \kappa_j,$$

with $\lambda_j = C \ell_T^{5/2} \ell_N^2 \ell_K \ell_{N^2K}^{1/2} \sigma_T^2 / \sqrt{T}$. Note that σ_T^2 denotes the largest variance among all $\epsilon_{j,t}$ and $x_{j,t}$, so we once again make the high level assumption that $\max_{j,t} \mathbb{E} x_{j,t}^2 \leq C$. To obtain these bounds, we need the additional assumption that the errors are gaussian, so $\epsilon_t \sim IIDN(\mathbf{0}, \Sigma_\epsilon)$, which implies Assumption 2.1. Additionally, they consider the case of exact sparsity, with $\sum_{k=1}^{KN} \mathbb{1} \left\{ |[A]_{j,k}| > 0 \right\} \leq s_{0,j}$. Finally, κ_j play a similar role to the compatibility constant in Assumption 2 of [Adamek et al. \(2022\)](#), and are bounded away from 0 when $\Lambda_{\min} \left(\frac{1}{T} \sum_{t=1}^T \mathbb{E} \mathcal{X}_t \mathcal{X}_t' \right) \geq 1/C$, see the discussion on page 7 of [Kock and Callot \(2015\)](#) for details. Regarding the growth rates of N and $s_{0,j}$, we take a similar example to Theorem 3 of [Kock and Callot \(2015\)](#), with $N \sim e^{(T^a)}$ and $s_{0,j} \leq CT^b$.

Corollary 2. *Let Assumptions 1, and 3-5 hold. Furthermore, assume $\max_{j,t} \mathbb{E} |x_{j,t}|^2 \leq C$, $\sum_{k=1}^{KN} \mathbb{1} \left\{ |[A]_{j,k}| > 0 \right\} \leq CT^b$ for some $b > 0$, and $\Lambda_{\min} \left(\frac{1}{T} \sum_{t=1}^T \mathbb{E} \mathcal{X}_t \mathcal{X}_t' \right) \geq 1/C$. Let $K \leq C$, $N \sim e^{(T^a)}$ for $a > 0$, $\psi_N = \log(\log(N)) = C \log(T)$, and $\lambda_j = C \frac{\ell_T^{5/2} T^{5a/2}}{\sqrt{T}}$. When $13a + 2b < 1$, $D_{N,T} \rightarrow 0$ with probability converging to 1 as $N, T \rightarrow \infty$.*

7 Conclusion

In this paper, we introduce a VAR multiplier bootstrap procedure which approximates the distribution of scaled high-dimensional means, using the lasso to estimate the VAR. We motivate the usefulness of this procedure as a tool for inference in high-dimensional time series, allowing for non-conservative simultaneous testing of a large set of hypotheses. We show that the bootstrap is consistent under two different moment assumptions on the errors: sub-gaussian moments, and a finite number of absolute moments. Under the former, N can grow at an exponential rate of T . Under the latter, N can only grow at a polynomial rate of T , with the growth rate of N limited by the number of absolute moments available.

We provide guidance for estimating the VAR bootstrap model by the lasso as a running example. We show that the lasso satisfies appropriate error bounds for consistency of the bootstrap distribution, under the assumption that the underlying VAR process is (row-wise) sparse. In our examples, we derive explicit limits on the growth rate of N relative to T thereby allowing for exact and weak sparsity of the VAR.

To establish the consistency of the VAR multiplier bootstrap, we derive a Gaussian approximation for the maximum mean of a linear process, which may be of independent interest. Our results

can be applied to more complex statistics than simple means, and we believe that extending this method to inference for linear model coefficients is an interesting avenue for future research.

Appendix A Preliminary Lemmas

Lemma A.1.

1. Under Assumption 2.1, $\max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{\psi_2} \leq d_N$ with $d_N = C\sqrt{\log(N)} \geq 1$.
2. Under Assumption 2.2, $\max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{L_m} \leq d_N$, with $d_N = CN^{1/m} \geq 1$.

Lemma A.2. Let Assumption 1 hold, and define

$$M_{N,T} := \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)\epsilon_t \right\|_{\infty} \leq y \right) - \mathbb{P}(\|z\|_{\infty} \leq y) \right|,$$

where $z \sim N(\mathbf{0}, \Sigma)$, $\Sigma = \mathcal{B}(1)\Sigma_{\epsilon}\mathcal{B}(1)'$.

1. Under Assumption 2.1

$$M_{N,T} \leq C \left(\frac{b_T \log(N)^{3/2} \log(T)}{\sqrt{T}} + \frac{b_T \log(N)^2}{\sqrt{T}} \right),$$

where $b_T = \tilde{S}^2 d_N^2$.

2. Under Assumption 2.2

$$M_{N,T} \leq C \left(\frac{b_T (\log N)^{3/2} \log(T)}{\sqrt{T}} + \frac{b_T^2 \log(N)^2 \log(T)}{T^{1-2/m}} + \left[\frac{b_T^m \log(N)^{3m/2-4} \log(T) \log(NT)}{T^{m/2-1}} \right]^{\frac{1}{m-2}} \right),$$

where $b_T = \tilde{S}^2 d_N^2$.

Lemma A.3. Define $\tilde{\mathcal{B}}(L) = \sum_{j=0}^{\infty} \sum_{k=j+1}^{\infty} \mathbf{B}_k$.

1. Under Assumption 2.1, for any $y > 0$

$$\mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L)\epsilon_T \right\|_{\infty} > y \right) = \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L)\epsilon_0 \right\|_{\infty} > y \right) \leq 2N \exp \left(-C \frac{y^2 T}{d_N^2 S_2} \right).$$

2. Under Assumption 2.2, for any $y > 0$

$$\mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L)\epsilon_T \right\|_{\infty} > y \right) = \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L)\epsilon_0 \right\|_{\infty} > y \right) \leq C \frac{N d_N^m S_m}{(y\sqrt{T})^m}.$$

Lemma A.4. Let a_j and b_j be non-negative sequences satisfying $\sum_{j=0}^{\infty} j^m a_j < \infty$ and $\sum_{j=0}^{\infty} j^m b_j < \infty$ for some $1 \leq m < \infty$,

$$\sum_{j=1}^{\infty} \sum_{s=0}^{j-1} a_s b_{j-1-s} = \left(\sum_{i=0}^{\infty} a_i \right) \left(\sum_{i=0}^{\infty} b_i \right),$$

and

$$\sum_{j=1}^{\infty} \sum_{s=0}^{j-1} j^m a_s b_{j-1-s} \leq 4^{m-1} \left(\sum_{i=0}^{\infty} (i^m + 1) a_i \right) \left(\sum_{i=0}^{\infty} (i^m + 1) b_i \right).$$

Lemma A.5. For $|\lambda| < 1$, and $1 \leq m < \infty$, $\sum_{k=1}^{\infty} k^m \lambda^{km} \leq C$.

Lemma A.6. Under Assumption 3, for any constant $1 \leq m < \infty$

$$\sum_{k=1}^{\infty} k^m \left\| \mathbb{A}^k \right\|_{\infty}^m \leq C \psi_N^m.$$

Additionally, on \mathcal{P} , when $\xi_{N,T} \psi_N^2 \rightarrow 0$, and N, T are sufficiently large

$$\sum_{k=1}^{\infty} k^m \left\| \hat{\mathbb{A}}^k \right\|_{\infty}^m \leq C \psi_N^m \text{ and } \sum_{k=1}^{\infty} k^m \left\| \hat{\mathbb{A}}^k - \mathbb{A}^k \right\|_{\infty}^m \leq C \xi_{N,T}^m \psi_N^{3m}.$$

Lemma A.7. Under Assumption 3, on \mathcal{P} , when $\xi_{N,T} \psi_N^2 \rightarrow 0$ and $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$, for sufficiently large N, T , $\rho(\hat{\mathbb{A}}) \leq \lambda'$, for some $\lambda < \lambda' < 1$.

Lemma A.8. Under Assumption 3, for any constant $1 \leq m < \infty$,

1. $\tilde{S} = \sum_{j=0}^{\infty} \left\| \mathbf{B}_j \right\|_{\infty} \leq C \psi_N$.
2. $\sum_{j=0}^{\infty} \left\| \mathbf{B}_j \right\|_{\infty}^m \leq C \psi_N^m$
3. $S_m = \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \left\| \mathbf{B}_k \right\|_{\infty} \right)^m \leq C \psi_N^m$,

Additionally, on \mathcal{P} , when $\xi_{N,T} \psi_N^2 \rightarrow 0$, $\frac{\log(\psi_N)}{\log(\xi_{N,T})} \rightarrow 0$, and N, T are sufficiently large

4. $\tilde{S}^* = \sum_{j=0}^{\infty} \left\| \hat{\mathbf{B}}_j \right\|_{\infty} \leq C \psi_N$.
5. $\sum_{j=0}^{\infty} \left\| \hat{\mathbf{B}}_j - \mathbf{B}_j \right\|_{\infty}^m \leq C \xi_{N,T}^m \psi_N^{3m}$,
6. $S_m^* = \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \left\| \hat{\mathbf{B}}_k \right\|_{\infty} \right)^m \leq C \psi_N^m$.

Lemma A.9. Define the set

$$\mathcal{R}_1 := \left\{ \max_{1 \leq j \leq N} \left| \frac{1}{T} \sum_{t=1}^T c_{j,t}^2 \right| \leq C d_N^2 \right\},$$

Under Assumption 2.1, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{R}_1) = 1$. Furthermore, define the set

$$\mathcal{R}_2 := \left\{ \max_{1 \leq j \leq N} \left| \frac{1}{T} \sum_{t=1}^T \epsilon_{j,t}^2 \right| \leq C d_N^2 \right\},$$

Under Assumption 2.2, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{R}_2) = 1$.

Lemma A.10. *On either $\mathcal{Q} \cap \mathcal{R}_1$ or $\mathcal{Q} \cap \mathcal{R}_2$,*

$$\left\| \frac{1}{T} \sum_{t=1}^T \hat{\epsilon}_t \epsilon'_t - \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t \right\|_{\max} \leq C \left(\phi_{N,T} + d_N^2 \sqrt{\phi_{N,T}} \right).$$

Lemma A.11. *Define the set*

$$\mathcal{S}_1 := \left\{ \left\| \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T} \sum_{t=1}^T \mathbb{E} \epsilon_t \epsilon'_t \right\|_{\max} \leq C \frac{\sqrt{\log(N)}}{\sqrt{T}} \right\},$$

Under Assumption 2.1, $\lim_{N,T} \mathbb{P}(\mathcal{S}_1) = 1$. Furthermore, define the set

$$\mathcal{S}_2 := \left\{ \left\| \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T} \sum_{t=1}^T \mathbb{E} \epsilon_t \epsilon'_t \right\|_{\max} \leq \frac{N^{4/m}}{T^{(m-2)/m} \eta_T^{-1}} \right\}$$

for some sequence $\eta_T \rightarrow 0$. Under Assumption 2.2, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{S}_2) = 1$.

Lemma A.12. *Define the set $\mathcal{U}_1 := \left\{ \max_{j,t} |\epsilon_{j,t}| \leq d_N \log(T) \right\}$. Under Assumption 2.1, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{U}_1) = 1$. Furthermore, define the set $\mathcal{U}_2 := \left\{ \max_{j,t} |\epsilon_{j,t}| \leq d_N T^{1/m} \eta_T^{-1} \right\}$, for some $\eta_T \rightarrow 0$. Under Assumption 2.2, $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{U}_2) = 1$.*

Lemma A.13.

1. On $\mathcal{U}_1 \cap \mathcal{Q}$, $\max_t \left\| \max_j \epsilon_{j,t}^* \right\|_{\psi_2}^* \leq d_N^*$, with $d_N^* = C \left(\sqrt{T \phi_{N,T}} + d_N \log(T) \right)$,
2. On $\mathcal{U}_2 \cap \mathcal{Q}$, $\max_t \left\| \max_j \epsilon_{j,t}^* \right\|_{L_m}^* \leq d_N^*$, with $d_N^* = C \left(\sqrt{T \phi_{N,T}} + d_N T^{1/m} \eta_T^{-1} \right)$.

Lemma A.14. *Let Assumption 1 hold, and define*

$$M_{N,T}^* := \sup_{y \in \mathbb{R}} \left| \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)^* \epsilon_t^* \right\|_{\infty} \leq y \right) - \mathbb{P}^*(\|z\|_{\infty} \leq y) \right|,$$

where $z \sim N(\mathbf{0}, \Sigma)$. On $\mathcal{T}_1 \cap \mathcal{U}_1 \cap \mathcal{Q}$

$$M_{N,T}^* \leq C \left\{ \log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \sqrt{\frac{\log(N)}{T}} + \xi_{N,T} \psi_N^2 \right] + (\tilde{S}^* d_N^*)^2 \left[\frac{\log(N)^{3/2} \log(T)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)^2}{T} \right] + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \right\}.$$

On $\mathcal{T}_2 \cap \mathcal{U}_2 \cap \mathcal{Q}$

$$M_{N,T}^* \leq C \left\{ \log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{\frac{m-2}{m}}} + \xi_{N,T} \psi_N^2 \right] \right. \\ \left. + (\tilde{S}^* d_N^*)^2 \left[\frac{\log(N)^{3/2} \left(\log(T) + (\tilde{S}^* d_N^*)^{\frac{1}{m-1}} \right)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)}{T^{\frac{m-2}{m}}} \right] + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \right\}.$$

Lemma A.15. Define $\tilde{\mathcal{B}}^*(L) = \sum_{j=0}^{\infty} \sum_{k=j+1}^{\infty} \mathbf{B}_k^*$. Under Assumption 3, on \mathcal{P} , the following holds

1. Under Assumption 2.1, for any $y > 0$

$$\mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_{\infty} > \eta_T \right) \leq 2N \exp \left(-C \frac{\eta_T^2 T}{K \psi_N^2 d_N^2 S_2} \right).$$

2. Under Assumption 2.2, for any $y > 0$

$$\mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_{\infty} > \eta_T \right) \leq C \frac{NK S_m \psi_N^m d_N^m}{(\eta_T \sqrt{T})^m}.$$

Appendix B Proofs

Proof of Lemma A.1. Following Lemma 2.2.2 of van der Vaart and Wellner (1996),³

$$\max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{\psi_2} \leq C \sqrt{\log(N)} \max_{j,t} \|\epsilon_{j,t}\|_{\psi_2},$$

and by the statement on page 96 of van der Vaart and Wellner (1996),

$$\max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{L_m} \leq N^{1/m} \max_{j,t} \|\epsilon_{j,t}\|_{L_m}. \quad \square$$

Proof of Lemma A.2. Note that $\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \epsilon_t$ is a scaled sum of iid random variables, and the proof will proceed by applying the Gaussian approximation in Corollary 2.1 of Chernozhukov et al. (2020). In particular, we will use either the second or third clause of this corollary, depending on whether we use Lemma A.1.1 or Lemma A.1.2.

First, using Lemma A.1.1 we use the second clause, which needs their conditions (E.2) and (M).

For (E.2), we have by Lemma A.1.1 that

$$\left\| \frac{x_{j,t}}{\sigma_j} \right\|_{\psi_2} = \left\| \frac{\mathcal{B}(1)_j \epsilon_t}{\sigma_j} \right\|_{\psi_2} \leq \left\| \frac{\|\mathcal{B}(1)_j\|_1 \max_j |\epsilon_{j,t}|}{\sigma_j} \right\|_{\psi_2} \leq \frac{\|\mathcal{B}(1)_j\|_1}{|\sigma_j|} \left\| \max_j |\epsilon_{j,t}| \right\|_{\psi_2} \leq C \tilde{S} d_N,$$

where $\mathcal{B}(1)_j$ denotes the j th row of $\mathcal{B}(1)$. The last inequality comes from bounding $\sigma_j^2 \geq \Lambda_{\min}(\Sigma) \geq 1/C$ by Assumption 1, and

³We take $\psi(x) = e^{x^2} - 1$ (see the explanation of their page 97), and note that $\sqrt{\log(1+N)} \leq C\sqrt{\log N}$ when $N > 1$.

$$\|\mathcal{B}(1)_j\|_1 = \left\| \sum_{j=0}^{\infty} \mathbf{b}_{j,k} \right\|_1 \leq \sum_{j=0}^{\infty} \|\mathbf{b}_{j,k}\|_1 \leq \sum_{j=0}^{\infty} \|\mathbf{B}_k\|_{\infty} = \tilde{S},$$

where $\mathbf{b}_{j,k}$ is the j th row of \mathbf{B}_k . For (M),

$$\mathbb{E} \left| \frac{x_{j,t}}{\sigma_j} \right|^4 = \left\| \frac{\mathcal{B}(1)_j \boldsymbol{\epsilon}_t}{\sigma_j} \right\|_{L_4}^4 \leq C \left\| \frac{\mathcal{B}(1)_j \boldsymbol{\epsilon}_t}{\sigma_j} \right\|_{\psi_2}^4 \leq C \tilde{S}^4 d_N^4,$$

by equation (2.15) in [Vershynin \(2019\)](#). To satisfy the second clause of Corollary 2.1 in [Chernozhukov et al. \(2020\)](#), we then need a sequence b_T such that $C\tilde{S}d_t \leq b_T$ and $C\tilde{S}^4 d_N^4 \leq b_T^2$. Note that $\tilde{S} \geq 1$ since $\mathbf{B}_0 = \mathbf{I}$, and $d_N \geq 1$ by assumption, so these inequalities are satisfied when $b_T \sim \tilde{S}^2 d_N^2$. It therefore follows that

$$M_{N,T} \leq C \left(\frac{b_T (\log N)^{3/2} \log T}{\sqrt{T} \Lambda_{\min}(\tilde{\boldsymbol{\Sigma}})} + \frac{b_T (\log N)^2}{\sqrt{T} \sqrt{\Lambda_{\min}(\tilde{\boldsymbol{\Sigma}})}} \right),$$

where $\tilde{\boldsymbol{\Sigma}}$ is the correlation matrix of \mathbf{x}_t . To show that $\Lambda_{\min}(\tilde{\boldsymbol{\Sigma}})$ is bounded away from 0, write $\tilde{\boldsymbol{\Sigma}} = \mathbf{D}\boldsymbol{\Sigma}\mathbf{D}$, where $\mathbf{D} = \text{diag}(1/\sigma_1, \dots, 1/\sigma_N)$. Since \mathbf{D} and $\boldsymbol{\Sigma}$ are symmetric and positive definite by Assumption 1, we have $\Lambda_{\min}(\tilde{\boldsymbol{\Sigma}}) \geq \Lambda_{\min}(\mathbf{D})^2 \Lambda_{\min}(\boldsymbol{\Sigma})$. The eigenvalues of a diagonal matrix are just its diagonal entries, which are bounded away from 0 since the variances σ_j are bounded, and $\Lambda_{\min}(\boldsymbol{\Sigma})$ is bounded away from 0; both by Assumption 1. The result of the first statement then follows.

Second, using Lemma A.1.2, we use the third clause of Corollary 2.1 in [Chernozhukov et al. \(2020\)](#), which needs their conditions (E.3) and (M). For (E.3),

$$\begin{aligned} \left\| \max_{1 \leq j \leq N} \left| \frac{x_{j,t}}{\sigma_j} \right| \right\|_{L_m} &\leq \max_j |1/\sigma_j| \left\| \max_j x_{j,t} \right\|_{L_m} \\ &\leq C \left\| \max_j \mathcal{B}(1)_j \boldsymbol{\epsilon}_t \right\|_{L_m} \leq C \|\mathcal{B}(1)\|_{\infty} \left\| \max_j |\epsilon_{j,t}| \right\|_{L_m} \leq C \tilde{S} d_N. \end{aligned}$$

For (M),

$$\mathbb{E} \left| \frac{x_{j,t}}{\sigma_j} \right|^4 = \left\| \frac{\mathcal{B}(1)_j \boldsymbol{\epsilon}_t}{\sigma_j} \right\|_{L_4}^4 \leq C \tilde{S}^4 d_N^4.$$

Similarly to before, we need the sequence b_T to satisfy $\tilde{S}d_N \leq b_T$, and $\tilde{S}^4 d_N^4 \leq b_T^2$, which is satisfied

when taking $b_T \sim \tilde{S}^2 d_N^2$. Therefore

$$M_{N,T} \leq C \left(\frac{b_T (\log N)^{3/2} \log T}{\sqrt{T} \Lambda_{\min}(\tilde{\Sigma})} + \frac{b_T^2 (\log N)^2 \log T}{T^{1-2/m} \Lambda_{\min}(\tilde{\Sigma})} + \left[\frac{b_T^m (\log N)^{3m/2-4} (\log T) \log(NT)}{T^{m/2-1} \Lambda_{\min}(\tilde{\Sigma})^{m/2}} \right]^{\frac{1}{m-2}} \right),$$

and the result of the second statement follows. \square

Proof of Lemma A.3.

$$\begin{aligned} \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathbf{B}}(L) \boldsymbol{\epsilon}_0 \right\|_{\infty} > \eta_T \right) &= \mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \left[\tilde{\mathbf{B}}(L) \right]_{p,\cdot} \boldsymbol{\epsilon}_0 \right| > \eta_T \right) \\ &= \mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \left[\sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) L^j \right]_{p,\cdot} \boldsymbol{\epsilon}_0 \right| > \eta_T \right) \\ &= \mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right| > \eta_T \right), \end{aligned} \quad (\text{B.1})$$

where $\mathbf{b}_{p,k}$ is the p th row of \mathbf{B}_k .

By Lemma A.1.1, we proceed from eq. (B.1) with the union bound and Hoeffding's inequality (see Theorem 2.6.2 in Vershynin (2019))

$$\begin{aligned} \mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \sum_{j=1}^{\infty} \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right| > y \right) &\leq \sum_{p=1}^N \mathbb{P} \left(\left| \sum_{j=1}^{\infty} \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right| > y \sqrt{T} \right) \\ &\leq \sum_{p=1}^N 2 \exp \left(-C \frac{[y \sqrt{T}]^2}{\sum_{j=1}^{\infty} \left\| \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right\|_{\psi_2}^2} \right). \end{aligned}$$

Using Lemma A.1.1 and arguments similar to those in the proof of Lemma A.2, we can bound

$$\left\| \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right\|_{\psi_2} \leq d_N \sum_{k=j}^{\infty} \|\mathbf{B}_k\|_{\infty},$$

and therefore

$$\mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \sum_{j=1}^{\infty} \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right| > y \right) \leq 2N \exp \left(-C \frac{y^2 T}{d_N^2 S_2} \right),$$

so the first statement follows. For the second statement, by Lemma A.1.2, we proceed from eq. (B.1)

with the union bound and Markov's inequality

$$\begin{aligned} & \mathbb{P} \left(\max_{1 \leq p \leq N} \frac{1}{\sqrt{T}} \left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right| > y \right) \\ & \leq \sum_{p=1}^N \mathbb{P} \left(\left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right| > y\sqrt{T} \right) \leq \sum_{p=1}^N \frac{\mathbb{E} \left[\left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right|^m \right]}{(y\sqrt{T})^m}. \end{aligned} \quad (\text{B.2})$$

We continue with the Marcinkiewicz–Zygmund inequality, which gives for independent, mean zero random variables X_j that

$$\mathbb{E} \left(\left| \sum_{j=1}^{\infty} X_j \right|^m \right) \leq C \mathbb{E} \left[\left(\sum_{j=1}^{\infty} |X_j|^2 \right)^{m/2} \right].$$

Let $X_j = \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j}$. Using the M-Z inequality and C_r inequality

$$\begin{aligned} & \mathbb{E} \left[\left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right|^m \right] = \mathbb{E} \left[\left| \sum_{j=1}^{\infty} \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right|^m \right] = \mathbb{E} \left[\left| \sum_{j=1}^{\infty} X_j \right|^m \right] \\ & \stackrel{M-Z}{\leq} C \mathbb{E} \left[\left(\sum_{j=1}^{\infty} |X_j|^2 \right)^{m/2} \right] \stackrel{C_r}{\leq} C \times 2^{m/2-1} \sum_{j=1}^{\infty} \mathbb{E} (|X_j|^m). \end{aligned}$$

Using Lemma A.1.2 and arguments similar to those in Lemma A.2, we can bound

$$\|X_j\|_{L_m} = \left\| \left(\sum_{k=j}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{1-j} \right\|_{L_m} \leq d_N \sum_{k=j}^{\infty} \|\mathbf{B}_k\|_{\infty},$$

and

$$\mathbb{E} \left[\left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right|^m \right] \leq d_N^m \sum_{j=1}^{\infty} \left(\sum_{k=j}^{\infty} \|\mathbf{B}_k\|_{\infty} \right)^m = d_N^m S_m.$$

Continuing from eq. (B.2), we therefore obtain

$$\sum_{p=1}^N \frac{\mathbb{E} \left[\left| \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \mathbf{b}_{p,k} \right) \boldsymbol{\epsilon}_{-j} \right|^m \right]}{(y\sqrt{T})^m} \leq \sum_{p=1}^N C \frac{d_N^m S_m}{(y\sqrt{T})^m} = C \frac{N d_N^m S_m}{(y\sqrt{T})^m} \square$$

Proof of Theorem 1. We first write the Beveridge-Nelson decomposition of the process

$$\mathbf{x}_t = \mathcal{B}(L)\boldsymbol{\epsilon}_t = \mathcal{B}(1)\boldsymbol{\epsilon}_t - (1-L)\tilde{\mathcal{B}}(L)\boldsymbol{\epsilon}_t,$$

where $\tilde{\mathcal{B}}(L) = \sum_{j=0}^{\infty} \tilde{\mathbf{B}}_j L^j$, $\tilde{\mathbf{B}}_j = \sum_{k=j+1}^{\infty} \mathbf{B}_k$, such that

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t = \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \boldsymbol{\epsilon}_t - \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_T + \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_0.$$

Define

$$x_T^{(\max)} = \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty}, \quad \epsilon_T^{(\max)} = \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \boldsymbol{\epsilon}_t \right\|_{\infty}, \quad z_T^{(\max)} = \|\mathbf{z}\|_{\infty},$$

$$F_{1,T}(y) := \mathbb{P}\left(x_T^{(\max)} \leq y\right) \quad F_{2,T}(y) := \mathbb{P}\left(\epsilon_T^{(\max)} \leq y\right)$$

$$G_T(y) := \mathbb{P}\left(z_T^{(\max)} \leq y\right) \quad r_T := x_T^{(\max)} - \epsilon_T^{(\max)}$$

Then

$$\begin{aligned} |r_T| &= \left| \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} - \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \boldsymbol{\epsilon}_t \right\|_{\infty} \right| \\ &\leq \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t - \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1) \boldsymbol{\epsilon}_t \right\|_{\infty} = \left\| -\frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_T + \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_0 \right\|_{\infty} \\ &\leq \left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_T \right\|_{\infty} + \left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L) \boldsymbol{\epsilon}_0 \right\|_{\infty} = R_T + R_0. \end{aligned}$$

By Lemma A.3 we have $\mathbb{P}(R_0 > \eta_{T,1}) = \mathbb{P}(R_T > \eta_{T,1}) \leq 2N \exp\left(-C \frac{\eta_{T,1}^2 T}{d_N^2 S_2}\right)$, and this allows us to bound

$$\begin{aligned} \mathbb{P}(|r_T| > 2\eta_{T,1}) &\leq \mathbb{P}(R_T + R_0 > 2\eta_{T,1}) = \mathbb{P}(R_T + R_0 > 2\eta_{T,1} | R_T \leq \eta_{T,1}) \mathbb{P}(R_T \leq \eta_{T,1}) \\ &\quad + \mathbb{P}(R_T + R_0 > 2\eta_{T,1} | R_T > \eta_{T,1}) \mathbb{P}(R_T > \eta_{T,1}) \\ &\leq \mathbb{P}(\eta_{T,1} + R_0 > 2\eta_{T,1}) \mathbb{P}(R_T > \eta_{T,1}) \leq 4N \exp\left(-C \frac{\eta_{T,1}^2 T}{d_N^2 S_2}\right) =: \eta_{T,2}. \end{aligned}$$

Continue with

$$\begin{aligned} &|F_{1,T}(y) - G_T(y)| \\ &\leq \left| \mathbb{P}\left(\epsilon_T^{(\max)} + r_T \leq y \mid |r_T| \leq 2\eta_{T,1}\right) \mathbb{P}(|r_T| \leq 2\eta_{T,1}) - \mathbb{P}\left(z_T^{(\max)} \leq y\right) \right| \\ &\quad + \mathbb{P}\left(x_T^{(\max)} \leq y \mid |r_T| > 2\eta_{T,1}\right) \mathbb{P}(|r_T| > 2\eta_{T,1}) \\ &\leq \left| \mathbb{P}\left(\epsilon_T^{(\max)} \leq y + 2\eta_{T,1}\right) - \mathbb{P}\left(y_T^{(\max)} \leq y\right) \right| + 4\eta_{T,2} \\ &\leq \underbrace{\left| \mathbb{P}\left(\epsilon_T^{(\max)} \leq y + 2\eta_{T,1}\right) - \mathbb{P}\left(z_T^{(\max)} \leq y + 2\eta_{T,1}\right) \right|}_{A_{T,1}(y+2\eta_{T,1})} \\ &\quad + \underbrace{\left| \mathbb{P}\left(z_T^{(\max)} \leq y + 2\eta_{T,1}\right) - \mathbb{P}\left(z_T^{(\max)} \leq y\right) \right|}_{A_{T,2}(y)} + 4\eta_{T,2}. \end{aligned}$$

Note that $\sup_{y \in \mathbb{R}} A_{T,1}(y + 2\eta_{T,1}) = M_{N,T}$ which can be bounded by Lemma A.2, and $\sup_{y \in \mathbb{R}} A_{T,2}(y)$

can be bounded by Lemma A.1 in Chernozhukov et al. (2017), which states that for centered gaussian vectors $\mathbf{z} \in \mathbb{R}^N$ with variances uniformly bounded away from 0 (as is the case here by Assumption 1), for all $\mathbf{y} \in \mathbb{R}^N$ and $a > 0$

$$\mathbb{P}(\mathbf{z} \leq \mathbf{y} + a) - \mathbb{P}(\mathbf{z} \leq \mathbf{y}) \leq Ca\sqrt{\log(N)}.$$

Note that this applies to $\|\mathbf{z}\|_\infty$ as well, since

$$\mathbb{P}(\|\mathbf{z}\|_\infty \leq y + a) - \mathbb{P}(\|\mathbf{z}\|_\infty \leq y) = 2[\mathbb{P}(\mathbf{z} \leq \mathbf{y} + a) - \mathbb{P}(\mathbf{z} \leq \mathbf{y})],$$

when \mathbf{y} has each element equal to y , and if the bound holds for all $\mathbf{y} \in \mathbb{R}^N$, it also holds for the supremum over $y \in \mathbb{R}$. We therefore have the bound

$$\sup_{y \in \mathbb{R}} |F_{1,T}(y) - G_T(y)| \leq M_{N,T} + C_1 \left[\eta_{T,1} \sqrt{\log N} + N \exp \left(-C_2 \frac{\eta_{T,1}^2 T}{d_N^2 S_2} \right) \right].$$

In order for this expression to converge, we need to choose $\eta_{T,1}$ converging to 0 fast enough such that $\eta_{T,1} \sqrt{\log(N)} \rightarrow 0$, but slow enough such that $N \exp \left(-C_2 \frac{\eta_{T,1}^2 T}{d_N^2 S_2} \right) \rightarrow 0$. One such choice is $\eta_{T,1} = \sqrt{\log(N \log(N))} \frac{d_N^2 S_2}{C_2 T}$ (assuming $N > 1$), which lets us bound

$$\begin{aligned} C_1 \left[\eta_{T,1} \sqrt{\log N} + N \exp \left(-C_2 \frac{\eta_{T,1}^2 T}{d_N^2 S_2} \right) \right] &\leq C \left[\frac{d_N \sqrt{S_2}}{\sqrt{T}} \sqrt{\log(N) \log(N \log(N))} + \frac{1}{\log(N)} \right] \\ &\leq C \left[\frac{\log(N) d_N \sqrt{S_2}}{\sqrt{T}} + \frac{1}{\log(N)} \right], \end{aligned}$$

and the result of the first statement follows.

For the second statement, by Lemma A.1.2, we may follow the same steps as above, taking $\eta_{T,2} := 2C \frac{Nd_N^m S_m}{(\eta_{T,1} \sqrt{T})^m}$ by the second clause of Lemma A.3. We then have the bound

$$\sup_{y \in \mathbb{R}} |F_{1,T}(y) - G_T(y)| \leq M_{N,T} + C_1 \left[\eta_{T,1} \sqrt{\log N} + \frac{Nd_N^m S_m}{(\eta_{T,1} \sqrt{T})^m} \right].$$

In this case, we can easily solve for the optimal rate of convergence for $\eta_{T,1}$, which has both terms converging at the same rate, $\eta_{T,1} = \left(\frac{Nd_N^m S_m}{\sqrt{T}^m \sqrt{\log(N)}} \right)^{\frac{1}{m+1}}$. We then have

$$\eta_{T,1} \sqrt{\log N} = \frac{Nd_N^m S_m}{(\eta_{T,1} \sqrt{T})^m} = (Nd_N^m S_m)^{\frac{1}{m+1}} \left(\frac{\sqrt{\log(N)}}{\sqrt{T}} \right)^{\frac{m}{m+1}},$$

and the result of the second statement follows. \square

Proof of Lemma A.4. First, rewrite the sums using indicator functions

$$\begin{aligned}
& \sum_{j=1}^{\infty} \sum_{s=0}^{j-1} j^m a_s b_{j-1-s} = \sum_{j=1}^{\infty} \sum_{s=0}^{\infty} j^m a_s b_{j-1-s} \mathbf{1}_{(s \leq j-1)} = \sum_{j=1}^{\infty} \sum_{s=0}^{j-1} \sum_{k=0}^{\infty} j^m a_s b_k \mathbf{1}_{(s \leq j-1)} \mathbf{1}_{(k=j-1-s)} \\
& = \sum_{j=1}^{\infty} \sum_{s=0}^{\infty} \sum_{k=0}^{\infty} j^m a_s b_k \mathbf{1}_{(s \leq k+s)} \mathbf{1}_{(j=k+s+1)} = \sum_{j=1}^{\infty} \sum_{s=0}^{\infty} \sum_{k=0}^{\infty} (k+s+1)^m a_s b_k \mathbf{1}_{(0 \leq k)} \mathbf{1}_{(j=k+s+1)} \\
& = \sum_{s=0}^{\infty} \sum_{k=0}^{\infty} (k+s+1)^m a_s b_k \sum_{j=1}^{\infty} \mathbf{1}_{(j=k+s+1)} = \sum_{s=0}^{\infty} \sum_{k=0}^{\infty} (k+s+1)^m a_s b_k.
\end{aligned}$$

Since $(k+l+1) \leq (k+1)(l+1) \implies (k+l+1)^m \leq (k+1)^m(l+1)^m$, we can bound this sum as follows

$$\sum_{s=0}^{\infty} \sum_{k=0}^{\infty} (k+s+1)^m a_s b_k \leq \sum_{s=0}^{\infty} \sum_{k=0}^{\infty} (k+1)^m (s+1)^m a_s b_k = \left(\sum_{i=0}^{\infty} (i+1)^m a_i \right) \left(\sum_{i=0}^{\infty} (i+1)^m b_i \right),$$

and the second result follows by the C_r inequality $(i+1)^m \leq 2^{m-1}(i^m + 1)$. The first result follows by the same arguments. \square

Proof of Lemma A.5. We can show this by the ratio test: $\left| \frac{(k+1)^m \lambda^{(k+1)m}}{k^m \lambda^{km}} \right| = \left(\frac{k+1}{k} \right)^m |\lambda|^m \rightarrow |\lambda|^m$ as $k \rightarrow \infty$. $|\lambda|^m < 1$ for all $1 \leq m < \infty$, since $|\lambda| < 1$, so the series converges absolutely, and the result follows. \square

Proof of Lemma A.6. For the first statement, note that $\sum_{j=0}^{\infty} \|\mathbb{A}^j\|_{\infty}^m = \psi_N^m \sum_{j=0}^{\infty} \lambda^{jm} = \frac{\psi_N^m}{1-\lambda^m} \leq C\psi_N^m$ and then $\sum_{j=0}^{\infty} j^m \|\mathbb{A}^j\|_{\infty}^m \leq C\psi_N^m$ by Lemma A.5. For the second statement, By the proof of Lemma 11 in [Krampe et al. \(2021\)](#),

$$\begin{aligned}
\hat{\mathbb{A}}^j - \mathbb{A}^j &= \sum_{s=0}^{j-1} \hat{\mathbb{A}}^s (\hat{\mathbb{A}} - \mathbb{A}) \mathbb{A}^{j-1-s}. \\
\sum_{j=0}^{\infty} \|\hat{\mathbb{A}}^j\|_{\infty}^m &\leq C \sum_{j=0}^{\infty} \|\mathbb{A}^j\|_{\infty}^m + C \sum_{j=1}^{\infty} \|\hat{\mathbb{A}}^j - \mathbb{A}^j\|_{\infty}^m \leq C\psi_N^m + C \sum_{j=1}^{\infty} \sum_{s=0}^{j-1} \|\hat{\mathbb{A}}^s (\hat{\mathbb{A}} - \mathbb{A}) \mathbb{A}^{j-1-s}\|_{\infty}^m \\
&\leq C\psi_N^m + C \|\hat{\mathbb{A}} - \mathbb{A}\|_{\infty}^m \sum_{j=1}^{\infty} \sum_{s=0}^{j-1} \|\hat{\mathbb{A}}^s\|_{\infty}^m \|\mathbb{A}^{j-1-s}\|_{\infty}^m.
\end{aligned}$$

By Lemma A.4, taking $a_j = \|\hat{\mathbb{A}}^j\|_{\infty}^m$ and $b_j = \|\mathbb{A}^j\|_{\infty}^m$, and on \mathcal{P} ,

$$\begin{aligned}
\sum_{j=0}^{\infty} \|\hat{\mathbb{A}}^j\|_{\infty}^m &= C\psi_N^m + \|\hat{\mathbb{A}} - \mathbb{A}\|_{\infty}^m \left(\sum_{i=0}^{\infty} \|\hat{\mathbb{A}}^i\|_{\infty}^m \right) \left(\sum_{i=0}^{\infty} \|\mathbb{A}^i\|_{\infty}^m \right) \\
&\leq C_1\psi_N^m + \|\hat{\mathbb{A}} - \mathbb{A}\|_{\infty}^m \left(\sum_{i=0}^{\infty} \|\hat{\mathbb{A}}^i\|_{\infty}^m \right) C_2\psi_N^m \leq C_1\psi_N^m + C_2\xi_{N,T}^m \psi_N^{2m} \left(\sum_{i=0}^{\infty} \|\hat{\mathbb{A}}^i\|_{\infty}^m \right).
\end{aligned}$$

By the assumption that $\xi_{N,T}\psi_T^2 \rightarrow 0$, for sufficiently large N, T we have $1 - C_2\xi_{N,T}^m\psi_N^{2m} \geq 1/C$, so

$$\sum_{j=0}^{\infty} \left\| \hat{\mathbb{A}}^j \right\|_{\infty}^m \leq \frac{C_1\psi_N^m}{1 - C_2\xi_{N,T}^m\psi_N^{2m}} \leq C\psi_N^m.$$

Similarly, we bound

$$\begin{aligned} \sum_{j=0}^{\infty} j^m \left\| \hat{\mathbb{A}}^j \right\|_{\infty}^m &\leq C_1 \sum_{j=0}^{\infty} j^m \left\| \mathbb{A}^j \right\|_{\infty}^m + C_2 \sum_{j=1}^{\infty} j^m \left\| \hat{\mathbb{A}}^j - \mathbb{A}^j \right\|_{\infty}^m \\ &\leq C_1\psi_N^m + C_2\xi_{N,T}^m\psi_N^m \sum_{j=1}^{\infty} \sum_{s=0}^{j-1} j^m \left\| \hat{\mathbb{A}}^s \right\|_{\infty}^m \left\| \mathbb{A}^{j-1-s} \right\|_{\infty}^m \\ &\leq C_1\psi_N^m + C_2\xi_{N,T}^m\psi_N^m \left(\sum_{i=0}^{\infty} (i^m + 1) \left\| \hat{\mathbb{A}}^i \right\|_{\infty}^m \right) \left(\sum_{i=0}^{\infty} (i^m + 1) \left\| \mathbb{A}^i \right\|_{\infty}^m \right) \\ &\leq C_1\psi_N^m + C_2\xi_{N,T}^m\psi_N^{2m} \left(\sum_{i=0}^{\infty} (i^m + 1) \left\| \hat{\mathbb{A}}^i \right\|_{\infty}^m \right) \\ &\leq C_1\psi_N^m + C_2\xi_{N,T}^m\psi_N^{2m} \left(\sum_{i=0}^{\infty} i^m \left\| \hat{\mathbb{A}}^i \right\|_{\infty}^m + \psi_N^m \right), \end{aligned}$$

such that

$$\sum_{j=0}^{\infty} j^m \left\| \hat{\mathbb{A}}^j \right\|_{\infty}^m \leq \frac{C_1\psi_N^m + C_2\xi_{N,T}^m\psi_N^{3m}}{1 - C_2\xi_{N,T}^m\psi_N^{2m}} \leq C\psi_N^m.$$

Finally,

$$\begin{aligned} \sum_{k=1}^{\infty} k^m \left\| \hat{\mathbb{A}}^k - \mathbb{A}^k \right\|_{\infty}^m &= \sum_{k=1}^{\infty} k^m \left\| \sum_{s=0}^{k-1} \hat{\mathbb{A}}^s (\hat{\mathbb{A}} - \mathbb{A}) \mathbb{A}^{k-1-s} \right\|_{\infty}^m \\ &\leq C\xi_{N,T}^m\psi_N^m \sum_{k=1}^{\infty} \sum_{s=0}^{k-1} k^m \left\| \hat{\mathbb{A}}^s \right\|_{\infty}^m \left\| \mathbb{A}^{k-1-s} \right\|_{\infty}^m \\ &\leq C\xi_{N,T}^m\psi_N^m \left(\sum_{i=0}^{\infty} (i^m + 1) \left\| \hat{\mathbb{A}}^i \right\|_{\infty}^m \right) \left(\sum_{i=0}^{\infty} (i^m + 1) \left\| \mathbb{A}^i \right\|_{\infty}^m \right) \\ &\leq C\xi_{N,T}^m\psi_N^{3m}. \quad \square \end{aligned}$$

Proof of Lemma A.7. Note that since $\rho(\mathbb{A}) = \lim_{k \rightarrow \infty} \left\| \mathbb{A}^k \right\|_{\infty}^{1/k} \leq \lim_{k \rightarrow \infty} (\psi_N \lambda^k)^{1/k} = \lambda$, it is also the case that for every \mathbb{A} , there exists some K^* such that for all $k \geq K^*$, $(\psi_N \lambda^k)^{1/k} \leq \lambda + (1 - \lambda)/3$. Rearranging, $\psi_N^{1/k} \leq \frac{1+2\lambda}{3\lambda} \implies k \geq \frac{\log(\psi_N)}{\log((1+2\lambda)/(3\lambda))}$, where $\log((1+2\lambda)/(3\lambda)) > 0$. So one such value is $K^* = \log(\psi_N)/C$. We then bound

$$\begin{aligned} \rho(\hat{\mathbb{A}}) &\leq \left\| \hat{\mathbb{A}}^{K^*} \right\|_{\infty}^{1/K^*} \leq \left(\left\| \hat{\mathbb{A}}^{K^*} - \mathbb{A}^{K^*} \right\|_{\infty} + \left\| \mathbb{A}^{K^*} \right\|_{\infty} \right)^{1/K^*} \leq \left\| \hat{\mathbb{A}}^{K^*} - \mathbb{A}^{K^*} \right\|_{\infty}^{1/K^*} + (\psi_N \lambda^{K^*})^{1/K^*} \\ &\leq \left\| \hat{\mathbb{A}}^{K^*} - \mathbb{A}^{K^*} \right\|_{\infty}^{1/K^*} + \frac{1+2\lambda}{3}. \end{aligned}$$

By Lemma A.6,

$$\begin{aligned} \left\| \hat{\mathbb{A}}^{K^*} - \mathbb{A}^{K^*} \right\|_{\infty}^{1/K^*} &\leq C_1 (\xi_{N,T} \psi_N^3)^{C_2/\log(\psi_N)} = C_1 \xi_{N,T}^{C_2/\log(\psi_N)} (\psi_N)^{3C_2/\log(\psi_N)} \\ &= C_1 \exp \left[C_2 \frac{\log(\xi_{N,T})}{\log(\psi_N)} \right] \exp \left[3C_2 \frac{\log(\psi_N)}{\log(\psi_N)} \right] = C_1 \exp \left[C_2 \frac{\log(\xi_{N,T})}{\log(\psi_N)} \right]. \end{aligned}$$

By assumption, we have $\frac{\log(\xi_{N,T})}{\log(\psi_N)} \rightarrow -\infty$, which implies that $\exp \left[C_2 \frac{\log(\xi_{N,T})}{\log(\psi_N)} \right] \rightarrow 0$, so for sufficiently large N, T , $C_1 \exp \left[C_2 \frac{\log(\xi_{N,T})}{\log(\psi_N)} \right] \leq \frac{1-\lambda}{3}$. The result then follows, with $\lambda' = \frac{2+\lambda}{3}$. \square

Proof of Lemma A.8. First, note that Assumption 3 also implies that the VAR process in eq. (1) is stationary, since $\rho(\mathbb{A}) = \lim_{k \rightarrow \infty} \|\mathbb{A}^k\|_{\infty}^{1/k} \leq \lim_{k \rightarrow \infty} (\psi_N \lambda^k)^{1/k} = \lambda$, and it can therefore be written as a VMA in eq. (3), with $\mathbf{B}_k = \mathbf{J} \mathbb{A}^k \mathbf{J}'$, where $\mathbf{J}_{N \times KN} = (\mathbf{I}, \mathbf{0}, \dots, \mathbf{0})$. We then have

$$\|\mathbf{B}_k\|_{\infty} = \|\mathbf{J} \mathbb{A}^k \mathbf{J}'\|_{\infty} \leq \|\mathbf{J}\|_{\infty} \|\mathbb{A}^k\|_{\infty} \|\mathbf{J}'\|_{\infty} = \|\mathbb{A}^k\|_{\infty},$$

so $\|\mathbf{B}_k\|_{\infty} \leq \psi_N \lambda^k$, and $\sum_{j=0}^{\infty} \|\mathbf{B}_j\|_{\infty} \leq \psi_N \sum_{j=0}^{\infty} \lambda^j = \frac{1}{1-\lambda} \psi_N$. By the arguments in the proof of Lemma 2.1 in Phillips and Solo (1992),

$$S_m = \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \|\mathbf{B}_k\|_{\infty} \right)^m \leq C \sum_{k=1}^{\infty} k^m \|\mathbf{B}_k\|_{\infty}^m \leq C \psi_N^m \sum_{k=1}^{\infty} k^m \lambda^{km} = C \psi_N^m.$$

using Lemma A.5, so the third statement follows. By the same arguments, we also have

$$\sum_{k=0}^{\infty} \|\mathbf{B}_k\|_{\infty}^m \leq 1 + \sum_{k=1}^{\infty} k^m \|\mathbf{B}_k\|_{\infty}^m \leq C \psi_N^m,$$

and the second statement follows. It also implies the first statement, by taking $m = 1$. When $\rho(\hat{\mathbb{A}}) < 1$ (guaranteed here by Lemma A.7 when $\frac{\log(\psi_N)}{\log(\xi_{N,T})} \rightarrow 0$), the estimated VAR process is also stationary and may be inverted into a VMA process with coefficient matrices $\hat{\mathbf{B}}_k = \mathbf{J} \hat{\mathbb{A}}^k \mathbf{J}'$, so we have $\|\hat{\mathbf{B}}_k - \mathbf{B}_k\|_{\infty} = \|\mathbf{J}(\hat{\mathbb{A}}^k - \mathbb{A}^k) \mathbf{J}'\|_{\infty} \leq \|\hat{\mathbb{A}}^k - \mathbb{A}^k\|_{\infty}$. Next, using the C_r inequality,

$$\begin{aligned} \sum_{j=0}^{\infty} \left(\sum_{k=j+1}^{\infty} \|\hat{\mathbf{B}}_k\|_{\infty} \right)^m &\leq C \sum_{k=1}^{\infty} k^m \|\hat{\mathbf{B}}_k\|_{\infty}^m \leq C \sum_{k=1}^{\infty} k^m \left(\|\hat{\mathbf{B}}_k - \mathbf{B}_k\|_{\infty} + \|\mathbf{B}_k\|_{\infty} \right)^m \\ &\stackrel{C_r}{\leq} C \sum_{k=1}^{\infty} k^m \left(2^{m-1} \|\hat{\mathbf{B}}_k - \mathbf{B}_k\|_{\infty}^m + 2^{m-1} \|\mathbf{B}_k\|_{\infty}^m \right) \\ &= C \sum_{k=1}^{\infty} k^m \|\hat{\mathbf{B}}_k - \mathbf{B}_k\|_{\infty}^m + C \sum_{k=1}^{\infty} k^m \|\mathbf{B}_k\|_{\infty}^m. \end{aligned}$$

We already showed that $\sum_{k=1}^{\infty} k^m \|\mathbf{B}_k\|_{\infty}^m \leq C \psi_N^m$, so we continue with the first term of this bound.

By Lemma A.6, we have

$$\sum_{k=1}^{\infty} k^m \|\hat{\mathbf{B}}_k - \mathbf{B}_k\|_{\infty}^m \leq \sum_{k=1}^{\infty} k^m \|\hat{\mathbb{A}}^k - \mathbb{A}^k\|_{\infty}^m \leq C \xi_{N,T}^m \psi_N^{3m},$$

so $S_m^* \leq C\xi_{N,T}^m \psi_N^{3m} + C\psi_N^m \leq C\psi_N^m$, for sufficiently large N, T , and the sixth statement follows. By the argument above, we also have the fifth statement

$$\sum_{k=0}^{\infty} \left\| \hat{\mathbf{B}}_k - \mathbf{B}_k \right\|_{\infty}^m \leq 1 + \sum_{k=1}^{\infty} k^m \left\| \hat{\mathbf{B}}_k - \mathbf{B}_k \right\|_{\infty}^m \leq C\xi_{N,T}^m \psi_N^{3m}.$$

The fourth statement follows by

$$\tilde{S}^* = \sum_{j=0}^{\infty} \left\| \hat{\mathbf{B}}_j \right\|_{\infty} \leq \sum_{j=0}^{\infty} \left\| \hat{\mathbf{B}}_j - \mathbf{B}_j \right\|_{\infty} + \sum_{j=0}^{\infty} \left\| \mathbf{B}_j \right\|_{\infty} \leq C\xi_{N,T} \psi_N^3 + C\psi_N \leq C\psi_N. \quad \square$$

Proof of Lemma A.9. By Markov's inequality and Lemma A.1.1, which implies $\mathbb{E} \exp(\max_j \epsilon_{j,t}^2/d_N^2) \leq 2$, we have that

$$\begin{aligned} \mathbb{P} \left(\max_j \sum_{t=1}^T \epsilon_{j,t}^2 > Ty \right) &= \mathbb{P} \left(\exp \left(\max_j \sum_{t=1}^T \epsilon_{j,t}^2/d_N^2 \right) > \exp(Ty/d_N^2) \right) \\ &\leq \frac{\mathbb{E} \exp \left(\max_j \sum_{t=1}^T \epsilon_{j,t}^2/d_N^2 \right)}{\exp(Ty/d_N^2)} \leq \frac{\prod_{t=1}^T \mathbb{E} \exp \left(\max_j \epsilon_{j,t}^2/d_N^2 \right)}{\exp(Ty/d_N^2)} \leq \frac{2^T}{\exp(Ty/d_N^2)}. \end{aligned}$$

Therefore

$$\mathbb{P} \left(\max_j \frac{1}{T} \sum_{t=1}^T \epsilon_{j,t}^2 \leq y \right) \geq 1 - \frac{2^T}{\exp(Ty/d_N^2)},$$

and we need to choose y such that this converges to 1. In particular, we take $y = Cd_N^2$, and the first statement follows.

For the second statement, we use Markov's and C_r inequality, and Lemma A.1.2

$$\mathbb{P} \left(\max_j \sum_{t=1}^T \epsilon_{j,t}^2 > Ty \right) \leq \frac{\mathbb{E} \left| \max_j \sum_{t=1}^T \epsilon_{j,t}^2 \right|^{m/2}}{(Ty)^{m/2}} \leq \frac{2^{m/2-1} \mathbb{E} \sum_{t=1}^T \left| \max_j \epsilon_{j,t} \right|^m}{(Ty)^{m/2}} \leq C \frac{Td_N^m}{(Ty)^{m/2}}.$$

Therefore

$$\mathbb{P} \left(\max_j \frac{1}{T} \sum_{t=1}^T \epsilon_{j,t}^2 \leq y \right) \geq 1 - C \frac{Td_N^m}{(Ty)^{m/2}},$$

which converges to 1 when $y \geq C \frac{d_N^2}{T^{(m-2)/m}} \eta_T^{-1}$, for some sequence $\eta_T \rightarrow 0$. $\frac{1}{T^{(m-2)/m}} \eta_T^{-1} \rightarrow 0$ when η_T converges sufficiently slowly, so we may take $y = Cd_N^2$ and the second statement follows. \square

Proof of Lemma A.10. We have that

$$\begin{aligned} \left\| \frac{1}{T} \sum_{t=1}^T \hat{\epsilon}_t \hat{\epsilon}'_t - \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t \right\|_{\max} &= \left\| \frac{1}{T} \sum_{t=1}^T [(\hat{\epsilon}_t - \epsilon_t)(\hat{\epsilon}'_t - \epsilon'_t) + (\hat{\epsilon}_t - \epsilon_t)\epsilon'_t + \epsilon_t(\hat{\epsilon}'_t - \epsilon'_t)] \right\|_{\max} \\ &\leq \left\| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_t - \epsilon_t)(\hat{\epsilon}'_t - \epsilon'_t) \right\|_{\max} + 2 \left\| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_t - \epsilon_t)\epsilon'_t \right\|_{\max}. \end{aligned}$$

By the Cauchy-Schwarz inequality,

$$\begin{aligned}
\left\| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_t - \epsilon_t) (\hat{\epsilon}'_t - \epsilon'_t) \right\|_{\max} &= \max_{r,s} \left| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_{r,t} - \epsilon_{r,t}) (\hat{\epsilon}_{s,t} - \epsilon_{s,t}) \right| \\
&\leq \max_{r,s} \left\{ \frac{1}{T} \left(\sum_{t=1}^T |\hat{\epsilon}_{r,t} - \epsilon_{r,t}|^2 \right)^{1/2} \left(\sum_{t=1}^T |\hat{\epsilon}_{s,t} - \epsilon_{s,t}|^2 \right)^{1/2} \right\} \\
&= \frac{1}{T} \max_r \left(\sum_{t=1}^T |\hat{\epsilon}_{r,t} - \epsilon_{r,t}|^2 \right) \stackrel{\mathcal{Q}}{\leq} \frac{1}{T} \max_r \|\hat{\epsilon}_r - \epsilon_r\|_2^2 \leq \phi_{N,T}.
\end{aligned}$$

Then

$$\begin{aligned}
\left\| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_t - \epsilon_t) \epsilon'_t \right\|_{\max} &= \max_{s,r} \left| \frac{1}{T} \sum_{t=1}^T (\hat{\epsilon}_{r,t} - \epsilon_{r,t}) \epsilon_{s,t} \right| \\
&\leq \max_{s,r} \left| \frac{1}{T} \left(\sum_{t=1}^T |\hat{\epsilon}_{r,t} - \epsilon_{r,t}|^2 \right)^{1/2} \left(\sum_{t=1}^T |\epsilon_{s,t}|^2 \right)^{1/2} \right| \\
&\leq \max_r \left| \left(\sum_{t=1}^T |\hat{\epsilon}_{r,t} - \epsilon_{r,t}|^2 \right)^{1/2} \right| \max_s \left| \frac{1}{T} \left(\sum_{t=1}^T |\epsilon_{s,t}|^2 \right)^{1/2} \right| \\
&= \frac{1}{\sqrt{T}} \max_r \|\hat{\epsilon}_r - \epsilon_r\|_2 \max_s \left| \frac{1}{T} \sum_{t=1}^T \epsilon_{s,t}^2 \right|^{1/2} \stackrel{\mathcal{Q}, \mathcal{R}_1}{\leq} C \sqrt{\phi_{N,T}} d_N
\end{aligned}$$

and the first statement follows. The second statement follows by identical steps except the last, where we use the set \mathcal{R}_2 to bound $\max_r \left| \frac{1}{T} \sum_{t=1}^T \epsilon_{r,t}^2 \right|^{1/2} \leq C d_N$. \square

Proof of Lemma A.11. By the union bound

$$\mathbb{P} \left(\left\| \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T} \sum_{t=1}^T \mathbb{E} \epsilon_t \epsilon'_t \right\|_{\max} \leq y \right) \geq 1 - \sum_{1 \leq s, r \leq N} \mathbb{P} \left(\left| \frac{1}{T} \sum_{t=1}^T [\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}] \right| > y \right).$$

Note that by Lemma 2.7.7 and Exercise 2.7.10 of Vershynin (2019) we have that under Assumption 2.1, $\epsilon_{r,t} \epsilon_{s,t}$ is sub-exponential with $\|\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}\|_{\psi_1} \leq C \|\epsilon_{r,t} \epsilon_{s,t}\|_{\psi_1} \leq \|\epsilon_{r,t}\|_{\psi_2} \|\epsilon_{s,t}\|_{\psi_2} \leq C$. Furthermore, by Theorem 2.8.1 of Vershynin (2019), we have Bernstein's inequality

$$\begin{aligned}
&\mathbb{P} \left(\left| \sum_{t=1}^T (\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}) \right| > T y \right) \\
&\leq 2 \exp \left(-C \min \left\{ \frac{T^2 y^2}{\sum_{t=1}^T \|\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}\|_{\psi_1}^2}, \frac{T y}{\max_t \|\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}\|_{\psi_1}} \right\} \right)
\end{aligned}$$

If we separately bound the terms in the minimum, $\sum_{t=1}^T \|\epsilon_{r,t} \epsilon_{s,t} - \mathbb{E} \epsilon_{r,t} \epsilon_{s,t}\|_{\psi_1}^2 \leq CT$, and

$\max_t \|\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t}\|_{\psi_1} \leq C$, so this simplifies to

$$\mathbb{P}\left(\left|\sum_{t=1}^T (\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t})\right| > Ty\right) \leq 2 \exp(-C \min\{Ty^2, Ty\}).$$

since we will choose $y \rightarrow 0$, the first term is smaller, and we obtain the bound $2 \exp(-CTy^2)$, and

$$\mathbb{P}\left(\left\|\frac{1}{T}\sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T}\sum_{t=1}^T \mathbb{E}\epsilon_t \epsilon'_t\right\|_{\max} \leq y\right) \geq 1 - C_1 N^2 \exp(-C_2 Ty^2).$$

We then find y by bounding $N^2 \exp(-C_2 \frac{Ty^2}{d_N^4}) \leq N^{-1} \implies y \geq C \frac{\sqrt{\log(N)}}{\sqrt{T}}$, and the first result follows.

For the second result,

$$\begin{aligned} \mathbb{P}\left(\left|\frac{1}{T}\sum_{t=1}^T [\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t}]\right| > y\right) &\leq \frac{\mathbb{E}\left|\sum_{t=1}^T [\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t}]\right|^{m/2}}{T^{m/2} y^{m/2}} \\ &\leq \frac{\sum_{t=1}^T \mathbb{E}\|\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t}\|_{L_{m/2}}^{m/2}}{T^{m/2} y^{m/2}}. \end{aligned}$$

By triangle and Jensen's, and Cauchy-Schwarz inequalities, and Assumption 2.2

$$\|\epsilon_{r,t}\epsilon_{s,t} - \mathbb{E}\epsilon_{r,t}\epsilon_{s,t}\|_{L_{m/2}}^{m/2} \leq C \|\epsilon_{r,t}\epsilon_{s,t}\|_{L_{m/2}}^{m/2} \leq C \|\epsilon_{r,t}\|_{L_m}^{m/2} \|\epsilon_{s,t}\|_{L_m}^{m/2} \leq C,$$

and

$$\mathbb{P}\left(\left\|\frac{1}{T}\sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T}\sum_{t=1}^T \mathbb{E}\epsilon_t \epsilon'_t\right\|_{\max} \leq y\right) \geq 1 - N^2 \frac{T}{T^{m/2} y^{m/2}}.$$

This probability then converges to 1 when $y \sim \frac{N^{4/m}}{T^{(m-2)/m}} \eta_T^{-1}$, and the second result follows. \square

Proof of Theorem 2. For $N \times N$ matrices $\mathbf{A}, \mathbf{B}, \mathbf{C}$,

$$\begin{aligned} \|\mathbf{A}\mathbf{B}\mathbf{C}'\|_{\max} &= \max_{1 \leq r, s \leq N} \|\mathbf{a}_r \mathbf{B} \mathbf{c}'_s\| = \max_{r, s} \left\| \sum_{1 \leq i, j \leq N} \mathbf{a}_{r,i} b_{i,j} \mathbf{c}'_{s,j} \right\| \leq \max_{i,j} |b_{i,j}| \max_{r,s} \left\{ \sum_{i,j} |a_{r,i}| |c_{s,j}| \right\} \\ &= \max_{i,j} |b_{i,j}| \max_{r,s} \{\|\mathbf{a}_r\|_1 \|\mathbf{c}_s\|_1\} \leq \|\mathbf{B}\|_{\max} \|\mathbf{A}\|_{\infty} \|\mathbf{C}\|_{\infty}. \end{aligned}$$

Using telescoping sums, sub-additivity of the $\|\cdot\|_{\max}$ norm, and the result above, we can rewrite

$$\begin{aligned} \|\hat{\Sigma} - \Sigma\|_{\max} &= \|\hat{\mathbf{B}}(1) \hat{\Sigma}_{\epsilon} \hat{\mathbf{B}}(1)' - \mathbf{B}(1) \Sigma_{\epsilon} \mathbf{B}(1)'\|_{\max} \\ &\leq \|\Delta \hat{\Sigma}_{\epsilon}\|_{\max} \|\Delta \hat{\mathbf{B}}(1)\|_{\infty}^2 + \|\Sigma_{\epsilon}\|_{\max} \|\Delta \hat{\mathbf{B}}(1)\|_{\infty}^2 + \|\Delta \hat{\Sigma}_{\epsilon}\|_{\max} \|\mathbf{B}(1)\|_{\infty}^2 \\ &\quad + 2 \|\Delta \hat{\Sigma}_{\epsilon}\|_{\max} \|\Delta \hat{\mathbf{B}}(1)\|_{\infty} \|\mathbf{B}(1)\|_{\infty} + 2 \|\Sigma_{\epsilon}\|_{\max} \|\Delta \hat{\mathbf{B}}(1)\|_{\infty} \|\mathbf{B}(1)\|_{\infty}, \end{aligned}$$

where $\Delta \hat{\Sigma}_{\epsilon} = \hat{\Sigma}_{\epsilon} - \Sigma_{\epsilon}$ and $\Delta \hat{\mathbf{B}}(1) = \hat{\mathbf{B}}(1) - \mathbf{B}(1)$. There are therefore 4 distinct expressions we

need to bound. On $\mathcal{Q} \cap \mathcal{R}_1 \cap \mathcal{S}_1$, by Lemma A.10

$$\begin{aligned} \left\| \Delta \hat{\Sigma}_\epsilon \right\|_{\max} &\leq \left\| \frac{1}{T} \sum_{t=1}^T \hat{\epsilon}_t \hat{\epsilon}'_t - \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t \right\|_{\max} + \left\| \frac{1}{T} \sum_{t=1}^T \epsilon_t \epsilon'_t - \frac{1}{T} \sum_{t=1}^T \mathbb{E} \epsilon_t \epsilon'_t \right\|_{\max} \\ &\leq C \left(\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{\sqrt{\log(N)}}{\sqrt{T}} \right). \end{aligned}$$

On $\mathcal{Q} \cap \mathcal{R}_2 \cap \mathcal{S}_2$

$$\left\| \Delta \hat{\Sigma}_\epsilon \right\|_{\max} \leq C \left(\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{(m-2)/m}} \eta_T^{-1} \right).$$

By Lemma A.8.6, on \mathcal{P} , when $\xi_{N,T} \psi_N^2 \rightarrow 0$ and N, T are sufficiently large

$$\left\| \Delta \hat{\mathcal{B}}(1) \right\|_{\infty} \leq \sum_{k=0}^{\infty} \left\| \hat{\mathbf{B}}_k - \mathbf{B}_k \right\|_{\infty} \leq C \xi_{N,T} \psi_N^3.$$

By Cauchy-Schwarz and Lemma A.1,

$$\left\| \Sigma_\epsilon \right\|_{\max} = \max_{r,s} \left| \frac{1}{T} \sum_{t=1}^T \mathbb{E} \epsilon_{r,t} \epsilon_{s,t} \right| \leq \max_{r,t} \|\epsilon_{r,t}\|_{L_2}^2 \leq C.$$

Under Assumption 3, by Lemma A.8.1

$$\left\| \mathcal{B}(1) \right\|_{\infty} \leq \sum_{k=0}^{\infty} \left\| \mathbf{B}_k \right\|_{\infty} = \tilde{S} \leq C \psi_N.$$

Plugging these in, we find

$$\begin{aligned} \left\| \hat{\Sigma} - \Sigma \right\|_{\max} &\leq C_1 \left\| \Delta \hat{\Sigma}_\epsilon \right\|_{\max} (\xi_{N,T}^2 \psi_N^6 + \psi_N^2 + \xi_{N,T} \psi_N^4) + C_2 (\xi_{N,T}^2 \psi_N^6 + \xi_{N,T} \psi_N^4) \\ &\leq C \left(\left\| \Delta \hat{\Sigma}_\epsilon \right\|_{\max} \psi_N^2 + \xi_{N,T} \psi_N^4 \right), \end{aligned}$$

since we assume $\xi_{N,T} \psi_N^2 \rightarrow 0$. Plugging in the respective bounds on $\left\| \Delta \hat{\Sigma}_\epsilon \right\|_{\max}$, we obtain the bounds on \mathcal{T}_1 and \mathcal{T}_2 . Since Lemma A.8 requires that N, T are sufficiently large for the inequality to hold, we have $\lim_{N,T \rightarrow \infty} \mathbb{P}(\mathcal{T}_i) = 1$ rather than simply $\mathbb{P}(\mathcal{T}_i) = 1$. \square

Proof of Lemma A.12. By the union bound and equation (2.14) in Vershynin (2019), and using Lemma A.1.1

$$\begin{aligned} \mathbb{P} \left(\max_{j,t} |\epsilon_{j,t}| \leq y \right) &= 1 - \sum_{t=1}^T \mathbb{P} \left(\max_j |\epsilon_{j,t}| > y \right) \geq 1 - \sum_{t=1}^T 2 \exp \left(-C y^2 / \left\| \max_j |\epsilon_{j,t}| \right\|_{\psi_2}^2 \right) \\ &\geq 1 - 2T \exp \left(\frac{-C y^2}{d_N^2} \right). \end{aligned}$$

This probability converges to 1 when taking $y = d_N \log(T)$, showing the first statement. By union

bound, Markov's inequality and Lemma A.1.2,

$$\mathbb{P} \left(\max_{j,t} |\epsilon_{j,t}| \leq y \right) = 1 - \sum_{t=1}^T \mathbb{P} \left(\max_j |\epsilon_{j,t}| > y \right) \geq 1 - \sum_{t=1}^T \frac{\mathbb{E} \left[\max_j |\epsilon_{j,t}|^m \right]}{y^m} \geq 1 - T d_N^m y^{-m}.$$

This probability converges to 1 when $y = d_N T^{1/m} \eta_T^{-1}$ for some $\eta_T \rightarrow 0$, showing the second statement. \square

Proof of Lemma A.13. By submultiplicativity of the Orlicz norm,

$$\max_t \left\| \max_j \epsilon_{j,t}^* \right\|_{\psi_2}^* = \max_t \left\| \max_j \hat{\epsilon}_{j,t} \gamma_t \right\|_{\psi_2}^* \leq \max_t \left\| \max_j \hat{\epsilon}_{j,t} \right\|_{\psi_2}^* \max_t \|\gamma_t\|_{\psi_2}^*.$$

Since γ_t is by construction independent of \mathbf{X} and identically Gaussian distributed, we have by Example 2.5.8 in Vershynin (2019) $\max_t \|\gamma_t\|_{\psi_2}^* = \max_t \|\gamma_t\|_{\psi_2} \leq C$.

$$\begin{aligned} \max_t \left\| \max_j \hat{\epsilon}_{j,t} \right\|_{\psi_2}^* &= \max_t \inf \left\{ \lambda > 0 : \mathbb{E}^* \exp \left(\left| \max_j \hat{\epsilon}_{j,t} \right|^2 / \lambda^2 \right) \leq 2 \right\} \\ &= \max_t \inf \left\{ \lambda > 0 : \exp \left(\left| \max_j \hat{\epsilon}_{j,t} \right|^2 / \lambda^2 \right) \leq 2 \right\} \\ &\leq \max_t \inf \left\{ \lambda > 0 : \exp \left(\max_j |\hat{\epsilon}_{j,t}|^2 / \lambda^2 \right) \leq 2 \right\} \\ &= \max_t \inf \left\{ \lambda > 0 : \max_j |\hat{\epsilon}_{j,t}| \leq \sqrt{\log(2)} \lambda \right\}. \end{aligned}$$

Therefore, up to a $\sqrt{\log(2)}$ constant, any bound on $\max_{j,t} |\hat{\epsilon}_{j,t}|$ is also a bound on $\max_t \left\| \max_j \hat{\epsilon}_{j,t} \right\|_{\psi_2}^*$. By triangle inequality, $\max_{j,t} |\hat{\epsilon}_{j,t}| \leq \max_{j,t} |\hat{\epsilon}_{j,t} - \epsilon_{j,t}| + \max_{j,t} |\epsilon_{j,t}|$, and we further bound the individual terms using \mathcal{Q}

$$\max_{j,t} |\hat{\epsilon}_{j,t} - \epsilon_{j,t}| \leq \max_j \sqrt{\sum_{t=1}^T |\hat{\epsilon}_{j,t} - \epsilon_{j,t}|^2} = \sqrt{T} \max_j \sqrt{\frac{1}{T} \|\hat{\epsilon}_j - \epsilon_j\|_2^2} \leq \sqrt{T \phi_{N,T}}.$$

Then, on \mathcal{U}_1 , $\max_{j,t} |\epsilon_{j,t}| \leq d_N \log(T)$, and the first statement follows.

For the second statement, since γ_t is again i.i.d. Gaussian, we have $\max_t \|\gamma_t\|_{L_m} \leq C$ for all $0 < m < \infty$, so

$$\begin{aligned} \max_t \left\| \max_j \epsilon_{j,t}^* \right\|_{L_m}^* &= \max_t \left(\mathbb{E}^* \max_j |\epsilon_{j,t}^*|^m \right)^{1/m} = \max_t \left(\mathbb{E}^* \max_j |\hat{\epsilon}_{j,t} \gamma_t|^m \right)^{1/m} \\ &= \max_t \left(\max_j |\hat{\epsilon}_{j,t}|^m \mathbb{E} |\gamma_t|^m \right)^{1/m} \leq C \max_{j,t} |\hat{\epsilon}_{j,t}|. \end{aligned}$$

We use the same arguments for bounding this term as for the first statement, using that on \mathcal{U}_2 , $\max_{j,t} |\epsilon_{j,t}| \leq d_N T^{1/m} \eta_T^{-1}$, and the second statement is obtained. \square

Proof of Lemma A.14. By Theorem in Chernozhukov et al. (2020), for all $\lambda > 0$

$$M_{N,T}^* \leq C \left\{ (\log(T)) \left(\Delta_0 + \sqrt{\Delta_1 \log(N)} + \frac{(\mathcal{M} \log(N))^2}{T \Lambda_{\min}(\tilde{\Sigma})} \right) + \sqrt{\frac{\Lambda_1 M(\lambda)}{T \Lambda_{\min}^2(\tilde{\Sigma})}} + \frac{\lambda \log(N)^{3/2}}{\sqrt{T \Lambda_{\min}(\tilde{\Sigma})}} \right\},$$

where

$$\Delta_0 = \frac{\log(N)}{\Lambda_{\min}(\tilde{\Sigma})} \|\Sigma - \Sigma^*\|_{\max},$$

$$\begin{aligned} \Sigma^* &= \mathbb{E}^* \left[\left(\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)^* \epsilon_t^* \right) \left(\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)^* \epsilon_t^* \right)' \right] = \mathcal{B}(1)^* \left(\frac{1}{T} \sum_{s,t} \mathbb{E}^* \epsilon_s^* \epsilon_t^{*'} \right) \mathcal{B}(1)^{*'} \\ &= \mathcal{B}(1)^* \left(\frac{1}{T} \sum_t \mathbb{E}^* \epsilon_t^* \epsilon_t^{*'} \right) \mathcal{B}(1)^{*'} = \mathcal{B}(1)^* \left(\frac{1}{T} \sum_t \epsilon_t^* \mathbb{E}(\gamma_t^2) \epsilon_t^{*'} \right) \mathcal{B}(1)^{*'} = \hat{\mathcal{B}}(1) \hat{\Sigma}_\epsilon \hat{\mathcal{B}}(1)', \end{aligned}$$

since conditionally on \mathbf{X} , ϵ_s^* and ϵ_t^* are independent for $s \neq t$. Furthermore,

$$\Delta_1 = \frac{(\log N)^2}{T^2 \Lambda_{\min}^2(\tilde{\Sigma})} \max_j \sum_{t=1}^T \mathbb{E}^* |\mathcal{B}(1)_j^* \epsilon_t^*|^4,$$

$$\mathcal{M} = \left(\mathbb{E}^* \left[\max_{j,t} |\mathcal{B}(1)_j^* \epsilon_t^*|^4 \right] \right)^{1/4},$$

$$\Lambda_1 = (\log(N))^2 \log(T) \log(NT),$$

and

$$M(\lambda) = \max_t \mathbb{E}^* [\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty \mathbb{1} \{\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda\}].$$

We now derive bounds for each of these expressions. By similar arguments to those in the proof of Lemma A.2, by Assumption 1, $\Lambda_{\min}(\tilde{\Sigma}) \geq 1/C$, and on \mathcal{T}_1 or \mathcal{T}_2 , we have respectively

$$\Delta_0 \leq C \log(N) \psi_N^2 \left[\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{\sqrt{\log(N)}}{\sqrt{T}} + \xi_{N,T} \psi_N^2 \right],$$

or

$$\Delta_0 \leq C \log(N) \psi_N^2 \left[\phi_{N,T} + d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{(m-2)/m}} \eta_T^{-1} + \xi_{N,T} \psi_N^2 \right].$$

For Δ_1

$$\frac{(\log N)^2}{T^2 \Lambda_{\min}^2(\tilde{\Sigma})} \max_j \sum_{t=1}^T \mathbb{E}^* |\mathcal{B}(1)_j^* \epsilon_t^*|^4 \leq C \frac{\log(N)^2 \tilde{S}^{*4} \left\| \max_j |\epsilon_{j,t}^*| \right\|_{L_4}^{*4}}{T},$$

so on $\mathcal{U}_1 \cap \mathcal{Q}$ or $\mathcal{U}_2 \cap \mathcal{Q}$, we have by Lemma A.13

$$\Delta_1 \leq C \frac{\log(N)^2 \tilde{S}^{*4} d_N^{*4}}{T}.$$

Note that d_N^* is different depending on which clause of Lemma A.13 we use. For \mathcal{M} we have

$$\left(\mathbb{E}^* \left[\max_{j,t} |\mathcal{B}(1)_j^* \epsilon_t^*|^4 \right] \right)^{1/4} \leq \tilde{S}^* \left\| \max_{j,t} |\epsilon_{j,t}^*| \right\|_{L_4}^* \leq \tilde{S}^* \left\| \max_{j,t} |\epsilon_{j,t}^*| \right\|_{L_m}^*,$$

so on $\mathcal{U}_1 \cap \mathcal{Q}$ or $\mathcal{U}_2 \cap \mathcal{Q}$, we have respectively

$$\mathcal{M} \leq \tilde{S}^* \sqrt{\log(T)} d_N^* \text{ or } \mathcal{M} \leq \tilde{S}^* T^{1/m} d_N^*.$$

For $M(\lambda)$, we have by Cauchy-Schwarz

$$\begin{aligned} \max_t \mathbb{E}^* \left[\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty \mathbf{1} \{ \|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda \} \right] &\leq \max_t \left\{ \left\| \|\mathcal{B}(1)^* \epsilon_t^*\|_\infty \right\|_{L_2}^* (\mathbb{P}^* (\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda))^{1/2} \right\} \\ &\leq \tilde{S}^* \max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{L_2}^* \max_t (\mathbb{P}^* (\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda))^{1/2}. \end{aligned}$$

On $\mathcal{U}_1 \cap \mathcal{Q}$, by equation (2.14) in Vershynin (2019),

$$\mathbb{P}^* (\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda) \leq 2 \exp \left(-C \frac{\lambda^2}{d_N^{*2} \tilde{S}^{*2}} \right),$$

and we may let $\lambda = C d_N^* \tilde{S}^* \sqrt{\log(d_N^* \tilde{S}^*)}$ such that $M(\lambda) \leq C$. On $\mathcal{U}_2 \cap \mathcal{Q}$, we use Hölder's inequality instead of Cauchy-Schwarz,

$$\max_t \mathbb{E}^* \left[\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty \mathbf{1} \{ \|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda \} \right] \leq \tilde{S}^* \max_t \left\| \max_j |\epsilon_{j,t}| \right\|_{L_m}^* \max_t (\mathbb{P}^* (\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda))^{\frac{m-1}{m}}.$$

By Markov's inequality

$$\mathbb{P}^* (\|\mathcal{B}(1)^* \epsilon_t^*\|_\infty > \lambda) \leq \frac{\mathbb{E}^* \left[\max_j |\epsilon_{j,t}^*| \right]}{\lambda / \tilde{S}^*} \leq \frac{d_N^* \tilde{S}^*}{\lambda}.$$

We then take $\lambda = C (d_N^* \tilde{S}^*)^{\frac{2m-1}{m-1}}$ such that $M(\lambda) \leq C$. The result then follows by plugging in the bounds on these terms, and using that $\phi_{N,T} \rightarrow 0$, $d_N \geq 1$, $\tilde{S}^* \geq 1$, $d_N^* \rightarrow \infty$ to omit asymptotically dominated terms. \square

Proof of Lemma A.15. The general strategy for this proof is similar to that of Lemma A.3, but we first need to properly treat ϵ_0^* .

$$\begin{aligned} \epsilon_0^* &= \mathbf{x}_0 - \sum_{k=1}^{K-1} \hat{A}_k \mathbf{x}_{-k} = \left(\sum_{j=0}^{\infty} \mathbf{B}_j \epsilon_{-j} \right) - \sum_{k=1}^{K-1} \hat{A}_k \left(\sum_{j=0}^{\infty} \mathbf{B}_j \epsilon_{-k-j} \right) \\ &= \left(\sum_{j=0}^{\infty} \mathbf{B}_j \epsilon_{-j} \right) - \sum_{j=0}^{\infty} \sum_{k=1}^{K-1} \left(\hat{A}_k \mathbf{B}_j \epsilon_{-k-j} \right) = \sum_{\ell=0}^{\infty} \left(\mathbf{B}_\ell - \sum_{j=1}^{K-1} \hat{A}_j \mathbf{B}_{\ell-j} \right) \epsilon_{-\ell} \\ &= \sum_{\ell=0}^{\infty} \left(\mathbf{B}_\ell - \hat{A}^\dagger \mathbf{B}_\ell^\dagger \right) \epsilon_{-\ell}, \end{aligned}$$

where $\mathbf{B}_j = \mathbf{0}$ for $j < 0$, and $\hat{\mathbf{A}}^\dagger = [\mathbf{A}_1, \dots, \mathbf{A}_{K-1}]$, and $\mathbf{B}_\ell^\dagger = \begin{bmatrix} \mathbf{B}_{\ell-1} \\ \vdots \\ \mathbf{B}_{\ell-K+1} \end{bmatrix}$. For convenience

in later arguments, we consider the m th power, and use mainly a combination of triangle and C_r inequalities

$$\begin{aligned} \left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \boldsymbol{\epsilon}_0^* \right\|_\infty^m &= \max_p \frac{1}{\sqrt{T}^m} \left| \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) (\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger) \right]_{p, \cdot} \boldsymbol{\epsilon}_{-j-\ell} \right|^m \\ &\leq C \max_p \frac{1}{\sqrt{T}^m} \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) (\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger) \right]_{p, \cdot} \right\|_1 \max_p |\boldsymbol{\epsilon}_{p, -j-\ell}|^m \\ &\leq C \frac{1}{\sqrt{T}^m} \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) (\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger) \right\|_\infty^m \max_p |\boldsymbol{\epsilon}_{p, -j-\ell}|^m. \end{aligned}$$

By submultiplicativity of $\|\cdot\|_\infty$

$$\begin{aligned} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) (\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger) \right\|_\infty^m &\leq \left\| \sum_{k=j+1}^{\infty} \mathbf{B}_k \right\|_\infty^m \left\| \mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right\|_\infty^m \\ &\leq C \left[\|\mathbf{B}_\ell\|_\infty^m + \|\hat{\mathbf{A}}^\dagger\|_\infty^m \|\mathbf{B}_\ell^\dagger\|_\infty^m \right] \left(\sum_{k=j+1}^{\infty} \|\mathbf{B}_k\|_\infty \right)^m. \end{aligned}$$

In the following steps, we will use that $\sum_{k=1}^{\infty} \|\mathbf{B}_k\|_\infty^m \leq C\psi_N^m$, by Lemma A.8.2, using Assumption 3.

Since $\hat{\mathbf{A}}^\dagger$ is a submatrix of $\hat{\mathbf{A}}$, by Assumption 3, on \mathcal{P} ,

$$\|\hat{\mathbf{A}}^\dagger\|_\infty^m \leq \|\hat{\mathbf{A}}\|_\infty^m \leq C \left[\|\hat{\mathbf{A}} - \mathbf{A}\|_\infty^m + \|\mathbf{A}\|_\infty^m \right] \leq C \left[\xi_{N,T}^m \psi_N^m + C\psi_N^m \right] \leq C\psi_N^m.$$

For N, T sufficiently large.

$$\begin{aligned} \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) (\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger) \right\|_\infty^m &\leq C \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left(\left[\|\mathbf{B}_\ell\|_\infty^m + \psi_N^m \|\mathbf{B}_\ell^\dagger\|_\infty^m \right] \left[\sum_{k=j+1}^{\infty} \|\mathbf{B}_k\|_\infty \right]^m \right) \\ &= C \left(\sum_{j=0}^{\infty} \left[\sum_{k=j+1}^{\infty} \|\mathbf{B}_k\|_\infty \right]^m \right) \left(\sum_{\ell=0}^{\infty} \left[\|\mathbf{B}_\ell\|_\infty^m + \psi_N^m \|\mathbf{B}_\ell^\dagger\|_\infty^m \right] \right) = CS_m \sum_{\ell=0}^{\infty} \left[\|\mathbf{B}_\ell\|_\infty^m + \psi_N^m \|\mathbf{B}_\ell^\dagger\|_\infty^m \right] \\ &\leq CS_m \left[\psi_N^m + \psi_N^m \sum_{\ell=0}^{\infty} \|\mathbf{B}_\ell^\dagger\|_\infty^m \right]. \end{aligned}$$

Due to the construction of \mathbf{B}_ℓ^\dagger , $\left\| \mathbf{B}_\ell^\dagger \right\|_\infty \leq \sum_{j=1}^{K-1} \|\mathbf{B}_{\ell-j}\|_\infty$, so by Assumption 3

$$\begin{aligned} \sum_{\ell=0}^{\infty} \left\| \mathbf{B}_\ell^\dagger \right\|_\infty^m &\leq C \sum_{\ell=0}^{\infty} \sum_{j=1}^{K-1} \|\mathbf{B}_{\ell-j}\|_\infty^m = C \left[\sum_{\ell=1}^{K-1} \sum_{j=1}^{\ell} \|\mathbf{B}_{\ell-j}\|_\infty^m + \sum_{\ell=K}^{\infty} \sum_{j=1}^{K-1} \|\mathbf{B}_{\ell-j}\|_\infty^m \right] \\ &\leq C \psi_N^m \left[\sum_{\ell=0}^{K-1} \sum_{j=1}^{\ell} \lambda^{m(\ell-j)} + \sum_{\ell=K}^{\infty} \sum_{j=1}^{K-1} \lambda^{m(\ell-j)} \right] \\ &= C \psi_N^m \left[\sum_{j=1}^{K-1} j \lambda^{m(K-1-j)} + \sum_{\ell=K}^{\infty} \sum_{j=1}^{K-1} \lambda^{m(\ell-j)} \right] \\ &= C \psi_N^m \left[\frac{\lambda^{Km} - K\lambda^m + K - 1}{(1 - \lambda^m)^2} + \frac{\lambda^m - \lambda^{Km}}{(1 - \lambda^m)^2} \right] = \frac{K - 1}{1 - \lambda^m}. \end{aligned}$$

so

$$\sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right\|_\infty^m \leq CK S_m \psi_N^m,$$

and

$$\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_\infty^m \leq C \frac{1}{\sqrt{T}^m} K S_m \psi_N^m \max_p |\epsilon_{p,-j-\ell}|^m.$$

From here, we can apply the same steps as in Lemma A.3. By the union bound and Hoeffding's inequality,

$$\begin{aligned} &\mathbb{P}^* \left(\max_p \frac{1}{\sqrt{T}} \left| \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right]_{p,\cdot} \epsilon_{-j-\ell} \right| > \eta_T \right) \\ &\leq \sum_{p=1}^N \mathbb{P}^* \left(\left| \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right]_{p,\cdot} \epsilon_{-j-\ell} \right| > \eta_T \sqrt{T} \right) \\ &\leq \sum_{p=1}^N 2 \exp \left(-C \frac{[\eta_T \sqrt{T}]^2}{\sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right]_{p,\cdot} \epsilon_{-j-\ell} \right\|_{\psi_2}^{*2}} \right). \end{aligned}$$

Following the arguments above, by Lemma A.1.1,

$$\begin{aligned} &\sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right]_{p,\cdot} \epsilon_{-j-\ell} \right\|_{\psi_2}^{*2} \\ &\leq \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right\|_\infty^2 \left\| \max_p |\epsilon_{p,-j-\ell}| \right\|_{\psi_2}^{*2} \leq CK S_2 \psi_N^2 d_N^2, \end{aligned}$$

since $\left\| \max_p |\epsilon_{p,-j-\ell}| \right\|_{\psi_2}^{*2} = \left\| \max_p |\epsilon_{p,-j-\ell}| \right\|_{\psi_2}^2$, as ϵ_t is independent of \mathbf{X} .

$$\mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_{\infty} > \eta_T \right) \leq 2N \exp \left(-C \frac{\eta_T^2 T}{K \psi_N^2 d_N^2 S_2} \right).$$

With Lemma A.1.2, the proof follows almost identically to Lemma A.3, taking in the M-Z inequality

$$y_{j,\ell} = \left[\left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right]_{p,} \epsilon_{-j-\ell}, \text{ and bounding}$$

$$\begin{aligned} \mathbb{E}^* \left(\left| \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} y_{j,\ell} \right|^m \right) &\leq C \mathbb{E}^* \left[\left(\sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} |y_{j,\ell}|^2 \right)^{m/2} \right] \leq C \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \|y_{j,\ell}\|_{L_m}^{*m} \\ &\leq C \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \left\| \left(\sum_{k=j+1}^{\infty} \mathbf{B}_k \right) \left(\mathbf{B}_\ell - \hat{\mathbf{A}}^\dagger \mathbf{B}_\ell^\dagger \right) \right\|_{\infty}^m \left\| \max_p |\epsilon_{p,-j-\ell}| \right\|_{L_m}^{*m} \\ &\leq CK S_m \psi_N^m d_N^m, \end{aligned}$$

and

$$\mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_{\infty} > \eta_T \right) \leq C \frac{NK S_m \psi_N^m d_N^m}{(\eta_T \sqrt{T})^m}. \quad \square$$

Proof of Theorem 3. This proof largely follows the same structure as the proof of Theorem 1. By Lemma A.7, the bootstrap process is invertible, and we write the Beveridge-Nelson decomposition of the process:

$$\mathbf{x}_t^* = \mathcal{B}(L)^* \epsilon_t^* = \mathcal{B}(1)^* \epsilon_t^* - (1-L) \tilde{\mathcal{B}}^*(L) \epsilon_t^*, \text{ where } \tilde{\mathcal{B}}^*(L) = \sum_{j=0}^{\infty} \tilde{\mathcal{B}}_j^* L^j, \tilde{\mathcal{B}}_j^* = \sum_{k=j+1}^{\infty} \mathbf{B}_k^*,$$

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* = \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)^* \epsilon_t^* - \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_T^* + \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^*.$$

Define

$$x_T^{(\max)*} = \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* \right\|_{\infty}, \quad \epsilon_T^{(\max)*} = \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathcal{B}(1)^* \epsilon_t^* \right\|_{\infty}, \quad z_T^{(\max)*} = \|z\|_{\infty},$$

$$F_{1,T}^*(y) := \mathbb{P} \left(x_T^{(\max)*} \leq y \right) \quad F_{2,T}^*(y) := \mathbb{P} \left(\epsilon_T^{(\max)*} \leq y \right)$$

$$G_T^*(y) := \mathbb{P} \left(z_T^{(\max)*} \leq y \right) \quad r_T^* := x_T^{(\max)*} - \epsilon_T^{(\max)*}$$

Then

$$|r_T^*| \leq \left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}(L)^* \epsilon_T^* \right\|_{\infty} + \left\| \frac{1}{\sqrt{T}} \tilde{\mathcal{B}}^*(L) \epsilon_0^* \right\|_{\infty} = R_T^* + R_0^*.$$

The main deviation from Theorem 1 is in the treatment of the terms R_T^* and R_0^* . For R_T^* , we may simply apply Lemma A.3 to the bootstrap quantity directly, using Lemma A.13 instead of

Lemma A.1: On $\mathcal{U}_1 \cap \mathcal{Q}$, by Lemma A.13.1, we have

$$\mathbb{P}^*(R_T^* > \eta_T) \leq 2N \exp\left(-C \frac{\eta_T^2 T}{d_N^{*2} S_2^*}\right).$$

Similarly, on $\mathcal{U}_2 \cap \mathcal{Q}$, by Lemma A.13.2

$$\mathbb{P}^*(R_T^* > \eta_T) \leq C \frac{N d_N^{*m} S_m^*}{(\eta_T \sqrt{T})^m}.$$

For R_0^* , we have a different bound by Lemma A.15: Under Assumption 3 and 2.1, on \mathcal{P}

$$\mathbb{P}^*(R_0^* > \eta_T) \leq 2N \exp\left(-C \frac{\eta_T^2 T}{K \psi_N^2 d_N^2 S_2}\right),$$

and under Assumption 3 and 2.2, on \mathcal{P}

$$\mathbb{P}^*(R_0^* > \eta_T) \leq C \frac{N K S_m \psi_N^m d_N^m}{(\eta_T \sqrt{T})^m}.$$

Under Assumption 2.1, we can bound

$$\begin{aligned} \mathbb{P}^*(|r_T^*| > 2\eta_{T,1}) &\leq \mathbb{P}^*(\eta_{T,1} + R_0^* > 2\eta_{T,1}) \times 1 + 1 \times \mathbb{P}^*(R_T^* > \eta_{T,1}) \\ &\leq C_1 N \left[\exp\left(-C_2 \frac{\eta_{T,1}^2 T}{K \psi_N^2 d_N^2 S_2}\right) + \exp\left(-C_3 \frac{\eta_{T,1}^2 T}{d_N^{*2} S_2^*}\right) \right] =: \eta_{T,2}. \end{aligned}$$

Continue with

$$\begin{aligned} |F_{1,T}^*(y) - G_T^*(y)| &\leq \underbrace{\left| \mathbb{P}^*\left(\epsilon_T^{(\max)*} \leq y + 2\eta_{T,1}\right) - \mathbb{P}^*\left(z_T^{(\max)*} \leq y + 2\eta_{T,1}\right) \right|}_{A_{T,1}^*(y + 2\eta_{T,1})} \\ &\quad + \underbrace{\left| \mathbb{P}^*\left(z_T^{(\max)*} \leq y + 2\eta_{T,1}\right) - \mathbb{P}^*\left(z_T^{(\max)*} \leq y\right) \right|}_{A_{T,2}^*(y)} + 4\eta_{T,2}. \end{aligned}$$

Note that $\sup_{y \in \mathbb{R}} A_{T,1}^*(y + 2\eta_{T,1}) = M_{N,T}^*$ which can be bounded by Lemma A.14, and $\sup_{y \in \mathbb{R}} A_{T,2}^*(y) \leq C\eta_{T,1} \sqrt{\log(N)}$ by Lemma A.1 in Chernozhukov et al. (2017). We therefore have the bound

$$\sup_{y \in \mathbb{R}} |F_{1,T}^*(y) - G_T^*(y)| \leq M_{N,T}^* + C_1 \left[\eta_{T,1} \sqrt{\log N} + N \exp\left(-C_2 \frac{\eta_{T,1}^2 T}{K \psi_N^2 d_N^2 S_2}\right) + N \exp\left(-C_3 \frac{\eta_{T,1}^2 T}{d_N^{*2} S_2^*}\right) \right].$$

Under Assumption 3, by Lemma A.8.3 and Lemma A.8.6, on \mathcal{P} , both S_2 and S_2^* can be bounded by $C\psi_N^2$, and since d_N^* grows faster than d_N , for sufficiently large N, T we have $d_N^*/d_N \geq 1$ and $C\psi_N^2 \geq 1$ (we also obviously have $K \geq 1$). We choose $\eta_{T,1} = \sqrt{\log(N \log(N)) \frac{K d_N^{*2} \psi_N^4}{C T}}$ which lets us

bound

$$\begin{aligned}
& C_1 \left[\eta_{T,1} \sqrt{\log N} + N \exp \left(-C_2 \frac{\eta_{T,1}^2 T}{K \psi_N^2 d_N^2 S_2} \right) + N \exp \left(-C_3 \frac{\eta_{T,1}^2 T}{d_N^{*2} S_2^*} \right) \right] \\
& \leq C_1 \left[\frac{\sqrt{K} d_N^* \psi_N^2}{\sqrt{T}} \sqrt{\log(N) \log(N \log(N))} + \frac{N}{(N \log(N))^{C_2 \frac{d_{T,1}^{*2}}{d_N^2}}} + \frac{N}{(N \log(N))^{C_3 K \psi_N^2}} \right] \\
& \leq C \left[\frac{\sqrt{K} \log(N) d_N^* \psi_N^2}{\sqrt{T}} + \frac{1}{\log(N)} \right],
\end{aligned}$$

and the result of the first statement follows.

Under Assumption 2.2, we may follow the same steps as above, taking

$$\eta_{T,2} := C \left[\frac{NK S_m d_N^m}{(\eta_{T,1} \sqrt{T})^m} + \frac{N S_m^* d_N^{*m}}{(\eta_{T,1} \sqrt{T})^m} \right] \leq C \frac{NK \psi_N^{2m} d_N^{*m}}{(\eta_{T,1} \sqrt{T})^m}.$$

We then have the bound

$$\begin{aligned}
\sup_{y \in \mathbb{R}} |F_{1,T}^*(y) - G_T^*(y)| & \leq M_{N,T}^* + C \left[\eta_{T,1} \sqrt{\log N} + \frac{NK \psi_N^{2m} d_N^{*m}}{(\eta_{T,1} \sqrt{T})^m} \right] \\
& \leq M_{N,T}^* + C (NK d_N^{*m} \psi_N^m)^{\frac{1}{m+1}} \left(\frac{\sqrt{\log(N)}}{\sqrt{T}} \right)^{\frac{m}{m+1}},
\end{aligned}$$

and the result of the second statement follows. \square

Proof of Theorem 4. With a simple telescopic sum argument

$$\begin{aligned}
& \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} \leq y \right) - \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* \right\|_{\infty} \leq y \right) \right| \\
& \leq \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} \leq y \right) - \mathbb{P}(\|\mathbf{z}\|_{\infty} \leq y) \right| + \sup_{y \in \mathbb{R}} \left| \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* \right\|_{\infty} \leq y \right) - \mathbb{P}(\|\mathbf{z}\|_{\infty} \leq y) \right| \\
& = \sup_{y \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t \right\|_{\infty} \leq y \right) - \mathbb{P}(\|\mathbf{z}\|_{\infty} \leq y) \right| + \sup_{y \in \mathbb{R}} \left| \mathbb{P}^* \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{x}_t^* \right\|_{\infty} \leq y \right) - \mathbb{P}^*(\|\mathbf{z}\|_{\infty} \leq y) \right| \\
& \leq J_{N,T} + J_{N,T}^*,
\end{aligned}$$

which are bounded by Theorems 1 and 3 respectively. The bounds provided by these theorems only hold under Assumptions 1 to 3, on the set $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{T}_i \cap \mathcal{U}_i$ ($i \in \{1, 2\}$, depending on which moment assumption we make in Assumption 2) and for sufficiently large N, T . The latter is satisfied as we look consider the asymptotic case as $N, T \rightarrow \infty$ in this theorem. Consider first the set \mathcal{T}_i . By Theorem 2, it holds with probability converging to 1 on the set $\mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_i \cap \mathcal{S}_i$. These sets then hold with probability converging to 1 individually by Assumption 4, Assumption 5, Lemma A.9, and Lemma A.11 respectively. By the union bound, we then have

$$\mathbb{P} \left(\mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_i \cap \mathcal{S}_i \right) \geq 1 - [\mathbb{P}(\mathcal{P}^c) + \mathbb{P}(\mathcal{Q}^c) + \mathbb{P}(\mathcal{R}_i^c) + \mathbb{P}(\mathcal{S}_i^c)] \rightarrow 1,$$

as $N, T \rightarrow \infty$. We therefore also have $\lim_{N, T \rightarrow \infty} \mathbb{P}(\mathcal{T}_i) = 1$, unconditionally. to see why, we may alternatively phrase the result of Theorem 2 as $\lim_{N, T \rightarrow \infty} \mathbb{P}(\mathcal{T}_i | \mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_i \cap \mathcal{S}_i) = 1$. We may then write the unconditional probability as

$$\begin{aligned} \lim_{N, T \rightarrow \infty} \mathbb{P}(\mathcal{T}_i) &= \lim_{N, T \rightarrow \infty} \underbrace{\mathbb{P}(\mathcal{T}_i | \mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_i \cap \mathcal{S}_i)}_{\rightarrow 1} \times \underbrace{\mathbb{P}(\mathcal{P} \cap \mathcal{Q} \cap \mathcal{R}_i \cap \mathcal{S}_i)}_{\rightarrow 1} \\ &\quad + \underbrace{\mathbb{P}(\mathcal{T}_i | \mathcal{P}^c \cup \mathcal{Q}^c \cup \mathcal{R}_i^c \cup \mathcal{S}_i^c)}_{\leq 1} \times \underbrace{\mathbb{P}(\mathcal{P}^c \cup \mathcal{Q}^c \cup \mathcal{R}_i^c \cup \mathcal{S}_i^c)}_{\rightarrow 0} = 1. \end{aligned}$$

We can apply the same logic to the bounds on $J_{N,T}$, $J_{N,T}^*$, and $M_{N,T}^*$ in Lemma A.14 (the bound on $M_{N,T}$ in Lemma A.2 holds deterministically), noting that we also have $\lim_{N, T \rightarrow \infty} \mathbb{P}(\mathcal{U}_i) = 1$ by Lemma A.12. Then if each bound holds with probability converging to 1, the bound obtained by combining them all holds with probability converging to 1 also.

Combining all bounds under Assumption 2.1, we obtain the bound

$$\begin{aligned} &C \left[\underbrace{\frac{b_T \log(N)^{3/2} \log(T)}{\sqrt{T}} + \frac{b_T \log(N)^2}{\sqrt{T}}}_{M_{N,T}} + \frac{\log(N) d_N \sqrt{S_2}}{\sqrt{T}} \right. \\ &+ \underbrace{\log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{\frac{m-2}{m}}} + \xi_{N,T} \psi_N^2 \right]}_{M_{N,T}^*} \\ &+ \underbrace{(\tilde{S}^* d_N^*)^2 \left[\frac{\log(N)^{3/2} \left(\log(T) + (\tilde{S}^* d_N^*)^{\frac{1}{m-1}} \right)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)}{T^{\frac{m-2}{m}}} \right]}_{M_{N,T}^*} + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \\ &\left. + \frac{\sqrt{K} \log(N) d_N^* \psi_N^2}{\sqrt{T}} + \frac{1}{\log(N)} \right] \end{aligned}$$

After plugging in $b_T = \tilde{S}^2 d_N^2$, $\tilde{S}, \tilde{S}^* \leq C \psi_N$, $S_2 \leq C \psi_N^2$, $d_N = C \sqrt{\log(N)}$, $d_N^* = \sqrt{T \phi_{N,T}} + d_N \log(T)$, $d_N^* \leq CT \phi_{N,T} + \log(N) \log(T)^2$, and eliminating dominated terms, we obtain the simpler bound

$$\begin{aligned} &C \left\{ \psi_N^2 \left[\frac{\log(N)^3}{\sqrt{T}} + \frac{\log(N)^{5/2} \log(T)^3}{\sqrt{T}} + \sqrt{\phi_{N,T}} \log(N)^{3/2} \log(T) + \phi_{N,T} \sqrt{T} \log(N)^{3/2} \log(T) \right. \right. \\ &\quad \left. \left. + \psi_N^2 \xi_{N,T} \log(N) \log(T) + \sqrt{K} \left(\sqrt{\phi_{N,T}} + \frac{\log(N)^{3/2} \log(T)}{\sqrt{T}} \right) \right] + \frac{1}{\log(N)} \right\} \\ &= C \left\{ \psi_N^2 \left[\frac{\ell_N^3}{\sqrt{T}} + \frac{\ell_N^{5/2} \ell_T^3}{\sqrt{T}} + \sqrt{\phi_{N,T}} \ell_N^{3/2} \ell_T + \phi_{N,T} \sqrt{T} \ell_N^{3/2} \ell_T \right. \right. \\ &\quad \left. \left. + \psi_N^2 \xi_{N,T} \ell_N \ell_T + \sqrt{K} \left(\sqrt{\phi_{N,T}} + \frac{\ell_N^{3/2} \ell_T}{\sqrt{T}} \right) \right] + \frac{1}{\ell_N} \right\}, \end{aligned}$$

where $\ell_T = \log(T)$, $\ell_N = \log(N)$. Combining all bounds under Assumption 2.1,

$$\begin{aligned}
& C \underbrace{\left\{ \frac{b_T (\log N)^{3/2} \log(T)}{\sqrt{T}} + \frac{b_T^2 \log(N)^2 \log(T)}{T^{1-2/m}} + \left[\frac{b_T^m \log(N)^{3m/2-4} \log(T) \log(NT)}{T^{m/2-1}} \right]^{\frac{1}{m-2}} \right\}}_{M_{N,T}} \\
& + (Nd_N^m S_m)^{\frac{1}{m+1}} \left(\frac{\sqrt{\log(N)}}{\sqrt{T}} \right)^{\frac{m}{m+1}} + \underbrace{\log(N) \log(T) \psi_N^2 \left[d_N \sqrt{\phi_{N,T}} + \frac{N^{4/m}}{T^{\frac{m-2}{m}}} + \xi_{N,T} \psi_N^2 \right]}_{M_{N,T}^*} \\
& + (\tilde{S}^* d_N^*)^2 \underbrace{\left[\frac{\log(N)^{3/2} \left(\log(T) + (\tilde{S}^* d_N^*)^{\frac{1}{m-1}} \right)}{\sqrt{T}} + \frac{\log(N)^2 \log(T)}{T^{\frac{m-2}{m}}} \right]}_{M_{N,T}^*} + \sqrt{\frac{\log(N)^2 \log(T) \log(NT)}{T}} \\
& + (NK d_N^m \psi_N^m)^{\frac{1}{m+1}} \left(\frac{\sqrt{\log(N)}}{\sqrt{T}} \right)^{\frac{m}{m+1}} \}.
\end{aligned}$$

Now plugging in $d_N = CN^{1/m}$, $d_N^* = C(\sqrt{T\phi_{N,T}} + (NT)^{1/m}\eta_T^{-1})$, we give the simpler bound

$$\begin{aligned}
& C\eta_T^{-1} \left\{ \psi_N^2 \left[\ell_N \ell_T \sqrt{\phi_{N,T}} + \ell_N \ell_T \xi_{N,T} \psi_N^2 + \left(T\phi_{N,T} + (NT)^{2/m} \right) \left(\frac{\ell_N^{3/2} \ell_T}{\sqrt{T}} + \frac{\ell_N^2 \ell_T^2}{T} \right) \right] \right. \\
& \left. + \frac{\psi_N N^{1/m} \ell_N^{3/2} \ell_T}{\sqrt{T}} + \frac{\psi_N^m N \ell_N^{3m/2-4} \ell_T \ell_{NT}}{T^{m/2-1}} + \left[\left(\frac{\sqrt{\ell_N}}{\sqrt{T}} \psi_N \right)^m NK \left(\sqrt{T\phi_{N,T}}^m + NT \right) \right]^{\frac{1}{m+1}} \right\}.
\end{aligned}$$

where $\ell_{NT} = \log(NT)$. \square

Proof of Corollary 1. Under this choice of growth rates, we may take $\lambda_j = T^{\frac{1}{4} + \frac{8a}{m} - \frac{m-2}{m}} \eta_T^{-1}$, such that $\max_j \frac{1}{T} \|\hat{\epsilon}_j - \epsilon_j\|_2^2 \leq C\lambda_j^{2-r} s_{r,j} = C\frac{T^{\frac{12a+3}{m}}}{T}$ and $\|\hat{\mathbf{A}} - \mathbf{A}\|_\infty = \max_j \|\hat{\beta}_j - \beta_j\|_1 \leq C\lambda_j s_{0,j} = C\frac{T^{\frac{4a+1}{m}}}{T^{1/4}}$. We also have $\psi_N = a \log(T) = C \log(T)$, and similarly for any other $\log(N)$ term. Therefore, we may take $\xi_{N,T} = C\frac{T^{\frac{4a+1}{m}}}{\log(T)T^{1/4}}$, $\phi_{N,T} = \frac{T^{\frac{12a+3}{m}}}{T}$. Plugging these into the bound of Theorem 4 and eliminating dominated terms, we see it converges to 0 when

$$\eta_T^{-1} \frac{\ell_T^{3/2} T^{\frac{26a+6}{m}}}{\sqrt{T}} \rightarrow 0.$$

Note that any $\log(T)$ term is dominated by a term polynomial in T , so this terms converges to 0 when $52a + 12 < m$. Note that this condition also satisfies equation C.5 in Adamek et al. (2022), which in this case requires only that $16a + 4 < m$.

Finally, note that the two rates required for Theorem 4 are also satisfied: $\xi_{N,T} \psi_N^2 = \xi_{N,T} = C\frac{T^{\frac{4a+1}{m}}}{T^{1/4}} \log(T) \rightarrow 0$, and $\frac{\log(\xi_{N,T})}{\log(\psi_N)} = \frac{([4a+1]/m-1/4) \log(T) - \log(\log(T))}{\log(\log(T))} \rightarrow -\infty$. \square

References

- Adamek, R., S. Smeeke, and I. Wilms (2022). Lasso inference for high-dimensional time series. *Journal of Econometrics*, Forthcoming.
- Bi, D., H. L. Shang, Y. Yang, and H. Zhu (2021). AR-sieve bootstrap for high-dimensional time series. arXiv e-print 2112.00414.
- Bickel, P. J., Y. Ritov, and A. B. Tsybakov (2009). Simultaneous analysis of lasso and dantzig selector. *Annals of Statistics* 37(4), 1705–1732.
- Chang, Y. and J. Y. Park (2003). A sieve bootstrap for the test of a unit root. *Journal of Time Series Analysis* 24(4), 379–400.
- Chernozhukov, V., D. Chetverikov, and K. Kato (2013). Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *Annals of Statistics* 41, 2786–2819.
- Chernozhukov, V., D. Chetverikov, and K. Kato (2017). Central limit theorems and bootstrap in high dimensions. *Annals of Probability* 45, 2309–2352.
- Chernozhukov, V., D. Chetverikov, K. Kato, and Y. Koike (2022). High-dimensional data bootstrap. arXiv e-print 2205.09691.
- Chernozhukov, V., D. Chetverikov, and Y. Koike (2020). Nearly optimal central limit theorem and bootstrap approximations in high dimensions. arXiv e-print 2012.09513.
- Giessing, A. and J. Fan (2020). Bootstrapping ℓ_p -statistics in high dimensions. arXiv e-print 2006.13099.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 6(2), 65–70.
- Kilian, L. and H. Lütkepohl (2017). *Structural vector autoregressive analysis*. Cambridge University Press.
- Kock, A. B. and L. Callot (2015). Oracle inequalities for high dimensional vector autoregressions. *Journal of Econometrics* 186, 325–344.
- Krampe, J., J.-P. Kreiss, and E. Paparoditis (2021). Bootstrap based inference for sparse high-dimensional time series models. *Bernoulli* 27(3), 1441–1466.

- Liu, L. and D. Zhang (2021). Robust estimation of high-dimensional vector autoregressive models. arXiv e-print 2109.10354.
- Masini, R. P., M. C. Medeiros, and E. F. Mendes (2021). Regularized estimation of high-dimensional vector autoregressions with weakly dependent innovations. *Journal of Time Series Analysis* 43(4), 532–557.
- Medeiros, M. C. and E. F. Mendes (2016). ℓ_1 -regularization of high-dimensional time-series models with non-gaussian and heteroskedastic errors. *Journal of Econometrics* 191, 255–271.
- Meyer, M. and J.-P. Kreiss (2015). On the vector autoregressive sieve bootstrap. *Journal of Time Series Analysis* 36, 377–397.
- Nicholson, W. B., I. Wilms, J. Bien, and D. S. Matteson (2020). High dimensional forecasting via interpretable vector autoregression. *Journal of Machine Learning Research* 21(166), 1–52.
- Paparoditis, E. (1996). Bootstrapping autoregressive and moving average parameter estimates of infinite order vector autoregressive processes. *Journal of Multivariate Analysis* 57(2), 277–296.
- Park, J. Y. (2002). An invariance principle for sieve bootstrap in time series. *Econometric Theory* 18(2), 469–490.
- Phillips, P. C. B. and V. Solo (1992). Asymptotics for linear processes. *Annals of Statistics* 20, 971–1001.
- Romano, J. P. and M. Wolf (2005). Exact and approximate stepdown methods for multiple hypothesis testing. *Journal of the American Statistical Association* 100(469), 94–108.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B* 58(1), 267–288.
- Trapani, L. (2013). On bootstrapping panel factor series. *Journal of Econometrics* 172(1), 127–141.
- van der Vaart, A. W. and J. A. Wellner (1996). *Weak convergence and empirical processes*. Springer New York, NY.
- Vershynin, R. (2019). *High-Dimensional Probability*. Cambridge University Press.
- Wu, W. B. (2005). Nonlinear system theory: Another look at dependence. *Proceedings of the National Academy of Sciences* 102(40), 14150–14154.

Zhang, D. and W. B. Wu (2017). Gaussian approximation for high dimensional time series. *Annals of Statistics* 45(5), 1895–1919.

Zhang, X. and G. Cheng (2018). Gaussian approximation for high dimensional vector under physical dependence. *Bernoulli* 24(4A), 2640–2675.