

# Non-Relativistic Particles Interact with Potential Under the effect of Quantum flux in a space-time with a distortion of a vertical line into a vertical spiral

PRABIR RUDRA<sup>(a)1</sup>, FAIZUDDIN AHMED<sup>(b)2</sup> and HOUCINE AOUNALLAH<sup>(c)3</sup>

<sup>1</sup> *Department of Mathematics, Asutosh College, Kolkata-700026, India*

<sup>2</sup> *Department of Physics, University of Science & Technology Meghalaya, Ri-Bhoi, Meghalaya-793101, India*

<sup>3</sup> *Department of Science and Technology. Echahid Cheikh Larbi Tebessi University - Tébessa, Algeria*

PACS 03.65.Ge – Non-relativistic wave equation

PACS 03.65.Pm – solutions of wave-equation: bound-state

PACS 61.72.Lk – Linear defects: dislocations, disclinations

PACS 02.30.Hq – Ordinary differential equations

**Abstract** – The quantum motions of the non-relativistic particles interact with potential in the presence of the Aharonov-Bohm (AB) flux field in a topological defect geometry associated to screw dislocation, which corresponds to a distortion of a vertical line into a vertical spiral are investigated. We derived the radial Schrödinger wave equation with anharmonic oscillator potential (superposition of harmonic oscillator plus inverse square potential) and obtain the eigenvalue solution through the Heun equation. The expressions for the ground state energy level and the radial wave function for the radial mode  $n = 1$  are analyzed. Then, we utilize this eigenvalue solution to molecular potential models (pseudoharmonic and shifted pseudoharmonic potentials) and presented the eigenvalue solutions. We study the quantum system with inverse square potential in the presence of the quantum flux field and obtained the ground state energy level and the radial wave function through the same procedure. We show that the eigenvalue solutions are influenced by the topological defect of screw dislocation characterized by the parameter  $\beta$  and the quantum flux field  $\Phi_{AB}$  which causes the shift in the energy spectrum and shows the gravitational analogue to the Aharonov-Bohm effect.

**I. Introduction.** – Our universe went through a phase transition process in the primordial era. This transition in phase was associated with the decoupling of the fundamental interactions which resulted in symmetry breaking. With the breaking of symmetry, weird geometrical forms are expected to come into existence. Consequently, such topological defects [1] are supposed to be observed in the universe in the form of exotic cosmological objects. Some of them may be listed as the domain wall [2], cosmic string [3–5], global monopole [6], etc. In practice, these objects are yet to be observed, so they can still be considered hypothetical objects, instead of reality. But some of them are so obvious that they are expected

to be observed in near future. Out of the probable candidates, the global monopole is the most promising one that is expected to be observed; hence, a lot of research has been dedicated to it in recent years [7–13]. In addition, exact solutions of the KG-equation have been extensively investigated in the background of global monopole [14–22].

In quantum mechanical systems, several authors have studied the non-relativistic Schrodinger wave equation using different potentials and obtained the eigenvalue solutions. Some of the potentials used are pseudoharmonic potential, Mie-type potential, Kratzer potential, Yukawa potential, Hulthen potential, Morse potential, hyperbolic Potential, non-central potential, Manning-Rosen (MR) potential, Rosen-Morse potential, Deng-Fan potential, Poschl-Teller potential (see, [23] for references). In the literature, we see that in order to obtain these eigen-

<sup>(a)</sup>E-mail: [prudra.math@gmail.com](mailto:prudra.math@gmail.com)

<sup>(b)</sup>E-mail: [faizuddinahmed15@gmail.com](mailto:faizuddinahmed15@gmail.com)

<sup>(c)</sup>E-mail: [houcine12400@gmail.com](mailto:houcine12400@gmail.com)

value solutions of the non-relativistic wave equation, different methods or mathematical techniques have been employed, such as the parametric Nikiforov-Uvarov method and its functional analysis, Super-symmetric quantum mechanics (SUSY), Asymptotic iteration method (AIM), Laplace's transformation method, through confluent hypergeometric and bi-confluent Heun equations method, etc. (see, [24] for references). It must be noted that all these investigations have been performed in Minkowski flat space. In addition, a few investigations have been done in the background of the topological defects produced by a point-like global monopole [25, 26]. In the present analysis, we are mainly interested in the anharmonic oscillator potential given by [23, 27–32]

$$V(r) = \left( \eta r^2 + \frac{\gamma}{r^2} + \delta \right), \quad (1)$$

where  $\delta, \gamma, \eta > 0$ . From this potential expression, one can recover the harmonic oscillator potential for  $\gamma \rightarrow 0, \delta \rightarrow 0$ . For  $\eta \rightarrow 0, \delta \rightarrow 0$ , we have the inverse square potential and, setting the potential strengths  $\eta, \gamma, \delta$ , one can recover pseudoharmonic potential (which we will discuss as an application).

Our motivation is the result presented in Ref. [33], where authors investigated the quantum effects of the harmonic oscillator in space-time with a distortion of a vertical line into a vertical spiral. As a natural extension, we study the quantum motions of the non-relativistic particles interacting with anharmonic oscillator potential (superposition of harmonic oscillator plus inverse square potential with a constant term) under the influence of the Aharonov-Bohm flux field in the same space-time background. We solve the radial wave equation through the Heun equation and obtain the eigenvalue solution of the quantum system. We show that the eigenvalue solution gets modified by the inverse square potential plus constant term in addition to the harmonic oscillator and the quantum flux field. We then utilize this eigenvalue solution to molecular potential models (pseudoharmonic potential, shifted pseudoharmonic potential) and presented the eigenvalue solutions. Finally, we study the same quantum system with only inverse square potential and the quantum flux field and obtain the eigenvalue solution using the same technique. Throughout the analysis, the system of units is chosen such that  $c = 1 = \hbar = G$ .

**II. Radial Solutions with Anharmonic oscillator potential Under AB-Flux Field in Topological Defect.** – In this section, we study the non-relativistic particles confined by the AB-flux field in a topological defect geometry with screw dislocation in the presence of an anharmonic oscillator potential and analyze the effects of the topological defect.

The Hamiltonian operator of a quantum system with interaction potential  $V(r)$  is described by [23–26, 34]

$$\hat{H} = -\frac{1}{2M} \left( \frac{1}{\sqrt{g}} D_i \left( \sqrt{g} g^{ij} D_j \right) \right) + V(r), \quad (2)$$

where  $M$  is the mass of the non-relativistic particle,  $D_i \equiv (\partial_i - i e A_i)$  with  $e$  is the electric charges, and  $A_i$  is the electromagnetic potential. In this analysis, we choose the electromagnetic three-vector potential  $\vec{A}$  with only the non-zero component [35]

$$A_r = 0 = A_z, \quad A_\phi = \frac{\Phi_{AB}}{2\pi}, \quad (3)$$

where,  $\Phi_{AB} = \text{const}$  is the Aharonov-Bohm magnetic flux,  $\Phi_0$  is the quantum of magnetic flux, and  $\Phi$  is the amount of magnetic flux which is a positive integer. It is mentioning that the Aharonov-Bohm effect has been investigated in several branches of physics by many authors (see, Ref. [20] and related references therein).

The topological defect geometry in the coordinates  $(r, \phi, z)$  is given by [33, 34, 36]

$$ds_{3D}^2 = dr^2 + r^2 d\phi^2 + 2\beta d\phi dz + dz^2, \quad (4)$$

The parameter  $\beta$  is a positive constant characterise the screw dislocation (torsion field) [33, 37, 38]. The ranges of this dislocation parameter are in the intervals  $0 < \beta < 1$ .

Writing the eigenvalue equation  $\hat{H} \Psi = E \Psi$  using Eq. (2) in the space-time background (4), we have

$$\begin{aligned} & -\frac{1}{2M} \left[ \frac{\partial^2 \Psi}{\partial r^2} + \frac{r}{r^2 - \beta^2} \frac{\partial \Psi}{\partial r} \right. \\ & \left. + \frac{1}{r^2 - \beta^2} \left( \frac{\partial}{\partial \phi} - i e A_\phi - \beta \frac{\partial}{\partial z} \right)^2 \right] + V(r) \Psi \\ & = E \Psi. \end{aligned} \quad (5)$$

In quantum mechanical system, the wave function  $\Psi(r, \phi, z)$  is always expressible in terms of different variables. Let us suppose that  $\Psi(r, \phi, z) = \psi(r) f(\phi) h(z)$ . Therefore, we choose a possible wave function in terms of a radial function  $\psi(r)$  given by

$$\Psi(r, \phi, z) = e^{i l \phi} e^{i k z} \psi(r), \quad (6)$$

where  $l = 0, \pm 1, \pm 2, \dots$  are the eigenvalues of the angular quantum operator and  $k > 0$  is an arbitrary constant.

Therefore, substituting the interactions potential (1), the electromagnetic three-vector potential component (3), and the wave function (6) into the Eq. (5), we obtain the following differential radial equation

$$\psi'' + \frac{r}{r^2 - \beta^2} \psi' + \left[ \Lambda - 2M\eta r^2 - \frac{2M\gamma}{r^2} - \frac{\iota^2}{r^2 - \beta^2} \right] \psi = 0, \quad (7)$$

where we have set the parameters

$$\begin{aligned} \Lambda &= 2M(E - \delta) - k^2, \quad \iota = (l - \Phi - \beta k), \\ \Phi &= \frac{\Phi_{AB}}{\Phi_0}, \quad \Phi = 2\pi e^{-1}. \end{aligned} \quad (8)$$

Now, we introduce a new variable via  $x = \frac{r^2}{\beta^2}$  in the Eq. (7), we obtain

$$4x\psi''(x) + \frac{4x-2}{x-1}\psi'(x) + \left[ \Lambda\beta^2 - \omega^2x - \frac{2M\gamma}{x} - \frac{l^2}{x-1} \right] \psi(x) = 0, \quad (9)$$

where  $\omega = \sqrt{2M\eta}\beta^2$ . If one choose  $\eta = \frac{1}{2}M\omega_0^2$ , where  $\omega_0$  is the oscillator frequency, then the parameter  $\omega \rightarrow M\omega_0\beta^2$ . In that case, the system we study here is called the harmonic oscillator problem interacts with inverse square potential in the background of topological defect.

In quantum system, it is required that the wave function  $\psi(x)$  must be regular everywhere. Therefore, the wave function in our case also should satisfies the boundary condition, that is,  $\psi(x) \rightarrow 0$  for both at  $x \rightarrow 0$  and  $x \rightarrow \infty$ . We choose a possible solution to the Eq (9) given by

$$\psi(x) = x^{\frac{1}{4} + \frac{j}{2}} e^{-\frac{\omega}{2}x} G(x), \quad (10)$$

where  $j = \sqrt{2M\gamma + \frac{1}{4}}$  and  $G(x)$  is the confluent Heun equation given by

$$G(x) = H_c \left( -\omega, j, -\frac{1}{2}, \frac{\Lambda\beta^2}{4}, \frac{3}{8} - \frac{l^2 + \Lambda\beta^2}{4}; x \right). \quad (11)$$

We assume  $G(x)$  in the following power series form

$$G(x) = \sum_{i=0}^{\infty} c_i x^i. \quad (12)$$

Substituting the wave function (10) and using (12) in the Eq. (9), we obtain the following recurrence relation

$$c_{i+2} = \frac{d_1}{d_3} c_{i+1} + \frac{d_2}{d_3} c_i \quad (13)$$

with the coefficient

$$c_1 = \frac{\left[ 2\omega(1+j) - l^2 - \Lambda\beta^2 + \frac{1}{2} + j \right]}{4(1+j)} c_0 \quad (14)$$

and we set the different parameters as

$$\begin{aligned} d_1 &= \left( n + \omega + \frac{3}{2} + j \right) (n+1) \\ &\quad - \frac{l^2 + \Lambda\beta^2 - \frac{1}{2} - j - 2\omega(1+j)}{4}, \\ d_2 &= -\omega n + \frac{\Lambda\beta^2 - \omega(3+2j)}{4}, \\ d_3 &= \left( n + \frac{3+2j}{2} \right) (n+2). \end{aligned} \quad (15)$$

Let us consider  $i = n-1$  when  $c_{n+1} = 0$ , we have from the relation (14)

$$c_n = \frac{4\omega(n-1) - \Lambda\beta^2 + \omega(3+2j)}{4n(n+\omega + \frac{1}{2} + j) - l^2 - \Lambda\beta^2 + \frac{1}{2} + j + 2\omega(1+j)} c_{n-1}$$

The condition  $c_{n+1} = 0$  is required to have a finite degree polynomial of the power series solution  $G(x) = c_0 + c_1x + \dots + c_nx^n$  so that the wave function  $\psi$  considered in (10) is regular everywhere.

Now, we can evaluate the individual energy levels the corresponding radial wave function for the radial mode  $n = 1, 2, \dots$ . For example, we consider the radial mode  $n = 1$  that corresponds to the ground state or the lowest state of the quantum syste. Thus, we obtain

$$c_1 = \frac{\omega(3+2j) - \Lambda\beta^2}{6 + 2\omega(j+3) + 5j - l^2 - \Lambda\beta^2 + \frac{1}{2}} c_0. \quad (16)$$

Comparing equation (14) with (16) and using equation (8), we obtain the ground state energy levels  $E_{l,1}$  given by

$$E_{l,1} = \frac{k^2}{2M} + \frac{\lambda}{2M} + \delta, \quad (17)$$

where

$$\lambda = \frac{1}{\beta^2} \left[ 3 - 2l^2 + 4\omega(2+j) + 2j \pm \sqrt{\Delta} \right] \quad (18)$$

and

$$\begin{aligned} \Delta &= 16l^2(1+j) + 16\omega(2+j) + 14\omega^2 - 44j \\ &\quad - 32M\gamma - 8. \end{aligned} \quad (19)$$

And that the ground state wave function is given by

$$\psi_{1,l}(x) = x^{\frac{1}{4} + \frac{\sqrt{2M\gamma + \frac{1}{4}}}{2}} e^{-\frac{\sqrt{2M\eta}\beta^2}{2}x} (c_0 + c_1x). \quad (20)$$

We can see that the energy eigenvalue depends on the screw dissociation parameter  $\beta$  and the geometric quantum phase which causes shifted the energy level and get them modified. Following the similar procedure, one can find other energy levels  $E_{2,l}, E_{3,l}, \dots$  and the radial wave function  $\psi_{1,l}, \psi_{2,l}, \dots$  for the radial mode  $n \geq 2$ . We see that the presence of the inverse square potential with constant term in addition to the harmonic oscillator like potential and the quantum flux field modified the eigenvalue solution in comparison to those result obtained in [33].

Below, we utilize the above eigenvalue solution to diatomic molecular potential models, such as pseudoharmonic potential and shifted pseudoharmonic potential and presented the energy levels and the wave function.

**III. Few Applications of the Eigenvalue Solutions to Molecular Potential.** – The above eigenvalue solution of the quantum system is utilize to diatomic molecular potential models, such as pseudoharmonic potential, shifted pseudoharmonic potential and analyze the effects of screw dislocation on the eigenvalue solutions.

#### Case A: Pseudoharmonic potential

The pseudoharmonic potential is recovered by setting the parameters  $\eta = \frac{D_e}{r_0^2}$ ,  $\gamma = D_e r_0^2$ , and  $\delta = -2D_e$  into

the potential expression (1) as follows

$$V(r) = D_e \left( \frac{r}{r_0} - \frac{r_0}{r} \right)^2, \quad (21)$$

where  $D_e$  is the dissociation energy of the molecule and  $r_0$  is the equilibrium separation between inter-nuclear. This potential has widely been used in atomic and molecular physics by many researchers [28–32].

The ground state energy level using this potential (21) becomes

$$E_{l,1} = \frac{k^2}{2M} + \frac{\chi}{2M} - 2D_e, \quad (22)$$

where

$$\begin{aligned} \chi &= \frac{1}{\beta^2} \left[ 3 - 2\iota^2 + 4\omega(2 + \sigma) + 2\sigma \pm \sqrt{\Pi} \right], \\ \Pi &= 16\iota^2(1 + \sigma) + 16\omega(2 + \sigma) + 12\omega^2 - 44\sigma \\ &\quad - 32M\gamma - 8, \\ \omega &= \sqrt{2MD_e} \frac{\beta^2}{r_0}, \quad \sigma = \sqrt{\frac{2MD_e}{r_0^2} + \frac{1}{4}}. \end{aligned} \quad (23)$$

### Case B: Shifted Pseudoharmonic potential

The shifted pseudoharmonic potential is recovered by setting the parameters  $\eta = \frac{D_e}{r_0^2}$ ,  $\gamma = D_e r_0^2$ , and  $\delta = 0$  into the potential expression (1) as follows [28, 29]

$$V(r) = D_e \left( \frac{r}{r_0} - \frac{r_0}{r} \right)^2 + 2D_e, \quad (24)$$

which is shifted by an amount twice the dissociation energy of the molecule in comparison to the pseudoharmonic potential.

In that case, the ground state energy levels will be

$$E_{l,1} = \frac{k^2}{2M} + \frac{\chi}{2M}, \quad (25)$$

where  $\chi$  is given in (23). One can see that the energy eigenvalue increases by the same amount as that of the potential in comparison to the pseudoharmonic potential.

### IV. Radial Solutions with Inverse Square potential Under AB-Flux Field in Topological Defect.

– In this section, we study the non-relativistic particles under the quantum flux field in a topological defect geometry in the presence of an inverse square potential  $V(r) = \frac{\gamma}{r^2}$  only.

Therefore, the radial Schrödinger equation with inverse square potential in the space-time background (4) and using (3) and (6) becomes

$$\psi'' + \frac{r}{r^2 - \beta^2} \psi' + \left[ \Theta - \frac{2M\gamma}{r^2} - \frac{\iota^2}{r^2 - \beta^2} \right] \psi = 0. \quad (26)$$

where  $\Theta = 2ME - k^2$ .

As done earlier, performing a change of variable via  $x = \frac{r^2}{\beta^2}$  in the Eq. (26), we obtain the following differential equation:

$$\begin{aligned} 4x\psi''(x) + \frac{4x-2}{x-1}\psi'(x) \\ + \left[ \Theta\beta^2 - \frac{2M\gamma}{x} - \frac{\iota^2}{x-1} \right] \psi(x) = 0. \end{aligned} \quad (27)$$

Here also, let us suppose a possible solution of the Eq. (27) given by

$$\psi(x) = x^{\frac{1}{4} + \frac{j}{2}} G(x), \quad (28)$$

where  $j$  is defined earlier and  $G(x)$  is the confluent Heun equation given here by

$$G(x) = H_c \left( 0, j, -\frac{1}{2}, \frac{\Theta\beta^2}{4}, \frac{3}{8} - \frac{\iota^2 + \Theta\beta^2}{4}; x \right). \quad (29)$$

Substituting the solution (28) and using the same power series of  $G(x)$  (12) in the Eq. (27), one can obtain the following recurrence relation

$$c_{i+2} = \frac{h_1}{h_3} c_{i+1} + \frac{h_2}{h_3} c_i \quad (30)$$

with the coefficient

$$c_1 = \frac{j + \frac{1}{2} - \Theta\beta^2 - \iota^2}{4(1+j)} c_0, \quad (31)$$

where we have set the parameters

$$\begin{aligned} h_1 &= \left( i + j + \frac{3}{2} \right) (i + 1) - \frac{\Theta\beta^2 + \iota^2 - \frac{1}{2} - j}{4}, \\ h_2 &= \frac{\Theta\beta^2}{4}, \quad h_3 = (i + 2 + j)(i + 2). \end{aligned} \quad (32)$$

Let us consider  $i = n - 1$  when  $c_{n+1} = 0$ , then we have from the relation (31)

$$c_n = -\frac{\Theta\beta^2}{(4n(n-1+j+\frac{3}{2}) - \Theta\beta^2 - \iota^2 + \frac{1}{2} + j)} c_{n-1}. \quad (33)$$

We obtain here the ground state energy level and the radial wave function for the radial mode  $n = 1$ . Setting  $n = 1$ , we obtain from (33)

$$c_1 = -\frac{\Theta\beta^2}{(5j - \Theta\beta^2 - \iota^2 + \frac{13}{2})} c_0. \quad (34)$$

Thus, by comparing Eq. (31) with Eq. (34), we obtain the ground state energy levels  $E_{l,1}$  given by

$$E_{l,1} = \frac{k^2}{2M} + \frac{\Theta}{2M}, \quad (35)$$

where

$$\Theta = \frac{(2j - 2\iota^2 + 3) \pm \sqrt{(\iota^2 + 4j\iota^2 - 4j^2 - 6j - 1)}}{2\beta^2}. \quad (36)$$

And that the ground state wave function is given by

$$\psi_{1,l}(x) = x^{\frac{1}{4} + \frac{\sqrt{2M\gamma + \frac{1}{4}}}{2}} (c_0 + c_1 x). \quad (37)$$

Equations (35)–(37) is the ground state energy level and the radial wave function of non-relativistic particles under the influence of the quantum flux in a space-time background with screw dislocation subject to an inverse square potential. Following the similar procedure, one can find other energy levels  $E_{2,l}, E_{3,l}, \dots$  and the radial wave function  $\psi_{1,l}, \psi_{2,l}, \dots$  for the radial mode  $n \geq 2$ . Thus, one can see that the eigenvalue solution depends on the screw dissociation parameter  $\beta$  and the quantum flux field.

Throughout the analysis, we see that the presence of a screw dislocation and the quantum flux field that was introduced through a minimal substitution in the wave equation changes the  $z$ -component of the angular momentum operator given by

$$\vec{L}_z^{eff} = -i \left( \frac{\partial}{\partial \phi} - i e A_\phi - \beta \frac{\partial}{\partial z} \right) \hat{z}. \quad (38)$$

Using the wave function (6), we have the shifted angular momentum quantum number  $l$  given by

$$l \rightarrow l_{eff} = \left( l - \frac{e \Phi_{AB}}{2\pi} - \beta k \right), \quad (39)$$

an effective angular momentum quantum number due to both the boundary conditions, which states that the total angle around the  $z$ -axis is less than  $2\pi$ , and the minimal coupling with the electromagnetic potential. Formula (39) suggests that when the particle circles about the  $z$ -axis, the wave-function changes according to

$$\begin{aligned} \Psi \rightarrow \Psi' &= Exp\left(2\pi i l_{eff}\right) \Psi \\ &= Exp\left[2\pi i \left(l - \frac{e \Phi_{AB}}{2\pi} - \beta k\right)\right] \Psi. \end{aligned} \quad (40)$$

Thus, one can easily show that,  $E_{1,l}(\Phi_{AB} \pm \Phi_0 \nu) = E_{1,l \mp \nu}(\Phi_{AB})$ , a periodic function of the geometric quantum phase with a periodicity  $\Phi_0$ , where  $\nu = 0, 1, 2, 3, \dots$ . This dependence of the eigenvalue solution on the quantum phase gives us an analogue to the Aharonov-Bohm effect [39, 40], quantum mechanical phenomena.

**V. Conclusions.** – In this analysis, we studied the non-relativistic particles with interaction potentials under the influence of the Aharonov-Bohm flux field in a space-time background with distortion of a vertical line into a vertical spiral. First we have chosen an anharmonic oscillator potential the superposition of harmonic oscillator plus inverse square including a constant term and derived the radial equation in the considered space-time background. We solved this equation through the Heun equation and obtained the ground state energy level  $E_{1,l}$  and the radial wave function  $\psi_{1,l}$  for the radial mode  $n = 1$  and others are in the same procedure. We have seen

that the energy level gets modified in comparison to those result obtained in [33] due to the presence of an inverse square potential and the magnetic flux field in addition to a harmonic oscillator-like potential. We then utilized this eigenvalue solution to the molecular potential model, such as pseudoharmonic potential and shifted pseudoharmonic potential and presented the ground state energy level and the wave function. Afterwards, we study the quantum system with only an inverse square potential under the influence of the same quantum flux field and following the previous procedure solved the radial wave equation. Here also, we have obtained the ground state energy levels  $E_{1,l}$  and the wave function  $\psi_{1,l}$  defined by the quantum number  $n = 1$  instead of  $n = 0$ . This effect arises due to the presence of interaction potential in the quantum system.

Throughout the analysis, we observed that the angular quantum number  $l$  is shifted or modified, that is,  $l \rightarrow l_{eff} = \left( l - \frac{e \Phi_{AB}}{2\pi} - \beta k \right)$ , a change in quantum number  $l$  that depends on the screw dislocation parameter  $\beta$  and the quantum flux field  $\Phi_{AB}$ . Thus, the energy eigenvalue depends on the geometric quantum phase and this dependence of the eigenvalue on the quantum phase gives us an analogue to the Aharonov-Bohm effect. This AB-effect is a quantum mechanical phenomenon where the particles are confined by the quantum flux field but experience no such external magnetic field. Several authors have investigated this AB-effect in the relativistic and non-relativistic quantum systems. It is worth mentioning that the geometrical approach used to describe space-time with the distortion of a vertical line into a vertical spiral can be useful in studies of condensed matter systems.

**Acknowledgement.** – P.R. acknowledges the Inter-University Centre for Astronomy and Astrophysics (IUCAA), Pune, India for granting visiting associateship.

**Data Availability Statement.** – No new data were analysed in this study.

**Conflict of Interest.** – There are no conflict of interests regarding the publication of this paper.

## REFERENCES

- [1] A. Vilenkin, E. P. S. Shellard :- Strings and Other Topological Defects (Cambridge University Press, Cambridge) (1994).
- [2] A. Vilenkin, Phys. Rep. **121**, 263 (1985).
- [3] A. Vilenkin, Phys. Lett. **B 133**, 177 (1983).
- [4] W. A. Hiscock, Phys. Rev. **D 31**, 3288 (1985).
- [5] B. Linet, Gen. Relativ. Gravit. **17**, 1109 (1985).
- [6] M. Barriola and A. Vilenkin, Phys. Rev. Lett. **63**, 341 (1989).
- [7] E. R. Bezerra de Mello and F. C. Carvalho, Class. Quantum Grav. **18**, 5455 (2001).
- [8] E. R. Bezerra de Mello and F. C. Carvalho, Class. Quantum Grav. **18**, 1637 (2001).
- [9] E. R. Bezerra de Mello, Class. Quantum Grav. **19**, 5141 (2002).



- [10] E. R. Bezerra de Mello, J. Spinelly and U. Freitas, Phys. Rev. **D 66**, 024018 (2002).
- [11] E. R. Bezerra de Mello and A. A. Saharian, Class. Quantum Grav. **29**, 135007 (2012).
- [12] T. R. P. Carames, J. C. Fabris, E. R. Bezerra de Mello and H. Belich, Eur. Phys. J. C **77**, 496 (2017).
- [13] E. R. Bezerra de Mello, Braz. J. Phys. **31**, 211 (2001).
- [14] A. L. Cavalcanti de Oliveira and E. R. Bezerra de Mello, Class. Quantum Grav. **23**, 5249 (2006).
- [15] A. Boumali and H. Aounallah, Adv. High Energy Phys. **2018**, 1031763 (2018).
- [16] H. Aounallah and A. Boumali, Phys. Part. Nucl. Lett. **16**, 195, (2019).
- [17] A. Boumali and H. Aounallah, Rev. Mex. Fis. **66**, 192 (2020).
- [18] E. A. F. Braganca, R. L. L. Vitoria, H. Belich and E. R. Bezerra de Mello, Eur. Phys. J. C **80**, 206 (2020).
- [19] M. de Montigny, J. Pinfeld, S. Zare and H. Hassanabadi, Eur. Phys. J. Plus **137**, 54 (2022).
- [20] F. Ahmed, Sci. Rep. **12**, 8794 (2022).
- [21] H. Aounallah, A. R. Soares and R. L. L. Vitoria, Eur. Phys. J. C **80**, 447 (2020).
- [22] A. R. Soares, R. L. L. Vitoria and H. Aounallah, Eur. Phys. J. Plus **136**, 966 (2021).
- [23] F. Ahmed, arXiv: 2209.13490 [quant-ph].
- [24] F. Ahmed, arXiv: 2210.04617 [hep-th].
- [25] F. Ahmed, Mol. Phys. **120**, Article: e2124935 (2022).
- [26] F. Ahmed, Phys. Scr. **98**, 015403 (2023).
- [27] V. Kumar, R. M. Singh, S. B. Bhardwaj, R. Rani and F. Chand, Mod. Phys. Lett. **A 37**, 2250010 (2022).
- [28] R. Sever, C. Tezcan, M. Aktas and O. Yesiltas, J. Math. Chem. **43**, 845 (2008).
- [29] R. Rani, S. B. Bhardwaj and F. Chand, Pramana-J. Phys. **91**, 46 (2018).
- [30] S. M. Ikhdaïr and R. Sever, J. Mol. Struct.(Theochem) **806**, 155 (2007).
- [31] A. Cetin, Phys. Lett. **A 372**, 3852 (2008).
- [32] S. M. Ikhdaïr and M. Hamzavi, Physica **B 407**, 4198 (2012).
- [33] W. C. F. da Silva, K. Bakke and R. L. L. Vitoria, Eur. Phys. J. C **79**, 657 (2019).
- [34] L. C. N. Santos, F. M. da Silva, C. E. Mota, V. B. Bezerra, Int. J. Geom. Meths. Mod. Phys. (2022), <https://doi.org/10.1142/S0219887823500676>.
- [35] R. L. L. Vitoria, Eur. Phys. J. C **79**, 844 (2019).
- [36] K. Valanis and V. Panoskaltisis, Acta mechanica **175**, 77 (2005).
- [37] K. Bakke, Eur. Phys. J. Plus **134**, 546 (2019).
- [38] W. C. F. da Silva and K. Bakke, Class. Quantum Grav. **36**, 235002 (2019).
- [39] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [40] M. Peshkin and A. Tonomura, *The Aharonov-Bohm Effect*, Lecture Notes Phys. **Vol. 340**, Springer-Verlag, Berlin, Germany (1989).