

GENERATORS OF THE MAPPING CLASS GROUP OF A NONORIENTABLE PUNCTURED SURFACE

TÜLİN ALTUNÖZ, MEHMETÇİK PAMUK, AND OĞUZ YILDIZ

ABSTRACT. Let $\text{Mod}(N_{g,p})$ denote the mapping class group of a nonorientable surface of genus g with p punctures. For $g \geq 14$, we show that $\text{Mod}(N_{g,p})$ can be generated by five elements or by six involutions.

1. INTRODUCTION

Let $N_{g,p}$ denote a nonorientable surface of genus g with p punctures specified by the set $P = \{z_1, z_2, \dots, z_p\}$ of p distinguished points. If p is zero, we omit it from the notation and denote the surface by N_g . The mapping class group $\text{Mod}(N_{g,p})$ of the surface $N_{g,p}$ is defined to be the group of the isotopy classes of self-diffeomorphisms of $N_{g,p}$ which preserve the set P . In this paper, we consider only connected surfaces and we are interested in finding minimal generating sets for $\text{Mod}(N_{g,p})$.

For an orientable surface, the mapping class group is defined as the group of isotopy classes of orientation-preserving diffeomorphisms. For orientable surfaces, it is known that the mapping class group is generated by Dehn twists and elementary braids. However, if the surface is nonorientable, Dehn twists do not generate the mapping class group. Lickorish was the first to consider the problem of finding generators of the group $\text{Mod}(N_g)$. He proved that $\text{Mod}(N_g)$ is generated by Dehn twists and the isotopy class of a homeomorphism he called the Y -homeomorphism [3]. Chillingworth [2] determined a finite set of generators for $\text{Mod}(N_g)$. The cardinality of this set depends on g . Korkmaz [4] extended Chillingworth's result to the case of punctured surfaces.

Since $\text{Mod}(N_g)$ is not abelian, a generating set must contain at least two elements. Szepietowski [6] proved that this group is generated by three elements for all $g \geq 3$. We [1] prove that, for $g \geq 19$ the mapping class group of a nonorientable surface of genus g , $\text{Mod}(N_g)$, can be generated by two elements, one of which is of order g .

In the presence of punctures, Szepietowski [5, Theorem 3] proved that $\text{Mod}(N_{g,p})$ is generated by involutions. We prove that for $g \geq 26$, $\text{Mod}(N_g)$ can be generated by three involutions if $g \geq 26$ [1]. Recently, Yoshihara [7] proved that, for $p \geq 0$, $\text{Mod}(N_{g,p})$ is generated by eight involutions if $g \geq 13$ is odd and by eleven involutions if $g \geq 14$ is even.

In this paper, we obtain the following main result (see Theorems 3.5 and 3.6):

Main Theorem. *For $g \geq 14$ and $p \geq 1$, $\text{Mod}(N_{g,p})$ is generated by five elements or six involutions.*

The paper is organized as follows: In Section 2, we present the necessary background and some results on mapping class groups. The proof of our main theorem is given in Section 3.

Acknowledgements. This work is supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK)[grant number 120F118].

2. BACKGROUND AND RESULTS ON MAPPING CLASS GROUPS

Let $\text{Mod}_0(N_{g,p})$ denote the subgroup of $\text{Mod}(N_{g,p})$ consisting of elements which fix the set P pointwise. This group is a normal subgroup of $\text{Mod}(N_{g,p})$ of index $p!$ and we have the following exact sequence:

$$1 \longrightarrow \text{Mod}_0(N_{g,p}) \longrightarrow \text{Mod}(N_{g,p}) \longrightarrow \text{Sym}_p \longrightarrow 1,$$

2000 *Mathematics Subject Classification*: 57M07, 20F05, 20F38

Keywords: Mapping class groups, punctured surfaces, involutions, generating sets

where Sym_p is the symmetric group on the set $\{1, 2, \dots, p\}$ and the last projection is given by the restriction of the isotopy class of a diffeomorphism to its action on the punctures.

Throughout the paper we do not distinguish a diffeomorphism from its isotopy class. For the composition of two diffeomorphisms, we use the functional notation; if f and g are two diffeomorphisms, then the composition fg means that g is applied first and then f .

For a two-sided simple closed curve a on $N_{g,p}$, we denote the right-handed Dehn twist t_a about a by the corresponding capital letter A .

Now, let us remind the following basic facts of Dehn twists that we use frequently in the remaining of the paper. Let a and b be two-sided simple closed curves on $N_{g,p}$ and $f \in \text{Mod}(N_{g,p})$.

- If a and b are disjoint, then $AB = BA$ (*Commutativity*).
- If $f(a) = b$, then $fAf^{-1} = B^\varepsilon$, where $\varepsilon = \pm 1$ depends on the orientation of a regular neighbourhood of $f(a)$ with respect to the chosen orientation. (*Conjugation*).

Throughout the paper we use repeatedly the conjugation relation and we denote fgf^{-1} by g^f for any $f, g \in \text{Mod}(N_{g,p})$.

We now define two types of diffeomorphisms of $\text{Mod}(N_{g,p})$; y homeomorphism (crosscap slide) or puncture slide. Let us consider the Klein bottle K depicted in Figure 1. A *crosscap transposition* u is defined to be the isotopy




FIGURE 1. The homeomorphisms u and $y = Au$.

classes of a diffeomorphism that interchanges two consecutive crosscaps and keeps the boundary of K fixed as shown on the left hand side of Figure 1. The diffeomorphism $y = Au$ whose effect on the interval c as in Figure 1 can be also obtained as sliding a Möbius band once along the core of another one and fixing each point of the boundary of K . This is called a *Y-homeomorphism* [3] (also called a *crosscap slide* [4]). Let us note that $A^{-1}u$ is a *Y-homeomorphism* i.e. the other choice of the orientation for a neighbourhood of the curve a also gives a *Y-homeomorphism*. Also note that y^2 is a Dehn twist about ∂K .

Lickorish [3] proved that $\text{Mod}(N_g)$ is generated by Dehn twists and a *Y* homeomorphism (one crosscap slide). Note that crosscap transpositions can be used instead of crosscap slides in generating sets of $\text{Mod}(N_g)$ since a crosscap transposition is equal to the product of a Dehn twist and a crosscap slide.

For the other type of diffeomorphisms of $\text{Mod}(N_{g,p})$, consider the Möbius band M with a puncture z and the one-sided simple closed curve a in Figure 2. We denote v_a the *puncture slide* obtained by sliding the puncture z




FIGURE 2. The puncture slide v_a .

once along a keeping each point of the boundary M fixed. The effect of the diffeomorphism v_a on the interval c is shown as Figure 2. Let us denote the puncture slide supported by the one-sided simple closed curve δ_i and the puncture z_j by $v_{i,j}$ as shown in Figure 2.




FIGURE 3. The puncture slide $v_{i,j}$.

Before we finish Preliminaries, let us give Korkmaz's generating set for $\text{Mod}_0(N_{g,p})$.

Let \mathcal{C} be the set of simple closed curves given by

$$\mathcal{C} = \{a_i, b_i, f_i, c_j, e_k, \text{ for } i = 1, \dots, r, \quad j = 1, \dots, r-1 \text{ and } k = 1, \dots, p-1 \text{ if } g = 2r+1,$$

and

$$\mathcal{C} = \{a_i, b_i, f_i, c_i, e_k, \text{ for } i = 1, \dots, r \text{ and } k = 1, \dots, p-1 \text{ if } g = 2r+2,$$

where the curves are shown in Figure 4.




FIGURE 4. The Dehn twist generators for $\text{Mod}(N_{g,p})$.

Theorem 2.1. [4] *For $g \geq 3$, the mapping class group $\text{Mod}_0(N_{g,p})$ is generated by*

- (i) $\{t_c, y, v_{g,i} : c \in \mathcal{C}, 1 \leq i \leq p\}$ if g is odd, and
- (ii) $\{t_c, y, v_{g-1,i}, v_{g,i} : c \in \mathcal{C}, 1 \leq i \leq p\}$ if g is even.

3. A GENERATING SET FOR $\text{Mod}(N_{g,p})$

We, first recall the following basic lemma from algebra.

Lemma 3.1. *Let G and K be groups. Suppose that the following short exact sequence holds,*

$$1 \longrightarrow N \xrightarrow{i} G \xrightarrow{\pi} K \longrightarrow 1.$$

Then the subgroup Γ contains $i(N)$ and has a surjection to K if and only if $\Gamma = G$.

For $G = \text{Mod}(N_{g,p})$ and $N = \text{Mod}_0(N_{g,p})$, the following short exact sequence holds:

$$1 \longrightarrow \text{Mod}_0(N_{g,p}) \longrightarrow \text{Mod}(N_{g,p}) \longrightarrow \text{Sym}_p \longrightarrow 1.$$

Therefore, the following useful result follows immediately from Lemma 3.1. Let Γ be a subgroup of $\text{Mod}(N_{g,p})$. If the subgroup Γ contains $\text{Mod}_0(N_{g,p})$ and has a surjection to Sym_p then $\Gamma = \text{Mod}(N_{g,p})$.




FIGURE 5. The reflections ρ_1 and ρ_2 on $N_{g,p}$ if $g = 2r + 2$.




FIGURE 6. The reflections ρ_1 and ρ_2 on $N_{g,p}$ if $g = 2r + 1$.

Some Dehn twists curves given in Theorem 2.1 are shown in Figure 9 when we use the model for the surface $N_{g,p}$ as a sphere with p punctures and g crosscaps.

Consider the models for $N_{g,p}$ such that it is invariant under the reflections ρ_1 and ρ_2 (see Figures 5 and 6). Note that $\text{Mod}(N_{g,p})$ contains the element $T = \rho_2\rho_1$ which satisfies the following:

- $T(b_i) = c_i$ and $T(c_j) = b_{j+1}$ for $i, j = 1, \dots, r - 1$,




FIGURE 7. The reflections ρ_3 and ρ_4 on $N_{g,p}$ if $p = 2b$ (when g is odd we remove the last crosscap).




FIGURE 8. The reflections ρ_3 and ρ_4 on $N_{g,p}$ if $p = 2b + 1$ (when g is odd we remove the last crosscap).

- $T(a_1) = b_1$. Also, note that $T^2(b_r) = a_1$ if $g = 2r + 1$ and $T(b_r) = c_r$, $T^2(c_r) = a_1$ if $g = 2r + 2$,
- $T(z_1) = z_2$, $T(z_2) = z_1$ and $T(z_k) = z_k$ for $k = 3, 4, \dots, p$,
- $T(\alpha_l) = \alpha_{l+1}$ for $l = 1, 2, \dots, g - 1$ and $T(\alpha_g) = \alpha_1$ where all curves are depicted in Figure 9.

We will use two different diffeomorphism of $\text{Mod}(N_{g,p})$, the reflections ρ_3 and ρ_4 depicted in Figures 7 and 8, in the remaining part of the paper.

Lemma 3.2. *For any $g = 2r + 1 \geq 15$ or $g = 2r + 2 \geq 14$, the group generated by the elements*

$$\{T, \rho_3, \rho_4, \Gamma_5 B_1 u_{r+5}, v_{r+3,p}\}$$

contains the Dehn twists A_i , B_i , F_i and C_j for $i = 1, \dots, r$ and $j = 1, \dots, r - 1$ (also the element C_r when g is even).

Proof. Let $G_1 := \Gamma_5 B_1 u_{r+5}$ and let G be the subgroup of $\text{Mod}(N_{g,p})$ generated by the elements T , ρ_3 , ρ_4 , G_1 and $v_{r+3,p}$.




FIGURE 9. The curves $a_i, b_i, c_i, \gamma_i, f_i$ and α_i (note that we do not have c_r when g is odd).

Let G_2 be the element obtained by the conjugation of G_1 by T^2 . It follows from

$$T^2(\gamma_5, b_1, \alpha_{r+5}) = (\gamma_7, b_2, \alpha_{r+7})$$

that

$$G_2 = G_1^{T^2} = \Gamma_7 B_2 u_{r+7} \in G.$$

Let G_3 denote the element $G_1^{G_1 G_2}$, which is contained in the subgroup G . Therefore,

$$G_3 = B_2 B_1 u_{r+5}$$

Let us give more details of this calculation since we use similar calculations in the rest of the proof. It can be verified that the diffeomorphism $G_1 G_2$ sends the curves $\{\gamma_5, b_1, \alpha_{r+5}\}$ to the curves $\{b_2, b_1, \alpha_{r+5}\}$, respectively. Then we get the element

$$\begin{aligned} G_3 &= G_1^{G_1 G_2} \\ &= (G_1 G_2)(\Gamma_5 B_1 u_{r+5})(G_1 G_2)^{-1} \\ &= B_2 B_1 u_{r+5}. \end{aligned}$$

Thus the subgroup G contains the element

$$G_1 G_3^{-1} = \Gamma_5 B_2^{-1}.$$

This implies that the element

$$B_3 \Gamma_7^{-1} = (B_2 \Gamma_5^{-1})^{T^2}$$

is also contained in G . Using these elements, we also obtain

$$\begin{aligned} G_4 &= (B_3 \Gamma_7^{-1}) G_2 = B_3 B_2 u_{r+7} \in G, \\ G_5 &= G_4^{T^{-3}} = C_1 A_1 u_{r+4} \in G, \\ G_6 &= G_4^{G_4 G_5} = B_3 C_1 u_{r+7} \in G. \end{aligned}$$

From these, we get the element $G_4G_6^{-1} = B_2C_1^{-1}$, which is in G . Hence, we have the following elements:

$$\begin{aligned} C_1B_1^{-1} &= (B_2C_1^{-1})^{T^{-1}}, \\ B_1B_2^{-1} &= (B_1C_1^{-1})(C_1B_2^{-1}), \\ C_1C_2^{-1} &= (B_1B_2^{-1})^T, \\ B_2C_2^{-1} &= (B_1C_1^{-1})^{T^2}, \\ \Gamma_5C_2^{-1} &= (\Gamma_5B_2^{-1})(B_2C_2^{-1}) \text{ and} \\ A_2A_1^{-1} &= \Gamma_1A_1^{-1} = (\Gamma_5C_2^{-1})^{T^{-4}}, \end{aligned}$$

which are all contained in G . Furthermore, the subgroup G contains the following elements:

$$\begin{aligned} A_1B_1^{-1} &= (C_1B_2^{-1})^{T^{-2}}, \\ A_1B_2^{-1} &= (A_1B_1^{-1})(B_1B_2^{-1}), \\ A_1C_1^{-1} &= (A_1B_1^{-1})(B_1B_2^{-1})(B_2C_1^{-1}), \\ C_1A_2^{-1} &= (C_1A_1^{-1})(A_1A_2^{-1}), \\ C_2A_1^{-1} &= (C_2C_1^{-1})(C_1A_2^{-1})(A_2A_1^{-1}), \\ C_2B_1^{-1} &= (C_2A_1^{-1})(A_1B_1^{-1}) \text{ and} \\ B_3B_1^{-1} &= (B_2C_1^{-1})^{T^2}(C_2B_1^{-1}). \end{aligned}$$

Using the lantern relation given in the proof of [1, Theorem 2.1], it can be verified that the subgroup G contains




FIGURE 10. The curves $d_{1,i}$ and $d_{2,i}$.

the elements A_1, A_2, A_3, B_i and C_i for $1 \leq i \leq r-1$ and B_r (also the element C_r when g is even).

We now inductively show that $A_j \in G$ for $j = 4, \dots, r$. To do this, we first use the diffeomorphism

$$(A_iB_{i+1}^{-1})(A_iC_i^{-1})(A_iC_{i+1}^{-1})(A_iB_{i+1}^{-1}),$$

which sends the curves (a_{i+1}, a_i) to $(d_{2,i}, a_i)$, where the curves are shown in Figure 10 for $i+2 \leq r$. This gives rise to the following identity:

$$((A_iB_{i+1}^{-1})(A_iC_i^{-1})(A_iC_{i+1}^{-1})(A_iB_{i+1}^{-1}))(A_{i+1}A_i^{-1})((A_iB_{i+1}^{-1})(A_iC_i^{-1})(A_iC_{i+1}^{-1})(A_iB_{i+1}^{-1}))^{-1} = D_{2,i}A_i^{-1}.$$

Thus, we have $D_{2,i}C_{i+1}^{-1} = (D_{2,i}A_i^{-1})(A_iC_{i+1}^{-1})$. Similarly, one can show that

$$(C_{i+1}B_i^{-1})(C_{i+1}A_i^{-1})(C_{i+1}C_i^{-1})(C_{i+1}B_i^{-1})(d_{2,i}, c_{i+1}) = (d_{1,i}, c_{i+1}),$$

which implies that

$$((C_{i+1}B_i^{-1})(C_{i+1}A_i^{-1})(C_{i+1}C_i^{-1})(C_{i+1}B_i^{-1}))(D_{2,i}C_{i+1}^{-1})((C_{i+1}B_i^{-1})(C_{i+1}A_i^{-1})(C_{i+1}C_i^{-1})(C_{i+1}B_i^{-1}))^{-1} = D_{1,i}C_{i+1}^{-1}.$$

We also have $D_{1,i}A_i^{-1} = (D_{1,i}C_{i+1}^{-1})(C_{i+1}A_i^{-1})$. By the lantern relation given in the proof of [1, Theorem 2.1], we get

$$A_{i+2} = (A_{i+1}C_i^{-1})(D_{1,i}C_{i+1}^{-1})(D_{2,i}A_i^{-1}).$$

Therefore, since A_1, A_2 and A_3 belong to G , we inductively obtain that all $A_j \in G$ for $j = 1, \dots, r$.

On the other hand, since $\rho_3(a_1) = f_1$ and $\rho_4(a_i) = f_i$ for $i = 2, \dots, r$,

$$F_1 = (A_1^{-1})^{\rho_3} \in G \text{ and } F_i = (A_i^{-1})^{\rho_4} \in G \text{ for } i = 2, \dots, r.$$

This completes the proof. \square

Let G be the subgroup of $\text{Mod}(N_{g,p})$ generated by the elements given explicitly in Lemma 3.2 with the given conditions.

Lemma 3.3. *The subgroup G contains the elements y and $v_{g,i}$ (and also $v_{g-1,i}$ if g is even) for $i = 1, 2, \dots, p$.*

Proof. Let us first prove that G contains a y -homeomorphism. By Lemma 3.2, the crosscap transposition

$$u_{r+5} = (B_1^{-1})(A_2^{-1})^{T^4}(\Gamma_5 B_1 u_{r+5}) \in G$$

since each factors on the right hand side are contained in G . Using the action of T , we get $u_i \in G$ for $i = 1, \dots, g-1$. In particular, $u_1 \in G$ and it is easy to see that

$$y = A_1 u_1 \in G.$$

In order to show that $v_{g,i} \in G$ for each $i = 1, 2, \dots, p$, observe that

$$v_{r+3,p}^{T^k} = v_{r+3+k,p} \in G \text{ for } k = 1, \dots, g-r-3.$$

In particular, $v_{g,p} \in G$. Using the element $\rho_3\rho_4$, we have the following elements:

$$v_{g,p}^{\rho_3\rho_4} = v_{g,1}, \quad v_{g,1}^{\rho_3\rho_4} = v_{g,2}, \quad \dots, \quad v_{g,p-2}^{\rho_3\rho_4} = v_{g,p-1},$$

which are all contained in G .

In the case of even g , similarly, using the elements $v_{g-1,p}$ and $\rho_3\rho_4$ that are contained in G , we get the elements

$$v_{g-1,p}^{\rho_3\rho_4} = v_{g-1,1} \in G, \quad v_{g-1,1}^{\rho_3\rho_4} = v_{g-1,2} \in G, \quad \dots, \quad v_{g-1,p-2}^{\rho_3\rho_4} = v_{g-1,p-1} \in G,$$

which completes the proof. \square

Lemma 3.4. *The group $\text{Mod}_0(N_{g,p})$ is contained in the group G .*

Proof. The group G contains the Dehn twist elements A_i, B_i, F_i and C_j for $i = 1, \dots, r$ and $j = 1, \dots, r-1$ (also C_r when g is even) by Lemma 3.2 and elements y and $v_{g,k}$ (and also $v_{g-1,k}$ if g is even) for $k = 1, 2, \dots, p$ by Lemma 3.3. It follows from Theorem 2.1 that it is enough to prove that $E_l \in G$ for $l = 1, 2, \dots, p-1$. Since the reflection ρ_4 maps a_1 to e_{p-1} , we have

$$(A_1^{-1})^{\rho_4} = E_{p-1} \in G.$$

The reflection ρ_3 maps the curve e_{p-1} to e_1 . Then we have

$$(E_{p-1}^{-1})^{\rho_3} = E_1 \in G.$$

Similarly, using the reflection ρ_4 , we get

$$(E_1^{-1})^{\rho_4} = E_{p-2} \in G.$$

Continuing in this way, we conclude that the elements E_1, E_2, \dots, E_{p-1} belong to G , which completes the proof. \square

Now, it is time to prove the main theorems.

Theorem 3.5. *The subgroup G is equal to the mapping class group $\text{Mod}(N_{g,p})$.*

Proof. By Lemma 3.4, the group $\text{Mod}_0(N_{g,p})$ is contained in the group G . Hence, by Lemma 3.1, we need to prove that G is mapped surjectively onto Sym_p . The elements $\rho_3\rho_4 \in G$ and $\rho_1 \in G$ have the images $(1, 2, \dots, p) \in \text{Sym}_p$ and $(1, 2) \in \text{Sym}_p$, respectively. This finishes the proof since these elements generate the whole group Sym_p . \square

Consider the surface $N_{g,p}$ of genus $g \geq 14$. Since

$$\rho_3(\gamma_5) = \gamma_5, \quad \rho_3(b_1) = b_1, \quad \text{and} \quad \rho_3(\alpha_{r+5}) = \alpha_{r+5},$$

with reverse orientation, the element $\rho_3\Gamma_5B_1u_{r+5}$ is an involution. Similarly, if $g = 2r + 1$ or $2r + 2$, it follows from $v_{p,r+3}^{\rho_1} = v_{p,r+3}^{-1}$ that the element $\rho_1v_{p,r+3}$ is an involution.

Theorem 3.6. *For any $g = 2r + 1 \geq 15$ or $g = 2r + 2 \geq 14$, the mapping class group $\text{Mod}(N_{g,p})$ is generated by the involutions*

$$\{\rho_1, \rho_2, \rho_3, \rho_4, \rho_3\Gamma_5B_1u_{r+5}, \rho_1v_{p,r+3}\}$$

Proof. The proof follows from Theorem 3.5 and by the fact that $T = \rho_2\rho_1$. □

REFERENCES

- [1] T. Altunöz, M. Pamuk and O. Yildiz: *Generating the mapping class group of a nonorientable surface by two elements or by three involutions*, Bull Braz Math Soc, New Series **53**, (2022), 1145–1156.
- [2] D. R. J. Chillingworth: *A finite set of generators for the homeotopy group of a non-orientable surface*, Proc. Cambridge Philos. Soc. **65**, (2) (1969), 409–430.
- [3] W. B. R. Lickorish: *Homeomorphisms of non-orientable two-manifolds*, Proc. Camb. Philos. Soc. **59**, (1963), 307–317.
- [4] M. Korkmaz: *Mapping class groups of nonorientable surfaces*, Geometriae Dedicata **89**(1), (2002), 107–131.
- [5] B. Szepietowski: *Involutions in mapping class groups of non-orientable surfaces*, Collect. Math. **55**, (3) (2004), 253–260.
- [6] B. Szepietowski: *The mapping class group of a nonorientable surface is generated by three elements and by four involutions*, Geom. Dedicata **117**, (1) (2006), 1–9.
- [7] K. Yoshihara: *Generating the mapping class group of a non-orientable punctured surface by involutions*, ArXiv math.GT/2212.09924, v1 20 Dec 2022.

FACULTY OF ENGINEERING, BAŞKENT UNIVERSITY, ANKARA, TURKEY

Email address: tulinaltunoz@baskent.edu.tr

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, ANKARA, TURKEY

Email address: mpamuk@metu.edu.tr

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, ANKARA, TURKEY

Email address: oguzyildiz16@gmail.com