

RAMANUJAN-TYPE CONGRUENCES FOR PARTITION k -TUPLES WITH 5-CORES

MANJIL P. SAIKIA, ABHISHEK SARMA, AND PRANJAL TALUKDAR

ABSTRACT. We prove several Ramanujan-type congruences modulo powers of 5 for partition k -tuples with 5-cores, for $k = 2, 3, 4$. We also prove some new infinite families of congruences modulo powers of primes for k -tuples with p -cores, where p is a prime.

1. INTRODUCTION

A partition of a positive integer n is a finite non-increasing sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ such that $\sum_{i=1}^k \lambda_i = n$. The number of partitions of n is denoted by $p(n)$. Euler found the generating function of $p(n)$, given by

$$\sum_{n \geq 0} p(n)q^n = \frac{1}{(q; q)_{\infty}},$$

where

$$(a; q)_{\infty} := \prod_{i \geq 0} (1 - aq^i), \quad |q| < 1.$$

Thereafter, the subject got a massive push when Ramanujan found some amazing congruences that the partition function $p(n)$ satisfies; this ushered in an era of study of arithmetic properties of the partition function as well as other restricted types of partitions. Ramanujan proved the following identities

$$p(5n + 4) \equiv 0 \pmod{5}, \quad p(7n + 5) \equiv 0 \pmod{7}, \quad \text{and} \quad p(11n + 6) \equiv 0 \pmod{11}.$$

Over the years various authors have studied different classes of partitions in the hope of proving such Ramanujan-type congruences for newer partition types. In this paper we look at the class of partitions called k -tuple partitions with t -cores and prove some Ramanujan-type congruences that these partitions satisfy.

The Ferrers-Young diagram of a partition is a pattern of dots with λ_i dots in the i -th row for the partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ of n . The hook number of a dot is the number of dots directly below and to the right of the dot, including the dot itself. A partition is called a t -core partition for $t \geq 2$ if the partition has no hook number divisible by t . We further define a partition k -tuple of n to be the k -tuple of partitions $(\Lambda_1, \Lambda_2, \dots, \Lambda_k)$, where $\sum_{i=1}^k \Lambda_i = n$. Let $A_{t,k}(n)$ be the number of partition k -tuples of n with t -cores and its generating function is given by

$$(1.1) \quad \sum_{n \geq 0} A_{t,k}(n)q^n = \frac{(q^t; q^t)_{\infty}^{kt}}{(q; q)_{\infty}^k}.$$

These partitions and the generating function are the objects of study in this paper.

There have been several studies involving the congruence properties of $A_{t,k}(n)$. Dasappa [Ran16] proved the following result

$$(1.2) \quad A_{5,2}(5^{\alpha}n + 5^{\alpha} - 2) \equiv 0 \pmod{5^{\alpha}}, \quad \alpha \geq 1.$$

2020 *Mathematics Subject Classification.* 11P81, 11P83.

Key words and phrases. integer partitions, Ramanujan-type congruences.

In a similar vein, recently Majid and Fathima [MF22] proved the following result

$$(1.3) \quad A_{5,3}(5^\alpha n + 5^\alpha - 3) \equiv 0 \pmod{5^\alpha}, \quad \alpha \geq 1.$$

Both of these results were proved using elementary techniques which involved using dissection formulas and induction. We extend these results in the following theorem.

Theorem 1.1. *For all $n \geq 0$ and $\alpha \geq 1$, we have*

$$(1.4) \quad A_{5,4}(5^{\alpha+1}n + 5^{\alpha+1} - 4) \equiv 0 \pmod{5^{\alpha+4}}.$$

We further find some new infinite family of congruences for $A_{t,k}(n)$ for some general values of k and t , as stated in the following results.

Theorem 1.2. *Let $p \geq 5$ be a prime and let $r \in \mathbb{N}$ with $1 \leq r \leq p-1$, be such that $24r+1$ is a quadratic nonresidue modulo p . Then, for all $n \geq 0$, $i \geq 1$ and $N \geq 1$, we have*

$$A_{p,p^{N_{i-1}}}(pn+r) \equiv 0 \pmod{p^N}.$$

Theorem 1.3. *Let $p \geq 5$ be a prime and let $r \in \mathbb{N}$ with $1 \leq r \leq p-1$, be such that $8r+1$ is a quadratic nonresidue modulo p . Then, for all $n \geq 0$, $i \geq 1$ and $N \geq 1$, we have*

$$A_{p,p^{N_{i-3}}}(pn+r) \equiv 0 \pmod{p^N}.$$

Theorem 1.4. *Let $p \geq 5$ be a prime and let $r \in \mathbb{N}$ with $1 \leq r \leq p-1$, be the unique value such that $8r+1 \equiv 0 \pmod{p}$. Then, for all $n \geq 0$ and $i \geq 1$, we have*

$$A_{p,pi-3}(pn+r) \equiv 0 \pmod{p}.$$

We also prove some new individual congruences for $A_{5,t}(n)$ for some specific values of t .

Theorem 1.5. *For all $n \geq 0$, the following congruences are true*

$$(1.5) \quad A_{5,2}(25n+23) \equiv 0 \pmod{25},$$

$$(1.6) \quad A_{5,2}(125n+123) \equiv 0 \pmod{125},$$

$$(1.7) \quad A_{5,3}(25n+22) \equiv 0 \pmod{5},$$

$$(1.8) \quad A_{5,3}(125n+122) \equiv 0 \pmod{25},$$

$$(1.9) \quad A_{5,4}(25n+21) \equiv 0 \pmod{3125},$$

$$(1.10) \quad A_{5,4}(125n+121) \equiv 0 \pmod{15625}.$$

Next, we present a congruence result for $A_{t,k}(n)$ modulo powers of primes, which can be also viewed as an existence result for infinite family of congruences.

Theorem 1.6. *Let p be a prime, $k \geq 1$, $j \geq 0$, $N \geq 1$, $M \geq 1$, and r be integers such that $1 \leq r \leq p^M - 1$. If for all $n \geq 0$,*

$$A_{p,k}(p^M n + r) \equiv 0 \pmod{p^N},$$

then for all $n \geq 0$ we have

$$A_{p,p^{M+N-1}i+k}(p^M n + r) \equiv 0 \pmod{p^N}.$$

The following is an easy corollary.

Corollary 1.7. *For all $i \geq 0$ and $n \geq 0$, we have*

$$\begin{aligned}
(1.11) \quad & A_{5,25i+2}(25n+23) \equiv 0 \pmod{5}, \\
(1.12) \quad & A_{5,125i+2}(25n+23) \equiv 0 \pmod{25}, \\
(1.13) \quad & A_{5,3125i+2}(125n+123) \equiv 0 \pmod{125}, \\
(1.14) \quad & A_{5,25i+3}(25n+22) \equiv 0 \pmod{5}, \\
(1.15) \quad & A_{5,625i+3}(125n+122) \equiv 0 \pmod{25}, \\
(1.16) \quad & A_{5,5i+4}(5n+3,4) \equiv 0 \pmod{5}, \\
(1.17) \quad & A_{5,125i+4}(25n+21) \equiv 0 \pmod{25}, \\
(1.18) \quad & A_{5,25i+4}(25n+21) \equiv 0 \pmod{5}, \\
(1.19) \quad & A_{5,125i+4}(25n+21) \equiv 0 \pmod{25}, \\
(1.20) \quad & A_{5,625i+4}(25n+21) \equiv 0 \pmod{125}, \\
(1.21) \quad & A_{5,3125i+4}(25n+21) \equiv 0 \pmod{625}, \\
(1.22) \quad & A_{5,15625i+4}(25n+21) \equiv 0 \pmod{3125}, \\
(1.23) \quad & A_{5,390625i+4}(125n+121) \equiv 0 \pmod{15625}.
\end{aligned}$$

Proof. The proofs of the above congruences follow from Theorem 1.5 and Theorem 1.6 and are similar in nature. Hence, here we only present the proof of (1.17).

The case for $i = 0$ is true by (1.9). Using Theorem 1.6, and the case for $i = 0$, we deduce that

$$\sum_{n \geq 0} A_{5,125i+4}(25n+21)q^n \equiv \sum_{n \geq 0} A_{5,4}(25n+21)q^n \equiv 0 \pmod{25},$$

which completes the proof. \square

The rest of the paper is organized as follows: in Section 2 we state some preliminary results that we require for our proofs, Theorem 1.1 is then proved in Section 3, Theorems 1.2, 1.3 and 1.4 are proved in Section 4, Theorem 1.5 is proved in Section 5, Theorem 1.6 is proved in Section 6 and finally we close the paper with some concluding remarks and conjectures in Section 7.

2. PRELIMINARY RESULTS

For the sake of brevity, we use the notation $f_k := (q^k; q^k)_\infty$ throughout the rest of the paper. We also let

$$R(q) = \frac{(q; q^5)_\infty (q^4; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty}.$$

First, we recall the following 5-dissections of $\frac{1}{f_1}$ and f_1 . We have [Ber06, Eq. (7.4.14)]

$$\begin{aligned}
(2.1) \quad \frac{1}{f_1} = \frac{f_{25}^5}{f_5^6} & \left(\frac{1}{R^4(q^5)} + \frac{q}{R^3(q^5)} + \frac{2q^2}{R^2(q^5)} + \frac{3q^3}{R(q^5)} + 5q^4 - 3q^5 R(q^5) + 2q^6 R^2(q^5) \right. \\
& \left. - q^7 R^3(q^5) + q^8 R^4(q^5) \right),
\end{aligned}$$

and [Ber06, Theorem 7.4.4]

$$(2.2) \quad f_1 = f_{25} \left(\frac{1}{R(q^5)} - q - q^2 R(q^5) \right).$$

Lemma 2.1. *Let $\sum_{n \geq 0} P_4(n)q^n = \frac{1}{f_1^4}$. Then we have*

$$\sum_{n \geq 0} P_4(5n+1)q^n = 4 \frac{f_5^2}{f_1^6} + 550q \frac{f_5^8}{f_1^{12}} + 12500q^2 \frac{f_5^{14}}{f_1^{18}} + 78125q^3 \frac{f_5^{20}}{f_1^{24}}.$$

Proof. Using equation (2.1), extracting the terms involving q^{5n+1} , dividing by q and then replacing q^5 by q , we arrive at

(2.3)

$$\sum_{n \geq 0} P_4(5n+1)q^n = \frac{f_5^{20}}{f_1^{24}} \left(\frac{4}{R^{15}(q)} + \frac{418q}{R^{10}(q)} + \frac{1840q^2}{R^5(q)} + 1015q^3 - 1840q^4 R^5(q) + 418q^5 R^{10}(q) - 4q^6 R^{15}(q) \right).$$

We use the following formula [Ber06, Theorem 7.4.4]

(2.4)

$$\frac{1}{R(q)^5} - 11q - q^2 R(q)^5 = \frac{f_1^6}{f_5^6},$$

to obtain from equation (2.3)

(2.5)

$$\sum_{n \geq 0} P_4(5n+1)q^n = 4 \frac{f_5^2}{f_1^6} + 550q \frac{f_5^8}{f_1^{12}} + 12500q^2 \frac{f_5^{14}}{f_1^{18}} + 78125q^3 \frac{f_5^{20}}{f_1^{24}}.$$

□

Lemma 2.2. *If $\sum_{n \geq 0} Q_4(n)q^n = f_5^2 f_1^{14}$, then we have $\sum_{n \geq 0} Q_4(5n+4)q^n = -15625q^2 f_5^{14} f_1^2$.*

Proof. Using equation (2.2), we obtain

$$\begin{aligned} \sum_{n \geq 0} Q_4(n)q^n &= f_5^2 f_1^{14} (1/R^{14}(q) - 14q/R^{13}(q) + 77q^2/R^{12}(q) - 182q^3/R^{11}(q) + 910q^5/R^9(q) \\ &\quad - 1365q^6/R^8(q) - 1430q^7/R^7(q) + 5005q^8/R^6(q) - 10010q^{10}/R^4(q) \\ &\quad + 3640q^{11}/R^3(q) + 14105q^{12}/R^2(q) - 6930q^{13}/R(q) - 15625q^{14} \\ &\quad + 6930q^{15}R(q) + 14105q^{16}R^2(q) - 3640q^{17}R^3(q) - 10010q^{18}R^4(q) \\ &\quad + 5005q^{20}R^6(q) + 1430q^{21}R^7(q) - 1365q^{22}R^8(q) - 910q^{23}R^9(q) \\ &\quad + 182q^{25}R^{11}(q) + 77q^{26}R^{12}(q) + 14q^{27}R^{13}(q) + q^{28}R^{14}(q)). \end{aligned}$$

Extracting the terms involving q^{5n+4} and then dividing by q^4 and replacing q^5 by q , we get

$$\sum_{n \geq 0} Q_4(5n+4)q^n = -15625q^2 f_5^{14} f_1^2.$$

□

Lemma 2.3. *If $\sum_{n \geq 0} Q_5(n)q^n = q f_5^8 f_1^8$, then we have $\sum_{n \geq 0} Q_5(5n+4)q^n = -125q f_5^8 f_1^8$.*

Lemma 2.4. *If $\sum_{n \geq 0} Q_6(n)q^n = q^2 f_5^{14} f_1^2$, then we have $\sum_{n \geq 0} Q_6(5n+4)q^n = -f_5^2 f_1^{14}$.*

The proofs of Lemma 2.3 and Lemma 2.4 are exactly similar to the proof of Lemma 2.2. So, we leave them to the reader.

We also recall that

(2.6)

$$f_1 = \sum_{m=-\infty}^{\infty} (-1)^m q^{m(3m-1)/2},$$

and [Ber91, p.39, Entry 24(iii)]

(2.7)

$$f_1^3 = \sum_{m \geq 0} (-1)^m (2m+1) q^{m(m+1)/2}.$$

We know that for all primes p and integers $k \geq 1$, we have

(2.8)

$$f_1^{p^k} \equiv f_p^{p^{k-1}} \pmod{p^k}.$$

3. PROOF OF THEOREM 1.1

We prove Theorem 1.1 using elementary q -series techniques, reminiscent of the proof of the result of Majid and Fathima [MF22]. But, before that we need the following result.

Theorem 3.1. *For all integers $\alpha \geq 0$, we have*

$$(3.1) \quad \sum_{n \geq 0} A_{5,4}(5^{\alpha+1}n + 5^{\alpha+1} - 4)q^n = A_\alpha f_5^2 f_1^{14} + B_\alpha q f_5^8 f_1^8 + C_\alpha q^2 f_5^{14} f_1^2 + D_\alpha q^3 \sum_{n \geq 0} A_{5,4}(n)q^n,$$

where $A_0 = 4$, $B_0 = 550$, $C_0 = 12500$, $D_0 = 78125$, and for any integer $n \geq 1$, A_n , B_n , C_n , and D_n are defined as

$$(3.2) \quad A_n = -C_{n-1} + 4D_{n-1},$$

$$(3.3) \quad B_n = -125B_{n-1} + 550D_{n-1},$$

$$(3.4) \quad C_n = -15625A_{n-1} + 12500D_{n-1},$$

$$(3.5) \quad D_n = D_0^{n+1}.$$

Proof. From equation (1.1), we have

$$\sum_{n \geq 0} A_{5,4}(n)q^n = \frac{f_5^{20}}{f_1^4}.$$

From equation (2.5), we have

$$(3.6) \quad \begin{aligned} \sum_{n \geq 0} A_{5,4}(5n+1)q^n &= 4f_5^2 f_1^{14} + 550q f_5^8 f_1^8 + 12500q^2 f_5^{14} f_1^2 + 78125q^3 \frac{f_5^{20}}{f_1^4} \\ &= 4f_5^2 f_1^{14} + 550q f_5^8 f_1^8 + 12500q^2 f_5^{14} f_1^2 + 78125q^3 \sum_{n \geq 0} A_{5,4}(n)q^n. \end{aligned}$$

Equation (3.6), is the case for $\alpha = 0$. Now assume that the result holds for all values up to $\alpha + 1$ ($\alpha \geq 0$). Replacing n by $5n + 4$, and by using Lemmas 2.2, 2.3, 2.4, and equation (3.6), we have

$$\begin{aligned} &\sum_{n \geq 0} A_{5,4}(5^{\alpha+2}n + 5^{\alpha+2} - 4)q^n \\ &= A_\alpha(-15625q^2 f_1^2 f_5^{14}) + B_\alpha(-125q f_1^8 f_5^8) + C_\alpha(-f_1^{14} f_5^2) + D_\alpha(4f_5^2 f_1^{14} + 550q f_5^8 f_1^8 \\ &\quad + 12500q^2 f_5^{14} f_1^2 + 78125q^3 \sum_{n \geq 0} A_{5,4}(n)q^n) \\ &= (-C_\alpha + 4D_\alpha) f_1^{14} f_5^2 + (-125B_\alpha + 550D_\alpha) q f_1^8 f_5^8 \\ &\quad + (-15625A_\alpha + 12500D_\alpha) q^2 f_1^2 f_5^{14} + D_\alpha 78125q^3 \sum_{n \geq 0} A_{5,4}(n)q^n \\ &= A_{\alpha+1} f_1^{14} f_5^2 + B_{\alpha+1} q f_1^8 f_5^8 + C_{\alpha+1} q^2 f_1^2 f_5^{14} + D_{\alpha+1} q^3 \sum_{n \geq 0} A_{5,4}(n)q^n. \end{aligned}$$

Hence, the result is true by induction. □

We can finally prove Theorem 1.1 now.

Proof of Theorem 1.1. From equations (3.2), (3.3), (3.4) and (3.5), we see that

$$\begin{aligned} A_1 &\equiv 0 \pmod{5^5}, & B_1 &\equiv 0 \pmod{5^6}, & C_1 &\equiv 0 \pmod{5^6}, & D_1 &\equiv 0 \pmod{5^7}, \\ A_2 &\equiv 0 \pmod{5^6}, & B_2 &\equiv 0 \pmod{5^7}, & C_2 &\equiv 0 \pmod{5^7}, & D_2 &\equiv 0 \pmod{5^8}, \\ &\vdots & &\vdots & &\vdots & &\vdots \\ A_\alpha &\equiv 0 \pmod{5^{\alpha+4}}, & B_\alpha &\equiv 0 \pmod{5^{\alpha+5}}, & C_\alpha &\equiv 0 \pmod{5^{\alpha+5}}, & D_\alpha &\equiv 0 \pmod{5^{\alpha+6}}. \end{aligned}$$

Now, it is easy to see that equation (3.1) implies Theorem 1.1. □

4. PROOFS OF THEOREMS 1.2, 1.3 AND 1.4

Proof of Theorem 1.2. From the generating function of $A_{p,p^{N_{i-1}}}(n)$ and equation (2.8), we have

$$\sum_{n \geq 0} A_{p,p^{N_{i-1}}}(n)q^n = \frac{f_p^{p(p^{N_{i-1}})}}{f_1^{p^{N_{i-1}}}} \equiv \frac{f_p^{p(p^{N_{i-1}})}}{f_p^{p^{N_{i-1}}}} f_1 \pmod{p^N}.$$

With the help of equation (2.6), we obtain

$$\sum_{n \geq 0} A_{p,p^{N_{i-1}}}(n)q^n \equiv \frac{f_p^{p(p^{N_{i-1}})}}{f_p^{p^{N_{i-1}}}} \left(\sum_{m=-\infty}^{\infty} (-1)^m q^{m(3m-1)/2} \right) \pmod{p^N}.$$

For some m and n , we are interested in finding out whether $m(3m-1)/2 = pn+r$. This is equivalent to asking whether $24pn+24r+1 = (6m-1)^2$, which implies $24r+1 \equiv (6m-1)^2 \pmod{p}$. However $24r+1$ is a quadratic nonresidue modulo p . It follows that

$$A_{p,p^{N_{i-1}}}(pn+r) \equiv 0 \pmod{p^N}.$$

□

Proof of Theorem 1.3. Like before, we have

$$\sum_{n \geq 0} A_{p,p^{N_{i-3}}}(n)q^n = \frac{f_p^{p(p^{N_{i-3}})}}{f_1^{p^{N_{i-3}}}} \equiv \frac{f_p^{p(p^{N_{i-3}})}}{f_p^{p^{N_{i-1}}}} f_1^3 \pmod{p^N}.$$

With the help of equation (2.7), we obtain

$$(4.1) \quad \sum_{n \geq 0} A_{p,p^{N_{i-3}}}(n)q^n \equiv \frac{f_p^{p(p^{N_{i-3}})}}{f_p^{p^{N_{i-1}}}} \left(\sum_{m \geq 0} (-1)^m (2m+1) q^{m(m+1)/2} \right) \pmod{p^N}.$$

For some m and n , we are interested in finding out whether $m(m+1)/2 = pn+r$. This is equivalent to asking whether $8pn+8r+1 = (2m+1)^2$, which implies $8r+1 \equiv (2m+1)^2 \pmod{p}$. However $8r+1$ is a quadratic nonresidue modulo p . It follows that

$$A_{p,p^{N_{i-3}}}(pn+r) \equiv 0 \pmod{p^N}.$$

□

Proof of Theorem 1.4. Due to equations (2.7) and (4.1), we must determine whether $pn+r = m(m+1)/2$ for some integers m and n . Completing the square and considering the result modulo p gives $(2m+1)^2 \equiv 8r+1 \equiv 0 \pmod{p}$. Therefore, p divides $(2m+1)^2$, implying that p divides $2m+1$. Since the coefficient of $q^{m(m+1)/2}$ in the series representation in equation (2.7) is exactly $2m+1$, it follows that the coefficient we are interested in is congruent to 0 modulo p . □

5. PROOF OF THEOREM 1.5

The proofs of the congruences are similar in nature. So, we only present proofs of (1.9) and (1.10). For others, we just give the corresponding generating functions.

First, we prove (1.9). We have,

$$\sum_{n \geq 0} A_{5,4}(n)q^n = \frac{f_5^{20}}{f_1^4} = f_5^{20} \sum_{n \geq 0} P_4(n)q^n.$$

Then, extracting the terms involving q^{5n+1} and dividing by q and then replacing q^5 by q , we obtain

$$\sum_{n \geq 0} A_{5,4}(5n+1)q^n = f_1^{20} \sum_{n \geq 0} P_4(5n+1)q^n.$$

With the help of Lemma 2.1, we get

$$\begin{aligned} \sum_{n \geq 0} A_{5,4}(5n+1)q^n &= f_1^{20} \left(4 \frac{f_5^2}{f_1^6} + 550q \frac{f_5^8}{f_1^{12}} + 12500q^2 \frac{f_5^{14}}{f_1^{18}} + 78125q^3 \frac{f_5^{20}}{f_1^{24}} \right) \\ &= 4f_5^2 f_1^{14} + 550q f_5^8 f_1^8 + 12500q^2 f_5^{14} f_1^2 + 78125q^3 \frac{f_5^{20}}{f_1^4}. \end{aligned}$$

Using equations (2.1) and (2.2), extracting the terms involving q^{5n+4} , dividing by q and then replacing q^5 by q , we arrive at

$$\begin{aligned} \sum_{n \geq 0} A_{5,4}(25n+21)q^n &= 3125 \left(-4f_1^{14} f_5^2 + 100 \frac{f_5^{20}}{f_1^4 R^{15}(q)} + q \left(10450 \frac{f_5^{20}}{f_1^4 R^{10}(q)} - 22f_1^8 f_5^8 \right) \right. \\ &\quad + q^2 \left(46000 \frac{f_5^{20}}{f_1^4 R^5(q)} - 62500 f_1^2 f_5^{14} \right) + 25375q^3 \frac{f_5^{20}}{f_1^4} \\ &\quad \left. - 46000q^4 \frac{f_5^{20} R^5(q)}{f_1^4} + 10450q^5 \frac{f_5^{20} R^{10}(q)}{f_1^4} - 100q^6 \frac{f_5^{20} R^{15}(q)}{f_1^4} \right), \end{aligned}$$

which on usage of (2.4) reduces to

$$\sum_{n \geq 0} A_{5,4}(25n+21)q^n = 3125 \left(96f_1^{14} f_5^2 + 13728q f_1^8 f_5^8 + 312480q^2 f_1^2 f_5^{14} + 1953125q^3 \frac{f_5^{20}}{f_1^4} \right),$$

which implies (1.9).

Proceeding in a similar way, we can also deduce

$$\begin{aligned} \sum_{n \geq 0} A_{5,4}(125n+121)q^n &= 15625 \left(1500004f_1^{14} f_5^2 + 214500550q f_1^8 f_5^8 + 4882512500q^2 f_1^2 f_5^{14} \right. \\ &\quad \left. + 30517578125q^3 \frac{f_5^{20}}{f_1^4} \right) \end{aligned}$$

which proves (1.10).

Now, we give the generating functions which will complete the proofs of the other congruences stated in the result:

$$\begin{aligned} \sum_{n \geq 0} A_{5,2}(25n+23)q^n &= 25 \left(48f_1^4 f_5^4 + 625q \frac{f_5^{10}}{f_1^2} \right), \\ \sum_{n \geq 0} A_{5,2}(125n+123)q^n &= 125 \left(1202f_1^4 f_5^4 + 15625q \frac{f_5^{10}}{f_1^2} \right), \\ \sum_{n \geq 0} A_{5,3}(25n+22)q^n &= 5 \left(5838f_1^9 f_5^3 + 233250q f_1^3 f_5^9 + 1953125q^2 \frac{f_5^{15}}{f_1^3} \right), \\ \sum_{n \geq 0} A_{5,3}(125n+122)q^n &= 25 \left(3643791f_1^9 f_5^3 + 145754625q f_1^3 f_5^9 + 1220703125q^2 \frac{f_5^{15}}{f_1^3} \right). \end{aligned}$$

This completes the proof of Theorem 1.5.

6. PROOF OF THEOREM 1.6

Without loss of generality, we may assume that $r = \sum_{j=0}^{M-1} p^j r_j$ for $0 \leq r_j \leq p-1$, as $\sum_{j=0}^{M-1} p^j r_j$ can take any value between 1 and $p^M - 1$. For integers $M \geq 1$ (sufficiently large) and $N \geq 1$, we have

$$\begin{aligned} \sum_{n \geq 0} A_{p,p^{M+N-1}i+k}(n)q^n &= \frac{f_p^{p(p^{M+N-1}i+k)}}{f_1^{p^{M+N-1}i+k}} = \frac{f_p^{p^{M+N}i}}{f_1^{p^{M+N-1}i}} \sum_{n=0}^{M-1} A_{p,k}(n)q^n \\ &\equiv f_p^{p^{M+N-2}(p^2-1)i} \sum_{n=0}^{M-1} A_{p,k}(n)q^n \pmod{p^N}. \end{aligned}$$

Extracting the terms that involve q^{pn+r_0} from the above identity, we obtain

$$\begin{aligned} \sum_{n \geq 0} A_{p,p^{M+N-1}i+k}(pn+r_0)q^n &\equiv f_1^{p^{M+N-2}(p^2-1)i} \sum_{n=0}^{M-1} A_{p,k}(pn+r_0)q^n \\ &\equiv f_p^{p^{M+N-3}(p^2-1)i} \sum_{n=0}^{M-1} A_{p,k}(pn+r_0)q^n \pmod{p^N}. \end{aligned}$$

Now, extracting the terms that involve $q^{p^2n+r_1}$ from the above identity, we obtain

$$\begin{aligned} \sum_{n \geq 0} A_{p,p^{M+N-1}i+k}(p^2n+r_0+pr_1)q^n &\equiv f_1^{p^{M+N-3}(p^2-1)i} \sum_{n \geq 0} A_{p,k}(p^2n+r_0+pr_1)q^n \\ &\equiv f_p^{p^{M+N-4}(p^2-1)i} \sum_{n \geq 0} A_{p,k}(p^2n+r_0+pr_1)q^n \pmod{p^N}. \end{aligned}$$

From the above identity, we extract the terms that contain $q^{p^2n+r_2}$, and from the resulting identity, we again extract the terms that contain $q^{p^2n+r_3}$ and so on. It can be seen that after the M -th extraction using this iterative scheme, we arrive at

$$\begin{aligned} (6.1) \quad \sum_{n \geq 0} A_{p,p^{M+N-1}i+k}(p^M n + r_0 + pr_1 + \cdots + p^{M-1}r_{M-1})q^n \\ \equiv f_1^{p^{N-1}(p^2-1)i} \sum_{n \geq 0} A_{p,k}(p^M n + r_0 + pr_1 + \cdots + p^{M-1}r_{M-1})q^n \pmod{p^N}. \end{aligned}$$

Therefore, if we assume that $A_{p,k}(p^M n + r_0 + pr_1 + \cdots + p^{M-1}r_{M-1}) = A_{p,k}(p^M n + r) \equiv 0 \pmod{p^N}$, then from the above identity, we have

$$A_{p,p^{M+N-1}i+k}(p^M n + r) \equiv 0 \pmod{p^N}.$$

This completes the proof of Theorem 1.6.

We have the following easy corollary, which follows from equation (6.1) when $M = 1$.

Corollary 6.1. *Let p be a prime, $k \geq 1$, $j \geq 0$, $N \geq 1$, and r be integers such that $1 \leq r \leq p-1$. Then for all $n \geq 0$, we have*

$$\sum_{n \geq 0} A_{p,p^N i+k}(pn+r)q^n \equiv f_1^{p^{N-1}(p^2-1)i} \sum_{n \geq 0} A_{p,k}(pn+r)q^n \pmod{p^N}.$$

7. CONCLUDING REMARKS

- (1) We have found several congruences modulo powers of 5, individual as well as infinite families similar to those stated in Theorem 1.6, for $A_{5,k}(n)$ for higher values of k . The proofs of these are routine excises similar to the proofs of (1.9) and (1.10) hence, they are not proved here.

For instance, the following congruences are true:

$$\begin{aligned}
(7.1) \quad & A_{5,6}(25n + 14, 19, 24) \equiv 0 \pmod{25}, \\
(7.2) \quad & A_{5,6}(125n + 119) \equiv 0 \pmod{125}, \\
(7.3) \quad & A_{5,7}(25n + 13, 18, 23) \equiv 0 \pmod{25}, \\
(7.4) \quad & A_{5,7}(125n + 118) \equiv 0 \pmod{125}, \\
(7.5) \quad & A_{5,25i+6}(25n + 14, 19, 24) \equiv 0 \pmod{5}, \\
(7.6) \quad & A_{5,125i+6}(25n + 14, 19, 24) \equiv 0 \pmod{25}, \\
(7.7) \quad & A_{5,3125i+6}(125n + 119) \equiv 0 \pmod{125}, \\
(7.8) \quad & A_{5,25i+7}(25n + 13, 18, 23) \equiv 0 \pmod{5}, \\
(7.9) \quad & A_{5,125i+7}(25n + 13, 18, 23) \equiv 0 \pmod{25}, \\
(7.10) \quad & A_{5,3125i+7}(125n + 118) \equiv 0 \pmod{125}.
\end{aligned}$$

- (2) Looking at the sequence of results in equations (1.2), (1.3) and (1.9), it suggests the possibility of there being other such structures modulo higher powers of 5, and more generally for $A_{t,k}(n)$ modulo higher powers of t . Experiments also suggest some sort of cyclic behaviour, which we do not conjecture here, but it would be interesting to know what can be said about $A_{5,5i+3}(5^\alpha n + 5^\alpha - 3) \pmod{5^\alpha}$ and $A_{5,5i+2}(5^\alpha n + 5^\alpha - 2) \pmod{5^\alpha}$?
- (3) Experiments suggest some additional infinite family of congruences modulo powers of 5 which are stronger results than those indicated by Theorem 1.6. We present them here as conjectures:

$$\begin{aligned}
(7.11) \quad & A_{5,5i+1}(25n + 24) \equiv 0 \pmod{25}, \\
(7.12) \quad & A_{5,25i+2}(25n + 23) \equiv 0 \pmod{25}, \\
(7.13) \quad & A_{5,125i+2}(125n + 123) \equiv 0 \pmod{125}, \\
(7.14) \quad & A_{5,125i+3}(125n + 122) \equiv 0 \pmod{25}, \\
(7.15) \quad & A_{5,625i+4}(125n + 121) \equiv 0 \pmod{3125}, \\
(7.16) \quad & A_{5,3125i+4}(125n + 121) \equiv 0 \pmod{15625}, \\
(7.17) \quad & A_{5,25i+6}(25n + 14, 19) \equiv 0 \pmod{25}, \\
(7.18) \quad & A_{5,25i+6}(125n + 119) \equiv 0 \pmod{125}, \\
(7.19) \quad & A_{5,25i+7}(25n + 13, 18, 23) \equiv 0 \pmod{25}, \\
(7.20) \quad & A_{5,125i+7}(125n + 118) \equiv 0 \pmod{125}.
\end{aligned}$$

- (4) Many other congruences modulo higher powers of 5 may exist for higher subsequences. It may be interesting to study those congruences.

ACKNOWLEDGEMENTS

The second author is partially supported by the institutional fellowship for doctoral research from Tezpur University, Napaam, India. The third author was partially supported by the Council of Scientific & Industrial Research (CSIR), Government of India under the CSIR-JRF scheme. The author thanks the funding agency.

REFERENCES

- [Ber91] Bruce C. Berndt. *Ramanujan's notebooks. Part III*. Springer-Verlag, New York, 1991.
- [Ber06] Bruce C. Berndt. *Number theory in the spirit of Ramanujan*, volume 34 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2006.
- [MF22] N. V. Majid and S. N. Fathima. On a Ramanujan-type congruence for partition triples with 5-cores. *J. Integer Seq.*, 25(6):Art. 22.6.2, 7, 2022.
- [Ran16] Dasappa Ranganatha. On a Ramanujan-type congruence for bipartitions with 5-cores. *J. Integer Seq.*, 19(8):Art. 16.8.1, 5, 2016.

DEPARTMENT OF HUMANITIES AND BASIC SCIENCES, INDIAN INSTITUTE OF INFORMATION TECHNOLOGY (IIIT) MANIPUR, MANTRIPUKHRI, IMPHAL 795002, INDIA

Email address: manjil@saikia.in

DEPARTMENT OF MATHEMATICAL SCIENCES, TEZPUR UNIVERSITY, NAPAAM 784028, ASSAM, INDIA

Email address: abhiraj002@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES, TEZPUR UNIVERSITY, NAPAAM 784028, ASSAM, INDIA

Email address: msp21105@tezu.ac.in