

Upper bounds for eigenvalue multiplicities of almost cyclic elements in irreducible representations of simple algebraic groups

A.E. Zalesski

Abstract

We study the irreducible representations of simple algebraic groups in which some non-central semisimple element has at most one eigenvalue of multiplicity greater than 1. We bound the multiplicity of this eigenvalue in terms of the rank of the group. ¹ ²

Dedicated to the memory of Irina Suprunenko

1 Introduction

In this paper we study a special case of a general problem on eigenvalue multiplicities of semisimple elements in irreducible representations of algebraic groups. This problem is connected with a similar problem on weight multiplicities. In [22] we determine the irreducible representations of simple algebraic groups G whose all non-zero weights are of multiplicity 1, and compute the zero weight multiplicity in each case. Note that weights are one-dimensional representations of a maximal torus T , say, of G , and the multiplicity of a weight ω , say, in a representation ρ of G is the multiplicity of ω as a constituent of $\rho(T)$. Turning to individual non-central semisimple elements $g \in G$, an analog of this result could be (i) obtaining a list of the irreducible representations ρ of G such that at most one eigenvalue multiplicity of $\rho(g)$ is greater than 1, and then (ii) determining the multiplicity of the exceptional eigenvalue. An essential step of understanding problem (i) is made in [24], where it is proved that all non-zero weights of ρ must be of multiplicity 1. This does not mean however that the eigenvalue multiplicities of $\rho(g)$ are the same as the weight multiplicities. This can be seen on examples with G to be of classical type and ρ the natural representation of G . Moreover, in these examples $m_g(\rho)$, the maximum eigenvalue multiplicity of $\rho(g)$, is almost arbitrary (subject to the obvious condition $m_g(\rho) \leq \dim \rho$), whereas all weight multiplicities of ρ are equal to 1. However, it is shown below that

¹2020 Mathematics subject classification: 20G05, 15A18

²Keywords: simple algebraic groups, representations, regular elements, eigenvalue multiplicities.

$m_g(\rho)$ can be bounded in terms of the rank of G . An initial step in this direction is made in [24], where, for a non-central non-regular semisimple element $g \in G$, it is shown that the natural representation of a classical group G is essentially the only one in which at most one eigenvalue multiplicity of $\rho(g)$ is greater than 1; that result is quoted below as Theorem 2.18. To be more precise, ρ is a twist of the natural representation. (We call a representation ρ' a twist of ρ if $\rho'(g) = \rho(\sigma(g))$ for $g \in G$ and a surjective homomorphism $\sigma : G \rightarrow G$.) Therefore, the current paper mainly deals with regular semisimple elements.

The main result of this paper is the following

Theorem 1.1. *Let G be a simple algebraic group of rank n and V an irreducible G -module. Let $s \in G$ be a regular semisimple element and let $m_s(V)$ be the maximum of the eigenvalue multiplicities of s on V . Suppose that at most one eigenvalue multiplicity of s on V exceeds 1. Then $m_s(V) \leq n + 1$.*

In fact, we can give more details as follows:

Theorem 1.2. *Let G be a simple algebraic group of rank n and V an irreducible G -module. Let $s \in G$ be a regular semisimple element and let $m_s(V)$ be the maximum of the eigenvalue multiplicities of s on V . Suppose that at most one eigenvalue multiplicity of s on V exceeds 1 and $m_s(V) > 1$. Then either $m_s(V) = 2$ or one of following holds.*

- (1) $G = A_n$ and $m_s(V) \leq n + 1$;
- (2) V is a twist of a G -module from Table 1 and $m_s(V)$ equals the weight zero multiplicity of V recorded there; in particular $n - 2 \leq m_s(V) \leq n + 1$;
- (3) $G = E_6$ and $m_s(V) \leq 3$.

The bound in item (1) of Theorem 1.2 is sharp for $n > 1$, at least for $p \neq 0$, see Example 3.37. In this example for $G = A_n$ we construct a tensor-decomposable G -module V with $m_s(V) = n + 1$. For E_6 see Example 3.17.

Corollary 1.3. *Let G be a simple algebraic group in defining characteristic $p \geq 0$, $s \in G$ a regular semisimple element and let V be an irreducible G -module with highest weight $\omega \neq 0$. Suppose that at most one eigenvalue multiplicity of s on V exceeds 1. Then $m_s(V) < \dim V/2$, unless $G = A_1$, $\dim V = 3, p \neq 2$ or $\dim V = 4$ and either $\omega = 3p^i\omega_1, p \neq 2, 3$ or $\omega = (p^i + p^j)\omega_1$ with $1 \leq i < j$. (If $p = 0$ and $n = 4$ then $\omega = 3\omega_1$.)*

The fact that the multiplicity $m_s(V)$ in question is of certain significance can be observed from [8], in which the authors study irreducible subgroups X of $GL_n(q) = GL(V)$, containing a semisimple element g of prime order acting irreducibly on $(\text{Id} - g)V$. They assume that $\dim(\text{Id} - g)V > \dim V/2$, equivalently, $m_s(V) = \dim V - \dim(\text{Id} - g)V < \dim V/2$, and we complement this result by dropping the latter assumption for X a quasi-simple group of Lie type in defining characteristic $p|q$. More precisely, we use Corollary 1.3 to prove the following result. In the statement below $\bar{V} = V \otimes F$, where F is an algebraically closed field containing \mathbb{F}_q .

Theorem 1.4. *Let G be a quasi-simple group of Lie type in defining characteristic $p > 0$, and let V be a non-trivial irreducible $\mathbb{F}_q G$ -module where $p \nmid q$. Let $s \in G$ be a regular semisimple element. Suppose that at most one eigenvalue multiplicity of s on \bar{V} exceeds 1. Then $m_s(\bar{V}) < \dim V/2$, unless $G = SL_2(q')$ or $PSL_2(q')$ for some p -power q' , and either $m_s(\bar{V}) = \dim V/2$ is even or $\dim V = 3, p \neq 2$ and $s^2 \in Z(G)$.*

Conversely, if V is a vector space over \mathbb{F}_q and $\dim V \equiv 0 \pmod{4}$ or $p \neq 2$ and $\dim V = 3$ then there exists a group $G \cong SL_2(q')$, a semisimple element $s \in G$ and irreducible representation $\rho : SL_2(q') \rightarrow GL(V)$ such that at most one eigenvalue multiplicity of $\rho(s)$ on \bar{V} exceeds 1 and $m_s(\bar{V}) = \max\{\dim V/2, 2\}$.

In Theorem 1.4 saying that an element $g \in G$ is regular means that g is regular as an element of a simple algebraic group \mathbf{G} from which G is obtained as the fixed point set of a Steinberg (Frobenius) endomorphism of \mathbf{G} , see [15, §21.1]. For a semisimple element $g \in G$ this is equivalent to saying that $C_G(g)$ has no non-identity unipotent element. If $g \in G$ is semisimple and not regular then we have a result similar to that of [8]. To state it, it is convenient to introduce a notion of a nearly natural module of a classical group as follows.

Let G be a quasi-simple group of classical Lie type in defining characteristic $p > 0$. Let V be an irreducible $\mathbb{F}_q G$ -module for some p -power q and F an algebraically closed field containing \mathbb{F}_q , and $\bar{V} = V \otimes F$. Let $G = \mathbf{G}^\sigma = \{g \in \mathbf{G} : \sigma(g) = g\}$, where σ is a Steinberg endomorphism of \mathbf{G} . We say that V is *nearly natural* if the composition factor dimensions of G on \bar{V} are the same as the dimension of the natural \mathbf{G} -module. (See *Notation* for the formal definition of the natural module.) This notion unfortunately depends on the Lie type of \mathbf{G} . For instance, the dimensions of the natural module for $A_3 \cong D_3$ are 4 and 6, respectively, and these for $B_2 \cong C_2$ with $p \neq 2$ are 5, 4, respectively.

The following statement is a special case of the main theorem of [8].

Theorem 1.5. *Let G be a quasi-simple group of Lie type in defining characteristic $p > 0$ and q a p -power. Let $s \in G$ be a non-central semisimple element and let $\rho : G \rightarrow GL_n(q)$, $n > 9$, be an irreducible representation such that $\rho(g)$ is similar to $\text{diag}(\text{Id}_m, M)$, where $M \in GL_{n-m}(q)$ is irreducible. Suppose that $m < n/2$. Then G is classical and ρ is nearly natural.*

Note that the cases with $n \leq 9$ are listed in [8, Table 6].

Theorem 1.4 deals with regular elements $g \in G$. The case where g is not regular follows from Theorem 2.18 obtained in [24]:

Theorem 1.6. *Let G be a quasi-simple group of Lie type in defining characteristic p and $g \in G$ a non-central semisimple element. Let $\rho : G \rightarrow GL_n(q)$ be an irreducible representation. Suppose that g is not regular and at most one eigenvalue multiplicity of $\rho(g)$ exceeds 1. Then ρ is nearly natural for G or for a group $G_1 \cong G$. (Here $G_1 = Spin_6^+(q)$ if $G = SL_4(q)$; $G_1 = Spin_6^-(q)$ if $G = SU_4(q)$; $G_1 = C_2(q)$ if $G = B_2(q)$; $G_1 = B_2(q)$ if $G = C_2(q)$.)*

Notation The cardinality of a finite set A is denoted by $|A|$. The ring of integers is denoted by \mathbb{Z} , and by \mathbb{Z}^+ the subset of non-negative integers. If $0 \neq a, b$ are natural

number then we write $a|b$ to say that b is a multiple of a and we denote by (a, b) the greatest common divisor of a, b . We fix F for an algebraically closed field of characteristic $p \geq 0$; the multiplicative group of F is denoted by F^\times . \mathbb{C} is the field of complex numbers. We write $Z(X)$ for the center of a group X . For $x \in X$ the group generated by x is denoted by $\langle x \rangle$, by $C_X(x)$ the centralizer of x in X and by $|x|$ the order of x . A diagonal $(n \times n)$ -matrix with sequent diagonal entries d_1, \dots, d_n is denoted by $\text{diag}(d_1, \dots, d_n)$.

For our convenience we call an element $g \in GL_n(F)$ *cyclic* if g is diagonalizable and all eigenvalues of g are of multiplicity 1. This is equivalent to saying that the characteristic polynomial of g has no multiple roots. We call g *almost cyclic* if g is diagonalizable and at most one of the eigenvalues of g is of multiplicity greater than 1. This is meaningful if $g \in GL_n(F_1)$, where F_1 is a subfield of F . (See [6] and [7] for more comments on these notions.)

In Section 2,3 G is a non-trivial simply connected simple linear algebraic group over F (unless otherwise explicitly stated); sometimes we write $G(F)$ to make it precise. (In Section 4 we use \mathbf{G} a simple algebraic group and G for a finite group of Lie type.) All G -modules considered are rational finite-dimensional FG -modules. For a G -module V (or a rational representation ρ of G) we write $V \in \text{Irr}(G)$ (or $\rho \in \text{Irr}(G)$) to mean that V (or ρ) is irreducible. If H is a subgroup of G then $V|_H$ means the restriction of a G -module V to H .

We fix a maximal torus T in G and a Borel subgroup containing T , which in turn define the roots and simple roots of G , as well as the weights of G -modules and representations. The set $\Omega = \text{Hom}(T, F^\times)$ (the rational homomorphisms of T to F^\times) is called the *weight lattice*, which is free \mathbb{Z} -module of finite rank called *the rank of G* . The elements of Ω are called *T -weights of G* . As T is fixed we write "weights" in place of " T -weights". For $t \in T$ and $\mu \in \Omega(V)$, $\mu(t)$ is an eigenvalue of t on V ; then $(-\mu)(t) = t^{-1}$ and $(\mu + \nu)(t) = \mu(t)\nu(t)$ for $\nu \in \Omega(V)$.

The T -weights of a G -module V are the irreducible constituents of the restriction of V to T . When T is fixed (which we assume throughout the paper), we omit the indication to T and write "weights" in place of " T -weights". The set of weights of V is denoted by $\Omega(V)$. If μ is a weight of V , the μ -weight space is $\{v \in V : tv = \mu(t)v \text{ for all } t \in T\}$; the dimension of the μ -weight space is called the *multiplicity of μ on V* . The Weyl group of G is denoted by $W(G)$; in fact $W(G) = N_G(T)/T$ so the conjugation action of $N_G(T)$ on T yields an action of $W(G)$ on T and on the T -weights. The W -orbit of $\mu \in \Omega$ is denoted by $W\mu$.

We use G_{reg} to denote the set of regular elements of G , in particular, T_{reg} is the set of regular elements of T . As every semisimple element of G is conjugate to an element of T , in the most of our results the assumption $g \in G$ semisimple is equivalent to saying $g \in T$, the latter is more rigorous when we deal with $\mu(g)$ for a weight μ .

With an algebraic group is associated the Lie algebra of G denoted here by $L(G)$. The group G acts on $L(G)$ in the natural way making it a G -module. If G is simple (or reductive) then the non-zero weights of T on $L(G)$ are called *roots*. We denote the set of roots by Φ or $\Phi(G)$. The \mathbb{Z} -span of Φ is called the *root lattice* and denoted by R or $R(G)$. In $\Phi(G)$ one

can choose a \mathbb{Z} -basis Π of R whose elements are denoted by $\alpha_1, \dots, \alpha_n$ and called *simple roots* (here n is the rank of G). The ordering of simple roots and the Dynkin diagrams of G are as in [3]. The weights in R are called *radical*.

One defines a non-degenerate symmetric bilinear form on Ω , which we express as (μ, ν) . The elements ω_i satisfying $(\omega_j, \alpha_i) = 2(\alpha_i, \alpha_i)\delta_{ij}$ (the Kronecker delta) belong to Ω and called *fundamental weights* [3, Ch. VI, §1, no.10]. These form a \mathbb{Z} -basis of Ω so every $\nu \in \Omega$ can be express in the form $\sum a_i \omega_i$ with $a_1, \dots, a_n \in \mathbb{Z}$; the set of ν with $a_1, \dots, a_n \geq 0$ is denoted by Ω^+ . The weights of Ω^+ are called *dominant*. Suppose that $p > 0$; a (dominant) weight (a_1, \dots, a_n) is called *p-restricted* if $0 \leq a_i < p$ for all $i = 1, \dots, n$; the set of them is denoted by Ω_p^+ . For uniformity, we often do not separate the cases with $p = 0$ and $p > 0$; by convention, a *p-restricted* weight is simply a dominant weight when $p = 0$. An irreducible G -module is called *p-restricted* if so is the highest weight of it.

The weight with $a_1 = \dots = a_n = 0$ is denoted by ω_0 and called the *zero weight* (also the weight 0). We often simplify $\mu \neq \omega_0$ by writing $\mu \neq 0$. We can (and sometimes do) write the elements of Ω as (a_1, \dots, a_n) and in some cases, especially when we compare representations of $G(\mathbb{C})$ with $G(F)$, saying that weights of $G(\mathbb{C})$ and $G(F)$ are the same means the coincidence of the set of strings (a_1, \dots, a_n) for these groups. We set $\Omega^+(V) = \Omega^+ \cap \Omega(V)$, so $\Omega^+(V)$ is the set of dominant weights of V .

There is a standard partial ordering of elements of Ω ; specifically, for $\mu, \mu' \in \Omega$ we write $\mu \preceq \mu'$ and $\mu' \succeq \mu$ if and only if $\mu - \mu' \in R^+$. (Sometimes we write $\mu \prec \mu'$ and $\mu' \succ \mu$ to indicate that $\mu \neq \mu'$.) Every non-trivial irreducible G -module has a weight ω such that $\mu \prec \omega$ for every $\mu \in \Omega(V)$ with $\mu \neq \omega$. This is called the *highest weight of V*. There is a bijection between Ω^+ and $\text{Irr}(G)$, so for $\omega \in \Omega^+$ we denote by V_ω the irreducible G -module with highest weight ω . The zero weight space of V is sometimes denoted by V^T . In general, for $S \subseteq G$ we set $V^S = \{v \in V : sv = v \text{ for all } s \in S\}$. $L(G)$ is a G -module called *adjoint*; it is irreducible if $p = 0$; if $p > 0$ it may be reducible but has a highest weight denoted here by ω_a . We denote by V_a the composition factor of $L(G)$ with highest weight ω_a .

Every $\omega = \sum a_i \omega_i \in \Omega^+$ has a unique p -adic expansion $\omega = \lambda_0 + p\lambda_1 + \dots + p^k \lambda_k$ for some integer $k \geq 0$, where $\lambda_0, \dots, \lambda_k \in \Omega_p^+$ are p -restricted dominant weights. This yields a formula $V_\omega = V_{\lambda_0} \otimes V_{\lambda_1}^{Fr} \otimes \dots \otimes V_{\lambda_k}^{Fr^k}$, which we refer as the Steinberg expansion of V .

Our notation for simple algebraic groups is standard, in particular those of types A_n, B_n, C_n, D_n are called *classical*, E_6, E_7, E_8, F_4, G_2 are called *exceptional*, the subscript is the rank of the group. Here $n \geq 1$ for A_n , and we usually assume $n > 1$ for B_n, C_n , and $n > 3$ for D_n , due to isomorphisms $D_3 \cong A_3$, $D_2 \cong A_1 \times A_1$ and $A_1 \cong B_1 \cong C_1$. For brevity we write $G = A_n$ to say that G is a simple simply connected algebraic group of type A_n , and similarly for the other types. There is also an isomorphism $B_2 \cong C_2$, and we usually choose C_2 for the standard representative of this algebraic group. (In Section 4 we allow to view A_3 as D_3 for uniformity of some statements.) If $p = 2$ then for every n there are surjective algebraic group homomorphisms $B_n \rightarrow C_n$ and $C_n \rightarrow B_n$ with trivial kernel (so these are abstract group isomorphisms); for our purpose the choice between these two groups is not essential, so we choose to work with C_n .

For an algebraic group G of classical type the module (and the representation) with

highest weight ω_1 is called *natural*.

In our reasonings we regularly use so called "Bourbaki weights", which are elements of a \mathbb{Z} -lattice containing $\Omega(G)$ with basis $\varepsilon_1, \varepsilon_2, \dots$; the explicit expressions of the fundamental weights and the simple roots of G in terms of ε_i 's is given in [3, Planches 1 – XIII].

If $\sigma : G \rightarrow G$ is a surjective algebraic group homomorphism and ϕ is a representation of G then the σ -twist ϕ^σ of ϕ is defined as the mapping $g \rightarrow \phi(\sigma(g))$ for $g \in G$. Of fundamental importance is the Frobenius mapping $Fr : G \rightarrow G$ arising from the mapping $x \rightarrow x^p$ ($x \in F$) when $p > 0$. If V is a G -module then the modules V^{Fr^k} ($k \geq 1$ an integer) are called *Frobenius twists of V* ; if V is irreducible with highest weight ω then the highest weight of V^{Fr^k} is $p^k\omega$ for some integer $k \geq 0$.

A surjective algebraic group homomorphism $\sigma : G \rightarrow G$ is called *Steinberg* if the subgroup $G^\sigma = \{g \in G : \sigma(g) = g\}$ is finite. Equivalently, σ is Steinberg if and only if $\sigma^m = Fr^k$ for some integers $m, k > 0$, see [15, p. 183]. If G is semisimple and σ is Steinberg then G^σ is called a finite group of Lie type.

Let G be an algebraic group, V a G -module and let T be a maximal torus of G . We say that $s \in T$ *separates a weight ω of V* if the eigenvalue $\omega(s)$ equals the multiplicity of ω . In addition, we say that s *separates the weights of V* if every eigenspace of s on V is a weight space. It is well known that s is regular if and only if it separates weight zero of the adjoint module. If s separates all weights of the adjoint module, we say that s is *strictly regular*.

The set of eigenvalues of an element $g \in G$ on a G -module V (disregarding the multiplicities) is denoted by $E_V(g)$.

2 Preliminaries

In this section G is a simple simply connected linear algebraic group defined over a field of characteristic $p \geq 0$. For our purpose we can assume that $G = G(F)$, the group of F -points of G , where F is an algebraically closed field of characteristic p .

Recall that a diagonalizable matrix is called *cyclic* if all of its eigenspaces are one-dimensional, and *almost cyclic* if at most one of its eigenspaces has dimension greater than 1.

In Lemma 2.1 tM means the transpose of a matrix M .

Lemma 2.1. (1) *Let $M \in M_{r \times r}(F)$ be an almost cyclic diagonal matrix such that M is conjugate to ${}^tM^{-1}$ and let $e \in F$ be an eigenvalue of M . If $e \notin \{\pm 1\}$, then the multiplicity of e equals 1.*

(2) *Let G be a semisimple algebraic group, $s \in G$ a semisimple element, and V is a G -module. Suppose that V is self-dual and s is almost cyclic on V . Then all eigenvalues of s on V , not equal to 1 or -1 , occur with multiplicity 1. In particular, this holds if V is irreducible and G is simple of type A_1, B_n, C_n, D_n (n even), E_n, F_4 and G_2 .*

Proof. The first claim is clear, whereas the second one follows from a well known fact that every irreducible representation of G listed is self-dual. \square

Lemma 2.2. *Let V, V_1, V_2 be G -modules. Suppose that $\Omega(V) = \Omega(V_1) + \Omega(V_2)$ and s is separates the weights of V . Then s separates the weights of V_1 and of V_2 .*

Proof. Suppose the contrary, let, say, s does not separate the weights of V_1 . Then there are distinct weights λ, μ of V_1 such that $\lambda(s) = \mu(s)$. Let ν be a weight of V_2 . Then $(\lambda + \nu)(s) = (\mu + \nu)(s)$, and $\lambda + \nu, \mu + \nu$ are distinct weights of V . This is a contradiction. \square

Lemma 2.3. *Let $M = M_1 \otimes M_2$ be a Kronecker product of diagonal non-scalar matrices M_1, M_2 of sizes $m \leq n$, respectively.*

(1) *If M_1, M_2 are cyclic then the eigenvalue multiplicities of M do not exceed m .*

(2) *Suppose that M is almost cyclic. Then M_1 and M_2 are cyclic.*

(3) *Suppose that M is almost cyclic and M_i is similar to M_i^{-1} for $i = 1, 2$. Then the eigenvalue multiplicities of M do not exceed 2. In addition, if e is an eigenvalue of M of multiplicity 2 then $e \in \{\pm 1\}$.*

Proof. (1) Suppose the contrary, and let $r > m$ be the multiplicity of some eigenvalue e of M . Let $a_1, \dots, a_m, b_1, \dots, b_n$ be the eigenvalues of M_1, M_2 , respectively. Then there exist a_i, b_j, b_k ($1 \leq i \leq m, 1 \leq j < k \leq n$) such that $a_i b_j = a_i b_k$ and hence $b_j = b_k$. This is a contradiction.

(2) Suppose that M_1 has an eigenvalue e , say, of multiplicity $r > 1$. Let b_1, b_2 be distinct eigenvalues of M_2 . Then eb_1, eb_2 are distinct eigenvalues of M , each of multiplicity greater than 1. So M is not almost cyclic, thus establishing the claim.

(3) Suppose the contrary, and let e be an eigenvalue of M of multiplicity at least 2. By (2), M_1 and M_2 are cyclic so $e = a_i b_i$ for $i = 1, 2$ and some (distinct) eigenvalues a_i of M_1 and b_i of M_2 . Then $e^{-1} = a_i^{-1} b_i^{-1}$ is an eigenvalue of M of the same multiplicity as that of e . As M is almost cyclic, $e = e^{-1}$ and $e \in \{\pm 1\}$. Then $a_1 b_1 = a_2^{-1} b_2^{-1}$, whence $a_1 b_2 = a_2^{-1} b_1^{-1}$. If $a_2^{-1} \neq a_1$ then $a_1 b_2$ is an eigenvalue of M of multiplicity at least 2; this is not equal to $e = a_1 b_1$, hence s is not almost cyclic on M , a contradiction. So $a_2 = a_1^{-1}$. If the multiplicity of e is at least 3, say, $e = a_3 b_3$ then, as above, this implies $a_3 = a_1^{-1}$. Then $a_3 = a_2$, contrary to (2). \square

Lemma 2.4. *Let V, M be non-trivial G -modules and $s \in T$.*

(1) *If s is cyclic on V then s is regular.*

(2) *If s is almost cyclic on $V \otimes M$ then s is regular.*

Proof. (1) Let ρ be the representation afforded by V . If s is cyclic on V then $C_{GL(V)}(\rho(s))$ is a torus. Then so is $C_G(s)^\circ$, giving the claim.

(2) By Lemma 2.3(2), s is cyclic on V , so the result follows from (1). \square

In Lemma 2.5 below and elsewhere, for $A, B \subset \Omega$ we define $A + B = \{a + b : a \in A, b \in B\}$.

Lemma 2.5. *Let V, V_1, V_2 be non-trivial G -modules. Let $s \in G$ be a non-central semisimple element and assume that s is almost cyclic on V .*

(1) Suppose that $V = V_1 \otimes V_2$. Then the matrices of s on V_1 and V_2 are cyclic, all weights of V_1 and V_2 are of multiplicity 1, and s is regular.

(2) Suppose that $\Omega(V_1) + \Omega(V_2) = \Omega(V)$. Then s separates the weights of V_1 and V_2 .

(3) Suppose that $\Omega(V_1) + \Omega(V_2) = \Omega(V)$ and the weights of V are of multiplicity 1. Then the multiplicity of every eigenvalue of s on V does not exceed $|\Omega(V_1)|$, the number of weights of V_1 .

Proof. Assertion (1) follows from Lemma 2.3, except the last claim, so suppose that s is not regular. Then $C_G(s)$ contains a unipotent element $u \neq 1$. As u stabilizes every eigenspace of s on V_1 , at least one of them is of dimension greater than 1, contradicting that s is cyclic on V_1 .

(2) Suppose the contrary, that the weights of V_1 , say, are not separated by s , so there exist distinct weights $\mu_1, \mu_2 \in \Omega(V_1)$ such that $\mu_1(s) = \mu_2(s)$. Then for every $\lambda, \mu \in \Omega(V_2)$, $\mu_i + \lambda, \mu_i + \mu \in \Omega(V)$ for $i = 1, 2$ and $(\mu_1 + \lambda)(s) = (\mu_2 + \lambda)(s)$ and $(\mu_1 + \mu)(s) = (\mu_2 + \mu)(s)$. This implies $\lambda(s) = \mu(s)$ as s is almost cyclic on V . Since μ is an arbitrary weight of V_2 , we conclude that s is scalar on V_2 , and hence $s \in Z(G)$, a contradiction.

(3) Suppose the contrary. Let e be an eigenvalue of s on V whose multiplicity exceeds $m = |\Omega(V_1)|$. As all weight spaces of V have dimension at most one, there are at least $m + 1$ distinct weights $\sigma_1, \dots, \sigma_{m+1}$ of V such that $e = \sigma_1(s) = \dots = \sigma_{m+1}(s)$. We have $\sigma_i = \mu_i + \lambda_i$ for $\mu_i \in \Omega(V_1)$, $\lambda_i \in \Omega(V_2)$ for $i = 1, \dots, m + 1$. By hypothesis, $\mu_i = \mu_j$ for some $1 \leq i \neq j \leq m + 1$, which yields both $\lambda_i \neq \lambda_j$ and $\lambda_i(s) = \lambda_j(s)$, contradicting (2). \square

Obviously, $\Omega(V_1) + \Omega(V_2) = \Omega(V)$ if $V = V_1 \otimes V_2$; however in many instances the former equality holds for $V \neq V_1 \otimes V_2$, see Lemma 2.12(2) below. The next example illustrates the use of Lemma 2.5(3).

Example 2.6. (1) Let $G = A_n$, $p = 0$, $V_1 = V_{\omega_1}$, $V_2 = V_{m\omega_1}$, $V = V_{(m+1)\omega_1}$. Then all weight multiplicities of V are known to be equal to 1 and $\Omega(V_1) + \Omega(V_2) = \Omega(V)$ by [4, Ch. VIII, §7, Proposition 10]. So if s is almost cyclic on V then the multiplicity of any eigenvalue of s on V is at most $n + 1$.

(2) Let $G = A_n$, $V_1 = V_{\omega_1}$, $V_2 = V_{\omega_n}$ and $V = V_{\omega_1 + \omega_n}$. Then $\Omega(V) = \Omega(V_1) + \Omega(V_2)$. As V satisfies the hypotheses of Lemma 2.13(2), some s is almost cyclic on V . Here the zero weight of V is of multiplicity $n + 1$, so the bound in Lemma 2.5(3) is sharp.

For reader's convenience we record here the following well known observations on the natural modules for classical groups. Note that we add the case $n = 3$ for D_n for our convenience here.

Lemma 2.7. [24, Lemma 6] *Let $G \in \{A_n, B_n, p \neq 2, C_n, n > 1, D_n, n \geq 3\}$, $V = V_{\omega_1}$ and let $s \in G$ be a non-central semisimple element.*

(1) *If s is cyclic on V , then s is regular. For $G = A_n$ or C_n , the converse is true.*

(2) *If $G = B_n$, then s is regular if and only if the multiplicity of the eigenvalue -1 on V is at most 2 and the other eigenvalue multiplicities are equal to 1.*

(3) If $G = D_n$, then s is regular if and only if the multiplicities of the eigenvalues 1 and -1 on V are at most 2 and the other eigenvalue multiplicities are equal to 1. In addition, if s is not almost cyclic on V , then s is not almost cyclic on V_{ω_2} .

(4) If s is regular, then s is almost cyclic on V unless $G = D_n, p \neq 2$ and $1, -1$ are eigenvalues of s on V , each of multiplicity 2.

Lemma 2.8. *Let $G = B_2, p \neq 2$, and let $s \in T_{reg}$. Then s is cyclic on V_{ω_2} and almost cyclic on V_{ω_1} .*

Proof. This follows from [25, Lemma 3.10](1). (Note that there is an algebraic group homomorphism $h : Spin_5(F) \rightarrow Sp_4(F)$; under this homomorphism the natural module for B_2 has highest weight ω_2 as a C_2 -module. The natural module for C_2 viewed as a module for B_2 has highest weight ω_2 .) \square

The following lemma is frequently used with no explicit reference, especially when S consists of a single element.

Lemma 2.9. [19, Ch. II, §4, Theorem 4.1] *Let G be a connected algebraic group, T a maximal torus and $S \subset T$ a subset. Then the following conditions are equivalent:*

- (1) $C_G(S)^\circ = T$;
- (2) if $gs = sg$ for $g \in G$ and all $s \in S$ then g is semisimple;
- (3) $\alpha(S) \neq 1$ for every root α of G .

The following theorem proved in [24] is of fundamental significance for this paper, this is used many times, often without an explicit reference.

Theorem 2.10. [24, Theorem 2] *Let G be a simply connected simple algebraic group and ϕ a non-trivial irreducible representation of G . Then the following statements are equivalent:*

- (1) there exists a non-central semisimple element $s \in G$ such that at most one eigenvalue multiplicity of $\phi(s)$ exceeds 1;
- (2) All non-zero weights of ϕ are of multiplicity 1.

Recall the notation $e(G)$, which denotes the maximum of the squares of the ratios of the lengths of the roots in $\Phi(G)$. We will rely on the following important result [17, Theorem 1]:

Theorem 2.11. *Assume $p = 0$ or $p > e(G)$. Let λ be a p -restricted dominant weight. Then $\Omega(V_\lambda) = \{w(\mu) \mid \mu \in \Omega^+, \mu \preceq \lambda, w \in W\}$.*

An application of Theorem 2.11 then gives:

Lemma 2.12. [25, Lemma 2.6] *Assume $p = 0$ or $p > e(G)$. Let $\lambda, \mu \in \Omega^+$ with λ p -restricted, and V_λ , respectively, V_μ the associated irreducible FG -modules. Then the following hold.*

- (1) If $\mu \prec \lambda$ then $\Omega(V_\mu) \subseteq \Omega(V_\lambda)$.
- (2) If $\lambda + \mu$ is p -restricted then $\Omega(V_{\lambda+\mu}) = \Omega(V_\lambda \otimes V_\mu) = \Omega(V_\lambda) + \Omega(V_\mu)$.
- (3) If λ is radical then some root is a weight of V_λ ; otherwise $\Omega(V_\lambda)$ contains some minuscule weight.

Lemma 2.13. *Let V be a non-trivial G -module. Let $T' \subset T$ be the subset of all $t \in T$ that separate the weights of V .*

(1) *T' is nonempty Zarisky open in T .*

(2) *Suppose that at most one weight of V has multiplicity greater than 1. Then all elements of T' are almost cyclic on V .*

Proof. (1) Let μ, ν be weights of V . Then $T_{\mu, \nu} := \{x \in T \mid \mu(x) = \nu(x)\}$ is a Zarisky closed subset $T_{\mu, \nu}$ of T . The set of elements of T that do not separate some pair of weights of V , being the finite union of all $T_{\mu, \nu}$ with $\mu \neq \nu$, is a proper closed subset of T . So $T' = T \setminus (\cup T_{\mu, \nu})$ is nonempty open.

(2) If $s \in T'$ then $\mu(s) \neq \nu(s)$ whenever $\mu \neq \nu$ are weights of V . Therefore, the eigenvalues of s on V are exactly $\mu(s)$, where μ runs over the weights of V , and the multiplicity of $\mu(s)$ equals that of μ , whence the claim. \square

We often use the following lemma with no explicit reference.

Lemma 2.14. (1) *Let V, V_1 be FG -modules (for some group G) that are Galois conjugates of each other. Let $s \in G$ be a p' -element. Then $m_s(V) = m_s(V_1)$.*

(2) *Let G be an algebraic group and $s \in G$ a semisimple element. Let V be a G -module and V_1 a Frobenius twist of V . Then $\dim V^s = \dim V_1^s$ and $m_s(V) = m_s(V_1)$.*

Proof. (1) is obvious. (2) A Frobenius twist, as it is defined in *Notation*, arises from a so called standard Frobenius mapping $Fr : G \rightarrow G$; the latter can be defined in terms of the mapping $h_k : (a_{ij}) \rightarrow (a_{ij}^{p^k})$ for every matrix $(a_{ij}) \in GL_n(F)$ as follows, see [5, §1.17]. A mapping $Fr : G \rightarrow G$ is called standard Frobenius if there exists a faithful rational representation $\rho : G \rightarrow GL_n(F)$ such that $\rho(Fr(g)) = h_k(\rho(g))$ for every $g \in G$. If (a_{ij}) is a diagonal matrix then $h_k : (a_{ij}) = (a_{ij})^{p^k}$. As s is semisimple, so is $\rho(s)$, and hence $x\rho(s)x^{-1}$ is diagonal for some $x \in GL_n(F)$. So $h_k(x\rho(s)x^{-1}) = h_k(x)\rho(Fr(g))h_k(x)^{-1} = (x\rho(s)x^{-1})^{p^k} = x\rho(s^{p^k})x^{-1}$, whence $\rho(Fr(s))$ is a conjugate of $\rho(s^{p^k}) = \rho(s)^{p^k}$ for every representation ρ . In particular, this holds for the representation afforded by V . Since $\rho(s)$ is semisimple, the eigenvalue 1 multiplicity of $\rho(s)^{p^k}$ coincides with that of $\rho(s)$. If e_1, e_2 are distinct eigenvalues of $\rho(s)$ then $e_1^{p^k}, e_2^{p^k}$ are distinct eigenvalues of $\rho(s)^{p^k}$. So the result follows. \square

Note that we cannot drop "Frobenius" from the assumption in (2); indeed, let G be a simple algebraic group of type D_4 , and the highest weights of V, V' are ω_1, ω_4 , respectively. Then V' is a twist of V (by a graph automorphism of order 3) but, if $p \neq 3, 5$ then $m_s(V) = 2, m_s(V') = 0$ for a suitable element of order 15.

Lemma 2.15. *Let G be a simple algebraic group in characteristic p and V a non-trivial p -restricted irreducible G -module. Let $p \notin e(G)$. If $0 \in \Omega(V)$ then one of the following holds:*

- (1) $\Phi(G) \subseteq \Omega(V)$;
- (2) V is the short root module;
- (3) G is of type C_n and $\omega = \omega_i$ for $i > 2$ even.

Proof. If all root lengths are the same then the result is known, see for instance [21, Prop 2.3]. So we are left with groups B_n, C_n, F_4, G_2 and we have to show that the maximal root ω_a lies in $\Omega(V)$. We proceed case-by-case. Let $\omega = \sum a_i \omega_i$ be the highest weight of V .

Let G be of type B_n . Then $2\omega_1 \succ \omega_a = \omega_2$ and $2\omega_n \succ \omega_{n-1} \succ \omega_{n-2} \succ \cdots \succ \omega_2 \succ \omega_1$, see [3, Planche II]. As $\omega \succ 0$, we have a_n to be even. Therefore, $\omega \succeq (a_1 + \cdots + a_{n-1} + \frac{a_n}{2})\omega_1$. If $a_1 + \cdots + a_{n-1} + \frac{a_n}{2} > 1$ then $\omega \succeq 2\omega_1 \succ \omega_2$, as desired. If $a_1 + \cdots + a_{n-1} + \frac{a_n}{2} = 1$ and $\omega \neq \omega_1$ then $\omega \in \{\omega_2, \dots, \omega_{n-1}, 2\omega_n\}$. So the result again follows.

Let G be of type C_n . Then $\omega_a = 2\omega_1$, $\omega_i \succeq \omega_2$ for i even, and $\omega_i \succeq \omega_1$ for i odd. Therefore, $\omega \succeq (\sum_{i \text{ odd}} a_i)\omega_1 + (\sum_{i \text{ even}} a_i)\omega_2$. Here $\sum_{i \text{ odd}} a_i$ is even as $\omega \succ 0$; if this is non-zero then the result follows. Otherwise, since $2\omega_2 \succ 2\omega_1$, the result again follows unless $\omega = \omega_i$ for i even. Moreover, $2\omega_1 = \omega_a$ is not a subdominant weight of ω_i as one can conclude by comparing the α_1 -coefficients of the expansions of ω_i 's in terms of simple roots in [3, Planche III].

Let G be of type F_4 . Then $\omega_2 \succ \omega_3 \succ \omega_1 = \omega_a \succ \omega_4$ and $2\omega_4 \succ \omega_1$, see [3, Planche XIII]. So $\omega \succeq (a_1 + a_2 + a_3 + a_4)\omega_4 \succeq 2\omega_4 \succ \omega_1$, unless $a_1 + a_2 + a_3 + a_4 = 1$. If $a_1 + a_2 + a_3 > 0$ then the result follows, otherwise $a_1 = a_2 = a_3 = 0$, $a_4 = 1$, that is, $\omega = \omega_4$ is a short root.

Let G be of type G_2 . Then $2\omega_1 \succ \omega_a = \omega_2 \succ \omega_1$ so $\omega \succeq (a_1 + a_2)\omega_1$. If $a_2 > 0$ or $a_2 = 0, a_1 \geq 2$ then the result follows, otherwise $\omega = \omega_1$, which is a short root. \square

Lemma 2.16. *Let G be a simple algebraic group and V a G -module. Let $s \in T_{reg}$. Suppose that $\mu(s) = \nu(s)$ for distinct weights $\mu, \nu \in \Omega(V)$ and $\mu - \alpha, \nu - \alpha \in \Omega(V)$ for some $\alpha \in \Phi(G)$. Then s is not almost cyclic on V .*

Proof. Suppose the contrary. Observe that $\mu(s) = \nu(s)$ implies $(\mu - \alpha)(s) = (\nu - \alpha)(s)$. If s is almost cyclic on V then $(\mu - \alpha)(s) = \mu(s)$, whence $\alpha(s) = 1$. By Lemma 2.9(3), this is a contradiction as s is regular. \square

Lemma 2.17. *Let $s \in G = SL_n(F)$ be a strictly regular element. Then s is cyclic on V_{ω_2} .*

Proof. The roots of G are $\varepsilon_i - \varepsilon_j$ with $i \neq j$, $i, j \in \{1, \dots, n\}$. Let $s = \text{diag}(a_1, \dots, a_n)$. As s is strictly regular, $(\varepsilon_i - \varepsilon_j)(s) \neq (\varepsilon_k - \varepsilon_l)(s)$, equivalently, $(\varepsilon_i + \varepsilon_l)(s) \neq (\varepsilon_j + \varepsilon_k)(s)$, whenever $i \neq j, k \neq l$ and $(i, j) \neq (k, l)$.

The weights of V_{ω_2} are $\varepsilon_i + \varepsilon_l$ for $i < l$, so if s is not cyclic on V_{ω_2} then $(\varepsilon_i + \varepsilon_l)(s) = (\varepsilon_j + \varepsilon_k)(s)$ for some $i < l, j < k$, and hence $(\varepsilon_i - \varepsilon_j)(s) \neq (\varepsilon_k - \varepsilon_l)(s)$. If $i = j, k \neq l$ here then $(\varepsilon_i - \varepsilon_j)(s) = 1$ so $(\varepsilon_k - \varepsilon_l)(s) = 1$, so s is not regular, a contradiction. Similarly, $k \neq l$. \square

For reader's convenience we quote here [24, Theorem 2]:

Theorem 2.18. *Let G be a simply connected simple algebraic group defined over a field of characteristic $p \geq 0$ and let $s \in G$ be a non-regular non-central semisimple element. Let V be a non-trivial irreducible G -module. Then either s is not almost cyclic on V or one of the following holds:*

- (1) G is classical, that is, of Lie type A_n, B_n ($p \neq 2$), C_n or D_n and $\dim V = n+1, 2n+1, 2n, 2n$, respectively;

- (2) $G = B_n, p = 2$ and $\dim V = 2n$;
- (3) $G = A_3$ and $\dim V = 6$;
- (4) $G = C_2 \cong B_2$ and $\dim V = 5$.

3 Eigenvalue multiplicities: a proof of Theorem 1.2

In this section we bound the eigenvalue multiplicities of a regular semisimple element $s \in G$ on a G -module V provided s is almost cyclic on V . Of course, in this case there is at most one eigenvalue of multiplicity greater than 1, so we bound the multiplicity of this single eigenvalue.

3.1 A uniform bound

Lemma 3.1. *Let G be a simple algebraic group in characteristic $p \notin e(G)$ and let $0 \neq \omega \in \Omega^+$. If $p > 0$, suppose that $\omega \neq p\omega'$ for some $\omega' \in \Omega^+$. Let $V = V_\omega$ and $\lambda \in \Omega(V)$. Then there is $\alpha \in \{\pm\Pi\}$ such that $\lambda - \alpha \in \Omega(V)$.*

Proof. If $p = 0$ then the result is true by [4, Ch. VIII, §7, Prop. 5]. If $p \neq e(G)$ and ω is p -restricted then the result follows from that for $p = 0$ and Lemma 2.12. Suppose that ω is not p -restricted. Then $V = M \otimes V'$, where M, V' are irreducible G -modules and M is a p -restricted. Every weight of V is of shape $\mu + \nu$ with $\mu \in \Omega(M)$ and $\nu \in \Omega(V')$. By the above $\mu - \alpha \in \Omega(V)$ for some $\alpha \in \{\pm\Pi\}$. So $\lambda - \alpha = (\mu - \alpha) + \nu \in \Omega(V)$, as required. \square

Theorem 3.2. *Assume $p = 0$ or $p > e(G)$. Let $S \subset T$ be a subgroup such that $C_G(S) = T$, and V an irreducible G -module with p -restricted highest weight $\omega \neq 0$. Assume in addition that some homogeneous component U of $V|_S$ is of dimension greater than 1 and all other homogeneous component of $V|_S$ are 1-dimensional. Then*

- (1) $V^T \neq 0$, then $V^S = V^T$ and $\dim V^S \leq n + 1$;
- (2) $\dim U \leq 2n$.

Proof. By [24, Theorem 5], all non-zero weights of V are of multiplicity 1. Therefore, ω appears in Tables 1,2, and in particular $\dim V^T \leq n + 1$. By Lemma 2.9 we have $\alpha(S) \neq 1$ for every $\alpha \in \Phi(G)$.

(1) Here we have $V^T \subseteq V^S$. Suppose the contrary, that $V^T \neq V^S$, so $\dim V^S > 1$. Note that V^S is a sum of weight spaces of V , so $\mu(S) = 1$ for some non-zero weight of V .

Suppose first that ω is a root. Then all non-zero weights of V are also roots. (This is well known.) Since $\alpha(S) \neq 1$ for every $\alpha \in \Phi(G)$, we conclude that $V^S = V^T$, a contradiction.

Next suppose that ω is not a root. By the above, μ is not a root. By replacing S by a conjugate if necessary, we may assume that μ is dominant. Let $\beta \in \{\pm\Pi\}$ be such that $\mu + \beta$ is also a weight of V . Then $\mu + \beta \neq \beta$, while $\mu + \beta$ and β restricts to the same (non-zero) character of S . So $\beta(S) = 1$; by the above, β is not a weight of V . But $0 \prec \omega$ so Lemma 2.12(3) implies that some root is a weight of V . We deduce that $\Phi(G)$ has two root lengths and either the long roots or the short roots are not weights of V . As $p \notin e(G)$,

by Lemma 2.15, either ω is a short root, a contradiction, or $G = C_n$ and $\omega = \omega_i$ with i even. In the latter case $\omega_j \in \Omega(V)$ for $i - j > 0$ even, and these are the only dominant weights of $\Omega(V)$. So $\mu = \omega_r$ with $i - r > 0$ even. Here $r > 2$ as ω_2 is a root. Then $r - 2 \neq 0$ and $\omega_r - \omega_{r-2} = \varepsilon_r + \varepsilon_{r-1} \in \Phi(G)$. Set $\beta = -\varepsilon_r - \varepsilon_{r-1}$. As β is a short root, it lies in the W -orbit of $\omega_2 = \varepsilon_1 + \varepsilon_2$, and hence $\beta \in \Omega(V)$. However, by the above, we have a contradiction.

(2) In view of (1) we can assume that $\dim V^T \leq 1$, so all weights of V . If $U = V^S$ then the claim follows from (1). So we now assume that $U \cap V^S = 0$ hence U is the sum of T -weight spaces which are of dimension 1 by Theorem 2.10. Assume for a contradiction that there exist distinct T -weights μ_1, \dots, μ_{2n+1} such that $\mu_j|_S = \eta$ for some character η of S and all $1 \leq j \leq 2n+1$. For each j there exists $\gamma_j \in \Pi^\pm$ such that $\mu_j + \gamma_j$ is a weight of V . Then there exists $i \neq j$ such that $\gamma_i = \gamma_j$. But then $\mu_i + \gamma_i$ and $\mu_j + \gamma_j$ are distinct weights of V and have equal restrictions to S . By the hypothesis, this then means that $\mu_i|_S = \eta|_S = (\mu_j + \gamma_j)|_S$, whence $\gamma_i(S) = 1$, a contradiction. \square

3.2 $G \neq A_n$: Special cases

A more detailed case-by-case analysis allows us to improve the bound $\dim V_S(\eta) \leq 2n$ in Theorem 3.2 to $\dim V_S(\eta) \leq n+1$. We prefer to assume $S = \langle s \rangle$ to be cyclic, the general case is similar. We first perform this for groups $G \neq A_n$, and then consider the case of $G = A_n, n > 1$, which turns out to be more complex. We denote by $m_s(V)$ the maximum of eigenvalue multiplicities of $s \in G$ on V . In this section we obtain a sharp bound $m_s(V) \leq 2$ in a number of special cases.

Recall (Theorem 2.10) that the non-zero weights of an irreducible G -module V are of multiplicity 1 whenever a semisimple element $s \in G$ is almost cyclic on V . Such modules are classified in [22] and the highest weights of V , for V p -restricted, are reproduced in Tables 1,2 at the end of this papers.

Lemma 3.3. *Let $G = B_n, p \neq 2$, or $C_n, p = 2$, and let $s \in T_{reg}$. Suppose that s is almost cyclic on V_{ω_n} . Then $m_s(V_{\omega_n}) \leq 2$.*

Proof. The weights of V are $\frac{1}{2}(a_1\varepsilon_1 + \dots + a_n\varepsilon_n)$ for $G = B_n$ and $a_1\varepsilon_1 + \dots + a_n\varepsilon_n$ for $G = C_n$, where $a_1, \dots, a_n \in \{\pm 1\}$. Let $\mu, \nu \in \Omega(V)$ be distinct weights such that $\mu(s) = \nu(s)$. Note that $\pm\varepsilon_i$, respectively, $\pm 2\varepsilon_i$ for $i = 1, \dots, n$ are roots of G of type B_n , respectively, C_n . If $\mu \neq -\nu$ then, obviously, there is i such that $\mu - \alpha, \nu - \alpha \in \Omega(V)$ for $\alpha \in \{\pm\varepsilon_i : 1 \leq i \leq n\}$, respectively, $\alpha \in \{\pm 2\varepsilon_i : 1 \leq i \leq n\}$. This contradicts Lemma 2.16.

Now suppose the contrary, and let $m_s(V_{\omega_n}) > 2$. Then there are distinct weights λ, μ, ν of V such that $\lambda(s) = \mu(s) = \nu(s)$. By the above, $\lambda = -\mu, \lambda = -\nu$ and $\nu = -\mu$. This is a contradiction. \square

Lemma 3.4. *Let $G = B_n, n > 2, p \neq 2$, or $C_n, p = 2$, and let V be a non-trivial irreducible p -restricted G -module of highest weight ω . Let $s \in T_{reg}$. Suppose that s is almost cyclic on V . Then either $\omega \in \{\omega_1, \omega_n\}$ and $m_s(V) \leq 2$ or $\omega = \omega_2$ and $n - 1 \leq m_s(V) \leq n$ or $G = B_n, \omega = 2\omega_1$ and $n \leq m_s(V) \leq n + 1$.*

Proof. By Theorem 2.10, all non-zero weights of V are of multiplicity 1, and hence, by Theorem 2.10 and Tables 1,2, $\omega \in \{\omega_1, \omega_n, \omega_2, 2\omega_1\}$, the latter for $p \neq 2$. For $\omega = \omega_1$ see Lemma 2.18, for $\omega = \omega_n$ see Lemma 3.3. If $\omega = \omega_2$ or $2\omega_1$ with $p \neq 2$ then $\dim V^T \geq 2$ by Table 1. Then $m_s(V) = \dim V^s$, and $V^T = V^s$ by Theorem 3.2. So the result follows from Table 1. \square

The case $G = B_2 \cong C_2$, $p \neq 2$, will be considered later in Lemma 3.12.

Lemma 3.5. *Let G be an algebraic group of type D_n , $n > 3$, and let $V = V_\omega$, where $\omega \in \{\omega_1, \omega_{n-1}, \omega_n\}$. Let $\mu, \nu \in \Omega(V)$. Then $\mu - \alpha, \nu - \alpha \in \Omega(V)$ for some root α of G , unless n is even and $\mu = -\nu$, or n is odd and $\mu = -\nu \pm \varepsilon_i$ for some $i \in \{1, \dots, n\}$.*

Proof. (i) Suppose that $\omega = \omega_1$. Then the weights of V are $\pm\varepsilon_1, \dots, \pm\varepsilon_n$. We can assume that $\mu = \varepsilon_i$ for $i \in \{1, \dots, n\}$. If $\nu = \varepsilon_j$ ($1 \leq j \leq n, j \neq i$) then set $\alpha = \varepsilon_i + \varepsilon_j$. If $i = j$ then set $\alpha = \varepsilon_i + \varepsilon_j$ for any $j \neq i$. If $\nu = -\varepsilon_j$, $j \neq i$, then set $\alpha = \varepsilon_i - \varepsilon_j$. One can check that $\mu - \alpha, \nu - \alpha \in \Omega(V)$ in each case.

(ii) Suppose that $\omega = \omega_n$ or ω_{n-1} . Then the weights of V are $\frac{1}{2}(\sum a_i \varepsilon_i)$ with $a_i \in \{1, -1\}$, and the number of indices i with $a_i = -1$ is even if $\omega = \omega_n$, otherwise it is odd. Let $\mu = \frac{1}{2}(\sum a_i \varepsilon_i)$, $\nu = \frac{1}{2}(\sum b_i \varepsilon_i)$ be weights of V , and let $L = \{i : a_i = b_i\}$. If $|L| > 1$ then $\alpha = a_i \varepsilon_i + a_j \varepsilon_j \in \Phi(G)$ for $i, j \in L, i \neq j$ and $\mu - \alpha, \nu - \alpha \in \Phi(G)$, as required.

Suppose that $|L| = 1$. Let $i \in L, j \notin L, 1 \leq j \leq n$. Then $\mu + \nu = \pm\varepsilon_i$ and n is odd, as otherwise $-\nu \in \Omega(V)$ and $\mu - (-\nu) = \pm\varepsilon_i$ is not a radical weight.

Suppose that $|L| = 0$. Then $\mu = -\nu$ and n is even. \square

Corollary 3.6. *Let G be an algebraic group of type D_n , $n > 3$, and let $V = V_\omega$, where $\omega \in \{\omega_1, \omega_{n-1}, \omega_n\}$. Let $\mu_i \in \Omega(V)$ for $i = 1, 2, 3$. Then $\mu_i - \mu_j$ is a root for some $i, j \in \{1, 2, 3\}$.*

Proof. Suppose the contrary. If n is even then $\mu_i = -\mu_j$ for every pair $i, j \in \{1, 2, 3\}$ with $i \neq j$ by Lemma 3.5. This is obviously false. Suppose that n is odd. Then, by Lemma 3.5, $\mu_i + \mu_j = \pm\varepsilon_{k(i,j)}$ with $k(i, j) \in \{1, \dots, n\}$ for every pair $i, j \in \{1, 2, 3\}$ with $i \neq j$. If $k(1, 2) \neq k(1, 3)$ then $\mu_3 - \mu_2 = (\mu_1 - \mu_2) - (\mu_1 - \mu_3) = \pm\varepsilon_{k(1,2)} \mp \varepsilon_{k(1,3)}$ is a root, a contradiction. So $k(1, 2) = k(1, 3)$, and similarly, $k(1, 2) = k(2, 3)$. This means that $\mu_i + \mu_j = a(i, j)\varepsilon_k$ for some fixed $k \in \{1, \dots, n\}$ and $a(i, j) \in \{1, -1\}$ for every pair $i, j \in \{1, 2, 3\}$ with $i \neq j$. By reordering of μ_1, μ_2, μ_3 we can assume that $\mu_1 + \mu_2 = \mu_1 + \mu_3$, whence $\mu_2 = \mu_3$, a contradiction. \square

Lemma 3.7. *Let $G = D_n$, $n > 3$, let $s \in T_{reg}$. Suppose that s is almost cyclic on $V = V_{\omega_i}$ for $i \in \{n-1, n\}$. Then $m_s(V) \leq 2$.*

Proof. Suppose the contrary, and let μ_1, μ_2, μ_3 be distinct weights of V such that $\mu_1(s) = \mu_2(s) = \mu_3(s)$. By Corollary 3.6, for some $i, j \in \{1, 2, 3\}$ we have $\mu_i - \mu_j = \alpha$, where α is a root. Then $\alpha(s) = 1$, which is a contradiction as s is regular. \square

Lemma 3.8. *Let $G = D_n, n > 3$, let M be an irreducible G -module with highest weight $w \in \{\omega_1, \omega_n, \omega_{n-1}\}$ and let V an arbitrary irreducible G -module. Let $s \in T_{reg}$. Suppose that s is almost cyclic on $M \otimes V$. Then $m_s(M \otimes V) \leq 2$.*

Proof. Suppose the contrary, and let $\lambda_i, i = 1, 2, 3$ be distinct weights of $M \otimes V$ such that $\lambda_i(s) = e \in E(M \otimes V)$. Then $\lambda_i = \mu_i + \nu_i$, where μ_i, ν_i are weights of M, V , respectively. By Lemma 2.3(2), s is cyclic on M and V . By Corollary 3.6, there exists a root α of G such that $\mu_i - \alpha, \mu_j - \alpha \in \Omega(M)$ for some $i, j \in \{1, 2, 3\}$. Then $\mu_i - \alpha + \nu_i, \mu_j - \alpha + \nu_j \in \Omega(V)$, and $(\mu_i - \alpha + \nu_i)(s) = (\mu_j - \alpha + \nu_j)(s) = e \cdot \alpha(s)^{-1}$. As s is almost cyclic on V , we have $\alpha(s) = 1$, which is a contradiction as s is regular. \square

Lemma 3.9. *Let $G = C_3, p \neq 2$. Let $s \in T_{reg}$. Suppose that s is almost cyclic on $V = V_{\omega_3}$. Then $m_s(V) \leq 2$.*

Proof. Note that every weight of V has multiplicity 1 (see Table 2) and $\Omega(V) = X \cup Y$, where $X = \{\pm\varepsilon_1, \pm\varepsilon_2, \pm\varepsilon_3\}$ and $Y = \{\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3\}$. Suppose the contrary, and let $\mu_i, i = 1, 2, 3$, be distinct weights of V such that $\mu_i(s) = e$. By Lemma 2.1, $e \in \{\pm 1\}$. If $\mu_i \in X$ for some i then $(-\mu_i)(s) = e$, which is false as $X = \Omega(V_{\omega_1})$ and then s is not regular by Lemma 2.7(1). So $\mu_i \in Y$ for every $i = 1, 2, 3$. Then $\mu_i = a_{i1}\varepsilon_1 + a_{i2}\varepsilon_2 + a_{i3}\varepsilon_3$ for $a_{ik} \in \{\pm 1\}$ for $1 \leq i, k \leq 3$. By reordering μ_i we can assume $a_{11} = a_{21}$, and then $\mu_1 - \alpha, \mu_2 - \alpha \in \Omega(V)$ for $\alpha \in \{\pm 2\varepsilon_1\}$. This contradicts Lemma 2.16. \square

Lemma 3.10. *Let $G = C_n, n > 2, p = 3$, and $\omega \in \{\omega_{n-1}, \omega_n\}$. Let $s \in T_{reg}$ be almost cyclic on V_ω . Then $m_s(V) \leq 2$.*

Proof. The non-zero weights of V are $\pm\varepsilon_{i_1} \pm \dots \pm \varepsilon_{i_r}$, for $r \in \{1, \dots, n\}$, where $r - n$ is odd if $\omega = \omega_{n-1}$, and even if $\omega = \omega_n$.

Suppose the contrary, let $m_s(V) > 2$ and let $\lambda, \mu \in \Omega(V)$, $\lambda(s) = \mu(s) = e$, $\lambda \neq \mu$, where we can assume that $\lambda \neq -\mu$. By Lemma 2.1, $e \in \{\pm 1\}$. Set $\lambda = \sum a_i \varepsilon_i$, $\mu = \sum b_i \varepsilon_i$, where $a_i, b_i \in \{0, 1, -1\}$.

Observe that λ, μ are non-zero. Indeed, otherwise, $\lambda(s) = \mu(s) = 1$ so $\dim V^s \geq 2$. Then $m_s(V) = \dim V^s$ and $\dim V^T = \dim V^s$ by Theorem 3.2(1). However, $\dim V^T = 1$, which is a contradiction.

If $a_i = b_i \neq 0$ for some i then $\lambda - 2a_i \varepsilon_i, \mu - 2a_i \varepsilon_i \in \Omega(V)$ and $(\lambda - 2a_i \varepsilon_i)(s) = (\mu - 2a_i \varepsilon_i)(s)$. As $2a_i \varepsilon_i$ is a root, we have a contradiction by Lemma 2.16.

If $a_i = -b_i \neq 0$ for some i then we replace λ by $-\lambda$; as $e \in \{\pm 1\}$, we have $(-\lambda)(s) = \lambda(s)$, so we arrive at the case considered above.

So $a_i b_i = 0$ for every $i = 1, \dots, n$. Using the Weyl group, we can assume that $\lambda = \varepsilon_1 + \dots + \varepsilon_{r-1} - \varepsilon_r$, $\mu = \varepsilon_{r+1} + \dots + \varepsilon_{r'}$ for some $0 < r < r' \leq n$. Then $\lambda + (\varepsilon_r - \varepsilon_{r+1}), \mu + (\varepsilon_r - \varepsilon_{r+1}) \in \Omega(V)$, and $(\lambda + (\varepsilon_r - \varepsilon_{r+1}))(s) = (\mu + (\varepsilon_r - \varepsilon_{r+1}))(s)$. As above, this contradicts Lemma 2.16. \square

Lemma 3.11. *Let $G = C_n, n \geq 2, p > 3$, and $\omega \in \{\omega_{n-1} + \frac{p-3}{2}\omega_n, \frac{p-1}{2}\omega_n\}$. Let $s \in T_{reg}$ be almost cyclic on V_ω . Then $m_s(V_\omega) \leq 2$.*

Proof. The weights of V_ω are $l_1\varepsilon_1 + \cdots + l_n\varepsilon_n$, where $-(p-1)/2 \leq l_1, \dots, l_n \leq (p-1)/2$, see [26, proof of Lemma 1.6]; moreover, if μ is of this form then μ is a weight of V_ω with $\omega \in \{\omega_{n-1} + \frac{p-3}{2}\omega_n, \frac{p-1}{2}\omega_n\}$. Let λ, μ, ν be three distinct weights of V_ω such that $\lambda(s) = \mu(s) = \nu(s)$. As $(-\lambda)(s) = (-\mu)(s)$, we have $\lambda(s) = \mu(s) = \nu(s) \in \{\pm 1\}$. Let

$$\lambda = l_1\varepsilon_1 + \cdots + l_n\varepsilon_n, \quad \mu = m_1\varepsilon_1 + \cdots + m_n\varepsilon_n, \quad \nu = k_1\varepsilon_1 + \cdots + k_n\varepsilon_n.$$

By Lemma 2.16, either $\lambda - 2\varepsilon_i$ or $\mu - 2\varepsilon_i$ is not a weight of V_ω for every $i = 1, \dots, n$. Observe that $\lambda - 2\varepsilon_i$, respectively, $\mu - 2\varepsilon_i$ is a weight of $\Omega(V_\omega)$ unless $l_i \leq 1 - \frac{p-1}{2}$, respectively, or $m_i \leq 1 - \frac{p-1}{2}$, and similarly for the pairs λ, ν and μ, ν . So there are at least two pairs with i -th coefficients at most $1 - \frac{p-1}{2}$. By reordering λ, μ, ν we can assume that $l_i, m_i \leq 1 - \frac{p-1}{2}$. Then $-\lambda, -\mu$ are weights of V_ω , $(-\lambda)(s) = (-\mu)(s)$ and $-l_i, -m_i \geq \frac{p-1}{2} - 1$. Then the i -th coefficient of $-\lambda - 2\varepsilon_i$ equals $-l_i - 2 \geq \frac{p-1}{2} - 3 \geq -\frac{p-1}{2}$ for $p > 3$. So $-\lambda - 2\varepsilon_i \in \Omega(V_\omega)$; similarly, $-\mu - 2\varepsilon_i \in \Omega(V_\omega)$. Then Lemma 2.16 yields a contradiction. \square

Lemma 3.12. *Let $G = B_2 \cong C_2$, $s \in T_{reg}$ and let V be a non-trivial irreducible p -restricted G -module of highest weight ω . If s is almost cyclic on V then $m_s(V) \leq 2$.*

Proof. Let $G = C_2$. As s is almost cyclic on V , ω occurs in Tables 1,2. For those in Table 1 we have $\dim V^T = 2$, and hence $m_s(V) = \dim V^s = \dim V^T$ by Theorem 3.2; then $\dim V^T = 2$ by Table 1. Those in Table 2 are ω_1, ω_2 and, for $p > 3$, also $\omega_1 + \frac{p-3}{2}\omega_2, \frac{p-1}{2}\omega_2$. In the two last cases the result is contained in Lemma 3.11. The module V_{ω_1} is the natural module for G , and $m_s(V) = 0$ by Lemma 2.7(1). The module V_{ω_2} is the natural module for $B_2 \cong C_2$, so the result follows from Lemma 2.7(2). \square

Remark 3.13. If $\dim V = 4$ then $m_s(V) = 0$. Indeed, if $G = C_2$ then either V is a Frobenius twist of the natural module or $p = 2$ and V is a Frobenius twist of V_{ω_2} . The former case is ruled out as s is regular (see Lemma 2.7), the latter case can be checked straightforwardly. (Say, let $s = \text{diag}(a, b, b^{-1}, a^{-1}) \in Sp_4(q)$, where the diagonal entries are distinct; then the eigenvalues of s on V_{ω_2} are $a^{\pm 1}b^{\pm 1}$; if $m_s(V) > 0$ then $a^i b^j = 1$ for some $i, j \in \{\pm 1\}$, which is a contradiction.) If $G = B_2$ then $\omega = p^k\omega_2$; as V_{ω_2} is the natural module for C_2 , it follows that $m_s(V) = 0$ for a regular element $s \in G$.




Figure 1: Weight diagram of V_{ω_1} for $G = E_6$

Lemma 3.14. *Let $G = E_6$ and let $\mu_1, \mu_2, \mu_3, \mu_4$ be distinct weights of V_{ω_1} . Then $\mu_i - \mu_j$ is a root for some $i, j \in \{1, 2, 3, 4\}$, $i \neq j$.*

Proof. All weights of V_{ω_1} lie in the Weyl group orbit of ω_1 , so we can assume that $\mu_1 = \omega_1$. One observes that there are exactly 15 roots $\alpha \in \Phi(G)$ such that $\omega_1 - \alpha \in \Omega(V_{\omega_1})$. (This can be easily deduced from the weight diagram of V_{ω_1} borrowed from [16, Figure 20] using the list of the roots provided in [3]. In fact, these 15 weights are located at the left part of the diagram.)

Suppose the contrary. Then μ_2, μ_3, μ_4 are located at the remaining part of the diagram. Then one easily checks that $\mu_i - \mu_j$ is a root for some $i, j \in \{2, 3, 4\}$. So the result follows. \square

Lemma 3.15. *Let $G = E_6$, $V = V_{\omega_1}$ and let M be an irreducible G -module. Let $s \in T_{reg}$. Suppose that s is almost cyclic on $M \otimes V$. Then $m_s(M \otimes V) \leq 3$.*

Proof. Suppose the contrary, and let e be an eigenvalue of s on V of multiplicity at least 4.

(i) Suppose first that M is trivial. Let μ_i , $i = 1, 2, 3, 4$, be distinct weights of V such that $\mu_i(s) = e$ for every i . By Lemma 3.14, there are distinct $i, j \in \{1, 2, 3, 4\}$ and $\alpha \in \Phi(G)$ such that $\mu_i - \alpha, \mu_j - \alpha \in \Omega(V_{\omega_1})$. This contradicts Lemma 2.16.

(ii) Suppose that M is not trivial. By Lemma 2.3(2), s is cyclic on M and V . Then $e = a_i b_i$ for $i = 1, 2, 3, 4$, where a_1, a_2, a_3, a_4 are distinct eigenvalues of s on V and b_1, b_2, b_3, b_4 are distinct eigenvalues of s on M . We have $a_i = \mu_i(s)$ for some distinct weights $\mu_1, \mu_2, \mu_3, \mu_4$ of V . As above, there is a root $\alpha \in \Phi$ and $i, j \in \{1, 2, 3, 4\}$, $i \neq j$, such that $\mu_i - \alpha, \mu_j - \alpha \in \Omega(V_{\omega_1})$. Then $c_i = (\mu_i - \alpha)(s)$ and $c_j := (\mu_j - \alpha)(s)$ are eigenvalues of s on $V = V_{\omega_1}$, which are distinct as s is cyclic on V . In addition, $c_i \neq a_i$, $c_j \neq a_j$ as otherwise $\alpha(s) = 1$, which is false as s is regular. So $c_i b_i = c_j b_j \neq a_i b_i = a_j b_j$. This is a contradiction as s is almost cyclic on $M \otimes V$. \square

The following example shows that the bound in Lemma 3.15 attains.

Example 3.16. Let $G = E_6$. Then G contains a subgroup H , say, of type F_4 . Let $\eta_1, \eta_2, \eta_3, \eta_4$ be the fundamental weights of F_4 . Let V be an irreducible G -module of highest weight ω_1 . It is well known that $V|_H$ has a composition factor U of highest weight η_4 whose dimension is $\dim V - 1$ or $\dim V - 2$, the latter for $p = 3$. Then the other composition factors of $V_{\omega_1}|_H$ are trivial. Let $s \in F_4$ be a regular semisimple element in general position, that is, the cyclic group $\langle s \rangle$ is dense in a maximal torus S , say, of F_4 . Then s is regular and separates the weights of U , and hence s is almost cyclic on V . In addition, $\dim U^s = \dim V^s = 2$ or 1 , the latter for $p = 3$. It follows that $V^s = 3$.

Lemma 3.17. *Let $G = E_6$, let V be an irreducible G -module and $s \in T_{reg}$. Suppose that s is almost cyclic on V . Then $m_s(V) \leq 3$ unless the highest weight of V is $p^t \omega_2$.*

Proof. Let ω be the highest weight of V . Recall that the non-zero weights of V must be of multiplicity 1 (Theorem 2.10) and s is regular by Theorem 2.18. Therefore, either $\omega = p^t \omega'$, where $\omega' \in \{\omega_1, \omega_6, \omega_2\}$ by [22, Theorem 2] or $V = V' \otimes M$, where M is a non-trivial irreducible G -module and the highest weight of V' is $p^t \omega_1$ or $p^t \omega_6$ (here $\omega' \neq \omega_2$

by Lemma 2.3(2)). So the result follows from Lemma 3.15. (If $\omega' = \omega_2$ then $m_s(V) = 6$ by Theorem 3.2(1) and the fact that V_{ω_2} is the adjoint G -module.) \square

Lemma 3.18. *Let $G = E_7$, $V = V_{\omega_7}$ and $s \in T_{reg}$. Suppose that s is almost cyclic on V . Then $m_s(V) \leq 2$.*

Proof. Let H be a subgroup of G isomorphic to a simple algebraic group of type A_7 , so the maximal tori of H are maximal in G . Hence we can assume that $T \subset H$. Let η_1, \dots, η_7 be the fundamental weights of H . Then the restriction $V|_H$ is $V_{\eta_2} + V_{\eta_6}$. The result follows from the following lemma: \square

Lemma 3.19. *Let H be a simple algebraic group of type A_{n-1} , $n \geq 4$. Let $U := V_{\eta_2} \oplus V_{\eta_{n-2}}$. Let $s \in T_{reg}$. Suppose that s is almost cyclic on U . Then $m_s(U) \leq 2$.*

Proof. We can assume that $g = \text{diag}(a_1, \dots, a_n) \in SL_n(F)$, where $a_1, \dots, a_n \in F$ and $a_1 \cdots a_n = 1$. Then the eigenvalues of s on U are $a_i a_j, a_k^{-1} a_l^{-1}$ for $1 \leq i, j, k, l \leq n$ and $i < j, k < l$. Suppose that s is not cyclic on U . Then either $a_i a_j = a_k a_l$ for $(i, j) \neq (k, l)$ or $a_i a_j = a_k^{-1} a_l^{-1}$ for some i, j, k, l . Then $(a_i a_j)^{-1}$ is an eigenvalue of s of multiplicity at least 2, so $a_i a_j = (a_i a_j)^{-1}$ as s is almost cyclic. Then $a_i^2 = a_j^{-2}$ and $a_j = \theta a_i^{-1}$ with $\theta = 1$ or -1 . It follows that θ is an eigenvalue of s on U of multiplicity at least 2. We show that the multiplicity of θ is exactly 2. Suppose that contrary. Then $\theta = a_k a_l$ for some $k < l$ with $(k, l) \neq (i, j)$. Then $a_j a_l = a_i^{-1} a_k^{-1}$ and these are eigenvalues of s on U . As s is almost cyclic, we must have $a_j a_l = \theta = a_k a_l$, whence $a_j = a_k$. So s is not regular, a contradiction. \square

Lemma 3.20. *Let $G = E_7$, let V be an irreducible G -module and $s \in T_{reg}$. Suppose that s is almost cyclic on V . Then $m_s(V) \leq 2$ unless the highest weight of V is $p^t \omega_1$.*

Proof. Let ω be the highest weight of V . The non-zero weights of V are of multiplicity 1 (Theorem 2.10) and s is regular by Theorem 2.18. Therefore, either $\omega \in \{p^t \omega_7, p^t \omega_1\}$ by [22, Theorem 2] or $V = V' \otimes M$, where M is a non-trivial irreducible G -module and the highest weight of V' is $p^t \omega_7$ (this cannot be $p^t \omega_1$ by Lemma 2.3(2)). So the result follows from Lemma 3.18. (If $\omega = p^t \omega_2$ then $m_s(V) = 7$ by Theorem 3.2(1) and the fact that V_{ω_1} is the adjoint G -module.) \square

Lemma 3.21. *Let $G = F_4$, $p = 3$, let V be a p -restricted irreducible G -module with highest weight ω_4 and $s \in T_{reg}$. Suppose that s is almost cyclic on V . Then $m_s(V) \leq 2$.*

Proof. In this case $\dim V^T = 1$ and hence $V^s = V^T$ by Theorem 3.2. Therefore, if $m_s(V) > 1$ and e is the eigenvalue of s on V of multiplicity $m_s(V)$ then $e = -1$ by Lemma 2.1(2).

The non-zero weights of $V = V_{\omega_4}$ are short roots, which form a single orbit of the Weyl group of G . So we can assume that $\omega_4(s) = -1 = (-\omega_4)(s)$. Let $\mu \succ 0$ for $\mu \in \Omega(V)$. One can check that $\omega_4 - \mu$ is a root. If $m_s(V) > 2$ then $\mu(s) = -1 = (-\mu)(s)$ for some $0 \neq \mu \in \Omega(V)$, $\mu \neq \pm \omega_4$. So we can assume that $\mu \succ 0$. Then $\omega_4 - \mu$ is a root and $(\omega_4 - \mu)(s) = 1$, This is a contradiction as s is regular. \square

The case of $G = G_2$ is essentially treated in [25]:

Lemma 3.22. [25, Proposition 4.10] *Let $G = G_2$, $V_i = V_{\omega_i}$ for $i = 1, 2$ and let $s \in T_{reg}$.*

(1) *Suppose that s is not cyclic on V_1 . Then $p \neq 2$ and the eigenvalues of s on V_1 are $\{1, -1, -1, -b, b^{-1}, b, -b^{-1}\}$, where $b \in F, b^2 \neq 1$. In particular, s is almost cyclic on V_1 if and only if $b^2 \neq -1$. In addition, $\{\alpha(s) : \alpha \text{ a long root}\} = \{b, b^{-1}, -b, -b^{-1}, b^2, b^{-2}\}$.*

(2) *Let $p = 3$. Then s is almost cyclic on V_2 and cyclic on V_1 . More precisely, if s is not cyclic on V_2 then the eigenvalues of s on V_2 are $\{1, -1, -1, a^3, -a^3, a^{-3}, -a^{-3}\}$ for some $a \in F, a^{12} \neq 1$, and those on V_1 are $\{1, a, a^{-1}, -a, -a^{-1}, -a^2, -a^{-2}\}$.*

Corollary 3.23. *Theorem 1.2 is true for G of type G_2 .*

Proof. Suppose first that V is p -restricted. Then, by Theorem 2.10, we need to deal with FG -modules V of highest weight $\omega \in \{\omega_1, \omega_2\}$. The case $\omega = \omega_1$ is treated in Lemma 3.22(1). Let $\omega = \omega_2$. The case with $p = 3$ is examined in Lemma 3.22(2). Let $p \neq 3$. Then $\dim V^T = 2$ and hence $\dim V^s \geq 2$. If $\dim V^s > 2$ then s is not regular as the non-zero weights of V are roots. This contradicts Theorem 2.18.

Suppose first that V is not p -restricted. Then the result follows from Lemma 2.3(3). \square

3.3 Groups of type A_n

Let G be a simple algebraic group of type A_n and $\omega_1, \dots, \omega_n$ the fundamental weights of G . As above, for a dominant weight ω we denote by V_ω an irreducible G -module with highest weight ω .

3.4 Tensor-indecomposable modules

Lemma 3.24. *Let $V = V_{\omega_i}$, $i \leq (n+1)/2$, and $n+1 < (k+1)i$ for some integer $k > 0$. Suppose that $s \in T_{reg}$ and s is almost cyclic on V . Then $m_s(V) \leq k$. In particular, $m_s(V) \leq (n+1)/2$ for $i > 1$.*

Proof. The weights of V are $\varepsilon_{j_1} + \dots + \varepsilon_{j_i}$ for $1 \leq j_1 < \dots < j_i \leq n+1$. If $\lambda = \varepsilon_{j_1} + \dots + \varepsilon_{j_i}$ then we set $\text{supp}(\lambda) = \{j_1, \dots, j_i\}$. Suppose the contrary, and let μ_1, \dots, μ_{k+1} be distinct weights of V such that $\mu_1(s) = \dots = \mu_{k+1}(s)$. As $n+1 < (k+1)i$, it follows that there is $d \in \{1, \dots, n+1\}$ and a pair μ_r, μ_t with $1 \leq r < t \leq k+1$ such that $d \in \text{supp}(\mu_r) \cap \text{supp}(\mu_t)$. In addition, $|\text{supp}(\mu_r) \cup \text{supp}(\mu_t)| \leq 2i-1$ and $2i-1 \leq n$ by assumption. So $l \notin \text{supp}(\mu_r) \cup \text{supp}(\mu_t)$ for some $l \in \{1, \dots, n+1\}$. Set $\alpha = \varepsilon_d - \varepsilon_l$. It follows that $\mu_r - \alpha, \mu_t - \alpha$ are weights of V . This contradicts Lemma 2.16.

For the additional claim, take $k = (n+1)/2$. Then $(k+1)i > n+1$ as $i > 1$. So the claim follows from the above. \square

Lemma 3.25. *Let $G = A_n, n > 2$, $M = V_{\omega_i}$ with $2 \leq i \leq (n+1)/2$, $M_1 = V_{\omega_1}$, $M_2 = V_{\omega_{i-1}}$. Let $\lambda, \lambda' \in \Omega(M_2)$, $\lambda \neq \lambda'$. Then there are distinct weights $\mu, \mu' \in \Omega(M_1)$ such that $\lambda + \mu, \lambda + \mu', \lambda' + \mu, \lambda' + \mu' \in \Omega(M)$. In addition, $\Omega(M) \subset \Omega(M_1) + \Omega(M_2)$.*

Proof. The weights of M_1 are $\varepsilon_1, \dots, \varepsilon_{n+1}$ and those of M_2 are $\varepsilon_{j_1} + \dots + \varepsilon_{j_{i-1}}$ for $1 \leq j_1 < \dots < j_{i-1} \leq n+1$. We can assume (using the Weyl group) that $\lambda = \varepsilon_1 + \dots + \varepsilon_{i-1}$ and $\lambda' = \varepsilon_{j_1} + \dots + \varepsilon_{j_{i-1}}$ with $j_1 < \dots < j_{i-1} \leq 2i-2 \leq n-1$. Then we take $\mu = \varepsilon_n, \mu' = \varepsilon_{n+1}$.

For the additional claim, we have $\Omega(V_{\omega_1}) + \Omega(V_{\omega_{i-1}}) = \Omega(V_{\omega_1 + \omega_{i-1}}) \supset \Omega(V_{\omega_i})$ as $\omega_i < \omega_1 + \omega_{i-1}$, see Lemma 2.12. Whence the result. \square

Lemma 3.26. *Let $G = A_n, n > 2$, and let $s \in T_{reg}$.*

(1) *If s is almost cyclic on V_{ω_i} for $2 \leq i \leq (n+1)/2$, then s is cyclic on V_{ω_j} for every $j < i$.*

(2) *If s is almost cyclic on $V_{a\omega_1}$ for $1 < a \leq p-1$, then s is cyclic on $V_{b\omega_1}$ for $1 \leq b < a$.*

Proof. Set $M_1 = V_{\omega_1}$, and in case (1) set $M = V_{\omega_i}, M_2 = V_{\omega_{i-1}}$, in case (2) set $M = V_{a\omega_1}, M_2 = V_{(a-1)\omega_1}$.

(i) s separates the weights of M_2 .

In case (2) this follows from Lemma 2.12(2) and Lemma 2.5(1).

In case (1) suppose the contrary. Then there are distinct weights λ_1, λ_2 of M_2 such that $\lambda_1(s) = \lambda_2(s)$. By Lemma 3.25, there are distinct weights μ_1, μ_2 of M_1 such that $\lambda_k + \mu_l$ are weights of M for $k, l \in \{1, 2\}$. Then $(\lambda_1 + \mu_1)(s) = (\lambda_2 + \mu_1)(s)$ and $(\lambda_1 + \mu_2)(s) = (\lambda_2 + \mu_2)(s)$. As s is almost cyclic on M , it follows that $(\lambda_1 + \mu_1)(s) = (\lambda_1 + \mu_2)(s)$, and then $\mu_1(s) = \mu_2(s)$. This is a contradiction, as every regular semisimple element is cyclic on $V_{\omega_1} = M_1$.

(ii) s is cyclic on M_2 .

Indeed, the weights of V_{ω_j} are well known to be of multiplicity 1, so the claim follows from (i).

(iii) The result follows by induction on i in case (1) and on a in case (2). \square

We use Lemma 3.25 to prove

Lemma 3.27. *Let $G = A_n, n > 1, M_1 = V_{\omega_1}, M = V_{a\omega_i + (p-1-a)\omega_{i+1}}$ with $1 \leq i \leq (n+1)/2, M_2 = V_{\omega_{i-1} + (a-1)\omega_i + (p-1-a)\omega_{i+1}}$ with $1 \leq a \leq p-1$. (If $i = 1$ then ω_{i-1} is interpreted as the zero weight.)*

(1) *Let $\lambda, \lambda' \in \Omega(M_2), \lambda \neq \lambda'$. Then there are distinct weights $\mu, \mu' \in \Omega(M_1)$ such that $\lambda + \mu, \lambda + \mu', \lambda' + \mu, \lambda' + \mu' \in \Omega(M)$. In addition, $\Omega(M) \subseteq \Omega(M_1) + \Omega(M_2)$.*

(2) *If s is almost cyclic on M then s separates the weights of M_2 .*

Proof. (1) We have $\Omega(M) = \Omega(V_{\omega_i}) + \Omega(V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}})$ by Lemma 2.12, and $\Omega(V_{\omega_i}) \subset \Omega(V_{\omega_1}) + \Omega(V_{\omega_{i-1}})$ by Lemma 3.25. So

$$\Omega(M) \subset \Omega(M_1) + \Omega(V_{\omega_{i-1}}) + \Omega(V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}}) = \Omega(M_1) + \Omega(M_2)$$

as $\Omega(M_2) = \Omega(V_{\omega_{i-1}}) + \Omega(V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}})$ again by Lemma 2.12.

Let $i > 1$. Then we can write $\lambda = \beta + \gamma, \lambda' = \beta' + \gamma'$ with $\beta, \beta' \in \Omega(V_{\omega_{i-1}}), \gamma, \gamma' \in \Omega(V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}})$. By Lemma 3.25, there are weights $\mu, \mu' \in \Omega(V_{\omega_1})$ such that $\mu + \beta, \mu' + \beta, \mu + \beta', \mu' + \beta' \in \Omega(V_{\omega_i})$. As $\Omega(V_{\omega_i}) + \Omega(V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}}) = \Omega(V_{a\omega_i + (p-1-a)\omega_{i+1}})$, we have $\mu + \beta + \gamma, \mu' + \beta + \gamma, \mu + \beta' + \gamma', \mu' + \beta' + \gamma' \in \Omega(M)$, and the result follows.

Let $i = 1$ so $V_{\omega_{i-1}}$ is the trivial module. Then $\Omega(M) = \Omega(M_1) + \Omega(M_2)$, and the claim is obvious.

(2) We mimic the proof of Lemma 3.26. Suppose the contrary, and let λ_1, λ_2 be distinct weights of M_2 such that $\lambda_1(s) = \lambda_2(s)$. By the above, there are distinct weights μ_1, μ_2 of M_1 such that $\lambda_j + \mu_k$ are weights of M for $i, k \in \{1, 2\}$. Then $(\lambda_1 + \mu_j)(s) = (\lambda_2 + \mu_j)(s)$ for $j = 1, 2$. As s is almost cyclic on M , it follows that $(\lambda_1 + \mu_1)(s) = (\lambda_1 + \mu_2)(s)$, and then $\mu_1(s) = \mu_2(s)$. This is a contradiction, as s is regular. \square

Proposition 3.28. *For $G = A_n$ let $M \in \text{Irr}(G)$ be p -restricted and $s \in T_{\text{reg}}$. Suppose that s is almost cyclic on M . Then $m_s(M) \leq n + 1$.*

Proof. Suppose first that the weight zero multiplicity of M is greater than 1. Suppose the contrary; as s is almost cyclic, only the eigenvalue 1 multiplicity is greater than 1. Then the result follows from Theorem 3.2(1) and Table 1.

Suppose that the weight zero multiplicity of M is at most 1. By Theorem 2.10, all weight multiplicities of M equals 1. By Table 1, the highest weight ω of M is in the following list: $0, \omega_i$ ($1 \leq i \leq n$), $a\omega_1, a\omega_n, a\omega_i + (p-1-a)\omega_{i+1}$, where $1 \leq i \leq n-1$ and $1 \leq a < p$.

Note that we assume $\omega \neq 0, \omega_1, \omega_n$ as these cases are trivial. Since the list of the eigenvalue multiplicities of M is the same as for the dual of M , it suffices to deal with $\omega_i, a\omega_1, a\omega_i + (p-1-a)\omega_{i+1}$ for $i \leq (n+1)/2$.

Suppose the contrary. Let $M_1 = V_{\omega_1}$ and

$$M_2 = \begin{cases} V_{\omega_{i-1}} & \text{if } \omega = \omega_i, i > 1; \\ V_{(a-1)\omega_1} & \text{if } \omega = a\omega_1, a > 1; \\ V_{\omega_{i-1} + (a-1)\omega_i + (p-1-a)\omega_{i+1}} & \text{if } \omega = a\omega_i + (p-1-a)\omega_{i+1}. \end{cases}$$

(If $i = 1$ then ω_{i-1} is understood to be the zero weight.) By Lemmas 3.26 and 3.27(2), s separates the weights of M_2 .

Note that M is a constituent of $M_1 \otimes M_2$. Indeed, V_{ω_i} is a constituent of $V_{\omega_1} \otimes V_{\omega_{i-1}}$ and M_2 is a constituent of $V_{\omega_{i-1}} \otimes V_{(a-1)\omega_i + (p-1-a)\omega_{i+1}}$, whence the claim.

Furthermore, $\Omega(M) \subseteq \Omega(M_1) + \Omega(M_2)$ (see Lemma 3.27). As all eigenvalue multiplicities of M equal 1, there are $n+2$ distinct weights ν_1, \dots, ν_{n+2} of M such that $\nu_i(s) = e$. Then we can write $\nu_i = \lambda_i + \mu_i$, where λ_i, μ_i are weights of M_1, M_2 , respectively, for $i = 1, \dots, n+2$. Then $e = \lambda_i(s)\mu_i(s)$. As $M_1 = V_{\omega_1}$ has exactly $n+1$ distinct weights, we have $\lambda_i = \lambda_j$ for some $i, j \in \{1, \dots, n+1\}$ with $i \neq j$. Then $\mu_i \neq \mu_j$ as otherwise $\nu_i = \lambda_i + \mu_i = \lambda_j + \mu_j = \lambda_j + \mu_j = \nu_j$, which is false. However, $\lambda_j(s)\mu_j(s) = e = \lambda_i(s)\mu_i(s) = \lambda_j(s)\mu_i(s)$, whence $\mu_i(s) = \mu_j(s)$. This is a contradiction as s separates the weights of M_2 . \square

The bound in Proposition 3.28 is not probably sharp in general, at least, for $G = A_n$ we have no example of a p -restricted G -module M with $m_s(M) > n$ for s regular and almost cyclic on M . Moreover, the equality $m_s(M) = n$ probably holds only for M adjoint, and we have no sharp bound for other p -restricted modules. In Lemmas 3.24 and 3.29 we consider some special cases. Example 3.30 shows that, for $G = A_n$, $n > 3$ odd, there exists a regular

semisimple element $s \in G$ and a p -restricted irreducible G -module V which is not a twist of the natural or adjoint G -module such that s is almost cyclic on V and $m_s(V) = (n+1)/2$. (If V is not a twist of a p -restricted G -module then there are examples with $m_s(V) = n+1$, see Example 3.37.)

Lemma 3.29. *Let $G = A_n = SL_{n+1}(F)$. If $s \in T_{reg}$ then $m_s(V_{2\omega_1}) \leq (n+3)/2$. If s is strictly regular or if s is almost cyclic on $V_{2\omega_1}$ then $m_s(V_{2\omega_1}) \leq (n+3)/2$.*

Proof. By [25, Theorem 1.2], if s is almost cyclic on $V_{2\omega_1}$ then s is strictly regular. Suppose that $\mu_1, \dots, \mu_l \in \Omega(V_{2\omega_1})$ be distinct weights such that $\mu_1(s) = \dots = \mu_l(s) = e$. The weights of V are $2\varepsilon_i$ ($i = 1, \dots, n+1$) and $\varepsilon_i + \varepsilon_j$ ($i, j \in \{1, \dots, n+1\}, i < j$); this can be recorded as $\varepsilon_i + \varepsilon_j$ ($i, j \in \{1, \dots, n+1\}, i \leq j$). Suppose first that $\mu_i, \mu_j, \mu_k \in \{2\varepsilon_1, \dots, 2\varepsilon_{n+1}\}$ for some distinct $i, j, k \in \{1, \dots, l\}$. By reordering of μ_1, \dots, μ_l we can assume that $\mu_1, \mu_2, \mu_3 \in \{2\varepsilon_1, \dots, 2\varepsilon_{n+1}\}$, and let $\mu_i = 2\varepsilon_{r(i)}$ for $i = 1, 2, 3$. Then for $i, j \in \{1, 2, 3\}$ we have $1 = (\mu_i - \mu_j)(s) = (2\varepsilon_{r(i)} - 2\varepsilon_{r(j)})(s) = \varepsilon_{r(i)}(s)^2 \varepsilon_{r(j)}(s)^{-2} = (\varepsilon_{r(i)}(s) \varepsilon_{r(j)}(s))^{-1}$, whence $\varepsilon_{r(i)}(s) \varepsilon_{r(j)}(s)^{-1} = \pm 1$. In fact, $\varepsilon_{r(i)}(s) \varepsilon_{r(j)}(s)^{-1} = -1$, as $\varepsilon_{r(i)}(s) = \varepsilon_{r(j)}(s)$ implies $s \notin T_{reg}$. So $\varepsilon_{r(i)}(s) = -\varepsilon_{r(j)}(s)$. However, this cannot happen for three choices $i, j \in \{1, 2, 3\}, i \neq j$. Therefore, at most two weights μ_i ($1 \leq i \leq l$) are in the set $2\varepsilon_1, \dots, 2\varepsilon_{n+1}$.

If s is strictly regular then $\mu_i, \mu_j \in \{2\varepsilon_1, \dots, 2\varepsilon_{n+1}\}$ implies $\varepsilon_{r(i)}(s) = -\varepsilon_{r(j)}(s)$ and then $(\varepsilon_{r(i)} - \varepsilon_{r(j)})(s) = \varepsilon_{r(i)}(s) \varepsilon_{r(j)}(s)^{-1} = -1 = (\varepsilon_{r(j)} - \varepsilon_{r(i)})(s)$, a contradiction as $\varepsilon_{r(i)} - \varepsilon_{r(j)}$ and $\varepsilon_{r(j)} - \varepsilon_{r(i)}$ are distinct roots.

Note that $\Omega(V_{\omega_2}) \subset \Omega(V_{2\omega_1})$. If each μ_1, \dots, μ_l lies in $\Omega(V_{2\omega_1})$ then the result follows from Lemma 3.24. Moreover, the result follows if the supports of some $\mu_i, \mu_j \in \Omega(V_{\omega_2})$ are not disjoint.

We assume that either $\mu_1 = 2\varepsilon_1, \mu_2, \dots, \mu_l \in \Omega(V_{\omega_2})$, or $\mu_1 = 2\varepsilon_1, \mu_2 = 2\varepsilon_2$ and $\mu_3, \dots, \mu_l \in \Omega(V_{\omega_2})$. In addition, $\text{supp}(\mu_i) \cap \text{supp}(\mu_j) = \emptyset$ whenever $i, i > 1$ in the former case and $i, j > 2$ in the latter one. Therefore, $2l - 1 \leq n + 1$, resp., $2(l - 2) + 2 \leq n + 1$, whence $l \leq (n + 2)/2$, resp., $l \leq (n + 3)/2$. \square

Example 3.30. (1) Let $G = SL_{2m}(F)$ and $s = \text{diag}(a_1, \dots, a_m, a_m^{-1}, \dots, a_1^{-1}) \in G$, where the elements $a_1, \dots, a_m \in F$ are algebraically independent and non-zero. Let $\omega = \omega_2$. Then $V_\omega^T = 0$ and $\dim V_\omega^s = m = (n + 1)/2$, where $n = 2m - 1$ is the rank of G .

If $s = \text{diag}(-a_1, \dots, -a_m, a_m^{-1}, \dots, a_1^{-1})$ then -1 has multiplicity $m = (n + 1)/2$ and the other eigenvalues of s on V_{ω_2} are of multiplicity 1.

(2) Let $G = SL_{2m+1}(F)$, $p \neq 2$ and $s = \text{diag}(1, a_1, \dots, a_m, a_m^{-1}, \dots, a_1^{-1}) \in G$, where the elements $a_1, \dots, a_m \in F$ are as above. Let $\omega = 2\omega_1$. Then $V_\omega^T = 0$ and $\dim V_\omega^s = m + 1 = (n + 2)/2$, where $n = 2m$ is the rank of G . In this case $s \in H \cong B_m$.

(3) Let $G = SL_{2m}(F)$ and $s = \text{diag}(b, -b, b^2 a_1, a_1^{-1}, \dots, b^2 a_{m-1}, a_{m-1}^{-1}) \in G$, where the elements $a_1, \dots, a_{m-1} \in F$ are as above and $b^{n+1} = (-1)^{(n+1)/2}$. Then the eigenvalue b^2 has multiplicity $m + 1 = (n + 3)/2$ on $V_{2\omega_1}$ and the other eigenvalues have multiplicity 1.

3.5 Tensor-decomposable modules: special cases

Proposition 3.28 states that Theorem 1.2 is true for p -restricted irreducible A_n -modules. We next prove it for tensor-decomposable modules.

Lemma 3.31. [24, Lemma 7] *Let $s \in T$ be a non-central element. Let ω be a dominant weight which is not p -restricted and not of the form $p^k\mu$ for μ a p -restricted dominant weight. If s is almost cyclic on V_ω then all weights of V_ω are of multiplicity 1.*

For a G -module V and $s \in G$ denote by $\deg_V(s)$ the degree of the minimum polynomial of $\rho(s)$, where ρ is the representation afforded by V . If s is semisimple then $\deg_V(s) = |E_V(s)|$ is the number of distinct eigenvalues of s on V . Note that s is cyclic on V if and only if $\deg_V(s) = \dim V$.

For $A, B \subset F^\times$ we set $AB = \{ab : a \in A, b \in B\}$.

Lemma 3.32. *Let M, M_1, M_2, V be non-trivial G -modules and $s \in T$. Suppose that s is not scalar on both M_1, M_2 , $E_M(s) = E_{M_1}(s)E_{M_2}(s)$ and s is almost cyclic on $M \otimes V$.*

- (1) $\deg_{M_i \otimes V}(s) = \deg_{M_i}(s) \cdot \dim V$ for $i = 1, 2$.
- (2) the eigenvalue multiplicities of s on $M \otimes V$ do not exceed $d = \dim M_1$.

Proof. (1) We can assume $i = 1$. By Lemma 2.3, s is cyclic on M and V . Then $\dim V = \deg_V(s)$. Suppose the contrary, that $\deg_{M_1 \otimes V}(s) < \deg_{M_1}(s) \cdot \dim V$. Then there are distinct eigenvalues $\mu, \mu' \in E_{M_1}(s)$ and $\nu, \nu' \in E_V(s)$ such that $\mu\nu = \mu'\nu'$. Then $\mu\lambda\nu = \mu'\lambda\nu' \in E_{M \otimes V}(s)$ for every $\lambda \in E_{M_2}(s)$. So s is not almost cyclic on $M \otimes V$ unless $\lambda(s)$ is scalar, which is not the case by assumption.

(2) Let $E_{M_1}(s) = \{\mu_1, \dots, \mu_d\}$, $E_{M_2}(s) = \{\lambda_1, \dots, \lambda_r\}$, and $E_V(s) = \{\nu_1, \dots, \nu_t\}$. Then the eigenvalues of s on $M \otimes V$ are $\mu_i\lambda_j\nu_k$, that is, $E_{M \otimes V}(s) = E_{M_1}(s)E_{M_2}(s)E_V(s)$. Suppose the contrary, that the multiplicity of some eigenvalue e of s on $M \otimes V$ is greater than n . As s is cyclic on M and V , there are distinct eigenvalues $\sigma_1, \dots, \sigma_{d+1} \in E_s(M)$ and distinct eigenvalues $\nu_1, \dots, \nu_{d+1} \in E_V(s)$ such that $e = \sigma_i\nu_i$ for $i = 1, \dots, d+1$. By assumption, $\sigma_i = \mu'_i\lambda'_i$ for $i = 1, \dots, d+1$ for some $\mu'_i \in E_{M_1}(s)$ and $\nu'_i \in E_{M_2}(s)$. Therefore, $\mu'_i = \mu'_j$ for some $i, j \in \{1, \dots, d+1\}$; by reordering μ_1, \dots, μ_d we can assume $\mu'_i = \mu'_j = \mu_1$. Then $\sigma_i = \mu_1\lambda'_i$, $\sigma_j = \mu_1\lambda'_j$, and $\lambda'_i \neq \lambda'_j$ as $\sigma_i \neq \sigma_j$. Then $e = \mu_1\lambda'_i\nu_i = \mu_1\lambda'_j\nu_j$, whence $\lambda'_i\nu_i = \lambda'_j\nu_j$. As $\lambda'_i \neq \lambda'_j$, we conclude that $\deg_{M_2}(s) \otimes V < \deg_{M_2}(s) \deg_V(s)$, contrary to (1). \square

To illustrate the use of Lemma 3.32, we state the following:

Corollary 3.33. *Let $G = A_n$, $\omega = a\omega_1$ or $a\omega_1 + (p-1-a)\omega_2$, where $1 < a < p$. Let $M = V_\omega$ and let V be a non-trivial irreducible G -module. Let $s \in (T \setminus Z(G))$. Suppose that s is almost cyclic on $M \otimes V$. Then the eigenvalue multiplicities of s on $M \otimes V$ do not exceed $n+1$.*

Proof. Set $M_1 = V_{\omega_1}$ and $M_2 = V_{(a-1)\omega_1}$ or $V_{(a-1)\omega_1 + (p-1-a)\omega_2}$, respectively. Then $\Omega(M) = \Omega(M_1) + \Omega(M_2)$ by Lemma 2.12. So the result follows from Lemma 3.32. \square

Note that Corollary 3.33 is not implied by Proposition 3.28. In addition, here and in Lemmas 3.34, 3.35 below we do not assume s to be regular as this automatically holds due to Lemma 2.4(2) when $s \notin Z(G)$.

Lemma 3.34. *Let M, M_1, M_2, V be non-trivial G -modules. Let $s \in (T \setminus Z(G))$. Suppose that*

(i) *for every $f, f' \in E_{M_2}(s)$ there exist distinct eigenvalues $e, e' \in E_{M_1}(s)$ such that $ef, ef', e'f, e'f' \in E_M(s)$;*

(ii) *$E_M(s) \subseteq E_{M_1}(s) \cdot E_{M_2}(s)$;*

(iii) *$M \otimes V$ is irreducible and s is almost cyclic on $M \otimes V$.*

Then $\deg_{M_2 \otimes V}(s) = \deg_{M_2}(s) \cdot \dim V$, and $m_s(M \otimes V) \leq \dim M_1$.

Proof. By Lemma 2.3, s is cyclic on V , so $\dim V = \deg_V(s)$. To prove the first statement of the lemma, suppose (the contrary), that $\deg_{M_2 \otimes V}(s) < \deg_{M_2}(s) \cdot \dim V$. Then there are distinct eigenvalues $f, f' \in E_{M_2}(s)$ and eigenvalues $v, v' \in E_V(s)$ such that $fv = f'v'$.

By (i), there are distinct eigenvalues $e, e' \in E_{M_1}(s)$ such that $ef, ef', e'f, e'f' \in E_M(s)$, and hence $efv = ef'v', e'fv = e'f'v' \in E_{M \otimes V}(s)$. So $a := efv$ and $b := e'fv$ are distinct eigenvalues of s on $M \otimes V$, each of multiplicity at least 2. As s is almost cyclic on $M \otimes V$, we must have $a = b$, whence $e = e'$, a contradiction.

Turn to the second statement. Set $d = \dim M_1$. Suppose the contrary, that the multiplicity of some eigenvalue u of s on $M \otimes V$ is greater than d . By Lemma 2.3, s is cyclic on M and V , so there are distinct eigenvalues w_1, \dots, w_{d+1} of s on M and distinct eigenvalues v_1, \dots, v_{d+1} of s on V such that $u = w_i v_i$ for $i = 1, \dots, d+1$. Furthermore, by (ii), $w_i = e_i f_i$ for $i = 1, \dots, d+1$ for some eigenvalues $e_i \in E_{M_1}(s)$ and some eigenvalues $f_i \in E_{M_2}(s)$. Therefore, $e_i = e_j$ for some $i, j \in \{1, \dots, d+1\}$. Then $w_j = e_i f_j$, where $f_i \neq f_j$ as $w_i \neq w_j$, and $u = e_i f_i v_i = e_i f_j v_j$, whence $f_i v_i = f_j v_j$. Then $\deg_{M_2 \otimes V}(s) < \deg_{M_2}(s) \deg_V(s)$ contradicting the first statement of the lemma. \square

Lemma 3.35. *Let $G = A_n$ and $s \in (T \setminus Z(G))$. Let $M = V_{\omega_i}$ for $1 < i < n$ or $V_{a\omega_i + (p-1-a)\omega_{i+1}}$ for $1 < i < n-1$, and let V be a non-trivial G -module. Suppose that $M \otimes V$ is irreducible and s is almost cyclic on $M \otimes V$. Then $m_s(M \otimes V) \leq n+1$.*

Proof. We can assume (by replacing M by its dual) that $i \leq (n+1)/2$. Set $M_1 = V_{\omega_1}$.

Suppose first that $M = V_{\omega_i}$. Set $M_2 = V_{\omega_{i-1}}$. The result follows from Lemma 3.34 as soon as we show that the assumptions (i), (ii) of that lemma hold in this case (as (iii) is in the assumption). By Lemma 2.12(2), $\Omega(M) \subseteq \Omega(M_1) + \Omega(M_2)$, whence (ii). To prove (i), let $f, f' \in E_{M_2}(s)$. Then there are weights $\lambda, \lambda' \in \Omega(M_2)$ such that $\lambda(s) = f, \lambda'(s) = f'$. By Lemma 3.25, there are weights $\mu, \mu' \in \Omega(M_1)$ such that $\lambda + \mu, \lambda + \mu', \lambda' + \mu, \lambda' + \mu' \in \Omega(M)$. Set $e = \mu(s), e' = \mu'(s)$. To get (i), we need $e \neq e'$. Suppose $e = e'$; then $(\lambda + \mu)(s) = (\lambda + \mu')(s)$ and $\lambda + \mu \neq \lambda + \mu'$. This is a contradiction as s is cyclic on M by Lemma 2.3.

Suppose that $M = V_{a\omega_i + (p-1-a)\omega_{i+1}}$. Then we set $M_2 = V_{\omega_{i-1} + (a-1)\omega_i + (p-1-a)\omega_{i+1}}$. The result follows from Lemma 3.34 if the assumptions (i), (ii) of that lemma hold in this case (since (iii) is in the assumption). Lemma 3.27 yields (ii).

To prove (i), let $f, f' \in E_{M_2}(s)$. Then there are weights $\lambda, \lambda' \in \Omega(M_2)$ such that $\lambda(s) = f, \lambda'(s) = f'$. By Lemma 3.27, there are weights $\mu, \mu' \in \Omega(M_1)$ such that $\lambda + \mu, \lambda + \mu', \lambda' + \mu, \lambda' + \mu' \in \Omega(M)$. Set $e = \mu(s), e' = \mu'(s)$. To get (i), we need $e \neq e'$. Suppose $e = e'$; then $(\lambda + \mu)(s) = (\lambda + \mu')(s)$ and $\lambda + \mu \neq \lambda + \mu'$. This is a contradiction as s is cyclic on M by Lemma 2.3. \square

Theorem 3.36. *Let G be of type A_n and N an irreducible G -module. Let $s \in T_{reg}$. Suppose that s is almost cyclic on N . Then $m_s(N) \leq n + 1$.*

Proof. Let $N = N_1 \otimes \cdots \otimes N_k$, where N_1, \dots, N_k are tensor-indecomposable non-trivial irreducible G -modules. Then every N_i for $i = 1, \dots, k$ is a Frobenius twist of an infinitesimally irreducible G -module. The problem is well known to be equivalent to the case where one of N_i is infinitesimally irreducible and we can assume that this is N_1 by reordering the terms.

If $k = 1$ then the result is contained in Proposition 3.28. Let $k > 1$. Then, by Lemma 2.3, s is cyclic on every N_i . Therefore, all weights of N_i are of multiplicity 1. By [18, 6.1], N_i is a Frobenius twist of an irreducible G -module, whose highest weight belongs to the list $\{\omega_j, 1 \leq j \leq n, a\omega_1, a\omega_n \ (1 < a < p), a\omega_i + (p - 1 - a)\omega_{i+1}, i = 1, \dots, n - 1, 0 \leq a < p\}$. Set $M = N_1, V = N_2 \otimes \cdots \otimes N_k$, so V_2 is irreducible and non-trivial, whereas the highest weight ω of N_1 lies in the above list. Then the result follows from Corollary 3.33 and Lemma 3.35. \square

One can expect that the upper bound for $m_s(V)$ for V tensor-decomposable must be significantly lower than for V tensor-indecomposable. The following example makes evident that this is not so. In fact, we see that the bound in Theorem 3.36 is sharp.

Example 3.37. Let F be an algebraically closed field of characteristic $p > 0$, $G = A_n$, $n \geq 2$, and V an irreducible G -module with highest weight $\omega_1 + p\omega_n$. Then there exists $s \in G$ such that s is almost cyclic on V and $m_s(V) = n + 1$ for some $s \in G$.

Indeed, set $s = \text{diag}(a, a^p, \dots, a^{p^n}) \in SL_{n+1}(F)$, where $a \in F$ is a primitive $((p^{n+1} - 1)/(p - 1))$ -root of unity. Then the matrix of s on V is $s \otimes s^{-p}$ and the eigenvalues of this matrix are $a^{p^i - p^{j+1}}$ with $0 \leq i, j \leq n$. If $i = j + 1$ then this equals 1, as well as if $i = 0, j = n$ as then $a^{1 - p^{n+1}} = 1$. So 1 is an eigenvalue of $s \otimes s^{-p}$ of multiplicity at least $n + 1$. In fact, this is exactly $n + 1$ as $a^{p^i - p^{j+1}} = 1$ implies $p^i - p^{j+1} \equiv 0 \pmod{p^n + p^{n-1} + \cdots + p + 1}$ which is false unless $i = j + 1$ or $i = 0, j = n$. We show that all other eigenvalues are of multiplicity 1. Suppose the contrary. Then $a^{p^i - p^{j+1}} = a^{p^k - p^{l+1}}$ for some choice of $0 \leq i, j, k, l \leq n$ and $i \neq j + 1, k \neq l + 1$ and $(i, j) \neq (0, n) \neq (k, l)$. As above, we observe that $i \neq k, j \neq l$. We have $p^i - p^{j+1} - p^k + p^{l+1} \equiv 0 \pmod{p^n + p^{n-1} + \cdots + p + 1}$. Let $m = \min(i, j + 1, k, l + 1)$. If $m > 0$ then $p^{i-m} - p^{j+1-m} - p^{k-m} + p^{l+1-m} \equiv 0 \pmod{p^n + p^{n-1} + \cdots + p + 1}$ and now $d := \max(i - m, j + 1 - m, k - m, l + 1 - m)$ does not exceed n . If $d < n$ then the absolute value of the left hand side is at most $2p^{n-1}$ and the congruence does not hold. (If $p = 2$ then $2^n + 2^{n-1} + \cdots + 1 = 2^{n+1} - 1$ and the congruence fails too.) Suppose that $d = n$. Then $m = 1$ and $\max(j, l) = n$. We can assume $l = n$ and then $j < n$. Then the absolute

value of $p^{i-m} - p^{j+1-m} - p^{k-m} + p^n$ does not exceed $p^n + p^{n-1} < p^n + p^{n-1} + \cdots + p + 1$, hence the congruence above cannot hold.

So $m = 0$. As above, the congruence fails unless possibly $d \in \{n, n+1\}$.

Suppose that $d = n$. Then the absolute value of $p^i - p^{j+1} - p^k + p^{l+1}$ does not exceeds $bp^n - p^c - 1$, where $b \in \{0, 2\}, 0 \leq c \leq n$. Then the congruence cannot hold.

Suppose that $d = n+1$. Then we can replace p^d by 1, and then the absolute value of $p^i - p^{j+1} - p^k + p^{l+1}$ does not exceeds $2 + p^n + p^{n-1}$. If $n > 2$ then the congruence above cannot hold. Let $n = 2$. We can assume that $l = 2$ and then $s \otimes s^{-p} = \text{diag}(1, 1, 1, a^{1-p}, a^{2+p}, a^{1+2p}, a^{p-1}, a^{-1-2p}, a^{-2-p})$. One checks straightforwardly that this matrix is almost cyclic.

Proof of Theorem 1.2. If G is of type A_n then the result is contained in Theorem 3.36. For G of type E_6 and E_7 see Lemmas 3.17 and 3.20, respectively. The case with $G = G_2$ is covered by Corollary 3.23.

Suppose first that $V = V_1 \otimes V_2$ is tensor-decomposable, that is, both V_1, V_2 are non-trivial. Then s is cyclic on V_1 and on V_2 (Lemma 2.3(3)). If G is not of type $A_n, n > 1, D_n, n$ odd, or E_6 then every semisimple element of G is real and the result follows from Lemma 2.3(3).

Let $G = D_n, n > 3$ odd. Let $\omega = \sum p^i \mu_i$ be the highest weight of V , where μ_i 's are p -restricted. Then $V = \otimes_i V_{p^i \mu_i}$. By Lemma 3.31, the weights of $V_{p^i \mu_i}$, and hence of V_{μ_i} are of multiplicity 1, so, by Table 2, $\mu_i \in \{\omega_1, \omega_{n-1}, \omega_n\}$. Then $m_s(V) \leq 2$ by Lemma 3.8.

Next suppose that V is tensor-indecomposable. By Steinberg's theorem, V is a Frobenius twist of a p -restricted G -module. So we can assume that V itself is p -restricted.

Suppose first that ω occurs in Table 1. Then $\dim V^T > 1$ and $\dim V^T = \dim V^s = m_s(V)$ by Theorem 3.2(1), so (2) follows from the data in Table 1.

Suppose that ω does not occur in Table 1. By Theorem 2.10, ω occurs in Table 2. If $(G, p, \omega) = (F_4, 3, \omega_4)$ then the result follows from Lemma 3.21. This completes considerations of groups of the exceptional Lie types.

So we are left with classical groups (other than of type A_n). The case with $G = D_n, n \geq 4$ is settled in Lemma 3.7 for $\omega = \omega_{n-1}$ or ω_n , for $\omega = \omega_1$ see Lemma 2.7(3). This implies the result for an arbitrary p -restricted ω in view of Theorem 2.10 and Table 2. A similar argument works for $G = B_n, n > 2, p \neq 2$, with use of Lemmas 3.11 and 3.4. The case $B_2 \cong C_2$ is considered in Lemma 3.12.

The case $G = C_n, p = 2$ has been examined in Lemma 3.4. Suppose that G is of type C_n with $p \neq 2$. By Theorem 2.10 and Tables 1,2 we have to inspect the cases with $\omega \in \{\omega_1, \omega_{n-1} + \frac{p-3}{2}\omega_n, \frac{p-1}{2}\omega_n\}$ as well as $n = 3, p \neq 2, \omega = \omega_3$. The case with $\omega = \omega_1$ is trivial, the other cases are dealt with in Lemmas 3.11, 3.10 and the remaining case (where $G = C_3, p \neq 2, \omega = \omega_3$) is settled in Lemma 3.9. \square

One easily observes that the exceptions in Theorem 1.2 are genuine. It suffices to show this for $\omega \in \{3\omega_1, p \neq 2, (1+p^k)\omega_1\}$. Let $\text{diag}(a, a^{-1}) \in SL_2(F)$ be the matrix of s on V_{ω_1} . Then the matrices of s on $V_{2\omega_1}, V_{3\omega_1}$ and $V_{(1+p^k)\omega_1}$ are $\text{diag}(a^2, 1, a^{-2}), \text{diag}(a^3, a, a^{-1}, a^{-3})$ and $\text{diag}(a^{p+1}, a^{p-1}, a^{-p-1}, a^{-p+1})$, respectively. If $a^2 = -\text{Id}, a^3 = 1 \neq a$ then s is almost

cyclic on $V_{2\omega_1}, V_{3\omega_1}$, respectively, and $m_s(V_{2\omega_1}) = m_s(V_{3\omega_1}) = 2$. If a is a primitive $(p+1)$ -root of unity then s is almost cyclic on $V_{(p+1)\omega_1}$ and $m_s(V_{(p+1)\omega_1}) = 2$.

Proof of Theorem 1.1. The result follows from Theorem 1.2. □

Proof of Corollary 1.3. Suppose the contrary, that $m_s(V) \geq \dim V/2$, or, equivalently, $\dim V/m_s(V) \leq 2$. By Theorem 1.2, either $m_s(V) \leq 2$, hence $\dim V \leq 4$, or V is as in item (1), (2) or (3) of Theorem 1.2.

Suppose first that $\dim V \leq 4$. The case $\dim V = 2$ is ruled out as $m_s(V) = 0$ in this case. If $\dim V = 3$ then $G = A_1, p \neq 2, \omega = 2p^i\omega_1$ or $G = A_2, \omega \in \{p^i\omega_1, p^i\omega_2\}$. In the former case $m_V(s) = 2$ for $s \in SL_2(F)$ implies $s^2 = -\text{Id}$ as recorded in the corollary. The latter case is ruled out by Lemma 2.7 or straightforwardly. Let $\dim V = 4$. Then $G \in \{A_1, A_3, B_2, C_2\}$. In fact, if $G = A_3$ then V is a twist of the natural G -module, on which $m_s(V) \leq 1$ for a regular semisimple element $s \in G$ (Lemma 2.7). The cases with $G = B_2$ and C_2 are ruled out by Remark 3.13. Let $G = A_1$. Then ω is well known to be either $3p^i\omega_1$ ($p \neq 2, 3$), or $(p^i + p^j)\omega_1$ as recorded in the statement.

The result is obvious if item (3) of Theorem 1.2 holds as $\dim V \geq 27$ for $G = E_6$.

Suppose that (2) holds. Then $m_s(V) = \dim V^T$ and inspection of $\dim V/\dim V^T$ in Table 1 yields a contradiction.

Suppose that item (1) of Theorem 1.2 holds so G is of type A_n . We can assume $n > 1$ as otherwise $m_s(V) \leq 2$ by Theorem 1.1 and $\dim V \leq 4$; this case has been examined above. In addition, we can assume that V is not a twist of the adjoint G -module as this is included in case (2), unless $n = 2, p = 3$. In the latter case $\dim V = 7$ and $m_s(V) \leq 2$ by Theorem 1.1.

By the above, we can assume that $m_s(V) > 2$ and $\dim V > 5$.

So we assume that $n > 1$. If V is a twist of the irreducible G -module with highest weight ω_1 then we get a contradiction with Lemma 2.7.

If V is *not* a twist of the irreducible G -module with highest weight ω_1 then $\dim V \geq n(n+1)/2$ so $\dim V/m_s(V) \geq n/2$ and $n/2 > 2$ for $n > 4$. Suppose that $n \leq 4$ and $\dim V \leq 2(n+1)$. If $n = 2$, this implies $p \neq 2$ and $\omega \in \{2\omega_1, 2\omega_2\}$. However, in these cases $\dim V = 6$ and $m_s(V) \leq 2$ by Lemma 3.29, so this case is ruled out. By [14, Appendices A6, A7, A8], the highest weight ω of V is $p^t\omega_2$ if $n = 3$ and $\{p^t\omega_2$ or $p^t\omega_3\}$ if $n = 4$ for some integer $t \geq 0$. So either $G = SL_4(F), V = V_{\omega_2}$ or $G = SL_5(F), V = V_{\omega_2}$ or V_{ω_3} . As $V_{\omega_2}, V_{\omega_3}$ are dual for $n = 4$, it suffices to examine $\omega = \omega_2$ in both the cases. If $n = 3$ then V is a twist of the natural module for $D_3 \cong A_3$ on which $m_V(s) \leq 2$ as s is regular (see Lemma 2.7). If $n = 4$ then $m_s(V_{\omega_2}) \leq 2$ by Lemma 3.24 and $\dim V_{\omega_2} = 10$, so this case is ruled out. This completes the proof. □

4 An application to a recognition problem for linear groups over finite fields

There is a problem in the theory of linear groups over finite fields aimed to determining the subgroups H of $GL_n(q)$ containing a matrix $\text{diag}(\text{Id}_m, M)$, where $M \in GL_{n-m}(q)$ is irreducible. The latest and most significant contribution to this project is made in [8], where the authors solve this problem for $m < n/2$. The result of [8] can be also interpreted as a classification of subgroups of $GL_n(q)$ containing a cyclic Sylow ℓ -subgroup of $GL_n(q)$ for some prime divisor ℓ of $GL_n(q)$, see [1]. Observe that the case where H is irreducible in $GL_n(q)$ is the bulk of the problem.

In this section we contribute to the problem in the case where H is a quasi-simple group of Lie type in defining characteristic $p > 0$ for $p|q$. Our purpose is to consider the complimentary case with $m \geq n/2$. Observe that the case with $p \nmid q$ is considered in [7].

Note that specifying the main theorem of [8] to the case where H is a quasi-simple group of Lie type leads one to Examples 2.1, 2.4 and 2.9 in [8]. Then it is easy to observe that Examples 2.1, 2.4 yield nearly natural representations of H (whereas Examples 2.9 are irregular cases arising for $n \leq 9$). The converse is also true, that is, if H is a quasi-simple group of Lie type in defining characteristic $p|q$ and $\rho : H \rightarrow GL_n(q)$ is a nearly natural irreducible representation then $\rho(H)$ is as one of these examples. However, we do not provide here the proof of this fact.

4.1 Representation of $SL_2(q)$ containing almost cyclic elements

Our primary goal here is to give an example with $m = n/2$ (see Lemma 4.5), which is essential for what follows.

Lemma 4.1. *Let $\mathbf{G} = SL_2(F)$, let ρ be an irreducible representation of \mathbf{G} with highest weight $(1 + p^i)\omega_1$ for some integer $i \geq 1$. Let $s \in \mathbf{G}$ be a non-central semisimple element such that $\rho(s)$ is almost cyclic and $m_s(\rho) > 1$. Then $\rho(s) \in PSL_2(p^i) \subset GL_4(p^i)$ and s is conjugate in \mathbf{G} to an element of $SL_2(p^i)$, unless $p \neq 2$ and s^2 is conjugate to an element of $SL_2(p^i)$. In addition, the group $\rho(SL_2(p^{2i}))$ is irreducible whereas $\rho(SL_2(p^i))$ is reducible.*

Proof. Let $s = \text{diag}(a, a^{-1}) \in SL_2(F)$. Then $\rho(s) = \text{diag}(a^{p^i+1}, a^{p^i-1}, a^{-p^i+1}, a^{-p^i-1})$, and some of these eigenvalues coincide. Here $a \in F^\times$, $a^2 \neq 1$ as $s \notin Z(\mathbf{G})$. This implies $a^p \neq a^{-p}$, $a^{p^i+1} \neq a^{p^i-1}$ and $a^{-p^i+1} \neq a^{-p^i-1}$. So either $a^{2(p^i+1)} = 1$ or $a^{2(p^i-1)} = 1$, and hence either $s^{p^i+1} \in Z(\mathbf{G})$ or $s^{p^i-1} \in Z(\mathbf{G})$. If $p = 2$ then $Z(\mathbf{G}) = 1$, and the group $SL_2(2^i)$ contains elements of order $|s|$; these are obviously conjugate to elements of $\langle s \rangle$. Let $p > 2$. If $a^{p^i+1} = 1$ or $a^{p^i-1} = 1$ then, similarly, s is conjugate to an element of $SL_2(p^i)$. Let $a^{p^i+1} = -1$ or $a^{p^i-1} = -1$; then $\rho(s) = (-1, -1, -a^{-2}, -a^2) \in GL_4(p^i)$ as $s^2 \in SL_2(p^i)$. So $\rho(s) \in GL_4(p^i)$, as claimed.

Turn to the additional claim of the lemma. If $q = p^{2i}$ then the weight $(1 + p^i)\omega_1$ is q -restricted, so ρ remains irreducible under restriction to $SL_2(q)$ (see [9, §2.11]). If $q = p^i$ then $\rho|_{SL_2(q)} \cong (\tau \otimes \tau)|_{SL_2(q)}$, where τ is the irreducible representation of \mathbf{G} with highest weight ω_1 . It is well known that $\tau \otimes \tau$ is reducible. \square

Lemma 4.2. *Let $\mathbf{G} = SL_2(F)$, $G = SL_2(q)$, $q = p^d > p$, and let ρ be an irreducible representation of \mathbf{G} with highest weight $(1 + p^i)\omega_1$ for some integer $i \geq 1$. Suppose that $\rho(G)$ is conjugate in $GL_4(F)$ to an irreducible subgroup of $GL_4(p^m)$, where m is minimal possible. Then $d \nmid i$ and either d is even, $i \pmod{d} = d/2$ and $m = d/2$ or $i \pmod{d} \neq d/2$ and $m = d$.*

Proof. Obviously, $\rho(G) \subset GL_4(q)$. Suppose that the group $\rho(G)$ is conjugate to a subgroup of $GL_4(F_0)$, where F_0 is a subfield of \mathbb{F}_q and $|F_0| = p^m$. Then $m|d$. (As \mathbb{F}_q is a vector space over F_0 , we have $|\mathbb{F}_q| = |F_0|^j$ for some integer $j > 0$, so $q = p^d = (p^m)^j$, whence $mj = d$.) As $\rho(G)$ is irreducible, i is not a multiple of d .

Suppose that $F_0 \neq \mathbb{F}_q$, so $m < d$.

By Zsigmondy's theorem (see [11, 5.2.14]), if $q^2 = p^{2d} > p^2$ (this holds by assumption) and $q^2 \neq 64$ then there is a prime $\ell | (q^2 - 1) = p^{2d} - 1$ such that $(\ell, p^j - 1) = 1$ for every $j < 2d$. Therefore, if $F_0 \neq \mathbb{F}_q$ (hence $m < d$) and $q \neq 8$ then $(\ell, p^{2m} - 1) = 1$. Moreover, $(\ell, p^{4m} - 1) = 1$. Indeed, otherwise ℓ divides $(p^{4m} - 1, p^{2d} - 1) = p^{(4m, 2d)} - 1$ by [10, Hilfsatz 2(a)]. As $(4m, 2d) \leq 2d$, the claim is true unless $(4m, 2d) = 2d$, whence $d|2m$, and hence $2m = d$. Therefore, \mathbb{F}_q is the quadratic extension of F_0 . Let $q^2 = 64$ so $G = SL_2(8)$, $d = 3$ and $|s| = 9$. Then $\rho(G) \subset GL_4(8)$. As $2^m | 8$, it follows that either $2^m = 8$, as required, or $m = 1$; in the latter case $\rho(G) \subset GL_4(2)$. However, $GL_4(2)$ has no element of order 9, so we have a contradiction.

Finally, we show that i is a multiple of $d/2$. Let $\gamma : F \rightarrow F$ be the mapping defined by $x \rightarrow x^{p^m}$ for $x \in F$. This extends to the mapping $\tau : GL_4(F) \rightarrow GL_4(F)$. Then $\tau(y) = y$ for $y \in GL_4(F_0)$ and τ acts as an automorphism of order 2 on $GL_4(q^r)$. Then the highest weight of the representation $\rho^\tau : h \rightarrow \tau(\rho(h))$ for $h \in SL_2(F)$ is $p^m(1 + p^i)\omega_1$. As τ acts trivially on $GL_4(F_0)$ and hence on $\rho(G)$, it follows that i is a multiple of $p^m = p^{d/2}$, as claimed. \square

Corollary 4.3. *Let $K = GL_4(p^m) = GL(V)$, where $m > 1$. Then for every $j|2m$, $j > 1$, there is an absolutely irreducible subgroup $G_j \cong PSL_2(p^j)$ of K . In addition we have:*

- (1) *if $p \neq 3$ and $t \in G_j$, $j > 1$, is of order $(p+1)/(2, p+1)$ then t is almost cyclic and $m_t(V) = 2$;*
- (2) *if $p = 3$, j is even and $t \in G_j$ of order 4 then t is almost cyclic and $m_t(V) = 2$.*

Proof. Let \mathbf{G} be as in Lemma 4.1, and let ρ be an irreducible representation of \mathbf{G} with highest weight $(1+p)\omega_1$. Let $s \in SL_2(p)$ be of order $p+1$. Then $t := \rho(s) \in G_1 \subset GL_4(p)$ is almost cyclic and $|t| = (p+1)/(2, p+1)$. Then $G_j := \rho(SL_2(p^j)) \cong PSL_2(p^j)$ is irreducible for $j > 1$ by Lemma 4.1. Whence the result.

If $p = 3$ then $SL_2(9)$ has an element of order 8, so $t = \rho(s) \in G_2$ is almost cyclic with $m_t(V) = 2$ (as -1 is an eigenvalue of t of multiplicity 2). \square

Remark 4.4. It is not true that the irreducible subgroups of $GL_4(F)$ isomorphic to $G_j \cong PSL_2(p^j)$ are conjugate even if they contain elements s with $m_s(V) = 2$. Indeed, let ρ_i be an irreducible representation of $\mathbf{G} \cong SL_2(F)$ with highest weight $(1 + p^i)\omega_1$, $i|j$, $i < j$. Set $G_{ij} = \rho_i(SL_2(p^j))$. Then G_{ij} is irreducible and $G_{ij} \cong G_j$. Let $s_i \in SL_2(p^j)$ be of

order $p^i + 1$ and $t_i = \rho_i(s_i) \in G_{ij}$. Then t_i has eigenvalue 1. Let $i' | j$, and $i \neq i' < j$. Then $\rho_{i'}(s_i) \in G_{ij}$ does not have eigenvalue 1. Therefore, ρ_i and $\rho_{i'}$ are not equivalent and, moreover, G_{ij} and $G_{i'j}$ are not conjugate in $GL_4(F)$.

Lemma 4.5. *Let $H = GL_{4r}(q) = GL(V)$, $q^r \neq 3$. Then there exists an irreducible subgroup $Y \cong PSL_2(q^{2r})$ of H and $h \in PSL_2(q^r) \subset Y$ such that $|h|$ is a prime power, h is almost cyclic on V and the multiplicity of eigenvalue 1 of h equals $2r$. Moreover, $V = V_1 + V_2$, where V_1 is the 1-eigenspace of h , $hV_2 = V_2$ and h acts irreducibly on V_2 .*

Proof. In assumptions of Lemma 4.1 set $G = SL_2(q^{2r})$ and $i = r$. Let $s \in SL_2(q^r) \subset G$ be such that $|s|$ divides $q^r + 1$ and does not divide $q^{2t} - 1$ for any positive integer $t < r$. Set $h = \rho(s)$. By Lemma 4.1, $\rho(G)$ is irreducible and contained in $GL_4(q^r)$ (see Lemma 4.2). Let V' be the underlying space of $GL_4(q^r)$. Then $h = \rho(s)$ is conjugate in $GL_4(q^r)$ to a matrix $\text{diag}(1, A, 1)$, where $A \in GL_2(q^r)$, and A is irreducible (otherwise the eigenvalues of A would lie in \mathbb{F}_{q^r}). So $V' = V'_1 + V'_2$, where V'_1 is the 1-eigenspace of h on V' , $sV'_2 = V'_2$ and h acts irreducibly on V'_2 .

Next embed $GL_4(q^r)$ into $GL_{4r}(q)$ by viewing \mathbb{F}_{q^r} as a vector space over \mathbb{F}_q . Then V'_1, V'_2 and V' can be viewed as vector spaces V_1, V_2, V , say, over \mathbb{F}_q , and we have $\dim V_1 = \dim V_2 = 2r$. We claim that s remains irreducible on V_2 . Indeed, if h is reducible on V_2 then $|h|$ divides $q^j - 1$ for $j < 2r$ and also $q^{2r} - 1$, hence $q^{(2r,j)} - 1$ by [10, Hilfsatz 2(a)]. So we can assume $j | 2r$, hence $j \leq r$.

We claim that there is a prime ℓ such that the ℓ -part of $q^r + 1$ does not divide $q^j - 1$ for $j \leq r$. Indeed, by Zsigmondy's theorem (see [11, 5.2.14]), if $q^2 = p^{2d} > p^2$ and $q^2 \neq 64$ then there is a prime $\ell | (q^2 - 1) = p^{2d} - 1$ such that $\ell \nmid (p^j - 1)$ for every $j < 2d$. If $q^2 = 64$ then take $\ell^2 = 9$. If $q = p$ then let ℓ be the maximal prime dividing $q + 1$. If $\ell \neq 2$ then $\ell \nmid (q - 1)$ as desired. Otherwise, $\ell = 2$ and $q + 1$ is a 2-power, and the claim follows.

Then we take s of order equal to the ℓ -part of $q^r + 1$. As the order of h equal $|s|/(2, |s|)$, we conclude that h is irreducible on V_2 unless $q + 1 = 4$; this is not the case by assumption. This implies the matrix of s on V_2 to be cyclic. This is well known, one can argue as follows. Let $S = \langle s \rangle$ and let R be the regular $\mathbb{F}_{q^r}S$ -module. Then there is a surjective module homomorphism $\eta : R \rightarrow V_2$. Then η extends to FG -module homomorphism $FG := R \otimes F \rightarrow V_2 \otimes F$. As all eigenvalues of s on R are of multiplicity 1, so are those on $V_2 \otimes F$. \square

4.2 Regular semisimple elements in representations of quasi-simple finite groups of Lie type

Recall that the notions of a cyclic and almost cyclic elements are meaningful for elements $g \in GL_n(F_1)$ for any subfield F_1 of F , that is, we simply apply the definition to g as an element of $GL_n(F)$.

Lemma 4.6. *Let G be a finite quasi-simple group of Lie type in defining characteristic $p > 0$, let V be a non-trivial irreducible \mathbb{F}_qG -module for some p -power q and $\bar{V} = V \otimes F$. Let $s \in G$ be a regular semisimple element and let $m_s(V)$ be the maximal eigenvalue multiplicities of s on \bar{V} . Suppose that s is almost cyclic on V and $m_s(V) \geq \dim V/2$. Then*

G is isomorphic to $SL_2(q')$, or $PSL_2(q')$ for some p -power q' . Let ρ be the representation afforded by V . Then we can assume that $G \cong SL_2(q')$ and one of the following holds:

- (1) $\dim V = 3$, $\rho(G) \cong PSL_2(q')$, q is a power of q' , $p \neq 2$ and $s^2 \in Z(G)$;
- (2) $\dim V = 4$, $p \neq 2, 3$, $\rho(G) \cong SL_2(q')$, q is a power of q' and $s^3 \in Z(G)$;
- (3) $\dim V = 4$, $\rho(G) \cong PSL_2(q')$, $q' > p$, q^2 is a q' -power, and if $q' = 3^l$ then l is not a prime;
- (4) $\dim V = 4r$, $r > 1$, $\rho(G) \cong PSL_2(q')$, q^{2r} is a power of q' , and $m_s(V) = \dim V/2$.

Proof. (i) Suppose first that V is absolutely irreducible. Then V extends to a \mathbf{G} -module \bar{V} , where \mathbf{G} is a simply connected simple algebraic group such that $G = \mathbf{G}^\sigma$ and si is some Steinberg endomorphism of \mathbf{G} . By Corollary 1.3, \mathbf{G} is of type A_1 , so $\mathbf{G} \cong SL_2(F)$ and $3 \leq \dim \bar{V} = \dim V \leq 4$. We keep ρ for the corresponding representation of \mathbf{G} .

Then $G \cong SL_2(q')$ for some p -power q' . Let $s \in G$ and let $\text{diag}(a, a^{-1})$ with $a^2 \neq 1$ be a conjugate of s in $SL_2(F)$.

If $\dim \bar{V} = 3$ then $p \neq 2$ and \bar{V} is a Frobenius twist of the irreducible \mathbf{G} -module with highest weight $2\omega_1$; the matrix of $\rho(s)$ is $\text{diag}(a^2, 1, a^{-2})$, which is cyclic unless $a^2 = a^{-2}$, and hence $s^2 = -\text{Id} \in Z(G)$. One observes that such s is conjugate to an element of $SL_2(p)$. So $s \in SL_2(q')$ whenever $SL_2(q') \subseteq SL_2(q)$, equivalently, q is a q' -power.

Let $\dim \bar{V} = 4$. Then \bar{V} is a Frobenius twist of the irreducible \mathbf{G} -module with highest weight $3\omega_1$ (for $p \neq 3$) or $(1+p^i)\omega_1$ for some $i > 0$. By Lemma 2.14, we can assume that the highest weight of \bar{V} is $3\omega_1$ or $(1+p^i)\omega_1$. In the former case $\rho(G) \cong SL_2(q')$ and the matrix of s on \bar{V} is $\text{diag}(a^3, a, a^{-1}, a^{-3})$. If this matrix is not cyclic then either $a^6 = 1, p \neq 3$, and hence $s^3 \in Z(G)$, as required, or $a^4 = 1$ in which case s is not almost cyclic. In the latter case $q' > p$, $\rho(G) \cong PSL_2(q')$ and the matrix of s on \bar{V} is $\text{diag}(a^{p^i+1}, a^{p^i-1}, a^{-p^i+1}, a^{-p^i-1})$. If this matrix is not cyclic then either $a^{2p^i+2} = 1$ or $a^{2p^i-2} = 1$ or $a^{2p^i} = 1$; the latter implies $a^2 = 1$, which is false as s is regular. If $a^{2p^i+2} = a^{2p^i-2}$ then $a^4 = 1$, whence $p \neq 2$ and $a^2 = -1$. One checks that s is not almost cyclic on \bar{V} in this case. In two other cases s is almost cyclic on \bar{V} . By Lemma 4.2, $\rho(G)$ is contained in $GL_4(q')$ or $GL_4(\sqrt{q'})$, whence q^2 is a power of q' . If $p \neq 3$ then $SL_2(q')$ has an element s in question for every q' (see Corollary 4.3), so (3) holds in this case (provided $q' > p$).

Let $p = 3$ and let $q = 3^m$ for some $m > 1$. By Corollary 4.3, if q' is a square then $SL_2(q')$ contains an element s with properties required. Suppose that q' is not a square. Then $q' = 3^l$ with l odd, and $l > 1$ (as $q' > p = 3$). By Lemma 4.2, q' is a minimal 3-power x such that $\rho(SL_2(q')) \subseteq GL_4(x)$. So $GL_4(q') \subseteq GL_4(q)$, and hence q is a power of q' .

Suppose that l is a prime. Then s is not contained in a subgroup $SL_2(3)$ (otherwise $\rho(s)$ is not almost cyclic), and hence s is not contained in any proper subgroup $SL_2(q_1)$ of $SL_2(q')$ for any 3-power q_1 . By Lemma 4.1, s is contained in a subgroup isomorphic to $SL_2(p^i)$, whence p^i is a power of q' . Then $\rho(SL_2(q'))$ is reducible. (In particular, if m is a prime then q' is a square in this case.)

Suppose that l is not a prime, and let l' be a prime with $l'|l$. Let $i = l'$ and let $s \in SL_2(q')$ be of order $3^{l'} + 1$. Then the group $\rho(SL_2(q'))$ is irreducible by Steinberg's theorem (see [9, §2.11]), $\rho(s)$ is almost cyclic and $m_s(\bar{V}) = 2$ by Lemma 4.1. This completes the proof of the case with $\dim \bar{V} = 4$.

(ii) Suppose that V is not absolutely irreducible. Then, by Schur's lemma, the centralizer of $\rho(G)$ in $\text{End}(V)$ is a field F_0 , say, and $|F_0| = q^r$ for some $r > 1$. It follows that $\rho(G) \subset GL_{n/r}(q^r) \subset GL(V)$ ([11, Lemma 2.10.1]).

Therefore, $\bar{V} = V \otimes F$ as a $GL_{n/r}(q^r)$ -module is a direct sum $\bar{V}_1 \oplus \cdots \oplus \bar{V}_r$, where $\bar{V}_1, \dots, \bar{V}_r$ are irreducible $FGL_{n/r}(q^r)$ -modules that are Galois conjugate to each other (by $\text{Gal}(\mathbb{F}_{q^r}/\mathbb{F}_q)$), and each $\bar{V}_1, \dots, \bar{V}_r$ is absolutely irreducible as $\mathbb{F}_{q^r}G$ -module, see [11, Lemma 2.10.2]. Clearly, s is almost cyclic on each $\bar{V}_1, \dots, \bar{V}_r$. As these are Galois conjugate to each other, it follows that $m_s(V) \leq m_s(\bar{V}_1) + \cdots + m_s(\bar{V}_r) = r \cdot m_s(\bar{V}_1)$.

As above, each $\bar{V}_1, \dots, \bar{V}_r$ extends to a \mathbf{G} -module. By Corollary 1.3, $m_s(\bar{V}_i) < \dim \bar{V}_i/2$ for every $i = 1, \dots, r$ (which is a contradiction so the result follows), unless possibly \mathbf{G} is of type A_1 and $\dim \bar{V}_1 = 3, 4$. So we can assume that $\mathbf{G} \cong SL_2(F)$ and then $G = SL_2(q')$ for some p -power q' .

Since s is almost cyclic on \bar{V} and $m_s(\bar{V}) > 1$, a common eigenvalue of s on \bar{V}_j, \bar{V}_l for $j, l \in \{1, \dots, r\}, j \neq l$ is 1 or -1 (indeed, if $\eta \neq \pm 1$ is a common eigenvalue of s on \bar{V}_j and \bar{V}_l then so is η^{-1} , and s is not almost cyclic on \bar{V}). Moreover, both 1 and -1 cannot occur as eigenvalues of s on \bar{V} .

If $\dim \bar{V}_1 = 3$ and $m_s(\bar{V}_1) > 0$ then $s^2 \in Z(\mathbf{G})$, and s is almost cyclic on V implies $\dim V = 3$, which is a contradiction as $r > 1$.

Let $\dim \bar{V}_1 = 4$. If the highest weight of \bar{V}_1 is $3p^i\omega_1$ with $p \neq 2, 3$ then $s^3 \in Z(\mathbf{G})$. $s^3 = 1$. Then s is conjugate to an element of a subgroup $SL_2(p)$. The restrictions of \bar{V}_j to $SL_2(p)$ for $j = 1, \dots, r$ yield isomorphic modules. As s is almost cyclic and $r > 1$, this is a contradiction.

Suppose that the highest weight of \bar{V}_1 is $(1 + p^i)\omega_1$. Then $\rho_j(\mathbf{G}) \cong PSL_2(F)$ for every $j = 1, \dots, r$, so $Z(G)$ acts trivially on V .

By the above, \bar{V}_1 is an irreducible $FGL_{n/r}(q^r)$ -modules and absolutely irreducible as $\mathbb{F}_{q^r}G$ -module, that is, $\rho_1(G) = \rho_1(SL_2(q'))$ is absolutely irreducible subgroup of $GL_4(q^r)$. By (3), q^{2r} is a power of q' , as claimed in (4). In addition, $m_s(\bar{V}_1) \in \{0, 2\}$, so $m_s(\bar{V}) \in \{0, 2r\}$ as required.

By Lemma 4.5, if $q^{2r} = q'$ then $GL_{4r}(q)$ contains an irreducible subgroup isomorphic to $PSL_2(q')$ in which elements of order $q^r + 1$ are almost cyclic on V and $m_s(V) = \dim V/2$; moreover, s is irreducible on $(1 - s)V$. So the case (4) is genuine. \square

Remark 4.7. One can wish to further refine item (4) of Lemma 4.6 by determining q' and the irreducible groups $X \cong PSL_2(q') \subset GL_{4r}(q)$ up to conjugacy. However, we shall not work in this direction. Observe that, in notation of Lemma 4.6(4), $\rho(G)$ remains irreducible in some larger groups $GL_{4r}(q^t)$. For instance, let t be an arbitrary prime such that $(4r, t) = 1$. If ρ is reducible in $GL_{4r}(q^t)$ then $\rho(PSL_2(q')) = \rho_1(PSL_2(q')) \oplus \cdots \oplus \rho_t(PSL_2(q'))$, where $t|4r$. This is a contradiction.

Proof of Theorem 1.4. The main assertion of the theorem follows from Lemma 4.6.

Prove the converse. If $4|\dim V$ then this follows from Lemma 4.5. If $\dim V = 3, p \neq 2$ then consider an irreducible representation ρ of $SL_2(F)$ with highest weight $2\omega_1$, and set

$\rho' = \rho|_{SL_2(q')}$. Then for $s \in SL_2(q')$ with $s^2 = -\text{Id}$ we have $\rho'(s) = \text{diag}(-1, 1, -1)$ under a suitable basis of V . \square

Corollary 4.8. *In assumptions of Lemma 4.6 suppose that $q = 2$. Then item (3) or (4) Lemma 4.6 holds and $q' = 2^{2r}$. In addition, $\rho(G) \subset SO_{4r}^-(q)$.*

Proof. It is obvious that (1) and (2) of Lemma 4.6 do not hold. Let q' be as in Lemma 4.6. If (3) holds then $q'|4$ and $q > 2$, so $q' = 4$ as required. Let (4) holds. Suppose the contrary, that $q' < 2^{2r}$. We have seen in the proof of Lemma 4.6 that $V \otimes F = \overline{V}_1 \oplus \cdots \oplus \overline{V}_r$, where $\overline{V}_1, \dots, \overline{V}_r$ are irreducible $SL_2(F)$ -modules of dimension 4, s is almost cyclic on each \overline{V}_l and $m_s(\overline{V}_l) = 2$ for $l = 1, \dots, r$. Let $\omega = (2^j + 2^i)\omega_1$ be the highest weight of V_1 , where $1 \leq j < i$. By Lemma 2.14, we can assume here that $j = 1$. By Lemma 4.1, s is conjugate to an element of $SL_2(2^i)$ and $\rho_l(s)$ is conjugate to an element of $GL_4(2^i)$, where ρ_l is the representation afforded by \overline{V}_l . As s is almost cyclic on V , the modules $\overline{V}_l|_{SL_2(2^i)}$ for $l = 1, \dots, r$ are pairwise non-isomorphic. In addition, these are twisted (or conjugate) to each other by elements of the Galois group $\text{Gal}(\mathbb{F}_{2^r}/\mathbb{F}_2)$. It follows that $i = r$ and $\omega = (1 + 2^r)\omega_1$. So q' is a power of 2^r and 2^{2r} is a power of q' by Lemma 4.6(4). Here $q' \neq 2^r$ as $\rho_l(G)$ is irreducible (see Lemma 4.1). So $q' = 2^{2r}$ and $G = SL_2(2^{2r})$ as claimed. In addition, $\rho_1(G) \subset O_4^-(2^r)$, and hence $\rho(G) \subset O_{4r}^-(2)$ by [10, Hilfsatz 1(d) and Bemerkung at page 148]. \square

4.3 Non-regular semisimple elements in representations of quasi-simple finite groups of Lie type

In order to deal uniformly with $p = 2$ and $p > 2$ we denote by $SO_8(F)$ the connected component of the orthogonal group $O_8(F)$.

Lemma 4.9. *Let $s \in \mathbf{G} = Spin_8(F)$ be a non-central semisimple element, and V the natural G -module. Suppose that s is almost cyclic on V .*

(1) *If λ is an eigenvalue of s on V of multiplicity $m_s(V) > 1$ then $\lambda \in \{1, -1\}$ and either 1 or -1 is not an eigenvalue of s .*

(2) *Let $G \cong {}^3D_4(q)$ and $s \in G$. Then s is regular.*

Proof. Set $\mathbf{H} = SO_8(F)$ and let $h := \mathbf{G} \rightarrow \mathbf{H}$ be the representation of \mathbf{G} with highest weight ω_1 .

(1) By Lemma 2.1, λ^{-1} is an eigenvalue of s on V of multiplicity equal to that of λ . As s is almost cyclic on V , we have $\lambda \in \{1, -1\}$. The 1-eigenspace and the -1 -eigenspace are well known to be non-degenerate and of even dimension. As s is almost cyclic on V , both 1, -1 cannot be eigenvalues of s . It follows that $C_{\mathbf{H}}(s)$ is connected, see [27, Lemma 2.2].

(2) Let $V_1 \subset V$ be the eigenspace of dimension $m_s(V)$. We have $V = V_1 \oplus V_2$ for some s -stable subspace V_2 of V , and s is cyclic on V_2 . Then $C_{\mathbf{H}}(h(s))$ contains the subgroup $\mathbf{Y}_1 := \text{diag}(SO(V_1), \text{Id})$.

Suppose the contrary, that s is not regular. Then $C_G(s)$ and hence $C_{\mathbf{H}}(h(s))$ contains a unipotent element $u \neq 1$. If $\dim V_1 = 2$ then u is a transvection, however, \mathbf{H} contains no transvection. So $\dim V_1 \in \{6, 4\}$.

It is known that $C_{\mathbf{G}}(s)$ is a reductive group and contains a maximal torus T of \mathbf{G} [19, Theorem 5.8].

Note that $\mathbf{G} = Spin_8(F)$ is the universal group of type D_4 . Let σ be the Steinberg endomorphism of \mathbf{G} such that $G = \mathbf{G}^\sigma := \{g \in \mathbf{G} : \sigma(g) = g\}$. In particular, $\sigma(s) = s$ and hence $\sigma(C_{\mathbf{G}}(s)) = C_{\mathbf{G}}(s)$. Then $C_G(s) = C_{\mathbf{G}}(s)^\sigma$ is a group of maximal rank in G in terms of [13]. Maximal subgroups M of maximal rank in G are determined in [13], and those containing a non-trivial unipotent element are listed in [13, Table 5.1]. This tells us that M contains a normal subgroup M_1 isomorphic either to $SL_2(q) \circ SL_2(q^3)$ or to $SL_3(q)$ or to $SU_3(q)$, and the quotient M/M_1 is abelian.

Let \mathbf{Y} is a semisimple part of $C_{\mathbf{G}}(s)$. Then $h(\mathbf{Y}) = \mathbf{Y}_1$.

Suppose first that $\dim V_1 = 6$. Then $\mathbf{Y} \cong D_3 \cong A_3 \cong SL_4(F)$.

Obviously, the restriction of σ to \mathbf{Y} is a Steinberg endomorphism of \mathbf{Y} . Steinberg endomorphisms of simple algebraic groups are classified, see [15, Theorem 22.5 and Table 22.1]. In particular, $C_{\mathbf{G}}(s)^\sigma$ contains $\mathbf{Y}^\sigma = SL_4(q)$ or $SU_4(q)$. However, none of these groups is contained in M .

Suppose that $\dim V_1 = 4$. Set $\mathbf{X} = C_{\mathbf{G}}(\mathbf{Y})^\circ$ so $\mathbf{X}V_i = V_i$ for $i = 1, 2$ and \mathbf{X} acts trivially on V_1 . Clearly, $\sigma(\mathbf{X}) = \mathbf{X}$. Set $\mathbf{K} = \mathbf{X}\mathbf{Y}$; then $\sigma(\mathbf{K}) = \mathbf{K}$ and the rank of \mathbf{K} equals 4. Therefore, $K := \mathbf{K}^\sigma$ is a subgroup of maximal rank in G and hence $K \subseteq M$ by the above.

Observe that $\mathbf{X} \cong \mathbf{Y} \cong Spin_4(F) \cong \mathbf{X}_1 \circ \mathbf{X}_2$, where $\mathbf{X}_1 \cong \mathbf{X}_2 \cong SL_2(F)$. As $\sigma(\mathbf{X}) = \mathbf{X}$, we have either (i) $\sigma(\mathbf{X}_i) = \mathbf{X}_i$ for $i = 1, 2$ or (ii) $\sigma(\mathbf{X}_1) = \mathbf{X}_2$ and $\sigma(\mathbf{X}_2) = \mathbf{X}_1$. We show that (ii) does not hold.

Indeed, by Steinberg's famous theorem [15, Theorem 21.5], a surjective endomorphism of \mathbf{G} is either Steinberg or an automorphism. It follows that σ^i is Steinberg for every $i > 0$. In particular, σ^2 stabilizes \mathbf{X}_1 and \mathbf{X}_2 . Moreover, σ^3 acts on maximal tori T of \mathbf{H} and of \mathbf{K} by sending $t \in T$ to t^{q^3} , and hence stabilizes \mathbf{X}_1 and \mathbf{X}_2 too. For a moment view σ as the abstract group isomorphism $\mathbf{H} \rightarrow \mathbf{H}$. Then $\sigma = \sigma^3\sigma^{-2}$ must stabilize \mathbf{X}_1 and \mathbf{X}_2 .

So (i) holds. Similarly, $\mathbf{Y} \cong Spin_4(F) \cong \mathbf{Y}_1 \circ \mathbf{Y}_2$, where $\mathbf{Y}_1 \cong \mathbf{Y}_2 \cong SL_2(F)$ and σ stabilizes both \mathbf{Y}_1 and \mathbf{Y}_2 . Therefore, each group $\mathbf{X}_1^\sigma, \mathbf{X}_2^\sigma, \mathbf{Y}_1^\sigma$ and \mathbf{Y}_2^σ contains a subgroup isomorphic to $SL_2(p)$, and hence K contains a subgroup isomorphic to some central product of four copies of $SL_2(p)$. However, one easily observes that M contains no subgroup of this kind. This contradiction completes the proof. \square

The following result extends Theorem 2.18 to finite groups of Lie type. Observe that we use isomorphisms $SL_4(q) \cong Spin_6^+(q)$ and $SU_4(q) \cong Spin_6^-(q)$ to interpret an irreducible 6-dimensional module for the first group as a twist of the natural module for the second one. Similarly, we use isomorphisms $Sp_4(q) \cong SO_5(q)$ to interpret the irreducible module with highest weight ω_2 for one of these groups as the natural module for another one.

Theorem 4.10. *Let G be a universal quasi-simple group of Lie type and ϕ a non-trivial irreducible F -representation of G . Let $s \in G$ be a non-regular non-central semisimple element. Suppose that $\phi(g)$ is almost cyclic. Then $G \in \{SL_{n+1}(q), n > 1, SU_{n+1}(q), n > 1, Spin_{2n+1}(q), n \geq 3, q \text{ odd}, Sp_{2n}(q), n \geq 2, Spin_{2n}^\pm(q), n \geq 3\}$ and $\dim \phi$ is equal to $n + 1, n + 1, 2n + 1, 2n, 2n$, respectively.*

Proof. It is well known that ϕ extends to a representation of a simply connected simple algebraic group \mathbf{G} such that $G = \mathbf{G}^\sigma$ for a suitable Steinberg endomorphism of \mathbf{G} . We have $s \in \mathbf{G}$. By Theorem 2.18, $\mathbf{G} \in \{SL_{n+1}(F), Spin_{2n+1}(F), n \geq 3, p > 2, Sp_{2n}(F), n \geq 2, Spin_{2n}(F), n \geq 3\}$ and $\dim \phi = n + 1, 2n + 1, 2n, 2n$, respectively. By the classification theorem of Steinberg endomorphisms of simple algebraic groups, $G = \mathbf{G}^\sigma \in \{SL_{n+1}(q), SU_{n+1}(q), Spin_{2n+1}(q), n \geq 3, q \text{ odd}, Sp_{2n}(q), n \geq 2, Spin_{2n}^\pm(q), n \geq 3, {}^3D_4(q), {}^2C_2(q)\}$. As $s \in G$ is non-regular, we exclude from this list the groups $SL_2(q) \cong SU_2(q)$ and ${}^2C_2(q)$, as they have no non-regular non-central element. The group ${}^3D_4(q)$ must be dropped in view of Lemma 4.9. Indeed, every irreducible representation of $G = {}^3D_4(q)$ of degree 8 is a twist of the natural representation of G with an automorphism τ , say, of G . Obviously, if $\tau(s) = t$ and $\rho(t)$ is almost cyclic then so is $\rho^\tau(s) = \rho(\tau(s))$. So Lemma 4.9(2) extends to every irreducible representation of G of degree 8. \square

Proof of Theorem 1.6. If ρ is viewed as a representation into $GL_n(F)$ then $\rho = \rho_1 \oplus \cdots \oplus \rho_k$, where ρ_1, \dots, ρ_k are Galois conjugate irreducible representations of G into $GL_{n/k}(F)$, see [11, Lemma 2.10.2]. Each $\rho_i, 1 \leq i \leq k$, extends to a representation $\tau_i : \mathbf{G} \rightarrow GL_{n/k}(F)$, where \mathbf{G} is a simple simply connected algebraic group such that $G = \mathbf{G}^\sigma$ for a Steinberg endomorphism σ of \mathbf{G} ([9, §2.11]). Obviously, $\tau_i(g)$ is almost cyclic for $i = 1, \dots, k$. Then the result follows from the definition of a nearly natural representation of G and Theorem 4.10. \square

Acknowledgement. This paper was initiated while visiting Donna Testerman following on our collaboration on [TZ2], [TZ21] and [TZ22]. The author would like to acknowledge the hospitality of the Ecole Polytechnique Fédérale de Lausanne. At the final stage a part of this work was performed during the author's visit to the Isaac Newton Institute for Mathematical Sciences at Cambridge, UK (June-July 2022) in the framework of the research program "Groups, representations and applications".

References

- [1] J. Bamberg and T. Penttila, Overgroups of cyclic Sylow subgroups of linear groups, *Comm. Algebra* 36(2008), 2503 – 2543.
- [2] A. Baranov and A. Zalesski, Weight zero in tensor-decomposable irreducible representations of simple algebraic groups, *J. Pure and Applied Algebra* 225(2021), 106768
- [3] N. Bourbaki, *Groupes et algèbres de Lie*, Masson, Chapitres 4,5 et 6, Paris, 1981.
- [4] N. Bourbaki, *Groupes et algèbres de Lie*, Chapitres 7 et 8, Hermann, Paris, 1975.
- [5] R. Carter, *Finite groups of Lie type: Conjugacy classes and complex characters*, Wiley, Chichester, 1985.

- [6] L. Di Martino and A.E. Zalesski, Almost cyclic elements in Weil representations of classical groups, *Comm. Algebra* 46(2018), 2767 – 2810.
- [7] L. Di Martino, M. Pellegrini and A.E. Zalesski, Semisimple almost cyclic elements in cross-characteristic representations of finite groups of Lie type, *J. Group Theory* 23(2020), 235 – 285.
- [8] R. Guralnick, T. Pentilla, C. Praeger and J. Saxl, Linear groups having certain large prime divisors, *Proc. London Math. Soc.* (3) 78(1999), 167 – 214.
- [9] J. Humphreys, *Modular representations of finite groups of Lie type*, Cambridge Univ. Press, Cambridge, 2006.
- [10] B. Huppert, Singer zyklen in classischen Gruppen, *Math. Z.* 117(1970), 141 – 150.
- [11] P. Kleidman and M. Liebeck, *Subgroup of classical groups*, Cambridge Univ. Press, Cambridge, 1990 (London Math. Soc. Lecture notes no.129).
- [12] M.W. Liebeck, On the orders of maximal subgroups of the finite classical groups, *Proc. London Math. Soc.* 50(1985), 426 – 446.
- [13] M.W. Liebeck, J. Saxl and G.M. Seitz, Subgroups of maximal rank in finite exceptional groups of Lie type, *Proc. London Math. Soc.* 65(1992), 297 – 325.
- [14] F. Lübeck, Small degree representations of finite Chevalley groups in defining characteristic, *LMS J. Comput. Math.* 4(2001), 135 – 169.
- [15] G. Malle and D. Testerman, *Linear algebraic groups and finite groups of Lie type*, Cambridge Univ. Press, Cambridge, 2011.
- [16] E. Plotkin, A. Semenov, N. Vavilov, Visual basic representations: the atlas, *Intern. J. Algebra and Computation* 8(1), 1998, no. 1, 61 – 95.
- [17] A.A. Premet, Weights of infinitesimally irreducible representations of Chevalley groups over a field of prime characteristic, *Math. USSR Sb.* **61** (1988), 167 – 183.
- [18] G.M. Seitz, ‘*The Maximal Subgroups of Classical Algebraic Groups*’, *Memoirs Amer. Math. Soc.*, no. **365**, Amer. Math. Soc., Providence, 1987.
- [19] T. Springer and R. Steinberg, Conjugacy classes, In: A. Borel et al., *Seminar on algebraic groups and related finite groups*, Lect. Notes in Math. vol. 131, Springer Verlag, Berlin, 1970, 167 – 266.
- [20] R. Steinberg, *Lectures on Chevalley groups*, Amer. Math. Soc. Univ. Lect. Series, vol. 66, Providence, Rhode Island, 2016.
- [21] I.D. Suprunenko and A.E. Zalesski, Fixed vectors for elements in modules for algebraic groups, *Intern. J. Algebra and Comp.* 17(2007), 1249 – 1261.

- [22] D. Testerman and A. Zalesski, Subgroups of simple algebraic groups containing maximal tori and representations with multiplicity 1 non-zero weights, *Transformation Groups* 20(2015), 831 – 861.
- [23] D. Testerman and A. Zalesski, Irreducible representations of simple algebraic groups in which a unipotent element is represented by a matrix with single non-trivial Jordan block, *J. Group Theory* 21(2018), 1 – 20.
- [24] D.M. Testerman and A.E. Zalesski, Spectra of non-regular elements in irreducible representations of simple algebraic groups, *North-Western European J. Math.* 7(2021), 185 – 212.
- [25] D.M. Testerman and A.E. Zalesski, Almost cyclic elements in irreducible representations of simple algebraic groups, arXiv:2203.02900v1 [math.RT], 6 March 2022; *Transformation Groups* (2023), to appear
- [26] A. Zalesski, A fragment of the decomposition matrix of special unitary group over a finite field, *Math. USSR, Izvestija* 36(1991), 23 – 39.
- [27] A. Zalesski, On the Steinberg character of an orthogonal group over a finite field, *J. Algebra and Appl.* 13(2014), 1250038-1 — 1250038-25.

Table 1: The irreducible representations of G whose non-zero weights are of multiplicity 1 and weight 0 is not one-dimensional

G	highest weight	weight 0 multiplicity	dimension
$A_n, n > 1, p \nmid (n+1)$	$\omega_1 + \omega_n$	n	$n^2 + 2n$
$(n, p) \neq (2, 3), p \mid (n+1)$	$\omega_1 + \omega_n$	$n - 1$	$n^2 + 2n - 1$
$A_3, p > 3$	$2\omega_2$	2	20
$B_n, n > 2, p \neq 2$	ω_2	n	$2n^2 + n$
$p \mid (2n+1)$	$2\omega_1$	n	$2n^2 + 3n - 1$
$p \nmid (2n+1)$	$2\omega_1$	$n+1$	$2n^2 + 3n$
$B_2, p \neq 2$	$2\omega_2$	2	10
$p \neq 2, 5$	$2\omega_1$	2	14
$C_n, n > 2,$ $(n, p) \neq (3, 3)$	$2\omega_1$	n	$2n^2 + n$
	ω_2	$n - 1$ if $p \nmid n$	$2n^2 - n - 1$
		$n - 2$ if $p \mid n,$	$2n^2 - n - 2$
$C_2, p \neq 2$	$2\omega_1$	2	10
$p \neq 2, 5$	$2\omega_2$	2	14
$C_4, p \neq 2, 3$	ω_4	2	36
$D_n, n > 3, p \neq 2$	$2\omega_1$	$n - 2$ if $p \mid n,$	$2n^2 + n - 2$ if $p \mid n,$
		$n - 1$ if $p \nmid n$	$2n^2 + n - 1$ if $p \nmid n$
	ω_2	n	$2n^2 - n - 1$
$p = 2$	ω_2	$n - (2, n)$	$2n^2 - n - (2, n)$
$E_6, p = 3$	ω_2	5	77
$p \neq 3$		6	78
$E_7, p \neq 2$	ω_1	7	133
$p = 2$		6	132
E_8	ω_8	8	248
$F_4, p = 2$	ω_1	2	26
$p \neq 2$		4	52
$p \neq 3$	ω_4	2	26
$G_2, p \neq 3$	ω_2	2	14

Table 2: Non-trivial p -restricted G -modules whose weights are one-dimensional

G	highest weight
A_1	$a\omega_1, \quad 1 \leq a < p$
$A_n, n > 1$	$a\omega_1, b\omega_n, \quad 0 \leq a, b < p, \quad \omega_i, \quad 1 < i < n$ $c\omega_i + (p-1-c)\omega_{i+1}, \quad i = 1, \dots, n-1, \quad 0 \leq c < p$
$B_n, n \geq 2$	ω_1, ω_n
$B_2, p \neq 2, 3$	$\omega_1, \omega_2, \frac{p-3}{2}\omega_1 + \omega_2, \frac{p-1}{2}\omega_1$
$C_n, n > 1, p = 2$	ω_1, ω_n
$C_n, n \geq 3, p \neq 2$	$\omega_1, \omega_{n-1} + \frac{p-3}{2}\omega_n, \frac{p-1}{2}\omega_n$
$C_2, p \neq 2, 3$	$\omega_1, \omega_2, \omega_1 + \frac{p-3}{2}\omega_2, \frac{p-1}{2}\omega_2$
$C_3, p \neq 2$	ω_3
$D_n, n > 3$	$\omega_1, \omega_{n-1}, \omega_n$
E_6	ω_1, ω_6
E_7	ω_7
$F_4, p = 3$	ω_4
$G_2, p \neq 3$	ω_1
$G_2, p = 3$	ω_1, ω_2

Academy of Sciences of Belarus, Department of Physics, Mathematics and Informatics,
66 Prospekt Nezaleznasti, Minsk, 220000, Belarus.
e-mail: alexandre.zalesski@gmail.com