

EXACT BOREL SUBALGEBRAS OF PATH ALGEBRAS OF QUIVERS OF DYNKIN TYPE A

MARKUS THURESSON

Abstract

Hereditary algebras are quasi-hereditary with respect to any adapted partial order on the indexing set of the isomorphism classes of their simple modules. For any adapted partial order on $\{1, \dots, n\}$, we compute the quiver and relations for the Ext-algebra of standard modules over the path algebra of a uniformly oriented linear quiver with n vertices. Such a path algebra always admits a regular exact Borel subalgebra in the sense of König and we show that there is always a regular exact Borel subalgebra containing the idempotents e_1, \dots, e_n and find a minimal generating set for it. For a quiver Q and a deconcatenation $Q = Q^1 \sqcup Q^2$ of Q at a sink or source v , we describe the Ext-algebra of standard modules over KQ , up to an isomorphism of associative algebras, in terms of that over KQ^1 and KQ^2 . Moreover, we determine necessary and sufficient conditions for KQ to admit a regular exact Borel subalgebra, provided that KQ^1 and KQ^2 do. We use these results to obtain sufficient and necessary conditions for a path algebra of a linear quiver with arbitrary orientation to admit a regular exact Borel subalgebra.

2020 Mathematics Subject Classification: 16D90, 16E30, 16G20

Keywords: Quasi-hereditary algebra; Modules with standard filtration; Ext-algebra; A-infinity algebra; Catalan number

1. INTRODUCTION

Highest weight categories were first introduced in [CPS88] with the purpose of axiomatizing certain phenomena arising naturally in the representation theory of complex semisimple Lie algebras. This prompted the definition of a quasi-hereditary algebra (see [Sco87]), which was shown by the authors to be precisely such a finite-dimensional algebra whose module category is equivalent to a highest weight category. Since their introduction, examples of quasi-hereditary algebras have been found to be abundant in representation theory. Classical examples include hereditary algebras, algebras of global dimension two, Schur algebras and blocks of BGG category \mathcal{O} . The main protagonists of the representation theory of quasi-hereditary algebras are the standard modules. These are certain quotients of the indecomposable projective modules which depend on a chosen partial order on the indexing set of the isomorphism classes of simple modules. For blocks of BGG category \mathcal{O} , this order is the Bruhat order. Associated to the standard modules is the category $\mathcal{F}(\Delta)$: the full subcategory of the module category consisting of those modules which admit a filtration whose subquotients are standard modules. Given a quasi-hereditary algebra, natural objectives of its representation theory are understanding the standard modules and the associated category $\mathcal{F}(\Delta)$.

In the endeavor of understanding $\mathcal{F}(\Delta)$ for a general quasi-hereditary algebra, a breakthrough was made by König, Külshammer and Ovsienko in [KKO14]. They showed that any quasi-hereditary algebra A is Morita equivalent to an algebra Λ , which admits a very particular subalgebra B , called a regular exact Borel subalgebra. This subalgebra has the surprising property that the image of its module category under the functor $\Lambda \otimes_B _$ is equivalent to $\mathcal{F}(\Delta)$.

In general, the problem of determining Λ and B for an arbitrary quasi-hereditary algebra A may be very hard. One way to do this involves determining an A_∞ -structure on $\text{Ext}_A^*(\Delta, \Delta)$, the algebra of extensions between standard modules over A , which can be arduous. Some examples of where this has been done can be found in [KS12; Thu22]. However, in the situations studied in this paper, the aforementioned A_∞ -structures can be easily computed. Moreover, we are able to use criteria from [Con21] to check when our algebras themselves contain regular exact Borel subalgebras.

A given associative algebra can have different quasi-hereditary structures, depending on the partial order mentioned above. In particular, two different orders may yield identical quasi-hereditary structures. This phenomenon motivates the definition of the essential order, an order with the property that two quasi-hereditary structures coincide if and only if they generate the same essential order. The algebras considered in this article are hereditary, and hereditary algebras are precisely those algebras which are quasi-hereditary with respect to any total order on an indexing set of the isomorphism classes of their simple modules. A natural question to ask is then how many different quasi-hereditary structures there are. When the algebra in question admits a duality on its module category which preserves simple modules, the structure is essentially unique, as shown by Coulembier in [Cou20]. This contrasts the algebras studied in this article with the examples from [KS12; Thu22], as the algebras appearing there admit a simple-preserving duality.

In a recent paper by Flores, Kimura and Rognerud, [FKR22], the authors showed that when A is the path algebra of a uniformly oriented linear quiver, the different quasi-hereditary structures are counted by the Catalan numbers. Moreover, they counted the quasi-hereditary structures of path algebras of more complicated quivers by means of a combinatorial process called “deconcatenation”. The main goal of the present article is to use the combinatorial techniques of Flores, Kimura and Rognerud to further study the quasi-hereditary structures of some of the algebras considered in [FKR22]. In particular, we want to describe the Ext-algebras of their standard modules and their regular exact Borel subalgebras.

The following is a brief description of the main results of the present article. For a quasi-hereditary algebra A , denote by Δ the direct sum of standard modules, one from each isomorphism class, and denote by $\text{Ext}_A^*(\Delta, \Delta)$ the algebra of extensions between standard modules.

(A) Let A_n be the path algebra of the following quiver:

$$1 \longrightarrow 2 \longrightarrow \dots \longrightarrow n-1 \longrightarrow n$$

For any partial order \trianglelefteq on $\{1, \dots, n\}$ such that (A_n, \trianglelefteq) is quasi-hereditary, we construct a graded quiver Q and an admissible ideal $I \subset KQ$, such that there is an isomorphism of graded associative algebras $\text{Ext}_{A_n}^*(\Delta, \Delta) \cong KQ/I$. According to results from [FKR22], each quasi-hereditary structure corresponds to a unique binary tree, and we show that this tree encodes all necessary information about extensions between the standard modules. For all details, we refer to Section 3.

- (B) Let A_n be as in (A). For any partial order \trianglelefteq on $\{1, \dots, n\}$ such that (A_n, \trianglelefteq) is quasi-hereditary, we check that A_n admits a regular exact Borel subalgebra. We show that A_n always admits a regular exact Borel subalgebra B which contains the idempotents e_1, \dots, e_n and find a minimal generating set for B . For all details, we refer to Section 3.2.
- (C) Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation of the quiver Q at a sink or source v . Put $A = KQ$ and $A^\ell = KQ^\ell$, for $\ell = 1, 2$. We describe $\text{Ext}_A^*(\Delta, \Delta)$ up to isomorphism in terms of the Ext-algebras of standard modules over A^ℓ , via a certain “gluing” process.
- (D) Given that A^ℓ admits a regular exact Borel subalgebra B^ℓ , for $\ell = 1, 2$, we show the following:
 - (i) If v is a source, A admits a regular exact Borel subalgebra.
 - (ii) If v is a sink, then A admits a regular exact Borel subalgebra if and only if v is minimal or maximal with respect to the essential order on Q_0 .

In the cases where A admits a regular exact Borel subalgebra, we construct a regular exact Borel subalgebra of A from B^1 and B^2 , using a similar “gluing” as in (C).

This article is organized in the following way: In Section 2 we give the necessary background on quasi-hereditary algebras and fix some notation. In Section 3, we recall the results of [FKR22] on the quasi-hereditary structures of A_n and expand upon them, proving (A). In Subsection 3.1, we compute the quiver and relations of the Ringel dual of A_n . Subsection 3.2 is devoted to the description of the regular exact Borel subalgebras of A_n . Subsection 3.3 briefly discusses A_∞ -structures on $\text{Ext}_{A_n}^*(\Delta, \Delta)$.

In Section 4, we give the background on deconcatenations from [FKR22] and prove (C). Subsection 4.1 discusses how regular exact Borel subalgebras behave under deconcatenations and contains the proof of (D). In Subsection 4.2, we apply our results to the case where Q is a linear quiver with arbitrary orientation.

2. NOTATION AND BACKGROUND

Throughout, let K be an algebraically closed field. For a quiver $Q = (Q_0, Q_1)$, denote by KQ path algebra of Q . Let $I \subset KQ$ be an admissible ideal and let A be the quotient $A = KQ/I$. We take A -module to mean finite-dimensional left A -module if nothing else is stated. For an arrow $\alpha \in Q_1$, denote by $s(\alpha)$ and $t(\alpha)$ the starting and terminal vertex of α , respectively. We adopt the convention of writing composition of arrows from right to left. That is, for vertices and arrows arranged as

$$x \xrightarrow{\alpha} y \xrightarrow{\beta} z,$$

we write the composition “first α , then β ” as $\beta\alpha$. We extend the notation of starting and terminal vertex to paths in Q , so if $p = \alpha_n \dots \alpha_1$, then we put $s(p) = s(\alpha_1)$ and $t(p) = t(\alpha_n)$.

The isomorphism classes of the simple A -modules are indexed by the vertex set, Q_0 , of Q . Denote by $L(i)$ the simple A -module associated to the vertex i . Denote by $P(i)$ and $I(i)$ the projective cover and injective envelope of $L(i)$, respectively. For an A -module M and a simple module $L(i)$, denote by $[M : L(i)]$ the Jordan-Hölder multiplicity of $L(i)$ in M . We denote the category of finite-dimensional left A -modules by $A\text{-mod}$.

1. Definition. [CPS88] Let A be a finite-dimensional algebra. Let $\{1, \dots, n\}$ be an indexing set for the isomorphism classes of simple A -modules and let \trianglelefteq be a partial order on $\{1, \dots, n\}$. Let the *standard module* at i , denoted by $\Delta(i)$, be the largest quotient of $P(i)$, whose composition factors $L(j)$ are such that $j \trianglelefteq i$. The algebra A is said to be *quasi-hereditary* with respect to \trianglelefteq if the following hold.

- (QH1) There is a surjection $P(i) \twoheadrightarrow \Delta(i)$ whose kernel admits a filtration with subquotients $\Delta(j)$, where $j \triangleright i$.
- (QH2) There is a surjection $\Delta(i) \twoheadrightarrow L(i)$ whose kernel admits a filtration with subquotients $L(j)$, where $j \triangleleft i$.

Equivalently, let the *costandard module* at i , denoted by $\nabla(i)$, be the largest submodule of $I(i)$, whose composition factors are such that $j \trianglelefteq i$. The algebra A is said to be *quasi-hereditary* with respect to \trianglelefteq if the following hold.

(QH1)' There is an injection $\nabla(i) \hookrightarrow I(i)$ whose cokernel admits a filtration with subquotients $\nabla(j)$, where $j \triangleright i$.

(QH2)' There is an injection $L(i) \hookrightarrow \nabla(i)$ whose cokernel admits a filtration with subquotients $L(j)$, where $j \triangleleft i$.

For a quasi-hereditary algebra A , it is natural to consider two particular subcategories of its module category, $A\text{-mod}$. These are $\mathcal{F}(\Delta)$, the full subcategory of $A\text{-mod}$ consisting of those modules which admit a filtration by standard modules, and $\mathcal{F}(\nabla)$, the full subcategory of $A\text{-mod}$ consisting of those modules which admit a filtration by costandard modules.

For an A -module $M \in \mathcal{F}(\Delta)$ (or $\mathcal{F}(\nabla)$), the number of occurrences of a particular standard module $\Delta(i)$ (or costandard module $\nabla(i)$) as a subquotient of a filtration of M is well-defined, and we denote this number by $(M : \Delta(i))$ (or $(M : \nabla(i))$).

In general, refining the partial order \trianglelefteq may produce different standard and costandard modules. Traditionally, only so-called adapted orders on $\{1, \dots, n\}$ are considered in order to avoid this.

2. Definition. [DR92] Let A be a finite-dimensional algebra. Let $\{1, \dots, n\}$ be an indexing set for the isomorphism classes of simple A -modules and let \trianglelefteq be a partial order on $\{1, \dots, n\}$. We say that \trianglelefteq is *adapted* to A if and only if for any A -module M such that

$$\text{top } M \cong L(i) \quad \text{and} \quad \text{soc } M \cong L(j),$$

where i and j are incomparable, there is $1 \leq k \leq n$ such that $i \triangleleft k$, $j \triangleleft k$ and $[M : L(k)] > 0$.

3. Lemma. [Con16] If (A, \trianglelefteq) is a quasi-hereditary algebra, then \trianglelefteq is adapted to A .

Two quasi-hereditary structures (A, \trianglelefteq_1) and (A, \trianglelefteq_2) are said to be equivalent if the sets of standard (and costandard) modules with respect to \trianglelefteq_1 and \trianglelefteq_2 coincide. We denote this relationship by $\trianglelefteq_1 \sim \trianglelefteq_2$. Then, more precisely:

$$\trianglelefteq_1 \sim \trianglelefteq_2 \iff \Delta_1(i) = \Delta_2(i) \wedge \nabla_1(i) = \nabla_2(i), \quad \forall i \in Q_0.$$

Equivalence of different quasi-hereditary structures is also captured precisely by the *essential order*.

4. Definition. [Cou20, Definition 1.2.5] Let (A, \trianglelefteq) be a quasi-hereditary algebra. Define the *essential order* \trianglelefteq^e of \trianglelefteq as the partial order transitively generated by the relations

$$i \trianglelefteq^e j \iff [\Delta(j) : L(i)] > 0 \quad \text{or} \quad (P(i) : \Delta(j)) > 0.$$

The essential order is related to equivalence of quasi-hereditary structures via

$$\trianglelefteq_1 \sim \trianglelefteq_2 \iff \trianglelefteq_1^e = \trianglelefteq_2^e.$$

2.1. Gluing of subalgebras. At various points in the present article, there will appear algebras which, intuitively, arise from gluing two subspaces of some ambient algebra at a shared dimension. Let X and Y be subspaces of some algebra A , which are closed under multiplication. Assume that $\dim X \cap Y = 1$ and choose complements X' and Y' of $X \cap Y$ in X and Y , respectively. Assume moreover that $X' \cdot Y' = Y' \cdot X' = 0$ and that $1_A \in \text{span}(X, Y)$. Put

$$C = X' \oplus Y' \oplus X \cap Y.$$

Then, C is a subalgebra of A . We call C the *gluing* of X and Y and write $C = X \diamond Y$. Note that when X and Y are graded, there is a natural grading on $X \diamond Y$ induced by the gradings on X and Y .

3. THE PATH ALGEBRA OF \mathbb{A}_n

In this section, we consider the path algebra of the uniformly oriented linear quiver

$$\mathbb{A}_n: 1 \xrightarrow{\alpha_1} 2 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_{n-2}} n-1 \xrightarrow{\alpha_{n-1}} n .$$

Throughout the section, we put $A_n = K\mathbb{A}_n$. The simple modules over A_n are indexed by the vertices $1, \dots, n$. Moreover, the algebra A_n is hereditary, and therefore, it is quasi-hereditary with respect to any adapted order $(\{1, \dots, n\}, \preceq)$ to A_n [DR92]. Recall that the indecomposable A_n -modules, up to isomorphism, are given by the interval modules, which are modules having Loewy diagrams of the following form:

$$M(i, j): \begin{array}{c} i \\ \downarrow \\ i+1 \\ \downarrow \\ \vdots \\ \downarrow \\ j-1 \\ \downarrow \\ j \end{array}$$

With this notation, we have

$$L(i) = M(i, i), \quad P(i) = M(i, n) \quad \text{and} \quad I(i) = M(1, i), \quad \forall 1 \leq i \leq n.$$

Homomorphisms and extensions between interval modules are well understood, and we summarize this information in the following proposition. Note that [OT10] uses different conventions than the present article.

5. Proposition. [OT10, Theorem 3.6] Let $M(i_1, j_1)$ and $M(i_2, j_2)$ be interval modules. Then, we have

(i)

$$\dim \text{Hom}_{A_n}(M(i_1, j_1), M(i_2, j_2)) = \begin{cases} 1 & \text{if } i_2 \leq i_1 \leq j_2 \leq j_1; \\ 0 & \text{otherwise.} \end{cases}$$

(ii)


$$\dim \text{Ext}_{A_n}^1(M(i_1, j_1), M(i_2, j_2)) = \begin{cases} 1 & \text{if } i_1 + 1 \leq i_2 \leq j_1 + 1 \leq j_2; \\ 0 & \text{otherwise.} \end{cases}$$

6. Definition. A *binary tree* T is either the empty set or a triple (s, L, R) , where s is a singleton set, called the *root* of T , and L and R are two binary trees, called the *left* and *right* subtrees of s , respectively. The empty set has one *leaf*, and the set of *leaves* of $T = (s, L, R)$ is the (disjoint) union of the sets of leaves of L and R .


A *binary search tree* is a binary tree, whose vertices are labeled by integers, such that if a vertex x is labeled by k , then the vertices of the left subtree of x are labeled by integers less than k , and the vertices of the right subtree of x are labeled by integers greater than k .

If T is a binary tree with n vertices, there exists a unique labelling of the vertices of T by the integers $1, \dots, n$, turning T into a binary search tree. The procedure by which this labelling is obtained is known as the *in-order algorithm*, which recursively visits the left subtree, then the root, then the right subtree. The first vertex visited is labeled by 1, the second by 2, and so on.

7. **Example.** Consider the following binary tree with 6 vertices.



With the in-order algorithm, the vertices of the tree are labeled as follows, creating a binary search tree.



Any binary search tree T with n vertices induces a partial order on the set of its vertices $\{1, \dots, n\}$. We denote this partial order by \trianglelefteq_T . It is defined in the following way.

$$i \triangleleft_T j \iff i \text{ labels a vertex in the subtree of the vertex labeled by } j.$$

In the above example, the partial order \trianglelefteq_T would be given by:

$$1 \triangleleft_T 2 \triangleleft_T 3, \quad 5 \triangleleft_T 4 \triangleleft_T 6 \triangleleft_T 3.$$

At this point, it is natural to ask for which binary trees T we obtain a quasi-hereditary algebra (A_n, \trianglelefteq_T) .

Recall that, since A_n is hereditary, A_n is quasi-hereditary with respect to any partial order which is adapted to A_n . It turns out that any partial order \trianglelefteq with respect to which A_n is quasi-hereditary, is equivalent (that is, produces the same set of standard and costandard modules) to a partial order produced by a binary tree. Conversely, for any binary tree T , the order \trianglelefteq_T makes A_n quasi-hereditary. More precisely, we have the following.

8. **Proposition.** [FKR22, Proposition 4.4] Let n be a natural number and let \mathcal{T} be the set of binary trees with n vertices. Denote by \mathcal{A} the set of adapted orders to A_n . For any partial order \trianglelefteq in \mathcal{A} , denote by $\overline{\trianglelefteq}$ the equivalence class of \trianglelefteq with respect to the relation \sim . Then, the map from \mathcal{T} to \mathcal{A}/\sim defined by

$$T \mapsto \overline{\trianglelefteq_T}$$

is a bijection.

Now, given that any quasi-hereditary structure on A_n may be associated to a binary tree, it is natural to describe this structure in terms of the associated binary tree. The following proposition is stated in the proof of [FKR22, Proposition 4.4]. For the convenience of the reader, we give a proof.

9. **Proposition.** Let T be a binary search tree and let \trianglelefteq_T be the associated adapted order to A_n . Then, we have the following.

- (i) The composition factors of the standard module $\Delta(i)$ are indexed by the labels of the vertices in the right subtree of the vertex labeled by i .
- (ii) The composition factors of the costandard module $\nabla(i)$ are indexed by the labels of the vertices in the left subtree of the vertex labeled by j .

Proof. By definition, the standard module $\Delta(i)$ is a quotient of the indecomposable projective module $P(i)$. Since $P(i) = M(i, n)$, this implies that there exists an integer s_i , which satisfies $i \leq s_i \leq n$, such that $\Delta(i) = M(i, s_i)$. The composition factors of $\Delta(i)$ are then $L(i), \dots, L(s_i)$, all occurring with Jordan-Hölder multiplicity 1. Denote by v_i the vertex labeled by i .

The composition factors $L(j)$ of $\Delta(i)$ must satisfy $v_j \triangleleft_T v_i$. By definition of \triangleleft_T , the vertices v_j such that $v_j \triangleleft_T v_i$ are exactly the vertices of the left and right subtree of v_i .

By construction of \triangleleft_T , the vertices in the left subtree of v_i are labeled by integers less than i , which shows that the corresponding simple modules may not be composition factors of $\Delta(i)$. Since $\Delta(i)$ is the maximal quotient of $P(i)$ such that its composition factors $L(j)$ satisfy $v_j \triangleleft_T v_i$, we are done.

The argument for the form of the costandard modules is similar. □

For a vertex v of T , denote by $\ell(v)$ and $r(v)$ the vertices immediately down and to the left or down and to the right of v , respectively. If such vertices do not exist, we write $\ell(v) = \emptyset$ or $r(v) = \emptyset$.

10. Proposition. Let T be a binary search tree and let \triangleleft_T be the associated adapted order to A_n .

- (i) Suppose that i labels the vertex $\ell(v)$ and that j labels the vertex v . Then, we have

$$\dim \text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = 1.$$

- (ii) Suppose that i labels the vertex $r(v)$ and that j labels the vertex v . Then, we have

$$\dim \text{Hom}_{A_n}(\Delta(i), \Delta(j)) = 1.$$

Proof. Put $\Delta(i) = M(i, s_i)$ and $\Delta(j) = M(j, s_j)$.

- (i) Note that the first vertex visited by the in-order algorithm, after visiting the right subtree of $\ell(v)$, is v . Therefore, by Proposition 9, we have $j = s_i + 1$. The condition of Proposition 5, part (ii), is then

$$i + 1 \leq s_i + 1 \leq s_i + 1 \leq s_j,$$

which is clearly satisfied.

- (ii) With the in-order algorithm, the vertex $r(v)$ is visited after v . Then, using Proposition 9, we know that if $\Delta(i) = M(i, s_i)$, then $\Delta(j) = M(i + k, s_i)$, for some $k \leq s_i - i$. The condition of Proposition 5, part (i), is then

$$i \leq i + k \leq s_i \leq s_i,$$

which is clearly satisfied. □

11. Lemma. (i) Let v_i and v_j be vertices labeled by i and j , respectively. Assume that v_j is in the left subtree of v_i . Then, we have

$$\text{Hom}_{A_n}(\Delta(i), \Delta(j)) = \text{Hom}_{A_n}(\Delta(j), \Delta(i)) = 0.$$

- (ii) Let v_i and v_j be vertices labeled by i and j , respectively. Assume that v_j is in the right subtree of v_i . Then, we have

$$\text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = \text{Ext}_{A_n}^1(\Delta(j), \Delta(i)) = 0.$$

- (iii) Let v_i and v_j be vertices, labeled by i and j , respectively. Assume that v_i is not in the subtree of v_j and that v_j is not in the subtree of v_i . Then, we have

$$\text{Hom}_{A_n}(\Delta(i), \Delta(j)) = \text{Hom}_{A_n}(\Delta(j), \Delta(i)) = \text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = \text{Ext}_{A_n}^1(\Delta(j), \Delta(i)) = 0.$$

Proof. By the proof of Proposition 9, we have $\Delta(i) = M(i, s_i)$ and $\Delta(j) = M(j, s_j)$, where the integers

$$i + 1, i + 2, \dots, s_i$$

label the vertices in the right subtree of v_i and the integers

$$j + 1, j + 2, \dots, s_j$$

label the vertices in the right subtree of v_j .

(i) According to the in-order algorithm, the integers $j + 1, \dots, s_j$ are less than i . We immediately have

$$\text{Hom}_{A_n}(\Delta(i), \Delta(j)) = 0,$$

since $v_j \triangleleft_T v_i$ and A_n is quasi-hereditary. Moreover by Proposition 5, part (ii), we have

$$\text{Hom}_{A_n}(\Delta(j), \Delta(i)) = 0,$$

since the condition $i \leq j \leq s_i \leq s_j$ is not satisfied, because $j < i$.

(ii) Since v_j is in the right subtree of v_i , we have $s_j \leq s_i$. We immediately have

$$\text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = 0,$$


since $v_j \triangleleft_T v_i$ and A_n is quasi-hereditary. Moreover, by Proposition 5, part (i), we have

$$\text{Ext}_{A_n}^1(\Delta(j), \Delta(i)) = 0,$$

since the condition $i + 1 \leq j \leq s_i + 1 \leq s_j$ is not satisfied, because $s_j \leq s_i$.

(iii) This is immediate, since A_n is quasi-hereditary and the vertices v_i and v_j are incomparable with respect to \trianglelefteq_T . \square

12. **Example.** Consider the following binary search tree, T , with vertices labeled according to the in-order algorithm.



The partial order \trianglelefteq_T is given by:

$$1 \triangleleft_T 2 \triangleleft_T 4, \quad 3 \triangleleft_T 2 \triangleleft_T 4, \quad \text{and} \quad 6 \triangleleft_T 5 \triangleleft_T 4.$$

For the standard and costandard modules, we have

$$\begin{aligned} \Delta(1) \cong L(1), \quad \Delta(2) \cong M(2, 3), \quad \Delta(3) \cong L(3), \quad \Delta(4) \cong P(4), \quad \Delta(5) \cong P(6), \quad \Delta(6) \cong L(6), \\ \nabla(1) \cong L(1), \quad \nabla(2) \cong M(1, 2), \quad \nabla(3) \cong L(3), \quad \nabla(4) \cong M(1, 4), \quad \nabla(5) \cong L(5), \quad \text{and} \quad \nabla(6) \cong L(6), \end{aligned}$$

which we see by applying Proposition 9. Moreover, we have non-split short exact sequences

$$\Delta(2) \hookrightarrow I(3) \twoheadrightarrow \Delta(1) \quad \text{and} \quad \Delta(4) \hookrightarrow P(2) \twoheadrightarrow \Delta(2),$$

giving us the extensions guaranteed by part (i) of Proposition 10.

Next, we see there are natural monomorphisms $\Delta(3) \hookrightarrow \Delta(2)$ and $\Delta(6) \hookrightarrow \Delta(5) \hookrightarrow \Delta(4)$, giving us the homomorphisms guaranteed by part (ii) of Proposition 10.

Lastly, we have

$$\begin{aligned} \dim \operatorname{Hom}_{A_n}(\Delta(1), \Delta(2)) &= \dim \operatorname{Hom}_{A_n}(\Delta(2), \Delta(1)) = \dim \operatorname{Hom}_{A_n}(\Delta(1), \Delta(3)) = \dim \operatorname{Hom}_{A_n}(\Delta(3), \Delta(1)) = 0; \\ \dim \operatorname{Hom}_{A_n}(\Delta(2), \Delta(4)) &= \dim \operatorname{Hom}_{A_n}(\Delta(4), \Delta(2)) = \dim \operatorname{Hom}_{A_n}(\Delta(1), \Delta(4)) = \dim \operatorname{Hom}_{A_n}(\Delta(4), \Delta(1)) = 0; \\ \dim \operatorname{Hom}_{A_n}(\Delta(2), \Delta(5)) &= \dim \operatorname{Hom}_{A_n}(\Delta(5), \Delta(2)) = \dim \operatorname{Hom}_{A_n}(\Delta(2), \Delta(6)) = \dim \operatorname{Hom}_{A_n}(\Delta(6), \Delta(2)) = 0; \\ \dim \operatorname{Hom}_{A_n}(\Delta(1), \Delta(5)) &= \dim \operatorname{Hom}_{A_n}(\Delta(5), \Delta(1)) = \dim \operatorname{Hom}_{A_n}(\Delta(1), \Delta(6)) = \dim \operatorname{Hom}_{A_n}(\Delta(6), \Delta(1)) = 0, \end{aligned}$$


as prescribed by Lemma 11.

13. **Lemma.** Let $v, r(v)$ and $\ell(r(v))$ be vertices labeled by i, j and k , respectively. Then, the multiplication map

$$\operatorname{Hom}_{A_n}(\Delta(j), \Delta(i)) \times \operatorname{Ext}_{A_n}^1(\Delta(k), \Delta(j)) \rightarrow \operatorname{Ext}_{A_n}^1(\Delta(k), \Delta(i))$$

is the zero map.

Proof. Note that the vertices i, j and k are configured in the following way:




The multiplication map in question produces an extension in the space $\operatorname{Ext}_{A_n}^1(\Delta(k), \Delta(i))$. But since $\ell(r(v))$ is a vertex in the right subtree of v , this space is zero, by part (ii) of Lemma 11. \square

14. **Lemma.** Let $v, \ell(v)$ and $r(\ell(v))$ be vertices labeled by i, j and k , respectively. Then, the multiplication map

$$\operatorname{Ext}_{A_n}^1(\Delta(j), \Delta(i)) \times \operatorname{Hom}_{A_n}(\Delta(k), \Delta(j)) \rightarrow \operatorname{Ext}_{A_n}^1(\Delta(k), \Delta(i))$$

is non-zero.

Proof. We start by observing that the vertices labeled by i, j and k are configured in the following way:



According to the in-order algorithm, we have $j < k < i$. Note that, since A_n is hereditary, a minimal projective resolution of a standard module $\Delta(x)$ is of the form

$$\ker p_x \longrightarrow P(x) \xrightarrow{p_x} \Delta(x) .$$

Consider the following picture:

$$\begin{array}{ccccc} \ker p_k & \longrightarrow & P(k) & \xrightarrow{p_k} & \Delta(k) \\ \downarrow f & & \downarrow & & \\ \ker p_j & \longrightarrow & P(j) & \xrightarrow{p_j} & \Delta(j) \\ \downarrow g & & \downarrow & & \\ \ker p_i & \longrightarrow & P(i) & \xrightarrow{p_i} & \Delta(i) \end{array}$$

An extension from $\Delta(k)$ to $\Delta(i)$ is represented by a chain map, which has one (possibly) non-zero component, namely the map $g \circ f$. Now, assume that $\Delta(k) = M(k, s_k)$ and $\Delta(i) = M(i, s_i)$. Then, we have $\ker p_k = P(s_k + 1)$ and $\ker p_i = P(s_i + 1)$. Since $k < i$ and $s_k < s_i$, we have $\text{Hom}_{A_n}(P(k), P(i)) = \text{Hom}_{A_n}(\ker p_k, \ker p_i) = 0$. Therefore, the chain map, if it is non-zero, cannot be null-homotopic. By construction, f is the unique (up to a scalar) non-zero map from $M(s_k + 1, n)$ to $M(s_j + 1, n)$. Next, note that in our configuration, the first vertex visited by the in-order algorithm, after visiting the entire right subtree of $\ell(v)$, is v . This implies that $s_j + 1 = i$, so that the map g in the above picture is some scalar multiple of the identity homomorphism on $P(i)$. This shows that the composition $g \circ f$, and hence the chain map representing our extension, is non-zero. \square

For two interval modules $M(i_1, j_1)$ and $M(i_2, j_2)$, the space $\text{Hom}_{A_n}(M(i_1, j_1), M(i_2, j_2))$ is one-dimensional if it is non-zero. Fix a basis of this space, corresponding to the following homomorphism of representations:

$$\begin{array}{ccccccc} K & \longleftarrow & \dots & \longleftarrow & K & \longleftarrow & \dots & \longleftarrow & K \\ & & & & \downarrow 1 & & \downarrow 1 & & \downarrow 1 \\ & & & & K & \longleftarrow & \dots & \longleftarrow & K & \longleftarrow & \dots & \longleftarrow & K \end{array}$$

Note that this choice of basis also gives an obvious choice of basis for the spaces $\text{Ext}_{A_n}^1(M(i_1, j_1), M(i_2, j_2))$. Suppose now that we have chosen such basis vectors

$$y \in \text{Hom}_{A_n}(\Delta(i), \Delta(j)); \quad x \in \text{Ext}_{A_n}^1(\Delta(j), \Delta(k)); \quad z \in \text{Ext}_{A_n}^1(\Delta(i), \Delta(k));$$

and that i, j and k are as in the statement of Lemma 14. Then, we have $xy = z$.

15. Theorem. Let T be a binary search tree with n vertices, with the vertices labeled according to the in-order algorithm, and let \preceq_T be the associated adapted order to A_n . Construct a quiver Q in the following way:


- (i) The vertices of Q are $1, 2, \dots, n$.
- (ii) We draw an edge between i and j if and only if the integers i and j label either a set of vertices $\{v, \ell(v)\}$ or $\{v, r(v)\}$. If i labels $\ell(v)$ and j labels v , the orientation of the edge is from $\ell(v)$ to v , of degree 1, and is denoted by ε_i^j . If i labels $r(v)$ and j labels v , the orientation of the edge is from $r(v)$ to v , of degree 0, and is denoted by f_i^j .

Let $I \subset KQ$ be the ideal generated by the following elements:


- (•) $f_j^k \varepsilon_i^j$, for all i, j and k ;
- (•) $\varepsilon_j^k \varepsilon_i^j$, for all i, j and k ;

Then, there is an isomorphism of graded algebras $KQ/I \cong \text{Ext}_{A_n}^*(\Delta, \Delta)$.

16. Example. Consider the binary search tree



Then, the corresponding quiver Q is



and the ideal $I = \langle \varepsilon_2^4 \varepsilon_1^2 \rangle$. The full arrows, ε_1^2 and ε_2^4 , are of degree 1, while the dashed arrows, f_3^2 , f_5^4 and f_6^5 are of degree 0.

Proof of Theorem 15. Fix basis vectors $\varphi_i^j \in \text{Hom}_{A_n}(\Delta(i), \Delta(j))$ and $e_i^j \in \text{Ext}_{A_n}^1(\Delta(i), \Delta(j))$ of all non-zero spaces of the form $\text{Hom}_{A_n}(\Delta(i), \Delta(j))$ and $\text{Ext}_{A_n}^1(\Delta(i), \Delta(j))$, as discussed prior to Theorem 15. Define a map $\Phi : KQ/I \rightarrow \text{Ext}_{A_n}^*(\Delta, \Delta)$ by

$$\varepsilon_i^j \mapsto e_i^j, \quad f_i^j \mapsto \varphi_i^j, \quad e_i \mapsto 1_{\Delta(i)},$$

and extend by linearity. By applying Lemma 13 and Lemma 11, along with the fact that there are no extensions of degree greater than 1, since A_n is hereditary, we see that $I \subset \ker \Phi$, so that Φ is well-defined.

To check that Φ is a homomorphism of algebras, it is enough to check compatibility of Φ with products of the form $\varepsilon_j^k f_i^j$. We have:

$$\Phi(\varepsilon_j^k f_i^j) = \Phi(\varepsilon_i^k) = e_i^k = e_j^k \varphi_i^j = \Phi(\varepsilon_j^k) \Phi(f_i^j).$$

As the image of Φ contains bases of $\text{Hom}_{A_n}(\Delta(i), \Delta(j))$ and $\text{Ext}_{A_n}^1(\Delta(i), \Delta(j))$ for all i and j , Φ is surjective.

Next, we compare dimensions. Consider the degree 0 part of the space $e_j KQ/I e_i$. The dimension of this space is equal to the number of paths of degree 0 from i to j in Q . Clearly, such a path must be a product of the form $f_{k_s}^j f_{k_{s-1}}^j \dots f_{k_1}^{k_2} f_i^{k_1}$. Since vertices in Q have out-degree at most 1, it follows that such a path is unique. So the dimension of the degree 0 part $e_j KQ/I e_i$ is either 0 or 1. By construction of Q , it is now clear that this dimension coincides with $\dim \text{Hom}_{A_n}(\Delta(i), \Delta(j))$. Next, consider the degree 1 part of the space $e_j KQ/I e_i$. Similarly, the dimension of this space is either 0 or 1, depending on the number of paths from i to j in Q . Such a path is either of the form ε_i^j or $\varepsilon_k^j f_i^k$. Again, by construction of Q , we see that this dimension coincides with $\text{Ext}_{A_n}^1(\Delta(i), \Delta(j))$. Therefore, Φ is a surjective linear map between vector spaces of the same dimension, hence an isomorphism. \square

3.1. Ringel dual. We recall that, for any quasi-hereditary algebra A , there exists a module T , called the characteristic tilting module, whose additive closure equals $\mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. The characteristic tilting module decomposes as

$$T = \bigoplus_{i=1}^n T(i),$$

where each $T(i)$ is indecomposable. The Ringel dual is then, by definition, $R(A) = \text{End}(T)^{\text{op}}$. In this section, we compute quiver and relations of the Ringel dual of (A_n, \triangleleft_T) using the associated binary search tree. The following statement is part of the proof of Theorem 4.6 in [FKR22]. For the convenience of the reader, we give a proof.

17. Proposition. Let ν be a vertex labeled by i . The indecomposable summand $T(i)$ of the characteristic tilting module is the interval module $M(t_i, s_i)$, where t_i is the least integer labelling a vertex in the left subtree of ν and s_i is the greatest integer labelling a vertex in the right subtree of ν .

Proof. To prove that $T(i) = M(t_i, s_i)$, it suffices to show that $\Delta(i)$ is isomorphic to a submodule of $T(i)$, that the cokernel of this monomorphism admits a filtration by standard modules, and that $T(i)$ is contained in $\mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$ [Rin91, Proposition 2]. We proceed by induction on the size of the subtree of v . The basis of the induction is clear, since if $\ell(v) = r(v) = \emptyset$, we have $\Delta(i) = \nabla(i) = L(i) = T(i)$. Since $t_i \leq i$, it is clear that $\Delta(i)$ embeds into $M(t_i, s_i)$. Clearly, the cokernel of this embedding is the module $M(t_i, i-1)$.

Suppose that j labels the vertex $\ell(v)$. The composition factors of $M(t_i, i-1)$ are labeled by the vertices in the subtree of $\ell(v)$, so by the induction hypothesis, $M(t_i, i-1) = T(j)$. In other words, we have a short exact sequence

$$0 \rightarrow \Delta(i) \hookrightarrow M(t_i, s_i) \twoheadrightarrow T(j) \rightarrow 0.$$

Since $T(j) \in \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$, we have $M(t_i, s_i) \in \mathcal{F}(\Delta)$. By a similar argument, we see that there is a short exact sequence

$$0 \rightarrow T(k) \hookrightarrow T(i) \twoheadrightarrow \nabla(i) \rightarrow 0,$$


where k labels the vertex $r(v)$. Then, since $T(k) \in \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$, we get that $M(t_i, s_i) \in \mathcal{F}(\nabla)$. Now, $M(t_i, s_i)$ is a module such that $\Delta(i)$ embeds into it, the cokernel of this embedding is contained in $\mathcal{F}(\Delta)$, and $M(t_i, s_i) \in \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. This confirms that $M(t_i, s_i) = T(i)$. \square

18. Corollary. Let the vertices v , $\ell(v)$ and $r(v)$ be labeled by i, j and k , respectively. Then, for any v , there is an epimorphism $p_i^j: T(i) \twoheadrightarrow T(j)$ and a monomorphism $q_k^i: T(k) \hookrightarrow T(i)$.

19. Proposition. Let the vertices $\ell(\ell(v)), \ell(v), v, r(v)$ and $r(r(v))$ be labeled by x, j, i, k and y , respectively. Let p_i^j, p_j^x, q_k^i and q_y^k be the homomorphisms from Corollary 18. Then, we have:

- (i) $p_j^x p_i^j \neq 0$ and $q_k^i q_y^k \neq 0$;
- (ii) $p_i^j q_k^i = 0$;
- (iii) $\text{Hom}_{A_n}(T(j), T(i)) = 0$;
- (iv) $\text{Hom}_{A_n}(T(i), T(k)) = 0$.

Proof. The vertices are configured in the following way:



- (i) This is immediate, as $p_j^x p_i^j$ is the composition of two epimorphisms and $q_k^i q_y^k$ is the composition of two monomorphisms.
- (ii) Applying Proposition 17, we see that $T(k)$ and $T(j)$ have no common composition factors, which implies that $\text{Hom}_{A_n}(T(k), T(j)) = 0$, yielding the statement.
- (iii) Put $T(j) = M(t_j, s_j)$ and $T(i) = M(t_i, s_i)$, where $t_j = t_i$ and $s_j < s_i$. Then, using Proposition 5, the condition for $\text{Hom}_{A_n}(T(j), T(i))$ being non-zero is

$$t_i \leq t_j \leq s_i \leq s_j,$$

which is not satisfied, since $s_j < s_i$.

- (iv) Similar to (iii). \square


With these observations established, we are ready to describe the Ringel dual $R(A_n)$.

20. Theorem. Let T be a binary search tree with n vertices, with the vertices labeled according to the in-order algorithm, and let \preceq_T be the associated adapted order to A_n . Construct a quiver Q in the following way:


- (i) The vertices of Q are $1, \dots, n$.
- (ii) We draw an edge between i and j if and only if the integers i and j label either a set of vertices $\{v, \ell(v)\}$ or $\{v, r(v)\}$. If i labels v and j labels $\ell(v)$, the edge is denoted by f_i^j and its orientation is from i to j . If i labels v and j labels $r(v)$, the edge is denoted by g_j^i and its orientation is from j to i .

Let $I \subset KQ$ be the ideal generated by the elements $f_i^j g_k^i$, for all i, j and k . Then, there is an isomorphism of algebras $KQ/I \cong \text{End}(T)$, and consequently, $R(A_n) \cong KQ/I^{\text{op}}$.

21. **Example.** Consider the binary search tree



Then, the corresponding quiver Q is



and the ideal $I \subset KQ$ is generated by the elements $f_4^2 g_5^4$ and $f_2^1 g_3^2$.

Proof of Theorem 20. Recall the discussion prior to Theorem 15, which implies that for composable homomorphisms between indecomposable summands of T , we may choose basis vectors

$$x \in \text{Hom}_{A_n}(T(i), T(j)); \quad y \in \text{Hom}_{A_n}(T(j), T(k)); \quad z \in \text{Hom}_{A_n}(T(i), T(k))$$

such that $xy = z$. Define a map $\tilde{\Psi} : KQ \rightarrow \text{End}(T)$ by

$$f_i^j \mapsto p_i^j, \quad g_i^j \mapsto q_i^j, \quad e_i \mapsto 1_{T(i)}$$

and extend by linearity. Using Proposition 19, part (ii), we see that $I \subset \ker \tilde{\Psi}$, so that $\tilde{\Psi}$ factors uniquely through the quotient KQ/I . This gives us a well-defined homomorphism of algebras $\Psi : KQ/I \rightarrow \text{End}(T)$. Since the homomorphisms p_i^j, q_k^i and $1_{T(i)}$ generate $\text{End}(T)$, Ψ is surjective. It is now easy to see that $\dim KQ/I = \dim \text{End}(T)$, so that Ψ is a surjective linear map between spaces of the same dimension, hence an isomorphism. \square

3.2. **Regular exact Borel subalgebras of A_n .** In this section, we investigate when (A_n, \trianglelefteq_T) admits a regular exact Borel subalgebra. Recall the following.

22. **Definition.** [Kön95; BKK20, Definition 3.4] Let $(\Lambda, \trianglelefteq)$ be a quasi-hereditary algebra with n simple modules, up to isomorphism. Then, a subalgebra $B \subset \Lambda$ is called an *exact Borel subalgebra* provided that

- (i) B also has n simple modules up to isomorphism and (B, \trianglelefteq) is quasi-hereditary with simple standard modules,
- (ii) the functor $\Lambda \otimes_B _$ is exact, and
- (iii) there are isomorphisms $\Lambda \otimes_B L_B(i) \cong \Delta_\Lambda(i)$.

If, in addition, the map $\text{Ext}_B^k(L_B(i), L_B(j)) \rightarrow \text{Ext}_\Lambda^k(\Lambda \otimes_B L_B(i), \Lambda \otimes_B L_B(j))$ induced by the functor $\Lambda \otimes_B _$ is an isomorphism for all $k \geq 1$ and $i, j \in \{1, \dots, n\}$, $B \subset \Lambda$ is called a *regular exact Borel subalgebra*. If this map is an isomorphism for all $k \geq 2$ and an epimorphism for $k = 1$, $B \subset \Lambda$ is called a *homological Borel exact Borel subalgebra*.

23. Theorem. [Con21, Theorem D, part 5] Every quasi-hereditary algebra equivalent to (A_n, \triangleleft_T) has a regular exact Borel subalgebra if and only if $\text{rad} \Delta(i)$ is contained in $\mathcal{F}(\nabla)$, for all $i \in \{1, \dots, n\}$.

24. Proposition. (A_n, \triangleleft_T) has a regular exact Borel subalgebra.

Proof. Let the vertex v be labeled by i and consider $\Delta(i) = M(i, s_i)$. Then, $\text{rad} \Delta(i) = M(i+1, s_i)$. Let the vertex $r(v)$ be labeled by k . According to the in-order algorithm, the first vertex visited immediately after v is the left-most vertex in the subtree of $r(v)$. This shows, in our earlier notation, that $t_k = i+1$. This fact implies that

$$M(i+1, s_i) = M(t_k, s_i) = M(t_k, s_k) = T(k),$$

according to the proof of Proposition 17. Since $T(k) \in \mathcal{F}(\nabla)$, the statement follows from Theorem 23. \square

Thanks to the observations made earlier in the current section, it is fairly easy to find the quiver, Q_B , of a regular exact Borel subalgebra $B \subset A_n$. Recall that, if $B \subset A_n$ is a regular exact Borel subalgebra, then there are isomorphisms of vector spaces

$$\text{Ext}_B^k(L(i), L(j)) \cong \text{Ext}_{A_n}^k(\Delta(i), \Delta(j)),$$

for all $k \geq 1$ and $i, j \in \{1, \dots, n\}$. In particular, the spaces above have the same dimension. In our setup, these spaces are zero for $k > 1$. It is well-known that

$$\dim \text{Ext}_B^1(L(i), L(j)) = \text{number of arrows from } i \text{ to } j \text{ in } Q_B.$$

This means that we can determine the quiver Q_B by computing extensions between standard modules over A_n . To this end, we apply Theorem 15, resulting in the following.

25. Proposition. Let $B \subset A_n$ be a regular exact Borel subalgebra. Then, there is an isomorphism of algebras $B \cong KQ_B$, where Q_B is the following quiver.

- (i) The vertex set of Q_B is $\{1, \dots, n\}$.
- (ii) There is an arrow $i \rightarrow j$ if and only if $\dim \text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = 1$.

Next, we wish to determine how to realize B as a subalgebra of A_n . In this endeavour, we are aided by the following results.

26. Theorem (Wedderburn-Mal'tsev). [Wed08; Mal42] Let A be a finite-dimensional algebra over an algebraically closed field. Then, there is a semi-simple subalgebra S of A such that

$$A = \text{rad } A \oplus S$$

as a vector space. Moreover, if T is another semisimple subalgebra of A such that $A = \text{rad } A \oplus T$, then there is an inner automorphism σ of A such that $\sigma(S) = T$.

We remark that if S is a semi-simple subalgebra as in the theorem and T is another semi-simple subalgebra with $\dim_K S = \dim_K T$, then necessarily $A = \text{rad } A \oplus T$, as any semi-simple subalgebra intersects $\text{rad } A$ trivially. Consequently, S and T are conjugate.

Let $A = KQ/I$ be the path algebra of some quiver modulo an admissible ideal and let $S \subset A$ be the subalgebra $S = \langle e_1, \dots, e_n \rangle$. Then, S is semi-simple and $A = \text{rad } A \oplus S$. If $\{d_1, \dots, d_n\}$ is some other complete set of primitive orthogonal idempotents, then $T = \langle d_1, \dots, d_n \rangle$ is another semi-simple subalgebra of A with $\dim_K S = \dim_K T$, so S and T must be conjugate. We conclude that for any two complete sets of primitive orthogonal idempotents in A , there is an inner automorphism of A mapping one to the other.

27. Theorem. [Zha00, Proposition 3.4] Let A be a finite-dimensional quasi-hereditary algebra and let $B \subset A$ be an exact Borel subalgebra. Then, $C = \sigma(B)$ is again an exact Borel subalgebra.

28. Theorem. [Zha00, Theorem 3.6] Let A be a finite-dimensional quasi-hereditary algebra and let $B \subset A$ be an exact Borel subalgebra. Then, $B \subset A$ is unique up to an inner automorphism of A , that is, if $B' \subset A$ is another exact Borel subalgebra, then B and B' are conjugate.

Let $\{d_1, \dots, d_n\}$ be a complete set of primitive orthogonal idempotents in B . For any complete set of primitive orthogonal idempotents $\{f_1, \dots, f_n\}$ of A_n , there is an inner automorphism σ of A_n mapping $\{d_1, \dots, d_n\}$ to $\{f_1, \dots, f_n\}$. But by Theorem 27, $\sigma(B)$ is an exact Borel subalgebra of A_n , and by construction it contains the idempotents $\{f_1, \dots, f_n\}$. We conclude that for any complete set of primitive orthogonal idempotents $\{f_1, \dots, f_n\}$ in A_n , there is an exact Borel subalgebra C containing them. In particular, there is an exact Borel subalgebra containing the idempotents e_1, \dots, e_n .

29. Example. Consider A_2 , the path algebra of $1 \xrightarrow{\alpha} 2$ and let $B \subset A_2$ be a regular exact Borel subalgebra. Fix the order $1 \triangleleft 2$. Then we have

$$\Delta(1) \cong L(1) \quad \text{and} \quad \Delta(2) \cong L(2).$$

Therefore, the quiver of B coincides with the quiver of A_2 , and the (unique) regular exact Borel subalgebra is $B = A_2$. If we instead fix the order $2 \triangleleft 1$, we have

$$\Delta(1) \cong P(1) \quad \text{and} \quad \Delta(2) \cong P(2).$$

Since there are no non-split extensions between projective modules, the quiver of B has no arrows. Thus, we have $B \cong K \times K$. The obvious choice for B is then the subalgebra $B = \langle e_1, e_2 \rangle$. Suppose $xe_1 + ye_2 + z\alpha \in A_2$ is an idempotent. Then we have:

$$(xe_1 + ye_2 + z\alpha)^2 = x^2e_1 + y^2e_2 + z(x+y)\alpha = xe_1 + ye_2 + z\alpha$$

This implies that all possible complete sets of primitive orthogonal idempotents are $\{e_1 + z\alpha, e_2 - z\alpha\}$, where $z \in K$ is some scalar. Thus, we obtain exact Borel subalgebras $B_z = \langle e_1 + z\alpha, e_2 - z\alpha \rangle$. Since $z\alpha \in \text{rad } A_2$, the element $1 - z\alpha$ is invertible (with inverse $1 + z\alpha$). We see that

$$(1 + z\alpha)e_1(1 - z\alpha) = 1 + z\alpha \quad \text{and} \quad (1 + z\alpha)e_2(1 - z\alpha) = 1 - z\alpha,$$

so for all $z \neq 0$, the subalgebras B and B_z are conjugate.

Next, we want to argue as in the discussion after Theorem 28 to conclude that there is always a regular exact Borel subalgebra containing the idempotents e_1, \dots, e_n . That argument relies on [Zha00, Proposition 3.4], which states that inner automorphisms of quasi-hereditary algebras preserve exact Borel subalgebras. Our next goal is to extend this statement to regular and homological exact Borel subalgebras.

For any automorphism σ of A , there is an endofunctor Φ_σ of $A\text{-mod}$ defined by taking an A -module V to the module having the same underlying vector space, but with the action defined by $a \cdot_\sigma v := \sigma(a) \cdot v$. For ease of notation, we denote $\Phi_\sigma(V)$ by ${}_\sigma V$, indicating that the action of A is twisted by the automorphism σ . If $B \subset A$ is a subalgebra, then so is $C := \sigma(B)$. It is clear that Φ_σ restricts, in a natural way, to a functor $C\text{-mod} \rightarrow B\text{-mod}$.

Let b_1, \dots, b_n be some complete set of primitive orthogonal idempotents for $B \subset A$. Then, the set $\{b_1, \dots, b_n\}$ indexes the isomorphism classes of simple B -modules, and we denote by $L_B(i)$, where $1 \leq i \leq n$, the simple module having as a basis the idempotent b_i and the action defined by letting b_i act as the identity and other basis elements as 0. Since $\sigma : B \rightarrow C$ is an isomorphism, $\sigma(b_1), \dots, \sigma(b_n)$ is a complete set of primitive orthogonal idempotents for C , and putting $\sigma(b_i) = c_i$, for $1 \leq i \leq n$, we denote by $L_C(i)$ the simple C -module having as a basis the idempotent c_i and the action defined by letting c_i act as the identity and other basis elements as 0. With this notation, consider the module ${}_\sigma L_C(i)$ and fix the basis c_i . On this basis, the idempotent acts as $b_i \cdot_\sigma c_i := \sigma(b_i) \cdot c_i = c_i \cdot c_i = c_i$. Thus, with this notation, we have that ${}_\sigma L_C(i) = L_B(i)$.

30. Lemma. Let A be an algebra and let $B \subset A$ be a subalgebra. Let σ be an inner automorphism of A , given by $\sigma(x) = u^{-1}xu$ for some invertible element u and $x \in A$. Put $C = \sigma(B)$.

- (i) There is an isomorphism of A - A -bimodules $A \cong_\sigma A$ given by left multiplication with u^{-1} .
- (ii) For any C -module M , there is an isomorphism of left A -modules ${}_\sigma A \otimes_C M \rightarrow A \otimes_B {}_\sigma M$.

(iii) For any C -module M there is an isomorphism of left A -modules $A \otimes_C M \rightarrow A \otimes_B \sigma M$.

Proof. (i) That the map $x \mapsto u^{-1}(x)$ is a linear bijection on A is clear, so we check that it is a homomorphism of bimodules. Indeed,

$$u^{-1}(axb) = u^{-1}a u u^{-1}xb = \sigma(a)u^{-1}(x)b = a \cdot_{\sigma} u^{-1}(x) \cdot b.$$

(ii) Define a map $\varphi : {}_{\sigma}A \otimes_C M \rightarrow A \otimes_B \sigma M$ on generators by $a \otimes m \mapsto \sigma^{-1}(a) \otimes m$ and extend by linearity. Then, φ is well-defined:

$$\begin{aligned} \varphi(ac \otimes_C m) &= \sigma^{-1}(ac) \otimes_B m = \sigma^{-1}(a)\sigma^{-1}(c) \otimes_B m = \sigma^{-1}(a) \otimes_B \sigma^{-1}(c) \cdot_{\sigma} m \\ &= \sigma^{-1}(a) \otimes_B c \cdot m = \varphi(a \otimes_C c \cdot m), \end{aligned}$$

for any $c \in C$. Moreover, φ is a homomorphism of left A -modules:

$$\varphi(x \cdot_{\sigma} (a \otimes_C m)) = \varphi(\sigma(x)a \otimes_C m) = \sigma^{-1}(\sigma(x)a) \otimes_B m = x\sigma^{-1}(a) \otimes_B m = x \cdot (\sigma^{-1}(a) \cdot_B m) = x \cdot \varphi(a \otimes_C m)$$

for any $x \in A$. Define a map $\psi : A \otimes_B \sigma M \rightarrow {}_{\sigma}A \otimes_C M$ on generators by $a \otimes m \mapsto \sigma(a) \otimes m$ and extend by linearity. Then,

$$\psi\varphi(a \otimes_C m) = \psi(\sigma^{-1}(a) \otimes_B m) = a \otimes_C m \quad \text{and} \quad \varphi\psi(a \otimes_B m) = \varphi(\sigma(a) \otimes_C m) = a \otimes_B m,$$

so φ and ψ are mutually inverse bijections. We conclude that φ is an isomorphism.

(iii) Combine (i) and (ii). □

31. Theorem. Let A be a basic quasi-hereditary algebra and let $B \subset A$ be a regular (respectively, homological) exact Borel subalgebra. Let σ be an inner automorphism of A . Then, $\sigma(B)$ is again a regular (respectively, homological) exact Borel subalgebra of A .

Proof. Due to Theorem 27, we know that C is an exact Borel subalgebra. We interpret elements of the spaces $\text{Ext}_B^k(L_B(i), L_B(j))$ and $\text{Ext}_C^k(L_C(i), L_C(j))$ as exact sequences of length $k+2$, modulo equivalence. Since $A \otimes_B _$ and $A \otimes_C _$ are exact functors, applying them to exact sequences in $\text{Ext}_B^k(L_B(i), L_B(j))$ and $\text{Ext}_C^k(L_C(i), L_C(j))$ produces exact sequences in the spaces

$$\text{Ext}_A^k(A \otimes_B L_B(i), A \otimes_B L_B(j)) \quad \text{and} \quad \text{Ext}_A^k(A \otimes_C L_C(i), A \otimes_C L_C(j)).$$

Functoriality ensures that these maps are well-defined. Denote them by F_B and F_C , respectively. We denote by $\sigma _$ the map $\text{Ext}_C^k(L_C(i), L_C(j)) \rightarrow \text{Ext}_B^k(L_B(i), L_B(j))$ defined by taking an exact sequence

$$L_C(j) \rightarrow M_k \rightarrow \cdots \rightarrow M_1 \rightarrow L_C(i) \in \text{Ext}_C^k(L_C(i), L_C(j))$$

to the sequence

$$\begin{aligned} \sigma L_C(j) \rightarrow \sigma M_k \rightarrow \cdots \rightarrow \sigma M_1 \rightarrow \sigma L_C(i) \\ = \\ L_B(j) \rightarrow \sigma M_k \rightarrow \cdots \rightarrow \sigma M_1 \rightarrow L_B(i). \end{aligned}$$

Consider the following diagram.

$$\begin{array}{ccc} \text{Ext}_B^k(L_B(i), L_B(j)) & \xleftarrow{\sigma _} & \text{Ext}_C^k(L_C(i), L_C(j)) \\ \downarrow F_B & & \downarrow F_C \\ \text{Ext}_A^k(A \otimes_B L_B(i), A \otimes_B L_B(j)) & \xleftarrow{\quad \quad \quad} & \text{Ext}_A^k(A \otimes_C L_C(i), A \otimes_C L_C(j)) \\ & \searrow \sim & \swarrow \sim \\ & \text{Ext}_A^k(\Delta(i), \Delta(j)) & \end{array}$$

Let $L_C(j) \rightarrow M_k \rightarrow \cdots \rightarrow M_1 \rightarrow L_C(i)$ be an exact sequence, interpreted as an element of the space $\text{Ext}_C^k(L_C(i), L_C(j))$. Applying first σ_- and then F_B , we obtain

$$A \otimes_B L_B(j) \rightarrow A \otimes_B \sigma M_k \rightarrow \cdots \rightarrow A \otimes_B \sigma M_1 \rightarrow A \otimes_B L_B(i).$$

If we instead apply F_C to our original sequence $L_C(j) \rightarrow M_k \rightarrow \cdots \rightarrow M_1 \rightarrow L_C(i)$, we get

$$A \otimes_C L_C(j) \rightarrow A \otimes_C M_k \rightarrow \cdots \rightarrow A \otimes_C M_1 \rightarrow A \otimes_C L_C(i).$$

Applying Lemma 30, we know there is an isomorphism of left A -modules, given by the composite

$$A \otimes_C M \rightarrow {}_\sigma A \otimes_C M \rightarrow A \otimes_B \sigma M.$$

This means that we may choose the dashed arrow in the diagram to be the map

$$A \otimes_C L_C(j) \rightarrow A \otimes_C M_k \rightarrow \cdots \rightarrow A \otimes_C M_1 \rightarrow A \otimes_C L_C(i)$$

\mapsto

$$A \otimes_B L_B(j) \rightarrow A \otimes_B \sigma M_k \rightarrow \cdots \rightarrow A \otimes_B \sigma M_1 \rightarrow A \otimes_B L_B(i),$$

which is an isomorphism and makes the top square commute, by construction. It follows that F_C is an isomorphism. For the second statement, the same argument as above ensures that the maps $\text{Ext}_C^k(L_C(i), L_C(j)) \rightarrow \text{Ext}_A^k(A \otimes L_C(i), A \otimes L_C(j))$ are isomorphisms for $k \geq 2$. For $k = 1$, the above diagram implies that we can solve for F_C to see that F_C is the composition of three epimorphisms, hence an epimorphism. \square

32. Corollary. Let $A = KQ/I$ be a basic quasi-hereditary algebra containing a regular exact Borel subalgebra. Then, all exact Borel subalgebras of A are regular. In particular, A admits a regular exact Borel subalgebra containing the idempotents e_1, \dots, e_n .

Proof. By assumption, there exists a regular exact Borel subalgebra B of A . Let C be an exact Borel subalgebra of A . By Theorem 28, there exists an inner automorphism σ of A such that $\sigma(B) = C$, and since inner automorphisms preserve regular exact Borel subalgebras, by Theorem 31, C is a regular exact Borel subalgebra. In the discussion following Theorem 28, we saw that A admits an exact Borel subalgebra containing the idempotents e_1, \dots, e_n . By the above argument, it is regular. \square

33. Proposition. Let $B \subset A_n$ be a regular exact Borel subalgebra containing the idempotents e_1, \dots, e_n and suppose that $\Delta(i) = M(i, s_i)$.

- (i) If $i \leq s_i < n$, then $\alpha_j \dots \alpha_i \notin B$, for any $i < j < s_i$, and $\alpha_{s_i} \dots \alpha_i \in B$.
- (ii) If $s_i = n$, that is, if $\Delta(i)$ is projective, then $\alpha_j \dots \alpha_i \notin B$, for any $i \leq j < n$.

Proof. (i) Assume towards a contradiction that $\alpha_j \dots \alpha_i \in B$ for some $i < j < s_i$. Let $p \in A_n$ be a path and consider the generator $p \otimes e_i$ of $A_n \otimes_B L(i)$. If $s(p) \neq i$, then

$$e_{j+1} \cdot (p \otimes e_i) = e_{j+1} \cdot (p \otimes e_{s(p)} e_i) = 0,$$

since $e_{s(p)} \in B$. Similarly, if $t(p) \neq j+1$, then

$$e_{j+1} \cdot (p \otimes e_i) = e_{j+1} e_{t(p)} p \otimes e_i = 0.$$

If $s(p) = i$ and $t(p) = j+1$, then $p = \alpha_j \dots \alpha_i$, which we assume is contained in B . We get

$$\alpha_j \dots \alpha_i \otimes e_i = e_j \otimes \alpha_j \dots \alpha_i e_i = 0,$$

so that $e_{j+1} \cdot (A_n \otimes_B L(i)) = 0$. However, $e_{j+1} \cdot \Delta(i) \neq 0$, which is our desired contradiction.

For the second statement, assume towards a contradiction that $\alpha_{s_i} \dots \alpha_i \notin B$. Then, $\alpha_{s_i} \dots \alpha_i \otimes e_i$ is a non-zero element of $A_n \otimes_B L(i)$. To see this, note that $\alpha_j \dots \alpha_i \notin B$ for any $i < j < s_i$, by the first part. It follows that also $e_{s_i+1} \cdot (\alpha_{s_i} \dots \alpha_i \otimes e_i) \neq 0$. Now, we see that $e_{s_i+1} \cdot \Delta(i) = 0$, while $e_{s_i+1} \cdot A_n \otimes_B L(i)$, leading to a similar contradiction as above.

- (ii) Similar to the first part of (i). We remark that this is a separate case only because the first statement of part (ii) does not make sense when $\Delta(i)$ is projective. \square


34. Proposition. Let $B \subset A_n$ be a regular exact Borel subalgebra containing the idempotents e_1, \dots, e_n . Let S_B be the set of paths which are contained in B by Proposition 33 together with the idempotents e_1, \dots, e_n . Then, S_B is a minimal generating set for B .

Proof. It is well known that any basic and connected finite-dimensional algebra over an algebraically closed field K is isomorphic to the quotient of the path algebra of some quiver by some admissible ideal. We recall the construction of this isomorphism. For details, we refer to [ASS06, Theorem 3.7].

Let $\Phi : KQ_B \rightarrow B$ be the isomorphism mentioned above. Then, Φ is defined on the trivial paths in Q_B by $\Phi(e_i) = e_i$, for $1 \leq i \leq n$. Note that this makes sense, as we may talk about the idempotents e_i as elements of KQ_B and of B .

Consider the set of arrows $i \rightarrow j$ in Q_B . As discussed prior to Proposition 25, the cardinality of this set equals $\dim \text{Ext}_{A_n}^1(\Delta(i), \Delta(j))$, which may be 0 or 1, according to Theorem 15. Therefore, this set is either empty or contains a single arrow, which we denote by x_i^j . The arrow x_i^j should be mapped to an element $y \in \text{rad} B$, so that the residue class $y + \text{rad}^2 B$ forms a basis of $e_j \text{rad} B / \text{rad}^2 B e_i$. This is achieved by defining $\Phi(x_i^j) = \alpha_{j-1} \dots \alpha_i$. Since we know that Φ is an isomorphism, we know that the idempotents e_i , together with paths of the form $\Phi(x_i^j)$, constitute a minimal generating set for B .

Recall that the arrow $x_i^j \in Q_B$ exists precisely when $\dim \text{Ext}_{A_n}^1(\Delta(i), \Delta(j)) = 1$. Consider the following picture. The dashed edge between b and c signifies that c is a vertex in the right subtree of b (not necessarily the vertex $r(b)$), such that there exists a non-zero homomorphism from $\Delta(c)$ to $\Delta(b)$.




We know, using Theorem 15, that possible non-zero extensions between standard modules appear as either $\text{Ext}_{A_n}^1(\Delta(b), \Delta(a))$ or as $\text{Ext}_{A_n}^1(\Delta(c), \Delta(a))$, where a, b and c are configured as above. Since b labels $\ell(a)$, the vertex a is the first vertex visited immediately after visiting the entire right subtree of a . Therefore, $a = s_b + 1$. Since c is in the right subtree of b , we have $s_b = s_c$, and therefore $a = s_c + 1$. Plugging this into the above, we see that the generators of B are of the form

$$\Phi(x_i^{s_i+1}) = \alpha_{s_i} \dots \alpha_i,$$


whence it follows that S_B is a minimal generating set. □

35. Corollary. Let B be the exact Borel subalgebra $B \subset A_n$ containing the idempotents e_1, \dots, e_n . Then, the exact Borel subalgebras are precisely those given as $C = \langle u^{-1} S_B u \rangle$, where $u \in A_n$ is some invertible element.

36. Example. We consider the binary search tree




We saw, in the example following Theorem 15, that the Ext-algebra of standard modules over A_n is given by the quiver



modulo the relations $I = \langle \varepsilon_2^4 \varepsilon_1^2 \rangle$. Here, there are three non-zero spaces of extensions, namely

$$\dim \text{Ext}_{A_n}^1(\Delta(1), \Delta(2)) = \dim \text{Ext}_{A_n}^1(\Delta(2), \Delta(4)) = \dim \text{Ext}_{A_n}^1(\Delta(3), \Delta(4)) = 1.$$

Letting $B \subset A_n$ be the regular exact Borel subalgebra containing e_1, \dots, e_n , we see that the quiver of B is



The arrows a , b and c correspond to the paths $\alpha_1, \alpha_3 \alpha_2$ and α_3 , respectively. A generating set S for B is then

$$S = \{e_1, e_2, e_3, e_4, e_5, e_6, \alpha_1, \alpha_3, \alpha_3 \alpha_2\}.$$

3.3. A_∞ -structure on $\text{Ext}_{A_n}^*(\Delta, \Delta)$. The notion of an A_∞ -algebra is meant to capture the idea of an algebra which is not strictly associative, but associative only up to a system of “higher homotopies”. There is also the natural “multi-object” version, called an A_∞ -category, which we define below. In this section we are particularly interested in $\text{Ext}_{A_n}^*(\Delta, \Delta)$, which always carries the structure of an A_∞ -algebra. However, we will view $\text{Ext}_{A_n}^*(\Delta, \Delta)$ as an A_∞ -category with n objects, given by the standard modules. This will be useful since it enables us to enforce compatibility of the higher homotopies with the natural idempotents of $\text{Ext}_{A_n}^*(\Delta, \Delta)$: the identity homomorphisms $1_{\Delta(i)}$.

37. Definition. [Kel01] Let K be a field. An A_∞ -category \mathcal{A} consists of the following:

- (i) a class of objects $\text{Ob}(\mathcal{A})$;
- (ii) for each pair of objects A and B , a \mathbb{Z} -graded vector space $\text{Hom}_{\mathcal{A}}(A, B)$;
- (iii) for all $n \geq 1$ and objects A_0, \dots, A_n , a homogeneous linear map

$$m_n : \text{Hom}_{\mathcal{A}}(A_{n-1}, A_n) \otimes \text{Hom}_{\mathcal{A}}(A_{n-2}, A_{n-1}) \otimes \dots \otimes \text{Hom}_{\mathcal{A}}(A_0, A_1) \rightarrow \text{Hom}_{\mathcal{A}}(A_0, A_n)$$

of degree $2 - n$ such that

$$\sum_{r+s+t=n} (-1)^{r+st} m_{r+1+t}(1^{\otimes r} \otimes m_s \otimes 1^{\otimes t}) = 0.$$

Viewing $\text{Ext}_{A_n}^*(\Delta, \Delta)$ as an A_∞ -category as outlined above amounts to the following: if we have extensions ε_i arranged as

$$\Delta(i_1) \xrightarrow{\varepsilon_1} \Delta(i_2) \xrightarrow{\varepsilon_2} \dots \longrightarrow \Delta(i_\ell) \xrightarrow{\varepsilon_\ell} \Delta(i_{\ell+1})$$

then $m_\ell(\varepsilon_\ell, \dots, \varepsilon_1)$ is an extension from $\Delta(i_1)$ to $\Delta(i_{\ell+1})$. Writing out the A_∞ -relations for the first few n , we get:

$n = 1$: $m_1 m_1 = 0$, meaning that A is a complex.

$n = 2$: $m_1 m_2 = m_2(m_1 \otimes 1 + 1 \otimes m_1)$, meaning that m_1 is a derivation with respect to m_2 .

$n = 3$: $m_2(1 \otimes m_2 - m_2 \otimes 1) = m_1 m_3 + m_3(m_1 \otimes 1 \otimes 1 + 1 \otimes m_1 \otimes 1 + 1 \otimes 1 \otimes m_1)$, meaning that m_2 is associative up to a homotopy given by m_3 .

When evaluating at actual elements, even more signs appear according to the Koszul sign rule:

$$(f \otimes g)(x \otimes y) = (-1)^{|g||x|} f(x) \otimes g(y).$$


Here, f and g are homogeneous maps and x and y are homogeneous elements. The vertical bars denote degree. A graded category A such that any A_∞ -structure on it satisfies $m_n = 0$ for $n \geq 3$ is called *intrinsically formal*.

38. Proposition. $\text{Ext}_{A_n}^*(\Delta, \Delta)$ is intrinsically formal.


Proof. Let $\varphi_1, \dots, \varphi_\ell$ be homogeneous extensions in $\text{Ext}_{A_n}^*(\Delta, \Delta)$. Put $i = \sum_{j=1}^\ell \deg \varphi_j$. Since m_ℓ is of degree $2 - \ell$, the extension $m_\ell(\varphi_\ell, \dots, \varphi_1)$ is of degree $2 - \ell + i$. Because A_n is hereditary, the extension $m_\ell(\varphi_\ell, \dots, \varphi_1)$ is of degree 0 or 1 if it is non-zero. This implies that there are the following cases:

$$2 - \ell + i = \begin{cases} 0, & \text{if } i = \ell - 2 \\ 1, & \text{if } i = \ell - 1. \end{cases}$$

Assume that $i = \ell - 1$ and that $m_\ell(\varphi_\ell, \dots, \varphi_1)$ produces an extension in $\text{Ext}_{A_n}^1(\Delta(c), \Delta(a))$. We know that if such an extension is non-zero, the vertices a and c must be configured in the following way: here the dashed edge represents a path which is the concatenation of edges between a vertex v and $r(v)$.



Since $i = \ell - 1$, there is precisely one argument φ_j of $m_\ell(\varphi_\ell, \dots, \varphi_1)$ which is a homomorphism rather than a proper extension. Consider the following picture, where the dashed edge represents the homomorphism $\varphi_j : \Delta(x) \rightarrow \Delta(y)$ and the wavy arrows from c to x and from y to a represent the compositions $\varphi_{j-1}, \dots, \varphi_1$ and $\varphi_\ell, \dots, \varphi_{j+1}$, respectively.



Note that either of the wavy edges may represent a path of length > 1 , rather than actual edges. Moreover, depending on j , the edge between c and x or the edge between y and a may be “empty”. We see that it is impossible to have the allowed configuration from the previous picture.

If $i = \ell - 2$, the outcome of $m_\ell(\varphi_\ell, \dots, \varphi_1)$ is an element of $\text{Hom}_{A_n}(\Delta(c), \Delta(a))$. Since $\ell \geq 3$, there is at least one argument φ_j of $m_\ell(\varphi_\ell, \dots, \varphi_1)$ which is a proper extension. According to the allowed configuration above, this means that c is in the left subtree of a . Using Theorem 15, we get that $\text{Hom}_{A_n}(\Delta(c), \Delta(a)) = 0$. \square

4. EXT-ALGEBRAS OF STANDARD MODULES OVER DECONCATENATIONS

In the previous section, we studied the path algebra of \mathbb{A}_n with linear orientation. To extend this study to arbitrary orientations, we apply the method of “deconcatenation” of a quiver Q at a sink or a source. In [FKR22], the authors use this method to study the different quasi-hereditary structures of the path algebra KQ in terms of the quasi-hereditary structures on path algebras of certain subquivers of Q . In the present section, we extend this study to the Ext-algebra of standard modules.

39. Definition. [FKR22, Definition 3.1] Let Q be a finite connected quiver and let v be a vertex of Q which is a sink or a source. A *deconcatenation* of Q at the vertex v is a union $Q^1 \sqcup \cdots \sqcup Q^\ell$ of full subquivers Q^i of Q such that the following hold.

- (i) Each Q^i is a connected full subquiver of Q with $v \in Q_0^i$.
- (ii) For all $i, j \in \{1, \dots, \ell\}$, such that $i \neq j$, there holds $Q_0 = (Q_0^1 \setminus \{v\}) \sqcup \cdots \sqcup (Q_0^\ell \setminus \{v\}) \sqcup \{v\}$ and $Q_0^i \cap Q_0^j = \{v\}$.
- (iii) For $x \in Q_0^i \setminus \{v\}$ and $y \in Q_0^j \setminus \{v\}$, where $i, j \in \{1, \dots, \ell\}$ are such that $i \neq j$, there are no arrows between x and y in Q .

40. Example. Consider the quiver $Q = 1 \longleftarrow 2 \longleftarrow 3 \longrightarrow 4 \longrightarrow 5$. We see that Q has a deconcatenation at the source 3:

$$(1 \longleftarrow 2 \longleftarrow 3) \sqcup (3 \longrightarrow 4 \longrightarrow 5).$$

We follow the notation of [FKR22] and put

$$A^\ell = A / \langle e_u \mid u \in Q_0 \setminus Q_0^\ell \rangle.$$

Then, A surjects onto A^ℓ , and consequently, there is a fully faithful and exact functor $F^\ell : A^\ell\text{-mod} \rightarrow A\text{-mod}$. Regarding $A^\ell\text{-mod}$ as a full subcategory of $A\text{-mod}$ via the functor F^ℓ , we remark that an A -module M is an A^ℓ -module if and only if $e_u M = 0$ for any $u \in Q_0 \setminus Q_0^\ell$.

41. Lemma. Let M be an A^ℓ -module, let N be an A -module and consider the quotient $\overline{N} = N / \sum e_u \cdot N$, where the sum ranges over all $u \in Q_0 \setminus \{v\}$. Then, \overline{N} is an A^ℓ -module and there are isomorphisms of vector spaces

$$\text{Hom}_A(M, \overline{N}) \cong \text{Hom}_{A^\ell}(M, \overline{N}) \cong \text{Hom}_A(M, N).$$

Proof. The first isomorphism is immediate from the fact that F^ℓ is fully faithful. To see the second isomorphism, put $e^\ell = \sum e_u$ where the sum is as in the statement of the Lemma. Then, there is an isomorphism of A -modules

$$\overline{N} \cong \text{Hom}_A(A / \langle e^\ell \rangle, N).$$

Next, note that by tensor-Hom adjunction, there is an adjoint pair of functors $(F^\ell, \text{Hom}_A(A / \langle e^\ell \rangle, _))$. Then,

$$\text{Hom}_A(M, N) \cong \text{Hom}_A(F^\ell(M), N) \cong \text{Hom}_{A^\ell}(M, \text{Hom}_A(A / \langle e^\ell \rangle, N)) \cong \text{Hom}_{A^\ell}(M, \overline{N}).$$

□

42. Lemma. [FKR22, Lemma 3.3] Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or source v and let $\ell = 1, 2$.

- (i) For all $i \in Q_0^\ell$, we have $L(i) \cong L^\ell(i)$.
- (ii) For all $i \in Q_0^\ell \setminus \{v\}$, we have $P(i) \cong P^\ell(i)$ and $I(i) \cong I^\ell(i)$.
- (iii) If v is a sink, we have $L(v) \cong P(v) \cong P^\ell(v)$, for $\ell = 1, 2$.
- (iv) If v is a source, we have $L(v) \cong I(v) \cong I^\ell(v)$, for $\ell = 1, 2$.
- (v) For any non-zero A -module M , if both $\text{top } M$ and $\text{soc } M$ are simple, then, we have either $M \in A^1\text{-mod}$ or $M \in A^2\text{-mod}$.
- (vi) Let M be an A^ℓ -module and let $i \in Q_0$ be a vertex. If $[M : L(i)] \neq 0$, then $i \in Q_0^\ell$.

Given a partial order \trianglelefteq on Q_0 and a deconcatenation $Q_1 \sqcup Q_2$, of Q at a sink or source v , we may form the restriction $\trianglelefteq|_{Q_0^\ell}$ to obtain a partial order $\trianglelefteq^\ell = \trianglelefteq|_{Q_0^\ell}$ on Q_0^ℓ .

43. Lemma. [FKR22, Lemma 3.4] Let $Q_1 \sqcup Q_2$ be a deconcatenation of Q at a sink or source v and let \trianglelefteq be a partial order on Q_0 . Denote by Δ and ∇ the sets of standard and costandard A -modules associated to \trianglelefteq , respectively. Denote by Δ^ℓ and ∇^ℓ the sets of standard and costandard A^ℓ -modules, respectively.

- (i) For any $i \in Q_0^\ell \setminus \{v\}$, there are isomorphisms of A -modules $\Delta(i) \cong \Delta^\ell(i)$ and $\nabla(i) \cong \nabla^\ell(i)$, for $\ell = 1, 2$.
- (ii) If v is a sink, there are isomorphisms of A -modules $L(v) \cong \Delta(v) \cong \Delta^\ell(v)$.
- (iii) If v is a source, there are isomorphisms of A -modules $L(v) \cong \nabla(v) \cong \nabla^\ell(v)$.
- (iv) If A is quasi-hereditary with respect to \trianglelefteq , then A^ℓ is quasi-hereditary with respect to \trianglelefteq^ℓ , for $\ell = 1, 2$.

Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation of Q at a source or sink v and let \trianglelefteq^ℓ be a partial order on Q_0^ℓ for $\ell = 1, 2$. Put $\bar{1} = 2$ and $\bar{2} = 1$. Define a partial order $\trianglelefteq = \trianglelefteq (\trianglelefteq^1, \trianglelefteq^2)$ on Q_0 as follows. For $i, j \in Q_0$, we say that $i \triangleleft j$ if one of the following hold.

- (i) We have $i, j \in Q_0^\ell$ and $i \triangleleft^\ell j$, for some ℓ .
- (ii) We have $i \in Q_0^\ell, j \in Q_0^{\bar{\ell}}, i \triangleleft^\ell v$ and $v \triangleleft^{\bar{\ell}} j$.

44. Lemma. [FKR22, Lemma 3.5] Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or source v . Let \trianglelefteq^ℓ be a partial order on Q_0^ℓ and denote by Δ^ℓ and ∇^ℓ the sets of standard and costandard A^ℓ -modules, associated to \trianglelefteq^ℓ , for $\ell = 1, 2$, respectively. Denote by Δ and ∇ the sets of standard and costandard A -modules, respectively, associated to $\trianglelefteq = \trianglelefteq (\trianglelefteq^1, \trianglelefteq^2)$. Then, we have the following.

- (i) For any $i \in Q_0^\ell \setminus \{v\}$, there are isomorphisms of A -modules $\Delta(i) \cong \Delta^\ell(i)$ and $\nabla(i) \cong \nabla^\ell(i)$.
- (ii) If v is a sink, there are isomorphisms of A -modules $L(v) \cong \Delta(v) \cong \Delta^\ell(v)$.
- (iii) If v is a source, there are isomorphisms of A -modules $L(v) \cong \nabla(v) \cong \nabla^\ell(v)$.
- (iv) If \trianglelefteq^ℓ defines a quasi-hereditary structure on A^ℓ for $\ell = 1$ and $\ell = 2$, then $\trianglelefteq = \trianglelefteq (\trianglelefteq^1, \trianglelefteq^2)$ defines a quasi-hereditary structure on A .

We have the following observation about homomorphism spaces between indecomposable projective modules.

45. Lemma. Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at the sink or source v . Then, for $i \in Q_0^1 \setminus \{v\}$ and $j \in Q_0^2 \setminus \{v\}$, there holds $\text{Hom}_A(P(i), P(j)) = \text{Hom}_A(P(j), P(i)) = 0$.

46. Proposition. Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at the sink or source v . Then, for $i \in Q_0^1 \setminus \{v\}$ and $j \in Q_0^2 \setminus \{v\}$, there holds $\text{Ext}_A^k(\Delta(i), \Delta(j)) = \text{Ext}_A^k(\Delta(j), \Delta(i)) = 0$, for all $k \geq 0$.

Proof. Let $P^\bullet \rightarrow \Delta(i)$ be a minimal projective resolution. We claim that each term P^k of the resolution P^\bullet decomposes into a direct sum of indecomposable projective modules as

$$P^k = \bigoplus_{\ell=1}^n P(\ell)^{m_{\ell,k}},$$

where, if $m_{\ell,k} > 0$, then $\ell \in Q_0^1 \setminus \{v\}$. We proceed by induction. The basis is clear, as $P^0 = P(i)$. Consider the following picture.

$$\begin{array}{ccccccc} \dots & \xrightarrow{d_{k+2}} & P^{k+1} & \xrightarrow{d_{k+1}} & P^k & \xrightarrow{d_k} & \dots \\ & & & \searrow & \swarrow & & \\ & & & & \ker d_k & & \end{array}$$

Assume that the module P^k is of the form indicated above. By construction, the module P^{k+1} is a projective cover of the module $\ker d_k$. Therefore, no direct summand of P^{k+1} is contained in the kernel of the epimorphism $P^{k+1} \rightarrow \ker d_k$. Consequently, no direct summand of P^{k+1} is contained in the kernel of the map $d_{k+1} : P^{k+1} \rightarrow P^k$. Fixing some decomposition of the modules P^{k+1} and P^k into direct sums of indecomposable projective modules, the map d_{k+1} is given by a matrix, whose entries are homomorphisms

$f \in \text{Hom}_A(P(a), P(b))$, where a and b are such that $P(a)$ is a direct summand of P^{k+1} and $P(b)$ is a direct summand of P^k . Since $\text{Hom}_A(P(a), P(b)) \cong e_a A e_b$, and $b \in Q_0^1 \setminus \{v\}$ by the inductive assumption, if $f \neq 0$, we must have $a \in Q_0^1 \setminus \{v\}$. This proves the claim. Next, apply the functor $\text{Hom}_A(_, \Delta(j))$ to the projective resolution P^\bullet . The resulting complex looks as follows:

$$0 \rightarrow \text{Hom}_A(P(i), \Delta(j)) \rightarrow \text{Hom}_A(P^1, \Delta(j)) \rightarrow \text{Hom}_A(P^2, \Delta(j)) \rightarrow \dots$$

By our previous claim, we have

$$P^k = \bigoplus_{\ell=1}^n P(\ell)^{m_{\ell,k}},$$

where we may now assume that $\ell \in Q_0^1 \setminus \{v\}$. By additivity of the Hom-functor, we have

$$\text{Hom}_A(P^k, \Delta(j)) \cong \bigoplus_{\ell=1}^n \text{Hom}_A(P(\ell), \Delta(j))^{m_{\ell,k}}.$$

Consider the following picture.

$$\begin{array}{ccc} P(\ell) & & \\ \exists h \downarrow & \searrow f & \\ P(j) & \xrightarrow{p_j} & \Delta(j) \end{array}$$

By projectivity, any homomorphism f factors as $f = p_j \circ h$. But, according to Lemma 45, we have

$$\text{Hom}_A(P(\ell), P(j)) = 0,$$

since $\ell \in Q_0^1 \setminus \{v\}$ and $j \in Q_0^2 \setminus \{v\}$. This proves that $\text{Ext}_A^k(\Delta(i), \Delta(j)) = 0$. It is clear that we may argue in the same way to get the conclusion $\text{Ext}_A^k(\Delta(j), \Delta(i)) = 0$. \square

47. Lemma. Assume that v is a source. Let $i \in Q_0^\ell \setminus \{v\}$ and let $P^\bullet \rightarrow \Delta^\ell(i)$ be a minimal projective resolution of $\Delta^\ell(i)$ as an A -module. Then no term P^k of the projective resolution P^\bullet contains the projective module $P^\ell(v)$ as a direct summand.

Proof. We proceed by induction. The basis is clear, as $P^0 = P^\ell(i)$. Consider the following picture.

$$\begin{array}{ccccccc} \dots & \xrightarrow{d_{k+2}} & P^{k+1} & \xrightarrow{d_{k+1}} & P^k & \xrightarrow{d_k} & \dots \\ & & & \searrow & \nearrow & & \\ & & & \text{ker } d_k & & & \end{array}$$

Assume that the statement holds for the module P^k and assume towards a contradiction that the module P^{k+1} contains a direct summand $P^\ell(v)$. By construction, P^{k+1} is a projective cover of $\text{ker } d_k$, thus we conclude that the module $\text{topker } d_k$ contains a direct summand $L^\ell(v)$. This fact, in turn, implies that there is a direct summand $P^\ell(t)$ of P^k , such that $[P^\ell(t) : L^\ell(v)] > 0$. Since a basis of $P^\ell(t)$ is given by paths in A^ℓ starting in t , this implies that there is a path in A^ℓ from t to v . This is our desired contradiction, as v was assumed to be a source. \square

48. Proposition. Let $Q^1 \sqcup Q^2$ be a deconcatenation at a sink or source v .

- (i) If $i, j \in Q_0^\ell \setminus \{v\}$, there holds $\text{Ext}_{A^\ell}^k(\Delta^\ell(i), \Delta^\ell(j)) \cong \text{Ext}_A^k(\Delta(i), \Delta(j))$, for all $k \geq 0$.
- (ii) If v is a sink, then $\text{Ext}_{A^\ell}^k(\Delta^\ell(v), \Delta^\ell(j)) \cong \text{Ext}_A^k(\Delta(v), \Delta(j))$, for all $k \geq 0$.
- (iii) If v is a sink, then $\text{Ext}_{A^\ell}^k(\Delta^\ell(i), \Delta^\ell(v)) \cong \text{Ext}_A^k(\Delta(i), \Delta(v))$, for all $k \geq 0$.

Proof. Let $P^\bullet \rightarrow \Delta^\ell(i)$ and $Q^\bullet \rightarrow \Delta^\ell(j)$ be minimal projective resolutions. By Lemma 42, there is an isomorphism of A -modules $P^\ell(x) \cong P(x)$, for all $x \in Q_0^\ell \setminus \{v\}$. Note that this isomorphism is the one induced by the functor F^ℓ , that is, $F^\ell(P^\ell(x)) \cong P(x)$. Similarly, we have $F^\ell(\Delta^\ell(i)) \cong \Delta(i)$ and $F^\ell(\Delta^\ell(j)) \cong \Delta(j)$.

- (i) The functor $F^\ell : A^\ell\text{-mod} \rightarrow A\text{-mod}$ is fully faithful and exact. By Lemma 47, the terms of the projective resolution $P^\bullet \rightarrow \Delta^\ell(i)$ do not contain any direct summands isomorphic to $P^\ell(v)$. We conclude that $F^\ell(P^\bullet) \rightarrow F^\ell(\Delta^\ell(i)) \cong \Delta(i)$ is a minimal projective resolution. Similarly, we have that $F^\ell(Q^\bullet) \rightarrow F^\ell(\Delta^\ell(j)) \cong \Delta(j)$ is a minimal projective resolution. The statement follows.
- (ii) If v is a sink, we have $P^\ell(v) \cong L(v) \cong P(v)$ as A -modules, by Lemma 42, and $\Delta^\ell(v) \cong L(v) \cong \Delta(v)$ as A -modules, by Lemma 43. Thus, the statement follows by applying the functor F^ℓ to the minimal projective resolutions of $\Delta^\ell(v)$ and $\Delta^\ell(j)$.
- (iii) Similar to (ii). □

We recall the notation of [FKR22], where $\bar{1} = 2$ and $\bar{2} = 1$.

49. Lemma. Let $Q_1 \sqcup Q_2$ be a deconcatenation of Q at the source v and let $i \in Q_0^\ell \setminus \{v\}$. Then, there are isomorphisms of vector spaces

$$\text{Hom}_A(P(i), P^\ell(v)) \cong \text{Hom}_A(P(i), P(v)) \quad \text{and} \quad \text{Hom}_A(P^\ell(v), P(i)) \cong \text{Hom}_A(P(v), P(i)), \text{ for } \ell = 1, 2.$$

Proof. The first statement is precisely Lemma 41 applied to the modules $P(i)$ and $P^\ell(v)$. Note that in the notation of Lemma 41, we have $P^\ell(v) = \overline{P(v)}$. For the second statement, let $p : P(v) \rightarrow P^\ell(v)$ denote the natural epimorphism. Then, there is a short exact sequence

$$0 \longrightarrow \ker p \longrightarrow P(v) \xrightarrow{p} P^\ell(v) \longrightarrow 0.$$

Note that $P^\ell(v)$ is an A^ℓ -module while $\ker p$ is an $A^{\bar{\ell}}$ -module. Applying $\text{Hom}_A(_, P(i))$ to this sequence, we obtain:

$$0 \longrightarrow \text{Hom}_A(P^\ell(v), P(i)) \longrightarrow \text{Hom}_A(P(v), P(i)) \longrightarrow \text{Hom}_A(\ker p, P(i))$$

The space $\text{Hom}_A(\ker p, P(i))$ is zero because of our previous observation, so we have

$$\text{Hom}_A(P^\ell(v), P(i)) \cong \text{Hom}_A(P(v), P(i))$$

since the Hom-functor is left exact. □

50. Lemma. Let v be a source, let $P^\bullet \rightarrow \Delta^1(v)$ be a minimal projective resolution with terms P^k , $k \geq 0$ and let $Q^\bullet \rightarrow \Delta^2(v)$ be a minimal projective resolution with terms Q^k , $k \geq 0$. Then, there is a minimal projective resolution $R^\bullet \rightarrow \Delta(v)$, with terms $R^k = P^k \oplus Q^k$, for $k \geq 1$, and $R^0 = P(v)$.

Proof. We proceed by induction. For the case $k = 1$, consider the projection $p : P(v) \rightarrow \Delta(v)$. The module $\ker p$ has a basis consisting of paths q in A , such that $s(q) = v$ and $t(q) = j$ with $j \not\prec v$. Let x_1, \dots, x_n be those paths that satisfy $t(x_i) \in Q_0^1$ and let y_1, \dots, y_m be those paths that satisfy $t(y_j) \in Q_0^2$. Put $X = \text{span}\{x_1, \dots, x_n\}$ and $Y = \text{span}\{y_1, \dots, y_m\}$. Clearly, $\ker p = X \oplus Y$ as vector spaces. Since the action of A on X and Y is given by left multiplication, we see that, in fact, $\ker p = X \oplus Y$ as A -modules.

It is clear that, as an A^1 -module, X is isomorphic to the kernel of the projection $p^1 : P^1(v) \rightarrow \Delta^1(v)$, so P^1 is a projective cover of X . Similarly, as an A^2 -module, Y is isomorphic to the kernel of the projection $p^2 : P^2(v) \rightarrow \Delta^2(v)$, so Q^1 is a projective cover of Y . When $k = 2$, we consider the picture:

$$\begin{array}{ccccc} R^2 & \longrightarrow & P^1 \oplus Q^1 & \xrightarrow{(d^1, e^1)} & P(v) \\ & & \searrow & & \uparrow \\ & & \ker d^1 \oplus \ker e^1 & & X \oplus Y \end{array}$$

It is clear that we may view the map from $P^1 \oplus Q^1$ to $P(v)$ as (d^1, e^1) . It follows that its kernel is the direct sum $\ker d^1 \oplus \ker e^1$, which has a projective cover $R^2 = P^2 \oplus Q^2$. This finishes the basis of the induction. Consider

the following picture:

$$\begin{array}{ccccc}
 R^{k+1} & \xrightarrow{f^{k+1}} & P^k \oplus Q^k & \xrightarrow{\begin{pmatrix} d^k & 0 \\ 0 & e^k \end{pmatrix}} & P^{k-1} \oplus Q^{k-1} \\
 \downarrow & \nearrow & & & \\
 \ker d^k \oplus \ker e^k & & & &
 \end{array}$$

That the right matrix above is diagonal follows from Lemma 45 and the induction hypothesis. Then, the kernel of the map $\begin{pmatrix} d^k & 0 \\ 0 & e^k \end{pmatrix}$ is isomorphic to $\ker d^k \oplus \ker e^k$, and a projective cover is therefore $P^{k+1} \oplus Q^{k+1}$. This shows that $R^{k+1} = P^{k+1} \oplus Q^{k+1}$ and $f^{k+1} = \begin{pmatrix} d^{k+1} & 0 \\ 0 & e^{k+1} \end{pmatrix}$. \square

51. Proposition. Let $Q_1 \sqcup Q_2$ be a deconcatenation of Q at the source v and let $i \in Q_0^\ell \setminus \{v\}$. Then, there are isomorphisms of vector spaces

$$\text{Ext}_{A^\ell}^k(\Delta^\ell(i), \Delta^\ell(v)) \cong \text{Ext}_A^k(\Delta(i), \Delta(v)) \quad \text{and} \quad \text{Ext}_{A^\ell}^k(\Delta^\ell(v), \Delta^\ell(i)) \cong \text{Ext}_A^k(\Delta(v), \Delta(i)),$$

for all $k \geq 0$.

Proof. Let $P^\bullet \rightarrow \Delta^\ell(i)$ and $Q^\bullet \rightarrow \Delta^\ell(v)$ be minimal projective resolutions. Let $\varepsilon^\ell : P^\bullet \rightarrow Q^\bullet[k]$ be a chain map.

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & P^{k+2} & \xrightarrow{d^{k+2}} & P^{k+1} & \xrightarrow{d^{k+1}} & P^k & \longrightarrow & \dots \\
 & & \downarrow \varepsilon_2^\ell & & \downarrow \varepsilon_1^\ell & & \downarrow \varepsilon_0^\ell & & \\
 \dots & \longrightarrow & Q^2 & \xrightarrow{\delta^2} & Q^1 & \xrightarrow{\delta^1} & P^\ell(v) & &
 \end{array}$$

According to Lemma 50, there is a minimal projective resolution $T^\bullet \rightarrow \Delta(v)$ with terms

$$T^k = \begin{cases} Q^k \oplus R^k, & \text{if } k \geq 1 \\ P(v), & \text{if } k = 0, \end{cases}$$

where the modules R^k constitute a minimal projective resolution of $\Delta^\ell(v)$. Define a map $\varepsilon : P^\bullet \rightarrow T^\bullet[k]$, as having components given by

$$\varepsilon_n = \begin{cases} \begin{pmatrix} \varepsilon_n^\ell \\ 0 \end{pmatrix}, & \text{if } n \geq 1 \\ \Phi(\varepsilon_0^\ell), & \text{if } n = 0. \end{cases}$$

Here, $\Phi : \text{Hom}_A(P^k, P^\ell(v)) \rightarrow \text{Hom}_A(P^k, P(v))$ is the isomorphism obtained in the proof of Lemma 49, so that ε_0 is uniquely defined by the equation $p\varepsilon_0 = \varepsilon_0^\ell$, where $p : P(v) \rightarrow P(v)^\ell$ is the natural projection.

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & P^{k+2} & \xrightarrow{d^{k+2}} & P^{k+1} & \xrightarrow{d^{k+1}} & P^k & \longrightarrow & \dots \\
 & & \downarrow \varepsilon_2 & & \downarrow \varepsilon_1 & & \downarrow \varepsilon_0 & & \\
 \dots & \longrightarrow & Q^2 \oplus R^2 & \xrightarrow{\begin{pmatrix} \delta^2 & 0 \\ 0 & \partial^2 \end{pmatrix}} & Q^1 \oplus R^1 & \longrightarrow & P(v) & &
 \end{array}$$

We claim that ε is a chain map. Considering the following diagram, where $n \geq 1$.

$$\begin{array}{ccc}
 P^{k+n+1} & \xrightarrow{d^{k+n+1}} & P^{k+n} \\
 \varepsilon_{n+1} \downarrow & & \downarrow \varepsilon_n \\
 Q^{n+1} \oplus R^{n+1} & \xrightarrow{\begin{pmatrix} \delta^{n+1} & 0 \\ 0 & \partial^{n+1} \end{pmatrix}} & Q^n \oplus R^n
 \end{array}$$

We have

$$\begin{pmatrix} \delta^{n+1} & 0 \\ 0 & \partial^{n+1} \end{pmatrix} \varepsilon_{n+1} = \begin{pmatrix} \delta^{n+1} & 0 \\ 0 & \partial^{n+1} \end{pmatrix} \begin{pmatrix} \varepsilon_{n+1}^\ell \\ 0 \end{pmatrix} = \begin{pmatrix} \delta^{n+1} \varepsilon_{n+1}^\ell \\ 0 \end{pmatrix} = \begin{pmatrix} \varepsilon_n^\ell d^{k+n+1} \\ 0 \end{pmatrix} = \begin{pmatrix} \varepsilon_n^\ell \\ 0 \end{pmatrix} d^{k+n+1} = \varepsilon_n d^{k+n+1}$$

since $\delta^{n+1} \varepsilon_{n+1}^\ell = \varepsilon_n^\ell d^{k+n+1}$ by virtue of ε^ℓ being a chain map. It remains to check the following square.

$$\begin{array}{ccc} P^{k+1} & \xrightarrow{d^{k+1}} & P^k \\ \varepsilon_1 \downarrow & & \downarrow \varepsilon_0 \\ Q^1 \oplus R^1 & \longrightarrow & P(v) \end{array}$$

In the proof of Lemma 50, we saw that the kernel of the projection $P(v) \rightarrow \Delta(v)$ is isomorphic to $X \oplus Y$, where X is isomorphic to the kernel of the projection $P^1(v) \rightarrow \Delta^1(v)$ and Y is isomorphic to the kernel of the projection $P^2(v) \rightarrow \Delta^2(v)$. Consider the following pictures:

$$\begin{array}{ccccc} Q^1 & \xrightarrow{\delta^1} & P^1(v) & \longrightarrow & \Delta^1(v) \\ & \searrow & \nearrow & & \\ & & X & & \\ R^1 & \xrightarrow{\partial^1} & P^2(v) & \longrightarrow & \Delta^2(v) \\ & \searrow & \nearrow & & \\ & & Y & & \end{array}$$

By construction, δ^1 and ∂^1 are radical maps. Since $\text{rad } P^1(v)$ and $\text{rad } P^2(v)$ are isomorphic to submodules of $P(v)$, we may view δ^1 and ∂^1 as maps $\delta^1 : Q^1 \rightarrow P(v)$ and $\partial^1 : R^1 \rightarrow P(v)$, respectively. Then the bottom map in the above square can be written as $(\delta^1, \partial^1) : P^1 \oplus R^1 \rightarrow P(v)$. Consider the following picture:

$$\begin{array}{ccccc} P^{k+1} & \xrightarrow{d^{k+1}} & P^k & \xrightarrow{\varepsilon_0^\ell} & P^\ell(v) \\ \varepsilon_1 \downarrow & & \downarrow \varepsilon_0 & \nearrow p & \\ Q^1 \oplus R^1 & \xrightarrow{(\delta^1, \partial^1)} & P(v) & & \end{array}$$

By construction, the triangle on the right commutes. Since ε^ℓ is a chain map, we have

$$(\delta^1, \partial^1) \varepsilon_1 = (\delta^1, \partial^1) \begin{pmatrix} \varepsilon_1^\ell \\ 0 \end{pmatrix} = \delta^1 \varepsilon_1^\ell = \varepsilon_0^\ell d^{k+1}.$$

Moreover, the image of $\delta^1 \varepsilon_1^\ell = \varepsilon_0^\ell d^{k+1}$ is contained in the submodule $\text{rad}(A^\ell) \cdot P(v)$, on which p acts as the identity. This implies that the perimeter of the diagram commutes, and since p is an isomorphism on $\text{rad}(A^\ell) \cdot P(v)$, the square commutes. Let $e^\ell : P^\bullet \rightarrow Q^\bullet[k]$ be another chain map. We claim that if ε^ℓ and e^ℓ are homotopic, so are ε and e , where $e : P^\bullet \rightarrow T^\bullet[k]$ is the chain map with components

$$e_n = \begin{cases} \begin{pmatrix} e_n^\ell \\ 0 \end{pmatrix}, & \text{if } n \geq 1 \\ \Phi(e_0^\ell), & \text{if } n = 0. \end{cases}$$

$$\begin{array}{ccccc} P^{k+n+1} & \xrightarrow{d^{k+n+1}} & P^{k+n} & \xrightarrow{d^{k+n}} & P^{k+n-1} \\ \downarrow & \nearrow h_{n+1} & \downarrow e_n^\ell & \nearrow h_n & \downarrow \\ Q^{n+1} & \xrightarrow{\delta^{n+1}} & Q^n & \xrightarrow{\delta^n} & Q^{n-1} \end{array}$$

Assume that $h_n d^{k+n} + \delta^{n+1} h_{n+1} = \varepsilon_n^\ell - e_n^\ell$, for $n \geq 1$. Consider the picture:

$$\begin{array}{ccccc}
P^{k+n+1} & \xrightarrow{d^{k+n+1}} & P^{k+n} & \xrightarrow{d^{k+n}} & P^{k+n-1} \\
\downarrow & \nearrow \begin{pmatrix} h_{n+1} \\ 0 \end{pmatrix} & \downarrow \begin{pmatrix} e_n^\ell \\ 0 \end{pmatrix} & \begin{pmatrix} \varepsilon_n^\ell \\ 0 \end{pmatrix} & \nearrow \begin{pmatrix} h_n \\ 0 \end{pmatrix} & \downarrow \\
Q^{n+1} \oplus R^{n+1} & \xrightarrow{\begin{pmatrix} \delta^{n+1} & 0 \\ 0 & \delta^{n+1} \end{pmatrix}} & Q^n \oplus R^n & \xrightarrow{\begin{pmatrix} \delta^n & 0 \\ 0 & \delta^n \end{pmatrix}} & Q^{n-1} \oplus R^{n-1}
\end{array}$$

Then, we have

$$\begin{pmatrix} \delta^{n+1} & 0 \\ 0 & \delta^{n+1} \end{pmatrix} \begin{pmatrix} h_{n+1} \\ 0 \end{pmatrix} + \begin{pmatrix} h_n \\ 0 \end{pmatrix} d^{k+n} = \begin{pmatrix} \delta^{n+1} h_{n+1} \\ 0 \end{pmatrix} + \begin{pmatrix} h_n d^{k+n} \\ 0 \end{pmatrix} = \begin{pmatrix} \varepsilon_n^\ell - e_n^\ell \\ 0 \end{pmatrix}.$$

It remains to check the case $n = 0$. Consider the following picture.

$$\begin{array}{ccccc}
P^{k+1} & \xrightarrow{d^{k+1}} & P^k & \xrightarrow{d^k} & P^{k-1} \\
\downarrow & \nearrow h_1 & \downarrow \begin{matrix} e_0^\ell \\ \varepsilon_0^\ell \end{matrix} & \nearrow h_0 & \downarrow \\
Q^1 & \xrightarrow{\delta^1} & P^\ell(v) & \xrightarrow{\quad} & 0
\end{array}$$

Assume that $h_0 \circ d^k + \delta^1 h_1 = \varepsilon_0^\ell - e_0^\ell$ as maps $P^k \rightarrow P^\ell(v)$. To finish our homotopy, we take the map $\Phi(h_0)$, where

$$\Phi : \text{Hom}_A(P^k, P^\ell(v)) \rightarrow \text{Hom}_A(P^k, P(v))$$

is the isomorphism constructed in the proof of Lemma 49.

$$\begin{array}{ccccc}
P^{k+1} & \xrightarrow{d^{k+1}} & P^k & \xrightarrow{d^k} & P^{k-1} \\
\downarrow & \nearrow \begin{pmatrix} h_1 \\ 0 \end{pmatrix} & \downarrow \begin{matrix} e_0 \\ \varepsilon_0 \end{matrix} & \nearrow \Phi(h_0) & \downarrow h_0 \\
Q^1 \oplus R^1 & \xrightarrow{(\delta^1, \delta^1)} & P(v) & \xrightarrow{p} & P^\ell(v)
\end{array}$$

Note that the maps ε_0, e_0 and $\Phi(h_0)$ are defined by the equations

$$p\varepsilon_0 = \varepsilon_0^\ell, \quad pe_0 = e_0^\ell, \quad \text{and} \quad p\Phi(h_0) = h_0.$$

Using this, we obtain $\delta^1 h_1 + p\Phi(h_0)d^k = p(\varepsilon_0 - e_0)$. Since the image of $\delta^1 h_1$ is contained in the submodule $\text{rad } P^\ell(v)$, which in a natural way is a submodule of $P(v)$ on which $p : P(v) \rightarrow P^\ell(v)$ acts as the identity, we have $\delta^1 h_1 = p\delta^1 h_1$ as maps $P^k \rightarrow P^\ell(v)$. This means that we have the equality

$$p(\delta^1 h_1 + \Phi(h_0)d^k) = p(\varepsilon_0 - e_0)$$

as maps $P^k \rightarrow P^\ell(v)$. Now, since P^k is an A^ℓ -module, we may argue as in the proof of Lemma 49, to conclude that the images of $\delta^1 h_1, \Phi(h_0), \varepsilon_0$ and e_0 are all contained in the submodule $\text{rad}(A^\ell) \cdot P(v) \subset P(v)$, on which p acts as the identity. This implies that

$$\varepsilon_0 - e_0 = \delta^1 h_1 + \Phi(h_0)d^k = (\delta^1, \delta^1) \begin{pmatrix} h_1 \\ 0 \end{pmatrix} + \Phi(h_0)d^k,$$

confirming the claim that ε and e are homotopic. Thus, we have shown that the map

$$\Upsilon : \text{Ext}_{A^\ell}^k(\Delta^\ell(i), \Delta^\ell(v)) \rightarrow \text{Ext}_A^k(\Delta(i), \Delta(v)),$$

defined by $\varepsilon^\ell \mapsto \varepsilon$, is well-defined.

Next, note that, since $i \in Q_0^\ell \setminus \{v\}$, the terms of P^\bullet do not contain indecomposable direct summands isomorphic to $P^\ell(v)$, according to Lemma 47. Then, since the indecomposable direct summands of R^s are of the

form $P(j)$, for $j \in Q_0^\ell$, we have $\text{Hom}_A(P^{k+s}, R^s) = 0$, for any $s \geq 1$, according to Lemma 45. Thus, we have an isomorphism of vector spaces

$$\text{Hom}_{A^\ell}(P^{k+s}, Q^s) \cong \text{Hom}_A(P^{k+s}, T^s),$$

for all $s \geq 1$, given by $f \mapsto \begin{pmatrix} f \\ 0 \end{pmatrix}$ (note that $T^s = Q^s \oplus R^s$ for $s \geq 1$). When $s = 0$, we have the isomorphism $\text{Hom}_{A^\ell}(P^k, Q^0) \cong \text{Hom}_A(P^k, T^0)$, according to Lemma 49, since $Q^0 = P^\ell(v)$ and $T^0 = P(v)$. This means that we have an isomorphism of vector spaces

$$\xi^s : \text{Hom}_{A^\ell}(P^{k+s}, Q^s) \cong \text{Hom}_A(P^{k+s}, T^s),$$

for all $s \geq 0$. We note that, by construction, $(Y(\varepsilon^\ell))_n = \xi^n(\varepsilon_n^\ell)$ for all $n \geq 0$, which shows that Y is a linear isomorphism. This proves the first statement. The second statement is proved similarly. \square

Let $\text{Ext}_{A^\ell}^*(\Delta^\ell, \Delta^\ell)$ denote the Ext-algebra of standard modules over A^ℓ , for $\ell = 1, 2$. Similarly, let $\text{Ext}_A^*(\Delta, \Delta)$ denote the Ext-algebra of standard modules over A . Fix the basis $1_{\Delta(v)}$ of the space $\text{End}_A(\Delta(v))$.

52. Theorem. Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or source v . Then, there is an isomorphism of graded algebras $\text{Ext}_A^*(\Delta, \Delta) \cong \text{Ext}_{A^1}^*(\Delta^1, \Delta^1) \diamond \text{Ext}_{A^2}^*(\Delta^2, \Delta^2)$.

Proof. We start by noting that, according to Propositions 48 and 51, there holds

$$\text{Ext}_{A^\ell}^k(\Delta^\ell(i), \Delta^\ell(j)) \cong \text{Ext}_A^k(\Delta(i), \Delta(j)),$$

for all $i, j \in Q_0^\ell$ and all $k \geq 0$. Put $C = \text{Ext}_{A^1}^*(\Delta^1, \Delta^1) \diamond \text{Ext}_{A^2}^*(\Delta^2, \Delta^2)$. By definition of the multiplication on C , for any $x \in \text{Ext}_{A^1}^*(\Delta^1, \Delta^1)$ and $y \in \text{Ext}_{A^2}^*(\Delta^2, \Delta^2)$ such that $\deg x > 0$ and $\deg y > 0$, there holds $x \cdot_C y = y \cdot_C x = 0$. Similarly, according to Proposition 46, we have

$$\text{Ext}_{A^1}^{>0}(\Delta^1, \Delta^1) \cdot \text{Ext}_{A^2}^{>0}(\Delta^2, \Delta^2) = \text{Ext}_{A^2}^{>0}(\Delta^2, \Delta^2) \cdot \text{Ext}_{A^1}^{>0}(\Delta^1, \Delta^1) = 0.$$

Together, these facts imply that there is a homomorphism of graded algebras $\Phi : C \rightarrow \text{Ext}_A^*(\Delta, \Delta)$. Note that, because of Proposition 46, the subspaces $\Phi(\text{Ext}_{A^\ell}^*(\Delta^\ell, \Delta^\ell))$ generate $\text{Ext}_A^*(\Delta, \Delta)$, showing that Φ is surjective. Lastly, note that $\dim_K C = \dim_K \text{Ext}_A^*(\Delta, \Delta)$, so that Φ is a surjective linear map between vector spaces of the same dimension, hence a linear isomorphism. \square

53. Example. We consider the deconcatenation $(1 \leftarrow 2 \leftarrow 3) \sqcup (3 \rightarrow 4 \rightarrow 5)$ of the quiver

$$Q = 1 \leftarrow 2 \leftarrow 3 \rightarrow 4 \rightarrow 5.$$

We draw the Loewy diagrams of the projective and standard modules over A^1 .

$$P^1(1) : 1, \quad P^1(2) : \begin{array}{c} 2 \\ \downarrow \\ 1 \end{array}, \quad P^1(3) : \begin{array}{c} 3 \\ \downarrow \\ 2 \\ \downarrow \\ 1 \end{array}, \quad \Delta^1(1) \cong P^1(1), \quad \Delta^1(2) \cong P^1(2), \quad \Delta^1(3) \cong P^1(3).$$

As the standard modules over A^1 are projective, there are no non-split extensions between them. We see that there are homomorphisms $f : \Delta^1(1) \rightarrow \Delta^1(2)$, $g : \Delta^1(2) \rightarrow \Delta^1(3)$ and $h : \Delta^1(1) \rightarrow \Delta^1(3)$, such that $h = g \circ f$. The quiver of the Ext-algebra $\text{Ext}_{A^1}^*(\Delta^1, \Delta^1)$ is then

$$1 \xrightarrow{f} 2 \xrightarrow{g} 3,$$

with $\deg f = \deg g = 0$, subject to no additional relations.

Next, we draw the Loewy diagrams of the projective and standard modules over A^2 .

$$P^2(3): \begin{array}{c} 3 \\ \downarrow \\ 4 \\ \downarrow \\ 5 \end{array}, \quad P^2(4): \begin{array}{c} 4 \\ \downarrow \\ 5 \end{array}, \quad P^2(5): 5, \quad \Delta^2(3): 3, \quad \Delta^2(4): 4, \quad \Delta^2(5): 5.$$

As the standard modules over A^2 are simple, we see that we have non-split extensions $\alpha \in \text{Ext}_{A^2}^1(\Delta^2(3), \Delta^2(4))$ and $\beta \in \text{Ext}_{A^2}^1(\Delta^2(4), \Delta^2(5))$, such that $\beta\alpha = 0$. The quiver of the Ext-algebra $\text{Ext}_{A^2}^*(\Delta^2, \Delta^2)$ is then

$$3 \xrightarrow{\alpha} 4 \xrightarrow{\beta} 5,$$

subject to the relation $\beta\alpha = 0$. Finally, we draw the Loewy diagrams of the projective and standard modules over A .

$$P(1): 1, \quad P(2): \begin{array}{c} 2 \\ \downarrow \\ 1 \end{array}, \quad P(3): \begin{array}{c} 2 \\ \downarrow \\ 1 \end{array}, \quad P(4): \begin{array}{c} 4 \\ \downarrow \\ 5 \end{array}, \quad P(5): 5,$$

$$\Delta(1) \cong P(1), \quad \Delta(2) \cong P(2), \quad \Delta(3): \begin{array}{c} 3 \\ \downarrow \\ 2 \\ \downarrow \\ 1 \end{array}, \quad \Delta(4) \cong L(4), \quad \Delta(5) \cong P(5).$$

We see that there are homomorphisms $f': \Delta(1) \rightarrow \Delta(2)$, $g': \Delta(2) \rightarrow \Delta(3)$ and $h': \Delta(1) \rightarrow \Delta(3)$, such that $h' = g' \circ f'$. Additionally, we see that there are non-split extensions $\alpha' \in \text{Ext}_A^1(\Delta(3), \Delta(4))$ and $\beta' \in \text{Ext}_A^1(\Delta(4), \Delta(5))$, such that $\beta'\alpha' = 0$. The composition $\alpha'g'$ produces an extension in $\text{Ext}_A^1(\Delta(2), \Delta(4))$, and is therefore equal to zero, according to Lemma 48. Therefore, the Ext-algebra $\text{Ext}_A^*(\Delta, \Delta)$ is given by the quiver

$$1 \xrightarrow{f'} 2 \xrightarrow{g'} 3 \xrightarrow{\alpha'} 4 \xrightarrow{\beta'} 5,$$

subject to the relations $\beta'\alpha' = 0$ and $\alpha'g' = 0$. We observe that this algebra coincides with the algebra

$$C = \text{Ext}_{A^1}^*(\Delta^1, \Delta^1) \diamond \text{Ext}_{A^2}^*(\Delta^2, \Delta^2).$$

4.1. Exact Borel subalgebras under deconcatenations. Consider a deconcatenation $Q^1 \sqcup Q^2$ of Q at the sink or source ν . Let $B^1 \subset A^1$ be a subalgebra having a basis $\{b_1, \dots, b_n, e_\nu\}$ and let $B^2 \subset A^2$ be a subalgebra with basis $\{c_1, \dots, c_m, e_\nu\}$. Then $B = B^1 \diamond B^2$ is a subalgebra of A .

54. Proposition. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at the source ν and assume that we have regular exact Borel subalgebras $B^1 \subset A^1$ and $B^2 \subset A^2$. Then, A admits a regular exact Borel subalgebra.

Proof. For any $i \in Q_0^\ell \setminus \{\nu\}$, we have $\Delta(i) \cong \Delta^\ell(i)$. Let $0 \subset M_n \subset \dots \subset M_0 = \text{rad } \Delta^\ell(i)$ be a filtration with costandard subquotients. We claim that this is also a filtration of $\text{rad } \Delta(i)$. Consider a subquotient of the filtration

$$M_k / M_{k+1} \cong \nabla^\ell(j).$$

Then, $\nabla^\ell(j) \cong \nabla(j)$ for all $j \in Q_0^\ell$. For $j \neq \nu$, we appeal to Lemma 43. For $j = \nu$, this is automatic as $\nabla(\nu)$ is simple by virtue of ν being a source. Left to check is that we have a ∇ -filtration of $\text{rad } \Delta(\nu)$. Since ν is a source, we have a direct sum decomposition:

$$\text{rad } \Delta(\nu) = \text{rad } \Delta^1(\nu) \oplus \text{rad } \Delta^2(\nu).$$

Since A^1 has a regular exact Borel subalgebra, $\text{rad } \Delta^1(\nu)$ has a filtration whose subquotients are members of the set

$$\{\nabla(i) \mid i \in Q_0^1 \setminus \{\nu\}\}.$$

To see that $\nabla(\nu)$ does not occur as a composition factor, note that $[\text{rad } \Delta(\nu) : L(\nu)] = 0$. Indeed, an occurrence of a composition factor $L(\nu)$ would imply that $[P(\nu) : L(\nu)] \geq 2$, contradicting the fact that ν is a source. Similarly, $\text{rad } \Delta^2(\nu)$ has a filtration whose subquotients are members of the set

$$\{\nabla(i) \mid i \in Q_0^2 \setminus \{\nu\}\}.$$

We conclude that $\text{rad } \Delta(\nu) \in \mathcal{F}(\nabla)$, proving the statement. \square

55. Proposition. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at the sink ν . Assume further that we have regular exact Borel subalgebras $B^1 \subset A^1$, $B^2 \subset A^2$ and that the vertex ν is minimal with respect to the essential order on Q_0 . Then, there exists a regular exact Borel subalgebra $B \subset A$.

Proof. For any $i \in Q_0^\ell \setminus \{\nu\}$, we have $\Delta(i) \cong \Delta^\ell(i)$. Let $0 \subset M_n \subset \dots \subset M_0 = \text{rad } \Delta^\ell(i)$ be a filtration with costandard subquotients. We claim that this is also a filtration of $\text{rad } \Delta(i)$. Consider a subquotient of the filtration

$$M_k / M_{k+1} \cong \nabla^\ell(j).$$

Then $\nabla^\ell(j) \cong \nabla(j)$ for all $j \in Q_0^\ell \setminus \{\nu\}$, according to Lemma 43. For $j = \nu$, we claim that when ν is a sink, ν is minimal in the essential order on Q_0 if and only if $\nabla(\nu)$ is simple. Indeed, if ν is minimal, then $\nabla(\nu)$ is clearly simple. Conversely, assume that $\nabla(\nu)$ is simple and that there exists another vertex k such that $k \triangleleft^e \nu$. Then, we have

$$[\Delta(\nu) : L(k)] > 0 \quad \text{or} \quad (P(k) : \Delta(\nu)) > 0.$$

The first condition cannot be satisfied, since $\Delta(\nu)$ is simple by virtue of ν being a sink. According to [DR92, Lemma 2.5], we have

$$(P(k) : \Delta(\nu)) = [\nabla(\nu) : L(k)],$$

so this condition may not be satisfied either, since $\nabla(\nu)$ was assumed to be simple. Therefore, for $i \in Q_0^\ell \setminus \{\nu\}$, we have $\text{rad } \Delta(i) \in \mathcal{F}(\nabla)$, since A^1 and A^2 admit regular exact Borel subalgebras, by Theorem 23. Since ν is a sink, $\Delta(\nu) \cong L(\nu)$ which implies that $\text{rad } \Delta(\nu) = 0$, so there is nothing more to show. \square

56. Lemma. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at the sink ν . Then, for all $i \in Q_0$, we have $[\nabla(i) : L(\nu)] > 0$ if and only if $i = \nu$.

Proof. By definition, $\nabla(i)$ is a submodule of $I(i)$, so $[\nabla(i) : L(\nu)] > 0$ if and only if there is a path from ν to i , not passing through a vertex greater than i in Q_0 . This is a contradiction, since ν is assumed to be a sink. \square

From Lemma 56, it is also clear that for any module $M \in \mathcal{F}(\nabla)$, the costandard module $\nabla(\nu)$ appears as a subquotient in the filtration of M if and only if $[M : L(\nu)] > 0$.

Now we are ready to provide a necessary and sufficient condition for A to admit a regular exact Borel subalgebra $B \subset A$, where $A = KQ$ and $Q = Q^1 \sqcup Q^2$ is a deconcatenation at a sink ν .

57. Proposition. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at the sink ν and assume that $B^1 \subset A^1$ and $B^2 \subset A^2$ are regular exact Borel subalgebras. Moreover, assume that ν is not minimal with respect to the essential order on Q_0 . Then, the following are equivalent.

- (i) There exists a regular exact Borel subalgebra $C \subset A$.
- (ii) $[\text{rad } \Delta(i) : L(\nu)] = 0$, for all $i \in Q_0$.
- (iii) The vertex ν is maximal with respect to the essential order on Q_0 .

Proof. We first show the equivalence (i) \iff (ii). We assume that (ii) holds and claim that $\text{rad } \Delta(i) \in \mathcal{F}(\nabla)$, for all $i \in Q_0$. If $i = \nu$, we have $\Delta(\nu) \cong L(\nu)$ and consequently $\text{rad } \Delta(\nu) = 0$, so there is nothing to prove. Assume that $i \neq \nu$, where $i \in Q_0^\ell$. Then, $\Delta^\ell(i) \cong \Delta(i)$. According to the remark after Lemma 56, any ∇ -filtration of $\text{rad } \Delta^\ell(i)$

does not contain $\nabla^\ell(v)$ as a subquotient. Therefore, any ∇ -filtration of $\text{rad } \Delta^\ell(i)$ is automatically a ∇ -filtration of $\text{rad } \Delta(i)$, since all occurring subquotients in the filtration of $\text{rad } \Delta^\ell(i)$ are of the form $\nabla^\ell(j)$ with $j \neq v$.

Conversely, if $C \subset A$ is a regular exact Borel subalgebra, then $\text{rad } \Delta(i) \in \mathcal{F}(\nabla)$ for $1 \leq i \leq n$. Let i be such that $[\text{rad } \Delta(i) : L(v)] > 0$ and assume without loss of generality that $i \in Q_0^1 \setminus \{v\}$. Then, $\nabla(v)$ occurs in the ∇ -filtration of $\text{rad } \Delta(i)$, according to the remark after Lemma 56. Note that since v is assumed not to be minimal in the essential order, $\nabla(v)$ is not simple. Since Q^2 is connected, v has at least one neighbour, $j \in Q_0^2 \setminus \{v\}$. If $j \triangleleft^\ell v$, then $L(j)$ is a composition factor in $\nabla(v)$, which implies that $[\text{rad } \Delta(i) : L(j)] > 0$ since $\nabla(v)$ occurs in the ∇ -filtration of $\text{rad } \Delta(i)$. This is a contradiction, since $\text{rad } \Delta(i)$ is an A^1 -module. If $v \triangleleft^\ell j$, then $[\text{rad } \Delta(j) : L(v)] > 0$, so that $\nabla(v)$ is a subquotient of $\text{rad } \Delta(j)$. This contradicts $\nabla(v)$ being a subquotient of $\text{rad } \Delta(i)$, unless $\nabla(v)$ is simple. However, we assumed that v is not minimal in the essential order, guaranteeing it is not simple. If v and j are incomparable, we consider the module M , given by the following Loewy diagram:

$$\begin{array}{c} j \\ \downarrow \\ v \end{array}$$

Since A^2 is quasi-hereditary, \triangleleft^ℓ is adapted to A^2 , according to Lemma 3. Then, by definition, there is a vertex k such that $v \triangleleft^\ell k$, $j \triangleleft^\ell k$ and $[M : L(k)] \neq 0$. This is a contradiction, since M has no other composition factors besides $L(v)$ and $L(j)$. This proves (i) \iff (ii).

Next, we prove (ii) \iff (iii). Assume (ii) and that v is not maximal. Then there is another vertex k , such that $v \triangleleft^\ell k$. Then, by definition of the essential order, we have

$$[\Delta(k) : L(v)] > 0 \quad \text{or} \quad (P(v) : \Delta(k)) > 0.$$

The first condition can not be satisfied, as it contradicts (ii), and neither may the second, since $P(v)$ is simple by virtue of v being a sink. Conversely, assume that v is maximal and consider $P(i)$, for $i \in Q_0$. If $[P(i) : L(v)] = 0$, there is nothing to show, as $\text{rad } \Delta(i)$ is a submodule of a quotient of $P(i)$. Otherwise, if $[P(i) : L(v)] > 0$, there still holds $[\Delta(i) : L(v)] = 0$, since by definition, the composition factors $L(t)$ of $\Delta(i)$ satisfy $t \triangleleft i$. \square

58. Proposition. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at a sink or a source v . Assume that there are regular exact Borel subalgebras $B^1 \subset A^1$ and $B^2 \subset A^2$ containing the sets of idempotents $\{e_i \mid i \in Q_0^1\}$ and $\{e_j \mid j \in Q_0^2\}$, respectively, and that A admits a regular exact Borel subalgebra. Then $B^1 \diamond B^2 \subset A$ is a regular exact Borel subalgebra of A .

Proof. Consider the regular exact Borel subalgebra B^ℓ , where $\ell = 1, 2$. We have seen that $B^\ell = KQ_{B^\ell}$, where Q_{B^ℓ} is the following quiver.

- (i) The set of vertices of Q_{B^ℓ} is $Q_{B^\ell} = Q_0^\ell$.
- (ii) There is an arrow $i \rightarrow j$ in Q_{B^ℓ} if and only if $\dim \text{Ext}_{A^\ell}(\Delta^\ell(i), \Delta^\ell(j)) = 1$.

Let C be a regular exact Borel subalgebra. Then, $C = KQ_C$ for some quiver Q_C with vertex set Q_0 . We observe that, by Proposition 48 and 51, we have

$$\dim \text{Ext}_{A^\ell}(\Delta^\ell(i), \Delta^\ell(j)) = \dim \text{Ext}_A(\Delta(i), \Delta(j)),$$

for $i, j \in Q_0^\ell$. Using that B^1 and B^2 are regular exact Borel subalgebras, we get

$$\dim \text{Ext}_{B^\ell}^1(L(i), L(j)) = \dim \text{Ext}_{A^\ell}^1(\Delta^\ell(i), \Delta^\ell(j)) = \dim \text{Ext}_A^1(\Delta(i), \Delta(j)) = \dim \text{Ext}_C^1(L(i), L(j)).$$

This shows that for any $i, j \in Q_0^\ell$, there is an arrow $i \rightarrow j$ in the quiver of C if and only if there is an arrow $i \rightarrow j$ in the quiver of B^ℓ . Next, note that if $i \in Q_0^1 \setminus \{v\}$ and $x \in Q_0^2 \setminus \{v\}$, then there are no arrows between i and x in Q_C , because

$$\dim \text{Ext}_C^1(L(i), L(x)) = \dim \text{Ext}_A^1(\Delta(i), \Delta(x)) = 0,$$

according to Proposition 46. This means that the quivers of C and $B^1 \diamond B^2$ coincide, so that C and $B^1 \diamond B^2$ are isomorphic. This shows that $B^1 \diamond B^2$ has n simple modules up to isomorphism and is quasi-hereditary with simple standard modules.

Next, we check that the functor $A \otimes_{B^1 \diamond B^2} _$ is exact. To this end, we claim that A is projective as a right $B^1 \diamond B^2$ -module. Consider the decomposition of right A^1 -modules

$$A^1 \cong \bigoplus_{i \in Q_0^1} P^1(i).$$

Since $A^1 \otimes_{B^1} _$ is an exact functor, A^1 is projective as a right B^1 -module. Because $B^1 \subset A^1$ is a subalgebra, the decomposition above is also a decomposition of right B^1 -modules. This, in turn, implies that the summands $P^1(i)$ above are projective as right B^1 -modules. Because $B^1 \diamond B^2$ surjects onto B^1 , each B^1 -module has the natural structure of a $B^1 \diamond B^2$ -module (see the remark prior to Lemma 42). Then, the above decomposition is also a decomposition of right $B^1 \diamond B^2$ -modules, since the action of $\text{rad} B^2$ on each summand is trivial. Now, consider the decomposition

$$A \cong \bigoplus_{i \in Q_0^1 \setminus \{v\}} P(i) \oplus \bigoplus_{j \in Q_0^2 \setminus \{v\}} P(j) \oplus P(v)$$

of right A -modules. By the above argument, the summands $P(i)$ in the first term are projective as right $B^1 \diamond B^2$ -modules (note that for $i \in Q_0^1 \setminus \{v\}$, we have $P(i) \cong P^1(i)$). Similarly, the summands of the second term are projective right $B^1 \diamond B^2$ -modules. Left to check is that $P(v)$ is a projective right $B^1 \diamond B^2$ -module.

Let v be a sink. Then $P(v) \cong L(v)$ and there is nothing to show. Let v be a source. As noted earlier, $P^\ell(v)$ is projective as a right B^ℓ -module. Since v is a source also in the quiver of B^ℓ , the module $P_{B^\ell}^\ell(v)$ is the unique indecomposable projective B^ℓ -module having $L(v)$ as a composition factor. This implies that there is an isomorphism f^ℓ of right B^ℓ -modules:

$$f^\ell : P^\ell(v) \rightarrow P_{B^\ell}^\ell(v) \oplus M^\ell,$$

where M^ℓ is a direct sum of indecomposable projective B^ℓ -modules, with some multiplicities. Note that M^ℓ does not contain $P_{B^\ell}^\ell(v)$ as a direct summand. We claim that there is an isomorphism of right $B^1 \diamond B^2$ -modules

$$f : P(v) \rightarrow P_{B^1 \diamond B^2}(v) \oplus M^1 \oplus M^2.$$

Define f on paths in $P(v)$ by

$$p \mapsto \begin{cases} f^1(p) & \text{if } p \in A^1 \setminus \text{span}(e_v) \\ f^2(p) & \text{if } p \in A^2 \setminus \text{span}(e_v) \\ e_v & \text{if } p = e_v. \end{cases}$$

It is clear that f is bijective and a homomorphism of $B^1 \diamond B^2$ -modules, hence an isomorphism.

Next, we check that $A \otimes_{B^1 \diamond B^2} L(i) \cong \Delta(i)$, for all $i \in Q_0$. When $i \in Q_0^\ell \setminus \{v\}$, we know that $\Delta^\ell(i) \cong \Delta(i)$. In particular, $\Delta(i)$ is an A^ℓ -module. Let $\varphi : A^\ell \otimes_{B^\ell} L(i) \rightarrow \Delta(i)$ be an isomorphism. The module $A \otimes_{B^1 \diamond B^2} L(i)$ is generated by elements of the form $p \otimes e_i$, where p is a path in A . Note that, if p is a path in A^ℓ not equal to e_v , then

$$p \otimes e_i = p e_{s(p)} \otimes e_i = p \otimes e_{s(p)} e_i = 0.$$

To see this, note that p is a path in A^ℓ and $i \in Q_0^\ell \setminus \{v\}$, so that $i \neq s(p)$. Then, we may define a homomorphism

$$\tilde{\varphi} : A \otimes_{B^1 \diamond B^2} L(i) \rightarrow \Delta(i)$$

on the non-zero generators of $A \otimes_{B^1 \diamond B^2} e_i$ by

$$p \otimes_{B^1 \diamond B^2} e_i \mapsto \varphi(p \otimes_{B^\ell} e_i),$$

which is clearly an isomorphism. It remains to check that $A \otimes_{B^1 \diamond B^2} L(v) \cong \Delta(v)$. When v is a sink, we have $\Delta(v) \cong L(v)$, so the assertion is clear, given that $A^\ell \otimes_{B^\ell} L(v) \cong \Delta(v)$. Assume that v is a source and let

$$f_1 : A^1 \otimes_{B^1} L(v) \rightarrow \Delta^1(v), \quad \text{and} \quad f_2 : A^2 \otimes_{B^2} L(v) \rightarrow \Delta^2(v)$$

be isomorphisms. The module $A \otimes_{B^1 \diamond B^2} L(v)$ is generated by elements of the form $p \otimes e_v$, where $p \in A$ is some path. Note that with the exception of e_v , every such p is contained in A^1 or in A^2 . Define a map

$$f: A \otimes_{B^1 \diamond B^2} L(v) \rightarrow \Delta(v)$$

on generators by

$$p \otimes e_v \mapsto \begin{cases} f_1(p \otimes e_v) & \text{if } p \in A^1 \setminus \text{span}(e_v), \\ f_2(p \otimes e_v) & \text{if } p \in A^2 \setminus \text{span}(e_v), \\ e_v \otimes e_v & \text{if } p = e_v. \end{cases}$$

Note that, if $p \in A^1$ and $q \in A^2$, then $p \otimes e_v$ and $q \otimes e_v$ are either zero or linearly independent, except for when $p = q = e_v$. Then, f is a well-defined homomorphism because f_1 and f_2 are. Similarly, f is injective because f_1 and f_2 are and the intersection of their images equals $\text{span}(e_v)$. For surjectivity, note that as a vector space, we have $\Delta(v) = \text{span}(e_v) \oplus \text{rad} \Delta^1(v) \oplus \text{rad} \Delta^2(v)$.

To finish the proof, note that since $B^1 \diamond B^2$ and C are both exact Borel subalgebras, they are conjugate by Theorem 28. By Theorem 31, $B^1 \diamond B^2$ is regular since C is. \square

We conclude this section with an easy observation about the behavior of (certain) A_∞ -structures under deconcatenations.

59. Proposition. Let $Q = Q^1 \sqcup Q^2$ be a deconcatenation at a sink or a source v and assume that $\text{Ext}_{A^\ell}(\Delta^\ell, \Delta^\ell)$ is intrinsically formal, for $\ell = 1, 2$. Then, $\text{Ext}_A^*(\Delta, \Delta)$ is intrinsically formal.

Proof. Let m_n denote the higher multiplications on $\text{Ext}_A^*(\Delta, \Delta)$ and consider $m_n(\varphi_n, \dots, \varphi_1)$, where $\varphi_i \in \text{Ext}_A^*(\Delta, \Delta)$ for $1 \leq i \leq n$. Suppose that $m_n(\varphi_n, \dots, \varphi_1) \in \text{Ext}_A^k(\Delta(a), \Delta(b))$, for some $a, b \in Q_0$ and $k \geq 0$. If $a, b \in Q_0^\ell$, then $m_n(\varphi_n, \dots, \varphi_1) = 0$, because $\text{Ext}_{A^\ell}^*(\Delta^\ell, \Delta^\ell)$ is intrinsically formal. If $a \in Q_0^\ell \setminus \{v\}$ and $b \in Q_0^\ell \setminus \{v\}$, then $m_n(\varphi_n, \dots, \varphi_1) = 0$, according to Proposition 46. If $a = b = v$, then $m_n(\varphi_n, \dots, \varphi_1) = 0$, since standard modules have no non-split self-extensions. \square

4.2. Regular exact Borel subalgebras for path algebras of linear quivers. Let Q be the linear quiver

$$1 \text{ --- } 2 \text{ --- } \dots \text{ --- } n-1 \text{ --- } n$$

where the edges may be of either orientation, and put $\Lambda = KQ$. Denote by \mathbf{A}_m and \mathbf{B}_ℓ the special cases

$$1 \longrightarrow \dots \longrightarrow m \quad \text{and} \quad 1 \longleftarrow \dots \longleftarrow \ell,$$

respectively. By Proposition 24, $K\mathbf{A}_m$ admits a regular exact Borel subalgebra, regardless of the partial order on the vertices. We remark that all the statements from Section 2, describing the structure of $K\mathbf{A}_m$ in terms of its binary search tree, remain true for $K\mathbf{B}_\ell$, if we swap ‘‘left’’ and ‘‘right’’. For instance, over $K\mathbf{B}_\ell$, the composition factors of a standard module $\Delta_{K\mathbf{B}_\ell}(i)$ are found in the *left* subtree of the vertex labeled by i , rather than the right. From this, it follows that also $K\mathbf{B}_\ell$ admits a regular exact Borel subalgebra, regardless of the partial order on the vertices. Now, consider $\Lambda = KQ$. The quiver Q admits an iterated deconcatenation

$$Q = Q^1 \sqcup \dots \sqcup Q^s,$$

where, for each $1 \leq r \leq s$, the quiver Q^r is either \mathbf{A}_m or \mathbf{B}_ℓ . Recall that the partial order on the set of vertices of Q is constructed in the following way. When $Q = Q^1 \sqcup Q^2$, with $Q_0^1 \cap Q_0^2 = \{v\}$ and $i, j \in Q_0$, we say that $i \triangleleft j$ if one of the following hold.

- (i) We have $i, j \in Q_0^\ell$ and $i \triangleleft^\ell j$, for some ℓ .
- (ii) We have $i \in Q_0^\ell, j \in Q_0^{\bar{\ell}}, i \triangleleft^\ell v$ and $v \triangleleft^{\bar{\ell}} j$.

It is clear that, in this setting, v is maximal in Q_0 if and only if it is maximal in both Q_0^ℓ and $Q_0^{\bar{\ell}}$. We conclude that we have a special case of Proposition 55 as follows.

60. **Theorem.** The algebra Λ admits a regular exact Borel subalgebra $C \subset \Lambda$ if and only if each vertex $v \in Q_0$ which is a sink at which a deconcatenation occurs is minimal or maximal with respect to the essential order on Q_0 .

61. **Example.** Consider the quiver

$$Q = 1 \longrightarrow 2 \longrightarrow 3 \longleftarrow 4 \longleftarrow 5 \longrightarrow 6 \longrightarrow 7.$$


Deconcatenating Q as far as possible we find that

$$Q = (1 \xrightarrow{x} 2 \xrightarrow{y} 3) \sqcup (3 \xleftarrow{a} 4 \xleftarrow{b} 5) \sqcup (5 \xrightarrow{u} 6 \xrightarrow{v} 7).$$

Suppose the order on Q_0 is the one given by the following orders on Q_0^1, Q_0^2 and Q_0^3 .

$$1 \triangleleft_T 3, 2 \triangleleft_T 3, \quad 4 \triangleleft_T 5 \triangleleft_T 3, \quad 6 \triangleleft_T 5 \triangleleft_T 7,$$

corresponding to binary search trees



We compute the standard modules.

$$\Delta(1) \cong L(1), \quad \Delta(2) \cong L(2), \quad \Delta(3) \cong L(3), \quad \Delta(4) \cong L(4), \quad \Delta(5) \cong M(4,5), \quad \Delta(6) \cong L(6), \quad \Delta(7) \cong L(7)$$

The only standard module with non-trivial radical is $\Delta(5)$ with $\text{rad} \Delta(5) \cong L(4) \cong \nabla(4)$, so $\text{rad} \Delta(i) \in \mathcal{F}(\nabla)$ for all $1 \leq i \leq 7$. Then, A^1, A^2 and A^3 have regular exact Borel subalgebras given by

$$(1 \xrightarrow{x} 2 \quad 3), \quad (3 \quad 4 \xrightarrow{b} 5), \quad (5 \xrightarrow{vu} 6 \xrightarrow{v} 7).$$

We note that, since 3 is maximal in the first and second quivers, A has a regular exact Borel subalgebra, by Proposition 58, given by the following quiver.

$$1 \xrightarrow{x} 2 \quad 3 \quad 4 \xrightarrow{b} 5 \xrightarrow{vu} 6 \xrightarrow{v} 7.$$


Similarly to the above, we now wish to extend the statement of Proposition 38 to the algebra Λ discussed in this section.

62. **Proposition.** $\text{Ext}_\Lambda^*(\Delta, \Delta)$ is intrinsically formal.


Proof. This follows immediately from Proposition 38 and Proposition 59. □

In view of the situations considered in [FKR22], it would be interesting to do a similar investigation with quivers of Dynkin type \mathbb{D} and \mathbb{E} .

4.3. **A nonlinear example.** Consider the algebra A , given by the quiver




subject to the relations $ae = bd$, $ad = 0$ and $be = 0$. With the usual order $1 < 2 < 3 < 4$, the standard modules over A are



The only standard module with non-trivial radical is $\Delta(3)$, whose radical is isomorphic to the costandard module $\nabla(2)$. Therefore, A admits a regular exact Borel subalgebra $B \subset A$, which we find is given by the quiver




Consider now the algebra Λ given by the quiver



subject to the relations

$$ae = bd, \quad a'e' = b'd', \quad ad = 0, \quad a'd' = 0, \quad be = 0, \quad \text{and} \quad b'e' = 0,$$

with the order on the vertices being $1 < 2 < 3 < 4, \quad 7 < 6 < 5 < 4$. Then, 4 is maximal and Proposition 57 applies, and Λ has a regular exact Borel subalgebra $C \subset \Lambda$, given by the quiver



according to Proposition 58.

REFERENCES

- [ASS06] Ibrahim Assem, Andrzej Skowronski, and Daniel Simson. *Elements of the Representation Theory of Associative Algebras: Techniques of Representation Theory*. Vol. 1. London Mathematical Society Student Texts. Cambridge University Press, 2006.
- [BKK20] Tomasz Brzeziński, Steffen Koenig, and Julian Külshammer. “From quasi-hereditary algebras with exact Borel subalgebras to directed bocses”. In: *Bulletin of the London Mathematical Society* 52.2 (2020), pp. 367–378.
- [CPS88] E. Cline, B. Parshall, and L. Scott. “Finite-dimensional algebras and highest weight categories”. In: *J. Reine Angew. Math.* 391 (1988), pp. 85–99. ISSN: 0075-4102.
- [Con21] Teresa Conde. “All quasihereditary algebras with a regular exact Borel subalgebra”. In: *Advances in Mathematics* 384 (2021), p. 107751. ISSN: 0001-8708.
- [Con16] Teresa Conde. “On certain strongly quasi-hereditary algebras. PhD Thesis.” University of Oxford, 2016.
- [Cou20] Kevin Coulembier. “The classification of blocks in BGG category \mathcal{O} ”. In: *Mathematische Zeitschrift* 295 (2020), pp. 821–837.
- [DR92] Vlastimil Dlab and Claus Michael Ringel. “The module theoretical approach to quasi-hereditary algebras”. In: *Representations of Algebras and Related Topics*. Ed. by H. Tachikawa and S. Editors Brenner. London Mathematical Society Lecture Note Series. Cambridge University Press, 1992, 200–224.
- [FKR22] Manuel Flores, Yuta Kimura, and Baptiste Rognerud. “Combinatorics of quasi-hereditary structures”. In: *Journal of Combinatorial Theory, Series A* 187 (2022), p. 105559. ISSN: 0097-3165.
- [Kel01] Bernhard Keller. “Introduction to A -infinity algebras and modules”. In: *Homology, Homotopy and Applications* 3.1 (2001), pp. 1–35.
- [KS12] Angela Klamt and Catharina Stroppel. “On the Ext algebras of parabolic Verma modules and A infinity-structures”. English. In: *Journal of Pure and Applied Algebra* 216.2 (Feb. 2012), pp. 323–336. ISSN: 0022-4049.
- [KKO14] Steffen Koenig, Julian Külshammer, and Sergiy Ovsienko. “Quasi-hereditary algebras, exact Borel subalgebras, A -infinity-categories and boxes”. In: *Advances in Mathematics* 262 (2014), pp. 546 – 592. ISSN: 0001-8708.
- [Kön95] Steffen König. “Exact Borel subalgebras of quasi-hereditary algebras, I”. In: *Mathematische Zeitschrift* 220.1 (1995), pp. 399–426.
- [Mal42] A.I. Mal'tsev. “On the representation of an algebra as a direct sum of the radical and a semi-simple subalgebra”. In: *Dokl. Akad. Nauk SSSR* (1942), pp. 42–45.
- [OT10] Steffen Oppermann and Hugh Thomas. “Higher dimensional cluster combinatorics and representation theory”. In: *Journal of the European Mathematical Society* 14 (Jan. 2010).
- [Rin91] Claus Michael Ringel. “The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences.” In: *Mathematische Zeitschrift* 208.2 (1991), pp. 209–224.
- [Sco87] Leonard L. Scott. “Simulating Algebraic Geometry with Algebra, I: The Algebraic Theory of Derived Categories”. In: 1987.
- [Thu22] Markus Thuresson. “The Ext-algebra of standard modules over dual extension algebras”. In: *Journal of Algebra* 606 (2022), pp. 519–564. ISSN: 0021-8693.
- [Wed08] John H. Maclagan Wedderburn. “On Hypercomplex Numbers”. In: *Proceedings of The London Mathematical Society* (1908), pp. 77–118.
- [Zha00] Yuehui Zhang. “Strong Exact Borel Subalgebras of Quasi-Hereditary Algebras”. In: *Journal of Algebra - JALGEBRA* 231 (Sept. 2000), pp. 463–472.