
ON THE ENTROPY OF AN EXCHANGEABLE GRAPH

Anda Skeja
Institute of Mathematics
EPFL
anda.skeja@epfl.ch

Sofia C. Olhede
Institute of Mathematics
EPFL
sofia.olhede@epfl.ch

ABSTRACT

Model complexity remains a key feature of any proposed data generating mechanism. Measures of complexity can be extended to complex patterns such as signals in time and graphs. In this paper, we are concerned with the well-studied class of exchangeable graphs. Exchangeability for graphs implies a distributional invariance under node permutation and is a suitable default model that can widely be used for network data. For this well-studied class of graphs, we make a choice to quantify model complexity based on the (Shannon) entropy, resulting in graphon entropy. We estimate the entropy of the generating mechanism of a given graph, instead of choosing a specific graph descriptor suitable only for one graph generating mechanism. In this manner, we naturally consider the global properties of a graph and capture its important graph-theoretic and topological properties. Under an increasingly complex set of generating mechanisms, we propose a set of estimators of graphon entropy as measures of complexity for real-world graphs. We determine the large-sample properties of such estimators and discuss their usage for characterizing evolving real-world graphs.

Keywords Graph entropy, graph complexity, exchangeability, graphon entropy, entropy estimation.

1 Introduction

How complex is the generating mechanism of an observed graph, and when can its complexity be determined with some degree of certainty? This question has vexed numerous researchers and produced a number of different answers [9, 10, 28, 29, 46, 49–51]. There is ongoing debate on the quantification of complexity. One of the most commonly adopted methods is to define the complexity of a graph as a summary of its underlying structure [1, 29], [20], [13], [5]. Most of these answers are phrased in terms of *graph entropy*, and we will highlight when this is defined in terms of the generating mechanism of the graph, and when it is simply specified for a given graph. Entropy is a concept derived from information theory, and authors have considered its estimation for scalar random variables [7]. Once the object under study is more complicated than a scalar, more sophisticated measures can be introduced; see for example, [8, 25], which in turn reflect the nature of the generating mechanism of the observations as well as any additional random variation. Philosophically, many graphs can arise from the same generating mechanism, and we want to understand the complexity of that mechanism, not just of a single observed typical example.

Entropy as a topological measure of information content of deterministic graphs was first introduced in [41] in 1955, and this work was picked up in [31, 33–36, 44]. A slightly different approach was introduced in [26], where graph entropy indicates the number of bits that must be transmitted to clear up a graph vertex’s ambiguity. Yet another entropy measure was proposed for graphs, which is the so-called chromatic entropy [2]. This is the minimum entropy for any coloring in a graph. These measures do not involve probability distributions, but rather are calculated directly from a single deterministic graph itself. A probabilistic approach was proposed in [32], which used orbits of the automorphism group. However, specifying the whole automorphism group is computationally intractable for all graphs, but of moderate size. A review of the above-mentioned graph entropy measures is covered in [19]. The concept of entropy for network ensembles, and a definition and characterization for *structural entropy*, or as the author defines the entropy for a given degree sequence, was introduced in [10]. The Shannon entropy for network ensembles and its connection with the Gibbs and von Neumann entropies was discussed in [4]. The von Neumann entropy for graphs was introduced in [15]. In a recent study, [29] show that the von Neumann entropy of a graph is well-approximated by the Shannon entropy of the normalized degree sequence of the graph. We note related work that used entropy measures to

characterize complexity quantification [4, 5, 11], specific graph generating mechanisms [40], measuring the amount of information encoded in the graph [9, 52], measuring the order present in a graph and model choice [10, 40], detecting differences and similarities between graphs [5, 30, 46], and solving inference problems involving real-world graphs [6]. These measures are generally introduced for specific parametric graph models.

The aim of this paper is to estimate the entropy of the model that generated the graph, rather than trying to determine the entropy of a single graph realization. This different way of thinking, in contrast to attempting to quantify the complexity of a single given graph, will guide our endeavors. To understand an arbitrary exchangeable graph (an array satisfying a set of permutation invariances), we start from its representation using the Aldous–Hoover theorem (see, for example, [39] for a review). This representation is phrased in terms of a graph limit function, or equivalently, a so-called graphon. To make our methods as general as feasible, we assume that we study exchangeable binary arrays with 1-d continuous graph limits and that the arrays are disassociated. This former is not an extremely restrictive assumption (see, e.g., [24], for further discussion), while the latter is all we can estimate from one realized graph. If we observe more than one graph, we can non-parametrically determine the distribution of the observed entropies. We estimate the graphon using a stochastic block model following [12, 38] and, in turn, use it to produce a non-parametric plug-in estimator of the entropy. Alternative estimators are proposed if additional assumptions are made, such as the graphon being constant, separable, or low-rank.

It is natural to assume that some intrinsic properties of graphs are reflected in their entropy measures. Our measure of complexity captures important topological and graph-theoretic properties of complex graphs, the most important of which are linked to symmetry, connectedness, and density. Additionally, the graphon entropy is also insensitive to local features, such as the number of nodes, when the marginal edge probability is kept constant or only depends on the marginal edge probability, if not. As stated in [51], a measure of complexity that aims to be objective and universal should incorporate symmetries as simplifying factors. Owing to the inherent nature of our estimator, we naturally take into account symmetries. Symmetry is a structural characteristic of a graph. The structural information of a graph, is often computed given a partition of vertices (see, for example, [32–35, 41, 44]). A problem with allocating vertices to groups is the computational tractability of determining that partition, particularly if the whole automorphism group must be specified. This is an inevitable computational bottleneck, especially for all but moderate node numbers.

We propose a set of estimators of the entropy of a graph model given an observation of a graph. Our description in terms of a block model has an intrinsic resolution [48], but a whole set of resolutions all describe the intrinsic underlying graphon function [47]. We use the optimal block size as determined by [38]. This allows us to determine a non-parametric entropy estimator, appropriate to use in large graphs. We approximate the underlying graphon using block approximation, and for fixed block assignments, we show that the entropy estimator is asymptotically normal.

The paper is divided into four sections: First, we introduce the framework of exchangeable arrays, and their entropy in Section 2. We then discuss how the underlying graphon function can be estimated in Section 3, and follow by producing a plug-in estimator of the entropy. For an appropriate instantiation of this plug-in, we prove asymptotic normality (under modeling assumptions) specified in Section 3. To illustrate our obtained results, we implement a simulation study in Section 4, and study the complexity of Norwegian board of directors in Section 5. Finally, we offer some conclusions in Section 6.

2 Entropy of an exchangeable graph

Entropy was initially defined by [43] for quantifying the information produced by a source. Let X be a random variable taking values in \mathcal{X} , and be distributed according to $p : \mathcal{X} \rightarrow [0, 1]$. Then, the entropy of X is given by

$$\mathcal{H}(X) = \mathbf{E} \{ -\log(p(X)) \}, \quad (1)$$

where we will let \log denote the natural logarithm, and \mathbf{E} the expected value. Extending the concept to processes has already been considered for observed sequences in [8, 25]. For random graphs with no replicated observations, we have to define a model to make sense of observed variability. Similar to [8, 25], which considered stationary processes and random fields, we also restrict our interest to arrays satisfying certain permutation invariances, so-called exchangeable arrays [27, 39]. Let A be a two-dimensional $n \times n$ random array where $n \in \mathbb{N}$. We shall now define exchangeability for such arrays.

Definition 1. *Jointly Exchangeable Array* Let A be a two-dimensional $n \times n$ symmetric random array with binary entries. We say that A is jointly exchangeable if for any value of n , and any permutation $\pi : \{1, \dots, n\} \mapsto \{1, \dots, n\}$

$$\Pr\{A_{ij} = a_{ij} \ 1 \leq i, j \leq n\} = \Pr\{A_{ij} = a_{\pi(i)\pi(j)} \ 1 \leq i, j \leq n\}, \quad (2)$$

for $a_{ij} \in \{0, 1\}$.

As a consequence of exchangeability, we can represent A using Aldous–Hoover’s theorem [39] in terms of a *graph limit* or a *graphon* function, which we write as $f(x, y; \alpha)$, where α is a random variable.

Theorem 1 (Aldous–Hoover [23]). *Let A be a jointly exchangeable random array and ξ have independent $U(0, 1)$ entries. Then there is a function $f(\xi_i, \xi_j; \alpha)$ such that*

$$\Pr\{A_{ij} = 1 \mid \xi, \alpha\} = f(\xi_i, \xi_j; \alpha). \quad (3)$$

To represent any infinite array, we need the additional random variable α , but as we shall assume, we only have one realization of A . Thus, α cannot be estimated, and we shall suppress that dependence, assuming the graph is disassociated, and writing the graph limit as $f(x, y)$.

Finally, it has become popular to define a scaling $0 < \rho_n < 1$ such that for the $n \times n$ array, where n is large but finite, we have

$$\Pr\{A_{ij} = 1 \mid \xi, \alpha\} = \rho_n f(\xi_i, \xi_j; \alpha). \quad (4)$$

The entropy of a graphon is defined by [22].

Definition 2. *Entropy of an Exchangeable Array We define the binary entropy function $h : [0, 1] \rightarrow \mathbb{R}_+$ to be*

$$h(x) = -x \log(x) - (1 - x) \log(1 - x), \quad (5)$$

for $x \in [0, 1]$ where $h(0)$ and $h(1)$ are defined by continuity. The entropy of an exchangeable array A with graphon $f(x, y)$ is then [22] defined to be

$$\mathcal{H}(f) = \iint_{[0,1]^2} h(f(x, y)) \, dx \, dy. \quad (6)$$

We have stated the graph limit model without scaling the graph limit uniformly, as in [12, 14]. To capture some aspects of sparse graphs for large node numbers, we will include the uniform scaling by the marginal probability of observing an edge, which we write as $\rho_n > 0$. The definition of entropy remains the same, but the graph limit is now scaled by ρ_n .

3 Entropy estimation under different modeling assumptions

Estimating the graphon has remained a problem of intense scrutiny [12, 17, 38]. We form a plug-in estimator of the entropy from an estimator of the graph limit function $f(x, y)$, where the latter is written as $\hat{f}(x, y)$. This procedure then eventually yields the plug-in estimator

$$\hat{\mathcal{H}}(f) = \mathcal{H}(\hat{f}). \quad (7)$$

What assumptions can we then make to motivate a choice of graphon estimator? We shall list some potential choices for the nature of the graph limit below in Definitions 3–6. Each of these models come with a natural assumption of homogeneity and/or sparsity, and therefore, a natural estimation procedure, that is consistent.

Definition 3. *Constant graphon. We say that the exchangeable array A is generated by a constant graphon if $f(x, y) \propto 1$. This is often referred to as the Erdős–Renyi random graph.*

Definition 4. *Separable graphon. We say that the exchangeable array A is generated by a separable graphon if there is a function $g(x)$ such that $f(x, y) \propto g(x)g(y)$. This is equivalent to the Configuration model.*

Definition 5. *Smooth graphon. We say that the exchangeable array A is generated by a smooth graphon if $x, y \in [0, 1]$ and if $f(x, y)$ is a α -Hölder function. The restrictiveness of assuming the latent variable x being 1-d is limited, as discussed in [24].*

Definition 6. *Low-rank graphon. We say that the exchangeable array A is generated by a low-rank graphon if for a fixed J if $f(x, y)$ takes the form*

$$f(x, y) = \sum_{j=1}^J \lambda_j g_j(x) g_j(y), \quad (8)$$

for non-negative functions $g_j(x)$ and weights $\lambda_j \geq 0$.

Using each of these graphon models, we propose different estimators for the generating mechanism of A . What other assumptions may have been made? Low rank of the matrix $P_\xi = \mathbf{E}\{A \mid \xi\}$ has been used before [17]. Thus, we make an assumption about $f(x, y)$, say assumption $j \in \{1, 2, 3, 4\}$, so our function is $f_j(\cdot, \cdot)$ which encourages us to use a particular choice of estimator, resulting in $\hat{f}_j(\cdot, \cdot)$. From that estimator, then we compute

$$\hat{\mathcal{H}}_j(f) = \mathcal{H}(\hat{f}_j), \quad j \in \{1, \dots, J\}. \quad (9)$$

We begin by describing in detail how to form these estimators.

3.1 Estimating Constant Edge Probability Random Graphs

The simplest graph is generated by assigning a single edge probability $0 < \rho < 1$ to each edge, and we let each edge be drawn independently. Then,

$$A_{ij} \stackrel{iid}{\sim} \text{Bernoulli}(\rho), \quad 1 \leq i < j \leq n. \quad (10)$$

This can be represented using the graphon model with $f(x, y) = 1$ and $\rho_n = \rho$. The entropy of this model then becomes

$$\mathcal{H}_1(\rho) = -\rho \log(\rho) - (1 - \rho) \log(1 - \rho). \quad (11)$$

We see directly that as long as ρ does not take the value zero or infinity, the entropy is finite and is maximized at $\rho = 1/2$ (as is well-known). To form an estimator of this quantity, we can use a plug-in estimator after estimating ρ from the observed edge variables $\{A_{ij}\}_{i < j}$. This is formed as

$$\hat{\rho} = \frac{\sum_{i < j} A_{ij}}{\binom{n}{2}}. \quad (12)$$

We note directly the first two moments from $\hat{\rho}$;

$$\mathbf{E}\{\hat{\rho}\} = \rho, \quad \text{Var}\{\hat{\rho}\} = \frac{\rho(1 - \rho)}{\binom{n}{2}}. \quad (13)$$

The plug-in estimator of the entropy is then

$$\hat{\mathcal{H}}_1(\rho) = -\hat{\rho} \log(\hat{\rho}) - (1 - \hat{\rho}) \log(1 - \hat{\rho}). \quad (14)$$

Furthermore, as the A_{ij} are identically distributed and independent with this model, we can use a central limit theorem on $\sqrt{\rho n^2}(\hat{\rho} - \rho)$ to deduce asymptotic normality of $\hat{\rho}$, and the Delta method to determine the asymptotic normality of $\sqrt{\rho n^2}(\hat{\mathcal{H}}_1(\rho) - \mathcal{H}_1(\rho))$. Thus we deduce that (using the Delta method for fixed ρ)

$$\sqrt{\rho n^2}\{\hat{\mathcal{H}}_1(\rho) - \mathcal{H}_1(\rho)\} \xrightarrow{L} N(0, 2\rho(1 - \rho) \log^2(\rho(1 - \rho))).$$

This gives the exact distribution of the entropy for the constant edge probability random graph. If ρ varies with n , then there are levels of ρ for which the convergence in law still holds if the scaling is adjusted on both sides of the convergence sign. We observe symmetry in the case of $\rho \rightarrow 0$ and $\rho \rightarrow 1$, as is to be determined. There is nothing more to know about the entropy for the constant edge probability random graph for large graphs; as ρ tends to unity or zero, the entropy naturally becomes deterministic.

3.2 Estimating the Entropy of a Separable Graphon Model

The second simplest random graph is generated by assigning a separable graphon function, that is $f(x, y) = g(x)g(y)$, for some non-negative function g . Of course, we can estimate g to a sampled resolution determined by the number of nodes contained in our observed graph. We define the estimator of

$$\hat{g}(\xi_i) = C d_i, \quad (15)$$

where d_i is the i th degree of our observed graph, and ξ is defined in Theorem 1. To adhere to the same normalization as the graphon, we have to select C as $(n + 1)/\sqrt{\|d\|_1}$. C is then naturally both data- and sample size-dependent. Thus we have

$$\hat{g}(\xi_i) = \frac{(n + 1)}{\sqrt{\|d\|_1}} d_i, \quad i = 1, \dots, n. \quad (16)$$

A useful property of this estimator is its computational tractability. It can be computed by adding n numbers together, even if n such numbers have to be computed.

First, we note that

$$\frac{\hat{\rho}_n}{\rho_n} \xrightarrow{P} 1, \quad (17)$$

and so we can use Slutsky's theorem to deal with the remaining random variation.

We can now estimate $g(x)$ directly using (16). We then define the estimated entropy using a plug-in estimator and obtain the following estimator of the entropy. Following [22] we have that the entropy of an n -node graph generated from the separable graphon model takes the form:

$$\mathcal{H}_2(f) | \xi = -\frac{1}{\binom{n}{2}} \sum_{i < j} \{ \rho_n g(\xi_i) g(\xi_j) \log(\rho_n g(\xi_i) g(\xi_j)) + (1 - \rho_n g(\xi_i) g(\xi_j)) \log(1 - \rho_n g(\xi_i) g(\xi_j)) \},$$

$$\mathcal{H}_2(f) = \mathbf{E}_\xi \{ \mathcal{H}_2(f) | \xi \}.$$

Before we estimate $\mathcal{H}_2(f)$, we estimate $g(x)$ by using the degrees of A . We therefore form a plug-in estimator of $\mathcal{H}_2(f)$ using (15).

$$\widehat{\mathcal{H}}_2(f) = -\frac{1}{\binom{n}{2}} \sum_{i < j} \{ \hat{\rho}_n \hat{g}(\xi_i) \hat{g}(\xi_j) \log(\hat{\rho}_n \hat{g}(\xi_i) \hat{g}(\xi_j)) + (1 - \hat{\rho}_n \hat{g}(\xi_i) \hat{g}(\xi_j)) \log(1 - \hat{\rho}_n \hat{g}(\xi_i) \hat{g}(\xi_j)) \}.$$

This now leaves us with the task to determine the asymptotic distribution of $\widehat{\mathcal{H}}_2(f)$.

Theorem 2. [Asymptotic Normality of $\widehat{\mathcal{H}}_2(f)$] Assume that A_{ij} , for $1 \leq i, j \leq n$, is generated via Equation (4) using $f(x, y) = g(x)g(y)$ and $\rho_n = \omega(n^{-1} \log^3 n)$. As n grows sufficiently large, then the distribution of $\widehat{\mathcal{H}}_2(f)$ is given as

$$\frac{1}{\sqrt{\rho_n \log \rho_n}} \left\{ \widehat{\mathcal{H}}_2(f) - \mathbf{E} \widehat{\mathcal{H}}_2(f) \right\} \xrightarrow{\mathcal{L}} \mathbf{N} \left(0, \sigma_{\widehat{\mathcal{H}}_2}^2 \right), \quad (18)$$

where $\sigma_{\widehat{\mathcal{H}}_2}^2 \in \mathbb{R}^+$ is given in Equation (25).

Proof. See Appendix A. □

This estimator uses n degrees of freedom to estimate the function evaluated at the set of values $\{\xi_i\}_i$. The final averaging step makes it easier for us to derive the distribution of $\widehat{\mathcal{H}}_2(f)$, as in a sense, the quantity is obtained by averaging over all observations.

3.3 Estimating the Entropy of Smooth Graphons

We have looked at estimating the graph limit of a graph under strong simplifying assumptions. The next will be based on Aldous–Hoyer’s theorem and estimating $f(x, y; \alpha)$ for a single fixed drawn α . This will be implemented using the network histogram [38]. The network histogram is calculated from a stochastic block model fit to the observed graph.

Definition 7 (k -term Stochastic Block Model). Let z be an n -length vector taking values in $\{1, \dots, k\}^n$. The stochastic block model for an $n \times n$ array A is specified in terms of connectivity matrix Θ as

$$\Pr\{A_{ij} = 1 | z\} = \theta_{z_i, z_j}, \quad 1 \leq i < j \leq n.$$

The entropy of stochastic block model ensembles has been studied by [40] from a statistical mechanics approach. We define the entropy of the stochastic block model (with group a having h_a nodes in it, where the subscript is suppressed if all the blocks have the same size) by

$$\mathcal{H}(\theta) = -\sum_{a, b} \frac{h_a h_b}{n^2} \{ \theta_{ab} \log(\theta_{ab}) + (1 - \theta_{ab}) \log(1 - \theta_{ab}) \}. \quad (19)$$

If we approximate the graphon function by a block constant function [38], then it is straightforward to make a plug-in estimator of the graphon entropy. For convenience, we also define the aggregate block width function

$$H_a = \sum_{k=1}^a h_k, \quad a \in \{2, \dots, k\}, \quad H_1 = h_1.$$

We, therefore, note that a block constant approximation to the graphon function takes the form starting from (6) and defining the rectangular region ω_{ab} (for cases where n/k is an integer) by

$$\bar{f}(x, y) = -\sum_{a, b} \theta_{ab} \frac{h_a h_b}{n^2} \mathcal{I}((x, y) \in \omega_{ab}) \quad (20)$$

$$\omega_{ab} = \left\{ (x, y) : \frac{H_{a-1}}{n} \leq x < \frac{H_a}{n}, \frac{H_{b-1}}{n} \leq y < \frac{H_b}{n} \right\}.$$

With this choice of block constant function we note that the entropy equates to that of a stochastic block model and takes the form

$$\mathcal{H}(\bar{f}) = - \sum_{a,b} \frac{h_a h_b}{n^2} \{ \theta_{ab} \log(\theta_{ab}) + (1 - \theta_{ab}) \log(1 - \theta_{ab}) \}. \quad (21)$$

For details, see Appendix B.

Several methods are available for estimating z the label vector or cluster vector. Common methods include spectral clustering [45], likelihood methods [3, 12, 16, 18, 53], and modularity maximization [37]. We define the plug-in estimator of entropy $\widehat{\mathcal{H}}_3(f)$ by replacing θ_{ab} by its estimator $\widehat{\theta}_{ab}$.

Theorem 3. *Assume that A is generated as a realization from Definition 5 and $f(x, y) \in \mathcal{F}_\alpha(M)$ is a α -Hölder continuous function with $\rho_n = \Theta(1)$ bounded away from zero and one, and $k \leq \log n/n$. Then for any $C' > 0$, there exists a constant $C > 0$ only depending on C' and M , such that*

$$\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| \leq C \left(n^{-2\alpha/1+\alpha} + \frac{\log n}{n} \right),$$

with probability at least $1 - \exp(-C'n)$, uniformly over $f \in \mathcal{F}_\alpha(M)$.

Proof. See Appendix C. □

Theorem 4. *Assume that A is generated as a realization from Definition 7 and $\min_{a,b} \theta_{ab} = \omega \left(\frac{\log h_{ab}}{h_{ab}} \right)$, then for a given class assignment z , we have*

$$\frac{n^2}{\sigma_{\widehat{\mathcal{H}}_3}} \{ \widehat{\mathcal{H}}_3(f) - \mathbf{E} \widehat{\mathcal{H}}_3(f) \} \xrightarrow{\mathcal{L}} N(0, 1),$$

where $\sigma_{\widehat{\mathcal{H}}_3}^2$ is given in Equation (32).

Proof. See Appendix D. □

So in the case of dense graphs, we have shown how the block model can be used to estimate the entropy of an arbitrary smooth function $f(x, y)$.

3.4 Estimating Low-Rank Graphons

Finally, there are methods that approximate $\mathbf{E}\{A | \xi\} = P$ providing an entry-specific estimator, such as the U-SVT of [17]. These can also be converted to entropy estimates by plugging in estimates for P in

$$\mathcal{H}_4(P) = - \sum_{i,j} \frac{1}{n^2} \{ p_{ij} \log(p_{ij}) + (1 - p_{ij}) \log(1 - p_{ij}) \}. \quad (22)$$

We produce an estimator of this by replacing $\mathcal{H}_4(P)$ with $\mathcal{H}_4(\widehat{P})$, and use the symmetry of \widehat{P} to determine all n^2 entries from the initial $\binom{n}{2}$ estimates.

4 Simulation Study

To study the properties of estimators of the graphon entropy we shall study a few scaled graphon examples, namely for $a_0, a_1, a_2 \in \mathbb{R}^+$ and $I_x(\alpha, \beta)$ the regularized incomplete beta function

$$\begin{aligned} f_1(x, y) &= 4xy \\ f_2(x, y) &= a_0 + 4a_1 I_x^{-1}(\alpha_1, \alpha_1) I_y^{-1}(\alpha_1, \alpha_1) + 4a_1 I_x^{-1}(\alpha_1, \alpha_1) I_y^{-1}(\alpha_1, \alpha_1). \end{aligned}$$

To get a better sense of the marginal properties of this choice of graphon we compute

$$\int_0^1 f_2(x, y) dy = a_0 + 2a_1. \quad (23)$$

We note that this marginal is flat, so no information can be garnered from it (or the graph degrees), unlike $f_1(x, y)$. We propose to use three different entropy estimators suitable for the Erdős–Renyi graph $\widehat{\mathcal{H}}_1(f)$, the separable graphon estimator of $\widehat{\mathcal{H}}_2(f)$ and the block model approximation to the continuous graphon $\widehat{\mathcal{H}}_3(f)$.

It is standard to consider the case of scaled graphons [14]. We cannot define the entropy of an unscaled graphon function as the input of the entropy needs to lie between zero and one. For ρ_n scaling with n , the entropy does not converge to a single large sample limit. However, we can still study the mean square error between $\widehat{\mathcal{H}}_3(f)$ and $\mathbf{E}\widehat{\mathcal{H}}_3(f)$, as this will also scale with n . We will define

$$sRMSE = \frac{RMSE}{\sqrt{\rho_n \log \rho_n}} = \frac{\sqrt{\frac{1}{N} \sum_{i=1}^N (\widehat{\mathcal{H}}_{3,i} - \mathbf{E}\widehat{\mathcal{H}}_{3,i})^2}}{\sqrt{\rho_n \log \rho_n}},$$

where the $\widehat{\mathcal{H}}_{3,i}$ is computed for the i th graph realization out of $N > 1$, and where each $\widehat{\mathcal{H}}_{3,i}$ implicitly depends on n , unless $\rho_n \propto 1$ when we divide by unity. We study this root mean square error because $\mathbf{E}\widehat{\mathcal{H}}_3$ does not converge to a single limit with changing values of ρ_n over n due to the combination of the linear and logarithmic terms. This, in a sense, enables us to say that $\widehat{\mathcal{H}}_3 \approx \mathbf{E}\widehat{\mathcal{H}}_{3,i}$ unless the random variation around this value is too large. The simulation study confirms that after splitting our root mean square error into a deterministic factor that varies with ρ_n ; the remaining variation is the same for the same value of n , and is decreasing with n , so the estimated entropy is getting “close” to its expectation. We now refer to Table 1 to study the performance of the proposed estimators across simulation studies. For the simple separable graphon (f_1) (as expected), the separable graphon estimator performs best, as the block model allows for a more flexible approximation, but exhibits finite sample size bias. The variances are comparable between the separable graphon fit and the graphon fit, but the bias thus beats out the graphon. The constant graphon performs best on variance but is, of course, (in general) biased. Looking at a more complex graphon structure, namely f_2 , we can

	Constant graphon	Separable graphon	Graphon
RMSE	0.100352	0.012595	0.014522
Bias Squared	0.009933	1.81561×10^{-6}	5.38146×10^{-5}
Variance	0.000139	0.0001600	0.0001602

Table 1: Comparison of the proposed estimator for the separable graphon, f_1 , $n = 600$ $\rho_n \propto 1$, over 100 repetitions.

study the mean square error in Table 2. This is not a separable graphon, so the entropy estimator based on assuming

	Constant graphon	Separable graphon	Graphon
RMSE	0.049566	0.047929	0.0029468
Bias Squared	0.002456	0.002296	7.41×10^{-7}
Variance	2.63×10^{-7}	6.45×10^{-7}	8.022×10^{-6}

Table 2: Comparison of the proposed estimator for the non-separable graphon, f_2 , $n = 600$, $\rho_n \propto 1$, over 100 repetitions.

separability of the graphon is no longer the best performing method. The graphon estimator (and thus also the plug-in estimator of the entropy) in this case has a smaller bias, and a somewhat larger variance, but is an improvement on the other methods.

Studying the behavior of the entropy estimator over a range of sample sizes, we have plotted the root mean square scaled by $\sqrt{\rho_n \log(\rho_n)}$, as this illustrates the large sample behavior of the entropy estimator, see Figure 1. Normally when estimating the graphon for the entire unit square, methods work well as long as $\rho_n = \Omega(\log(n)^{2+\epsilon}/n)$, for $\epsilon > 0$. Here we are not estimating a function, but only a single scalar of the entropy, and so the last averaging over the n^2 symmetric edge variables, ensure a better convergence rate with reasonable performance even for very sparse graphs.

5 Study of Complexity of Boards from Norwegian Corporations

In addition to the simulation study, we would also like to see what entropy can tell us about a real-world graph.

We shall here study a graph constructed from observations of the behavior of corporate boards in Norway. The data set is publicly available at ¹. This dataset corresponds to a record of ties corresponding to sitting on the same board of Norwegian public limited companies and is studied by [42]. The data were put together from recordings of the Norwegian Business Register (5 August 2009). The aforementioned study aimed to understand the effects of a new legislation on gender representation. The law required having at least 40% representation of each gender on the boards of public limited companies by 2008 [42]. It is stated in [42] that the law succeeded in raising the participation of women on boards, thereby expanding the pool of women who could serve as directors.

¹graphs.skewed.de/net/board_directors




Figure 1: Log-log plot of the decay of $sRMSE(\hat{\mathcal{H}}_3(f))$ as a function of $n \in \{200, 400, \dots, 1000\}$ in the dense and sparse regime, for 100 trials.




Figure 2: Average proportion of women and entropy change over time

The average proportion of women on boards is shown in Fig. 2 (a). This ratio demonstrates a substantial increase in the representation of women on boards during the observation period. The time from the announcement of the law’s enforcement until it took effect, or between mid-2005 and 2008, had the biggest growth in particular. However, similarly, Figure 2 (b) shows that the highest decrease in entropy took place in the period May 2005 to December 2008. We obtained Figure 2 (b) by first estimating the graphon underlying the graph via the network histogram algorithm [38] for each point in time and then computing the entropy across observed time points as described in Equation 22. This shows us that our measure captures changes occurring in a real complex graph. The reason for the decrease in entropy in parallel with the increase in the average proportion of women on boards could be attributed to the findings of [42], who state that women have more social capital than men, and that the average betweenness score of women is 70% higher than that of men. Thus, female director boards in the mentioned dataset tend to form connected components, which in turn serve to make the graph more structured.

6 Conclusion

In this paper, we determined a method to estimate the entropy of an arbitrary exchangeable graph. Our methodology can be applied to any exchangeable graph; we approximate such using a block model approximation, thereby reducing variability. We were able to show that a sequence of estimators of increasing complexity could be constructed, and determined consistency and sampling characteristics. Additionally, we showed that our entropy estimator demonstrates promising results in both simulated and real networks. This is very important because our method can have extensive applications in inference problems for complex networks. Our main contribution, compared to previous approaches is treating large graphs without imposing strong assumptions. We achieved this by taking into account graph-theoretic properties, without depending on granular characterization or structure-dependent formulae. Our measure is insensitive to local properties of the network, and computationally cheaper compared to the algorithmic complexity defined in [51].

Acknowledgments

This work was supported by the European Research Council under Grant CoG 2015-682172NETS, within the Seventh European Union Framework Program. The first author also acknowledges support via the Excellence Scholarship from the Swiss Confederation.

Appendices

A. Proof of Theorem 2

Proof. The plug-in estimator for the entropy of the Configuration Model is given as follows:

$$\widehat{\mathcal{H}}_2(\rho_n f) = \frac{1}{\binom{n}{2}} \sum_{i < j} -\{[\rho_n \hat{g}(\xi_i) \hat{g}(\xi_j) \log(\rho_n \hat{g}(\xi_i) \hat{g}(\xi_j))] + [(1 - \rho_n \hat{g}(\xi_i) \hat{g}(\xi_j)) \log(1 - \rho_n \hat{g}(\xi_i) \hat{g}(\xi_j))]\}.$$

$\widehat{\mathcal{H}}_2(\rho_n f)$ has *almost* the form of U-statistic given as:

$$U_n = \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(\xi_i, \xi_j),$$

with

$$h(\xi_i, \xi_j) = -\{[\rho_n \hat{g}(\xi_i) \hat{g}(\xi_j) \log(\rho_n \hat{g}(\xi_i) \hat{g}(\xi_j))] + [(1 - \rho_n \hat{g}(\xi_i) \hat{g}(\xi_j)) \log(1 - \rho_n \hat{g}(\xi_i) \hat{g}(\xi_j))]\}.$$

We say *almost* as $\hat{g}(\cdot)$ is a random function. We then define:

$$\widehat{\mathcal{H}}_2(\rho_n f) = \widehat{\mathcal{H}}_2(\rho_n f) - \mathcal{E}(\rho_n f) + \mathcal{E}(\rho_n f),$$

where

$$\mathcal{E}(\rho_n f) = \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} \widetilde{h}(\xi_i, \xi_j),$$

with

$$\widetilde{h}(\xi_i, \xi_j) = -\{\rho_n g(\xi_i) g(\xi_j) \log(\rho_n g(\xi_i) g(\xi_j))\} + \{(1 - \rho_n g(\xi_i) g(\xi_j)) \log(1 - \rho_n g(\xi_i) g(\xi_j))\}.$$

We now need to show that $\widehat{\mathcal{H}}_2(\mathcal{G}) - \mathcal{E}(\mathcal{G}) = o_P(1)$. We give two auxiliary lemmas that will help us achieve our goal.

Lemma 1. *Let $\widehat{g}(\xi_i) = g(\xi_i) + Y_i$, where $Y_i = o_P(1)$.*

Proof.

$$\begin{aligned} \widehat{g}(\xi_i) &= g(\xi_i) + Y_i = \frac{d_i}{\sqrt{\|d\|_1}} \\ &= \frac{d_i}{\sqrt{\mathbf{E}[\|d\|_1]}} \underbrace{\left\{ \sqrt{\frac{\mathbf{E}[\|d\|_1]}{\|d\|_1}} \right\}}_{\xrightarrow{P} 1} \end{aligned} \quad (24)$$

We already know that $\mathbf{E}[Y_i] = 0$, so for determining the order of Y_i we calculate $\text{Var}(Y_i)$ and then use Chebyshev's inequality.

$$\begin{aligned} \text{Var}(Y_i) &= \text{Var}(\widehat{g}(\xi_i) - g(\xi_i)) \\ &= \text{Var}_\xi \mathbf{E}_{A|\xi} [\widehat{g}(\xi_i) - g(\xi_i) | \xi] + \mathbf{E}_\xi \text{Var}_{A|\xi} \left(\frac{d_i}{\sqrt{\|d\|_1}} | \xi \right) \\ &= \mathbf{E}_\xi \text{Var}_{A|\xi} \left(\frac{d_i}{\sqrt{\|d\|_1}} | \xi \right) \end{aligned}$$

Rewriting (24) we get:

$$\begin{aligned}
& \frac{d_i}{\sqrt{\mathbf{E}[\|d\|_1]}} \left\{ \sqrt{\frac{\mathbf{E}[\|d\|_1]}{\|d\|_1}} \right\} \\
& \xrightarrow{P} 1 \\
& = \frac{d_i}{\sqrt{\mathbf{E}[\|d\|_1|\xi]} \underbrace{\sqrt{1 + \frac{\|d\|_1 - \mathbf{E}[\|d\|_1|\xi]}{\mathbf{E}[\|d\|_1|\xi]}}}_{\sqrt{1+Z_n}}}
\end{aligned}$$

To determine the order of Z_n with $\mathbf{E}[Z_n] = 0$, we again need to calculate $\text{Var}(Z_n)$. We start by noting:

$$\begin{aligned}
& \text{Var}(\|d\|_1 - \mathbf{E}[\|d\|_1|\xi]) \\
& = \text{Var}(\|d\|_1|\xi) \\
& = \text{Var} \sum_i \sum_{j \neq i} A_{ij} | \xi \\
& = \sum_i \sum_{j \neq i} \rho_n g(\xi_i) g(\xi_j) (1 - \rho_n g(\xi_i) g(\xi_j)) \\
& \leq g_{max}^2 \rho_n n(n-1),
\end{aligned}$$

uniformly in ξ where $g_{max} = \sup_x g(x)$. For the denominator we have:

$$\mathbf{E}[\|d\|_1|\xi] = \sum_{i=1}^n \sum_{j \neq i} \rho_n g(\xi_i) g(\xi_j) \geq \rho_n n(n-1) g_{min}^2$$

Hence, for $\text{Var}(Z_n)$ we obtain

$$\text{Var}(Z_n) \leq \frac{g_{max}^2 \rho_n n(n-1)}{\rho_n^2 n^2 (n-1)^2 g_{min}^2} = \frac{g_{max}^2}{\rho_n n(n-1) g_{min}^2}.$$

As $Z_n = o_P(1)$, by the Stochastic Taylor Expansion we get

$$Y_i = \frac{d_i}{\sqrt{\mathbf{E}[\|d\|_1|\xi]}} \left\{ 1 + \frac{1}{2} Z_n + O_P(\sqrt{\text{Var}(Z_n)}) \right\}.$$

Assuming Z_n is uniformly integrable,

$$\text{Var}(Y_i) = \text{Var}(Y_i|\xi_i) = \text{Var}\left(\frac{d_i}{\sqrt{\mathbf{E}[\|d\|_1|\xi]}} \mid \xi\right).$$

We already know the order of $\sqrt{\mathbf{E}[\|d\|_1|\xi]}$, so now we calculate $\text{Var}(d_i|\xi)$ as follows

$$\begin{aligned}
\text{Var}(d_i|\xi) & = \text{Var}\left(\sum_{j \neq i} A_{ij} \mid \xi\right) = \sum_{j \neq i} \text{Var}(A_{ij}|\xi) \\
& = \sum_{j \neq i} \rho_n g(\xi_i) g(\xi_j) (1 - \rho_n g(\xi_i) g(\xi_j)) \\
& \leq \rho_n (n-1) g_{max}^2.
\end{aligned}$$

So,

$$\text{Var}(Y_i) \leq \frac{1}{\rho_n g_{min}^2 n(n-1)} \rho_n (n-1) g_{max}^2 = \mathcal{O}\left(\frac{1}{n}\right),$$

which in limit tends to zero, i.e.,

$$\lim_{n \rightarrow \infty} \text{Var}(Y_i) \rightarrow 0.$$

We need $k = O(\sqrt{n})$, so we choose k to be $O(\sqrt{n})$. Then, by Chebyshev's inequality we have:

$$\lim_{n \rightarrow \infty} P(|\sum_{i,j} Y_{ij}| > \epsilon) \leq \frac{1}{n^2} \frac{1}{E[d_{ij}]^2};$$

which corresponds to

$$\lim_{n \rightarrow \infty} P(|\sum_{i,j} Y_{ij}| > \epsilon) = 0;$$

Therefore, $Y_i = o_p(1)$. □

Lemma 2.

$$H_2(n, f) = E(n, f) + E(n, f)$$

Proof. As stated before, we know that $g(i) = g(i) + Y_i$ with $Y_i = o_p(1)$. Rewriting $H_2(n, f)$ in terms of $g(i) = g(i) + Y_i$, we get:

$$\begin{aligned} H_2(n, f) &= \frac{1}{2} \sum_{i,j} \frac{(g(i) + Y_i)(g(j) + Y_j) \log((g(i) + Y_i)(g(j) + Y_j))}{\text{term (1)}} \\ &= \frac{1}{2} \sum_{i,j} \frac{((1 + \frac{Y_i}{g(i)})(1 + \frac{Y_j}{g(j)})) \log(1 + \frac{Y_i}{g(i)} + \frac{Y_j}{g(j)})}{\text{term (2)}} \end{aligned}$$

After simple algebra and using identities for the logarithmic function; we arrive to the conclusion that:

$$H_2(n, f) = E(n, f) + E(n, f);$$

with

$$E(n, f) = O_p(Y_i^4);$$

As $Y_i = o_p(1)$, and every term in $E(n, f)$ has at least one factor of Y_i , then $E(n, f) = o_p(1)$. □

So, the distribution of $H_2(n, f)$ is determined only by $E(n, f)$ and its mean and variance are calculated via the classic theory of U-statistics as follows:

$$= \frac{1}{n} \log \frac{1}{n} \int_0^1 \int_0^1 g(x) \log(g(x)) dx + \frac{1}{n} \int_{[0;1]^2} (1 - g(i)g(j)) \log(1 - g(i)g(j)) dg_i dg_j$$

and

$$\text{Var}(H_2) = \frac{1}{n} \int_0^1 \int_0^1 (1 - g(x)g(y)) \log^2(1 - g(x)g(y)) dx dy + \frac{1}{n} \int_0^1 g(x) \log^2(g(x)) dx - \left(\int_0^1 g(x) \log(g(x)) dx \right)^2 \quad (25)$$

with

$$h_k = \text{Var}(h_k(i; j));$$

where

$$h_k(x_1, \dots, x_k) = E[h(X_1, \dots, X_m) | X_1 = x_1, \dots, X_k = x_k];$$

So, $E[h_1(i; j)] = E[E[h(i; j) | i, j]] = E[h(i; j)]$:

Then,

$$\begin{aligned} h_1 &= \text{Var}(h_1(i; j)) \\ &= E[h_1^2(i; j)] - E^2[h_1(i; j)] \\ &= \int_0^1 \int_0^1 g(y) \log^2(g(y)) g(y) dy + \int_0^1 g(j) \log^2(g(j)) dj - \left(\int_0^1 g(y) \log(g(y)) dy \right)^2 \\ &= \int_0^1 \log^2(g(x)) dx + \frac{1}{\log(2)} \int_0^1 g(x) \log(g(x)) dx - \left(\int_0^1 g(x) \log(g(x)) dx \right)^2 \end{aligned}$$

and

$$\begin{aligned}\zeta_2 &= \text{Var}(\tilde{h}_2(\xi_i, \xi_j)) \\ &= g_2^2(\rho_n^2 \log^2(\rho_n) - 2\rho_n^2 \log \rho_n + \rho_n) + g_3^2(2\rho_n^2 \log \rho_n - 2\rho_n^3) + g_2\{\rho_n^2 \int g^2(\xi_i) \log^2 g(\xi_i) d\xi_i\} \\ &\quad + 4\rho_n \int g^2(\xi_i) \log g(\xi_i) d\xi_i + g_4\rho_n^4 - \{\rho_n \log \rho_n + 2\rho_n \int_0^1 g(x) \log g(x) dx - \frac{1}{\log(2)}\rho_n - \mathcal{O}(\rho_n^2)\}^2,\end{aligned}$$

where we define the function p -norm to be

$$g_p = \|g\|_p = \int_0^1 g^p(x) dx,$$

for $p = 1, 2, 3, 4$. □

B. Entropy of a block-constant graphon

Given the block constant graphon function

$$f(x, y) = \sum_{a,b} \theta_{ab} \mathcal{I}(H(a-1) < x < H(a)) \mathcal{I}(H(b-1) < y < H(b)),$$

its entropy is given as

$$\mathcal{H}(f) = \iint_{[0,1]^2} h(f(x, y)) dx dy,$$

where $h(\cdot)$ is the binary entropy function defined in (2). Define the region

$$\omega_{ab} = \{(x, y) \mid H(a-1) < x < H(a), \quad H(b-1) < y < H(b)\},$$

then we get

$$\begin{aligned}\mathcal{H}(f) &= - \iint_{[0,1]^2} \left\{ \sum_{a,b} \theta_{ab} \mathcal{I}((x, y) \in \omega_{ab}) \log(\theta_{ab}) + \sum_{a,b} \mathcal{I}((x, y) \in \omega_{ab}) (1 - \theta_{ab}) \log((1 - \theta_{ab})) \right\} dx dy \\ &= - \sum_{a', b'} \iint_{\omega_{a', b'}} \left\{ \sum_{a,b} \mathcal{I}((x, y) \in \omega_{ab}) [\theta_{ab} \log(\theta_{ab}) + (1 - \theta_{ab}) (1 - \log(\theta_{ab}))] \right\} dx dy \\ &= - \sum_{a,b} \frac{h_a h_b}{n^2} [\theta_{ab} \log(\theta_{ab}) + (1 - \theta_{ab}) \log(1 - \theta_{ab})].\end{aligned}$$

C. Proof of Theorem 3

Proof. We note that the Kullback-Liebler divergence between two Bernoulli distributions $f(x, y)$ and $\hat{f}(x, y) = f(x, y) + \delta$, with $0 < f(x, y) < 1$ and $-f(x, y) \leq \delta \leq 1 - f(x, y)$ for $x, y \in [0, 1]$ is given as

$$D(f(x, y) \parallel \hat{f}(x, y)) = f(x, y) \log f(x, y) - f(x, y) \log \hat{f}(x, y) + (1 - f(x, y)) \log(1 - f(x, y)) - (1 - f(x, y)) \log(1 - \hat{f}(x, y)). \quad (26)$$

For brevity, we will let f and \hat{f} denote $f(x, y)$ and $\hat{f}(x, y)$, respectively. Using (26), we get the following

$$\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| = \left| \iint_{[0,1]^2} D(f \parallel \hat{f}) - D(\hat{f} \parallel f) dx dy + \iint_{[0,1]^2} f \log \hat{f} + (1 - f) \log(1 - \hat{f}) - \hat{f} \log f - (1 - \hat{f}) \log(1 - f) dx dy \right|$$

By the triangle inequality and f being Riemann-integrable, we then obtain,

$$\begin{aligned}\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| &\leq \iint_{[0,1]^2} \left| D(f \parallel \hat{f}) \right| dx dy + \iint_{[0,1]^2} \left| D(\hat{f} \parallel f) \right| dx dy \\ &\quad + \iint_{[0,1]^2} \left| f \log \hat{f} + (1 - f) \log(1 - \hat{f}) \right| dx dy + \iint_{[0,1]^2} \left| \hat{f} \log f + (1 - \hat{f}) \log(1 - f) \right| dx dy.\end{aligned}$$

We now use $\log(1+x) \leq x$ for $x > -1$ and $\log(1-x) \leq -x - \frac{x^2}{2}$ to obtain

$$\iint_{[0,1]^2} \left| f \log \hat{f} + (1-f) \log(1-\hat{f}) \right| dx dy \leq \iint_{[0,1]^2} \left| D(f \parallel \hat{f}) \right| dx dy + \iint_{[0,1]^2} \frac{\delta^2}{|1-f|} dx dy$$

Similarly, for the last term we have

$$\iint_{[0,1]^2} \left| \hat{f} \log f + (1-\hat{f}) \log(1-f) \right| dx dy \leq \iint_{[0,1]^2} \left| D(\hat{f} \parallel f) \right| dx dy$$

Rewriting what we had so far, we obtain

$$\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| \leq \underbrace{2 \iint_{[0,1]^2} \left| D(f \parallel \hat{f}) \right| dx dy}_{(1)} + \underbrace{2 \iint_{[0,1]^2} \left| D(\hat{f} \parallel f) \right| dx dy}_{(2)} + \underbrace{C_{min} \iint_{[0,1]^2} (\hat{f} - f)^2 dx dy}_{(3)}, \quad (27)$$

where $C_{min} = 1/[\min(f_{min}, 1 - f_{max})]$. In order to bound (1), (2) and (3) we use results from Lemma C.9 of [47] to obtain the following

$$\iint_{[0,1]^2} \left| D(f \parallel \hat{f}) \right| \leq \left(\frac{C_{min}}{2} + \frac{C_{min}^3}{3} \right) \iint_{[0,1]^2} (\hat{f} - f)^2 dx dy,$$

and

$$\iint_{[0,1]^2} \left| D(\hat{f} \parallel f) \right| \leq \left(\frac{C_{min}}{2} + \frac{2C_{min}^3}{3} + \frac{C_{min}^4}{3} \right) \iint_{[0,1]^2} (\hat{f} - f)^2 dx dy. \quad (28)$$

As the final step, we refer to Equation (27) and use Theorem 2.3 from [21] as

$$\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| \leq D \iint_{[0,1]^2} (\hat{f} - f)^2 dx dy,$$

where $D > 0$, and conclude that for any $C' > 0$, there exists a constant $C > 0$ only depending on C' and M , such that

$$\left| \widehat{\mathcal{H}}_3(f) - \mathcal{H}_3(f) \right| \leq C \left(n^{-2\alpha/1+\alpha} + \frac{\log n}{n} \right),$$

with probability at least $1 - \exp(-C'n)$, uniformly over $f \in \mathcal{F}_\alpha(M)$. \square

D. Proof of Theorem 4

Proof. We note that for any fixed class assignment z , $\widehat{\theta}_{ab} = \bar{A}_{ab}$ (see Lemma 3 and its proof in Appendix D1) is a sum of h_{ab} independent Bernoulli random variables, therefore it is a Poisson-Binomial random variable with mean $\bar{\theta}_{ab}$, where for $1 \leq i, j \leq n$, $p_{ij} = \rho_n f(\xi_i, \xi_j)$ we have $\bar{\theta}_{ab} = \frac{1}{h_{ab}} \sum_{i < j} p_{ij} \mathcal{I}(z_i = a) \mathcal{I}(z_j = b)$, and $h_{ab} = h_a h_b$ for $a \neq b$. By the Lyapunov Central Limit Theorem, we then have

$$h_{ab}(\widehat{\theta}_{ab} - \bar{\theta}_{ab}) | z \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma_{ab}^2). \quad (29)$$

Let $g(x) = -\{x \log x + (1-x) \log(1-x)\}$ for $x \in (0, 1)$. As $g'(x)$ exists and is non-zero, by the Delta method we have

$$h_{ab}\{g(\widehat{\theta}_{ab}) - g(\bar{\theta}_{ab})\} | z \xrightarrow{\mathcal{L}} \mathcal{N}(0, [g'(\bar{\theta}_{ab})]^2 \sigma_{ab}^2). \quad (30)$$

Summing (30) over a, b and dividing by n^2 , we get

$$\sum_{a,b} \frac{h_{ab}}{n^2} \{g(\widehat{\theta}_{ab}) - g(\bar{\theta}_{ab})\} = \widehat{\mathcal{H}}_3(f) - \mathbf{E}\widehat{\mathcal{H}}_3(f)$$

As the distribution of a finite sum of normal random variables is again normal, we can conclude that

$$\frac{n^2}{\sigma_{\widehat{\mathcal{H}}_3}} \{\widehat{\mathcal{H}}_3(f) - \mathbf{E}\widehat{\mathcal{H}}_3(f)\} | z \xrightarrow{\mathcal{L}} \mathcal{N}(0, 1), \quad (31)$$

where

$$\sigma_{\widehat{\mathcal{H}}_3}^2 = \sum_{a,b} [g'(\bar{\theta}_{ab})]^2 \sigma_{ab}^2. \quad (32)$$

\square

D1. Proof of Lemma 3

Lemma 3. Let G be a graph on n nodes generated by the stochastic block model with parameters k, z, Θ , and A the $n \times n$ adjacency matrix. Then, for a choice of z , the maximum likelihood estimator of θ_{ab} for $1 \leq a, b \leq k$ is given as:

$$\hat{\theta}_{ab} = \bar{A}_{ab} = \frac{\sum_{i < j} A_{ij} \mathcal{I}(z_i = a) \mathcal{I}(z_j = b)}{\sum_{i < j} \mathcal{I}(z_i = a) \mathcal{I}(z_j = b)}$$

Proof. We formulate the likelihood dependent on connection probability θ_{ab} and a group label $z_i \in \{1, \dots, k\}$, for $i = 1, \dots, n$ for k groups. Then,

$$\mathbb{P}(A) = \prod_{i < j} \theta_{z_i z_j}^{A_{ij}} (1 - \theta_{z_i z_j})^{1 - A_{ij}},$$

and the log-likelihood is given as

$$\ell(\theta, z) = \sum_{i < j} A_{ij} \log(\theta_{z_i z_j}) + (1 - A_{ij}) \log(1 - \theta_{z_i z_j}).$$

We define, $\sum_i \mathcal{I}(z_i = a) = h_a$, and $h_{ab} = h_a h_b$ for $a \neq b$ so that with

$$\ell(\theta, z) = \sum_{a \leq b} h_{ab} \bar{A}_{ab}(z) \log \theta_{ab} + (h_{ab} - h_{ab} \bar{A}_{ab}(z)) \log(1 - \theta_{ab}),$$

with $\bar{A}_{ab}(z) = \sum_{i < j} h_{ab}^{-1} A_{ij} \mathcal{I}(z_i = a) \mathcal{I}(z_j = b)$. For a fixed choice of z , by maximizing the likelihood we find that

$$\begin{aligned} \frac{\partial \ell(\theta, z)}{\partial \theta_{ab}} &= h_{ab} \frac{\bar{A}_{ab}(z)}{\theta_{ab}} - h_{ab} \frac{1 - \bar{A}_{ab}(z)}{1 - \theta_{ab}} \\ \hat{\theta}_{ab} &= \bar{A}_{ab}(z), \quad 1 \leq a \leq b \leq k. \end{aligned}$$

Indeed, $\hat{\theta}_{ab}$ is a maximum profile likelihood estimator as

$$\frac{\partial^2 \ell(z, \theta_{ab})}{\partial \theta_{ab}^2} = -h_{ab} \frac{\bar{A}_{ab}(z)}{\theta_{ab}^2} - h_{ab} \frac{1 - \bar{A}_{ab}(z)}{(1 - \theta_{ab})^2} < 0.$$

□

References

- [1] ALBERT, R., AND BARABÁSI, A.-L. Statistical mechanics of complex networks. *Reviews of modern physics* 74, 1 (2002), 47.
- [2] ALON, N., AND ORLITSKY, A. Source coding and graph entropies. *IEEE Transactions on Information Theory* 42, 5 (1996), 1329–1339.
- [3] AMINI, A. A., CHEN, A., BICKEL, P. J., AND LEVINA, E. Pseudo-likelihood methods for community detection in large sparse networks. *The Annals of Statistics* 41, 4 (2013), 2097–2122.
- [4] ANAND, K., AND BIANCONI, G. Entropy measures for networks: Toward an information theory of complex topologies. *Physical Review E* 80, 4 (2009), 045102.
- [5] ANAND, K., BIANCONI, G., AND SEVERINI, S. Shannon and von neumann entropy of random networks with heterogeneous expected degree. *Physical Review E* 83, 3 (2011), 036109.
- [6] ANAND, K., KRIUKOV, D., AND BIANCONI, G. Entropy distribution and condensation in random networks with a given degree distribution. *Physical Review E* 89, 6 (2014), 062807.
- [7] ANTOS, A., AND KONTOYIANNIS, I. Convergence properties of functional estimates for discrete distributions. *Random Structures & Algorithms* 19, 3-4 (2001), 163–193.
- [8] BERCHER, J., AND VIGNAT, C. Estimating the entropy of a signal with applications. *IEEE transactions on signal processing* 48, 6 (2000), 1687–1694.
- [9] BIANCONI, G. The entropy of randomized network ensembles. *EPL (Europhysics Letters)* 81, 2 (Dec 2007), 28005.
- [10] BIANCONI, G. Entropy of network ensembles. *Physical Review E* 79, 3 (2009), 036114.
- [11] BIANCONI, G. Statistical mechanics of multiplex networks: Entropy and overlap. *Physical Review E* 87, 6 (2013), 062806.
- [12] BICKEL, P. J., AND CHEN, A. A nonparametric view of network models and Newman–Girvan and other modularities. *Proceedings of the National Academy of Sciences* 106, 50 (2009), 21068–21073.
- [13] BOCCALETTI, S., LATORA, V., MORENO, Y., CHAVEZ, M., AND HWANG, D.-U. Complex networks: Structure and dynamics. *Physics reports* 424, 4-5 (2006), 175–308.
- [14] BOLLOBÁS, B., AND RIORDAN, O. Metrics for sparse graphs. *arXiv preprint arXiv:0708.1919* (2007).
- [15] BRAUNSTEIN, S. L., GHOSH, S., AND SEVERINI, S. The laplacian of a graph as a density matrix: a basic combinatorial approach to separability of mixed states. *Annals of Combinatorics* 10, 3 (2006), 291–317.
- [16] CELISSE, A., DAUDIN, J.-J., AND PIERRE, L. Consistency of maximum-likelihood and variational estimators in the stochastic block model. *Electronic Journal of Statistics* 6 (2012), 1847–1899.
- [17] CHATTERJEE, S. Matrix estimation by universal singular value thresholding. *The Annals of Statistics* 43, 1 (2015), 177–214.
- [18] CHOI, D. S., WOLFE, P. J., AND AIROLDI, E. M. Stochastic blockmodels with a growing number of classes. *Biometrika* 99, 2 (2012), 273–284.
- [19] DEHMER, M., AND MOWSHOWITZ, A. A history of graph entropy measures. *Information Sciences* 181, 1 (2011), 57–78.
- [20] DOROGOVTSSEV, S. N., AND MENDES, J. F. Evolution of networks. *Advances in physics* 51, 4 (2002), 1079–1187.
- [21] GAO, C., LU, Y., AND ZHOU, H. H. Rate-optimal graphon estimation. *The Annals of Statistics* 43, 6 (2015), 2624–2652.
- [22] HATAMI, H., JANSON, S., AND SZEGEDY, B. Graph properties, graph limits, and entropy. *Journal of Graph Theory* 87, 2 (2018), 208–229.
- [23] JANSON, S., AND DIACONIS, P. Graph limits and exchangeable random graphs. *Rendiconti di Matematica e delle sue Applicazioni. Serie VII* (2008), 33–61.
- [24] JANSON, S., AND OLHEDE, S. Can smooth graphons in several dimensions be represented by smooth graphons on $[0, 1]^d$? *Examples and Counterexamples 1* (2021), 100011.
- [25] KONTOYIANNIS, I., ALGOET, P. H., SUHOV, Y. M., AND WYNER, A. J. Nonparametric entropy estimation for stationary processes and random fields, with applications to english text. *IEEE Transactions on Information Theory* 44, 3 (1998), 1319–1327.

- [26] KÖRNER, J. Coding of an information source having ambiguous alphabet and the entropy of graphs. In *6th Prague conference on information theory* (1973), pp. 411–425.
- [27] LAURITZEN, S. L. Exchangeable rasch matrices. *Rend. Mat. Appl.*(7) 28, 1 (2008), 83–95.
- [28] LI, A., AND PAN, Y. Structural information and dynamical complexity of networks. *IEEE Transactions on Information Theory* 62, 6 (2016), 3290–3339.
- [29] LIU, X., FU, L., WANG, X., AND ZHOU, C. On the similarity between von neumann graph entropy and structural information: Interpretation, computation, and applications. *IEEE Transactions on Information Theory* 68, 4 (2022), 2182–2202.
- [30] MENICETTI, G., BIANCONI, G., CASTELLANI, G., GIAMPIERI, E., AND REMONDINI, D. Multiscale characterization of ageing and cancer progression by a novel network entropy measure. *Molecular BioSystems* 11, 7 (2015), 1824–1831.
- [31] MOWSHOWITZ, A. Entropy and the complexity of graphs: I. an index of the relative complexity of a graph. *The bulletin of mathematical biophysics* 30, 1 (1968), 175–204.
- [32] MOWSHOWITZ, A. Entropy and the complexity of graphs: I. an index of the relative complexity of a graph. *The bulletin of mathematical biophysics* 30, 1 (1968), 175–204.
- [33] MOWSHOWITZ, A. Entropy and the complexity of graphs: Ii. the information content of digraphs and infinite graphs. *The Bulletin of mathematical biophysics* 30, 2 (1968), 225–240.
- [34] MOWSHOWITZ, A. Entropy and the complexity of graphs: Iii. graphs with prescribed information content. *The bulletin of mathematical biophysics* 30, 3 (1968), 387–414.
- [35] MOWSHOWITZ, A. Entropy and the complexity of graphs: Iv. entropy measures and graphical structure. *The bulletin of mathematical biophysics* 30, 4 (1968), 533–546.
- [36] MOWSHOWITZ, A., AND DEHMER, M. Entropy and the complexity of graphs revisited. *Entropy* 14, 3 (2012), 559–570.
- [37] NEWMAN, M. E., AND GIRVAN, M. Finding and evaluating community structure in networks. *Physical review E* 69, 2 (2004), 026113.
- [38] OLHEDE, S. C., AND WOLFE, P. J. Network histograms and universality of blockmodel approximation. *Proceedings of the National Academy of Sciences* 111, 41 (Oct 2014), 14722–14727.
- [39] ORBANZ, P., AND ROY, D. M. Bayesian models of graphs, arrays and other exchangeable random structures. *IEEE transactions on pattern analysis and machine intelligence* 37, 2 (2014), 437–461.
- [40] PEIXOTO, T. P. Entropy of stochastic blockmodel ensembles. *Physical Review E* 85, 5 (2012), 056122.
- [41] RASHEVSKY, N. Life, information theory, and topology. *The bulletin of mathematical biophysics* 17, 3 (1955), 229–235.
- [42] SEIERSTAD, C., AND OPSAHL, T. For the few not the many? the effects of affirmative action on presence, prominence, and social capital of women directors in norway. *Scandinavian journal of management* 27, 1 (2011), 44–54.
- [43] SHANNON, C. E. A mathematical theory of communication. *ACM SIGMOBILE mobile computing and communications review* 5, 1 (2001), 3–55.
- [44] TRUCCO, E. A note on the information content of graphs. *The bulletin of mathematical biophysics* 18, 2 (1956), 129–135.
- [45] VON LUXBURG, U. A tutorial on spectral clustering. *Statistics and computing* 17, 4 (2007), 395–416.
- [46] WEST, J., BIANCONI, G., SEVERINI, S., AND TESCHENDORFF, A. E. Differential network entropy reveals cancer system hallmarks. *Scientific reports* 2, 1 (2012), 1–8.
- [47] WOLFE, P. J., AND OLHEDE, S. C. Nonparametric graphon estimation. *arXiv preprint arXiv:1309.5936* (2013).
- [48] YOUNG, J.-G., ST-ONGE, G., DESROSIERS, P., AND DUBÉ, L. J. Universality of the stochastic block model. *Physical Review E* 98, 3 (2018), 032309.
- [49] ZENIL, H., KIANI, N. A., AND TEGNÉR, J. Low-algorithmic-complexity entropy-deceiving graphs. *Physical Review E* 96, 1 (2017), 012308.
- [50] ZENIL, H., KIANI, N. A., AND TEGNÉR, J. A review of graph and network complexity from an algorithmic information perspective. *Entropy* 20, 8 (2018), 551.

- [51] ZENIL, H., SOLER-TOSCANO, F., DINGLE, K., AND LOUIS, A. A. Correlation of automorphism group size and topological properties with program-size complexity evaluations of graphs and complex networks. *Physica A: Statistical Mechanics and its Applications* 404 (2014), 341–358.
- [52] ZHAO, K., KARSAI, M., AND BIANCONI, G. Models, entropy and information of temporal social networks. In *Temporal Networks*. Springer, 2013, pp. 95–117.
- [53] ZHAO, Y., LEVINA, E., AND ZHU, J. Consistency of community detection in networks under degree-corrected stochastic block models. *The Annals of Statistics* 40, 4 (2012), 2266–2292.