

On the Maximum Atom-Bond Sum-Connectivity Index of Graphs

Tariq Alraqad*, Hicham Saber, Akbar Ali

*Department of Mathematics, Faculty of Science,
University of Ha'il, Ha'il, Saudi Arabia*

t.alraqad@uoh.edu.sa, Hicham.saber7@gmail.com, akbarali.maths@gmail.com

Abstract

Let G be a graph with m edges, namely e_1, \dots, e_m . For $i \in \{1, \dots, m\}$, let d_{e_i} be the number of edges adjacent with e_i . The sum of the numbers $\sqrt{1 - 2(d_{e_i} + 2)^{-1}}$ over $1 \leq i \leq m$ is known as the atom-bond sum-connectivity (ABS) index of G . A pendant vertex of G is a vertex having degree 1. A subset S of the vertex set of G is said to be independent if the vertices of S are pairwise non-adjacent in G . The maximum number among cardinalities of all independent sets of G is known as the independence number of G . The least number of colors required to color the vertices of a graph, so that no two adjacent vertices have the same color, is termed as the chromatic number. This paper characterizes the graphs attaining the greatest values of the ABS index over the classes of graphs of a given order and with a fixed (i) chromatic number (ii) independence number (iii) number of pendant vertices.

keywords: topological index; atom-bond sum-connectivity; graph

Mathematics Subject Classification: 05C07, 05C90.

1 Introduction

In this paper, just finite and simple graphs are taken into account. The sets of edges and vertices of a graph G are denoted, respectively, by $E(G)$ and $V(G)$. The degree of a vertex $v \in V(G)$ is indicated by $d_v(G)$ or just d_v if the graph being discussed is unambiguous. We utilize the conventional notation and nomenclature of (chemical) graph theory, and we refer readers to the relevant books, for example, [1, 2].

Graphs are being used to model chemical structures by replacing atoms and bonds of the structures with vertices and edges, respectively. In this way, it is possible to study the chemical structures using the concepts of graph theory. Such a field of study is usually

*Corresponding author.

referred to as the chemical graph theory. Graph invariants that adopt quantitative values are widely termed as topological indices in chemical graph theory.

The connectivity index (also known as the Randić index) [3], a well-known topological index, was devised in the 1970s by the chemist Milan Randić under the name “branching index” [4]. Soon after its discovery, the connectivity index quickly found a variety of uses [5–7] in chemistry and consequently it became one of the most applied and well-researched topological indices. For a graph G , the connectivity index is defined as

$$R(G) = \sum_{vw \in E(G)} \frac{1}{\sqrt{d_v d_w}}.$$

The Randić index has been modified in several ways. Here, we mention two topological indices which were introduced by taking into consideration the definition of the Randić index, namely the “sum-connectivity (SC) index” [8] and the “atom-bond connectivity (ABC) index” [9]. These indices have the following definitions for a graph G :

$$SC(G) = \sum_{vw \in E(G)} \frac{1}{\sqrt{d_v + d_w}}$$

and

$$ABC(G) = \sum_{ab \in E(G)} \sqrt{\frac{d_v + d_w - 2}{d_v d_w}}.$$

Detail regarding the mathematical aspects of the SC and ABC indices may be found in the review papers [10] and [11], respectively.

By utilizing the definitions of the ABC and SC indices, a novel topological index – the atom-bond sum-connectivity (ABS) index – has recently been proposed in [12]. For a graph G , this index is defined as

$$ABS(G) = \sum_{uv \in E(G)} \left(\frac{d_u + d_v - 2}{d_u + d_v} \right)^{\frac{1}{2}}.$$

In the paper [12], graphs possessing the maximum and minimum values of the ABS index were characterized over the classes of graphs and (chemical) trees of a given order; such kind of extremal results regarding unicyclic graphs were found in [13], where also chemical applications of the ABS index were reported. The preprint [14] is concerned with the problems of determining graphs possessing the minimum ABS index among all trees of a fixed order and/or a given number of pendant vertices; see also [15] where one of these two problems is attacked independently.

A pendant vertex in a graph is a vertex of degree 1. The least number of colors required to color the vertices of a graph, so that no two adjacent vertices have the same color, is termed as the chromatic number. A subset S of the vertex set of G is said to be independent if the vertices of S are pairwise non-adjacent in G . The maximum number among cardinalities of all independent sets of G is known as the independence number of G and it is denoted by $\alpha(G)$. This paper characterizes the graphs attaining the greatest values of the ABS index over the classes of graphs of a given order and with a fixed (i) chromatic number (ii) independence number (iii) number of pendant vertices.

2 Results

In order to avoid trivialities, throughout this section, we consider only connected graphs. To prove our results, we need few technical lemmas.

Lemma 1 (see [12]) *Let u and v be non-adjacent vertices in a connected graph G . If $G + uv$ is the graph obtained from G by adding the edge uv in G then*

$$ABS(G + uv) > ABS(G).$$

Lemma 2 *Let*

$$f(x, y) = \left(\frac{x + y - 2}{x + y} \right)^{\frac{1}{2}},$$

where $\min\{x, y\} \geq 1$. For every positive real number s , define the function $g_s(x, y) = f(x + s, y) - f(x, y)$. Then f is strictly increasing in x and in y . The function g_s is strictly decreasing and convex in x and in y .

Proof. The first and second partial derivatives function $\frac{\partial f}{\partial x}$ of f with respect to x and y are calculated as

$$\frac{\partial f}{\partial x}(x, y) = \frac{\partial f}{\partial y}(x, y) = (x + y - 2)^{-\frac{1}{2}}(x + y)^{-\frac{3}{2}},$$

$$\frac{\partial^2 f}{\partial x^2}(x, y) = \frac{\partial^2 f}{\partial y^2}(x, y) = -\frac{1}{2}(x + y - 2)^{-\frac{3}{2}}(x + y)^{-\frac{3}{2}} - \frac{3}{2}(x + y - 2)^{-\frac{1}{2}}(x + y)^{-\frac{5}{2}}$$

Clearly $\frac{\partial f}{\partial x}(x, y) > 0$, whenever $x > 1$, and thus f is strictly increasing in x and in y . Since $\frac{\partial^2 f}{\partial x^2}(x, y) < 0$, whenever $x > 1$, we get $\frac{\partial f}{\partial x}(x, y)$ is strictly decreasing in x when $x \geq 1$. This implies that $\frac{\partial g_s}{\partial x}(x, y) = \frac{\partial f}{\partial x}(x + s, y) - \frac{\partial f}{\partial x}(x, y) < 0$ when $x \geq 1$, and thus $g_s(x, y) = f(x + s, y) - f(x, y)$, is strictly decreasing in x when $x \geq 1$. Additional, $\frac{\partial^2 f}{\partial x^2}(x, y)$

is strictly increasing $x \geq 1$. So $\frac{\partial^2 g_s}{\partial x^2}(x, y) = \frac{\partial^2 f}{\partial x^2}(x+s, y) - \frac{\partial^2 f}{\partial x^2}(x, y) > 0$, and hence $g_s(x, y)$, is convex in x when $x \geq 1$. \square

Lemma 3 *Let M and N be real numbers satisfying $1 \leq M \leq N$. Then for every positive real number s , the function $h_s(x) = g_s(x, M) - g_s(x, N)$ is increasing in x when $x \geq 1$.*

Proof. When $x \geq 1$, we have $\frac{\partial^2 f}{\partial x \partial y}$ is strictly increasing in x . So $\frac{\partial^2 g_s}{\partial x \partial y}(x, y) = \frac{\partial^2 f}{\partial x \partial y}(x+s, y) - \frac{\partial^2 f}{\partial x \partial y}(x, y) > 0$. Thus $\frac{\partial g_s}{\partial x}$ is increasing in y , and hence $h'_s(x) = \frac{\partial g_s}{\partial x}(x, N) - \frac{\partial g_s}{\partial x}(x, M) > 0$. Therefore $h_s(x)$ is increasing in x when $x \geq 1$. \square

The next theorem gives a sharp upper bound on the *ABS* value of all connected graphs of a fixed order and fixed chromatic number. A graph whose vertex set can be partitioned into r sets V_1, V_2, \dots, V_r in such a way that all the vertices in every V_i (with $1 \leq i \leq r$) are pairwise non-adjacent is known as an r -partite graph, where $r \geq 2$ and the sets V_1, V_2, \dots, V_r are called the partite sets. If, in addition, every vertex of partite set V_i is adjacent to all the vertices of the other partite sets for $i = 1, 2, \dots, r$, then the graph is called the complete r -partite graph. We denote, by $T_{n, \chi}$, the complete χ -partite graph of order n such that $|n_i - n_j| \leq 1$, where n_i , with $i = 1, 2, \dots, \chi$, is the number of vertices in the i -th partite set of $T_{n, \chi}$.

Theorem 1 *If G be a connected graph of order $n \geq 5$ and having chromatic number $\chi \geq 3$, Then*

$$ABS(G) \leq \frac{r(r-1)q^2}{2} \sqrt{\frac{n-q-1}{n-q}} + r(\chi-r)q(q+1) \sqrt{\frac{2n-2q-3}{2n-2q-1}} + \frac{(\chi-r)(\chi-r-1)(q+1)^2}{2} \sqrt{\frac{n-q-2}{n-q}}, \quad (1)$$

where q and r are non negative integers such that $n = q\chi + r$ and $r < \chi$. Moreover, the equality holds in (1) if and only if $G \cong T_{n, \chi}$.

Proof. Let G be a graph having the maximum *ABS* in the class of all connected graphs of a fixed order n and with a fixed chromatic number χ , where $3 \leq \chi \leq n-1$ and $n \geq 5$. Note that the vertex set $V(G)$ of G can be partitioned into χ independent subsets, say V_1, V_2, \dots, V_χ such that $|V_i| = n_i$ for $i = 1, 2, \dots, \chi$, provided that $n_1 \leq n_2 \leq \dots \leq n_\chi$. Consequently, G is isomorphic to a χ -partite graph and hence, by Lemma 1, it must be isomorphic to the complete χ -partite graph $K_{n_1, n_2, \dots, n_\chi}$. To complete the proof, we have

to show that $n_\chi - n_1 \leq 1$. Contrarily, assume that $n_\chi - n_1 \geq 2$. Let $G' \cong K_{n'_1, n'_2, \dots, n'_\chi}$, where $n'_1 = n_1 + 1$, $n'_\chi = n_\chi - 1$, and $n'_i = n_i$ for every $i \in \{2, \dots, \chi - 1\}$

$$\begin{aligned}
ABS(G') - ABS(G) &= (n_1 + 1)(n_\chi - 1)f(n - n_1 - 1, n - n_\chi + 1) - n_1 n_\chi f(n - n_1, n - n_\chi) \\
&\quad + \sum_{i=2}^{\chi-1} [n_i(n_1 + 1)f(n - n_1 - 1, n - n_i) - n_1 n_i f(n - n_1, n - n_i)] \\
&\quad + \sum_{i=2}^{\chi-1} [n_i(n_\chi - 1)f(n - n_\chi + 1, n - n_i) - n_\chi n_i f(n - n_\chi, n - n_i)] \\
&= (n_\chi - n_1 - 1)f(n - n_1, n - n_\chi) \\
&\quad + \sum_{i=2}^{\chi-1} n_i [f(n - n_1 - 1, n - n_i) - f(n - n_\chi + 1, n - n_i)] \\
&\quad + \sum_{i=2}^{\chi-1} n_i [n_\chi g_1(n - n_\chi, n - n_i) - n_1 g_1(n - n_1 - 1, n - n_i)]
\end{aligned}$$

Since $n - n_1 - 1 \geq n - n_\chi + 1$, from Lemma 2 we get that for each $i = 2, \dots, \chi - 1$, $f(n - n_1 - 1, n - n_i) - f(n - n_\chi + 1, n - n_i) \geq 0$ and $n_\chi g_1(n - n_\chi, n - n_i) - n_1 g_1(n - n_1 - 1, n - n_i) > n_1 [g_1(n - n_\chi, n - n_i) - g_1(n - n_1 - 1, n - n_i)] \geq 0$. So $ABS(G') - ABS(G)$, a contradiction. Thus $n_\chi - n_1 \leq 1$. \square

The next theorem gives a sharp upper bound on the ABS value of all connected graphs of a fixed order and fixed independent number.

Theorem 2 *If G is a connected graph of order n and independent number α then*

$$ABS(G) \leq \alpha \sqrt{(n - \alpha)(n - \alpha - 1)} + \frac{1}{2}(n - \alpha) \sqrt{(n - \alpha - 1)(n - \alpha - 2)},$$

with equality holds if and only if $G \cong N_\alpha + K_{n-\alpha}$

Proof. Let G be a connected graph which has the maximum ABS value among all connected graphs of order n and independent number α . Let S be an independent set in G with $|S| = \alpha$. Assume that there is a vertex $u \in S$ that is not adjacent to a vertex $v \in V(G) - S$. Then $G + uv$ has order n and independent number α and $ABS(G + uv) > ABS(G)$, a contradiction. Thus, each vertex in S is adjacent to every vertex in $G - S$. Furthermore, every pair of vertices in $G - S$ are adjacent, yielding $G[V(G) - S] \cong K_{n-\alpha}$. Thus $G \cong N_\alpha + K_{n-\alpha}$. Therefore, $ABS(G) = \alpha \sqrt{(n - \alpha)(n - \alpha - 1)} + \frac{1}{2}(n - \alpha) \sqrt{(n - \alpha - 1)(n - \alpha - 2)}$ \square

The next theorem gives a sharp upper bound on the ABS value of all connected graphs of a fixed order and fixed number of pendant vertices. We denote by S_{n-1} the star graph of order n and by $S_{m,n-m}$ the double star graph of order n where the internal vertices have degrees m and $n-m$. We also denote by K_m^p the graph of order $m+p$ and p pendant vertices such that the induced subgraph on the internal vertices is complete graph and all pendant vertices are adjacent to the same internal vertex.

Theorem 3 *Let G be a graph of order n having p pendant vertices.*

1. *if $p = n - 1$ then $G \cong ABS(S_{n-1})$, and thus $ABS(G) = \frac{(n-1)\sqrt{n-2}}{n}$.*
2. *if $p = n - 2$ then $ABS(G) \leq \frac{1}{\sqrt{3}} + \frac{\sqrt{n-2}}{n} + \frac{(n-3)\sqrt{n-3}}{n-1}$ with equality holds if and only if $G \cong S_{2,n-2}$*
3. *if $p \leq n - 3$ then*

$$ABS(G) \leq p\sqrt{\frac{n-2}{n}} + (n-p-1)\sqrt{\frac{2n-2p-3}{2n-2p-1}} + \frac{1}{2}\sqrt{n-p-1}(n-p-2)^{\frac{3}{2}}$$

with equality holds if and only if $G \cong K_{n-p}^p$.

Proof. (1) Straightforward.

(2) Let u, v be the internal vertices of G . We may assume that there are t pendant vertices adjacent to u and $p-t$ pendant vertices adjacent to v . Thus

$$\begin{aligned} ABS(G) &= tf(1, d_u) + (p-t)f(1, d_v) + f(d_u, d_v) \\ &= tf(1, t+1) + (p-t)f(1, p-t+1) \end{aligned}$$

Consider the function $h(t) = tf(1, t+1) + (p-t)f(1, p-t+1)$.

$$h'(t) = \frac{M - N}{(t+2)(p-t+2)\sqrt{(t+2)(p-t+2)}},$$

where $M = ((p-1)t - t^2 + 3p + 6)\sqrt{pt - t^2 + 2t}$ and $N = (p-t+3)(t+2)\sqrt{(p-t)(t+2)}$. Clearly, both $M > 0$ and $N > 0$ when $1 \leq t \leq p-1$. Thus the sign of $h'(t)$ is determined by the sign of $(M - N)(M + N) = M^2 - N^2$. Now

$$M^2 - N^2 = (2t - p)(3tp(p-t) + 10t(p-t) + 8p^2 + 48p + 72).$$

Hence $h'(t) < 0$ when $1 \leq t < p/2$ and $h'(t) > 0$ when $p/2 < t \leq p-1$. Thus $h(t)$ has maximum value at $t = 1$ or $t = p-1$. Thus

$$ABS(G) \leq h(1) = h(p-1) = \frac{1}{\sqrt{3}} + \frac{\sqrt{p}}{p+2} + \frac{(p-1)\sqrt{p-1}}{p+1}$$

with equality holds if and only if $G \cong S_{2,p} = S_{2,n-2}$.

(3) Let P be the set of pendant vertices in G . If there are two non adjacent vertices $u, v \in V(G) \setminus P$ then $G + \{uv\} \in \Gamma_{n,p}$ and by Lemma 1, $ABS(G + \{uv\}) > ABS(G)$, a contradiction. Thus the induced subgraph $G[V(G) \setminus P]$ is K_{n-p} . Label the vertices of $G[V(G) \setminus P]$ by v_1, \dots, v_{n-p} , and for each $i = 1, \dots, n-p$, let $a_i = |N(v_i) \cap P|$ so that $a_1 \geq a_2 \geq \dots \geq a_{n-p}$. To obtain the desired result we want need to show that $a_1 = p$ and $a_2 = \dots = a_{n-p} = 0$. So seeking a contradiction assume that $a_i \geq 1$ for some $i \geq 2$. Then $a_1 \geq a_2 \geq 1$. Let $x \in P \cap N(v_2)$ and take $G' = G - \{xv_2\} + \{xv_1\}$. Note that for each $i = 1, \dots, n-p$, $deg_G(v_i) = a_i + n - p - 1$. Then

$$\begin{aligned}
ABS(G') - ABS(G) &= f(1, a_1 + n - p) - f(1, a_2 + n - p - 1) \\
&\quad + a_1(f(1, a_1 + n - p) - f(1, a_1 + n - p - 1)) \\
&\quad - (a_2 - 1)(f(1, a_2 + n - p - 1) - f(1, a_2 + n - p - 2)) \\
&\quad - \sum_{i=3}^{n-p} (f(a_i + n - p - 1, a_1 + n - p) - f(a_i + n - p - 1, a_1 + n - p - 1)) \\
&\quad - \sum_{i=3}^{n-p} (f(a_i + n - p - 1, a_2 + n - p - 1) - f(a_i + n - p - 1, a_2 + n - p - 2)) \\
&= f(1, a_1 + n - p) - f(1, a_2 + n - p - 1) \\
&\quad + a_1 g_1(1, a_1 + n - p - 1) - (a_2 - 1) g_1(1, a_2 + n - p - 2) \\
&\quad - \sum_{i=3}^{n-p} (g_1(a_1 + n - p - 1, a_i + n - p - 1) - g_1(a_2 + n - p - 2, a_i + n - p - 1))
\end{aligned}$$

Now for each $i = 3, \dots, n-p$, $a_i \geq 0$ and so by Lemma 3,

$$\begin{aligned}
g_1(a_1 + n - p - 1, a_i + n - p - 1) - g_1(a_2 + n - p - 2, a_i + n - p - 1) &\geq \\
g_1(a_1 + n - p - 1, n - p - 1) - g_1(a_2 + n - p - 2, n - p - 1). &
\end{aligned}$$

Thus

$$\begin{aligned}
ABS(G') - ABS(G) &\geq f(1, a_1 + n - p) - f(1, a_2 + n - p - 1) \\
&\quad + a_1 g_1(a_1 + n - p - 1, 1) - (a_2 - 1) g_1(a_2 + n - p - 2, 1) \\
&\quad + (n - p - 2)(g_1(a_1 + n - p - 1, n - p - 1) - g_1(a_2 + n - p - 2, n - p - 1)) \\
&= w(a_1) - w(a_2 - 1)
\end{aligned}$$

Where

$$w(t) = f(t + n - p, 1) + t g_1(t + n - p - 1, 1) + (n - p - 2) g_1(t + n - p - 1, n - p - 1)$$

Our next aim is to show that $w(t)$ is increasing in t . Note that

$$w'(t) = \frac{\partial f}{\partial t}(t+n-p, 1) + g_1(t+n-p-1, 1) + t \frac{\partial g_1}{\partial t}(t+n-p-1, 1) \\ + (n-p-2) \frac{\partial g_1}{\partial t}(t+n-p-1, n-p-1)$$

Since $\frac{\partial g_1}{\partial x}(x, y)$ is increasing in y when $y \geq 1$, we get

$$\frac{\partial g_1}{\partial t}(t+n-p-1, n-p-1) \geq \frac{\partial g_1}{\partial t}(t+n-p-1, 1),$$

So

$$w'(t) \geq \frac{\partial f}{\partial t}(t+n-p, 1) + g_1(t+n-p-1, 1) + (t+n-p-1) \frac{\partial g_1}{\partial t}(t+n-p-1, 1) \\ = L - K,$$

where

$$L = \frac{(t+n-p)^2 + (t+n-p) - 1}{(t+n-p-1)^{\frac{1}{2}}(t+n-p+1)^{\frac{3}{2}}} \text{ and } K = \frac{(t+n-p)^2 - (t+n-p) - 1}{(t+n-p-2)^{\frac{1}{2}}(t+n-p)^{\frac{3}{2}}}.$$

Since

$$L^2 - K^2 = \frac{2(t+n-p)^5 - 3(t+n-p)^4 - 8(t+n-p)^3 + 3(t+n-p)^2 + 4(t+n-p) + 1}{(t+n-p-1)(t+n-p+1)^3(t+n-p-2)(t+n-p)^3} > 0,$$

we get $w'(t) \geq L - K > 0$ and thus $w(t)$ is increasing in t as desired. This implies that $ABS(G') - ABS(G) > w(a_1) - w(a_2 - 1) > 0$, a contradiction. So $a_1 = p$ and $a_i = 0$ for all $i = 2, \dots, n-p-1$, and hence $G \cong K_{n-p}^p$.

□

References

- [1] J. A. Bondy, U. S. R. Murty, Graph Theory, Springer, London, 2008.
- [2] S. Wagner, H. Wang, Introduction to Chemical Graph Theory, CRC Press, Boca Raton, 2018.
- [3] I. Gutman, Degree based topological indices, Croat. Chem. Acta 86 (2013) 351–361.
- [4] M. Randić, On characterization of molecular branching, J. Am. Chem. Soc. 97 (1975) 6609–6615.

- [5] L. B. Kier, L. H. Hall, *Molecular Connectivity in Chemistry and Drug Research*, Academic Press, New York, 1976.
- [6] L. B. Kier, L. H. Hall, *Molecular Connectivity in Structure–Activity Analysis*, Wiley, New York, 1986.
- [7] M. Randić, M. Novič, D. Plavšić, *Solved and Unsolved Problems in Structural Chemistry*, CRC Press, Boca Raton, 2016.
- [8] B. Zhou, N. Trinajstić, On a novel connectivity index, *J. Math. Chem.* 46 (2009) 1252–1270.
- [9] E. Estrada, L. Torres, L. Rodríguez, I. Gutman, An atom-bond connectivity index: modelling the enthalpy of formation of alkanes, *Indian J. Chem. Sec. A* 37 (1998) 849–855.
- [10] A. Ali, L. Zhong, I. Gutman, Harmonic index and its generalization: extremal results and bounds, *MATCH Commun. Math. Comput. Chem.* 81 (2019) 249–311.
- [11] A. Ali, K. C. Das, D. Dimitrov, B. Furtula, Atom-bond connectivity index of graphs: a review over extremal results and bounds, *Discrete Math. Lett.* 5 (2021) 68–93.
- [12] A. Ali, B. Furtula, I. Redžepović, I. Gutman, Atom-bond sum-connectivity index, *J. Math. Chem.* 60 (2022) 2081–2093.
- [13] A. Ali, I. Gutman, I. Redžepović, Atom-bond sum-connectivity index of unicyclic graphs and some applications, *Electron. J. Math.* 5 (2023) 1–7.
- [14] T. A. Alraqad, I. Ž. Milovanović, H. Saber, A. Ali, J. P. Mazorodze, Minimum atom-bond sum-connectivity index of trees with a fixed order and/or number of pendent vertices, arXiv:2211.05218 [math.CO].
- [15] V. Maitreyi, S. Elumalai, S. Balachandran, The minimum ABS index of trees with given number of pendent vertices, arXiv:2211.05177 [math.CO].