

A structure theorem for homology 4-manifolds with $g_2 \leq 5$

Biplab Basak and Sourav Sarkar

Department of Mathematics, Indian Institute of Technology Delhi, New Delhi 110016, India.

February 5, 2023

Abstract

Numerous structural findings of homology manifolds have been derived in various ways in relation to g_2 -values. The homology 4-manifolds with $g_2 \leq 5$ are characterized combinatorially in this article. It is well-known that all homology 4-manifolds for $g_2 \leq 2$ are polytopal spheres. We demonstrate that homology 4-manifolds with $g_2 \leq 5$ are triangulated spheres and are derived from triangulated 4-spheres with $g_2 \leq 2$ by a series of connected sum, bistellar 1- and 2-moves, edge contraction, edge expansion, and edge flipping operations. The inequality is also quite pronounced for these combinatorial classes.

MSC 2020: Primary 57Q25; Secondary 05E45, 55U10, 57Q05, 57Q15.

Keywords: Homology 4-Manifold, g_2 -value, Edge contraction, Edge expansion.

1 Introduction

The g -conjecture [15] proposed a detailed characterization of f -vectors of simplicial polytopes as well as a search for potential f -vectors of other types of simplicial complexes. The components of the g -vector of any simplicial d -polytope are shown to be non-negative in [8, 9]. This naturally raises the issue of the general complexes' geometric arrangement in relation to the g -values. It is trivial that $g_1(\Delta) = 0$ holds if and only if Δ is a simplex.

Several classification results have been discovered based on the third component g_2 of the g -vector. The well-known Lower Bound Theorem (LBT), established by Barnette [4, 5], states that if Δ is the boundary complex of a simplicial $(d+1)$ -polytope, or more broadly a finite triangulation of a connected compact d -manifold without boundary, where $d \geq 2$, then $g_2(\Delta) \geq 0$. Any triangulation of a 3-manifold with $g_2 \leq 9$ is a triangulated 3-sphere, and it is stacked when $g_2 = 0$ [25]. Moreover, the inequality $g_2 \leq 9$ for triangulated 3-spheres is sharp, as there are triangulations of 3-dimensional handles (both orientable and non-orientable) with $g_2 = 10$. The inequality $g_2(\Delta) \geq 0$ has been established in [3, 11, 12] for the class of all normal pseudomanifolds of dimensions ≥ 3 . As a step toward equality on g_2 , for $d \geq 3$, the stacked d -spheres have been shown to be the only normal d -pseudomanifolds with $g_2 = 0$ (cf. [12, 23]). Some structural results on normal pseudomanifolds can be found in [2, 7].

The combinatorial characterization of homology d -manifolds with $g_2 \leq 2$ are known due to Nevo and Novinsky [16] for $g_2 = 1$ and due to Zheng [27] for $g_2 = 2$.

¹*E-mail addresses:* biplab@iitd.ac.in (B. Basak), Sourav.Sarkar@maths.iitd.ac.in (S. Sarkar).

Proposition 1.1. [16] *Let $d \geq 3$ and Δ be a prime homology d -sphere with $g_2(\Delta) = 1$. Then Δ is combinatorially isomorphic to either the join of the boundary complex of two simplices whose dimensions add up to $(d+1)$ or the join of a cycle and a boundary complex of a $(d-1)$ -simplex.*

Proposition 1.2. [27] *Let $d \geq 4$, and Δ be a prime homology d -manifold with $g_2 = 2$. Then Δ can be obtained from a polytopal d -sphere with $g_2 = 0$ or 1 , by centrally retriangulating along some stacked subcomplex.*

A homology manifold with $g_2 \leq 2$ is the connected sum of a finite number of prime homology manifolds with g_2 less than or equal to 2. Therefore, a homology d -manifold with $g_2 \leq 2$ is a triangulated d -sphere due to the following result.

Proposition 1.3. [18] *Let $d \geq 3$, and Δ be a d -dimensional homology manifold which is not prime. Then Δ is either a connected sum of homology manifolds or the result of a handle addition on a homology manifold.*

In this article, we aim to find out an upper bound, say b , on g_2 such that if Δ is a homology 4-manifold with $g_2(\Delta) \leq b$, then Δ is a triangulated 4-sphere. Moreover, we shall provide a combinatorial structure for such triangulated spheres. Our general approaches are based on the rigidity theory of graphs and some basic combinatorial operations on complexes.

Theorem 1.4. *If Δ is a prime homology 4-manifold with $g_2 \leq 5$, then Δ is a triangulated sphere, which is obtained from a triangulated 4-sphere with $g_2 \leq 2$ by a sequence of the following operations:*

- (i) *a bistellar 1-move and an edge contraction,*
- (ii) *an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Moreover, the inequality is sharp for triangulated 4-spheres.

Remark 1.5. If Δ is a prime homology 4-manifold with $g_2(\Delta) \leq 2$ then Δ is a triangulated sphere (cf. Propositions 1.1 and 1.2). Therefore, Theorem 1.4 implies that all the prime homology 4-manifolds with $g_2 \leq 5$ are triangulated spheres. Let K be a homology 4-manifold (which may or may not be prime) with $g_2(K) \leq 5$. Since the application of handle addition to a 4-dimensional manifold increases the g_2 value by 15 in the resulting complex, K is a connected sum of a finite number of prime homology 4-manifolds with g_2 less than or equal to 5. Therefore, K is also a triangulated 4-sphere.

Remark 1.6. In [10] Brehm and Kühnel proved that the 9-vertex combinatorial triangulation of the complex projective space $\mathbb{C}\mathbb{P}_9^2$ has the 1-skeleton as a complete graph. As a result, $g_2(\mathbb{C}\mathbb{P}_9^2) = 6$. Thus, the upper bound for $g_2(\Delta)$ for which the homology 4-manifold Δ is a triangulated 4-sphere is sharp.

2 Preliminaries

All the simplices taken in this article are geometric, and the simplicial complexes taken are finite. We say a simplicial complex Δ is pure if all the maximal simplices of Δ are of the

same dimension. Any simplex present in a simplicial complex Δ is called a face of Δ , and the maximal faces of Δ are called facets. The set of all vertices (or 0-simplices) of Δ is denoted by $V(\Delta)$, and the collection of all vertices and edges of Δ is called the graph of Δ and is denoted by $G(\Delta)$. For any subset $V' \subseteq V(\Delta)$, we shall take $\Delta[V'] := \{\sigma \in \Delta : V(\sigma) \subseteq V'\}$. A simplex σ is called a missing simplex of Δ if $\partial\sigma \subseteq \Delta$ but $\sigma \notin \Delta$. A pure simplicial complex is called prime if it has no missing facets.

The join $\sigma\tau$ (or, $\sigma\star\tau$) of two simplices σ and τ is the simplex obtained by the convex hull of vertices in $V(\sigma) \cup V(\tau)$. The simplicial complexes Δ_1 and Δ_2 are said to be independent if $\sigma\tau$ is a simplex of dimension $(i+j+1)$, for every pair of simplices $\sigma \in \Delta_1$ and $\tau \in \Delta_2$ of dimensions i and j , respectively. The join of two independent simplicial complexes Δ_1 and Δ_2 is the simplicial complex $\Delta_1 \star \Delta_2 := \Delta_1 \cup \Delta_2 \cup \{\sigma\tau : \sigma \in \Delta_1, \tau \in \Delta_2\}$. For any pair (σ, Δ) , where σ is a simplex and Δ is a simplicial complex, by $\sigma \star \Delta$ we mean the simplicial complex $\{\alpha : \alpha \leq \sigma\} \star \Delta$. The link of any face σ in Δ is defined as the collection $\{\gamma \in \Delta : \gamma \cap \sigma = \emptyset \text{ and } \gamma\sigma \in \Delta\}$ and is denoted by $lk_\Delta\sigma$, and the star of σ is defined as $\{\alpha : \alpha \leq \sigma\beta, \beta \in lk_\Delta\sigma\}$ and is denoted by $st_\Delta\sigma$. The number of vertices in $lk_\Delta u$ is called the degree of that vertex and is denoted by $d(u, \Delta)$ or simply $d(u)$.

One of the most common enumerative tool of any d -dimensional simplicial complex Δ is its f -vector $\{f_0, f_1, \dots, f_d\}$, where f_i is the number of i dimensional simplices present in Δ . We also define the h -vector of Δ as (h_0, h_1, \dots, h_d) , where

$$h_i = \sum_{j=0}^i (-1)^{i-j} \binom{d+1-j}{i-j} f_{j-1},$$

and we define $g_i := h_i - h_{i-1}$. In particular, $g_2 = f_1 - (d+1)f_0 + \binom{d+2}{2}$ and $g_3 = f_2 - df_1 + \binom{d+1}{2}f_0 - \binom{d+2}{3}$. In [20], a relation between g_i 's of all pure simplicial complexes and their vertex links is given, which was first stated by McMullen in [14] for shellable complexes.

Lemma 2.1. [20] *If Δ is a d -dimensional pure simplicial complex and k is a positive integer, then*

$$\sum_{v \in \Delta} g_k(lk_\Delta v) = (k+1)g_{k+1}(\Delta) + (d+2-k)g_k(\Delta).$$

Let Δ be a pure simplicial complex of dimension d . For a fixed field \mathbb{F} , we say that Δ is a d -dimensional \mathbb{F} -homology sphere if $\tilde{H}_i(lk_\Delta\sigma, \mathbb{F}) \cong \tilde{H}_i(\mathbb{S}^{d-|\sigma|}, \mathbb{F})$, for every simplex σ of dimension $i \geq -1$. Similarly, Δ is a d -dimensional \mathbb{F} -homology manifold if the link of every vertex in Δ is a $(d-1)$ -dimensional \mathbb{F} -homology sphere. A strongly connected and pure simplicial complex Δ of dimension d is called a normal d -pseudomanifold if every $(d-1)$ -face of Δ is contained in exactly two facets in Δ and the links of all the simplices of dimension $\leq (d-2)$ are connected. The following result is easy to verify.

Lemma 2.2. *Let $d \geq 4$ and Δ be a prime normal d -pseudomanifold such that $g_2(\Delta) \geq 1$. If u is a vertex in Δ such that $lk_\Delta u$ is prime, then $g_2(lk_\Delta u) \geq 1$.*

Given two positive integers a and i , there is a unique way to write

$$a = \binom{a_i}{i} + \binom{a_{i-1}}{i-1} + \dots + \binom{a_j}{j},$$

where $a_i > a_{i-1} > \dots > a_j \geq j \geq 1$. Then $a^{<i>}$, the i 'th Macaulay pseudo-power of a is defined as

$$a^{<i>} := \binom{a_i+1}{i+1} + \binom{a_{i-1}+1}{i} + \dots + \binom{a_j+1}{j+1}.$$

Lemma 2.3. [22] *Let Δ be a normal d -pseudomanifold, where $d \geq 3$. Then $g_3(\Delta) \leq g_2(\Delta)^{\langle 2 \rangle}$.*

Lemma 2.4. *Let Δ be a homology 4-manifold with $1 \leq g_2(\Delta) \leq 5$. Then $g_3(\Delta) = 0$.*

Proof. Recall that $g_3(\Delta) = h_3 - h_2$. Since Δ is a homology 4-manifold, it follows from [13, 17] that $h_{5-i}(\Delta) - h_i(\Delta) = (-1)^i \binom{5}{i} (\chi(\Delta) - \chi(\mathbb{S}^4))$. In particular, for $i = 2$, $h_3(\Delta) - h_2(\Delta) = 10(\chi(\Delta) - \chi(\mathbb{S}^4))$, i.e., $g_3(\Delta)$ is an integer multiple of 10. Since $g_2(\Delta) \leq 5$, it follows from Lemma 2.3 that $g_3(\Delta) \leq 7$. Therefore, by combining these two we have $g_3(\Delta) \leq 0$. On the other hand, Lemma 2.1 implies,

$$\sum_{u \in \Delta} g_2(\text{lk}_\Delta u) = 3g_3(\Delta) + 4g_2(\Delta). \quad (1)$$

If $g_3(\Delta) \leq -10$, then it follows from Equation (1) that, $\sum_{u \in \Delta} g_2(\text{lk}_\Delta u) \leq -30 + 20 = -10$, which is not possible. Therefore $g_3(\Delta) = 0$. \square

In [16, 27], the characterizations of all the homology d -manifolds with $1 \leq g_2 \leq 2$ are given. Topologically, these are polytopal spheres. The combinatorial structure of the homology 3-manifolds with $g_2 \leq 2$ are given by the following results.

Proposition 2.5. [16] *Let Δ be a prime homology 3-manifold with $g_2 = 1$. Then Δ is combinatorially isomorphic to $C \star \partial\sigma^2$, where C is an n -cycle, $n \geq 3$ and σ^2 is a 2-simplex.*

Proposition 2.6. [27] *Let Δ be a prime homology 3-manifold with $g_2 = 2$. Then Δ is one of the following:*

- (i) Δ is obtained from a triangulated 3-sphere with $g_2 = 1$ by stellar subdivision at a ridge,
- (ii) Δ is an octahedral 3-sphere, i.e., $\Delta = \partial(a_1b_1) \star \partial(a_2b_2) \star \partial(a_3b_3) \star \partial(a_4b_4)$.

2.1 Rigidity of graphs and Pseudomanifolds

Let G be a graph, where $V = \{v_1, \dots, v_n\}$ denote the vertex set. A d -embedding of G is a map $f : V \rightarrow \mathbb{R}^d$. A framework of G is a pair (G, f) , where f is a d -embedding of G for some $d \geq 1$. In this context, we always use the Euclidean norm.

Definition 2.7. *A d -embedding f of a graph $G = (V, E)$ is called **rigid** if there is an $\epsilon > 0$ such that for any d -embedding g of G satisfying (i) $\|f(v) - g(v)\| < \epsilon$ for every vertex $v \in V$, and (ii) $\|f(v) - f(u)\| = \|g(v) - g(u)\|$ for every edge $vu \in E$, one has $\|f(v) - f(u)\| = \|g(v) - g(u)\|$ for any pair of vertices v, u in V .*

An embedding f that is not rigid, is called **flexible**. If a d -embedding is rigid, then the corresponding framework is called a rigid framework. We denote the set of all d -embeddings of a graph G by $\mathcal{E}(G)$, and the set of all rigid d -embeddings of a graph G by $R\mathcal{E}(G)$. Note that $\mathcal{E}(G)$ is a topological vector space of dimension $d \times |V|$ and $R\mathcal{E}(G)$ is a subspace of $\mathcal{E}(G)$.

Definition 2.8. *A graph G is called generically d -rigid if the set of all rigid d -embeddings $R\mathcal{E}(G)$ is an **open** and **dense** subset of the set of all d -embeddings $\mathcal{E}(G)$.*

Proposition 2.9. [1, 16, Gluing Lemma] *Let G_1 and G_2 be two generically d -rigid graphs such that $G_1 \cap G_2$ contains at least d -vertices. Then $G_1 \cup G_2$ is a generically d -rigid.*

Lemma 2.10. *Let G be a generically d -rigid graph, and H_1 and H_2 be two generically d -rigid sub-graphs of G . Then there is a rigid d -embedding $\Psi : V(G) \rightarrow \mathbb{R}^d$ whose restrictions on $V(H_1)$ and $V(H_2)$ are rigid d -embeddings of H_1 and H_2 , respectively.*

Proof. Let $V(G) = \{v_1, \dots, v_n\}$ be the set of vertices of G . Since G is a generically d -rigid, the set of all rigid d -embeddings $R\mathcal{E}(G)$ is an open and dense subset of the set of all d -embeddings $\mathcal{E}(G)$. Let $f \in R\mathcal{E}(G)$. Note that $\mathcal{E}(G)$ is isomorphic to the Euclidean space $\mathbb{R}^{d \times |V|}$. Since $R\mathcal{E}(G)$ is an open subset of $\mathcal{E}(G)$, there is an open set $U = U_1 \times \dots \times U_{|V|}$ where each $U_i \subseteq \mathbb{R}^d$ (i corresponds to the vertex v_i) such that $f \in U \subseteq R\mathcal{E}(G) \subseteq \mathcal{E}(G)$. Without loss of generality, let $U_1, \dots, U_{|H_1|}$ correspond to the vertices of H_1 . Let f_{H_1} be the restriction of f on $V(H_1)$. Then f_{H_1} is a d -embedding of H_1 , and further, $f_{H_1} \in W = U_1 \times U_2 \times \dots \times U_{|H_1|}$. Since H_1 is a generically d -rigid, $R\mathcal{E}(H_1)$ is an open and dense subset of the set of all d -embeddings $\mathcal{E}(H_1)$. Since W is an open subset of $\mathcal{E}(H_1)$, $R\mathcal{E}(H_1) \cap W \neq \emptyset$. Let $\phi \in R\mathcal{E}(H_1) \cap W$. Then ϕ is a rigid d -embedding of H_1 . Define a map $\Phi : V(G) \rightarrow \mathbb{R}^d$ in the following way,

$$\begin{aligned} \Phi(v_i) &= \phi(v_i) & \text{if } 1 \leq i \leq |H_1|, \\ &= f(v_i) & \text{elsewhere.} \end{aligned}$$

Then $\Phi \in U$, and therefore, Φ is a generically d -rigid embedding of G such that the restriction of Φ on $V(H_1)$ is a rigid d -embedding of H_1 .

Let $Z \subseteq (R\mathcal{E}(H_1) \cap W) \times U_{|H_1|+1} \times \dots \times U_{|V|}$ be a basic open set in $\mathbb{R}^{d \times |V|}$ containing Φ , where $Z = Z_1 \times \dots \times Z_{|V|}$. Then any element h in Z is a rigid d -embedding of G , whose restriction on $V(H_1)$ is a rigid d -embedding of H_1 . Let Z_{i_1}, \dots, Z_{i_k} corresponds the vertices of H_2 and take $W_1 = Z_{i_1} \times \dots \times Z_{i_k}$. Then the restriction of Φ on W_1 is a d -embedding of H_2 . Since H_2 is generically d -rigid and W_1 is an open set containing a d -embedding of H_2 , we have $R\mathcal{E}(H_2) \cap W_1 \neq \emptyset$. Let $\psi \in R\mathcal{E}(H_2) \cap W_1$, i.e., ψ is a rigid d -embedding of H_2 . Now define a map $\Psi : V(G) \rightarrow \mathbb{R}^d$ in the following way,

$$\begin{aligned} \Psi(v_i) &= \psi(v_i) & \text{if } i_1 \leq i \leq i_k, \\ &= \Phi(v_i) & \text{elsewhere.} \end{aligned}$$

The map Ψ is an element of Z , and the restriction of Ψ on $V(H_2)$ is a rigid d -embedding of H_2 . This completes the proof. \square

Corollary 2.11. *Let G be a generically d -rigid graph and H_1, \dots, H_k be generically d -rigid subgraphs of G . Then there is a rigid d -embedding Ψ of G such that the restriction of Ψ on $V(H_i)$ is a rigid d -embedding of H_i , for $1 \leq i \leq k$.*

For a d -embedding f of a graph G , a stress with respect to f is a function $\omega : E \rightarrow \mathbb{R}$ such that for every vertex $v \in V$,

$$\sum_{uv \in E} \omega(uv)[f(v) - f(u)] = 0.$$

The collection of all stresses corresponding to a given d -embedding f forms a real vector space called the stress space of f . Let us denote the stress space of Δ corresponding to f

by $\mathcal{S}(\Delta_f)$. This stress space can be described in terms of matrices as well. Let $\text{card}(V) = n$ and $\text{card}(E) = m$. Let $\text{Rig}(G, f)$ be the $dn \times m$ matrix whose rows are labeled by edges and the columns are grouped into blocks of size d labeled by vertices. In the rows corresponding to the edge $e = uv$, the d -vector occupied in the block column corresponding to the vertex u (resp. v) will be $f(u) - f(v)$ (resp. $f(v) - f(u)$). The rest of the columns in that row will be occupied by the zero vector. The matrix $\text{Rig}(G, f)$ is called the Rigidity matrix of the d -embedding f associated with the graph G . It is easy to see that the stress space of a d -embedding f of G is same as left kernel of the Rigidity matrix $\text{Rig}(G, f)$. If G is a generically d -rigid graph, then the rigidity matrix is independent of the choice of d -embedding [12], and in this case, we write the rigidity matrix of a graph G as $\text{Rig}(G, d)$ and the corresponding stress space as $\mathcal{S}(\Delta)$.

Proposition 2.12. [1, 12] *Let $G = (V, E)$ be a generically $(d + 1)$ -rigid graph. Then $g_2(G)$ is the dimension of the left kernel of the rigidity matrix $\text{Rig}(G, d)$.*

Proposition 2.13. [24, 26, Cone Lemma] *Let G be a given graph and $C(G, u)$ be the cone over G with a new vertex u . Then,*

- (i) *G is generically d -rigid if and only if $C(G, u)$ is generically $(d + 1)$ -rigid.*
- (ii) *The left null spaces of $\text{Rig}(G, d)$ and $\text{Rig}(C(G, u), d + 1)$ are isomorphic real vector spaces. Moreover, if $g_2(G) \neq 0$ then u participates in a generic stress of $C(G, u)$.*

A simplicial complex Δ is called generically d -rigid if the graph (or 1-skeleton) $G(\Delta)$ is a generically d -rigid. The concept of rigidity have been used in the structural study of simplicial complex more precisely in normal pseudomanifolds.

Proposition 2.14. [11, 12] *Let $d \geq 3$ and Δ be a normal d -pseudomanifold. Then,*

- (i) *Δ is generically $(d + 1)$ -rigid, and*
- (ii) *for every face σ of co-dimension 3 or more, $g_2(\text{lk}_\Delta \sigma) \leq g_2(\Delta)$.*

Proposition 2.15. *Let Δ be a normal d -pseudomanifold with $g_2(\Delta) = p$ and that $g_2(\text{lk}_\Delta v) \geq 1$, for every vertex $v \in \Delta$. Let u be a vertex such that $g_2(\Delta[V(\text{st}_\Delta u)]) = (p - 1)$. Then the graph $G(\Delta[V(\Delta \setminus \text{st}_\Delta u)])$ is complete.*

Proof. If possible, let v and z be two vertices in $V(\Delta \setminus \text{st}_\Delta u)$ such that $vz \notin \Delta$. Note that $\Delta[V(\text{st}_\Delta u)]$, $\text{st}_\Delta v$, and $\text{st}_\Delta z$ are generically $(d + 1)$ -rigid. It follows from Corollary 2.11 that there is a rigid $(d + 1)$ -embedding of Δ such that the restriction of the map on $V(\text{st}_\Delta u)$, $V(\text{st}_\Delta v)$ and $V(\text{st}_\Delta z)$ are rigid $(d + 1)$ -embeddings of $\Delta[V(\text{st}_\Delta u)]$, $\text{st}_\Delta v$ and $\text{st}_\Delta z$, respectively. Let f be such a rigid $(d + 1)$ embedding.

Since $g_2(\Delta[V(\text{st}_\Delta u)]) = p - 1$, the stress space $\mathcal{S}(\Delta[V(\text{st}_\Delta u)]_f)$ has dimension $(p - 1)$. Let $\{w_1, \dots, w_{p-1}\}$ be a basis for $\mathcal{S}(\Delta[V(\text{st}_\Delta u)]_f)$, where $w_i : E(\Delta[V(\text{st}_\Delta u)]) \rightarrow \mathbb{R}$ are stresses. Let us extend this stresses w_1, \dots, w_{p-1} to $\tilde{w}_1, \dots, \tilde{w}_{p-1}$, respectively on $E(\Delta)$ by taking $\tilde{w}_i(e) = 0$ for all $e \in E(\Delta) \setminus E(\Delta[V(\text{st}_\Delta u)])$. Note that $\tilde{w}_1, \dots, \tilde{w}_{p-1}$ are linearly independent in $\mathcal{S}(\Delta_f)$.

Since $g_2(\text{lk}_\Delta v) \geq 1$, by Cone Lemma, there is a non zero stress in $\mathcal{S}(\text{st}_\Delta v_f)$ whose support contains an edge incident to v . Let $w_p : E(\text{st}_\Delta v) \rightarrow \mathbb{R}$ be such a non zero stress. Now extend this stress w_p to a stress $\tilde{w}_p : E(\Delta) \rightarrow \mathbb{R}$ by taking $\tilde{w}_p(e) = 0$ for every edge $e \in E(\Delta) \setminus E(\text{st}_\Delta v)$. Then $\tilde{w}_1, \dots, \tilde{w}_{p-1}$ and \tilde{w}_p are independent in $\mathcal{S}(\Delta_f)$ and hence a basis of $\mathcal{S}(\Delta_f)$.

By the same arguments as in the previous paragraph, there is a non-zero stress $w_{p+1} : E(st_\Delta z) \rightarrow \mathbb{R}$ whose support is incident to z . Since $v \notin st_\Delta z$, the support of the stress is not included in v . Let us extend this stress w_{p+1} to $\tilde{w}_{p+1} : E(\Delta) \rightarrow \mathbb{R}$ by taking $\tilde{w}_{p+1}(e) = 0$ for every edge $e \in E(\Delta) \setminus E(st_\Delta z)$. Then \tilde{w}_{p+1} can not be written as a linear combination of $\tilde{w}_1, \dots, \tilde{w}_p$ as $\tilde{w}_i(e) = 0$ for every edge e incident to z . Therefore, $\tilde{w}_1, \dots, \tilde{w}_p$ and \tilde{w}_{p+1} are independent in $\mathcal{S}(\Delta_f)$, which contradicts the fact that $g_2(\Delta) = p$. Thus, $vz \in \Delta$ for any two vertices $v, z \in \Delta \setminus st_\Delta u$, i.e., $G(\Delta[V(\Delta \setminus st_\Delta u)])$ is complete. \square

Corollary 2.16. *Let Δ be a normal d -pseudomanifold such that $g_2(lk_\Delta v) \geq 1$, for every vertex $v \in \Delta$. If u is a vertex in Δ with $g_2(\Delta) = g_2(\Delta[V(st_\Delta u)])$, then $V(\Delta) = V(st_\Delta u)$.*

2.2 Combinatorial operations

Definition 2.17. Let Δ be a 4-dimensional normal pseudomanifold. Let ab be an edge in Δ such that $lk_\Delta ab = \partial(cd) \star \partial(xyz)$, where cd is a missing edge in Δ . Let $\Delta' := (\Delta \setminus ab \star \partial(cd) \star \partial(xyz)) \cup (\partial(ab) \star cd \star \partial(xyz))$. Then we say that Δ' is obtained from Δ by flipping the edge ab by cd . Note that $|\Delta|$ and $|\Delta'|$ are PL-homeomorphic and $g_2(\Delta) = g_2(\Delta')$.

Definition 2.18. Let Δ be a 4-dimensional normal pseudomanifold and σ be a $(4 - i)$ -simplex in Δ , $1 \leq i \leq 3$. Let $lk_\Delta \sigma = \partial\tau$, where τ is an i -simplex and $\tau \notin \Delta$. Let $\Delta' = (\Delta \setminus \sigma \star \partial\tau) \cup (\tau \star \partial\sigma)$. The complex Δ' is said to be obtained from Δ by a *bistellar i -move* with respect to the pair (σ, τ) . In general shall write this operation as *bistellar i -move*.

Definition 2.19. Let Δ be a normal 4-pseudomanifold and w be a vertex in Δ . Let S be a 2-dimensional sphere in $lk_\Delta w$ such that S separates $lk_\Delta w$ into two portions D_1 and D_2 , where D_1 is a triangulated 3-ball, and D_1 and D_2 have a common boundary complex S . Let $\Delta' := (\Delta \setminus st_\Delta w) \cup (uv \star S) \cup (u \star D_1) \cup (v \star D_2)$. Then $lk_{\Delta'} u \cap lk_{\Delta'} v = lk_{\Delta'} uv$ and Δ is obtained from Δ' by contracting the edge uv . The normal pseudomanifold Δ' is said to be *obtained from Δ by an edge expansion*.

In particular, if there is some vertex $x \in lk_\Delta(w)$ such that $lk_\Delta(xw)$ is the $(d - 2)$ -dimensional sphere S , then we say Δ' is obtained from Δ by *central retriangulation* of $st_\Delta(xw)$ with center u (or by stellar subdivision at the edge xw with the vertex u , see [21] for more details) and renaming w by v . Thus, the central retriangulation of the star of an edge (or the stellar subdivision at an edge) is a special case of an edge expansion.

Given two vertices u and v in a normal 4-pseudomanifold K such that either $lk(u, K)$ or $lk(v, K)$ is a triangulated sphere, we say the pair (u, v) satisfies the **link condition** if uv is an edge in K and $lk(u, K) \cap lk(v, K) = lk(uv, K)$ holds. In such cases, we can contract the edge uv . If K' is the resulting complex, then $|K'|$ is PL-homeomorphic to $|K|$ (cf. [6]).

Lemma 2.20. *Let Δ be a homology 4-manifold with $g_2(\Delta) = m$. Let u be a vertex in Δ such that $lk_\Delta u = C \star \partial(pqc)$, where C is an n -cycle. If $pqc \notin \Delta$, then Δ is obtained from a homology 4-manifold with $g_2 \leq m - (n - 2)$ by an edge expansion.*

Proof. Here $lk_\Delta u \cap lk_\Delta p = lk_\Delta up = C \star \partial(qc)$. Therefore, we can contract the edge up . Let Δ' be the resulting complex. Then $f_0(\Delta') = f_0(\Delta) - 1$, $f_1(\Delta') = f_1(\Delta) - (n + 3)$. Thus, $g_2(\Delta') = f_1(\Delta) - (n + 3) - 5(f_0(\Delta) - 1) + 15 = g_2(\Delta) - (n - 2)$. \square

Lemma 2.21. *Let Δ be a homology 4-manifold with $g_2(\Delta) = m$. Let u be a vertex in Δ such that $lk_\Delta u = C \star \partial(pqc)$, where C is an n -cycle, $n \geq 4$. If at least one vertex of C*

is incident to no diagonal edge of C , then Δ is obtained from a homology 4-manifold with $g_2 \leq m - 1$ by an edge expansion.

Proof. Let $x \in C$ be a vertex that is free from all the diagonal edges of C . Then $lk_\Delta u \cap lk_\Delta x = lk_\Delta ux = \partial(pqc) \star \partial(x_1x_2)$, where x_1, x_2 are two vertices of C adjacent to x in C . Let Δ' be the complex obtained by contracting the edge ux . Then $g_2(\Delta') = m - 1$, and Δ is obtained from Δ' by an edge expansion. \square

In view of Lemmas 2.20 and 2.21, from now onwards, we assume that if there is a vertex u in a homology 4-manifold Δ with $lk_\Delta u = C \star \partial(pqc)$, then Δ satisfies the following property:

$$(*) \quad \begin{cases} pqc \in \Delta, \\ \text{every vertex of } C \text{ is incident to at least one diagonal edge.} \end{cases}$$

The result in Proposition 1.3 implies that any homology d -manifold is obtained from prime homology d -manifolds by a sequence of operations of the form connected sum and handle addition. So characterizing any homology manifold is equivalent to characterizing the prime homology manifolds. Let Δ be a prime homology d -manifold and u be a vertex in Δ such that $lk_\Delta u$ is not prime. Then Δ is obtained by a bistellar 1-move and edge contraction due to the following result.

Lemma 2.22. *Let $d \geq 4$ and Δ be a prime homology d -manifold. Let u be a vertex in Δ such that $lk_\Delta u$ is not prime. Then Δ is obtained from a homology d -manifold Δ' with $g_2(\Delta') = g_2(\Delta) - 1$ by the application of a bistellar 1-move and an edge contraction.*

Proof. Let u be a vertex in Δ such that $lk_\Delta u$ is not prime. Then $lk_\Delta u = S_1 \#_\sigma S_2$, where S_1 and S_2 are homology $d - 1$ spheres and σ is a missing $(d - 1)$ simplex in $lk_\Delta u$. Since Δ is prime, $\sigma \notin \Delta$. Let $\Delta' = (\Delta \setminus \{\tau : u \leq \tau\}) \cup x_1 \star (S_1 \cup \{\sigma\}) \cup x_2 \star (S_2 \cup \{\sigma\})$, where x_1 and x_2 are two new vertices. Then $g_2(\Delta') = g_2(\Delta) - 1$, and Δ is obtained from Δ' by a bistellar 1-move and edge contraction. \square

The operation used in Lemma 2.22, i.e., replacing $\{\tau : u \leq \tau\}$ with $x_1 \star (S_1 \cup \{\sigma\}) \cup x_2 \star (S_2 \cup \{\sigma\})$ in Δ was defined in [19, 25]. We used the reverse of that operation here, i.e., a combination of bistellar 1-move and edge contraction.

Definition 2.23. *For $\alpha \in \mathbb{N}$, the class \mathcal{H}_α consists of all homology 4-manifolds Δ such that $g_2(\Delta) = \alpha$ and*

- (i) *if Δ contains the boundary complex $\partial\sigma^4$ of a 4-simplex σ^4 , then $\sigma^4 \in \Delta$, i.e., Δ is prime, and*
- (ii) *$lk_\Delta v$ is prime for each vertex $v \in \Delta$.*

In view of Lemma 2.22, to characterize all 4-dimensional homology manifolds, it is enough to study the class \mathcal{H}_α . Let $\Delta \in \mathcal{H}_\alpha$. It follows from Lemma 2.2 that $g_2(lk_\Delta v) \geq 1$, for any vertex $v \in \Delta$. In the remaining we shall focus on the classes \mathcal{H}_α , where $3 \leq \alpha \leq 5$.

Lemma 2.24. *Let $\Delta \in \mathcal{H}_\alpha$, $1 \leq \alpha \leq 5$. Then there is a vertex $u \in \Delta$ with $g_2(lk_\Delta u) \leq 2$.*

Proof. If possible, let $g_2(lk_\Delta v) \geq 3$ for every vertex $v \in \Delta$. Then by Lemmas 2.1 and 2.4, $3f_0(\Delta) \leq 0 + 4.5 = 20$, i.e., $f_0(\Delta) \leq 6$. Since Δ is a homology 4-manifold, Δ is the stacked sphere. This contradicts the fact that $g_2(\Delta) \geq 1$. Thus, the result follows. \square

Lemma 2.25. *Let $\Delta \in \mathcal{H}_\alpha$, $4 \leq \alpha \leq 5$. Let u be a vertex such that $lk_\Delta u$ is an octahedral 3-sphere. Then Δ is obtained by an edge expansion (central retriangulation) from a homology 4-manifold with $g_2 \leq 3$.*

Proof. Since $lk_\Delta u$ is an octahedral 3-sphere, $lk_\Delta u$ is prime, and $g_2(lk_\Delta u) = 2$. Since $4 \leq g_2(\Delta) \leq 5$, there are at least two antipodal vertices in $lk_\Delta u$ that are not adjacent in Δ . Let x and y be two vertices (antipodal in $lk_\Delta u$) such that $xy \notin G(\Delta)$. Now consider the octahedral 2-sphere $S = \Delta[V(lk_\Delta u) \setminus \{x, y\}]$ and take the 4-dimensional combinatorial ball $B = S \star (xy)$. Then $\partial B = lk_\Delta u$. Let Δ' be the homology 4-manifold obtained from Δ by replacing $st_\Delta u$ with B . Then $g_2(\Delta') = g_2(\Delta) - 2$ and Δ is obtained from Δ' by a central retriangulation of $st_{\Delta'} xy$. \square

3 4-dimensional Homology Manifolds with $g_2 = 3$

Lemma 3.1. *Let Δ be a homology 4-manifold with $g_2 = 3$. Then there is a vertex u in Δ with $g_2(lk_\Delta u) \leq 1$.*

Proof. It follows from Lemma 2.4 that $g_3(\Delta) = 0$. Suppose $g_2(lk_\Delta v) \geq 2$ for every vertex $v \in \Delta$. Then Lemma 2.1 implies $2f_0(\Delta) \leq 3.4 = 12$, i.e., $f_0(\Delta) \leq 6$. This contradicts the fact that $g_2(\Delta) = 3$. Thus, there exists a vertex $u \in \Delta$ with $g_2(lk_\Delta u) \leq 1$. \square

Lemma 3.2. *Let $\Delta \in \mathcal{H}_3$. If u is a vertex in Δ with $g_2(lk_\Delta u) = 1$ and $d(u) \geq 7$, then Δ does not satisfy (*).*

Proof. Let $lk_\Delta u = C \star \partial(pqc)$, where C is a cycle of length $n \geq 4$. Since $g_2(\Delta) = 3$, $lk_\Delta u$ contains at most two missing edges that are present in Δ . Suppose Δ satisfies (*). Since every vertex of C is incident to at least one diagonal edge, we have $n = 4$ and $lk_\Delta u$ contains exactly two missing edges that are present in Δ . Therefore $V(\Delta) = V(st_\Delta u)$. But $pqc \in \Delta$ implies $lk_\Delta pqc$ contains at least three vertices from C . Let $a \in C \cap lk_\Delta pqc$. Then $\partial(upqc) \subseteq lk_\Delta a$ but $upqc \notin \Delta$. This contradicts the fact that $lk_\Delta a$ is prime. Therefore, Δ does not satisfy (*). \square

Lemma 3.3. *Let $\Delta \in \mathcal{H}_3$. If u is a vertex in Δ with $g_2(lk_\Delta u) = 1$ and $d(u) = 6$, then Δ is obtained by an edge expansion from a triangulated sphere with $g_2 \leq 2$.*

Proof. Let $lk_\Delta u = \partial(abt) \star \partial(pqc)$. If at least one of abt and pqc is a missing triangle in Δ , then the result follows from Lemma 2.20. Suppose both of $abt, pqc \in \Delta$. By similar arguments as in Lemma 3.2, $\{a, b, t\} \cap V(lk_\Delta pqc) = \emptyset$. Therefore, $V(lk_\Delta pqc) \subseteq V(\Delta \setminus st_\Delta u)$. Similarly, $V(lk_\Delta abt) \subseteq V(\Delta \setminus st_\Delta u)$. Hence $d(x) \geq 8$, for every vertex x in $lk_\Delta u$.

Suppose $g_2(lk_\Delta x) \geq 2$ for every vertex $x \in lk_\Delta u$. Then the equations $\sum_{v \in \Delta} g_2(lk_\Delta v) = 3g_3(\Delta) + 4g_2(\Delta)$ and $g_3(\Delta) = 0$ implies $6 \times 2 + 1 < 0 + 3.4$, which is not possible. Thus, there is at least one vertex, say y , in $lk_\Delta u$ such that $d(y) \geq 8$ and $g_2(lk_\Delta y) = 1$. Now, the result follows from Lemmas 2.20, 2.21, and 3.2. \square

Theorem 3.4. *Let Δ be a prime homology 4-manifold with $g_2 = 3$. Then Δ is obtained from a triangulated sphere with $g_2 \leq 2$ by either an edge expansion or a bistellar 1-move and edge contraction.*

Proof. It follows from Lemma 3.1 that there is a vertex v in Δ with $g_2(lk_\Delta v) \leq 1$. Let u be a vertex in Δ such that $lk_\Delta u$ is not prime. Then it follows from Lemma 2.22 that Δ is obtained from a triangulated sphere with $g_2 \leq 2$ by a bistellar 1-move and edge contraction.

Suppose that $lk_{\Delta}x$ is prime for every vertex $x \in \Delta$. Then $\Delta \in \mathcal{H}_3$ and the result follows from Lemmas 2.20, 2.21, 3.2, and 3.3. \square

4 4-dimensional Homology Manifolds with $g_2 = 4$

Lemma 4.1. *Let $\Delta \in \mathcal{H}_4$. If u is a vertex in Δ with $g_2(lk_{\Delta}u) = 1$ and Δ satisfies $(*)$, then $f_0(lk_{\Delta}u) \leq 7$.*

Proof. If possible let $f_0(lk_{\Delta}u) \geq 8$. Let $lk_{\Delta}u = C \star \partial(pqc)$, where C is a cycle of length $n \geq 5$. Here, $pqc \in \Delta$, and every vertex of C is incident to at least one diagonal edge. Since $g_2(\Delta) = 4$, $lk_{\Delta}u$ contains exactly three missing edges that are present in Δ . Therefore, by Corollary 2.16, $V(\Delta) = V(st_{\Delta}u)$. Since $pqc \in \Delta$, $lk_{\Delta}(pqc)$ is a cycle containing at least three vertices of C . Let $a \in C \cap lk_{\Delta}pqc$. Then $\partial(upqc) \subseteq lk_{\Delta}a$ but $upqc \notin \Delta$. This contradicts the fact that $lk_{\Delta}a$ is prime. Therefore $f_0(lk_{\Delta}u) \leq 7$. \square

Lemma 4.2. *Let $\Delta \in \mathcal{H}_4$. Let u be a vertex with $g_2(lk_{\Delta}u) = 1$, $f_0(lk_{\Delta}u) = 7$, and Δ satisfies $(*)$. Then Δ is obtained by an edge expansion from a triangulated sphere with $g_2 \leq 3$.*

Proof. Let $lk_{\Delta}u = C \star \partial(pqc)$, where C is a cycle of length 4. Let us take $C = C(a, b, d, t)$ as shown in Figure 1. Here, $pqc \in \Delta$, and every vertices of C are incident to at least one diagonal edge. Further, by the same arguments as in the proof of Lemma 4.1, $V(lk_{\Delta}pqc) \subseteq V(\Delta \setminus st_{\Delta}u)$ and $lk_{\Delta}u$ contains exactly two missing edges that are present in Δ as shown in Figure 1 using dotted lines.




Figure 1

We claim that $d(k) \geq 8$ for every vertex $k \in C$. If possible, let $d(a) = 7$. If $lk_{\Delta}ad$ contains exactly 4 vertices, then it is a boundary complex of a 3-simplex. Since $ad \notin lk_{\Delta}pqc$, $lk_{\Delta}ad$ contains exactly two vertices from $\{p, q, c\}$. Without loss of generality, let $p, q \in lk_{\Delta}ad$. Then $lk_{\Delta}ad = \partial(btpq)$. Therefore $d \star \partial(btpq) \subseteq lk_{\Delta}a$. Since $lk_{\Delta}a$ is prime, $\partial(btpq) \subseteq lk_{\Delta}a$ implies $btpq \in lk_{\Delta}a$ and so $lk_{\Delta}a = \partial(dbtpq)$, which contradicts the fact that $c \in lk_{\Delta}a$. Therefore, $lk_{\Delta}ad$ contains exactly 5 vertices from the set $\{p, q, c, b, t\}$, and $lk_{\Delta}ad$ is any one of the following type:

- (i) $\partial(pqc) \star \partial(bt)$
- (ii) $\partial(btc) \star \partial(pq)$ or, $\partial(btp) \star \partial(cq)$ or, $\partial(btq) \star \partial(pc)$.

If $lk_{\Delta}ad$ is of type (i), then $lk_{\Delta}pq$ contains the triangles adt and adb . On the other hand, $lk_{\Delta}u = C \star \partial(pqc)$ implies $C \star u \subseteq lk_{\Delta}pq$. Since $lk_{\Delta}pq$ is closed, the maximal simplices of $lk_{\Delta}pq$ are the 2 simplices of $(u \star C) \cup \{adt, adb\}$, which contradicts the fact that $c \in lk_{\Delta}pq$. Let $lk_{\Delta}ad$ be of type (ii). Without loss of generality, let $lk_{\Delta}ad = \partial(btc) \star \partial(pq)$. Then

$adb, adt \in lk_{\Delta}pc$. Further, $lk_{\Delta}u = C \star \partial(pqc)$ implies the maximal simplices of $lk_{\Delta}pc$ are the 2 simplices of $(C \star u) \cup \{adt, adb\}$. This contradicts the fact that $q \in lk_{\Delta}pc$. Therefore $d(a) \geq 8$. By similar arguments, $d(k) \geq 8$ for every vertex $k \in C$.

From the last discussions we conclude that $d(x) \geq 8$ for each vertex $x \in lk_{\Delta}u$. Since $V(lk_{\Delta}pqc) \subseteq V(\Delta \setminus st_{\Delta}u)$, $\Delta \setminus st_{\Delta}u$ contains at least three vertices. Suppose $g_2(lk_{\Delta}x) \geq 2$ for every vertices $x \in lk_{\Delta}u$. Then $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = \sum_{x \in lk_{\Delta}u} g_2(lk_{\Delta}x) + g_2(lk_{\Delta}u) + \sum_{y \in \Delta \setminus st_{\Delta}u} g_2(lk_{\Delta}y) \geq 18$. On the other hand, $g_3(\Delta) = 0$ implies $\sum_{v \in \Delta} g_2(lk_{\Delta}v) \leq 4 \times g_2(\Delta) = 16$. This is a contradiction. Thus, there is a vertex $v \in lk_{\Delta}u$ such that $g_2(lk_{\Delta}v) = 1$ and $d(v) \geq 8$. Therefore, the result follows from Lemmas 2.20, 2.21, and 4.1. \square

Remark 4.3. Let $\Delta \in \mathcal{H}_4$. If u is a vertex in Δ with $g_2(lk_{\Delta}u) = 1$ and $d(u) \geq 7$, then from Lemmas 2.20, 2.21, 4.1, and 4.2, it follows that Δ is obtained from a triangulated 4-sphere with $g_2 \leq 3$ by an edge expansion.

Lemma 4.4. Let $\Delta \in \mathcal{H}_4$. Let u be a vertex with $g_2(lk_{\Delta}u) = 2$ and $G(\Delta[V(lk_{\Delta}u)]) = G(lk_{\Delta}u)$. If $lk_{\Delta}u$ is not an octahedral 3-sphere, then Δ is obtained from a triangulated sphere with $g_2 \leq 3$ by an edge expansion.

Proof. Since $g_2(lk_{\Delta}u) = 2$ and $lk_{\Delta}u$ is not an octahedral 3-sphere, it follows from Lemma 2.6 that $lk_{\Delta}u$ is obtained from a triangulated 3-sphere K with $g_2 = 1$ by subdividing a triangle. By Lemma 2.5, K is obtained from a prime triangulated 3-sphere $K' := C \star \partial(pqc)$ by finitely many facet subdivisions, where C is an n -cycle, $n \geq 3$. Therefore, if K is prime then $K = K'$. Let $lk_{\Delta}u$ be obtained from K by subdividing the triangle, say σ^2 , with a new vertex, say w . The circle C , or the triangle pqc , contains exactly one vertex of the triangle σ^2 , which we call the central vertex of σ^2 . Moreover, σ^2 is a proper face of each facet that participated in the subdivision proces; otherwise, it contradicts the fact that $lk_{\Delta}u$ is prime. Let $lk_K\sigma^2 = \{r, s\}$, where s is the vertex added at the last facet subdivision. We shall use the same constructions of K, K', σ^2 , and w throughout the article.

Let K be not prime. Since $G(lk_{\Delta}u) = G(\Delta[V(lk_{\Delta}u)])$, the vertex w is not incident to any missing edge in $lk_{\Delta}u$ which is present in Δ . Therefore, $lk_{\Delta}u \cap lk_{\Delta}w = lk_{\Delta}uw = \partial\sigma^2 \star \partial(rs)$, and we can contract the edge uw . Let Δ' be the resulting complex. Then $g_2(\Delta') = 3$, and Δ is obtained from Δ' by an edge expansion.

Suppose K is prime. Then $K = K' = C \star \partial(pqc)$, where C is a cycle of length n . Let $n \geq 4$. Since $lk_{\Delta}u$ has no missing edge that is present in Δ , there is a vertex $z \in C$, which is not adjacent to w . Thus, for the vertex z , $lk_{\Delta}u \cap lk_{\Delta}z = lk_{\Delta}uz = \partial(pqc) \star \partial(xy)$, where x, y are the neighbouring vertices of z in C . Thus, we can contract the edge zw , and hence we are done. Let $n = 3$ and $K = \partial(abt) \star \partial(pqc)$. Without loss of generality, let $\sigma^2 = pqt$. Here $wc \notin \Delta$. If pqc is a missing triangle in Δ , then the pair (u, c) satisfies the link condition. Let $pqc \in \Delta$. Then $lk_{\Delta}pqc$ contains at least two elements from $V(\Delta \setminus st_{\Delta}u)$, and so $d(x) \geq 7$ for each vertex in $\{a, b, t, p, q, c\}$. If $g_2(lk_{\Delta}y) = 1$ for some vertex $y \in \{a, b, t, p, q, c\}$, then we are done. Let $g_2(lk_{\Delta}x) \geq 2$ for every vertex $x \in \{a, b, t, p, q, c\}$. Then Lemma 2.1 implies that $V(\Delta \setminus st_{\Delta}u)$ contains at most one element. Therefore, each of the triangles abt and pqt is a missing triangle in Δ . Hence the pair (u, t) satisfies the link condition. Let Δ' be the complex obtained by contracting the edge ut . Then $g_2(\Delta') = 2$, and Δ is obtained from Δ' by an edge expansion. \square

Lemma 4.5. Let $\Delta \in \mathcal{H}_4$. Let u be a vertex with $g_2(lk_{\Delta}u) = 2$ and $lk_{\Delta}u$ be not an octahedral 3-sphere. If $lk_{\Delta}u$ contains at least one missing edge that is present in Δ , then Δ is obtained from a triangulated sphere with $g_2 \leq 3$ by an edge expansion.

Proof. Let K, K', σ^2 , and w be as in Lemma 4.4.

Case 1: Suppose K is not prime, and let $lk_K \sigma^2 = \{r, s\}$, where s is the vertex added to K' at the last facet subdivision. The circle C , or the triangle pqc , contains exactly one vertex of the triangle σ^2 , which we call the vertex as the central vertex of σ^2 . Note that if w is not incident to any missing edge of $lk_\Delta u$ that is present in Δ , then the result follows from a similar argument as in Lemma 4.4. For the remaining part, we shall assume that w is incident to a missing edge in $lk_\Delta u$ that is present in Δ .

Let $S = \{s_1, \dots, s_m\}$ be the set vertices added during the facet subdivisions of K' where $s = s_m$. If s_j is not incident to any missing edge of $lk_\Delta u$ that is present in Δ , then $lk_\Delta s_j \cap lk_\Delta u = lk_\Delta us_j = \partial(\sigma^2) \star \partial(yz)$, where y and z are the vertices incident to s_j in $lk_\Delta u$ other than $V(\sigma^2)$. Thus, we can contract the edge us_j , and we are done. So from now onwards we assume that each vertex in S is incident to a missing edge in $lk_\Delta u$ which is present in Δ .

Note that each vertex in $S \cup V(st_\Delta w)$ has degree at least 7. Now suppose $g_2(lk_\Delta x) \geq 2$ for every vertex $x \in S \cup V(st_\Delta w)$. Since $g_3(\Delta) = 0$, from Lemma 2.1, we have $f_0(\Delta) \leq 8$. But K is not prime, which implies $f_0(lk_\Delta u) \geq 8$, and hence $f_0(\Delta) \geq 9$. Therefore, $g_2(lk_\Delta y) = 1$ for some $y \in S \cup V(st_\Delta w)$. Thus, the result follows from Lemmas 2.20, 2.21, and Remark 4.3.

Case 2: Suppose K is prime, i.e., $K = C \star \partial(pqc)$, where C is a cycle of length $n \geq 3$. Let $n = 3$ and $K = \partial(abt) \star \partial(pqc)$. Without loss of generality, we assume that $\sigma^2 = pqt$. Then wc is the missing edge in $lk_\Delta u$, which is present in Δ . Therefore, $d(x) \geq 7$ for every vertex $x \in lk_\Delta u$. If $g_2(lk_\Delta y) = 1$, for some $y \in lk_\Delta u$, then the result follows from Lemmas 2.21, 2.20, and Remark 4.3. Assume that $g_2(lk_\Delta x) \geq 2$ for every vertex $x \in \Delta$. Then Lemma 2.1 implies $V(\Delta \setminus st_\Delta u) = \emptyset$. Therefore, abt, pqc , and pqt are missing triangles in Δ . Hence $lk_\Delta u \cap lk_\Delta t = C(p, w, q, c) \star \partial(ab) = lk_\Delta ut$. Therefore, we can contract the edge ut , and we are done.

Let $n \geq 4$ and say $C = C(t, a, b, \dots, d, t)$. We prove that if there is a vertex, say $z \in C$, which is not incident to a missing edge e of $lk_\Delta u$ that is present in Δ , then the pair (u, z) satisfies the link condition.

Let z be a vertex in C which is not incident to e . If z is not adjacent to w , then $lk_\Delta u \cap lk_\Delta z = lk_\Delta uz = \partial(pqc) \star \partial(xy)$, where x, y are the vertices in C adjacent to z . Now assume that zw is an edge in Δ . Let $\sigma^2 = abc$, where c is the central vertex of σ^2 . Without loss of generality, let $z = a$. Then $lk_\Delta u \cap lk_\Delta z = lk_\Delta uz = \partial(pqct) \# \partial(pqcw) \# \partial(pqwb)$ (since $wt, bt, pqc, pqw \notin lk_\Delta u$). It is yet to consider $\sigma^2 = pqt$, where t is the central vertex of σ^2 . If t is not incident to any missing edge e of $lk_\Delta u$ that is present in Δ , then pqt is a missing triangle in Δ . Therefore $lk_\Delta u \cap lk_\Delta t = C(c, p, w, q) \star \partial(ad) = lk_\Delta ut$ (since $wc, ad \notin lk_\Delta u$, and $pq \notin lk_\Delta t$). On the other hand, if a is not incident to e , then $lk_\Delta a \cap lk_\Delta u = lk_\Delta au = \partial(bpqc) \# \partial(pqct) \# \partial(pqtw)$ (since $pqt, pqc \notin lk_\Delta u$). The situations for all the remaining vertices are trivial. Thus, the pair (u, z) satisfies the link condition.

Let all the vertices of C be incident to at least one missing edge in $lk_\Delta u$ which is present in Δ . Therefore, $n = 4$ and $d(x) \geq 7$ for each vertex $x \in V(C) \cup V(pqc)$. Now $g_2(lk_\Delta x) \geq 2$ for every vertex $x \in V(C) \cup V(p, q, c)$ together with $g_3(\Delta) = 0$ contradicts Lemma 2.1. Hence there is a vertex y in $V(C) \cup V(pqc)$ with $g_2(lk_\Delta y) = 1$, and the result follows from Lemmas 2.21, 2.20, and Remark 4.3. \square

Lemma 4.6. *Let $\Delta \in \mathcal{H}_4$ and u be a vertex in Δ with $g_2(lk_\Delta u) = 1$. If $d(u) = 6$, then Δ is obtained by an edge expansion from a triangulated sphere with $g_2 \leq 3$.*

Proof. Let $lk_{\Delta}u = \partial(abt) \star \partial(pqc)$. If any one of abt or pqc is a missing triangle in Δ , then the result follows from Lemma 2.20. Let both the triangles be present in Δ . Then $V(lk_{\Delta}pqc)$ and $V(lk_{\Delta}abt)$ are subsets of $V(\Delta \setminus st_{\Delta}u)$. Therefore, the degree of each vertex of $lk_{\Delta}u$ is at least 9.

If $g_2(lk_{\Delta}v) \geq 3$ for every vertex $v \in lk_{\Delta}u$, then $\sum_{v \in \Delta} g_2(lk_{\Delta}v) > 3 \times 6 + 1 = 19$. But $g_3(\Delta) = 0$ and the equation $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = 3g_3(\Delta) + 4g_2(\Delta)$ implies $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = 16$. This presents a contradiction. Thus, there is at least one vertex, say x , in $lk_{\Delta}u$ with $g_2(lk_{\Delta}x) \leq 2$. Now, the result follows from Remark 4.3 and Lemmas 4.2 - 4.5. \square

Theorem 4.7. *Let Δ be a prime homology 4-manifold with $g_2 = 4$. Then Δ is obtained from a triangulated 4-sphere with $g_2 \leq 3$ by one of the following operations:*

- (i) *a bistellar 1-move and an edge contraction,*
- (ii) *an edge expansion.*

Proof. If there is a vertex u in Δ such that $lk_{\Delta}u$ is not prime, then the result follows from Lemma 2.22. Let the link of every vertex in Δ be prime. By Lemma 2.3, there is a vertex u in Δ such that $g_2(lk_{\Delta}u) \leq 2$. Now the result follows from Remark 4.3, Lemma 2.25, and all the discussed lemmas in Section 4. \square

5 4-dimensional Homology Manifolds with $g_2 = 5$

Lemma 5.1. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_{\Delta}u) = 1$ and Δ satisfies $(*)$. Then $f_0(lk_{\Delta}u) \leq 8$.*

Proof. If possible, let $f_0(lk_{\Delta}u) \geq 9$. Since $g_2(lk_{\Delta}u) = 1$, $lk_{\Delta}u = C \star \partial(pqc)$, where C is a cycle of length $n \geq 6$. Here, $pqc \in \Delta$, and each vertex of C is incident to at least one diagonal edge. Let there be exactly three missing edges, say e_1, e_2 , and e_3 in $lk_{\Delta}u$, which are present in Δ . Then $n = 6$, and there are only two types of positions of the edges e_1, e_2 , and e_3 , as shown in Figure 2. Note that no vertex of C lies in $lk_{\Delta}pqc$, otherwise the link of that vertex will not be prime. Therefore, the triangle pqc can never be in the link of the edges e_1, e_2 and e_3 . In the case of Figure 2 (i), the link of the edge sd contains no vertex of C , and so $lk_{\Delta}sd$ contains at least two vertices from $\Delta \setminus st_{\Delta}u$. The link of other two diagonal edges in Figure 2 (i) contains at most one vertex from C . Since $pqc \notin lk_{\Delta}e_i$, the link of these two edges must contain at least one vertex from $\Delta \setminus st_{\Delta}u$. By similar arguments, the link of every diagonal edges in Figure 2(ii) contains at least




Figure 2

two vertices from $\Delta \setminus st_{\Delta}u$. Thus, the number of edges between the vertices in $V(C)$ and

$V(\Delta \setminus st_\Delta u)$ is at least 8. Since no vertices of C lie in $lk_\Delta pqc$, $V(lk_\Delta pqc) \subseteq V(\Delta \setminus st_\Delta u)$. Therefore, the number of edges between $V(\Delta \setminus st_\Delta u)$ and $V(pqc)$ is at least 9. By Proposition 2.15, $G(\Delta \setminus st_\Delta u)$ is a complete graph. Let $f_0(\Delta \setminus st_\Delta u) = m$. Then the total number of edges in Δ is at least $\binom{m}{2} + 8 + 9 + f_1(st_\Delta u) + |\{e_1, e_2, e_3\}| = \binom{m}{2} + 56$. Therefore, $g_2(\Delta) \geq \frac{m(m-1)}{2} + 56 - 5(m+10) + 15$, i.e., $m^2 - 11m + 32 \leq 0$. But there is no positive integer m , for which the last inequality is satisfied. Therefore, the situations in Figure 2 are not possible.

If $lk_\Delta u$ contains four missing edges that are present in Δ , then $V(\Delta) = V(st_\Delta u)$. Then pqc is a missing triangle in Δ , a contradiction to the assumption. Thus, the result follows. \square

Lemma 5.2. *Let $\Delta \in \mathcal{H}_5$ and u be a vertex with $g_2(lk_\Delta u) = 1$. If $d(u) = 8$, and Δ satisfies (*), then Δ is obtained by an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Proof. Since $g_2(lk_\Delta u) = 1$ and $d(u) = 8$, $lk_\Delta u = C \star \partial(pqc)$, where C is a cycle of length 5. Here, $pqc \in \Delta$, and every vertex of C is incident to at least one diagonal edge. Then the number of missing edges in $lk_\Delta u$ that are present in Δ is at least three.

Let there be exactly three missing edges, say e_1, e_2 , and e_3 in $lk_\Delta u$, which are present in Δ . Then the only possible way the edges e_1, e_2 , and e_3 can be arranged in C is shown in Figure 3 by dotted lines. We denote the vertex set of the cycle C by $\{a, b, t, r, s\}$. Note




Figure 3

that $pqc \notin lk_\Delta e_i$, for every i . Therefore, $lk_\Delta sb$ contains at least one more vertex from $\Delta \setminus st_\Delta u$. If $lk_\Delta ar$ does not contain any vertex from $\Delta \setminus st_\Delta u$, then $lk_\Delta ar = \partial(st) \star \partial(pqc)$ (since $st \notin \Delta$ and $pqc \notin lk_\Delta a$). Let us flip the edge ar by st (replace $ar \star \partial(st) \star \partial(pqc)$ by $\partial(ar) \star st \star \partial(pqc)$) and let Δ' be the resulting complex. Then Δ' is a homology manifold with $g_2 = 5$ and $lk_\Delta u = lk_{\Delta'} u$. Though $lk_{\Delta'} u$ contains three missing edges that are present in Δ' , the vertex $r \in C$ is incident to none of the missing edges in Δ' . Hence, $lk_{\Delta'} u \cap lk_{\Delta'} r = lk_{\Delta'} ur = \partial(pqc) \star \partial(st)$, and so we can contract the edge ur . Let Δ'' be the resulting complex, then $g_2(\Delta'') = 4$, and Δ' is obtained from Δ'' by an edge expansion. Further, Δ is obtained from Δ' by flipping an edge. Similarly, if $lk_\Delta at$ does not contain any vertex from $\Delta \setminus st_\Delta u$, we use the same arguments to flip at by br , and hence the vertex t will serve the link condition in this case. Therefore, Δ is obtained by flipping an edge together with an edge expansion from a triangulated 4-sphere with $g_2 = 4$. It remains to be considered when $lk_\Delta e_i$ contains at least one vertex from $\Delta \setminus st_\Delta u$. Let $lk_\Delta e_i$ contain at least one vertex from $\Delta \setminus st_\Delta u$, for each i . Then $f_0(lk_\Delta x) \geq 8$, for each vertex $x \in lk_\Delta u$, and further, $f_0(lk_\Delta x) \geq 9$ for $x \in \{a, p, q, c\}$.

Let the number of vertices in $\Delta \setminus st_\Delta u$ be m . By Proposition 2.15, $G(\Delta \setminus st_\Delta u)$ is a complete graph. By hypothesis ($lk_\Delta e_i$ contains at least one vertex from $\Delta \setminus st_\Delta u$), the number of edges between vertices of $\Delta \setminus st_\Delta u$ and vertices of C is at least 5. On the

other hand, $V(lk_{\Delta}pqc) \subseteq V(\Delta \setminus st_{\Delta}u)$. So the number of edges between the vertices of pqc and the vertices in $\Delta \setminus st_{\Delta}u$ is at least 9, and equality occurs when $lk_{\Delta}pqc$ is a 3-cycle. Therefore, $f_1(\Delta) \geq \binom{m}{2} + 5 + 9 + f_1(st_{\Delta}u) + |\{e_1, e_2, e_3\}| = \binom{m}{2} + 48$. Moreover, if $V(\Delta) \setminus V(st_{\Delta}u \cup lk_{\Delta}pqc) \neq \emptyset$, then the number of edges will be more as we shall see when required.

We first claim that $lk_{\Delta}pqc$ is a 3-cycle. If $lk_{\Delta}pqc$ is a cycle of length greater than 3, then $f_1(\Delta) \geq \binom{m}{2} + 48 + 3k$, for some natural number k . However, $g_2(\Delta) = 5$ implies that, $5 \geq \binom{m}{2} + 48 + 3k - 5(9 + m) + 15$, i.e., $m^2 - 11m + 26 + 6k \leq 0$, where $k \geq 1$. But there is no such natural number m satisfying the last inequality. Hence $lk_{\Delta}pqc$ is a 3-cycle. Take $lk_{\Delta}pqc = \partial(xyz)$, for some vertices x, y , and z in $\Delta \setminus st_{\Delta}u$.

Let us calculate the possible values of m . Note that $g_2(\Delta) = 5$, and $f_1(\Delta) \geq \binom{m}{2} + 48$ implies $m^2 - 11m + 26 \leq 0$, i.e., $m(11 - m) \geq 28$, where $m \geq 3$. The only positive integers satisfying the last inequality are $m = 4, 5, 6$, and 7.

Let $m \leq 6$. If $xyz \notin \Delta$, then we apply a bistellar 2-move with respect to the pair (pqc, xyz) . Let Δ' be the resulting complex. By Lemma 2.20, Δ' is obtained by an edge expansion from a triangulated sphere with $g_2 = 2$, and Δ is obtained from Δ' by a bistellar 2-move. Now assume that $xyz \in \Delta$. Then $lk_{\Delta}xyz$ is a cycle. Further, $lk_{\Delta}xyz$ contains no vertex from $\{p, q, c\}$. If $lk_{\Delta}xyz$ contains a vertex from C , then there are at least 9 edges between the vertices of C and $V(\Delta \setminus st_{\Delta}u)$, and hence $f_1(\Delta) \geq \binom{m}{2} + 11 + 9 + f_1(st_{\Delta}u) + |\{e_1, e_2, e_3\}| = \binom{m}{2} + 52$, which contradicts the fact that $g_2 = 5$. Thus, $lk_{\Delta}xyz$ is a 3-cycle, say $\partial(x_1x_2x_3)$, where $\{x_1, x_2, x_3\} \subseteq V(\Delta \setminus st_{\Delta}u)$. Clearly $x_1x_2x_3$ is a missing triangle in Δ , as otherwise $lk_{\Delta}x_1x_2x_3$ will be a cycle containing vertices from $lk_{\Delta}u$, which gives a contradiction on $g_2(\Delta) = 5$. Now we apply bistellar 2-moves with respect to pairs $(x_1x_2x_3, xyz)$ and (pqc, xyz) . Let Δ'' be the resulting complex. By similar arguments, Δ is obtained by two repeated bistellar 2-moves and an edge expansion from a triangulated sphere with $g_2 \leq 2$.

Let $m = 7$. Since $G(\Delta \setminus st_{\Delta}u)$ is a complete graph and $lk_{\Delta}(pqc) = \partial(xyz)$, degree of each of the vertices x, y , and z is at least 9. Thus, Δ contains at least seven vertices of degree at least 9. Now $f_0(\Delta) = 16$ and the relation $\sum_{v \in \Delta} g_2(lk_{\Delta}v) \leq 20$ implies there is at least one vertex, say x , with $d(x) \geq 9$ and $g_2(lk_{\Delta}x) = 1$. Therefore, the proof follows from Lemmas 2.20, 2.21, and 5.1.

If $lk_{\Delta}u$ contains four missing edges that are present in Δ , then $V(\Delta) = V(st_{\Delta}u)$. Therefore, pqc is a missing triangle in Δ , which is a contradiction. \square

Lemma 5.3. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_{\Delta}u) = 1$. If $d(u) = 7$, and Δ satisfies (*), then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-moves (possibly zero times).*

Proof. Let $lk_{\Delta}u = C(a, b, d, t) \star \partial(pqc)$. Here, $pqc \in \Delta$, and every vertex of C is incident to at least one diagonal edge. Therefore, $lk_{\Delta}u$ contains exactly two diagonal edges. Let $G(\Delta[V(lk_{\Delta}u)]) = G(lk_{\Delta}u) \cup \{e_1, e_2\}$ where $e_1 = ad$ and $e_2 = bt$. If we can find a vertex x with $g_2(lk_{\Delta}x) = 1$ and $d(x) \geq 8$, then the result follows from Lemmas 2.20, 2.21, 5.1, and 5.2. By the same arguments as in Lemma 4.2, $lk_{\Delta}ad$ and $lk_{\Delta}bt$ contain at least one vertex from $V(\Delta \setminus st_{\Delta}u)$, so that $d(x) \geq 8$ for every vertex $x \in C$. Therefore, Δ contains at least seven vertices (the vertices of $lk_{\Delta}u$) of degree at least 8. If the g_2 value of any of these vertex links is 1, then we are done. So for the remaining part, we assume $g_2(lk_{\Delta}x) \geq 2$, for every vertex $x \in lk_{\Delta}u$. From the relations $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = 3g_3(\Delta) + 4g_2(\Delta)$ and $g_3(\Delta) = 0$, we get $f_0(\Delta \setminus st_{\Delta}u) \leq 5$.

Let $lk_{\Delta}pqc$ be a 3-cycle, say $\partial\tau$, where τ is a 2-simplex. If $\tau \in \Delta$, then $lk_{\Delta}\tau$ will be a cycle. Since no vertex from $\{p, q, c\}$ is in $lk_{\Delta}\tau$, the degree of each vertex in τ is at least 8. It follows that Δ contains at least one vertex, say x with $d(x) \geq 8$ and $g_2(lk_{\Delta}x) = 1$, and we are done. If $\tau \notin \Delta$ then replace the 4-dimensional ball $pqc \star \partial(\tau)$ by $\tau \star \partial(pqc)$. In the resulting complex Δ' , $lk_{\Delta'}u$ is the same as $lk_{\Delta}u$, but pqc is a missing triangle here. Then it follows from Lemma 2.20 that Δ is obtained by a bistellar 2-move from an edge expansion on a triangulated sphere with $g_2 = 3$.

Now we assume that $lk_{\Delta}pqc$ is a cycle of length greater than 3. Let $f_0(\Delta \setminus st_{\Delta}u) = 4$ and $lk_{\Delta}pqc = C(x_1, x_2, x_3, x_4)$. We claim that there is at least one vertex, say x_i with $d(x_i) = 6$. If possible let $d(x_i) \geq 7$ for $1 \leq i \leq 4$. Without loss of generality, let $x_1 \in lk_{\Delta}ad$. If $x_1x_3 \in \Delta$ then $d(x_1) \geq 8$. Then $f_1(\Delta) \geq \binom{8}{2} + 3 \cdot 7 + 8 - \binom{4}{2} = 51$. If $x_1x_3 \notin \Delta$ then $f_1(\Delta) \geq \binom{8}{2} + 4 \cdot 7 - (\binom{4}{2} - 1) = 51$. This contradicts the fact that $g_2(\Delta) = 5$. Thus, there is a vertex in $lk_{\Delta}pqc$ whose link contains exactly six vertices. Let the vertex be x_2 . Then $g_2(lk_{\Delta}x_2) = 1$ and x_2 is adjacent to at least two vertices in the cycle $lk_{\Delta}pqc$. Let $lk_{\Delta}x_2 = \partial\tau_1 \star \partial\tau_2$, where τ_1 and τ_2 are 2-simplices. If any one of them is a missing simplex in Δ , then the link condition is satisfied for a pair (u, x) , where x is any vertex in that triangle. So Δ is obtained by an edge expansion from a triangulated sphere with $g_2 = 4$. However, if both 2-simplices are present in Δ , then the link of both the vertices in $lk_{\Delta}pqc$ that are adjacent to x_2 has a degree of at least 8. As a result, we conclude that Δ has at least one vertex, say x , with $g_2(lk_{\Delta}x) = 1$ and $d(x) = 8$, and we are done.

Let $f_0(\Delta \setminus st_{\Delta}u) = 5$, $V(\Delta \setminus st_{\Delta}u) = \{x_1, x_2, x_3, x_4, x_5\}$, and $lk_{\Delta}pqc = C(x_1, x_2, x_3, x_4)$. In this case $g_2(lk_{\Delta}x_j) = 1$ for each j . We claim that there exists an i such that $d(x_i)$ is either 6 or at least 8. If possible let $d(x_i) = 7$ for $1 \leq j \leq 5$. Then $g_2(\Delta) = 5$ implies at most two edges are missing edges whose end points are in $\{x_1, x_2, x_3, x_4, x_5\}$. Let $x_1 \in lk_{\Delta}ad$ and $x_k \in lk_{\Delta}bt$. If $x_1 = x_k$, then $d(x_1) \geq 9$ and hence we are done. Let $k \in \{2, 3, 4\}$. Then x_1x_3, x_1x_5 , and x_kx_5 are missing, and we get a contradiction. Thus, $x_5 \in lk_{\Delta}bt$. Since x_1x_3 and x_1x_5 are missing, all the remaining vertices are adjacent to each other. Since $d(x_i) = 7$ for $1 \leq j \leq 5$, x_2 and x_4 are not adjacent to any vertices from $\{a, b, d, t\}$, x_3 is adjacent to exactly one vertex from $\{a, b, d, t\}$ and x_5 is adjacent to two vertices from $\{a, d, p, q, c\}$. Thus, we shall not get five vertices in both of the links $lk_{\Delta}(bx_5)$ and $lk_{\Delta}(tx_5)$, and we get a contradiction. Therefore, $lk_{\Delta}(ad)$ and $lk_{\Delta}(bt)$ does not contain any vertices from $\{x_1, x_2, x_3, x_4\}$, and x_5 is in both of the links $lk_{\Delta}(ad)$ and $lk_{\Delta}(bt)$. Since $lk_{\Delta}(bx_5)$ contains at least five vertices, we must have at least two edges either from b to a vertex from $\{x_1, x_2, x_3, x_4\}$ or from x_5 to a vertex from $\{p, q, c\}$. But, for those cases, we shall have more than two missing edges whose end points are in $\{x_1, x_2, x_3, x_4, x_5\}$, and we get a contradiction. Therefore, the claim is achieved. Now the result follows from similar arguments as in the last paragraph.

Let $f_0(\Delta \setminus st_{\Delta}u) = 5$ and $lk_{\Delta}pqc = C(x_1, x_2, x_3, x_4, x_5)$. In this case $g_2(lk_{\Delta}x_j) = 1$ for each j . We claim that there exists an i such that $d(x_i)$ is either 6 or at least 8. If possible let $d(x_i) = 7$ for $1 \leq j \leq 5$. Then $g_2(\Delta) = 5$ implies at most two edges are missing edges whose end points are in $\{x_1, x_2, x_3, x_4, x_5\}$. Let $x_1 \in lk_{\Delta}ad$ and $x_k \in lk_{\Delta}bt$. If $x_1 = x_k$ then $d(x_1) \geq 9$ and hence we are done. Let $k \in \{2, 3, 4, 5\}$. At least three edges from $x_1x_3, x_1x_4, x_kx_{k-2}$ and x_kx_{k+2} are missing, and we get a contradiction. Therefore, the claim is achieved. Now the result follows from similar arguments as in the third paragraph of this lemma. \square

Lemma 5.4. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_{\Delta}u) = 2$, and $lk_{\Delta}u$ contains at most one missing edge that is present in Δ . If $d(u) \geq 8$ and $lk_{\Delta}u$ is not an octahedral 3-sphere,*

then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion.

Proof. If $G(\Delta[V(lk_\Delta u)]) = G(lk_\Delta u)$, then the result follows from similar arguments as in Lemma 4.4. Now suppose $G(\Delta[V(lk_\Delta u)]) = G(lk_\Delta u) \cup \{e\}$. Let K, K', σ^2 , and w be as in Lemma 4.4.

Suppose K is not prime, and let $lk_K \sigma^2 = \{r, s\}$, where s is the vertex added to K' at the last facet subdivision. If e is not incident to w , then $lk_\Delta u \cap lk_\Delta w = \partial \sigma^2 \star \partial(rs) = lk_\Delta uw$. On the other hand, if e is not incident to s , then $lk_\Delta u \cap lk_\Delta s = \partial \sigma^2 \cap \partial(wx) = lk_\Delta us$, where $lk_K s = \partial(x \star \sigma^2)$. Thus, in both cases, we can contract an edge. Let Δ' be the resulting complex after contracting the edge. Then $g_2(\Delta') = 4$, and Δ is obtained from Δ' by an edge expansion.

If K is prime, then $K = K' = C \star \partial(pqc)$, where C is a cycle of length $n \geq 4$. Let y be a vertex in C which is not incident to e . By the same arguments as in Lemma 4.5, the pair (u, y) satisfies the link condition, and we are done. \square

Lemma 5.5. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_\Delta u) = 2$, and $G(\Delta[V(lk_\Delta u)]) = G(lk_\Delta u) \cup \{e_1, e_2\}$. If $d(u) \geq 8$ and $lk_\Delta u$ is not an octahedral 3-sphere, then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Proof. Let K, K', σ^2 and w be as in Lemma 4.4.

Case 1: Let K be not prime. Let s_1, \dots, s_m be the vertices that are added to K' during the subdivision of tetrahedrons to form K . By applying similar arguments as in Lemma 4.4 and 4.5, we can show that if at least one vertex in $\{w, s_1, \dots, s_m\}$ is not incident to some e_i , then Δ is obtained by an edge expansion from a triangulated sphere with $g_2 \leq 4$. For the remaining part we assume that each vertex in the set $\{w, s_1, \dots, s_m\}$ is incident to at least one edge in $\{e_1, e_2\}$. Therefore $m \leq 3$.

Let $K' = \partial(abt) \star \partial(pqc)$. Without loss of generality, let $\sigma^2 = pqt$. Clearly $lk_{K'}(pqt) = \{a, b\}$. Suppose $m = 1$ and let $pqta$ be the missing tetrahedron in K such that $lk_K s_1 = \partial(pqta)$. Thus, the degree of each vertex of $lk_\Delta u$ is at least 7. If $g_2(lk_\Delta x) = 1$ for some vertex in Δ , then the result follows from Lemma 5.3. Let $g_2(lk_\Delta x) \geq 2$ for every vertex in $lk_\Delta u$. Then the relation $\sum_{v \in \Delta} g_2(lk_\Delta v) = 4g_2(\Delta)$ implies $\sum_{v \in \Delta \setminus st_\Delta u} g_2(lk_\Delta v) \leq 2$. So $\Delta \setminus st_\Delta u$ contains at most two vertices.

In $lk_\Delta u$, the missing edges are wc, wa, s_1b , and s_1c . Now if abt is a missing triangle in Δ , then $lk_\Delta u \cap lk_\Delta b = \partial(wpqt) \# \partial(pqtc) \# \partial(pqca) = lk_\Delta ub$, since $aw, wc, pqc, pqt \notin lk_\Delta u$, and $at \notin lk_\Delta b$. On the other hand, if $pqc \notin \Delta$, then $lk_\Delta pu = (C(a, q, w, t) \star \partial(bs_1)) \#_{abt} \partial(abtc)$. We claim that $lk_\Delta u \cap lk_\Delta p = lk_\Delta pu$. Note that $abt, wa, wc, s_1b, s_1c \notin lk_\Delta u$ and $qc \notin lk_\Delta p$. Therefore, to conclude the claim, it is enough to prove $pqt \notin \Delta$. If $pqt \in \Delta$, then $lk_\Delta(pqt)$ is a circle which does not contain c (since $pqc \notin \Delta$). Moreover no vertex from $\{a, b, s_1, w\}$ is in $lk_\Delta(pqt)$, otherwise $\partial(upqt)$ will be a missing tetrahedron in the corresponding vertex link. Thus, $V(lk_\Delta pqt) \subseteq V(\Delta \setminus st_\Delta u)$, which is not possible. Therefore, $pqt \notin \Delta$, and we conclude that $lk_\Delta u \cap lk_\Delta p = lk_\Delta pu = (C(a, q, w, t) \star \partial(bs_1)) \#_{abt} \partial(abtc)$.

Let us assume that both $abt, pqc \in \Delta$. Suppose both wc and s_1c are present in Δ , i.e., $\{e_1, e_2\} = \{wc, s_1c\}$. Then abt is a missing triangle in Δ ; because if $abt \in \Delta$, then $lk_\Delta abt \subseteq \Delta \setminus st_\Delta u$, and hence $\Delta \setminus st_\Delta u$ contains at least three vertices, which is a contradiction. If at least one of wc and s_1c is a missing edge in Δ , then one of the links, $lk_\Delta abt$ and $lk_\Delta pqc$, contains at least two vertices, and the other contains at least one vertex from $V(\Delta \setminus st_\Delta u)$. Thus, $V(\Delta \setminus st_\Delta u)$ contains exactly two vertices, say x_1, x_2 and $g_2(lk_\Delta x_i) = 1$, for each i . By Proposition 2.15, the edge $x_1x_2 \in \Delta$. Therefore, at least one of x_1 , and x_2 is adjacent to

more than six vertices, i.e., there is a vertex $x \in \Delta$ with $g_2(lk_\Delta x) = 1$ and $d(x) \geq 7$. Hence the result follows from Lemmas 2.20, 2.21, 5.1, 5.2, and 5.3.

Let $m \geq 2$. If the pair (u, c) satisfies the link condition $lk_\Delta u \cap lk_\Delta c = lk_\Delta uc$, then we are done. Suppose that $lk_\Delta u \cap lk_\Delta c \neq \partial(abt) \star \partial(pq) = lk_\Delta uc$. Then c must be incident to at least one more vertex other than the vertices in $\{a, b, t, p, q, u\}$. Therefore, the degree of each vertex in $lk_\Delta u$ is at least 7. If $g_2(lk_\Delta x) \geq 2$ for every vertex in $lk_\Delta u$, then $f_0(\Delta) \leq 10$. Thus, there is no vertex in $\Delta \setminus st_\Delta u$. It contradicts the fact that $g_2(\Delta) = 5$. So $g_2(lk_\Delta x) = 1$ for some vertex $x \in lk_\Delta u$, and hence the result.

Let C be a cycle of length $n \geq 4$. Since $lk_\Delta u$ has at most two missing edges that are present in Δ and $\{w, s_1, \dots, s_m\} \subseteq V(\{e_1, e_2\})$, there is at least one vertex in the cycle C that is not incident to e_i 's. Hence the result follows from similar arguments as in Lemma 4.5.

Case 2: Let K be prime. Then $K = C \star \partial(pqc)$, where C is a cycle of length $n \geq 3$. Since two missing edges are there in $lk_\Delta u$ that are present in Δ , we have $n \geq 4$. If some vertex, say z of C is not incident to a missing edge e_i , then by the same arguments as in Lemma 4.5, (u, z) satisfies the link condition. Now, suppose each vertex of C is incident to at least one missing edge. Then $n = 4$ and w is not incident to any e_i 's. Let $C = C(a, b, d, t)$. If the central vertex of σ^2 belongs to C , then the pair (u, w) satisfies the link condition trivially.

It remains to consider $\sigma^2 = abc$, where c is the central vertex of σ^2 . Then $lk_\Delta uw = \partial(abc) \star \partial(pq)$. If $wpq \notin \Delta$, then the pair (u, w) satisfied the link condition. Let $wpq \in \Delta$, where $\sigma^2 = abc$. Then $lk_\Delta wpq$ contains at least three vertices. Note that $ac \notin lk_\Delta wpq$; otherwise, $a \in lk_\Delta pqc$ and, hence $\partial(upqc) \subseteq lk_\Delta a$. This contradicts the fact that $lk_\Delta a$ is prime. Since $wpq \notin lk_\Delta u$, it follows that $lk_\Delta wpq$ contains at least one more vertex other than a and b . Therefore, the degree of w is at least 7. Thus, $d(x) \geq 7$ for every $x \in st_\Delta u$. If $g_2(lk_\Delta x) = 1$ for some vertex in $lk_\Delta u$, then the result follows from Lemmas 5.1-5.3 and Lemmas 2.20-2.21. Let $g_2(lk_\Delta x) \geq 2$, for every vertex $x \in lk_\Delta u$. Then Lemma 2.1 and $g_3(\Delta) = 0$ imply that the number of vertices in $\Delta \setminus st_\Delta u$ is at most 2.

If $pqc \notin \Delta$ then $lk_\Delta c \cap lk_\Delta u = lk_\Delta cu = C(a, w, b, d, t) \star \partial(pq)$ holds. On the other hand, if $pqc \in \Delta$, then $lk_\Delta pqc$ contains at most one vertex from $lk_\Delta u$. Note that $lk_\Delta pqc$ cannot contain a vertex of $lk_\Delta u$ other than w . Therefore, $\Delta \setminus st_\Delta u$ contains exactly two vertices and $lk_\Delta pqc = ww_1w_2$, where $\{w_1, w_2\} = V(\Delta \setminus st_\Delta u)$. Clearly, $g_2(lk_\Delta w_i) = 1$, for each i . If $ww_1w_2 \in \Delta$ then $lk_\Delta ww_1w_2$ contains at least three vertices from C , and hence $d(w_i) \geq 7$. If ww_1w_2 is a missing triangle, then we shall replace the 4-dimensional ball $pqc \star \partial(ww_1w_2)$ with $\partial(pqc) \star ww_1w_2$. If Δ' is the resulting complex, then $g_2(\Delta') = 5$. Furthermore, $lk_\Delta u = lk_{\Delta'} u$, and pqc is a missing triangle in $lk_{\Delta'} u$. Thus, Δ is obtained by an edge expansion and bistellar 2-move from a triangulated sphere with $g_2 \leq 4$.

In both Case 1 and Case 2, whenever we have a link condition, we can apply an edge contraction there. So, Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion or a combination of an edge expansion and a bistellar 2-move. \square

Lemma 5.6. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_\Delta u) = 2$, and $G(\Delta[V(lk_\Delta u)]) = G(lk_\Delta u) \cup \{e_1, e_2, e_3\}$. If $lk_\Delta u$ is not an octahedral 3-sphere, then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion.*

Proof. By Corollary 2.16, $V(\Delta) = V(st_\Delta u)$. Let K, K', σ^2 , and w be as in Lemma 4.4.

Let K be not prime and s_1, \dots, s_m be the vertices added during facet subdivisions. Further, we assume that each vertex in $\{w, s_1, \dots, s_m\}$ is incident to at least one edge e_i ;

otherwise, the result follows from the same arguments as in Lemmas 4.4 and 4.5. Therefore $m \leq 5$.

Suppose C is a 3-cycle, i.e., $n = 3$ and take $C = C(a, b, t)$. Without loss of generality, let $\sigma^2 = pqt$ and $pqta$ be a missing tetrahedron in K . We first prove that abt is a missing triangle in Δ . If possible let $abt \in \Delta$. Note that no vertex from the set $\{p, q, c\}$ is in $lk_{\Delta}abt$; otherwise, it contradicts the fact that the link of each vertex of Δ is prime. Therefore, $V(lk_{\Delta}abt) \subseteq \{w, s_1, \dots, s_m\}$, and this implies $m \geq 2$. Since σ^2 participates in each facet subdivision, it follows that any vertex in $\{w, s_1, \dots, s_m\}$ is incident to at most one vertex from the set $\{a, b\}$. Moreover, $wa \notin lk_{\Delta}u$ and at most one s_j is adjacent to a (or b) in $lk_{\Delta}u$. Thus, $V(lk_{\Delta}abt) \subseteq \{w, s_1, \dots, s_m\}$ implies at least four missing edges are there in $lk_{\Delta}u$, which are present in Δ . This is a contradiction. Therefore, abt is a missing triangle in Δ .

Note that the vertex t is not incident to any missing edge e_i . By taking the appropriate subdivisions into account and the fact that no vertex from the set $\{a, b, w, s_1, \dots, s_m\}$ is in $lk_{\Delta}pqt$, it is immediate that the pair (u, t) satisfies the link condition.

Let $K' = C \star \partial(pqc)$ where C is a cycle of length $n \geq 4$. If there is some vertex, say z in C that is free from e_j , then by similar arguments as in Lemma 4.5, the pair (u, z) satisfies the link condition. Thus, every vertex of C is incident to some e_j . It follows that no vertex of pqc incident e_i 's, and hence pqc is a missing triangle in Δ . Thus, there is a vertex, say $x \in \{p, q, c\}$ such that the pair (u, x) satisfies the link condition.

Let K be prime and $K = C \star \partial(pqc)$, where C is a cycle of length n . Since the number of missing edge is 3, we have $n \geq 5$. If there is some vertex, say z in C , which is not incident to any missing edge e_i , then by similar arguments as in Lemma 4.5 the pair (u, z) satisfies the link condition. Let every vertex of C be incident to some e_i . Then $n = 5$. Since p, q, c are adjacent every vertex of C in $lk_{\Delta}u$, it follows that e_i is incident to none of p, q , and c . Therefore, pqc is a missing triangle in Δ and there is a vertex, say $x \in \{p, q, c\}$ such that (u, x) satisfies the link condition. Hence the proof. \square

Lemma 5.7. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_{\Delta}u) = 2$, and $d(u) = 7$. Then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Proof. Let K, K', σ^2 , and w be as in Lemma 4.4. Since $d(u) = 7$, K is prime. Let $K = \partial(abt) \star \partial(pqc)$, and $\sigma^2 = pqt$ as earlier. Then $lk_{\Delta}u$ has at most one missing edge. If any one of the triangles pqc and abt is missing in Δ , then there is a vertex, say x in that triangle such that the pair (u, x) satisfies the link condition. Suppose both the triangles abt and pqc are in Δ . Then the link of both the triangles contain at least two vertices from $V(\Delta \setminus st_{\Delta}u)$. Therefore, the degree of each vertex in $\{p, q, t, a, b\}$ is at least 9 and $d(c) \geq 8$. If $g_2(lk_{\Delta}x) \geq 3$ for every vertex $x \in \{p, q, c, a, b, t\}$, then $\sum_{v \in \Delta} g_2(lk_{\Delta}v) \geq 23$. But the relations $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = 3g_3(\Delta) + 4g_2(\Delta)$ and $g_3(\Delta) = 0$ implies $\sum_{v \in \Delta} g_2(lk_{\Delta}v) = 20$. Therefore, $lk_{\Delta}u$ contains a vertex, say x with $d(x) \geq 8$ and $g_2(lk_{\Delta}x) \leq 2$. Thus, the result follows from Lemmas 2.20, 2.21, 2.25, and 5.1-5.6. \square

Lemma 5.8. *Let $\Delta \in \mathcal{H}_5$. Let u be a vertex with $g_2(lk_{\Delta}u) = 1$ and $d(u) = 6$. Then Δ is obtained from a triangulated sphere with $g_2 \leq 4$ by an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Proof. Let $lk_{\Delta}u = \partial(abt) \star \partial(pqc)$. If any one of the triangles abt , and pqc is a missing triangle in Δ , then for every vertex x of that triangle, the pair (u, x) satisfies the link condition. Suppose both the triangles abt and pqc are in Δ . Then all the vertices of $lk_{\Delta}pqc$

and $lk_{\Delta}abt$ are in $\Delta \setminus st_{\Delta}u$. Also $d(x) \geq 9$ for every vertex $x \in lk_{\Delta}u$. If $g_2(lk_{\Delta}x) \geq 3$ for every vertex $x \in lk_{\Delta}u$, then $\sum_{v \in \Delta} g_2(lk_{\Delta}v) \leq \sum_{v \in \Delta \setminus st_{\Delta}u} g_2(lk_{\Delta}v) + 3 \times 6 + 1$. This implies $\Delta \setminus st_{\Delta}u$ contains at most two vertices, which contradicts the fact that $V(lk_{\Delta}pqc) \subseteq V(\Delta \setminus st_{\Delta}u)$. Thus, there is at least one vertex, say x in $lk_{\Delta}u$, such that $d(x) \geq 9$ and $g_2(lk_{\Delta}x) \leq 2$. Thus, the result follows from Lemmas 2.20, 2.21, 2.25, and 5.1-5.6. \square

Theorem 5.9. *Let Δ be a prime homology 4-manifold with $g_2 = 5$. Then Δ is a triangulated sphere which is obtained from a triangulated 4-sphere with $g_2 \leq 4$ by one of the following operations:*

- (i) *a bistellar 1-move and an edge contraction,*
- (ii) *an edge expansion together with flipping an edge (possibly zero times) or a bistellar 2-move (possibly zero times).*

Proof. If Δ contains at least one vertex, say u , such that $lk_{\Delta}u$ is not prime, then the result follows from Theorem 4.7.

If all the vertices of Δ have prime links, then there is a vertex, say u with $g_2(lk_{\Delta}u) \leq 2$ and the proof follows from Lemmas 2.20, 2.21, 2.25, and all the discussed lemmas in this section. \square

Proof of Theorem 1.4. The proof follows from Theorems 3.4, 4.7, 5.9, and Remark 1.6. \square

Acknowledgement: The first author is supported by Science and Engineering Research Board (CRG/2021/000859). The second author is supported by Prime Minister's Research Fellows (PMRF) Scheme.

References

- [1] L. Asimow, B. Roth, The rigidity of graphs. II, *J. Math. Anal. Appl.* **68** (1) (1979) 171–190.
- [2] B. Bagchi and B. Datta, A structure theorem for pseudomanifolds, *Discrete Math.* **188** (1998), no. 1-3, 41–60.
- [3] B. Bagchi and B. Datta, Lower bound theorem for normal pseudomanifolds, *Expo. Math.* **26** (2008), 327–351.
- [4] D. Barnette, A proof of the lower bound conjecture for convex polytopes, *Pacific J. Math.* **46** (1973), 349–354.
- [5] D. Barnette, Graph theorems for manifolds, *Israel J. Math.* **16** (1973), 62–72.
- [6] B. Basak, R. K. Gupta and S. Sarkar, A characterization of normal 3-pseudomanifolds with at most two singularities, 2021, 27 pages, arXiv: 2104.03751v2.
- [7] B. Basak and E. Swartz, Three-dimensional normal pseudomanifolds with relatively few edges, *Adv. Math.* **365** (2020) 107035, 1–25.
- [8] L. Billera, C. Lee, Sufficiency of McMullen's conditions for f -vectors of simplicial polytopes, *Bull. Amer. Math. Soc.* **2** (1980) 181–185.

- [9] L. Billera, C. Lee, A proof of the sufficiency of McMullen’s conditions for f -vectors of simplicial convex polytopes, *J. Combin. Theory Ser. A* **31** (1981) 237–255.
- [10] U. Brehm and W. Kühnel, Combinatorial manifolds with few vertices, *Topology* **26** (1987), no. 4, 465–473.
- [11] A. Fogelsanger, The generic rigidity of minimal cycles, Ph.D. thesis, Cornell University, 1988.
- [12] G. Kalai, Rigidity and the lower bound theorem I, *Invent. Math.* **88** (1987), 125–151.
- [13] V. Klee, A combinatorial analogue of Poincaré’s duality theorem, *Canadian J. Math.* **16** (1964), 517–531.
- [14] P. McMullen, The maximum numbers of faces of a convex polytope, *Mathematika* **17** (1970), 179–184.
- [15] P. McMullen, The numbers of faces of simplicial polytopes, *Israel J. Math.* **9** (1971), 559–570.
- [16] E. Nevo and E. Novinsky, A characterization of simplicial polytopes with $g_2 = 1$, *J. Combin. Theory Ser. A* **118** (2011), 387–395.
- [17] I. Novik and E. Swartz, Applications of Klee’s Dehn-Sommerville relations, *Discrete Comput. Geom.* **42** (2009), no. 2, 261–276.
- [18] I. Novik and E. Swartz, g -vectors of manifolds with boundary, *Algebraic Combinatorics* **3(4)** (2020), 887–911.
- [19] E. Swartz, Topological finiteness for edge-vertex enumeration, *Adv. Math.* **219** (2008), 1722–1728.
- [20] E. Swartz, Lower bounds for h -vectors of k -CM, independence and broken circuit complexes, *SIAM J. Discrete Math.* **18** (2004), 647–661.
- [21] E. Swartz, Face enumeration: From spheres to manifolds, *J. Eur. Math. Soc.*, **11** (2009), 449–485.
- [22] E. Swartz, Thirty-five years and counting, 2014, 29 pages, arXiv:1411.0987.
- [23] T.S. Tay, Lower-bound theorems for pseudomanifolds, *Discrete Comput. Geom.*, **13** (1995) 203–216.
- [24] T. Tay, N. White and W. Whiteley, Skeletal rigidity of simplicial complexes II, *European J. Combin.* **16** (1995), 503–525.
- [25] D. Walkup, The lower bound conjecture for 3- and 4-manifolds, *Acta Math.* **125** (1970), 75–107.
- [26] W. Whiteley, Cones, infinity and 1-story buildings, *Structural Topology* **8** (1983), 53–70.
- [27] H. Zheng, A characterization of homology manifolds with $g_2 \leq 2$, *J. Combin. Theory Ser. A* **153** (2018), 31–45.