

–  $\mathcal{Q}$ -Set & Friends –

## Regarding Singleton and Gluing Completeness

José G. Alvim\*    Hugo L. Mariano†    Caio de A. Mendes‡

February 9, 2023

### Abstract

This work is largely focused on extending D. Higgs'  $\Omega$ -sets to the context of quantales, following the broad program of [7], we explore the rich category of  $\mathcal{Q}$ -sets for strong, integral and commutative quantales, or other similar axioms. The focus of this work is to study the different notion of “completeness” a  $\mathcal{Q}$ -set may enjoy and their relations, completion functors, resulting reflective subcategories, their relations to relational morphisms.

We establish the general equivalence of singleton complete  $\mathcal{Q}$ -sets with functional morphisms and the category of  $\mathcal{Q}$ -sets with relational morphisms; we provide two characterizations of singleton completeness in categorical terms; we show that the singleton complete categorical inclusion creates limits.

---

\*<mailto:alvim@ime.usp.br>

†<mailto:hugomar@ime.usp.br>

‡<mailto:caio.mendes@usp.br>

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries: Quantales</b>	<b>3</b>
<b>3</b>	<b>Preliminaries on <math>\mathcal{Q}</math>-Sets</b>	<b>6</b>
3.1	Some Examples . . . . .	10
3.2	The Underlying Graph Functor . . . . .	13
<b>4</b>	<b>Gluing Completeness</b>	<b>14</b>
4.1	Some Results about Gluings and Compatible Families . . . . .	18
4.2	Gluing-Completion . . . . .	19
<b>5</b>	<b>Singleton/Scott-Completeness</b>	<b>27</b>
5.1	Basics about Singletons . . . . .	27
5.2	Scott Completion . . . . .	33
<b>6</b>	<b>Connection between Completeness Conditions</b>	<b>40</b>
6.1	Scott-Completeness and Relational Morphisms . . . . .	41

# 1 Introduction

## History & Motivation

The notion of sheaf on a topological space depends only on the (complete) lattice of the open sets of the space, thus it is straightforward to define sheaves for “spaces without points”, that is, for a category  $(H, \leq)$  for a given locale  $H$  (see [3]).

In the 1970s, the topos of sheaves over a locale (= complete Heyting algebra)  $\mathbb{H}$  was described, alternatively, as a category of  $\mathbb{H}$ -sets [5]. More precisely, in [3], there were three categories whose objects were locale valued sets that are equivalent to the category of sheaves over a locale  $\mathbb{H}$ . Two different notions of separability and completeness have been proposed. On the one hand, the traditional notions of these properties in  $\mathbf{Sh}(\mathbb{H})$  can be translated to appropriate definitions in  $\mathbb{H}$ -sets. In addition, Scott’s notion of singletons, a definition that is inspired from the ordinary singleton set, leads alternative notions of completeness and separability. Despite some folkloric misconceptions about those notions, a simple counter-example (in a finite boolean algebra) shows that these two definitions of completeness are not logically equivalent – and do not give rise to equivalent full subcategories either.

This wealth of definitions and notions, however, has had the unfortunate effect of muddying any discussion concerning those kinds of objects. A veritably deep folklore has taken root in the field which hinders careful thought and frightens – almost to the point of panic – anyone who is paying attention.

Nevertheless, there is a non-commutative and non-idempotent generalization of locales called “quantales”, introduced by C.J. Mulvey [10]. Quantales show up in logic [17], and in the study of  $C^*$ -algebras [14].

Many notions of sheaf over a quantal and quantal-valued set are studied in many works ([16], [2], [mulveyquantale], [9], [8], [4] [7], [6], [12], [13], [15]). In many cases, the base quantales are *right-sided and idempotent*. Herein we continue our study of quantale-valued set on *commutative and semicartesian* quantales, initiated in [1]. Our approach is similar to the last one but, since every idempotent semicartesian quantale is a locale (Proposition 2.1), our axioms and theirs are orthogonal in some sense.

The goal of the present work is to examine these two notions of completeness (and separability): (i) via (unique) gluing of compatible families (gluing completeness of  $\mathcal{Q}$ -sets); (ii) via (unique) representation of (*strict*) singletons (Scott completeness). their relations and the properties of the full subcategories of their examples.

## Main results and the paper’s structure

We have shown that:

1. Scott-completeness implies gluing-completeness;

2. Both full subcategories of gluing-complete and Scott-complete  $\mathcal{Q}$ -sets with functional morphisms are reflective;
3. For “strong” [7] quantales, it makes sense to speak of “the Scott completion” of a given  $\mathcal{Q}$ -set;
4. Every  $\mathcal{Q}$ -set is relationally isomorphic to its own Scott completion, completion is invariant under functional isomorphisms;
5. For strong quantales, the categories of Scott-complete  $\mathcal{Q}$ -sets with relational morphisms and the one with morphisms are isomorphic.
6. For strong quantales, that  $X$  being Scott-complete is equivalent to the its functional representable functor “being the same” as its relational representable functor.
7. For strong quantales, that  $X$  being Scott-complete is equivalent to its representable functor being “invariant” under the completion endofunctor.
8. For strong quantales, that the full inclusion of Scott-complete  $\mathcal{Q}$ -sets not only preserves (due to it being a right adjoint) but also *creates* limits. And specifically, that limits of complete  $\mathcal{Q}$ -sets can be computed “pointwise”.

## 2 Preliminaries: Quantales

### Definition 2.1:

A *quantale* is a type of structure  $\mathcal{Q} = (|\mathcal{Q}|, \leq, \otimes)$  for which  $(|\mathcal{Q}|, \leq)$  is a complete lattice;  $(|\mathcal{Q}|, \otimes)$  is a semigroup<sup>1</sup>; and, moreover,  $\mathcal{Q}$  is required to satisfy the following distributive laws: for all  $a \in \mathcal{Q}$  and  $B \subseteq \mathcal{Q}$ ,

$$a \otimes \left( \bigvee_{b \in B} b \right) = \bigvee_{b \in B} (a \otimes b)$$

$$\left( \bigvee_{b \in B} b \right) \otimes a = \bigvee_{b \in B} (b \otimes a)$$

We denote by  $E \mathcal{Q}$  the subset of  $\mathcal{Q}$  comprised of its idempotent elements.

### Remark 2.1:

1. In any quantale  $\mathcal{Q}$  the multiplication is increasing in both entries;
2. Since  $\perp$  is also the supremum of  $\emptyset$ , for any  $a$ ,  $a \otimes \perp = \perp = \perp \otimes a$
3. Since  $\top$  is  $\sup \mathcal{Q}$ , then  $\top \otimes \top = \sup_{a,b} a \otimes b = \mathbf{img} \otimes$

---

<sup>1</sup>*i.e.* the binary operation  $\otimes : \mathcal{Q} \times \mathcal{Q} \rightarrow \mathcal{Q}$  (called multiplication) is associative.

**Remark 2.2:**

If  $(\mathcal{Q}, \leq)$  is a complete lattice for which the binary infimum satisfies the above distributive laws, the resulting quantale has  $\top$  as its unit and is – in fact – a locale. Conversely, every locale is a unital quantale in such a manner.

**Definition 2.2:**

A quantale  $\mathcal{Q}$  is said to be

- *bidivisible* when

$$a \leq b \implies \exists \lambda, \rho : a \otimes \rho = b = \lambda \otimes a$$

left (right) divisibility means to drop the  $\rho$  ( $\lambda$ ) portion of the axiom.


- *integral* when  $\top \otimes a = a = a \otimes \top$ . we say it's right-sided when the right equality holds, and left-sided when the left equality holds.
- *unital* when  $\otimes$  has a unit;
- *semicartesian* when  $a \otimes b \leq a \wedge b$
- *commutative* when  $\otimes$  is;
- *idempotent* when  $a \otimes a = a$ ;
- *linear and strict* when  $\leq$  is a linear order and for

$$a \neq \perp \text{ and } a \otimes b = a \otimes c \text{ or } b \otimes a = c \otimes a \implies b = c$$

- *strong* when for any  $e$  and  $A$  [7, cf. p. 30],

$$e = e \otimes e \implies e \leq \bigvee_{a \in A} a \implies e \leq \bigvee_{a \in A} a \otimes a$$

We offer the following diagram to explain some of the relations between those definitions:



**Example 2.1:**

Locales are – perhaps – the best example of quantales that are commutative, idempotent, integral (and hence both semicartesian and right-sided), divisible and strong (both trivially). Among which, and of special significance to Sheaf Theory, are the locales of open subsets of a topological space  $X$ , where the order relation is given by the inclusion, the supremum is the union, and the finitary infimum is the intersection.

**Example 2.2:**

We list below some examples of unital quantales that are not locales:

1. The extended half-line  $[0, \infty]$  with order the inverse order  $- \geq -$ , and the usual sum of real numbers as the multiplication.  
Since the order relation is  $\geq$ , the top element is 0 and the bottom elements is  $\infty$ . We call this the Lawvere quantale due to its relation to Lawvere spaces (related to metric spaces);
2. The extended natural numbers  $\mathbb{N} \cup \{\infty\}$ , as a restriction of the the Lawvere quantale (related to distance on graphs);
3. The set  $\mathcal{J}(R)$  of ideals of a commutative and unital ring  $R$  with order  $\subseteq$ , and the multiplication as the multiplication of ideals. The supremum is the sum of ideals, the top element is  $R$  and the trivial ideal is the bottom;
4. The set  $\mathcal{RJ}(R)$  of right (or left) ideals of an unital ring  $R$  with the same order and multiplication of the above example. Then the supremum and the top and the bottom elements are also the same of  $\mathcal{J}(R)$ ;
5. The set of closed right (or left) ideals of a unital  $C^*$ -algebra, the order is the inclusion of closed right (or left) ideals, and the multiplication is the topological closure of the multiplication of the ideals.

For more details and examples we recommend [14].

**Remark 2.3:**

The first three examples we introduced in 2.2 are commutative and integral quantales. The last two examples are neither commutative nor semicartesian. The forth is not idempotent but the fifth is, and both are right-sided (resp. left-sided) quantales [14, cf.].

The main examples of strong quantales are Heyting algebras,  $([0, 1], \leq, \cdot)$ , strict linear quantales. Some MV-Algebras, like the Chang's  $([0, 1], \wedge, \vee, \oplus, \otimes, 0, 1)$  [chang]'s, are not strong.

**From now on, we assume all quantales to be commutative.**

**Definition 2.3:**

Let  $\mathcal{Q}$  be a a quantale, we define an alternative partial order  $\preceq$  given by

$$a \preceq b \iff a = a \otimes b$$

**Remark 2.4:**

1. Let  $\mathcal{Q}$  be a unital quantale, then it is integral iff it is semicartesian;
2.  $a \preceq b \leq c$  implies  $a \preceq c$ ;
3. If  $e \in \mathbf{E} \mathcal{Q}$ ,  $(e \preceq a) \iff (e \leq a)$ .

**Proposition 2.1:**

If  $\mathcal{Q}$  is semicartesian and idempotent, it is in fact a complete distributive lattice and  $\otimes = \wedge$ . In other words, it is a locale.

*Proof.* Suppose  $\mathcal{Q}$  is in fact idempotent, we have – because  $\otimes$  is increasing in both arguments – that

$$a \leq b \implies a \leq c \implies a \leq b \otimes c$$

Hence, if  $a$  is less than both  $b$  and  $c$ , then it must be smaller than  $b \otimes c$ ; but since  $\mathcal{Q}$  is semicartesian, by remark 2.4 above,  $\otimes \leq \wedge$ . This means that  $b \otimes c$  is a lowerbound for  $b$  and  $c$ , but what we had gotten from idempotency means it's the greatest such upper bound.

Thus the multiplication satisfies the universal property of infima. The above is just a particular case of [11, Proposition 2.1].  $\square$

### 3 Preliminaries on $\mathcal{Q}$ -Sets

**Remark 3.1:**

Hereon we are working exclusively with commutative semicartesian quantales, as opposed to ones without those properties.

Given a quantale  $\mathcal{Q}$ , one may form – roughly speaking – a  $\mathcal{Q}$ -space, wherein distances between points are measured by elements of  $\mathcal{Q}$  as opposed to – say –  $[0, \infty)$  as we often do. This definition is made precise in the notion of a  $\mathcal{Q}$ -set.

**Definition 3.1:**

A  $\mathcal{Q}$ -set is a set endowed with a  $\mathcal{Q}$ -distance operation usually denoted by  $\delta$ . *i.e.* a  $\mathcal{Q}$ -set is a set  $X$  with a map  $\delta : X^2 \rightarrow X$  satisfying:

1.  $\delta(x, y) = \delta(y, x)$ ;
2.  $\delta(x, y) \otimes \delta(y, z) \leq \delta(x, z)$ ;
3.  $\delta(x, x) \otimes \delta(x, y) = \delta(x, y)$ .

and it is usual to denote  $\delta(x, x)$  by simply the “extent of  $x$ ” written as  $\mathbf{E}x$ .

A couple of things might jump out to reader in the above definition. (i)  $\delta$  is symmetric, even though we have thrown out all but the vaguest notions

tying ourselves to metric spaces; (ii) Why is the triangle inequality upside down? (iii)  $E x \otimes \delta(x, y) = \delta(x, y)$ , why not just ask that  $E x = \top$ ?

Those questions are all valid – and answering the first and last ones differently has been done in the past and is the main difference between  $\mathcal{Q}$ -sets and  $\mathcal{Q}$ -enriched categories from a definitional perspective. The question of order being inverse is more one of sanity: since we treat a  $\mathcal{Q}$ -set as a set with  $\mathcal{Q}$ -valued equality, it makes sense to think that  $E x$  is the maximally valid equality to  $x$  and hence the triangular inequality needs to be turned upsidedown – and turned into the transitivity of equality.

**Remark 3.2:**

When we speak of properties of the type  $P(\vec{x}) \leq Q(\vec{x})$  in  $\mathcal{Q}$ -sets, it is often more insightful to think that the logically equivalent (but notationally less helpful) statement

$$P(\vec{x}) \rightarrow Q(\vec{x}) (= \top)$$

There are two main category structures that one can canonically endow the collection of all  $\mathcal{Q}$ -sets with. One is taking maps to be co-contractions (*i.e.* they make  $\delta$  bigger) – the other is to consider well behaved  $\mathcal{Q}$ -valued relations between the underlying sets.

**Definition 3.2:**

A functional morphism  $f : X \rightarrow Y$  is a function  $f$  between the underlying sets of  $X$  and  $Y$  such that  $f$  increases  $\delta$  and preserves  $E$ ; that is to say

$$\begin{aligned} \delta_X &\leq \delta_Y \circ (f \times f) \\ E_X &= E_Y \circ f \end{aligned}$$

A relational morphism  $\varphi : X \rightarrow Y$  is a function  $\varphi : |X| \times |Y| \rightarrow \mathcal{Q}$  satisfying

$$\begin{aligned} \delta(x, x') \otimes \varphi(x, y) &\leq \varphi(x', y) \\ \varphi(x, y) \otimes \delta(y, y') &\leq \varphi(x, y') \\ \varphi(x, y) \otimes \varphi(x, y') &\leq \delta(y, y') \\ \varphi(x, y) \otimes E y &= \varphi(x, y) \\ E x \otimes \varphi(x, y) &= \varphi(x, y) \\ \bigvee_{y \in Y} \varphi(x, y) &= E x \end{aligned}$$

The reader should beware we don't often distinguish between  $\delta_X$  and  $\delta_Y$  and instead rely on suggestively named variables so as to indicate their type and hence the  $\delta$  they refer to. In other words, the reader is expected to be familiar with Koenig lookup<sup>2</sup>.

We denote by  $\mathcal{Q}\text{-Set}_r$  the category  $\mathcal{Q}$ -sets and relational morphisms between them and by  $\mathcal{Q}\text{-Set}_f$  the category with the same objects but functional morphisms between them instead.

<sup>2</sup> Which, to quote a great website – [cpreference.com](http://cpreference.com) –



**Proposition 3.1:**

It should be observed that both notions of morphism actually do form a category. For functional morphisms, we take composition to be the usual function composition – since functional morphisms are obviously closed under it – and the identity function as the identity. For relational morphisms, the identity on  $X$  becomes  $\delta_X$  and composition is the (perhaps obvious) relational composition:

$$[\psi \circ \varphi](x, z) = \bigvee_{y \in Y} \varphi(x, y) \otimes \psi(y, z)$$

Since proving that functional morphisms do indeed form a category would be trivial, we shall instead prove a stronger result afterwards.

*Proof.* Firstly we should observe that  $\delta$  is indeed a relational morphism. This is easy enough, as the first three axioms are direct applications of the triangular inequality, the fourth and fifth are direct applications of the extension axiom of  $\mathcal{Q}$ -sets, and the last one is trivially true once we realize that

$$\delta(x, y) \leq \mathbb{E}x = \delta(x, x)$$

Once we have that  $\delta$  is indeed a morphism, we can wonder about the action of (pre-)composing with it. In which case we obtain

$$\begin{aligned} [\delta \circ \varphi](x, y) &= \bigvee_{y' \in Y} \varphi(x, y') \otimes \delta(y', y) \\ &\geq \varphi(x, y) \otimes \delta(y, y) \\ &= \varphi(x, y) \end{aligned}$$

on the other hand, we could have applied the appropriate relational morphism axiom inside the  $\bigvee$  thus getting that the composite was smaller than  $\varphi(x, y)$  instead. This proves they were in fact equal all along. The same goes for the other composite:

$$\begin{aligned} [\varphi \circ \delta](x, y) &= \bigvee_{x' \in X} \delta(x, x') \otimes \varphi(x', y) \\ &\geq \delta(x, x) \otimes \varphi(x, y) \\ &= \varphi(x, y) \end{aligned}$$

Hence it only remains to see that  $\circ$  as defined for relational morphisms is indeed a composition, in that it is associative and that the composite of two morphisms is a morphism. Firstly suppose that  $\varphi$  and  $\psi$  are composable

---

Argument-dependent lookup, also known as ADL, or Koenig lookup, is the set of rules for looking up the unqualified function names in function-call expressions, including implicit function calls to overloaded operators.

morphisms – we ought to show that their composite is also a morphism:

$$\begin{aligned}
[\psi \circ \varphi](x, z) \otimes \delta(z, z') &= \bigvee_y \varphi(x, y) \otimes \psi(y, z) \otimes \delta(z, z') \\
&\leq \bigvee_y \varphi(x, y) \otimes \psi(y, z') \\
&= [\psi \circ \varphi](x, z')
\end{aligned}$$

And likewise, *mutatis mutandis* one can show that axioms 1-2 and 4-5 hold. Axiom 6 can be seen to hold easily as well:

$$\begin{aligned}
\bigvee_z [\psi \circ \varphi](x, z) &= \bigvee_y \varphi(x, y) \otimes \bigvee_z \psi(y, z) \\
&= \bigvee_y \varphi(x, y) \otimes \mathbf{E} y \\
&= \bigvee_y \varphi(x, y) \\
&= \mathbf{E} x
\end{aligned}$$

Axiom 3 actually requires commutativity, which is unfortunately the first of many times it is very necessary to make use of it;

$$\begin{aligned}
[\psi \circ \varphi](x, z) \overset{\psi \circ \varphi}{\otimes} (x, z') &= \bigvee_y \bigvee_{y'} \varphi(x, y) \otimes \psi(y, z) \otimes \varphi(x, y') \otimes \psi(y', z) \\
&= \bigvee_y \bigvee_{y'} \psi(y, z) \otimes \varphi(x, y) \otimes \varphi(x, y') \otimes \psi(y', z) \\
&\leq \bigvee_y \bigvee_{y'} \psi(y, z) \otimes \delta(y, y') \otimes \psi(y', z) \\
&\leq \bigvee_y \bigvee_{y'} \psi(y, z) \otimes \psi(y, z) \\
&\leq \bigvee_y \bigvee_{y'} \delta(z, z') = \delta(z, z')
\end{aligned}$$

Associativity can obviously be seen to hold when we realize that one can rearrange the terms of  $(\psi \circ \varphi) \circ \chi$  and  $\psi \circ (\varphi \circ \chi)$  to be in the form of

$$\bigvee_{x, y} \chi(w, x) \otimes \varphi(x, y) \otimes \psi(y, z)$$

□

**Definition 3.3:**

Instead of proving the category axioms for functional morphisms we promised to prove a stronger result – which is incidently useful for another paper of ours

in the works – which is to prove that  $e$ -morphisms form a category (given a generic commutative unital quantale) and that functional morphisms form a wide subcategory of  $I$ -morphisms.

So, let us define  $e$ -morphisms: given an idempotent element  $e$  of  $\mathcal{Q}$ , a  $e$ -morphism is a functional morphism “up to error  $e$ ”:

$$e \otimes \delta(x, x') \leq \delta(f(x), f(x'))$$

$$E f(x) = e \otimes E f(x)$$

**Proposition 3.2:**

We claim that the collection of  $\langle e, \varphi \rangle$  where  $\varphi$  is an  $e$ -morphism constitutes a category under the obvious composition laws. Furthermore, the identity function is a  $I$ -morphism where  $I$  is the unit of the quantale, and further still:  $I$ -morphisms are closed under composition and form a subcategory which is definitionally equal to  $\mathcal{Q}\text{-Set}_f$ .

*Proof.* Firstly, the obvious composition takes an  $e$ -morphism  $f$  and an  $e'$ -morphism  $g$  to a  $(e \otimes e')$  morphism  $g \circ f$ . Associativity is due to functional (in **Set**, that is)  $\circ$  associativity and the fact that  $\otimes$  makes  $\mathcal{Q}$  a semigroup. The fact that  $g \circ f$  is a  $(e \otimes e')$ -morphism is rather obvious and the proof is omitted.

The identity is evidently a  $I$ -morphism – and of course that composing  $I$ -morphisms gives a  $I \otimes I = I$ -morphism.  $\square$

### 3.1 Some Examples

**Example 3.1** (Initial Object):

The empty set is – vacuously – a  $\mathcal{Q}$ -set, and since morphisms are functions, it also happens to be the initial object.

**Example 3.2** (Terminal Object):

The set of idempotent elements of  $\mathcal{Q}$ , denoted  $E\mathcal{Q}$  naturally has a structure of a  $\mathcal{Q}$ -set – given by  $\delta(a, b) = a \otimes b = a \wedge b^3$ . It is trivial to see that  $\otimes$  satisfies all  $\mathcal{Q}$ -set laws. More interestingly, however,  $E\mathcal{Q}$  must be the terminal object because  $Ee = e$ : since functional morphisms preserve extents, one has that  $E f(x) = E x$  – however,  $E f(x) = f(x)$  and thus  $f(x) = E x$ . This proves that there is at most one morphism  $X \rightarrow E\mathcal{Q}$ .

On the other hand,  $\delta(x, y) \leq E x \wedge E y$  which just happens to make  $x \mapsto E x$  a functional morphism. And thus,  $E\mathcal{Q}$  is a  $\mathcal{Q}$ -set.

**Remark 3.3:**

One cannot use the whole of  $\mathcal{Q}$  in  $E\mathcal{Q}$ 's stead, as  $\delta(x, y) \otimes E x$  would not hold. The only reason it holds in the above example is because for idempotents  $\otimes = \wedge$ . However, one can obtain a  $\mathcal{Q}$ -set where the underlying set is  $\mathcal{Q}$  itself

---

<sup>3</sup>This is the infimum in the subposet  $E\mathcal{Q}m$  that turns out to be a locale.

**Example 3.3:**

Much akin to how Lawvere’s quantale is a Lawvere space, a (integral and commutative) quantale  $\mathcal{Q}$  is a  $\mathcal{Q}$ -set. This is achieved with the following:

$$\delta(x, y) = (x \rightarrow y) \wedge (y \rightarrow x)$$

which is roughly equivalent to  $|x - y|$  for real numbers. This isn’t necessarily the best  $\mathcal{Q}$ -set structure we can give them, as  $\mathbb{E} x = \top$  for any  $x$ .

Ways to mitigate this phenomenon, which is specially strange for  $\perp$ , involve taking into account idempotents above  $x$ . An important quantallic property is the existence of an operation  $(\_)\text{-}$  taking an element  $x$  to the value  $\sup\{e \in \mathbb{E} \mathcal{Q} \mid e \preceq x\}$ . Multiplying  $\delta(x, y)$  by  $x\text{-} \otimes y\text{-}$  guarantees – for instance – that the above construction coincides with the terminal object when  $\mathcal{Q}$  is a locale.

Another way to correct this, is to incorporate  $\mathbb{E} \mathcal{Q}$  more directly, considering the space with underlying set  $\mathcal{Q} \times \mathbb{E} \mathcal{Q}$  and  $\delta$  given by

$$\delta((x, e), (y, a)) = a \otimes e \otimes [(x \rightarrow y) \wedge (y \rightarrow x)]$$

We write this  $\mathcal{Q}$ -set as  $\mathcal{Q}_{\mathbb{E}}$ .

**Example 3.4:**

A construction that is explored in this work’s sister-article [1] but deserves to be mentioned here in passing  $X \boxtimes X$ , given by the underlying set  $|X| \times |X|$  and with  $\delta$  given by the product of the  $\delta$  of the coordinates. The reason this construction is relevant here is because  $\delta : X \boxtimes X \rightarrow \mathcal{Q}_{\mathbb{E}}$  in a natural way:

$$(x, y) \mapsto (\delta(x, y), \mathbb{E} x \otimes \mathbb{E} y)$$

And this happens to be a functional morphism.

**Remark 3.4:**

Monomorphisms are always injective functions, and epimorphisms are always surjective.

**Example 3.5 (Regular Subobjects):**

A monomorphism is regular when it is an equalizer of a pair of parallel arrows. Suppose  $f, g : A \rightarrow B$ , one way to conceive of an equalizer is as being the maximal subobject of  $A$  making the diagram commute. It is quite trivial to see that subobjects in general *must* be subsets with a point-wise smaller  $\delta$ . Hence, the largest subobject of  $A$  that equalizes the pair is simply the subset of  $A$  where the functions agree on, with the largest possible  $\delta$ : that being  $A$ ’s.

Hence, we see a pattern where equalizers – in general – are simply subsets with  $\delta$  coming from a restriction of the original object. Importantly, though, we refer to “regular subobjects” as monomorphisms that preserve  $\delta$  – as they have been equivalently characterized.

The skeptical reader might question that we have merely shown that regular monos preserve  $\delta$ , as opposed to showing such to be a sufficient condition.

In which case, given the fact that monos are injective functions,  $\delta$ -preserving monos are simply subsets with its superset's  $\delta$ ; consider one such mono:  $m : A \hookrightarrow X$ , with  $A \subseteq X$ , one takes  $(X \amalg X) / \sim$  with  $\sim$  defined so as to identify both copies of  $A$ .

$$\delta(\llbracket(x, i)\rrbracket, \llbracket(y, j)\rrbracket) = \begin{cases} \delta(x, y), & i = j \\ \bigvee_{a \in A} \delta(x, a) \otimes \delta(a, y), & i \neq j \end{cases}$$

There are two obvious inclusions of  $X$  into this set, namely the upper branch and the lower branch. And they coincide exactly on the section corresponding to  $A$ , so the equalizer of those two arrows must be the subset inclusion of  $A$  into  $X$ , the  $\delta$  must be the biggest possible, which is the restriction of  $X$ 's so as to remain a morphism.

**Example 3.6:**

Suppose  $(X, d)$  is a pseudo-metric space, then  $(X, d)$  is a  $[0, \infty]$ -set where  $\delta(x, y) \neq \perp = \infty, \forall x, y \in X$ .

**Example 3.7:**

Given a nonempty index set  $I$ , we have a  $\mathcal{Q}$ -set  $\coprod_{i \in I} \top$ , given by

$$\delta((e, i), (e', i')) = \begin{cases} e \wedge e', & \text{if } i = i'; \\ \perp, & \text{otherwise.} \end{cases}$$

**Example 3.8:**

Given a commutative ring  $A$ , let the set of its (left) ideals be denoted  $\mathcal{I}_A$ .  $\mathcal{I}_A$  is a quantale. Given a left  $A$ -module  $M$ , we can endow it with the structure of a  $\mathcal{I}_A$ -set:

$$\delta(x, y) = \bigvee \{I \in \mathcal{I}_A \mid I \cdot x = I \cdot y\}$$

In fact, that supremum is attained at with a particular ideal. Moreover,  $Ex = A = \max \mathcal{I}_A$ .

**Example 3.9:**

Suppose that  $\mathcal{Q}$  is a quantale with ‘‘idempotent upper approximations’’<sup>4</sup>:

$$\forall q \in \mathcal{Q} : \exists q^+ \in \mathbf{E} \mathcal{Q} : q^+ = \min \{e \in \mathbf{E} \mathcal{Q} \mid q \preceq e\}$$

Then

$$\delta(x, y) = \begin{cases} x \otimes y, & x \neq y; \\ x^+, & x = y. \end{cases}$$

defines a  $\mathcal{Q}$ -set structure on  $\mathcal{Q}$  itself.

---

<sup>4</sup>In [15] are described sufficient conditions for  $\mathcal{Q}$  to have such a property.

## 3.2 The Underlying Graph Functor

### Definition 3.4:

There is a functor  $\mathfrak{G}_R : \mathcal{Q}\text{-Set}_f \rightarrow \mathcal{Q}\text{-Set}_r$  which is the underlying *graph* functor, because it takes functional morphisms to their graph relation. More precisely,  $\mathfrak{G}_R(X) = X$  and given  $f : X \rightarrow Y$

$$(\mathfrak{G}_R f)(x, y) = \delta(f(x), y)$$

### Proposition 3.3:

As defined above,  $\mathfrak{G}_R f$  is indeed a functional morphism and  $\mathfrak{G}_R$  is indeed a functor.

*Proof.* It is clear, from  $f$  being a functional morphism, that  $\mathfrak{G}_R f$  should at least satisfy the  $\delta$  and E axioms for relational morphism axioms; the  $(\Sigma)$  axiom holds because of the triangular inequality; the strictness axiom holds as taking  $b = f(a)$  gives E  $f(a)$  which is E  $a$ .

Regarding functoriality,  $\mathfrak{G}_R \text{id} = \delta$  and hence the identity in the relational category; moreover,

$$\begin{aligned} (\mathfrak{G}_R g) \circ (\mathfrak{G}_R f)(x, z) &= \bigvee_y (\mathfrak{G}_R f)(x, y) \otimes (\mathfrak{G}_R g)(y, z) \\ &= \bigvee_y \delta(f(x), y) \otimes \delta(g(y), z) \\ &= \delta(g \circ f(x), z) \\ &= \mathfrak{G}_R(g \circ f) \end{aligned}$$

□

## Interlude

In the general course of studying  $\Omega\text{-Set}$  for a frame/locale  $\Omega$  one is bound to eventually come across a certain notion called “completeness”. Completeness is, largely speaking, a property necessary to transition between  $\Omega\text{-Set}$ -world to  $\mathbf{Sh}\Omega$ -world [3, prop. 2.7.10, cf. def. 2.9.1]. This notion of completeness we call Scott-completeness and refers to a thing called singletons.

Sheaves, almost by definition, are about gluing partial data – in that morally speaking, compatible elements over a covering admit exactly one gluing which lies over the covered open. This gives a second natural notion of completeness – which we have come to call Gluing-completeness.

In these next sections, therefore, we shall give the precise definitions pertaining to both notions, make explicit some of the relations between them.

## 4 Gluing Completeness

**Definition 4.1** (Compatible Family):

Given a  $\mathcal{Q}$ -set  $X$ , a family  $A$  of its elements is said to be *compatible* when  $\forall a, b \in A$ ,

$$\delta(a, b) = E a \otimes E b$$

naturally, since extents are idempotent, and for idempotents  $\otimes = \wedge$ , this is the same as saying  $\delta(a, b) = E a \wedge E b$ .

Note that the empty family of a  $\mathcal{Q}$ -set is vacuously compatible.

In light of our usual interpretation for  $\delta$  being ‘the place where two fragments of data agree’, this says that  $a$  and  $b$  agree in the intersection of their extents – their domains of definition and degrees of certainty, so to speak. Hence, they are compatible.

**Definition 4.2** (Gluing Element):

Given a  $\mathcal{Q}$ -set  $X$  and a family of its elements  $A$ , we say that  $x_A \in X$  is a *gluing of  $A$*  or that *it glues  $A$*  when  $\forall a \in A$ ,

$$\delta(a, x_A) = E a$$

$$E x_A = \bigvee_{a \in A} E a$$

This definition then captures the idea that an amalgamation of local data agrees integrally with the data it aggregates, and – importantly – does not make choices or provide information not present in the family  $A$  it glues.

**Proposition 4.1:**

If a family  $A$  admits a gluing element, it is compatible. Moreover, any gluing elements are “morally unique” – that is to say, they are  $\delta$ -equivalent.

*Proof.* Suppose  $x_A$  glues  $A$ ; then let  $a$  and  $b$  be members of the family  $A$ ; observe that

$$\delta(x_A, a) = E a$$

$$\delta(x_A, b) = E b$$

hence

$$E a \otimes E b = \delta(x_A, a) \otimes \delta(x_A, b)$$

but then,

$$E a \otimes E b \leq \delta(a, b)$$

but we know the following always holds

$$\delta(a, b) \leq E a \otimes E b$$

therefore they are the same.

This suffices for compatibility, now for the second claim: consider  $y \in X$  and let us compare two possible gluings – say  $x$  and  $x'$ .

$$\begin{aligned}
\delta(x, y) &= \delta(x, y) \otimes \mathbf{E} x \\
&= \bigvee_{a \in A} \delta(x, y) \otimes \mathbf{E} a \\
&= \bigvee_{a \in A} \delta(x, y) \otimes \mathbf{E} a \otimes \mathbf{E} a \\
&= \bigvee_{a \in A} \delta(x, y) \otimes \delta(x', a) \otimes \delta(x, a) \\
&\leq \bigvee_{a \in A} \delta(x, y) \otimes \delta(x', x) \\
&\leq \delta(x', y)
\end{aligned}$$

since the same holds swapping  $x$  for  $x'$  and vice versa, it follows that  $\delta(x, y) = \delta(x', y)$  for *any*  $y$ ; hence, they are essentially the same.  $\square$

**Definition 4.3** ( $\delta$ -equivalence):

The above property is somewhat important, as – as often is the case – equality is *evil* and equivalence is *good*. Two elements  $x, y$  of a  $\mathcal{Q}$ -set  $X$  are said to be  $\delta$ -*equivalent* whenever either of the following equivalent conditions hold:

$$\begin{aligned}
\forall z \in X : \delta(x, z) &= \delta(y, z) \\
\delta(x, y) &= \mathbf{E} x = \mathbf{E} y
\end{aligned}$$

That the former implies the latter is rather obvious, simply substitute  $z$  for  $y$  and for  $x$ . That the latter yields the former is also simple to show:

$$\begin{aligned}
\delta(x, z) &= \delta(x, z) \otimes \mathbf{E} x \\
&= \delta(x, z) \otimes \delta(x, y) \\
&\leq \delta(y, z)
\end{aligned}$$

and *mutatis mutandis* one obtains the other required inequality.

**Remark 4.1:**

It is evident that this is an equivalence relation.

Two  $\delta$ -equivalent points cannot – for instance – be taken to non-equivalent points via a functional morphism; and relational morphisms can't distinguish them. This is codified in the following lemma



**Lemma 4.1:**

The relation of  $\delta$ -equivalence is ‘congruential’ for either type of morphism. In more precise terms: provided with equivalent elements  $x$  and  $x'$ ;

1. For functional morphisms, congruential means  $f(x) \sim f(x')$ ;
2. For relational morphisms, it means that both  $\varphi(x, y) = \varphi(x', y)$  and  $\psi(w, x) = \psi(w, x')$ .

*Proof.*

$$\begin{aligned} \mathbb{E} x = \mathbb{E} x' &= \delta(x, x') \leq \delta(f(x), f(x')) \\ &\leq \mathbb{E} f(x) = \mathbb{E} x \\ &= \mathbb{E} x' = \mathbb{E} f(x') \end{aligned}$$

As for relational morphisms,

$$\begin{aligned} \varphi(x, y) &= \varphi(x, y) \otimes \mathbb{E} x \\ &= \varphi(x, y) \otimes \delta(x, x') \\ &\leq \varphi(x' y) \end{aligned}$$

The dual inequality holds for the same reasons; and similarly, the same equality can be obtained for  $\psi$ .  $\square$

We can state the gluing completeness as follows

**Definition 4.4** (Gluing Completeness):

A  $\mathcal{Q}$ -set  $X$  is said to be *gluing complete* when every compatible family has exactly one gluing. This is equivalent to asking that all compatible families admit a gluing and for  $\delta$ -equivalence to be “extensionally equal to equality” – which sounds very pretentious.

The equivalence can be seen to be true by realizing that  $\{x, x'\}$  is glued by both  $x$  and  $x'$  if and only if they are  $\delta$ -equivalent.

**Definition 4.5** (Extensionality):

Extensionality is the name of the property of  $\delta$ -equivalence being extensionally equal to equality. That is, to say:

$$\delta(x, y) = \mathbb{E} x = \mathbb{E} y \implies x = y$$

or equivalently

$$\delta(\_, x) = \delta(\_, y) \implies x = y$$

**Example 4.1:**

It is quite easy to see that the terminal  $\top$   $\mathcal{Q}$ -set is gluing complete. Given that  $\mathbb{E} a \otimes \mathbb{E} b = \delta(a, b)$  always holds, any set of idempotent elements is compatible. Moreover, the gluing of such a set is its supremum, as can easily be seen:

$$x_A = \mathbb{E} x_A = \bigvee_{a \in A} \mathbb{E} a = \bigvee_{a \in A} a$$

**Example 4.2:**

The initial object, the empty set, is trivially extensional but it fails to satisfy the gluing condition as the empty family is also trivially *compatible* and there is no element in the empty set that actually glues it. Hence, completeness doesn't hold, whereas extensionality does.

**Example 4.3:**

In an extended pseudometric space  $(X, d)$  seen as a  $[0, \infty]$ -set, points at infinity (with minimal extent) are always compatible with every other point, and normal points are compatible if and only if they are morally the same (ie. their distance is 0).

Hence, one such extended pseudometric space satisfies the gluing condition if and only if it has a point at infinity, and is extensional if and only if it is morally a metric space (ie. it is a metric space if considered sans points at infinity) and it has exactly one point at infinity.

**Example 4.4:**

For nontrivial index sets  $I$  (namely: with more than one element) and  $\mathcal{Q}$ -sets  $X_i$  with elements  $\perp_i$  at infinity (null-extensional)  $\coprod_{i \in I} X$  is never extensional, as  $\delta((\perp_i, i), (\perp_j, j)) = \perp$  for multiple different  $\perp_i$ . In general, any  $\mathcal{Q}$ -set with more than exactly one element with null extent cannot be extensional.

**Example 4.5:** 1. If  $A$  is a commutative ring and  $M$  is a left  $A$ -module, then the set over  $\mathcal{Q} = Ideals(A)$ ,  $X_M = (M, \delta)$ , is such that for each  $x, y \in M$ ,  $x \sim_\delta y$  iff  $A.x = A.y$ , thus it is not an extensional  $\mathcal{Q}$ -set, in general: any member of a non-empty compatible family in  $(M, \delta)$  is a gluing for that family. Moreover,  $M$  is a divisible  $A$ -module iff  $\sim_\delta = \{(0, 0)\} \cup (M \setminus \{0\}) \times (M \setminus \{0\})$ .

2. If  $card(I) > 1$ , then the  $\mathcal{Q}$ -set  $I.1_{\mathcal{Q}}$ -set is not extensional, since  $\delta((\perp, i), (\perp, j)) = \perp$  and there are  $i, j \in I$  such that  $i \neq j$ . If  $I' \subseteq I$ , then the family  $\{(\perp, i) : i \in I'\}$  is compatible and, for any  $j \in I$ ,  $(\perp, j)$  is a gluing for that family.

3. If  $\mathcal{Q}$  is a integral quantale with upper idempotent approximation, then the associated  $\mathcal{Q}$ -set on  $\mathcal{Q}$  is extensional. A family  $S$  on that  $\mathcal{Q}$ -set is compatible iff for each  $x, y \in S$ ,  $x \otimes y = x^+ \otimes y^+$ .

**Proposition 4.2:**

The functor  $\mathcal{G}_r$  is faithful exactly on those pairs of objects where the codomain is extensional. This is to say: if  $Y$  is extensional, then the following map is injective for all  $X$

$$\mathcal{Q}\text{-Set}_f(X, Y) \xrightarrow{\mathcal{G}_r} \mathcal{Q}\text{-Set}_r(X, Y)$$

Moreover, extensional  $\mathcal{Q}$ -sets are the largest class of objects such that you can quantify universally over  $X$ .

*Proof.* First let's prove the latter claim: let  $Y$  be any  $\mathcal{Q}$ -set; and suppose  $y$  and  $y'$  are  $\delta$ -equivalent. Now let  $X = \{*\}$ , setting  $E* = E y$ .

There are two obvious maps  $X \rightarrow Y$ , namely  $* \mapsto y$  and  $* \mapsto y'$ . Call them  $f$  and  $g$  respectively. By the equivalent definition of  $\delta$ -equivalence,  $y$  and  $y'$  have the same  $\delta$ s, so

$$\mathfrak{G}_{\mathbb{R}}(* \mapsto y) = \delta(\_, y) = \delta(\_, y') = \mathfrak{G}_{\mathbb{R}}(* \mapsto y')$$

And hence, if  $y \neq y'$ , then  $\mathfrak{G}_{\mathbb{R}}$  must send different functional morphisms to the same relational one and therefore fail injectivity.

Now, that being said we still must prove that the functor is injective on **homs** when the codomain is extensional. But this is simple. Suppose  $\mathfrak{G}_{\mathbb{R}}(f) = \mathfrak{G}_{\mathbb{R}}(g)$  – and thus that for all  $x$  and  $y$ ,

$$\delta(f(x), y) = \delta(g(x), y)$$

by taking  $y$  to be  $f(x)$  we get that  $\delta(g(x), f(x)) = E f(x)$  and *mutatis mutandis* the same can be done to show that  $f(x)$  is always  $\delta$ -equivalent to  $g(x)$ . Because  $Y$  is extensional we have that  $g(x) = f(x)$  for all  $x$  and therefore they are they one and the same functional morphism.  $\square$

**Remark 4.2:**

Completeness, be it gluing-wise or singleton-wise, is a rather pointless notion in the category of relational morphisms; this is because every  $\mathcal{Q}\text{-Set}_r$  is equivalent to its full subcategories whose objects are complete in either notion.

Hence, any categorical property we *could ever prove* about those subcategories would also be true of the larger relational category.

In fact, completeness can – in a way – be understood as the property one can impose on objects so as to make their functional morphisms representable functor “the same” as their relational morphisms representable functor. This will be elaborated in a subsequent section.

In the light of the remark above, we shall focus ourselves on the functional morphisms category, since talking about complete  $\mathcal{Q}$ -sets under relational morphisms is just a roundabout way of talking about general  $\mathcal{Q}$ -sets under those same morphisms.

## 4.1 Some Results about Gluings and Compatible Families

**Lemma 4.2:**

Family compatibility is preserved by direct images of functional morphisms.

*Proof.* Given  $A$  compatible for  $X$  and  $f : X \rightarrow Y$ ; take any  $a, b \in A$ ; it suffices to show that  $\delta(f(a), f(b)) = E f(a) \otimes E f(b)$ .

$$\begin{aligned}
\delta(a, b) &\leq \delta(f(a), f(b)) \\
&= \delta(f(a), f(b)) \otimes E f(a) \\
&= \delta(f(a), f(b)) \otimes E f(a) \otimes E f(b) \\
&= \delta(f(a), f(b)) \otimes \delta(a, b) \\
&\leq \delta(a, b) \qquad \text{(semi-cartesian)}
\end{aligned}$$

□

**Lemma 4.3:**

Gluing elements are preserved by direct images of functional morphisms

*Proof.* Should  $x$  glue  $A$ , we already know that  $f[A]$  is compatible – in light of the previous lemma and the fact that a family that admits a gluing must be compatible – so it merely suffices to show that  $f(x)$  glues  $f[A]$ . Take  $a \in A$ ,

$$\begin{aligned}
E a &= \delta(x, a) \\
&\leq \delta(f(x), f(a)) \\
&\leq E f(a) = E a
\end{aligned}$$

Thus establishing the first condition.

$$\begin{aligned}
E f(x) &= E x \\
&= \bigvee_{a \in A} E a \\
&= \bigvee_{a \in A} E f(a) \\
&= \bigvee_{\alpha \in f[A]} E \alpha
\end{aligned}$$

Thus completing the proof. □

We shall see that there is a natural way of ‘making’ a  $\mathcal{Q}$ -set gluing complete. This what we shall focus ourselves on in the next subsection.

## 4.2 Gluing-Completion

Completion problems often come in the shape of “all  $X$  induce a  $Y$ , but not all  $Y$  come from an  $X$  – can we make it so  $X$  and  $Y$  are in correspondence?” and the solution to those kind of problems can often be found in looking at the collection of all  $Y$ s or quotients of it.

In that spirit, we shall give the collection of all compatible families the structure of a  $\mathcal{Q}$ -set; we shall embed the original  $\mathcal{Q}$ -set in it; and then we “throw away” redundant points in an appropriate quotient.

**Definition 4.6:**

Given a  $\mathcal{Q}$ -set  $X$ , we say  $\mathfrak{G}(X)$  is the  $\mathcal{Q}$ -set given by

$$|\mathfrak{G}(X)| = \{A \subseteq |X| : A \text{ is compatible}\}$$

$$\delta(A, B) = \bigvee_{\substack{a \in A \\ b \in B}} \delta(a, b)$$

Obviously, we claim that the above does indeed define a  $\mathcal{Q}$ -set.

*Proof.*

$$\begin{aligned} \delta(A, B) \otimes \delta(B, C) &= \bigvee_{a \in A} \bigvee_{b \in B} \bigvee_{\beta \in B} \bigvee_{c \in C} \delta(a, b) \otimes \delta(\beta, c) \\ &= \bigvee_{a \in A} \bigvee_{b \in B} \bigvee_{\beta \in B} \bigvee_{c \in C} \delta(a, b) \otimes \mathbf{E} b \otimes \mathbf{E} \beta \otimes \delta(\beta, c) \\ &= \bigvee_{a \in A} \bigvee_{b \in B} \bigvee_{\beta \in B} \bigvee_{c \in C} \delta(a, b) \otimes \delta(b, \beta) \otimes \delta(\beta, c) \\ &\leq \bigvee_{a \in A} \bigvee_{c \in C} \delta(a, c) \\ &= \delta(A, C) \end{aligned}$$

Which suffices for transitivity/triangular inequality. It is obvious that the definition is symmetric, so it only remains to show that extents behave as they should.

$$\begin{aligned} \delta(A, A) \otimes \delta(A, B) &= \bigvee_{a, a' \in A} \bigvee_{\alpha \in A, b \in B} \delta(a, a') \otimes \delta(\alpha, b) \\ &\geq \bigvee_{\alpha \in A, b \in B} \delta(\alpha, \alpha) \otimes \delta(\alpha, b) \\ &= \delta(A, B) \end{aligned}$$

and thus

$$\mathbf{E} A \otimes \delta(A, B) = \delta(A, B) \quad (\text{semi-cartesian})$$

Thus proving it is actually a  $\mathcal{Q}$ -set as desired.  $\square$

**Definition 4.7** (The  $\mathcal{Q}$ -set of Compatible Families):

We extend  $\mathfrak{G}^5$  from an object-map to a dignified functor, by taking functional morphisms to their direct image; this has the surreptitious implicit premise that direct images are indeed functional morphisms between the compatible families  $\mathcal{Q}$ -sets.

---

<sup>5</sup>Here it seems necessary to justify the use of a fraktur typeface; Capital  $\mathbf{G}$  is fine because we are taking spaces with all possible  $\mathbf{G}$ luings. But since they aren't extensional, our functor is still a bit 'broken'. You can call it a mnemonic, but it's a visual pun.

*Proof.*

$$\begin{aligned}\delta(A, B) &= \bigvee_{a,b} \delta(a, b) \\ &\leq \bigvee_{a,b} \delta(f(a), f(b)) \\ &= \delta(f[A], f[B])\end{aligned}$$

For extents it's even more immediate

$$E A = \bigvee_{a \in A} E a = \bigvee_{a \in A} E f(a) = E f[A]$$

□

As mentioned in the previous footnote,  $\mathfrak{G}$  doesn't take objects to gluing-complete  $\mathcal{Q}$ -sets; think of it in the same vein of object as Cauchy sequences sans the quotient. We first show that every compatible family over  $\mathfrak{G}(X)$  has a gluing.

**Lemma 4.4:**

If  $\mathcal{A}$  is a compatible family over  $\mathfrak{G}(X)$ , then  $\bigcup \mathcal{A}$  is compatible over  $X$ .

*Proof.* Take  $A, B \in \mathcal{A}$  and  $a \in A$  and  $b \in B$ ; it suffices to show that  $\delta(a, b) \geq E a \otimes E b$  since we know the dual inequality always holds.

The following is always true:

$$\delta(a, b) \geq \bigvee_{a' \in A} \bigvee_{b' \in B} \delta(a, a') \otimes \delta(b, b') \otimes \delta(a', b')$$

forcing us to accept the remaining argument

$$\begin{aligned}&= \bigvee_{a' \in A} \bigvee_{b' \in B} E a \otimes E a' \otimes E b \otimes E b' \otimes \delta(a', b') \\ &= E a \otimes E b \otimes \bigvee_{a', b'} \delta(a', b') \\ &= E a \otimes E b \otimes \delta(A, B) \\ &= E a \otimes E b \otimes E A \otimes E B\end{aligned}$$

since for extents  $\otimes = \wedge$ , and since  $E a \leq E A$  and likewise for  $b$  and  $B$ ,

$$= E a \otimes E b$$

□

**Proposition 4.3** (Compatible Families of  $\mathfrak{G}$  have Gluings):

We must give at least *one* gluing, but in general we should have many such gluings. That's a defect we shall correct by taking an appropriate quotient later.

*Proof.* Given  $\mathcal{A}$ , we claim that  $\bigcup \mathcal{A}$  is a canonical candidate for its gluing. We shall now verify that it indeed is up for the task. First, let  $A \in \mathcal{A}$  and denote by  $X$  the family given by  $\bigcup \mathcal{A}$ ; let us proceed.

$$\begin{aligned} \delta(X, A) &= \bigvee_{x \in X, a \in A} \delta(x, a) \\ &\geq \bigvee_{x \in A, a \in A} \delta(x, a) \\ &= \delta(A, A) = \mathbb{E} A \\ &\geq \delta(X, A) \end{aligned}$$

Remaining for us simply to show that the extent matches what we wanted.

$$\begin{aligned} \mathbb{E} X &= \bigvee_{x \in X} \mathbb{E} x \\ &= \bigvee_{A \in \mathcal{A}} \bigvee_{x \in A} \mathbb{E} x \\ &= \bigvee_{A \in \mathcal{A}} \mathbb{E} A \end{aligned}$$

□

The last piece of the puzzle of gluing completion is ensuring extensionality; and this is done by another functor, the  $\delta$ -quotient functor.

**Definition 4.8** ( $\delta$ -quotient):

We define the  $\neg/\delta$  functor naturally by quotienting together  $\delta$ -equivalent elements. It does indeed take  $\mathcal{Q}$ -sets to  $\mathcal{Q}$ -sets if we take  $\delta(\llbracket x \rrbracket, \llbracket y \rrbracket) = \delta(x, y)$ . It does indeed take morphisms to morphisms, due to  $\delta$  equivalence being congruential.

**Lemma 4.5:**

A compatible family of  $X/\delta$  has a gluing if and only if (any) section of that family has a gluing in  $X$ . That is to say, if  $A$  is our compatible family and  $\bar{A}$  is such that  $\mathfrak{u}[\bar{A}] = A$ , then  $A$  has a gluing if and only if  $\bar{A}$  does. Moreover, if any section has a gluing, then every section is also glued by it.

*Proof.* Suppose that  $A$  has a gluing, call it  $\llbracket x \rrbracket$ ; by definition,

$$\mathbb{E} x = \mathbb{E} \llbracket x \rrbracket = \bigvee_{\llbracket a \rrbracket \in A} \mathbb{E} \llbracket a \rrbracket = \bigvee_{\llbracket a \rrbracket \in A} \mathbb{E} a$$

So, if  $\bar{A}$  is a section of  $A$ , then

$$\begin{aligned} \mathbb{E} x &= \bigvee_{a \in \bar{A}} \mathbb{E} a \\ \delta(x, a) &= \delta(\llbracket x \rrbracket, \llbracket a \rrbracket) = \mathbb{E} \llbracket a \rrbracket = \mathbb{E} a \end{aligned}$$

Thus,  $x$  glues any section of  $A$ . Now suppose instead that  $x$  glues some section  $\bar{A}$  of  $A$ . Of course  $\llbracket x \rrbracket$  must now glue  $A$ :

$$\begin{aligned} E \llbracket x \rrbracket &= E x = \bigvee_{a \in \bar{A}} E a = \bigvee_{a \in \bar{A}} E \llbracket a \rrbracket = \bigvee_{\llbracket a \rrbracket \in A} E \llbracket a \rrbracket = E A \\ \delta(\llbracket a \rrbracket, \llbracket x \rrbracket) &= \delta(a, x) = E a = E \llbracket a \rrbracket \end{aligned}$$

□

**Remark 4.3:**

Evidently, every compatible family over  $X/\delta$  has a section that is a compatible family over  $X$ .

**Lemma 4.6:**

The functor  $\neg/\delta$  makes  $\mathcal{Q}$ -sets extensional (in fact it's “the best” extensional approximation).

*Proof.* Suppose

$$\delta(\llbracket x \rrbracket, \llbracket y \rrbracket) = E \llbracket x \rrbracket = E \llbracket y \rrbracket$$

Then  $\delta(x, y) = E x = E y$  and immediately we have  $x \sim y$  and as such,  $\llbracket x \rrbracket = \llbracket y \rrbracket$ .

The claim that it is “the best” at doing this is harder to state and prove, and it could/should be a theorem. □

**Theorem 4.1** (it became a theorem):

The functor  $\mathcal{Q}\text{-Set}_{f\text{Ext}} \hookrightarrow \mathcal{Q}\text{-Set}_f$  has a left adjoint and it is  $\neg/\delta$ .

*Proof.* As will become a trend throughout this article, we go about looking for units and counits and proving the triangle/zig-zag identities for those.

The counit is obvious, as applying  $\neg/\delta$  to an already extensional object yields an isomorphic object whose elements are singleton sets containing exactly the elements of the original  $\mathcal{Q}$ -set. So we simply unwrap them and that's the counit.

The unit is also simple, it's just the quotient map  $\mathfrak{u}$  – taking elements to their  $\delta$ -equivalence class.

In this case, there is no real point to showing the zig-zag identities hold; but the reader may if they feel so inclined. □

**Definition 4.9** (Gluing Completion):

The gluing completion functor  $\mathcal{G}$  is simply  $(\neg/\delta) \circ \mathfrak{E}$ .

**Proposition 4.4:**

The Gluing-completion of a  $\mathcal{Q}$ -set is indeed gluing-complete.

*Proof.* We know that every compatible family over  $\mathcal{G}X$  has at least one gluing, but we also know they are extensional; and thus we have that they are gluing complete. □



**Definition 4.10** ( $\mathcal{G}\mathcal{Q}\text{-Set}$ ):

Here we give the name  $\mathcal{G}\mathcal{Q}\text{-Set}$  to the full subcategory of  $\mathcal{Q}\text{-Set}_f$  whose objects are gluing complete. We drop the  $f$  subscript as this notion isn't particularly useful for relational morphisms anyway, so a it's pointless datum. In a later subsection the  $f$  subscript will reappear as will a  $r$  subscript, so that we may prove that

$$\mathcal{G}\mathcal{Q}\text{-Set}_r \simeq \mathcal{Q}\text{-Set}_r$$

In the light of the previous proposition, we are entitled to say that the signature of  $\mathcal{G}$  is actually

$$\mathcal{G} : \mathcal{Q}\text{-Set}_f \rightarrow \mathcal{G}\mathcal{Q}\text{-Set}$$

**Lemma 4.7:**

Equivalent compatible families (in the sense of the canonical  $\delta$  we have defined for the set of such families) have exactly the same gluings – and they are equivalent to a singleton set if and only if its element is one of their gluings.

*Proof.* It suffices that we show  $x$  glues  $A$  iff.  $\{x\} \sim A$  – as the  $\sim$  being an equivalence takes care of the rest of the claim for us.

So, let's suppose that  $x$  does indeed glue  $A$ ; it follows that

$$\delta(\{x\}, A) = \bigvee_{a \in A} \delta(x, a) = \bigvee_{a \in A} E a = E A = E\{x\}$$

Now instead suppose that  $\{x\} \sim A$ ; In particular  $E x = E\{x\}$  must be  $E A$  and thus  $\bigvee_{a \in A} E a$ . So that takes care of the last condition; the first condition for gluing still remains. But not for long:

$$\begin{aligned} E a &= E a \otimes E A \\ &= E a \otimes \delta(\{x\}, A) \\ &= \bigvee_{a' \in A} E a \otimes \delta(x, a') \\ &= \bigvee_{a' \in A} E a \otimes E a' \otimes \delta(x, a') \\ &= \bigvee_{a' \in A} \delta(a, a') \otimes \delta(x, a') \\ &\leq \delta(x, a) \end{aligned}$$

□

**Theorem 4.2** (The Gluing Completion Adjunction):

It's not particularly useful to have a functor that makes  $\mathcal{Q}$ -sets gluing-complete if that completion isn't actually “universal” in that it is the reflector of the subcategory we are interested in.

*Proof.* Again, we proceed by showing a unit-counit pair satisfying the zig-zag identities. For (my) sanity, let us name  $L = \mathcal{G}$  and  $R$  the fully faithful inclusion  $\mathcal{G}\mathcal{Q}\text{-Set} \hookrightarrow \mathcal{Q}\text{-Set}_f$ .

Let's find  $\eta : \mathbf{id} \rightarrow R \circ L$ ; what it must do is take an element  $x$  of  $X$  to something in the completion of  $X$ ; the completion of  $X$  is comprised of equivalence classes of compatible families, so it is enough that we find a compatible family to assign  $x$  to.  $\{x\}$  is the natural candidate as

$$\delta(x, y) = \delta(\{x\}, \{y\}) = \delta(\llbracket \{x\} \rrbracket, \llbracket \{y\} \rrbracket)$$

Of course, we have only defined components, and don't know if they form a natural transformation together; naturality holds quite trivially though:

$$\begin{array}{ccc}
 X & \xrightarrow{\eta} & \mathcal{G}(X) \\
 \downarrow f & & \downarrow \mathcal{G}(f) \\
 & \begin{array}{ccc}
 x & \longmapsto & \llbracket \{x\} \rrbracket \\
 \downarrow & & \downarrow \\
 f(x) & \longmapsto & ?
 \end{array} & \\
 \downarrow & & \downarrow \\
 Y & \xrightarrow{\eta} & \mathcal{G}(Y)
 \end{array}$$

We know that the action of  $\overline{\cdot}/\delta$  on an arrow is

$$f \xrightarrow{\overline{\cdot}/\delta} (\llbracket x \rrbracket \xrightarrow{\llbracket f \rrbracket} \llbracket f(x) \rrbracket)$$

we know this from regular old set theory, it's just how quotients work. The action of  $\mathfrak{G}$  on morphisms is

$$f \xrightarrow{\mathfrak{G}} (A \xrightarrow{f_*} f[A])$$

that is, obviously, just taking functions to direct images. And thus, we combine the two to get

$$\llbracket f_* \rrbracket (\llbracket \{x\} \rrbracket) = \llbracket f[\{x\}] \rrbracket = \llbracket \{f(x)\} \rrbracket$$

making the above square commute.

We should now provide a counit  $\varepsilon : L \circ R \rightarrow \mathbf{id}$ . One such function should take things from the completion of an already gluing-complete  $\mathcal{Q}$ -set back to the original  $\mathcal{Q}$ -set. The things  $\varepsilon$  operates over are of the form  $\llbracket A \rrbracket$  where  $A$  is a compatible family over – say –  $X$ .

The natural candidate is to take  $A$  to the unique element  $x_A$  in  $X$  (which was gluing-complete to begin with) that glues it. Considering the previous lemma's, it is evident that this definition does indeed yield a *map*, as any equivalent family will be glued by the same unique element. Moreover, since  $X$  was complete to begin with,  $\{x_A\} \in \llbracket A \rrbracket$  and hence the map will obviously be a morphism, as

$$\delta(\llbracket A \rrbracket, \llbracket B \rrbracket) = \delta(\llbracket x_A \rrbracket, \llbracket x_B \rrbracket) = \delta(x_A, x_B)$$

Remaining now only the need to show  $\varepsilon$  is natural transformation  $L \circ R \rightarrow \mathbf{id}$ ;

$$\begin{array}{ccc}
 \mathcal{G}(X) & \xrightarrow{\varepsilon} & X \\
 \downarrow \mathcal{G}(f) & & \downarrow f \\
 \mathcal{G}(Y) & \xrightarrow{\varepsilon} & Y
 \end{array}
 \quad
 \begin{array}{ccc}
 \llbracket A \rrbracket & \xrightarrow{\quad} & x_A \\
 \downarrow & & \downarrow \\
 \llbracket f[A] \rrbracket & \xrightarrow{\quad} & ?
 \end{array}$$

since gluing elements are preserved by functional morphisms, the square does indeed commute; hence, the family of maps is actually natural transformation and we can proceed in showing that the selected transformations satisfy the zig-zag identities. Recapping,

$$\begin{array}{c}
 x \xrightarrow{\eta_X} \llbracket \{x\} \rrbracket \\
 \llbracket A \rrbracket \xrightarrow{\varepsilon_{X'}} x_A
 \end{array}$$

and we aim to show that

$$\begin{array}{c}
 (\varepsilon L) \circ (L\eta) = \mathbf{id} \\
 (R\varepsilon) \circ (\eta R) = \mathbf{id}
 \end{array}$$

Given the below, it should then become obvious that we do indeed have witnesses to the adjunction we have set out to show to hold.

$$\begin{array}{c}
 L\eta : L \longrightarrow L \circ R \circ L \\
 (X \in \mathcal{Q}\text{-Set}) \mapsto (\llbracket A \rrbracket \xrightarrow{(L\eta)_X} \llbracket \{\llbracket A \rrbracket\} \rrbracket)
 \end{array}$$

$$\begin{array}{c}
 \varepsilon L : L \circ R \circ L \longrightarrow L \\
 (X \in \mathcal{Q}\text{-Set}) \mapsto (\llbracket \{\llbracket A \rrbracket\} \rrbracket \xrightarrow{(\varepsilon L)_X} \llbracket A \rrbracket)
 \end{array}$$

$$\begin{array}{c}
 R\varepsilon : R \circ L \circ R \longrightarrow R \\
 (K \in \mathcal{G}\mathcal{Q}\text{-Set}) \mapsto (\llbracket A \rrbracket \xrightarrow{(R\varepsilon)_K} x_A)
 \end{array}$$

$$\begin{array}{c}
 \eta R : R \longrightarrow R \circ L \circ R \\
 (K \in \mathcal{G}\mathcal{Q}\text{-Set}) \mapsto (x \xrightarrow{(\eta R)_K} \llbracket \{x\} \rrbracket)
 \end{array}$$

□

**Remark 4.4:**

The adjunction counit is very easily seen to be a natural *isomorphism* (because  $R$  was fully faithful), however the unit isn't: Consider the  $\mathcal{Q}$ -set given naturally by

$$X := \{\perp\} \amalg \{\perp\}$$

It obviously won't be extensional, because of this,  $\mathcal{G}(X) \cong \{\perp\}$  and there is no possible *functional* isomorphism between those, because functional isomorphisms are – in particular – also bijections. So the adjunction above is not an adjoint equivalence. For relational morphisms there are those kinds of isomorphisms, and so the adjunction unit will be a natural isomorphism too.

## 5 Singleton/Scott-Completeness

As previously mentioned, there is another notion completeness; it relates to a concept called singleton, which we will talk about now.

A singleton on a  $\mathcal{Q}$ -set  $X$ , morally speaking, is a  $\mathcal{Q}$ -valued characteristic function. Although it isn't *per se* a construction in the category  $\mathcal{Q}\text{-Set}$  – it does lead to categorically relevant results.

**Definition 5.1** (Singleton):

A *singleton* on  $X$  is a map between the underlying set and  $\mathcal{Q}$ :  $\sigma : |X| \rightarrow \mathcal{Q}$ . This map is expected to satisfy the following axioms

1.  $\sigma(x) \otimes \text{E } x = \sigma(x)$ ;
2.  $\sigma(x) \otimes \delta(x, y) \leq \sigma(y)$ ;
3.  $\sigma(x) \otimes \sigma(y) \leq \delta(x, y)$ ;
4.  $\sigma(x) \otimes \bigvee_y \sigma(y) = \sigma(x)$ ;

the first two being called the “subset axioms” the third being the “singleton condition” and the last axiom being called the “strictness condition”.

Intuitively, singletons represent a membership relation to a set – this is governed by the subset axioms; the singleton condition then states that the simultaneous membership of two different points implies their similitude; the last axiom, strictness, is more technical in nature.

It has been known in the literature for quite a few years [7, p. 30] – but its meaning is somewhat obscure, in that it is a necessary condition for a singleton to be representable (to be defined next), but it doesn't have a neat interpretation. The best we have arrived at is that strictness gives us that the supremum  $\sup_y \sigma(y)$  “behaves like an extent” in that it is idempotent.

### 5.1 Basics about Singletons

**Definition 5.2** (Representable Singleton):

A singleton  $\sigma$  is said to be representable (or represented by an element  $x_\sigma$ ) when

there is one such element  $x_\sigma$  such that

$$\sigma(y) = \delta(y, x_\sigma)$$

**Proposition 5.1:**

If  $A$  is a compatible family, then  $\delta(\{\_\}, A)$  is a singleton. In particular,  $\delta(\_, x)$  is always a singleton (by taking  $A = \{x\}$ ).

*Proof.* The subset axioms are immediately satisfied by the fact that  $\mathfrak{G}(X)$  is always a  $\mathcal{Q}$ -set. The Singleton condition is trivial too:

$$\sigma_A(x) \otimes \sigma_A(y) = \delta(\{x\}, A) \otimes \delta(\{y\}, A) \leq \delta(\{x\}, \{y\}) = \delta(x, y)$$

obviously, it is also strict considering that

$$\bigvee_{y \in X} \sigma_A(y) \geq \bigvee_{a \in A} \sigma_A(a) = \bigvee_{a \in A} E a = E A \geq \bigvee_{y \in X} \sigma_A(y)$$

□

**Remark 5.1:**

Quite obviously, if the singleton above is representable, then the representing element glues  $A$ . And if an element glues  $A$ , then it is bound to represent the singleton.

**Proposition 5.2:**

If  $A$  is a compatible family, and  $\sigma$  is some singleton over  $A$ 's  $\mathcal{Q}$ -set, we have that if  $x_A$  glues  $A$ , then

$$\sigma(x_A) = \bigvee_{a \in A} \sigma(a)$$

*Proof.*

$$\begin{aligned} \sigma(a) &= \sigma(a) \otimes E a \\ &= \sigma(a) \otimes \delta(a, x_A) \\ &\leq \sigma(x_A) \end{aligned}$$

$$\begin{aligned} \sigma(x_A) &= \sigma(x_A) \otimes E x_A \\ &= \sigma(x_A) \otimes \bigvee_{a \in A} E a \\ &= \sigma(x_A) \otimes \bigvee_{a \in A} \delta(a, x_A) \\ &= \bigvee_{a \in A} \sigma(x_A) \otimes \delta(a, x_A) \\ &\leq \bigvee_{a \in A} \sigma(a) \end{aligned}$$

□

**Definition 5.3** (Scott-Completeness):

We say that  $X$  is Scott-complete, or singleton-complete, when every one of its singletons have exactly one representing element.

**Example 5.1:**

The terminal object is Scott-complete. An indirect way of seeing this is to skip through the paper to where we prove that the inclusion of Scott-complete  $\mathcal{Q}$ -sets creates limits. A more direct answer comes from the fact that since  $\top \in \mathbf{E} \mathcal{Q}$  is the gluing of the compatible family  $\mathbf{E} \mathcal{Q}$ , thanks to prop. 5.2,  $\sigma(\top) = \bigvee_e \sigma(e)$ . And hence:

$$\sigma(e) = \sigma(e) \otimes e \leq \sigma(\top) \otimes e = \sigma(\top) \otimes \delta(e, \top) \leq \sigma(e)$$

And therefore,  $\sigma(\top)$  represents  $\sigma$ . Uniqueness is quite obvious: if for some  $x$ ,  $\forall e : \sigma(e) = e \otimes x$ , in particular  $\sigma(\top) = \top \otimes x = x$  by integrality.

**Example 5.2:** 1. If  $A$  is a commutative ring and  $M$  is a left  $A$ -module, then the set over  $\mathcal{Q} = \text{Ideals}(A)$ ,  $X_M = (M, \delta)$ , is such that for each  $x, y \in M$ ,  $x \sim_\delta y$  iff  $A.x = A.y$ , so in general, is not a extensional  $\mathcal{Q}$ -set.

**Remark 5.2:**

Scott-completeness is invariant under isomorphisms in  $\mathcal{Q}\text{-Set}_f$  – but not in  $\mathcal{Q}\text{-Set}_r$ , this will be explained in a later subsection.

The first claim is easy to show, as functional isomorphisms are simply bijections preserving  $\delta$ .

**Theorem 5.1:**

Scott-Completeness implies Gluing-Completeness

*Proof.* Take a compatible family  $A$ , consider the singleton defined in the previous proposition; suppose  $x_A$  represents  $\sigma_A$ ; thus,

$$\bigvee_{a \in A} \delta(x_A, a) = \sigma_A(x_A) = \delta(x_A, x_A) = \mathbf{E} x_A$$

$$\delta(a, x_A) = \sigma_A(a) = \bigvee_{a' \in A} \delta(a, a') = \mathbf{E} a$$

and hence  $x_A$  glues  $A$ . Now suppose that  $x_A$  glues  $A$ , then

$$\begin{aligned}\delta(y, x_A) &\geq \delta(x_A, a) \otimes \delta(a, y) \\ &\geq \mathbf{E} a \otimes \delta(a, y) = \delta(a, y)\end{aligned}$$


$$\begin{aligned}\delta(y, x_A) &\geq \bigvee_{a \in A} \delta(y, a) \\ &\geq \bigvee_{a \in A} \delta(a, x_A) \otimes \delta(y, x_A) \\ &= \delta(y, x_A) \otimes \bigvee_{a \in A} \delta(a, x_A) \\ &= \delta(y, x_A) \otimes \bigvee_{a \in A} \mathbf{E} a \\ &= \delta(y, x_A) \otimes \mathbf{E} x_A \\ &= \delta(y, x_A)\end{aligned}$$

hence  $x_A$  represents  $\sigma_A$ . Thus, if all singletons have exactly one representing element, then all compatible families have exactly one gluing.  $\square$

**Theorem 5.2:**

There are gluing complete  $\mathcal{Q}$ -sets over  $\mathcal{Q} = \mathcal{P}(2)$  which *are not* Scott-complete – so those two concepts *are not* logically equivalent.

*Proof.* Let  $\mathcal{Q}$  be the following partial order



First, let  $S \subseteq \mathcal{Q}$  and let  $\delta = \wedge$ ;  $S$  is gluing-complete if and only if  $S$  is complete as a lattice:

If  $A$  is a subset of  $S$ , it must be compatible (!) since  $\mathbf{E} x = x$  and  $\delta(x, y) = x \wedge y$  and thus  $\delta(x, y) = \mathbf{E} x \wedge \mathbf{E} y$ . Moreover, if  $x_A$  glues  $A$ , then

$$\mathbf{E} x_A = x_A = \bigvee_{a \in A} \mathbf{E} a = \bigvee_{a \in A} a$$

and hence  $x_A$  is  $\sup A$ . It's easy to verify that the supremum is – if it exists in  $S$  – the gluing of  $A$ .

So, the set  $S = \{\perp, a, \top\}$  is gluing complete, obviously. But the singleton  $\sigma(x) = \neg a \wedge x$  (which is the restriction of the singleton represented by  $\neg a$ ) has no representing element.

$$\sigma(\perp) = \perp \quad \sigma(a) = \perp \quad \sigma(\top) = \neg a$$

And so  $S$  is gluing-complete but not Scott-complete.  $\square$

**Definition 5.4** (The  $\mathcal{Q}$ -set of Singletons):

It is, in general, not possible to form a  $\mathcal{Q}$ -set of singletons over a  $\mathcal{Q}$ -set  $X$ . There is some trouble in defining  $\delta$  for singletons in a way akin to [3, thm. 2.9.5]. This is due to the fact that  $\bigvee_x \sigma(x) \otimes \sigma(x)$  not being “extent-like” necessarily.

We must – seemingly – add the axiom of Strength to our quantale  $\mathcal{Q}$  to guarantee we can form a  $\mathcal{Q}$ -set of singletons. This enables us to define the following:

$$\begin{aligned} |\Sigma(X)| &= \{\sigma \mid \sigma \text{ is a singleton over } X\} \\ \delta(\sigma, \xi) &= \bigvee_{x \in X} \sigma(x) \otimes \xi(x) \end{aligned}$$

**Proposition 5.3:**

Obviously, we must prove this is indeed a  $\mathcal{Q}$ -set.

*Proof.* Firstly, we know that  $\bigvee_x \sigma(x)$  is idempotent, and hence

$$\bigvee_x \sigma(x) \leq \bigvee_x \sigma(x)$$

since  $\mathcal{Q}$  is hypothesized to be strong, we have

$$\bigvee_x \sigma(x) \leq \bigvee_x \sigma(x) \otimes \sigma(x)$$

and since  $\mathcal{Q}$  is semi-cartesian

$$\bigvee_x \sigma(x) \otimes \sigma(x) \leq \bigvee_x \sigma(x) \quad (\text{semi-cartesian})$$

and therefore,  $E\sigma = \bigvee_x \sigma(x)$ . Now we move onto the  $\mathcal{Q}$ -set axioms.  $\delta$  as defined is obviously symmetric; let’s show extensionality:

$$\delta(\sigma, \xi) \otimes E\sigma = \bigvee_x \sigma(x) \otimes \xi(x) \otimes E\sigma$$

now recall  $\sigma$ ’s strictness condition

$$= \bigvee_x \sigma(x) \otimes \xi(x)$$

Now we need only to show triangle inequality/transitivity

$$\begin{aligned} \delta(\sigma, \xi) \otimes \delta(\xi, \psi) &= \bigvee_x \bigvee_y \sigma(x) \otimes \xi(x) \otimes \xi(y) \otimes \psi(y) \\ &\leq \bigvee_x \bigvee_y \sigma(x) \otimes \delta(x, y) \otimes \psi(y) \\ &\leq \bigvee_y \sigma(y) \otimes \psi(y) \\ &= \delta(\sigma, \psi) \end{aligned}$$



□

**Lemma 5.1:**

A relational morphism  $\varphi : X \rightarrow Y$  induces a functional map  $\check{\varphi} : X \rightarrow \Sigma(Y)$  given by

$$\check{\varphi}(x) = \varphi(x, \_)$$

*Proof.* We ought to show that the map does indeed take every  $x$  to a singleton  $\check{\varphi}(x)$  and that the map is indeed a morphism.

$$\begin{aligned} \delta(x, x') \otimes \check{\varphi}(x) &= \delta(x, x') \otimes \varphi(x, \_) \\ &\leq \varphi(x', \_) \end{aligned}$$

$$\begin{aligned} \mathbf{E} x \otimes \check{\varphi}(x) &= \mathbf{E} x \otimes \varphi(x, \_) \\ &= \varphi(x, \_) \end{aligned}$$

$$\begin{aligned} [\check{\varphi}(x)](y) \overset{\check{\varphi}(x)}{\otimes} (y') &= \varphi(x, y) \otimes \varphi(x, y') \\ &\leq \delta(y, y') \end{aligned}$$

$$\begin{aligned} [\check{\varphi}(x)](y) \otimes \bigvee_{y'} [\check{\varphi}(x)](y') &= \varphi(x, y) \otimes \bigvee_y \varphi(x, y') \\ &= \varphi(x, y) \otimes \mathbf{E} x \\ &= \varphi(x, y) \\ &= \check{\varphi}(x') \end{aligned}$$

Establishing that  $\check{\varphi}(x)$  is always a singleton over  $Y$ . Now, with regards to it being a morphism, just above we have also shown that  $\mathbf{E} \check{\varphi}(x) = \mathbf{E} x$ , so only one axiom remains:

$$\begin{aligned} \delta(\check{\varphi}(x), \check{\varphi}(x')) &= \bigvee_y \varphi(x, y) \otimes \varphi(x', y) \\ &\geq \bigvee_y \varphi(x, y) \otimes \varphi(x, y) \otimes \delta(x, x') \\ &\geq \delta(x, x') \otimes \bigvee_y \varphi(x, y) \otimes \varphi(x, y) \\ &= \delta(x, x') \otimes \mathbf{E} x \\ &= \delta(x, x') \end{aligned}$$

□

**Remark 5.3:**

It is not *necessary* for  $\mathcal{Q}$  to be strong for the question of Scott completeness

to make sense; but it is not apparent how one may perform *Scott-completion* without that added strength. We will talk about completion now.

**Definition 5.5:**

We denote by  $\Sigma\mathcal{Q}\text{-Set}$  the full subcategory of  $\mathcal{Q}\text{-Set}_f$  whose objects are Scott-complete.

**Definition 5.6** (Separability):

A  $\mathcal{Q}$ -set is said to be *separable* when the mapping  $x \mapsto \delta(\_, x)$  is *injective*; *ie.* it is the completeness condition sans surjectivity.

**Proposition 5.4:**

Separability is equivalent to Extensionality.

*Proof.* Immediate from what was proven in the definition of  $\delta$ -equivalence.  $\square$

## 5.2 Scott Completion

Scott-completion is, unsurprisingly, going to be a reflector functor of the from  $\mathcal{Q}\text{-Set}$  to the full subcategory  $\Sigma\mathcal{Q}\text{-Set}$ ; in this subsection we therefore will show that  $\Sigma(X)$  is indeed Scott-complete and that object assignment can be made functorial – in such a way to get a left adjoint to the fully faithful subcategory inclusion.

**Lemma 5.2** ([3] cf. p. 158):

If  $\sigma_x$  denotes  $\delta(\_, x)$ , then

$$\delta(\sigma_x, \xi) = \xi(x)$$

a result akin to the Yoneda lemma, wherein

$$\mathbf{hom}(\mathfrak{y}(x), F) \cong F(x)$$

*Proof.*

$$\begin{aligned} \delta(\sigma_x, \xi) &= \bigvee_y \delta(y, x) \otimes \xi(y) \\ &\leq \xi(x) \\ &= \mathbf{E}x \otimes \xi(x) \\ &= \delta(x, x) \otimes \xi(x) \\ &\leq \bigvee_z \delta(z, x) \otimes \xi(z) \\ &= \delta(\sigma_x, \xi) \end{aligned}$$

$\square$

**Remark 5.4:**

Notice that this means  $\delta(\sigma_x, \sigma_y) = \delta(x, y)$  and thus the mapping  $x \mapsto \sigma_x$  preserves  $\delta$  and hence is a morphism – it is in fact a regular monomorphism.

**Proposition 5.5:**

$\Sigma(X)$  is, indeed, Scott-complete.

*Proof.* In this proof, we shall refer to members of  $\Sigma(X)$  as 1-singletons and use lowercase  $\xi, \psi, \varphi$  and refer to members of  $\Sigma \circ \Sigma(X)$  as 2-singletons and use uppercase  $\Sigma$  *etc.*

To go about doing this we must show two things: that two 1-singletons cannot represent the same 2-singleton; and that every 2-singleton comes from a 1-singleton.

For the purpose of showing that  $\xi \mapsto \sigma_\xi$  is injective, suppose that  $\xi$  and  $\psi$  are such that  $\sigma_\xi = \sigma_\psi$ . Let  $x \in X$  and realize

$$\xi(x) = \delta(\sigma_x, \xi) = \sigma_\xi(\sigma_x) = \sigma_\psi(\sigma_x) = \delta(\sigma_x, \psi) = \psi(x)$$

thence they are extensionally equal.

Now take a 2-singleton  $\Sigma$ , we shall define a 1-singleton that we hope represents it; namely:

$$\varphi(x) = \Sigma(\sigma_x)$$

one way to become convinced that  $\varphi$  is indeed a singleton over  $X$  is that  $\sigma_{\underline{\quad}}$  preserves  $\delta$  and thus  $\Sigma \circ \sigma_{\underline{\quad}}$  is “more or less” the restriction of  $\Sigma$  to the set of representable singletons. The one axiom that doesn’t obviously hold is strictness – which we show after the following deduction:

$$\begin{aligned} \Sigma(\xi) &= \mathbb{E} \xi \otimes \Sigma(\xi) \\ &= \bigvee_x \xi(x) \otimes \xi(x) \otimes \Sigma(\xi) \\ &= \bigvee_x \xi(x) \otimes \delta(\sigma_x, \xi) \otimes \Sigma(\xi) \\ &\leq \bigvee_x \xi(x) \otimes \Sigma(\sigma_x) \\ &= \delta(\xi, \varphi) \end{aligned}$$

Hence, if we can show strictness, we are done. So take  $x \in X$  which we ought to show enjoys the property that

$$\varphi(x) \otimes \bigvee_y \varphi(y) = \varphi(x)$$

it is clear that it would suffice if we proved that  $\mathbb{E} \Sigma \leq \mathbb{E} \varphi$ . For that would suffice for them to be indeed equal; and them being equal mean we could then argue:

$$\varphi(x) \otimes \bigvee_y \varphi(y) = \varphi(x) \otimes \mathbb{E} \varphi = \Sigma(\sigma_x) \otimes \mathbb{E} \Sigma = \Sigma(\sigma_x) = \varphi(x)$$

since  $\Sigma$  was taken to be strict. And thus we shall prove instead simply that  $\mathbb{E} \Sigma \leq \mathbb{E} \varphi$ .

$$\begin{aligned}
\mathbf{E}\Sigma &= \bigvee_{\xi} \Sigma(\xi) \otimes \Sigma(\xi) \\
&= \bigvee_{\xi} \Sigma(\xi) && \text{(Strength)} \\
&= \bigvee_{\xi} \Sigma(\xi) \otimes \mathbf{E}\xi \\
&= \bigvee_x \bigvee_{\xi} \Sigma(\xi) \otimes \xi(x) \\
&= \bigvee_x \bigvee_{\xi} \Sigma(\xi) \otimes \delta(\xi, \sigma_x) \\
&\leq \bigvee_x \Sigma(\sigma_x) \\
&= \bigvee_x \Sigma(\sigma_x) \otimes \Sigma(\sigma_x) && \text{(Strength)} \\
&= \mathbf{E}\varphi
\end{aligned}$$

□

Having then proved that  $\Sigma(X)$  is indeed Scott-complete, we now must move into the associated completion functor.

**Definition 5.7:**

We extend the action of  $\Sigma$  by acting on functional morphisms as follows:

$$f : X \rightarrow Y$$

$$\begin{aligned}
\Sigma f : \Sigma(X) &\rightarrow \Sigma(Y) \\
\xi &\mapsto \bigvee_{x \in X} \delta(f(x), \_) \otimes \xi(x)
\end{aligned}$$

Morally speaking, if  $\xi(x)$  measures how much  $x$  is akin to some abstract point, then  $(\Sigma f)(\xi)$  measures how close a point in  $y$  is to the “image” of that abstract point under  $f$ .

**Proposition 5.6:**

Associated to the previous definition, we must prove that  $\Sigma f(\xi)$  is indeed a singleton; that this assignment is a morphism; and that the action on morphisms is functorial.

*Proof.* First of all, for notational sanity denote  $\Sigma f(\xi)$  by  $\xi_f$ .

$$\begin{aligned}\xi_f(y) \otimes \mathbb{E} y &= \bigvee_x \delta(f(x), y) \otimes \xi(x) \otimes \mathbb{E} y \\ &= \bigvee_x \delta(f(x), y) \otimes \xi(x) \\ &= \xi_f(y)\end{aligned}$$

$$\begin{aligned}\xi_f(y) \otimes \delta(y, y') &= \bigvee_x \delta(f(x), y) \otimes \delta(y, y') \otimes \xi(y) \\ &\leq \bigvee_x \delta(f(x), y') \otimes \xi(y) \\ &= \xi_f(y')\end{aligned}$$

$$\begin{aligned}\xi_f(y) \otimes \xi_f(y') &= \bigvee_{x, x'} \delta(f(x), y) \otimes \delta(f(x'), y') \otimes \xi(x) \otimes \xi(x') \\ &\leq \bigvee_{x, x'} \delta(f(x), y) \otimes \delta(f(x'), y') \otimes \delta(x, x') \\ &\leq \bigvee_{x, x'} \delta(f(x), y) \otimes \delta(f(x'), y') \otimes \delta(f(x), f(x')) \\ &\leq \bigvee_{x'} \delta(f(x'), y) \otimes \delta(f(x'), y') \\ &\leq \delta(y, y')\end{aligned}$$

$$\begin{aligned}\xi_f(y) \otimes \bigvee_{y'} \xi_f(y') &= \bigvee_{y'} \bigvee_{x'} \xi_f(y) \otimes \delta(f(x'), y) \otimes \xi(x') \\ &= \bigvee_{x'} \xi_f(y) \otimes \delta(f(x'), f(x')) \otimes \xi(x') \\ &= \bigvee_{x'} \xi_f(y) \otimes \mathbb{E} x' \otimes \xi(x') \\ &= \bigvee_{x'} \xi_f(y) \otimes \xi(x') \\ &= \bigvee_x \delta(f(x), y) \otimes \xi(x) \otimes \bigvee_{x'} \xi(x') \\ &= \bigvee_x \delta(f(x), y) \otimes \xi(x) \\ &= \xi_f(y)\end{aligned}$$

The second thing we ought to do is showing that the map  $\Sigma f$  is a morphism; that is rather easy actually. Let  $\psi_f$  denote the image of  $\psi$  under  $\Sigma f$  as we had

done with  $\xi$ .

$$\begin{aligned}
\delta(\xi_f, \psi_f) &= \bigvee_y \xi_f(y) \otimes \psi_f(y) \\
&= \bigvee_y \bigvee_{x, x'} \delta(f(x), y) \otimes \delta(f(x'), y) \otimes \xi(x) \otimes \varphi(x') \\
&\geq \bigvee_y \bigvee_x \delta(f(x), y) \otimes \delta(f(x), y) \otimes \xi(x) \otimes \varphi(x) \quad (\text{by letting } x = x') \\
&\geq \bigvee_x \mathbf{E} f(x) \otimes \mathbf{E} f(x) \otimes \xi(x) \otimes \varphi(x) \quad (\text{by letting } y = f(x)) \\
&= \bigvee_x \xi(x) \otimes \varphi(x) \\
&= \delta(\xi, \varphi)
\end{aligned}$$

$$\begin{aligned}
\xi_f(y) &\geq \xi_f(y) \otimes \bigvee_{y'} \xi_f(y') \quad (\text{semi-cartesian}) \\
&= \bigvee_{y'} \bigvee_{x'} \xi_f(y) \otimes \delta(f(x'), y') \otimes \xi(x') \\
&\geq \bigvee_{x'} \xi_f(y) \otimes \delta(f(x'), f(x')) \otimes \xi(x') \quad (\text{by letting } y' = f(x')) \\
&= \bigvee_{x'} \xi_f(y) \otimes \mathbf{E} f(x') \otimes \xi(x') \\
&= \xi_f(y) \otimes \bigvee_{x'} \xi(x') \\
&= \bigvee_x \delta(f(x), y) \otimes \xi(x) \otimes \bigvee_{x'} \xi(x') \\
&= \bigvee_x \delta(f(x), y) \otimes \xi(x) \\
&= \xi_f(y)
\end{aligned}$$

The final problem to be unravelled is functoriality. We must show that  $\mathbf{id}$  is sent to  $\mathbf{id}$  and composition is preserved.  $\Sigma \mathbf{id}$  is defined as

$$\begin{aligned}
[\Sigma \mathbf{id}(\xi)](x) &= \bigvee_{x'} \delta(x, x') \otimes \xi(x') \\
&\geq \mathbf{E} x \otimes \xi x = \xi(x) \\
&\geq \delta(x', x) \otimes \xi(x), \quad \forall x'
\end{aligned}$$

and hence by universality of the supremum, we have the desired equality:

$$(\Sigma \mathbf{id})(\xi) = \xi$$

Now, take  $X \xrightarrow{f} Y \xrightarrow{g} Z$

$$\begin{aligned}
[\Sigma(f \circ g)(\xi)](z) &= \bigvee_x \delta(g \circ f(x), z) \otimes \xi(x) \\
&= \bigvee_x \bigvee_y \delta(f(x), y) \otimes \xi(x) \otimes \delta(g(y), z) \\
&= \bigvee_y \delta(g(y), z) \otimes \left( \bigvee_x \delta(f(x), y) \otimes \xi(x) \right) \\
&= \bigvee_y \delta(g(y), z) \otimes^{\Sigma f(\xi)} (y) \\
&= [(\Sigma g)((\Sigma f)(\xi))](z) \\
&= [(\Sigma g) \circ (\Sigma f)](z)
\end{aligned}$$

Thus, we have shown that  $\Sigma$  is a functor from  $\mathcal{Q}\text{-Set}_f$  to  $\Sigma\mathcal{Q}\text{-Set}$ .  $\square$

**Theorem 5.3:**

We aim to show that  $\Sigma$  indeed a completion, in the sense that it is the left adjoint to an full inclusion functor – of  $\Sigma\mathcal{Q}\text{-Set}$ ; yielding that if  $K$  is Scott-complete,

$$\mathcal{Q}\text{-Set}(X, K) \cong \mathcal{Q}\text{-Set}(\Sigma X, K)$$

*Proof.* Again, we shall proceed by finding units and counits. Once more let us rename  $\Sigma$  by  $L$  dub the inclusion functor simply  $R$ , after their role in the adjunction. We ought to find  $\eta$  and  $\varepsilon$  as we had in Gluing-completion proof.

The unit,  $\eta$  is straightforward, it is the associated singleton morphism  $x \mapsto \delta(\_, x)$  – which we know to be a morphism already, and hence we need only to show naturality from  $\mathbf{id}$  to  $R \circ L$ .

$$\begin{array}{ccc}
X & \xrightarrow{\eta} & \Sigma(X) \\
\downarrow f & & \downarrow \Sigma(f) \\
Y & \xrightarrow{\eta} & \Sigma(Y)
\end{array}$$

$$\begin{array}{ccc}
x & \longmapsto & \delta(\_, x) \\
\downarrow & & \downarrow \\
f(x) & \longmapsto & ?
\end{array}$$

$$\Sigma f(\delta(\_, x)) = \bigvee_{x'} \delta(\_, f(x')) \otimes \delta(x', x) = \delta(\_, f(x))$$

and hence we have naturality. It should be remarked that  $\Sigma f(\sigma_x) = \sigma_{f(x)}$  as just proved above.

The counit is also rather straightforward; we start with a Scott complete  $X$ , we forget all about completeness and perform a completion. Since we know that  $x \mapsto \sigma_x$  is bijective (by definition of completeness) and that this mapping preserves  $\delta$ , we know that  $X$  is isomorphic to  $\Sigma(X)$ . So we need only show that this map is a *natural* isomorphism. Here  $\varepsilon$  takes a singleton (which is always representable) to its unique representing element:

$$\begin{array}{ccc}
\Sigma(X) & \xrightarrow{\varepsilon} & X \\
\downarrow \Sigma(f) & & \downarrow f \\
\Sigma(Y) & \xrightarrow{\varepsilon} & Y
\end{array}
\quad
\begin{array}{ccc}
\delta(\_, x) & \dashrightarrow & x \\
\downarrow & & \downarrow \\
\delta(\_, f(x)) & \dashrightarrow & ?
\end{array}$$

Which is obviously commutative. Recapping the definitions,

$$\begin{aligned}
\eta_X : X &\rightarrow \Sigma(X) \\
x &\mapsto \delta(\_, x)
\end{aligned}$$

$$\begin{aligned}
L\eta : L &\longrightarrow L \circ R \circ L \\
(X \in \mathcal{Q}\text{-Set}) &\mapsto (\xi \xrightarrow{(L\eta)_X} \bigvee_x \delta(\_, \eta(x)) \otimes \xi(x))
\end{aligned}$$

As we know from the Yoneda-esque lemma, the above can be simplified into the following:

$$\begin{aligned}
L\eta : L &\longrightarrow L \circ R \circ L \\
(X \in \mathcal{Q}\text{-Set}) &\mapsto (\xi \xrightarrow{(L\eta)_X} \bigvee_x \_ (x) \otimes \xi(x))
\end{aligned}$$



which we know to be simply  $\delta(\_, \xi)$ . As for  $\varepsilon L$ , we first recall what is the representing element of a 2-singleton  $\Sigma$ . It is what we had called  $\varphi$  – which was the action of  $\Sigma$  on representable singletons, or more helpfully, the composite  $\Sigma(\sigma\_)$  or even better:  $\Sigma \circ \eta$ .

$$\begin{aligned} \varepsilon L : L \circ R \circ L &\longrightarrow L \\ (X \in \mathcal{Q}\text{-Set}) &\mapsto (\Sigma \xrightarrow{(\varepsilon L)_X} \Sigma \circ \eta) \end{aligned}$$

$$\begin{aligned} R\varepsilon : R \circ L \circ R &\longrightarrow R \\ (X \in \mathcal{G}\mathcal{Q}\text{-Set}) &\mapsto (\delta(\_, x) \xrightarrow{(R\varepsilon)_X} x) \end{aligned}$$

$$\begin{aligned} \eta R : R &\longrightarrow R \circ L \circ R \\ (X \in \mathcal{G}\mathcal{Q}\text{-Set}) &\mapsto (x \xrightarrow{(\eta R)_X} \delta(\_, x)) \end{aligned}$$

Tracing  $x$  along  $(R\varepsilon) \circ (\eta R)$  we find

$$x \xrightarrow{(\eta R)_X} \delta(\_, x) \xrightarrow{(R\varepsilon)_X} x$$

and now, tracing  $\xi$  along  $(\varepsilon L) \circ (L\eta)$ , we get

$$\xi \xrightarrow{(L\eta)_X} \delta(\_, \xi) \xrightarrow{(\varepsilon L)_X} \delta(\sigma\_, \xi)$$

which, again, by the Yoneda-esque lemma is  $\xi(\_)$  which is extensionally equal to  $\xi$ , of course. And thus, our two composites were equal to the appropriate identities; thus establishing the adjunction.

We already knew the counit was an isomorphism, but since it is the counit of an adjunction with a fully faithful right adjoint that would have proved it as well.  $\square$


## 6 Connection between Completeness Conditions

### Theorem 6.1:

Let  $(X, \delta)$  be an extensional  $\mathcal{Q}$ -set. The following conditions are equivalent:

1.  $(X, \delta)$  is Scott-complete.
2.  $(X, \delta)$  is gluing-complete and, for each singleton  $\sigma$  over  $(X, \delta)$ , it holds the condition below:
  - $(*_\sigma)$  For each  $x \in X$ , there is  $y \in X$  such that  $\sigma(x) \leq \sigma(y) = E y$ .

We have already mentioned that Scott-completeness implies Gluing-completeness; and hence there is another fully faithful subcategory inclusion at play, forming the triangle:



The question, visually posed in the diagram by the gray dotted arrow, is “is there a functor left adjoint to that lonely inclusion?” The, answer, not surprisingly, is yes. And it is given by the only composite going in that direction. Now, this isn’t very spectacular, and it comes directly from **hom** isomorphism and the categories involved being full *etc.*

The more interesting categorical property is the characterization of Scott-completeness; which we shall work towards in this next subsection.

### 6.1 Scott-Completeness and Relational Morphisms

Scott-completeness is deeply tied to relational morphisms – which might be surprising since we have deliberately not dealt with  $\mathcal{Q}\text{-Set}_r$  in the context of completeness thus far. The reason is that singleton completeness is *too* deeply tied to relational morphisms.

**Theorem 6.2:**

$$\Sigma\mathcal{Q}\text{-Set}_r \simeq \mathcal{Q}\text{-Set}_r$$

*Proof.* We proceed by providing a functor fully faithful functor that is essentially surjective; in this case, it is easier to show that the full subcategory inclusion is essentially surjective, since by necessity is fully faithful.

This then amounts to us providing an isomorphism between any object and a Scott-complete one. In this case we have an obvious candidate, that being  $\Sigma(X)$ . Define

$$\begin{aligned}
 \varphi : X &\rightarrow \Sigma(X) \\
 (x, \xi) &\mapsto \xi(x)
 \end{aligned}$$

$$\begin{aligned}
 \varphi^{-1} : \Sigma(X) &\rightarrow X \\
 (\xi, x) &\mapsto \xi(x)
 \end{aligned}$$

We ought to prove that both of them are morphisms first, but should the reader grant us a moratorium on that we could argue that

$$\begin{aligned}\varphi \circ \varphi^{-1}(\xi, \psi) &= \bigvee_x \xi(x) \otimes \psi(x) \\ &= \delta(\xi, \psi)\end{aligned}$$

$$\begin{aligned}\varphi^{-1} \circ \varphi(x, x') &= \bigvee_{\xi} \xi(x) \otimes \xi(x') \\ &\geq \delta(x, x) \otimes \delta(x, x') \\ &= \delta(x, x')\end{aligned}$$

hence, their composite is pointwise greater than the identity; but we know that under strong quantales, the pointwise order on **homs** is in fact discrete. So they are actually equal.

It remains to be seen that as defined, both  $\varphi$  and its supposed inverse are actually morphisms. To it, then. It should be obvious why the extensionality axioms should hold:

$$\begin{aligned}\xi(x) \otimes \mathbf{E} \xi &= \xi(x) \otimes \bigvee_{x'} \xi(x') \\ &= \xi(x) \\ &= \xi(x) \otimes \mathbf{E} x\end{aligned}$$

The  $\delta$  laws also come for free:

$$\delta(x, x') \otimes \xi(x) \leq \xi(x) \quad (\text{subset axiom})$$

$$\begin{aligned}\delta(\xi, \psi) \otimes \xi(x) &= \delta(\xi, \psi) \otimes \delta(\sigma_x, \xi) \\ &\leq \delta(\sigma_x, \psi) \\ &= \psi(x)\end{aligned}$$

The singleton condition holds trivially as well – due to how  $\delta(\xi, \psi)$  is defined.

$$\begin{aligned}\varphi(x, \xi) \otimes \varphi(x, \psi) &= \xi(x) \otimes \psi(x) \\ &\leq \bigvee_x \xi(x) \otimes \psi(x) \\ &= \delta(\xi, \psi)\end{aligned}$$

$$\begin{aligned}\varphi^{-1}(\xi, x) \otimes \varphi^{-1}(\xi, x') &= \xi(x) \otimes \xi(x') \quad (\xi \text{ is a singleton}) \\ &\leq \delta(x, x')\end{aligned}$$

remaining only to show strictness

$$\begin{aligned} \bigvee_{\xi} \varphi(x, \xi) &= \bigvee_{\xi} \xi(x) && \text{(now let } \xi = \sigma_x) \\ &\geq \delta(x, x) = \mathbb{E} x \\ &\geq \xi(x) \quad \forall \xi \end{aligned}$$

and hence  $\varphi$  is indeed strict. Now for  $\varphi^{-1}$ :

$$\begin{aligned} \bigvee_x \varphi^{-1}(\xi, x) &= \bigvee_x \xi(x) \\ &= \mathbb{E} \xi \end{aligned}$$

□

**Theorem 6.3:**

$$\Sigma\mathcal{Q}\text{-Set}_f \cong \Sigma\mathcal{Q}\text{-Set}_r$$

Recall the underlying graph functor  $\mathcal{G}_r$ ; we propose that its restriction to  $\Sigma\mathcal{Q}\text{-Set}_f$  has an inverse, and we give it explicitly now:

$$\mathcal{F}_N : \Sigma\mathcal{Q}\text{-Set}_r \rightarrow \Sigma\mathcal{Q}\text{-Set}_f$$

is a functor that “remembers” that relational morphisms between  $\mathcal{Q}$ -sets are always functional. It’s action on objects is the identity. As we had mentioned in the previous subsection, relational morphisms  $\varphi : X \rightarrow Y$  induce a functional morphism  $\check{\varphi} : X \rightarrow \Sigma(Y)$  given by  $x \mapsto \varphi(x, \_)$ .

Since  $Y$  is Scott complete, it is naturally isomorphic to  $\Sigma(Y)$  and hence each  $\check{\varphi}(x)$  corresponds to some  $y_{\check{\varphi}(x)}$  which is the unique point in  $y$  representing  $\check{\varphi}(x)$ . We claim that this mapping is a functional morphism  $X \rightarrow Y$  (easy) and that this action on morphisms is functorial, and assign it to  $\mathcal{F}_N$ ; we further claim that  $\mathcal{F}_N$  is the inverse of  $\mathcal{G}_r \upharpoonright_{\Sigma\mathcal{Q}\text{-Set}_f}^{\Sigma\mathcal{Q}\text{-Set}_r}$  – which is a nasty thing to typeset, so we’ll just say  $\mathcal{G}_r$ .

*Proof.* There is quite a lot to unpack; formally, we say that if we define  $\mathcal{F}_N(\varphi) = \varepsilon \circ \check{\varphi}$ , then this action is both functorial and makes  $\mathcal{F}_N$  and  $\mathcal{G}_r$  mutual inverses — where  $\varepsilon$  is the counit of the adjunction, which takes singletons over Scott-complete  $\mathcal{Q}$ -sets to their representing elements.

Let’s write, for a given  $\varphi$ ,  $\mathcal{F}_N(\varphi)$  as  $f_\varphi$  for simplicity sake. First, note that  $\varphi(x, f_\varphi(x)) = \mathbb{E} x$  — as  $\varphi(x, \_) = \delta(\_, f_\varphi(x))$  and  $\mathbb{E} f_\varphi(x) = \mathbb{E} x$ . Obviously,  $\mathcal{F}_N$  preserves identities, as

$$\begin{aligned} f_{\text{id}} &= \varepsilon(x \mapsto (\_, x)) \\ &= x \mapsto x \\ &= \text{id} \end{aligned}$$

functoriality can now be seen to hold: given  $x$ , consider the singleton

$$\begin{aligned}
\bigvee_y \varphi(x, y) \otimes \psi(y, \_ ) &\geq \varphi(x, f_\varphi(x)) \otimes \psi(f_\varphi(x), \_ ) \\
&= \mathbf{E} f_\varphi(x) \otimes \psi(f_\varphi(x), \_ ) \\
&= \psi(f_\varphi(x), \_ ) \\
&\geq \psi(y, \_ ) \otimes \delta(f_\varphi(x), y) && \text{(for any } y) \\
&= \psi(y, \_ ) \otimes \varphi(x, y)
\end{aligned}$$

and hence  $[\psi \circ \varphi](x, \_ ) = \psi(f_\varphi(x), \_ )$ , from which we can conclude that

$$f_{\psi \circ \varphi} = f_\psi \circ f_\varphi$$

We ought to show that  $\mathcal{G}_R \circ \mathcal{F}_N(\varphi) = \varphi$  and  $\mathcal{F}_N \circ \mathcal{G}_R(f) = f$ . Let's go about doing it in the order we've written. Take some relational morphism  $\varphi : X \rightarrow Y$  and let's consider the following:

$$\begin{aligned}
[\mathcal{G}_R \circ \mathcal{F}_N(\varphi)](x, y) &= [\mathcal{G}_R(\varepsilon \circ \check{\varphi})](x, y) \\
&= \delta(\varepsilon \circ \check{\varphi}(x), y) \\
&= \delta(\check{\varphi}(x), \sigma_y) \\
&= \check{\varphi}(x)(y) \\
&= \varphi(x, y)
\end{aligned}$$

Now take some functional morphism  $f : X \rightarrow Y$

$$\begin{aligned}
[\mathcal{F}_N \circ \mathcal{G}_R(f)](x) &= [\mathcal{F}_N(\delta(f(\_1), \_2))](x) \\
&= [f(\_1)](x) \\
&= f(x)
\end{aligned}$$

And therefore, we have shown that the composites are indeed actually the identities and  $\mathcal{F}_N$  witnesses the fact that  $\Sigma\mathcal{Q}\text{-Set}_r \cong \Sigma\mathcal{Q}\text{-Set}_f$ .  $\square$

**Lemma 6.1:**

$$\mathcal{G}_R \circ \Sigma \cong \mathcal{G}_R$$

Where  $\mathcal{G}_R \circ \Sigma$  really ought to be the composite of  $\Sigma$  with the appropriate restriction of  $\mathcal{G}_R$  to the full subcategory that is the image of  $\Sigma$ . Or, alternatively,  $\mathcal{G}_R \circ i \circ \Sigma$ .

*Proof.* We must provide an  $\alpha_X$  for every  $X$  such that

$$\begin{array}{ccc}
X & \xrightarrow{\alpha} & i \circ \Sigma(X) \\
\downarrow \mathcal{G}_R(f) & & \downarrow \mathcal{G}_R \circ i \circ \Sigma(f) \\
Y & \xrightarrow{\alpha} & i \circ \Sigma(X)
\end{array}$$

which simplifies to finding  $\alpha_X$  such that the following commutes, thus justifying our lack of precision in the “clean” statement of the lemma:

$$\begin{array}{ccc} X & \xrightarrow{\alpha} & \Sigma(X) \\ \downarrow \mathcal{G}_R(f) & & \downarrow \mathcal{G}_R \circ \Sigma(f) \\ Y & \xrightarrow{\alpha} & \Sigma(X) \end{array}$$

if we then let  $\alpha$  be the relational isomorphism we already know exists between  $X$  and  $\Sigma(X)$  there is a good chance naturality holds by magic. We need to verify that

$$\alpha \circ [\mathcal{G}_R(f)] = [\mathcal{G}_R \circ \Sigma(f)] \circ \alpha$$

We do so by *cheating*, and proving instead that

$$\alpha \circ [\mathcal{G}_R(f)] \geq [\mathcal{G}_R \circ \Sigma(f)] \circ \alpha$$

and then we remember that for relational morphisms  $\leq$  is = because  $\mathcal{Q}$  is **strong**.

$$\begin{aligned} [\mathcal{G}_R \circ \Sigma(f)] \circ \alpha(x, \psi) &= \bigvee_{\xi} \xi(x) \otimes \delta(\Sigma f(\xi), \psi) \\ &= \bigvee_{\xi} \xi(x) \otimes \bigvee_y \bigvee_x \delta(f(x), y) \otimes \xi(x) \otimes \psi(y) \\ &= \bigvee_y \bigvee_{x'} \bigvee_{\xi} \xi(x) \otimes \delta(f(x'), y) \otimes \xi(x') \otimes \psi(y) \\ &\leq \bigvee_y \bigvee_{x'} \delta(x, x') \otimes \delta(f(x'), y) \otimes \psi(y) \\ &\leq \bigvee_y \bigvee_{x'} \delta(f(x), f(x')) \otimes \delta(f(x'), y) \otimes \psi(y) \\ &\leq \bigvee_y \delta(f(x), y) \otimes \psi(y) \\ &= \alpha \circ [\mathcal{G}_R(f)](x, \psi) \end{aligned}$$

□

Now we are ready to state a more categorical description of Scott-completeness, merely in terms of representable functors *etc.*

**Theorem 6.4:**

Let  $\uparrow \mathcal{G}_R$  denote the functor between the presheaf categories of  $\mathcal{Q}\text{-Set}_r$  and  $\mathcal{Q}\text{-Set}_f$  as below:

$$\begin{array}{ccc} \mathcal{Q}\text{-Set}_f & \xrightarrow{\mathcal{G}_R} & \mathcal{Q}\text{-Set}_r \\ \downarrow \text{PSh} & & \downarrow \text{PSh} \\ \text{PSh}(\mathcal{Q}\text{-Set}_f) & \xleftarrow{\uparrow \mathcal{G}_R} & \text{PSh}(\mathcal{Q}\text{-Set}_r) \end{array}$$

Given by precomposing presheaves with the functor  $\mathcal{G}_r$  so as to change their domains appropriately. Then

$$[\mathfrak{y}_f(X) \cong \mathfrak{y}_r(X) \upharpoonright \mathcal{G}_r] \iff X \text{ is Scott-complete}$$

*Proof.* If  $X$  is indeed complete, then we know that the following are natural isomorphisms in  $Y$

$$\begin{aligned} \mathfrak{y}_f(X) &= \mathfrak{y}_f(R(X)) \\ &\cong \mathfrak{y}_f(X) \circ \Sigma \end{aligned} \quad (\text{adjunction})$$

and, as both entries are now complete,

$$= \Sigma\mathcal{Q}\text{-Set}_f(\Sigma(\_), X)$$

We also know that  $\mathcal{G}_r$  is an isomorphism between the subcategories  $\Sigma\mathcal{Q}\text{-Set}_f$  and  $\Sigma\mathcal{Q}\text{-Set}_r$ ; in particular we have

$$\Sigma\mathcal{Q}\text{-Set}_r(\mathcal{G}_r(\_1), \mathcal{G}_r(\_2)) \cong \Sigma\mathcal{Q}\text{-Set}_f(\_1, \_2)$$

if we fix  $\_2 = X$ , we can forget about the  $\mathcal{G}_r$  on the second coordinate (because it's there just to act on morphisms), thus getting

$$\Sigma\mathcal{Q}\text{-Set}_r(\mathcal{G}_r(\_), X) \cong \Sigma\mathcal{Q}\text{-Set}_f(\_, X)$$

Resuming our isomorphism chain:

$$\begin{aligned} \Sigma\mathcal{Q}\text{-Set}_f(\Sigma(\_), X) &\cong \Sigma\mathcal{Q}\text{-Set}_r(\mathcal{G}_r \circ \Sigma(\_), X) \\ &= \mathfrak{y}_r(X) \circ \mathcal{G}_r \circ \Sigma \end{aligned}$$

For our claim to hold, we would need a natural isomorphism  $\mathcal{G}_r \rightarrow \mathcal{G}_r \circ \Sigma$  – the reader should be happy to remember that this isomorphism has already been proven in an earlier lemma. We can then append one last isomorphism then:

$$\mathfrak{y}_r(X) \circ \mathcal{G}_r \circ \Sigma \cong \mathfrak{y}_r \circ \mathcal{G}_r$$

We can then put it all together and show there is some twisted little natural isomorphism

$$\mathfrak{y}_f(X) \cong \mathfrak{y}_r \circ \mathcal{G}_r = \mathfrak{y}_r(X) \upharpoonright \mathcal{G}_r$$

The converse implication is, thankfully, easier; suppose there is indeed an isomorphism as above; we ought to show that  $X$  was Scott-complete to begin with.

$$\begin{aligned} \mathfrak{y}_f(i \circ \Sigma(X)) &\cong \mathfrak{y}_r(i \circ \Sigma(X)) \circ \mathcal{G}_r && (\text{just proven}) \\ &\cong \mathfrak{y}_r(X) \circ \mathcal{G}_r && (X \cong_r i \circ \Sigma(X)) \\ &\cong \mathfrak{y}_f(X) && (\text{hypothesis}) \end{aligned}$$

thence, by Yoneda we obtain that

$$\Sigma(X) \cong_f X$$

But Scott-completeness is invariant under functional isomorphisms. So  $X$  was Scott-complete to begin with and thus ends our proof.  $\square$

We then give a useful tool to show Scott-completeness in terms of an object's representable functor. Namely

**Theorem 6.5:**

$$X \text{ is Scott-complete} \iff \mathcal{Q}\text{-Set}_f(\_, X) \cong \mathcal{Q}\text{-Set}_f(\Sigma(\_), X)$$

here again we actually wrote something slightly informal and we really mean that,  $\mathcal{Q}\text{-Set}_f(i \circ \Sigma(\_), X)$  is isomorphic to  $\mathcal{Q}\text{-Set}_f(\_, X)$ . We can rewrite it as

$$\mathfrak{y}_f(X) \cong \mathfrak{y}_f(X) \circ i \circ \Sigma$$

*Proof.* Scott-completeness implies the existence of the isomorphism since  $X$  as an object of  $\mathcal{Q}\text{-Set}_f$  is just  $i(X)$  for  $X$  an object of  $\Sigma\mathcal{Q}\text{-Set}$ , and hence adjunction applies:

$$\begin{aligned} \mathcal{Q}\text{-Set}_f(\_, X) &= \mathcal{Q}\text{-Set}_f(\_, i(X)) \\ &\cong \Sigma\mathcal{Q}\text{-Set}(\Sigma(\_), X) \\ &= \mathcal{Q}\text{-Set}_f(i \circ \Sigma(\_), i(X)) \\ &= \mathcal{Q}\text{-Set}_f(\Sigma(\_), X) \end{aligned}$$

Now, provided with the isomorphism but no guarantee that  $X$  is complete, we have to find sufficient reason for  $X$  to be complete.

$$\begin{aligned} \mathfrak{y}_f(X) &\cong \mathfrak{y}_f(X) \circ i \circ \Sigma \\ &= \Sigma\mathcal{Q}\text{-Set}_f(\Sigma(\_), X) \\ &\cong \Sigma\mathcal{Q}\text{-Set}_r(\mathfrak{G}_r \circ \Sigma(\_), \mathfrak{G}_r(X)) \\ &= \Sigma\mathcal{Q}\text{-Set}_r(\mathfrak{G}_r \circ \Sigma(\_), X) \\ &\cong \Sigma\mathcal{Q}\text{-Set}_r(\mathfrak{G}_r(\_), X) \\ &= \mathfrak{y}_r(X) \upharpoonright \mathfrak{G}_r \end{aligned}$$

$\square$

**Proposition 6.1:**

The true force of the theorem above is that  $i$  in  $\Sigma \dashv i$  not only preserves limits but indeed it creates them.

*Proof.* Take a diagram  $D$  on  $\Sigma\mathcal{Q}\text{-Set}$ ,

$$\begin{aligned} \mathfrak{y}_f(\lim i \circ D) &\cong \lim \mathfrak{y}_f(i \circ D) \\ &\cong \lim \mathfrak{y}_f(i \circ D) \circ i \circ \Sigma \\ &\cong \mathfrak{y}_f(\lim i \circ D) \circ i \circ \Sigma \end{aligned}$$



This shows that the external (ie. in  $\mathcal{Q}\text{-Set}_f$ ) the limit of complete  $\mathcal{Q}$ -sets is itself complete.

Since any cone for  $D$  – say, with vertex  $X$  – would be a cone for  $i \circ D$  (as  $i$  is simply an inclusion, they literally are) – by universality there is exactly one morphism  $\lim i \circ D \rightarrow X$  making everything commute.

Since the external limit is now known to be complete, by necessity, it also is the limit in  $\Sigma\mathcal{Q}\text{-Set}$  as well. Hence,  $i$  both preserves all limits from its domain, and reflects all limits in its codomain.  $\square$

## References

- [1] Alvim, J. G, Mendes, C. de A., and Mariano, H. L. “ $\mathcal{Q}\text{-Set}$  & Friends: Categorical Constructions and Properties”. 2023.
- [2] F. Borceux and G. van den Bossche. “Quantales and their sheaves”. In: *Order* 3 (1986), pp. 61–87.
- [3] Francis Borceux. *Handbook of Categorical Algebra: Volume 3, Sheaf Theory*. Encyclopedia of Mathematics and its Applications. Cambridge: Cambridge University Press, 1994.
- [4] Marcelo E Coniglio and Francisco Miraglia. “Non-commutative topology and quantales”. In: *Studia Logica* 65.2 (2000), pp. 223–236.
- [5] Michael P Fourman and Dana S Scott. “Sheaves and logic”. In: *Applications of sheaves*. Springer, 1979, pp. 302–401.
- [6] Hans Heymans and Isar Stubbe. “Grothendieck quantaloids for allegories of enriched categories”. In: *Bulletin of the Belgian Mathematical Society-Simon Stevin* 19.5 (2012), pp. 859–888.
- [7] Ulrich Höhle. “Monoidal closed categories, weak topoi and generalized logics”. In: *Fuzzy Sets and Systems* 42.1 (1991). Mathematical Aspects of Fuzzy Sets, pp. 15–35. ISSN: 0165-0114. DOI: [https://doi.org/10.1016/0165-0114\(91\)90086-6](https://doi.org/10.1016/0165-0114(91)90086-6). URL: <https://www.sciencedirect.com/science/article/pii/0165011491900866>.
- [8] Francisco Miraglia and Ugo Solitro. “Sheaves over right sided idempotent quantales”. In: *Logic Journal of IGPL* 6.4 (1998), pp. 545–600.
- [9] C. J. Mulvey and M. Nawaz. “Quantales: Quantal sets”. In: *Non-Classical Logics and their Applications to Fuzzy Subsets: A Handbook of the Mathematical Foundations of Fuzzy Set Theory*. Ed. by Ulrich Höhle and Erich Peter Klement. Dordrecht: Springer Netherlands, 1995, pp. 159–217. ISBN: 978-94-011-0215-5. DOI: 10.1007/978-94-011-0215-5\_8. URL: [https://doi.org/10.1007/978-94-011-0215-5\\_8](https://doi.org/10.1007/978-94-011-0215-5_8).
- [10] Chris Mulvey. “J. 1986”. In: *Suppl. Rend. Circ. Mat. Palermo Ser. II* 12 (), pp. 99–104.
- [11] *nLab Quantale entry*. <https://ncatlab.org/nlab/show/quantale>. accessed on 2022-10-25.
- [12] Pedro Resende. “Groupoid sheaves as quantale sheaves”. In: *Journal of Pure and Applied Algebra* 216.1 (2012), pp. 41–70.

- [13] David Reyes and Pedro H. Zambrano. “Co-quantale valued logics”. In: *arxiv preprint*, <https://arxiv.org/abs/2102.06067> (2021).
- [14] K.I. Rosenthal. *Quantales and Their Applications*. Pitman research notes in mathematics series. Longman Scientific & Technical, 1990. ISBN: 9780582064232. URL: <https://books.google.com.br/books?id=PrkrAAAAYAAJ>.
- [15] Ana Luiza Tenório, Caio de Andrade Mendes, and Hugo Luiz Mariano. “Introducing sheaves over commutative semicartesian quantales”. In: *arXiv preprint arXiv:2204.08351* (2022).
- [16] R. F. C. Walters. “Sheaves and Cauchy-complete categories”. In: *Cahiers de Topologie et Géométrie Différentielle Catégoriques* 22.3 (1981), pp. 283–286.
- [17] David N. Yetter. “Quantales and (noncommutative) linear logic”. In: *Journal of Symbolic Logic* 55.1 (1990), pp. 41–64. DOI: 10.2307/2274953.