

# Explicit formulas for hypermaps and maps

Ricky Xiao-Feng Chen \*

School of Mathematics, Hefei University of Technology

Hefei, Anhui 230601, P. R. China

xiaofengchen@hfut.edu.cn

## Abstract

The study of hypermaps and maps is ubiquitous, as they are closely connected with geometry, mathematical physics, free probability and algebra. A universal recursion generalizing a fundamental identity of Frobenius which enumerates factorizations of a permutation in group algebra theory has been discovered by the author recently. Here we apply the recursion to study hypermaps as well as maps and obtain a plethora of results in a unified way. For instance, we succinctly provide a short proof of the celebrated Harer-Zagier formula and present a general explicit formula for one-face hypermaps. As special cases of the latter, we obtain simple explicit formulas for the numbers of ways of expressing a long cycle as a product of a permutation of cycle-type  $[1^p, n-p]$  and, respectively,  $[p, n-p]$  and a permutation with  $m$  cycles for any  $p$  and  $m$ . To the best of our knowledge, only the cases for  $p = 0, 1$  are known before.

**Keywords:** Group characters, Permutation products, Hypermaps, Harer-Zagier formula, Polynomiality, Stirling numbers

Mathematics Subject Classifications 2020: 05E10, 20B30, 14H10

## 1 Introduction

The study of hypermaps and maps is ubiquitous, as they are closely connected with geometry, mathematical physics, free probability and algebra. Maps are a subclass of hypermaps. A map of genus  $g$  is a 2-cell embedding of a connected graph (consisting of vertices and multiedges connecting the vertices) in an orientable topological surface of genus  $g$  such that all faces are homomorphic to an open disk. The genus  $g$  of the topological surface relates to the number  $v$  of vertices,  $e$  of edges and  $f$  of faces via the Euler characteristic formula:

$$v - e + f = 2 - 2g.$$

---

\*ORCID: 0000-0003-1061-3049

An end of an edge is called a half edge or a dot, so there are  $2n$  half edges in a graph with  $n$  edges. It is long known that an embedding is completely determined by the cyclic orders of the half edges around the vertices (see e.g., [10]). Consequently, there is a well established algebraic formulation of maps which works as follows. If we assign the numbers in  $[2n] = \{1, 2, \dots, 2n\}$  to the half edges, then two permutations on  $[2n]$  immediately arise: the cycles of one permutation come from the half edges around the vertices of the graph and the other consists of  $n$  cycles where the two half edges of an edge give a length-two cycle. The product of these two permutations actually encode the faces of the embedding. As such, enumeration of maps can be translated into permutation product problems. The genus  $g$  of the underlying surface is then encoded in the numbers of cycles of the involved permutations. Hypermaps are merely generalization of maps by including hyperedges [24], that is, the cycles of the second permutation could be of arbitrary lengths. If one of the half edges is particularly distinguished, the hypermaps are called rooted. All hypermaps are rooted in this work.

The enumeration of maps may date back to Tutte in the sixties in a series of papers about planar maps (i.e., genus zero maps). Explicit formulas for maps of arbitrary genus were first obtained by Walsh and Lehman [23]. In the study of moduli space of curves, the celebrated Harer-Zagier formula and recurrence satisfied by the numbers of maps were obtained via integral over matrices [16]. Making use of the permutation product model, Jackson [17] first provided an approach to compute these numbers using the symmetric group character theory, and Zagier [25] later also provided a group character approach but shorter. Very recently, Pittel [20] gave another derivation of the Harer-Zagier formula via group characters in the form of Fourier transform. Combinatorial approaches attacking this topic have also received lots of interests. We refer to Bernardi [2], Chapuy [3], Goupil and Schaeffer [13], Goulden and Nica [14], Morales and Vassilieva [19], Chen and Reidys [7], Chen [5] and the references therein.

A universal recursion generalizing a fundamental identity of Frobenius which enumerates factorizations of a permutation in group algebra theory has been discovered by the author recently. Many permutation product problems may be studied by exploring the universal recursion. Along this line, we focus on the application of the recursion to the enumeration of hypermaps and maps.

The organization of the paper is as follows. In Section 2, we introduce relevant notation and present a brief review of the universal recursion about permutation products. We enumerate hypermaps of an arbitrary genus  $g$  which have one face and hyperedges of a given cycle-type  $(\alpha_1, \dots, \alpha_d)$  in Section 3. As a consequence, we obtain a plethora of results in a unified way. For instance, we succinctly provide a short proof of the celebrated Harer-Zagier formula and present a general explicit formula for one-face hypermaps. As special cases of the latter, we obtain simple explicit formulas for the numbers of ways of expressing a long cycle as a product of a permutation of cycle-type  $[1^p, n - p]$  and, respectively,  $[p, n - p]$  and a permutation with  $m$  cycles for any  $p$  and  $m$ . To the best of our knowledge, only the cases for  $p = 0, 1$  are known before. For example, the case for  $p = 0$  is given by the Zagier-Stanley formula [22, 25], and the case for  $p = 1$  can be derived from a result of Bocarra [1]. In addition, we show that the hypermap numbers are polynomials in  $\alpha_i$ 's. All coefficients there can be extracted if desired.

## 2 Review of the universal recursion

Denote by  $\mathfrak{S}_n$  the symmetric group on  $[n] := \{1, 2, \dots, n\}$ . A permutation  $\pi$  can be written as a product of its disjoint cycles, i.e., the orbits of the cyclic group generated by  $\pi$  acting on  $[n]$ . The length distribution of the cycles of  $\pi$  is called the cycle-type of  $\pi$  and denoted by  $ct(\pi)$ . One usually writes  $ct(\pi)$  as an (integer) partition of  $n$ . A partition  $\lambda$  of  $n$ , denoted by  $\lambda \vdash n$ , is usually represented by a nonincreasing positive integer sequence  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  such that  $\lambda_1 + \lambda_2 + \dots + \lambda_k = n$ . The number  $k$  is called the length of  $\lambda$ , denoted by  $\ell(\lambda)$ . Then,  $ct(\pi) = \lambda$  means the multiset consisting of the lengths of the cycles of  $\pi$  is the same as the multiset  $\{\lambda_1, \dots, \lambda_k\}$ . Another representation of  $\lambda$  is in the form  $\lambda = [1^{m_1}, 2^{m_2}, \dots, n^{m_n}]$  indicating that there are  $m_i$  of  $i$ 's in the partition. We usually discard the entry  $i^{m_i}$  in case  $m_i = 0$  and write the entry  $i^1$  simply as  $i$ . Obviously,  $\sum_{i=1}^n im_i = n$  and  $\ell(\lambda) = \sum_{i=1}^n m_i$ . One more relevant quantity, denoted by  $Aut(\lambda)$ , is the number of permutations of the entries of  $\lambda$  that fix  $\lambda$ . Clearly,  $Aut(\lambda) = \prod_{i=1}^n m_i!$ . In this paper, we will use the two representations of partitions interchangeably, whichever is more convenient.

It is well known that a conjugacy class of  $\mathfrak{S}_n$  contains permutations of the same cycle-type. So, the conjugacy classes can be indexed by partitions of  $n$ . Let  $\mathcal{C}_\lambda$  denote the one indexed by  $\lambda$ . If  $\pi \in \mathcal{C}_\lambda$ , then the number of cycles contained in  $\pi$  is  $\ell(\pi) = \ell(\lambda)$ . Moreover, the number of elements contained in  $\mathcal{C}_\lambda$  is well known to be  $|\mathcal{C}_\lambda| = \frac{n!}{z_\lambda}$  where

$$z_\lambda = \prod_{i=1}^n i^{m_i} m_i!$$

Recall the generating function of the signless Stirling numbers of the first kind  $\left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right]$  is given by:

$$\sum_{k=1}^n (-1)^{n-k} \left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right] x^k = x(x-1) \cdots (x-n+1).$$

The number  $\left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right]$  counts permutations on  $[n]$  having exactly  $k$  cycles.

From the representation theory of the symmetric group  $\mathfrak{S}_n$ , we know that the number of irreducible representations is the same as the number of its conjugacy classes. Consequently, we can index the irreducible representations by the partitions of  $n$  as well. We write the character associated to the irreducible representation indexed by  $\lambda$  as  $\chi^\lambda$  and the dimension of the irreducible representation as  $f^\lambda$ . Suppose  $C$  is a conjugacy class of  $\mathfrak{S}_n$  indexed by  $\alpha$  and  $\pi \in C$ . We view  $\chi^\lambda(\pi)$ ,  $\chi^\lambda(\alpha)$ ,  $\chi^\lambda(C)$  as the same, and we trust the context to prevent confusion.

Let

$$m_{\lambda,m} = \prod_{u \in \lambda} \frac{m + c(u)}{h(u)}, \quad c_{\lambda,m} = \sum_{d=0}^m (-1)^d \binom{m}{d} m_{\lambda,m-d}.$$

where for  $u = (i, j) \in \lambda$ , i.e., the  $(i, j)$  cell in the Young diagram of  $\lambda$ ,  $c(u) = j - i$  and  $h(u)$  is the hook length of the cell  $u$ .

**Theorem 2.1** (Chen [9]). *Let  $\xi_{n,m}(C_1, \dots, C_t)$  be the number of tuples  $(\sigma_1, \sigma_2, \dots, \sigma_t)$  such that the permutation  $\pi = \sigma_1 \sigma_2 \cdots \sigma_t$  has  $m$  cycles, where  $\sigma_i$  belongs to a conjugacy class  $C_i$ . Then we have*

$$\xi_{n,m}(C_1, \dots, C_t) = \sum_{k \geq 0} (-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix} W_{n,m+k}(C_1, \dots, C_t), \quad (1)$$

where

$$W_{n,m+k}(C_1, \dots, C_t) = \frac{\prod_{i=1}^t |C_i|}{(m+k)!} \sum_{\lambda \vdash n} \epsilon_{\lambda, m+k} \{f^\lambda\}^{-t+1} \prod_{i=1}^t \chi^\lambda(C_i). \quad (2)$$

The summation in eq. (2) can be simplified if one of the conjugacy classes corresponds to permutations with only one cycle. This is presented in the following proposition.

**Proposition 2.1** (Chen [9]).

$$W_{n,m}(\mathcal{C}_{(n)}, C_1, \dots, C_t) = \frac{(n-1)! \prod_{i=1}^t |C_i|}{m!} \sum_{j=0}^{n-1} (-1)^j \frac{\binom{n-1-j}{n-m}}{\binom{n-1}{j}^{t-1}} \prod_{i=1}^t \chi^{[1^j, n-j]}(C_i). \quad (3)$$

## 3 Hypermaps and maps

### 3.1 One-face hypermaps

Continuing the introduction in Section 1, a (rooted) one-face hypermap with  $n$  half edges is a triple of permutations  $(s, \gamma, \pi)$  on  $[n]$  such that  $s = \gamma\pi$  and  $s$  is a fixed permutation in  $\mathcal{C}_{(n)}$ , say the cyclic permutation  $(1 \ 2 \ \cdots \ n)$ . If  $n = 2k$  and  $\gamma \in \mathcal{C}_{[2^k]}$ , the hypermap is really a rooted one-face map. The (combinatorial) genus  $g$  of the hypermap is determined by

$$2 - 2g = \ell(s) + \ell(\gamma) + \ell(\pi) - n.$$

We are now interested in enumerating all one-face hypermaps  $(s, \gamma, \pi)$  of  $n$  half edges and genus  $g$  where the cycle-type of the hyperedge  $\gamma$  is  $(\gamma_1, \dots, \gamma_d)$ . By definition of genus,  $\ell(\pi) = m = 1 - 2g + n - d$ . It is more convenient to use  $u_{n,m}(\gamma)$  to denote the desired number and make the genus  $g$  implicit in our setting.

**Theorem 3.1** (Hypermap formula). *Let  $\gamma = (\gamma_1, \dots, \gamma_d)$  and  $m = 1 - 2g + n - d$ . Then,*

$$\mu_{n,m}(\gamma) = |\mathcal{C}_\gamma| \sum_{k=0}^{n-m} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \sum_{\substack{k_1 + \dots + k_d = n-m-k+1 \\ k_1 > 0, \dots, k_d > 0}} \prod_{i=1}^d \binom{\gamma_i}{k_i}. \quad (4)$$

*Proof.* In the light of Theorem 2.1, this comes down to computing the  $W$ -number for  $t = 2$ ,  $C_1 = \mathcal{C}_{(n)}$ ,  $C_2 = \mathcal{C}_\gamma$  with  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_d)$ , and we have

$$\begin{aligned}
W_{n,m+k} &= \frac{(n-1)! |\mathcal{C}_\gamma|}{(m+k)!} \sum_{j=0}^{n-1} (-1)^j \binom{n-1-j}{n-m-k} \chi^{[1^j, n-j]}(C_2) \\
&= \frac{(n-1)! |\mathcal{C}_\gamma|}{(m+k)!} \sum_{j=0}^{n-1} [y^{n-m-k}] (1+y)^{n-1-j} (-1)^j \chi^{[1^j, n-j]}(C_2) \\
&= \frac{(n-1)! |\mathcal{C}_\gamma|}{(m+k)!} [y^{n-m-k+1}] \prod_{i=1}^d \{(1+y)^{\gamma_i} - 1\} \\
&= \frac{(n-1)! |\mathcal{C}_\gamma|}{(m+k)!} \sum_{\substack{k_1 + \dots + k_d = n-m-k+1, \\ k_1 > 0, \dots, k_d > 0}} \prod_{i=1}^d \binom{\gamma_i}{k_i}.
\end{aligned}$$

Note that  $\mu_{n,g}(\gamma)$  is  $\frac{1}{(n-1)!}$  times the corresponding  $\xi$ -number. The rest is clear, and the proof is complete.  $\square$

Immediately from eq. (4), we obtain the following unexpected simple formula.

**Corollary 3.1** (Genus-zero hypermaps). *Let  $\gamma = [1^{a_1}, \dots, n^{a_n}]$ . Then, the number of genus zero hypermap with hyper-edge type  $\gamma$  is given by*

$$\frac{n!}{a_1! \cdots a_n! [n+1-\ell(\gamma)]!}. \quad (5)$$

*Proof.* When  $g = 0$ , the number  $m = n+1-d$  where  $d = \ell(\gamma)$ . As a result, in the summations in eq. (4), only the case  $k = 0$  and  $k_i = 1$  matters. Therefore,

$$\mu_{n,m}(\gamma) = \frac{n!}{1^{a_1} \cdots n^{a_n} a_1! \cdots a_n!} \frac{1}{(n+1-d)!} \gamma_1 \cdots \gamma_d = \frac{n!}{a_1! \cdots a_n! [n+1-d]}!$$

and the proof follows.  $\square$

In the following, we apply the notation:

$$\begin{aligned}
\tilde{\mu}_{n,m}(\gamma) &= \frac{m!}{n!} \mu_{n,m}(\gamma), \\
\tilde{S}_{m,i}(l) &= \sum_{j=1}^i \binom{i}{j} \frac{(m+j-i)! S(l, m+j-i)}{l!}.
\end{aligned}$$

Let  $\gamma = [1^{a_1}, \dots, n^{a_n}]$  and suppose  $a_i > 0$ . We also define

$$\gamma^{(i)} := [1^{a_1}, \dots, (i-1)^{a_{i-1}}, i^{a_i-1}, (i+1)^{a_{i+1}}, \dots, n^{a_n}].$$

Now, we are in a position to present the dimension-reduction formula for hypermaps.

**Theorem 3.2.** *Let  $\gamma = [1^{a_1}, \dots, n^{a_n}]$ . Suppose  $a_i > 0$ . Then,*

$$\tilde{\mu}_{n,m}(\gamma) = - \sum_{l>m} \tilde{S}_{m,1}(l) \tilde{\mu}_{n,l}(\gamma) + \frac{1}{ia_i} \sum_{l>0} \tilde{S}_{m,i}(l) \tilde{\mu}_{n-i,l}(\gamma^{(i)}). \quad (6)$$

In view of the dimension-reduction formula, all one-face hypermap numbers are eventually determined by the hypermap numbers of the form  $\mu_{n,m}([n])$ , i.e., the Zagier-Stanley formula (Cor. 3.2). Theorem 3.2 also allows us to compute one-face hypermap numbers  $\mu_{n,m}(\gamma)$  for all  $n, m$  and  $\gamma$  in an efficient recursive way.

We point out that there is a recursion for  $\mu_{n,m}(\gamma)$  in Chen and Reidys [7]. However, that recursion is recursive in both the parameters  $g$  (i.e.,  $m$ ) and  $\gamma$ . Moreover, a recursion for the numbers refining  $\mu_{n,m}(\gamma)$  (i.e., tracking the cycle-type of  $\pi$  as well) is also presented there. Note that hypermaps can be translated into bicolored (bipartite) maps, see Walsh [24]. As such, there is a summation formula for  $u_{n,m}(\gamma)$  in Goupil and Schaeffer [13, Theorem 4.1]. Comparatively speaking, our formula eq. (4) may be arguably simpler. A generating function for the refinement numbers is given in Morales and Vassilieva [19].

Simple formulas for  $\mu_{n,m}(\gamma)$  without complicated summations (e.g., over partitions) are rare. To the best of our knowledge, only the cases for  $\gamma = (n)$ ,  $\gamma = (n-1, 1)$  and  $\gamma = [2^n]$  have relatively simple explicit formulas. However, by exploring our general formula eq. (4), we may derive some new simple explicit formulas. We next present two such results as an illustration.

**Theorem 3.3.** *Let  $\gamma = [1^p, n-p]$  for  $p \geq 0$ . For  $m > 0$ , the following is true:*

$$\mu_{n,m}(\gamma) = \frac{1 + (-1)^{n-p-m}}{(n+1)(n-p)} \binom{n+1}{p} \begin{bmatrix} n+1-p \\ m \end{bmatrix}. \quad (7)$$

*Proof.* We first compute their generating function as follows:

$$\begin{aligned} & 1 + \sum_n \sum_{m>0} \frac{\mu_{n,m}(\gamma)}{|\mathcal{C}_\gamma|} x^m y^{n+1-p} \\ &= 1 + \sum_n \sum_{m>0} \sum_{k=0}^{n-m-p} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \binom{n-p}{n-m-k+1-p} x^m y^{n+1-p} \\ &= 1 + \sum_n \sum_{i \geq 1} \binom{x}{i} \binom{n-p}{n-i+1-p} y^{n+1-p} - \sum_n \binom{x}{n+1-p} y^{n+1-p} \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} y^i \sum_n \binom{n-p}{i-1} y^{n+1-p-i} - (1+y)^x + \delta_{p,1} \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} y^i \frac{1}{(1-y)^i} - (1+y)^x + \delta_{p,1} \\ &= (1-y)^{-x} - (1+y)^x + \delta_{p,1}. \end{aligned}$$

The coefficient of the term  $y^{n+1-p}$  on the RHS is

$$\binom{-x}{n+1-p} (-1)^{n+1-p} - \binom{x}{n+1-p}.$$

The coefficient of the term  $x^m$  in the last formula is given by

$$\frac{\begin{bmatrix} n+1-p \\ m \end{bmatrix} - (-1)^{n+1-p-m} \begin{bmatrix} n+1-p \\ m \end{bmatrix}}{(n+1-p)!}.$$

Hence, we obtain

$$\mu_{n,m}(\gamma) = \frac{\begin{bmatrix} n+1-p \\ m \end{bmatrix} - (-1)^{n+1-p-m} \begin{bmatrix} n+1-p \\ m \end{bmatrix}}{(n+1-p)!} \frac{n!}{p!(n-p)},$$

completing the proof.  $\square$

**Theorem 3.4.** Let  $m > 0$  and  $\gamma = [p, n-p]$  for  $p \geq 1$ . If  $p \neq n-p$ , then

$$\mu_{n,m}(\gamma) = \frac{n!}{p(n-p)} \sum_{j=1}^n \frac{(-1)^{n-j} + (-1)^{j-m+1}}{j!} \binom{p}{n+1-j} \begin{bmatrix} j \\ m \end{bmatrix}. \quad (8)$$

If  $n = 2p$ , we have

$$\mu_{n,m}(\gamma) = \frac{n!}{2p^2} \sum_{j=1}^n \frac{(-1)^{n-j} + (-1)^{j-m+1}}{j!} \binom{p}{n+1-j} \begin{bmatrix} j \\ m \end{bmatrix}. \quad (9)$$

*Proof.* We First compute

$$\begin{aligned} & 1 + \sum_n \sum_{m>0} \sum_{k=0}^{n-m} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \sum_{j=1}^{n-m-k} \binom{p}{j} \binom{n-p}{n-m-k+1-j} x^m y^{n+1} \\ &= 1 + \sum_n \sum_{m>0} \sum_{k=0}^{n-m} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \left\{ \sum_{j \geq 1} \binom{p}{j} \binom{n-p}{n-m-k+1-j} - \binom{p}{n-m-k+1} \right\} x^m y^{n+1} \\ &= 1 + \sum_n \sum_{m>0} \left\{ \sum_{i \geq 1} \frac{(-1)^{i-m} \begin{bmatrix} i \\ m \end{bmatrix}}{i!} \left\{ \sum_{j \geq 1} \binom{p}{j} \binom{n-p}{n-i+1-j} - \binom{p}{n-i+1} \right\} \right. \\ & \quad \left. + \frac{(-1)^{n+1-m} \begin{bmatrix} n+1 \\ m \end{bmatrix}}{(n+1)!} \right\} x^m y^{n+1} \end{aligned}$$

Summing over  $m > 0$  first and using the generating function for the signless Stirling numbers of the first kind, the last expression equals

$$\begin{aligned} & 1 + \sum_n \sum_{i \geq 1} \binom{x}{i} \left\{ \sum_{j \geq 1} \binom{p}{j} \binom{n-p}{n-i+1-j} - \binom{p}{n-i+1} \right\} y^{n+1} + \sum_n \binom{x}{n+1} y^{n+1} \\ &= 1 + \sum_n \sum_{i \geq 1} \binom{x}{i} \left\{ \sum_{j \geq 1} \binom{p}{j} \binom{n-p}{n-i+1-j} \right\} y^{n+1} - \{(1+y)^x - 1\} (1+y)^p + (1+y)^x - 1 \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} \left\{ \sum_{j \geq 1} \binom{p}{j} y^{i+j} \frac{1}{(1-y)^{i+j-p}} \right\} - \{(1+y)^x - 1\} (1+y)^p + (1+y)^x - 1 \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} y^i \frac{1}{(1-y)^{i-p}} \{(1-y)^{-p} - 1\} - \{(1+y)^x - 1\} (1+y)^p + (1+y)^x - 1 \\ &= 1 + \{(1-y)^{-x} - 1\} (1-y)^p \{(1-y)^{-p} - 1\} - \{(1+y)^x - 1\} (1+y)^p + (1+y)^x - 1 \\ &= 1 + \{(1-y)^{-x} - 1\} \{1 - (1-y)^p\} - \{(1+y)^x - 1\} \{(1+y)^p - 1\}. \end{aligned}$$

In view of Theorem 2.1, the last formula and

$$1 + \sum_n \sum_{m>0} \frac{\mu_{n,m}(\gamma)}{|\mathcal{C}_\gamma|} x^m y^{n+1}$$

may differ by a polynomial in  $x$  and  $y$  where the highest degree of  $y$  is at most  $p + 1$ . This is caused by the fact that we summed over all nonnegative integers for  $n$ . Nevertheless, for our purpose here, it is satisfactory enough. The coefficient of the term  $y^{n+1}$  on the RHS of the second last formula is

$$\sum_{j=1}^n \binom{-x}{j} (-1)^j (-1)^{n-j} \binom{p}{n+1-j} - \sum_{j=1}^n \binom{x}{j} \binom{p}{n+1-j}.$$

The coefficient of the term  $x^m$  in the last formula is next obtained as

$$\begin{aligned} & \sum_{j=1}^n \binom{p}{n+1-j} \frac{(-1)^{n-j} [j]_m - (-1)^{j-m} [j]_m}{j!} \\ &= \sum_{j=1}^n \frac{(-1)^{n-j} + (-1)^{j-m+1}}{j!} \binom{p}{n+1-j} [j]_m. \end{aligned}$$

Multiplying the last number by  $|\mathcal{C}_\gamma|$  complete the proof.  $\square$

In eq. (7), setting  $p = 0$ , we recover the Zagier-Stanley formula below.

**Corollary 3.2** (Zagier-Stanley formula [22, 25]). *The number of ways of expressing a fixed  $n$ -cycle as a product of an  $n$ -cycle and a permutation with  $m$  cycles is given by*

$$\left[ \begin{matrix} n+1 \\ m \end{matrix} \right] / \binom{n+1}{2}$$

if  $n - m$  is even, and 0 otherwise.

Stanley [22] once asked for combinatorial proofs of Corollary 3.2. Various proofs of combinatorial nature have been presented next. For instance, Cori, Marcus and Schaeffer [4], Chen and Reidys [7], and Chen [5]. In particular, a versatile combinatorial approach was presented in Chen [5] to deal with a variety of problems concerning products of two long cycles. In Féray and Vassilieva [12], a refinement of the problem considered in Corollary 3.2 was studied, that is, enumerating the pairs of long cycles whose product has a given cycle-type. A simple quantitative relation was obtained in [12] and was given an alternative simpler proof in [6]. It would be very interesting to know any combinatorial proofs of our general cases, eq. (7) and eq. (8).

In either eq. (7) or eq. (8), setting  $p = 1$ , we obtain the following formula which can be also obtained by applying a result of Bocarra [1]. The latter states that there are  $2(n-2)!$  ways for writing any odd permutation as a product of an  $n$ -cycle and a permutation of  $(n-1)$ -cycle.

**Corollary 3.3.** *The number of ways of expressing a fixed  $n$ -cycle as a product of a permutation in  $\mathcal{C}_{(n-1,1)}$  and a permutation with  $m$  cycles is given by*

$$\frac{2}{n-1} \left[ \begin{matrix} n \\ m \end{matrix} \right].$$

if  $n - m$  is odd, and 0 otherwise.



**Theorem 3.5.** Let  $m > 0$ , and  $\gamma = [1^t, p, n - p - t]$  for  $p \geq 1$  and  $t \geq 0$ . Then,  $\frac{\mu_{n,m}(\gamma)}{|\mathcal{C}_\gamma|}$  is given by

$$\sum_{j=1}^{n-t} \frac{(-1)^{n-j-t} - (-1)^{j-m}}{j!} \binom{p}{n+1-j-t} \begin{bmatrix} j \\ m \end{bmatrix} + \frac{(-1)^{n-t-m} \begin{bmatrix} n+1-t \\ m \end{bmatrix}}{(n+1-t)!} - \frac{(-1)^{n-m} \begin{bmatrix} n+1 \\ m \end{bmatrix}}{(n+1)!} \delta_{t,0}.$$

When  $t = 0$ , the above theorem reduces to Theorem 3.4.

Finally, we have the following polynomial property for hypermap numbers.

**Theorem 3.6 (Polynomiality).** For fixed  $g, n$  and  $d$ , the weighted number  $\frac{\text{Aut}(\gamma)}{n!} \mu_{n,m}(\gamma)$  for  $m = n - 2g + d - 1$  is a symmetric polynomial in  $\gamma_1, \gamma_2, \dots, \gamma_d$  with the highest degree  $2g$ .

*Proof.* According to eq. (4), we first have

$$\frac{\text{Aut}(\gamma)}{n!} \mu_{n,g}(\gamma) = \frac{1}{\gamma_1 \cdots \gamma_d} \sum_{k=0}^{n-m} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \sum_{\substack{k_1 + \dots + k_d = n-m-k+1, \\ k_1 > 0, \dots, k_d > 0}} \prod_{i=1}^d \binom{\gamma_i}{k_i}.$$

Clearly, each summand for  $k$  is either zero or a polynomial in  $\gamma_1, \dots, \gamma_d$  where the total degree is  $n - m - k + 1$  and the power of  $\gamma_i$  for any  $1 \leq i \leq d$  is at least one. Consequently,  $\frac{\text{Aut}(\gamma)}{n!} \mu_{n,g}(\gamma)$  is a symmetric polynomial in  $\gamma_1, \gamma_2, \dots, \gamma_d$  with the highest degree being  $(n - m + 1 + d) - d = 2g$ . This completes the proof.  $\square$

## 3.2 The Harer-Zagier formula

In this part, we focus on the case for  $\gamma = [2^n]$ . From eq. (4), we immediately obtain a new formula counting the number of one-face maps by genus.

**Theorem 3.7 (Map formula).** For any  $n$  and  $g$ , the number  $\xi_{n,g}$  of one-face maps of  $n$  edges and genus  $g$  is given by

$$\xi_{n,g} = (2n-1)!! \sum_{k \geq 0} \frac{(-1)^k \begin{bmatrix} m+k \\ m \end{bmatrix}}{(m+k)!} \binom{n}{m+k-1} 2^{m+k-1}, \quad (10)$$

where  $m = n + 1 - 2g$ . In particular,  $\xi_{n,0} = \frac{1}{n+1} \binom{2n}{n}$  is the famous  $n$ -th Catalan number.

*Proof.* In this case, the number  $m = n + 1 - 2g$ , and

$$\begin{aligned} W_{2n,m+k} &= \frac{(2n-1)!(2n)!}{(m+k)!n!2^n} [y^{2n-m-k+1}] \{y^2 + 2y\}^n \\ &= \frac{(2n-1)!(2n)!}{(m+k)!n!2^n} \binom{n}{x} 2^x \end{aligned}$$

for  $x + 2(n-x) = 2n - m - k + 1$ . Solve to obtain  $x = k + m - 1$ , and the proof follows.  $\square$

**Corollary 3.4** (Harer-Zagier formulas). *For  $n > 0$  and  $g \geq 0$ , we have*

$$\sum_{g \geq 0} \xi_{n,g} x^{n+1-2g} = (2n-1)!! \sum_{i \geq 1} \binom{x}{i} \binom{n}{i-1} 2^{i-1}, \quad (11)$$

$$1 + 2 \sum_{n \geq 0} \sum_{g \geq 0} \frac{\xi_{n,g}}{(2n-1)!!} x^{n+1-2g} y^{n+1} = \left( \frac{1+y}{1-y} \right)^x. \quad (12)$$

*Proof.* Based on eq. (10), we have

$$\begin{aligned} \sum_{m > 0} \frac{\mu_{n,m}}{(2n-1)!!} x^m &= \sum_{m > 0} \sum_{k=0}^{n-m} \frac{(-1)^k \binom{m+k}{m}}{(m+k)!} \binom{n}{m+k-1} 2^{m+k-1} x^m \\ &= \sum_{m > 0} \sum_{i \geq 1} \frac{(-1)^{i-m} \binom{i}{m}}{(i)!} \binom{n}{i-1} 2^{i-1} x^m \\ &= \sum_{i \geq 1} \binom{x}{i} \binom{n}{i-1} 2^{i-1}, \end{aligned}$$

whence the first formula. We proceed to compute as follows.

$$\begin{aligned} 1 + 2 \sum_{n \geq 0} \sum_{m > 0} \frac{\mu_{n,m}}{(2n-1)!!} x^m y^{n+1} &= 1 + \sum_{n \geq 0} \sum_{i \geq 1} \binom{x}{i} \binom{n}{i-1} 2^i y^{n+1} \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} 2^i y^i \sum_{n \geq i-1} \binom{n}{i-1} y^{n+1-i} \\ &= 1 + \sum_{i \geq 1} \binom{x}{i} 2^i y^i \frac{1}{(1-y)^i} \\ &= \left( \frac{1+y}{1-y} \right)^x. \end{aligned}$$

This completes the proof of the corollary.  $\square$

As for the numbers  $\xi_{n,g}$ , the celebrated Harer-Zagier recurrence [16] provides a three term recurrence recursive in both  $n$  and  $g$  while Chapuy [3] has obtained the first recurrence recursive only in  $g$ . Our recurrence here is apparently also only recursive in  $g$ . The first explicit formula for  $\xi_{n,g}$  was obtained by Walsh and Lehman [23]. Chapuy's recursion can of course yield an explicit formula. Some relation between these previous formulas, like deriving one from another, and some new expressions can be found in [8]. More generally, for  $\gamma = [p^n]$ , we have

$$\begin{aligned} W_{np,m} &= \frac{(np-1)!(np)!}{m!n!p^n} [y^{np-m+1}] \{(1+y)^p - 1\}^n \\ &= \frac{(np-1)!(np)!}{m!n!p^n} \sum_{i=0}^n (-1)^i \binom{n}{i} \binom{p(n-i)}{np-m+1}. \end{aligned}$$

In particular, for the extreme case  $m = n(p-1) + 1$ , we have the well-known (generalized) Catalan number expression [15, 17].

$$\mu_{np,m}([p^n]) = \frac{1}{(np-1)!} W_{np,m} = \frac{1}{(np-1)!(n(p-1)+1)!n!} = \frac{1}{n(p-1)+1} \binom{np}{n}.$$

## Acknowledgements

The work was partially supported by the Anhui Provincial Natural Science Foundation of China (No. 2208085MA02) and Overseas Returnee Support Project on Innovation and Entrepreneurship of Anhui Province (No. 11190-46252022001).

## References

- [1] G. Boccara, Nombre de représentations d'une permutation comme produit de deux cycles de longueurs données, *Discrete Math.* 29 (1980), 105–134.
- [2] O. Bernardi, An analogue of the Harer-Zagier formula for unicellular maps on general surfaces, *Adv. Appl. Math.* 48 (2012) 164–180.
- [3] G. Chapuy, A new combinatorial identity for unicellular maps, via a direct bijective approach, *Adv. Appl. Math.* 47 (2011) 874–893.
- [4] R. Cori, M. Marcus, G. Schaeffer, Odd permutations are nicer than even ones, *European J. Combin.* 33(7) (2012), 1467–1478.
- [5] R.X.F. Chen, A versatile combinatorial approach of studying products of long cycles in symmetric groups, *Adv. Appl. Math.* 133 (2022) 102283.
- [6] R. X. F. Chen, Combinatorially refine a Zagier-Stanley result on products of permutations, *Discrete Math.* 343(8) (2020), Article 111912.
- [7] R.X.F. Chen, C.M. Reidys, Plane permutations and applications to a result of Zagier-Stanley and distances of permutations, *SIAM J. Discrete Math.* 30(3) (2016) 1660–1684.
- [8] R.X.F. Chen, C.M. Reidys, New formulas counting one-face maps and Chapuy's recursion, arXiv:1510.05038.
- [9] R.X.F. Chen, A universal recursion with an application to studying the structure of triple Hurwitz numbers, submitted.
- [10] J. Edmonds, A combinatorial representation for polyhedral surfaces, *Notices Amer. Math. Soc.* 7 (1960), A646.
- [11] V. Féray, A. Rattan, On products of long cycles: short cycle dependence and separation probabilities, *J. Algebraic Comb.* 42 (2015), 183–224.
- [12] V. Féray, E. A. Vassilieva, Bijective enumeration of some colored permutations given by the product of two long cycles, *Discrete Math.* 312(2) (2012), 279–292.
- [13] A. Goupil, G. Schaeffer, Factoring  $n$ -cycles and counting maps of given genus, *European J. Combin.* 19(7) (1998) 819–834.

- [14] I.P. Goulden, A. Nica, A direct bijection for the Harer-Zagier formula, *J. Combin. Theory Ser. A* 111 (2005) 224–238.
- [15] I.P. Goulden, D.M. Jackson, The combinatorial relationship between trees, cacti and certain connection coefficients for the symmetric group, *European J. Combin.* 13 (1992) 357–365.
- [16] J. Harer, D. Zagier, The Euler characteristics of the moduli space of curves, *Invent. Math.* 85 (1986) 457–485.
- [17] D.M. Jackson, Counting cycles in permutations by group characters, with an application to a topological problem, *Trans. Amer. Math. Soc.* 299(2) (1987) 785–801.
- [18] D.M. Jackson, Some combinatorial problems associated with products of conjugacy classes of the symmetric group, *J. Combin. Theory Ser. A* 49 (1988) 363–369,
- [19] A.H. Morales, E.A. Vassilieva, Bijective Enumeration of Bicolored Maps of Given Vertex Degree Distribution, *Discrete Math. Theoret. Comput. Sci.* (2009) 661–672.
- [20] B. Pittel, Another Proof of the Harer-Zagier Formula, *Electron. J. Combin.* 23 (2016) 1–21.
- [21] R. P. Stanley, Factorization of permutations into  $n$ -cycles, *Discrete Math.* 37 (1981), 255–261.
- [22] R. P. Stanley, Two enumerative results on cycles of permutations, *European J. Combin.* 32 (2011), 937–943.
- [23] T.R.S. Walsh, A.B. Lehman, Counting rooted maps by genus I, *J. Combin. Theory Ser. B* 13 (1972) 192–218.
- [24] T.R.S. Walsh, Hypermaps versus bipartite maps, *J. Combin. Theory Ser. B* 18 (1975) 155–163.
- [25] D. Zagier, On the distribution of the number of cycles of elements in symmetric groups, *Nieuw Arch. Wisk.* (4) 13 (1995) 489–495.