

DECOMPOSITIONS AND TERRACINI LOCI OF CUBIC FORMS OF LOW RANK

LUCA CHIANTINI AND FULVIO GESMUNDO

ABSTRACT. We study Waring rank decompositions for cubic forms of rank $n + 2$ in $n + 1$ variables. In this setting, we prove that if a concise form has more than one non-redundant decomposition of length $n + 2$, then all such decompositions share at least $n - 3$ elements, and the remaining elements lie in a special configuration. Following this result, we give a detailed description of the $(n + 2)$ -th Terracini locus of the third Veronese embedding of n -dimensional projective space.

1. INTRODUCTION

For an integer n , let V be a complex vector space of dimension $n + 1$ and let $\mathbb{P}^n = \mathbb{P}V$ be the projective space of lines in V . Identify V with the space of linear forms on V^* and let S^dV be the space of homogeneous polynomials of degree d on V^* . The Waring rank of a form $F \in S^dV$ is

$$R(F) = \min\{r : F = L_1^d + \cdots + L_r^d \text{ for some } L_1, \dots, L_r \in V\}.$$

The d -th Veronese embedding of $\mathbb{P}V$ is the map $v_d : \mathbb{P}V \rightarrow \mathbb{P}S^dV$ defined by $v_d([L]) = [L^d]$. If A is a set of r points in $\mathbb{P}V$, we say that A has length r , and we write $\ell(A) = r$. Geometrically, the Waring rank of F can be equivalently defined as

$$R(F) = \min\{r : [F] \in \langle v_d(A) \rangle \text{ for a set } A \subseteq \mathbb{P}V \text{ with } \ell(A) = r\}.$$

Here $\langle - \rangle$ denotes the (projective) linear span. A finite set A with $\ell(A) = r$ such that $[F] \in \langle v_d(A) \rangle$ is a *decomposition* of F of length r . If $A = \{[L_1], \dots, [L_r]\}$, this is equivalent to the existence of an expression

$$(1) \quad F = \alpha_1 L_1^d + \cdots + \alpha_r L_r^d$$

for some $\alpha_1, \dots, \alpha_r \in \mathbb{C}$. We say that the decomposition is *non-redundant* (or *irredundant*) if there is no proper subset $A' \subsetneq A$ such that $[F] \in \langle v_d(A') \rangle$; this is equivalent to saying that L_1^d, \dots, L_r^d are linearly independent and no coefficient α_i in (1) is 0. We say that the decomposition is *minimal* if its length is the minimal possible, namely $r = R(F)$. If F is a form of rank r which has a unique minimal decomposition of length r , we say that F is *identifiable*.

We say that $F \in S^dV$ is *concise* if there is no proper subspace $V' \subsetneq V$ such that $F \in S^dV'$; equivalently, there is no change of coordinates in V after which F can be written in fewer than $n + 1$ variables. It is a classical fact that F is concise if and only if its first order partial derivatives are linearly independent; equivalently F is concise if and only if its partial derivatives of order $d - 1$ span V ; we refer to [Car06] for a proof of this fact. If F is concise, then every decomposition A of F satisfies $\langle A \rangle = \mathbb{P}V$; in other

2020 *Mathematics Subject Classification.* 14N07, 14N05, 15A69.

Key words and phrases. Waring rank, tensor decomposition, Hilbert function, Terracini locus.

words, no decomposition of F is contained in a hyperplane. In particular $R(F) \geq n + 1$. A partial converse of this property in low rank is given in [Lemma 2.7](#).

For a subset $A \subseteq \mathbb{P}V$ with $\ell(A) = r$, we say that A is linearly independent if $\dim\langle A \rangle = \ell(A) - 1$, which is the maximum possible value; here \dim is the dimension in projective space. The Kruskal rank of A is the largest integer k such that every subset of k elements of A is linearly independent. We say that A is in *linear general position* (LGP) if any subset of elements of A is linearly independent, namely the Kruskal rank of A is $\min\{\ell(A), n + 1\}$, which is the maximum possible value.

In this work, we study non-redundant decompositions of length $n + 2$ for concise cubic forms in $n + 1$ variables. This is the first case where Kruskal's criterion does not guarantee uniqueness of the decomposition and in fact there are examples of forms having multiple decompositions, see e.g. [\[Der13\]](#). However, our main result shows that if the form is concise, then non-uniqueness of a decomposition can only occur under very special conditions, and in particular very low Kruskal rank.

Theorem 1.1. *Let V be a vector space of dimension $n + 1$, with $n \geq 1$. Let $F \in S^3V$ be concise with $R(F) \leq n + 2$. Then one of the following occurs:*

- (I) *there is a unique non-redundant decomposition A of length $n + 2$ for F ; in this case the Kruskal rank of A is at least 4;*
- (II) *there are infinitely many non-redundant decompositions of F of length $n + 2$; any two decompositions A, B of length $n + 2$ intersect in at least $n - 3$ points; moreover $(A \setminus B) \cup (B \setminus A)$ lies in the union of two lines; in particular the Kruskal rank of any decomposition is 2;*
- (III) *there are infinitely many non-redundant decompositions of F of length $n + 2$; any two decompositions A, B of length $n + 2$ intersect in at least $n - 2$ points; moreover $(A \setminus B) \cup (B \setminus A)$ is contained in the union of two planes; in particular the Kruskal rank of any decomposition is at most 3.*

[Theorem 1.1](#) is built on an induction argument for $n \geq 4$. The base of the induction is given by the cases $n = 1, 2, 3$, where special configurations of points yield some pathological behaviour. The general cases are obtained as extensions of a special configuration on a subspace.

As a consequence of [Theorem 1.1](#), in [Section 5](#), we obtain a detailed description of the $(n + 2)$ -th Terracini locus of the Veronese variety $v_3(\mathbb{P}^n)$ in the concise setting.

The study of identifiability of homogeneous polynomials, and more generally of tensors, is a classical topic in algebraic geometry and invariant theory. Generic forms in S^dV are identifiable only for few combination of degree d and number of variables $n + 1$, namely $(d, n) = (2k - 1, 1), (3, 3), (5, 2)$: identifiability in these cases was known classically [\[Syl52, Hil\]](#); the full proof that these are the only cases where generic identifiability holds is recent [\[GM19\]](#). On the other hand, generic forms of subgeneric rank are identifiable with only few exceptions [\[BCO14, COV17\]](#). In general, however, little is known about the explicit genericity condition providing identifiability. The most general tool in this context is Kruskal's criterion [\[Kru77\]](#), which guarantees uniqueness of decomposition under the hypothesis of large enough *Kruskal rank*; a variant of Kruskal criterion is given in [\[LP21\]](#). A geometric study of the subvarieties of non-identifiable forms, in special cases, is given in [\[ACV18, AC20, CO21\]](#), which are based on properties of the Hilbert function of points [\[Chi19\]](#) and on apolarity theory [\[IK99\]](#).

Acknowledgements. This work is partially supported by the Thematic Research Programme “Tensors: geometry, complexity and quantum entanglement”, University of Warsaw, Excellence Initiative – Research University and the Simons Foundation Award No. 663281 granted to the Institute of Mathematics of the Polish Academy of Sciences for the years 2021-2023. We thank IMPAN and the AGATES organizers for providing an excellent research environment throughout the semester.

2. PRELIMINARIES

Let $X \subseteq \mathbb{P}^N$ be an irreducible projective variety. The r -th secant variety of X is

$$\sigma_r(X) = \overline{\{p \in \mathbb{P}^N : p \in \langle x_1, \dots, x_r \rangle \text{ for some } x_1, \dots, x_r \in X\}};$$

here the closure can be taken equivalently in the Zariski or the Euclidean topology. The r -th symmetric product of X is $X^{(r)} = X^{\times r} / \mathfrak{S}_r$, where the symmetric group \mathfrak{S}_r acts by permuting the factors of $X^{\times r}$. One can verify that $X^{(r)}$ is a projective variety. The r -th abstract secant variety of X is

$$A\sigma_r(X) = \overline{\{(x_1, \dots, x_r), p\} : p \in \langle x_1, \dots, x_r \rangle\}} \subseteq X^{(r)} \times \mathbb{P}^N.$$

There are two natural projections $\pi_X : A\sigma_r(X) \rightarrow X^{(r)}$, which is surjective, and $\pi_\sigma : A\sigma_r(X) \rightarrow \mathbb{P}^N$, which surjects onto the secant variety $\sigma_r(X)$. It is easy to verify that $A\sigma_r(X)$ is irreducible and $\dim A\sigma_r(X) = r \dim X + (r - 1)$; in fact, $A\sigma_r(X)$ is birational to a projective bundle over $X^{(r)}$ whose fibers are copies of \mathbb{P}^{r-1} . We say that $\sigma_r(X)$ is non-defective, or that X is not r -defective, if $\dim \sigma_r(X) = \min\{N, r \dim X + (r - 1)\}$; if $\sigma_r(X) \subsetneq \mathbb{P}^N$, this is equivalent to the condition that π_σ has generically finite fibers.

Definition 2.1. Let $X \subseteq \mathbb{P}^N$ be an irreducible variety and let $r \geq 1$ be an integer such that $N \geq r \dim X + (r - 1)$. The r -th Terracini locus of X is

$$\mathbb{T}_r(X) = \overline{\left\{ (x_1, \dots, x_r) : \begin{array}{l} x_i \in X^{\text{sm}} \text{ are linearly independent,} \\ T_{x_1}X, \dots, T_{x_r}X \text{ are not linearly independent} \end{array} \right\}} \subseteq X^{(r)},$$

where X^{sm} is the open set of the smooth points of X and $T_x X$ denotes the affine tangent space to X at x .

Terracini’s Lemma [BCC⁺18, Lemma 1] guarantees that if $\sigma_r(X)$ is non-defective and it does not fill the space, then $\mathbb{T}_r(X)$ is a proper subvariety of $X^{(r)}$.

The preimage $\pi_X^{-1}(\mathbb{T}_r(X)) \subseteq A\sigma_r(X)$ is contained in the locus where the differential of π_σ drops rank. If a point $q \in \mathbb{P}^N$ is in the image $\pi_\sigma \pi_X^{-1}(\mathbb{T}_r(X)) \subseteq \sigma_r(X)$ then either $\pi_\sigma^{-1}(q)$ is positive dimensional, or q is a cuspidal singularity of $\sigma_r(X)$.

By definition, the Terracini locus is the sets of points $\{x_1, \dots, x_r\}$ with the property that they impose independent conditions on $\mathcal{O}_X(1)$ but the double points $\{2x_1, \dots, 2x_r\}$ do not impose independent conditions on $\mathcal{O}_X(1)$. In fact, with this formulation, one can give the definition of Terracini locus for an abstract variety and any (ample) line bundle, see [BC21].

An important tool in the study of sets of point in projective space is given by the Hilbert function. Let $\mathbb{C}[V] = \text{Sym}(V^*)$ be the homogeneous coordinate ring of $\mathbb{P}V$, which is a polynomial ring in $n+1$ variables. If $Z \subseteq \mathbb{P}V$ is a subvariety with homogeneous ideal $I(Z)$, the Hilbert function of Z is

$$\begin{aligned} h_Z : \mathbb{N} &\rightarrow \mathbb{N} \\ t &\mapsto \dim(\mathbb{C}[V]_t / I(Z)_t). \end{aligned}$$

If Z is a set of points, $Z = \{[v_1], \dots, [v_r]\}$ for $v_j \in V$, let $\text{ev}_i : \mathbb{C}[V] \rightarrow \mathbb{C}$ be the evaluation map at v_i . For every t , the restriction $\text{ev}_i : \mathbb{C}[V]_t \rightarrow \mathbb{C}$ is linear, and the Hilbert function of Z is characterized as

$$h_Z(t) = \text{rank}((\text{ev}_1, \dots, \text{ev}_r) : \mathbb{C}[V] \rightarrow \mathbb{C}^r).$$

The *first difference* of the Hilbert function of Z is $Dh_Z(t) = h_Z(t) - h_Z(t-1)$. It is often identified simply with the sequence of its non-zero values, which is called the *h-vector* of Z . We record three main properties of the Hilbert function and the h-vector, which will be useful in the following. We refer to [Chi19] for details and proofs.

Proposition 2.2. *Let $Z' \subseteq Z \subseteq \mathbb{P}V$ be finite sets of points. Then*

- (i) $Dh_{Z'}(t) \leq Dh_Z(t)$ for every t ; in particular $h_{Z'}(t) \leq h_Z(t)$ for every t .
- (ii) There is an integer $\tau \geq 0$ such that $h_Z(t)$ is strictly increasing for $t \leq \tau$ and $h_Z(t) = \ell(Z)$ for $t \geq \tau$. In particular $Dh_Z(t) > 0$ if $t \leq \tau$ and $Dh_Z(t) = 0$ if $t > \tau$.
- (iii) If $Dh_Z(t) \leq t$, then $Dh_Z(t+1) \leq Dh_Z(t)$.

Understanding what functions h can occur as Hilbert functions of sets of points is the object of a long line of research. Macaulay characterized the *maximal growth* of h_Z . In [BGM94], strong consequences of Macaulay's result are given; we state here a restricted version of [BGM94, Theorem 3.6], extending a result from [Dav85] in the case of \mathbb{P}^2 .

Theorem 2.3. *Let $Z \subseteq \mathbb{P}V$ be a finite set of points. If $s := Dh_Z(t_0) = Dh_Z(t_0+1) \leq t_0$ for some t_0 then there exists a reduced curve $C \subseteq \mathbb{P}V$ of degree s such that, setting $Z' = Z \cap C$, one has*

- $Dh_{Z'}(t) = Dh_C(t)$ for $t \leq t_0 + 1$;
- $Dh_{Z'}(t) = Dh_Z(t)$ for $t \geq t_0$.

Intuitively, Theorem 2.3 says that if Dh_Z is constant and small on an interval, then a subset of points of Z lies on a low degree curve. This is useful to deduce pathologies of certain sets of points and prove that under suitable genericity assumptions only certain Hilbert functions are possible.

Sets of points defining non-redundant decompositions satisfy particular conditions that are reflected in their Hilbert function. An important one concerns the Cayley-Bacharach property. A finite set $Z \subseteq \mathbb{P}V$ satisfies the Cayley-Bacharach property in degree t , denoted $CB(t)$, if, for every $p \in Z$, every form of degree d vanishing on $Z \setminus \{p\}$ vanishes at p as well; in other words, Z satisfy $CB(t)$ if for every $p \in Z$, $I(Z \setminus \{p\})_t = I(Z)_t$.

The role of the Cayley-Bacharach property in the study of decompositions of homogeneous polynomials is stated in the following result.

Proposition 2.4 ([AC20], Proposition 2.25). *Let $F \in S^dV$ and let $A, B \subseteq \mathbb{P}V$ be non-redundant disjoint decompositions of F . The $A \cup B$ satisfies the Cayley-Bacharach property in degree d .*

In particular, the failure to satisfy the Cayley-Bacharach property allows one to rule out the existence of certain decompositions. This will be achieved using the fact that the Cayley-Bacharach property poses strong conditions on the Hilbert function of a set of points. More precisely, the following holds, see [ACV18, Thm. 4.9].

Theorem 2.5. *Let $Z \subseteq \mathbb{P}V$ be a set of points satisfying $CB(t)$. Then, for every $s \leq t+1$*

$$Dh_Z(0) + \dots + Dh_Z(s) \leq Dh_Z(t+1) + \dots + Dh_Z(t+1-s).$$

2.1. First results. In this section we prove some simple technical results toward the proof [Theorem 1.1](#). The first immediate observation is that $n + 2$ points in LGP in $\mathbb{P}V$ can be normalized via the action of $\mathrm{GL}(V)$. We record a slightly more general fact in the following result, see e.g. [[Har92](#), Ch. 10].

Lemma 2.6. *Let V be a vector space with $\dim V = n + 1$ and let e_0, \dots, e_n be a basis of V . Let*

$$\Omega = \{A \subseteq \mathbb{P}V : \ell(A) = n + 2, \langle A \rangle = \mathbb{P}V\} \subseteq (\mathbb{P}V)^{(n+2)}.$$

Then Ω is Zariski open in $(\mathbb{P}V)^{(n+2)}$; the action of $\mathrm{GL}(V)$ on Ω has exactly n orbits K_2, \dots, K_{n+1} . The elements of K_r are subsets having Kruskal rank exactly r and they are all equivalent to

$$A_r = \{[e_0], \dots, [e_n], [e_0 + \dots + e_{r-1}]\}.$$

Moreover, $K_r \subseteq \overline{K_{r+1}}$ and for every r , $\overline{K_r}$ is irreducible of dimension $n(n + 1) + r - 1$.

Proof. The first part of the statement is classical. If $A \in (\mathbb{P}V)^{(n+2)}$ is a set of points having Kruskal rank r and such that $\langle A \rangle = \mathbb{P}V$, then we may assume $A = \{[v_0], \dots, [v_n], [v_{n+1}]\}$ with v_0, \dots, v_n linearly independent and $v_{n+1} \in \langle v_0, \dots, v_{r-1} \rangle$. Since v_0, \dots, v_n are linearly independent, there is an element $g \in \mathrm{GL}(V)$ such that $gv_i = e_i$ for $i = 0, \dots, n$; as a result $gv_{n+1} = \lambda_0 e_0 + \dots + \lambda_{r-1} e_{r-1}$ for some $\lambda_j \in \mathbb{C} \setminus \{0\}$. Define an element $h \in \mathrm{GL}(V)$ by $he_i = \lambda_i^{-1} e_i$ for $i \leq r - 1$ and $he_i = e_i$ for $i \geq r$. We conclude that $hgA = A_r$.

The condition that $K_r \subseteq \overline{K_{r+1}}$ is clear; moreover $\overline{K_r}$ is irreducible, because it is the closure of an orbit under the action of $\mathrm{GL}(V)$. Finally, to compute the dimension of $\overline{K_r}$, consider the diagram

$$\begin{array}{ccc} & (\mathbb{P}V)^{n+2} & \\ \pi_{\mathfrak{S}} \swarrow & & \searrow \pi_{n+2} \\ (\mathbb{P}V)^{(n+2)} & & (\mathbb{P}V)^{n+1} \end{array}$$

where $\pi_{\mathfrak{S}}$ is the projection onto the quotient $(\mathbb{P}V)^{(n+2)} = (\mathbb{P}V)^{n+2} / \mathfrak{S}_{n+2}$ and π_{n+2} is the projection onto the first $n + 1$ factors. The preimage $\pi_{\mathfrak{S}}(K_r)$ is the union several irreducible components, all isomorphic and all surjecting onto K_r ; one of these components is

$$J_r = \left\{ ([v_0], \dots, [v_{n+1}]) \in (\mathbb{P}V)^{n+2} : \begin{array}{l} \mathbb{P}V = \langle [v_0], \dots, [v_n] \rangle \\ v_{n+1} \in \langle v_0, \dots, v_{r-1} \rangle \end{array} \right\}$$

Since $\pi_{\mathfrak{S}}$ is generically finite, $\dim K_r = \dim J_r$. The restriction $\pi_{n+2} : J_r \rightarrow (\mathbb{P}V)^{n+1}$ is dominant; the fiber over a generic element $([v_0], \dots, [v_n])$ is the linear span $\mathbb{P}\langle v_0, \dots, v_{r-1} \rangle$ which has dimension $r - 1$. We conclude

$$\dim K_r = \dim J_r = \dim((\mathbb{P}V)^{n+1}) + \dim \mathbb{P}^{r-1}(r - 1) = n(n + 1) + r - 1.$$

This concludes the proof. \square

We observe that forms admitting non-redundant decompositions of length $n + 2$ are necessarily concise:

Lemma 2.7. *Let $F \in S^d V$ be a homogeneous form with $\dim V = n + 1$. Let $A \subseteq \mathbb{P}V$ be a set of $n + 2$ points with $\langle A \rangle = \mathbb{P}V$. If A is a non-redundant decomposition of F then F is concise.*

Proof. By Lemma 2.6, we may assume $A \in K_r$ for some r and we can easily reduce to the case $r = n$. Write $L = x_0 + \cdots + x_n$ and assume

$$F = \alpha_0 x_0^d + \cdots + \alpha_n x_n^d + \alpha_{n+1} L^d$$

where $\alpha_j \neq 0$ for every j .

Let H be the space of $(d-1)$ -th order partial derivatives of F . Up to scaling, we have

$$\frac{\partial^{d-1}}{\partial x_0^{d-2} \partial x_1} F = \alpha_{n+1} L;$$

hence $L \in H$. Moreover, $\frac{\partial^{d-1}}{\partial x_j^d} F = \alpha_j x_j + \alpha_{n+1} L$, showing $x_j \in H$ for every j . This shows $H = V$, therefore F is concise. \square

Similarly, forms admitting non-redundant decompositions of length exactly $n+1$ are concise, and their decomposition is unique. The uniqueness of the decomposition can be obtained via an easy calculation with the Hilbert function, but it is also a consequence of the classical Kruskal criterion for tensors, [Kru77].

Lemma 2.8. *Let $F \in S^d V$ be a homogeneous form, with $\dim V = n+1$ with $d \geq 3$. Let $A \subseteq \mathbb{P}V$ be a non-redundant decomposition of F with A linearly independent in $\mathbb{P}V$ and $\ell(A) = n+1$. Then F is concise, $R(F) = n+1$ and A is the unique minimal decomposition of F .*

Proof. Since A is linearly independent with $\ell(A) = n+1$, we may normalize it via the action of $\mathrm{GL}(V)$ and assume $A = \{[x_0], \dots, [x_n]\}$. Therefore $\langle v_d(A) \rangle = \langle [x_0^d], \dots, [x_n^d] \rangle$ and $F = x_0^d + \cdots + x_n^d$. This shows that F is concise with $R(F) = n+1$.

The uniqueness of the decomposition follows by Kruskal's criterion, see [Kru77]. The proof follows from the immediate fact that if $A \subseteq \mathbb{P}V$ is a set of linearly elements with $\ell(A) = n+1$, then A is in LGP and in particular it has maximal Kruskal ranks. \square

A stronger result holds for forms admitting two non-redundant decompositions of length $n+1$ and $n+2$, respectively.

Lemma 2.9. *Let $F \in S^d V$ be a concise form with $d \geq 3$. Let $A \subseteq \mathbb{P}V$, $B \subseteq \mathbb{P}V$ be non-redundant decompositions of F of length $n+1$ and $n+2$ respectively. Then $d = 3$, $\ell(A \cap B) \geq n-1$ and $(A \setminus B) \cup (B \setminus A)$ is collinear.*

Proof. We proceed by induction on n . If $n = 1$ the statement is clearly true. Indeed, every concise $F \in S^d V$ has at most one decomposition of length 2 and a two-parameter family of decompositions of length 3. All decompositions are collinear because $\mathbb{P}V$ is a line.

Assume $n \geq 2$. First assume $A \cap B = \emptyset$. Let $Z = A \cup B$. Since F is concise, we have $Dh_A = (1, n)$, which implies $Dh_Z(0) + Dh_Z(1) = n+1$; by the Cayley-Bacharach property of Z , we have $Dh_Z(d+1) + Dh_Z(d) \geq n+1$. This implies $Dh_Z(2) \leq 1$, which arithmetically is a contradiction by Proposition 2.2.

Therefore $A \cap B \neq \emptyset$. Without loss of generality, assume $A = \{[x_0], \dots, [x_n]\}$ and $F = x_0^d + \cdots + x_n^d$. Since $A \cap B \neq \emptyset$, assume $[x_n] \in B$ and write $B = \{[L_0], \dots, [L_n], [x_n]\}$, so that $F = L_0^d + \cdots + L_n^d + \beta x_n^d$ for some nonzero $\beta \in \mathbb{C}$. Consider

$$(2) \quad F' = F - \beta x_n^d = x_0^d + \cdots + x_{n-1}^d + (1 - \beta) x_n^d = L_0^d + \cdots + L_n^d.$$

Let $B' = \{[L_0], \dots, [L_n]\}$; clearly $[x_n] \notin B'$. If $\beta \neq 1$, then F' is concise and (2) gives two non-redundant decompositions F' of length $n+1$. By Lemma 2.8, the two

decompositions are the same, in contradiction with the fact that $[x_n] \notin B'$. Therefore $\beta = 1$ and F' is non-concise. Moreover, if B' was linearly independent in $\mathbb{P}V$, [Lemma 2.8](#) would imply that F' is concise; hence B' is not linearly independent.

Let $V' = \langle x_0, \dots, x_{n-1} \rangle$. Notice $V' = \langle B' \rangle$ and F' is concise in $S^d V'$. Let $A' = \{[x_0], \dots, [x_{n-1}]\}$. Now F' is concise and it has two decompositions A', B' of length n and $n+1$ respectively. By the inductive hypothesis, we deduce $d = 3$, $\ell(A' \cap B') \geq n-2$ and B' has three collinear points. Since $A = A' \cup \{[x_n]\}$ and similarly for B , we deduce that the statement holds for A and B as well. This concludes the proof. \square

In the study of identifiability of forms, it is often useful to consider the case where two potential decompositions of a form are disjoint. We illustrate a general method to reduce the case of non-disjoint decompositions to the one of disjoint decompositions:

Remark 2.10. Let $F \in S^d V$ and suppose $A, B \subseteq \mathbb{P}V$ give two non-redundant decompositions of F . Write $A = \{[L_1], \dots, [L_r]\}$ and $B = \{[M_1], \dots, [M_s]\}$. Then

$$F = \alpha_1 L_1^d + \dots + \alpha_r L_r^d = \beta_1 M_1^d + \dots + \beta_s M_s^d.$$

After possibly reordering the elements of A and B , assume $L_i = M_i$ for $i = 1, \dots, p$, where $p = \ell(A \cap B)$. In this case, define

$$F' = F - (\alpha_1 L_1^d + \dots + \alpha_p L_p^d) = (\beta_1 - \alpha_1) M_1^d + \dots + (\beta_p - \alpha_p) M_p^d + \beta_{p+1} M_{p+1}^d + \dots + \beta_s M_s^d;$$

note

$$F' = \alpha_{p+1} L_{p+1}^d + \dots + \alpha_r L_r^d.$$

Let $A' = A \setminus B$; let $B' = \{[M_i] : \alpha_i \neq \beta_i, i = 1, \dots, p\} \cup \{[M_{p+1}], \dots, [M_s]\}$; then A', B' are two disjoint non-redundant decompositions of F' . Moreover, if A is a minimal decomposition for F , then A' is a minimal decomposition for F' .

We conclude this section with a classical result for which we do not know an explicit reference, in the form that we need in the following. The result dates back to [\[Syl52\]](#); analogous statements, in a more general setting, are discussed in [\[BBCG19, Section 7.1\]](#).

Proposition 2.11. *Let F be a concise form in $S^d V$, with $\dim V = n + 1$. Let A, B be non-redundant decompositions of F of length r and s respectively. Then $r + s \geq d + 2n$.*

Proof. Let p be the length of $A \cap B$. If $p = 0$, set $F' = F$. If $p > 0$, consider the form F' with disjoint decompositions A', B' of length r', s' respectively, as constructed in [Remark 2.10](#). Notice that $r - r' = p \geq s - s'$; specifically $s - s'$ is the number of elements of $A \cap B$ appearing with the same coefficient in the two decompositions of F . Observe $\dim \langle B' \rangle \geq n - (s - s')$: indeed F is concise, which implies that B spans $\mathbb{P}V$; since $\dim \langle B \setminus B' \rangle \leq s - s' - 1$, we deduce $\dim \langle B' \rangle \geq n - (s - s')$.

Set $Z = A' \cup B'$ and let $n' = \dim \langle Z' \rangle$. We have $n' \geq \dim \langle B' \rangle \geq n - (s - s')$, so $n - n' \leq s - s' \leq r - r'$. Since Z is union of two disjoint decompositions of a form of degree d , we have $Dh_{Z'}(d+1) \geq 1$, which implies $Dh_{Z'}(t) \geq 1$ for $t \leq d+1$. Moreover, $Dh_{Z'}(1) = \dim \langle Z' \rangle = n'$. Finally, [Proposition 2.4](#) implies that the Cayley-Bacharach property holds for Z' in degree d , and [Theorem 2.5](#) implies $Dh_{Z'}(d+1) + Dh_{Z'}(d) \geq Dh_{Z'}(0) + Dh_{Z'}(1) \geq 1 + n'$.

We deduce $Dh_{Z'} = (1, n', \delta_2, \dots, \delta_{d+1}, \dots)$ with $\delta_t \geq 1$ for $t \leq d+1$ and $\delta_{d+1} + \delta_d \geq 1 + n'$. We conclude $r' + s' = \ell(Z') \geq \sum_{t=0}^{d+1} \delta_t \geq 2 + 2n' + (d-2) = d + 2n'$. Therefore

$$r + s = (r - r') + (s - s') + (r' + s') \geq (n - n') + (n - n') + d + 2n' = d + 2n$$

as desired. \square

The bound of [Proposition 2.11](#) is sharp. An example where the bound is attained can be constructed, for every n and d , as follows. When $d = 2$, general forms of maximal rank have many decompositions of length $n + 1$, which attains the bound.

For higher values of d , write $V = V' \oplus V''$ where with $\dim V' = 2$, $\dim V'' = n - 1$. Let $F' \in S^d V'$ be a binary form having two minimal disjoint decompositions A', B' of length r', s' with $r' + s' = d + 2$; it is a classical result that such forms exist [[Sy152](#), [CS11](#)]. Let $F'' = L_1^d + \cdots + L_{n-1}^d$ be a sum of $n - 1$ d -powers of generic linear forms. Then $F = F' + F''$ is concise and it has two distinct non-redundant decompositions $A = A' \cup \{L_1, \dots, L_{n-1}\}$ and $B = B' \cup \{L_1, \dots, L_{n-1}\}$ with $\ell(A) = r' + (n - 1)$, $\ell(B) = s' + (n - 1)$ and we have $\ell(A) + \ell(B) = r' + s' + 2(n - 1) = d + 2n$.

3. FIRST CASES

In this section, we characterize decompositions of length $n + 2$ in \mathbb{P}^n for $n = 1, 2$.

Case $n = 1$.

Let V be a vector space with $\dim V = 2$. The generic Waring rank in $\mathbb{P}S^3V$ is 2. In this case [Theorem 1.1](#) is easily verified. Let $F \in S^3V$ be a concise form. If $R(F) = 2$, then F has a unique decomposition of length 2 by Kruskal criterion, and in particular [Lemma 2.8](#); moreover F has a 2-dimensional family of decompositions of length 3. If $R(F) = 3$, then $[F]$ is an element of $\tau(v_3(\mathbb{P}^1)) \setminus v_3(\mathbb{P}^1)$, where $\tau(v_3(\mathbb{P}^1)) = \{[L_1^2 L_2] \in \mathbb{P}S^3V : [L_1], [L_2] \in \mathbb{P}^1\}$; in this case F has a two dimensional family of decompositions of length 3.

In both cases, the two dimensional family of decompositions can be described as follows. Let Λ be a generic plane passing through $[F]$. Since $v_3(\mathbb{P}V)$ is a curve of degree 3, its intersection with Λ consists of three points counted with multiplicity. The genericity assumption guarantees that the three intersection points are distinct, $v_3(\mathbb{P}V) \cap \Lambda = \{[L_1^3], [L_2^3], [L_3^3]\}$ for some $L_1, L_2, L_3 \in V$. Moreover, any three points on $v_3(\mathbb{P}V)$ are linearly independent, therefore $\Lambda = \langle [L_1^3], [L_2^3], [L_3^3] \rangle$. This shows that $A = \{[L_1], [L_2], [L_3]\}$ is a decomposition of length 3. The genericity assumption on Λ guarantees it is non-redundant, otherwise one would obtain several decompositions of length 2, in contradiction with [Lemma 2.8](#).

In other words, every generic enough plane Λ through $[F]$ determines uniquely a non-redundant decomposition of length 3. Planes through $[F]$ form a hyperplane in $\mathbb{P}S^3V^*$, hence this is a 2-dimensional family.

Since $\mathbb{P}V = \mathbb{P}^1$, all decompositions are defined by collinear points. In particular, both case (II) and case (III) of [Theorem 1.1](#) trivially hold.

Case $n = 2$.

Let V be a vector space with $\dim V = 3$. The generic Waring rank in $\mathbb{P}S^3V$ is 4. Let $F \in S^3V$ be a concise form. It is known that $3 \leq R(F) \leq 5$ [[LT10](#)] and we are interested in the dense subset of forms satisfying $R(F) \leq 4$. Notice $\dim A\sigma_4(v_3(\mathbb{P}^2)) = 12$. Therefore the projection map $A\sigma_4(v_3(\mathbb{P}^2)) \rightarrow \mathbb{P}S^3V$ has generic fibers of dimension 2. This guarantees that if $R(F) \leq 4$, then F has at least a 2-dimensional family of decompositions of length 4.

In particular, case (III) of [Theorem 1.1](#) trivially holds. We also provide a geometric characterization of the decompositions.

Theorem 3.1. *Let $F \in S^3V$ be a concise form with $\dim V = 3$. Let A, B be non-redundant decompositions of F , with $\ell(A) = \ell(B) = 4$. Then one of the following holds:*

- $A \cap B = \emptyset$ and there is a conic passing through $A \cup B$; if A has three collinear points, so does B .
- $\ell(A \cap B) \geq 1$ and all non-redundant decompositions of F of length 4 have three collinear points, on a line Λ .

Proof. The statement follows from [Proposition 3.2](#) and [Proposition 3.3](#) below. [Proposition 3.2](#) analyzes the case where A and B are disjoint and [Proposition 3.3](#) the one where they are not. \square

Proposition 3.2. *Let $F \in S^3V$ be a concise form with $\dim V = 3$. Let A, B be non-redundant decompositions of F , with $\ell(A) = \ell(B) = 4$. If A and B are disjoint then there is a unique conic C vanishing on $A \cup B$. Moreover, if Λ is a line with $\ell(A \cap \Lambda) \geq 3$, then $C = \Lambda \cup \Lambda'$ for a line Λ' and $\ell(B \cap \Lambda') = 3$.*

Proof. Since F is concise, there are no linear forms vanishing on A . Therefore the h-vector of A is $Dh_A = (1, 2, 1)$ and there is a pencil of conics vanishing on A .

Let $Z = A \cup B$; since $A \cap B = \emptyset$, $\ell(Z) = 8$. Since $[F] \in \langle v_3(A) \rangle \cap \langle v_3(B) \rangle$, we have that $v_3(Z)$ is not linearly independent, namely $\dim \langle v_3(Z) \rangle < \ell(Z) - 1 = 7$. In particular, $Dh_Z(4) > 0$. Since $Dh_A = (1, 2, 1)$, we have $Dh_Z(0) + Dh_Z(1) \geq 3$. Since A, B are non-redundant, they satisfy the Cayley-Bacharach property in degree 3 by [Proposition 2.4](#): therefore [Theorem 2.5](#) provides $Dh_Z(3) + Dh_Z(4) \geq 3$, so $Dh_Z(2) \leq 2$. If $Dh_Z(2) \leq 1$, by [Proposition 2.2](#), we have $Dh_Z(i) \leq 1$ for $i \geq 2$, which provides a contradiction; therefore $Dh_Z(2) = 2$. We obtain $Dh_Z = (1, 2, 2, 2, 1)$. In particular, $h_Z(2) = 1 + 2 + 2 = 5$, which implies that there is a unique conic vanishing on Z .

Now suppose A contains three collinear points, lying on a line $\Lambda \subseteq \mathbb{P}V$. Since $A \subseteq C$, we deduce that the conic C passing through Z is reducible with $C = \Lambda \cup \Lambda'$ for some other line $\Lambda' \subseteq \mathbb{P}V$. We are going to show that $B \cap \Lambda'$ consists of three points. Suppose by contradiction $B \cap \Lambda'$ contains fewer than three points; since $B \subseteq C$, we deduce that $B \cap \Lambda$ contains at least two points. Notice $h_Z(3) = 1 + 2 + 2 + 2 = 7 = \ell(Z) - 1$: this guarantees $\dim(\langle v_3(A) \rangle \cap \langle v_3(B) \rangle) = 0$ therefore the two linear spans only intersect at $[F]$. However,

$$\ell(\Lambda \cap Z) = \ell(\Lambda \cap A) + \ell(\Lambda \cap B) \geq 3 + 2 = 5;$$

therefore $v_3(\Lambda \cap Z)$ is linearly dependent. This implies $\langle v_3(A \cap \Lambda) \rangle \cap \langle v_3(B \cap \Lambda) \rangle \neq \emptyset$, in contradiction with the fact that the two spans only intersect at $[F]$, which is not an element of $\langle v_3(\Lambda) \rangle$ because F is concise. This shows that $B \cap \Lambda'$ contains three points. \square

The case where A and B are not disjoint is characterized in the following result.

Proposition 3.3. *Let $F \in S^3V$ be a concise form with $\dim V = 3$. Let A, B be non-redundant decompositions of F , with $\ell(A) = \ell(B) = 4$. If $\ell(A \cap B) \geq 1$, then there is a line $\Lambda \subseteq \mathbb{P}V$ such that $A \setminus \Lambda = B \setminus \Lambda$ consists of exactly one point. In particular, both A and B have three collinear points on Λ .*

Proof. Let $p = \ell(A \cap B)$. Let F', A', B' be constructed from F, A, B as in [Remark 2.10](#). Then A' is a non-redundant decomposition of F' of length $4 - p \leq 3$ and B' is a non-redundant decomposition of F' of length at most 4.

Suppose A' is not contained in a line. In particular, $\ell(A') = 3$ and F' is concise by [Lemma 2.8](#). Moreover A' is the unique decomposition of length 3. This contradicts the existence of B' .

Therefore there exists $V' \subseteq V$, with $\dim V' = 2$ and $A' \subseteq \Lambda = \mathbb{P}V'$. In particular $F' \in S^3V'$ is not concise in S^3V . If F' is not concise in S^3V' , then $F' = L^3$ for some $L \in V'$; therefore $\{[L]\}$ is a decomposition of F' and by [Proposition 2.11](#), any other non-redundant decomposition of F' must have length at least 4. On the other hand $\ell(B') \leq 3$, because if $\ell(B') = 4$, then $B = B'$, and F' would be concise by [Lemma 2.7](#). Therefore, the condition $\ell(B') \leq 3$ provides a contradiction with [Proposition 2.11](#). This guarantees F' is concise in S^3V' .

To conclude, we analyze two possibilities. If $\ell(A') = 2$, then A' is the unique decomposition of F' of length 2 and $\ell(B') = 3$. In this case $\ell(A \cap B) = 2$ and it consists of one point on Λ and one not in Λ , because B' contains three of the four points of B and $B' \subseteq \Lambda$. In particular $A \setminus \Lambda = B \setminus \Lambda = \{[L]\}$ with $F = F' + L^3$. If $\ell(A') = 3 = \ell(B') = 3$, then $A \cap B = A \setminus \Lambda = B \setminus \Lambda = \{[L]\}$ with $F = F' + L^3$. \square

4. THE GENERAL CASE

In this section, we give the proof of [Theorem 1.1](#). This will follow from an induction argument, where the cases $n = 1, 2, 3$ give the base of the induction. We provide first a result which analyzes the case $n = 3$.

If $\dim V = 4$, the generic Waring rank in $\mathbb{P}S^3V$ is 5. Sylvester's Pentahedral Theorem [[Syl52](#), [Cle61](#)] guarantees that a generic form in S^3V has a unique decomposition. We first study the situation of two disjoint decompositions.

Proposition 4.1. *Let $\dim V = 4$. Let $F \in S^3V$ be a concise form. Let $A, B \subseteq \mathbb{P}V$ be two disjoint non-redundant decompositions of F with $\ell(A), \ell(B) = 5$. Then $A \cup B$ is contained in the union of two lines.*

Proof. Since $A \cap B = \emptyset$, the union $Z = A \cup B$ is a set of length 10. Since F is concise, the decomposition A is not contained in a hyperplane; therefore $Dh_A(0) = 1$, $Dh_A(1) = 3$. Since A, B are non-redundant, by [Proposition 2.4](#) Z satisfies the Cayley-Bacharach property in degree 3. Therefore $Dh_Z(4) + Dh_Z(3) \geq Dh_Z(0) + Dh_Z(1) \geq 4$. We deduce $Dh_Z(2) \leq 2$, which implies $Dh_Z(3), Dh_Z(4) \leq 2$ by [Proposition 2.2\(iii\)](#). This guarantees $Dh_Z = (1, 3, 2, 2, 2)$.

We apply [Theorem 2.3](#): we have $Dh_Z(3) = Dh_Z(4) = 2 \leq 2$. Therefore there is a curve C of degree 2 such that, setting $Z' = Z \cap C$, one has $Dh_{Z'}(t) = Dh_C(t)$ for $t \leq 4$. Since $\deg(C) = 2$, C is either a plane conic or a union of two lines. If C is a plane conic, we have $Dh_{Z'}(t) = (1, 2, 2, 2, 2, \delta_5)$, with $\delta_5 = 0, 1$; in this case $Z' = Z \cap C$ contains at least 9 points, therefore either $A \subseteq C$ or $B \subseteq C$, in contradiction with the conciseness of F , which implies $\langle A \rangle = \langle B \rangle = \mathbb{P}V$. Therefore C is union of two lines, and we have $Dh_{Z'} = (1, 3, 2, 2, 2)$; this guarantees $Z = Z' \subseteq C$, as desired. \square

On the other hand, when $n \geq 4$, two distinct decompositions always intersect, as shown in the next result.

Proposition 4.2. *Let $n \geq 4$ and let V be a vector space with $\dim V = n + 1$. Let $F \in S^3V$ be a concise form. Let A be a non-redundant decomposition of F of length $n + 2$ and let B be a non-redundant decomposition of F of length $s \leq n + 2$. Then $A \cap B \neq \emptyset$.*

Proof. Assume $B \cap A = \emptyset$, so that $Z = A \cup B$ is a set of length at most $2n + 4$. Since F is concise, then A is not contained in a hyperplane; therefore $Dh_A(0) = 1$, $Dh_A(1) = n$. Since A, B are non-redundant, Z satisfies the Cayley-Bacharach property; therefore $Dh_Z(4) + Dh_Z(3) \geq Dh_Z(0) + Dh_Z(1) \geq n + 1$. We deduce $Dh_Z(2) \leq 2$, which implies $Dh_Z(3), Dh_Z(4) \leq 2$; this provides a contradiction, for $n \geq 4$. \square

The proof of [Theorem 1.1](#) will follow because in the setting of [Proposition 4.2](#) one can guarantee that the intersection between two decompositions must be surprisingly large, similarly to [Lemma 2.9](#).

Proposition 4.3. *Let V be a vector space with $\dim V = n + 1$. Let $F \in S^3V$ be a concise form. Let A be a non-redundant decomposition of F of length $n + 2$ and let B be a non-redundant decomposition of F of length $s \leq n + 2$. Then one of the following holds:*

- $\ell(A \cap B) \geq n - 2$, and the points of $(A \setminus B) \cup (B \setminus A)$ are contained in the union of two planes;
- $\ell(A \cap B) \geq n - 3$ and the points of $(A \setminus B) \cup (B \setminus A)$ are contained in the union of two lines.

Proof. If $s = n + 1$, then the statement follows immediately from [Lemma 2.9](#). Therefore assume $s = n + 2$.

Let $A = \{[L_1], \dots, [L_{n+2}]\}$, $B = \{[M_1], \dots, [M_{n+2}]\}$. We proceed by induction on n . The case $n = 1$ and $n = 2$ are clearly verified. So assume $n \geq 3$.

If $n = 3$ and $A \cap B = \emptyset$, then by [Proposition 4.1](#) condition (ii) is verified. If $n > 3$, then $A \cap B \neq \emptyset$ by [Proposition 4.2](#). Therefore assume $A \cap B \neq \emptyset$ with $[L_{n+2}] = [M_{n+2}] \in A \cap B$.

Write, as usual,

$$\begin{aligned} F &= \alpha_1 L_1^3 + \dots + \alpha_{n+1} L_{n+1}^3 + \alpha_{n+2} L_{n+2}^3 = \\ &= \beta_1 M_1^3 + \dots + \beta_{n+1} M_{n+1}^3 + \beta_{n+2} L_{n+2}^3. \end{aligned}$$

Let $F' = F - \beta_{n+2} L_{n+2}^3$. Then F' has a decomposition $B' = B \setminus \{[L_{n+2}]\}$ of length $n + 1$.

If B' is linearly independent, by [Lemma 2.8](#), we deduce that F' is concise and B' is its unique decomposition of length $n + 1$. If $\alpha_{n+2} \neq \beta_{n+2}$, then A defines a non-redundant decomposition of F' of length $n + 2$; in this case [Lemma 2.9](#) applies, and we deduce $\ell(A \cap B') \geq n - 1$ and $A \setminus B'$ is collinear. Since $A \setminus B' = A \setminus B$, condition (ii) holds. If $\alpha_{n+2} = \beta_{n+2}$, then $A' = A \setminus \{[L_{n+2}]\}$ is a decomposition of F' of length $n + 1$; by Kruskal's criterion [Lemma 2.8](#), we conclude $A' = B'$ which implies $A = B$ and the statement follows.

Now assume B' is not linearly independent, which implies that F' is not concise. Let $V' \subseteq V$ be the subspace in which F' is concise. Since F is concise, we have $\dim V' = \dim V - 1 = n$, and $\langle B' \rangle = V'$. Now, if $\alpha_{n+2} \neq \beta_{n+2}$, A defines a non-redundant decomposition of F' of length $n + 2$. Moreover, $\langle A \rangle = V$. By [Lemma 2.7](#), this would imply that F' is concise, which is a contradiction. Therefore $\alpha_{n+2} = \beta_{n+2}$.

We reduced to the following setting. The form F' is concise in S^3V' , and A', B' are two non-redundant decompositions of F' of length $n + 1$. Therefore the induction hypothesis applies. We deduce that condition (i) or condition (ii) hold for A' and B' . Since $A = A' \cup \{[L_{n+2}]\}$ and $B = B' \cup \{[L_{n+2}]\}$, we deduce that condition (i) or condition (ii) holds for A and B as well. \square

The proof of [Theorem 1.1](#) is obtained directly from the results of this section.

Proof of [Theorem 1.1](#). If F has a unique decomposition A of length $n + 2$, then the Kruskal rank of A must be at least 4. Indeed, let $A' \subseteq A$ be a minimal linearly dependent set, so that the Kruskal rank of A equals $\ell(A') - 1$. Let $A = A' \cup A''$: A'' is linearly independent and $\langle v_3(A') \rangle \cap \langle v_3(A'') \rangle = \emptyset$. Correspondingly $F = F' + F''$, with A' non-redundant decomposition of F' and A'' non-redundant decomposition of F'' . If $\ell(A') \leq 4$, then F' has many decompositions of length $\ell(A')$, by the discussion on the case $n = 2$. This implies that A is not the unique decomposition of F , in contradiction with the assumption.

If F does not have a unique decomposition, then either case (II) or case (III) occur, by [Proposition 4.3](#). \square

We record two easy consequence of [Theorem 1.1](#).

Corollary 4.4. *Let $F \in S^3V$ be a concise form admitting at least two distinct non-redundant decompositions of length $n + 2$. Then the Kruskal rank of any non-redundant decomposition of length $n + 2$ is at most 3. Moreover, the family of decompositions of F of length $n + 2$ has dimension at least 2.*

Proof. Since F has at least two non-redundant decompositions of length $n + 2$, either case (II) or case (III) of [Theorem 1.1](#) is verified. This guarantees every non-redundant decomposition of F of length $n + 2$ has Kruskal rank at most 3.

Write $A = \{[L_0], \dots, [L_{n+1}]\}$ for a non-redundant decomposition of length $n + 2$ and assume $[L_0], \dots, [L_3]$ are linearly dependent. Write $F = L_0^3 + \dots + L_{n+1}^3$ and define $F' = L_0^3 + \dots + L_3^3$. Then $F' \in S^3V'$ with $\dim V' = 3$ and it has a non-redundant decomposition of length 4. From the discussion on the case $n = 2$, we deduce that F' has a 2-dimensional family of decomposition of length 4. For every such decomposition B' , define $B = B' \cup \{[L_4], \dots, [L_{n+1}]\}$. Then B is a non-redundant decomposition of F , showing that the family of decompositions of F has dimension at least 2. \square

As a byproduct of these results, we completely characterize the genericity condition of Sylvester's Pentahedral Theorem.

Theorem 4.5. *Let $\dim V = 4$ and let $F \in S^3V$ be a concise form. Then the following are equivalent:*

- (i) F has at least two decompositions as sum of five powers of linear forms;
- (ii) F has a 2-dimensional family of decompositions as sum of five powers of linear forms;
- (iii) $F = F' + L^3$ for $F' \in S^3V'$ where $V' \subseteq V$ is a subspace of dimension 3 and L is a linear form with $L \notin V'$.

Proof. (i) \Rightarrow (iii). Since the decomposition is not unique, either condition (I) or condition (II) of [Theorem 1.1](#) hold. If condition (II) holds, then $F = F' + L^3$ as desired. If condition (I) holds, then $F = F'' + L_1^3 + L_2^3$ where $F'' \in S^3V''$ for a subspace $V'' \subseteq V$ of dimension 2; setting $F' = F'' + L_1^3$ we obtain the desired expression.

(iii) \Rightarrow (ii) Let $F = F' + L^3$. Every non-redundant decomposition of F' provides a non-redundant decomposition of F . Since $\dim V' = 3$, F' has at least a 2-dimensional family of decompositions.

(ii) \Rightarrow (i) This is trivially satisfied. \square

Corollary 4.6. *Let $\dim V = 4$. A form $F \in S^3V$ has a unique decomposition as sum of five powers of linear forms if and only if*

$$[F] \notin \mathbf{J}(\text{Sub}_3, v_3(\mathbb{P}V))$$

where Sub_3 denotes the variety of non-concise forms and $\mathbf{J}(-, -)$ denotes the geometric join of two varieties.

The variety Sub_3 is called *subspace variety* in [Lan12, Ch. 3] and it is defined by the vanishing of 4×4 minors of the first partial derivative map, see, e.g., [Car06]. We refer to [Har92, Ch. 8] for the definition and the basic properties of the join of two varieties.

5. CHARACTERIZATION OF THE TERRACINI LOCUS

In this section, we characterize a subvariety of the $(n+2)$ -th Terracini locus of the third Veronese embedding of \mathbb{P}^n . Define the *r-th concise Terracini locus* to be the subvariety of the Terracini locus arising from points in \mathbb{P}^n whose span is the entire space. More precisely, for a vector space V of dimension $n+1$, define

$$\mathbb{T}_r^{\text{con}}(v_3(\mathbb{P}V)) = \overline{\mathbb{T}_r(v_3(\mathbb{P}V)) \cap \{v_3(A) : A \in \mathbb{P}V^{(r)} \text{ and } \langle A \rangle = \mathbb{P}V\}}.$$

It is clear that if $\mathbb{T}_r^{\text{con}}(v_3(\mathbb{P}V))$ is non-empty, then it is (union of) irreducible components of $\mathbb{T}_r(v_3(\mathbb{P}V))$. Theorem 1.1 allows us to describe $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ as the closure of the set $v_3(K_3) \subseteq \mathbb{P}V^{(n+2)}$, where K_3 is the orbit, for the action of $\text{GL}(V)$, described in Lemma 2.6. In particular, this guarantees that it is irreducible.

Theorem 5.1. *The concise Terracini locus $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ is the closure of*

$$(3) \quad v_3(K_3) = \{v_3(A) \in v_3(\mathbb{P}V)^{(n+2)} : A \text{ concise with four coplanar points}\}.$$

In particular, $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ is irreducible of dimension $n(n+1) + 2$.

The proof of Theorem 5.1 is based on Lemma 2.6 and two technical results presented below, Lemma 5.2 and Lemma 5.3.

Lemma 5.2. *Let $d \geq 3$. Let $\mathbb{P}V_1, \mathbb{P}V_2 \subseteq \mathbb{P}V$ be disjoint linear spaces. For $i = 1, 2$, let*

$$H_i = \langle T_{[M^d]}v_d(\mathbb{P}V) : M \in V_i \rangle.$$

Then $\mathbb{P}H_1 \cap \mathbb{P}H_2 = \emptyset$.

Proof. We have $T_{[M^d]}v_d(\mathbb{P}V) = V \cdot M^{d-1}$. Therefore

$$H_i = \langle T_{[M^d]}v_d(\mathbb{P}V) : M \in V_i \rangle = V \cdot S^{d-1}V_i;$$

since $V_1 \cap V_2 = 0$ and $d \geq 3$, we deduce $\mathbb{P}H_1 \cap \mathbb{P}H_2 = 0$. \square

The next result is essentially contained in [CC10, Lemma 5.10].

Lemma 5.3. *Let $V' \subseteq V$ be a proper subspace of V and let $L_1, \dots, L_k \in V'$. If the tangent spaces to $v_d(\mathbb{P}V)$ at $[L_1^d], \dots, [L_k^d]$ are linearly dependent, so are the tangent spaces to $v_d(\mathbb{P}V')$ to $[L_1^d], \dots, [L_k^d]$.*

Proof. If $[L_1^{d-1}], \dots, [L_k^{d-1}]$ are linearly dependent then the statement is clearly satisfied. Therefore suppose they are independent. If $T_{[L_1^d]}v_d(\mathbb{P}V), \dots, T_{[L_k^d]}v_d(\mathbb{P}V)$ are linearly dependent, then there exist $M_1, \dots, M_k \in V$, not all zero, such that

$$M_1L_1^{d-1} + \dots + M_kL_k^{d-1} = 0.$$

In fact, we show $M_j \in V'$ for every j . To see this, let V'' be a complement to V' , so that $V = V' \oplus V''$; write $M_j = M'_j + M''_j$ for some $M'_j \in V'$, $M''_j \in V''$. We have

$$\begin{aligned} M_1 L_1^{d-1} + \cdots + M_k L_k^{d-1} &= \\ (M'_1 L_1^{d-1} + \cdots + M'_k L_k^{d-1}) + (M''_1 L_1^{d-1} + \cdots + M''_k L_k^{d-1}) &= 0. \end{aligned}$$

Since $L_j \in V'$, the two summands in the expression above are elements of $S^{d-1}V'$ and $V'' \cdot S^{d-1}V'$ respectively; since these two subspaces are linearly independent, each summand must vanish. Since $V' \cap V'' = 0$, the elements of $V'' \cdot S^{d-1}V'$ can be regarded as elements of $V'' \otimes S^{d-1}V' \subseteq S^d V$; in particular, we have $M''_1 \otimes L_1^{d-1} + \cdots + M''_k \otimes L_k^{d-1} = 0$. If some M''_j is nonzero, we deduce that $L_1^{d-1}, \dots, L_k^{d-1}$ are linearly dependent, which contradicts the initial assumption. Therefore $M''_j = 0$ for every j and $M_j = M'_j$. \square

We can now provide a complete proof of [Theorem 5.1](#).

Proof of Theorem 5.1. Let $\Omega \subseteq \mathbb{P}V^{(n+2)}$ be the subset of sets of points spanning the entire $\mathbb{P}V$. By definition of concise Terracini locus, $v_3(\Omega) \cap \mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ is Zariski open in $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ and non-empty if $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V))$ is non-empty. We prove that $v_3(\Omega) \cap \mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V)) = v_3(K_3)$.

Let $A \in \Omega$ and write $A = \{[L_1], \dots, [L_{n+2}]\}$. Let r be the Kruskal rank of A : by [Lemma 2.6](#), $A \in K_r$ and it can be normalized as $A = \{[x_0], \dots, [x_n], [x_0 + \cdots + x_{r-1}]\}$. Write $T_j = T_{[x_j]}v_3(\mathbb{P}V)$, $T_+ = T_{[x_0 + \cdots + x_{r-1}]}v_3(\mathbb{P}V)$; let $V' = \langle x_0, \dots, x_{r-1} \rangle$ and let $T'_j = T_{[x_j]}v_3(\mathbb{P}V')$, $T'_+ = T_{[x_0 + \cdots + x_{r-1}]}v_3(\mathbb{P}V')$.

If $v_3(A)$ is an element of the Terracini locus, then T_0, \dots, T_n, T_+ are linearly dependent. By [Lemma 5.2](#), the same must hold for the tangent spaces T_0, \dots, T_{r-1}, T_+ . Further, by [Lemma 5.3](#), the same must hold for $T'_0, \dots, T'_{r-1}, T'_+$. In particular, it is enough to $A' \subseteq A$ defined by $A' = \{[x_0], \dots, [x_{r-1}], [x_0 + \cdots + x_{r-1}]\} \in (\mathbb{P}V')^{(r+2)}$.

We deduce that $v_3(A')$ belongs to the concise Terracini locus $\mathbb{T}_{r+2}^{\text{con}}(v_3(\mathbb{P}V'))$. By the action of $\text{GL}(V')$, the same holds for every set of $r+2$ points in linear general position in $\mathbb{P}V'$. Passing to the closure, we obtain $v_3(\mathbb{P}V')^{(r+2)}$ is entirely contained in the Terracini locus. By Terracini's Lemma, we deduce that all fibers of $\pi_\sigma : A\sigma_{r+2}(v_3(\mathbb{P}V')) \rightarrow \mathbb{P}S^3V'$ have positive dimension. [Theorem 1.1](#) guarantees that this is the case only if $r \leq 3$. This shows that $A \subseteq K_3$, or equivalently that it has four coplanar points. We obtained $v_3(\Omega) \cap \mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}V)) \subseteq v_3(K_3)$.

The reverse inclusion holds because of [Theorem 1.1](#) and this concludes the proof of the equality in [\(3\)](#).

Finally, by [Lemma 2.6](#), we conclude that $\mathbb{T}_{n+2}^{\text{con}}(v_3(\mathbb{P}^n)) = \overline{v_3(K_3)}$ is irreducible, and of dimension $n(n+1) + 2$. \square

REFERENCES

- [AC20] E. Angelini and L. Chiantini. On the identifiability of ternary forms. *Lin. Alg. Appl.*, 599:36–65, 2020. doi:10.1016/j.laa.2020.03.042.
- [ACV18] E. Angelini, L. Chiantini, and N. Vannieuwenhoven. Identifiability beyond Kruskal's bound for symmetric tensors of degree 4. *Rend. Lincei - Mat. e Appl.*, 29(3):465–485, 2018. doi:10.4171/RLM/817.
- [BBCG19] E. Ballico, A. Bernardi, M. Christandl, and F. Gesmundo. On the partially symmetric rank of tensor products of W-states and other symmetric tensors. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 30:93–124, 2019. doi:10.4171/RLM/837.
- [BC21] E. Ballico and L. Chiantini. On the Terracini locus of projective varieties. *Milan J. Math.*, 89(1):1–17, 2021. doi:10.1007/s00032-020-00324-5.

- [BCC⁺18] A. Bernardi, E. Carlini, M. Catalisano, A. Gimigliano, and A. Oneto. The Hitchhiker Guide to: Secant Varieties and Tensor Decomposition. *Mathematics*, 6(12):314, 2018. doi:10.3390/math6120314.
- [BCO14] C. Bocci, L. Chiantini, and G. Ottaviani. Refined methods for the Identifiability of Tensors. *Ann. di Mat. Pura ed Appl.*, 193(6):1691–1702, 2014.
- [BGM94] A. Bigatti, A. V. Geramita, and J. C. Migliore. Geometric consequences of extremal behavior in a theorem of Macaulay. *Trans. Amer. Math. Soc.*, 346(1):203–235, 1994. doi:10.1090/S0002-9947-1994-1272673-7.
- [Car06] E. Carlini. Reducing the number of variables of a polynomial. In *Algebraic geometry and geometric modeling*, page 237–247. Springer, 2006. doi:978-3-540-33275-6_15.
- [CC10] L. Chiantini and C. Ciliberto. On the dimension of secant varieties. *J. Europ. Math. Soc.*, 12:1267–1291, 2010. doi:10.4171/JEMS/229.
- [Chi19] L. Chiantini. Hilbert functions and tensor analysis. In *Quantum Physics and Geometry*, pages 125–151. Springer, 2019. doi:10.1007/978-3-030-06122-7_6.
- [Cle61] A. Clebsch. Zur Theorie der algebraischen Flächen. *J. Reine Angew. Math.*, 58:93–108, 1861. doi:10.1515/9783112368480-003.
- [CO21] L. Chiantini and G. Ottaviani. A footnote to a footnote to a paper of B. Segre. *arXiv:2103.04659*, 2021.
- [COV17] L. Chiantini, G. Ottaviani, and N. Vannieuwenhoven. On generic identifiability of symmetric tensors of subgeneric rank. *Trans. Amer. Math. Soc.*, 369(6):4021–4042, 2017. doi:10.1090/tran/6762.
- [CS11] G. Comas and M. Seiguer. On the rank of a binary form. *Found. Comp. Math.*, 11(1):65–78, 2011. doi:10.1007/s10208-010-9077-x.
- [Dav85] E. D. Davis. Complete intersections of codimension 2 in \mathbb{P}^r : the Bezout-Jacobi-Segre theorem revisited. *Rend. Sem. Mat. Univ. Politec. Torino*, 43(2):333–353, 1985.
- [Der13] H. Derksen. Kruskal’s uniqueness inequality is sharp. *Lin. Alg. Appl.*, 438(2):708–712, 2013. doi:10.1016/j.laa.2011.05.041.
- [GM19] F. Galuppi and M. Mella. Identifiability of homogeneous polynomials and Cremona transformations. *J. Reine Angew. Math.*, 2019(757):279–308, 2019. doi:10.1515/crelle-2017-0043.
- [Har92] J. Harris. *Algebraic geometry. A first course*, volume 133 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1992.
- [Hil] D. Hilbert. Lettre adresseé à M. Hermite. *Gesam. Abh.*, II:148–153.
- [IK99] A. Iarrobino and V. Kanev. *Power sums, Gorenstein algebras, and determinantal loci*, volume 1721 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1999.
- [Kru77] J. B. Kruskal. Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics. *Lin. Alg. Appl.*, 18(2):95–138, 1977. doi:10.1016/0024-3795(77)90069-6.
- [Lan12] J. M. Landsberg. *Tensors: Geometry and Applications*, volume 128 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [LP21] B. Lovitz and F. Petrov. A generalization of Kruskal’s theorem on tensor decomposition. *arXiv:2103.15633*, 2021.
- [LT10] J. M. Landsberg and Z. Teitler. On the ranks and border ranks of symmetric tensors. *Found. Comp. Math.*, 10(3):339–366, 2010. doi:10.1007/s10208-009-9055-3.
- [Syl52] J. J. Sylvester. On the principles of the calculus of forms. *Cambridge and Dublin Math. J.*, 7:52–97, 1852.

(L. Chiantini) DIPARTIMENTO DI INGEGNERIA DELL’INFORMAZIONE E SCIENZE MATEMATICHE, UNIVERSITÀ DI SIENA, ITALY

Email address: luca.chiantini@unisi.it

(F. Gesmundo) SAARLAND UNIVERSITY, SAARBRÜCKEN 66123, GERMANY

Email address: gesmundo@cs.uni-saarland.de