

## CYCLIC FROBENIUS ALGEBRAS.

V. M. BUCHSTABER<sup>s</sup>, A. V. MIKHAILOV<sup>h</sup>

Let  $\mathcal{A}$  be an associative  $\mathbb{C}$ -algebra with identity 1, and let  $\mathcal{M}$  be some  $\mathbb{C}$ -linear space,  $\dim \mathcal{M} \geq 1$ .

**Definition.** A cyclic Frobenius algebra (CF-algebra)  $\mathcal{A}$  is an algebra  $\mathcal{A}$  with a  $\mathbb{C}$ -bilinear skew-symmetric form  $\Phi(\cdot, \cdot): \mathcal{A} \otimes_{\mathbb{C}} \mathcal{A} \rightarrow \mathcal{M}$  such that  $\Phi(A, BC) + \Phi(B, CA) + \Phi(C, AB) = 0$  where  $A, B, C \in \mathcal{A}$ .

Examples: 1)  $\mathcal{M} = \mathcal{A}$  and  $\Phi(A, B) = AB - BA = [A, B]$ ; 2)  $\mathcal{A}$  is a commutative algebra with Poisson bracket  $\{\cdot, \cdot\}$ ,  $\mathcal{M} = \mathcal{A}$  and  $\Phi(A, B) = \{A, B\}$ ; 3) CF-algebras, where  $\mathcal{M}$  is a field, obtained from the constructions of algebras from [1], [5].

Let us introduce the  $\mathbb{C}$ -linear subspace  $\text{Span} \subset \mathcal{A}$ , spanned by all commutators  $[A, B] \in \mathcal{A}$ , and the projection  $\pi: \mathcal{A} \rightarrow \mathcal{A}/\text{Span}$ . Set  $A \approx B$  if  $A - B \in \text{Span}$ .

**Lemma 1.** Let  $\varphi(\cdot, \cdot): \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$  be a  $\mathbb{C}$ -bilinear skew-symmetric form such that  $\varphi(A, BC) \approx \varphi(A, B)C + B\varphi(A, C)$ . Then  $\mathcal{A}$  is a CF-algebra with  $\mathcal{M} = \mathcal{A}/\text{Span}$  and  $\Phi = \pi\varphi$ .

Let  $\mathcal{K} = \{B \in \mathcal{A} : \Phi(A, B) = 0, \text{ for all } A \in \mathcal{A}\}$ . From Definition it follows that  $\mathcal{K}$  is the subring in  $\mathcal{A}$ ,  $1 \in \mathcal{K}$ , and  $\Phi(A, bC) = \Phi(Ab, C)$  for all  $A, C \in \mathcal{A}$  and  $b \in \mathcal{K}$ .

Let  $\mathfrak{A} = \mathbb{C}\langle u_0, u_1, \dots \rangle = \bigoplus \mathfrak{A}_k$ ,  $k \geq 0$ , be a free associative graded algebra with derivation  $D$ ,  $|u_k| = k + 2$ ,  $D(u_k) = u_{k+1}$ ,  $k = 0, 1, \dots$ . We introduce the algebra  $\mathfrak{A}^D = \{A = \sum_{i \leq m} a_i D^i, a_m \neq 0, m \in \mathbb{Z}, a_i \in \mathfrak{A}_{|a_m|+m-i}\}$ , where  $[D, u_k] = u_{k+1}$ ,  $[D^{-1}, u_k] = \sum_{i \geq 1} (-1)^i u_{k+i} D^{-i-1}$ . Let  $A_+ + A_- = A \in \mathfrak{A}^D$ , where  $A_+ = \sum a_i D^i$ ,  $0 \leq i \leq m$ , for  $m \geq 0$ , and  $A_+ = 0$ , for  $m < 0$ . We have  $\text{res}[D, A] = D(\text{res}A)$ , where  $\text{res}A = a_{-1}$ .

We introduce a homogeneous  $\mathbb{C}$ -bilinear skew-symmetric form  $\sigma(\cdot, \cdot): \mathfrak{A}^D \otimes \mathfrak{A}^D \rightarrow \mathfrak{A}$ ,  $|\sigma(A, B)| = |A| + |B|$ , by

$$\sigma(aD^n, bD^m) = \begin{cases} \frac{1}{2} \binom{n}{n+m+1} \sum_{s=0}^{n+m} (-1)^s (a^{(s)} b^{(n+m-s)} + b^{(n+m-s)} a^{(s)}), & \text{for } n+m \geq 0, nm < 0, \\ 0, & \text{otherwise.} \end{cases}$$

**Lemma 2.** For any  $A, B \in \mathfrak{A}^D$ , we have  $D(\sigma(A, B)) = \text{res}[A, B] - \Delta(A, B)$ , where

$$\Delta(aD^n, bD^m) = \begin{cases} \frac{1}{2} \binom{n}{n+m+1} \left( [a, b^{(n+m+1)}] + (-1)^{n+m} [b, a^{(n+m+1)}] \right), & \text{for } n+m \geq -1, \\ 0, & \text{otherwise.} \end{cases}$$

**Lemma 3.** Under the canonical projection  $\pi: \mathfrak{A} \rightarrow \mathfrak{A}/\text{Span} = \mathcal{M} = \bigoplus \mathcal{M}_k$ ,  $k \geq 0$ , the operator  $D$  gives monomorphisms  $\bar{D}: \mathcal{M}_k \rightarrow \mathcal{M}_{k+1}$ ,  $k > 0$ .

**Theorem 1.** The algebra  $\mathfrak{A}^D$  is a CF-algebra with the form  $\Phi = \pi\sigma: \mathfrak{A}^D \otimes \mathfrak{A}^D \rightarrow \mathcal{M}$  such that  $\mathcal{K} = \mathfrak{A}$  and  $\Phi(D^n, D^{-n}) = n$ ,  $n \in \mathbb{Z}$ .

---

AVM acknowledges support from EPSRC Grant EP/V050451/1.

**Corollary 1.** *On the set of generators  $\{D^n, n \in \mathbb{Z}\}$  of the free left  $\mathfrak{A}$ -module  $\mathfrak{A}^D$ , the form  $\Phi$  is non-degenerate.*

Let  $L = D^2 - u$  and  $\mathcal{L} = D + \sum I_i D^{-i}$ ,  $i \geq 1$ ,  $I_i \in \mathfrak{A}_{i+1}$ , where  $\mathcal{L}^2 = L$ . We obtain a sequence of series  $\mathcal{L}^{2k-1}$ ,  $k \in \mathbb{N}$ . Set

$$\sigma_{2k-1, 2n-1} = \sigma(\mathcal{L}_+^{2k-1}, \mathcal{L}^{2n-1}) \in \mathfrak{A}_{2n+2k-2}, \quad \rho_{2n} = \sigma_{1, 2n-1} = \text{res } \mathcal{L}^{2n-1}, \quad n > 0. \quad (1)$$

From the properties of the form  $\sigma(\cdot, \cdot)$ , we obtain  $\sigma_{2k-1, 2n-1} = \sigma_{2n-1, 2k-1}$ ,  $k, n \in \mathbb{N}$ .

Let us introduce derivations  $D_{2k-1}$ ,  $k \in \mathbb{N}$ , of the algebra  $\mathfrak{A}$  such that

$$D_1 = D; \quad [D, D_{2k-1}] = 0; \quad D_{2k-1}(u) = -2D(\rho_{2k}), \quad k \in \mathbb{N}. \quad (2)$$

**Corollary 2.**  $[D_{2k-1}, D_{2n-1}] = 0$ .

Let  $u = u(t_1, t_3, \dots)$ . Set  $\partial_{t_{2k-1}}(u) = D_{2k-1}(u)$ .

**Theorem 2.** *The system of equations  $\partial_{t_{2k-1}}(u) = -2D(\rho_{2k}(u))$ ,  $k \in \mathbb{N}$ , where  $\rho_2(u) = -\frac{1}{2}u$ , coincides with the KdV hierarchy on  $\mathfrak{A}$*

$$4\partial_{t_3}(u) = D(u_2 - 3u^2); \quad 16\partial_{t_5}(u) = D(u_4 - 5u_2u - 5uu_2 - 5u_1^2 + 10u^3); \quad \dots$$

The *proof* follows from the above constructions (cf. [2]–[6]).

Following [3], for  $N \in \mathbb{N}$ , set  $F_{2N+2}(u) = \rho_{2N+2} + \sum_{k=0}^{N-1} \alpha_{2(N-k+1)} \rho_{2k}$ , where  $\rho_0 = 1$  and  $\alpha_4, \dots, \alpha_{2N+2}$  are free parameters. Equation  $F_{2N+2}(u) = 0$  is called the  $N$ -th Novikov equation. Let  $J(F)$  be the two-sided  $D$ -differential ideal in  $\mathfrak{A}$  generated by the polynomial  $F_{2N+2}(u)$ . Since  $2^{2k+1} \rho_{2k+2} = u_{2k} + f(u_{2k-2}, \dots, u)$  and  $u_{k+1} = D^k(u)$ , on the factor-algebra  $\mathfrak{A}/J(F) = \mathbb{C}\langle u, \dots, u_{2N-1} \rangle$ , the KdV hierarchy (see Theorem 2) reduces to the  $N$ -th Novikov hierarchy, where the first system represents the  $N$ -th Novikov equation in the form  $D(u_k) = u_{k+1}$  for  $0 \leq k < 2N - 1$ ,  $D(u_{2N-1}) = -f(u_{2k-2}, \dots, u)$ . It was shown in [3] that, in terms of the form  $\Phi$ , (see Theorem 1), the polynomials

$$H_{2n+1, 2N+1} = \sigma_{2n+1, 2N+1} + \sum_{k=1}^{N-1} \alpha_{2(N-k+1)} \sigma_{2n+1, 2k-1}, \quad n = 1, \dots, N, \quad (3)$$

define first integrals  $\widehat{H}_{2n+1, 2N+1} = \pi(H_{2n+1, 2N+1})$  of the  $N$ -th Novikov hierarchy:

$$\pi(\partial_{t_{2k-1}}(\widehat{H}_{2n+1, 2N+1})) = 0, \quad k, n \in \{1, \dots, N\}.$$

**Theorem 3.** *In the quantum case, the  $N$ -th Novikov hierarchy (see [3]) is written in the form of compatible systems of Heisenberg equations. The polynomials (3) are quantum commuting Hamiltonians of the hierarchy and are self-adjoint if the variables  $u_k$  are self-adjoint and the values of the parameters  $\alpha_4, \dots, \alpha_{2N+2}$  are real.*

We would like to thank S.P. Novikov and V.N. Rubtsov for stimulative discussions of the results of this paper.

## REFERENCES

- [1] M. Aguiar, *On the associative analog of Lie bialgebras.*, J. Algebra 244:2 (2001), 492–532.
- [2] V. M. Buchstaber, A. V. Mikhailov, *Integrable polynomial Hamiltonian systems and symmetric powers of plane algebraic curves.*, Russian Math. Surv., 76:4(460)(2021), 587–652.
- [3] V. M. Buchstaber, A. V. Mikhailov, *KdV hierarchies and quantum Novikov's equations.*, arXiv:2109.06357 v2 [nlin.SI].
- [4] I. M. Gelfand, L. A. Dikii, *Asymptotic behaviour of the resolvent of Sturm-Liouville equations and the algebra of the Korteweg-de Vries equations.*, Russian Math. Surveys, 30:5 (1975), 77–113.

- [5] A. V. Odesskii, V. N. Rubtsov, V. V. Sokolov, *Bi-Hamiltonian ordinary differential equations with matrix variables*, Theor. Math. Phys., 171:1 (2012), 26–32.
- [6] V. V. Sokolov, *Algebraic Structures in Integrability*, World Sci. Publ., Hackensack, NJ, 2020.

<sup>§</sup> STEKLOV MATHEMATICAL INSTITUTE, MOSCOW, DEPARTMENT OF MECHANICS AND MATHEMATICS,  
MOSCOW STATE UNIVERSITY, RUSSIA  
*Email address:* buchstab@mi-ras.ru

<sup>‡</sup> UNIVERSITY OF LEEDS, LEEDS, UK  
*Email address:* a.v.mikhailov@leeds.ac.uk