

Singleton-Optimal LRCs and Perfect LRCs via Cyclic and Constacyclic Codes*

WeiJun Fang^{†,1,2,3}, Fang-Wei Fu⁴, Bin Chen⁵, and Shu-Tao Xia⁶

¹Key Laboratory of Cryptologic Technology and Information Security,
Ministry of Education, Shandong University, Qingdao, China

²School of Cyber Science and Technology, Shandong University,
Qingdao, China

³Quancheng Laboratory, Jinan, China

⁴Chern Institute of Mathematics and LPMC, Nankai University,
Tianjin, China

⁵Department of Computer Science and Technology, Harbin Institute of
Technology, Shenzhen, China

⁶Tsinghua Shenzhen International Graduate School, Tsinghua
University, Shenzhen, China

Abstract

Locally repairable codes (LRCs) have emerged as an important coding scheme in distributed storage systems (DSSs) with relatively low repair cost by accessing fewer non-failure nodes. Theoretical bounds and optimal constructions of LRCs have been widely investigated. Optimal LRCs via cyclic and constacyclic codes provide significant benefit of elegant algebraic structure and efficient encoding procedure. In this paper, we continue to consider the constructions of optimal LRCs via cyclic and constacyclic codes with long code length. Specifically, we first obtain two classes of q -ary cyclic Singleton-optimal $(n, k, d = 6; r = 2)$ -LRCs with length $n = 3(q+1)$ when $3 \mid (q-1)$ and q is even, and length $n = \frac{3}{2}(q+1)$ when $3 \mid (q-1)$ and $q \equiv 1 \pmod{4}$, respectively. To the best of our knowledge, this is the first construction of q -ary cyclic

*This paper was presented in part at the 2021 IEEE International Symposium on Information Theory (ISIT) [11]

[†]Corresponding author: fwj@sdu.edu.cn

Singleton-optimal LRCs with length $n > q + 1$ and minimum distance $d \geq 5$. On the other hand, an LRC achieving the Hamming-type bound is called a perfect LRC. By using cyclic and constacyclic codes, we construct two new families of q -ary perfect LRCs with length $n = \frac{q^m - 1}{q - 1}$, minimum distance $d = 5$ and locality $r = 2$.

Keywords: Local repairable codes, Singleton-optimal LRCs, perfect LRCs, cyclic codes, constacyclic codes

1 Introduction

Distributed storage systems (DSSs) have become a popular way to store large amounts of data in modern data center, e.g., Windows [21] and Facebook [31]. By distributing huge data over multiple storage nodes, DSSs can not only reduce the storage cost, but also improve the reliability and scalability. However, node failures become a norm in DSSs rather than an exception. Thus erasure codes that optimize the repair bandwidth and repair degree (the number of nodes used in a repairing process) for recovering a failed node have been adopted in DSSs. In [14], Gopalan *et al.* first introduced the locality in an erasure code to reduce the repair degree, that is, a failed node can be recovered by accessing few available nodes. Let C be a q -ary $[n, k, d]$ linear code, which is just a subspace of \mathbb{F}_q^n of dimension k with minimum Hamming distance d . Formally, C is called a *locally repairable code* (LRC) with *locality* r if for any $i \in [n] \triangleq \{1, 2, \dots, n\}$, there exists a subset $R_i \subset [n]$ with $i \in R_i$ and $|R_i| \leq r + 1$ such that the i -th symbol c_i can be recovered by $\{c_j\}_{j \in R_i \setminus \{i\}}$, i.e., c_i can be represented as a linear combination of $\{c_j\}_{j \in R_i \setminus \{i\}}$. R_i is called the *repair group* of the i -th symbol. An LRC is called to have *disjoint local repair groups* if for any $1 \leq i \neq j \leq n$, we have $|R_i| = r + 1$ and either $R_i \cap R_j = \emptyset$ or $R_i = R_j$. There are some restrictions among the parameters n, k, d, r . One of these bounds is the famous Singleton-type bound derived in [14], which said that for any q -ary $[n, k, d]$ -LRCs with locality r ,

$$d \leq n - k + 2 - \lceil k/r \rceil. \quad (1)$$

Note that when $r = k$, (1) reduces to the classical Singleton bound.

Definition 1 (Singleton-optimal LRCs). *A q -ary $[n, k, d]$ linear code C with locality r is called a Singleton-optimal $(n, k, d; r)$ -LRC if it achieves the bound (1) with equality.*

From the Singleton-type bound (1), a Singleton-optimal $(n, k, d; r)$ -LRC has to satisfy that $\frac{k}{n} = \frac{r}{r+1} - \frac{d-2}{n}$ (suppose $r \mid k$). Thus for fixed d and r , the longer code length of a Singleton-optimal LRC leads to larger code rate. The construction of Singleton-optimal LRCs with long code length has been a hot topic in coding theory in recent years (see [6, 7, 8, 16, 24, 25, 26, 28, 33, 34, 38]). In [33], Tamo and Barg first gave a breakthrough construction of Singleton-optimal LRCs via subcodes of Reed-Solomon codes whose code length can go up to the alphabet size. Due to their efficient encoding and decoding performance, cyclic codes and constacyclic codes have been widely used in the constructions of optimal LRCs (see [6, 8, 28, 34]). For $n \mid (q + 1)$, by using the

techniques of cyclic and constacyclic codes, Chen *et al.* [6, 8] have constructed the q -ary Singleton-optimal LRCs for all possible parameters. In [25], Jin *et al.* also presented a construction of q -ary optimal LRCs with length $n = q + 1$ via the automorphism groups of rational function fields. For minimum distance $d = 3$ or 4, Luo *et al.* [28] proposed a construction of cyclic optimal LRCs with unbounded length. For minimum distance $d \geq 5$, Guruswami *et al.* [16] proved that the code length of a q -ary optimal LRC must be at most roughly $O(dq^3)$ and presented an algorithmic construction of q -ary optimal LRCs of length $\Omega_{d,r}(q^{1+1/\lfloor \frac{d-3}{2} \rfloor})$ for $r \geq d - 2$. By employing algebraic structures of elliptic curves, Li *et al.* [26] constructed a family of q -ary optimal LRCs with length up to $q + 2\sqrt{q}$ and some particular localities. For $d = 5, 6$ and $r \geq d - 2$, Jin [24] presented an explicit construction of q -ary optimal LRCs of length $\Omega_r(q^2)$ via binary constant weight codes. In [38], Xing and Yuan generalized Jin's results and presented a construction of q -ary optimal LRCs for general $d \geq 7$ via hypergraph theory. In [7], Chen *et al.* considered the case of $r < d - 2$. In particular, for $d = 6$ and $r = 2$, the authors established an equivalent connection between the existence of Singleton-optimal LRCs and the existence of lines in finite projective plane with certain properties. And they also provided some new constructions and improved bounds on Singleton-optimal LRCs. However, as we have seen, for $d \geq 5$, the longest code length of known q -ary cyclic Singleton-optimal LRCs is $q + 1$ until now. So it is quite interesting to construct q -ary cyclic Singleton-optimal LRCs with length $n > q + 1$ and $d \geq 5$.

On the other hand, the Singleton-type bound (1) is not always achievable in some cases (see [32, 35]), and there are only a few Singleton-optimal LRCs over binary or ternary fields (see [17, 18]). Some other theoretical bounds on the parameters of LRCs were also investigated in the literature. Cadambe and Mazumdar [4] first derived a theoretical bound of LRCs depending on the field size, which is known as C - M bound. Some well-known binary codes, such as Hamming codes, Simplex codes etc., are optimal with respect to the C - M bound (see [22]). In [1], Agarwal *et al.* presented several new combinatorial bounds on LRC codes including the locality-aware sphere packing and Plotkin bounds. Recently, Tebbi *et al.* [36] also obtained a linear programming bound for LRCs with arbitrary code parameters. However, it is not easy to give an explicit form of the C - M bound or the bound in [36] in general. In [19], Hao *et al.* generalized the C - M bound and obtained a Griesmer-type bound. In [37], Wang *et al.* derived a new explicit bound on the size of LRCs via a *sphere-packing based approach*. Precisely, they proved that

$$A_q(n, d; r) \leq \frac{q^{\dim(\mathcal{V})}}{B_{\mathcal{V}}(\lfloor (d-1)/2 \rfloor)}, \quad (2)$$

where $A_q(n, d; r)$ is the maximal number of codewords of a q -ary LRC of length n , minimum distance d , locality r and with disjoint local repair groups, \mathcal{V} is the \mathbb{F}_q -linear space containing all of $(n, d; r)$ -LRCs with disjoint local repair groups and $B_{\mathcal{V}}(\lfloor \frac{d-1}{2} \rfloor) \triangleq |\{\mathbf{v} \in \mathcal{V} : wt(\mathbf{v}) \leq \lfloor \frac{d-1}{2} \rfloor\}|$. Note that when \mathcal{V} is the whole space \mathbb{F}_q^n , the bound (2) reduces to the well-known classical Hamming bound. On the other hand, as the authors mentioned in [37], they obtained the bound (2) by using the essential idea of the Hamming bound, not just its expression. Thus, in this paper, we regard bound (2) as a *Hamming-type bound* for LRCs with disjoint local repair groups. Moreover, in [37, Theorem 11], taking \mathcal{V} as the

dual space of the locality rows of the parity-check matrix, they proved that

$$B_{\mathcal{V}}(\lfloor \frac{d-1}{2} \rfloor) = \sum_{0 \leq i_1 + \dots + i_\ell \leq \lfloor \frac{d-1}{2} \rfloor} \prod_{j=1}^{\ell} \beta(r, i_j),$$

where $\beta(r, i) = \frac{1}{q}((q-1)^i + (-1)^i(q-1)) \binom{r+1}{i}$. Thus we can rewrite bound (2) as a more explicit form, i.e.,

$$q^k \leq \frac{q^{\frac{rn}{r+1}}}{\sum_{0 \leq i_1 + \dots + i_\ell \leq \lfloor \frac{d-1}{2} \rfloor} \prod_{j=1}^{\ell} \beta(r, i_j)}. \quad (3)$$

It can be seen that for $d = 5$ or 6 , bound (3) becomes

$$q^k \leq \frac{q^{\frac{r}{r+1}n}}{\frac{rn}{2}(q-1) + 1}. \quad (4)$$

Definition 2 (Perfect LRCs, [10, 12]). *A q -ary $(n, k, d; r)$ -LRC with disjoint local repair groups is called a perfect LRC if it achieves the bound (2) (or (3)) with equality.*

As we known, Hamming codes, Binary and Ternary Golay codes are the all nontrivial perfect linear codes in classical coding theory, so it is expected to determine perfect LRCs for all possible parameters. From the Hamming-type bound (3), we know that if there exists a q -ary $(n, k, d; r)$ -perfect LRC with disjoint repair groups, then $\sum_{0 \leq i_1 + \dots + i_\ell \leq \lfloor \frac{d-1}{2} \rfloor} \prod_{j=1}^{\ell} \beta(r, i_j)$

should be exactly a power of q . So it is quite difficult to construct perfect LRCs and there are few works on constructions of perfect LRCs. In [15], Goparaju *et al.* have presented a family of binary perfect $(n = 2^u - 1, k = \frac{2}{3}n - u, d = 6; r = 2)$ -LRCs for even u , which is the only one known class of cyclic perfect LRCs until now. Some new classes of perfect LRCs have been presented in [10, 12] via the techniques of finite geometry and finite fields.

In this paper, firstly, we will give two new constructions of cyclic Singleton-optimal LRCs with longer length. Precisely, for $3 \mid (q-1)$ and q is even, we give a construction of q -ary cyclic optimal LRCs with length $n = 3(q+1)$, minimum distance $d = 6$ and locality $r = 2$; For $3 \mid (q-1)$ and $q \equiv 1 \pmod{4}$, we also obtain a family of q -ary cyclic Singleton-optimal LRCs with length $n = \frac{3}{2}(q+1)$, minimum distance $d = 6$ and locality $r = 2$. To the best of our knowledge, this is the first construction of q -ary cyclic Singleton-optimal LRCs with length $n > q+1$ and minimum distance $d \geq 5$. Secondly, for even $q \geq 4$ and even m with $3 \mid (q+1)$ and $\gcd(m, q-1) = 1$, we construct a family of cyclic $(n = \frac{q^m-1}{q-1}, k = \frac{2}{3}n - m, d = 5; r = 2)$ -perfect LRCs. Moreover, for general q with $3 \mid (q+1)$ and even m , we construct a family of constacyclic $(n = \frac{q^m-1}{q-1}, k = \frac{2}{3}n - m, d = 5; r = 2)$ -perfect LRCs. To the best of our knowledge, this is the first construction of q -ary (consta)cyclic perfect LRCs for general q .

The rest of this paper is organized as follows. In Section 2, we briefly review some basic results about LRCs and (consta)cyclic codes. In Section 3, we provide our new constructions of cyclic Singleton-optimal LRCs. In Section 4, we give two new constructions of cyclic and constacyclic perfect LRCs, respectively. We conclude this paper in Section 5.

2 Preliminaries

In this section, we briefly review some basic results about cyclic codes, constacyclic codes and LRCs.

Suppose q is a prime power and \mathbb{F}_q is a finite field with q elements. Denote $[n] \triangleq \{1, 2, \dots, n\}$. For a vector $\mathbf{v} = (v_1, v_2, \dots, v_n) \in \mathbb{F}_q^n$, the support of \mathbf{v} is defined as $\text{supp}(\mathbf{v}) \triangleq \{i \in [n] : v_i \neq 0\}$. The Hamming weight $wt(\mathbf{v})$ of \mathbf{v} is defined as the size of $\text{supp}(\mathbf{v})$, i.e., $wt(\mathbf{v}) = |\text{supp}(\mathbf{v})|$. The Hamming distance between \mathbf{u} and \mathbf{v} is defined as $d(\mathbf{u}, \mathbf{v}) \triangleq wt(\mathbf{u} - \mathbf{v})$. A q -ary linear code C of length n and dimension k is just a k -dimensional subspace of \mathbb{F}_q^n . The minimum distance of C is defined as $d \triangleq \min_{\mathbf{c} \in C, \mathbf{c} \neq \mathbf{0}} wt(\mathbf{c})$.

An $[n, k, d]_q$ -linear code C is a linear code of length n , dimension k and minimum distance d over \mathbb{F}_q .

We recall some basic results about constacyclic codes, which are generalizations of cyclic codes. The readers may refer some standard textbooks on coding theory for more details on cyclic codes ([23, 29]) and some references on constacyclic codes ([2, 3, 5, 9, 13, 30, 39]).

Definition 3 (λ -constacyclic codes and cyclic codes). *Suppose $\lambda \in \mathbb{F}_q^*$. A q -ary linear code C of length n is called a λ -constacyclic code if $(c_0, c_1, \dots, c_{n-1}) \in C$ implies that $(\lambda c_{n-1}, c_0, \dots, c_{n-2}) \in C$. In particular, if $\lambda = 1$, a λ -constacyclic code C is called a cyclic code.*

If we identify a vector $\mathbf{c} = (c_0, c_1, \dots, c_{n-1})$ with a polynomial $c(x) = c_0 + c_1x + \dots + c_{n-1}x^{n-1}$, then a q -ary λ -constacyclic code C of length n can be identified with an ideal of the ring $\mathbb{F}_q[x]/(x^n - \lambda)$. Note that $\mathbb{F}_q[x]/(x^n - \lambda)$ is a principal domain, thus every ideal of $\mathbb{F}_q[x]/(x^n - \lambda)$ is generated by a monic polynomial $g(x)$ with $g(x) \mid (x^n - \lambda)$. We also define the Hamming weight of $c(x)$ as $wt(c(x)) \triangleq wt(\mathbf{c})$, the Hamming weight of the corresponding vector \mathbf{c} .

Definition 4 (Generator polynomial). *Suppose $\lambda \in \mathbb{F}_q^*$ and C is a λ -constacyclic code of length n generated by a monic polynomial $g(x)$ with $g(x) \mid (x^n - \lambda)$, then $g(x)$ is called the generator polynomial of C , and we denote $C = \langle g(x) \rangle$. It is well-known that $\dim(C) = n - \deg(g(x))$.*

For any polynomial $f(x) = f_0 + f_1x + \dots + f_mx^m \in \mathbb{F}_q[x]$, where $f_0, f_m \neq 0$, define its reciprocal polynomial $\tilde{f}(x) \triangleq f_m + f_{m-1}x + \dots + f_0x^m = x^m f(x^{-1})$. For convenience, we denote $\bar{f}(x) = f_0^{-1} \tilde{f}(x)$, which is a monic polynomial.

The following lemma can be easily derived from the definitions.

Lemma 1. *For any polynomial $f(x), f_1(x) \in \mathbb{F}_q[x]$ with $f(0)$ and $f_1(0)$ are nonzeros, we have*

- (i) $\deg(f(x)) = \deg(\bar{f}(x))$ and $wt(f(x)) = wt(\bar{f}(x))$.
- (ii) $f_1(x) \mid f(x)$ if and only if $\bar{f}_1(x) \mid \bar{f}(x)$.

Similar to cyclic codes, the dual code of a λ -constacyclic code is a λ^{-1} -constacyclic code. Precisely,

Lemma 2 ([9, 13, 39]). *Let $\lambda \in \mathbb{F}_q^*$ and $C = \langle g(x) \rangle$ be a λ -constacyclic code of length n , where $g(x)$ is the generator polynomial of C . Denote $h(x) = \frac{x^n - \lambda}{g(x)}$. Then the dual code C^\perp of C is a λ^{-1} -constacyclic code and $C^\perp = \langle \bar{h}(x) \rangle$.*

We now always assume that $\gcd(n, q) = 1$. Let \mathbb{F}_{q^m} be the splitting field of $x^n - \lambda$ over \mathbb{F}_q . Suppose $\theta \in \mathbb{F}_{q^m}$ such that $\theta^n = \lambda$. Let s be the order of q modulo n , i.e., the least number i such that $n \mid (q^i - 1)$. Then we can choose a primitive n -th root of unity $\alpha \in \mathbb{F}_{q^s}$ and the roots of $x^n - \lambda$ are $\theta, \theta\alpha, \dots, \theta\alpha^{n-1}$.

The BCH bound and Hartmann-Tzeng bound [20] are two well-known lower bounds on the minimum distance of cyclic codes. In [2], the authors presented the BCH bound for constacyclic codes. The Hartmann-Tzeng bound for constacyclic codes has been obtained in [30]. We give an alternative proof of the Hartmann-Tzeng bound for constacyclic codes for completeness.

Lemma 3 (BCH Bound for Constacyclic Codes [2]). *Let $C = \langle g(x) \rangle$ be a λ -constacyclic code of length n , where $g(x) \mid (x^n - \lambda)$ is the generator polynomial of C . Suppose u and b are integers with $\gcd(b, n) = 1$. If $g(\theta\alpha^{u+ib}) = 0$, for $i = 0, 1, \dots, \delta - 2$, where $\delta \geq 2$, then the minimum distance $d(C)$ of C is at least δ .*

Lemma 4 (Hartmann-Tzeng Bound for Constacyclic Codes [30]). *Let $C = \langle g(x) \rangle$ be a λ -constacyclic code of length n , where $g(x) \mid (x^n - \lambda)$ is the generator polynomial of C . Suppose u, b_1 and b_2 are integers with $\gcd(b_1, n) = \gcd(b_2, n) = 1$. If $g(\theta\alpha^{u+i_1b_1+i_2b_2}) = 0$, for $i_1 = 0, 1, \dots, \delta - 2$; $i_2 = 0, 1, \dots, \gamma$, where $\delta \geq 2$ and $\gamma \geq 0$, then the minimum distance $d(C)$ of C is at least $\delta + \gamma$.*

Proof. Firstly, by the conditions, we know the minimum distance $d(C) \geq \delta$ from Lemma 3. Let $c(x) \in C$ be a nonzero codeword with $wt(c(x)) = w$. We suppose $\delta \leq w \leq \delta + \gamma - 1$. Since C is a λ -constacyclic code, without loss of generality, we may assume that

$$c(x) = 1 + \sum_{j=1}^{w-1} c_j x^{k_j},$$

where $c_i \in \mathbb{F}_q^*$ and $0 < k_1 < k_2 < \dots < k_{w-1} < n$. Denote

$$S_i = c(\theta\alpha^i) - 1 = \sum_{j=1}^{w-1} c_j (\theta\alpha^i)^{k_j}.$$

Note that $g(x) \mid c(x)$, hence $c(\theta\alpha^{u+i_1b_1+i_2b_2}) = 0$, for $i_1 = 0, 1, \dots, \delta - 2$; $i_2 = 0, 1, \dots, \gamma$. Thus

$$S_{u+i_1b_1+i_2b_2} = -1, \text{ for } i_1 = 0, 1, \dots, \delta - 2; i_2 = 0, 1, \dots, \gamma.$$

Let

$$f_1(x) = \prod_{j_1=1}^{\delta-2} (x - \alpha^{k_{j_1} b_1}) = \sum_{i_1=0}^{\delta-2} \sigma_{i_1} x^{i_1},$$

and

$$f_2(x) = \prod_{j_2=\delta-1}^{w-1} (x - \alpha^{k_{j_2} b_2}) = \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} x^{i_2}.$$

Since $\gcd(b_1, n) = \gcd(b_2, n) = 1$ and $0 < k_1 < k_2 < \dots < k_{w-1} < n$, we have $\alpha^{k_{j_1} b_1} \neq 1$ and $\alpha^{k_{j_2} b_2} \neq 1$ for any $1 \leq j_1, j_2 \leq w-1$. Thus

$$f_1(1)f_2(1) = \prod_{j_1=1}^{\delta-2} (1 - \alpha^{k_{j_1} b_1}) \prod_{j_2=\delta-1}^{w-1} (1 - \alpha^{k_{j_2} b_2}) \neq 0.$$

On the other hand,

$$\begin{aligned} f_1(1)f_2(1) &= \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{i_1=0}^{\delta-2} \sigma_{i_1} = - \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{i_1=0}^{\delta-2} S_{u+i_1 b_1+i_2 b_2} \sigma_{i_1} \\ &= - \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{i_1=0}^{\delta-2} \sigma_{i_1} \sum_{j=1}^{w-1} c_j (\theta \alpha^{u+i_1 b_1+i_2 b_2})^{k_j} \\ &= - \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{j=1}^{w-1} c_j (\theta \alpha^{u+i_2 b_2})^{k_j} \sum_{i_1=0}^{\delta-2} \sigma_{i_1} (\alpha^{k_j b_1})^{i_1} \\ &= - \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{j=1}^{w-1} c_j (\theta \alpha^{u+i_2 b_2})^{k_j} f_1(\alpha^{k_j b_1}) \\ &= - \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} \sum_{j=\delta-1}^{w-1} c_j (\theta \alpha^{u+i_2 b_2})^{k_j} f_1(\alpha^{k_j b_1}) \\ &= - \sum_{j=\delta-1}^{w-1} c_j \theta^{k_j} \alpha^{u k_j} f_1(\alpha^{k_j b_1}) \sum_{i_2=0}^{w-\delta-1} \tau_{i_2} (\alpha^{k_j b_2})^{i_2} \\ &= - \sum_{j=\delta-1}^{w-1} c_j \theta^{k_j} \alpha^{u k_j} f_1(\alpha^{k_j b_1}) f_2(\alpha^{k_j b_2}) = 0, \end{aligned}$$

where the 6-th equality holds since $f_1(\alpha^{k_j b_1}) = 0$ for any $1 \leq j \leq \delta-2$, and the last equality holds since $f_2(\alpha^{k_j b_2}) = 0$ for any $\delta-1 \leq j \leq w-1$. Thus it leads to a contradiction. So we have $w \geq \delta + \gamma$, hence $d(C) \geq \delta + \gamma$. The lemma is proved. \square

Recall that C is called an r -LRC if and only if each symbol c_i of any codeword \mathbf{c} can be represented as a linear combination of $\{c_j\}_{j \in R_i \setminus \{i\}}$ for some $R_i \subset [n]$ with $i \in R_i$ and $|R_i| \leq r+1$. It can be equivalently defined in a viewpoint of dual codes:

Proposition 1 ([19]). *Let C be a linear code of length n . Then the i -th symbol of C has locality r if and only if there exists a dual codeword $\mathbf{h} \in C^\perp$ such that*

$$i \in \text{supp}(\mathbf{h}) \text{ and } |\text{supp}(\mathbf{h})| \leq r+1.$$

In this paper, an $(n, k, d; r)$ -LRC means a linear code of length n , dimension k , minimum distance d and locality r .

Due to their nice algebraic structures, cyclic and constacyclic codes have been widely used in the constructions of optimal LRCs (or (r, δ) -LRCs), e.g., [6, 8, 33]. The following proposition presented a sufficient condition to ensure that a constacyclic code is an LRC.

Proposition 2 ([6]). *Suppose $(r+1) \mid n$, $\gcd(n, q) = 1$, $\lambda = \theta^n \in \mathbb{F}_q^*$ and α is a primitive n -th root of unity. Let $C = \langle g(x) \rangle$ be a constacyclic code of length n , where $g(x) \mid (x^n - \lambda)$ is the generator polynomial of C . If $g(\theta\alpha^j\alpha^{i(r+1)}) = 0$, for $i = 0, 1, \dots, \frac{n}{r+1} - 1$ and some integer j , then C has locality r .*

Proof. Since $g(\theta\alpha^j\alpha^{i(r+1)}) = 0$, for $i = 0, 1, \dots, \frac{n}{r+1} - 1$, we have

$$\prod_{i=0}^{\frac{n}{r+1}-1} (x - \theta\alpha^j\alpha^{i(r+1)}) \mid g(x),$$

i.e., $(x^{\frac{n}{r+1}} - (\theta\alpha^j)^{\frac{n}{r+1}}) \mid g(x)$. Denote $\xi = \theta\alpha^j$, then $\xi^n = \lambda$ and $x^n - \lambda = x^n - \xi^n = (x^{\frac{n}{r+1}} - \xi^{\frac{n}{r+1}})f(x)$, where $f(x) = \sum_{i=0}^r x^{i\frac{n}{r+1}}\xi^{(r-i)\frac{n}{r+1}}$. Then $wt(f(x)) = r+1$ and

$$h(x) = \frac{x^n - \lambda}{g(x)} \mid \frac{x^n - \lambda}{x^{\frac{n}{r+1}} - \xi^{\frac{n}{r+1}}} = f(x).$$

From Lemma 1, we have $\bar{h}(x) \mid \bar{f}(x)$. Note that $C^\perp = \langle \bar{h}(x) \rangle$ by Lemma 2 and $wt(\bar{f}(x)) = wt(f(x)) = r+1$ by Lemma 1. Thus $\bar{f}(x)$ is a codeword of C^\perp of weight $r+1$. According to Proposition 1, the first symbol has r -locality. Then we can deduce that all symbols of C have r -locality since the constacyclicity of C . \square

3 Cyclic Singleton-Optimal LRCs

3.1 Cyclic Singleton-Optimal LRCs for even q

In this subsection, for q is even and $3 \mid (q-1)$, we will give a construction of q -ary cyclic Singleton-optimal LRCs with length $n = 3(q+1)$, minimum distance $d = 6$ and locality $r = 2$. Note that $n \mid (q^2 - 1)$, thus we can choose $\alpha \in \mathbb{F}_{q^2}$, which is a primitive n -th root of unity.

Theorem 1. *With the above notation, let*

$$L = \{1, \alpha^3, \alpha^6, \dots, \alpha^{3q}\}$$

and

$$D = \{\alpha, \alpha^{q-1}, \alpha^q, \alpha^{q+1}, \alpha^{q+2}\}.$$

Set

$$g(x) = \prod_{a \in L \cup D} (x - a)$$

and $C = \langle g(x) \rangle$. Then C is a q -ary cyclic Singleton-optimal LRC with length $n = 3(q+1)$, minimum distance $d = 6$ and locality $r = 2$.

Proof. Since $\alpha^{3(q+1)} = 1$, we have

$$\prod_{a \in L} (x - a) = \prod_{i=0}^q (x - \alpha^{3i}) = x^{q+1} - 1.$$

Since $3 \mid (q-1)$, we have $L \cap D = \{\alpha^{q-1}, \alpha^{q+2}\}$, thus

$$g(x) = (x^{q+1} - 1)(x - \alpha)(x - \alpha^q)(x - \alpha^{q+1}).$$

Since $\alpha^{q+1} \in \mathbb{F}_q$, thus $g(x) \in \mathbb{F}_q[x]$ and $k = \dim(C) = n - \deg(g(x)) = 2q - 1$. According to the set L , C has locality $r = 2$ by Proposition 2. Since $\alpha^{q-1}, \alpha^q, \alpha^{q+1}, \alpha^{q+2}$ are roots of $g(x)$, we have $d \geq 5$ by Lemma 3. On the other hand, by the Singleton-type bound (1), we have

$$d \leq n - k + 2 - \lceil k/r \rceil = 3(q+1) - (2q-1) + 2 - q = 6.$$

Thus we only need to prove that $d \neq 5$. By contradiction, we suppose $d = 5$. Let

$$c(x) = c_0 + c_1x^{k_1} + c_2x^{k_2} + c_3x^{k_3} + c_4x^{k_4}$$

be a codeword of C with $c_0, c_1, c_2, c_3, c_4 \in \mathbb{F}_q^*$ and $k_0 = 0 < k_1, k_2, k_3, k_4 \leq n-1$, where k_1, k_2, k_3, k_4 are pairwise distinct. Then $g(x) \mid c(x)$. Let

$$I = \{\ell : 0 \leq \ell \leq 4 \text{ and } (q+1) \mid k_\ell\}.$$

Since $k_0 = 0 < k_1, k_2, k_3, k_4 \leq n-1 = 3(q+1) - 1$, we have $1 \leq |I| \leq 3$. Note that α is a primitive n -th root of unity, we can show that

$$\sum_{i=0}^q (\alpha^{3i})^{k_\ell} = \begin{cases} 1, & \text{for } \ell \in I, \\ 0, & \text{for } \ell \notin I. \end{cases} \quad (5)$$

For any $0 \leq i \leq q$, we have $c(\alpha^{3i}) = 0$. Thus by Eq. (5)

$$\begin{aligned} 0 &= \sum_{i=0}^q c(\alpha^{3i}) = \sum_{i=0}^q \sum_{\ell=0}^4 c_\ell (\alpha^{3i})^{k_\ell} \\ &= \sum_{\ell=0}^4 c_\ell \sum_{i=0}^q (\alpha^{3i})^{k_\ell} = \sum_{\ell \in I} c_\ell, \end{aligned}$$

which implies that $|I| \geq 2$.

Case 1: $|I| = 3$. Without loss of generality, we suppose

$$I = \{0, 1, 2\}, k_1 = q+1, k_2 = 2(q+1), \text{ and } (q+1) \nmid k_3.$$

Thus $c_0 + c_1 + c_2 = 0$. By $c(1) = 0$, we have

$$c_0 + c_1 + c_2 + c_3 + c_4 = 0 \Rightarrow c_3 + c_4 = 0.$$

By $c(\alpha^3) = 0$, we have

$$\begin{aligned} c_0 + c_1(\alpha^3)^{q+1} + c_2(\alpha^3)^{2(q+1)} + c_3(\alpha^3)^{k_3} + c_4(\alpha^3)^{k_4} &= 0 \\ \Rightarrow c_3\alpha^{3k_3} + c_4\alpha^{3k_4} &= 0 \Rightarrow \alpha^{3k_3} = \alpha^{3k_4} \\ &\Rightarrow (q+1) \mid (k_4 - k_3). \end{aligned}$$

By $c(\alpha) = 0$, we have

$$\begin{aligned} c_0 + c_1\alpha^{q+1} + c_2\alpha^{2(q+1)} + c_3\alpha^{k_3} + c_4\alpha^{k_4} &= 0 \\ \Rightarrow c_3\alpha^{k_3} + c_4\alpha^{k_4} \in \mathbb{F}_q &\Rightarrow c_3\alpha^{k_3}(1 - \alpha^{k_4-k_3}) \in \mathbb{F}_q. \end{aligned}$$

Since $(q+1) \mid (k_4 - k_3)$ and $n \nmid (k_4 - k_3)$, we have $\alpha^{k_4-k_3} \neq 1 \in \mathbb{F}_q$. Thus

$$\alpha^{k_3} \in \mathbb{F}_q^* \Rightarrow \alpha^{k_3(q-1)} = 1 \Rightarrow n \mid k_3(q-1).$$

Since q is even, $\gcd(\frac{n}{3}, q-1) = \gcd(q+1, q-1) = 1$. Thus $\frac{n}{3} \mid k_3$, which is a contradiction.

Case 2: $|I| = 2$. Without loss of generality, we suppose

$$I = \{0, 1\} \text{ and } (q+1) \mid k_1, (q+1) \nmid k_2, k_3, k_4.$$

Thus $c_0 + c_1 = 0$. Similarly, from $c(1) = 0$, we have $c_2 + c_3 + c_4 = 0$. From $c(\alpha^3) = c(\alpha^6) = 0$, we deduce that

$$\begin{aligned} c_2\alpha^{3k_2} + c_3\alpha^{3k_3} + c_4\alpha^{3k_4} &= 0, \\ c_2\alpha^{6k_2} + c_3\alpha^{6k_3} + c_4\alpha^{6k_4} &= 0. \end{aligned}$$

Thus

$$\det \begin{pmatrix} 1 & 1 & 1 \\ \alpha^{3k_2} & \alpha^{3k_3} & \alpha^{3k_4} \\ \alpha^{6k_2} & \alpha^{6k_3} & \alpha^{6k_4} \end{pmatrix} = 0,$$

which implies that $\alpha^{3k_2} = \alpha^{3k_3}$, or $\alpha^{3k_2} = \alpha^{3k_4}$, or $\alpha^{3k_3} = \alpha^{3k_4}$. Without loss of generality, we assume that $\alpha^{3k_2} = \alpha^{3k_3}$. We next prove that $\alpha^{3k_2} = \alpha^{3k_3} = \alpha^{3k_4}$, otherwise, suppose that $\alpha^{3k_2} = \alpha^{3k_3} \neq \alpha^{3k_4}$. Then for $\ell = 1, 2, 3$,

$$\sum_{i=0}^q \alpha^{3i(k_\ell - k_4)} = 0.$$

From $c(\alpha^{3i}) = 0$ ($i = 0, 1, 2, \dots, q$), we have

$$\begin{aligned} 0 &= \sum_{i=0}^q \alpha^{-3ik_4} c(\alpha^{3i}) \\ &= \sum_{i=0}^q \alpha^{-3ik_4} \sum_{\ell=0}^4 c_\ell \alpha^{3ik_\ell} \\ &= \sum_{\ell=0}^4 c_\ell \sum_{i=0}^q \alpha^{3i(k_\ell - k_4)} \\ &= c_4, \end{aligned}$$

which is a contradiction. So $\alpha^{3k_2} = \alpha^{3k_3} = \alpha^{3k_4} \Rightarrow (q+1) \mid (k_2 - k_4)$ and $(q+1) \mid (k_3 - k_4)$. From $c(\alpha) = 0$, we have

$$c_0 + c_1\alpha^{k_1} + c_2\alpha^{k_2} + c_3\alpha^{k_3} + c_4\alpha^{k_4} = 0.$$

Since $\alpha \in \mathbb{F}_{q^2}$ and $(q+1) \mid k_1$, we have $c_0 + c_1\alpha^{k_1} \in \mathbb{F}_q$. Since $c_0 + c_1 = 0$, $c_0 + c_1\alpha^{k_1} = c_0(1 - \alpha^{k_1}) \in \mathbb{F}_q^*$. Thus

$$c_2\alpha^{k_2} + c_3\alpha^{k_3} + c_4\alpha^{k_4} \in \mathbb{F}_q^*,$$

i.e.,

$$c_4\alpha^{k_4}\left(1 + \frac{c_3}{c_4}\alpha^{k_3-k_4} + \frac{c_2}{c_4}\alpha^{k_2-k_4}\right) \in \mathbb{F}_q^*.$$

Since $(q+1) \mid (k_2 - k_4)$ and $(q+1) \mid (k_3 - k_4)$, we have $\alpha^{k_3-k_4}, \alpha^{k_2-k_4} \in \mathbb{F}_q$, hence $\alpha^{k_4} \in \mathbb{F}_q^*$. Then $n \mid k_4(q-1)$. Note that $\gcd(\frac{n}{3}, q-1) = \gcd(q+1, q-1) = 1$, thus $(q+1) \mid k_4$, which is a contradiction.

In summary, $d = 6$ and hence C is a q -ary cyclic Singleton-optimal LRC with length $n = 3(q+1)$, minimum distance $d = 6$ and locality $r = 2$. \square

Remark 1. *To the best of our knowledge, Theorem 1 is the first construction of q -ary cyclic Singleton-optimal LRCs with length $n > q+1$ and minimum distance $d \geq 5$.*

Example 1. *Suppose $q = 4$ and $n = 3(q+1) = 15$. Let ω be a primitive element of \mathbb{F}_4 and $\alpha \in \mathbb{F}_{16}$ which is a root of the primitive polynomial $x^2 + x + \omega \in \mathbb{F}_4[x]$. Then $\alpha + \alpha^4 = 1$ and $\alpha^5 = \omega$. Let*

$$\begin{aligned} g(x) &= (x^5 - 1)(x - \alpha)(x - \alpha^4)(x - \alpha^5) \\ &= (x^5 - 1)(x^2 - x + \omega)(x - \omega) \in \mathbb{F}_4[x] \end{aligned}$$

and C be the cyclic code of length 15 with the generator polynomial $g(x)$. With the help of Magma, we can calculate that $d = 6$, which coincides with Theorem 1.

The condition of q is even is necessary in Theorem 1. Indeed, we show that when q is odd, the minimum distance of the LRCs constructed in Theorem 1 is equal to 5.

Proposition 3. *Suppose q is odd and $3 \mid (q-1)$. Let C be the cyclic code of length $n = 3(q+1)$ constructed as in Theorem 1. Then the minimum distance of C is 5.*

Proof. From the proof of Theorem 1, the generator polynomial of C is

$$g(x) = (x^{q+1} - 1)(x - \alpha)(x - \alpha^q)(x - \alpha^{q+1}),$$

where $\alpha \in \mathbb{F}_{q^2}$ is a primitive n -th root of unity. Denote $\pi = \alpha^{\frac{q+1}{2}} \in \mathbb{F}_q$. Then $\pi^3 = -1$ and $\pi^2 - \pi + 1 = 0$. Let

$$c(x) = 1 - 2x^{q+1} + x^{2(q+1)} - \frac{3}{2\pi - 1}x^{\frac{q+1}{2}} + \frac{3}{2\pi - 1}x^{\frac{3(q+1)}{2}} \in \mathbb{F}_q[x].$$

Then $wt(c(x)) = 5$. From α is a primitive n -th root of unity, for $i = 0, 1, \dots, q$, we have

$$c(\alpha^{3i}) = 1 - 2 + 1 - \frac{3}{2\pi - 1}(-1)^i + \frac{3}{2\pi - 1}(-1)^{3i} = 0.$$

Moreover,

$$\begin{aligned}
c(\alpha) &= 1 - 2\pi^2 + \pi^4 - \frac{3}{2\pi-1}\pi + \frac{3}{2\pi-1}\pi^3 \\
&= 1 - 2(\pi-1) - \pi - \frac{3(\pi+1)}{2\pi-1} \\
&= \frac{3((1-\pi)(2\pi-1) - (\pi+1))}{2\pi-1} \\
&= \frac{-6(\pi^2 - \pi + 1)}{2\pi-1} = 0, \\
c(\alpha^q) &= (c(\alpha))^q = 0,
\end{aligned}$$

and from $3 \mid (q-1)$, we have

$$\begin{aligned}
c(\alpha^{q+1}) = c(\pi^2) &= 1 - 2\pi^{2(q+1)} + \pi^{4(q+1)} - \frac{3}{2\pi-1}\pi^{q+1} + \frac{3}{2\pi-1}\pi^{3(q+1)} \\
&= 1 + 2\pi + \pi^2 - \frac{3}{2\pi-1}\pi^2 + \frac{3}{2\pi-1} \\
&= 3\pi - \frac{3\pi-6}{2\pi-1} = \frac{3\pi(2\pi-1) - (3\pi-6)}{2\pi-1} \\
&= \frac{6(\pi^2 - \pi + 1)}{2\pi-1} = 0.
\end{aligned}$$

Thus $g(x) \mid c(x)$ and hence $c(x)$ is a nonzero codeword of C , which implies that $d \leq 5$. Since $\alpha^{q-1}, \alpha^q, \alpha^{q+1}, \alpha^{q+2}$ are roots of $g(x)$, by the BCH bound, $d \geq 5$. Thus $d = 5$. The proof is completed. \square

Example 2. Suppose $q = 7$ and $n = 3(q+1) = 24$. Let $\alpha \in \mathbb{F}_{49}$ be a primitive 24-th root of unity. Let $g(x) = (x^8-1)(x-\alpha)(x-\alpha^7)(x-\alpha^8) = x^{11} + 3x^{10} + 6x^9 + 3x^8 + 6x^3 + 4x^2 + x + 4 \in \mathbb{F}_7[x]$ and C be the cyclic code of length 24 with generator polynomial $g(x)$. With the help of Magma, we can calculate that $d = 5$, which coincides with Proposition 3.

3.2 Cyclic Singleton-optimal LRCs for $q \equiv 1 \pmod{4}$

In Subsection 3.1, we give a construction of q -ary cyclic Singleton-optimal LRCs of length $n = 3(q+1)$ for q is even. And we also have proved that this construction is not optimal for q is odd in Proposition 3. In this subsection, we construct a family q -ary Singleton-optimal LRCs of length $n = \frac{3(q+1)}{2}$ for $q \equiv 1 \pmod{4}$ and $3 \mid (q-1)$.

Theorem 2. Suppose $q \equiv 1 \pmod{4}$ and $3 \mid (q-1)$. Let $n = \frac{3(q+1)}{2}$ and $\alpha \in \mathbb{F}_{q^2}$ be a primitive n -th root of unity. Let

$$g(x) = (x^{\frac{n}{3}} - 1)(x - \alpha)(x - \alpha^q)(x - \alpha^{q+1}).$$

Then the cyclic code $C \triangleq \langle g(x) \rangle$ is a q -ary cyclic Singleton-optimal LRC with length n , minimum distance $d = 6$ and locality $r = 2$.

Proof. From $(x^{\frac{n}{3}} - 1) \mid g(x)$, we know that $1, \alpha^3, \alpha^6, \dots, \alpha^{n-3}$ are roots of $g(x)$. By Proposition 2, C has locality $r = 2$. Since $3 \mid (q - 1)$ and $\alpha^{q-1}, \alpha^q, \alpha^{q+1}, \alpha^{q+2}$ are roots of $g(x)$, we have $d \geq 5$ by Lemma 3. The dimension k of C is equal to $n - \deg(g(x)) = q - 2$. By the Singleton-type bound,

$$d \leq \frac{3(q+1)}{2} - (q-2) + 2 - \lceil \frac{q-2}{2} \rceil = 6.$$

Thus we only need to prove that $d \neq 5$. By contradiction, let

$$c(x) = c_0x^{k_0} + c_1x^{k_1} + c_2x^{k_2} + c_3x^{k_3} + c_4x^{k_4}$$

be a codeword of C with $c_0, c_1, c_2, c_3, c_4 \in \mathbb{F}_q^*$ and $0 = k_0 < k_1 < k_2 < k_3 < k_4 \leq n - 1$. Let

$$I = \{\ell : 0 \leq \ell \leq 4 \text{ and } \frac{n}{3} \mid k_\ell\}.$$

Then $1 \leq |I| \leq 3$. Similar to the proof of Theorem 1, we have

$$\sum_{\ell \in I} c_\ell = 0,$$

and hence $|I| \geq 2$.

Case 1: $|I| = 3$. Without loss of generality, we suppose

$$I = \{0, 1, 2\}, k_1 = \frac{n}{3}, k_2 = \frac{2n}{3} \text{ and } \frac{n}{3} \nmid k_3, k_4.$$

Thus $c_0 + c_1 + c_2 = 0$. By $c(1) = 0$, we have

$$c_0 + c_1 + c_2 + c_3 + c_4 = 0 \Rightarrow c_3 + c_4 = 0.$$

By $c(\alpha^3) = 0$, we have

$$\begin{aligned} c_0 + c_1(\alpha^3)^{\frac{n}{3}} + c_2(\alpha^3)^{\frac{2n}{3}} + c_3(\alpha^3)^{k_3} + c_4(\alpha^3)^{k_4} &= 0, \\ \Rightarrow c_3\alpha^{3k_3} + c_4\alpha^{3k_4} = 0 &\Rightarrow \alpha^{3k_3} = \alpha^{3k_4} \Rightarrow \frac{n}{3} \mid (k_4 - k_3). \end{aligned}$$

By $c(\alpha) = 0$, we have

$$c_0 + c_1\alpha^{\frac{n}{3}} + c_2\alpha^{\frac{2n}{3}} + c_3\alpha^{k_3} + c_4\alpha^{k_4} = 0.$$

Note that $(\alpha^{\frac{n}{3}})^{q-1} = 1$, thus $\alpha^{\frac{n}{3}} \in \mathbb{F}_q$.

$$\Rightarrow c_3\alpha^{k_3} + c_4\alpha^{k_4} \in \mathbb{F}_q \Rightarrow c_3\alpha^{k_3}(1 - \alpha^{k_4-k_3}) \in \mathbb{F}_q.$$

Since $\frac{n}{3} \mid (k_4 - k_3)$ and $n \nmid (k_4 - k_3)$, we have $\alpha^{k_4-k_3} \neq 1$ and $\alpha^{k_4-k_3} - 1 \in \mathbb{F}_q$. Thus

$$\alpha^{k_3} \in \mathbb{F}_q \Rightarrow n \mid k_3(q-1).$$

Note that $\gcd(\frac{n}{3}, q-1) = \gcd(\frac{q+1}{2}, q-1) = 1$ by $q \equiv 1 \pmod{4}$. Thus $\frac{n}{3} \mid k_3$, which is a contradiction.

Case 2: $|I| = 2$. The proof of this case is completely similar to the above case and the proof of Theorem 1, so we omit it.

The theorem is proved. □

Remark 2. Suppose q is odd and $3 \mid (q-1)$. If $q+1 = 2^a \cdot b$, where b is an odd, then let $n = \frac{3(q+1)}{2^a} = 3b$ and $\alpha \in \mathbb{F}_{q^2}$ be a primitive n -th root of unity. Let

$$g(x) = (x^{\frac{n}{3}} - 1)(x - \alpha)(x - \alpha^q)(x - \alpha^{q+1}).$$

Similar to the proof of Theorem 2, we can show that $C \triangleq \langle g(x) \rangle$ is a q -ary cyclic Singleton-optimal LRC with length n , minimum distance $d = 6$ and locality $r = 2$.

Example 3. Suppose $q = 13$ and $n = 3(q+1)/2 = 21$. Let $\alpha \in \mathbb{F}_{13^2}$ be a primitive 21-th root of unity. Let $g(x) = (x^7 - 1)(x - \alpha)(x - \alpha^{13})(x - \alpha^{14}) = x^{10} + 9x^9 + 3x^8 + 10x^7 + 12x^3 + 4x^2 + 10x + 3 \in \mathbb{F}_{13}[x]$ and C be the cyclic code of length 21 with generator polynomial $g(x)$. With the help of Magma, we can calculate that $d = 6$, which coincides with Proposition 3.

4 Cyclic and Constacyclic Perfect LRCs

In [10], we first introduced the definition of perfect LRCs. By using the techniques of finite geometry and finite fields, we proposed two general constructions of perfect LRCs in [10]. In this section, we will give two constructions of perfect LRCs via cyclic and constacyclic codes.

4.1 Cyclic Perfect LRCs

Before give our construction of cyclic perfect LRCs, we need an auxiliary lemma.

Lemma 5 ([27]). Let $a \in \mathbb{F}_q$ and p be the characteristic of \mathbb{F}_q . Then the trinomial $x^p - x - a$ is irreducible in $\mathbb{F}_q[x]$ if and only if $\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a) \neq 0$.

Suppose $q \geq 4$ is even with $3 \mid (q+1)$, and m is even with $\gcd(m, q-1) = 1$. Let $n = \frac{q^m-1}{q-1}$, then $n = q^{m-1} + q^{m-2} + \dots + 1 \equiv 0 \pmod{q+1}$. Now, our construction of cyclic perfect LRCs is presented as follows.

Theorem 3. Suppose $q \geq 4$ is even with $3 \mid (q+1)$, and m is even with $\gcd(m, q-1) = 1$. Let $\alpha \in \mathbb{F}_{q^m}$ be a primitive n -th root of unity. Set

$$L = \{1, \alpha^3, \alpha^6, \dots, \alpha^{n-3}\}$$

and

$$D = \{\alpha, \alpha^q, \dots, \alpha^{q^{m-1}}\}.$$

Let $g(x) = \prod_{a \in L \cup D} (x - a)$ and $C = \langle g(x) \rangle$. Then C is a q -ary cyclic ($n = \frac{q^m-1}{q-1}, k = \frac{2}{3}n - m, d = 5; r = 2$)-perfect LRC.

Proof. The elements of L and D are roots of $x^n - 1$, and

$$\prod_{a \in L} (x - a) = \prod_{i=0}^{\frac{n}{3}-1} (x - \alpha^{3i}) = x^{\frac{n}{3}} - 1.$$

Then

$$g(x) = \prod_{a \in L \cup D} (x - a) = (x^{\frac{n}{3}} - 1)(x - \alpha)(x - \alpha^q) \cdots (x - \alpha^{q^{m-1}}).$$

Thus $g(x) \in \mathbb{F}_q[x]$ and $g(x) \mid (x^n - 1)$. So C is a cyclic code over \mathbb{F}_q of length n and the dimension of C is $k = n - \deg(g(x)) = \frac{2}{3}n - m$.

Note that $\alpha^0, \alpha^1, \alpha^q, \alpha^{q+1}$ are roots of $g(x)$ and $\gcd(q, n) = 1$. By the Hartmann-Tzeng bound (let $\lambda = 1, \delta = 3, \gamma = 1, u = 0, b_1 = 1, b_2 = q$ in Lemma 4), we have $d \geq 3 + 1 = 4$. Next we prove that $d \neq 4$. Otherwise, suppose there exists a codeword

$$c(x) = c_0 + c_1x^{k_1} + c_2x^{k_2} + c_3x^{k_3} \in C$$

with $c_0, c_1, c_2, c_3 \in \mathbb{F}_q^*$ and $0 = k_0 < k_1, k_2, k_3 \leq n - 1$, where k_1, k_2, k_3 are pairwise distinct positive integers. Then $g(x) \mid c(x)$. Let

$$I = \{\ell : 0 \leq \ell \leq 3 \text{ and } \frac{n}{3} \mid k_\ell\}.$$

Then $0 \in I$. Since $k_1, k_2, k_3 \leq n - 1$, $|I| \leq 3$. Note that

$$\sum_{i=0}^{\frac{n}{3}-1} (\alpha^{3i})^{k_\ell} = \begin{cases} 1, & \text{for } \ell \in I, \\ 0, & \text{for } \ell \notin I. \end{cases}$$

From $c(\alpha^{3i}) = 0$, $i = 0, 1, \dots, \frac{n}{3} - 1$, we have

$$\begin{aligned} 0 &= \sum_{i=0}^{\frac{n}{3}-1} c(\alpha^{3i}) = \sum_{i=0}^{\frac{n}{3}-1} \sum_{\ell=0}^3 c_\ell (\alpha^{3i})^{k_\ell} \\ &= \sum_{\ell=0}^3 c_\ell \sum_{i=0}^{\frac{n}{3}-1} (\alpha^{3i})^{k_\ell} = \sum_{\ell \in I} c_\ell, \end{aligned}$$

which implies that $|I| \geq 2$.

Case 1: $|I| = 3$. Without loss of generality, we suppose

$$I = \{0, 1, 2\}, k_1 = \frac{n}{3} \text{ and } k_2 = \frac{2n}{3}.$$

Thus $c_0 + c_1 + c_2 = 0$. By $c(1) = 0$, we have $c_0 + c_1 + c_2 + c_3 = 0 \Rightarrow c_3 = 0$, which is a contradiction.

Case 2: $|I| = 2$. Without loss of generality, we suppose

$$I = \{0, 1\}, \frac{n}{3} \mid k_1 \text{ and } \frac{n}{3} \nmid k_2, k_3.$$

Thus $c_0 + c_1 = 0$. Similarly, from $c(1) = 0$, we have $c_2 + c_3 = 0$. From $c(\alpha^3) = 0$, we have

$$\begin{aligned} c_0 + c_1 + c_2\alpha^{3k_2} + c_3\alpha^{3k_3} &= 0 \\ \Rightarrow \alpha^{3k_2} = \alpha^{3k_3} &\Rightarrow \frac{n}{3} \mid (k_3 - k_2). \end{aligned}$$

From $c(\alpha) = 0$, we have $c_0 + c_1\alpha^{k_1} + c_2\alpha^{k_2} + c_3\alpha^{k_3} = 0 \Rightarrow c_0(1 + \alpha^{k_1}) + c_2\alpha^{k_2}(1 + \alpha^{k_3-k_2}) = 0 \Rightarrow \frac{\alpha^{k_2}(1 + \alpha^{k_3-k_2})}{(1 + \alpha^{k_1})} = \frac{c_0}{c_2} \in \mathbb{F}_q \Rightarrow$

$$\left(\frac{\alpha^{k_2}(1 + \alpha^{k_3-k_2})}{(1 + \alpha^{k_1})} \right)^q = \frac{\alpha^{k_2}(1 + \alpha^{k_3-k_2})}{(1 + \alpha^{k_1})}.$$

Since $3 \mid (q+1)$ and $\frac{n}{3} \mid k_1, (k_3 - k_2)$, we have $\alpha^{(q+1)k_1} = \alpha^{(q+1)(k_3-k_2)} = 1$. Thus

$$\left(\frac{\alpha^{k_2}(1 + \alpha^{k_3-k_2})}{(1 + \alpha^{k_1})} \right)^q = \frac{\alpha^{qk_2}(1 + \alpha^{-(k_3-k_2)})}{(1 + \alpha^{-k_1})} = \frac{\alpha^{qk_2+k_1}(1 + \alpha^{k_3-k_2})}{\alpha^{k_3-k_2}(1 + \alpha^{k_1})}.$$

Then $\alpha^{k_2} = \frac{\alpha^{qk_2+k_1}}{\alpha^{k_3-k_2}} \Rightarrow n \mid ((q-1)k_2 + k_1 + k_3 - k_2) \Rightarrow \frac{n}{3} \mid (q-1)k_2$. Note that $\gcd(n, q-1) = \gcd(m, q-1) = 1$, thus $\frac{n}{3} \mid k_2$, which is a contradiction. Therefore $d \geq 5$.

According to the set L , C has locality $r = 2$ by Proposition 2. Note that $\frac{q^{\frac{r}{2}+1}n}{\frac{r}{2}(q-1)+1} = \frac{q^{\frac{2}{3}n}}{q^m} = q^k$, which achieves the bound (4) with equality. Thus we only need to show that $d(C) = 5$, i.e., find a codeword of C of Hamming weight 5.

Denote $\omega = \alpha^{\frac{n}{3}}$ and $\beta = \alpha^{\frac{n}{q+1}}$. Then $\omega^3 = 1, \beta^{q+1} = 1 \Rightarrow \omega^2 = \omega + 1$ and $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Thus there exist $a, b \in \mathbb{F}_q^*$ such that

$$\beta^2 + a\beta + b = 0, \quad (6)$$

i.e., $(\frac{\beta}{a})^2 + \frac{\beta}{a} + \frac{b}{a^2} = 0$. Thus $x^2 + x + \frac{b}{a^2}$ is irreducible. By Lemma 5, $\text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\frac{b}{a^2}) = 1$. Since $3 \mid (q+1)$, we can derive that $q = 2^\ell$ for some odd ℓ . Thus

$$\text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(1 + \frac{b}{a^2}) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(1) + \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\frac{b}{a^2}) = 1 + 1 = 0.$$

By Lemma 5 again, there exists $c_0 \in \mathbb{F}_q^*$ such that $(\frac{c_0}{a})^2 + \frac{c_0}{a} + 1 + \frac{b}{a^2} = 0$, i.e.,

$$c_0^2 + ac_0 + a^2 + b = 0. \quad (7)$$

Note that

$$\begin{aligned} (\beta + c_0)^3 &= \beta^3 + \beta^2c_0 + \beta c_0^2 + c_0^3 \\ &= \beta(a\beta + b) + (a\beta + b)c_0 + \beta c_0^2 + c_0^3 \\ &= a(a\beta + b) + b\beta + (a\beta + b)c_0 + \beta c_0^2 + c_0^3 \\ &= (c_0^2 + ac_0 + a^2 + b)\beta + ab + bc_0 + c_0^3 \\ &= (a + c_0)b + c_0^3. \end{aligned}$$

Thus

$$(\beta + c_0)^3 + a^3 = (a + c_0)(b + c_0^2 + ac_0 + a^2) = 0,$$

i.e., $(\frac{a}{\beta+c_0})^3 = 1 \Rightarrow \frac{a}{\beta+c_0} = \omega$ or ω^2 . Without loss of generality, we suppose $\frac{a}{\beta+c_0} = \omega$ (the proof of the case of ω^2 is similar). We prove that $c_0 \neq a$. Otherwise, $c_0 = a$, then from Eq. (7), we have $b = a^2$. Then $\beta^2 + a\beta + a^2 = 0$ by Eq. (6). One can deduce that

$$\left(\frac{\beta}{a}\right)^3 = 1 \Rightarrow \beta^{3(q-1)} = 1 \Rightarrow (q+1) \mid 3(q-1) \Rightarrow (q+1) \mid 3,$$

which contradicts with $q \geq 4$. Now let

$$c(x) = c_0 + ax^{\frac{n}{3}} + (c_0 + a)x^{\frac{2n}{3}} + x^{\frac{n}{q+1}} + x^{\frac{n}{q+1} + \frac{2n}{3}}$$

which is a polynomial over \mathbb{F}_q of weight 5. Note that for $i = 0, 1, \dots, \frac{n}{3} - 1$,

$$c(\alpha^{3i}) = c_0 + a + (c_0 + a) + \alpha^{\frac{3in}{q+1}} + \alpha^{\frac{3in}{q+1} + 2in} = 0.$$

And

$$\begin{aligned} c(\alpha) &= c_0 + a\omega + (c_0 + a)\omega^2 + \beta + \beta\omega^2 \\ &= c_0 + a\omega + (c_0 + a)(1 + \omega) + \beta + \beta(1 + \omega) \\ &= a + c_0\omega + \beta\omega = 0. \end{aligned}$$

Thus $g(x) \mid c(x)$, i.e, $c(x)$ is a codeword of C of weight 5. Thus $d = 5$.

The Theorem is proved. \square

Example 4. Suppose $q = 8$, $m = 4$, $n = \frac{q^m - 1}{q - 1} = 585$ and $\alpha \in \mathbb{F}_{8^4}$ is a primitive element. Let $g(x) = (x^{195} - 1)(x - \alpha)(x - \alpha^8)(x - \alpha^{64})(x - \alpha^{256}) \in \mathbb{F}_8[x]$ and $C = \langle g(x) \rangle$. By the Magma software, we can calculate that $d(C) = 5$, which coincides with Theorem 3.

4.2 Constacyclic Perfect LRCs

For q is even, we have constructed a family of q -ary cyclic perfect LRCs in Subsection 4.1. For general q , we consider the construction of q -ary perfect LRCs via constacyclic codes. Suppose $3 \mid (q + 1)$ and m is even. Let $n = \frac{q^m - 1}{q - 1}$. Then it can be shown that $3 \mid n$. Let π be a primitive element of \mathbb{F}_{q^m} ,

$$\theta = \pi^{-q+2} \text{ and } \lambda = \theta^n.$$

Then $\lambda^{q-1} = \theta^{q^m-1} = 1$, hence $\lambda \in \mathbb{F}_q^*$. Let

$$\alpha = \pi^{q-1},$$

then α is a primitive n -th root of unity and

$$x^n - \lambda = x^n - \theta^n = \prod_{i=0}^{n-1} (x - \theta\alpha^i)$$

Theorem 4. Suppose $q > 2$, $3 \mid (q + 1)$ and m is even. Let $n = \frac{q^m - 1}{q - 1}$. Denote

$$L = \{\theta, \theta\alpha^3, \theta\alpha^6, \dots, \theta\alpha^{n-3}\},$$

$$D = \{\theta\alpha, (\theta\alpha)^q, \dots, (\theta\alpha)^{q^{m-1}}\}.$$

Let $g(x) = \prod_{a \in L \cup D} (x - a)$ and $C = \langle g(x) \rangle$. Then C is a λ -constacyclic $(n = \frac{q^m - 1}{q - 1}, k = \frac{2}{3}n - m, d = 5; r = 2)$ -perfect LRC.

Proof. From the above, we know that the elements of L and D are roots of $x^n - \lambda$, and

$$\prod_{a \in L} (x - a) = \prod_{i=0}^{\frac{n}{3}-1} (x - \theta\alpha^{3i}) = x^{\frac{n}{3}} - \theta^{\frac{n}{3}}.$$

Thus

$$g(x) = \prod_{a \in L \cup D} (x - a) = (x^{\frac{n}{3}} - \theta^{\frac{n}{3}})(x - \theta\alpha)(x - (\theta\alpha)^q) \cdots (x - (\theta\alpha)^{q^{m-1}}).$$

Since $3 \mid (q+1)$, thus $3 \mid (-q+2)$ and $(\theta^{\frac{n}{3}})^{q-1} = (\pi^{\frac{-q+2}{3}})^{q^{m-1}} = 1$, hence $\theta^{\frac{n}{3}} \in \mathbb{F}_q^*$. Note that

$$\theta\alpha = \pi^{-q+2}\pi^{q-1} = \pi.$$

Then $g(x) \in \mathbb{F}_q[x]$ and $g(x) \mid (x^n - \lambda)$. Hence $\dim(C) = \frac{2}{3}n - \deg(g(x)) = \frac{2}{3}n - m$. Note that

$$(\theta\alpha)^q = \pi^q = \pi^{-q+2}\pi^{2(q-1)} = \theta\alpha^2.$$

Thus $\theta, \theta\alpha, \theta\alpha^2, \theta\alpha^3$ are roots of $g(x)$. By the BCH bound for constacyclic codes (see Lemma 3), the minimum distance $d(C) \geq 5$. According to the set L , C has locality $r = 2$ by Proposition 2. Note that $\frac{q^{\frac{r+1}{2}n}}{\frac{r}{2}(q-1)+1} = \frac{q^{\frac{3}{2}n}}{q^m} = q^k$, which achieves the bound (4) with equality. Thus we only need to show that $d(C) = 5$, i.e., find a codeword of C of Hamming weight 5.

Note that $q > 2$ and $3 \mid (q+1)$, thus $q \geq 5$. We can choose $a \in \mathbb{F}_q^*$, such that $a \neq -1, 2$ and $2a - 1 \neq 0$ (for example, $a = 1$ if q is odd; $a \neq 0, -1$, if q is even). Let $\omega = \alpha^{\frac{n}{3}}$, then $\omega^3 = 1$. Choose $\bar{k}_3 \in \mathbb{Z}$, such that

$$\pi^{\bar{k}_3} = \frac{a + \omega}{1 - \omega}.$$

Let $\bar{k}_3 = sn + k_3$, where $0 \leq k_3 < n$. We next prove that $\frac{n}{3} \nmid \bar{k}_3$. Otherwise, we have $(\pi^{\bar{k}_3})^{3(q-1)} = 1$. Then

$$\begin{aligned} \left(\frac{a + \omega}{1 - \omega}\right)^{3q} &= \left(\frac{a + \omega}{1 - \omega}\right)^3 \\ \Rightarrow \left(\frac{a + \omega^q}{1 - \omega^q}\right)^3 &= \left(\frac{a + \omega}{1 - \omega}\right)^3. \end{aligned}$$

Since $3 \mid (q+1)$, $\omega^q = \omega^{-1}$. Thus

$$\begin{aligned} \left(\frac{a\omega + 1}{\omega - 1}\right)^3 &= \left(\frac{a + \omega}{1 - \omega}\right)^3 \\ \Rightarrow (a\omega + 1)^3 + (a + \omega)^3 &= 0, \\ \Rightarrow \omega(\omega + 1)(a + 1)(a - 2)(2a - 1) &= 0, \\ \Rightarrow a = -1 \text{ or } 2, \text{ or } 2a - 1 = 0, \end{aligned}$$

which is a contradiction. Thus $\frac{n}{3} \nmid k_3$, hence $k_3 \neq 0, \frac{n}{3}, \frac{2}{3}n$. Let

$$k_4 = k_3 + \frac{n}{3} \pmod{n} \text{ with } 0 \leq k_4 < n.$$

Then $k_4 \neq 0, \frac{n}{3}, \frac{2}{3}n$ and $\alpha^{k_4-k_3} = \alpha^{\frac{n}{3}} = \omega \Rightarrow \alpha^{3k_4} = \alpha^{3k_3}$. Let $b = \pi^{sn} \in \mathbb{F}_q^*$, then $\pi^{k_3} = \frac{a+\omega}{b(1-\omega)}$. Now, let

$$c(x) = 1 + a\theta^{-\frac{n}{3}}x^{\frac{n}{3}} - (1+a)\theta^{-\frac{2}{3}n}x^{\frac{2}{3}n} + bx^{k_3} - b\theta^{k_3-k_4}x^{k_4} \in \mathbb{F}_q[x].$$

Then $wt(c(x)) = 5$, for $i = 0, 1, \dots, \frac{n}{3} - 1$, we have

$$c(\theta\alpha^{3i}) = 1 + a - (1+a) + b\theta^{k_3}\alpha^{3ik_3} - b\theta^{k_3-k_4}\theta^{k_4}\alpha^{3ik_4} = 0,$$

and

$$\begin{aligned} c(\theta\alpha) &= 1 + a\alpha^{\frac{n}{3}} - (1+a)\alpha^{\frac{2}{3}n} + b\pi^{k_3} - b\pi^{k_3}\alpha^{k_4-k_3} \\ &= 1 + a\omega - (1+a)\omega^2 + b\pi^{k_3}(1-\omega) \\ &= -(a+\omega) + b\pi^{k_3}(1-\omega) = 0. \end{aligned}$$

Hence $g(x) \mid c(x)$, i.e., $c(x)$ is a codeword of C of Hamming weight 5. \square

Example 5. Suppose $q = 5, m = 6, n = \frac{q^m-1}{q-1} = 3906$ and π is a primitive element of \mathbb{F}_{5^6} . Let $\theta = \pi^{-3}, \lambda = \theta^n = 3 \in \mathbb{F}_5$ and $\alpha = \pi^4$. Let $g(x) = (x^{1302} - \theta\alpha)(x - (\theta\alpha))(x - (\theta\alpha)^5)(x - (\theta\alpha)^{25})(x - (\theta\alpha)^{125})(x - (\theta\alpha)^{625}) = x^{1308} + x^{1306} + 4x^{1305} + x^{1304} + 2x^{1302} + 2x^6 + 2x^4 + 3x^3 + 2x^2 + 4 \in \mathbb{F}_5[x]$ and $C = \langle g(x) \rangle$. With the help of Magma, we can calculate that $d(C) = 5$, which coincide with Theorem 4.

5 Conclusion

In this paper, we consider new constructions of Singleton-optimal LRCs and perfect LRCs via cyclic and constacyclic codes. Firstly, we obtain two classes of q -ary cyclic $(n, k, d = 6; r = 2)$ -Singleton-optimal LRCs with $n = 3(q+1)$ and $n = \frac{3(q+1)}{2}$, respectively. To the best of our knowledge, this is the first construction of q -ary Singleton-optimal LRCs of length $n > q+1$ and minimum distance $d \geq 5$ via cyclic codes. Secondly, by using cyclic and constacyclic codes, we give two constructions of perfect LRCs. Cyclic and constacyclic codes have nice algebraic structures and efficient encoding and decoding algorithm. Thus it is interesting to find more new constructions of Singleton-optimal LRCs and perfect LRCs with longer code length via cyclic and constacyclic codes, and investigate the theoretical bounds for cyclic and constacyclic LRCs in the future.

Acknowledgements

This research is supported in part by National Key Research and Development Program of China under Grant Nos. 2022YFA1004900, 2021YFA1001000, 2018YFA0704703 and

2022YFA1005001, the National Natural Science Foundation of China under Grant Nos. 62201322, 62171248, 12141108, 61971243 and 12226336, the Natural Science Foundation of Shandong (ZR2022QA031), the Natural Science Foundation of Tianjin (20JCZDJC00610), the Fundamental Research Funds for the Central Universities, Nankai University, and the Nankai Zhide Foundation.

References

- [1] A. Agarwal, A. Barg, S. Hu, A. Mazumdar, and I. Tamo, Combinatorial alphabet-dependent bounds for locally recoverable codes, *IEEE Trans. Inf. Theory* 64(5) (2018) 3481-3492.
- [2] N. Aydin, I. Siap, D.J. Ray-Chaudhuri, The structure of 1-generator quasi-twisted codes and new linear codes, *Des. Codes Cryptogr.* 24(3) (2001) 313–326.
- [3] T. Blackford, Negacyclic duadic codes, *Finite Fields Appl.* 14(4) (2008) 930–943.
- [4] V. R. Cadambe and A. Mazumdar, Bounds on the size of locally recoverable codes, *IEEE Trans. Inf. Theory* 61(11) (2015) 5787-5794.
- [5] B. Chen, Y. Fan, L. Lin and H. Liu, Constacyclic codes over finite fields, *Finite Fields Appl.* 18(6) (2012) 1217– 1231.
- [6] B. Chen, W. Fang, S.-T. Xia, and F.-W. Fu, Constructions of optimal (r, δ) locally repairable codes via constacyclic codes, *IEEE Trans. Commun.* 67(8) (2019) 5253-5263.
- [7] B. Chen, W. Fang, S.-T. Xia, J. Hao and F.-W. Fu, Improved bounds and Singleton-optimal constructions of locally repairable codes with minimum distance 5 and 6, *IEEE Trans. Inf. Theory* 67(1) (2021) 217-231.
- [8] B. Chen, S.-T. Xia, J. Hao, and F.-W. Fu, Constructions of optimal cyclic (r, δ) locally repairable codes, *IEEE Trans. Inf. Theory* 64(4) (2018) 2499-2511.
- [9] H.Q. Dinh, Constacyclic codes of length p^s over $\mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$, *J. Algebra*, 324 (2010) 940-950.
- [10] W. Fang, B. Chen, S.-T. Xia and F.-W. Fu, Perfect LRCs and k -optimal LRCs, in *Proc. Int. Symp. Inf. Theory (ISIT)* (2020) 600-604.

- [11] W. Fang, B. Chen, S.-T. Xia and F.-W. Fu, Singleton-optimal LRCs and perfect LRCs via cyclic Codes, in Proc. Int. Symp. Inf. Theory (ISIT) (2021) 3261-3266.
- [12] W. Fang, B. Chen, S.-T. Xia, F.-W. Fu and X. Chen, Perfect LRCs and k-optimal LRCs. Des. Codes Cryptogr. (2022). <https://doi.org/10.1007/s10623-022-01148-7>.
- [13] W. Fang, J. Wen and F.-W. Fu, A q -polynomial approach to constacyclic codes, Finite Fields Appl. 47 (2017) 161–182.
- [14] P. Gopalan, C. Huang, H. Simitci and S. Yekhanin, On the locality of codeword symbols, IEEE Trans. Inf. Theory 58(11) (2012) 6925-6934.
- [15] S. Goparaju and R. Calderbank, Binary cyclic codes that are locally repairable, in Proc. Int. Symp. Inf. Theory (ISIT) (2014) 676-680.
- [16] V. Guruswami, C. Xing and C. Yuan. How long can optimal locally repairable codes be ? IEEE Trans. Inf. Theory 65(6) (2019) 3662-3670.
- [17] J. Hao, S.-T. Xia and B. Chen, Some results on optimal locally repairable codes, in Proc. IEEE Int. Symp. Inf. Theory (ISIT) (2016) 440-444.
- [18] J. Hao, S.-T. Xia and B. Chen, On optimal ternary locally repairable codes, in Proc. IEEE Int. Symp. Inf. Theory (ISIT) (2017) 171-175.
- [19] J. Hao, S.-T. Xia, K.W. Shum, B. Chen, F.-W. Fu and Y. Yang, Bounds and constructions of locally repairable codes: Parity-check matrix approach, IEEE Trans. Inf. Theory 66(12) (2020) 7465-7474.
- [20] C.R.P. Hartmann and K.K. Tzeng, Generalizations of the BCH bound, Inform. and Control 20 (1972) 489–498.
- [21] C. Huang, H. Simitci, Y. Xu, A. Ogus, B. Calder, P. Gopalan, J. Li and S. Yekhanin, Erasure coding in windows azure storage, in: USENIX Annual Technical Conference (ATC) (2012) 15-26.
- [22] P. Huang, E. Yaakobi, H. Uchikawa and P. H. Siegel, Binary linear locally repairable codes, IEEE Trans. Inf. Theory 62(11) (2016) 6268-6283.
- [23] W.C. Huffman and V. Pless, *Fundamentals of Error-Correcting Codes*. Cambridge Univ. Press, UK, 2003.

- [24] L. Jin, Explicit construction of optimal locally recoverable codes of distance 5 and 6 via binary constant weight codes, *IEEE Trans. Inf. Theory* 65(8), (2019) 4658-4663.
- [25] L. Jin, L. Ma and C. Xing, Construction of optimal locally repairable codes via automorphism groups of rational function fields, *IEEE Trans. Inf. Theory* 66(1) (2020) 210-221.
- [26] X. Li, L. Ma and C. Xing, Optimal locally repairable codes via elliptic curves, *IEEE Trans. Inf. Theory* 65(1) (2019) 108-117.
- [27] R. Lidl and H. Niederreiter, *Finite Fields*. Cambridge, U.K.: Cambridge Univ. Press, 1997.
- [28] Y. Luo, C. Xing and C. Yuan, Optimal locally repairable codes of distance 3 and 4 via cyclic codes, *IEEE Trans. Inf. Theory* 65(2) (2019) 1048-1053.
- [29] F.J. MacWilliams and N.J.A. Sloane, *The Theory of Error-Correcting Codes*. Amsterdam, The Netherlands: North-Holland, 1977.
- [30] D. Radkova and A.J. van Zanten, Bounds for the minimum distance in constacyclic codes, in *Proceedings Eleventh International Workshop on Algebraic and Combinatorial Coding Theory*, Pamporovo, Bulgaria (2008) 236-242.
- [31] M. Sathiamoorthy, M. Asteris, D.S. Papailiopoulos, A.G. Dimakis, R. Vadali, S. Chen and D. Borthakur, XORing elephants: novel erasure codes for big data, in *Proceedings of VLDB Endowment (PVLDB)* (2013) 325-336.
- [32] W. Song, S. Day, C. Yuen and T. Li, Optimal locally repairable linear codes, *IEEE J. Selected Areas Comm.* 32 (2014) 6925-6934.
- [33] I. Tamo and A. Barg, A family of optimal locally recoverable codes, *IEEE Trans. Inf. Theory* 60(8) (2014) 4661-4676.
- [34] I. Tamo, A. Barg, S. Goparaju and R. Calderbank, Cyclic LRC codes and their subfield subcodes, in *Proc. IEEE Int. Symp. Inf. Theory (ISIT)* (2015) 1262-1266.
- [35] I. Tamo, D.S. Papailiopoulos and A.G. Dimakis, Optimal locally repairable codes and connections to matroid theory, *IEEE Trans. Inf. Theory* 62(12) (2016) 6661-6671.

- [36] A. Tebbi, T. Chan and C. Sung, Linear programming bounds for distributed storage codes, *Adv. in Math. of Commun.* 14(2) (2019) 333-357.
- [37] A. Wang, Z. Zhang and D. Lin, Bounds and constructions for linear locally repairable codes over binary fields, *IEEE Trans. Inf. Theory* 65(7) (2019) 4167-4179.
- [38] C. Xing and C. Yuan, Construction of (r, δ) locally recoverable codes and connection with graph theory, *IEEE Trans. Inf. Theory* 68(7) (2022) 4320 - 4328.
- [39] Y. Yang and W. Cai, On the self-dual constacyclic codes over the finite fields, *Des. Codes Cryptogr.* 74 (2015) 355–364.