

Estimate distillable entanglement and quantum capacity by squeezing useless entanglement

Chengkai Zhu,^{1,*} Chenghong Zhu,^{1,*} and Xin Wang^{1,†}

¹*Institute for Quantum Computing, Baidu Research, Beijing 100193, China*

(Dated: March 16, 2023)

Entanglement distillation is crucial in quantum information processing. But it remains challenging to estimate the distillable entanglement and its closely related essential quantity, the quantum capacity of a noisy quantum channel. In this work, we propose methods for evaluating both quantities by squeezing out useless entanglement within a state or a quantum channel, whose contributions are expected to be ignored for the distillable entanglement or the quantum capacity, respectively. We first consider a general resource measure called the reverse divergence of resources to quantify the minimum divergence between a target state and the set of free states. We then introduce the reverse max-relative entropy of entanglement and apply it to establish efficiently computable upper bounds on the distillable entanglement. We also extend the reverse divergence of resources to quantum channels and derive upper bounds on the quantum capacity. We further apply our method to investigate purifying the maximally entangled states under practical noises, such as depolarizing and amplitude damping noises, and notably establish improvements in estimating the one-way distillable entanglement. Our bounds also offer useful benchmarks for evaluating the quantum capacities of qubit quantum channels of interest, including the Pauli channels and the random mixed unitary channels.

CONTENTS

I. Introduction	2
A. Background	2
B. Main contributions	3
II. Reverse divergence of resources	3
A. Preliminaries	3
B. Reverse divergence of resources	4
III. Applications on distillable entanglement	6
A. Upper bound on the one-way distillable entanglement	6
B. Continuity bounds of the one-way distillable entanglement	8
C. Examples of less-entangled states	9
D. Extending the method to the two-way distillable entanglement	12
IV. Applications on quantum channel capacity	13
A. Quantum capacity of qubit channels	15
V. Concluding remarks	18
Acknowledgements.	18
References	18
A. Properties of the reverse max-relative entropy of resources	22

* These two authors contributed equally.

† wangxin73@baidu.com

B. Dual SDP for the squeezed parameter of states and channels	22
C. Two-way distillable entanglement	23
D. Proof of Proposition 3 and Proposition 4	25
E. Proof of Proposition 10	27

I. INTRODUCTION

A. Background

Quantum entanglement is the most nonclassical manifestation of quantum mechanics and has become a key ingredient in many quantum information processing tasks, e.g., teleportation [1], superdense coding [2], and quantum cryptography [3, 4]. More specifically, the entanglement resource is usually assumed to be copies of the maximally entangled states. In a practical scenario, noises inevitably occur in quantum states, resulting in some mixed entangled states, which leads to the need for distillation or purification. A natural question is how to obtain the maximally entangled states from a source of less entangled states using well-motivated operations, known as the *entanglement distillation*.

One fundamental measure for characterizing the entanglement distillation is the *one-way distillable entanglement* [5], denoted by $E_{D,\rightarrow}$, which is also one of the most important entanglement measures motivated by operational tasks. It captures the highest rate at which one can obtain the maximally entangled states from less entangled states by one-way local operations and classical communication (LOCC):

$$E_{D,\rightarrow}(\rho_{AB}) = \sup\{r : \lim_{n \rightarrow \infty} [\inf_{\Lambda} \|\Lambda(\rho_{AB}^{\otimes n}) - \Phi(2^{rn})\|_1] = 0\}, \quad (1)$$

where Λ ranges over one-way LOCC operations and $\Phi(d) = 1/d \sum_{i,j=1}^d |ii\rangle\langle jj|$ is the standard $d \otimes d$ maximally entangled state. Likewise, the *two-way distillable entanglement* $E_{D,\leftrightarrow}(\rho_{AB})$ is defined by the supremum over all achievable rates under two-way LOCC. We have for all bipartite states ρ_{AB} that $E_{D,\rightarrow}(\rho_{AB}) \leq E_{D,\leftrightarrow}(\rho_{AB})$. Notably, the distillable entanglement turns out to be closely related to the central idea of *quantum capacity* in quantum communication tasks, which is central to quantum Shannon theory. Consider modelling the noise in transmitting quantum information from Alice to Bob as a quantum channel $\mathcal{N}_{A \rightarrow B}$. The quantum capacity $Q(\mathcal{N}_{A \rightarrow B})$ is the maximal achievable rate at which Alice can reliably transmit quantum information to Bob by asymptotically many uses of the channel. By the state–channel duality, if the distillation protocol of the Choi state [6] $J_{AB}^{\mathcal{N}}$ of $\mathcal{N}_{A \rightarrow B}$ yields the maximally entangled states at a positive rate, then Alice may apply the standard teleportation scheme to send arbitrary quantum states to Bob at the same rate. Thus, one has $Q(\mathcal{N}_{A \rightarrow B}) \geq E_{D,\rightarrow}(J_{AB}^{\mathcal{N}})$ since classical forward communication in teleportation does not affect the channel capacity. For the teleportation-simulable channels [1, 7, 8], the equality here holds. Many recent works [9–15] have studied the properties of quantum communication over quantum channels, e.g., non-additivity and superactivation, which determines the complexity of calculating the quantum capacity.

Despite many efforts that have been made in the past two decades, computing $E_{D,\rightarrow}(\cdot)$ and $Q(\cdot)$ still generally remains a challenging task. Even for the qubit isotropic states and the depolarizing channels, it remains unsolved. Therefore, numerous studies try to estimate them by deriving lower and upper bounds (see, e.g., [5, 16–21] for the distillable entanglement, e.g., [22–25] for the quantum capacity). For the distillable entanglement, a well-known lower bound dubbed *Hashing bound* is established by Devetak and Winter [5], i.e., $E_{D,\rightarrow}(\rho_{AB}) \geq I_c(A)B_\rho$, where $I_c(A)B_\rho$ is the coherent information of the bipartite quantum state ρ_{AB} . Considering upper bounds, the Rains bound [16] is arguably the best-known efficiently computable bound for the two-way distillable entanglement of general states, which can consequently be

an upper bound of $E_{D,\rightarrow}(\cdot)$. Recent works [19, 21] utilize the techniques of finding upper bounds by constructing meaningful extended states. For quantum capacity, many useful upper bounds for general quantum channels are studied for benchmarking arbitrary quantum noise [26–33]. When considering some specific classes of quantum channels, useful upper bounds are also developed to help us better understand quantum communication via these channels [21, 23–25, 34–36].

In specific, due to the regularization in the characterizations of $E_{D,\rightarrow}(\cdot)$ and $Q(\cdot)$, one main strategy to establish efficiently computable upper bounds on them is to develop single-letter formulae. For example, one common approach is to decompose a state (resp. a quantum channel) into degradable parts and anti-degradable parts [24], or use approximate degradability (anti-degradability) [27]. Another recent fruitful technique called flag extension optimization [21, 23, 25] relies on finding a degradable extension of the state or the quantum channel. However, the performance of these methods is limited by the absence of a good decomposition strategy. It is unknown how to partition a general state or quantum channel to add flags or how to construct a proper and meaningful convex decomposition on them. Thus, the flag extension optimization is only effective for the states and channels with symmetry or known structures.

B. Main contributions

This work considers a family of resource measures called *reverse divergence of resources*. With a specific construction, we define the *reverse max-relative entropy of entanglement* for quantum states, which has applications for estimating the distillable entanglement. In the meantime, we introduce *reverse max-relative entropy of anti-degradability* for quantum channels as a generalization of the concept of that for states, which can be applied to bound the quantum capacity. All these bounds can be efficiently computed via semidefinite programming [37]. Furthermore, drawing on the idea of [27], we thoroughly analyze different continuity bounds on the one-way distillable entanglement of a state in terms of its anti-degradability. Finally, we investigate the distillation of the maximally entangled states under practical noises and focus on the quantum capacity of qubit channels. We show that the bound obtained by the reverse max-relative entropy of entanglement outperforms other known bounds in a high-noise region, including the Rains bound and the above continuity bounds. The upper bound offered by the reverse max-relative entropy of anti-degradability also provides an alternative interpretation of the no-cloning bound of the Pauli channel [34], and notably outperforms the continuity bounds on random unital qubit channels.

The rest of this paper is structured as follows. We start with some preliminaries used throughout the paper in section II A. In section II B, we introduce our conceptual contribution, a family of resource measures called the reverse divergence of resources. In section III, we present the application of this concept on the distillable entanglement. Specifically, in subsection III A, we show how the reverse max-relative entropy of entanglement can be used to derive a useful upper bound on the one-way distillable entanglement. We derive different continuity bounds in subsection III B, with which we compare our bounds and the Rains bound by practical distillable states in subsection III C. We also extend our method to the two-way distillable entanglement in subsection III D. In section IV, we show the application of our method in deriving upper bounds on quantum capacity. We establish some analytical results for Pauli channels and compare our bounds with continuity bounds in subsection IV A for random mixed unitary channels. In section V, we conclude with a summary and some outlooks for future research.

II. REVERSE DIVERGENCE OF RESOURCES

A. Preliminaries

Let \mathcal{H} be a finite-dimensional Hilbert space, and $\mathcal{L}(\mathcal{H})$ be the set of linear operators acting on it. We consider two parties Alice and Bob with Hilbert space $\mathcal{H}_A, \mathcal{H}_B$, whose dimensions are d_A, d_B , respectively.

A linear operator $\rho \in \mathcal{L}(\mathcal{H})$ is called a density operator if it is Hermitian and positive semidefinite with trace one. We denote the trace norm of ρ as $\|\rho\|_1 = \text{Tr} \sqrt{\rho^\dagger \rho}$ and let $\mathcal{D}(\mathcal{H})$ denote the set of density operators. We call a linear map CPTP if it is both completely positive and trace-preserving. A CPTP map that transforms linear operators in $\mathcal{L}(\mathcal{H}_A)$ to linear operators in $\mathcal{L}(\mathcal{H}_B)$ is also called a quantum channel, denoted as $\mathcal{N}_{A \rightarrow B}$. For a quantum channel $\mathcal{N}_{A \rightarrow B}$, its Choi-Jamiołkowski state is given by $J_{AB}^{\mathcal{N}} \equiv \sum_{i,j=0}^{d_A-1} |i\rangle\langle j| \otimes \mathcal{N}_{A \rightarrow B}(|i\rangle\langle j|)$, where $\{|i\rangle\}_{i=0}^{d_A-1}$ is an orthonormal basis in \mathcal{H}_A . The von Neumann entropy of a state ρ_A is $S(A)_\rho := -\text{Tr}(\rho_A \log \rho_A)$ and the coherent information of a bipartite state ρ_{AB} is defined by $I_c(A)B)_\rho := S(B)_\rho - S(AB)_\rho$. The entanglement of formation of a state ρ_{AB} is given by

$$E_F(\rho_{AB}) = \min_{\{p_i, |\phi_i\rangle\}} \sum_i p_i S(A)_{\phi_i}, \quad (2)$$

where $\rho_{AB} = \sum_i p_i |\phi_i\rangle\langle \phi_i|_{AB}$ and the minimization ranges over all pure state decomposition of ρ_{AB} . We introduce the generalized divergence $\mathbf{D}(\rho_A \|\sigma_A)$ as a map $\mathbf{D} : \mathcal{D}(\mathcal{H}_A) \times \mathcal{D}(\mathcal{H}_A) \mapsto \mathbb{R} \cup \{+\infty\}$ that obeys:

1. Faithfulness: $\mathbf{D}(\rho_A \|\sigma_A) = 0$ iff $\rho_A = \sigma_A$.
2. Data processing inequality: $\mathbf{D}(\rho_A \|\sigma_A) \geq \mathbf{D}[\mathcal{N}_{A \rightarrow A'}(\rho_A) \|\mathcal{N}_{A \rightarrow A'}(\sigma_A)]$, where $\mathcal{N}_{A \rightarrow A'}$ is an arbitrary quantum channel.

The generalized divergence is intuitively some measure of distinguishability of the two states, e.g., Bures metric, quantum relative entropy. Another example of interest is the *sandwiched Rényi relative entropy* [38, 39] of ρ, σ that is defined by

$$D_\alpha(\rho \|\sigma) := \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right)^\alpha \right], \quad (3)$$

if $\text{supp}(\rho) \subset \text{supp}(\sigma)$ and it is equal to $+\infty$ otherwise, where $\alpha \in (0, 1) \cup (1, \infty)$. In the case that $\alpha \rightarrow \infty$, one can find the *max-relative entropy* [40] of ρ with respect to σ by

$$D_{\max}(\rho \|\sigma) = \inf \{ \lambda \in \mathbb{R} : \rho \leq 2^\lambda \sigma \}. \quad (4)$$

B. Reverse divergence of resources

In the usual framework of quantum resource theories [41], there are two main notions: i) subset \mathcal{F} of free states, i.e., the states that do not possess the given resource; ii) subset \mathcal{O} of free operations, i.e., the quantum channels that are unable to generate the resource. Meanwhile, two axioms for a quantity being a resource measure $\mathcal{R}(\cdot)$ are essential:

- 1). Vanishing for free states: $\rho \in \mathcal{F} \Rightarrow \mathcal{R}(\rho) = 0$.
- 2). Monotonicity: $\mathcal{R}(\mathcal{O}(\rho)) \leq \mathcal{R}(\rho)$ for any free operation \mathcal{O} . $\mathcal{R}(\cdot)$ is called a *resource monotone*.

Let us define a family of resource measures called *reverse divergence of resources*:

$$\mathcal{R}_{\mathcal{F}}(\rho_{AB}) := \min_{\tau \in \mathcal{F}} \mathbf{D}(\tau_{AB} \|\rho_{AB}), \quad (5)$$

where \mathcal{F} is some set of free states. By the definition of the reverse divergence of resources in Eq. (5), one can easily check it satisfies condition 1). Whenever the free state set \mathcal{F} is closed, by the data-processing inequality of $\mathbf{D}(\cdot \|\cdot)$, condition 2) will be satisfied. Thus $\mathcal{R}_{\mathcal{F}}(\cdot)$ is a resource measure. Specifically, in the resource theory of entanglement, some representative free state sets are the separable states (SEP) and the states having a positive partial transpose (PPT). Examples of free sets of operations are LOCC and PPT. We




Fig 1. **Interpretation of the reverse max-relative entropy of resources.** $\mathcal{D}(\mathcal{H})$ is the set of density operators. \mathcal{F} is a set of free states, $\mathcal{R}_{\max, \mathcal{F}}(\rho)$ is the minimum value of λ for which $(1 - 2^{-\lambda})\omega + 2^{-\lambda}\tau = \rho$, for some choice of density operators $\omega \in \mathcal{D}(\mathcal{H})$ and $\tau \in \mathcal{F}$. Equivalently, this convex decomposition of ρ is characterized by the weight of resource and the free component discussed in Eq. (7). It can be interpreted as the case that the red segment occupies the least proportion while maintaining ω in $\mathcal{D}(\mathcal{H})$. The dotted line depicts the set distance between ρ and the free states set \mathcal{F} .

note that the "reverse" here means minimizing the divergence over a free state set in the first coordinate, rather than the second one which has helped one define the *relative entropy of entanglement* [42, 43] and the *max-relative entropy of entanglement* [40]. For some divergence of particular interest, e.g., the quantum relative entropy $D(\cdot||\cdot)$, relevant discussion of the coordinate choices can be traced back to work in [43]. In [44], the authors further studied properties of the quantity $\min_{\tau \in \mathcal{F}} D(\tau||\rho)$. Here, we try to further investigate meaningful applications of some reverse divergence of resources.

In the following, we consider the generalized divergence as the max-relative entropy and study a measure called *reverse max-relative entropy of resources*,

$$\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB}) := \min_{\tau \in \mathcal{F}} D_{\max}(\tau_{AB}||\rho_{AB}), \quad (6)$$

where \mathcal{F} is some set of free states. If there is no such state $\tau_{AB} \in \mathcal{F}$ that satisfies $\tau_{AB} \leq 2^\lambda \rho_{AB}$ for any $\lambda \in \mathbb{R}$, $\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})$ is set to be 0. This measure bears many nice properties. First, it can be efficiently computed via semidefinite programming (SDP) in many cases which will be discussed in the latter sections. Second, Eq. (6) gives the closest free state $\tau_{AB} \in \mathcal{F}$ to ρ_{AB} , w.r.t. the max-relative entropy. Third, $\mathcal{R}_{\max, \mathcal{F}}(\cdot)$ is subadditive w.r.t the tensor product of states. In fact, $\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})$ is closely related to the *weight of resource* $W(\rho_{AB})$ [45–48] and the *free component* $\Gamma(\rho_{AB})$ [49], both of which have fruitful properties and applications [50–52], as follows

$$2^{-\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})} = 1 - W(\rho_{AB}) = \Gamma(\rho_{AB}). \quad (7)$$

We note that each part of Eq. (7) quantifies the largest weight where a free state can take in a convex decomposition of ρ_{AB} . When moving on to operational tasks that the free state can be ignored, what is left in a convex decomposition becomes our main concern. Optimization of the weight in the decomposition can be visualized as *squeezing out* all free parts of the given state. Thus, we further introduce the *F-squeezed state* of ρ_{AB} as follows.

Definition 1 For a bipartite quantum state ρ_{AB} and a free state set \mathcal{F} , If $\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})$ is non-zero, the *F-squeezed state* of ρ_{AB} is defined by

$$\omega_{AB} = \frac{\rho_{AB} - 2^{-\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})} \cdot \tau_{AB}}{1 - 2^{-\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB})}}, \quad (8)$$




Fig 2. **Illustration of the \mathcal{F} -squeezed state of ρ_{AB} and Theorem 1.** The brown region corresponds to the sub-state τ_{AB} belonging to the free state set \mathcal{F} , which is squeezed out via a convex decomposition of ρ_{AB} . ω_{AB} is the \mathcal{F} -squeezed state.

where τ_{AB} is the closest free state to ρ_{AB} in terms of the max-relative entropy, i.e., the optimal solution in Eq. (6). If $\mathcal{R}_{\max, \mathcal{F}}(\rho_{AB}) = 0$, the \mathcal{F} -squeezed state of ρ_{AB} is itself.

In the following sections, we will illustrate the applications of the reverse max-relative entropy of resources as well as the squeezing idea on example tasks. One is giving upper bounds on the distillable entanglement of arbitrary quantum states. The other is to derive upper bounds on the quantum capacity of channels.

III. APPLICATIONS ON DISTILLABLE ENTANGLEMENT

In this section, we investigate the information-theoretic application of the reverse max-relative entropy of resources in deriving efficiently computable upper bounds on the distillable entanglement. To showcase the advantage of our bounds, we compare the results with different continuity bounds and the Rains bound on the maximally entangled states with practical noises.

A. Upper bound on the one-way distillable entanglement

Recall that the one-way distillable entanglement has a regularized formula [5]:

$$E_{D, \rightarrow}(\rho_{AB}) = \lim_{n \rightarrow \infty} \frac{1}{n} E_{D, \rightarrow}^{(1)}(\rho_{AB}^{\otimes n}), \quad (9)$$

where $E_{D, \rightarrow}^{(1)}(\rho_{AB}) := \max_T I_C(A' MB)_{T(\rho_{AB})}$, and the maximization ranges over all quantum instruments $T : A \rightarrow A'M$ on Alice's system. The regularization in Eq. (9) for $E_{D, \rightarrow}(\rho_{AB})$ is intractable to compute in most cases. However, there are some categories of states whose $E_{D, \rightarrow}$ can be reduced to single-letter formulae. Two important classes are called *degradable states* and *anti-degradable states*.

Let ρ_{AB} be a bipartite state with purification $|\phi\rangle_{ABE}$. ρ_{AB} is called degradable if there exists a CPTP map $\mathcal{M}_{B \rightarrow E}$ such that $\mathcal{M}_{B \rightarrow E}(\rho_{AB}) = \text{Tr}_B(\phi_{ABE})$, and is called anti-degradable if there exists a CPTP map $\mathcal{M}_{E \rightarrow B}$ such that $\mathcal{M}_{E \rightarrow B}(\rho_{AE}) = \text{Tr}_E(\phi_{ABE})$. Equivalently, a state is anti-degradable if and only if it has a symmetric extension [53], thus is also called a *two-extendible state*. For the degradable states, it is shown that [19]

$$E_{D, \rightarrow}^{(1)}(\rho_{AB}^{\otimes n}) = n E_{D, \rightarrow}^{(1)}(\rho_{AB}) = n I_C(A)_{B\rho}, \forall n \in \mathbb{N}, \quad (10)$$

resulting in $E_{D,\rightarrow}(\rho_{AB}) = I_c(A)B)_\rho$. For the anti-degradable states, consisting of a compact and convex set, it always holds

$$E_{D,\rightarrow}^{(1)}(\rho_{AB}) = E_{D,\rightarrow}(\rho_{AB}) = 0. \quad (11)$$

Moreover, $E_{D,\rightarrow}(\cdot)$ is convex on decomposing a state into degradable and anti-degradable parts [19]. To better exploit this convexity, we introduce the *reverse max-relative entropy of unextendible entanglement* to help identify the anti-degradable (two-extendible) part of a given bipartite state ρ_{AB} :

$$\mathcal{R}_{\max, \text{ADG}}(\rho_{AB}) := \min_{\tau \in \text{ADG}} D_{\max}(\tau_{AB} || \rho_{AB}), \quad (12)$$

where ADG is the set of all anti-degradable states. In this resource theory, the resource we consider is the extendible entanglement, and the free states are bipartite states that are possibly shareable between A and a third party E , where E is isomorphic to B . Notably, the extendibility of entanglement is a key property in entanglement theory with many existing applications [20, 54]. Here, combined with the idea of entanglement of formation, $\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})$ can be applied to derive an upper bound on the one-way distillable entanglement of an arbitrary state ρ_{AB} as shown in Theorem 1.

Theorem 1 *For any bipartite state ρ_{AB} , it satisfies*

$$E_{D,\rightarrow}(\rho_{AB}) \leq E_{\text{rev}}^u(\rho_{AB}) := [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \cdot E_F(\omega_{AB}), \quad (13)$$

where ω_{AB} is the ADG-squeezed state of ρ_{AB} , $\mathcal{R}_{\max, \text{ADG}}(\cdot)$ is the reverse max-relative entropy of unextendible entanglement, and $E_F(\cdot)$ is the entanglement of formation.

Proof Suppose the ADG-squeezed state of ρ_{AB} is ω_{AB} and the optimal solution in Eq. (6) for ρ_{AB} is τ_{AB} . It follows

$$\rho_{AB} = [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \omega_{AB} + 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})} \tau_{AB}, \quad (14)$$

where τ_{AB} is anti-degradable. Suppose ω_{AB} admit a pure state decomposition $\omega_{AB} = \sum_i p_i |\omega_i\rangle\langle\omega_i|_{AB}$. Since any pure state is degradable, we now can decompose ρ_{AB} into degradable parts and anti-degradable parts as

$$\rho_{AB} = [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \sum_i p_i |\omega_i\rangle\langle\omega_i|_{AB} + 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})} \tau_{AB}, \quad (15)$$

According to the essential convexity of $E_{D,\rightarrow}(\cdot)$ on decomposing a state into degradable and anti-degradable parts proved by Leditzky et al. [19], we have

$$\begin{aligned} E_{D,\rightarrow}(\rho_{AB}) &\leq [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \cdot \sum_i p_i E_{D,\rightarrow}(\omega_i) + 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})} E_{D,\rightarrow}(\tau_{AB}) \\ &\leq [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \cdot \sum_i p_i I_c(A)B)_{\omega_i}, \end{aligned} \quad (16)$$

where the second inequality is due to the fact in Eq. (10) and Eq. (11) of degradable states and anti-degradable states, respectively. After taking the minimization over all possible decomposition of ω_{AB} , we arrived at

$$E_{D,\rightarrow}(\rho_{AB}) \leq [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \cdot \min_i \sum_i p_i I_c(A)B)_{\omega_i} \quad (17)$$

$$= [1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}] \cdot E_F(\omega_{AB}). \quad (18)$$

■

Remark 1 The bound $E_{\text{rev}}^u(\rho_{AB})$ has a cartoon illustration as shown in Fig. 2. The main insight of it is to squeeze out as much of the free or useless part, the anti-degradable state here, as possible. We point out that squeezing *all* useless parts out does not necessarily give the tightest upper bound in terms of the one-way distillable entanglement, e.g., isotropic state [19]. Instead of squeezing out all the useless parts, there may be room for exploring more appropriate partitions when we try to decompose a specific quantum state. However, the approach we present in Theorem 1 is an invaluable method for general states as shown in subsection III C and can be seen as a variant of the continuity bound in terms of the anti-degradability of the state.

Corollary 2 For any bipartite state ρ_{AB} , it satisfies

$$E_{D,\rightarrow}(\rho_{AB}) \leq \hat{E}_{\text{rev}}^u(\rho_{AB}) := [1 - 2^{-\mathcal{R}_{\text{max,ADG}}(\rho_{AB})}] \cdot \sum_i \lambda_i S(B)_{\psi_i}, \quad (19)$$

where $\omega_{AB} = \sum_i \lambda_i |\psi_i\rangle\langle\psi_i|$ is the spectral decomposition of the ADG-squeezed state ω_{AB} of ρ_{AB} .

Corollary 2 is followed by the fact that $E_F(\omega_{AB})$ has a trivial upper bound as $E_F(\omega_{AB}) \leq \sum_i \lambda_i S(B)_{\psi_i}$. We note that any other good upper bound on the entanglement of formation can also be applied to Theorem 1. In particular, the bound $\hat{E}_{\text{rev}}^u(\rho_{AB})$ is efficiently computable since $\mathcal{R}_{\text{max,ADG}}(\rho_{AB})$ and $2^{-\mathcal{R}_{\text{max,ADG}}(\rho_{AB})}$ can be efficiently computed via an SDP. By Slater's condition [55], the following two optimization programs satisfy the strong duality, and both evaluate to $1 - 2^{-\mathcal{R}_{\text{max,ADG}}(\rho_{AB})}$. We denote P_{BE} as a permutation operator on the system B and E and remain the derivation of the dual program in Appendix B.

Primal Program	Dual Program
$\min_{\omega_{AB}, \tau_{AB}, \tau_{ABE}} \text{Tr}[\omega_{AB}],$	$\max_{M_{AB}, N_{AB}, K_{AE}} \text{Tr}[M_{AB}\rho_{AB}],$
$\text{s.t. } \rho_{AB} = \omega_{AB} + \tau_{AB},$	$\text{s.t. } M_{AB} \leq I_{AB},$
$\omega_{AB} \geq 0, \tau_{AB} \geq 0, \tau_{ABE} \geq 0,$	$M_{AB} + N_{AB} + K_{AB} \leq 0,$
$\text{Tr}_E[\tau_{ABE}] = \text{Tr}_B[\tau_{ABE}] = \tau_{AB},$	$N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger \geq 0.$

It is worth mentioning that this new bound is related to the bound $E_{\text{DA}}(\cdot)$ proposed in [19] utilizing the convexity of $E_{D,\rightarrow}(\cdot)$ on decomposing a state into degradable and anti-degradable parts. Remarkably, such decomposition is challenging to construct since the degradable state set is not necessarily convex. Thus it is difficult to compute $E_{\text{DA}}(\cdot)$ in [19] in practice due to the hardness of tracking all possible decompositions. Whereas, $\hat{E}_{\text{rev}}^u(\rho_{AB})$ overcomes this difficulty and is efficiently computable. It outperforms the known upper bounds for many maximally entangled states under practical noises in a high-noise region, as shown in subsection III C. Furthermore, our method is flexible in selecting other decomposition strategies, i.e., the object functions in the SDP, other than trivially calculating the ratio at the sub-state decomposition.

B. Continuity bounds of the one-way distillable entanglement

Note that the insight of the bound above is considering the distance of a given state between the anti-degradable states set. With different distance measures, the authors in [27] derived continuity bounds on quantum capacity in terms of the (anti)degradability of the channel. Thus for self-contained, we introduce some distance measures between a state and the set ADG and prove the continuity bounds for the state version as a comparison with $\hat{E}_{\text{rev}}^u(\rho_{AB})$.

Definition 2 Let ρ_{AB} be a bipartite quantum state, the anti-degradable set distance is defined by

$$d_{\text{set}}(\rho_{AB}) := \min_{\sigma_{AB} \in \text{ADG}} \frac{1}{2} \|\rho_{AB} - \sigma_{AB}\|_1, \quad (20)$$

where the minimization ranges over all anti-degradable states on system AB .

Analogous to the anti-degradable set distance, one also has the anti-degradable map distance as follows.

Definition 3 ([19]) Let ρ_{AB} be a bipartite quantum state with purification ϕ_{ABE} , the anti-degradable map distance is defined by

$$d_{\text{map}}(\rho_{AB}) := \min_{\mathcal{D}: E \rightarrow B} \frac{1}{2} \|\rho_{AB} - \mathcal{D}(\rho_{AE})\|_1, \quad (21)$$

where $\rho_{AE} = \text{Tr}_B(\phi_{ABE})$ and the minimization ranges over all CPTP maps \mathcal{D} .

Both parameters can be computed via SDP, ranging from $[0, 1]$, and are equal to 0 iff ρ_{AB} is anti-degradable. Similar to the idea in [27] for channels and the proof techniques in [19], we utilize the continuity bound of the conditional entropy in Lemma S4 proved by Winter [56] to derive two continuity upper bounds on the one-way distillable entanglement, concerning the distance measures above. The proofs can be found in Appendix D. We denote $h(p) = -p \log p - (1-p) \log(1-p)$ as the binary entropy and $g(p) := (1+p)h(\frac{p}{1+p})$ as the bosonic entropy.

Proposition 3 For any bipartite state ρ_{AB} with an anti-degradable set distance ε_{set} , it satisfies

$$E_{D, \rightarrow}(\rho_{AB}) \leq E_{\text{SCB}}(\rho_{AB}) := 2\varepsilon_{\text{set}} \log(|A|) + g(\varepsilon_{\text{set}}). \quad (22)$$

Proposition 4 For any bipartite state ρ_{AB} with an anti-degradable map distance ε_{map} , it satisfies

$$E_{D, \rightarrow}(\rho_{AB}) \leq E_{\text{MCB}}(\rho_{AB}) := 4\varepsilon_{\text{map}} \log(|B|) + 2g(\varepsilon_{\text{map}}). \quad (23)$$

C. Examples of less-entangled states

We now compare the performance of different continuity bounds and the Rains bound with $\hat{E}_{\text{rev}}^u(\cdot)$ by some concrete examples. Due to noise and other device imperfections, one usually obtains some less entangled states in practice rather than the maximally entangled ones. Such a disturbance can be characterized by some CPTP maps appearing in each party. Thus for the task of the distillation of the maximally entangled states under practical noises, we consider Alice and Bob are sharing pairs of maximally entangled states affected by bi-local noisy channels, i.e.,

$$\rho_{A'B'} = \mathcal{N}_{A \rightarrow A'} \otimes \mathcal{N}_{B \rightarrow B'}(\Phi_{AB}). \quad (24)$$

a. Qubit system Suppose Alice's qubit is affected by the qubit amplitude damping channel with Kraus operators $K_0 = |0\rangle\langle 0| + \sqrt{1-\gamma}|1\rangle\langle 1|$, $K_1 = \sqrt{\gamma}|0\rangle\langle 1|$, and Bob's qubit is affected by the depolarizing channel $\mathcal{D}(\rho) = (1-p)\rho + pI_2/2$. Set $\gamma = 0.1$ and the noise parameter of depolarizing noise varies in the range $p \in [0.15, 0.3]$. $E_{D, \rightarrow}(\rho_{A'B'})$ has upper bounds as functions of p shown in Fig. 3 (a).




Fig 3. Upper bounds on the one-way distillable entanglement of less entangled states in a qubit, qutrit, and qudit system. Panel (a-c) depict the cases where the maximally entangled states are affected by bi-local noises in different systems as stated in Eq. (24). The x -axis represents the change of the depolarizing noise p . Panel (d) represents the analytic bound in a qubit system in which both parties are affected by a local depolarizing noise as stated in Proposition 5. For all panels, the state's coherent information I_c provides a lower bound. R is the Rains bound. \hat{E}_{rev}^u is the upper bound derived in Corollary 2. E_{SCB} and E_{MCB} are continuity bounds derived in Proposition 3 and 4, respectively. It shows that \hat{E}_{rev}^u outperforms all other upper bounds on these less entangled states.

b. Qutrit system For the system with a dimension $d > 2$, we consider the multilevel versions of the amplitude damping channel (MAD) [57] as a local noise for Alice. The Kraus operators of a MAD channel in a d -dimensional system are defined by

$$\begin{aligned} \hat{K}_{ij} &\equiv \sqrt{\gamma_{ji}} |i\rangle\langle j|, \quad \forall i, j \text{ s.t. } 0 \leq i \leq d-1, \\ \hat{K}_0 &\equiv |0\rangle\langle 0| + \sum_{1 \leq j \leq d-1} \sqrt{1 - \xi_j} |j\rangle\langle j|, \end{aligned} \quad (25)$$

with γ_{ji} real quantities describing the decay rate from the j -th to the i -th level that fulfill the conditions

$$\begin{cases} 0 \leq \gamma_{ji} \leq 1, & \forall i, j \text{ s.t. } 0 \leq i < j \leq d-1, \\ \xi_j \equiv \sum_{0 \leq i < j} \gamma_{ji} \leq 1, & \forall j = 1, \dots, d-1. \end{cases} \quad (26)$$

Suppose Alice qutrit is affected by a MAD channel with $\gamma_{10} = \gamma_{20} = 0.1, \gamma_{21} = 0$. Bob's qutrit is affected by a qutrit depolarizing channel with the noise parameter p . Then $E_{D, \rightarrow}(\rho_{A'B'})$ has upper bounds

as functions of p shown in Fig. 3 (b).

c. *Qudit system* For the qudit system, we consider Alice's qudit is affected by a MAD channel with $d = 4$ defined in Eq. (25) and Eq. (26), where $\gamma_{10} = \gamma_{20} = \gamma_{30} = \gamma_{21} = 0.1, \gamma_{31} = \gamma_{32} = 0$. Let Bob's qudit be affected by a qudit depolarizing channel with noise parameter p , then $E_{D,\rightarrow}(\rho_{A'B'})$ has upper bounds as functions of p shown in Fig. 3 (c).

d. *Noisy Bell state* Specifically for a Bell state destroyed by bi-local depolarizing noises, i.e., $\rho_{A'B'} = \mathcal{D}_{A\rightarrow A'} \otimes \mathcal{D}_{B\rightarrow B'}(\Phi_{AB})$, we give an analytical upper bound on $E_{D,\rightarrow}(\rho_{A'B'})$ as shown in Proposition 5.

Proposition 5 *Let Φ_{AB} be a Bell state and $\mathcal{D}_{A\rightarrow A'}, \mathcal{D}_{B\rightarrow B'}$ be depolarizing channels on system A and B, respectively. The state $\rho_{A'B'} = \mathcal{D}_{A\rightarrow A'} \otimes \mathcal{D}_{B\rightarrow B'}(\Phi_{AB})$ either is anti-degradable with $E_{D,\rightarrow}(\rho_{A'B'}) = 0$ or satisfies*

$$2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} = 3(p_A + p_B - p_{APB}) \quad (27)$$

and

$$E_{D,\rightarrow}(\rho_{A'B'}) \leq 1 - 3(p_A + p_B - p_{APB}), \quad (28)$$

where p_A, p_B are noise parameters of $\mathcal{D}_{A\rightarrow A'}$ and $\mathcal{D}_{B\rightarrow B'}$, respectively.

Proof In the beginning, we note that if $1 - 3(p_A + p_B - p_{APB}) \leq 0$, the state $\rho_{A'B'}$ is anti-degradable according to Lemma S5, which gives $E_{D,\rightarrow}(\rho_{A'B'}) = 0$. Then if $1 - 3(p_A + p_B - p_{APB}) > 0$, we will first prove $2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \geq 3(p_A + p_B - p_{APB})$ use the primal SDP in (III A). We show that $\{\hat{\omega}_{A'B'}, \hat{\tau}_{A'B'E}\}$ is a feasible solution to the primal SDP, where

$$\hat{\omega}_{A'B'} = \frac{1 - 3(p_A + p_B - p_{APB})}{2} (|00\rangle\langle 00| + |11\rangle\langle 11| + |00\rangle\langle 11| + |11\rangle\langle 00|), \quad (29)$$

$$\hat{\tau}_{A'B'E} = \begin{pmatrix} 4\zeta & 0 & 0 & 0 & 0 & 2\zeta & 2\zeta & 0 \\ 0 & \zeta & \zeta & 0 & 0 & 0 & 0 & 2\zeta \\ 0 & \zeta & \zeta & 0 & 0 & 0 & 0 & 2\zeta \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2\zeta & 0 & 0 & 0 & 0 & \zeta & \zeta & 0 \\ 2\zeta & 0 & 0 & 0 & 0 & \zeta & \zeta & 0 \\ 0 & 2\zeta & 2\zeta & 0 & 0 & 0 & 0 & 4\zeta \end{pmatrix}, \quad (30)$$

with $\zeta = \frac{p_A + p_B - p_{APB}}{4}$. When the Bell state shared by Alice and Bob is affected by bi-local depolarizing channels $\mathcal{D}_{A\rightarrow A'}$ and $\mathcal{D}_{B\rightarrow B'}$ with noise parameters p_A and p_B , respectively, the resulting state is

$$\begin{aligned} \rho_{A'B'} &= \mathcal{D}_{A\rightarrow A'} \otimes \mathcal{D}_{B\rightarrow B'}(\Phi_{AB}) \\ &= \frac{2 - (p_A + p_B) + p_{APB}}{4} (|00\rangle\langle 00| + |11\rangle\langle 11|) + \frac{p_A + p_B - p_{APB}}{4} (|01\rangle\langle 01| + |10\rangle\langle 10|) \\ &\quad + \frac{1 - (p_A + p_B) + p_{APB}}{2} (|00\rangle\langle 11| + |11\rangle\langle 00|). \end{aligned} \quad (31)$$

Then the unnormalized state $\hat{\tau}_{A'B'} = \rho_{A'B'} - \hat{\omega}_{A'B'}$ is

$$\begin{aligned} \hat{\tau}_{A'B'} &= \frac{5(p_A + p_B - p_{APB})}{4} (|00\rangle\langle 00| + |11\rangle\langle 11|) + \frac{p_A + p_B - p_{APB}}{4} (|01\rangle\langle 01| + |10\rangle\langle 10|) \\ &\quad + (p_A + p_B - p_{APB}) (|00\rangle\langle 11| + |11\rangle\langle 00|). \end{aligned} \quad (32)$$

It is straightforward to check that $\hat{\tau}_{A'B'E} \geq 0$ and $\text{Tr}_E[\hat{\tau}_{A'B'E}] = \text{Tr}_B[\hat{\tau}_{A'B'E}] = \hat{\tau}_{A'B'}$. Thus, we have proven that $\{\hat{\omega}_{A'B'}, \hat{\tau}_{A'B'E}\}$ is a feasible solution to the primal SDP, implying that

$$1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \leq \text{Tr}[\hat{\omega}_{A'B'}] = 1 - 3(p_A + p_B - p_{APB}) \quad (33)$$

and $2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \geq 3(p_A + p_B - p_{APB})$.

Second, we will use the dual SDP in (III A) to prove $2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \leq 3(p_A + p_B - p_{APB})$. We show that $\{M_{AB}, N_{AB}, K_{AB}\}$ is a feasible solution to the dual problem, where

$$M_{AB} = \begin{pmatrix} -1 & 0 & 0 & 2 \\ 0 & -3 & 0 & 0 \\ 0 & 0 & -3 & 0 \\ 2 & 0 & 0 & -1 \end{pmatrix}, N_{AB} = K_{AB} = -\frac{1}{2}M_{AB}. \quad (34)$$

To check the constraints, we have $M_{AB} \leq I_{AB}$, $M_{AB} + N_{AB} + K_{AB} = 0$ and $N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger \geq 0$. By direct calculation, we have $\text{Tr}[M_{AB}\rho_{A'B'}] = 1 - 3(p_A + p_B - p_{APB})$. Thus, we have proven $\{M_{AB}, N_{AB}, K_{AB}\}$ is a feasible solution, which implies that

$$1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \geq \text{Tr}[M_{AB}\rho_{A'B'}] = 1 - 3(p_A + p_B - p_{APB}), \quad (35)$$

and $2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} \leq 3(p_A + p_B - p_{APB})$. Combining the dual part with the primal part, we arrive at

$$2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{A'B'})} = 3(p_A + p_B - p_{APB}) \quad (36)$$

Since the normalized state $\omega_{A'B'}$ of $\hat{\omega}_{A'B'}$ is a maximally entangled state with $E_{D, \rightarrow}(\omega_{A'B'}) = 1$, we have $E_{D, \rightarrow}(\rho_{A'B'}) \leq 1 - 3(p_A + p_B - p_{APB})$. ■

As shown in Fig 3, we plot four different upper bounds and the Hashing bound in different dimensional systems in a high-noise region. We can see that $\hat{E}_{\text{rev}}^u(\rho_{AB})$ is lower than the other two continuity bounds and the Rains bound in the orange dotted line, acting the tightest among all upper bounds. We demonstrate the advantage of our analytic bound in Proposition 5 by comparing it with the continuity bounds and the Rains bound in Fig. 3 (d), where the depolarizing noise parameter p_B is set to 0.1 and the noise parameter p_A is varied.

D. Extending the method to the two-way distillable entanglement

Similar to the *reverse max-relative entropy of unextendible entanglement*, for a given bipartite state ρ_{AB} , we introduce the *reverse max-relative entropy of NPT entanglement* as

$$\mathcal{R}_{\max, \text{PPT}}(\rho_{AB}) := \min_{\tau \in \text{PPT}} D_{\max}(\tau_{AB} || \rho_{AB}), \quad (37)$$

where the minimization ranges over all PPT states. Based on the convexity of $E_{D, \leftrightarrow}(\cdot)$ on decomposing a state into the maximally correlated (MC) states and the PPT states [19] (see Appendix C for more details), the reverse max-relative entropy of NPT entanglement can be applied to derive an upper bound on the two-way distillable entanglement as Theorem 6.

Theorem 6 *For any bipartite state ρ_{AB} , it satisfies*

$$E_{D, \leftrightarrow}(\rho_{AB}) \leq E_{\text{rev}}^{\text{npt}}(\rho_{AB}) := [1 - 2^{-\mathcal{R}_{\max, \text{PPT}}(\rho_{AB})}] \cdot E_F(\omega_{AB}), \quad (38)$$

where ω_{AB} is the PPT-squeezed state of ρ_{AB} and $\mathcal{R}_{\max, \text{PPT}}(\cdot)$ is the reverse max-relative entropy of NPT entanglement.

It also follows an efficiently computable relaxation as $\hat{E}_{\text{rev}}^{\text{npt}}(\rho_{AB}) = [1 - 2^{-\mathcal{R}_{\text{max,PPT}}(\rho_{AB})}] \sum_i \lambda_i S(B)_{\psi_i}$, where $\omega_{AB} = \sum_i \lambda_i |\psi_i\rangle\langle\psi_i|$ is the spectral decomposition of the PPT-squeezed state ω_{AB} of ρ_{AB} .

In fact, $\hat{E}_{\text{rev}}^{\text{npt}}(\cdot)$ can be interpreted as an easily computable version of the bound $E_{\text{MP}}(\cdot)$ in [19], utilizing the convexity of $E_{D,\leftrightarrow}(\rho_{AB})$ on the convex decomposition of ρ_{AB} into MC states and PPT states. Since the set of all MC states is not convex, tracking all possible decompositions to compute $E_{\text{MP}}(\cdot)$ is generally hard. However, $\hat{E}_{\text{rev}}^{\text{npt}}(\cdot)$ is efficiently computable by SDP and tightens the upper bound of the example states illustrated in [19], whose details can be found in Appendix C. We note that $R(\rho_{AB}) \leq E_{\text{MP}}(\rho_{AB}) \leq \hat{E}_{\text{rev}}^{\text{npt}}(\rho_{AB})$ where $R(\cdot)$ is the Rains bound for the two-way distillable entanglement. Nevertheless, $\hat{E}_{\text{rev}}^{\text{npt}}(\cdot)$ connects the reverse max-relative entropy of NPT entanglement with the entanglement of formation, and we believe such connection would shed light on the study of other quantum resource theories as well.

IV. APPLICATIONS ON QUANTUM CHANNEL CAPACITY

For a general quantum channel $\mathcal{N}_{A \rightarrow B}$, its quantum capacity has a regularized formula proved by Lloyd, Shor, and Devetak [58–60]:

$$Q(\mathcal{N}) = \lim_{n \rightarrow \infty} \frac{1}{n} Q^{(1)}(\mathcal{N}^{\otimes n}), \quad (39)$$

where $Q^{(1)}(\mathcal{N}) := \max_{|\phi\rangle_{A'A}} I_c(A'B)_{(\mathcal{I} \otimes \mathcal{N})(\phi)}$ is the channel coherent information. Similar to the one-way distillable entanglement of a state, the regularization in Eq. (39) makes the quantum capacity of a channel intractable to compute generally. Substantial efforts have been made to establish upper bounds. One routine is inspired by the Rains bound in entanglement theory [16]. Tomamichel et al. introduced Rains information [61] for a quantum channel as an upper bound on the quantum capacity. Then some efficiently computable relaxations or estimations are given in [30, 33]. Another routine is to consider the (anti)degradability of the quantum channels and to construct flag extensions, which gives the currently tightest upper bound for quantum channels with symmetry or known structures [21, 23, 25, 27].

A channel $\mathcal{N}_{A \rightarrow B}$ is called *degradable* if there exists a CPTP map $\mathcal{D}_{B \rightarrow E}$ such that $\mathcal{N}^c = \mathcal{D} \circ \mathcal{N}$, and is called *anti-degradable* if there exists a CPTP map $\mathcal{A}_{E \rightarrow B}$ such that $\mathcal{N} = \mathcal{A} \circ \mathcal{N}^c$. It is known that \mathcal{N} is (anti)degradable if and only if its Choi state $J_{\mathcal{N}}$ is (anti)degradable. The quantum capacity of an anti-degradable channel is zero and the coherent information of a degradable channel is additive which leads to $Q(\mathcal{N}) = Q^{(1)}(\mathcal{N})$. Also, the quantum capacity admits convexity on decomposing a channel into a convex combination of degradable channels and anti-degradable channels [24] which can give a simple upper bound on $Q(\mathcal{N})$. Concerning the (anti)degradability of a channel, the authors in [27] called a channel ε -*degradable channel* if there is a CPTP map $\mathcal{D}_{B \rightarrow E}$ such that $\|\mathcal{N}^c - \mathcal{D} \circ \mathcal{N}\|_{\diamond} \leq 2\varepsilon$. A channel is called ε -*anti-degradable channel* if there is a CPTP map $\mathcal{A}_{E \rightarrow B}$ such that $\|\mathcal{N} - \mathcal{A} \circ \mathcal{N}^c\|_{\diamond} \leq 2\varepsilon$. Based on these, one has continuity bounds of the quantum capacity as follows.

Theorem 7 ([27]) *Given a quantum channel $\mathcal{N}_{A \rightarrow B}$, if it is ε -degradable, then it satisfies $Q(\mathcal{N}) \leq Q^{(1)}(\mathcal{N}) + \varepsilon \log(d_E - 1) + h(\varepsilon) + 2\varepsilon \log d_E + g(\varepsilon)$. If $\mathcal{N}_{A \rightarrow B}$ is ε -anti-degradable, it satisfies $Q(\mathcal{N}) \leq \varepsilon \log(|B| - 1) + 2\varepsilon \log |B| + h(\varepsilon) + g(\varepsilon)$.*

With a similar spirit of the reverse max-relative entropy of unextendible entanglement in Eq. (12), for a quantum channel $\mathcal{N}_{A \rightarrow B}$, we define the *reverse max-relative entropy of anti-degradability of the channel* as

$$\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N}_{A \rightarrow B}) := \min_{\mathcal{N}'_{A \rightarrow B} \in \mathcal{C}_{\text{ADG}}} D_{\text{max}}(\mathcal{N}'_{A \rightarrow B} || \mathcal{N}_{A \rightarrow B}), \quad (40)$$

where \mathcal{C}_{ADG} is the set of all anti-degradable channels and the max-relative entropy of $\mathcal{N}'_{A \rightarrow B}$ with respect to $\mathcal{N}_{A \rightarrow B}$ is defined by

$$D_{\text{max}}(\mathcal{N}'_{A \rightarrow B} || \mathcal{N}_{A \rightarrow B}) := \inf\{\lambda \in \mathbb{R} : J_{AB}^{\mathcal{N}'} \leq 2^\lambda J_{AB}^{\mathcal{N}}\}. \quad (41)$$

If there is no such a channel $\mathcal{N}'_{A \rightarrow B} \in \mathcal{C}_{\text{ADG}}$ that satisfies $J_{AB}^{\mathcal{N}'} \leq 2^\lambda J_{AB}^{\mathcal{N}}$, $\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N}_{A \rightarrow B})$ is set to be 0. Similar to the state case, $\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N}_{A \rightarrow B})$ has a geometric implication analogous to the distance between $\mathcal{N}_{A \rightarrow B}$ to the set of all anti-degradable channels. We can introduce the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$ as follows.

Definition 4 For a quantum channel $\mathcal{N}_{A \rightarrow B}$ and the anti-degradable channel set \mathcal{C}_{ADG} , if $\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})$ is non-zero, the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$ is defined by

$$\mathcal{S}_{A \rightarrow B} = \frac{\mathcal{N}_{A \rightarrow B} - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})} \cdot \mathcal{N}'_{A \rightarrow B}}{1 - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}} \quad (42)$$

where $\mathcal{N}'_{A \rightarrow B}$ is the closest anti-degradable channel to $\mathcal{N}_{A \rightarrow B}$ in terms of the max-relative entropy, i.e., the optimal solution in Eq. (40). If $\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})$ is zero, the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$ is itself.

Notably, $\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})$, or $2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}$ equivalently, can be efficiently computed via SDP. The following two optimization programs satisfy the strong duality, and both evaluate to $1 - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}$. We remain the derivation of the dual program in Appendix B.

<u>Primal Program</u>	<u>Dual Program</u>
$\min_{\Gamma_{AB}^S, \Gamma_{AB}^{\mathcal{N}'}, \gamma_{ABE}} \text{Tr}[\Gamma_{AB}^S],$	$\max_{M_{AB}, N_{AB}, K_{AB}, R_A} \text{Tr}[M_{AB} J_{AB}^{\mathcal{N}}],$
$\text{s.t. } J_{AB}^{\mathcal{N}} = \Gamma_{AB}^S + \Gamma_{AB}^{\mathcal{N}'},$	$\text{s.t. } M_{AB} \leq (1 - \text{Tr } R_A / d_A) I_{AB} + R_A \otimes I_B,$
$\Gamma_{AB}^S \geq 0, \Gamma_{AB}^{\mathcal{N}'} \geq 0, \gamma_{ABE} \geq 0,$	$M_{AB} + N_{AB} + K_{AB} \leq 0,$
$\text{Tr}_B[\Gamma_{AB}^{\mathcal{N}'}] = \text{Tr}[\Gamma_{AB}^{\mathcal{N}'}] / d_A \cdot I_A,$	$N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E) P_{BE}^\dagger \geq 0.$
$\text{Tr}_E[\gamma_{ABE}] = \text{Tr}_B[\gamma_{ABE}] = \Gamma_{AB}^{\mathcal{N}'},$	

The conceptual idea we used here is similar to that for the state case in Eq. (6) and Definition 1, which is to squeeze or isolate out as much part of anti-degradable channel as possible via a convex decomposition of the original channel. The insight here is that one can ignore the contribution from the anti-degradable part for the quantum capacity, and the quantum capacity admits convexity on the decomposition into degradable and anti-degradable parts. In this way, the following Theorem 8 gives an upper bound $Q_{\text{sqz}}(\mathcal{N})$ for the quantum capacity of \mathcal{N} .

Theorem 8 Given a quantum channel $\mathcal{N}_{A \rightarrow B}$, if it has an ADG-squeezed channel $\mathcal{S}_{A \rightarrow B}$, we denote $\hat{\mathcal{S}}_{A \rightarrow BB'}$ as an extended channel of $\mathcal{S}_{A \rightarrow B}$ such that $\text{Tr}_{B'}[\hat{\mathcal{S}}_{A \rightarrow BB'}(\rho_A)] = \mathcal{S}_{A \rightarrow B}(\rho_A), \forall \rho_A \in \mathcal{D}(\mathcal{H}_A)$. Then it satisfies

$$Q(\mathcal{N}) \leq Q_{\text{sqz}}(\mathcal{N}) := [1 - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}] \cdot \min \left\{ Q^{(1)}(\hat{\mathcal{S}}) \mid \hat{\mathcal{S}}_{A \rightarrow BB'} \text{ is degradable} \right\}, \quad (43)$$

where the minimization is over all possible extended channels of $\mathcal{S}_{A \rightarrow B}$. If there is no such a degradable $\hat{\mathcal{S}}_{A \rightarrow BB'}$ exists, the value of this bound is set to be infinity.

Proof By the definition of the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$, we have

$$\mathcal{N}_{A \rightarrow B} = [1 - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}] \cdot \mathcal{S}_{A \rightarrow B} + 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})} \cdot \mathcal{N}'_{A \rightarrow B} \quad (44)$$

where $\mathcal{N}'_{A \rightarrow B}$ is anti-degradable. We write an extended channel of $\mathcal{N}'_{A \rightarrow B}$ as $\hat{\mathcal{N}}'_{A \rightarrow BB'}(\rho_A) = \mathcal{N}'_{A \rightarrow B}(\rho_A) \otimes |0\rangle\langle 0|_{B'}$, which is obviously anti-degradable. Then we can construct a quantum channel $\hat{\mathcal{N}}_{A \rightarrow BB'}$ as

$$\hat{\mathcal{N}}_{A \rightarrow BB'} = [1 - 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})}] \cdot \hat{\mathcal{S}}_{A \rightarrow BB'} + 2^{-\tilde{\mathcal{R}}_{\text{max,ADG}}(\mathcal{N})} \cdot \hat{\mathcal{N}}'_{A \rightarrow BB'}, \quad (45)$$

such that $\text{Tr}_{B'}[\widehat{\mathcal{S}}_{A \rightarrow BB'}(\rho_A)] = \mathcal{S}_{A \rightarrow B}(\rho_A)$ for any state ρ_A and $\widehat{\mathcal{S}}_{A \rightarrow BB'}$ is degradable. This means after discarding the partial environment B' , the receiver can obtain the original quantum information sent through $\mathcal{N}_{A \rightarrow B}$. In this case, $\widehat{\mathcal{N}}$ can certainly convey more quantum information than the \mathcal{N} , i.e., $Q(\mathcal{N}) \leq Q(\widehat{\mathcal{N}})$. Note that the quantum capacity admits convexity on the decomposition into degradable parts and anti-degradable parts [24]. We conclude that

$$\begin{aligned} Q(\mathcal{N}) &\leq Q(\widehat{\mathcal{N}}) \leq [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot Q(\widehat{\mathcal{S}}) + 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})} \cdot Q(\widehat{\mathcal{N}}') \\ &= [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot Q^{(1)}(\widehat{\mathcal{S}}), \end{aligned} \quad (46)$$

where the equality is followed by the quantum capacity is additive on degradable channels and is zero for anti-degradable channels. Considering the freedom of the choice of $\widehat{\mathcal{S}}_{A \rightarrow BB'}$, we obtain

$$Q_{\text{sqz}}(\mathcal{N}) := [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot \min \left\{ Q^{(1)}(\widehat{\mathcal{S}}) \mid \widehat{\mathcal{S}}_{A \rightarrow BB'} \text{ is degradable} \right\} \quad (47)$$

as an upper bound on $Q(\mathcal{N})$. \blacksquare

Remark 2 Theorem 8 can be seen as a channel version of Theorem 1. However, in order to utilize the convexity of the quantum capacity after the squeezing process, it is challenging to decompose the ADG-squeezed channel into the sum of degradable ones. An alternative approach here is to use the idea of the extension channel. For the qubit channels specifically, this bound is efficiently computable and effective, as shown in subsection IV A.

A. Quantum capacity of qubit channels

For a quantum channel with dimension two in both input and output systems, we prove that the ADG-squeezed channel is always degradable. Thus, we give an efficiently computable upper bound on the quantum capacity using the idea of the reverse max-relative entropy of anti-degradability.

Proposition 9 *For any qubit channel $\mathcal{N}_{A \rightarrow B}$, it is either anti-degradable or satisfies*

$$Q(\mathcal{N}) \leq [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot \max_{p \in [0,1]} I_c \left(p|0\rangle\langle 0| + (1-p)|1\rangle\langle 1|, \mathcal{S}_{A \rightarrow B} \right), \quad (48)$$

where $I_c(\rho, \mathcal{N}) \equiv H(\mathcal{N}(\rho)) - H(\mathcal{N}^c(\rho))$ and $\mathcal{S}_{A \rightarrow B}$ is the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$.

Proof By the definition of the ADG-squeezed channel of $\mathcal{N}_{A \rightarrow B}$, we have

$$\mathcal{N}_{A \rightarrow B} = [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot \mathcal{S}_{A \rightarrow B} + 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})} \cdot \mathcal{N}'_{A \rightarrow B} \quad (49)$$

where $\mathcal{N}'_{A \rightarrow B}$ is anti-degradable and $\mathcal{S}_{A \rightarrow B}$ is not anti-degradable. If $\mathcal{S}_{A \rightarrow B}$ is also not degradable, we can further decompose $\mathcal{S}_{A \rightarrow B}$ into $\mathcal{S} = c\mathcal{S}_0 + (1-c)\mathcal{S}_1$ such that \mathcal{S}_0 is degradable and \mathcal{S}_1 is anti-degradable since the extreme points of the set of all qubit channels have been shown to be degradable or anti-degradable channels [27, 62]. This conflicts with the definition of $2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}$, which implies $\mathcal{S}_{A \rightarrow B}$ is degradable. Thus,

$$\begin{aligned} Q(\mathcal{N}) &\leq [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot Q(\mathcal{S}) + 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})} \cdot Q(\mathcal{N}') \\ &= [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot Q^{(1)}(\mathcal{S}) \\ &= [1 - 2^{-\widetilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}] \cdot \max_{p \in [0,1]} I_c \left(p|0\rangle\langle 0| + (1-p)|1\rangle\langle 1|, \mathcal{S}_{A \rightarrow B} \right). \end{aligned} \quad (50)$$

Note that the last equality is because $\mathcal{S}_{A \rightarrow B}$ is degradable, and diagonal input states outperform non-diagonal states during the optimization of the channel coherent information [24]. \blacksquare




Fig 4. Upper bounds on the quantum capacity of the random mixed unitary channels. In Panel (a) and (b), we compare our bound Q_{sqz} in Proposition 9 with other upper bounds on the quantum capacity of 1000 randomly generated mixed unitary channels with fixed parameters $p_0 = 0.58, p_1 = 0.22, p_2 = 0.15, p_3 = 0.05$. The x -axis represents the distance between two bounds, and the y -axis represents the distribution of these channels. $Q_{\epsilon\text{-adg}}$ is the continuity bound in Theorem 7 regarding the anti-degradability. $\hat{R}_{\alpha(10)}$ is the bound in [33]. Panel (c) and (d) depict the same comparison for another 1000 randomly generated mixed unitary channels with fixed parameters $p_0 = 0.6, p_1 = 0.2, p_2 = 0.1, p_3 = 0.1$. In (a) and (c), we see that our bound always outperforms $Q_{\epsilon\text{-adg}}$. In (b) and (d), we can see for many cases, our bound is tighter than $\hat{R}_{\alpha(10)}$.

a. Mixed unitary channels To compare the performance of our method with some best-known computable bounds, e.g., the continuity bound in Theorem 7 and the bound \hat{R}_{α} [33] generalized from the max-Rain information [30], we consider the mixed unitary channel $\mathcal{U}_{A \rightarrow B}(\cdot)$ as follows:

$$\mathcal{U}(\rho) = \sum_{i=0}^k p_i U_i \rho U_i^\dagger, \quad (51)$$

where $\sum_{i=0}^k p_i = 1$ and U_i are unitary operators on a qubit system. In specific, we choose some fixed set of parameters p_i and sample 1000 channels with randomly generated unitaries according to the Haar measure. We compute the distance between Q_{sqz} and other bounds, then have statistics on the distribution of these channels according to the distance value. The distribution results are depicted in Fig. 4 where the purple region corresponds to the cases Q_{sqz} is tighter, and the pink region corresponds to the cases Q_{sqz} is looser. We can see that in Fig. 4(a) and Fig. 4(c), Q_{sqz} always outperforms the continuity bound of anti-degradability and in Fig. 4(b) and Fig. 4(d), our bound is tighter than $\hat{R}_{\alpha(10)}$ for many cases.

b. Pauli channels A representative qubit channel is the Pauli channel describing bit-flip errors and phase-flip errors with certain probabilities in qubits. A qubit Pauli channel $\Lambda(\cdot)$ is defined as:

$$\Lambda(\rho) = p_0 \rho + p_1 X \rho X + p_2 Y \rho Y + p_3 Z \rho Z, \quad (52)$$

where X, Y, Z are the Pauli operators and $\sum_{i=0}^3 p_i = 1$ are probability parameters. Note that for the quantum capacity, we only need to consider the cases where p_0 dominates. Since if, for example, bit flip error X happens with probability larger than p_0 , one can first apply a X flip, mapping that channel




Fig 5. **Upper bounds on quantum capacity of the covariant Pauli channels.** The parameters are $p_3 = 0.05, p_0 \in [0.75, 0.85]$. The single-shot coherent information $Q^{(1)}$ [63, 64] provides a lower bound on the quantum capacity. $\hat{R}_{\alpha(10)}$ is the upper bound in [33]. $Q_{\epsilon-\text{adg}}$ is the continuity bound of anti-degradability. Q_{flag} is the upper bound by flag extension method in [65]. Q_{sqz} is our bound in Corollary 12 and Q_{nc} is the no-cloning bound [34]. It shows that Q_{sqz} can recover the no-cloning bound and outperform other computable bounds.

back into the case where $p_0 > p_1$. After utilizing our method on Pauli channels, the ADG-squeezed parameter is characterized in Proposition 10, whose proof can be found in Appendix E. Thus, combined with Proposition 9, we can recover the no-cloning bound [34] on the quantum capacity of qubit Pauli channels.

Proposition 10 *For a qubit Pauli channel $\Lambda(\cdot)$ with $p_0 \geq p_i > 0 (i = 1, 2, 3)$, it is either anti-degradable or satisfies*

$$2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} = (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2 \quad (53)$$

with an ADG-squeezed channel as the identity channel.

Theorem 11 ([34]) *For a qubit Pauli channel $\Lambda(\cdot)$, its quantum capacity is either vanishing or satisfies*

$$Q(\Lambda) \leq 1 - (\sqrt{p_1} + \sqrt{p_2})^2 - (\sqrt{p_2} + \sqrt{p_3})^2 - (\sqrt{p_1} + \sqrt{p_3})^2 \quad (54)$$

One recent work [65] studies the capacities of a subclass of Pauli channels called the *covariant Pauli channel*, where the parameters are set $p_1 = p_2$ with $p_0 + 2p_1 + p_3 = 1$, i.e., $\Lambda_{\text{cov}}(\rho) = p_0\rho + p_1(X\rho X + Y\rho Y) + p_3Z\rho Z$. Applying Theorem 11 on the covariant Pauli channels, we can bound their quantum capacity as follows.

Corollary 12 *For a covariant Pauli channel $\Lambda_{\text{cov}}(\cdot)$, it is either anti-degradable with a zero quantum capacity or satisfies $Q(\Lambda_{\text{cov}}) \leq Q_{\text{sqz}}(\Lambda_{\text{cov}})$, where*

$$Q_{\text{sqz}}(\Lambda_{\text{cov}}) = 3p_0 + p_3 - \sqrt{8(p_3 - p_0p_3 - p_3^2)} - 2. \quad (55)$$

In Fig. 5, we compare our bound with the upper bounds given in [65] and the continuity bound of anti-degradability in Theorem 7. It can be seen that our bound in the orange line, coinciding with the no-cloning bound, outperforms previous bounds, and thus can better characterize the quantum capacity of $\Lambda_{\text{cov}}(\cdot)$ when it is close to being anti-degradable.

V. CONCLUDING REMARKS

We have introduced a resource measure called reverse divergence of resources for quantum states. Specifically, by choosing target free state sets and max-relative entropy, we have employed the reverse max-relative entropy of entanglement which is also related to the "weight of resource" in general resource theory. From a conceptual and technical side, these concepts help us to quantify how much useless entanglement we can squeeze out from a state or a channel, which is meaningful in the distillable entanglement and the quantum capacity, respectively. As these quantities can be efficiently determined by semidefinite programming, they are useful in deriving efficiently computable upper bounds on the distillable entanglement and the quantum capacity.

To better explore entanglement distillation, we have derived different continuity bounds on the one-way distillable entanglement in terms of the anti-degradability of the state. In particular, our bound derived from the reverse max-relative entropy of unextendible entanglement outperforms the continuity bounds and the Rains bound in estimating the one-way distillable entanglement for the maximally entangled states under certain noises of interest. We also introduced the reverse max-relative entropy of NPT entanglement and established connections to prior results on the two-way distillable entanglement. Furthermore, we have established an upper bound on the quantum capacity based on the reverse max-relative entropy of anti-degradability of quantum channels. Our method delivers improved results for random mixed unitary qubit channels, making it an improved benchmark for the quantum capacities of these noisy channels. Also, the analytical bound obtained from our method also recovers the no-cloning bound on Pauli channels [34].

These results open a novel way to connect valuable quantum resource measures with quantum communication tasks. Except for the existing applications of the reverse max-relative entropy of resources [47–52, 66], we expect the reverse divergence of resources will find more applications in quantum resource theories in both asymptotic and non-asymptotic regimes [67–72]. One may further consider choosing other divergences, e.g., the *sandwiched Rényi relative entropy* to solve problems in other tasks possibly. In addition, the reverse divergence of resources may potentially be helpful to the study of other quantum resource theories, including asymmetry [73], thermodynamics [74], coherence [75], nonlocality [76, 77], non-stabilizerness [78–80], and non-Gaussianity [81–83]. It is also an interesting problem to explore the idea of squeezing useless resources to estimate resource purification rates in quantum resource theories.

ACKNOWLEDGEMENTS.

Part of this work was done when C. Z. and C. Z. were research interns at Baidu Research. We would like to thank Bartosz Regula and Ludovico Lami for helpful comments.

-
- [1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels, *Physical Review Letters* **70**, 1895 (1993).
 - [2] C. H. Bennett and S. J. Wiesner, Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states, *Physical Review Letters* **69**, 2881 (1992).
 - [3] C. H. Bennett and G. Brassard, Quantum cryptography: Public key distribution and coin tossing, in *International Conference on Computers, Systems & Signal Processing, Bangalore, India, Dec 9-12, 1984* (1984) pp. 175–179.

- [4] A. K. Ekert, Quantum cryptography based on Bell's theorem, *Physical Review Letters* **67**, 661 (1991).
- [5] I. Devetak and A. Winter, Distillation of secret key and entanglement from quantum states, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **461**, 207 (2005), arXiv:0306078 [quant-ph].
- [6] M.-D. Choi, Completely positive linear maps on complex matrices, *Linear Algebra and its Applications* **10**, 285 (1975).
- [7] S. L. Braunstein and H. J. Kimble, Teleportation of Continuous Quantum Variables, *Physical Review Letters* **80**, 869 (1998).
- [8] R. F. Werner, All teleportation and dense coding schemes, *Journal of Physics A: Mathematical and General* **34**, 7081 (2001), arXiv:0003070 [quant-ph].
- [9] D. Elkouss and S. Strelchuk, Superadditivity of Private Information for Any Number of Uses of the Channel, *Physical Review Letters* **115**, 040501 (2015), arXiv:1502.05326.
- [10] F. Leditzky, D. Leung, V. Siddhu, G. Smith, and J. A. Smolin, Generic nonadditivity of quantum capacity in simple channels, *arXiv preprint arXiv:2202.08377* (2022), arXiv:2202.08377.
- [11] F. Leditzky, D. Leung, V. Siddhu, G. Smith, and J. A. Smolin, The platypus of the quantum channel zoo, *arXiv preprint arXiv:2202.08380* (2022), arXiv:2202.08380.
- [12] G. Smith and J. Yard, Quantum communication with zero-capacity channels, *Science* **321**, 1812 (2008).
- [13] F. Leditzky, D. Leung, and G. Smith, Dephasing Channel and Superadditivity of Coherent Information, *Physical Review Letters* **121**, 160501 (2018), arXiv:1806.08327.
- [14] M. Tomamichel, M. Berta, and J. M. Renes, Quantum Coding with Finite Resources, *Nature Communications* **7**, 11419 (2015), arXiv:1504.04617.
- [15] V. Siddhu, Entropic singularities give rise to quantum transmission, *Nature Communications* **12**, 5750 (2021), arXiv:2003.10367.
- [16] E. M. Rains, A semidefinite program for distillable entanglement, *IEEE Transactions on Information Theory* **47**, 2921 (2000), arXiv:0008047 [quant-ph].
- [17] X. Wang and R. Duan, Improved semidefinite programming upper bound on distillable entanglement, *Physical Review A* **94**, 050301 (2016).
- [18] M. Hayashi, *Igarss 2014*, 1 (Springer, 2006) pp. 1–5, arXiv:arXiv:1011.1669v3.
- [19] F. Leditzky, N. Datta, and G. Smith, Useful states and entanglement distillation, *IEEE Transactions on Information Theory* **64**, 4689 (2017), arXiv:1701.03081.
- [20] E. Kaur, S. Das, M. M. Wilde, and A. Winter, Extendibility Limits the Performance of Quantum Processors, *Physical Review Letters* **123**, 070502 (2019), arXiv:1803.10710.
- [21] X. Wang, Pursuing the Fundamental Limits for Quantum Communication, *IEEE Transactions on Information Theory* **67**, 4524 (2021), arXiv:1912.00931.
- [22] Y. Ouyang, Channel covariance, twirling, contraction, and some upper bounds on the quantum capacity, *Quantum Information and Computation* **14**, 917 (2011), arXiv:1106.2337.
- [23] F. Kianvash, M. Fanizza, and V. Giovannetti, Bounding the quantum capacity with flagged extensions, *Quantum* **6**, 647 (2022), arXiv:2008.02461.
- [24] M. M. Wolf and D. Pérez-García, Quantum capacities of channels with small environment, *Physical Review A* **75**, 012303 (2007).
- [25] M. Fanizza, F. Kianvash, and V. Giovannetti, Quantum Flags and New Bounds on the Quantum Capacity of the Depolarizing Channel, *Physical Review Letters* **125**, 020503 (2020), arXiv:1911.01977.
- [26] A. Holevo and R. Werner, Evaluating capacities of bosonic Gaussian channels, *Physical Review A* **63**, 032312 (2001).
- [27] D. Sutter, V. B. Scholz, A. Winter, and R. Renner, Approximate Degradable Quantum Channels, *IEEE Transactions on Information Theory* **63**, 7832 (2014), arXiv:1412.0980.
- [28] A. Müller-Hermes, D. Reeb, and M. M. Wolf, Positivity of linear maps under tensor powers, *Journal of Mathematical Physics* **57**, 015202 (2016), arXiv:1502.05630.
- [29] X. Wang and R. Duan, A semidefinite programming upper bound of quantum capacity, in *2016 IEEE International Symposium on Information Theory (ISIT)*, Vol. 2016-Augus (IEEE, 2016) pp. 1690–1694.
- [30] X. Wang, K. Fang, and R. Duan, Semidefinite Programming Converse Bounds for Quantum Communication, *IEEE Transactions on Information Theory* **65**, 2583 (2019), arXiv:1709.00200.
- [31] R. Piarczyk, Z. Zhao, Y. Ouyang, V. Vedral, and J. F. Fitzsimons, Causal Limit on Quantum Communication, *Physical Review Letters* **123**, 150502 (2019), arXiv:1804.02594.
- [32] S. Pirandola, R. Laurenza, C. Ottaviani, and L. Banchi, Fundamental limits of repeaterless quantum communications, *Nature Communications* **8**, 15043 (2017).

- [33] K. Fang and H. Fawzi, Geometric Renyi Divergence and its Applications in Quantum Channel Capacities, arXiv:1909.05758 (2019), [arXiv:1909.05758](#).
- [34] N. J. Cerf, Pauli Cloning of a Quantum Bit, [Physical Review Letters](#) **84**, 4497 (2000).
- [35] G. Smith, J. A. Smolin, and A. Winter, The Quantum Capacity With Symmetric Side Channels, [IEEE Transactions on Information Theory](#) **54**, 4208 (2008).
- [36] L. Gao, M. Junge, and N. LaRacuente, Capacity bounds via operator space methods, [Journal of Mathematical Physics](#) **59**, 122202 (2018), [arXiv:1509.07294](#).
- [37] L. Vandenberghe and S. Boyd, Semidefinite Programming, [SIAM Review](#) **38**, 49 (1996).
- [38] M. M. Wilde, A. Winter, and D. Yang, Strong converse for the classical capacity of entanglement-breaking and hadamard channels via a sandwiched rényi relative entropy, [Communications in Mathematical Physics](#) **331**, 593 (2014).
- [39] M. Müller-Lennert, F. Dupuis, O. Szehr, S. Fehr, and M. Tomamichel, On quantum rényi entropies: A new generalization and some properties, [Journal of Mathematical Physics](#) **54**, 122203 (2013).
- [40] N. Datta, Min- and max-relative entropies and a new entanglement monotone, [IEEE Transactions on Information Theory](#) **55**, 2816 (2009).
- [41] E. Chitambar and G. Gour, Quantum resource theories, [Reviews of Modern Physics](#) **91**, 10.1103/revmodphys.91.025001 (2019).
- [42] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Quantifying entanglement, [Physical Review Letters](#) **78**, 2275 (1997).
- [43] V. Vedral, M. B. Plenio, K. Jacobs, and P. L. Knight, Statistical inference, distinguishability of quantum states, and quantum entanglement, [Physical Review A](#) **56**, 4452 (1997).
- [44] J. Eisert, K. Audenaert, and M. B. Plenio, Remarks on entanglement measures and non-local state distinguishability, [Journal of Physics A: Mathematical and General](#) **36**, 5605 (2003).
- [45] A. C. Elitzur, S. Popescu, and D. Rohrlich, Quantum nonlocality for each pair in an ensemble, [Physics Letters A](#) **162**, 25 (1992).
- [46] M. Lewenstein and A. Sanpera, Separability and entanglement of composite quantum systems, [Phys. Rev. Lett.](#) **80**, 2261 (1998).
- [47] A. F. Ducuara and P. Skrzypczyk, Operational Interpretation of Weight-Based Resource Quantifiers in Convex Quantum Resource Theories, [Physical Review Letters](#) **125**, 110401 (2020), [arXiv:1909.10486](#).
- [48] R. Uola, T. Bullock, T. Kraft, J.-P. Pellonpää, and N. Brunner, All Quantum Resources Provide an Advantage in Exclusion Tasks, [Physical Review Letters](#) **125**, 110402 (2020), [arXiv:1909.10484](#).
- [49] K. Fang and Z.-W. Liu, No-Go Theorems for Quantum Resource Purification: New Approach and Channel Theory, [PRX Quantum](#) **3**, 010337 (2022), [arXiv:2010.11822](#).
- [50] B. Regula and R. Takagi, Fundamental limitations on distillation of quantum channel resources, [Nature Communications](#) **12**, 10.1038/s41467-021-24699-0 (2021).
- [51] B. Regula, L. Lami, and M. M. Wilde, Overcoming entropic limitations on asymptotic state transformations through probabilistic protocols, arXiv preprint arXiv:2209.03362 (2022).
- [52] B. Regula, L. Lami, and M. M. Wilde, Postselected quantum hypothesis testing, arXiv preprint arXiv:2209.10550 (2022).
- [53] G. O. Myhr, *Symmetric extension of bipartite quantum states and its use in quantum key distribution with two-way postprocessing*, Ph.D. thesis (2011).
- [54] K. Wang, X. Wang, and M. M. Wilde, Quantifying the unextendibility of entanglement, [arXiv:1911.07433](#) (2019), [arXiv:1911.07433](#).
- [55] G. Giorgi and T. H. Kjeldsen, *Traces and emergence of nonlinear programming* (Springer Science & Business Media, 2013).
- [56] A. Winter, Tight uniform continuity bounds for quantum entropies: Conditional entropy, relative entropy distance and energy constraints, [Communications in Mathematical Physics](#) **347**, 291 (2016).
- [57] S. Chessa and V. Giovannetti, Quantum capacity analysis of multi-level amplitude damping channels, [Communications Physics](#) **4**, 1 (2021).
- [58] S. Lloyd, Capacity of the noisy quantum channel, [Physical Review A](#) **55**, 1613 (1997).
- [59] P. W. Shor, The quantum channel capacity and coherent information, in *lecture notes, MSRI Workshop on Quantum Computation* (2002).
- [60] I. Devetak, The Private Classical Capacity and Quantum Capacity of a Quantum Channel, [IEEE Transactions on Information Theory](#) **51**, 44 (2005).

- [61] M. Tomamichel, M. M. Wilde, and A. Winter, Strong converse rates for quantum communication, *IEEE Transactions on Information Theory* **63**, 715 (2014), [arXiv:1406.2946](#).
- [62] T. S. Cubitt, M. B. Ruskai, and G. Smith, The structure of degradable quantum channels, *Journal of Mathematical Physics* **49**, 102104 (2008).
- [63] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Mixed-state entanglement and quantum error correction, *Physical Review A* **54**, 3824 (1996).
- [64] J. Bausch and F. Leditzky, Error thresholds for arbitrary pauli noise, *SIAM Journal on Computing* **50**, 1410 (2021).
- [65] A. Poshtvan and V. Karimipour, Capacities of the covariant pauli channel, *Physical Review A* **106**, 062408 (2022).
- [66] A. F. Ducuara, P. Lipka-Bartosik, and P. Skrzypczyk, Multiobject operational tasks for convex quantum resource theories of state-measurement pairs, *Physical Review Research* **2**, [10.1103/physrevresearch.2.033374](#) (2020).
- [67] M. Tomamichel, *Quantum Information Processing with Finite Resources*, SpringerBriefs in Mathematical Physics, Vol. 5 (Springer International Publishing, Cham, 2016).
- [68] K. Fang, X. Wang, M. Tomamichel, and R. Duan, Non-asymptotic Entanglement Distillation, *IEEE Transactions on Information Theory* **65**, 6454 (2019), [arXiv:1706.06221](#).
- [69] N. Datta, Min- and Max-Relative Entropies and a New Entanglement Monotone, *IEEE Transactions on Information Theory* **55**, 2816 (2009).
- [70] L. Wang and R. Renner, One-Shot Classical-Quantum Capacity and Hypothesis Testing, *Physical Review Letters* **108**, 200501 (2012).
- [71] B. Regula, K. Fang, X. Wang, and M. Gu, One-shot entanglement distillation beyond local operations and classical communication, *New Journal of Physics* **21**, 103017 (2019), [arXiv:1906.01648](#).
- [72] X. Wang and M. M. Wilde, Resource theory of asymmetric distinguishability, *Physical Review Research* **1**, 033170 (2019), [arXiv:1905.11629](#).
- [73] I. Marvian and R. W. Spekkens, The theory of manipulations of pure state asymmetry: I. basic tools, equivalence classes and single copy transformations, *New Journal of Physics* **15**, 033001 (2013).
- [74] C. Sparaciari, L. Del Rio, C. M. Scandolo, P. Faist, and J. Oppenheim, The first law of general quantum resource theories, *Quantum* **4**, 259 (2020).
- [75] A. Streltsov, G. Adesso, and M. B. Plenio, Colloquium : Quantum coherence as a resource, *Reviews of Modern Physics* **89**, 041003 (2017), [arXiv:1609.02439](#).
- [76] R. Gallego, L. E. Würflinger, A. Acín, and M. Navascués, Operational framework for nonlocality, *Physical Review Letters* **109**, [10.1103/physrevlett.109.070401](#) (2012).
- [77] J. I. de Vicente, On nonlocality as a resource theory and nonlocality measures, *Journal of Physics A: Mathematical and Theoretical* **47**, 424017 (2014).
- [78] V. Veitch, S. A. Hamed Mousavian, D. Gottesman, and J. Emerson, The resource theory of stabilizer quantum computation, *New Journal of Physics* **16**, 013009 (2014), [arXiv:1307.7171](#).
- [79] M. Howard and E. Campbell, Application of a Resource Theory for Magic States to Fault-Tolerant Quantum Computing, *Physical Review Letters* **118**, 090501 (2017), [arXiv:1609.07488](#).
- [80] X. Wang, M. M. Wilde, and Y. Su, Quantifying the magic of quantum channels, *New Journal of Physics* **21**, 103002 (2019), [arXiv:1903.04483](#).
- [81] R. Takagi and Q. Zhuang, Convex resource theory of non-gaussianity, *Physical Review A* **97**, 062337 (2018).
- [82] L. Lami, B. Regula, X. Wang, R. Nichols, A. Winter, and G. Adesso, Gaussian quantum resource theories, *Physical Review A* **98**, [10.1103/physreva.98.022335](#) (2018).
- [83] F. Albarelli, M. G. Genoni, M. G. A. Paris, and A. Ferraro, Resource theory of quantum non-gaussianity and wigner negativity, *Physical Review A* **98**, [10.1103/physreva.98.052350](#) (2018).
- [84] J. Chen, Z. Ji, D. Kribs, N. Lütkenhaus, and B. Zeng, Symmetric extension of two-qubit states, *Physical Review A* **90**, [10.1103/physreva.90.032318](#) (2014).

Appendix A: Properties of the reverse max-relative entropy of resources

Proposition S1 (Subadditivity) For bipartite states ρ_0 and ρ_1 , it satisfies

$$\mathcal{R}_{\max, \mathcal{F}}(\rho_0) + \mathcal{R}_{\max, \mathcal{F}}(\rho_1) \geq \mathcal{R}_{\max, \mathcal{F}}(\rho_0 \otimes \rho_1). \quad (\text{A.1})$$

Proof Since the max-relative entropy is additive, w.r.t, tensor products, we have

$$\begin{aligned} \min_{\tau' \in \mathcal{F}} D_{\max}(\tau' || \rho_0 \otimes \rho_1) &\leq \min_{\tau_0, \tau_1 \in \mathcal{F}} D_{\max}(\tau_0 \otimes \tau_1 || \rho_0 \otimes \rho_1) \\ &= \min_{\tau_0 \in \mathcal{F}} D_{\max}(\tau_0 || \rho_0) + \min_{\tau_1 \in \mathcal{F}} D_{\max}(\tau_1 || \rho_1), \end{aligned} \quad (\text{A.2})$$

which completes the proof. \blacksquare

Proposition S2 (Monotonicity) Let Φ be a one-way LOCC, then for a bipartite state ρ , it satisfies

$$\mathcal{R}_{\max, \mathcal{F}}[\Phi(\rho)] \leq \mathcal{R}_{\max, \mathcal{F}}(\rho). \quad (\text{A.3})$$

Proof This can be directly obtained by the monotonicity of the max-relative entropy with respect to the action of channels. \blacksquare

Appendix B: Dual SDP for the squeezed parameter of states and channels

The primal SDP for calculating $2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})}$ of the state ρ_{AB} can be written as:

$$1 - 2^{-\mathcal{R}_{\max, \text{ADG}}(\rho_{AB})} = \min_{\omega_{AB}, \tau_{AB}, \tau_{ABE}} \text{Tr}[\omega_{AB}], \quad (\text{B.1a})$$

$$\text{s.t. } \rho_{AB} = \omega_{AB} + \tau_{AB}, \quad (\text{B.1b})$$

$$\omega_{AB} \geq 0, \tau_{AB} \geq 0, \tau_{ABE} \geq 0, \quad (\text{B.1c})$$

$$\text{Tr}_E[\tau_{ABE}] = \text{Tr}_B[\tau_{ABE}] = \tau_{AB}, \quad (\text{B.1d})$$

where Eq. (B.1d) corresponds to the anti-degradable condition of τ_{AB} . The Lagrange function of the primal problem is

$$\begin{aligned} L(\omega_{AB}, \tau_{AB}, \tau_{ABE}, M, N, K) &= \text{Tr}[\omega_{AB}] + \langle M, \rho_{AB} - \omega_{AB} - \tau_{AB} \rangle \\ &\quad + \langle N, \text{Tr}_E[\tau_{ABE}] - \tau_{AB} \rangle + \langle K, \text{Tr}_B[\tau_{ABE}] - \tau_{AB} \rangle \\ &= \langle M, \rho_{AB} \rangle + \langle I - M, \omega_{AB} \rangle + \langle -M - N - K, \tau_{AB} \rangle \\ &\quad + \langle N \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger, \tau_{ABE} \rangle, \end{aligned} \quad (\text{B.2})$$

where M_{AB}, N_{AB}, K_{AB} are Lagrange multipliers and P_{BE} is the permutation operator between B and E . The corresponding Lagrange dual function is

$$g(M, N, K) = \inf_{\omega_{AB}, \tau_{AB}, \tau_{ABE} \geq 0} L(\omega_{AB}, \tau_{AB}, \tau_{ABE}, M, N, K). \quad (\text{B.3})$$

Since $\omega_{AB} \geq 0, \tau_{AB} \geq 0, \tau_{ABE} \geq 0$, it must hold that $I - M_{AB} \geq 0, -M - N - K \geq 0, N \otimes I + P_{BE}(K \otimes I)P_{BE}^\dagger \geq 0$. Thus the dual SDP is

$$\begin{aligned} \max_{M_{AB}, N_{AB}, K_{AB}} \text{Tr}[M_{AB}\rho_{AB}], \\ \text{s.t. } M_{AB} \leq I_{AB}, \\ M_{AB} + N_{AB} + K_{AB} \leq 0, \\ N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger \geq 0. \end{aligned} \quad (\text{B.4})$$

The primal SDP for calculating $2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})}$ of the channel $\mathcal{N}_{A \rightarrow B}$ is:

$$1 - 2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\mathcal{N})} = \min_{\Gamma_{AB}^{\mathcal{S}}, \Gamma_{AB}^{\mathcal{N}'}, \gamma_{ABE}} \text{Tr}[\Gamma_{AB}^{\mathcal{S}}], \quad (\text{B.5a})$$

$$\text{s.t. } J_{AB}^{\mathcal{N}} = \Gamma_{AB}^{\mathcal{S}} + \Gamma_{AB}^{\mathcal{N}'}, \quad (\text{B.5b})$$

$$\Gamma_{AB}^{\mathcal{S}} \geq 0, \Gamma_{AB}^{\mathcal{N}'} \geq 0, \gamma_{ABE} \geq 0, \quad (\text{B.5c})$$

$$\text{Tr}_B[\Gamma_{AB}^{\mathcal{N}'}] = \text{Tr}[\Gamma_{AB}^{\mathcal{N}'}]/d_A \cdot I_A, \quad (\text{B.5d})$$

$$\text{Tr}_E[\gamma_{ABE}] = \text{Tr}_B[\gamma_{ABE}] = \Gamma_{AB}^{\mathcal{N}'}, \quad (\text{B.5e})$$

where Eq. (B.5e) corresponds to the anti-degradable condition of the unnormalized Choi state $\Gamma_{AB}^{\mathcal{N}'}$. The Lagrange function of the primal problem is

$$\begin{aligned} & L(\Gamma_{AB}^{\mathcal{S}}, \Gamma_{AB}^{\mathcal{N}'}, \gamma_{ABE}, M, N, K, R) \\ &= \text{Tr}[\Gamma_{AB}^{\mathcal{S}}] + \langle M, J_{AB}^{\mathcal{N}} - \Gamma_{AB}^{\mathcal{S}} - \Gamma_{AB}^{\mathcal{N}'} \rangle + \langle N, \text{Tr}_E[\gamma_{ABE}] - \Gamma_{AB}^{\mathcal{N}'} \rangle + \langle K, \text{Tr}_B[\gamma_{ABE}] - \Gamma_{AB}^{\mathcal{N}'} \rangle \\ & \quad + \langle R, \text{Tr}_B[\Gamma_{AB}^{\mathcal{S}}] - \text{Tr}[\Gamma_{AB}^{\mathcal{S}}]/d_A \cdot I_A \rangle \\ &= \langle M, J_{AB}^{\mathcal{N}} \rangle + \langle (1 - \text{Tr} R/d_A)I - M + R \otimes I_B, \Gamma_{AB}^{\mathcal{S}} \rangle + \langle -M - N - K, \Gamma_{AB}^{\mathcal{N}'} \rangle \\ & \quad + \langle N \otimes I_E + P_{BE}(K_{AE} \otimes I_B)P_{BE}^\dagger, \gamma_{ABE} \rangle, \end{aligned} \quad (\text{B.6})$$

where M_{AB}, N_{AB}, K_{AB} are Lagrange multipliers and P_{BE} is the swap operator between the system B and E . The corresponding Lagrange dual function is

$$g(M, N, K) = \inf_{\Gamma_{AB}^{\mathcal{S}}, \Gamma_{AB}^{\mathcal{N}'}, \gamma_{ABE} \geq 0} L(\Gamma_{AB}^{\mathcal{S}}, \Gamma_{AB}^{\mathcal{N}'}, \gamma_{ABE}, M, N, K). \quad (\text{B.7})$$

Since $\Gamma_{AB}^{\mathcal{S}} \geq 0, \Gamma_{AB}^{\mathcal{N}'} \geq 0, \gamma_{ABE} \geq 0$, it must hold that

$$\begin{aligned} (1 - \text{Tr} R_A/d_A)I_{AB} - M_{AB} + R_A \otimes I_B &\geq 0 \\ -M_{AB} - N_{AB} - K_{AB} &\geq 0 \\ N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger &\geq 0. \end{aligned} \quad (\text{B.8})$$

Thus the dual SDP is

$$\begin{aligned} & \max_{M_{AB}, N_{AB}, K_{AB}, R_A} \text{Tr}[M_{AB} J_{AB}^{\mathcal{N}}], \\ & \text{s.t. } M_{AB} \leq (1 - \text{Tr} R_A/d_A)I_{AB} + R_A \otimes I_B, \\ & \quad M_{AB} + N_{AB} + K_{AB} \leq 0, \\ & \quad N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger \geq 0. \end{aligned} \quad (\text{B.9})$$

Appendix C: Two-way distillable entanglement

We first start with the definition of the maximally correlated (MC) state.

Definition S1 A bipartite state ρ_{AB} on $\mathbb{C}^d \times \mathbb{C}^d$ is said to be maximally correlated (MC), if there exist bases $\{|i\rangle_A\}_{i=0}^{d-1}$ and $\{|i\rangle_B\}_{i=0}^{d-1}$ such that

$$\rho_{AB} = \sum_{i,j=0}^{d-1} \alpha_{ij} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B, \quad (\text{C.1})$$

where (α_{ij}) is a positive semidefinite matrix with trace 1.

We note that every pure state is an MC state. Then, by the following lemma, one can easily arrive at the upper bound in Theorem 6.

Lemma S3 *The two-way distillable entanglement is convex on convex combinations of MC and PPT states.*

We recall the example state given in [19]. Consider a d dimensional Hilbert space, the generalized Pauli operators X and Z are defined by their action on a computational basis $\{|k\rangle\}_{k=0}^{d-1}$ as:

$$X|k\rangle := |k+1(\text{mod } d)\rangle \quad Z|k\rangle := \omega^k|k\rangle, \quad (\text{C.2})$$

where $\omega := \exp(2\pi i/d)$ is a d -th root of unity. The generalized Pauli operators satisfy $XZ = \omega ZX$. Then the generalized Bell basis $\{|\Phi_{n,m}\rangle\}_{n,m=0,\dots,d-1}$ are defined as

$$|\Phi_{n,m}\rangle := (I_d \otimes X^m Z^n) |\Phi\rangle, \quad (\text{C.3})$$

where $|\Phi\rangle := \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |ii\rangle$. Now set $d = 3$ and denote $B = \{1, 6, 8\}$. After numbering $|\Phi_{n,m}\rangle = |\Phi_{3n+m+1}\rangle$, we define the state

$$\omega_{AB} = \sum_{i,j=1}^d \alpha_{i,j} |\Phi_{B[i]}\rangle \langle \Phi_{B[j]}|, \quad (\text{C.4})$$

where $\alpha = |\psi\rangle\langle\psi|$, $|\psi\rangle = \frac{1}{3}(|0\rangle + |1\rangle + |2\rangle)$. Then consider the states of the form

$$\rho_{AB} = (1-p)\omega_{AB} + p\tau_{AB}, \quad (\text{C.5})$$

where $p \in [0, 1]$, the state τ_{AB} is the following PPT entangled state with $a = \frac{1}{2}$:

$$\tau_{AB} = \frac{1}{8a+1} \begin{pmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & a & 0 \\ a & 0 & 0 & 0 & a & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{pmatrix}. \quad (\text{C.6})$$

We plot the different upper bounds on the two-way distillable entanglement $E_{D,\leftrightarrow}(\rho_{AB})$ in Fig. S1. It shows that our bound $\hat{E}_{\text{rev}}^{\text{mpt}}(\rho_{AB})$ is tighter than E_{W} and E_{MP} as a function of p .




Fig S1. **Upper bounds on the two-way distillable entanglement.** \hat{E}_{rev}^{npt} is our bound derived from the reverse max-relative entropy of NPT entanglement. E_W is the SDP bound in [17]. E_{MP} is the upper bound in [19] based on the direct division of ρ_{AB} into the MC states and the PPT states.

Appendix D: Proof of Proposition 3 and Proposition 4

Lemma S4 ([56]) For any states ρ_{AB} and σ_{AB} such that $\frac{1}{2}\|\rho_{AB} - \sigma_{AB}\|_1 \leq \varepsilon \leq 1$, it satisfies

$$|S(A|B)_\rho - S(A|B)_\sigma| \leq 2\varepsilon \log(|A|) + (1 + \varepsilon)h\left(\frac{\varepsilon}{1 + \varepsilon}\right). \quad (\text{D.1})$$

Proposition 3 For any bipartite state ρ_{AB} with an anti-degradable set distance ε_{set} , it satisfies

$$E_{D, \rightarrow}(\rho_{AB}) \leq 2\varepsilon_{\text{set}} \log(|A|) + (1 + \varepsilon_{\text{set}})h\left(\frac{\varepsilon_{\text{set}}}{1 + \varepsilon_{\text{set}}}\right). \quad (\text{D.2})$$

Proof Since ρ_{AB} has a anti-degradable set distance ε_{set} , we denote σ_{AB} the anti-degradable state with $\frac{1}{2}\|\rho_{AB} - \sigma_{AB}\|_1 \leq \varepsilon_{\text{set}}$. Let $T : A^n \rightarrow A'M$ be an instrument with isometry $U_n : A^n \rightarrow A'MN$ and denote $\Delta = 2\varepsilon_{\text{set}} \log(|A|) + (1 + \varepsilon_{\text{set}})h\left(\frac{\varepsilon_{\text{set}}}{1 + \varepsilon_{\text{set}}}\right)$. Then we have

$$E_{D, \rightarrow}^{(1)}(\rho_{AB}^{\otimes n}) = \max_{U_n} I_c(A'|B^n M)_{U_n \rho^{\otimes n} U_n^\dagger} \quad (\text{D.3a})$$

$$= \max_{U_n} -S(A'|B^n M)_{U_n \rho^{\otimes n} U_n^\dagger} \quad (\text{D.3b})$$

$$\leq \max_{U_n} -S(A'|B^n M)_{U_n (\sigma^{\otimes n} \rho^{\otimes n-1}) U_n^\dagger} + \Delta \quad (\text{D.3c})$$

$$\leq \max_{U_n} -S(A'|B^n M)_{U_n \sigma^{\otimes n} U_n^\dagger} + n\Delta \quad (\text{D.3d})$$

$$= n\Delta, \quad (\text{D.3e})$$

where Eq. (D.3c) follows by the fact that

$$\|\rho_0 \otimes \rho_1 - \sigma_0 \otimes \sigma_1\|_1 \leq \|\rho_0 - \sigma_0\|_1 + \|\rho_1 - \sigma_1\|_1, \quad (\text{D.4})$$

and Lemma S4. The inequality in Eq. (D.3d) follows by applying the same argument n times considering $\sigma^{\otimes i} \rho^{\otimes n-i}$ for $i = 1, \dots, n$. After dividing Eq. (D.3) by n and taking the limit $n \rightarrow \infty$, we arrive at

$$E_{D, \rightarrow}(\rho_{AB}) = \lim_{n \rightarrow \infty} \frac{1}{n} E_{D, \rightarrow}^{(1)}(\rho_{AB}^{\otimes n}) \leq \Delta. \quad (\text{D.5})$$

■

Proposition 4 For any bipartite state ρ_{AB} with an anti-degradable map distance ε_{map} , it satisfies

$$E_{D,\rightarrow}(\rho_{AB}) \leq 4\varepsilon_{\text{map}} \log(|B|) + 2(1 + \varepsilon_{\text{map}})h\left(\frac{\varepsilon_{\text{map}}}{1 + \varepsilon_{\text{map}}}\right). \quad (\text{D.6})$$

Proof Let ϕ_{ABE} be a purification of ρ_{AB} , $\mathcal{D} : E \rightarrow B$ be the CPTP map such that $\frac{1}{2}\|\rho_{AB} - \mathcal{D}(\rho_{AE})\|_1 \leq \varepsilon_{\text{map}}$ with an isometry $W : E \rightarrow B'G$. Let $T : A^n \rightarrow A'M$ be an instrument with isometry $U_n : A^n \rightarrow A'MN$ and denote $\Delta = 2\varepsilon_{\text{map}} \log(|B|) + (1 + \varepsilon_{\text{map}})h\left(\frac{\varepsilon_{\text{map}}}{1 + \varepsilon_{\text{map}}}\right)$. For $t = 1, 2, \dots, n$, we can define pure states

$$\begin{aligned} \psi_{A^n B^n B'_1 G_1 \dots B'_t G_t E_{t+1} \dots E_n}^t &= (W_1 \otimes \dots \otimes W_t) \phi_{ABE}^n \\ \theta_{A'MN B^n B'_1 G_1 \dots B'_t G_t E_{t+1} \dots E_n}^t, \quad \omega_{A'MN B^n E^n} &= U_n \psi^t, \quad \omega_{A'MN B^n E^n} = U_n \phi_{ABE}^n \end{aligned} \quad (\text{D.7})$$

We further define $\hat{\rho}_{AB'} = \mathcal{D}(\rho_{AE})$ which shares the same purification with ρ_{AB} listed above, thus an anti-degradable state. Then for $t = n$ we have $\theta^n = U_n(W_1 \otimes \dots \otimes W_n)U_n^\dagger \omega$, it yields

$$\begin{aligned} I_c(A')B^n M)_\omega &= I_c(A')B^n M)_{\theta^n_{A'MN B^n B'^n G^n}} \\ &= S(B^n M)_\theta - S(A'B^n M)_\theta \\ &= S(A'NB'^n G^n)_\theta - S(A'B^n M)_\theta \\ &= S(A'NB'^n G^n)_\theta - S(A'B'^n M)_\theta \\ &\quad + S(A'B'^n M)_\theta - S(A'B^n M)_\theta \\ &= S(G^n|A'B'^n M)_\theta + S(A'B'^n M)_\theta \\ &\quad - S(A'B^n M)_\theta \end{aligned} \quad (\text{D.8})$$

where we abbreviate $\theta = \theta^n_{A'MN B^n B'^n G^n}$. Applying the same technique in the proof of Theorem 2.12 in [19], we can bound

$$S(A'B'^n M)_\theta - S(A'B^n M)_\theta \leq n\Delta. \quad (\text{D.9})$$

Consequently, it follows that

$$\begin{aligned} I_c(A')B^n M)_\omega &\leq S(G^n|A'B'^n M)_\theta + n\Delta \\ &\leq S(G^n|B'^n)_\theta + n\Delta \\ &= S(G^n B'^n)_\theta - S(B'^n)_\theta + n\Delta \\ &\leq S(G^n B'^n)_\theta - S(B^n)_\theta + 2n\Delta \\ &\leq S(G^n B'^n)_\theta - S(A'MN G^n B'^n)_\theta + 2n\Delta \\ &\leq I(A'MN)G^n B'^n)_\theta + 2n\Delta \\ &= I(A^n)B'^n)_{\hat{\rho}_{AB'}^{\otimes n}} + 2n\Delta \\ &= n[I(A)B)_{\hat{\rho}_{AB'}} + 2\Delta] \\ &= 2n\Delta, \end{aligned} \quad (\text{D.10})$$

where the last equality is due to the anti-degradability of $\hat{\rho}_{AB'}$. After dividing Eq. (D.3) by n and taking the limit $n \rightarrow \infty$, we arrive at

$$E_{D,\rightarrow}(\rho_{AB}) = \lim_{n \rightarrow \infty} \frac{1}{n} E_{D,\rightarrow}^{(1)}(\rho_{AB}^{\otimes n}) \leq 2\Delta. \quad (\text{D.11})$$

■

Appendix E: Proof of Proposition 10

Lemma S5 ([84]) A two qubit state ρ_{AB} is anti-degradable if and only if,

$$\mathrm{Tr}(\rho_B^2) \geq \mathrm{Tr}(\rho_{AB}^2) - 4\sqrt{\det(\rho_{AB})}. \quad (\text{E.1})$$

Proposition 10 For a qubit Pauli channel $\Lambda(\cdot)$ with $p_0 \geq p_i > 0 (i = 1, 2, 3)$, it is either anti-degradable or satisfies

$$2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} = (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2 \quad (\text{E.2})$$

with an ADG-squeezed channel as the identity channel.

Proof We first will prove

$$2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \geq (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2, \quad (\text{E.3})$$

by using the SDP in Eq. (B.5). We show that

$$\Gamma_{AB}^{\mathcal{S}} = \frac{\alpha}{2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \quad (\text{E.4})$$

is a feasible solution where $\alpha = 1 - [(\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2]$. Note that the Choi state of the Pauli channel is

$$J_{AB}^{\Lambda} = \frac{1}{2} \begin{pmatrix} p_0 + p_3 & 0 & 0 & p_0 - p_3 \\ 0 & p_1 + p_2 & p_1 - p_2 & 0 \\ 0 & p_1 - p_2 & p_1 + p_2 & 0 \\ p_0 - p_3 & 0 & 0 & p_0 + p_3 \end{pmatrix}, \quad (\text{E.5})$$

and the unnormalized state $\Gamma_{AB}^{\mathcal{N}'} = J_{AB}^{\Lambda} - \Gamma_{AB}^{\mathcal{S}}$ is

$$\Gamma_{AB}^{\mathcal{N}'} = \frac{1}{2} \begin{pmatrix} p_0 + p_3 - \alpha & 0 & 0 & p_0 - p_3 - \alpha \\ 0 & p_1 + p_2 & p_1 - p_2 & 0 \\ 0 & p_1 - p_2 & p_1 + p_2 & 0 \\ p_0 - p_3 - \alpha & 0 & 0 & p_0 + p_3 - \alpha \end{pmatrix}. \quad (\text{E.6})$$

Recalling that $p_0 + p_1 + p_2 + p_3 = 1$, it is then straightforward to check that $\mathrm{Tr}_B[\Gamma_{AB}^{\mathcal{N}'}] = \mathrm{Tr}[\Gamma_{AB}^{\mathcal{N}'}]/d_A \cdot I_A$. The constraint in Eq. (B.5e) corresponds to the anti-degradable condition of $\Gamma_{AB}^{\mathcal{N}'}$. By direct calculation, we have

$$\begin{aligned} \mathrm{Tr}[(\Gamma_B^{\mathcal{N}'})^2] &= \frac{1}{2}(1 - \alpha)^2, \quad \det(\Gamma_{AB}^{\mathcal{N}'}) = p_1 p_2 p_3 (p_0 - \alpha), \\ \mathrm{Tr}[(\Gamma_{AB}^{\mathcal{N}'})^2] &= \alpha^2 - 2\alpha p_0 + p_0^2 + p_1^2 + p_2^2 + p_3^2. \end{aligned} \quad (\text{E.7})$$

Then Eq. (E.1) holds and $\Gamma_{AB}^{\mathcal{N}'}$ is anti-degradable by Lemme S5, which satisfy the constraint in Eq. (B.5e). Thus, we have proven that $\Gamma_{AB}^{\mathcal{S}}$ is a feasible solution to the primal SDP, implying that

$$1 - 2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \leq \mathrm{Tr}[\Gamma_{AB}^{\mathcal{N}'}] = 1 - [(\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2]. \quad (\text{E.8})$$

It yields $2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \geq (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2$.

Second, we will use the dual SDP in Eq. (B.9) to prove

$$2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \leq (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2. \quad (\text{E.9})$$

We show that $\{M_{AB}, N_{AB}, K_{AB}, R_A\}$ is a feasible solution to the dual problem, where

$$M_{AB} = \begin{pmatrix} \eta & 0 & 0 & -\eta + 1 \\ 0 & \xi & \zeta & 0 \\ 0 & \zeta & \xi & 0 \\ -\eta + 1 & 0 & 0 & \eta \end{pmatrix}, N_{AB} = K_{AB} = -\frac{1}{2}M_{AB}, R_A = 0, \quad (\text{E.10})$$

and

$$\eta = -\frac{\sqrt{p_1} + \sqrt{p_2}}{2\sqrt{p_3}}, \xi = -\frac{\sqrt{p_1} + \sqrt{p_3}}{2\sqrt{p_2}} - \frac{\sqrt{p_2} + \sqrt{p_3}}{2\sqrt{p_1}} - 1, \zeta = \frac{\sqrt{p_1} + \sqrt{p_3}}{2\sqrt{p_2}} - \frac{\sqrt{p_2} + \sqrt{p_3}}{2\sqrt{p_1}}. \quad (\text{E.11})$$

It is easy to check that when $p_0 \geq p_i > 0 (i = 1, 2, 3)$, we have $M_{AB} + N_{AB} + K_{AB} = 0$,

$$\begin{aligned} M_{AB} &\leq (1 - \text{Tr } R_A/d_A)I_{AB} + R_A \otimes I_B \Leftrightarrow I_{AB} - M_{AB} \geq 0, \text{ and} \\ N_{AB} \otimes I_E + P_{BE}(K_{AB} \otimes I_E)P_{BE}^\dagger &= \frac{1}{2}M_{AB} \otimes I_E + P_{BE}(\frac{1}{2}M_{AB} \otimes I_E)P_{BE}^\dagger \geq 0. \end{aligned} \quad (\text{E.12})$$

It also satisfies $\text{Tr}[M_{AB}J_{AB}^\Lambda] = 1 - [(\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2]$. Thus we have proven that $\{M_{AB}, N_{AB}, K_{AB}, R_A\}$ is a feasible solution to the dual SDP in Eq. (B.9), which implies

$$1 - 2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \geq \text{Tr}[M_{AB}J_{AB}^\Lambda] = 1 - [(\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2]. \quad (\text{E.13})$$

Then it yields $2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} \leq (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2$. Thus, we arrive at

$$2^{-\tilde{\mathcal{R}}_{\max, \text{ADG}}(\Lambda)} = (\sqrt{p_1} + \sqrt{p_2})^2 + (\sqrt{p_2} + \sqrt{p_3})^2 + (\sqrt{p_1} + \sqrt{p_3})^2. \quad (\text{E.14})$$

Since Γ_{AB}^S is the Bell state after normalization, we know the ADG-squeezed channel is the identity channel. ■