

# Deep Calibration With Artificial Neural Network: A Performance Comparison on Option Pricing Models

Young Shin Kim<sup>a</sup>, Hyangju Kim<sup>b,\*</sup>, Jaehyung Choi<sup>c,\*\*</sup>

<sup>a</sup>College of Business, Stony Brook University, New York, USA

<sup>b</sup>Citibank, Inc. New York, USA

<sup>c</sup>Goldman Sachs & Co., New York, USA

---

## Abstract

This paper explores Artificial Neural Network (ANN) as a model-free solution for a calibration algorithm of option pricing models. We construct ANNs to calibrate parameters for two well-known GARCH-type option pricing models: Duan's GARCH and the classical tempered stable GARCH that significantly improve upon the limitation of the Black-Scholes model but have suffered from computation complexity. To mitigate this technical difficulty, we train ANNs with a dataset generated by Monte Carlo Simulation (MCS) method and apply them to calibrate optimal parameters. The performance results indicate that the ANN approach consistently outperforms MCS and takes advantage of faster computation times once trained. The Greeks of options are also discussed.

*JEL classification:* C15, C63, C65, G130

*Keywords:* Deep calibration, Artificial Neural Network (ANN), Feedforward Neural Network (FNN), Option pricing models, GARCH model, Duan's GARCH model, Tempered stable model

---

\*Disclaimer: The views, information, or opinions expressed in this publication are solely those of Hyangju Kim and do not represent those of Citigroup, its respective affiliates, or employees.

\*\*Disclaimer: The opinions and statements expressed in this article are those of the authors. None of Goldman Sachs or their respective affiliates offer, or are otherwise the source of, any of the opinions or statements set out in this article (which may be different to views or opinions held or expressed by or within Goldman Sachs). The content of this article is for information purposes only and is not investment advice (or advice of any other kind). None of the authors, Goldman Sachs, or their respective affiliates, officers, employees, or representatives accepts any liability whatsoever in connection with any of the content of this article or for any action or inaction of any person taken in reliance upon such content (or any part thereof).

*Email addresses:* aaron.kim@stonybrook.edu (Young Shin Kim), hyangju.kim@gmail.com (Hyangju Kim), jj.jaehyung.choi@gmail.com (Jaehyung Choi)

## 1. Introduction

Option pricing has been a centric topic in quantitative finance for the past few decades, with remarkable growth in option market. Since the seminal work of Black and Scholes (1973) and Merton (1973), the Black-Scholes model has remained the most fundamental model for option pricing. However, its restrictive assumptions, such as constant volatility or Geometric Brownian Motion (GBM), have been criticized for not reflecting the empirical characteristics of financial markets.

Many subsequent models have since been proposed to relax the assumptions of the Black-Scholes model. One successful approach is employing stochastic volatility under the generalized autoregressive conditional heteroskedastic (GARCH) framework. The early attempt was introduced by Engle and Mustafa (1992) focusing on implied conditional volatilities. Subsequently, Duan (1995) developed a more rigorous framework of the GARCH option pricing model using the locally risk-neutral valuation relationship that one-period ahead conditional variance remains constant under both the risk-neutral measure and the physical measure. The model extended later to the jump-diffusion model in Duan *et al.* (2004) and Duan *et al.* (2006). Duan's GARCH model assumes that the residuals follow a normal distribution, however, empirical evidence shows that the assumption of a normally distributed residual in the model does not properly describe asset return dynamics (Duan (1999) and Menn and Rachev (2009)).

Another approach, known as the Lévy distribution, has been developed allowing for jumps, skewness, kurtosis, and heavy tails in underlying distribution to overcome the flaws of the GBM assumption. It is also referred to as a stable distribution, and its one well-known subclass is the classical tempered stable (CTS) distribution.<sup>1</sup> The CTS distribution has been researched in several directions such as portfolio management (Tsuchida *et al.* (2012); Beck *et al.* (2013); Georgiev *et al.*

---

<sup>1</sup>The other subclasses are the classical tempered stable process, generalized classical tempered stable process, modified tempered stable process, normal tempered stable process, Kim-Rachev tempered stable process, and rapidly decreasing tempered stable process.

(2015); Anand *et al.* (2016); Choi *et al.* (2021)) and momentum strategy (Choi *et al.* (2015)).

As a consolidation of these two approaches in the context of option pricing, Kim *et al.* (2008a), Kim *et al.* (2010a) and Kim *et al.* (2022) enhanced Duan's GARCH model by incorporating the classical tempered stable (CTS) distribution<sup>2</sup> and referred to it as the CTS-GARCH model. The CTS-GARCH model, which takes into account the non-normality in its innovation process, is considered one of the most advanced option pricing models as it addresses two key limitations of the Black-Scholes model simultaneously (Kim *et al.* (2010b)).

However, these models are mostly high-dimensional, and this becomes problematic in the stage of parameter calibration; parameters need to be tuned, so as to find the best model output closest to the market price. Hence the Black-Scholes model still has been a practically useful model even with the superior performance of multidimensional models.

Recently, machine learning has shed a different light on the curse of dimensionality as a non-parametric or model-free solution. This is a rapidly evolving area with the maturation of computation power and technical advances in algorithms.

One of the main pillars of machine learning is Artificial Neural Network (ANN). ANN, also referred to as neural network or multilayer perceptions, has been introduced by McCulloch and Pitts (1943). ANNs are composed of the input layer, multiple hidden layers, and the output layer where each hidden layer performs a vector-to-vector or vector-to-scalar calculation to make the best output layer approximation. The parallel structure of hidden layers allows using multiprocessors, hence ANN is considerably faster than traditional algorithms at computation speed. With this advantage, ANN has been developed rapidly and different types of ANN have been proposed such as multilayer perceptron neural network, convolutional neural network, radial basis function neural network, recurrent neural network, modular neural network, and so on.

---

<sup>2</sup>The CTS distribution has been studied under different names including the *truncated Lévy flight* by Koponen (1995), the *tempered stable* by Barndorff-Nielsen and Levendorskii (2001), Barndorff-Nielsen and Shephard (2001) and Cont and Tankov (2004), the *KoBoL* distribution by Boyarchenko and Levendorskiĭ (2000), and the *CGMY* by Carr *et al.* (2002). The KR distribution of Kim *et al.* (2008b) is an extension of the CTS distribution. Rosiński (2007) generalized CTS distribution, referred to as the tempered stable distribution.

Many studies in finance also sought to find its ANN applications. In particular, ANN receives attention as a promising alternative for chronic computational difficulties in option pricing models as it is built upon multidimensional nonlinear models. The early attempts were applying ANN as a functional approximator for the Black-Scholes formula suggested by Malliaris and Salchenberger (1993) and Hutchinson *et al.* (1994). Since then, over a hundred papers have studied ANN for option pricing with various parameter inputs and performance measures (Ruf and Wang (2019)).

While most of the studies utilize ANN to estimate the option price based on parameters from the market data and measure their performance by out-of-sample tests, a different approach was proposed by Bayer *et al.* (2019) and Alaya *et al.* (2021), recently. Rather than focusing on the approximation of the pricing formula, ANN is applied to the calibration stage and finds model parameters in this approach. Since training sets are generated by Monte Carlo Simulation (MCS) method, hence a more sound dataset avoiding market incompleteness is available. Utilizing enough large datasets is another advantage.

This paper also focuses on ANN as an alternative calibration method. We generate training sets having 100,000 simulated vectors of parameters and prices using the MCS method for each of Duan's GARCH and CTS-GARCH models, and then train ANNs that consist of three hidden layers with twenty nodes for each layer. Specifically, a feedforward neural network is used, which does not include any cycles or loops. The trained ANNs are utilized in the calibration for S&P 500 index call and put option prices from every second Wednesday of each month between June 2021 and May 2022. We find that this approach not only presents a superior performance to the previous MCS method but performs remarkably faster. Additionally, the option Greeks are also computed by using the ANN method.

The remainder of this paper is organized as follows. We discuss the GARCH option pricing models in Section 2. Section 3 presents the construction and training methods for ANN. The performance of ANN in terms of calibration is investigated using empirical data in Section 4. Finally, Section 5 provides our conclusions.

## 2. Preliminaries

In this section, we revisit Duan's GARCH and CTS-GARCH option pricing models that feature *infinitely divisible* innovations. An infinitely divisible random variable is a random variable that can be represented as an infinite sum of independent, identically distributed random variables. This concept plays a significant role in the study of stable distributions and option pricing models.

We begin by reviewing the CTS distribution as an example of an infinitely divisible distribution, followed by an overview of Duan's GARCH and CTS-GARCH option pricing models. Lastly, we outline European call and put option pricing using the MCS method for both GARCH option pricing models.

### 2.1. CTS Distribution

$X$  is referred to as the *Classical Tempered Stable distributed random variable* and denoted by  $X \sim \text{CTS}(\alpha, C, \lambda_+, \lambda_-, m)$  (Rachev *et al.* (2011) and Kim *et al.* (2010b)) if the characteristic function of the distribution is expressed as

$$\begin{aligned} E[e^{iuX}] &= \phi_{\text{CTS}}(u; \alpha, C, \lambda_+, \lambda_-, m) \\ &= \exp\left((m - C\Gamma(1 - \alpha)(\lambda_+^{\alpha-1} - \lambda_-^{\alpha-1}))iu - C\Gamma(-\alpha)\left((\lambda_+ - iu)^\alpha - \lambda_+^\alpha + (\lambda_- + iu)^\alpha - \lambda_-^\alpha\right)\right) \end{aligned} \quad (1)$$

where  $C_+, C_-, \lambda_+, \lambda_-$  are positive,  $0 < \alpha < 2$ ,  $m \in \mathbb{R}$  and  $\Gamma$  is the gamma function.

If we substitute  $C = (\Gamma(2 - \alpha)(\lambda_+^{\alpha-2} + \lambda_-^{\alpha-2}))^{-1}$  and  $m = 0$  for  $Z \sim \text{CTS}(\alpha, C, \lambda_+, \lambda_-, m)$ , then we have  $E[Z] = 0$  and  $\text{var}(Z) = 1$ . In this case, the random variable  $Z$  is called the *standard CTS distribution*, and  $Z \sim \text{stdCTS}(\alpha, \lambda_+, \lambda_-)$ . The characteristic function of  $Z$  is given by

$$\begin{aligned} E[e^{iuZ}] &= \phi_{\text{stdCTS}}(u; \alpha, \lambda_+, \lambda_-) \\ &= \exp\left(\frac{\lambda_+^{\alpha-1} - \lambda_-^{\alpha-1}}{(\alpha - 1)(\lambda_+^{\alpha-2} + \lambda_-^{\alpha-2})}iu + \frac{(\lambda_+ - iu)^\alpha - \lambda_+^\alpha + (\lambda_- + iu)^\alpha - \lambda_-^\alpha}{\alpha(\alpha - 1)(\lambda_+^{\alpha-2} + \lambda_-^{\alpha-2})}\right). \end{aligned} \quad (2)$$

We denote the function of log-Laplace transform of  $Z$  as follows (Kim *et al.* (2010b)):

$$\begin{aligned} l(x) &:= \log(\phi_{\text{stdCTS}}(-ix; \alpha, \lambda_+, \lambda_-)) \\ &= \frac{x(\lambda_+^{\alpha-1} - \lambda_-^{\alpha-1})}{(\alpha-1)(\lambda_+^{\alpha-2} + \lambda_-^{\alpha-2})} + \frac{(\lambda_+ - x)^\alpha - \lambda_+^\alpha + (\lambda_- + x)^\alpha - \lambda_-^\alpha}{\alpha(\alpha-1)(\lambda_+^{\alpha-2} + \lambda_-^{\alpha-2})}. \end{aligned} \quad (3)$$

## 2.2. Duan's GARCH and CTS-GARCH Option Pricing Models

Let  $(S_t)_{t \in \{0,1,\dots,T^*\}}$  be the underlying asset price process and  $(y_t)_{t \in \{0,1,2,\dots,T^*\}}$  be the underlying asset log return process, where  $y_t = \log(\frac{S_t}{S_{t-1}})$  with  $y_0 = 0$ , and  $T^* < \infty$  in the time horizon. Under the physical measure  $\mathbb{P} = \bigoplus_{t=1}^{T^*} \mathcal{P}_t$ ,  $(y_t)_{t \in \{0,1,2,\dots,T^*\}}$  is supposed to follow the GARCH model:

$$\begin{cases} y_{t+1} = \mu_{t+1} + \sigma_{t+1}\epsilon_{t+1} \\ \sigma_{t+1}^2 = \kappa + \xi\sigma_t^2\epsilon_t^2 + \zeta\sigma_t^2 \end{cases} \quad \text{for } t \in \{0, 1, 2, \dots, T^* - 1\}, \quad (4)$$

where  $\mu_{t+1}$  is the daily expected return and  $\epsilon_{t+1}$  follows an infinitely divisible distribution.  $S_0$ ,  $\epsilon_0$  and  $\sigma_0$  are real constants with  $\epsilon_0 = 0$ . The GARCH model involves parameters with  $\kappa$ ,  $\xi$ ,  $\zeta$  where  $\xi + \zeta < 1$ .

We next define  $(R_t)_{t \in \{1,2,\dots,T^*\}}$  and  $(d_t)_{t \in \{1,2,\dots,T^*\}}$  as the sequences of the daily risk-free rate of return and daily dividend rate of the underlying, respectively. Then, there is a risk-neutral measure  $\mathbb{Q} = \bigoplus_{t=1}^{T^*} \mathcal{Q}_t$  such that:

- $\eta_{t+1} = \theta_{t+1} + \epsilon_{t+1}$ ,  
where  $\theta_{t+1} = \frac{\mu_{t+1} - R_{t+1} + d_{t+1} + \omega_{t+1}}{\sigma_{t+1}}$  is the market price of risk and  $\omega_{t+1} = \log E_{\mathbb{Q}}[e^{\sigma_{t+1}\epsilon_{t+1}}]$ .
- $\eta_{t+1}$  is also infinitely divisible under the measure  $\mathbb{Q}$ .

By applying change of measures to Eq. (4), we obtain the risk-neutral price process under  $\mathbb{Q}$  as

$$\begin{cases} y_{t+1} = R_{t+1} - d_{t+1} - \omega_{t+1} + \sigma_{t+1}\eta_{t+1} \\ \sigma_{t+1}^2 = \kappa + \xi\sigma_t^2(\eta_t - \theta_t)^2 + \zeta\sigma_t^2 \end{cases} \quad \text{for } t \in \{0, 1, 2, \dots, T^* - 1\} \quad (5)$$

with  $\xi + \zeta < 1$ . To simplify the condition  $\xi + \zeta < 1$ , we define two parameters  $\psi$  and  $\gamma$  as  $\psi = \frac{\xi}{\zeta}$  and  $\gamma = \xi + \zeta$ , respectively. Then we have

$$\sigma_t^2 = \kappa + \frac{\gamma}{\psi + 1} (\psi \sigma_{t-1}^2 (\eta_{t-1} - \theta_t)^2 + \sigma_{t-1}^2), \quad (6)$$

where  $\kappa, \psi, \gamma > 0, \eta_0 = 0$  and  $\sigma_0 > 0$ .

Under the risk-neutral measure  $\mathbb{Q}$ , the underlying asset price is defined as  $S_t = S_0 e^{\sum_{j=1}^t y_j}$  for  $t \in \{1, 2, \dots, T-1, T, \dots, T^*\}$ . The European option with a payoff function  $H(S(T))$  at the maturity  $T$  with  $t \leq T \leq T^*$  is given by

$$E_{\mathbb{Q}} \left[ e^{-\sum_{j=t+1}^T R_j} H(S(T)) \mid \mathcal{F}_t \right] = E_{\mathbb{Q}} \left[ e^{-\sum_{j=t+1}^T R_j} H(S_t e^{\sum_{j=t+1}^T y_j}) \mid \mathcal{F}_t \right]. \quad (7)$$

For example, European vanilla call and put price with strike price  $K$  and time to maturity  $T$  at time  $t = 0$  are

$$\text{(Call)} = E_{\mathbb{Q}} \left[ e^{-\sum_{j=1}^T R_j} \max\{S_0 e^{\sum_{j=1}^T y_j} - K, 0\} \right] \quad (8)$$

and

$$\text{(Put)} = E_{\mathbb{Q}} \left[ e^{-\sum_{j=1}^T R_j} \max\{K - S_0 e^{\sum_{j=1}^T y_j}, 0\} \right], \quad (9)$$

respectively.

Moreover, let  $m$  be the moneyness defined as  $m = \frac{K}{S_0 e^{\sum_{t=1}^T R_t}}$ , then we have

$$\text{(Call)} = S_0 V_C \text{ and } \text{(Put)} = S_0 V_P, \quad (10)$$

where

$$V_C = E_{\mathbb{Q}} \left[ \max \left\{ \exp \left( \sum_{t=1}^T -d_t - w_t + \sigma_t \eta_t \right) - m, 0 \right\} \right] \quad (11)$$

and

$$V_P = E_{\mathbb{Q}} \left[ \max \left\{ m - \exp \left( \sum_{t=1}^T -d_t - w_t + \sigma_t \eta_t \right), 0 \right\} \right]. \quad (12)$$

Now, we present two popular examples of the GARCH option pricing model with infinitely divisible innovations:

- *Duan's GARCH Model* (Duan (1995)): If we assume that  $\eta_t$ 's follow the standard Gaussian distribution, which is infinitely divisible, we obtain the Duan's GARCH option pricing model where  $w_t = \frac{\sigma_t^2}{2}$ .
- *CTS-GARCH Model* (Kim *et al.* (2010a)): By assuming that  $\eta_t$ 's follow the standard CTS distribution, which is also infinitely divisible, we refer to the GARCH model as the CTS-GARCH option pricing model. Specifically, when we set  $\eta_t \sim \text{stdCTS}(\alpha, \lambda_+, \lambda_-)$  for all  $t \in \{1, 2, \dots\}$  under the measure  $\mathbb{Q}$ , we have

$$w_t = \log (\phi_{\text{stdCTS}}(-i\sigma_t; \alpha, \lambda_+, \lambda_-)) = l(\sigma_t) \quad (13)$$

by Eq. (3).

### 2.3. European Call and Put Option Pricing with the MCS Method

To simplify the model, we make the following assumptions for the remainder of this paper: the daily risk-free return  $R_t$  is a constant  $R$ ,  $\theta_t$  is a constant  $\theta$ , and  $d_t = 0$ . We define  $t \in \{1, 2, \dots, T, \dots, T^*\}$  for  $0 < T \leq T^*$  as a set of the time steps, where one step represents one business day. We also assume that there are 250 business days in a year, and we define the annual risk-free return as  $r = 250 \cdot R$ . The year-fraction time is denoted by  $\tau = \frac{T}{250}$ , and we define  $m$  as the moneyness,



where  $m = \frac{Ke^{-RT}}{S_0} = \frac{Ke^{-r\tau}}{S_0}$ .

Based on these assumptions, we can generate a set of infinitely divisible random numbers  $\{\eta_{t,n} : t \in \{1,2,\dots, T^*\}, n \in \{1, 2, \dots, N\}\}$  using the MCS method. Next, we apply the GARCH model to obtain the set of volatility  $\{\sigma_{t,n} : t \in \{1, 2, \dots, T^*\}, n \in \{1, 2, \dots, N\}\}$  defined as

$$\sigma_{t,n} = \sqrt{\kappa + \frac{\gamma}{\psi + 1} (\psi\sigma_{t-1,n}^2(\eta_{t-1} - \theta)^2 + \sigma_{t-1,n}^2)}, \quad (14)$$

where  $\kappa, \psi, \gamma > 0$ ,  $\eta_0 = 0$  and  $\sigma_{0,n} = \sigma_0$  for a constant  $\sigma_0$ .

For a given time to maturity  $T \leq T^*$  and a moneyness  $m$ , we approximate  $V_C$  and  $V_P$  as follows:

$$V_C \approx \hat{V}_C = \frac{1}{N} \sum_{n=1}^N \max \left\{ \exp \left( \sum_{t=1}^{250\tau} -w_{t,n} + \sigma_{t,n}\eta_{t,n} \right) - m, 0 \right\} \quad (15)$$

and

$$V_P \approx \hat{V}_P = \frac{1}{N} \sum_{n=1}^N \max \left\{ m - \exp \left( \sum_{t=1}^{250\tau} -w_{t,n} + \sigma_{t,n}\eta_{t,n} \right), 0 \right\}. \quad (16)$$

From (15) and (16), call and put option values in Duan's GARCH and CTS-GARCH models can be calculated as follows:

- *Duan's GARCH model:* We generate a set of standard Gaussian random numbers  $\{\eta_{t,n} : t \in \{1,2,\dots, T^*\}, n \in \{1,2,\dots, N\}\}$  and set  $w_{t,n} = \frac{\sigma_{t,n}^2}{2}$ . Then we obtain call and put option prices of Duan's GARCH model. In this case, we denote (15) and (16) as

$$V_C^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0) = \hat{V}_C \quad \text{and} \quad V_P^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0) = \hat{V}_P, \quad (17)$$

respectively. The call and put option prices can be approximated using MCS as

$$\text{Call}^{Duan}(S_0, K, \tau, r) \approx S_0 V_C^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0) \quad (18)$$

and

$$\text{Put}^{Duan}(S_0, K, \tau, r) \approx S_0 V_P^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0). \quad (19)$$

- *CTS-GARCH model:* We simulate a set of standard CTS random numbers  $\{\eta_{t,n} : t \in \{1, 2, \dots, T^*\}, n \in \{1, 2, \dots, N\}\}$  with parameters  $(\alpha, \lambda_+, \lambda_-)$ . By setting  $w_{t,n} = l(\sigma_{t,n})$ , we obtain call and put option prices of the CTS-GARCH model. Specifically, we use (15) and (16) to denote the call and put option values as

$$V_C^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \hat{V}_C \quad \text{and} \quad V_P^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \hat{V}_P, \quad (20)$$

respectively. Using MCS, we can obtain call and put option prices under the CTS-GARCH model such that

$$\text{Call}^{CTS}(S_0, K, \tau, r) \approx S_0 V_C^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) \quad (21)$$

and

$$\text{Put}^{CTS}(S_0, K, \tau, r) \approx S_0 V_P^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-). \quad (22)$$

### 3. Artificial Neural Network

In this section, we construct two ANNs to calculate call and put option prices under Duan's GARCH model and CTS-GARCH model. More precisely, we design multi-layer ANNs to gen-

erate similar results as the function values of  $V_C^{Duan}$ ,  $V_P^{Duan}$ ,  $V_C^{CTS}$ , and  $V_P^{CTS}$ . To facilitate the training, we take the logarithm of these four functions and define new functions as follows:

$$v_C^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \log (V_C^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)) \quad (23)$$

$$v_P^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \log (V_P^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)) \quad (24)$$

$$v_C^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \log (V_C^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)) \quad (25)$$

$$v_P^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-) = \log (V_P^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)) \quad (26)$$

### 3.1. Generating Training Set

We first generate training sets for the four functions  $v_C^{Duan}$ ,  $v_P^{Duan}$ ,  $v_C^{CTS}$ , and  $v_P^{CTS}$  using the MCS method as explained in the previous section. Since  $v_C^{Duan}$  and  $v_P^{Duan}$  have seven input parameters ( $m, \tau, \kappa, \psi, \gamma, \theta, \sigma_0$ ), we consider seven nodes in the input layer. For  $v_C^{CTS}$ , and  $v_P^{CTS}$ , three additional input parameters ( $\alpha, \lambda_+, \lambda_-$ ) are needed, in addition to the seven input parameters for  $v_C^{Duan}$  and  $v_P^{Duan}$ . The range of the input parameters can be found in Table 1. More details on training set generation process for each model are as follows:

- *Duan's GARCH Model:* We generate 100,000 seven-dimensional uniformly distributed random vectors for  $(m, \tau, \kappa, \psi, \gamma, \lambda, \sigma_0)$  with the boundary specified in Table 1. To avoid the clustering of the random vector, we use the Halton algorithm (Halton (1964)). Then we calculate 100,000 of  $v_C^{Duan}$  and  $v_P^{Duan}$ , respectively, using MCS. In MCS, we use 20,000 sample paths based on the Duan's GARCH model with the seven-dimensional model parameters.
- *CTS-GARCH Model:* We generate 100,000 ten-dimensional uniformly distributed random vectors for  $(m, \tau, \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)$  with the boundary specified in Table 1. Random numbers of  $\lambda_+$  and  $\lambda_-$  are generated by  $\lambda_+ = \tan(\frac{u_1\pi}{2})+0.1$  and  $\lambda_- = \lambda_+ = \tan(\frac{u_2\pi}{2})+0.1$ , respectively, for uniform random numbers  $u_1, u_2 \in (0, 1)$ . We also use the Halton algorithm to generate uniform random vectors, as we did in the case of Duan's GARCH model. Then we calculate 100,000 of  $v_C^{CTS}$  and  $v_P^{CTS}$ , respectively, using the MCS. In the MCS, we use

Parameter	Lower Bound	Upper Bound
$m$	0.5	1.5
$\tau$	0.4	1
$\kappa$	0	$1 \cdot 10^{-5}$
$\psi$	0.1	0.4
$\gamma$	0.5	0.9999
$\theta$	0	0.8
$\sigma_0$	$1 \cdot 10^{-6}$	0.04
$\alpha$	0.01	1.999
$\lambda_+$	0.1	$\infty$
$\lambda_-$	0.1	$\infty$

Table 1: The range of input parameters

20,000 sample paths based on the CTS-GARCH model with the ten-dimensional model parameters.

### 3.2. Training Multi-Layer ANNs

Using four training sets for  $v_C^{Duan}$ ,  $v_P^{Duan}$ ,  $v_C^{CTS}$ , and  $v_P^{CTS}$ , we train four multi-layer ANNs in this section. Each ANN consisted of three hidden layers with twenty nodes in each layer. The output is a single value, and the activation function of hidden layer nodes is the simple sigmoid function<sup>3</sup>, while the output activation function is the linear function<sup>4</sup>. The ANNs for  $v_C^{Duan}$  and  $v_P^{Duan}$  have seven input nodes, while the ANNs for  $v_C^{CTS}$  and  $v_P^{CTS}$  have ten input nodes. The architecture is depicted in Fig. 1. We denote the four ANNs corresponding to  $v_C^{Duan}$ ,  $v_P^{Duan}$ ,  $v_C^{CTS}$ , and  $v_P^{CTS}$  as

$$F_C^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0), \quad (27)$$

$$F_P^{Duan}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0), \quad (28)$$

$$F_C^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-), \quad (29)$$

$$\text{and } F_P^{CTS}(m, \tau; \kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-). \quad (30)$$

---

<sup>3</sup> $f(x) = \frac{1}{1+e^{-x}}$   
<sup>4</sup> $f(x) = x$




Fig. 1: ANN structure of the calibration

Additionally, we set the parameters  $\Theta_{Duan} = (\kappa, \psi, \gamma, \theta, \sigma_0)$  of Duan's GARCH model and  $\Theta_{CTS} = (\kappa, \psi, \gamma, \theta, \sigma_0, \alpha, \lambda_+, \lambda_-)$  of CTS-GARCH model. Then, we have

$$\text{Call}_{ANN}^{Duan}(S_0, K, \tau, r) = S_0 \exp \left( F_C^{Duan} \left( \frac{K e^{-r\tau}}{S_0}, \tau; \Theta_{Duan} \right) \right) \quad (31)$$

$$\text{Put}_{ANN}^{Duan}(S_0, K, \tau, r) = S_0 \exp \left( F_P^{Duan} \left( \frac{K e^{-r\tau}}{S_0}, \tau; \Theta_{Duan} \right) \right) \quad (32)$$

$$\text{Call}_{ANN}^{CTS}(S_0, K, \tau, r) = S_0 \exp \left( F_C^{CTS} \left( \frac{K e^{-r\tau}}{S_0}, \tau; \Theta_{CTS} \right) \right) \quad (33)$$

$$\text{Put}_{ANN}^{CTS}(S_0, K, \tau, r) = S_0 \exp \left( F_P^{CTS} \left( \frac{K e^{-r\tau}}{S_0}, \tau; \Theta_{CTS} \right) \right). \quad (34)$$

Subsequently, we obtain

$$\text{Call}^{Duan}(S_0, K, \tau, r) \approx \text{Call}_{ANN}^{Duan}(S_0, K, \tau, r), \quad (35)$$

$$\text{and Put}^{Duan}(S_0, K, \tau, r) \approx \text{Put}_{ANN}^{Duan}(S_0, K, \tau, r), \quad (36)$$

under Duan’s GARCH model, and

$$\text{Call}^{CTS}(S_0, K, \tau, r) \approx \text{Call}_{ANN}^{CTS}(S_0, K, \tau, r), \quad (37)$$

$$\text{and Put}^{CTS}(S_0, K, \tau, r) \approx \text{Put}_{ANN}^{CTS}(S_0, K, \tau, r), \quad (38)$$

under CTS-GARCH model.

We train the four ANNs using functions using the Deep Learning toolbox in Matlab<sup>TM</sup>. The default training algorithm for a function fitting network is Levenberg-Marquardt, as stated in the documentation<sup>5</sup>. The mean squared error (MSE) is presented as the performance measure of the network.

Fig. 2 exhibits MSE for the epoch during the training of the four ANNs:  $F_C^{Duan}$ ,  $F_P^{Duan}$ ,  $F_C^{CTS}$ , and  $F_P^{CTS}$ . The minimum values of MSEs for  $F_C^{Duan}$  and  $F_P^{Duan}$  are 0.0169 and 0.0061 at epochs 84 and 178, respectively. For  $F_C^{CTS}$  and  $F_P^{CTS}$ , the training was limited to 300 epochs, with minimum MSEs being 0.0541 and 0.0447 at epoch 300, respectively.

It is also noteworthy that the ANNs in this study are not intended for forecasting, but rather for finding better analytic approximations. Therefore, we do not utilize a validation set.

#### 4. Calibration

In this section, we discuss the calibration of parameters using two different methods: the ANN method and the MCS method, with the MCS method serving as a benchmark method. We calibrate the parameters  $\Theta_{Duan}$  and  $\Theta_{CTS}$  using call and put option data of the S&P 500 index. For this investigation, we have selected 12 Wednesdays from the second week of each month, ranging from June 2021 to May 2022. The time to maturity of the option contracts varies between 7 and 90 days, and we exclude the options with zero bid prices or zero ask prices. Furthermore, we only calibrate parameters for out-of-the-money (OTM) options for both calls and puts.

---

<sup>5</sup><https://www.mathworks.com/help/deeplearning/ref/fitnet.html>




Fig. 2: Number of epochs and mean squared error. Top left and right are the MSE of call and put prices for Duan's GARCH model. Bottom left and right are the MSE of the CTS-GARCH model.

We first specify ANNs for OTM options as

$$F_{OTM}^{Duan}(m, \tau; \Theta_{Duan}) = F_P^{Duan}(m, \tau; \Theta_{Duan}) \cdot 1_{m < 1} + F_C^{Duan}(m, \tau; \Theta_{Duan}) \cdot 1_{m \geq 1} \quad (39)$$

and

$$F_{OTM}^{CTS}(m, \tau; \Theta_{CTS}) = F_P^{CTS}(m, \tau; \Theta_{CTS}) \cdot 1_{m < 1} + F_C^{CTS}(m, \tau; \Theta_{CTS}) \cdot 1_{m \geq 1}. \quad (40)$$

We calibrate parameters using the relative root mean square error (rel-RMSE) minimization method as follows:

$$\min_{\Theta_{Duan}} \left( \sum_{K_n e^{-rT_n} < S_0} \left( \frac{F_{OTM}^{Duan} \left( \frac{K_n e^{-rT_n}}{S_0}, \frac{T_n}{250}; \Theta_{Duan} \right) - \log \left( \frac{P_{market}(K_n, T_n)}{S_0} \right)}{\log \left( \frac{P_{market}(K_n, T_n)}{S_0} \right)} \right)^2 + \sum_{K_n e^{-rT_n} \geq S_0} \left( \frac{F_{OTM}^{Duan} \left( \frac{K_n e^{-rT_n}}{S_0}, \frac{T_n}{250}; \Theta_{Duan} \right) - \log \left( \frac{C_{market}(K_n, T_n)}{S_0} \right)}{\log \left( \frac{C_{market}(K_n, T_n)}{S_0} \right)} \right)^2 \right)^{\frac{1}{2}} \quad (41)$$

and

$$\min_{\Theta_{CTS}} \left( \sum_{K_n e^{-rT_n} < S_0} \left( \frac{F_{OTM}^{CTS} \left( \frac{K_n e^{-rT_n}}{S_0}, \frac{T_n}{250}; \Theta_{CTS} \right) - \log \left( \frac{P_{market}(K_n, T_n)}{S_0} \right)}{\log \left( \frac{P_{market}(K_n, T_n)}{S_0} \right)} \right)^2 + \sum_{K_n e^{-rT_n} \geq S_0} \left( \frac{F_{OTM}^{CTS} \left( \frac{K_n e^{-rT_n}}{S_0}, \frac{T_n}{250}; \Theta_{CTS} \right) - \log \left( \frac{C_{market}(K_n, T_n)}{S_0} \right)}{\log \left( \frac{C_{market}(K_n, T_n)}{S_0} \right)} \right)^2 \right)^{\frac{1}{2}}, \quad (42)$$

where  $S_0$  is the S&P 500 index price of the given Wednesday, and  $C_{market}(K_n, T_n)$  and  $P_{market}(K_n, T_n)$  are mid-prices of observed bid and ask prices for the call and put options with strike price  $K_n$  and



time to maturity  $T_n$ <sup>6</sup>.

As a benchmark, we use the MCS method to calibrate the same parameters. Similar to the ANN cases, we define

$$v_{OTM}^{Duan}(m, \tau; \Theta_{Duan}) = v_C^{Duan}(m, \tau; \Theta_{Duan}) \cdot 1_{m \geq 1} + v_P^{Duan}(m, \tau; \Theta_{Duan}) \cdot 1_{m < 1} \quad (43)$$

and

$$v_{OTM}^{CTS}(m, \tau; \Theta_{CTS}) = v_C^{CTS}(m, \tau; \Theta_{CTS}) \cdot 1_{m \geq 1} + v_P^{CTS}(m, \tau; \Theta_{CTS}) \cdot 1_{m < 1}. \quad (44)$$

We calibrate parameters using the MCS method as described in (41) and (42) by replacing  $F_{OTM}^{Duan}$  and  $F_{OTM}^{CTS}$  with  $v_{OTM}^{Duan}$  and  $v_{OTM}^{CTS}$ , respectively. More details of the parameter calibration for the option pricing with the CTS-GARCH model are presented in Kim *et al.* (2010a), Kim *et al.* (2019), and Kim *et al.* (2022).

The calibrated parameters for Duan's GARCH model are provided in Table 2 and Table 3, while those for the CTS-GARCH model are provided in Table 4 and Table 5. Table 3 and 5 present the results obtained using the ANN method, while Table 2 and Table 4 present the results obtained using the MCS method. The rel-RMSE values are presented for performance analysis. Table 6 collects all rel-RMSEs for the two models (Duan's GARCH and CTS-GARCH) and both option pricing methods (MCS and ANN).

According to the table, we observe that:

- CTS-GARCH model with the ANN method shows the smallest rel-RMSE and therefore performed the best, with the exception of four cases on 11/10/2022, 12/8/2021, 3/9/2022, and 5/10/2022.

---

<sup>6</sup>To solve the nonlinear optimization problems, we used the function `lsqcurvefit()` in Matlab<sup>TM</sup> with Trust-Region-Reflective Least Squares algorithm. See Coleman and Li (1994, 1996) and <https://www.mathworks.com/help/optim/ug/lsqcurvefit.html> for the details.

- Duan’s GARCH model with the MCS method has the smallest rel-RMSE for the cases on 3/8/2022 and 5/10/2022.
- Duan’s GARCH model with the ANN method has the smallest rel-RMSE for the cases on 11/10/2022 and 12/8/2022.
- CTS-GARCH model with the MCS method has the largest rel-RMSE values in this investigation.

Compared to Duan’s GARCH model, CTS-GARCH model is more flexible as it has three more parameters. However, the complexity of the CTS-GARCH model makes model calibration with the MCS method inefficient. In this regard, the ANN offers an efficient alternative for model calibration, enabling the use of the CTS-GARCH model in practical applications.

In Fig. 3, the left column plots present the market prices and calibrated model prices of the Duan’s GARCH model and CTS-GARCH model for OTM calls and puts on 6/9/2021. The right column plots exhibit the implied volatility curves for the market prices and Duan’s GARCH and CTS-GARCH models with respect to the MCS method and ANN method. The first row plots of Fig. 3 show option prices and implied volatility for 9 days to maturity, the second row plots show 37 days to maturity, and the third row plots show 72 days to maturity. The curves suggest that the implied volatility curve of CTS-GARCH model prices with the ANN method is the closest to the market implied volatility curve compared to the other methods investigated in this study.

One of the advantages of using the ANN method is the ability to easily obtain the Greeks of options, which is not almost possible with the MCS method. For instance, Table 7 exhibits Delta ( $\Delta$ ), Gamma ( $\Gamma$ ), Theta ( $\Theta$ ) and Rho ( $\rho$ ) of at-the-money (ATM) calls and puts for the parameters calibrated on 6/9/2021 with maturities of 7, 37, and 72 days, respectively<sup>7</sup>. The Vega ( $\nu$ ) is not

---

<sup>7</sup>Delta ( $\Delta = \frac{\partial V}{\partial S}$ ): the sensitivity of an option’s price( $V$ ) changes relative to the underlying price( $S$ ) changes.

Gamma ( $\Gamma = \frac{\partial^2 V}{\partial S^2}$ ): the sensitivity of the Delta’s changes relative to the underlying price changes.

Theta ( $\Theta = \frac{\partial V}{\partial \tau}$ ): the sensitivity of the option price relative to the option’s time to maturity( $\tau$ ).

Rho ( $\rho = \frac{\partial V}{\partial r}$ ): the sensitivity of the option price relative to the risk-free rate of return ( $r$ ).

Vega ( $\nu = \frac{\partial V}{\partial \sigma}$ ): the sensitivity of an option’s price changes relative to the underlying asset’s volatility.

Date	$\theta$	$\kappa$	$\xi$	$\zeta$	$\sigma_0$	rel-RMSE
6/9/2021	0.9293	$1.68 \cdot 10^{-6}$	0.2778	0.5000	0.0085	0.3048
7/7/2021	0.8984	$1.21 \cdot 10^{-6}$	0.2931	0.5000	0.0068	0.3585
8/11/2021	0.9513	$1.05 \cdot 10^{-6}$	0.2847	0.5000	0.0057	0.3927
9/8/2021	1.0877	$9.88 \cdot 10^{-7}$	0.2492	0.5000	0.0065	0.4349
10/6/2021	0.9826	$2.70 \cdot 10^{-6}$	0.2584	0.5000	0.0094	0.2837
11/10/2021	0.9905	$1.65 \cdot 10^{-6}$	0.2681	0.5000	0.0075	0.2952
12/8/2021	0.8833	$2.27 \cdot 10^{-6}$	0.2943	0.5000	0.0081	0.2763
1/12/2022	0.8447	$2.13 \cdot 10^{-6}$	0.3000	0.5000	0.0073	0.2799
2/9/2022	0.8745	$2.94 \cdot 10^{-6}$	0.2869	0.5000	0.0091	0.2305
3/9/2022	1.0237	$5.27 \cdot 10^{-6}$	0.2464	0.5000	0.0158	0.1881
4/6/2022	1.0867	$1.47 \cdot 10^{-6}$	0.2434	0.5000	0.0092	0.3980
5/10/2022	1.0220	$3.97 \cdot 10^{-6}$	0.1918	0.5971	0.0176	0.2311

Table 2: Calibration of Duan’s GARCH model parameters using the MCS method

considered since the volatility is not constant but stochastic in this investigation. We calculate Greeks using the finance difference method for each variable.

It is widely acknowledged that the calculation time of the ANN method is remarkably faster than the MCS method. Table 8 presents the calculation time for OTM calls and puts using the MCS and ANN methods on Duan’s GARCH and CTS-GARCH models, respectively. For example, we observed 558 prices of calls and puts on 6/9/2021. The MCS and ANN methods respectively take 0.13 and 0.085 seconds to calculate 558 calls and puts on Duan’s GARCH model, and 0.3286 and 0.0187 seconds on the CTS-GARCH model. The number of call and put price observations (Obs.) varies each day and is shown in the Obs. column in Table 8. The calculation times using the ANN method of Duan’s GARCH model are about 9 times faster than the MCS method. Similarly, in the CTS-GARCH model case, the MCS method takes approximately 20 times longer than the ANN method.

## 5. Conclusion

In this paper, we review Duan’s GARCH model and the CTS-GARCH model for option pricing and investigate the use of ANNs to enhance calibration performance. To achieve this goal, we generate training sets for various model parameters and compute calls and puts prices using the MCS method. We then train a three-layer ANN with twenty nodes per layer using the generated

Date	$\theta$	$\kappa$	$\xi$	$\zeta$	$\sigma_0$	rel-RMSE
6/9/2021	1.1103	$4.64 \cdot 10^{-7}$	0.2258	0.5395	0.0064	0.1972
7/7/2021	1.0196	$6.32 \cdot 10^{-7}$	0.2354	0.5621	0.0049	0.2793
8/11/2021	1.1729	$1.74 \cdot 10^{-7}$	0.1994	0.5737	0.0056	0.3383
9/8/2021	1.1810	$2.89 \cdot 10^{-7}$	0.1927	0.5783	0.0064	0.4107
10/6/2021	0.9015	$8.75 \cdot 10^{-7}$	0.2115	0.6441	0.0074	0.2826
11/10/2021	1.1827	$4.99 \cdot 10^{-7}$	0.1968	0.5700	0.0066	0.2634
12/8/2021	1.1276	$7.31 \cdot 10^{-7}$	0.2238	0.5424	0.0060	0.2078
1/12/2022	0.6377	$4.46 \cdot 10^{-7}$	0.3000	0.6252	0.0047	0.3632
2/9/2022	0.8367	$6.29 \cdot 10^{-7}$	0.2081	0.6739	0.0076	0.2626
3/9/2022	0.9301	$1.64 \cdot 10^{-6}$	0.1905	0.6666	0.0121	0.2297
4/6/2022	1.0649	$5.37 \cdot 10^{-7}$	0.2088	0.5990	0.0070	0.3616
5/10/2022	1.0130	$1.67 \cdot 10^{-6}$	0.1506	0.7046	0.0141	0.2417

Table 3: Calibration of Duan’s GARCH model parameters using the ANN method

Date	$\theta$	$\kappa$	$\xi$	$\zeta$	$\sigma_0$	$\alpha$	$\lambda_+$	$\lambda_-$	rel-RMSE
6/9/2021	0.9386	$1.69 \cdot 10^{-6}$	0.2805	0.5050	0.0086	1.9998	14.0138	10.0760	0.8819
7/7/2021	0.9074	$1.23 \cdot 10^{-6}$	0.2960	0.5050	0.0068	1.9921	11.2482	11.9459	0.9014
8/11/2021	0.9608	$1.06 \cdot 10^{-6}$	0.2876	0.5050	0.0057	1.9998	11.9686	6.3593	0.9211
9/8/2021	1.0986	$9.97 \cdot 10^{-7}$	0.2517	0.5050	0.0066	1.9543	13.9703	87.5816	0.9113
10/6/2021	0.9923	$2.72 \cdot 10^{-6}$	0.2609	0.5050	0.0095	1.9833	51.2546	9.3704	0.8716
11/10/2021	1.0004	$1.67 \cdot 10^{-6}$	0.2708	0.5050	0.0076	1.8757	13.9376	11.8176	0.8938
12/8/2021	0.8921	$2.30 \cdot 10^{-6}$	0.2973	0.5050	0.0082	1.9999	6.0208	24.1959	0.8832
1/12/2022	0.8531	$2.15 \cdot 10^{-6}$	0.3030	0.5050	0.0073	1.7526	5.2020	27.2385	0.8971
2/9/2022	0.8832	$2.97 \cdot 10^{-6}$	0.2898	0.5050	0.0092	1.9999	11.2642	9.8365	0.8822
3/9/2022	1.0340	$5.32 \cdot 10^{-6}$	0.2488	0.5050	0.0160	0.8949	8.5344	28.3573	0.8142
4/6/2022	1.0976	$1.48 \cdot 10^{-6}$	0.2458	0.5050	0.0093	1.9999	2.4136	0.7439	0.8864
5/10/2022	1.0323	$4.01 \cdot 10^{-6}$	0.1937	0.6031	0.0178	1.9984	14.8261	15.2165	0.7809

Table 4: Calibration of CTS-GARCH model parameters using the MCS method

Date	$\theta$	$\kappa$	$\xi$	$\zeta$	$\sigma_0$	$\alpha$	$\lambda_+$	$\lambda_-$	rel-RMSE
6/9/2021	0.5526	$1.56 \cdot 10^{-8}$	0.0954	0.7906	0.0046	1.4202	0.1019	0.5492	0.1662
7/7/2021	2.5598	$4.65 \cdot 10^{-6}$	0.0631	0.5124	0.0065	1.0542	13.6212	0.1155	0.2372
8/11/2021	1.2173	$1.43 \cdot 10^{-8}$	0.1960	0.6349	0.0076	0.7306	46.0982	0.1629	0.2936
9/8/2021	1.1724	$6.67 \cdot 10^{-10}$	0.1582	0.7666	0.0068	0.8133	3.9428	0.1019	0.2781
10/6/2021	0.5119	$2.51 \cdot 10^{-8}$	0.0014	0.5016	0.0163	0.5610	13.7536	12.3532	0.2504
11/10/2021	1.9605	$4.93 \cdot 10^{-8}$	0.1048	0.5092	0.0079	0.6510	1.4461	4.1668	0.2701
12/8/2021	0.0281	$3.46 \cdot 10^{-6}$	0.2965	0.7125	0.0091	0.7452	31.5555	0.5439	0.2474
1/12/2022	0.6766	$2.27 \cdot 10^{-6}$	0.2979	0.5750	0.0075	0.5862	20.2640	3.5172	0.2233
2/9/2022	0.4874	$2.72 \cdot 10^{-6}$	0.2624	0.6696	0.0100	0.7577	27.9998	1.5475	0.2152
3/9/2022	0.5884	$2.91 \cdot 10^{-6}$	0.1191	0.8183	0.0134	0.6454	21.7075	4.0287	0.2684
4/6/2022	1.5418	$9.84 \cdot 10^{-6}$	0.1040	0.5271	0.0078	0.9238	21.6161	1.5811	0.3571
5/10/2022	0.8144	$3.15 \cdot 10^{-8}$	0.1669	0.7220	0.0178	0.7674	28.8533	4.4167	0.2756

Table 5: Calibration of CTS-GARCH model parameters using the ANN method

Date	Duan's GARCH Model		CTS-GARCH Model	
	MCS	ANN	MCS	ANN
6/9/2021	0.3048	0.1972	0.8819	<b>0.1662</b>
7/7/2021	0.3585	0.2793	0.9014	<b>0.2372</b>
8/11/2021	0.3927	0.3383	0.9211	<b>0.2936</b>
9/8/2021	0.4349	0.4107	0.9113	<b>0.2781</b>
10/6/2021	0.2837	0.2826	0.8716	<b>0.2504</b>
11/10/2021	0.2952	<b>0.2634</b>	0.8938	0.2701
12/8/2021	0.2763	<b>0.2078</b>	0.8832	0.2474
1/12/2022	0.2799	0.3632	0.8971	<b>0.2233</b>
2/9/2022	0.2305	0.2626	0.8822	<b>0.2152</b>
3/9/2022	<b>0.1881</b>	0.2297	0.8142	0.2684
4/6/2022	0.3980	0.3616	0.8864	<b>0.3571</b>
5/10/2022	<b>0.2311</b>	0.2417	0.7809	0.2756

Table 6: Relative RMSE values for model calibrations

Call/Put	$S_0$	$r$	$K$	$\tau$	Model	$\Delta$	$\Gamma$	$\Theta$	$\rho$
Call	4219.55	0.0020	4220	0.0360	Duan's GARCH	0.4109	$5.7074 \cdot 10^{-3}$	382.59	61.56
					CTS-GARCH	0.2933	$3.5068 \cdot 10^{-3}$	274.16	43.88
				0.1480	Duan's GARCH	0.6089	$3.2794 \cdot 10^{-3}$	311.14	370.61
					CTS-GARCH	0.4744	$2.4125 \cdot 10^{-3}$	312.57	288.33
				0.2880	Duan's GARCH	0.6656	$2.2740 \cdot 10^{-3}$	181.63	780.57
					CTS-GARCH	0.5042	$1.3049 \cdot 10^{-3}$	194.84	586.98
Put	4219.55	0.0020	4215	0.0360	Duan's GARCH	-0.2923	$2.6660 \cdot 10^{-3}$	439.87	-45.64
					CTS-GARCH	-0.2196	$1.7826 \cdot 10^{-3}$	389.37	-34.41
				0.1480	Duan's GARCH	-0.4410	$2.0529 \cdot 10^{-3}$	442.75	-288.81
					CTS-GARCH	-0.2956	$1.4555 \cdot 10^{-3}$	283.71	-194.58
				0.2880	Duan's GARCH	-0.4726	$1.5977 \cdot 10^{-3}$	152.80	-613.73
					CTS-GARCH	-0.2807	$1.0474 \cdot 10^{-3}$	178.97	-369.67

Table 7: Greeks for ATM calls and puts on 6/9/2021




Fig. 3: Calibrated call and put option prices and implied volatility curves at 6/9/2021

Date	Obs.	Duan's GARCH Model		CTS-GARCH Model	
		MCS (sec)	ANN (sec)	MCS (sec)	ANN (sec)
6/9/2021	558	0.1300	0.0185	0.3286	0.0187
7/7/2021	644	0.1387	0.0191	0.3411	0.0175
8/11/2021	706	0.1339	0.0169	0.3025	0.0172
9/8/2021	729	0.1518	0.0166	0.3363	0.0172
10/6/2021	845	0.1422	0.0178	0.3390	0.0178
11/10/2021	858	0.1463	0.0183	0.3229	0.0175
12/8/2021	843	0.1538	0.0174	0.3342	0.0182
1/12/2022	878	0.1598	0.0169	0.3356	0.0172
2/9/2022	928	0.2017	0.0193	0.3539	0.0207
3/9/2022	923	0.1625	0.0176	0.3351	0.0185
4/6/2022	818	0.1410	0.0165	0.3285	0.0189
5/10/2022	570	0.1211	0.0163	0.2978	0.0171

Table 8: Comparison of calculation time using the MCS and ANN methods for Duan's GARCH and CTS-GARCH models

training set. Additionally, we create four ANNs for calls and puts under both Duan's GARCH model and the CTS-GARCH model. Finally, we demonstrate the effectiveness of the trained ANNs by using them to calibrate market option prices. The results show that the ANN is significantly faster than the MCS method once it has been trained and it outperforms the MCS method in terms of calibration performance. Furthermore, the ANN method allows us to calculate Greeks, which is not available with the MCS method.

## References

- Alaya, M. B., Kebaier, A., and Sarr, D. (2021). Deep calibration of interest rates model. *arXiv preprint arXiv:2110.15133*.
- Anand, A., Li, T., Kurosaki, T., and Kim, Y. S. (2016). Foster–hart optimal portfolios. *Journal of Banking & Finance*, 68, 117–130.
- Barndorff-Nielsen, O. E. and Levendorskii, S. (2001). Feller processes of normal inverse Gaussian type. *Quantitative Finance*, 1, 318 – 331.
- Barndorff-Nielsen, O. E. and Shephard, N. (2001). Normal modified stable processes. *Economics Series Working Papers from University of Oxford, Department of Economics*, 72.
- Bayer, C., Horvath, B., Muguruza, A., Stemper, B., and Tomas, M. (2019). On deep calibration of (rough) stochastic volatility models. *arXiv preprint arXiv:1908.08806*.
- Beck, A., Kim, Y. S. A., Rachev, S., Feindt, M., and Fabozzi, F. (2013). Empirical analysis of ARMA-GARCH models in market risk estimation on high-frequency U.S. data. *Studies in Nonlinear Dynamics and Econometrics*, 17(2), 167–177.
- Black, F. and Scholes, M. (1973). The pricing of options and corporate liabilities. *The Journal of Political Economy*, 81(3), 637–654.
- Boyarchenko, S. I. and Levendorskii, S. Z. (2000). Option pricing for truncated Lévy processes. *International Journal of Theoretical and Applied Finance*, 3, 549–552.
- Carr, P., Geman, H., Madan, D., and Yor, M. (2002). The fine structure of asset returns: An empirical investigation. *Journal of Business*, 75(2), 305–332.
- Choi, J., Kim, H., and Kim, Y. S. (2021). Diversified reward-risk parity in portfolio construction. *arXiv preprint arXiv:2106.09055*.
- Choi, J., Kim, Y. S., and Mitov, I. (2015). Reward-risk momentum strategies using classical tempered stable distribution. *Journal of Banking & Finance*, 58, 194–213.
- Coleman, T. and Li, Y. (1994). On the convergence of reflective newton methods for large-scale nonlinear minimization subject to bounds. *Mathematical Programming*, 67(2).
- Coleman, T. and Li, Y. (1996). An interior, trust region approach for nonlinear minimization subject to bounds. *SIAM Journal on Optimization*, 6.
- Cont, R. and Tankov, P. (2004). *Financial Modelling with Jump Processes*. Chapman & Hall / CRC.
- Duan, J.-C. (1995). The GARCH option pricing model. *Mathematical Finance*, 5(1), 13–32.
- Duan, J.-C. (1999). Conditionally fat-tailed distributions and the volatility smile in options. *Rotman School of Management, University of Toronto, Working Paper*.



- Duan, J.-C., Ritchken, P., and Sun, Z. (2004). Jump starting GARCH: Pricing and hedging options with jumps in returns and volatilities. University of Toronto and Case Western Reserve University.
- Duan, J.-C., Ritchken, P., and Sun, Z. (2006). Approximating GARCH-jumps models, jump-diffusion processes, and option pricing. *Mathematical Finance*, 16(1), 21–52.
- Engle, R. F. and Mustafa, C. (1992). Implied arch models from options prices. *Journal of Econometrics*, 52(1-2), 289–311.
- Georgiev, K., Kim, Y. S., and Stoyanov, S. (2015). Periodic portfolio revision with transaction costs. *Mathematical Methods of Operations Research*, 81(3), 337–359.
- Halton, J. (1964). Algorithm 247: Radical-inverse quasi-random point sequence. *Communications of the ACM*, 7.
- Hutchinson, J. M., Lo, A. W., and Poggio, T. (1994). A nonparametric approach to pricing and hedging derivative securities via learning networks. *The Journal of Finance*, 49(3), 851–889.
- Kim, Y. S., Jiang, D., and Stoyanov, S. (2019). Long and short memory in the risk-neutral pricing process. *Journal of Derivatives*, 26(4), 71 – 88.
- Kim, Y. S., Rachev, S. T., Bianchi, M. L., and Fabozzi, F. J. (2008a). Financial market models with Lévy processes and time-varying volatility. *Journal of Banking and Finance*, 32, 1363–1378.
- Kim, Y. S., Rachev, S. T., Bianchi, M. L., and Fabozzi, F. J. (2008b). A new tempered stable distribution and its application to finance. In G. Bol, S. T. Rachev, and R. Wuerth (Eds.), *Risk Assessment: Decisions in Banking and Finance*, Physika Verlag, Springer. 77–110.
- Kim, Y. S., Rachev, S. T., Bianchi, M. L., and Fabozzi, F. J. (2010a). Tempered stable and tempered infinitely divisible GARCH models. *Journal of Banking and Finance*, 34, 2096–2109.
- Kim, Y. S., Rachev, S. T., Bianchi, M. L., and Fabozzi, F. J. (2010b). Tempered stable and tempered infinitely divisible GARCH models. *Journal of Banking & Finance*, 34(9), 2096–2109.
- Kim, Y. S., Roh, K.-H., and Douady, R. (2022). Tempered stable processes with time varying exponential tails. *Quantitative Finance*, 22, 541–561.
- Koponen, I. (1995). Analytic approach to the problem of convergence of truncated Lévy flights towards the Gaussian stochastic process. *Physical Review E*, 52, 1197–1199.
- Malliaris, M. and Salchenberger, L. (1993). A neural network model for estimating option prices. *Applied Intelligence*, 3(3), 193–206.
- McCulloch, W. S. and Pitts, W. (1943). A logical calculus of the ideas immanent in nervous activity. *The Bulletin of Mathematical Biophysics*, 5(4), 115–133.

- Menn, C. and Rachev, S. T. (2009). Smoothly truncated stable distributions, GARCH-models, and option pricing. *Mathematical Methods of Operations Research*, 69, 411–438.
- Merton, R. C. (1973). Theory of rational option pricing. *The Bell Journal of Economics and Management Science*, 141–183.
- Rachev, S. T., Kim, Y. S., Bianchi, M. L., and Fabozzi, F. J. (2011). *Financial models with Lévy processes and volatility clustering*. John Wiley & Sons.
- Rosiński, J. (2007). Tempering stable processes. *Stochastic Processes and Their Applications*, 117(6), 677–707.
- Ruf, J. and Wang, W. (2019). Neural networks for option pricing and hedging: a literature review. *arXiv preprint arXiv:1911.05620*.
- Tsuchida, N., Zhou, X., and Rachev, S. (2012). Mean-ent portfolio selection under maximum weight and turnover constraints based on fundamental security factors. *The Journal of Investing*, 21(1), 14–24.