

SUPERSINGULAR ISOGENY GRAPHS FROM ALGEBRAIC MODULAR CURVES

NADIR HAJOUJI

ABSTRACT. We describe algorithms for computing supersingular ℓ -isogeny graphs using algebraic models of modular curves. The algorithms we describe were used to compute all supersingular isogeny graphs for all pairs (p, ℓ) with $p \leq 2048$ and $\ell \leq 13$. In addition, we use our results to prove a conjecture posed by Nakaya.

1. INTRODUCTION

Given a pair of primes p, ℓ , with $p \neq \ell$, we would like to compute the following:

- (1) The set of supersingular j -invariants.
- (2) The supersingular ℓ -isogeny graphs.

There are already algorithms for computing these graphs, see e.g. [4], [9], [15]. These algorithms usually involve computations in the field of definition of the ℓ -torsion points of the supersingular curves. We will describe simpler algorithms using the modular curves $X_0(\ell)$ and $X_0(\ell)^+$. In addition, we will use our results about supersingular isogeny graphs to explain and generalize some of the observations in [12].

Our paper is structured as follows:

- In Section 2, we prove that every isogeny between supersingular curves is defined over \mathbb{F}_{p^2} . This result is needed to ensure we can compute isogeny graphs by checking only the \mathbb{F}_{p^2} of a suitable modular curve.
- In Section 3, we obtain models for the modular curves $X_0(\ell)$ and $X_0(\ell)^+$ for $\ell \in \{3, 5, 7, 11, 13\}$.
- In Section 4, we describe algorithms for computing supersingular isogeny graphs.
- In Section 5, we use the tools we built for computing supersingular isogeny graphs to prove new theorems.
 - We obtain a formula for the trace of the adjacency matrix of the ℓ -isogeny graph in 5.1.
 - In 5.2, we explain how to translate questions asked in [12] to questions about isogeny graphs. We use this perspective to prove and extend Conjecture 2 in that paper.

2. SUPERSINGULAR ELLIPTIC CURVES

2.1. Quadratic Twists. Let \mathbb{F}_q be a field of characteristic not equal to 2, let $d \in \mathbb{F}_q^\times$ be a nonsquare and let E/\mathbb{F}_q be an elliptic curve given by a Weierstrass equation:

$$E : y^2 = x^3 + ax^2 + bx + c$$

We define the quadratic twist of E/\mathbb{F}_q to be the elliptic curve:

$$E^{-1} : \quad dy^2 = x^3 + ax^2 + bx + c$$

There is an obvious injection:

$$E^{-1}(\mathbb{F}_q) \rightarrow E(\mathbb{F}_{q^2}) \quad (x, y) \mapsto (x, \sqrt{d}y)$$

We identify $E^{-1}(\mathbb{F}_q)$ with its image in $E(\mathbb{F}_{q^2})$. Note that $E^{-1}(\mathbb{F}_q)$ coincides with the kernel of the trace map $E(\mathbb{F}_{q^2}) \rightarrow E(\mathbb{F}_q)$. In particular, this means:

$$E^{-1}(\mathbb{F}_q) \cap E(\mathbb{F}_q) = \ker(\text{Tr}|_{E(\mathbb{F}_q)}) = E[2](\mathbb{F}_q)$$

Let E/\mathbb{F}_q be an elliptic curve and $\phi_n : E \rightarrow E$ the p^n -power Frobenius. We write $\text{Tr}(E, \phi_n)$ to denote the trace of ϕ_n on E . If E is clear from context, we write $\text{Tr}(\phi_n)$.

Lemma 2.1. *Let E/\mathbb{F}_q be an elliptic curve, let E^{-1}/\mathbb{F}_q be the quadratic twist and let $\phi : E \rightarrow E$ be the q -power Frobenius. Then:*

$$\text{Tr}(\phi, E^{-1}) = -\text{Tr}(\phi, E)$$

Proof. We will show that $E(\mathbb{F}_q) \cup E^{-1}(\mathbb{F}_q)$ coincides with the preimage of $\mathbb{P}_{\mathbb{F}_q}^1 \subset \mathbb{P}_{\mathbb{F}_{q^2}}^1$ under the double cover $E(\mathbb{F}_{q^2}) \rightarrow \mathbb{P}_{\mathbb{F}_{q^2}}^1$. This will allow us to compute $\#E(\mathbb{F}_q) + \#E^{-1}(\mathbb{F}_q)$ by determining the number of points in each fiber over each point in $\mathbb{P}_{\mathbb{F}_q}^1$. We will see that $\#E(\mathbb{F}_q) + \#E^{-1}(\mathbb{F}_q) = 2q + 2$, which allows us to deduce the lemma from the Weil conjectures.

Let $x_0 \in \mathbb{P}_{\mathbb{F}_q}^1$. Then exactly one of the following is true:

- (1) $x_0^3 + ax_0^2 + bx_0 + c = 0$. There are at most 3 points x_0 for which this is true, and whenever this happens, it means we have $(x_0, 0) \in E(\mathbb{F}_q) \cap E^{-1}(\mathbb{F}_q)$.
- (2) $x_0^3 + ax_0^2 + bx_0 + c = y_0^2$ for some $y_0 \in \mathbb{F}_q^\times$: in this case, we have $(x_0, y_0), (x_0, -y_0) \in E(\mathbb{F}_q)$.
- (3) $x_0^3 + ax_0^2 + bx_0 + c = dy_0^2$ for some $y_0 \in \mathbb{F}_q^\times$: in this case, we have $(x_0, y_0), (x_0, -y_0) \in E^{-1}(\mathbb{F}_q)$.

Thus, every point $[x_0 : 1] \in \mathbb{P}_{\mathbb{F}_q}^1$ contributes 2 to the sum $\#E(\mathbb{F}_q) + \#E^{-1}(\mathbb{F}_q)$. Together with the identity, they give us a total of:

$$\#E(\mathbb{F}_q) + \#E^{-1}(\mathbb{F}_q) = 2 \cdot \#\mathbb{P}^1(\mathbb{F}_q) = 2(q + 1)$$

which proves the lemma. □

2.2. Isogenies.

Theorem 2.2. *Let E/\mathbb{F}_{p^2} be a supersingular curve and $\psi : E \rightarrow E'$ an isogeny of degree prime to p . Then ψ is defined over \mathbb{F}_{p^2} .*

Proof. First, we prove the theorem for supersingular curves E/\mathbb{F}_p . Let $\phi : E \rightarrow E$ be the p -power Frobenius map. Since E is supersingular, the characteristic polynomial of ϕ is:

$$\chi_\phi(t) = t^2 + p$$

By the Cayley-Hamilton theorem, ϕ^2 acts on E as multiplication-by- $[-p]$.

Now, let $\psi : E \rightarrow E'$ be an isogeny. We can factor ψ as $\psi = \psi_0 \circ [m]$, where $\psi_0 : E \rightarrow E'$ is an isogeny with cyclic kernel, and $[m] : E \rightarrow E$ is the multiplication map. The latter is defined over \mathbb{Z} , so we just have to prove that ψ_0 is defined over \mathbb{F}_{p^2} .

Let $\langle P_0 \rangle \cong \mathbb{Z}/d\mathbb{Z}$ be the kernel of ψ_0 . Then:

$$\phi^2 \cdot \langle P_0 \rangle = \langle \phi^2 \cdot P_0 \rangle = \langle [-p] \cdot P_0 \rangle = \langle P_0 \rangle$$

Thus, $\langle P_0 \rangle$ is defined over \mathbb{F}_{p^2} , so ψ_0 is defined over \mathbb{F}_{p^2} as claimed. This completes the proof of the theorem for supersingular curves E/\mathbb{F}_p .

Next, let E/\mathbb{F}_{p^2} be a supersingular curve with $j(E) \notin \mathbb{F}_p$, and let $\phi_2 : E \rightarrow E$ be the p^2 -power Frobenius. Note that $Tr(\phi_2, E) \in \{-2p, 2p\}$ for all supersingular curves with $j(E) \notin \mathbb{F}_p$.¹

Let E_0/\mathbb{F}_{p^2} be any supersingular curve satisfying:

- $j(E_0) \in \mathbb{F}_p$.
- $Tr(\phi_2, E_0) = Tr(\phi_2, E)$.

Note that such a curve necessarily exists. First, we can always find a supersingular curve E/\mathbb{F}_p - this follows from Prop 14.18 in [6], or Theorem 2 in [12]. The trace of Frobenius of the base change E_0/\mathbb{F}_{p^2} is $2p$, so the trace of Frobenius on the quadratic twist of E_0/\mathbb{F}_{p^2} must be $-2p$ by Lemma 2.1. Thus, we take E_0 to be the base change of E_0/\mathbb{F}_p when $Tr(E, \phi_2) = -2p$ and we take E_0 to be the quadratic twist of the base change of E_0/\mathbb{F}_p otherwise.²

Fix an isogeny $\psi_0 : E_0 \rightarrow E$. Let $\psi : E \rightarrow E'$ be any isogeny. We know the theorem holds for E_0 , so the isogenies $\psi_0 : E_0 \rightarrow E$ and $\psi_1 = \psi \circ \psi_0 : E_0 \rightarrow E'$ must both be defined over \mathbb{F}_{p^2} . Thus ψ must be defined over \mathbb{F}_{p^2} . \square

3. ALGEBRAIC MODELS OF MODULAR CURVES

In this section, we derive formulas that are needed in Section 4. For each odd prime $\ell \leq 13$, we will obtain descriptions of $X_0(\ell)$, as well as formulae for the j -maps $j_\ell : X_0(\ell) \rightarrow X(1)$ and the Fricke involutions $F_\ell : X_0(\ell) \rightarrow X_0(\ell)$.

For $\ell > 3$, our starting point will be the “universal elliptic curve with a point P of order ≥ 4 ”.³ Explicitly, this is the family of elliptic curves over $\mathbb{A}^2 = \text{Spec } \mathbb{Z}[u, v]$ given by the equation:

$$\mathcal{E}_{u,v} : y^2 + (1 - u)xy - vy = x^3 - vx^2$$

Note that $(0, 0) \in \mathcal{E}_{u,v}(\mathbb{Z}[u, v])$ is a point of infinite order. If E/K is any elliptic curve, and $P \in E(K)$ is a point of order ≥ 4 , there is a unique point $(u_0, v_0) \in \mathbb{A}_K^2$ such that E is isomorphic to the fiber over (u_0, v_0) in $\mathcal{E}_{u,v} \times \text{Spec } K$, and in fact there is an isomorphism from E to the fiber that takes P to the point $(0, 0)$ on the fiber. We will describe the modular curves $X_1(\ell)$ by describing maps $i_\ell : X_1(\ell) \rightarrow \mathbb{A}^2$ from smooth curves with the property that $\mathcal{E} \times_{\mathbb{A}^2} X_1(\ell) \rightarrow X_1(\ell)$ is the universal curve with a point of order ℓ .

¹In fact, this is true for all except at most 4 supersingular curves. If $p \equiv 3 \pmod{4}$, there are two models for the curve with $j = 1728$ that are defined over \mathbb{F}_{p^2} with $Tr(\phi_2) \in \{\pm p\}$, and if $p \equiv 2 \pmod{3}$, there are two models for the curve with $j = 0$ with $Tr(\phi_2) = 0$. See [17],[13],[4] for additional details.

²Note that order matters - the base change of the quadratic twist of E_0 is not isomorphic to the quadratic twist of the base change of E_0 ; we have to quadratically twist using a nonsquare $d \in \mathbb{F}_{p^2}$ to obtain the model with trace $2p$.

³See, e.g., Ex. 8.13 in [14].

Next, define:

$$\alpha_{\mathcal{E}} : \mathbb{A}^2 \rightarrow \mathbb{A}^2 \quad \alpha_{\mathcal{E}}(u, v) = \left(-\frac{u^2v - u^2 + 3uv - 2v^2}{u^4}, -\frac{v(u^2 + u - v)^3}{u^8} \right)$$

If $(u_0, v_0) \in \mathbb{A}^2$ represents a pair (E, P_0) , then $\alpha_{\mathcal{E}}(u_0, v_0)$ represents $(E, 2P_0)$. In particular, $j_{\mathcal{E}} \circ \alpha_{\mathcal{E}} = j_{\mathcal{E}}$. We will restrict $\alpha|_{\mathcal{E}}$ to the modular curves $X_1(\ell)$ to obtain automorphisms $\alpha_{\ell} : X_1(\ell) \rightarrow X_1(\ell)$ of order $\frac{\ell-1}{2}$. We obtain $X_0(\ell)$ as a quotient $X_1(\ell) \rightarrow X_1(\ell)/\langle \alpha_{\ell} \rangle$, and we obtain the maps $j_{\ell} : X_0(\ell) \rightarrow X(1)$ by finding functions on $X_0(\ell)$ that pull back to $j_{\mathcal{E}} \circ i_{\ell}$ on $X_1(\ell)$.

Obtaining a formula for the Fricke involution $F_{\ell} : X_0(\ell) \rightarrow X_0(\ell)$ is straightforward: we simply look for involutions that permute the two cusps of $X_0(\ell)$. This determines F up to a scalar multiple. We can determine F completely by using Vélú's formula⁴ to determine the correct value of $j_{\ell} \circ F_{\ell}(t_0)$ at some point t_0 for which $j_{\ell} \circ F_{\ell}(t_0) \neq 0$.

In addition, we will obtain a description of the double cover $X_0(\ell) \rightarrow X_0(\ell)/\langle F_{\ell} \rangle =: X_0(\ell)^+$ in terms of the *Atkin modular polynomial*:

$$R_{\ell}(x, y) = x^2 - a_{\ell}(y)x + b_{\ell}(y)$$

We will describe the quotient map $X_0(\ell) \rightarrow X_0(\ell)^+$ explicitly, and give formulas for the polynomials a_{ℓ}, b_{ℓ} . Note that analogs of these polynomials $R_{\ell}(x, y)$ can be obtained whenever $X_0(\ell)^+$ has genus 0.

3.0.1. Remark. The set of primes ℓ for which $X_0(\ell)^+$ has genus 0 famously coincides with the set of primes that divide the size of the Monster group. Furthermore, these are precisely the primes for which every supersingular curve has a model defined over the ground field \mathbb{F}_p . We will refer to them as *supersingular primes*.

Now, the curve defined by $R_{\ell}(x, y)$ will have singularities at *some* of the points (x_0, y_0) where $a_{\ell}(y_0)^2 - 4b_{\ell}(y_0) = 0$. Note that x_0 is determined by y_0 in this case, since the quadratic has a repeated root. These points represent elliptic curves with an endomorphism of degree ℓ . We have to be careful dealing with these points, as *some* of them represent multiple edges on the isogeny graph. To avoid having to determine the correct multiplicity for each edge, we will simply compute these multiplicities in characteristic 0, and write down a polynomial $\Delta_{\ell}(j)$ whose roots are j -invariants of elliptic curves with an endomorphism of degree ℓ , and where the multiplicity of each root indicates the number of isogenies of degree ℓ for that j -invariant. The factors of these Δ_{ℓ} will be Hilbert class polynomials; we give explicit formula for the Hilbert class polynomials that appear as factors in the appendix.

3.1. $\ell = 3$. For $\ell = 3$, we can obtain the formulas we need without using \mathcal{E} .

Let E be an elliptic curve with a point P of order 3, and assume E is given by an equation:

$$y^2 = x^3 + b_2x^2 + b_4x + b_6$$

with $x(P) = 0$. The condition $[3]P = 0$ is equivalent to $x([2]P) = x(P)$, so we must have $x([2]P) = 0$. Now, a computation shows that $x([2]P) = \frac{b_4^2 - 4b_2b_6}{4b_8}$ so the condition $[3]P = 0$

⁴On the universal curve over $X_1(\ell)$.

implies $b_4^2 - 4b_2b_6 = 0$, which in turn implies that the quadratic:

$$b_2x^2 + b_4x + b_6$$

has a repeated root. This means we can find c_0, c_1 such that:

$$y^2 = x^3 + c_0(x + c_1)^2$$

Furthermore, replacing c_0, c_1 by tc_0, tc_1 for any nonzero t does not change the isomorphism type of the pair (E, P_0) , so we can use $\mathbb{P}^1 = \{[c_0 : c_1]\}$ as a model for $X_0(3)$. The map j_3 is:

$$j_3([c_0 : c_1]) = \frac{256c_0^3(c_0^3 - 6c_1)^3}{c_1^3(4c_0^3 - 27c_1)}$$

In these coordinates, the Fricke involution is given by⁵:

$$[c_0 : c_1] \mapsto [-27c_0 : 4c_0 + 27c_1]$$

To match the form of the equations we will obtain for $\ell > 3$, we use the change of variable to move the cusps⁶ to $0, \infty$. We can ignore the point at infinity and work with an affine coordinate t . In these coordinates:

$$j_3(t) = -\frac{(t-27)(t-3)^3}{t} \tag{J3}$$

and the Fricke involution is:

$$F_3(t) = \frac{729}{t} \tag{F3}$$

The coefficients of the Atkin modular polynomial $R_3(x, y)$ are easy to obtain in these coordinates:

$$a_3(y) = y(y^2 - 126y + 2944) \tag{RA3}$$

$$b_3(y) = y(y + 192)^3 \tag{RB3}$$

The discriminant of $R_3(x, y)$ is:

$$a_3(y)^2 - 4b_3(y) = y(y - 108)(y - 64)^2(y - 8)^2$$

We compute the j -invariants represented by $y = 0, 8, 64, 108$ and record the multiplicity of those j -invariants in the following polynomial:

$$\Delta_3(j) = j(j - 54000)(j - 8000)^2(j + 32768)^2 \tag{D3.1}$$

Using the notation explained in the appendix, we can rewrite the previous equation as:

$$\Delta_3(j) = H_{-3}(j)H_{-3.4}(j)H_{-2}(j)^2H_{-11}(j)^2 \tag{D3.2}$$

⁵See [7] for more on 3-isogenies in these coordinates.

⁶In other words, the points on $X_0(\ell)$ that represent singular curves.

3.2. $\ell = 5$. The curve $X_1(5)$ is abstractly isomorphic to \mathbb{P}^1 , and the rational map $i_5 : X_1(5) \rightarrow \mathbb{A}^2$ is essentially the diagonal embedding: if we fix an affine coordinate $t = \frac{t_0}{t_1}$ on \mathbb{P}^1 , then the map $i_5 : X_1(5) \rightarrow \mathbb{A}^2$ is given by $i_5(t) = (t, t)$. The universal elliptic curve with a point of order 5 is therefore given by:

$$y^2 + (1-t)xy - ty = x^3 - tx^2 \quad (\text{E5.1})$$

We compute $j_{5,1} := j_{\mathcal{E}} \circ i_5$:

$$j_{5,1}(t) = \frac{(t^4 - 12t^3 + 14t^2 + 12t + 1)^3}{t^5(t^2 - 11t - 1)} \quad (\text{J5.1})$$

Now, we can apply Vélu's formula ([16]) to the entire family of elliptic curves over $X_1(5)$ to obtain a new family over $X_1(5)$ given by a Weierstrass equation:

$$y^2 + a'_1xy + a'_3y = x^3 + a'_2x^2 + a'_4x + a'_6$$

where:

$$\begin{aligned} a'_1 &= 1 - t \quad (= a_1) \\ a'_2 &= -t \quad (= a_2) \\ a'_3 &= -t \quad (= a_3) \\ a'_4 &= -5t(t^2 + 2t - 1) \\ a'_6 &= -t(t^4 + 10t^3 - 5t^2 + 15t - 1) \end{aligned}$$

The j -invariant of this new family is:

$$j'_{5,1}(t) = \frac{(t^4 + 228t^3 + 494t^2 - 228t + 1)^3}{t(t^2 - 11t - 1)^5} \quad (\text{J5.1}')$$

We will use this when computing a formula for the Fricke involution.

3.2.1. $X_1(5) \rightarrow X_0(5)$. To obtain the map $X_1(5) \rightarrow X_0(5)$, we compute the quotient of $X_1(5)$ by the automorphism $\alpha_{\mathcal{E}}|_{X_1(5)}$. Observe that:

$$\alpha_{\mathcal{E}} \circ i_5(t) = \alpha_{\mathcal{E}}(t, t) = \left(\frac{-1}{t}, \frac{-1}{t} \right)$$

Thus, $\alpha_{\mathcal{E}}|_{X_1(5)} = \alpha_5(t)$, where:

$$\alpha_5(t) = \frac{-1}{t} \quad (\text{A5.1})$$

The function $t + \frac{-1}{t}$ is clearly invariant under α_5 , so we use it to construct a quotient map $X_1(5) \rightarrow X_0(5)$:

$$q_5(t) = \frac{t^2 - 11t - 1}{t} \quad (\text{Q5})$$

Note that the quotient map has been constructed to ensure the I_5 cusp gets sent to the point at infinity, and the I_1 cusp gets sent to the point 0 on $X_0(5)$.

To obtain the map $j_5 : X_0(5) \rightarrow X(1)$, we do the following:

- The poles of j_5 are at $0, \infty$ by construction: one can check that the map q_5 takes the cusps of $X_1(5)$ to $0, \infty$ on $X_0(5)$.

- We compute the zeros of $j_{5,1}$, and use the map q_5 to determine what the zeros of j_5 should be.
- Once we know the zeros and poles of j_5 , the function is determined up to a scalar multiple. We can ensure we have the correct scalar by evaluating $j_{5,1}$ at a random point which is not a zero or pole.

The end result is:

$$j_5(t) = \frac{(t^2 + 10t + 5)^3}{t} \quad (\text{J5})$$

The Fricke involution is:

$$F_5(t) = \frac{125}{t} \quad (\text{F5})$$

One can check that $j_{5,1}(t) = j_5 \circ q_5(t)$. Furthermore, note that $j'_{5,1}(t) = j_{5,0} \circ q_5(\frac{125}{t})$, so we no longer need to use Vélú's formula.

3.2.2. $X_0(5) \rightarrow X_0(5)^+$. It is easy to find a quotient map $X_0(5) \rightarrow X_0(5)/\langle F_5 \rangle$ - we will use q_5^+ :

$$q_5^+ : X_0(\ell) \rightarrow X_0(\ell)^+ \quad q_5^+(t) = t + 22 + \frac{125}{t} \quad (\text{Q5+})$$

Note that the constant term 22 is chosen so that the polynomials we obtain using q_5^+ match those in [12].

We obtain the coefficients of a_5, b_5 by matching coefficients with the system:

$$a_5(q_5^+(t)) = j_5(t) + (j_5 \circ F_5)(t)$$

$$b_5(q_5^+(t)) = j_5(t) \cdot (j_5 \circ F_5)(t)$$

The polynomials are given by:

$$a_5(y) = y^5 - 80y^4 + 1890y^3 - 12600y^2 + 7776y + 3456 \quad (\text{RA5})$$

$$b_5(y) = (y^2 + 216y + 144)^3 \quad (\text{RB5})$$

Finally, we compute the polynomial Δ_5 :

$$\Delta_5(j) = (j - 287496)^2(j - 1728)^2(j + 32768)^2(j + 884736)^2(j^2 - 1264000j - 681472000) \quad (\text{D5})$$

$$= H_{-1}(j)^2 H_{-4}(j)^2 H_{-11}(j)^2 H_{-19}(j)^2 H_{-5}(j)$$

3.3. $\ell = 7$. The map $i_7 : X_1(7) \rightarrow \mathbb{A}^2$ is given by:

$$i_7(t) = (t^2 - t, t^3 - t^2) \quad (\text{I7})$$

The universal elliptic curve with a point of order 7 is therefore given by the following equation:

$$y^2 + (1 + t - t^2)xy + t^2(1 - t)y = x^3 + t^2(1 - t)x^2 \quad (\text{E7.1})$$

We compute $j_{7,1} = j_{\mathcal{E}} \circ i_7$:

$$j_{7,1}(t) = \frac{(t^2 - t + 1)^3 (t^6 - 11t^5 + 30t^4 - 15t^3 - 10t^2 + 5t + 1)^3}{t^7(t - 1)^7(t^3 - 8t^2 + 5t + 1)} \quad (\text{J7.1.1})$$

We can compute the formula for $j'_{7,1}(t)$ by applying Vélú's formula to the universal elliptic curve, and computing the j -invariant of the result:

$$j'_{7,1}(t) = \frac{(t^2 - t + 1)^3 (t^6 + 229t^5 + 270t^4 - 1695t^3 + 1430t^2 - 235t + 1)^3}{t(t-1)(t^3 - 8t^2 + 5t + 1)^7} \quad (\text{J7.1.2})$$

3.3.1. $X_1(7) \rightarrow X_0(7)$. To obtain a model of $X_0(7)$, we take the quotient of $X_1(7)$ by the automorphism:

$$\alpha_7(t) = \frac{1}{1-t} \quad (\text{A7.1})$$

Finding a function invariant under this action is straightforward (e.g. we can take the sum of the orbit). We construct our quotient map $X_1(7) \rightarrow X_0(7)$ so that the I_7 singularity is at ∞ and the I_1 singularity is at 0:

$$q_7(t) = \frac{t^3 - 8t^2 + 5t + 1}{(1-t)t} \quad (\text{Q7})$$

The map $j_7 : X_0(7) \rightarrow X(1)$ is:

$$j_7(t) = -\frac{(t^2 - 13t + 49)(t^2 - 5t + 1)^3}{t} \quad (\text{J7})$$

The Fricke involution swaps the I_1, I_7 cusps of $X_0(7)$, so it has the form $t \mapsto \frac{a}{t}$ for some a . We choose a so that $j_7 \circ F_7 \circ q_7 = j'_{7,1}$, where $j'_{7,1}$ is the j -map J7.1.2 obtained using Vélú's formula:

$$F_7(t) = \frac{49}{t} \quad (\text{F7})$$

3.3.2. $X_0(7) \rightarrow X_0(7)^+$. The computations are similar to the $\ell = 5$ case. The quotient map $X_0(7) \rightarrow X_0(7)^+$ is:

$$q_7^+(t) = -t + 13 - \frac{49}{t} \quad (\text{Q7+})$$

The polynomials a_7, b_7 are:

$$a_7(y) = y^5 - 80y^4 + 1890y^3 - 12600y^2 + 7776y + 3456 \quad (\text{RA7})$$

$$b_7(y) = (y^2 + 216y + 144)^3 \quad (\text{RB7})$$

and the polynomial Δ_7 is:

$$\Delta_7(j) = \left(H_{-3}(j) \cdot H_{-12}(j) \cdot H_{-27}(j) \cdot H_{-6}(j) \cdot H_{-7}(j) \cdot H_{-19}(j) \right)^2 \quad (\text{D7})$$

3.4. $\ell = 11$. This is the first modular curve of positive genus, so we actually have to specify an equation. We obtained a nice model of $X_1(11)$, and a map $X_1(11) \rightarrow \mathbb{A}^2$, using methods described in [1]. The model for $X_1(11)$ is:

$$w^2 - w = z^3 - z^2 \quad (\text{X11.1})$$

The map $i_{11} : X_1(11) \rightarrow \mathbb{A}^2$ is given by:

$$(z, w) \mapsto \left(\frac{(w-1)(w+z-1)}{z}, \frac{w(w-1)(w+z-1)}{z} \right) \quad (\text{I11})$$

The formula $j_{11,1} = j_{\mathcal{E}} \circ i_{11}$ is too complicated to write out, but we will only need to compute it for a few values to obtain j_{11} .

3.4.1. *Quotient.* The equation for $X_1(11)$ coincides with the equation of the fiber over $t = 1$ on $X_1(5)$ in E5.1, which means that the point $(0, 0) \in X_1(11)$ has order 5. Furthermore, the automorphism $\alpha : X_1(11) \rightarrow X_1(11)$ given by $\alpha(P) = (0, 0) + P$ generates $\text{Aut}(X_1(11)/X_0(11))$, so the quotient map $X_1(11) \rightarrow X_0(11)$ is an isogeny of degree 5, and can be computed using Vélu's formula [16]. The equation for $X_0(11)$ we obtain is:

$$y^2 - y = x^3 - x^2 - 10x - 20 \quad (\text{X11.0})$$

The isogeny $X_1(11) \rightarrow X_0(11)$ is:

$$q_{11}(z, w) = \left(\frac{z^5 - 2z^4 + 3z^3 - 2z + 1}{(z-1)^2 z^2}, \frac{w(z^6 - 3z^5 + z^4 - 3z^3 + 6z^2 - 6z + 2) + (z^4 + z^3 - 3z^2 + 3z - 1)}{(z-1)^3 z^3} \right) \quad (\text{Q11})$$

We obtain j_{11} by computing the zeros and poles of $j_{\mathcal{E}} \circ i_{11}$, and using the map q_{11} to determine the zeros and poles of j_{11} . This determines $j_{11,0}$ up to a scalar multiple. We then compute $j_{11,1}$ at a point that is not a zero or pole to determine the correct scalar.

The curve $X_1(11)$ has 5 cusps of width 11 over the 5-torsion subgroup, and 5 cusps of width 1 elsewhere. The cusps of width 11 get mapped to the identity on $X_0(11)$, and the cusps of width 1 get mapped to the point $(16, 61) \in X_0(11)$. Thus, the j -map should have a poly of order 1 at $(16, 61)$ and a pole of order 11 at 0. The zeros can be obtained by finding the points on $X_1(11)$ with j -invariant 0, and computing their image under the isogeny.

The following function has the correct zeros and poles:

$$j_{11}(x, y) = \frac{(x^4 + 12x^3 - 8x^2y - 114x^2 + 40xy + 140x - 32y + 217)^3}{(-5x + y + 19)(4x^2 - xy - 29x + 4y + 51)^2} \quad (\text{J11})$$

Furthermore, $j_{11,0} \circ q_{11} = j_{\mathcal{E}} \circ i_{11}$ at points which are not zeros or poles.

Obtaining the Fricke involution is much easier. There is precisely only involution on $X_0(11)$ that swaps P_1 and 0: the map $P \mapsto P_1 - P$. Thus, we deduce that the Fricke involution on $X_0(11)$ is:

$$F_{11}(P) = (16, 61) - P \quad (\text{F11})$$

where the difference is computed using the group law on E .

3.4.2. $X_0(11) \rightarrow X_0(11)^+$. To compute the quotient of $X_0(11)$ by the Fricke involution, it is helpful to first do a change of variable⁷ on $X_0(11)$ so that the Fricke involution coincides with the negation map. Let $x_2(P) = x(P + [3]P_1)$ and $y_2(P) = y(P + [3]P_1)$. In terms of x_2, y_2 , the Fricke involution is $(x_2, y_2) \mapsto (x_2, 1 - y_2)$, and the map $j_{11,0}$ is given by:

$$j_{11}(x_2, y_2) = \frac{(6x_2 - y_2 - 35)^3(6x_2 + y_2 - 36)(32x_2^2 + 48x_2 + 120y_2 - 429)^3}{(5x_2 - y_2 - 19)^6(5x_2 + y_2 - 20)^2} \quad (\text{J11v2})$$

⁷We are simply going to translate every point on $X_0(11)$ by a multiple of P_1 .

Now, the sum and product of $j_{11}(x_2, y_2)$ and $j_{11}(x_2, 1 - y_2)$ do not depend on y_2 , so we can write them as rational functions of x_2 . In fact, they turn out to be polynomials in $\frac{5x_2-1}{x_2+2}$. Thus, we define:

$$q_{11}^+(P) = \frac{5x_2(P) - 1}{x_2(P) + 2} \quad (\text{Q11+})$$

We have:

$$\begin{aligned} a_{11}(y) = & y^{11} - 44y^{10} + 693y^9 - 4334y^8 + 4400y^7 + 42658y^6 \cdots \\ & - 44968y^5 - 178376y^4 - 58432y^3 + 86240y^2 + 67200y + 16000 \end{aligned} \quad (\text{RA.11})$$

and:

$$b_{11}(y) = (y^4 + 232y^3 + 1176y^2 + 1120y + 400)^3 \quad (\text{B11})$$

Finally, we compute $\Delta_{11}(j)$:

$$\Delta_{11}(j) = \left(H_{-2}(j) \cdot H_{-7}(j) \cdot H_{-28}(j) \cdot H_{-10}(j) \cdot H_{-11}(j) \cdot H_{-44}(j) \cdot H_{-19}(j) \cdot H_{-35}(j) \cdot H_{-43}(j) \right)^2 \quad (\text{D11})$$

Note that the formulae for $R_{11}(x, y)$ and $\Delta_{11}(j)$ also appear in [3], although they are obtained using different means.

3.5. $\ell = 13$. The curve $X_1(13)$ can be described by the following model [8]:

$$w^2 + wz^3 + wz^2 + w - z^2 - z = 0 \quad (\text{X13})$$

To compute j -invariants on this curve, we use the map $i_{13} : X_1(13) \rightarrow \mathbb{A}^2$:

$$i_{13}(z, w) = \left(\frac{wz(wz - w - 1)}{w + 1}, \frac{wz(1 - wz)(wz - w - 1)}{w + 1} \right) \quad (\text{I13})$$

We define $j_{13,1} := j_{\mathcal{E}} \circ i_{13}$.

3.5.1. *Quotient Map.* Because $X_1(13)$ has genus > 1 , it only has finitely many automorphisms. In fact, the full automorphism group of $X_1(13)$ is isomorphic to the dihedral group with 12 elements - see [8]. This is great news:

- The cyclic group of order 6 must correspond to $\text{Aut}(X_1(13)/X_0(13))$.
- Let σ be any involution of $X_1(13)$ not contained in the cyclic subgroup of order 6 Then σ must be a lift of the Fricke involution.

Explicit formulas for the generators of the automorphism group are computed in [8]. We only need the automorphism of order 6 to compute $X_0(13)$:

$$\alpha_{13}(z, w) = \left(\frac{-1}{1+z}, \frac{w-z}{z+z^2-w} \right) \quad (\text{A13.1})$$

Note that α_{13}^3 coincides with the hyperelliptic involution on $X_1(13)$. Thus, the quotient map $X_1(13) \rightarrow X_1(13)/\langle \alpha_{13}^3 \rangle \cong \mathbb{P}^1$ is simply given by $([z_0 : z_1], w) \mapsto [z_0 : z_1]$. The action of α_{13} on this partial quotient is $[z_0 : z_1] \mapsto [-z_0 : z_0 + z_1]$, or $z \mapsto \frac{-1}{1+z}$ in terms of the affine coordinate $z = \frac{z_0}{z_1}$. Thus, $X_0(13) \cong \mathbb{P}^1 / \langle z \mapsto \frac{-1}{1+z} \rangle$. We can define:

$$q_{13} : X_1(13) \rightarrow X_0(13) \quad q_{13}(z, w) = \frac{z^3 + 4z^2 + z - 1}{z(z+1)} \quad (\text{Q13})$$

$$j_{13}(t) = -\frac{(t^2 - 5t + 13)(t^4 - 7t^3 + 20t^2 - 19t + 1)^3}{t} \quad (\text{J13})$$

Finally, the Fricke involution on this model of $X_0(13)$ is given by:

$$F_{13}(t) = \frac{13}{t} \quad (\text{F13})$$

3.5.2. $X_0(13) \rightarrow X_0(13)^+$. The quotient map $X_0(13) \rightarrow X_0(13)^+$ is:

$$q_{13}^+(t) = -t + 5 - \frac{13}{t} \quad (\text{Q13+})$$

For this choice of q_{13}^+ , the polynomials a_{13}, b_{13} are:

$$\begin{aligned} a_{13}(y) &= y(y^4 - 13y^3 + 13y^2 + 91y + 40) \\ &\cdot (y^8 - 26y^7 + 195y^6 - 156y^5 - 2120y^4 + 624y^3 + 5070y^2 + 1664y - 896) \end{aligned} \quad (\text{RA13})$$

and:

$$b_{13}(y) = y^2(y^4 + 234y^3 + 1417y^2 + 1872y + 832)^3 \quad (\text{RB13})$$

Finally, we compute Δ_{13} :

$$\begin{aligned} \Delta_{13}(j) &= \\ &H_{13}(j) \cdot \left(H_{-1}(j) \cdot H_{-4}(j) \cdot H_{-9}(j) \cdot H_{-3}(j) \cdot H_{-12}(j) \cdot H_{-27}(j) \cdot H_{-48}(j) H_{-43}(j) H_{-51}(j) \right)^2 \end{aligned} \quad (\text{D13})$$

4. ALGORITHMS

In this section, we describe explicit algorithms for computing the supersingular ℓ -isogeny graph for primes $\ell \leq 13$.

4.1. Graph Method. We give an outline of the algorithm described in [4],[9] for computing supersingular isogeny graphs. We will mainly use this algorithm when $\ell = 2$ to obtain the set of supersingular j -invariants.

To get started, we need a model for *any* supersingular curve E_0/\mathbb{F}_{p^2} .

- We find a generator for each subgroup of order ℓ on E_0 .
- For each subgroup of order ℓ , we obtain a model for the isogenous curve using Vélú's formula. This means we now have $E'_0, E'_1, \dots, E'_\ell$.
- We now repeat step 1 for each of $E'_0, E'_1, \dots, E'_\ell$.

4.1.1. $\ell = 2$. In practice, the main challenge is finding generators for the subgroups of order ℓ . When $\ell = 2$, in the worst case scenario, we have to solve a cubic equation to obtain one of the points of order 2. By using carefully chosen models, we only have to find a root of a cubic polynomial once during the entire algorithm.

Every pair (E, P) , where P is a point of order 2, is isomorphic to a pair given by a model of the form:

$$y^2 = x(x^2 + ax + b) \quad P = (0, 0) \quad (\text{E2})$$

The isogenous curve $E/\langle P_0 \rangle$ is given by an equation of the same form:

$$y^2 = x(x^2 - 2ax + (a^2 - 4b))$$

Given a curve of this form, one of the 2-torsion subgroups is generated by $(0, 0)$, and the other two subgroups are generated by $(x_0, 0)$, where x_0 is a root of the quadratic:

$$x^2 + ax + b = 0$$

Altogether, this means we can obtain the complete 2-isogeny graph using Algorithm 1.

Algorithm 1 Supersingular Isogeny Graphs for $\ell = 2$

Require: Coefficients (a, b) of a model of the form E2 for a supersingular curve

Set Edges = $\{\}$

Set Models = $\{(a, b)\}$

Set $J = \{\}$.

while Models $\neq \emptyset$ **do**

 Set NewModels = $\{\}$.

for $(a_0, b_0) \in$ Models **do**

 Compute $j_2(a_0, b_0)$

if $j_2(a_0, b_0) \notin J$ **then**

 Add $j_2(a_0, b_0)$ to the set J .

 Find the roots x_1, x_2 of the quadratic $x^2 + a_0x + b_0$.

for $i \in \{1, 2\}$ **do**

 Compute a new equation for E_0 with $(x_i, 0)$ now at $(0, 0)$.

 Compute the coefficients (a_i, b_i) in equation for E with $(x_i, 0)$ at $(0, 0)$.

 Compute the coefficients $(a'_i, b'_i) = (-2a_i, a_i^2 - 4b_i)$ of the isogenous curve.

 Add the pair (a'_i, b'_i) to NewModels.

end for

end if

end for

 Set Models = NewModels.

end while

4.1.2. $\ell \in \{3, 5, 7, 13\}$. We can use our models for $X_0(\ell)$ to efficiently mimic what we did for $\ell = 2$, as long as we have the ability to factor polynomials over \mathbb{F}_{p^2} .

Instead of coefficients (a, b) , we use points $t \in X_0(\ell)$ to keep track of isomorphism classes of isogenies. We can obtain t' from t using the Fricke involution; the only issue is obtaining the points t_1, \dots, t_ℓ from a point t_0 , where the t_i represent different subgroups of order ℓ on the same elliptic curve. The key is solving the equation $j_\ell(t) - j_\ell(t_0)$. The numerator of this rational function is a polynomial of degree $\ell + 1$ in t , and has a root at $(t - t_0)$. The remaining points t_1, \dots, t_ℓ are the roots of the remaining factor of degree ℓ . For example:

$$j_3(t_0) - j_3(t) = \frac{(t - t_0)}{tt_0} \cdot (t^3t_0 + t^2t_0^2 - 36t^2t_0 + tt_0^3 - 36tt_0^2 + 270tt_0 - 729)$$

Thus, to obtain t_1, t_2, t_3 on $X_0(3)$, we need to solve:

$$(t^3 t_0 + t^2 t_0^2 - 36t^2 t_0 + t t_0^3 - 36t t_0^2 + 270t t_0 - 729) = 0$$

Note that these polynomials will always split completely over \mathbb{F}_{p^2} , as long as $j_\ell(t_0)$ is supersingular, but finding the roots requires more work than solving a quadratic polynomial.

4.1.3. *Implementation.* We implemented Algorithm 1 in Python, and use it to obtain the set of supersingular j -invariants. The code can be found in:

<https://github.com/nhajouji/supsingecs>.

Furthermore, for $\ell \in \{2, 3, 5, 7, 13\}$, the algorithm can easily be implemented in Sage, as the necessary formulas are already built in.⁸

4.2. **Parallelizable algorithms.** Next, we describe simple algorithms for computing the ℓ -isogeny graph for all odd primes $\ell \leq 13$. While these algorithms are less mathematically sophisticated, they present the following practical advantages:

- The algorithms are simple enough that they can be easily implemented on Python, without requiring any additional packages. A Python script that does this can be found at:
<https://github.com/nhajouji/supsingecs>.
- The computations in this algorithm can be done in parallel. In particular, the program can be dramatically sped up if one does the computations on a GPU.

4.2.1. $g(X_0(\ell)) = 0$. To compute the graph for $\ell \in \{3, 5, 7, 13\}$, we use Algorithm 2. This algorithm assumes that we've already computed the set S_p of supersingular j -invariants, e.g. using Algorithm 1. As long as we've already obtained this set, all we need is the formulas for j_ℓ that we obtained in the previous section (J3,J5,J7,J13), as well as the formula⁹ for the Fricke involution F_ℓ . Note that the cusps are at $0, \infty$, so we can evaluate $j_{\ell,0}(t)$ at any point in \mathbb{F}_{p^2} without having to worry about dividing by 0.

Algorithm 2 Supersingular Isogeny Graphs from $X_0(\ell)$ ($\ell \in \{3, 5, 7, 13\}$)

Require: $S_p = \{j \in \mathbb{F}_{p^2} : j \text{ supersingular}\}$.

Set Edges = $\{\}$

for $t \in \mathbb{F}_{p^2}^\times$ **do**

 Compute $j_{\ell,0}(t)$

if $j_{\ell,0}(t) \in S_p$ **then**

 Compute $j'_{\ell,0}(t)$.

 Record the edge $(j_{\ell,0}(t), j'_{\ell,0}(t))$.

end if

end for

Warning: We have to be careful when $p \not\equiv 1 \pmod{12}$: we might have multiple isogenies $E \rightarrow E'$ that are represented by a single point on $X_0(\ell)$ if $j(E) \in \{0, 1728\}$.

⁸https://doc.sagemath.org/html/en/reference/arithmetic_curves/sage/schemes/elliptic_curves/isogeny_sma

⁹The Fricke involution has the form $t \mapsto \frac{c}{t}$ for all of these values of ℓ , so all we really need to remember is the constant c . Similarly, the denominator of j_ℓ is t so we only need to remember the numerator.

- If $j_{\ell,0}(t) = 0$ and $j'_{\ell,0}(t) \neq 0$, the edge $(0, j'_{\ell,0}(t))$ should be recorded 3 times.
- If $j_{\ell,0}(t) = 1728$ and $j'_{\ell,0}(t) \neq 0$, the edge $(1728, j'_{\ell,0}(t))$ should be recorded twice.

4.2.2. $X_0(11)$. For $\ell = 11$, we can use the same ideas, but there are additional steps due to the fact that we can't simply list all points on a curve of positive genus the way we can with a curve of genus 0.

Algorithm 3 Supersingular Isogeny Graphs from $X_0(\ell)$ ($\ell = 11$)

Require: $S_p = \{j \in \mathbb{F}_{p^2} : j \text{ supersingular}\}$.
 Define a set Edges = $\{\}$
for $x_0 \in \mathbb{F}_{p^2} - \{5, 16\}$ **do**
 Find all points on $(x_0, y_i) \in X_0(11)(\mathbb{F}_{p^2})$.
 for $(x_0, y_i) \in X_0(11)(\mathbb{F}_{p^2})$ **do**
 Compute $j_{11}(x_0, y_i)$.
 if $j_{11}(x_0, y_i) \in S_p$ **then**
 Compute $j_{11} \circ F_{11}(x_0, y_i)$.
 Record the edge $(j_{11}(x_0, y_i), j_{11} \circ F_{11}(x_0, y_i))$.
 end if
 end for
end for

As with $\ell = 3, 5, 7, 13$, we have to count edges with the appropriate multiplicities when $j = 0, 1728$.

4.2.3. $X_0(\ell)$ *general*. We can adapt Algorithm 3 to work for all values of ℓ , as long as we have models for $X_0(\ell)$, together with formulas for j_ℓ, F_ℓ . Precisely, if $X_0(\ell)$ is the zero set of $G(z, w) = 0$, we can use Algorithm 4.

Algorithm 4 Supersingular Isogeny Graphs from $X_0(\ell)$ (General ℓ)

Require: $S_p = \{j \in \mathbb{F}_{p^2} : j \text{ supersingular}\}$.
 Define a set Edges = $\{\}$
for $z_0 \in \mathbb{F}_{p^2}$ **do**
 Find all roots w_0, w_1, \dots of $G(z_0, w) = 0$.
 for $(z_0, w_i) \in X_0(\ell)(\mathbb{F}_{p^2})$ **do**
 Compute $j_\ell(z_0, w_i)$.
 if $j_\ell(z_0, w_i) \in S_p$ **then**
 Compute $j_\ell \circ F_\ell(z_0, w_i)$.
 Record the edge.
 end if
 end for
end for

4.2.4. $g(X_0(\ell)^+) = 0$. We can also compute isogeny graphs by using $X_0(\ell)^+$ instead of $X_0(\ell)$. This is not particularly helpful when $X_0(\ell)$ already has genus 0- this algorithm is about as complicated as the algorithm for $\ell = 11$. However, we can use the same version of the algorithm for all supersingular primes ℓ - the only thing that changes when we vary ℓ is the pair of polynomials a_ℓ, b_ℓ we use. Furthermore, the polynomials $R_\ell(x, y)$ are easier to obtain via alternative means than j_ℓ when ℓ is a supersingular prime and $X_0(\ell)$ has positive genus.

Instead of evaluating j_ℓ , we evaluate the polynomials a_ℓ, b_ℓ at each point in $y_0 \in \mathbb{F}_{p^2}$. Each y_0 gives us a quadratic $x^2 - a_\ell(y_0)x + b_\ell(y_0)$, whose roots are ℓ -isogenous j -invariants. To compute the isogeny graph, we simply check whether the roots of each quadratic are supersingular, and if they are, we record them in the graph.

Now, we have to be careful when using $X_0(\ell)^+$ because the models we are using have singularities. For example, the order of vanishing of $R_5(x, y)$ at $(-884736, 36)$ is 2:

- The polynomial $R_5(x, 36)$ has a repeated root at -884736 . This means there is an isogeny of degree 5 from the curve with j -invariant 884736 to itself.
- In fact, there are *two* such isogenies - but they are both represented by the point $y = 36$. To see this, we can factor the polynomial $R_5(-884736, y)$ and check that the factor $(y - 36)$ appears with multiplicity 2.

Fortunately, we can use the factorizations of the polynomials Δ_ℓ to determine the correct number of endomorphisms of degree ℓ for each supersingular curve mod p . This is easy - we just need to compute the roots of each factor of Δ_ℓ in \mathbb{F}_{p^2} and check if they are supersingular.

We can simplify this step by grouping together factors whose roots are j -invariants of elliptic curves that are isogenous in characteristic 0. For $d \in \mathbb{N}$, write \mathcal{O}_d to denote the ring of integers of $\mathbb{Q}(\sqrt{-d})$ and $\text{Nrm}_d : \mathcal{O}_d \rightarrow \mathbb{Z}$ for the norm map. Define:

$$\text{supp}(\ell) = \{-d : \ell \in \text{Nrm}_d(\mathcal{O}_d)\}$$

Note that $\text{supp}(\ell)$ is finite, e.g. for the values of ℓ we are interested, we have:

$$\begin{aligned} \text{supp}(3) &= \{-2, -3, -11\} \\ \text{supp}(5) &= \{-1, -5, -11, -19\} \\ \text{supp}(7) &= \{-3, -6, -7, -19\} \\ \text{supp}(11) &= \{-2, -7, -10, -11, -19, -35, -43\} \\ \text{supp}(13) &= \{-1, -3, -13, -43, -51\} \end{aligned}$$

The values $-d \in \text{supp}(\ell)$ coincide with the subscripts that appear in the factors H_{-d} and H_{-dn^2} of Δ_ℓ . For each pair $\ell, -d$, where $-d \in \text{supp}(\ell)$, we write $\Delta_{\ell, -d}(j)$ to denote the product of the factors of Δ_ℓ whose roots are j -invariants of elliptic curves in the same isogeny as the elliptic curves whose j -invariants are roots of $H_{-d}(j)$.

For a pair p, ℓ , we set:

$$\begin{aligned} \text{supp}_p(\ell) &= \left\{ -d \in \text{supp}(\ell) : \left(\frac{-d}{p} \right) = -1 \right\} \\ \Delta_{\ell, p}(j) &= \prod_{-d \in \text{supp}_p(\ell)} \Delta_{\ell, -d}(j) \end{aligned}$$

Algorithm 5 uses $\Delta_{ell,p}(j)$ to compute the diagonal of the adjacency matrix of the isogeny graph.

Algorithm 5 Supersingular Isogeny Graphs from $X_0(\ell)^+$: Diagonal

Require: $\Delta_\ell(j) = \prod \Delta_{\ell,-d}(j)$.

Set Edges = {}.

for $-d \in \text{supp}_p(\ell)$ **do**

for Each factor $H_{-dn^2}^{e_n}$ of $\Delta_{\ell,-d}$ **do**

 Compute the set of roots $Z(H_{-dn^2}) \subset \mathbb{F}_{p^2}$.

for $j \in Z(H_{-dn^2})$ **do**

 Add e_n copies of (j, j) to Edges.

end for

end for

end for

To compute the rest of the isogeny graph, we use Algorithm 6.

Algorithm 6 Supersingular Isogeny Graphs from $X_0(\ell)^+$:Off-Diagonal

Require: $a_\ell(y), b_\ell(y)$.

Require: $S_p = \{j \in \mathbb{F}_{p^2} : j \text{ supersingular}\}$.

for $y_0 \in \mathbb{F}_{p^2}$ **do**

 Evaluate $a_\ell(y_0), b_\ell(y_0)$.

 Compute the discriminant $d_\ell(y_0) = a_\ell(y_0)^2 - 4b_\ell(y_0)$.

if $d_\ell(y_0) \neq 0$ and $\sqrt{d_\ell(y_0)} \in \mathbb{F}_{p^2}$ **then**

 Compute the roots j_0, j'_0 of $R_\ell(x, y_0) = 0$.

if $j_0, j_1 \in S_p$ **then**

 Add $(j_0, j'_0), (j_0, j'_0)$ to the set of edges.

end if

end if

end for

4.3. **Resultants.** For a prime p , we define:

$$ss_p(x) = \prod_{j \in S_p} x - j$$

The polynomial ss_p vanishes if and only if x is a supersingular j -invariant. For a pair of polynomials $f(x), g(x)$, we write $\text{Res}_x(f, g)$ to denote the resultant of f, g .

Whenever we have a modular curve of genus 0, we can obtain a characterization of the set of points on the modular curve that represent supersingular curves by computing the resultant of $ss_p(x)$ with a suitable polynomial.

- Let $\ell \in \{3, 5, 7, 13\}$. Then $j_\ell(t) \cdot t$ is a polynomial in t .

To obtain the set of points on $X_0(\ell)$ whose j -invariants are supersingular, we can compute:

$$\text{Res}_x(ss_p(x), t \cdot (j_\ell(t) - x))$$

The result is a polynomial in t whose roots are precisely the points on $X_0(\ell)$ that represent supersingular elliptic curves.

- We can do something similar for all supersingular primes using the polynomial $R_\ell(x, y)$. Precisely, we compute:

$$\text{Res}_x(ss_p(x), R_\ell(x, y))$$

to obtain a polynomial in y , whose roots are precisely the values y_0 for which the roots of $R_\ell(x, y_0)$ are supersingular.

The roots of the resultant will all lie in the image of $q_\ell^+(X_0(\ell)(\mathbb{F}_q))$, so the resultant will split completely over \mathbb{F}_{p^2} .¹⁰ To take full advantage of this, we need a way of factoring the resultant over \mathbb{F}_p as a product of polynomials of degree at most 2. Fortunately, most computer algebra systems can compute resultants and factor polynomials over \mathbb{F}_p .

5. APPLICATIONS

5.1. Trace Formula. Let $\ell \in \{3, 5, 7, 11, 13\}$, let $p \neq \ell$ be a prime, and let $\Gamma_{p,\ell}$ be the supersingular ℓ -isogeny graph. We write $Tr(\Gamma_{p,\ell})$ to denote the trace of the adjacency matrix of $\Gamma_{p,\ell}$. Algorithm 5 allows us to compute the diagonal of the adjacency graph directly, without needing to compute the rest of the isogeny graph, so we can use it to deduce a formula for $Tr(\Gamma_{p,\ell})$.

It's clear that $Tr(\Gamma_{p,\ell}) = \deg(\Delta_{p,\ell}(j))$. Furthermore, the degree of $\Delta_{p,\ell}$ is determined by the Jacobi symbols $\left(\frac{-2}{p}\right)$ for $-d \in \text{supp}(\ell)$. Thus, we can obtain a formula for the trace as:

$$Tr(\Gamma_{p,\ell}) = \sum_{-d \in \text{supp}(\ell)} m_{\ell,-d} \left(\frac{-d}{p}\right)$$

for some integers $m_{\ell,-d}$ that depend on ℓ .

For example, when $\ell \leq 13$, we obtain the following formulae:

$$\begin{aligned} \text{Tr}(\Gamma_{p,3}) &= 3 - \left(\left(\frac{-2}{p}\right) + \left(\frac{-3}{p}\right) + \left(\frac{-11}{p}\right) \right) \\ \text{Tr}(\Gamma_{p,5}) &= 5 - \left(2 \left(\frac{-1}{p}\right) + \left(\frac{-5}{p}\right) + \left(\frac{-11}{p}\right) + \left(\frac{-19}{p}\right) \right) \\ \text{Tr}(\Gamma_{p,7}) &= 7 - \left(3 \left(\frac{-3}{p}\right) + 2 \left(\frac{-6}{p}\right) + \left(\frac{-7}{p}\right) + \left(\frac{-19}{p}\right) \right) \\ \text{Tr}(\Gamma_{p,11}) &= 11 - \left(\left(\frac{-2}{p}\right) + 2 \left(\frac{-7}{p}\right) + 2 \left(\frac{-10}{p}\right) + 2 \left(\frac{-11}{p}\right) + \left(\frac{-19}{p}\right) + 2 \left(\frac{-35}{p}\right) + \left(\frac{-43}{p}\right) \right) \\ \text{Tr}(\Gamma_{p,13}) &= 13 - \left(4 \left(\frac{-1}{p}\right) + 5 \left(\frac{-3}{p}\right) + \left(\frac{-13}{p}\right) + \left(\frac{-43}{p}\right) + 2 \left(\frac{-51}{p}\right) \right) \end{aligned}$$

We expect that there are similar formulas for all primes ℓ , not just the supersingular primes.

¹⁰We discuss this in more detail in 5.2.

5.2. Nakaya's Conjectures. The resultants we used at the end of 4.3 play a central role in [12]. Consequently, we can translate many of the questions in that paper into questions about supersingular isogeny graph. In particular, we will use this perspective give a simple proof of Conjecture 2 in that paper.¹¹

The actual object of the conjectures in [12] is the polynomial:

$$ss_{p,\ell}^*(y) = \text{rad}(\text{Res}_x(ss_p(x), R_\ell(x, y)))$$

Here, $\text{rad}(-)$ indicates that we are ignoring the multiplicity of each irreducible factor. Now, the factors of $\text{Res}_x(ss_p(x), R_\ell(x, y))$ correspond to (pairs) of edges on the isogeny graph, so we can think of $ss_{p,\ell}^+(y)$ as representing the underlying undirected graph. This perspective makes it easier to explain some of the phenomena observed in [12].

As a warm-up, we will use Theorem 2.2 to explain why $ss_{p,\ell}^+(y)$ always splits over \mathbb{F}_{p^2} . This will answer Question 1 in [12].

Suppose $y_0 \in \mathbb{F}_{p^2}$ is a root of $ss_{p,\ell}^+(y) = 0$. That means that the roots x_1, x_2 of $R_\ell(x, y_0) = 0$ are supersingular j -invariants that are related by an isogeny of degree ℓ , so there exists $t_0 \in X_0(\ell)(\mathbb{F}_{p^2})$ such that $j_\ell(t_0) = x_1$ and $j_\ell \circ F_\ell(t_0) = x_2$, and $y_0 = q_\ell^+(t_0)$. Since $t_0 \in \mathbb{F}_{p^2}$ by Theorem 2.2, and q_ℓ^+ is defined over \mathbb{F}_p , it follows that $y_0 \in \mathbb{F}_{p^2}$. This proves that $ss_{p,\ell}^+$ splits over \mathbb{F}_{p^2} .

We can use similar ideas to prove that $ss_{p,\ell}$ has a non-linear factors whenever p is *not* a supersingular prime. This will allow us to reduce the proof of Conjecture 2 in [12] a finite computation. Furthermore, we will be able to prove additional results in the same vein.

Theorem 5.1. *Let ℓ be a supersingular prime, and let p be a prime which is **not** supersingular. Then $ss_{p,\ell}(y)^+$ has an irreducible factor of degree > 1 .*

Proof. Let $S_{p,1} = S_p \cap \mathbb{F}_p$ and $S_{p,2} = S_p \setminus \mathbb{F}_p$. Note that $S_{p,1}$ is always nonempty, because there are always supersingular curves defined over \mathbb{F}_p , and $S_{p,2}$ is nonempty because p is not supersingular. By connectedness of the ℓ -isogeny graph, we can find $j_1 \in S_{p,1}$ and $j_2 \in S_{p,2}$ such that the elliptic curves with those invariants are ℓ -isogenous. This means there exists $t_0 \in \mathbb{F}_{p^2}$ such that $j_\ell(t_0) = j_1$ and $j_\ell \circ F_\ell(t_0) = j_2$.

Let y_0 be the image of t_0 under the map $X_0(\ell) \rightarrow X_0(\ell)^+$. Then the roots of $R_\ell(x, y_0)$ are j_1, j_2 , so y_0 is a root of $\text{Res}_x(ss_p(x), R_\ell(x, y)) = 0$. Furthermore, the roots of $R(x, y_0)$ are not Galois conjugates, so $a_\ell(y), b_\ell(y) \notin \mathbb{F}_p$, so $y_0 \notin \mathbb{F}_p$. This means the minimal polynomial of y_0 , which has degree > 1 , divides $ss_{p,\ell}(y)$. □

Consequently, we see that $ss_{p,\ell}^+$ can only split over \mathbb{F}_p if p is also supersingular. Thus, to determine the complete set of primes p for which $ss_{p,\ell}^+$ splits over \mathbb{F}_p , we just have to compute $ss_{p,\ell}^+$ for the 15 supersingular primes. For $\ell = 5, 7, 11, 13$, we obtained the following sets of

¹¹We note that some of the conjectures were recently proven in [10] using different methods.

primes:

$$\ell = 5 : \quad p \in \{2, 3, 5, 7, 11, 19\}$$

$$\ell = 7 : \quad p \in \{2, 3, 5, 7, 17\}$$

$$\ell = 11 : \quad p \in \{2, 3, 5, 11\}$$

$$\ell = 13 : \quad p \in \{2, 3, 13\}$$

Now, Conjecture 2 predicts that the set of primes for $\ell = 5$ coincides with the set of primes that divide the Hamada-Norton group; and it predicts that the set of primes for $\ell = 7$ will coincide with the set of primes that divide the Held group. Our computation shows that these predictions are correct. The correspondence between sporadic simple groups and modular curves $X_0(\ell)$ is explained in Section 9 of [5]. We note that the sporadic group associated to $X_0(11)$ is the Mathieu group M_{12} , which has cardinality $2^6 \cdot 3^3 \cdot 5 \cdot 11$. As with $\ell = 5, 7$, the set of primes that divides M_{12} coincides with the set of primes for which $ss_{p,11}$ splits over \mathbb{F}_p . Finally, we mention that the correspondence between sporadic groups and modular curves is only known for $\ell \leq 11$.¹²

6. CONCLUSION

6.1. Implementation and tables. Tables containing supersingular isogeny graphs (for $p \leq 512$) and adjacency matrices (for $p \leq 2048$) were computed by implementing the algorithms in this paper using Python. The results and code can be found at: <https://github.com/nhajouji/supsingecs>.

6.2. Generalizations. The algorithms we have described for primes $\ell \leq 13$ can be extended to higher values of ℓ , as long as we have analogs of some of the formulas derived in 3.

- The algorithms that use $X_0(\ell)^+$ only work for supersingular primes ℓ . However, they can be used without modification, as long as we know the polynomials a_ℓ, b_ℓ .

We note that the polynomials a_ℓ, b_ℓ can be obtained using analytic methods, although they require knowledge of the Fourier coefficients of a Hauptmodul of $X_0(\ell)^+$. See [3].

- The algorithms using $X_0(\ell)$ can also be used for all values of ℓ , but we need models of the modular curve, together with formulas for the j -map and the Fricke involution.

The complexity of these algorithms will hinge on how hard it is to find all points on $X_0(\ell)$. For example $X_0(\ell)$ is hyperelliptic¹³, we can essentially mimic Algorithm 3.

However, we can go beyond the set of supersingular primes ℓ : for example, if $X_0(\ell)$ has gonality ≤ 4 ,¹⁴ we can use the cubic/quartic formula to compute the fibers of $X_0(\ell) \rightarrow \mathbb{P}^1$, and obtain an algorithm with the same complexity as 3.

The curve $X_0(43)$ is 3-gonal, and $X_0(\ell)$ has gonality 4 for the following values of ℓ ¹⁵:

$$67, 73, 103, 107$$

¹²This is due to the fact that $PSL(2, p)$ only has a subgroup of index p when $p \in \{2, 3, 5, 7, 11\}$; see [5] for further details.

¹³Note that $X_0(\ell)$ is hyperelliptic precisely when ℓ is a supersingular prime.

¹⁴The idea of using gonality of the modular curve as a threshold came from [15].

¹⁵See [11].

We can also obtain efficient algorithms when $X_0(\ell)$ is bielliptic. The primes values ℓ for which $X_0(\ell)$ is bielliptic are:¹⁶:

$$37, 53, 61, 79, 83, 89, 101, 131$$

6.3. Acknowledgements. The author would like to thank Christian Bueno for helpful discussions.

APPENDIX A. HILBERT CLASS POLYNOMIALS

The formulae for Δ_ℓ in Section 3 are expressed as products of polynomials $H_{-dn^2}(j)$. We give explicit formulas for those factors in this section.

The polynomials labelled $H_{-d}(j)$, where d is a squarefree integer, are the Hilbert class polynomials for $\mathbb{Q}(\sqrt{-d})$. For example, when d is a Heegner number, we have:

$$H_{-1}(j) = j - 1728 \tag{H1}$$

$$H_{-2}(j) = j - 8000 \tag{H2}$$

$$H_{-3}(j) = j \tag{H3}$$

$$H_{-7}(j) = j + 3375 \tag{H7}$$

$$H_{-11}(j) = j + 32768 \tag{H11}$$

$$H_{-19}(j) = j + 884736 \tag{H19}$$

$$H_{-43}(j) = j + 884736000 \tag{H43}$$

$$H_{-67}(j) = j + 147197952000 \tag{H67}$$

$$H_{-163}(j) = j + 262537412640768000 \tag{H163}$$

Note that H67, H163 do not appear as factors of any of the Δ_ℓ , but they can be used to obtain a supersingular curve E/\mathbb{F}_p to start Algorithm 1.

The remaining Hilbert class polynomials of positive degree that appear in the Δ_ℓ are:

$$H_{-5}(j) = j^2 - 1264000j - 681472000 \tag{H5}$$

$$H_{-6}(j) = j^2 - 4834944j + 14670139392 \tag{H6}$$

$$H_{-10}(j) = j^2 - 425692800j + 9103145472000 \tag{H10}$$

$$H_{-13}(j) = j^2 - 6896880000j - 567663552000000 \tag{H13}$$

$$H_{-35}(t) = j^2 + 117964800j - 134217728000 \tag{H35}$$

$$H_{-51}(j) = j^2 + 5541101568j + 6262062317568 \tag{H51}$$

Note that additional Hilbert class polynomials can be obtained in the same way using the modular curves $X_0(\ell)$ for $\ell = 4, 6, 8, 9, 10, 12$. The following three have odd degree, which means they are guaranteed to have at least one root in \mathbb{F}_p (provided $-d$ is a nonsquare in \mathbb{F}_p):

$$H_{-23}(j) = j^3 + 3491750j^2 - 5151296875j + 12771880859375 \tag{H23}$$

$$H_{-31}(j) = j^3 + 39491307j^2 - 58682638134j + 1566028350940383 \tag{H31}$$

¹⁶See [2].

Finally, we note that there are a few factors that appear in Δ_ℓ which are not Hilbert class polynomials. We have labelled these as H_{-dn^2} to indicate that the roots of $H_{-dn^2}(j) = 0$ are n -isogenous to the roots of $H_{-d}(j) = 0$. Note that the roots of $H_{-dn^2}(j)$ are supersingular mod p if and only if $-d$ is a quadratic nonresidue mod p .

The polynomials of this form that appear in the Δ_i are:

$$H_{-4}(j) = j - 287496 \tag{H1.2}$$

$$H_{-9}(j) = j^2 - 153542016j - 1790957481984 \tag{H1.3}$$

$$H_{-12}(j) = j - 54000 \tag{H3.2}$$

$$H'_{-27}(j) = j - 12288000 \tag{H3.3}$$

$$H_{-48}(j) = j^2 - 2835810000j + 6549518250000 \tag{H3.4}$$

$$H_{-44}(j) = j^3 - 1122662608j^2 + 270413882112j - 653249011576832 \tag{H11.2}$$

REFERENCES

[1] Houria Baaziz. Equations for the modular curve $X_1(N)$ and models of elliptic curves with torsion points. 79(272):2371–2386, October 2010.

[2] Francesc Bars. Bielliptic modular curves. *Journal of Number Theory*, 76(1):154–165, May 1999.

[3] Alin Bostan, Salah Boukraa, Gilles Christol, Saoud Hassani, and J-M Maillard. Ising n -fold integrals as diagonals of rational functions and integrality of series expansions. *Journal of Physics A: Mathematical and Theoretical*, 46, 2012.

[4] Denis Charles and Kristin Lauter. Computing modular polynomials. *LMS J. Comput. Math.*, 8:195–204, 2005.

[5] J. H. Conway and S. P. Norton. Monstrous moonshine. *Bulletin of the London Mathematical Society*, 11(3):308–339, 1979.

[6] David A. Cox. *Primes of the form $x^2 + ny^2$* . A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1989. Fermat, class field theory and complex multiplication.

[7] Matt Delong. A formula for the Selmer group of a rational three-isogeny. *Acta Arithmetica*, 105:119–131, 1999.

[8] Daeyeol Jeon. Automorphism groups of hyperelliptic modular curves. *Proceedings of the Japan Academy, Series A, Mathematical Sciences*, 91(7), 2015.

[9] J.-F. Mestre. The method of graphs. Examples and applications. Class numbers and fundamental units of algebraic number fields, Proc. Int. Conf., Katata/Jap. 1986, 217-242 (1986)., 1986.

[10] Patrick Morton. Supersingular conjectures for the Fricke group. *International Journal of Number Theory*, 19(04):803–828, October 2022.

[11] Filip Najman and Petar Orlić. Gonality of the modular curve $X_0(n)$, 2022.

[12] Tomoaki Nakaya. The number of linear factors of supersingular polynomials and sporadic simple groups. *Journal of Number Theory*, 2018.

[13] René Schoof. Nonsingular plane cubic curves over finite fields. *J. Combin. Theory Ser. A*, 46(2):183–211, 1987.

[14] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1986.

[15] Andrew V. Sutherland. Computing hilbert class polynomials with the chinese remainder theorem. *Math. Comput.*, 80:501–538, 2009.

[16] J. Velu. Isogenies entre courbes elliptiques. *Comptes-Rendus de l'Academie des Sciences*, 273:238–241, 1971.

[17] William C. Waterhouse. Abelian varieties over finite fields. *Ann. Sci. École Norm. Sup. (4)*, 2:521–560, 1969.