

AFFINE GRASSMANNIANS FOR G_2

ZHIHAO ZHAO

ABSTRACT. We study affine Grassmannians for the exceptional group of type G_2 . This group can be given as automorphisms of octonion algebras (or para-octonion algebras). By using this automorphism group, we consider all maximal parahoric subgroups in G_2 , and give a description of affine Grassmannians for G_2 as functors classifying suitable orders in a fixed space.

CONTENTS

1. Introduction	1
2. Octonion Algebras and Para-Octonion Algebras	4
3. The Building of G_2	8
4. Affine Grassmannians for \mathcal{G}	11
5. Proof of the First Main Theorem	13
6. Affine Grassmannians for Other Parahoric Subgroups	16
References	22

1. INTRODUCTION

In this paper we give an explicit description of affine Grassmannians for the exceptional group G_2 . Generally, affine Grassmannians can be defined by loop groups. Let k be a field, with $\text{char}(k) \neq 2$, and let G be a linear algebraic group over $\text{Spec}(k((t)))$, where $k((t))$ is the Laurent power series with variable t . In [4], [5], Bruhat-Tits attach to any connected reductive group G over $\text{Spec}(k((t)))$ its building $\mathcal{B}(G)$, which is a polysimplicial complex equipped with an action of $G(k((t)))$. To any point $x \in \mathcal{B}(G)$, there is a unique smooth affine group scheme \mathcal{G}_x over $\text{Spec}(k[[t]])$, where $k[[t]]$ is the formal power series with variable t . We call it the parahoric subgroup associated to the point x . The $k[[t]]$ -points $\mathcal{G}_x(k[[t]])$ is (at least when G is simply connected) the stabilizer of x in $G(k((t)))$. Its generic fiber is equal to G . We define the algebraic loop group LG associated to G , which is the ind-scheme representing the functor:

$$R \mapsto LG(R) = G(R((t))),$$

for any k -algebra R . Since $R((t))$ is a $k((t))$ -algebra, this definition makes sense. We also define the positive loop group $L^+\mathcal{G}_x$ associated to \mathcal{G}_x , representing the functor:

$$R \mapsto L^+\mathcal{G}_x(R) = \mathcal{G}_x(R[[t]]).$$

for any k -algebra R . Thus, $L^+\mathcal{G}_x \subset LG$ is a subgroup functor, and the fpqc-quotient

$$\text{Gr}_{\mathcal{G}_x} := LG/L^+\mathcal{G}_x$$

is by definition the affine Grassmannian associated to \mathcal{G}_x . This definition is given by Pappas-Rapoport in [15], where they develop a theory of twisted loop groups and of their associated flag varieties. We refer [2], [3] and [15] for important results in the general theory of affine Grassmannians.

We now take G to be the exceptional group G_2 and study its corresponding affine Grassmannians. Our goal (see Theorem 1.1, 1.2) is to give a description of these affine Grassmannians in terms of lattices with certain extra conditions. A sketch of the history of this problem is the following: Lusztig in [13] first shows that affine Grassmannians for simple Lie algebras can be described in terms of lattices that are closed under the Lie bracket. Our goal here is more in line of which is known for classical groups. The prototype is the general linear group \mathbf{GL}_n . It is well known that affine Grassmannians for \mathbf{GL}_n can be viewed as a space of lattices in a fixed n -dimensional vector space. For other classical groups, Pappas-Rapoport give a description of affine Grassmannians and affine flag varieties for unitary groups in [15] using lattices (or lattice chains) which are self-dual for a hermitian form; Görtz considers the symplectic group case in [11]; Smithling works out the case of split orthogonal groups in [18]; In [22], the author describes affine Grassmannians for triality groups, which are groups of type 3D_4 . It turns out that our results here are similar to [22], and different from other classical group cases. One reason is that the exceptional group G_2 is constructed by octonion algebras, instead of usual vector spaces with quadratic forms.

The exceptional group G_2 can be viewed as an automorphism group of (split) octonion algebras. These algebras are neither commutative nor associative, and are equipped with a nonsingular quadratic forms q . Briefly speaking, an octonion algebra C is such an 8-dim algebra with identity element satisfying $q(x \cdot y) = q(x)q(y)$. Generally, algebras satisfying $q(x \cdot y) = q(x)q(y)$ are called composition algebras. An octonion algebra C is a composition algebra with highest dimension by the classification theorem (see Theorem 2.5 below). Springer in [19, §2.3] shows that the automorphism group $\mathbf{Aut}(C)$ is of type G_2 . We will see that there is a natural embedding of $\mathbf{Aut}(C)$ to the orthogonal group $\mathbf{SO}(C)$, since the multiplication of C determines the quadratic form q . In [9], Gan-Yu show that this embedding gives rise to a canonical embedding of buildings $\mathcal{B}(\mathbf{Aut}(C)) \rightarrow \mathcal{B}(\mathbf{SO}(C))$.

Another ingredient we need is a parahoric subgroup \mathcal{G}_x associated to $x \in \mathcal{B}(G_2)$. In [9], Gan-Yu study the building of G_2 , and give a nice classification of all maximal parahoric subgroups of G_2 (see [9, Theorem 9.5]). There are three types of maximal parahoric subgroups corresponding to three types of vertices in the building $\mathcal{B}(G_2)$ (see Theorem 3.6 below). The first type is the parahoric subgroup associated to hyperspecial points. By fixing the parahoric subgroup $\mathcal{G} = \mathcal{G}_x$, with the generic fiber $\mathcal{G}_\eta = G_2$, our first main theorem is the following:

Theorem 1.1. *There is an $L\mathcal{G}_\eta$ -equivariant isomorphism*

$$\mathrm{Gr}_{\mathcal{G}} \simeq \mathcal{F},$$

where the functor \mathcal{F} sends a k -algebra R to the set of finitely generated projective $R[[t]]$ -modules L (i.e., $R[[t]]$ -lattices) of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is self dual under the bilinear form $\langle \cdot, \cdot \rangle$, i.e., the form induces an isomorphism $L \simeq \mathrm{Hom}_{R[[t]]}(L, R[[t]])$.
- (2) L is an order in $C_s \otimes_{k((t))} R((t))$, i.e., $e \in L$, $L \star L \subset L$.

Here C_s is the split para-octonion algebra over $k((t))$ obtained from C , and \star is the multiplication in C_s . The element $e \in C_s$ is the para-unit. We refer §2 for more details about composition algebras. This theorem is shown in §5.

For the other two types of parahoric subgroups, we have similar results. By fixing the parahoric subgroups \mathcal{G}_2 (resp. \mathcal{G}_3) in $G = G_2$ (see Definition 6.3), our second main theorem is the following:

Theorem 1.2. (a). *There is an LG-equivariant isomorphism*

$$\mathrm{Gr}_{\mathcal{G}_2} \simeq \mathcal{F}_2$$

where the functor \mathcal{F}_2 sends a k -algebra R to the set of $R[[t]]$ -lattices L of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is an order in $C_s \otimes_{k((t))} R((t))$, i.e., $e \in L$, $L \star L \subset L$.
- (2) $L \subsetneq L^\vee \subsetneq t^{-1}L$.
- (3) $L^\vee \star L^\vee \subset t^{-1}L$.

(b). *There is an LG-equivariant isomorphism*

$$\mathrm{Gr}_{\mathcal{G}_3} \simeq \mathcal{F}_3$$

where the functor \mathcal{F}_3 sends a k -algebra R to the set of $R[[t]]$ -lattices L of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is an order in $C_s \otimes_{k((t))} R((t))$.
- (2) $L \subsetneq L^\vee \subsetneq t^{-1}L$.
- (3) $M = tL^\vee \star L^\vee + L$ is self-dual.

Here $L^\vee = \{x \in C_s \otimes_{k((t))} R((t)) \mid \langle x, L \rangle \subset R[[t]]\}$ is the dual lattice of L .

Octonions algebras play an important role in the study of other exceptional groups. We can construct Albert algebras by regarding their elements as 3×3 matrices (x_{ij}) with x_{ij} in the field k or in C . The automorphism group of an Albert algebra is of type F_4 , and the group of transformations that leave the cubic form \det invariant is of type E_6 (see [19, Ch. 7]). We hope to use the work here to obtain explicit descriptions of affine Grassmannians for other exceptional groups.

The organization of the paper is as follows: In §2, we review definitions and basic propositions of composition algebras. We are particular interested in composition algebras of dimension 8, which are octonion algebras and para-octonion algebras. We give a nice construction of octonion algebras, and consider their isotropic subspaces. In §3, we recall the building theory of group G_2 . We first show that there is an action of S_3 on the spin group of the (split) para-octonion algebra, and the subgroup of this spin group, which is fixed under S_3 , is the automorphism group G of type G_2 (see Proposition 3.3). By using this relation, we study the building of G and classify vertices in $\mathcal{B}(G)$. These vertices corresponds to the maximal parahoric subgroups. We refer [9] for more details. In §4, we construct affine Grassmannians for G_2 , and give our first main theorem. The proof of Theorem 1.1 is in §5. In the last section, we fix the other two maximal parahoric subgroups \mathcal{G}_2 (resp. \mathcal{G}_3) and construct affine Grassmannians $\mathrm{Gr}_{\mathcal{G}_2}$ (resp. $\mathrm{Gr}_{\mathcal{G}_3}$). By using the classification of quadratic forms over local fields (see [10]), we prove Theorem 1.2.

I thank G. Pappas for his valuable comments, patient help, and corrections that greatly improved this paper.

2. OCTONION ALGEBRAS AND PARA-OCTONION ALGEBRAS

In this section, we will introduce unital composition algebras and symmetric composition algebras. These algebras are often not commutative or associative. We refer [12, Ch. VIII], [19, Ch. 1] for general facts and more details. The octonion algebra (resp. para-Octonion algebra) is a unital composition algebra (resp. symmetric composition algebra) with dimension 8. We will see that they have a close connection with each other.

Let F be a field, and suppose $\text{char}(F) \neq 2$. In this and the following sections, by an F -algebra A we mean (unless further specified) a finite dimensional vector space over F equipped with an F -bilinear multiplication. The multiplication here is not necessarily associative, and we do not assume that A has an identity element. Consider an algebra A over F with a quadratic form $q : A \rightarrow F$. We always assume that q is nonsingular, i.e., the bilinear form $\langle \cdot, \cdot \rangle$ associated to q :

$$\langle x, y \rangle := q(x + y) - q(x) - q(y),$$

for all $x, y \in A$ has radical $\{0\}$.

Definition 2.1. *A composition algebra A over a field F with multiplication $x \cdot y$ is an algebra with a nonsingular quadratic form q on A satisfying:*

$$q(x \cdot y) = q(x)q(y).$$

Furthermore, if A has an identity element e , we call A a unital composition algebra.

Every element of a unital composition algebras satisfies a quadratic polynomial. This is the minimal polynomial if the element is not a scalar multiple of the identity. More precisely, we have:

Proposition 2.2. *Every element x of a unital composition algebra A satisfies*

$$x \cdot x - \langle x, e \rangle x + q(x)e = 0.$$

For all $x, y \in A$. By linearizing the above equation, we get:

$$x \cdot y + y \cdot x - \langle x, e \rangle y - \langle y, e \rangle x + \langle x, y \rangle e = 0.$$

Proof. See [19, Proposition 1.2.3]. □

We now introduce conjugation in A . Consider the mapping $\bar{\cdot} : A \rightarrow A$ given by

$$\bar{x} = \langle x, e \rangle e - x,$$

for all $x \in A$. We call \bar{x} the conjugate of x . It is easy to see that $\bar{\cdot}$ is a linear map with $\overline{\bar{x}} = x$ for all $x \in A$. So $\bar{\cdot}$ is an involution on A .

The following lemmas hold in every unital composition algebras and can be directly deduced by linearizing the quadratic form q and using composition condition (see [19, §1.2, 1.3] for the proof). We will use them later:

Lemma 2.3. *We have*

- (1) $\langle x \cdot z, y \cdot z \rangle = \langle x, y \rangle q(z), \quad \langle z \cdot x, z \cdot y \rangle = q(z) \langle x, y \rangle,$
- (2) $\langle x \cdot z, y \cdot w \rangle + \langle x \cdot w, y \cdot z \rangle = \langle x, y \rangle \langle z, w \rangle,$

for all $x, y, z \in A$.

Lemma 2.4. *We have*

- (1) $x \cdot (\bar{x} \cdot y) = q(x)y, \quad (x \cdot \bar{y}) \cdot y = q(y)x,$
- (2) $x \cdot (\bar{y} \cdot z) + y \cdot (\bar{x} \cdot z) = \langle x, y \rangle z,$
- (3) $(x \cdot \bar{y}) \cdot z + (x \cdot \bar{z}) \cdot y = \langle y, z \rangle x,$

for all $x, y, z \in A$.

Given a unital composition algebra A with dimension n , one can get a new composition algebra with dimension $2n$ by using Cayley-Dickson process (see [12, §33.C]). We now have the well-know classification of unital composition algebras:

Theorem 2.5. *Every unital composition algebra over F is obtained by the Cayley-Dickson process. The possible dimensions are 1, 2, 4 and 8. Composition algebras of dimension 1 or 2 are commutative and associative, those of dimension 4 are associative but not commutative, and those of dimension 8 are neither commutative nor associative.*

Proof. See [12, Theorem 33.17]. □

We call A a quadratic étale algebra if $\dim A = 2$, a quaternion algebra (resp. an octonion algebra) if $\dim A = 4$ (resp. $\dim A = 8$). The Cayley-Dickson process applied to an octonion algebra does not yield a composition algebra, so octonion algebras have highest dimension. We will use these algebras to construct the exceptional group G_2 .

Recall that an element $x \in A$ is called isotropic if $q(x) = 0$. The quadratic form q is called isotropic if there exist nonzero isotropic elements in A . From [12, Theorem 33.19], there is only one isomorphism class of octonion algebras with isotropic quadratic forms. We call C_s the split octonion algebra if they have isotropic quadratic form. The split octonion algebra C_s can be constructed from the split quaternion algebra, but we have a remarkable way to express its multiplication rule (see [20, §2] and [9, §5]):

Consider a free \mathbb{Z} -module W of rank 3, and let $W^\vee = \text{Hom}_{\mathbb{Z}}(W, \mathbb{Z})$ be the dual module of W . By fixing a basis in W , we set

$$W = \mathbb{Z}u_1 + \mathbb{Z}u_2 + \mathbb{Z}u_3, \quad W^\vee = \mathbb{Z}v_1 + \mathbb{Z}v_2 + \mathbb{Z}v_3,$$

where $v_i(u_j) = \delta_{ij}$ for $i, j \in \{1, 2, 3\}$. We also fix an isomorphism $\wedge^3 W$ with $\mathbb{Z} \simeq \mathbb{Z} \cdot (u_1 \wedge u_2 \wedge u_3)$. Then we get an isomorphism:

$$\wedge^2 W \simeq W^\vee \quad \text{given by } w_1 \wedge w_2 \mapsto (w_1 \wedge w_2, -),$$

where $(w_1 \wedge w_2, w) := w_1 \wedge w_2 \wedge w$ for any $w \in W$. Similarly, we also have an isomorphism $\wedge^2 W^\vee \simeq W$ by duality. For instance, $u_1 \wedge u_2$ is identified with v_3 , $v_1 \wedge v_3$ is identified with $-u_2$, etc.

Let Λ be the space of matrices of the form:

$$\begin{pmatrix} a & w \\ \varphi & b \end{pmatrix}, \quad \text{where } a, b \in \mathbb{Z}, w \in W, \text{ and } \varphi \in W^\vee.$$

This is a free \mathbb{Z} -module of rank 8. If we define addition by matrices addition, and multiplication by:

$$\begin{pmatrix} a_1 & w_1 \\ \varphi_1 & b_1 \end{pmatrix} \cdot \begin{pmatrix} a_2 & w_2 \\ \varphi_2 & b_2 \end{pmatrix} := \begin{pmatrix} a_1 a_2 - \varphi_2(w_1) & a_1 w_2 + b_2 w_1 + \varphi_1 \wedge \varphi_2 \\ a_2 \varphi_1 + b_1 \varphi_2 + w_1 \wedge w_2 & b_1 b_2 - \varphi_1(w_2) \end{pmatrix}.$$

Then Λ is an algebra over \mathbb{Z} . Moreover, consider the quadratic form q on Λ given by:

$$q : \begin{pmatrix} a & w \\ \varphi & b \end{pmatrix} \mapsto ab + \varphi(w).$$

It is easy to see that $q(x \cdot y) = q(x)q(y)$, so that Λ is a unital composition algebra over \mathbb{Z} . Here Λ has a standard basis $\{e_1, e_2, u_i, v_i\}_{i=1,2,3}$, where

$$e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad u_i = \begin{pmatrix} 0 & u_i \\ 0 & 0 \end{pmatrix}, \quad v_i = \begin{pmatrix} 0 & 0 \\ v_i & 0 \end{pmatrix}.$$

The identity element in Λ is the identity matrix $e = e_1 + e_2$. Therefore, the split octonion algebra C_s can be viewed as $C_s = \Lambda \otimes_{\mathbb{Z}} F$.

Although unital composition algebras have lots of good properties, most composition algebras do not admit an identity. In the rest of this section, we will discuss a special class of composition algebras, called symmetric composition algebras.

Definition 2.6. *A symmetric composition algebra (S, \star) is a composition algebra with multiplication $x \star y$, satisfying*

$$\langle x \star y, z \rangle = \langle y \star z, x \rangle,$$

for all $x, y, z \in S$.

Similar to Lemma 2.3, 2.4, the following results hold in every symmetric composition algebra (S, \star) (see [12, Lemma 34.1]):

Lemma 2.7. *We have*

- (1) $\langle x \star z, y \star z \rangle = \langle x, y \rangle q(z)$,
- (2) $\langle z \star x, z \star y \rangle = q(z) \langle x, y \rangle$,
- (3) $\langle x \star z, y \star w \rangle + \langle x \star w, y \star z \rangle = \langle x, y \rangle \langle z, w \rangle$,

for all $x, y, z \in S$.

Lemma 2.8. *We have*

- (1) $(x \star y) \star z + (z \star y) \star x = \langle x, z \rangle y$,
- (2) $x \star (y \star z) + z \star (y \star x) = \langle x, z \rangle y$,

for all $x, y, z \in S$.

In particular, we have $(x \star y) \star x = x \star (y \star x) = q(x)y$.

Note that we have $(x \star y) \star x = x \star (y \star x) = q(x)y$ for all $x, y \in S$ by Lemma 2.8. Generally, we can show that an algebra with quadratic form q satisfying this equation is a symmetric composition algebra.

It turns out that unital composition algebras have a close relation to symmetric composition algebras. Starting from a unital composition algebra (A, \cdot) over F , we can always define a symmetric composition algebra (A, \star) given by:

$$x \star y = \bar{x} \cdot \bar{y},$$

for all $x, y \in A$. Symmetric composition algebras do not have identity, but the identity element $e \in A$ plays a special role in the corresponding symmetric composition algebra (A, \star) : it is an idempotent ($e \star e = e$) and satisfies

$$e \star x = x \star e = -x,$$

for any $x \in A$, with $\langle x, e \rangle = 0$. We call an element which satisfies the above condition a para-unit. In particular, symmetric composition algebras corresponding to octonion algebras (resp. split octonion algebras) are called para-octonion algebras (resp. split para-octonion algebras), denoted by (C, \star) (resp. (C_s, \star)). Using the same symbols as above ($C_s \simeq \Lambda \otimes_{\mathbb{Z}} F$,

with basis $\{e_1, e_2, u_i, v_i\}_{i=1,2,3}$, we obtain the multiplication table of the split para-octonion algebra (C_s, \star) as Table 1 below (we write “.” instead of 0 for clarity).

		y						
\star	e_1	e_2	u_1	u_2	u_3	v_1	v_2	v_3
e_1	e_2	$-v_1$	$-v_2$	$-v_3$
e_2	.	e_1	$-u_1$	$-u_2$	$-u_3$.	.	.
u_1	$-u_1$.	.	v_3	$-v_2$	$-e_1$.	.
u_2	$-u_2$.	$-v_3$.	v_1	.	$-e_1$.
x	u_3	$-u_3$.	v_2	$-v_1$.	.	$-e_1$
v_1	.	$-v_1$	$-e_2$.	.	.	u_3	$-u_2$
v_2	.	$-v_2$.	$-e_2$.	$-u_3$.	u_1
v_3	.	$-v_3$.	.	$-e_2$	u_2	$-u_1$.

TABLE 1. The split para-octonion algebra multiplication $x \star y$

With respect to this basis, the corresponding bilinear form $\langle \cdot, \cdot \rangle$ is given by

$$\langle e_1, e_2 \rangle = 1, \quad \langle u_i, v_j \rangle = \delta_{ij},$$

where all others are equal to 0. The involution $\bar{\cdot}$ on (C_s, \star) is given by:

$$\bar{e}_1 = e_2, \quad \bar{e}_2 = e_1, \quad \bar{u}_i = -u_i, \quad \bar{v}_i = -v_i.$$

We prefer to use para-octonion algebras (C, \star) instead of octonion algebras (C, \star) in the following sections. One reason is that it is convenient to work with the spin group $\text{Spin}(C, \star)$. We refer Remark 3.2 for more details.

Remark 2.9. Recall that a subspace U in an algebra A with a quadratic form q is isotropic if there exists $x \in U$ such that $q(x) = 0$, and U is totally isotropic if $q(x) = 0$ for all $x \in U$. The maximal totally isotropic subspaces are totally isotropic subspaces with the highest dimension. All maximal totally isotropic subspaces have the same dimension, which is called the Witt index of q . The Witt index is at most equal to half dimension of A .

The maximal totally isotropic subspaces in the split octonion (para-octonion) algebra C_s over F is described by Blij-Springer in [21], and Matzri-Vishne translate the classification to arbitrary composition algebras in [14]. In C_s , Every totally isotropic subspace is of the form

$$x \star C_s, \text{ or } C_s \star x,$$

where x is an isotropic element. These spaces have dimension 4. Furthermore, $x \star C_s = y \star C_s$ if and only if $Fx = Fy$ (see [14, Theorem 3.1, Proposition 3.2]).

Let U be a totally isotropic subspace in C_s . Consider the intersection of maximal isotropic subspaces:

$$\mathcal{L}(U) := \bigcap_{x \in U} (C_s \star x), \quad \mathcal{R}(U) := \bigcap_{x \in U} (x \star C_s).$$

We have the following diagram:

$$\begin{array}{ccc} & Fx & \\ & \swarrow \mathcal{L} & \nwarrow \mathcal{L} \\ C_s \star x & \xrightarrow{\mathcal{R} \circ \mathcal{R}} & x \star C_s \end{array}$$

for any x isotropic element. We call this diagram the geometric triality graph (see [14, §6.2]).

3. THE BUILDING OF G_2

In this section, we will briefly recall the theory of Bruhat-Tits building, and we are particularly interested in the special case when the group is of type G_2 . The theory of buildings of reductive groups over local fields are given by Bruhat-Tits in their series of papers [4], [5], [6], [7], [8]. In [4], [5], the building of a classical group G is a polysimplicial complex equipped with an action of G , and can be given as a set of graded lattice chains satisfying certain conditions.

For a simply connected classical group G , let F be a discrete valuation field, and \mathcal{O}_F be the ring of integers with uniformizer $\pi \in \mathcal{O}_F$. The building $\mathcal{B}(G)$ can be realized geometrically as a set of graded lattice chains. To each point $x \in \mathcal{B}(G)$, the subgroup \mathcal{G}_x that stabilizes the graded lattice chain x is a smooth, connected, affine group scheme over $\text{Spec}(\mathcal{O}_F)$, and called the parahoric subgroup of $\mathcal{B}(G)$ associated to x . We will fix a parahoric subgroup of $\mathcal{B}(G)$ in the next section and use it to define our affine Grassmannians. We refer [6], [8] for the theory of buildings of classical groups, and [9] for the building of G_2 .

Let us first construct the exceptional group of type G_2 . Generally, let (V, q) be a finite dimensional vector space with a nonsingular quadratic form q over a field F , with $\text{char}(F)$ different from 2. Denote by $\langle \cdot, \cdot \rangle$ the bilinear form corresponding to q . The special orthogonal group $\text{SO}(V, q)$ is the subgroup of the special linear group $\text{SL}(V, q)$ that preserve the form q . The universal covering of $\text{SO}(V, q)$ is the spin group $\text{Spin}(V, q)$ defined by:

$$\text{Spin}(V, q) = \{x \in C_0(V, q)^* \mid xVx^{-1} = V, \tau(x)x = 1\}.$$

Here $C(V, q)$ is the Clifford algebra of (V, q) , which is the factor of the tensor algebra $T(V) = \bigoplus_{n \geq 0} V \otimes \cdots \otimes V$ by the ideal $I(q)$ generated by all elements of the form $x \otimes x - q(x) \cdot 1$ for $x \in V$. It is a $\mathbb{Z}/2\mathbb{Z}$ -graded algebra with $C(V, q) = C_0(V, q) \oplus C_1(V, q)$. We call $C_0(V, q)$ the even Clifford algebra, and $C_1(V, q)$ the odd Clifford algebra (see [12, Ch. IV]). There is a canonical involution of the Clifford algebra $\tau : C(V, q) \rightarrow C(V, q)$ given by $\tau(x_1 \cdots x_d) = x_d \cdots x_1$ for $x_1, \dots, x_d \in V$. Thus, the spin group $\text{Spin}(V, q)$ is a subgroup of $C_0(V, q)$. We have an exact sequence:

$$1 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow \text{Spin}(V, q) \rightarrow \text{SO}(V, q) \rightarrow 1.$$

In particular, when consider the split para-octonion algebra (C_s, \star) as an 8-dim vector space over F , we can get the corresponding special orthogonal group $\text{SO}(C_s, \star)$ (resp. the spin group $\text{Spin}(C_s, \star)$). The corresponding group scheme $\mathbf{SO}(C_s, \star)$ (resp. $\mathbf{Spin}(C_s, \star)$) over $\text{Spec}(F)$ are defined by:

$$\begin{aligned} \mathbf{SO}(C_s, \star)(R) &= \{g \in \text{SL}(C_{sR}, \star) \mid \langle g(x), g(y) \rangle = \langle x, y \rangle\}, \\ \mathbf{Spin}(C_s, \star)(R) &= \{x \in C_0(C_{sR}, \star)^* \mid xC_{sR}x^{-1} = C_{sR}, \tau(x)x = 1\}, \end{aligned}$$

for any $x, y \in C_{sR}$. Here R is an F -algebra, and $C_{sR} = C_s \otimes_F R$. The following description of $\mathbf{Spin}(C_s, \star)$ shows that the spin group $\mathbf{Spin}(C_s, \star)$ is a subgroup of the triple orthogonal group $\mathbf{SO}_8^{\times 3}$.

Theorem 3.1. *There is an isomorphism*

$$\mathbf{Spin}(C_s, \star) \simeq \mathbf{RT}(C_s, \star),$$

where $\mathbf{RT}(C_s, \star)$ is the related triple group scheme that represents the functor from F -algebras to groups that sends R to

$$\mathbf{RT}(C_s, \star)(R) := \{(g_1, g_2, g_3) \in \mathbf{SO}(C_s, \star)(R)^{\times 3} \mid g_i(x \star y) = g_{i+1}(x) \star g_{i+2}(y)\}.$$

for any $x, y \in C_s \otimes_F R, i = 1, 2, 3 \pmod 3$.

See the proof in [12, Proposition 35.8]. The triple $(g_1, g_2, g_3) \in \mathbf{SO}(C_s, \star)(R)^{\times 3}$ that satisfies $g_i(x \star y) = g_{i+1}(x) \star g_{i+2}(y)$ is called a related triple. For any given $g_1 \in \mathbf{SO}(C_s, \star)(R)$, there exists $g_2, g_3 \in \mathbf{SO}(C_s, \star)(R)$ such that (g_1, g_2, g_3) is a related triple (see [12, Proposition 35.4]).

Remark 3.2. We have a similar result for the spin group of the split octonion algebra (C_s, \cdot) : There is an isomorphism

$$\mathbf{Spin}(C_s, \cdot) \simeq \{(g_1, g_2, g_3) \in \mathbf{SO}(C_s, \cdot)(R)^{\times 3} \mid g_i(x \cdot y) = \overline{g_{i+1}(\bar{x}) \cdot g_{i+2}(\bar{y})}\}.$$

(See [1, Remark 3.1]). It is also easy to get the above isomorphism from Theorem 3.1 by regarding $x \star y = \bar{x} \cdot \bar{y}$. Since it is more clear to view the spin group as the subgroup of $\mathbf{SO}_8^{\times 3}$ preserving multiplication, we prefer to use para-octonion algebras in the following sections.

From Theorem 3.1, it is easy to see that there is an action ρ on $\mathbf{Spin}(C_s, \star)$ given by

$$\rho : (g_1, g_2, g_3) \mapsto (g_2, g_3, g_1).$$

Further, for any $g \in \mathbf{SO}(C_s, \star)$, let \hat{g} be the automorphism of (C_s, \star) given by $\hat{g}(x) = \overline{g(\bar{x})}$. Then \hat{g} is also an element in $\mathbf{SO}(C_s, \star)$, and there is an involution on $\mathbf{Spin}(C_s, \star)$ given by:

$$\theta : (g_1, g_2, g_3) \mapsto (\hat{g}_1, \hat{g}_3, \hat{g}_2).$$

The involution θ and the $\mathbb{Z}/3\mathbb{Z}$ -action ρ generate an action S_3 on $\mathbf{Spin}(C_s, \star)$, whose group of fixed points, is precisely the exceptional group G_2 :

Proposition 3.3. *For the given action of S_3 on $\mathbf{Spin}(C_s, \star)$, we have a simple, simply connected group scheme*

$$G = \mathbf{Spin}(C_s, \star)^{S_3} \simeq \mathbf{Spin}(C_s, \star)^{A_3} \simeq \mathbf{Aut}(C_s, \star),$$

where $\mathbf{Aut}(C_s, \star)$ is the automorphism group scheme

$$\mathbf{Aut}(C_s, \star)(R) := \{g \in \mathbf{SO}(C_s, \star)(R) \mid g(x \star y) = g(x) \star g(y)\},$$

for any F -algebra $R, x, y \in C_s \otimes_F R$. The group G is of type G_2 .

Proof. See [12, §35]. □

Since there is an action S_3 on $\mathbf{Spin}(C_s, \star)$, one can form the building $\mathcal{B}(\mathbf{Spin}(C_s, \star))$ equipped with an action S_3 . It is natural to ask whether the set of fixed points of $\mathcal{B}(\mathbf{Spin}(C_s, \star))$ under S_3 , is the building of $\mathcal{B}(G)$. When $p \neq 2, 3$, Prasad-Yu in [16] show that it is indeed true. In the general case, the result is given by Gan-Yu in [9]. So we have

$$\mathcal{B}(G) \simeq \mathcal{B}(\mathbf{Spin}(C_s, \star))^{S_3}.$$

Gan-Yu give an explicit description of the building $\mathcal{B}(G)$, and show that there is a bijection between the building $\mathcal{B}(G)$ and the set of graded lattice chains satisfying certain conditions.

Recall that a lattice L in (C_s, \star) is a finitely generated projective \mathcal{O}_F -submodule in C_s , such that $L \otimes_{\mathcal{O}_F} F = C_s$. A lattice chain $\{L_i\}_{i \in I}$ is a totally ordered set of lattices of C_s , such that

$$L_{i_0} \subsetneq L_{i_1} \cdots \subsetneq L_{i_n} \subsetneq \pi^{-1}L_{i_0},$$

for $I = \{i_0 \leq i_1 \leq \cdots \leq i_n\}$. A graded lattice chain is a pair $(\{L_i\}_{i \in I}, c)$ where $\{L_i\}_{i \in I}$ is a lattice chain, and c is a strictly decreasing map from $\{L_i\}_{i \in I}$ to \mathbb{R} such that

$$c(\lambda L) = \text{ord}(\lambda) + c(L),$$

for $\lambda \in F, L \in \{L_i\}_{i \in I}$. If for any lattice $L \in \{L_i\}_{i \in I}$, the dual space $L^\vee = \{x \in C_s \mid \langle x, L \rangle \subset \mathcal{O}_F\}$ is also in $\{L_i\}_{i \in I}$, then we call $\{L_i\}_{i \in I}$ a self-dual lattice chain.

An order in C_s is an \mathcal{O}_F -lattice which is closed under multiplication (i.e., $x \star y \in L$ for all $x, y \in L$), and contains the para-unit e . This order is said to be maximal if it is not contained in any larger one. Since we always assume $\text{char}(F) \neq 2$ in this paper, we have

Proposition 3.4. *Suppose that L is a lattice in C_s , which is closed under multiplication, then L is a maximal order if and only if L is self-dual with respect to $\langle \cdot, \cdot \rangle$.*

Proof. See [9, Proposition 5.1]. □

Remark 3.5. In [9], Gan-Yu define that an order L in the split octonion algebra (C_s, \cdot) is an \mathcal{O}_F -lattice which is a unital ring. We transfer this definition to the para-octonion algebra (C_s, \star) here. Notice that any order is closed under conjugation, so that the condition $x \cdot y \in L$ is equivalent to

$$x \star y = \bar{x} \cdot \bar{y} \in L$$

for all $x, y \in L$. Meanwhile, the unit e in (C_s, \cdot) becomes the para-unit e in (C_s, \star) . Hence, we define that an order in a para-octonion algebra is an \mathcal{O}_F -lattice which is closed under multiplication, and contains the para-unit e as above.

We are now ready to describe vertices in the building $\mathcal{B}(G)$, when $G = \mathbf{Aut}(C_s, \star)$. By fixing a maximal split torus $S \subset G$, we can identify the apartment $\mathcal{A}(S)$ with the real vector space $X_+(S) \otimes \mathbb{R}$. Here $X_+(S)$ is the cocharacter group. Having fixed the origin, Gan-Yu show that the closed chamber in $\mathcal{A}(S)$ is a triangle with three vertices. Each vertex is corresponding to a type of lattices (or lattice chains) in $\mathcal{B}(G)$.

Theorem 3.6. *There are three types of vertices in building $\mathcal{B}(G)$. The vertices of type i are in natural bijection with the orders satisfying certain conditions as follows:*

- *type 1: L is a maximal order in C_s , i.e., L is self-dual, and closed under multiplication. The corresponding graded lattice chain is a pair $(\{\pi^i L\}, c)$:*

$$\cdots \subsetneq \pi L \subsetneq L \subsetneq \pi^{-1} L \subsetneq \cdots, \text{ where } c(L) = 0.$$

- *type 2: L is an order satisfying $L \subsetneq L^\vee \subsetneq \pi^{-1} L$, $L^\vee \star L^\vee \subset \pi^{-1} L$. The corresponding graded lattice chain is the following:*

$$\cdots \subsetneq \pi L^\vee \subsetneq L \subsetneq L^\vee \subsetneq \pi^{-1} L \subsetneq \cdots, \text{ where } c(L) = 0, c(L^\vee) = -\frac{1}{2}.$$

- *type 3: L is an order satisfying $L \subsetneq L^\vee \subsetneq \pi^{-1} L$, with $M = \pi L^\vee \star L^\vee + L$ is self-dual. The corresponding graded lattice chain is the following:*

$$\cdots \subsetneq \pi L^\vee \subsetneq L \subsetneq M \subsetneq L^\vee \subsetneq \pi^{-1} L \subsetneq \cdots, \text{ where } c(L) = 0, c(M) = -\frac{1}{3}, c(L^\vee) = -\frac{2}{3}.$$

Proof. See [9, §9]. □

Example 3.7. Let $(C_s, \star) = \Lambda \otimes_{\mathbb{Z}} F$ be the split para-octonion algebra, with the basis $e_1, e_2, \{u_i, v_i\}_{i=1,2,3}$. The multiplication table with respect to this basis is in Table 1.

(1). The trivial example is the lattice

$$\mathbb{L}_1 = \mathcal{O}_F \langle e_1, e_2, u_1, u_2, u_3, v_1, v_2, v_3 \rangle.$$

It is easy to see that \mathbb{L}_1 is self-dual, and closed under multiplication table, so \mathbb{L}_1 is the vertex in $\mathcal{B}(G)$ of type 1. We call \mathbb{L}_1 the standard lattice of type 1.

(2). Let

$$\mathbb{L}_2 = \mathcal{O}_F \langle e_1, e_2, \pi u_1, u_2, u_3, v_1, \pi v_2, v_3 \rangle.$$

The dual lattice

$$\mathbb{L}_2^\vee = \mathcal{O}_F \langle e_1, e_2, u_1, \pi^{-1} u_2, u_3, \pi^{-1} v_1, v_2, v_3 \rangle.$$

It is easy to see that $\mathbb{L}_2 \subsetneq \mathbb{L}_2^\vee \subsetneq \pi^{-1} \mathbb{L}_2$, and

$$\mathbb{L}_2^\vee \star \mathbb{L}_2^\vee = \mathcal{O}_F \langle \pi^{-1} e_1, \pi^{-1} e_2, u_1, \pi^{-1} u_2, \pi^{-1} u_3, \pi^{-1} v_1, v_2, \pi^{-1} v_3 \rangle,$$

so that $\mathbb{L}_2^\vee \star \mathbb{L}_2^\vee = \pi^{-1} \mathbb{L}_2$. Hence the corresponding lattice chain of \mathbb{L}_2 is a vertex in $\mathcal{B}(G)$. We call \mathbb{L}_2 the standard lattice of type 2.

(3). Let

$$\mathbb{L}_3 = \mathcal{O}_F \langle e_1, e_2, \pi u_1, \pi u_2, u_3, v_1, v_2, \pi v_3 \rangle.$$

We leave readers to check

$$M = \pi \mathbb{L}_3^\vee \star \mathbb{L}_3^\vee + \mathbb{L}_3 = \mathcal{O}_F \langle e_1, e_2, u_1, u_2, \pi^{-1} u_3, v_1, v_2, \pi v_3 \rangle$$

is self-dual. We call \mathbb{L}_3 the standard lattice of type 3.

To each vertex x of type i in $\mathcal{B}(G)$, the stabilizer of the corresponding lattice (lattice chains) L_x in $G(F)$ is a maximal parahoric subgroup \mathcal{G}_x . From [9, Theorem 7.2], the hyperspecial points in $\mathcal{B}(G)$ corresponding to vertices of type 1. Now we can define the parahoric subgroup \mathcal{G} corresponding to the hyperspecial points: Recall that Λ is the free \mathbb{Z} -module of rank 8, with $C_s = \Lambda \otimes_{\mathbb{Z}} F$. Consider $\mathbb{L}_1 = \Lambda \otimes_{\mathbb{Z}} \mathcal{O}_F$ (the standard lattice) as a submodule in C_s . We denote by $\mathbf{SO}(\Lambda)$ the subscheme of points in the isomorphism group scheme $\text{Isom}(\Lambda)$ that preserve the form and whose determinant is equal to 1.

Definition 3.8. *The parahoric subgroup corresponding to hyperspecial points \mathcal{G} is an affine group scheme over $\text{Spec}(\mathcal{O}_F)$ that represents the functor from \mathcal{O}_F -algebras to groups that sends R to*

$$\mathcal{G}(R) = \{g \in \mathbf{SO}(\Lambda)(R) \mid g(x \star y) = g(x) \star g(y)\},$$

for all $x, y \in \mathbb{L}_1 \otimes_{\mathcal{O}_F} R$.

Similarly, we can use \mathbb{L}_2 (resp. \mathbb{L}_3) in Example 3.7 to define the parahoric subgroup \mathcal{G}_2 (resp. \mathcal{G}_3) (see Definition 6.3).

4. AFFINE GRASSMANNIANS FOR \mathcal{G}

In this section, we will define affine Grassmannians by loop groups. The affine Grassmannians is represented by an ind-scheme. We refer [2] [3] for important results on the structure of loop groups and associated affine Grassmannians.

Let k be a field, and let G_0 be an algebraic group over $\text{Spec}(k)$. The functor LG_0 is from the category of k -algebras to sets that sends R to

$$LG_0(R) = G_0(R((t))).$$

This functor is represented by an ind-scheme, called the loop group associated to G_0 . We also consider the positive loop group L^+G_0 , which is the functor on the category of k -algebras, given by

$$L^+G_0(R) = G_0(R[[t]]).$$

Then $L^+G_0 \subset LG_0$ is a subgroup functor, and the fpqc-quotient $\mathrm{Gr}_{G_0} = LG_0/L^+G_0$ is by definition the affine Grassmannian associated to G_0 . The fpqc-sheaf Gr_{G_0} is also represented by an ind-scheme.

In [15], Pappas-Rapoport develop this definition for “twisted” loop groups as follows: Consider G an algebraic group over $\mathrm{Spec}(k((t)))$. The algebraic loop group LG is the functor from the category of k -algebras to the sets given by

$$LG(R) = G(R((t))).$$

Since $R((t))$ is a $k((t))$ -algebra, this definition makes sense. Note that $R[[t]]$ is not a $k((t))$ -algebra, so we cannot use G to define positive loop groups now. By choosing a parahoric subgroup \mathcal{G} of G , we denote by $L^+\mathcal{G}$ the positive loop group that represents the functor from k -algebras to sets that sends R to

$$L^+\mathcal{G}(R) = \mathcal{G}(R[[t]]).$$

Notice that when $G = G_0 \otimes_k k((t))$, we recover the previous definition in the untwisted case. Since the generic fiber \mathcal{G}_η of \mathcal{G} is always equal to G , our affine Grassmannian $\mathrm{Gr}_\mathcal{G}$ associated to \mathcal{G} is the fpqc-sheaf $GL\mathcal{G}_\eta/L^+\mathcal{G} = LG/L^+\mathcal{G}$. We refer to [15] for more details.

A classical prototype is the affine Grassmannian for the general linear group GL_n . This $\mathrm{Gr}_{\mathrm{GL}_n}$ can be viewed as a functor from k -algebras to sets that sends R to

$$\{L \mid L \text{ is a finitely generated projective } R[[t]]\text{-modules in } R((t))^n, \text{ such that } L[t^{-1}] \simeq R((t))^n\}.$$

In the rest of this section, let k be a field with $\mathrm{char}(k) \neq 2$. The field $F = k((t))$ is a discrete valuation field with the ring of integers $\mathcal{O}_F = k[[t]]$, and uniformizer $t \in k[[t]]$. Let $(C_s, \star) = \Lambda \otimes_{\mathbb{Z}} F$ be the split para-octonion algebra over F , and $G = \mathbf{Aut}(C_s, \star)$ be the automorphism group scheme of split para-octonion algebra, which is a group of type G_2 . Consider the parahoric subgroup \mathcal{G} corresponding to hyperspecial points in $\mathcal{B}(G)$ as defined in Definition 3.8. Then the quotient fpqc-sheaf

$$\mathrm{Gr}_\mathcal{G} = L\mathcal{G}_\eta/L^+\mathcal{G}$$

is by definition the affine Grassmannian for G_2 .

Our goal in this section is to give an explicit description of the affine Grassmannian $\mathrm{Gr}_\mathcal{G}$. Inspired by the results from [9], we already know the k -points in $\mathrm{Gr}_\mathcal{G}$: since $L^+\mathcal{G}(k) = \mathcal{G}(k[[t]])$ is the stabilizer of the standard lattice \mathbb{L} , and $L\mathcal{G}_\eta(k) = G(k((t)))$ acts transitively on vertices of type 1, then there is a bijection between k -points of $\mathrm{Gr}_\mathcal{G}$ and the set of maximal orders in C_s . Our first main theorem is the following:

Theorem 4.1. *There is an $L\mathcal{G}_\eta$ -equivariant isomorphism*

$$\mathrm{Gr}_\mathcal{G} \simeq \mathcal{F}$$

where the functor \mathcal{F} sends a k -algebra R to the set of $R[[t]]$ -lattices L of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is self dual under the bilinear form $\langle \cdot, \cdot \rangle$, i.e., $L \simeq \mathrm{Hom}_{R[[t]]}(L, R[[t]])$.
- (2) L is an order, i.e., $e \in L$, $L \star L \subset L$.

This theorem is proven in the next section. It gives a bijection between R -points in the affine Grassmannian for \mathcal{G} and a certain set of $R[[t]]$ -lattices in $C_s \otimes_{k((t))} R((t))$ that are self-dual and closed under multiplication.

5. PROOF OF THE FIRST MAIN THEOREM

In [15], Pappas-Rapoport show that there is a bijection between the affine Grassmannians for unitary groups and the space of self-dual lattice chains for a hermitian form. Recall that $\mathbb{L}_1 = k[[t]]\langle e_1, e_2, u_i, v_i \rangle_{i=1,2,3}$ is the standard lattice of type 1 in the split para-octonion algebra $C_s = \Lambda \otimes_{\mathbb{Z}} k((t))$. Similar to the proof of [15, Theorem 4.1] in the unitary case, it suffices to check the following two statements:

- (i) For any R , the subgroup of $L\mathcal{G}_\eta(R)$ consisting of elements that stabilize the standard lattice $\mathbb{L}_1 \otimes_{k[[t]]} R[[t]]$ agrees with $L^+\mathcal{G}(R)$.
- (ii) For any $L \in \mathcal{F}(R)$ with (R, \mathcal{M}) a local Henselian ring with maximal ideal \mathcal{M} , there exists $g \in L\mathcal{G}_\eta(R)$ such that $L = g(\mathbb{L}_1 \otimes_{k[[t]]} R[[t]])$.

It is easy to see part (i). By definition, the $R[[t]]$ -points of \mathcal{G} is the stabilizer of the base change of the standard lattice $\mathbb{L}_1 \otimes_{k[[t]]} R[[t]]$ (we also call $\mathbb{L}_1 \otimes_{k[[t]]} R[[t]]$ the standard lattice in this section if there is no confusion). We will focus to prove part (ii). The key is to find a basis in $L \in \mathcal{F}(R)$, such that its multiplication table is the same as Table 1 (the multiplication table of the standard lattice $\mathbb{L}_1 \otimes_{k[[t]]} R[[t]]$). If this basis exists, then there is an element $g \in \mathrm{SL}_8$ such that $g(\mathbb{L}_1 \otimes_{k[[t]]} R[[t]]) = L$, with $g(x \star y) = g(x) \star g(y)$. By Lemma 2.8, the quadratic form q is determined by its multiplication, so that g also preserves the form q . Thus, $g \in \mathbf{SO}(C_s, \star)(R((t)))$, and so that $g \in G(R((t))) = L\mathcal{G}_\eta(R)$.

For any $L \in \mathcal{F}(R)$, by Proposition 3.4, L is a maximal order in $C_s \otimes_{k((t))} R((t))$, which contains a para-unit element $e \in L$. Since L is a self-dual lattice, we have $L \simeq \mathrm{Hom}_{R[[t]]}(L, R[[t]])$. So there exists a basis $\{x_i\}_{i=1}^8$ of L such that $\langle x_i, x_{9-j} \rangle = \delta_{ij}$. Consider $e = \sum a_i x_i$ for all $a_i \in R[[t]]$. We set:

$$e_1 = \sum_{i=1}^4 a_i x_i, \quad e_2 = \sum_{i=5}^8 a_i x_i.$$

Then $q(e_1) = q(e_2) = 0$, and $\langle e_1, e_2 \rangle = 1$ by $q(e) = 1$. Moreover, consider the minimal polynomial for e_1 in the split octonion algebra $C_s \otimes_{k((t))} R((t))$:

$$e_1 \cdot e_1 - \langle e_1, e \rangle e_1 + q(e_1)e = 0.$$

We get $e_1 \cdot e_1 = e_1$ by $\langle e_1, e \rangle = 1$ and $q(e_1) = 0$. Thus in the split para-octonion algebra $(C_s \otimes_{k((t))} R((t)), \star)$, we have

$$e_2 \star e_2 = \bar{e}_2 \cdot \bar{e}_2 = e_1,$$

by $\bar{e}_2 = \langle e_2, e \rangle e - e_2 = e_1$. Similarly, we have $e_1 \star e_1 = e_2$. For $e_1 \star e_2$, it is easy to see that

$$\begin{aligned} e_1 \star e_2 &= \bar{e}_1 \cdot \bar{e}_2 = e_2 \cdot e_1 \\ &= (e - e_1) \cdot e_1 \\ &= 0. \end{aligned}$$

Also $e_2 \star e_1 = 0$ by similar calculation. Therefore, the sublattice $R[[t]]e_1 \oplus R[[t]]e_2$ is a hyperbolic space satisfying:

$$\begin{aligned} e_1 \star e_1 &= e_2, & e_2 \star e_2 &= e_1, \\ e_1 \star e_2 &= e_2 \star e_1 = 0, \\ q(e_1) &= q(e_2) = 0, & \langle e_1, e_2 \rangle &= 1. \end{aligned}$$

Readers may notice that here e_1, e_2 play the same roles as for e_1, e_2 in the Table 1 (that is the reason we keep the same symbol). We claim that e_1, e_2 are primitive elements in L (elements

that extends to a basis of L): Since (R, \mathcal{M}) is a local ring, the canonical quotient map:

$$R[[t]] \rightarrow R \rightarrow \kappa = R/\mathcal{M}$$

gives a base change of $L \rightarrow L \otimes_{R[[t]]} \kappa$. The image of e_1, e_2 in $L \otimes_{R[[t]]} \kappa$ are non-zero elements by $\langle e_1, e_2 \rangle = 1$. By Nakayama's lemma, we can extend $\{e_1, e_2\}$ as a basis of L . So they are primitive elements. Choose $y_3, \dots, y_8 \in L$ such that $\{e_1, e_2, y_3, \dots, y_8\}$ is a basis of L . Consider the sublattice $L_0 \subset L$:

$$L_0 := \{x \in L \mid \langle x, e_1 \rangle = 0, \langle x, e_2 \rangle = 0\}.$$

For any $x \in L$, we write x as

$$x = b_1 e_1 + b_2 e_2 + \sum_{i=3}^8 b_i y_i,$$

for $b_i \in R[[t]]$. By replacing y_i by

$$y'_i = y_i - \langle y_i, e_2 \rangle e_1 - \langle y_i, e_1 \rangle e_2,$$

we get $\langle y'_i, e_1 \rangle = \langle y'_i, e_2 \rangle = 0$. So we can always assume $y_i \in L_0$. Hence $L = R[[t]]e_1 \oplus R[[t]]e_2 \oplus L_0$, and L_0 is a sublattice of rank 6. Notice that for any $x \in L_0$, we have $\langle x, e \rangle = \langle x, e_1 + e_2 \rangle = 0$. Thus $x \star e = e \star x = -x$ since e is the para-unit element. Set $L_i = L_0 \star e_i$ for $i = 1, 2$. By Remark 2.9, L_i is a totally isotropic subspace with rank ≤ 3 .

Lemma 5.1. *We have*

- (1) $L_i \subset L_0$ for $i = 1, 2$.
- (2) $e_i \star L_i = 0$, $L_i \star e_{i+1} = 0$ for $i = 1, 2 \pmod{2}$.

Proof. (1). For any $x \in L_1$, we can write x as $x_1 \star e_1$ for $x_1 \in L_0$. By the definition of symmetric composition algebra,

$$\begin{aligned} \langle x_1 \star e_1, e_1 \rangle &= \langle e_1 \star e_1, x_1 \rangle = \langle e_2, x_1 \rangle = 0, \\ \langle x_1 \star e_1, e_2 \rangle &= \langle e_1 \star e_2, x_1 \rangle = 0. \end{aligned}$$

So that $x \in L_0$, and $L_1 \subset L_0$. We have a similar result for L_2 .

(2). Again, we only consider L_1 case. For any $x = x_1 \star e_1 \in L_1$, we have $e_1 \star (x_1 \star e_1) = q(e_1)x_1 = 0$ by Lemma 2.8. Hence $e_i \star L_i = 0$. For $(x_1 \star e_1) \star e_2$, consider

$$(x_1 \star e_1) \star e_2 + (e_2 \star e_1) \star x_1 = \langle x_1, e_2 \rangle e_1.$$

Since $e_2 \star e_1 = 0$, and $\langle x_1, e_2 \rangle = 0$, we get $x \star e_2 = (x_1 \star e_1) \star e_2 = 0$ for all $x \in L_1$. \square

For any $x \in L_0$, we obtain

$$\begin{aligned} x &= \langle e_1, e_2 \rangle x \\ &= (e_1 \star x) \star e_2 + (e_2 \star x) \star e_1. \end{aligned}$$

by Lemma 2.8. Since $e_1 \star x, e_2 \star x \in L_0$, we get $L_0 = L_1 + L_2$. Each L_i is totally isotropic with rank ≤ 3 , so that $L_0 = L_1 \oplus L_2$. Since L_1, L_2 are totally isotropic, and $\langle \cdot, \cdot \rangle$ restricted to L_0 is nondegenerate, we see that L_1, L_2 are in duality: Consider the isomorphism $L_1 \rightarrow L_2^\vee$ given by $x \mapsto \langle x, - \rangle$, we have $L_1 \simeq \text{Hom}_{R[[t]]}(L_2, R[[t]])$. Set the basis for L_i :

$$L_1 = R[[t]]\langle u_1, u_2, u_3 \rangle, \quad L_2 = R[[t]]\langle v_1, v_2, v_3 \rangle.$$

with $\langle u_i, v_j \rangle = \delta_{ij}$.

Lemma 5.2. *We have*

$$\begin{aligned} u_i \star e_1 &= e_2 \star u_i = -u_i, \\ v_i \star e_2 &= e_1 \star v_i = -v_i, \end{aligned}$$

for $i = 1, 2, 3$.

Proof. We can directly get these equations from Lemma 5.1 (2): Since for any $x \in L_0$, we have $e \star x = x \star e = -x$. Thus $u_i \star e = u_i \star (e_1 + e_2) = -u_i$. By $u_i \star e_2 = 0$, we obtain $u_i \star e_1 = -u_i$. Similarly, we get the rest equations. \square

So far, we find the basis $e_1, e_2, \{u_i, v_i\}_{i=1,2,3}$ of L . To show that the multiplication table of this basis is the same as Table 1. It remains to check $u_i \star v_j$ (resp. $v_i \star u_j$), and $u_i \star u_j$ (resp. $v_i \star v_j$) for $i = 1, 2, 3$. Let us check $u_i \star v_j$ first. Consider

$$\begin{aligned} u_i \star v_j &= -(u_i \star e_1) \star v_j \\ &= (v_j \star e_1) \star u_i - \langle u_i, v_j \rangle e_1 \\ &= -\delta_{ij} e_1. \end{aligned}$$

The first row in the above equation comes from Lemma 5.2, and the second row comes from Lemma 2.8. Similarly, we also have

$$v_i \star u_j = -\delta_{ij} e_2.$$

for $i = 1, 2, 3$. Finally, for $u_i \star u_j$ (resp. $v_i \star v_j$), we claim that $u_i \star u_j \in L_0$. It is easy to see that $\langle u_i \star u_j, e_1 \rangle = \langle u_j \star e_1, u_i \rangle = -\langle u_j, u_i \rangle$, where $\langle u_j, u_i \rangle = 0$ since L_i is totally isotropic. Also, $\langle u_i \star u_j, e_2 \rangle = \langle u_j \star e_2, u_i \rangle = 0$ by $u_j \star e_2 = 0$. Thus, $u_i \star u_j \in L_0$. We can also get $v_i \star v_j \in L_0$ by the same way. Moreover, consider

$$(u_j \star u_i) \star e_2 + (e_2 \star u_i) \star u_j = \langle u_j, e_2 \rangle u_i = 0$$

by Lemma 2.8. We have

$$u_i \star u_j = -(e_2 \star u_i) \star u_j = (u_j \star u_i) \star e_2,$$

so that $u_i \star u_j \in L_0 \star e_2 = L_2$. Now consider $u_1 \star u_2$ for instance. Since

$$\begin{aligned} \langle u_1 \star u_2, u_1 \rangle &= -\langle u_1 \star u_2, u_1 \star e_1 \rangle = -q(u_1) \langle u_2, e_1 \rangle, \\ \langle u_1 \star u_2, u_2 \rangle &= -\langle u_1 \star u_2, e_2 \star u_2 \rangle = -q(u_2) \langle u_1, e_2 \rangle, \end{aligned}$$

by Lemma 2.7. We get $\langle u_1 \star u_2, u_1 \rangle = \langle u_1 \star u_2, u_2 \rangle = 0$ by L_1 totally isotropic, which implies

$$u_1 \star u_2 = \lambda v_3$$

for some $\lambda \in R[[t]]$. Multiplying v_1 on the right side, we get $(u_1 \star u_2) \star v_1 = \lambda(v_3 \star v_1)$. Since $(u_1 \star u_2) \star v_1 + (v_1 \star u_2) \star u_1 = \langle u_1, v_1 \rangle u_2$, and $v_1 \star u_2 = 0$, $\langle u_1, v_1 \rangle = 1$, we have $(u_1 \star u_2) \star v_1 = u_2$. Hence $v_3 \star v_1 = \lambda^{-1} u_2$. Therefore, $\lambda, \lambda^{-1} \in R[[t]]$ by $L \star L \subset L$, which implies λ is a unit in $R[[t]]$. We can assume $\lambda = 1$, so that

$$u_1 \star u_2 = v_3.$$

We perform similar calculations for the other $u_i \star u_j$ and $v_i \star v_j$. The multiplication tables are the following:

TABLE 2. $u_i * u_j$

*	u_1	u_2	u_3
u_1	0	v_3	$-v_2$
u_2	$-v_3$	0	v_1
u_3	v_2	$-v_1$	0

TABLE 3. $v_i * v_j$

*	v_1	v_2	v_3
v_1	0	u_3	$-u_2$
v_2	$-u_3$	0	u_1
v_3	u_2	$-u_1$	0

Therefore, we complete the multiplication table of L , which it is the same as Table 1. So there exist $g \in L\mathcal{G}_\eta(R)$, such that $L = g(\mathbb{L}_1 \otimes_{k[[t]]} R[[t]])$. That proves Theorem 4.1.

6. AFFINE GRASSMANNIANS FOR OTHER PARAHORIC SUBGROUPS

So far, we described the affine Grassmannians for \mathcal{G} , where \mathcal{G} is the maximal parahoric subgroups corresponding to hyperspecial points in the building $\mathcal{B}(G)$. We can view \mathcal{G} as the stabilizer subgroups of the standard lattice \mathbb{L}_1 (type 1). According to Theorem 3.6, there are two other maximal parahoric subgroups up to conjugation, corresponding to vertices of type 2 (resp. type 3) in the building $\mathcal{B}(G)$. It is natural to ask if we can describe the affine Grassmannians for these parahoric subgroups.

To answer this question, we first need to fix the parahoric subgroup that we want. The notation here is the same as that in §5. That is: $F = k((t))$ is a discrete valuation field with the ring of integers $\mathcal{O}_F = k[[t]]$. Let $(C_s, \star) = \Lambda \otimes_{\mathbb{Z}} F$ be the split para-octonion algebra over F . Consider the lattice $\mathbb{L}_2 \subset C_s$ in Example 3.7:

$$\mathbb{L}_2 = k[[t]]\langle e_1, e_2, \pi u_1, u_2, u_3, v_1, \pi v_2, v_3 \rangle,$$

and $\mathbb{L}_3 \subset C_s$:

$$\mathbb{L}_3 = k[[t]]\langle e_1, e_2, \pi u_1, \pi u_2, u_3, v_1, v_2, \pi v_3 \rangle.$$

It is easy to see that \mathbb{L}_2 satisfies $e \in \mathbb{L}_2$, $\mathbb{L}_2 \star \mathbb{L}_2 \subset \mathbb{L}_2$, $\mathbb{L}_2 \subsetneq \mathbb{L}_2^\vee \subsetneq t^{-1}\mathbb{L}_2$, and $\mathbb{L}_2^\vee \star \mathbb{L}_2^\vee \subset t^{-1}\mathbb{L}_2$. The corresponding graded lattice chain

$$(6.1) \quad \cdots \subsetneq t\mathbb{L}_2^\vee \subsetneq \mathbb{L}_2 \subsetneq \mathbb{L}_2^\vee \subsetneq t^{-1}\mathbb{L}_2 \subsetneq \cdots$$

is the vertex of type 2 in $\mathcal{B}(G)$. Similarly, the lattice \mathbb{L}_3 satisfies $e \in \mathbb{L}_3$, $\mathbb{L}_3 \star \mathbb{L}_3 \subset \mathbb{L}_3$, $\mathbb{L}_3 \subsetneq \mathbb{L}_3^\vee \subsetneq t^{-1}\mathbb{L}_3$, and $M = t\mathbb{L}_3^\vee \star \mathbb{L}_3^\vee + \mathbb{L}_3$ is self-dual. The corresponding graded lattice chain

$$(6.2) \quad \cdots \subsetneq t\mathbb{L}_3^\vee \subsetneq \mathbb{L}_3 \subsetneq M \subsetneq \mathbb{L}_3^\vee \subsetneq t^{-1}\mathbb{L}_3 \subsetneq \cdots$$

is the vertex of type 3 in $\mathcal{B}(G)$. We denote by $\mathbf{SO}(\mathbb{L}_2)$ (resp. $\mathbf{SO}(\mathbb{L}_3)$) the subscheme of points in the isomorphism group scheme $\text{Isom}(\mathbb{L}_2)$ (resp. $\text{Isom}(\mathbb{L}_3)$) that preserve the form and whose determinant is equal to 1.

Definition 6.3. *The parahoric subgroup \mathcal{G}_2 corresponding to vertices of type 2, is the stabilizer subgroup scheme of \mathbb{L}_2 in $G(k((t)))$. More precisely, \mathcal{G}_2 is an affine group scheme over $\text{Spec}(k[[t]])$ that represents the functor from $k[[t]]$ -algebras to groups that sends R to*

$$\mathcal{G}_2(R) = \{g \in \mathbf{SO}(\mathbb{L}_2)(R) \mid g(x \star y) = g(x) \star g(y)\},$$

for all $x, y \in \mathbb{L}_2 \otimes_{k[[t]]} R$. Similarly, the parahoric subgroup \mathcal{G}_3 corresponding to vertices of type 3 represents the functor that sends $k[[t]]$ -algebras R to

$$\mathcal{G}_3(R) = \{g \in \mathbf{SO}(\mathbb{L}_3)(R) \mid g(x \star y) = g(x) \star g(y)\},$$

for all $x, y \in \mathbb{L}_3 \otimes_{k[[t]]} R$.

Remark 6.4. For any $g \in \mathcal{G}_2(k[[t]])$, we have $g(\mathbb{L}_2) = \mathbb{L}_2$. It is easy to see that $g(\mathbb{L}_2^\vee) = \mathbb{L}_2^\vee$ since $\langle g(x), \mathbb{L}_2 \rangle = \langle x, g^{-1}(\mathbb{L}_2) \rangle = \langle x, \mathbb{L}_2 \rangle \in k[[t]]$. So that g preserves the graded lattice chain (6.1).

For $g \in \mathcal{G}_3(k[[t]])$, we get $g(\mathbb{L}_3^\vee) = \mathbb{L}_3^\vee$ similarly as above. and $g(M) = tg(\mathbb{L}_3^\vee \star \mathbb{L}_3^\vee) + g(\mathbb{L}_3)$. Since g is a change of basis of \mathbb{L}_3^\vee , we have $g(\mathbb{L}_3^\vee \star \mathbb{L}_3^\vee) = \mathbb{L}_3^\vee \star \mathbb{L}_3^\vee$. Thus $g(M) = M$, and g preserves the graded lattice chain (6.2).

The quotient fpqc-sheaves:

$$\mathrm{Gr}_{\mathcal{G}_2} = LG/L^+\mathcal{G}_2, \quad (\text{resp. } \mathrm{Gr}_{\mathcal{G}_3} = LG/L^+\mathcal{G}_3)$$

are by definition the affine Grassmannians for \mathcal{G}_2 (resp. \mathcal{G}_3). We want to give explicit descriptions of affine Grassmannians $\mathrm{Gr}_{\mathcal{G}_2}$ (resp. $\mathrm{Gr}_{\mathcal{G}_3}$) in this section.

Theorem 6.5. (a). *There is an LG-equivariant isomorphism*

$$\mathrm{Gr}_{\mathcal{G}_2} \simeq \mathcal{F}_2$$

where the functor \mathcal{F}_2 sends a k -algebra R to the set of $R[[t]]$ -lattices L of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is an order in $C_s \otimes_{k((t))} R((t))$, i.e., $e \in L$, $L \star L \subset L$.
- (2) $L \subsetneq L^\vee \subsetneq t^{-1}L$.
- (3) $L^\vee \star L^\vee \subset t^{-1}L$.

(b). *There is an LG-equivariant isomorphism*

$$\mathrm{Gr}_{\mathcal{G}_3} \simeq \mathcal{F}_3$$

where the functor \mathcal{F}_3 sends a k -algebra R to the set of $R[[t]]$ -lattices L of $C_s \otimes_{k((t))} R((t))$ such that

- (1) L is an order in $C_s \otimes_{k((t))} R((t))$.
- (2) $L \subsetneq L^\vee \subsetneq t^{-1}L$.
- (3) $M = tL^\vee \star L^\vee + L$ is self-dual.

Notice that for any k -algebra R , $\mathcal{F}_2(R) \cap \mathcal{F}_3(R) = \emptyset$. If $L \in \mathcal{F}_2(R)$, we have $L^\vee \star L^\vee \subset t^{-1}L$, and $M = tL^\vee \star L^\vee + L = L$ is not self-dual. So that $L \notin \mathcal{F}_3(R)$.

The proof of Theorem 6.5 is similar to what we did in §5, i.e., we want to check to following two statements:

- For any k -algebra R , the subgroup of $LG(R)$ consisting of elements that stabilize the lattice $\mathbb{L}_2 \otimes_{k[[t]]} R[[t]]$ (resp. $\mathbb{L}_3 \otimes_{k[[t]]} R[[t]]$) agrees with $L^+\mathcal{G}_2(R)$ (resp. $L^+\mathcal{G}_3(R)$).
- For any $L \in \mathcal{F}_2(R)$ (resp. $L \in \mathcal{F}_3(R)$) with (R, \mathcal{M}) a local henselian ring with maximal ideal \mathcal{M} , there exists $g \in LG(R)$ such that $L = g(\mathbb{L}_2 \otimes_{k[[t]]} R[[t]])$ (resp. $L = g(\mathbb{L}_3 \otimes_{k[[t]]} R[[t]])$).

The first statement comes directly from Definition 6.3. To prove the second statement, we use the same idea that we did in §5: Find a basis in $L \in \mathcal{F}_2(R)$ (resp. $L \in \mathcal{F}_3(R)$), such that its multiplication table is the same as the multiplication table of $\mathbb{L}_2 \otimes_{k[[t]]} R[[t]]$ (resp. $\mathbb{L}_3 \otimes_{k[[t]]} R[[t]]$). The hard part is that we cannot find a basis $\{x_i\}_{i=1}^8$ in L such that $\langle e_i, e_{9-j} \rangle = \delta_{ij}$, since L is not a self-dual lattice anymore. We now turn to find “standard forms” of these lattice chains.

By [17, Appendix, Lemma A.21], after a finite unramified base change F'/F , there exists a “standard form” of lattice $L \otimes_{\mathcal{O}_F} \mathcal{O}_{F'}$. Generally, let V be an F -vector space with dimension $d = 2n$ or $d = 2n + 1$ (we assume $d \geq 5$), equipped with a non-degenerate symmetric form $\langle \cdot, \cdot \rangle$.

Consider a lattice $L \subset V$ satisfying $L \subset L^\vee \subset t^{-1}L$ with $l + l' = d$. Here l (resp. l') is the length $l = \lg(L^\vee/L)$ (resp. $l' = \lg(t^{-1}L/L^\vee)$). We assume $l \geq 1$ and $l' \geq 1$, i.e., $L \neq L^\vee$ and $L^\vee \neq t^{-1}L$. Since octonion algebras have dimension 8, we only list results for even dimension case.

For $d = 2n$, after a finite unramified extension, we can find a basis $\{x_i\}$ of $L_{\mathcal{O}_{F'}} = L \otimes_{\mathcal{O}_F} \mathcal{O}_{F'}$ as in one of the following cases:

(1) Split form: when $l = 2r$, we have $L_{\mathcal{O}_{F'}} = \bigoplus_{i=1}^d \mathcal{O}_{F'} \cdot x_i$ with

$$\begin{aligned} \langle x_i, x_{d+1-j} \rangle &= \delta_{ij}, \text{ for } i \notin [n-r+1, n+r]; \\ \langle x_i, x_{d+1-j} \rangle &= t\delta_{ij}, \text{ for } i \in [n-r+1, n+r]. \end{aligned}$$

(2) Quasi-split form: when $l = 2r + 1$, we have $L_{\mathcal{O}_{F'}} = \bigoplus_{i=1}^d \mathcal{O}_{F'} \cdot x_i$ with

$$\begin{aligned} \langle x_i, x_{d+1-j} \rangle &= \delta_{ij}, \text{ for } i \notin [n-r, n+r+1]; \\ \langle x_i, x_{d+1-j} \rangle &= \pi\delta_{ij}, \text{ for } i \in [n-r, n+r+1] \setminus \{n, n+1\}; \\ \langle x_n, x_n \rangle &= \pi, \quad \langle x_{n+1}, x_{n+1} \rangle = 1, \quad \langle x_n, x_{n+1} \rangle = 0. \end{aligned}$$

Here π is a uniformizer of $\mathcal{O}_{F'}$. In the rest of this section, we omit the base change of L for simplicity. Since a finite étale extension of a power series ring \mathcal{O}_F is still a power series ring, we still denote $k[[t]]$ as a power series ring $\mathcal{O}_{F'}$, and t is a uniformizer of $k[[t]]$.

In our case, L is a lattice in C_s with $d = 8$. No matter $L \in \mathcal{F}_2(R)$ or $L \in \mathcal{F}_3(R)$, we always have $L \subsetneq L^\vee \subsetneq t^{-1}L$. Hence $l \geq 1$ and $l' \geq 1$. We claim that $L \subset C_s$ is of split form.

Recall that the discriminant dL of $(L, \langle \cdot, \cdot \rangle)$ is the determinant of the matrix with respect to the symmetric form $\langle \cdot, \cdot \rangle$. By change of basis in L , two matrices are congruent to each other with respect to different basis (i.e., $B' = T^T B T$, where T is the change of basis matrix), so we have

$$dL \in k[[t]]^*/k[[t]]^{*2}.$$

The discriminant is a class invariant of quadratic forms. From above, for a lattice $L \subset V$, with $\dim V = 2n$. If L is of split form, we have $dL = 1$; if L is of quasi-split form, $dL = t$. In our case, L is a lattice of the split para-octonion algebra C_s . Since C_s can be viewed as an orthogonal sum of hyperbolic subspaces, we have $dC_s = 1$. Thus, if L is of quasi-split form, we have $L \otimes_{k[[t]]} k((t)) \simeq C_s$, and

$$d(L \otimes_{k[[t]]} k((t))) = t \neq 1 \in k((t))^*/k((t))^{*2},$$

which is a contradiction. Therefore, we only need to consider the split form case. More precisely: we consider the cases when $l = 2, 4, 6$. Here $L = \bigoplus_{i=1}^8 k[[t]]x_i$, and the symmetric forms $\langle \cdot, \cdot \rangle$ with respect to basis $\{x_i\}$ are:

$$\begin{aligned} \langle \cdot, \cdot \rangle &= \text{antidiag}(1, 1, 1, t, t, 1, 1, 1), \quad \text{if } l = 2; \\ \langle \cdot, \cdot \rangle &= \text{antidiag}(1, 1, t, t, t, t, 1, 1), \quad \text{if } l = 4; \\ \langle \cdot, \cdot \rangle &= \text{antidiag}(1, t, t, t, t, t, t, 1), \quad \text{if } l = 6. \end{aligned}$$

Here $\text{antidiag}(\dots)$ means an antidiagonal matrix. Our main conclusion is the following:

- when $l = 2$, we claim that L is not closed under multiplication ($L \star L \not\subseteq L$). Hence $L \notin \mathcal{F}_2(R)$ and $L \notin \mathcal{F}_3(R)$.
- when $l = 4$, we claim that there exists a basis of L , such that its multiplication table is the same as multiplication table in $\mathbb{L}_2 \otimes_{k[[t]]} R[[t]]$. Hence there exists $g \in LG(R)$, such

that $L = g(\mathbb{L}_2 \otimes_{k[t]} R[[t]])$. Then we get:

$$L \star L = g((\mathbb{L}_2 \otimes_{k[t]} R[[t]]) \star (\mathbb{L}_2 \otimes_{k[t]} R[[t])) \subset g(\mathbb{L}_2 \otimes_{k[t]} R[[t]]) = L;$$

$$L^\vee \star L^\vee = g((\mathbb{L}_2^\vee \otimes_{k[t]} R[[t]]) \star (\mathbb{L}_2^\vee \otimes_{k[t]} R[[t])) \subset g(t^{-1}(\mathbb{L}_2^\vee \otimes_{k[t]} R[[t]))) = t^{-1}L,$$

since $L^\vee = g(\mathbb{L}_2^\vee \otimes_{k[t]} R[[t]])$. So that $L \in \mathcal{F}_2(R)$.

- when $l = 6$, there exists a basis of L such that its multiplication table is the same as table in $\mathbb{L}_3 \otimes_{k[t]} R[[t]]$. Similarly, we have $L = g(\mathbb{L}_3 \otimes_{k[t]} R[[t]])$ for some $g \in LG(R)$, and we get

$$L \star L \subset L;$$

$$M = tL^\vee \star L^\vee + L \text{ is self-dual,}$$

since $M = g(t(\mathbb{L}_3^\vee \otimes_{k[t]} R[[t]]) \star (\mathbb{L}_3^\vee \otimes_{k[t]} R[[t])) + (\mathbb{L}_3 \otimes_{k[t]} R[[t]))$. So that $L \in \mathcal{F}_3(R)$.

Our claims as above will finish the proof of Theorem 6.5, and the proof of our claims is similar to §5.

Let L be a lattice with $e \in L$, $L \subsetneq L^\vee \subsetneq t^{-1}L$. No matter $l = \lg(L^\vee/L) = 2, 4$ or 6 , there exists a basis $\{x_i\}$ of L , such that the symmetric form is an anti-diagonal matrix with respect to $\{x_i\}$. Consider $e = \sum_{i=1}^8 a_i x_i \in L$. We set:

$$e_1 = \sum_{i=1}^4 a_i x_i, \quad e_2 = \sum_{i=5}^8 a_i x_i.$$

Then $q(e_1) = q(e_2) = 0$. We get $\langle e_1, e_2 \rangle = 1$ by $q(e) = 1$, and we also have $e_1 \star e_1 = e_2$, $e_2 \star e_2 = e_1$, $e_1 \star e_2 = e_2 \star e_1 = 0$ by similar calculation that we did in §5. Thus, $R[[t]]e_1 \oplus R[[t]]e_2$ is a hyperbolic subspace. By Nakayama's lemma, $\{e_1, e_2\}$ are primitive elements in L . Choose elements $y_3, \dots, y_8 \in L$ such that $\{e_1, e_2, y_3, \dots, y_8\}$ is a basis of L , By replacing y_i to

$$y'_i = y_i - \langle y_i, e_2 \rangle e_1 - \langle y_i, e_1 \rangle e_2,$$

we can always assume that $y_i \in L_0$, where

$$L_0 = \{x \in L \mid \langle x, e_1 \rangle = \langle x, e_2 \rangle = 0\}.$$

Therefore, $L = R[[t]]e_1 \oplus R[[t]]e_2 \oplus L_0$. Set $L_i = L_0 \star e_i$ for $i = 1, 2$.

Lemma 6.6. *We have*

- (1) $L_i \subset L_0$ for $i = 1, 2$.
- (2) $e_i \star L_i = 0$, $L_i \star e_{i+1} = 0$ for $i = 1, 2 \pmod{2}$.

We omit the proof of Lemma 6.6, since it is the same as the proof of Lemma 5.1. For any $x \in L_0$, we get

$$x = \langle e_1, e_2 \rangle x = (e_2 \star x) \star e_1 + (e_1 \star x) \star e_2 \in L_1 + L_2,$$

by Lemma 2.8. Hence $L_0 = L_1 + L_2$. We again get $L_0 = L_1 \oplus L_2$ since L_1, L_2 are totally isotropic with rank ≤ 3 . When we restrict the symmetric form to L_0 , there exists a basis $\{u_i\}_{i=1}^3$ of L_1 (resp. $\{v_i\}_{i=1}^3$ of L_2) such that

$$\langle u_i, v_j \rangle = 0,$$

for $i \neq j$, since the symmetric form is an anti-diagonal matrix.

Lemma 6.7. *We have*

(1)

$$\begin{aligned} u_i \star e_1 &= e_2 \star u_i = -u_i; \\ v_i \star e_2 &= e_1 \star u_i = -v_i, \end{aligned}$$

(2)

$$\begin{aligned} u_i \star v_j &= -\langle u_i, v_j \rangle e_1; \\ v_j \star u_i &= -\langle u_i, v_j \rangle e_2, \end{aligned}$$

(3)

$$u_i \star u_j = -u_j \star u_i \in L_2, \quad v_i \star v_j = -v_j \star v_i \in L_1.$$

for $i, j = 1, 2, 3$.

Proof. The proof of Lemma 6.6 (1) is similar to Lemma 5.2. For (2), we have

$$\begin{aligned} u_i \star v_j &= -(u_i \star e_1) \star v_j \\ &= (v_j \star e_1) \star u_i - \langle u_i, v_j \rangle e_1 \\ &= -\langle u_i, v_j \rangle e_1. \end{aligned}$$

Similarly, we also have $v_j \star u_i = -\langle u_i, v_j \rangle e_2$. For (3), it is easy to see that $u_i \star u_j \in L_0$ since $\langle u_i \star u_j, e_1 \rangle = \langle u_i \star u_j, e_2 \rangle = 0$. Consider

$$(e_2 \star u_i) \star u_j + (u_j \star u_i) \star e_2 = \langle e_2, u_j \rangle u_i = 0,$$

and $(e_2 \star u_i) \star u_j = -u_i \star u_j$ by (1). We get $u_i \star u_j = (u_j \star u_i) \star e_2 \in L_0 \star e_2 = L_2$. Thus, $(u_j \star u_i) \star e_2 = -u_j \star u_i$ since $v_i \star e_2 = -v_i$, which gives us $u_i \star u_j = -u_j \star u_i$. Similarly, we have $v_i \star v_j = -v_j \star v_i$. \square

So far we only use that the symmetric form is an antidiagonal matrix. To prove our claims, we now need to consider the explicit number of $l = \lg(L^\vee/L)$.

(a) $l = 2$ case. When $l = 2$, the restriction of $\langle \cdot, \cdot \rangle$ to L_0 is :

$$\langle \cdot, \cdot \rangle|_{L_0} = \text{antidiag}(1, 1, t, t, 1, 1).$$

Since $L_0 = L_1 \oplus L_2$ and L_1, L_2 are totally isotropic, there exists a basis $\{u_i\}_{i=1}^3$ of L_1 (resp. $\{v_i\}_{i=1}^3$ of L_2) such that

$$\begin{aligned} \langle u_1, v_1 \rangle &= t, & \langle u_2, v_2 \rangle &= 1; \\ \langle u_3, v_3 \rangle &= 1, & \langle u_i, v_j \rangle &= 0, \end{aligned}$$

for $i \neq j$. By Lemma 6.6, Lemma 6.7, we almost complete the multiplication table of L . The only left part is to understand $u_i \star u_j$ (resp. $v_i \star v_j$).

Consider $u_1 \star u_2 \in L_2$. Since

$$\langle u_1 \star u_2, u_1 \rangle = -\langle u_1 \star u_2, u_1 \star e_1 \rangle = q(u_1) \langle u_2, e_1 \rangle = 0,$$

by Lemma 2.7. Similarly, we get $\langle u_1 \star u_2, u_2 \rangle = 0$. So we can assume $u_1 \star u_2 = \lambda v_3$. Suppose that $L \star L \subset L$. We have $\lambda \in R[[t]]$. Since

$$(u_1 \star u_2) \star v_1 + (v_1 \star u_2) \star u_1 = \langle u_1, v_1 \rangle u_2 = t u_2,$$

and $v_1 \star u_2 = 0$ by Lemma 6.7 (2), we get $(u_1 \star u_2) \star v_1 = t u_2$. Thus $\lambda(v_3 \star v_1) = t u_2$. Similarly, consider $(u_1 \star u_2) \star v_2 = \lambda(v_3 \star v_2)$ and $(u_1 \star u_2) \star v_3 = \lambda(v_3 \star v_3)$, we get

$$\lambda(v_3 \star v_2) = -u_1, \quad v_3 \star v_3 = 0.$$

Thus, $\lambda^{-1} \in R[[t]]$, which implies $\lambda \in R[[t]]^*$. We can assume $\lambda = 1$ and get

$$v_3 \star v_1 = tu_2, \quad v_3 \star v_2 = -u_1, \quad v_3 \star v_3 = 0.$$

Next we consider $u_2 \star u_3 \in L_2$, we can assume $u_2 \star u_3 = \mu v_1$ for some $\mu \in R[[t]]$, since $\langle u_2 \star u_3, v_2 \rangle = \langle u_2 \star u_3, v_3 \rangle = 0$. Since

$$(u_2 \star u_3) \star v_3 + (v_3 \star u_3) \star u_2 = \langle u_2, v_3 \rangle u_3,$$

and $(v_3 \star u_3) \star u_2 = u_2, \langle u_2, v_3 \rangle = 0$ by Lemma 6.7 (1), (2), we get $\mu(v_1 \star v_3) = -u_2$. But $v_1 \star v_3 = -v_3 \star v_1 = -tu_2$ by our discussion above. Therefore

$$-t\mu u_2 = -u_2.$$

Hence $\mu = t^{-1} \notin R[[t]]$, which is a contradiction. So that $L \star L \not\subseteq L$.

(b) $l = 4$ case. When $l = 4$, the restriction of $\langle \cdot, \cdot \rangle$ to L_0 is :

$$\langle \cdot, \cdot \rangle|_{L_0} = \text{antidiag}(1, t, t, t, 1).$$

There exists a basis $\{u_i\}_{i=1}^3$ of L_1 (resp. $\{v_i\}_{i=1}^3$ of L_2) such that

$$\begin{aligned} \langle u_1, v_1 \rangle &= t, & \langle u_2, v_2 \rangle &= t; \\ \langle u_3, v_3 \rangle &= 1, & \langle u_i, v_j \rangle &= 0, \end{aligned}$$

for $i \neq j$. Assume $L \star L \subset L$. By using the same discussion as we did in $l = 2$ case, we get the following multiplication table:

TABLE 4. $u_i \star u_j$

*	u_1	u_2	u_3
u_1	0	tv_3	$-v_2$
u_2	$-tv_3$	0	v_1
u_3	v_2	$-v_1$	0

TABLE 5. $v_i \star v_j$

*	v_1	v_2	v_3
v_1	0	tu_3	$-u_2$
v_2	$-tu_3$	0	u_1
v_3	u_2	$-u_1$	0

It is easy to see that L has the same multiplication table as $\mathbb{L}_2 \otimes_{k[[t]]} R[[t]]$. Thus, there exists $g \in LG(R)$, such that $L = g(\mathbb{L}_2 \otimes_{k[[t]]} R[[t]])$, and we have:

$$L^\vee \star L^\vee \subset t^{-1}L.$$

So that $L \in \mathcal{F}_2(R)$.

(c) $l = 6$ case. When $l = 6$, the restriction of $\langle \cdot, \cdot \rangle$ to L_0 is :

$$\langle \cdot, \cdot \rangle|_{L_0} = \text{antidiag}(t, t, t, t, t, t).$$

There exists a basis $\{u_i\}_{i=1}^3$ of L_1 (resp. $\{v_i\}_{i=1}^3$ of L_2) such that

$$\begin{aligned} \langle u_1, v_1 \rangle &= t, & \langle u_2, v_2 \rangle &= t; \\ \langle u_3, v_3 \rangle &= t, & \langle u_i, v_j \rangle &= 0, \end{aligned}$$

for $i \neq j$. Assume $L \star L \subset L$. Again, by using the same discussion as we did in $l = 2$ and $l = 4$ case, we get the following multiplication table:

TABLE 6. $u_i * u_j$

*	u_1	u_2	u_3
u_1	0	tv_3	$-tv_2$
u_2	$-tv_3$	0	tv_1
u_3	tv_2	$-tv_1$	0

TABLE 7. $v_i * v_j$

*	v_1	v_2	v_3
v_1	0	u_3	$-u_2$
v_2	$-u_3$	0	u_1
v_3	u_2	$-u_1$	0

Thus L has the same multiplication table as $\mathbb{L}_3 \otimes_{k[[t]]} R[[t]]$. Similarly, there exists $L = g(\mathbb{L}_3 \otimes_{k[[t]]} R[[t]])$ for some $g \in LG(R)$, and we have

$$M = tL^\vee \star L^\vee + L \text{ is self-dual.}$$

So that $L \in \mathcal{F}_3(R)$.

Therefore, for $L \in \mathcal{F}_2(R)$, we show that $l = lg(L^\vee/L) = 4$ (if $l = 2$, $L \star L \not\subseteq L$; if $l = 6$, $L \in \mathcal{F}_3(R)$), and there exists $g \in LG(R)$, such that $L = g(\mathbb{L}_2 \otimes_{k[[t]]} R[[t]])$. For $L \in \mathcal{F}_3(R)$, we show that $l = lg(L^\vee/L) = 6$, and there exists $g \in LG(R)$, such that $L = g(\mathbb{L}_3 \otimes_{k[[t]]} R[[t]])$.

REFERENCES

- [1] S. Alsaody and P. Gille. Isotopes of octonion algebras, \mathbf{G}_2 -torsors and triality. *Adv. Math.*, 343:864–909, 2019.
- [2] A. Beauville and Y. Laszlo. Un lemme de descente. *C. R. Acad. Sci. Paris Sér. I Math.*, 320(3):335–340, 1995.
- [3] A. Beilinson and V. Drinfeld. Quantization of Hitchin’s integrable system and Hecke eigensheaves. preprint.
- [4] F. Bruhat and J. Tits. Groupes réductifs sur un corps local. *Inst. Hautes Études Sci. Publ. Math.*, (41):5–251, 1972.
- [5] F. Bruhat and J. Tits. Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée. *Inst. Hautes Études Sci. Publ. Math.*, (60):197–376, 1984.
- [6] F. Bruhat and J. Tits. Schémas en groupes et immeubles des groupes classiques sur un corps local. *Bull. Soc. Math. France*, 112(2):259–301, 1984.
- [7] F. Bruhat and J. Tits. Groupes algébriques sur un corps local. Chapitre III. Compléments et applications à la cohomologie galoisienne. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 34(3):671–698, 1987.
- [8] F. Bruhat and J. Tits. Schémas en groupes et immeubles des groupes classiques sur un corps local. II. Groupes unitaires. *Bull. Soc. Math. France*, 115(2):141–195, 1987.
- [9] W.T. Gan and J.-K. Yu. Schémas en groupes et immeubles des groupes exceptionnels sur un corps local. I. Le groupe G_2 . *Bull. Soc. Math. France*, 131(3):307–358, 2003.
- [10] L.J. Gerstein. *Basic Quadratic Forms. Graduate studies in mathematics.* American Mathematical Society, 2008.
- [11] U. Görtz. On the flatness of local models for the symplectic group. *Adv. Math.*, 176(1):89–115, 2003.
- [12] M.-A. Knus, A. Merkurjev, M. Rost, and J.-P. Tignol. *The book of involutions*, volume 44 of *American Mathematical Society Colloquium Publications.* American Mathematical Society, Providence, RI, 1998. With a preface in French by J. Tits.
- [13] G. Lusztig. Singularities, character formulas, and a q -analog of weight multiplicities. In *Analysis and topology on singular spaces, II, III (Luminy, 1981)*, volume 101 of *Astérisque*, pages 208–229. Soc. Math. France, Paris, 1983.
- [14] E. Matzri and U. Vishne. Isotropic subspaces in symmetric composition algebras and Kummer subspaces in central simple algebras of degree 3. *Manuscripta Math.*, 137(3-4):497–523, 2012.
- [15] G. Pappas and M. Rapoport. Twisted loop groups and their affine flag varieties. *Adv. Math.*, 219(1):118–198, 2008. With an appendix by T. Haines and Rapoport.
- [16] G. Prasad and J.-K. Yu. On finite group actions on reductive groups and buildings. *Invent. Math.*, 147(3):545–560, 2002.
- [17] M. Rapoport and Th. Zink. *Period spaces for p -divisible groups*, volume 141 of *Annals of Mathematics Studies.* Princeton University Press, Princeton, NJ, 1996.

- [18] B. Smithling. Topological flatness of orthogonal local models in the split, even case. I. *Math. Ann.*, 350(2):381–416, 2011.
- [19] T.A. Springer and F.D. Veldkamp. Octonions, Jordan algebras and exceptional groups. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2000.
- [20] F. van der Blij and T. A. Springer. The arithmetics of octaves and of the group G_2 . *Nederl. Akad. Wet., Proc., Ser. A*, 62:406–418, 1959.
- [21] F. van der Blij and T. A. Springer. Octaves and triality. *Nieuw Arch. Wisk. (3)*, 8:158–169, 1960.
- [22] Z. Zhao. Affine Grassmannians for triality groups. *J. Algebra*, 606:298–322, 2022.

MORNINGSIDE CENTER OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, CHINA

E-mail address: zhihaozhao@amss.ac.cn