

**FOURTH ORDER HARDY-SOBOLEV EQUATIONS:
SINGULARITY AND DOUBLY CRITICAL EXPONENT**

HUSSEIN CHEIKH ALI

ABSTRACT. In dimension $N \geq 5$ and given $0 < s < 4$ and $\gamma \in \mathbb{R}$, we study the existence of nontrivial weak solutions for the doubly critical problem

$$\Delta^2 u - \frac{\gamma}{|x|^4} u = |u|^{2_s^* - 2} u + \frac{|u|^{2_s^* - 2} u}{|x|^s} \text{ in } \mathbb{R}_+^N, \quad u = \Delta u = 0 \text{ on } \partial\mathbb{R}_+^N,$$

where $2_s^* := \frac{2(N-s)}{N-4}$ is the critical Hardy-Sobolev exponent. For $N \geq 8$ and $0 < \gamma < \frac{(N^2-4)^2}{16}$, we show the existence of nontrivial solution using the Mountain-Pass theorem by Ambrosetti-Rabinowitz. The method used is based on the existence of extremals for certain Hardy-Sobolev embeddings that we prove in this paper.

1. INTRODUCTION

Let $\mathbb{R}_+^N := \{x \in \mathbb{R}^N \text{ such that } x_1 > 0\}$ be the half-space domain, $N \geq 5$ and $0 \leq s < 4$. In this work, we establish the existence of nontrivial weak solutions for the following doubly critical problem:

$$(1) \quad \begin{cases} \Delta^2 u - \frac{\gamma}{|x|^4} u = |u|^{2_s^* - 2} u + \frac{|u|^{2_s^* - 1} u}{|x|^s} & \text{in } \mathbb{R}_+^N, \\ u = \Delta u = 0 & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

where $\Delta = \operatorname{div}(\nabla)$ is the Laplacian, $\gamma \in \mathbb{R}$ and $2_s^* := \frac{2(N-s)}{N-4}$ is the critical Hardy-Sobolev exponent. In order to write a variational formulation, the relevant space is the following: given Ω be a smooth domain in \mathbb{R}^N , define

$$\tilde{H}_0(\Omega) = \text{completion of } \{u \in C_c^2(\bar{\Omega}) \text{ s.t. } u|_{\partial\Omega} \equiv 0\} \text{ for the norm } u \mapsto \|\Delta u\|_2.$$

Note that $\tilde{H}_0(\mathbb{R}^N)$ is the usual Beppo-Levi space $D^{2,2}(\mathbb{R}^N)$. We say that $u \in \tilde{H}_0(\mathbb{R}_+^N)$ is a weak solution of (1), if

$$\int_{\mathbb{R}_+^N} \Delta u \Delta \varphi \, dx - \gamma \int_{\mathbb{R}_+^N} \frac{u\varphi}{|x|^4} \, dx = \int_{\mathbb{R}_+^N} |u|^{2_s^* - 2} u \varphi \, dx + \int_{\mathbb{R}_+^N} \frac{|u|^{2_s^* - 2} u \varphi}{|x|^s} \, dx,$$

for all $\varphi \in \tilde{H}_0(\mathbb{R}_+^N)$. For any $u \in \tilde{H}_0(\mathbb{R}_+^N)$, Sobolev's embedding yields $u \in L^{2_s^*}(\mathbb{R}_+^N) \cap L^{2_s^*}(\mathbb{R}_+^N, |x|^{-s})$ (see (5) below), and therefore the definition of weak solution makes sense.

The existence of weak solution to (1) on \mathbb{R}^N has been studied by Filippucci-Pucci-Robert [10] for the p-Laplacian. Equations like (1) have been studied for the Fractional Laplacian (see Ghoussoub-Shakerian [14]) and the bi-laplacian in \mathbb{R}^N (see Bhakta [2, 3] and Bhakta-Musina [4]).

In the present paper, we tackle this type of nonlinear singular problems on \mathbb{R}_+^N when $0 \in \partial\mathbb{R}_+^N$. The case $0 \in \partial\Omega$, when Ω be a domain of \mathbb{R}^N , ($N \geq 3$) to

the equations (1) with the Laplace operator $-\Delta$ and one critical nonlinearity was initiated by Ghoussoub–Kang [13] and studied by Chern–Lin [9] and Ghoussoub–Robert [11],[12]) when Ω is a smooth domain. For non-smooth domains modeled on cones, we refer to the more recent works of Cheikh-Ali [5, 6].

Our main result is the following:

Theorem 1.1. *Let $N \geq 8$, $0 < s < 4$ and $0 < \gamma < \frac{(N^2-4)^2}{16}$. Then, there exists a nontrivial weak solution of (1).*

Let us discuss the hypothesis of the Theorem. Our problem depends of the constant γ in \mathbb{R} . From here, given an arbitrary domain $\Omega \subset \mathbb{R}^N$, $N \geq 5$, we define the Hardy-Rellich constant

$$(2) \quad \gamma_H(\Omega, X) := \inf \left\{ \frac{\int_{\Omega} |\Delta u|^2 dx}{\int_{\Omega} \frac{u^2}{|x|^4} dx}; u \in X \setminus \{0\} \right\},$$

for a suitable space X . There are several references around this constant. A first version of this Hardy inequality has been introduced by Rellich in 1953 [19, 20] (see also Mitidieri [17]), and reads

$$(3) \quad \gamma_H(\mathbb{R}^N, C_c^2(\mathbb{R}^N)) = \frac{N^2(N-4)^2}{16}.$$

As a consequence, for Ω be a bounded domain in \mathbb{R}^N with $0 \in \Omega$, Perez-Llamos-Primo [18] proved that

$$\gamma_H(\Omega, H^2(\Omega) \cap H_0^1(\Omega)) = \frac{N^2(N-4)^2}{16}.$$

In particular, the value of the constant is independent of the domain as long as 0 is an interior point.

The situation is different when $0 \in \partial\Omega$. Let us consider cones. For any regular domain Σ in the unit sphere \mathbb{S}^{N-1} , we define the cone

$$C_{\Sigma} := \{r\sigma \mid r > 0, \sigma \in \Sigma\}.$$

Caldirola-Musina [8] proved that

$$\gamma_H(C_{\Sigma}, X_0(C_{\Sigma})) = \text{dist} \left(-\frac{N(N-4)}{4}, \Lambda(\Sigma) \right)^2,$$

where

$$X_0(\Omega) := \{u \in C^2(\bar{\Omega}) \cap C_c^2(\mathbb{R}^N \setminus \{0\}) \text{ s.t. } u|_{\partial\Omega} \equiv 0\}.$$

and $\Lambda(\Sigma)$ of the Laplace–Beltrami operator on Σ . For instance:

- If $\Sigma = \mathbb{S}^{N-1}$, then $C_{\Sigma} = \mathbb{R}^N \setminus \{0\}$. Therefore, we have

$$\gamma_H(\mathbb{R}^N \setminus \{0\}, X_0(\mathbb{R}^N \setminus \{0\})) = \min_{k \in \{0, 1, \dots\}} |\gamma_N + k(N-2+k)|^2 = \frac{N^2(N-4)^2}{16}.$$

- If Σ is the half-sphere S_+^{N-1} , then $C_{\Sigma} = \mathbb{R}_+^N$. Therefore, we have

$$(4) \quad \gamma_H(\mathbb{R}_+^N, X_0(\mathbb{R}_+^N)) = \min_{k \in \{1, 2, \dots\}} |\gamma_N + k(N-2+k)|^2 = \frac{(N^2-4)^2}{16}.$$

In the sequel, we write for convenience

$$\gamma_{H,+} := \gamma_H(\mathbb{R}_+^N, \tilde{H}_0(\mathbb{R}_+^N)).$$

In order to prove Theorem 1.1, we will prove in Section 2 that

$$\gamma_{H,+} = \gamma_H(\mathbb{R}_+^N, X_0(\mathbb{R}_+^N)) = \frac{(N^2 - 4)^2}{16}.$$

In order to tackle the nonlinear problem (1), let us consider the Sobolev inequality

$$(5) \quad \left(\int_{\mathbb{R}^N} \frac{|u|^{2_s^*}}{|x|^s} dx \right)^{\frac{2}{2_s^*}} \leq C \int_{\mathbb{R}^N} |\Delta u|^2 dx \text{ for all } u \in C_c^\infty(\mathbb{R}^N).$$

Let us fix a domain $\Omega \subset \mathbb{R}^N$. Interpolating the Hardy inequality (2) and the Sobolev inequality (5) and using that $X_0(\Omega) \subset C_c^\infty(\mathbb{R}^N)$, given $s \in [0, 4]$, we get the existence of $C(N, s, \Omega) > 0$ such that

$$(6) \quad \left(\int_{\Omega} |x|^{-s} |u|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \leq C(N, s, \Omega) \int_{\Omega} |\Delta u|^2 dx \text{ for any } u \in C_c^\infty(\Omega)$$

Using again (2), for any $s \in [0, 4]$ and any $\gamma < \gamma_H(\Omega, \tilde{H}_0(\Omega))$, we get the Hardy-Sobolev inequality

$$(7) \quad \left(\int_{\Omega} |x|^{-s} |u|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \leq C \int_{\mathbb{R}^N} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx \text{ for all } u \in \tilde{H}_0(\Omega).$$

Let us define

$$(8) \quad Q_{\gamma,s}(\Omega) = \inf \left\{ I_{\gamma,s}^\Omega(u)/u \in \tilde{H}_0(\Omega) \setminus \{0\} \right\}, \text{ with } I_{\gamma,s}^\Omega(u) := \frac{\int_{\Omega} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx}{\left(\int_{\Omega} \frac{|u|^{2_s^*}}{|x|^s} dx \right)^{\frac{2}{2_s^*}}}.$$

In order to prove the existence of weak solutions for (1), we need extremals for the best constant $Q_{\gamma,s}(\mathbb{R}_+^N)$ in (8), that is $u \in \tilde{H}_0(\mathbb{R}_+^N) \setminus \{0\}$ such that $I_{\gamma,s}^{\mathbb{R}_+^N}(u) = Q_{\gamma,s}(\mathbb{R}_+^N)$. If $u \in \tilde{H}_0(\mathbb{R}_+^N) \setminus \{0\}$ is an extremal for $Q_{\gamma,s}(\mathbb{R}_+^N) > 0$, then, up to a constant factor, u is a solution to the following Euler-Lagrange equation:

$$(9) \quad \begin{cases} \Delta^2 u - \frac{\gamma}{|x|^4} u = \frac{|u|^{2_s^*-2} u}{|x|^s} & \text{in } \mathbb{R}_+^N, \\ u = \Delta u = 0 & \text{on } \partial\mathbb{R}_+^N. \end{cases}$$

The existence of extremals is mostly ruled by the following Theorem that we prove in Section 3:

Theorem 1.2. *For $s \in [0, 4]$ and $\gamma < \gamma_{H,+}$, we have that*

- (a) *If $\{s > 0\}$ or $\{s = 0, \gamma > 0, N \geq 8\}$, then there are extremals for $Q_{\gamma,s}(\mathbb{R}_+^N)$.*
- (b) *If $\{s = 0 \text{ and } \gamma \leq 0\}$, there are no extremals for $Q_{\gamma,s}(\mathbb{R}_+^N)$.*
- (c) *If there are no extremals for $Q_{\gamma,0}(\mathbb{R}_+^N)$, then $Q_{\gamma,0}(\mathbb{R}_+^N) = S_N$,*

where

$$(10) \quad S_N := \inf_{u \in \tilde{H}_0(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |\Delta u|^2 dx}{\left(\int_{\mathbb{R}^N} |u|^{2_0^*} dx \right)^{\frac{2}{2_0^*}}}.$$

The remaining case, that is $\{s = 0, \gamma > 0, \text{ and } N = 5, 6, 7\}$, is not clear. This limitation is due to the lack of localization of the L^2 -norm in the computation (70).

The study of fourth-order Hardy-Sobolev problem with a singularity on the boundary of a smooth domain is the object of the work [7].

With all these elements, we will construct weak solutions for the doubly critical problem (1) by finding critical points of corresponding functional on $\tilde{H}_0(\mathbb{R}_+^N)$. The method to obtain this critical points is via the Mountain-Pass Theorem of Ambrosetti and Rabinowitz. Since (1) is invariant under the conformal one parameter transformation group:

$$\left\{ \begin{array}{l} T_r : \tilde{H}_0(\mathbb{R}_+^N) \rightarrow \tilde{H}_0(\mathbb{R}_+^N) \\ u \mapsto T_r[u] := \{x \mapsto r^{\frac{N-4}{2}} u(rx)\} \end{array} \right\}, \text{ where } r > 0,$$

then the Mountain-Pass theorem will not yield critical points, but only Palais-Smale sequences. We will use the strategy of Filippucci-Pucci-Robert [10]. As in [10], we the main difficulty will be the asymptotic competition between the energy carried by the two critical nonlinearities. Hence, the crucial point here is to balance the competition to avoid the domination of one term over another. Otherwise, there is vanishing of the weakest one, and we get solution for the same equation but with only one critical nonlinearity. To deal with this problem, we will choose a Palais-Smale sequence at a minimax energy level. In such a way, after a careful analysis of concentration, we will show that there is a balance between the energies of the two nonlinearities mentioned above, and therefore none can dominate the other. This will yield a solution to (1).

2. PROFILE OF SOLUTIONS AND STUDY THE VALUE OF HARDY RELICH CONSTANT $\gamma_{H,+}$

First result, we are interested by determine the value of Hardy Rellich constant $\gamma_{H,+} := \gamma_H(\mathbb{R}_+^N, \tilde{H}_0(\mathbb{R}_+^N))$ when $0 \in \partial\mathbb{R}_+^N$.

Lemma 2.1. *We have that*

$$(11) \quad \gamma_{H,+} = \gamma_H(\mathbb{R}_+^N, X_0(\mathbb{R}_+^N)) = \frac{(N^2 - 4)^2}{16}.$$

Proof of Lemma 2.1: We prove the Lemma in two steps proving each an inequality.

Step 2.1. *We claim that*

$$(12) \quad \int_{\mathbb{R}_+^N} |\Delta v|^2 dx \geq \frac{(N^2 - 4)^2}{16} \int_{\mathbb{R}_+^N} \frac{v^2}{|x|^4} dx \text{ for all } v \in C_c^2(\overline{\mathbb{R}_+^N}) \text{ s.t. } u|_{\partial\mathbb{R}_+^N} = 0.$$

Proof of Step 2.1: Fix $v \in C_c^2(\overline{\mathbb{R}_+^N})$ such that $v|_{\partial\mathbb{R}_+^N} = 0$. We choose $\eta \in C^\infty(\mathbb{R}^N)$ such that $\eta(x)|_{B_1(0)} \equiv 0$, $\eta(x)|_{B_2(0)^c} \equiv 1$ and $0 \leq \eta \leq 1$. For $\epsilon > 0$, we set $v_\epsilon(x) := \eta_\epsilon(x)v(x)$, where $\eta_\epsilon(x) := \eta(\frac{x}{\epsilon})$ for all $x \in \mathbb{R}^N$. We have that $v_\epsilon \in X_0(\mathbb{R}_+^N)$. It follows from Caldiroli-Musina [8] (see (4)) that

$$(13) \quad \int_{\mathbb{R}_+^N} |\Delta v_\epsilon|^2 dx \geq \frac{(N^2 - 4)^2}{16} \int_{\mathbb{R}_+^N} \frac{v_\epsilon^2}{|x|^4} dx.$$

Step 2.1.1 We claim that

$$(14) \quad \int_{\mathbb{R}_+^N} |\Delta v_\epsilon|^2 dx = \int_{\mathbb{R}_+^N} |\eta_\epsilon|^2 |\Delta v|^2 dx + o(1) \text{ as } \epsilon \rightarrow 0.$$

Proof of this claim: For convenience $A_{+,\epsilon} := \mathbb{R}_+^N \cap (B_{2\epsilon}(0) \setminus B_\epsilon(0))$. Using the definition of v_ϵ yields

$$(15) \quad \int_{\mathbb{R}_+^N} |\Delta v_\epsilon|^2 dx = \int_{\mathbb{R}_+^N} |\eta_\epsilon|^2 |\Delta v|^2 dx + R_\epsilon, \text{ with}$$

$$(16) \quad R_\epsilon := \int_{A_{+,\epsilon}} \left[|\Delta \eta_\epsilon|^2 v^2 + 4 \nabla \eta_\epsilon \cdot \nabla v \Delta(\eta_\epsilon) v + 2 \Delta(\eta_\epsilon) \eta_\epsilon v \Delta v + 4 (\nabla \eta_\epsilon \cdot \nabla v)^2 + 4 \nabla \eta_\epsilon \cdot \nabla v \eta_\epsilon \Delta v \right] dx.$$

We claim that

$$(17) \quad R_\epsilon = O \left(\int_{A_{+,\epsilon}} \frac{|v|^2}{|x|^4} dx + \left(\int_{A_{+,\epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+,\epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{4}} \right).$$

Proof. We estimate each term of R_ϵ . Since $v|_{\partial \mathbb{R}_+^N} = 0$, integrating by parts yield

$$\begin{aligned} \int_{A_{+,\epsilon}} \nabla \eta_\epsilon \cdot \nabla v \Delta(\eta_\epsilon) v dx &= O \left(- \int_{A_{+,\epsilon}} (|\Delta \eta_\epsilon|^2 + \nabla \eta_\epsilon \cdot \nabla(\Delta \eta_\epsilon)) v^2 dx \right. \\ &\quad \left. + \int_{\mathbb{R}_+^N \cap \partial(B_{2\epsilon}(0) \setminus B_\epsilon(0))} v^2 \Delta \eta_\epsilon \partial_\nu \eta_\epsilon d\sigma \right), \end{aligned}$$

where ν is the outer normal vector of $B_{2\epsilon}(0) \setminus B_\epsilon(0)$. Since $\partial_\nu \eta_\epsilon = 0$ on $\mathbb{R}_+^N \cap \partial(B_{2\epsilon}(0) \setminus B_\epsilon(0))$, we have

$$(18) \quad \begin{aligned} \int_{A_{+,\epsilon}} \nabla \eta_\epsilon \cdot \nabla v \Delta(\eta_\epsilon) v dx &= O \left(\frac{1}{\epsilon^4} \|(\Delta \eta)^2 + \nabla \eta \cdot \nabla(\Delta \eta)\|_\infty \int_{A_{+,\epsilon}} v^2 dx \right) \\ &= O \left(\int_{A_{+,\epsilon}} \frac{|v|^2}{|x|^4} dx \right). \end{aligned}$$

By Hölder's inequality and $v \in C_c^2(\overline{\mathbb{R}_+^N})$, we get

$$(19) \quad \begin{aligned} \int_{A_{+,\epsilon}} \Delta(\eta_\epsilon) \eta_\epsilon v \Delta v dx &= O \left(\left(\int_{A_{+,\epsilon}} |\Delta v|^2 dx \right)^{\frac{1}{2}} \left(\int_{A_{+,\epsilon}} |\Delta(\eta_\epsilon) \eta_\epsilon v|^2 dx \right)^{\frac{1}{2}} \right) \\ &= O \left(\left(\int_{A_{+,\epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{2}} \right). \end{aligned}$$

It follows from the Cauchy-Schwarz and Hölder inequalities and integrations by parts that

$$\begin{aligned}
\int_{A_{+, \epsilon}} (\nabla \eta_\epsilon \cdot \nabla v)^2 dx &= O \left(\int_{A_{+, \epsilon}} |\nabla \eta_\epsilon|^2 |\nabla v|^2 dx \right) \\
&= O \left(\int_{A_{+, \epsilon}} v \Delta v |\nabla \eta_\epsilon|^2 dx + \int_{A_{+, \epsilon}} v \nabla v \nabla (|\nabla \eta_\epsilon|^2) dx \right) \\
(20) \qquad \qquad \qquad &= O \left(\left(\int_{A_{+, \epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{2}} + \int_{A_{+, \epsilon}} \frac{|v|^2}{|x|^4} dx \right).
\end{aligned}$$

Using again Hölder's inequality yields

$$\begin{aligned}
\int_{A_{+, \epsilon}} \nabla \eta_\epsilon \cdot \nabla v \eta_\epsilon \Delta v dx &= O \left(\left(\int_{A_{+, \epsilon}} |\nabla \eta_\epsilon|^2 |\nabla v|^2 dx \right)^{\frac{1}{2}} \right) \\
(21) \qquad \qquad \qquad &= O \left(\left(\int_{A_{+, \epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+, \epsilon}} \frac{|v|^2}{|x|^4} dx \right)^{\frac{1}{4}} \right).
\end{aligned}$$

We inject (18), (19), (20) and (21) in (16), we obtain (17). \square

It follows from (17) that $R_\epsilon = o(1)$ as $\epsilon \rightarrow 0$. Therefore, by (15) we obtain as $\epsilon \rightarrow 0$ that (14). This ends the proof of Step 2.1. \square

Using again the inequality (13) and by (14), we find that

$$\int_{\mathbb{R}_+^N} |\eta_\epsilon|^2 |\Delta v|^2 dx + o(1) \geq \frac{(N^2 - 4)^2}{16} \int_{\mathbb{R}_+^N} \frac{|\eta_\epsilon|^2 v^2}{|x|^4} dx.$$

Therefore, passing $\epsilon \rightarrow 0$, we get (12). This proves Step 2.1. \square

Step 2.2. *We claim that*

$$(22) \qquad \int_{\mathbb{R}_+^N} |\Delta u|^2 dx \geq \frac{(N^2 - 4)^2}{16} \int_{\mathbb{R}_+^N} \frac{u^2}{|x|^4} dx \text{ for all } u \in \tilde{H}_0(\mathbb{R}_+^N).$$

Proof of Step 2.2: We fix $u \in \tilde{H}_0(\mathbb{R}_+^N)$. We then take a sequence $(u_n)_n$ such that $u_n \in C_c^2(\overline{\mathbb{R}_+^N})$ and $u_n(x) = 0$ for all $x \in \partial \mathbb{R}_+^N$ and $\lim_{n \rightarrow +\infty} u_n = u$ for the norm $\|\Delta \cdot\|_2$. Therefore,

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{u_n^2}{|x|^4} dx = \int_{\mathbb{R}_+^N} \frac{u^2}{|x|^4} dx \text{ and } \lim_{n \rightarrow +\infty} \int_{\mathbb{R}_+^N} |\Delta u_n|^2 dx = \int_{\mathbb{R}_+^N} |\Delta u|^2 dx.$$

It then follows from (12) that

$$\int_{\mathbb{R}_+^N} |\Delta u_n|^2 dx \geq \frac{(N^2 - 4)^2}{16} \int_{\mathbb{R}_+^N} \frac{u_n^2}{|x|^4} dx.$$

Letting $n \rightarrow +\infty$ and we get (22). This proves Step 2.2. \square

Step 2.3. *We claim that (22) is optimal.*

Proof of Step 2.3: This will be achieved via test-function estimates. We define

$$v(x) := x_1|x|^{-\frac{N-2}{2}} \text{ for all } x \in \mathbb{R}_+^N \setminus \{0\}.$$

We have that $v \in C^2(\overline{\mathbb{R}_+^N})$, $v = 0$ on $\partial\mathbb{R}_+^N$ and

$$-\Delta v(x) = \frac{N^2 - 4}{16} \frac{v(x)}{|x|^2}.$$

Let $\varphi, \psi \in C^\infty(\mathbb{R}^n)$ such that

$$\begin{cases} \varphi(0) = 0, |\varphi(x)| \leq c|x| \text{ if } |x| < 1 \text{ and } \varphi(x) = 1 \text{ if } |x| \geq 1; \\ \psi(x) = 1 \text{ if } |x| < 1 \text{ and } \psi(x) = 0 \text{ if } |x| > 2, \end{cases}$$

for some constant $c > 0$. For $0 < \epsilon \ll 1$, we define the function $v_\epsilon \in \tilde{H}_0(\mathbb{R}_+^N)$ as follows:

$$v_\epsilon(x) := \varphi\left(\frac{x}{\epsilon}\right) \psi(\epsilon x) v(x) = \begin{cases} \varphi\left(\frac{x}{\epsilon}\right) v(x) & \text{if } |x| < \epsilon, \\ v(x) & \text{if } \epsilon \leq |x| < \frac{1}{\epsilon}, \\ \psi(\epsilon x) v(x) & \text{if } |x| \geq \frac{1}{\epsilon}. \end{cases}$$

It follows from the definition of v_ϵ that

$$(23) \quad \int_{\mathbb{R}_+^N \setminus \overline{B_{\epsilon^{-1}}(0)}} |\Delta v_\epsilon|^2 dx = O(1) \text{ and } \int_{\mathbb{R}_+^N \cap B_\epsilon(0)} |\Delta v_\epsilon|^2 dx = O(1) \text{ as } \epsilon \rightarrow 0.$$

It remains just one factor to calculate, we have

$$(24) \quad \int_{B_{\epsilon^{-1}}(0) \setminus \overline{B_\epsilon(0)}} |\Delta v_\epsilon|^2 dx = \frac{(N^2 - 4)^2}{16} \int_{B_{\epsilon^{-1}}(0) \setminus \overline{B_\epsilon(0)}} \frac{v_\epsilon^2}{|x|^4} dx.$$

Moreover, we calculate that

$$(25) \quad \begin{aligned} \int_{B_{\epsilon^{-1}}(0) \setminus \overline{B_\epsilon(0)}} \frac{v_\epsilon^2}{|x|^4} dx &= \int_{B_{\epsilon^{-1}}(0) \setminus \overline{B_\epsilon(0)}} x_1^2 |x|^{-N-2} dx \\ &= 2w(2) \ln\left(\frac{1}{\epsilon}\right), \end{aligned}$$

where $w(2) := \int_{\mathbb{S}_+^{N-1}} x_1^2 d\sigma$. It follows from (23), (24) and (25) that

$$(26) \quad \int_{\mathbb{R}_+^N} |\Delta v_\epsilon|^2 dx = 2w(2) \frac{(N^2 - 4)^2}{16} \ln\left(\frac{1}{\epsilon}\right) + O(1),$$

when $\epsilon \rightarrow 0$. Using again the definition of v_ϵ yields,

$$(27) \quad \int_{\mathbb{R}_+^N \setminus \overline{B_{\epsilon^{-1}}(0)}} \frac{v_\epsilon^2}{|x|^4} dx = O(1) \quad \text{and} \quad \int_{\mathbb{R}_+^N \cap B_\epsilon(0)} \frac{v_\epsilon^2}{|x|^4} dx = O(1).$$

Therefore, it follows from (25), we get as $\epsilon \rightarrow 0$ that

$$(28) \quad \int_{\mathbb{R}_+^N} \frac{v_\epsilon^2}{|x|^4} dx = 2w(2) \ln\left(\frac{1}{\epsilon}\right) + O(1).$$

Combining (26) and (28), we get

$$\frac{\int_{\mathbb{R}_+^N} |\Delta v_\epsilon|^2 dx}{\int_{\mathbb{R}_+^N} \frac{v_\epsilon^2}{|x|^4} dx} = \frac{(N^2 - 4)^2}{16} + o(1),$$

as $\epsilon \rightarrow 0$. Since $v_\epsilon \in \tilde{H}_0(\mathbb{R}_+^N)$, we get that $\gamma_H(\mathbb{R}_+^N, \tilde{H}_0(\mathbb{R}_+^N)) \leq \frac{(N^2-4)^2}{16}$. This completes the proof of Lemma 2.1 \square

To conclude this section, we discuss the model solutions for the homogeneous equation.

Lemma 2.2. *Given $\alpha \in \mathbb{R}$, for $x \in \mathbb{R}_+^N$, $N \geq 5$, we define $v_\alpha(x) := x_1|x|^{-\alpha}$. Then for $-N^2 \leq \gamma < \gamma_{H,+}$, we have that*

$$(29) \quad \begin{cases} \Delta^2 v_\alpha - \frac{\gamma}{|x|^4} v_\alpha = 0 & \text{in } \mathbb{R}_+^N, \\ v_\alpha = \Delta v_\alpha = 0 & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

if and only if $\alpha \in \{\alpha_-(\gamma), \alpha_+(\gamma), \beta_-(\gamma), \beta_+(\gamma)\}$ where

$$(30) \quad \alpha_\pm(\gamma) := \frac{N-2}{2} \pm \frac{1}{2} \sqrt{N^2 + 4 - 4\sqrt{N^2 + \gamma}},$$

and,

$$(31) \quad \beta_\pm(\gamma) := \frac{N-2}{2} \pm \frac{1}{2} \sqrt{N^2 + 4 + 4\sqrt{N^2 + \gamma}}.$$

Proof of Lemma 2.2: First, it follows from $-\Delta v = \alpha(N - \alpha) \frac{x_1|x|^{-\alpha}}{|x|^2}$ on \mathbb{R}_+^N that

$$\Delta^2 v - \frac{\gamma}{|x|^4} v = \{\alpha(N - \alpha)(\alpha + 2)(N - \alpha - 2) - \gamma\} \frac{v}{|x|^4}.$$

To get our result, we want to solve the following equation:

$$(32) \quad \alpha^4 - 2(N-2)\alpha^3 + (N^2 - 6N + 4)\alpha^2 + (2N^2 - 4N)\alpha - \gamma = 0.$$

We denote $a = 1$, $b = -2(N-2)$, $c = N^2 - 6N + 4$ and $d = 2N^2 - 4N$. Since $b^3 - 4abc + 8a^2d = 0$ then we can transform (32) to biquadratic equation. We take $\alpha := t - \frac{b}{4a}$, and it follows from (32) that

$$(33) \quad t^4 - [(N+2)^2 + (N-2)^2] \frac{t^2}{4} + \frac{(N+2)^2(N-2)^2}{16} - \gamma = 0.$$

It's easy to find the roots of (33). Since $-N^2 \leq \gamma < \gamma_{H,+}$, we find that

$$t_\pm(\gamma) := \pm \frac{1}{2} \sqrt{N^2 + 4 \pm 4\sqrt{N^2 + \gamma}}.$$

Therefore, since $\alpha = t + \frac{N-2}{2}$ we get (30). Then, our function v is a solution of equation (29) when $\alpha \in \{\alpha_\pm(\gamma), \beta_\pm(\gamma)\}$. This ends the proof of Lemma 2.2. \square

Remarks about this Lemma: First, we denote that

$$\beta_-(\gamma) \leq \alpha_-(\gamma) \leq \alpha_+(\gamma) \leq \beta_+(\gamma).$$


If $\gamma = \gamma_{H,+}$, then we have that $\alpha_\pm(\gamma) = \frac{N-2}{2}$. Hence, for $\gamma \in [0, \gamma_{H,+})$, we find that

$$\alpha_-(\gamma) \in \left[0, \frac{N-2}{2}\right) \text{ and } \alpha_+(\gamma) \in \left(\frac{N-2}{2}, N-2\right].$$

Moreover, notice that if $\gamma \in [0, \gamma_{H,+})$, then

$$\beta_-(\gamma) \in (\beta_-(\gamma_{H,+}), -2] \text{ and } \beta_+(\gamma) \in [N, \beta_+(\gamma_{H,+})).$$

We also remark that $x_1|x|^{-\beta_-}$ is bounded and $x_1|x|^{-\alpha_-}$ is the singular solution that is locally in $\tilde{H}_0(\mathbb{R}_+^N)$. The following graph concerns the localizations of α_\pm, β_\pm when $\gamma \in [0, \gamma_{H,+})$:



3. EXISTENCE OF EXTREMALS FOR $Q_{\gamma,s}(\mathbb{R}_+^N)$: PROOF OF THEOREM 1.2

We fix $\gamma < \gamma_{H,+} = \frac{(N^2-4)^2}{16}$ and $0 \leq s < 4$. Recall that

$$(34) \quad Q_{\gamma,s}(\mathbb{R}_+^N) = \inf_{u \in \tilde{H}_0(\mathbb{R}_+^N) \setminus \{0\}} \frac{\int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{|u|^2}{|x|^4} \right) dx}{\left(\int_{\mathbb{R}_+^N} \frac{|u|^{2^*_s}}{|x|^s} dx \right)^{\frac{2}{2^*_s}}} > 0.$$

In order to prove the existence of extremals for (34), we proceed as in Ghoussoub-Robert [12] (see Filippucci-Pucci-Robert [10]): these proofs rely on Lions's proof of the existence of extremals for the Sobolev inequality [16]. We let $(\bar{u}_m)_{m \in \mathbb{N}} \in \tilde{H}_0(\mathbb{R}_+^N)$ be a minimizing sequence for $Q_{\gamma,s}(\mathbb{R}_+^N)$ in (34) such that

$$\int_{\mathbb{R}_+^N} \frac{|\bar{u}_m|^{2^*_s}}{|x|^s} dx = 1 \text{ and } \lim_{m \rightarrow +\infty} \left(\int_{\mathbb{R}_+^N} |\Delta \bar{u}_m|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|\bar{u}_m|^2}{|x|^4} dx \right) = Q_{\gamma,s}(\mathbb{R}_+^N).$$

For all $m \in \mathbb{N}$, since $\int_{\mathbb{R}_+^N} \frac{|\bar{u}_m|^{2^*_s}}{|x|^s} dx = 1$, then, up to considering a subsequence, there exists $\rho_m > 0$ such that $\int_{\mathbb{R}_+^N \cap B_{\rho_m}(0)} \frac{|\bar{u}_m|^{2^*_s}}{|x|^s} dx = \frac{1}{2}$. We define

$$u_m(x) := \rho_m^{\frac{N-4}{2}} \bar{u}_m(\rho_m x) \text{ for any } x \in \mathbb{R}_+^N.$$

It is easy to check that $u_m \in \tilde{H}_0(\mathbb{R}_+^N)$ for all $m \in \mathbb{N}$, and

$$(35) \quad \lim_{m \rightarrow +\infty} \left(\int_{\mathbb{R}_+^N} |\Delta u_m|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx \right) = Q_{\gamma,s}(\mathbb{R}_+^N),$$

and

$$(36) \quad \int_{\mathbb{R}_+^N} \frac{|u_m|^{2^*_s}}{|x|^s} dx = 1 \text{ and } \int_{\mathbb{R}_+^N \cap B_1(0)} \frac{|u_m|^{2^*_s}}{|x|^s} dx = \frac{1}{2},$$

for all $m \in \mathbb{N}$. Since $\gamma < \gamma_{H,+}$, there exists $C > 0$ such that

$$(37) \quad \int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{|u|^2}{|x|^4} \right) dx \geq C \int_{\mathbb{R}_+^N} |\Delta u|^2 dx \text{ for all } u \in \tilde{H}_0(\mathbb{R}_+^N).$$

Therefore, with (35), there exists $C > 0$ such that $C \|u_m\|^2 \leq Q_{\gamma,s}(\mathbb{R}_+^N) + o(1)$ as $m \rightarrow +\infty$. Hence, $(u_m)_{m \in \mathbb{N}}$ is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$. As a consequence, up to the extraction of a subsequence, there exists $u \in \tilde{H}_0(\mathbb{R}_+^N)$ such that

$$\begin{cases} u_m \rightharpoonup u \text{ weakly in } \tilde{H}_0(\mathbb{R}_+^N), \\ u_m \rightarrow u \text{ strongly in } L_{loc}^q(\mathbb{R}^N) \text{ for any } 1 \leq q < 2_0^* := \frac{2N}{N-4}. \end{cases}$$

We define now the measures ν_m, λ_m on \mathbb{R}^N as

$$(38) \quad \nu_m := \frac{|u_m|^{2^*_s}}{|x|^s} \mathbf{1}_{\mathbb{R}_+^N} dx \text{ and } \lambda_m := \left(|\Delta u_m|^2 - \frac{\gamma}{|x|^4} u_m^2 \right) \mathbf{1}_{\mathbb{R}_+^N} dx.$$

Using (36) and (35) yield

$$(39) \quad \int_{\mathbb{R}^N} d\nu_m = 1 \text{ and } \lim_{m \rightarrow +\infty} \int_{\mathbb{R}^N} d\lambda_m = Q_{\gamma,s}(\mathbb{R}_+^N).$$

Up to extracting a subsequence, there exist two measures λ, ν on \mathbb{R}^N such that

$$(40) \quad \lambda_m \rightharpoonup \lambda \text{ and } \nu_m \rightharpoonup \nu \text{ weakly in the sense of measure as } m \rightarrow +\infty.$$

We now apply Lions's first concentration-compactness Lemma [16] to the sequence of measures $(\nu_m)_{m \in \mathbb{N}}$. Indeed, up to a subsequence, three situations may occur:

- (a) (Compactness) There exists a sequence $(x_m)_{m \in \mathbb{N}}$ in \mathbb{R}^N such that for any $\epsilon > 0$ there exists $R_\epsilon > 0$ with the property that

$$\int_{B_{R_\epsilon}(x_m)} d\nu_m \geq 1 - \epsilon \text{ for all } m \in \mathbb{N} \text{ large.}$$

- (b) (Vanishing) For all $R > 0$ there holds

$$\lim_{m \rightarrow +\infty} \left(\sup_{x \in \mathbb{R}^N} \int_{B_R(x)} d\nu_m \right) = 0.$$

- (c) (Dichotomy) There exists $\alpha \in (0, 1)$ such that for any $\epsilon > 0$ there exists $R_\epsilon > 0$ and a sequence $(x_m^\epsilon)_{m \in \mathbb{N}} \in \mathbb{R}^N$, with the following property: give $R' > R_\epsilon$, there are non-negative measures ν_m^1 and ν_m^2 such that

$$\begin{aligned} 0 \leq \nu_m^1 + \nu_m^2 \leq \nu_m, \text{ Supp}(\nu_m^1) \subset B_{R_\epsilon}(x_m^\epsilon), \text{ Supp}(\nu_m^2) \subset \mathbb{R}^N \setminus B_{R'}(x_m^\epsilon), \\ \nu_m^1 = \nu_m \big|_{B_{R_\epsilon}(x_m^\epsilon)}, \quad \nu_m^2 = \nu_m \big|_{\mathbb{R}^N \setminus B_{R'}(x_m^\epsilon)} \\ \lim_{m \rightarrow +\infty} \sup \left(\left| \alpha - \int_{\mathbb{R}^n} d\nu_m^1 \right| + \left| (1 - \alpha) - \int_{\mathbb{R}^n} d\nu_m^2 \right| \right) \leq \epsilon. \end{aligned}$$

Step 3.1. We claim that point (a) (Compactness) holds. In particular, we have that $\int_{\mathbb{R}^N} d\nu = 1$.

Proof of Step 3.1: Indeed, it follows from (36) that point (b), does not hold. Assume by contradiction that point (c) holds, that there exists $\alpha \in (0, 1)$ such that (c) above holds. Taking $\epsilon = (m + 1)^{-1}$, we can assume that, up to a subsequence, there exists $(R_m)_{m \in \mathbb{N}}$ in \mathbb{R}_+ , $(x_m)_{m \in \mathbb{N}}$ in \mathbb{R}^N and two sequence of non-negative measures, $(\nu_m^1)_{m \in \mathbb{N}}$ and $(\nu_m^2)_{m \in \mathbb{N}}$ such that $\lim_{m \rightarrow +\infty} R_m = \infty$ and

$$(41) \quad \begin{cases} 0 \leq \nu_m^1 + \nu_m^2 \leq \nu_m & , \quad \text{Supp}(\nu_m^1) \subset B_{R_m}(x_m), \text{ Supp}(\nu_m^2) \subset \mathbb{R}^N \setminus B_{2R_m}(x_m), \\ \nu_m^1 = \nu_m \big|_{B_{R_m}(x_m)} & , \quad \nu_m^2 = \nu_m \big|_{\mathbb{R}^N \setminus B_{2R_m}(x_m)}, \\ \lim_{m \rightarrow +\infty} \int_{\mathbb{R}^N} d\nu_m^1 = \alpha & , \quad \lim_{m \rightarrow +\infty} \int_{\mathbb{R}^N} d\nu_m^2 = 1 - \alpha. \end{cases}$$

We define $D_m := B_{2R_m}(x_m) \setminus B_{R_m}(x_m)$, it follows from (39) and (41) that

$$(42) \quad \lim_{m \rightarrow +\infty} \int_{D_m} d\nu_m = 0.$$

Taking $\eta \in C_c^\infty(\mathbb{R}^N)$ be such that $\eta|_{B_1(0)} \equiv 1$, $\eta|_{B_2(0)^c} \equiv 0$ and $0 \leq \eta \leq 1$. For $m \in \mathbb{N}$, we define $\eta_m(x) := \eta(R_m^{-1}(x - x_m))$ for all $x \in \mathbb{R}^N$. Using (41) and (42)

$$(43) \quad \begin{aligned} 1 &= \left(\int_{\mathbb{R}^N} \eta_m^{2_s^*} d\nu_m^1 + \int_{\mathbb{R}^N} (1 - \eta_m)^{2_s^*} d\nu_m^2 \right)^{\frac{2}{2_s^*}} + o(1) \\ &\leq \left(\int_{\mathbb{R}^N} \eta_m^{2_s^*} d\nu_m^1 \right)^{\frac{2}{2_s^*}} + \left(\int_{\mathbb{R}^N} (1 - \eta_m)^{2_s^*} d\nu_m^2 \right)^{\frac{2}{2_s^*}} + o(1). \end{aligned}$$

On the other hand, it follows from the (34) that

$$(44) \quad \begin{aligned} &\left(\int_{\mathbb{R}^N} \eta_m^{2_s^*} d\nu_m^1 \right)^{\frac{2}{2_s^*}} + \left(\int_{\mathbb{R}^N} (1 - \eta_m)^{2_s^*} d\nu_m^2 \right)^{\frac{2}{2_s^*}} + o(1) \\ &\leq \left(\int_{\mathbb{R}_+^N} \frac{|\eta_m u_m|^{2_s^*}}{|x|^s} dx \right)^{\frac{2}{2_s^*}} + \left(\int_{\mathbb{R}_+^N} \frac{|(1 - \eta_m) u_m|^{2_s^*}}{|x|^s} dx \right)^{\frac{2}{2_s^*}} + o(1) \\ &\leq Q_{\gamma, s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} \left(|\Delta(\eta_m u_m)|^2 - \gamma \frac{|\eta_m u_m|^2}{|x|^4} \right) dx \\ &+ Q_{\gamma, s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} \left(|\Delta((1 - \eta_m) u_m)|^2 - \gamma \frac{|(1 - \eta_m) u_m|^2}{|x|^4} \right) dx + o(1). \end{aligned}$$

Step 3.1.1: We claim that, as $m \rightarrow +\infty$,

$$(45) \quad \int_{\mathbb{R}_+^N} |\Delta(\eta_m u_m)|^2 dx = \int_{\mathbb{R}_+^N} |\eta_m|^2 |\Delta u_m|^2 dx + o(1),$$

$$(46) \quad \int_{\mathbb{R}_+^N} |\Delta((1 - \eta_m) u_m)|^2 dx = \int_{\mathbb{R}_+^N} |(1 - \eta_m)|^2 |\Delta u_m|^2 dx + o(1).$$

Proof of the claim: We write for convenience $A_{+,m} := \mathbb{R}_+^N \cap (B_{2R_m}(x_m) \setminus B_{R_m}(x_m))$.

Since $(u_m)_{m \in \mathbb{N}}$ is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$, similarly to (15) and (17), we get

$$(47) \quad \begin{aligned} \int_{\mathbb{R}_+^N} |\Delta(\eta_m u_m)|^2 dx &= \int_{\mathbb{R}_+^N} |\eta_m|^2 |\Delta u_m|^2 dx + O \left(\int_{A_{+,m}} \frac{|u_m|^2}{|x - x_m|^4} dx \right. \\ &\left. + \left(\int_{A_{+,m}} \frac{|u_m|^2}{|x - x_m|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+,m}} \frac{|u_m|^2}{|x - x_m|^4} dx \right)^{\frac{1}{4}} \right). \end{aligned}$$

We claim that,

$$(48) \quad \lim_{m \rightarrow +\infty} \int_{A_{+,m}} \frac{u_m^2}{|x - x_m|^4} dx = 0.$$

Proof. Indeed, by Hölder's inequality and since $N(1 - \frac{2}{2_s^*}) + \frac{2s}{2_s^*} = 4$, we get that

$$\begin{aligned}
\int_{A_{+,m}} \frac{u_m^2}{|x-x_m|^4} dx &\leq R_m^{-4} \left[\int_{A_{+,m}} dx \right]^{1-\frac{2}{2^*}} \left[\int_{A_{+,m}} u_m^{2^*} dx \right]^{\frac{2}{2^*}} \\
&\leq c w_{N-1}^{1-\frac{2}{2^*}} R_m^{\frac{2s}{2^*}-4} \left[\int_{R_m}^{2R_m} r^{N-1} dr \right]^{1-\frac{2}{2^*}} \left[\int_{A_{+,m}} \frac{u_m^{2^*}}{|x|^s} dx \right]^{\frac{2}{2^*}} \\
(49) \quad &\leq c w_{N-1}^{1-\frac{2}{2^*}} \left[\int_{A_{+,m}} \frac{u_m^{2^*}}{|x|^s} dx \right]^{\frac{2}{2^*}},
\end{aligned}$$

where w_{N-1} is the volume of the canonical $(N-1)$ -sphere. Using (36) yields

$$\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{|u_m|^{2^*}}{|x|^s} dx = 1 + \lim_{m \rightarrow +\infty} \int_{A_{+,m}} \frac{u_m^{2^*}}{|x|^s} dx.$$

In addition, by (41) we infer that $\lim_{m \rightarrow +\infty} \|u_m\|_{2^*,|x|^{-s}}^2 = 0$. Therefore, it follows from (49) that (48). This ends the proof of this claim. \square

Combining (47) and (48), we get the result. Similarly we prove (46). This ends the proof of Step 3.1.1. \square

It follows from (44), (45), (46) and (39) that

$$\begin{aligned}
&\left(\int_{\mathbb{R}^N} \eta_m^{2^*} d\nu_m^1 \right)^{\frac{2}{2^*}} + \left(\int_{\mathbb{R}^N} (1-\eta_m)^{2^*} d\nu_m^2 \right)^{\frac{2}{2^*}} \\
&\leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} \eta_m^2 \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx \\
&\quad + Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} (1-\eta_m)^2 \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx + o(1) \\
&\leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}^N} (\eta_m^2 + (1-\eta_m)^2) d\lambda_m + o(1) \\
&\leq 1 + 2Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} \eta_m(1-\eta_m) \frac{|u_m|^2}{|x|^4} dx + o(1) \\
&\leq 1 + O \left(\int_{A_{+,m}} \frac{|u_m|^2}{|x-x_m|^4} dx \right) + o(1).
\end{aligned}$$

Letting $m \rightarrow +\infty$ and using (41), (43) and (48) yields $\alpha^{\frac{2}{2^*}} + (1-\alpha)^{\frac{2}{2^*}} = 1$. This is impossible when $\alpha \in (0,1)$ and $2^* > 2$. This contradiction proves Step 3.1. \square

Step 3.2. *There exists $I \subset \mathbb{N}$ at most countable, and a family $\{x_i\}_{i \in I} \in \mathbb{R}^N$ such that*

$$(50) \quad \nu = \frac{|u|^{2^*}}{|x|^s} \mathbf{1}_{\mathbb{R}_+^N} dx + \sum_{i \in I} \nu_i \delta_{x_i},$$

where $\nu_i := \nu(x_i) > 0$ for all $i \in I$. In particular, $\{x_i, i \in I\} \subset \{0\}$ when $s > 0$. Moreover, there exists a bounded non-negative measure $\lambda_0 \geq 0$ with no atoms (that

is $\lambda_0(\{x\}) = 0$ for all $x \in \mathbb{R}^N$) and

$$(51) \quad \lambda = \lambda_0 + \left(|\Delta u|^2 - \gamma \frac{|u|^2}{|x|^4} \right) \mathbf{1}_{\mathbb{R}_+^N} dx + \sum_{i \in I} \lambda_i \delta_{x_i},$$

with $\lambda_i = \lambda(\{x_i\}) > 0$ and $\lambda_i \geq Q_{\gamma,s}(\mathbb{R}_+^N) \nu_i^{\frac{2}{2^*}}$.

Proof of Step 3.2: For $s = 0$, (50) is a consequence Lions's second concentration-compactness Lemma [16]. Take now $s > 0$ so that $2_s^* < \frac{2N}{N-4}$, then $u_m \rightarrow u$ strongly in $L_{loc}^{2_s^*}(\mathbb{R}_+^N)$. Therefore, we obtain that

$$(52) \quad \nu = \frac{|u|^{2_s^*}}{|x|^s} \mathbf{1}_{\mathbb{R}_+^N} dx + \nu(\{0\}) \delta_0.$$

This proves (50) in the case $s \geq 0$.

We now prove (51) of Step 3.2. We start by the following claim.

Step 3.2.1: We claim that

$$(53) \quad (\nu(\{x\}))^{\frac{2}{2_s^*}} \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \lambda(x) \text{ for all } x \in \mathbb{R}^N.$$

Proof of the claim: Indeed, $\varphi \in C^\infty(\mathbb{R}^n)$ be such that $\varphi(x) = 1$ for $x \in B_1(0)$, $\varphi(x) = 0$ for $x \in \mathbb{R}^N \setminus B_2(0)$ and $0 \leq \varphi \leq 1$. Given $y \in \mathbb{R}^N$ and $\delta > 0$, we define $\varphi_\delta(x) = \varphi(\frac{x-y}{\delta})$ for all $x \in \mathbb{R}^N$. Since $\varphi_\delta u_m \in \tilde{H}_0(\mathbb{R}_+^N)$, the definition (8) yields

$$(54) \quad \left(\int_{\mathbb{R}_+^N} \frac{|\varphi_\delta u_m|^{2_s^*}}{|x|^s} dx \right)^{\frac{2}{2_s^*}} \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}_+^N} \left(|\Delta(\varphi_\delta u_m)|^2 - \gamma \frac{|\varphi_\delta u_m|^2}{|x|^4} \right) dx.$$

As in the last proof of Step 3.1.1 (see (47)), we have that

$$\int_{\mathbb{R}_+^N} |\Delta(\varphi_\delta u_m)|^2 dx = \int_{\mathbb{R}_+^N} |\varphi_\delta|^2 |\Delta u_m|^2 dx + R_{m,\delta} + o(1),$$

where $o(1) \rightarrow 0$ as $m \rightarrow +\infty$ and,

$$R_{m,\delta} := O \left(\int_{A_{+,\delta}} \frac{u_m^2}{|x-y|^4} dx + \left(\int_{A_{+,\delta}} \frac{u_m^2}{|x-y|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+,\delta}} \frac{|u_m|^2}{|x-y|^4} dx \right)^{\frac{1}{4}} \right).$$

where $A_{+,\delta} := \mathbb{R}_+^N \cap (B_{2\delta}(y) \setminus B_\delta(y))$. Therefore, for all $\delta > 0$, using (54) yields

$$\left(\int_{\mathbb{R}^N} |\varphi_\delta|^{2_s^*} d\nu_m \right)^{\frac{2}{2_s^*}} \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}^N} \varphi_\delta^2 d\lambda_m + R_{m,\delta} + o(1),$$

letting $m \rightarrow +\infty$ and then $\delta \rightarrow 0$, we get that (53). \square

Up to extraction, let λ' be the weak limit of $|\Delta u_m|^2 \mathbf{1}_{\mathbb{R}_+^N} dx$ as $m \rightarrow +\infty$. Since $u_m \rightharpoonup u$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$, we get that $\lambda' \geq |\Delta u|^2 \mathbf{1}_{\mathbb{R}_+^N} dx$. Hence

$$(55) \quad \lambda' = \lambda_0 + |\Delta u|^2 \mathbf{1}_{\mathbb{R}_+^N} dx + \sum_{j \in J} \lambda'(\{z_j\}) \delta_{z_j} + \lambda'_0 \delta_0,$$

where $\lambda_0 \geq 0$ with no atoms, z_j 's, $j \in J$ countable, and are the atoms of λ' . As above (52), we have that there exists $L \geq 0$ such that

$$(56) \quad \frac{|u_m|^2}{|x|^4} \mathbf{1}_{\mathbb{R}_+^N} dx \rightharpoonup \frac{|u|^2}{|x|^4} \mathbf{1}_{\mathbb{R}_+^N} dx + L \delta_0,$$

It follows from (55) and (56) that

$$(57) \quad \lambda = \lambda_0 + \left(|\Delta u|^2 - \gamma \frac{|u|^2}{|x|^4} \right) \mathbf{1}_{\mathbb{R}_+^N} dx - \gamma L \delta_0 + \sum_{j \in J} \lambda'(\{z_j\}) \delta_{z_j}.$$

First, using (57) and (53) yields

$$(58) \quad 0 < (\nu(\{0\}))^{\frac{2}{2^*}} Q_{\gamma,s}(\mathbb{R}_+^N) \leq \lambda(\{0\}) = (\lambda'_0(\{0\}) - \gamma L).$$

On the other hand, for $x_j \neq 0$ we have

$$(59) \quad 0 < (\nu(\{x_j\}))^{\frac{2}{2^*}} Q_{\gamma,s}(\mathbb{R}_+^N) \leq \lambda(\{x_j\}) = \lambda'(\{x_j\}).$$

From (57), (58) and (59) we get the result (51). This proves Step 3.2. \square

Step 3.3. We claim that one and only one of the two following situations occur:

$$\begin{aligned} & \text{either } \left\{ \nu = \frac{|u|^{2^*}}{|x|^s} \mathbf{1}_{\mathbb{R}_+^N} dx \text{ and } \int_{\mathbb{R}_+^N} \frac{|u|^{2^*}}{|x|^s} dx = 1 \right\} \\ & \text{or } \left\{ \text{there exists } x_0 \in \mathbb{R}^N \text{ such that } \nu = \delta_{x_0} \text{ and } u = 0 \right\}. \end{aligned}$$

Proof of Step 3.3: Indeed, it follows from Step 3.1 that,

$$(60) \quad \begin{aligned} 1 &= \left(\int_{\mathbb{R}_+^N} d\nu \right)^{\frac{2}{2^*}} = \left(\int_{\mathbb{R}_+^N} \frac{|u|^{2^*}}{|x|^s} dx + \sum_{i \in I} \nu^i \int_{\mathbb{R}^N} \delta_{x_i} dx \right)^{\frac{2}{2^*}} \text{ from (50)} \\ &= \left(\int_{\mathbb{R}_+^N} \frac{|u|^{2^*}}{|x|^s} dx + \sum_{i \in I} \nu^i \right)^{\frac{2}{2^*}} \leq \|u\|_{2^*,|x|^{-s}}^2 + \sum_{i \in I} \nu_i^{\frac{2}{2^*}}. \end{aligned}$$

Now, using again (34) and (53) yields,

$$(61) \quad \|u\|_{2^*,|x|^{-s}}^2 + \sum_{i \in I} \nu_i^{\frac{2}{2^*}} \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \left(\int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx + \sum_{i \in I} \lambda^i \right) \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \int_{\mathbb{R}^N} d\lambda$$

from (51). Combining (60) and (61), we have that $\int_{\mathbb{R}^N} d\lambda \geq Q_{\gamma,s}(\mathbb{R}_+^N)$.

We claim now that $\int_{\mathbb{R}^N} d\lambda \leq Q_{\gamma,s}(\mathbb{R}_+^N)$. Indeed, we let $f \in C^\infty(\mathbb{R}^N)$ be such that $f(x) = 0$ for $x \in B_1(0)$, $f(x) = 1$ for $x \in \mathbb{R}^N \setminus B_2(0)$ and $0 \leq f \leq 1$. Given $\rho > 0$, we let $f_\rho(x) = f(\rho^{-1}x)$ for all $x \in \mathbb{R}^N$. So $(1 - f_\rho^2)u_m \in \tilde{H}_0(\mathbb{R}_+^N)$ and therefore

$$\begin{aligned} \int_{\mathbb{R}^N} (1 - f_\rho^2) d\lambda_m &= \int_{\mathbb{R}^N} d\lambda_m - \int_{\mathbb{R}_+^N} \left(f_\rho^2 |\Delta u_m|^2 - \gamma \frac{|f_\rho u_m|^2}{|x|^4} \right) dx \\ &= \int_{\mathbb{R}^N} d\lambda_m - \int_{\mathbb{R}_+^N} \left(|\Delta(f_\rho u_m)|^2 - \gamma \frac{|f_\rho u_m|^2}{|x|^4} \right) dx \\ &\quad + \int_{\mathbb{R}_+^N} (|\Delta(f_\rho u_m)|^2 - f_\rho^2 |\Delta u_m|^2) dx \\ &\leq \int_{\mathbb{R}^N} d\lambda_m + \int_{\mathbb{R}_+^N} (|\Delta(f_\rho u_m)|^2 - f_\rho^2 |\Delta u_m|^2) dx \text{ from (37)} \\ &\leq Q_{\gamma,s}(\mathbb{R}_+^N) + R_{m,\rho} + o(1) \text{ from (47),} \end{aligned}$$

where $o(1) \rightarrow 0$ as $m \rightarrow +\infty$, and

$$R_{m,\rho} := O \left(\int_{A_{+,\rho}} \frac{u_m^2}{|x-y|^4} dx + \left(\int_{A_{+,\rho}} \frac{u_m^2}{|x-y|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+,\rho}} \frac{|u_m|^2}{|x-y|^4} dx \right)^{\frac{1}{4}} \right).$$

where $A_{+,\rho} := \mathbb{R}_+^N \cap (B_{2\rho}(0) \setminus B_\rho(0))$. Therefore, letting $m \rightarrow +\infty$, and then $\rho \rightarrow +\infty$, and we then get this claim. This implies that $\int_{\mathbb{R}^N} d\lambda = Q_{\gamma,s}(\mathbb{R}_+^N)$. Therefore, it follows from (60) and (61) that $\|u\|_{2_s^*,|x|^{-s}}^2 + \sum_{i \in I} \nu_i^{\frac{2}{2_s^*}} = 1$. By convexity, we have that one and only one term in (50) is nonzero, then there exist $i_0 \in I$ such that $x_0 := x_{i_0}$ and

$$\left\{ \nu^{i_0} = 1 \text{ and } \int_{\mathbb{R}_+^N} \frac{|u|^{2_s^*}}{|x|^s} dx = 0 \right\} \text{ or } \left\{ \nu^{i_0} = 0 \text{ and } \int_{\mathbb{R}_+^N} \frac{|u|^{2_s^*}}{|x|^s} dx = 1 \right\},$$

with the equation (50), the exist $x_0 \in \mathbb{R}^N$ such that we get the claim of Step 3.3. \square

Step 3.4. Suppose that $u \not\equiv 0$. We claim that that u is an extremal for $Q_{\gamma,s}(\mathbb{R}_+^N)$.

Proof of Step 3.4: Since $u \not\equiv 0$, it follows from the previous Step that we have $\|u\|_{2_s^*,|x|^{-s}}^2 = 1$. Using again the Hardy-Sobolev inequality (34) yields,

$$Q_{\gamma,s}(\mathbb{R}_+^N) \leq \int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{u}{|x|^4} \right) dx.$$

On the other hand, we have $u_m \rightharpoonup u$ as $m \rightarrow +\infty$ and we get that

$$\int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx \leq \liminf_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 - \gamma \frac{u_m^2}{|x|^4} \right) dx = Q_{\gamma,s}(\mathbb{R}_+^N).$$

Therefore, we get the equality $I_{\gamma,s}^{\mathbb{R}_+^N}(u) = Q_{\gamma,s}(\mathbb{R}_+^N)$. That is u is an extremal for $Q_{\gamma,s}(\mathbb{R}_+^N)$. We obtain the result of Step 3.4. \square

Step 3.5. We suppose that $u \equiv 0$. Then, we have

$$s = 0, \lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx = 0 \text{ and } |\Delta u_m|^2 dx \rightharpoonup Q_{\gamma,0}(\mathbb{R}_+^N) \delta_{x_0},$$

as $m \rightarrow +\infty$ in the sense of measures.

Proof of step 3.5: Since $u \equiv 0$, and it follows from the Step 3.3 that there exists $x_0 \in \mathbb{R}^N$ such that $\nu = \delta_{x_0}$.

We claim that $x_0 \neq 0$. Indeed, if $x_0 = 0$, we get that $\int_{B_{1/2}(0)} d\nu = 1$ which contradicts (36). Therefore $x_0 \neq 0$. Since $u_m \rightharpoonup 0$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$, then for any $1 \leq q < \frac{2N}{N-4}$, we have $u_m \rightarrow 0$ strongly in $L_{loc}^q(\mathbb{R}_+^N)$.

We claim that $s = 0$. Indeed, we argue by contradiction and assume that $s > 0$, then $2_s^* < \frac{2N}{N-4}$. Let $r > 0$, since $x_0 \neq 0$ and $u_m \rightarrow 0$ strongly in $L_{loc}^{2_s^*}(\mathbb{R}_+^N)$. Hence, we have $\lim_{m \rightarrow +\infty} \int_{B_r(x_0) \cap \mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx = 0$, and, it follows from (40) and $\nu = \delta_{x_0}$ that $\lim_{m \rightarrow +\infty} \int_{B_r(x_0) \cap \mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx = 1$, for all $r > 0$ enough, a contradiction to our assumption.

Therefore $s = 0$, and we prove the rest of this Step. Let $\rho > 0$ and $f \in C^\infty(\mathbb{R}^N)$ be such that $f(x) = 0$ for $x \in B_\rho(x_0)$, $f(x) = 1$ for $x \in \mathbb{R}^N \setminus B_{2\rho}(x_0)$ and $0 \leq f \leq 1$. We now define, $\varphi := 1 - f^2$ and $\psi := f\sqrt{2 - f^2}$. Clearly $\varphi, \psi \in C^\infty(\mathbb{R}^N)$ and $\varphi^2 + \psi^2 = 1$. It follows from (34) and (47) that

$$\begin{aligned} Q_{\gamma,0}(\mathbb{R}_+^N) \left(\int_{\mathbb{R}_+^N} |\varphi u_m|^{2_0^{**}} dx \right)^{\frac{2}{2_0^{**}}} &\leq \int_{\mathbb{R}_+^N} \left(|\Delta(\varphi u_m)|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|\varphi u_m|^2}{|x|^4} dx \right) dx \\ &\leq \int_{\mathbb{R}_+^N} \varphi^2 \left(|\Delta u_m|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx \right) dx + R_{m,\rho} + o(1), \end{aligned}$$

where $o(1) \rightarrow 0$ as $m \rightarrow +\infty$, and

$$R_{m,\rho} := O \left(\int_{A_{+,\rho}} \frac{u_m^2}{|x-y|^4} dx + \left(\int_{A_{+,\rho}} \frac{u_m^2}{|x-y|^4} dx \right)^{\frac{1}{2}} + \left(\int_{A_{+,\rho}} \frac{|u_m|^2}{|x-y|^4} dx \right)^{\frac{1}{4}} \right).$$

where $A_{+,\rho} := \mathbb{R}_+^N \cap (B_{2\rho}(x_0) \setminus B_\rho(x_0))$. Using $u_m \rightarrow 0$ in $L_{loc}^2(\mathbb{R}^N)$ yields $R_{m,\rho} = o(1)$ as $m \rightarrow +\infty$. And, so by $\varphi^2 = 1 - \psi^2$

$$\begin{aligned} Q_{\gamma,0}(\mathbb{R}_+^N) \left(\int_{\mathbb{R}_+^N} |\varphi u_m|^{2_0^{**}} dx \right)^{\frac{2}{2_0^{**}}} &\leq \int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx \right) dx \\ (62) \quad &- \int_{\mathbb{R}_+^N} \psi^2 \left(|\Delta u_m|^2 dx - \gamma \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx \right) dx + o(1), \end{aligned}$$

as $m \rightarrow +\infty$. It follows from (40) and Step 3.3 that

$$(63) \quad \left(\int_{\mathbb{R}_+^N} |\varphi u_m|^{2_0^{**}} dx \right)^{\frac{2}{2_0^{**}}} = \left(|\varphi(x_0)|^{2_0^*} + o(1) \right)^{\frac{2}{2_0^{**}}} = 1 + o(1) \text{ as } m \rightarrow +\infty.$$

Plugging (63) into (62) we get that

$$\begin{aligned} Q_{\gamma,0}(\mathbb{R}_+^N) + o(1) &\leq \int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx \\ (64) \quad &- \int_{\mathbb{R}_+^N} \psi^2 \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx + o(1), \end{aligned}$$

as $m \rightarrow +\infty$. From (35), we have

$$(65) \quad \int_{\mathbb{R}_+^N} \psi^2 \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx \leq o(1) \text{ as } m \rightarrow +\infty.$$

As in the proof of (45), we obtain that $\int_{\mathbb{R}_+^N} |\Delta(\psi u_m)|^2 dx = \int_{\mathbb{R}_+^N} \psi^2 |\Delta u_m|^2 dx + o(1)$ as $m \rightarrow +\infty$. Plugging this expansion into (65) yields

$$\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} (|\Delta(\psi u_m)|^2 - \gamma |x|^{-4} |\psi u_m|^2) dx = 0.$$

Hence, by the coercivity (37), we get

$$(66) \quad \lim_{m \rightarrow +\infty} \|\Delta(\psi u_m)\|_2 = 0.$$

With the result of the Lemma 2.1, we have that

$$\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{|\psi u_m|^2}{|x|^4} dx = 0.$$

We then have $\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N \setminus B_{2\rho}(x_0)} \frac{|u_m|^2}{|x|^4} dx = 0$. Moreover, taking $\rho > 0$ small enough, since $x_0 \neq 0$ and $u_m \rightarrow 0$ in $L^2_{loc}(\mathbb{R}_+^N)$ around $x_0 \neq 0$ yields

$$\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{|u_m|^2}{|x|^4} dx = 0,$$

which implies by (35) that $\lim_{m \rightarrow +\infty} \|\Delta u_m\|^2 = Q_{\gamma,0}(\mathbb{R}_+^N)$. Hence, using (66) yields the third part of claim. Step 3.5 is proved. \square

Step 3.6. We now claim that, if $u \equiv 0$, then $s = 0$ and $Q_{\gamma,s}(\mathbb{R}_+^N) = S_N$, where S_N is defined in (10).

Proof of step 3.6: We have already seen that $s = 0$. Since $u_m \in \tilde{H}_0(\mathbb{R}_+^N) \subset \tilde{H}_0(\mathbb{R}^N)$, we have that

$$\begin{aligned} S_N \left(\int_{\mathbb{R}^N} |u_m|^{2^{**}(0)} dx \right) &\leq \int_{\mathbb{R}^N} |\Delta u_m|^2 dx \\ &\leq \int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 - \gamma \frac{u_m^2}{|x|^4} \right) dx \\ &\quad + \int_{\mathbb{R}^N \setminus \mathbb{R}_+^N} |\Delta u_m|^2 dx + \gamma \int_{\mathbb{R}_+^N} \frac{u_m^2}{|x|^4} dx, \end{aligned}$$

so with (35) and by the result of Step 3.5, we get

$$S_N \left(\int_{\mathbb{R}^N} |u_m|^{2^{**}(0)} dx \right) \leq Q_{\gamma,0}(\mathbb{R}_+^N) + o(1).$$

It follows then from (36) that $S_N \leq Q_{\gamma,0}(\mathbb{R}_+^N) + o(1)$. Letting $m \rightarrow +\infty$ we obtain that $S_N \leq Q_{\gamma,0}(\mathbb{R}_+^N)$. Conversely, it follows from the computations of Proposition 5.1 below that $Q_{\gamma,0}(\mathbb{R}_+^N) \leq S_N$. Hence, we have $Q_{\gamma,0}(\mathbb{R}_+^N) = S_N$. This proves Step 3.6 \square

Step 3.7. We assume that $s = 0$ and $\gamma \leq 0$. Then, we have that $Q_{\gamma,0}(\mathbb{R}_+^N) = S_N$, where S_N is defined in (10).

Proof of step 3.7: Indeed, since $\gamma \leq 0$, it follows from the Proposition 5.1 when $\Omega = \mathbb{R}_+^N$ is smooth domain that $Q_{\gamma,0}(\mathbb{R}_+^N) = S_N$. \square

Step 3.8. Taking $\{s = 0, \gamma > 0 \text{ and } N \geq 8\}$, we claim that

$$(67) \quad Q_{\gamma,0}(\mathbb{R}_+^N) < S_N.$$

Therefore, we get that $Q_{\gamma,0}(\mathbb{R}_+^N)$ is attained.

Proof of step 3.8: We fix $x_0 \in \mathbb{R}_+^N$ such that $x_0 \neq 0$. We define $U(x) := (1 + |x|^2)^{-\frac{N-4}{2}}$ for all $x \in \mathbb{R}^N$. It follows from Lions [15, 16] that $U \in \tilde{H}_0(\mathbb{R}^N) = D^{2,2}(\mathbb{R}^N)$ is an extremal for (10), that is

$$S_N := \frac{\int_{\mathbb{R}^N} |\Delta U|^2 dx}{\left(\int_{\mathbb{R}^N} |U|^{2^*_0} dx \right)^{\frac{2}{2^*_0}}}.$$

Let $\eta \in C_c^\infty(\mathbb{R}_+^N)$ and $0 < \delta < |x_0|/2$ be such that $\eta(x) = 1$ for $x \in B_\delta(x_0)$. We consider the test function

$$(68) \quad U_\epsilon(x) := \eta(x)u_\epsilon(x) \text{ for all } x \in \mathbb{R}_+^N \text{ and } \epsilon > 0,$$

$$(69) \quad \text{where } u_\epsilon(x) := \epsilon^{-\frac{N-4}{2}} U\left(\frac{x-x_0}{\epsilon}\right) = \left(\frac{\epsilon}{\epsilon^2 + |x-x_0|^2}\right)^{\frac{N-4}{2}} \text{ for all } x \in \mathbb{R}^N.$$

Step 3.8.1: We claim as $\epsilon \rightarrow 0$ that:

$$(70) \quad \int_{\mathbb{R}_+^N} \frac{U_\epsilon^2}{|x|^4} dx = \begin{cases} \frac{\epsilon^4}{2} |x_0|^{-4} \int_{\mathbb{R}^N} U^2 dx + o(\epsilon^4) & \text{if } N \geq 9, \\ \epsilon^4 \ln\left(\frac{1}{\epsilon}\right) |x_0|^{-4} + o(\epsilon^4 \ln\left(\frac{1}{\epsilon}\right)) & \text{if } N = 8, \\ O(\epsilon^{N-4}) & \text{if } N = 5, 6, 7. \end{cases}$$

Proof of the claim: Indeed, for $\delta > 0$ we begin by noticing that

$$(71) \quad \int_{\mathbb{R}_+^N} \frac{U_\epsilon^2}{|x|^4} dx = |x_0|^{-4} I_{1,\epsilon} + I_{2,\epsilon} + O(\epsilon^{N-4}) \text{ as } \epsilon \rightarrow 0,$$

where:

$$(72) \quad I_{1,\epsilon} := \int_{B_\delta(x_0) \cap \mathbb{R}_+^N} u_\epsilon^2 dx \text{ and } I_{2,\epsilon} := \int_{B_\delta(x_0) \cap \mathbb{R}_+^N} \left(\frac{1}{|x|^4} - \frac{1}{|x_0|^4}\right) u_\epsilon^2 dx.$$

First, we claim that

$$(73) \quad I_{1,\epsilon} = \begin{cases} \frac{\epsilon^4}{2} \int_{\mathbb{R}^N} U^2 dx + O(\epsilon^{N-4}) & \text{if } N \geq 9, \\ \epsilon^4 \ln\left(\frac{1}{\epsilon}\right) + O(\epsilon^4) & \text{if } N = 8, \\ O(\epsilon^{N-4}) & \text{if } N = 5, 6, 7, \end{cases}.$$

Proof. For $N \geq 9$, $U \in L^2(\mathbb{R}^N)$, and we get as $\epsilon \rightarrow 0$ that

$$I_{1,\epsilon} = \frac{\epsilon^4}{2} \int_{\mathbb{R}^N} U^2 dx + O(\epsilon^{N-4}).$$

Take now the case $N = 8$. It follows from the change of variable to polar coordinates that

$$\begin{aligned} I_{1,\epsilon} &= \epsilon^4 w_7 \left[O(1) + \int_1^{\delta\epsilon^{-1}} \frac{1}{r} dr + \int_1^{\delta\epsilon^{-1}} r^7 \left[\frac{1}{(1+r^2)^4} - \frac{1}{r^8} \right] dr \right] \\ &= \epsilon^4 w_7 \ln\left(\frac{\delta}{\epsilon}\right) + O(\epsilon^4). \end{aligned}$$

As one checks, we have that $I_{1,\epsilon} = O(\epsilon^{N-4})$ for $N = 5, 6, 8$. This proves (73). \square

Next, we claim as $\epsilon \rightarrow 0$ that

$$(74) \quad I_{2,\epsilon} = \begin{cases} o(\epsilon^4) & \text{if } N \geq 9, \\ o(\epsilon^4 \ln\left(\frac{1}{\epsilon}\right)) & \text{if } N = 8, \\ O(\epsilon^{N-4}) & \text{if } N = 5, 6, 7. \end{cases}$$

Proof. It follows from the definition of u_ϵ that

$$|I_{2,\epsilon}| \leq \epsilon^{N-4} \int_{B_\delta(0) \cap \mathbb{R}_+^N} \frac{|A(x)|}{(\epsilon^2 + |x|^2)^{N-4}} dx,$$

where $A(x) := \frac{1}{|x+x_0|^4} - \frac{1}{|x_0|^4}$. Fix $\alpha \in (0, \delta)$, with a change of variables we write

$$\epsilon^{N-4} \int_{(B_\delta(0) \setminus B_\alpha(0)) \cap \mathbb{R}_+^N} \frac{|A(x)|}{(\epsilon^2 + |x|^2)^{N-4}} dx \leq \epsilon^{N-4} \|A\|_\infty \frac{\delta^N w_{N-1}}{\alpha^{2(N-4)}} := C_{1,\alpha} \epsilon^{N-4}.$$

On the other hand, we have that

$$(75) \quad \epsilon^{N-4} \int_{B_\alpha(0) \cap \mathbb{R}_+^N} \frac{|A(x)|}{(\epsilon^2 + |x|^2)^{N-4}} dx \leq \epsilon^4 \left(\sup_{|x| \leq \alpha} |A(x)| \right) w_{N-1} \int_0^{\alpha \epsilon^{-1}} \frac{r^{N-1}}{(1+r^2)^{N-4}} dr.$$

From here, we distinguish our proof to three cases:

Case 1: If $N \geq 9$, we have $U \in L^2(\mathbb{R}^N)$, and then there exists $C_{2,\alpha} > 0$ such that

$$\epsilon^{N-4} \int_{B_\alpha(0) \cap \mathbb{R}_+^N} \frac{|A(x)|}{(\epsilon^2 + |x|^2)^{N-4}} dx \leq C \epsilon^4 \left(\sup_{|x| \leq \alpha} |A(x)| \right).$$

Take now $\theta > 0$. Then there exists $\alpha_0 > 0$ such that for all $\alpha < \alpha_0$ we have that $C \left(\sup_{|x| \leq \alpha} |A(x)| \right) \leq \frac{\theta}{2}$. On the other hand, we have $\lim_{\epsilon \rightarrow 0} C_{1,\alpha} \epsilon^{N-8} = 0$, then there exists $\epsilon_0 := \epsilon(\theta) > 0$ such that $C_{1,\alpha} \epsilon^{N-8} \leq \frac{\theta}{2}$. Then, we have $I_{2,\epsilon} = o(\epsilon^4)$ as $\epsilon \rightarrow 0$.

Case 2: The proof of case $N = 8$ is similar to the proof of (73). \square

Plugging (73) and (74) into (71), we obtain that (70). This proves Step 3.8.1. \square

For $N \geq 5$, it is also classical as $\epsilon \rightarrow 0$ that:

$$(76) \quad \int_{\mathbb{R}_+^N} |\Delta U_\epsilon(x)|^2 dx = \int_{\mathbb{R}^N} |\Delta U|^2 dx + O(\epsilon^{N-4}),$$

$$(77) \quad \int_{\mathbb{R}_+^N} |U_\epsilon|^{2^*_0} dx = \int_{\mathbb{R}^N} |U|^{2^*_0} dx + O(\epsilon^N).$$

Combining (74), (76) and (77), we have that

$$I_{\gamma,0}^{\mathbb{R}_+^N}(U_\epsilon) = \begin{cases} S_N - \gamma |x_0|^{-4} c \epsilon^4 + o(\epsilon^{N-4}) & \text{if } N \geq 9, \\ S_N - \gamma |x_0|^{-4} c \epsilon^4 \ln\left(\frac{1}{\epsilon}\right) + o(\epsilon^4 \ln\left(\frac{1}{\epsilon}\right)) & \text{if } N = 8, \end{cases}$$

where c is a positive constant. Since $\gamma > 0$ and $x_0 \neq 0$, then $Q_{\gamma,0}(\mathbb{R}_+^N) < S_N$. Therefore, it follows from the Step 1.2 that $u \neq 0$, and we have u is a extremal for $Q_{\gamma,0}(\mathbb{R}_+^N)$. This ends Step 3.8. \square

All these cases end the proof of Theorem 1.2.

4. PROOF OF THEOREM 1.1

In this section, we use the existence of extremals for Hardy-Sobolev inequality, established in Section 3 to prove that there exists a nontrivial weak solution for double critical equation (1).

For any function $G \in C^1(E, \mathbb{R})$ where $(E, \|\cdot\|)$ is a Banach space, we say that $(u_m)_{m \in \mathbb{N}} \in E$ is a Palais-Smale sequence of G if there exists $\beta \in \mathbb{R}$ such that

$$G(u_m) \rightarrow \beta \text{ and } G'(u_m) \rightarrow 0 \text{ in } E' \text{ as } m \rightarrow +\infty.$$

Here, we say that the Palais-Smale sequence is at level β . The main tool is the Mountain-Pass Lemma of Ambrosetti-Rabinowitz [1]:

Theorem 4.1 (Mountain-Pass Lemma [1]). *We consider $G \in C^1(X, \mathbb{R})$ where $(X, \|\cdot\|)$ is a Banach space. We assume that $G(0) = 0$ and that*

- *There exists $\lambda, r > 0$ such that $G(u) \geq \lambda$ for all $u \in X$ such that $\|u\| = r$,*
- *There exists u_0 in X such that $\limsup_{t \rightarrow +\infty} G(tu_0) < 0$.*

We consider $t_0 > 0$ sufficiently large such that $\|t_0 u_0\| > r$ and $G(t_0 u_0) < 0$, and

$$\beta = \inf_{c \in \Gamma} \sup_{t \in [0, 1]} G(c(t)),$$

where $\Gamma := \{c \in C^0([0, 1], X) \text{ s.t. } c(0) = 0, c(1) = t_0 u_0\}$.

Then, there exists a Palais-Smale sequence at level β for G . Moreover, we have that $\beta \leq \sup_{t \geq 0} G(tu_0)$.

We define the energy functional noted by E

$$E(u) := \frac{1}{2} \int_{\mathbb{R}_+^N} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx - \frac{1}{2_s^*} \int_{\mathbb{R}_+^N} \frac{|u|^{2_s^*}}{|x|^s} dx - \frac{1}{2_0^*} \int_{\mathbb{R}_+^N} |u|^{2_0^*} dx,$$

for any $u \in \tilde{H}_0(\mathbb{R}_+^N)$. Any weak solution to (1) is a critical point of E . In the sequel, since $\gamma < \gamma_{H,+}$, then (37) holds.

Proposition 4.1. *We assume that (37) holds. Fix $u_0 \in \tilde{H}_0(\mathbb{R}_+^N)$ such that $u_0 \geq 0$, $u_0 \not\equiv 0$. Then there exists a sequence $(u_m)_{m \in \mathbb{N}} \in \tilde{H}_0(\mathbb{R}_+^N)$ that is a Palais-Smale sequence for E at level β such that $0 < \beta \leq \sup_{t \geq 0} E(tu_0)$.*

Proof of Proposition 4.1: Indeed, clearly $E \in C^1(\tilde{H}_0(\mathbb{R}_+^N))$. Note that $E(0) = 0$. It follows from (37) and the Sobolev and Hardy-Sobolev embeddings that there exist $c_0, c_1, c_2 > 0$ such that

$$(78) \quad E(u) \geq c_0 \|u\|^2 - c_1 \|u\|^{2_s^*} - c_2 \|u\|^{2_0^*} \text{ for all } u \in \tilde{H}_0(\mathbb{R}_+^N).$$

Define $f(r) = r^2 [c_0 - c_1 r^{2_s^* - 2} - c_2 r^{2_0^* - 2}] := r^2 g(r)$ and since $2_s^*, 2_0^* > 2$ we have $g(r) \rightarrow c_0$ as $r \rightarrow 0$. Then there exists $r_0 > 0$ such that $r < r_0$, we have $g(r) > \frac{c_0}{2}$. Therefore, for all $u \in \tilde{H}_0(\mathbb{R}_+^N)$ such that $\|u\| = \frac{r_0}{2}$ and by (78), we have $E(u) \geq \frac{c_0 r_0^2}{8} := \lambda$. We fix $u_0 \in \tilde{H}_0(\mathbb{R}_+^N)$, $u_0 \not\equiv 0$, and

$$\begin{aligned} E(tu_0) &= \frac{t^2}{2} \int_{\mathbb{R}_+^N} \left(|\Delta u_0|^2 - \frac{\gamma}{|x|^4} u_0^2 \right) dx \\ &\quad - \frac{t^{2_s^*}}{2_s^*} \int_{\mathbb{R}_+^N} \frac{|u_0|^{2_s^*}}{|x|^s} dx - \frac{t^{2_0^*}}{2_0^*} \int_{\mathbb{R}_+^N} |u_0|^{2_0^*} dx \\ &:= \frac{t^2}{2} R_1 - \frac{t^{2_s^*}}{2_s^*} R_2 - \frac{t^{2_0^*}}{2_0^*} R_3 \leq t^{2_s^*} \left(\frac{t^{2-2_s^*}}{2} R_1 - R_2 \right), \end{aligned}$$

where $R_1, R_2 > 0$ and $R_3 \geq 0$. Since $2_s^* > 2$, we have $E(tu_0) \rightarrow -\infty$ as $t \rightarrow +\infty$. Then $\limsup_{t \rightarrow +\infty} E(tu_0) < 0$. We consider $t_0 > 0$ large such that $\|t_0 u_0\| > r$ and $E(t_0 u_0) < 0$. For $t \in [0, 1]$, we have $E(c(t)) \geq \lambda$ and then there exists

$$\beta := \inf_{c \in \Gamma} \sup E(c(t)) \geq \lambda > 0.$$

We apply Theorem 4.1 to get the expected Palais-Smale sequence. This ends the proof of Proposition 4.1. \square

Proposition 4.2. *Suppose $0 < \gamma < \gamma_{H,+}$, $0 \leq s < 4$ and $N \geq 8$. We assume that (37) holds. Then there exists a sequence $(u_m)_{m \in \mathbb{N}} \in \tilde{H}_0(\mathbb{R}_+^N)$ that is a Palais-Smale sequence for E at level β such that*

$$(79) \quad 0 < \beta < \beta^* := \min \left\{ \frac{2}{N} Q_{\gamma,0}(\mathbb{R}_+^N)^{\frac{N}{4}}, \frac{4-s}{2(N-s)} Q_{\gamma,s}(\mathbb{R}_+^N)^{\frac{N-s}{4-s}} \right\}.$$

Proof of Proposition 4.1: From Theorem 1.2, we know that there exists an extremal $u_0 \in \tilde{H}_0(\mathbb{R}_+^N)$ for $Q_{\gamma,0}(\mathbb{R}_+^N)$ whenever $\gamma > 0$ and $N \geq 8$. It follows then from Proposition 4.1 that there exists a sequence $u_m \in \tilde{H}_0(\mathbb{R}_+^N)$ a Palais-Smale sequence for E at level β such that

$$\beta \leq \sup_{t \geq 0} E(tu_0) \leq \sup_{t \geq 0} f_1(t),$$

where:

$$f_1(t) := \frac{t^2}{2} \int_{\mathbb{R}_+^N} \left(|\Delta u_0|^2 - \frac{\gamma}{|x|^4} u_0^2 \right) dx - \frac{t^{2_0^*}}{2_0^*} \int_{\mathbb{R}_+^N} |u_0|^{2_0^*} dx \text{ for all } t > 0.$$

Simple computations yield that $f_1(t)$ attains its maximum at the point

$$t_{max} = \left(\frac{\int_{\mathbb{R}_+^N} \left(|\Delta u_0|^2 - \frac{\gamma}{|x|^4} u_0^2 \right) dx}{\int_{\mathbb{R}_+^N} |u_0|^{2_0^*} dx} \right)^{\frac{1}{2_0^*-2}}.$$

Therefore, using $2_0^* = \frac{2N}{N-4}$ and u_0 is an extremal for $Q_{\gamma,0}(\mathbb{R}_+^N)$ yields,

$$\sup_{t \geq 0} f_1(t) = \left[\frac{1}{2} - \frac{1}{2_0^*} \right] \left(\frac{\int_{\mathbb{R}_+^N} \left(|\Delta u_0|^2 - \frac{\gamma}{|x|^4} u_0^2 \right) dx}{\left(\int_{\mathbb{R}_+^N} |u_0|^{2_0^*} dx \right)^{\frac{2}{2_0^*}}} \right)^{\frac{2_0^*}{2_0^*-2}} = \frac{2}{N} Q_{\gamma,0}(\mathbb{R}_+^N)^{\frac{N}{4}}.$$

Thus, $\beta \leq \sup_{t \geq 0} f_1(t) = \frac{2}{N} Q_{\gamma,0}(\mathbb{R}_+^N)^{\frac{N}{4}}$. We now prove that this inequality is strict. Assume by contradiction that

$$\sup_{t \geq 0} E(tu_0) = \sup_{t \geq 0} f_1(t),$$

and we consider $t_1 > 0$ (resp. $t_2 > 0$) where $\sup_{t \geq 0} E(tu_0)$ (resp. $\sup_{t \geq 0} f_1(t)$) is attained. We obtain that

$$f_1(t_1) - \frac{t_1^{2_s^*}}{2_s^*} \int_{\mathbb{R}_+^N} \frac{|u_0|^{2_s^*}}{|x|^s} dx = f_1(t_2),$$

this give us $f_1(t_1) - f_1(t_2) > 0$ (because $t_1 > 0$). Contradiction with t_2 is a maximum point of $f_1(t)$. Therefore, we have $\beta < \frac{2}{N} Q_{\gamma,0}(\mathbb{R}_+^N)^{\frac{N}{4}}$. Similar, we can get $\beta < \frac{4-s}{2(N-s)} Q_{\gamma,s}(\mathbb{R}_+^N)^{\frac{N-s}{4-s}}$ whenever $s > 0$. Thus, we can define β^* as in (79) such that $0 < \beta < \beta^*$. This proves Proposition 4.1. \square

Proposition 4.3. *Suppose $0 < \gamma < \gamma_{H,+}$, $0 \leq s < 4$ and $N \geq 8$. We assume that $(u_m)_{m \in \mathbb{N}}$ is a Palais-Smale sequence of G at energy level $\beta \in (0, \beta^*)$. If $u_m \rightharpoonup 0$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$, then there exists $\epsilon := \epsilon(N, \gamma, s, \beta) > 0$ such that for every $r > 0$ such that*

$$\text{either } \lim_{m \rightarrow +\infty} \sup \int_{B_r(0)} |u_m|^{2_0^*} dx = \lim_{m \rightarrow +\infty} \sup \int_{B_r(0)} \frac{|u_m|^{2_s^*}}{|x|^s} dx = 0;$$

$$\text{or } \lim_{m \rightarrow +\infty} \sup \int_{B_r(0)} |u_m|^{2_0^*} dx \geq \epsilon \text{ and } \lim_{m \rightarrow +\infty} \sup \int_{B_r(0)} \frac{|u_m|^{2_s^*}}{|x|^s} dx \geq \epsilon.$$

Proof of Proposition 4.3: Indeed, the proof of this proposition is divided into several steps :

Step 4.1. For $s \in (0, 4)$. Let K be an arbitrary compact set in $\mathbb{R}_+^N \setminus \{0\}$, we claim

$$\begin{aligned} \lim_{m \rightarrow +\infty} \int_K \frac{u_m^{2_s^*}}{|x|^s} dx &= \lim_{m \rightarrow +\infty} \int_K \frac{u_m^2}{|x|^4} dx = 0, \\ \lim_{m \rightarrow +\infty} \int_K |\Delta u_m|^2 dx &= \lim_{m \rightarrow +\infty} \int_K |u_m|^{2_0^*} dx = 0. \end{aligned}$$

Proof of Step 4.1: Note that $u_m \rightharpoonup 0$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$ implies that $u_m \rightarrow 0$ strongly in $L_{loc}^q(\mathbb{R}_+^N)$ for $1 \leq q < 2_0^*$. Therefore, since $2 < 2_s^* < 2_0^*$ because $0 < s < 4$ and the fact $|x|^{-1}$ is bounded on K , we get

$$(80) \quad \int_K \frac{u_m^{2_s^*}}{|x|^s} dx = o(1) \text{ and } \int_K \frac{u_m^2}{|x|^4} dx = o(1) \text{ as } m \rightarrow +\infty.$$

We take $\eta \in C_c^\infty(\mathbb{R}_+^N \setminus \{0\})$ such that $\eta = 1$ in K and $0 \leq \eta \leq 1$.

Step 4.1.1 We claim as $m \rightarrow +\infty$ that

$$(81) \quad \int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx \leq \left(\int_{\mathbb{R}_+^N} |\eta u_m|^{2_0^*} dx \right)^{\frac{2}{2_0^*}} \left(\int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx \right)^{\frac{2_0^*-2}{2_0^*}} + o(1).$$

Proof of Step 4.1.1 : Indeed, using $\lim_{m \rightarrow +\infty} \langle E'(u_m), \eta^2 u_m \rangle = 0$ yields,

$$(82) \quad \begin{aligned} o(1) &= \langle E'(u_m), \eta^2 u_m \rangle = \int_{\mathbb{R}_+^N} \langle \Delta u_m, \Delta(\eta^2 u_m) \rangle dx - \gamma \int_{\mathbb{R}_+^N} \frac{|\eta u_m|^2}{|x|^4} dx \\ &\quad - \int_{\mathbb{R}_+^N} \frac{\eta^2 |u_m|^{2_s^*}}{|x|^s} dx - \int_{\mathbb{R}_+^N} \eta^2 |u_m|^{2_0^*} dx. \end{aligned}$$

Regarding the first term, we have

$$(83) \quad \begin{aligned} \int_{\mathbb{R}_+^N} \langle \Delta u_m, \Delta(\eta^2 u_m) \rangle dx &= \int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx + \int_{\mathbb{R}_+^N} u_m \Delta u_m \Delta(\eta^2) dx \\ &\quad + 2 \int_{\mathbb{R}_+^N} \Delta u_m \langle \nabla(\eta^2), \nabla u_m \rangle dx. \end{aligned}$$

From Hölder's inequality and since $\nabla u_m \rightarrow 0$ in $L_{loc}^2(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$, we get

$$(84) \quad \begin{aligned} \int_{\mathbb{R}_+^N} \Delta u_m \langle \nabla(\eta^2), \nabla u_m \rangle dx &= O \left(\|\nabla(\eta_m^2)\|_\infty \int_{\text{supp}(\nabla(\eta^2))} |\Delta u_m| |\nabla u_m| dx \right) \\ &= O \left(\|\nabla(\eta^2)\|_\infty \|u_m\| \left(\int_{\text{supp}(\nabla(\eta^2))} |\nabla u_m|^2 dx \right)^{\frac{1}{2}} \right) \\ &= o(1) \text{ as } m \rightarrow +\infty. \end{aligned}$$

Also, since $u_m \rightarrow 0$ in $L^2_{loc}(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$, we obtain that

$$(85) \quad \int_{\mathbb{R}_+^N} u_m \Delta u_m \Delta(\eta^2) dx = o(1) \text{ as } m \rightarrow +\infty.$$

Plugging (84) and (85) in (83) yields

$$\int_{\mathbb{R}_+^N} \langle \Delta u_m, \Delta(\eta^2 u_m) \rangle dx = \int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx + o(1) \text{ as } m \rightarrow +\infty.$$

We write $D := \text{supp}(\eta)$. It follows from (82) that

$$(86) \quad o(1) = \int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx - \gamma \int_D \frac{|\eta u_m|^2}{|x|^4} dx - \int_D \frac{\eta^2 |u_m|^{2_s^*}}{|x|^s} dx - \int_{\mathbb{R}_+^N} \eta^2 |u_m|^{2^{**}(0)} dx.$$

Similarly to (80) to get

$$\lim_{m \rightarrow +\infty} \int_D \frac{|\eta u_m|^2}{|x|^4} dx = \lim_{m \rightarrow +\infty} \int_D \frac{\eta^2 |u_m|^{2_s^*}}{|x|^s} dx = 0.$$

Therefore, by (86) and using again the Hölder's inequality, we find as $m \rightarrow +\infty$ that

$$\int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx \leq \left(\int_{\mathbb{R}_+^N} |\eta u_m|^{2_0^*} dx \right)^{\frac{2}{2_0^*}} \left(\int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx \right)^{\frac{2_0^*-2}{2_0^*}} + o(1).$$

This proves Step 4.1.1. □

Step 4.1.2 We claim that

$$(87) \quad \int_{\mathbb{R}_+^N} |\Delta(\eta u_m)|^2 dx = \int_{\mathbb{R}_+^N} |\eta \Delta u_m|^2 dx + o(1) \text{ as } m \rightarrow +\infty.$$

Proof of Step 4.1.2: Indeed, using the following inequality for $X, Y \in \mathbb{R}$

$$\left| |X+Y|^2 - |X|^2 \right| \leq C \left(|X||Y| + |Y|^2 \right),$$

with $X := \eta \Delta u_m$ and $Y := u_m \Delta \eta + 2 \nabla \eta \nabla u_m$, and we obtain that

$$(88) \quad \int_{\mathbb{R}_+^N} \left(|\Delta(\eta u_m)|^2 - |\eta \Delta u_m|^2 \right) dx = O \left(\int_{\mathbb{R}_+^N} |\eta \Delta u_m| |u_m \Delta \eta + 2 \nabla \eta \nabla u_m| dx \right. \\ \left. + \int_{\mathbb{R}_+^N} |u_m \Delta \eta + 2 \nabla \eta \nabla u_m|^2 dx \right).$$

Using Hölder's inequality and $u_m \rightarrow 0$ in $L^2_{loc}(\mathbb{R}_+^N)$ and $H^1_{loc}(\mathbb{R}_+^N)$ as $m \rightarrow +\infty$

$$(89) \quad \int_{\mathbb{R}_+^N} |\eta \Delta u_m| |u_m \Delta \eta + 2 \nabla \eta \nabla u_m| dx = O \left(\|\eta\|_\infty \|\Delta \eta\|_\infty \|u_m\| \left(\int_{\text{supp}(\eta) \cap \text{supp}(\Delta \eta)} u_m^2 dx \right)^{\frac{1}{2}} \right. \\ \left. + \|\eta\|_\infty \|\nabla \eta\|_\infty \|u_m\| \left(\int_{\text{supp}(\eta) \cap \text{supp}(\nabla \eta)} |\nabla u_m|^2 dx \right)^{\frac{1}{2}} \right) \\ = o(1) \text{ as } m \rightarrow +\infty.$$

In a similar way, we have

$$(90) \quad \int_{\mathbb{R}_+^N} |u_m \Delta \eta + 2 \nabla \eta \nabla u_m|^2 dx = o(1) \text{ as } m \rightarrow +\infty.$$

Injecting (89) and (90) in (88) and we have (87). This proves of Step 4.1.2. \square

Using (81) and (87) yields as $m \rightarrow +\infty$.

$$(91) \quad \int_{\mathbb{R}_+^N} |\Delta(\eta u_m)|^2 dx \leq \left(\int_{\mathbb{R}_+^N} |\eta u_m|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \left(\int_{\mathbb{R}_+^N} |u_m|^{2_s^*} dx \right)^{\frac{2_s^*-2}{2_s^*}} + o(1).$$

Now, since $\eta u_m \in \tilde{H}_0(\mathbb{R}_+^N)$, and we go back to the definition of $Q_{\gamma,0}(\mathbb{R}_+^N)$, and $\lim_{m \rightarrow +\infty} \int_{\text{supp}(\eta)} \frac{|\eta u_m|^2}{|x|^4} dx = 0$, we have that

$$(92) \quad \left(\int_{\mathbb{R}_+^N} |\eta u_m|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \|\eta u_m\|^2 + o(1) \text{ as } m \rightarrow +\infty.$$

It follows from (91) and (92) that

$$(93) \quad \left[1 - Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \left(\int_{\mathbb{R}_+^N} |u_m|^{2_s^*} dx \right)^{\frac{2_s^*-2}{2_s^*}} \right] \|\eta u_m\|^2 \leq o(1) \text{ as } m \rightarrow +\infty.$$

Since $E_{2_0^*}(u_m) = \beta$ and $E'_{2_0^*}(u_m) = o(1)$ as $m \rightarrow +\infty$, we have that

$$(94) \quad \begin{aligned} \beta + o(1) &= E_{2_0^*}(u_m) - \frac{1}{2} \langle E'_{2_0^*}(u_m), u_m \rangle \\ &= \left[\frac{1}{2} - \frac{1}{2_s^*} \right] \int_{\mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx + \left[\frac{1}{2} - \frac{1}{2_0^*} \right] \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx. \end{aligned}$$

Therefore, since $2_s^* > 2$ when $0 \leq s < 4$, we obtain that

$$(95) \quad \int_{\mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx \leq 2\beta \left[\frac{N-s}{4-s} \right] + o(1) \text{ and } \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx \leq \frac{N}{2}\beta + o(1) \text{ as } m \rightarrow +\infty.$$

Therefore, It follows from (93) and $\frac{2_0^*-2}{2_0^*} = \frac{4}{N}$ that

$$(96) \quad \left[1 - Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \left(\frac{N}{2}\beta \right)^{\frac{4}{N}} \right] \|\eta u_m\|^2 \leq o(1) \text{ as } m \rightarrow +\infty.$$

Since $\beta \in (0, \beta^*)$, and we have $\left[1 - Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \left(N \frac{\beta}{2} \right)^{\frac{4}{N}} \right] > 0$. Moreover, using inequality (96) yields $\lim_{m \rightarrow +\infty} \|\eta u_m\|^2 = 0$. But $\eta = 1$ in the compact K , then $\lim_{m \rightarrow +\infty} \int_K |\Delta u_m|^2 dx = 0$, from this and by the Sobolev inequality, we obtain that $\lim_{m \rightarrow +\infty} \int_K |u_m|^{2_s^*} dx = 0$. The proof of the Step (4.1) is complete. \square

For $R > 0$, we define

$$I_{1,R} := \lim_{m \rightarrow +\infty} \sup \int_{B_R(0)} |u_m|^{2_0^*} dx, \quad I_{2,R} := \lim_{m \rightarrow +\infty} \sup \int_{B_R(0)} \frac{|u_m|^{2_s^*}}{|x|^s} dx,$$

and

$$I_{3,R} := \lim_{m \rightarrow +\infty} \sup \int_{B_R(0)} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx.$$

Step 4.2. For $R > 0$, we claim that

$$I_{1,R}^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{3,R} ; I_{2,R}^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{3,R} \text{ and } I_{3,R} \leq I_{1,R} + I_{2,R}.$$

Proof of Step 4.2: Indeed, for $R > 0$ we take a cut-off function $\zeta \in C_c^\infty(\mathbb{R}_+^N)$ such that $\zeta = 1$ in $B_R(0)$ and $0 \leq \zeta \leq 1$. Since $\zeta u_m \in \tilde{H}_0(\mathbb{R}_+^N)$ and by the definition of $Q_{\gamma,0}(\mathbb{R}_+^N)$, we get

$$(97) \quad \left(\int_{\mathbb{R}_+^N} |\zeta u_m|^{2^*} dx \right)^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \left[\int_{B_R(0)} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx + \int_{\text{supp}(\zeta) \setminus B_R(0)} \left(|\Delta(\zeta u_m)|^2 - \gamma \frac{|\zeta u_m|^2}{|x|^4} \right) dx \right].$$

It follows from $\text{supp}(\zeta) \setminus B_R(0) \subset \mathbb{R}_+^N \setminus \{0\}$ and Step 4.1 that

$$\lim_{m \rightarrow +\infty} \int_{\text{supp}(\zeta) \setminus B_R(0)} \left(|\Delta(\zeta u_m)|^2 - \gamma \frac{|\zeta u_m|^2}{|x|^4} \right) dx = 0.$$

Therefore, using (97) and since $\zeta = 1$ in $B_R(0)$ yields

$$\left(\int_{B_R(0)} |u_m|^{2^*} dx \right)^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \int_{B_R(0)} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx + o(1),$$

as $m \rightarrow +\infty$, and we have that $I_{1,R}^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{3,R}$. The proof of $I_{2,R}^{\frac{2}{2^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{3,R}$ is similar.

Since $\zeta^2 u_m \in \tilde{H}_0(\mathbb{R}_+^N)$ and with $\lim_{m \rightarrow +\infty} \langle E'(u_m), \eta^2 u_m \rangle = 0$, we have

$$(98) \quad \begin{aligned} o(1) &= \langle E'(u_m), \eta^2 u_m \rangle \\ &= \int_{\mathbb{R}_+^N} \langle \Delta u_m, \Delta(\zeta^2 u_m) \rangle dx - \gamma \int_{\mathbb{R}_+^N} \frac{|\zeta u_m|^2}{|x|^4} dx \\ &\quad - \int_{\mathbb{R}_+^N} \frac{\zeta^2 |u_m|^{2^*}}{|x|^s} dx - \int_{\mathbb{R}_+^N} \zeta^2 |u_m|^{2^*} dx. \end{aligned}$$

It is similar of the proof of Step 4.1, we have as $m \rightarrow +\infty$ that

$$\int_{\mathbb{R}_+^N} \langle \Delta u_m, \Delta(\zeta^2 u_m) \rangle dx = \int_{\mathbb{R}_+^N} |\Delta(\zeta^2 u_m)|^2 dx + o(1).$$

Therefore, by (98), since $\zeta = 1$ in $B_R(0)$, $\text{supp}(\zeta) \setminus B_R(0) \subset \mathbb{R}_+^N \setminus \{0\}$ and with the result of Step 4.1, we have that

$$\begin{aligned} \int_{B_R(0)} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx &\leq \int_{\mathbb{R}_+^N} \left(|\Delta(\zeta^2 u_m)|^2 - \gamma \frac{|\zeta u_m|^2}{|x|^4} \right) dx \\ &= \int_{B_R(0)} \frac{|u_m|^{2^*}}{|x|^s} dx + \int_{B_R(0)} |u_m|^{2^*} dx + o(1). \end{aligned}$$

Taking $m \rightarrow +\infty$ on both sides yields $I_{3,R} \leq I_{1,R} + I_{2,R}$. This proves Step 4.2. \square
 Now, I will to complete the proof of the Proposition 4.3. Using Step 4.2 yields

$$I_{1,R}^{\frac{2}{2_0^*}} \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{1,R} + Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{2,R},$$

this give us

$$(99) \quad I_{1,R}^{\frac{2}{2_0^*}} \left[1 - Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{1,R}^{\frac{4}{N}} \right] \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{2,R}.$$

It follows from (95) and the definition of $I_{1,R}$ that $I_{1,R} \leq \frac{N}{2}\beta$. Therefore, by (99)

$$I_{1,R}^{\frac{2}{2_0^*}} \left[1 - Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} \left(\frac{N}{2}\beta \right)^{\frac{4}{N}} \right] \leq Q_{\gamma,0}(\mathbb{R}_+^N)^{-1} I_{2,R}.$$

Since $\beta < \beta^* < \frac{2}{N} Q_{\gamma,0}(\mathbb{R}_+^N)^{\frac{4}{N}}$, then there exists a constant $C_1(N, \gamma, \beta) > 0$ such that $I_{1,R}^{\frac{2}{2_0^*}} \leq C_1(N, \gamma, \beta) I_{2,R}$. Similar, then there exists $C_2(N, \gamma, s, \beta) > 0$ such that $I_{2,R}^{\frac{2}{2_s^*}} \leq C_2(N, \gamma, s, \beta) I_{1,R}$. Combining these two inequality we find that

$$I_{2,R}^{\frac{2}{2_s^*}} \left[1 - C_2(N, \gamma, s, \beta) C_1(N, \gamma, \beta)^{\frac{2_0^*}{2}} I_{2,R}^{\frac{N}{2(N-4)}} \right] \leq 0.$$

Therefore, we have $I_{2,R} = 0$ or there exists $\epsilon := \epsilon(N, \gamma, s, \beta)$ such that $I_{2,R} \geq \epsilon$. Similarly, we have $I_{1,R} = 0$ or there exists ϵ such that $I_{1,R} \geq \epsilon$. This ends of the proof of Propostion 4.3. \square

End of proof of Theorem 1.1 : Indeed, we let $(u_m)_{m \in \mathbb{N}}$ be the Palais-Smale sequence for E that was constructed in Proposition 4.1. First, we claim that

$$(100) \quad \lim_{m \rightarrow +\infty} \sup \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx > 0.$$

Indeed, otherwise $\lim_{m \rightarrow +\infty} \sup \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx = 0$. Using again $\lim_{m \rightarrow +\infty} \langle E'(u_m), u_m \rangle = 0$ yields,

$$\int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 - \gamma \frac{|u_m|^2}{|x|^4} \right) dx = \int_{\mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx + o(1).$$

Therefore, we go back to the definition of $Q_{\gamma,s}(\mathbb{R}_+^N)$

$$\|u\|_{2_s^*,s}^2 \leq Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \|u\|_{2_s^*,s}^{2_s^*} + o(1).$$

This give us,

$$\|u\|_{2_s^*,s}^2 \left[1 - Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \|u\|_{2_s^*,s}^{2_s^*-2} \right] \leq o(1).$$

It follows then from the left inequality of (95) that

$$\|u\|_{2_s^*,s}^2 \left[1 - Q_{\gamma,s}(\mathbb{R}_+^N)^{-1} \left(2 \frac{N-s}{4-s} \beta \right)^{\frac{2_s^*-2}{2_s^*}} \right] \leq o(1)$$

Since $0 < \beta < \beta^*$, we have the quantity between the brackets is positive. Thus, we get $\lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \frac{|u_m|^{2_s^*}}{|x|^s} dx = 0$. Therefore, using (94) yields $\beta = 0$ which contradicts the fact that $\beta \in (0, \beta^*)$. This proves the claim. \square

Next, we claim that the sequence $(u_m)_{m \in \mathbb{N}}$ is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$.

Indeed, since u_m is a Palais-Smale sequence for E and using (37) yields

$$\begin{aligned} \beta + o(1) &= E(u_m) - \frac{1}{2_s^*} \langle E'(u_m), u_m \rangle \\ &= \left[\frac{1}{2} - \frac{1}{2_s^*} \right] \int_{\mathbb{R}_+^N} \left(|\Delta u_m|^2 - \gamma \frac{u_m^2}{|x|^4} \right) dx + \left[\frac{1}{2_s^*} - \frac{1}{2_0^*} \right] \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx \\ &\geq C \left[\frac{1}{2} - \frac{1}{2_s^*} \right] \int_{\mathbb{R}_+^N} |\Delta u_m|^2 dx, \end{aligned}$$

where C is a positive constant. It follows then from $2 < 2_s^* < 2_0^*$ that u_m is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$. This proves the claim. \square

Since u_m is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$, then there exists $u \in \tilde{H}_0(\mathbb{R}_+^N)$ such that $u_m \rightharpoonup u$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$. If $u \neq 0$, we get that u is a nontrivial weak solution of (1).

If $u \equiv 0$, we have $u_m \rightharpoonup 0$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$. We claim that, for small enough $\epsilon' > 0$, there exists another Palais-Smale sequence $(v_m)_{m \in \mathbb{N}}$ satisfying the properties of Proposition 4.1 and

$$\int_{B_1(0)} |v_m|^{2_0^*} dx = \epsilon' ; v_m \text{ is bounded in } \tilde{H}_0(\mathbb{R}_+^N) \text{ for all } m \in \mathbb{N}.$$

Indeed, by (100), we can take $c := \limsup_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} |u_m|^{2_0^*} dx$. We set $\epsilon_0 := \min\{c, \frac{\epsilon}{2}\}$, where $\epsilon > 0$ is the same wick we obtain from Proposition 4.1. Therefore, for any $\epsilon' \in (0, \epsilon_0)$, there exists a sequence $(r_m)_{m \in \mathbb{N}} > 0$ such that up to a subsequence $\int_{B_{r_m}(0)} |u_m|^{2_0^*} dx = \epsilon'$. Define now $v_m(x) := r_m^{\frac{N-4}{2}} u_m(r_m x)$ for all $x \in \mathbb{R}_+^N$. With change of variable, we write

$$(101) \quad \int_{B_1(0)} |v_m|^{2_0^*} dx = \int_{B_{r_m}(0)} |u_m|^{2_0^*} dx = \epsilon'.$$

As one checks, $(v_m)_{m \in \mathbb{N}}$ is also a Palais-Smale sequence for E that satisfies the properties of Proposition 4.1. Using the definition of v_m and the boundedness of the sequence u_m yields v_m is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$. This ends the prove of Claim. \square

Hence, we can assume that there exists $v \in \tilde{H}_0(\mathbb{R}_+^N)$ such that, up to a subsequence $v_m \rightharpoonup v$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$.

We claim now that v is a nontrivial weak solution of problem (1).

Indeed, if $v \equiv 0$. It follows from the result of Proposition 4.3 that

$$\text{either } \limsup_{m \rightarrow +\infty} \int_{B_1(0)} |v_m|^{2_0^*} dx = 0 \text{ or } \limsup_{m \rightarrow +\infty} \int_{B_1(0)} |v_m|^{2_0^*} dx \geq \epsilon.$$

Since $\epsilon' \in (0, \frac{\epsilon}{2})$, this is contradiction with (101). Then $v \neq 0$. \square

Since $(v_m)_{m \in \mathbb{N}}$ is a sequence Palais-Smale for E , we have

$$(102) \quad \begin{aligned} o(1) &= \langle E'(v_m), \varphi \rangle \\ &= \int_{\mathbb{R}_+^N} \langle \Delta v_m, \Delta \varphi \rangle dx - \gamma \int_{\mathbb{R}_+^N} \frac{v_m \varphi}{|x|^4} dx \\ &\quad - \int_{\mathbb{R}_+^N} |v_m|^{2_0^*-2} v_m \varphi dx - \int_{\mathbb{R}_+^N} \frac{|v_m|^{2_s^*-2} v_m \varphi}{|x|^s} dx, \end{aligned}$$

for all $\varphi \in C_c^\infty(\mathbb{R}_+^N)$. Using $v_m \rightharpoonup v$ weakly in $\tilde{H}_0(\mathbb{R}_+^N)$ yields

$$(103) \quad \lim_{m \rightarrow +\infty} \int_{\mathbb{R}_+^N} \langle \Delta v_m, \Delta \varphi \rangle dx = \int_{\mathbb{R}_+^N} \langle \Delta v, \Delta \varphi \rangle dx \text{ for all } \varphi \in C_c^\infty(\mathbb{R}_+^N).$$

Since v_m is bounded in $\tilde{H}_0(\mathbb{R}_+^N)$, we get that v_m , $|v_m|^{2_0^*-2} v_m$ and $|v_m|^{2_s^*-2} v_m$ are bounded in $L^2(|x|^{-4}, \mathbb{R}_+^N)$, $L^{\frac{2_0^*}{2_0^*-1}}(\mathbb{R}_+^N)$ and $L^{\frac{2_s^*}{2_s^*-1}}(|x|^{-s}, \mathbb{R}_+^N)$ respectively. Therefore, we get that

$$\begin{cases} v_m \rightharpoonup v \text{ weakly in } L^2(|x|^{-4}, \mathbb{R}_+^N), \\ |v_m|^{2_0^*-2} v_m \rightharpoonup |v|^{2_0^*-2} v \text{ weakly in } L^{\frac{2_0^*}{2_0^*-1}}(\mathbb{R}_+^N), \\ |v_m|^{2_s^*-2} v_m \rightharpoonup |v|^{2_s^*-2} v \text{ weakly in } L^{\frac{2_s^*}{2_s^*-1}}(|x|^{-s}, \mathbb{R}_+^N). \end{cases}$$

Moreover, passing the $m \rightarrow +\infty$ in (102) and using (103) yields

$$\begin{aligned} o(1) &= \int_{\mathbb{R}_+^N} \langle \Delta v, \Delta \varphi \rangle dx - \gamma \int_{\mathbb{R}_+^N} \frac{v\varphi}{|x|^4} dx \\ &\quad - \int_{\mathbb{R}_+^N} |v|^{2_0^*-2} v\varphi dx - \int_{\mathbb{R}_+^N} \frac{|v|^{2_s^*-2} v\varphi}{|x|^s} dx \text{ for all } \varphi \in C_c^\infty(\mathbb{R}_+^N). \end{aligned}$$

Thus, v is a weak solution of (1). This completes the proof of Theorem 1.1. \square

5. APPENDIX

Proposition 5.1. *Let $\Omega \subset \mathbb{R}^N$ be a smooth domain such that $0 \in \partial\Omega \neq \emptyset$ (No bound- edness is assumed). If $\gamma \leq 0$ and $s = 0$, we have that $Q_{\gamma,0}(\Omega) = S_N$, where S_N is defined in (10) and there is no extremal.*

Proof of Proposition 5.1: Indeed, we call back $2_0^* = \frac{2N}{N-4}$. Since $\gamma \leq 0$, we find

$$\frac{\int_{\Omega} \left(|\Delta u|^2 - \gamma \frac{u^2}{|x|^4} \right) dx}{\left(\int_{\Omega} |u|^{2_0^*} dx \right)^{\frac{2}{2_0^*}}} \geq \frac{\int_{\Omega} |\Delta u|^2 dx}{\left(\int_{\Omega} |u|^{2_0^*} dx \right)^{\frac{2}{2_0^*}}} \geq S_N,$$

and we have that $Q_{\gamma,0}(\Omega) \geq S_N$. Fix $x_0 \in \Omega$ such that $x_0 \neq 0$. We define $(U_\epsilon)_\epsilon$ as in (68). It follows from (70) that $\lim_{\epsilon \rightarrow 0} \int_{\Omega} \frac{U_\epsilon^2}{|x|^4} dx = 0$. Moreover, with (76) and (77), we get that

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{\Omega} |\Delta U_\epsilon|^2 dx}{\left(\int_{\Omega} |U_\epsilon|^{2_0^*} dx \right)^{\frac{2}{2_0^*}}} = S_N.$$

We then get $Q_{\gamma,0}(\Omega) \leq S_N$. This proves that $Q_{\gamma,0}(\Omega) = S_N$. If there was an extremal for $Q_{\gamma,0}(\Omega)$, it would also be a extremal for S_N , with their support is the whole of \mathbb{R}^N , contradicting since the extremal has support in $\Omega \neq \mathbb{R}^N$ smooth. This proves Proposition 5.1. \square

REFERENCES

- [1] A. Ambrosetti and P.H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis **14** (1973), 349–381.
- [2] M. Bhakta, *Semilinear elliptic equation with biharmonic operator and multiple critical nonlinearities*, Adv. Nonlinear Stud. **15** (2015), no. 4, 835–848.
- [3] ———, *Caffarelli-Kohn-Nirenberg type equations of fourth order with the critical exponent and Rellich potential*, J. Math. Anal. Appl. **433** (2016), no. 1, 681–700.

- [4] M. Bhakta and R. Musina, *Entire solutions for a class of variational problems involving the biharmonic operator and Rellich potentials*, *Nonlinear Anal.* **75** (2012), no. 9, 3836–3848.
- [5] H. Cheikh Ali, *Hardy-Sobolev inequalities with singularities on non smooth boundary: Hardy constant and extremals. Part I: Influence of local geometry.*, *Nonlinear analysis* **182** (2019), 316–349.
- [6] ———, *Hardy-Sobolev inequalities with singularities on non smooth boundary. Part 2: Influence of the global geometry in small dimensions.*, *Journal of differential equations* **270** (2021), 185–216.
- [7] ———, *Hardy-Sobolev singular of fourth order with a singularity on a smooth boundary.* in progress.
- [8] P. Caldiroli and R. Musina, *Rellich inequalities with weights*, *Calculus of Variations and Partial Differential Equations* **45** (2012), 147–164.
- [9] J. L. Chern and C. S. Lin, *Minimizers of Caffarelli-Kohn-Nirenberg inequalities with the singularity on the boundary*, *Arch. Ration. Mech. Anal.* **197** (2010), no. 2, 401–432.
- [10] R. Filippucci, P. Pucci, and F. Robert, *On a p -Laplace equation with multiple critical nonlinearities*, *J. Math. Pures Appl.* (9) **91** (2009), no. 2, 3200–3225.
- [11] N. Ghoussoub and F. Robert, *The effect of curvature on the best constant in the Hardy-Sobolev inequalities*, *Gafa, Geom. funct. anal.* **10** (2006), no. 16, 1201–1245.
- [12] ———, *Hardy-singular boundary mass and Sobolev-critical variational problems*, *Anal. PDE* **10** (2017), no. 5, 1017–1079.
- [13] N. Ghoussoub and X. S. Kang, *Hardy-Sobolev critical elliptic equations with boundary singularities*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **21** **6** (2004), 767–793.
- [14] N. Ghoussoub and S. Shakerian, *Borderline Variational Problems Involving Fractional Laplacians and Critical Singularities*, *Advanced Nonlinear Studies* **15** (2015), no. 3, 527–555.
- [15] P. L. Lions, *Applications de la méthode de concentration-compacité à l'existence de fonctions extrémales.*, *C. R. Acad. Sci. Paris Sér. I Math.* **296** (1983), no. 15, 645–648.
- [16] ———, *The concentration-compactness principle in the calculus of variations.*, The limit case. I, *Rev. Mat. Iberoamericana* **1** (1985), 145–201.
- [17] E. Mitidieri, *A simple approach to Hardy inequalities*, translation in *Math. Notes* **67** (2000), no. 3-4, 479–486.
- [18] M. Pérez-Llanos and A. Primo, *Semilinear biharmonic problems with a singular term*, *Journal of Differential Equations* **257** (2014), 3200–3225.
- [19] F. Rellich, *Halbeschränkte Differentialoperatoren höherer Ordnung*, In: J.C.H. Gerretsen, J. de Groot (Eds.): *Proceedings of the International Congress of Mathematicians 1954*, Volume III (pp. 243–250) Groningen: Noordhoff 1956.
- [20] ———, *Perturbation theory of eigenvalue problems*, *Gordon and Breach*, New York, 1969.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ LIBRE DE BRUXELLES, CP 214, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIUM
Email address: Hussein.cheikh-ali@ulb.ac.be