

LOCAL STRUCTURE OF HOMOGENEOUS ANR-SPACES

V. VALOV

ABSTRACT. We investigate to what extent finite-dimensional homogeneous locally compact ANR-spaces have common properties with Euclidean manifolds. Specially, the local structure of homogeneous ANR-spaces is described. Using that description, we provide a positive solution of the problem whether every finite-dimensional homogeneous metric ANR-compactum X is dimensionally full-valued, i.e. $\dim X \times Y = \dim X + \dim Y$ for any metric compactum Y .

1. INTRODUCTION

By a *space* we mean a locally compact separable metric space, and *maps* are continuous mappings. Reduced Čech homology $H_n(X; G)$ and cohomology groups $H^n(X; G)$ with coefficient from an abelian group G are considered everywhere below. A space is said to be *homogeneous* provided that, for every two points $x, y \in X$ there exists a homeomorphism h mapping X onto itself with $h(x) = y$. Homogeneity of X implies *local homogeneity*, that is for every $x, y \in X$ there exists a homeomorphism h mapping a neighborhood of x onto a neighborhood of y such that $h(x) = y$.

One of the reasons to investigate the homogeneous ANR-spaces is the well-known Bing-Borsuk [1] conjecture that every finite-dimensional homogeneous metric ANR-compactum is an Euclidean manifold. According to Jacobsche [14], a positive solution of that conjecture in dimension three implies the celebrated Poincaré conjecture. J. Bryant and S. Ferry [4] announced the existence of many counter-examples to that conjecture for every $n \geq 6$, but to the best of author's knowledge, the Bryant-Ferry paper is not published yet. Despite that the status of the Bing-Borsuk conjecture is unknown, it is still interesting to

Date: April 24, 2023.

2000 Mathematics Subject Classification. Primary 55M15; Secondary 55M10.

Key words and phrases. absolute neighborhood retracts, cohomological dimension, cohomology groups, homogeneous spaces.

The author was partially supported by NSERC Grant 261914-19.

investigate the extend to which finite-dimensional homogeneous *ANR*-spaces have common properties with Euclidean manifolds. We show that homogeneous *ANR*-spaces have indeed some properties typical for Euclidean manifolds.

The local structure of homogeneous *ANR*-compacta X with a finite cohomological dimension $\dim_G X = n$ was established in [29, Theorem 1.1], where G is a countable principal ideal domain with unity and $n \geq 2$. In the present paper we improve the results in [29] by considering locally compact homogeneous *ANR*-spaces and replacing the principal ideal domains with countable groups. Moreover, we also provide some new properties of homogeneous *ANR*'s or locally homogeneous *ANR*'s.

The next theorem describes the local structure of homogeneous *ANR*-spaces and shows their similarity with Euclidean manifolds:

Theorem 1.1. *Let X be a homogeneous connected *ANR*-space with $\dim_G X = n \geq 2$, where G is a countable group. Then every point $x \in X$ has a base \mathcal{B}_x of connected open sets $U \subset X$ each having a compact closure and satisfying the following conditions:*

- (1) $\text{Int} \bar{U} = U$, the boundary $\text{bd} \bar{U}$ of \bar{U} is connected and its complement in X has exactly two components;
- (2) $H^{n-1}(\text{bd} \bar{U}; G) \neq 0$, $H^{n-1}(\bar{U}; G) = 0$ and \bar{U} is an $(n-1)$ -cohomology membrane spanned on $\text{bd} \bar{U}$ for any non-zero $\gamma \in H^{n-1}(\text{bd} \bar{U}; G)$;
- (3) $\text{bd} \bar{U}$ is an $(n-1, G)$ -bubble.

Recall that for any nontrivial abelian group G the Čech cohomology group $H^n(X; G)$ is isomorphic to the group of pointed homotopy classes of maps from X to $K(G, n)$, where $K(G, n)$ is a *CW*-complex of type (G, n) , see [17]. The cohomological dimension $\dim_G X$ is the largest number n such that there exists a closed set $A \subset X$ with $H^n(X, A; G) \neq 0$. Equivalently, for a metric space X we have $\dim_G X \leq n$ if and only if for any closed pair $A \subset B$ in X the homomorphism $j_{B,A}^n : H^n(B; G) \rightarrow H^n(A; G)$, generated by the inclusion $A \hookrightarrow B$, is surjective, see [10]. This means that every map from A to $K(G, n)$ can be extended over B . If X is a finite-dimensional space, then for every G we have $\dim_G X \leq \dim_{\mathbb{Z}} X = \dim X$ [21]. On the other hand, there is a compactum X [8] with $\dim X = \infty$ and $\dim_{\mathbb{Z}} X = 3$. A space A is a (k, G) -bubble if $H^k(A; G) \neq 0$ but $H^k(B; G) = 0$ for every closed proper subset B of A .

The following property of homogeneous *ANR*'s is essential, see [1], [2]: If (K, A) is a pair of closed subsets of X such that K is an $(n-1)$ -cohomological membrane for some $\gamma \in H^{n-1}(A; G)$, then $(K \setminus A) \cap$

$\overline{X \setminus K} = \emptyset$. We call this property the *n-cohomology membrane property*. Recall that K is said to be an *k-cohomology membrane spanned on a closed set $A \subset K$ for an element $\gamma \in H^k(A; G)$* if γ is not extendable over K , but it is extendable over every proper closed subset of K containing A . Here, $\gamma \in H^k(A; G)$ is not extendable over K means that γ is not contained in the image $j_{K,A}^k(H^k(K; G))$.

Everywhere below, \mathcal{B}_x stands for the family of all open neighborhoods of a point $x \in X$. We say that a space X has an *n-dimensional G-obstruction* at a point $x \in X$ [21] if there is $W \in \mathcal{B}_x$ such that the homomorphism $j_{U,W}^n : H^n(X, X \setminus U; G) \rightarrow H^n(X, X \setminus W; G)$ is nontrivial for every $U \in \mathcal{B}_x$ with $U \subset W$. Kuzminov [21], [22], proved that every compactum X with $\dim_G X = n$ contains a compact set Y with $\dim_G Y = n$ such that X has an *n-dimensional G-obstruction* at any point of Y . According to the next theorem, *n-dimensional homogeneous spaces have an obstruction at every point and the corresponding homomorphisms are surjective*.

Theorem 1.2. *Let X be a locally homogeneous ANR-space such that $\dim_G X = n$, where G is a countable group. Then the following holds:*

- (i) *X has the n-cohomology membrane property;*
- (ii) *X has an n-dimensional G-obstruction at every $x \in X$. Moreover, there is $W \in \mathcal{B}_x$ such that the homomorphism $j_{U,V}^n$ is surjective for any $U, V \in \mathcal{B}_x$ with $\overline{U} \subset V \subset \overline{V} \subset W$.*

Here is another common property of locally homogeneous ANR's and Euclidean manifolds.

Corollary 1.3. *Let X be as in Theorem 1.2. If $U \subset X$ is open and $f : U \rightarrow X$ is an injective map, then $f(U)$ is also open in X .*

One of the important questions concerning homogeneous ANR's is whether every finite-dimensional homogeneous ANR-compactum is dimensionally full-valued. Recall that a locally compact space X is *dimensionally full-valued* if $\dim X \times Y = \dim X + \dim Y$ for any compact space Y . Let us note that there are metric ANR-compacta which are not dimensionally full-valued, see [7] and [9]. The question for the dimensional full-valuedness of homogeneous ANR-compacta goes back to [3] and was also discussed in [5] and [13]. A positive answer of that question for 3-dimensional spaces was given in [29]. Now, we provide a complete solution.

Theorem 1.4. *Let X be a finite-dimensional locally homogeneous ANR-space. Then the following holds:*

- (i) *X is dimensionally full-valued;*

- (ii) *Providing X is homogeneous and connected, every $x \in X$ has a neighborhood W_x such that $\text{bd} \overline{U}$ is dimensionally full-valued for all $U \in \mathcal{B}_x$ with $\overline{U} \subset W_x$.*

2. THE COHOMOLOGY MEMBRANE PROPERTY

If, in the definition of the n -cohomology membrane property, we additionally require K to be a compact set contractible in a proper subset of X , then we say that X has the *weak n -cohomology membrane property*. We will see that the proof of Theorem 1.1 is based mainly on that property. In this section we prove that any locally homogeneous space X with $\dim_G X = n$ has the n -cohomology membrane property, and provide some implications of that property.

A space X is called a (k, G) -carrier of a nontrivial element $\gamma \in H^k(X; G)$ if $j_{X,B}^k(\gamma) = 0$ for every closed proper subset $B \subset X$.

Lemma 2.1. *Let $A \subset X$ be a compact set and γ be a non-zero element of $H^n(A; G)$.*

- (i) *If γ is not extendable over a compact set $P \subset X$ containing A , then there exists an n -cohomology membrane $K \subset P$ for γ spanned on A ;*
(ii) *There is a closed set $B \subset A$ such that B is a carrier of $j_{A,B}^n(\gamma)$.*

Proof. (i) Consider the family \mathcal{F} of all closed subsets F of P containing A such that γ is not extendable over F . Let $\{F_\tau\}$ be a decreasing subfamily of \mathcal{F} and $F_0 = \bigcap_\tau F_\tau$. Suppose γ is extendable to $\tilde{\gamma} \in H^n(F_0; G)$. Then, considering $\tilde{\gamma}$ as a map from F_0 into $K(G, n)$ and having in mind that $K(G, n)$ is an absolute neighborhood for metrizable spaces [19], we can extend $\tilde{\gamma}$ over a neighborhood W of F_0 in P . But that is impossible since W contains some F_τ . Hence, by Zorn's lemma, \mathcal{F} has a minimal element K which is an n -cohomology membrane for γ spanned on A .

(ii) Now, let \mathcal{F} be the family of closed subsets F of A such that $j_{A,F}^n(\gamma) \neq 0$ and $\{F_\tau\}$ be a decreasing subfamily of \mathcal{F} with $F_0 = \bigcap_\tau F_\tau$. If $\gamma_0 = j_{A,F_0}^n(\gamma) = 0$, then there is a homotopy $H : F_0 \times [0, 1] \rightarrow K(G, n)$ connecting the constant map and γ_0 . Using again that $K(G, n)$ is an absolute neighborhood extensor for metric spaces, we find a closed neighborhood W of F_0 in A and a homotopy $\tilde{H} : W \times [0, 1] \rightarrow K(G, n)$ connecting $j_{A,W}^n(\gamma)$ and the constant map. This is a contradiction because W contains some F_τ . Hence, \mathcal{F} has a minimal element B which is a carrier of $j_{A,B}^n(\gamma)$. \square

We also need the following version of Effros' theorem [11] for locally compact spaces (see [20, Proposition 1.4]):

Theorem 2.2. *Let X be a homogeneous space with a metric ρ , $a \in X$ and $\varepsilon > 0$. Then there is $\delta > 0$ such that for every $x \in X$ with $\rho(x, a) < \delta$ there exists a homeomorphism $h : X \rightarrow X$ with $h(a) = x$ and $\rho(h(y), y) < \varepsilon$ for all $y \in X$.*

Proposition 2.3. *Let X be a homogeneous ANR-space with $\dim_G X = n$, where G is an arbitrary group. Then $H^n(P; G) = 0$ for any proper compact set $P \subset X$.*

Proof. This was established in [30] in case X is compact using the following proposition, see [30, Proposition 2.3]: If X is a metric compactum with $\dim_G X = n$, $A \subset X$ is carrier of a nontrivial element of $H^n(X; G)$ and there is a map $f : X \rightarrow X$ homotopic to the identity id_X of X , then $A \subset f(A)$. This statement remains true if X is an ANR-space with $\dim_G X = n$, A is a carrier of a nontrivial element of $H^n(Q; G)$, where $Q \subset X$ is compact, and there exists a homotopy $F : A \times [0, 1] \rightarrow X$ with $F(x, 0) = x$ and $F(x, 1) = f(x)$, $x \in A$.

Fix a metric ρ on X generating its topology and suppose there exists a proper compact set $P \subset X$ with $H^n(P; G) \neq 0$. So, by Lemma 2.1(ii), P contains a compact set K such that K is a carrier for $j_{P,K}^n(\alpha)$, where $\alpha \in H^n(P; G)$ is nontrivial. Take two open sets $U, V \subset X$ containing P and having compact closures in X with $\overline{U} \subset V$. Since $X \in \text{ANR}$, there is an open cover ω of X such that any two ω -close maps $f_1, f_2 : \overline{U} \rightarrow X$ are homotopic. Restricting ω on \overline{V} , we find $\varepsilon > 0$ such that any maps $f_1, f_2 : \overline{U} \rightarrow X$ with $\rho(f_1(x), f_2(x)) < \varepsilon$, $x \in \overline{U}$, are homotopic. Moreover, we can assume $\varepsilon < \min\{\rho(P, X \setminus U), \rho(\overline{U}, X \setminus V)\}$. Next, take a point a from the boundary $\text{bd}K$ of K in X and a point $b \notin K$ with $\rho(a, b) < \delta$, where $\delta > 0$ is the Effros' number corresponding to ε and the point a . Accordingly, there is homeomorphism $h : X \rightarrow X$ such that $h(a) = b$ and $\rho(x, h(x)) < \varepsilon$ for all $x \in X$. So, $h(K) \subset U$ and $h(\overline{U}) \subset V$. Since h generates an isomorphism $h^* : H^n(h(P); G) \rightarrow H^n(P; G)$, the set $A = h(K)$ is a carrier for $j_{h(P), h(K)}^n(\beta)$, where $\beta = (h^*)^{-1}(\alpha)$. Consider the map $g = h^{-1}|_{\overline{U}} : \overline{U} \rightarrow h^{-1}(\overline{U})$. Observe that $\rho(g(x), x) < \varepsilon$ for all $x \in \overline{U}$. Hence, g and the identity $\text{id}_{\overline{U}}$ on \overline{U} are homotopic. In particular, there is a homotopy $F : g(A) \times [0, 1] \rightarrow X$ such that $F(x, 0) = x$ and $F(x, 1) = g(x)$, $x \in g(A)$. Therefore, we can apply the modification of [30, Proposition 2.3] stated above to conclude that $A \subset g(A)$. Because $b = h(a) \in A$, the last inclusion implies $b \in g(A) = K$, a contradiction. \square

Remark. *Proposition 2.3 remains true without homogeneity of X provided P is a closed set contractible in X*

This is true if $H^n(X; G) = 0$ because $\dim_G X \leq n$. If $H^n(X; G) \neq 0$ this is also true. Indeed, since $\dim_G X \leq n$, every $\alpha \in H^n(P; G)$ is extendable to $\tilde{\alpha} \in H^n(X; G)$. On the other hand P is contractible in X , so every such $\tilde{\alpha}$ is zero.

A compact pair $A \subset K$ is called a *k-homological membrane spanned on A* for a nontrivial element $\gamma \in H_k(A; G)$ if $i_{A,K}^k(\gamma) = 0$ but $i_{A,B}^k(\gamma) \neq 0$ for any proper closed subset B of K containing A . Here, H_* denotes the Čech homology and $i_{A,K}^k : H_k(A; G) \rightarrow H_k(K; G)$ is the homomorphism induced by the inclusion $A \hookrightarrow K$. According to Bing-Borsuk [1], if $A \subset K$ is a compact pair and $\gamma \in H_k(A; G)$ is a nontrivial element homologous to zero in K , then there is a closed set $B \subset K$ containing A such that $A \subset B$ a *k-homological membrane for γ spanned on A*. Bing-Borsuk [1] considered the Vietoris homology which is isomorphic with the Čech homology in the realm of metric compacta. So, homology membranes with respect to Čech homology always exist in the class of metric compacta.

Proposition 2.4. *Any locally homogeneous ANR-space with $\dim_G X = n$, where G is a countable group, has the weak n -cohomological membrane property.*

Proof. Suppose there exists a compact pair $A \subset K$ such that K is contractible in a proper subset of X and K is an $(n-1)$ -cohomological membrane for some nontrivial $\gamma \in H^{n-1}(A; G)$, but $(K \setminus A) \cap \overline{X \setminus K} \neq \emptyset$. Since G is countable, the homology group $H_{n-1}(Y; G^*)$ is isomorphic to $H^{n-1}(Y; G)^*$ for any metric compactum Y , see [16]. Here, G^* and $H^{n-1}(Y; G)^*$ are the character groups of G and $H^{n-1}(Y; G)$, considered as discrete groups. Then K is an $(n-1)$ -homological membrane spanned on A for some nontrivial $\beta \in H_{n-1}(A; G^*)$, see the proof of [29, Proposition 2.1]. Hence, following the proof of [1, Theorem 8.1], we can find a compact set $P \subset X$ contractible in X with $H_n(P; G^*) \neq 0$. So, $H^n(P; G) \neq 0$, which contradicts the remark after Proposition 2.3. \square

We also need the following properties of cohomology membranes, see Corollary 2.2 and Lemma 2.4 from [29], respectively.

Lemma 2.5. *For any space X with the weak n -cohomology membrane property the following conditions hold:*

- (i) *If K is an $(n-1)$ -cohomology membrane spanned on a set $A \subset K$ for some $\gamma \in H^{n-1}(A; G)$, where K is a compactum contractible in a proper subset of X , then $K \setminus A$ is a connected open subset of X ;*
- (ii) *Let $A \subset P$ be a compact pair such that P is contractible in a proper subset of X and there exists a non-zero $\gamma \in H^{n-1}(A; G)$*

not extendable over P . Then A separates every set $\Gamma \subset X$ containing P as a proper subset.

Let (X, ρ) be a metric space. We say that ρ is *convex* if for each $x, y \in X$ there exists an arc $A \subset X$ with end-points x and y such that A with the restriction of the metric ρ is isometric to the interval $[0, \rho(x, y)]$ in the real line (where the real line is considered with its usual metric). According to [27], every connected locally connected space admits a convex metric.

Proposition 2.6. *Let $\{G_i\}_{i=1}^k$ be a finite family of countable groups and X be a locally homogeneous ANR-space with $\dim_{G_i} X = n_i$. Then we have:*

- (1) *Every closed set $P \subset X$ with $\dim_{G_i} P = n_i$, $i \leq k$, has a non-empty interior;*
- (2) *Every $x \in X$ has a neighborhood W such that any connected $U \in \mathcal{B}_x$ with $U = \text{int} \bar{U} \subset W$ satisfies the following for all i :*
 - (i) *$H^{n_i}(\bar{U}; G_i) = 0$ and $H^{n_i-1}(\text{bd} \bar{U}; G_i) \neq 0$ contains an element not extendable over \bar{U} ;*
 - (ii) *\bar{U} is an $(n_i - 1)$ -cohomology membrane spanned on $\text{bd} \bar{U}$ for any nontrivial $\alpha_i \in H^{n_i-1}(\text{bd} \bar{U}; G_i)$ not extendable over \bar{U} .*

Proof. (i) This was established in [29, Corollary 2.3] in case X is compact homogeneous ANR. For the covering dimension \dim and locally homogeneous ANR's it was established in [25, Theorem A] (see also [24] for a property stronger than locally homogeneity). Suppose $P \subset X$ is a closed set with $\dim_{G_i} P = n_i$ for some i . Then P is the union of countably many compact sets F_j each contractible in a proper subset of X . By the countable sum theorem for \dim_{G_i} , we have $\dim_{G_i} F_j = n$ for at least one j . So, we can assume that P is a compact set contractible in a proper subset of X . Since $\dim_{G_i} P = n_i$, there is a closed set $A \subset P$ and an element $\gamma_i \in H^{n_i-1}(A; G_i)$ not extendable over P . Then, according to Lemma 2.1(i), there exists a closed set $K \subset P$ such that K is an $(n_i - 1)$ -cohomology membrane for γ_i spanned on A . Because X has the weak n_i -cohomological membrane property, $(K \setminus A) \cap \overline{X \setminus K} = \emptyset$, which implies $K \setminus A$ is open in X . Finally, the inclusion $K \setminus A \subset P$ completes the proof.

(ii) Since X is a countable union of compact sets each contractible in a proper subset of X , as in the previous paragraph, for every i there exists a compact pair $A_i \subset K_i$ such that $\dim_{G_i} K_i = n_i$, K_i is an $(n_i - 1)$ -cohomology membrane for some $\gamma_i \in H^{n_i-1}(A_i; G_i)$ spanned on A_i and K_i is contractible in a proper subset of X . Then each $K_i \setminus A_i$ is a connected open set in X , see Lemma 2.5(i). Let $x_i \in K_i \setminus A_i$ and let

$W_i \in \mathcal{B}_{x_i}$ with $\overline{W_i} \subset K_i \setminus A_i$. Because of the local homogeneity of X , we can assume that all W_i are homeomorphic.

Let $U_i \in \mathcal{B}_{x_i}$ be connected with $U_i = \text{int}\overline{U_i} \subset W_i$. Such sets U_i exist. Indeed, since X is locally connected, each its component of connectedness X_c is open, and according to [27], there is a convex metric generating the topology of X_c . On the other hand, if d is a convex metric, then every open ball $B(x, \delta) = \{y \in X_c : d(x, y) < \delta\}$ is connected and $\text{int}\overline{B(x, \delta)} = B(x, \delta)$.

Since each $\overline{U_i}$ is contractible in a proper subset of X , $H^{n_i}(\overline{U_i}; G_i) = 0$. We claim that $H^{n_i-1}(\text{bd}\overline{U_i}; G) \neq 0$ and contains an element not extendable over $\overline{U_i}$. Indeed, $K_i \setminus U_i$ is a proper closed subset of K_i containing A_i . So, γ_i can be extended to $\tilde{\gamma}_i \in H^{n_i-1}(K_i \setminus U_i; G_i)$. Then $\gamma_{U_i} = j_{K_i \setminus U_i, \text{bd}\overline{U_i}}^{n_i-1}(\tilde{\gamma}_i)$ is a non-zero element of $H^{n_i-1}(\text{bd}\overline{U_i}; G)$ (otherwise γ_i would be extendable over K_i), and γ_{U_i} is not extendable over $\overline{U_i}$. Let show that $\overline{U_i}$ is an (n_i-1) -cohomology membrane spanned on $\text{bd}\overline{U_i}$ for every $\alpha_i \in H^{i-1}(\text{bd}U_i; G_i)$ not extendable over $\overline{U_i}$. By Lemma 2.1, for any such α_i there is an (n_i-1) -cohomology membrane $K_{\alpha_i} \subset \overline{U_i}$ for α_i spanned on $\text{bd}\overline{U_i}$. Hence, $(K_{\alpha_i} \setminus \text{bd}\overline{U_i}) \cap \overline{X} \setminus \overline{K_{\alpha_i}} = \emptyset$. In particular, $K_{\alpha_i} \setminus \text{bd}\overline{U_i}$ is open in U_i . Thus, $K_{\alpha_i} = \overline{U_i}$, otherwise U_i would be the union of the non-empty disjoint open sets $U_i \setminus K_{\alpha_i}$ and $(K_{\alpha_i} \setminus \text{bd}\overline{U_i})$. Finally, since X is locally homogeneous, every $x \in X$ has a neighborhood W satisfying condition (2). \square

Lemma 2.7. *Let X be a locally homogeneous ANR-space and $x \in X$. If G is a countable group and $H^{n-1}(\text{bd}\overline{U}; G) \neq 0$ for all sufficiently small neighborhoods $U \in \mathcal{B}_x$, then $\dim_G X \geq n$.*

Proof. Suppose $\dim_G X \leq n-1$. Since the interior of $\text{bd}\overline{U}$ is empty, $\dim_G \text{bd}\overline{U} \leq n-2$ for all $U \in \mathcal{B}_x$. This, according to the definition of cohomological dimension, implies $H^{n-1}(\text{bd}\overline{U}; G) = 0$, a contradiction. \square

3. PROOF OF THEOREM 1.1

The next lemma was established in [6, Theorem 8] for locally connected continua. The same proof works for locally connected and connected spaces X by passing to the one-point compactification of X .

Lemma 3.1. *Let X be a connected and locally connected space and $\{K_\alpha : \alpha \in \Lambda\}$ be an uncountable collection of disjoint continua such that for each α the set $X \setminus K_\alpha$ has more than one component. Then there exists $\alpha_0 \in \Lambda$ such that $X \setminus K_{\alpha_0}$ has exactly two components.*

The proof of Theorem 1.1 follows from Proposition 2.6 and Proposition 3.2 below.

Proposition 3.2. *Let X be a homogeneous connected ANR-space with $\dim_G X = n \geq 2$, where G is a countable group. Then every point $x \in X$ has a basis \mathcal{B}_x of open connected sets U each with a compact closure satisfying the following conditions:*

- (i) $H^{n-1}(\overline{U}; G) = 0$ and $X \setminus \text{bd } \overline{U}$ has exactly two components;
- (ii) $\text{bd } \overline{U}$ is an $(n-1, G)$ -bubble.

Proof. By Proposition 2.6, every point $x \in X$ has a neighborhood W_x with a compact closure such any connected neighborhood $U = \text{int } \overline{U} \subset W_x$ of x satisfies the following condition: $H^n(\overline{U}; G) = 0$, $H^{n-1}(\text{bd } \overline{U}; G)$ contains elements γ not extendable over \overline{U} and for each such γ the set \overline{U} is an $(n-1)$ -cohomology membrane for γ spanned on $\text{bd } \overline{U}$. We assume also that \overline{W}_x is a connected set contractible in a proper subset of X and there is $\alpha_x \in H^{n-1}(\text{bd } \overline{W}_x; G)$ such that \overline{W}_x is an $(n-1)$ -cohomology membrane for α_x spanned on $\text{bd } \overline{W}_x$. We fix $x \in X$ and let \mathcal{B}'_x be the family of all open connected neighborhoods U of x such that $U = \text{int } \overline{U}$ and \overline{U} is contractible in W_x .

Claim 1. For every $U \in \mathcal{B}'_x$ there exists a non-zero $\gamma_U \in H^{n-1}(\text{bd } \overline{U}; G)$ such that γ_U is extendable over $\overline{W}_x \setminus U$ and \overline{U} is an $(n-1)$ -cohomology membrane for γ_U spanned on $\text{bd } \overline{U}$.

Indeed, since \overline{W}_x is an $(n-1)$ -cohomology membrane for α_x spanned on $\text{bd } \overline{W}_x$, α_x is not extendable over \overline{W}_x , but it is extendable over every closed proper subset of \overline{W}_x containing $\text{bd } \overline{W}_x$. So, α_x is extendable to an element $\tilde{\alpha}_x \in H^{n-1}(\overline{W}_x \setminus U; G)$ and the element $\gamma_U = j_{\overline{W}_x \setminus U, \text{bd } \overline{U}}^{n-1}(\tilde{\alpha}_x) \in H^{n-1}(\text{bd } \overline{U}; G)$ is not extendable over \overline{U} (otherwise α_x would be extendable over \overline{W}_x), in particular $\gamma_U \neq 0$. Therefore, according to the choice of W_x , \overline{U} is an $(n-1)$ -cohomology membrane for γ_U spanned on $\text{bd } \overline{U}$.

Let \mathcal{B}''_x be the family of all $U \in \mathcal{B}'_x$ satisfying the following condition: $\text{bd } \overline{U}$ contains a continuum F_U such that $X \setminus F_U$ has exactly two components and F_U is an $(n-1, G)$ -carrier of $j_{\text{bd } \overline{U}, F_U}^{n-1}(\gamma_U)$.

Claim 2. \mathcal{B}''_x is a local base at x .

Since X is arc connected and locally arc connected, there is a convex metric d generating the topology of X , see [27]. We fix $W_0 \in \mathcal{B}'_x$ and for every $\delta > 0$ denote by $B(x, \delta)$ the open ball in X with a center x and a radius δ . There exists $\varepsilon_x > 0$ such that $B(x, \delta) \subset W_0$ for all $\delta \leq \varepsilon_x$. We already observed in the proof of Proposition 2.6 that

any $B(x, \delta)$ is connected and $\text{int}(\overline{B(x, \delta)}) = B(x, \delta)$. Moreover, $\overline{B(x, \delta)}$ is contractible in W_x . Hence, all $U_\delta = B(x, \delta)$, $\delta \leq \varepsilon_x$, belong to \mathcal{B}'_x . Consequently, by Claim 1, for every δ there exists a non-zero $\gamma_\delta \in H^{n-1}(\text{bd } \overline{U}_\delta; G)$ such that \overline{U}_δ is an $(n-1)$ -cohomology membrane for γ_δ spanned on $\text{bd } \overline{U}_\delta$ and γ_δ is extendable over $\overline{W}_x \setminus U_\delta$. By Lemma 2.1(ii), there exists a closed subset F_δ of $\text{bd } \overline{U}_\delta$ which is a carrier of $\gamma_\delta^* = j_{\text{bd } \overline{U}_\delta, F_\delta}^{n-1}(\gamma_\delta)$. Since $n \geq 2$ and F_δ is a carrier of $\gamma_\delta^* \in H^{n-1}(F_\delta; G)$, F_δ is a continuum, see [29, Lemma 2.7]. Let us show that the family $\{F_\delta : \delta \leq \varepsilon_x\}$ is uncountable. Since the function $f: X \rightarrow \mathbb{R}$, $f(y) = d(x, y)$, is continuous and W_0 is connected, $f(W_0)$ is an interval containing $[0, \varepsilon_x]$ and $f^{-1}([0, \varepsilon_x]) = B(x, \varepsilon_x) \subset W_0$. So, $f^{-1}(\delta) = \text{bd } \overline{U}_\delta \neq \emptyset$ for all $\delta \leq \varepsilon_x$. Hence, the family $\{F_\delta : \delta \leq \varepsilon_x\}$ is uncountable and consist of disjoint continua. Moreover, γ_δ^* is a non-zero element of $H^{n-1}(F_\delta; G)$ not extendable over \overline{W}_x because F_U is contractible in \overline{W}_x . Thus, by Lemma 2.5(ii), F_δ separates X . So, each $X \setminus F_\delta$ has at least two components. Then, by Lemma 3.1, there exists $\delta_0 \leq \varepsilon_x$ such that $X \setminus F_{\delta_0}$ has exactly two components. Therefore, $U_{\delta_0} = B(x, \delta_0) \in \mathcal{B}''_x$ and it is contained in W_0 . This completes the proof of Claim 2.

Now, let $\tilde{\mathcal{B}}_x$ be the family of all $U \in \mathcal{B}''_x$ with $H^{n-1}(\text{bd } \overline{U}; G) \neq 0$ such that both U and $X \setminus \overline{U}$ are connected.

Claim 3. $\tilde{\mathcal{B}}_x$ is a local base at x .

Let U_0 be an arbitrary neighborhood of x such that \overline{U}_0 is contractible in W_x . We are going to find a member of $\tilde{\mathcal{B}}_x$ contained in U_0 . To this end, let $\varepsilon = d(x, X \setminus U_0)$. According to Theorem 2.2 there is $\eta > 0$ corresponding to $\varepsilon/2$ and the point x (i.e., for every $y \in X$ with $d(y, x) < \eta$, there exists a homeomorphism $h: X \rightarrow X$ with $h(y) = x$ and $d(h(z), z) < \varepsilon/2$ for all $z \in X$). Now, choose a connected neighborhood W of x with $\overline{W} \subset B(x, \varepsilon/2)$ and $\text{diam}(\overline{W}) < \eta$. Finally, take $U \in \mathcal{B}''_x$ such that \overline{U} is contractible in W . By Claim 2, there exists a continuum $F_U \subset \text{bd } \overline{U}$ such that $X \setminus F_U$ has exactly two components and F_U is an $(n-1, G)$ -carrier for $\gamma_U^* = j_{\text{bd } \overline{U}, F_U}^{n-1}(\gamma_U)$. If $F_U = \text{bd } \overline{U}$, then U is the desired member of $\tilde{\mathcal{B}}_x$. Indeed, since $X \setminus \text{bd } \overline{U} = U \cup X \setminus \overline{U}$ with $U \cap (X \setminus \overline{U}) = \emptyset$ and U is connected, then $X \setminus \overline{U}$ should be also connected (recall that $X \setminus F_U$ has exactly two components).

Suppose that F_U is a proper subset of $\text{bd } \overline{U}$. Because F_U (as a subset of \overline{U}) is contractible in a compact set $\Gamma \subset W$, γ_U^* is not extendable over Γ . Thus, we can apply Lemma 2.5(ii) to conclude that F_U separates \overline{W} . So, $\overline{W} \setminus F_U = V_1 \cup V_2$ for some open, non-empty disjoint subsets $V_1, V_2 \subset \overline{W}$. Since U is a connected subset of $\overline{W} \setminus F_U$, U is contained

in one of the sets V_1, V_2 , say $U \subset V_1$. Hence, $F_U \cup \overline{V}_2 \subset \overline{W}_x \setminus U$. Since γ_U is extendable over $\overline{W}_x \setminus U$ (see Claim 1), γ_U^* is also extendable over $\overline{W}_x \setminus U$, in particular γ_U^* is extendable over $F_U \cup \overline{V}_2$. On the other hand, γ_U^* is not extendable over \overline{W} because F_U is contractible in \overline{W} . The last fact, together with the equality $(F_U \cup \overline{V}_1) \cap (F_U \cup \overline{V}_2) = F_U$, yields that γ_U^* is not extendable over $F_U \cup \overline{V}_1$. Let $\beta = j_{F_U, F'}^{n-1}(\gamma_U^*)$, where $F' = \overline{V}_1 \cap F_U$ (observe that $F' \neq \emptyset$ because \overline{W} is connected). If F' is a proper subset of F_U , then $\beta = 0$ since F_U is a carrier for γ_U^* . So, β would be extendable over \overline{V}_1 , which implies γ_U^* is extendable over $F_U \cup \overline{V}_1$, a contradiction. Therefore, $F' = F_U \subset \overline{V}_1$ and γ_U^* is not extendable over \overline{V}_1 . Consequently, there exists an $(n-1)$ -cohomology membrane $P_\beta \subset \overline{V}_1$ for γ_U^* spanned on F_U . By Lemma 2.5(i), $V = P_\beta \setminus F_U$ is a connected open set in X whose boundary is the set $F'' = X \setminus P_\beta \cap P_\beta \setminus F_U \subset F_U$. As above, using that γ_U^* is not extendable over P_β and $j_{F_U, Q}^{n-1}(\gamma_U^*) = 0$ for any proper closed subset $Q \subset F_U$, we can show that $F'' = F_U$ and $\text{bd } \overline{V} = F_U$. Summarizing the properties of V , we have that \overline{V} is contractible in W_x (because so is \overline{U}_0), $V = \text{int}(\overline{V})$ (because $F_U = \text{bd } \overline{V}$) and V is connected. Moreover, since $X \setminus F_U$ is the union of the open disjoint non-empty sets V and $X \setminus P_\beta$ such that V is connected and $X \setminus F_U$ has exactly two components, $X \setminus \overline{V}$ is also connected. Finally, because F_U is an $(n-1, G)$ -carrier for the nontrivial γ_U^* , $H^{n-1}(\text{bd } \overline{V}; G) \neq 0$. Thus, if V contains x , then V is a member of $\tilde{\mathcal{B}}_x$.

If V does not contain x , we take a point $y \in V$ with $d(x, y) < \eta$. This is possible because $V \subset \overline{W}$ and $\text{diam}(\overline{W}) < \eta$. So, according to the choice of η , there is a homeomorphism h on X such that $h(y) = x$ and $d(z, h(z)) < \varepsilon/2$ for all $z \in X$. Then $h(V) \subset U_0$ (this inclusion follows from the choice of ε and the fact that h is $(\varepsilon/2)$ -close to the identity on X). So, $\overline{h(V)}$ is contractible in W_x . Since the remaining properties from the definition of $\tilde{\mathcal{B}}_x$ are invariant under homeomorphisms, $h(V)$ is the required member of $\tilde{\mathcal{B}}_x$, which provides the proof of Claim 3.

The next claim completes the proof of Proposition 3.2.

Claim 4. $H^{n-1}(\overline{U}; G) = 0$ and $\text{bd } \overline{U}$ is an $(n-1, G)$ -bubble for every $U \in \tilde{\mathcal{B}}_x$.

Because each \overline{U} , $U \in \tilde{\mathcal{B}}_x$, is contractible in \overline{W}_x , any nontrivial $\gamma \in H^{n-1}(\overline{U}; G)$ cannot be extendable over \overline{W}_x . Since \overline{W}_x is contractible in a proper subset of X , by Lemma 2.5(ii), \overline{U} would separate X provided $H^{n-1}(\overline{U}; G) \neq 0$. On the other hand, each $X \setminus \overline{U}$ is connected. Therefore, $H^{n-1}(\overline{U}; G) = 0$ for all $U \in \tilde{\mathcal{B}}_x$. Suppose

there exists a proper closed subset $F \subset \text{bd } \overline{U}$ and a nontrivial element $\alpha \in H^{n-1}(F; G)$. Since $H^{n-1}(\overline{U}; G) = 0$, α is not extendable over \overline{U} . Hence, there is an $(n-1)$ -cohomology membrane $K_\alpha \subset \overline{U}$ for α spanned on F . Therefore, $(K_\alpha \setminus F) \cap \overline{X \setminus K_\alpha} = \emptyset$. In particular, $K_\alpha \setminus F$ is open in $\overline{U} \setminus F$. On the other hand, $K_\alpha \setminus F$ is also closed in $\overline{U} \setminus F$. Because $\overline{U} \setminus F$ is connected (recall that U is a dense connected subset of $\overline{U} \setminus F$), we obtain $K_\alpha = \overline{U}$. Finally, observe that any point from $\text{bd } \overline{U} \setminus F$ belongs to $(K_\alpha \setminus F) \cap \overline{X \setminus K_\alpha}$, a contradiction. Therefore, $\text{bd } \overline{U}$ is an $(n-1, G)$ -bubble. \square

4. PROOF OF THEOREM 1.2 AND COROLLARY 1.3

Proof of Theorem 1.2. (i) Suppose K is an $(n-1)$ -cohomology membrane spanned on A for some $\gamma \in H^{n-1}(A; G)$ and there is a point $a \in (K \setminus A) \cap \overline{X \setminus K}$. Take $V \in \mathcal{B}_a$ with $\overline{V} \cap A = \emptyset$, where \mathcal{B}_a is a local base at a satisfying the hypotheses of Proposition 2.6(2). Since $K \setminus V$ is a proper subset of K containing A , γ is extendable to $\gamma^* \in H^{n-1}(K \setminus V; G)$. Then $\alpha_V = j_{K \setminus V, K \cap \text{bd } \overline{V}}^{n-1}(\gamma^*)$ is not extendable over $K \cap \overline{V}$ (otherwise γ would be extendable over K). In particular, α_V a non-zero element of $H^{n-1}(K \cap \text{bd } \overline{V}; G)$. Because $\text{bd } \overline{V}$ has an empty interior in X , $\dim_G \text{bd } \overline{V} \leq n-1$, see Proposition 2.6(1). Consequently, α_V can be extended to $\tilde{\alpha}_V \in H^{n-1}(\text{bd } \overline{V}; G)$. Moreover, since α_V is not extendable over $K \cap \overline{V}$, $\tilde{\alpha}_V$ is not extendable over \overline{V} . Finally, using that \overline{V} is an $(n-1)$ -cohomology membrane spanned on $\text{bd } \overline{V}$ for $\tilde{\alpha}_V$ and, since $a \in (K \setminus A) \cap \overline{X \setminus K}$ implies that $\text{bd } \overline{V} \cup (K \cap \overline{V})$ is a proper closed subset of \overline{V} containing $\text{bd } \overline{V}$, we can find $\beta \in H^{n-1}(\text{bd } \overline{V} \cup (K \cap \overline{V}); G)$ extending $\tilde{\alpha}_V$. Thus, α_V is extendable over $K \cap \overline{V}$, a contradiction. Therefore, X has the n -cohomology membrane property.

(ii) We need the following result [22, Theorem 1]: For every metric compactum K with $\dim_G K = n$ there exist a closed set $Y \subset K$ and a dense set $D \subset Y$ such that $\dim_G Y = n$ and K has an n -dimensional G -obstruction at every $y \in D$. In our situation we take a compact set $K \subset X$ with $\dim_G K = n$ and find corresponding sets $D \subset Y \subset K$. Since $\dim_G Y = n$, $\text{int} Y \neq \emptyset$ (Proposition 2.6) and there is $y \in D \cap \text{int} Y$. Because K has an n -dimensional G -obstruction at y , we can find a neighborhood $W \subset K$ of y such that for any open in K neighborhood V of y contained in W the homomorphism $H^n(K, K \setminus V; G) \rightarrow H^n(K, K \setminus W; G)$ is not trivial. This implies that for every open in K neighborhoods U, V of y with $\overline{U} \subset V$ the homomorphism $H^n(K, K \setminus U; G) \rightarrow H^n(K, K \setminus V; G)$ is also nontrivial. Since $y \in \text{int} Y$, we can suppose that $W \in \mathcal{B}_y$ such that \overline{W} is contractible in

X and satisfying the hypotheses of Proposition 2.6(2). Then, by the excision axiom, for every $U, V \in \mathcal{B}_y$ with $\bar{U} \subset V \subset \bar{V} \subset W$, the groups $H^n(K, K \setminus U; G)$ and $H^n(K, K \setminus V; G)$ are isomorphic to $H^n(X, X \setminus U; G)$ and $H^n(X, X \setminus V; G)$, respectively. So, $j_{U,V} : H^n(X, X \setminus U; G) \rightarrow H^n(X, X \setminus V; G)$ is a nontrivial homomorphism for all $U, V \in \mathcal{B}_y$ with $\bar{U} \subset V \subset \bar{V} \subset W$. Finally, because X is locally homogeneous, it has an n -dimensional G -obstruction at every $x \in X$.

To prove the second half of condition (ii), let $U, V \in \mathcal{B}_x$ with $\bar{U} \subset V \subset \bar{V} \subset W$.

Claim 5. The homomorphism $j_{\bar{W} \setminus U, \bar{W} \setminus V}^{n-1} : H^{n-1}(\bar{W} \setminus U; G) \rightarrow H^{n-1}(\bar{W} \setminus V; G)$ is surjective and $H^{n-1}(\bar{W} \setminus V; G) \neq 0$.

Indeed, consider the Mayer-Vietoris exact sequence below, where the coefficient group G is suppressed,

$$H^{n-1}(\bar{W} \setminus U) \xrightarrow{\varphi} H^{n-1}(\bar{V} \setminus U) \oplus H^{n-1}(\bar{W} \setminus V) \xrightarrow{\psi} H^{n-1}(\text{bd } \bar{V}) \dots$$

The maps φ and ψ are defined by $\varphi(\gamma) = (j_{\bar{W} \setminus U, \bar{V} \setminus U}^{n-1}(\gamma), j_{\bar{W} \setminus U, \bar{W} \setminus V}^{n-1}(\gamma))$ and $\psi((\beta, \alpha)) = j_{\bar{V} \setminus U, \text{bd } \bar{V}}^{n-1}(\beta) - j_{\bar{W} \setminus V, \text{bd } \bar{V}}^{n-1}(\alpha)$. For every $\alpha \in H^{n-1}(\bar{W} \setminus V; G)$ the element $\beta'_\alpha = j_{\bar{W} \setminus V, \text{bd } \bar{V}}^{n-1}(\alpha) \in H^{n-1}(\text{bd } \bar{V}; G)$ is extendable to an element $\beta_\alpha \in H^{n-1}(\bar{V} \setminus U; G)$. Indeed, there are two possibilities: either β'_α is extendable over \bar{V} or it is not extendable over \bar{V} . The first case obviously implies that β'_α is extendable over $\bar{V} \setminus U$. In the second case, by Proposition 2.6(2), \bar{V} is an $(n-1)$ -cohomological membrane spanned on $\text{bd } \bar{V}$ for β'_α . Then, since $\bar{V} \setminus U$ is a proper closed subset of \bar{V} , β'_α is extendable over $\bar{V} \setminus U$. Hence, $\psi(\beta_\alpha, \alpha) = 0$ for any $\alpha \in H^{n-1}(\bar{W} \setminus V; G)$. Consequently, there is $\gamma_\alpha \in H^{n-1}(\bar{W} \setminus U; G)$ with $\varphi(\gamma_\alpha) = (\beta_\alpha, \alpha)$. In particular, $j_{\bar{W} \setminus U, \bar{W} \setminus V}^{n-1}(\gamma_\alpha) = \alpha$, which shows the surjectivity of $j_{\bar{W} \setminus U, \bar{W} \setminus V}^{n-1}$. The nontriviality of $H^{n-1}(\bar{W} \setminus V; G)$ follows from the Mayer-Vietoris exact sequence

$$\rightarrow H^{n-1}(\bar{V}; G) \oplus H^{n-1}(\bar{W} \setminus V; G) \rightarrow H^{n-1}(\text{bd } \bar{V}; G) \rightarrow H^n(\bar{W}; G) \rightarrow$$

Indeed, \bar{W} being contractible in X implies that $H^n(\bar{W}; G) = 0$ (see the remark after Proposition 2.3). Hence, the homomorphism $j_{\bar{V}, \text{bd } \bar{V}}^{n-1}$ would be surjective provided $H^{n-1}(\bar{W} \setminus V; G) = 0$. This would imply that every non-trivial element of $H^{n-1}(\text{bd } \bar{V}; G)$ is extendable over \bar{V} , which contradicts Proposition 2.6(2).

To complete the proof, consider the commutative diagram whose rows are parts of exact sequences

$$\begin{array}{ccccc}
H^{n-1}(\overline{W}\setminus U; G) & \xrightarrow{\delta_U} & H^n(\overline{W}, \overline{W}\setminus U; G) & \xrightarrow{i_U} & H^n(\overline{W}; G) \\
\downarrow j_{\overline{W}\setminus U, \overline{W}\setminus V}^{n-1} & & \downarrow j'_{U, V} & & \downarrow \text{id} \\
H^{n-1}(\overline{W}\setminus V; G) & \xrightarrow{\delta_V} & H^n(\overline{W}, \overline{W}\setminus V; G) & \xrightarrow{i_V} & H^n(\overline{W}; G).
\end{array}$$

Since $H^n(\overline{W}; G) = 0$, both δ_U and δ_V are surjective. This, combined with the surjectivity of $j_{\overline{W}\setminus U, \overline{W}\setminus V}^{n-1}$ and nontriviality of $H^{n-1}(\overline{W}\setminus V; G)$, implies that $j'_{U, V}$ is also surjective. Finally, by the excision axiom the groups $H^n(\overline{W}, \overline{W}\setminus U; G)$ and $H^n(\overline{W}, \overline{W}\setminus V; G)$ are isomorphic to $H^n(X, X\setminus U; G)$ and $H^n(X, X\setminus V; G)$, respectively. Therefore, the homomorphism $j_{U, V}^n : H^n(X, X\setminus U; G) \rightarrow H^n(X, X\setminus V; G)$ is surjective.

Proof of Corollary 1.3. The conclusion of this corollary implies the well-known invariance of domains property, which was established in [23] and [25], respectively, for compact homogeneous and locally homogeneous *ANR*-spaces X with $\dim X < \infty$. For homogeneous *ANR*-compacta X with $\dim_G X < \infty$, where G is a countable principal ideal domain, it was established in [29]. The arguments from [29] also work in our situation.

5. PROOF OF THEOREM 1.4.

(i) It suffices to show that the one-point compactification $bX = X \cup \{b\}$ of X is dimensionally full-valued. To this end, we use the following result of Dranishnikov [7, Theorem 12.3-12.4]: If Y is a finite-dimensional *ANR*-compactum, then $\dim_{\mathbb{Z}_{(p)}} Y = \dim_{\mathbb{Z}_p} Y$ for all prime p . Moreover, $\dim_{\mathbb{Q}} Y \leq \dim_G Y$ for any group $G \neq 0$ and there exists a prime number p with $\dim Y = \dim_{\mathbb{Z}_{(p)}} Y = \dim_{\mathbb{Z}_p} Y$. The proof presented in [7] works also when Y is the one-point compactification of an *ANR*-space. Here $\mathbb{Z}_p = \mathbb{Z}/p\mathbb{Z}$ is the cyclic group and $\mathbb{Z}_{(p)} = \{m/n : n \text{ is not divisible by } p\} \subset \mathbb{Q}$, where \mathbb{Q} is the field of rational numbers. We also consider the quotient group $\mathbb{Z}_{p^\infty} = \mathbb{Q}/\mathbb{Z}_{(p)}$. It is well known [22] that the so called Bockstein basis consists of the groups $\sigma = \{\mathbb{Q}, \mathbb{Z}_p, \mathbb{Z}_{(p)}, \mathbb{Z}_{p^\infty} : p \in \mathcal{P}\}$, \mathcal{P} is the set of all primes. For any group (not necessarily countable) there exists a collection $\sigma(G) \subset \sigma$ such that $\dim_G X = \sup\{\dim_H X : H \in \sigma(G)\}$ for any space X .

Let X be a locally homogeneous *ANR*-space with $\dim X = n$. According to the mentioned above Dranishnikov's result, there exists $p \in \mathcal{P}$ with $\dim_{\mathbb{Z}_{(p)}} bX = \dim_{\mathbb{Z}_p} bX = n$. If $\dim_{\mathbb{Z}_{p^\infty}} bX = n$, we are

done. Indeed, according to [7, Lemma 2.6], bX is p -regular, i.e.

$$\dim_{\mathbb{Z}_{(p)}} bX = \dim_{\mathbb{Z}_{p^\infty}} bX = \dim_{\mathbb{Z}_p} bX = \dim_{\mathbb{Q}} bX = n.$$

Then, applying again Dranishnikov's result [7, Theorem 12.3], we obtain $\dim_{\mathbb{Q}} bX \leq \dim_G bX \leq n$ for any group $G \neq 0$. Hence, $\dim_G bX = \dim bX = n$ for all nontrivial groups G , and by [21, Theorem 11], bX is dimensionally full-valued. Therefore, the next claim completes the proof.

Claim 6. If $\dim_{\mathbb{Z}_{(p)}} bX = \dim_{\mathbb{Z}_p} bX = n$ for some prime number p , then $\dim_{\mathbb{Z}_{p^\infty}} bX = n$.

Suppose $\dim_{\mathbb{Z}_{p^\infty}} bX \leq n-1$. According to the Bockstein inequalities (see, for example [7], or [21]), we have $\dim_{\mathbb{Z}_p} bX \leq \dim_{\mathbb{Z}_{p^\infty}} bX + 1$, which in our case implies $\dim_{\mathbb{Z}_{p^\infty}} bX = \dim_{\mathbb{Z}_{p^\infty}} X = n-1$. Since $\dim_{\mathbb{Z}_p} (bX)^2 = 2 \dim_{\mathbb{Z}_p} X = 2n$, using the next equality (see [21])

$$\dim_{\mathbb{Z}_{p^\infty}} (bX)^2 = \max\{2 \dim_{\mathbb{Z}_{p^\infty}} bX, \dim_{\mathbb{Z}_p} (bX)^2 - 1\},$$

we obtain $\dim_{\mathbb{Z}_{p^\infty}} X^2 = \dim_{\mathbb{Z}_{p^\infty}} (bX)^2 = 2n-1$. Then, by Proposition 2.6(2), for every $(x, y) \in X^2$ there are neighborhoods U_x and V_y of x and y with compact closures such that $W = U \times V \subset U_x \times V_y$ yields $H^{2n-2}(\text{bd } \overline{W}; \mathbb{Z}_{p^\infty}) \neq 0$ for any $W \in \mathcal{B}_{(x,y)}$. Since $\text{bd } \overline{W} = (\overline{U} \times \text{bd } \overline{V}) \cup (\text{bd } \overline{U} \times \overline{V})$, we have the Meyer-Vietoris exact sequence (in all exact sequences below the coefficient groups \mathbb{Z}_{p^∞} are suppressed)

$$H^{2n-3}(\Gamma_U \times \Gamma_V) \rightarrow H^{2n-2}(\Gamma_W) \rightarrow H^{2n-2}(\overline{U} \times \Gamma_V) \oplus H^{2n-2}(\Gamma_U \times \overline{V}) \rightarrow$$

Here, $\Gamma_U = \text{bd } \overline{U}$, $\Gamma_V = \text{bd } \overline{V}$ and $\Gamma_W = \text{bd } \overline{W}$. The Künneth formulas provide the following exact sequences

$$\sum_{i+j=2n-2} H^i(\overline{U}) \otimes H^j(\Gamma_V) \rightarrow H^{2n-2}(\overline{U} \times \Gamma_V) \rightarrow \sum_{i+j=2n-1} H^i(\overline{U}) * H^j(\Gamma_V),$$

$$\sum_{i+j=2n-2} H^i(\Gamma_U) \otimes H^j(\overline{V}) \rightarrow H^{2n-2}(\Gamma_U \times \overline{V}) \rightarrow \sum_{i+j=2n-1} H^i(\Gamma_U) * H^j(\overline{V})$$

and

$$\sum_{i+j=2n-3} H^i(\Gamma_U) \otimes H^j(\Gamma_V) \rightarrow H^{2n-3}(\Gamma_U \times \Gamma_V) \rightarrow \sum_{i+j=2n-2} H^i(\Gamma_U) * H^j(\Gamma_V).$$

Since $\dim_{\mathbb{Z}_p} X = n-1$, by Proposition 2.6(2), we can assume that the groups $H^{n-1+i}(\Gamma_U; \mathbb{Z}_{p^\infty})$, $H^{n-1+i}(\overline{U}; \mathbb{Z}_{p^\infty})$, as well as the groups $H^{n-1+i}(\Gamma_V; \mathbb{Z}_{p^\infty})$ and $H^{n-1+i}(\overline{V}; \mathbb{Z}_{p^\infty})$ are trivial for all $U \subset U_x$, $V \subset V_y$ and $i \geq 0$. Hence, the groups $H^{2n-2}(\overline{U} \times \Gamma_V; \mathbb{Z}_{p^\infty})$, $H^{2n-2}(\Gamma_U \times \overline{V}; \mathbb{Z}_{p^\infty})$ and $H^{2n-3}(\Gamma_U \times \Gamma_V; \mathbb{Z}_{p^\infty})$ are also trivial, which implies the triviality of

the group $H^{2n-2}(\Gamma_W; Z_{p^\infty})$, a contradiction. Therefore, $\dim Z_{p^\infty} bX = n$.

(ii) To prove the second half of Theorem 1.4, denote by \widehat{H}_* the exact homology developed in [26] for locally compact spaces. The homological dimension $h \dim_G Y$ of a space Y is the largest number n such that $\widehat{H}_n(Y, \Phi; G) \neq 0$ for some closed set $\Phi \subset Y$. According to [15] and [26], $h \dim_G Y$ is the greatest m such that the local homology group $\widehat{H}_m(Y, Y \setminus y; G) = \varinjlim_{y \in U} \widehat{H}_m(Y, Y \setminus U; G)$ is not trivial for some $y \in Y$. Moreover, for any field F we have $h \dim_F Y = \dim_F Y$, see [15]. Therefore, since $\dim_{\mathbb{Q}} X = \dim X = n$, we have $h \dim_{\mathbb{Q}} X = n$ and $\widehat{H}_n(X, X \setminus x; \mathbb{Q}) \neq 0$ for all $x \in X$. This means that every $x \in X$ has a neighborhood W such that $\widehat{H}_n(X, X \setminus U; \mathbb{Q}) \neq 0$ for all open sets $U \subset W$ containing x . We assume that W satisfies conditions (1) – (3) from Theorem 1.1 such that \overline{W} is contractible in a proper set in X and $H^{n-1}(\text{bd } \overline{U}; \mathbb{Z}) \neq 0$ for all neighborhoods U of x with $U \subset W$, see Proposition 2.6(ii). Let V be a neighborhood of x with $\overline{V} \subset W$.

Claim 7. The pair $\overline{W} \setminus V \subset \overline{W}$ is an $(n-1)$ -homology membrane spanned on $\overline{W} \setminus V$ for any nontrivial $\gamma \in H_{n-1}(\overline{W} \setminus V; \mathbb{Q})$.

Since $\widehat{H}_n(X, X \setminus V; \mathbb{Q}) \neq 0$, by the excision axiom $\widehat{H}_n(\overline{W}, \overline{W} \setminus V; \mathbb{Q}) \neq 0$. Consider the exact sequences (see [26] for the existence of such sequences)

$$\rightarrow \widehat{H}_n(\overline{W}; \mathbb{Q}) \rightarrow \widehat{H}_n(\overline{W}, \overline{W} \setminus V; \mathbb{Q}) \rightarrow \widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q}) \rightarrow \widehat{H}_{n-1}(\overline{W}; \mathbb{Q})$$

and

$$0 \rightarrow \text{Ext}(H^{n+1}(\overline{W}; \mathbb{Z}), \mathbb{Q}) \rightarrow \widehat{H}_n(\overline{W}; \mathbb{Q}) \rightarrow \text{Hom}(H^n(\overline{W}; \mathbb{Z}), \mathbb{Q}) \rightarrow 0$$

$$0 \rightarrow \text{Ext}(H^n(\overline{W}; \mathbb{Z}), \mathbb{Q}) \rightarrow \widehat{H}_{n-1}(\overline{W}; \mathbb{Q}) \rightarrow \text{Hom}(H^{n-1}(\overline{W}; \mathbb{Z}), \mathbb{Q}) \rightarrow 0.$$

Since $H^{n+1}(\overline{W}; \mathbb{Z}) = H^n(\overline{W}; \mathbb{Z}) = H^{n-1}(\overline{W}; \mathbb{Z}) = 0$, both $\widehat{H}_n(\overline{W}; \mathbb{Q})$ and $\widehat{H}_{n-1}(\overline{W}; \mathbb{Q})$ are trivial. Therefore, the group $\widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q})$ is nontrivial. Moreover, the exact sequence

$$\text{Ext}(H^n(\overline{W} \setminus V; \mathbb{Z}), \mathbb{Q}) \rightarrow \widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q}) \rightarrow \text{Hom}(H^{n-1}(\overline{W} \setminus V; \mathbb{Z}), \mathbb{Q}) \rightarrow 0$$

shows that $\widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q})$ is isomorphic to $\text{Hom}(H^{n-1}(\overline{W} \setminus V; \mathbb{Z}), \mathbb{Q})$ because $H^n(\overline{W} \setminus V; \mathbb{Z}) = 0$ according to the remark after by Proposition 2.3.

To complete the proof of Claim 7, we need to show that if B is a proper closed subset of \overline{W} containing $\overline{W} \setminus V$, then $i_{\overline{W} \setminus V, B}^{n-1}(\gamma) \neq 0$ for every nontrivial $\gamma \in H_{n-1}(\overline{W} \setminus V; \mathbb{Q})$. To this end, consider the exact

sequence

$$0 \rightarrow \text{Ext}(H^n(B; \mathbb{Z}), \mathbb{Q}) \rightarrow \widehat{H}_{n-1}(B; \mathbb{Q}) \rightarrow \text{Hom}(H^{n-1}(B; \mathbb{Z}), \mathbb{Q}) \rightarrow 0.$$

Since B is contractible in X (as a subset of W), $H^n(B; \mathbb{Z}) = 0$. Hence, $\widehat{H}_{n-1}(B; \mathbb{Q})$ is isomorphic to $\text{Hom}(H^{n-1}(B; \mathbb{Z}), \mathbb{Q})$. Because $B = \overline{W} \setminus \Gamma$ for some open set $\Gamma \subset V$, passing to a smaller subset of Γ , we can assume that Γ is a neighborhood of some $y \in V$ such that $\overline{\Gamma} \subset V$ and Γ satisfies conditions (1) – (3) from Theorem 1.1. Then, the arguments from the proof of Claim 5 show that $H^{n-1}(B; \mathbb{Z}) = H^{n-1}(\overline{W} \setminus \Gamma; \mathbb{Z})$ is not trivial and there is a surjective homomorphism $H^{n-1}(B; \mathbb{Z}) \rightarrow H^{n-1}(\overline{W} \setminus V; \mathbb{Z})$. Therefore, there exists an injective homomorphism from $\widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q})$ into $\widehat{H}_{n-1}(B; \mathbb{Q})$. Because we already saw that $\widehat{H}_{n-1}(\overline{W}; \mathbb{Q}) = 0$, this shows that $\overline{W} \setminus V \subset \overline{W}$ is an $(n-1)$ -homology membrane spanned on $\overline{W} \setminus V$ for any nontrivial $\gamma \in \widehat{H}_{n-1}(\overline{W} \setminus V; \mathbb{Q})$, but with respect to the exact homology \widehat{H}_* . To complete the proof of Claim 7, we observe that for every pair $A \subset Y$ of a space Y and its closed set A , any integer k and a group G there is an epimorphism $T_{Y,A}^k : \widehat{H}_k(Y, A; G) \rightarrow H_k(Y, A; G)$, which is an isomorphism when G is a field, see [26, Theorem 4].

Suppose U is a neighborhood of x with $\overline{U} \subset V$. Since the pair $\overline{W} \setminus V \subset \overline{W}$ is an $(n-1)$ -homology membrane spanned on $\overline{W} \setminus V$ for any nontrivial $\gamma \in H_{n-1}(\overline{W} \setminus V; \mathbb{Q})$, according to [1], $H_{n-1}(\text{bd } \overline{U}; \mathbb{Q}) \neq 0$ and \overline{U} is an $(n-1)$ -homology membrane spanned on $\text{bd } \overline{U}$ for some nontrivial $\gamma' \in H_{n-1}(\text{bd } \overline{U}; \mathbb{Q})$. Most important for us is the nontriviality of $H_{n-1}(\text{bd } \overline{U}; \mathbb{Q})$ because it implies also the nontriviality of $H_{n-1}(\text{bd } \overline{U}; \mathbb{Z})$, see [28, Proposition 4.5]. Finally, since $\dim \text{bd } \overline{U} = n-1$, [18, Corollary] shows that $\text{bd } \overline{U}$ is dimensionally full-valued. \square

Acknowledgments. The author would like to express his gratitude to A. Dranishnikov, K. Kawamura, A. Koyama and E. Tymchatyn for their helpful comments.

REFERENCES

- [1] R. H. Bing and K. Borsuk, *Some remarks concerning topological homogeneous spaces*, Ann. of Math. **81** (1965), no. 1, 100–111.
- [2] K. Borsuk, *Theory of retracts*, Monografie Matematyczne **44**, PWN, Warsaw, 1967.
- [3] J. Bryant, *Reflections on the Bing-Borsuk conjecture*, Abstract of the talks presented at the 19th Annual Workshop in Geometric Topology, June 13-15, 2002, 2-3.
- [4] J. Bryant and S. Ferry, *An alpha approximation theorem for homology manifolds*, preprint.

- [5] M. Cárdenas, F. Lasheras, A. Quintero and D. Repovš, *On manifolds with nonhomogeneous factors*, Centr. Eur. J. Math. **10** (2012), 857–862.
- [6] J. Choi, *Properties of n -bubbles in n -dimensional compacta and the existence of $(n - 1)$ -bubbles in n -dimensional clc^n compacta*, Topology Proc. **23** (1998), 101–120.
- [7] A. Dranishnikov, *Cohomological dimension theory of compact metric spaces*, Topology Atlas invited contributions, <http://at.yorku.ca/topology.taic.html> (see also arXiv:math/0501523).
- [8] A. Dranishnikov, *On a problem of P.S. Alexandrov*, Math. USSR Sbornik **63** (1988), 412–426.
- [9] A. Dranishnikov, *Homological dimension theory*, Russian Math. Surveys **43** (1988), 11–63.
- [10] J. Dydak, *Cohomological dimension and metrizable spaces*, Trans. Amer. Math. Soc. **337** (1993), 219–234.
- [11] E. G. Effros, *Transformation groups and C^* -algebras*, Ann. of Math. **81** (1965), 38–55.
- [12] S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton Univ. Press, Princeton, New Jersey, 1952.
- [13] V. Fedorchuk, *On homogeneous Pontryagin surfaces*, Dokl. Akad. Nauk **404** (2005), 601–603 (in Russian).
- [14] W. Jacobsche, *The Bing-Borsuk conjecture is stronger than Poincaré conjecture*, Fund. Math. **106** (1980), 127–134.
- [15] A. Harlap, *Local homology and cohomology, homological dimension, and generalized manifolds*, Mat. Sb. **96(138)** (1975), 347–373 (in Russian).
- [16] W. Hurewicz and H. Wallman, *Dimension theory*, Princeton University Press, Princeton, 1948.
- [17] P. Huber, *Homotopical cohomology and Čech cohomology*, Math. Annalen **144** (1961), 73–76.
- [18] Y. Kodama, *On a problem of Alexandroff concerning the dimension of product spaces II*, J. Math. Soc. Japan **11** (1959), 94–111.
- [19] Y. Kodama, *Note on an absolute neighborhood extensor for metric spaces*, J. Math. Soc. Japan **8** (1956), 206–215.
- [20] P. Krupski, *Recent results on homogeneous curves and ANR's*, Topology Proc. **16** (1991), 109–118.
- [21] V. Kuz'minov, *Homological dimension theory*, Russian Math. Surveys **23** (1968), no. 1, 1–45.
- [22] V. Kuz'minov, *On continua V^n* , Dokl. Akad. Nauk SSSR **139** (1961), no. 1, 24–27 (in Russian).
- [23] J. Lysko, *Some theorems concerning finite-dimensional homogeneous ANR-spaces*, Bull. Polish Acad. Sci. **24** (1976), 491–496.
- [24] D. Montgomery, *Local homogeneous spaces*, Ann. of Math. **52** (1950), 261–271.
- [25] H. Seidel, *Locally homogeneous ANR-spaces*, Aech. Math. **44** (1985), 79–81.
- [26] E. Sklyarenko, *Homology theory and the exactness axiom*, Uspehi Mat. Nauk **24** (1969) no. 5(149), 87–140 (in Russian).
- [27] A. Tominaga and T. Tanaka, *Convexification of locally connected generalized continua*, J. Sci. Hiroshima Univ. Ser. A **19** (1955), 301–306.

- [28] V. Valov, *Homological dimension and homogeneous ANR spaces*, Top. Appl. **221** (2017), 38–50.
- [29] V. Valov, *Local cohomological properties of homogeneous ANR compacta*, Fund. Math. **223** (2016), 257–270.
- [30] V. Valov, *Homogeneous ANR-spaces and Alexandroff manifolds*, Top. Appl. **173** (2014), 227–233.

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. BOX 5002, NORTH BAY, ON, P1B 8L7, CANADA

Email address: `veskov@nipissingu.ca`