ON THE GENERAL RANKS OF QP REPRESENTATIONS

JIARUI FEI

ABSTRACT. We propose an algorithm to compute the general rank from a principal component $PC(\delta)$ of representations to another one $PC(\epsilon)$ for a quiver with potential. We give sufficient conditions for the algorithm to work. In particular, the algorithm works effectively if any of δ and ϵ is reachable. We find an explicit mutation formula for the general rank, and discover several related mutation invariants.

1. INTRODUCTION

Schofield introduced the general rank for quiver representations in his theory of general representations [23]. It was further pursued by Crawley-Boevey in [3]. Let Q be a finite quiver without oriented cycles, and α, β be some dimension vectors of Q. There is an open subset Uof $\operatorname{rep}_{\alpha}(Q) \times \operatorname{rep}_{\beta}(Q)$ and a dimension vector γ such that for all $(M, N) \in U$, $\operatorname{Hom}_Q(M, N)$ has the minimal dimension and $\{\phi \in \operatorname{Hom}_Q(M, N) \mid \operatorname{rank} \phi = \gamma\}$ is open and non-empty in $\operatorname{Hom}_Q(M, N)$. The dimension vector γ is called the general rank from $\operatorname{rep}_{\alpha}(Q)$ to $\operatorname{rep}_{\beta}(Q)$.

A remarkable property of γ is the following

$$\operatorname{ext}_Q(\alpha,\beta) = -\langle \alpha - \gamma, \beta - \gamma \rangle_Q = \operatorname{ext}_Q(\alpha - \gamma, \beta - \gamma).$$
(1.1)

where $\operatorname{ext}_Q(\alpha,\beta)$ is the generic (minimal) value of $\operatorname{dim}\operatorname{Ext}_Q^1(M,N)$ on $\operatorname{rep}_{\alpha}(Q) \times \operatorname{rep}_{\beta}(Q)$, and $\langle -, -\rangle_Q$ is the Euler form of Q. Schofield further used this property to deduce an algorithm to compute $\operatorname{ext}_Q(\alpha,\beta)$. This algorithm is one of the key ingredients in Derksen-Weyman's proof of the saturation conjecture [4]. The algorithm was reformulated in terms of tropical F-polynomials and generalized to finite-dimensional algebras in [9].

Fock and Goncharov formulated their duality pairing conjecture for cluster varieties in [11]. We gave their duality pairing a representation-theoretic interpretation in [9]. We show that the duality pairing holds for generic bases if $f_{\check{\epsilon}}(\delta) = \check{f}_{\delta}(\check{\epsilon})$ holds for any pair $(\delta, \check{\epsilon})$ of weight vectors in \mathbb{Z}^{Q_0} , where $f_{\check{\epsilon}}$ and \check{f}_{δ} are the generic tropical *F*-polynomials (see Appendix for the definitions) of a relevant quiver with potential (Q, S). As pointed out in [9], more optimistically one may expect that

$$f_{\check{\epsilon}}(\delta) = f_{\delta}(\check{\epsilon}) = \hom(\delta,\check{\epsilon}) \text{ for any pair } (\delta,\check{\epsilon}).$$

This turns out to be equivalent to the saturation conjecture for nondegenerate quivers with potentials (see Definition 7.2 and Proposition 7.3).

One would hope by studying the general ranks of representations of quivers with potentials, the duality pairing conjecture and the saturation conjecture for nondegenerate QPs can be settled. This is our original motivation for this study. Let us first explain what we mean by the general ranks in the setting of quivers with potentials. The definition of the general rank has a straightforward generalization (see Lemma 5.1) if we replace the representation spaces of a quiver Q by some irreducible components in the representation variety of any basic algebra,

²⁰²⁰ Mathematics Subject Classification. Primary 16G10; Secondary 13F60.

Key words and phrases. General Rank, Representation, Quiver with Potential, Mutation, Jacobian Algebra.

in particular the Jacobian algebra J of (Q, \mathcal{S}) [5]. For quivers with potentials, the irreducible components of interests for us are those hit by general presentations [7]. More precisely, for any $\delta \in \mathbb{Z}^{Q_0}$ we define the presentation space

$$\operatorname{PHom}_J(\delta) := \operatorname{Hom}_J(P([-\delta]_+), P([\delta]_+)).$$

Here $P(\beta) = \bigoplus_{u \in Q_0} \beta(u) P_u$ and P_u is the indecomposable projective representation corresponding to u. The vector δ is called the weight vector or the δ -vector of the presentation space. There is an open subset U of $PHom_J(\delta)$ such that the cokernels of presentations in U lie in an irreducible component $PC(\delta)$ of the representation variety of J (see Definition 3.6). A general representation in $PC(\delta)$ is called a general representation of weight δ . As shown in [21], such a component is exactly the *strongly reduced component* introduced in [13]. The general rank we consider in this article is the one from the component $PC(\delta)$ to another one $PC(\epsilon)$, denoted by $rank(\delta, \epsilon)$.

However, we find it difficult to prove a variation of (1.1) in the setting of quivers with potentials (see Question 5.23). Most of geometric arguments in Schofield's proof break down here, which seems to be the largest obstacle in generalizing Schofield's theory. On the other hand, we get some new weapon from the theory of cluster algebras and quivers with potentials. Thanks to Derksen-Weyman-Zelevinsky's mutation, we are able to give an algorithm for computing the general ranks. Our algorithm is expected to work for any pair (δ, ϵ) but we can only show it works under some conditions. Those conditions are trivially satisfied if one of δ and ϵ is *extended-reachable*, that is, can be obtained by a sequence of mutations and τ . The sufficient conditions are close related to a conjecture, which implies the duality pairing conjecture and the saturation conjecture for nondegenerate QPs.

For any weight vector ϵ of (Q, S), we define a pair of operators, which plays an important role not only in our algorithm but also in the crystal structure of upper cluster algebras [10].

Definition 1.1. For any weight vector ϵ of (Q, S), we define the two operators r_{ϵ} and l_{ϵ} on the set of weight vectors of (Q, S) as follows:

$$r_{\epsilon}(\delta) = \delta + \epsilon + \operatorname{rank}(\epsilon, \tau \delta)B;$$

$$l_{\epsilon}(\delta) = \delta - \check{\epsilon} + \operatorname{rank}(\delta, \epsilon)B,$$

where τ (defined before Theorem 3.11) is related to the AR-translation and B is the skew-symmetric matrix attached to Q.

By the mutation μ_u of δ for $u \in Q_0$, we mean the formula (3.1). To make these mutation operators enter the picture, we study when the operators r_{ϵ} and l_{ϵ} commute with mutations. We say (δ, ϵ) has completely extremal rank if any of the following occurs:

$$\operatorname{rank}(\delta, \epsilon) = 0, \quad \operatorname{rank}(\delta, \epsilon) = \underline{\dim}(\delta), \quad \operatorname{rank}(\delta, \epsilon) = \underline{\dim}(\epsilon),$$

where $\underline{\dim}(\delta)$ denotes the dimension vector of a general representation in $PC(\delta)$. It turns out that if there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ has completely extremal rank, then $r_{\epsilon}(\delta)$ and $l_{\epsilon}(\delta)$ commute with any sequence of mutations and τ^{i} (Lemmas 5.7, 5.9 and 5.11). This would imply the following theorem.

Theorem 1.2 (Theorem 5.16). Let ϵ and δ be two weight vectors for a quiver with potential (Q, S).

(1) Suppose that there is an extended sequence of mutation $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\epsilon), \mu_{\mathbf{u}}(\tau\delta))$ has completely extremal rank r. Then there is an exact sequence

$$\cdots \to \tau^{-1}M \to \tau^{-1}R \to \tau^{-1}E \xrightarrow{h_{-1}} M \to R \to E \xrightarrow{h_0} \tau M \to \tau R \to \tau E \xrightarrow{h_1} \tau^2 M \to \cdots,$$

where R is general of weight $r_{\epsilon}(\delta)$, (E, M) is general as a pair of weights δ and ϵ , and h_i is a general homomorphism in $\text{Hom}_J(\tau^i E, \tau^{i+1}M)$.

- (2) Suppose that there is an extended sequence of mutation $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ has completely extremal rank r. Then there is an exact sequence
- $\cdots \to \tau^{-1}L \to \tau^{-1}M \xrightarrow{g_{-1}} \tau^{-1}E \to L \to M \xrightarrow{g_0} E \to \tau L \to \tau M \xrightarrow{g_1} \tau E \to \tau^2 L \to \cdots,$

where L is general of weight $l_{\epsilon}(\delta)$, (M, E) is general as a pair of weight δ and ϵ , and g_i is a general homomorphism in $\text{Hom}_J(\tau^i M, \tau^i E)$.

Here, (E, M) is general as a pair of weight δ and ϵ means that (E, M) can be chosen in an open subset of $PC(\delta) \times PC(\epsilon)$.

If such a sequence of mutations exists (conjecturally always exists), then we get an algorithm to compute the general rank from δ to ϵ (see Algorithm 5.20 for details). The algorithm leads to the following explicit mutation formula for the general rank.

Theorem 1.3 (Theorem 6.1). Let $\gamma_r = \operatorname{rank}(\epsilon, \tau \delta)$ and $\gamma_l = \operatorname{rank}(\delta, \epsilon)$. We denote $\gamma'_r = \operatorname{rank}(\epsilon', \tau \delta')$ and $\gamma'_l = \operatorname{rank}(\delta', \epsilon')$, where $\delta' = \mu_u(\delta)$ and $\epsilon' = \mu_u(\epsilon)$.

(1) If r_{ϵ} commutes with μ_u , then

$$\gamma_r'(v) = \begin{cases} \gamma_r(v) & \text{for all } v \neq u \\ \gamma_r[b_u]_+ - \gamma_r(u) + [\delta(u)]_+ + [\epsilon(u)]_+ - [r_\epsilon(\delta)(u)]_+ & \text{for all } v = u \end{cases}$$

(2) If l_{ϵ} commutes with μ_u , then

$$\gamma_l'(v) = \begin{cases} \gamma_l(v) & \text{for all } v \neq u, \\ \gamma_l[b_u]_+ - \gamma_l(u) + [\delta(u)]_+ + [-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+ & \text{for all } v = u. \end{cases}$$

With these explicit formulas, we find some interesting mutation-invariants.

Proposition 1.4 (Proposition 6.3). Suppose that l_{ϵ} and r_{ϵ} commutes the mutation μ_u . Let $\gamma_r = \operatorname{rank}(\epsilon, \tau \delta)$ and $\gamma_l = \operatorname{rank}(\delta, \epsilon)$. Then the following are mutation-invariant:

$$h_l(\delta, \epsilon) := \check{\epsilon}(\gamma_l) - \hom(\delta, \epsilon) + \hom(l_{\epsilon}(\delta), \epsilon);$$

$$e_r(\delta, \epsilon) := \epsilon(\gamma_r) - e(\delta, \epsilon) + e(r_{\epsilon}(\delta), \epsilon).$$

Here hom (δ, ϵ) is the generic value of dim Hom $_J(M, N)$ for $(M, N) \in PC(\delta) \times PC(\epsilon)$, and $e(\delta, \epsilon)$ is certain generic value on PHom $_J(\delta) \times PHom_J(\epsilon)$ introduced in [7] (see also below Lemma 3.1).

If ϵ is extended-reachable, then Theorem 5.16 and Algorithm 5.20 take a particularly nice form (Corollaries 5.27 and 5.28). In this case, the operators r_{ϵ} and l_{ϵ} are inverse of each other (Proposition 5.30). We also find an interesting property of these operators related to the canonical decomposition of presentations [7]. We write $\epsilon = \epsilon_1 \oplus \epsilon_2$ if a general presentation of weight ϵ is a direct sum of a presentation of weight ϵ_1 and a presentation of weight ϵ_2 .

Proposition 1.5 (Proposition 5.6). For any two weight vectors ϵ_1 and ϵ_2 of a QP, the following are equivalent:

(1) $\epsilon = \epsilon_1 \oplus \epsilon_2;$ (2) $r_{\epsilon} = r_{\epsilon_1} r_{\epsilon_2} = r_{\epsilon_2} r_{\epsilon_1};$ (3) $l_{\epsilon} = l_{\epsilon_1} l_{\epsilon_2} = l_{\epsilon_2} l_{\epsilon_1}.$

1.1. Organization. In Section 2 we briefly review the theory of quivers with potentials. In Section 3 after a brief review of the theory of general presentations [7], we prove that general presentations behaves well under the Auslander-Reiten translation (Theorem 3.11), which we think is mildly new. In Section 4 we review the theory of 2-Calabi-Yau triangulated categories developed by C. Amiot, B. Keller, Y. Palu, P. Plamondon and so on. Notably is the Palu's formula on index (Lemma 4.8), which is important to us. In Section 5 we introduce the pair of operators r_{ϵ} and l_{ϵ} , and prove our first main result (Theorem 5.16). In Section 6 we study the properties around the general ranks, including our second main result (Theorem 6.1), and some mutation invariants (Proposition 6.3). In Appendix we explain the relationship between the saturation conjecture for nondegenerate QPs and the Fock-Goncharnov's duality pairing conjecture for the generic bases, and explain why Conjecture 5.22 implies them.

1.2. About Notations. By a quiver Q we mean a quadruple $Q = (Q_0, Q_1, t, h)$ where Q_0 is a finite set of vertices, Q_1 is a finite set of arrows, and t and h are the tail and head functions $Q_1 \to Q_0$.

All modules are right modules, and all vectors are row vectors. For direct sum of n copies of M, we write nM instead of the traditional $M^{\oplus n}$. We write hom, ext and e for dim Hom, dim Ext, and dim E. The superscript * is the trivial dual for vector spaces.

the category of finite-dimensional representations of ${\cal A}$
the space of α -dimensional representations of A
the simple representation supported on the vertex \boldsymbol{u}
the projective cover of S_u
the injective envelope of S_u
the dimension vector of M

2. A REVIEW ON REPRESENTATION THEORY OF QUIVERS WITH POTENTIALS

2.1. Decorated Representations and Presentations. Let Q be a finite quiver with no loops. For such a quiver, we associate a skew-symmetric matrix B_Q given by

$$B_Q(u, v) = |\operatorname{arrows} u \to v| - |\operatorname{arrows} v \to u|.$$

Following [5], we define a potential S on a quiver Q as a (possibly infinite) linear combination of oriented cycles in Q. More precisely, a *potential* is an element of the *trace space* $\operatorname{Tr}(\widehat{kQ}) := \widehat{kQ}/[\widehat{kQ},\widehat{kQ}]$, where \widehat{kQ} is the completion of the path algebra kQ and $[\widehat{kQ},\widehat{kQ}]$ is the closure of the commutator subspace of \widehat{kQ} . The pair (Q, S) is a *quiver with potential*, or QP for short. For each arrow $a \in Q_1$, the cyclic derivative ∂_a on \widehat{kQ} is defined to be the linear extension of

$$\partial_a(a_1\cdots a_d) = \sum_{k=1}^d a^*(a_k)a_{k+1}\cdots a_da_1\cdots a_{k-1}.$$

For each potential S, its Jacobian ideal ∂S is the closed (two-sided) ideal in \widehat{kQ} generated by all $\partial_a S$. The Jacobian algebra J(Q, S) is $\widehat{kQ}/\partial S$. A QP is Jacobi-finite if its Jacobian algebra is finite-dimensional.

Definition 2.1. A decorated representation of the Jacobian algebra J is a pair $\mathcal{M} = (M, M^{-})$, where $M \in \operatorname{rep} J$, and M^{-} is a finite-dimensional k^{Q_0} -module.

By abuse of language, we also say that \mathcal{M} is a representation of (Q, \mathcal{S}) . When appropriate, we will view an ordinary representation M as the decorated representation (M, 0).

Following [7] we call a homomorphism between two projective representations, a projective presentation (or presentation in short). As a full subcategory of the category of complexes in rep J, the category of projective presentations is Krull-Schmidt as well. Sometimes it is convenient to view a presentation $P_- \rightarrow P_+$ as elements in the homotopy category $K^b(\text{proj}-J)$ of bounded complexes of projective representations of J. Our convention is that P_- sits in degree -1 and P_+ sits in degree 0.

We denote by P_u (resp. I_u) the indecomposable projective (resp. injective) representation of J corresponding to the vertex u of Q. For $\beta \in \mathbb{Z}_{>0}^{Q_0}$ we write $P(\beta)$ for $\bigoplus_{u \in Q_0} \beta(u) P_u$.

Definition 2.2. ¹ The δ -vector (or weight vector) of a presentation

$$d: P(\beta_{-}) \to P(\beta_{+})$$

is the difference $\beta_+ - \beta_- \in \mathbb{Z}^{Q_0}$. When working with injective presentations

$$\check{d}: I(\check{\beta}_+) \to I(\check{\beta}_-),$$

we call the vector $\check{\beta}_+ - \check{\beta}_-$ the $\check{\delta}$ -vector of \check{d} .

The δ -vector is just the corresponding element in the Grothendieck group of $K^b(\text{proj}-J)$.

Let ν be the Nakayama functor $\operatorname{Hom}_J(-, J)^*$. There is a map still denoted by ν sending a projective presentation to an injective one

$$P_- \to P_+ \mapsto \nu(P_-) \to \nu(P_+)$$

Note that if there is no direct summand of the form $P_i \to 0$, then $\ker(\nu d) = \tau \operatorname{coker}(d)$ where τ is the classical Auslander-Reiten translation.

Let $\mathcal{R}ep(J)$ be the set of decorated representations of J up to isomorphism. There is a bijection between two additive category $\mathcal{R}ep(J)$ and $K^2(\text{proj}-J)$ mapping any representation M to its minimal presentation in rep J, and the simple representation S_u^- of k^{Q_0} to $P_u \to 0$. Now we can naturally extend the classical AR-translation to decorated representations:



Note that this definition agrees with the one in [7]. Suppose that \mathcal{M} corresponds to a projective presentation $d_{\mathcal{M}}$. The δ -vector $\delta_{\mathcal{M}}$ of \mathcal{M} is by definition the δ -vector of $d_{\mathcal{M}}$. If working with the injective presentations, we can define the $\check{\delta}$ -vector $\check{\delta}_{\mathcal{M}}$ of \mathcal{M} . It is known [6] that $\delta_{\mathcal{M}}$ and $\check{\delta}_{\mathcal{M}}$ are related by

$$\delta_{\mathcal{M}} = \delta_{\mathcal{M}} + (\underline{\dim}M)B_Q. \tag{2.1}$$

Definition 2.3 ([6, 7]). Given any projective presentation $d : P_- \to P_+$ and any $N \in \operatorname{rep}(A)$, we define $\operatorname{Hom}(d, N)$ and $\operatorname{E}(d, N)$ to be the kernel and cokernel of the induced map:

$$0 \to \operatorname{Hom}(d, N) \to \operatorname{Hom}_J(P_+, N) \to \operatorname{Hom}_J(P_-, N) \to \operatorname{E}(d, N) \to 0.$$
(2.2)

¹The δ -vector is the same one defined in [7], but is the negative of the **g**-vector defined in [6].

Similarly for an injective presentation $\check{d}: I_+ \to I_-$, we define $\operatorname{Hom}(M, \check{d})$ and $\check{E}(M, \check{d})$ to be the kernel and cokernel of the induced map $\operatorname{Hom}_J(M, I_+) \to \operatorname{Hom}_J(M, I_-)$. It is clear that

$$\operatorname{Hom}(d, N) = \operatorname{Hom}_J(\operatorname{coker}(d), N)$$
 and $\operatorname{Hom}(M, d) = \operatorname{Hom}_J(M, \operatorname{ker}(d)).$

We set $\operatorname{Hom}_J(\mathcal{M}, \mathcal{N}) = \operatorname{Hom}(d_{\mathcal{M}}, N) = \operatorname{Hom}(M, \check{d}_{\mathcal{N}}), E_J(\mathcal{M}, \mathcal{N}) := E(d_{\mathcal{M}}, N) \text{ and } \check{E}_J(\mathcal{M}, \mathcal{N}) := \check{E}(M, \check{d}_{\mathcal{N}}).$

Note that according to this definition, we have that $\operatorname{Hom}_J(\mathcal{M}, \mathcal{N}) = \operatorname{Hom}_J(\mathcal{M}, \mathcal{N}).^2$ We also set $\operatorname{E}(d_{\mathcal{M}}, d_{\mathcal{N}}) = \operatorname{E}_J(\mathcal{M}, \mathcal{N})$ and $\check{\operatorname{E}}(\check{d}_{\mathcal{M}}, \check{d}_{\mathcal{N}}) = \check{\operatorname{E}}_J(\mathcal{M}, \mathcal{N})$. We refer readers to [7] for an interpretation of $\operatorname{E}_J(\mathcal{M}, \mathcal{N})$ in terms of the presentations $d_{\mathcal{M}}$ and $d_{\mathcal{N}}$. We call \mathcal{M} or $d_{\mathcal{M}}$ rigid if $\operatorname{E}_J(\mathcal{M}, \mathcal{M}) = 0$.

Lemma 2.4 ([6, Corollary 10.8 and Proposition 7.3], [7, Corollary 7.6]). We have the following equalities:

- (1) $\operatorname{E}_J(\mathcal{M},\mathcal{N}) = \operatorname{Hom}_J(\mathcal{N},\tau\mathcal{M})^*$ and $\operatorname{\check{E}}_J(\mathcal{M},\mathcal{N}) = \operatorname{Hom}_J(\tau^{-1}\mathcal{N},\mathcal{M})^*$.
- (2) $E_J(\mathcal{M}, \mathcal{M}) = \check{E}_J(\mathcal{M}, \mathcal{M}) = E_J(\tau \mathcal{M}, \tau \mathcal{M}).$

2.2. Mutation of QPs. In [5] and [6], the mutation of quivers with potentials is invented to model the cluster algebras. The *mutation* μ_u of a QP (Q, S) at a vertex u is defined as follows. The first step is to define the following new QP $\tilde{\mu}_u(Q, S) = (\tilde{Q}, \tilde{S})$. We put $\tilde{Q}_0 = Q_0$ and \tilde{Q}_1 is the union of three different kinds

- all arrows of Q not incident to u,
- a composite arrow [ab] from t(a) to h(b) for each a, b with h(a) = t(b) = u,
- an opposite arrow a^* (resp. b^*) for each incoming arrow a (resp. outgoing arrow b) at u.

The new potential on Q is given by

$$\widetilde{\mathcal{S}} := [\mathcal{S}] + \sum_{h(a)=t(b)=u} b^* a^* [ab],$$

where [S] is obtained by substituting [ab] for each words ab occurring in S. Finally we define $(Q', S') = \mu_u(Q, S)$ as the reduced part ([5, Definition 4.13]) of (\tilde{Q}, \tilde{S}) . For this last step, we refer readers to [5, Section 4,5] for details. A sequence of vertices is called *admissible* for (Q, S) if its mutation along this sequence is defined. If all sequences are admissible for (Q, S) then we call (Q, S) nondegenerate.

Now we start to define the mutation of decorated representations of J := J(Q, S). Consider the triangle of linear maps with $\beta_u \gamma_u = 0$ and $\gamma_u \alpha_u = 0$.



We first define a decorated representation $\widetilde{\mathcal{M}} = (\widetilde{M}, \widetilde{M}^+)$ of $\widetilde{\mu}_u(Q, \mathcal{S})$. We set

$$\widetilde{M}(v) = M(v), \quad \widetilde{M}^+(v) = M^+(v) \quad (v \neq u);$$

$$\widetilde{M}(u) = \frac{\ker \gamma_u}{\operatorname{im} \beta_u} \oplus \operatorname{im} \gamma_u \oplus \frac{\ker \alpha_u}{\operatorname{im} \gamma_u} \oplus M^+(u), \quad \widetilde{M}^+(u) = \frac{\ker \beta_u}{\ker \beta_u \cap \operatorname{im} \alpha_u}$$

²This definition is slightly different from the one in [6], which involves the decorated part.

We then set $\widetilde{M}(a) = M(a)$ for all arrows not incident to u, and $\widetilde{M}([ab]) = M(ab)$. It is defined in [5] a choice of linear maps $\widetilde{M}(a^*)$ and $\widetilde{M}(b^*)$ making \widetilde{M} a representation of $(\widetilde{Q}, \widetilde{S})$. We refer readers to [6, Section 10] for details. Finally, we define $\mathcal{M}' = \mu_u(\mathcal{M})$ to be the *reduced part* ([5, Definition 10.4]) of $\widetilde{\mathcal{M}}$.

We say a decorated representation \mathcal{M} of (Q, \mathcal{S}) negative reachable if there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $\mu_{\mathbf{u}}(\mathcal{M})$ is negative, i.e., $\mu_{\mathbf{u}}(\mathcal{M})$ has only the decorated part. Similarly we say \mathcal{M} positive reachable if there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $\mu_{\mathbf{u}}(\mathcal{M})$ is a projective representation.

Let us recall several formula relating the δ -vector of \mathcal{M} and its mutation $\mu_u(\mathcal{M})$. We will use the notation $[b]_+$ for max(b, 0).

Lemma 2.5 ([6, Lemma 5.2]). Let $\delta = \delta_{\mathcal{M}}$ and $\delta' = \delta_{\mu_{\mathbf{u}}(\mathcal{M})}$. We use the similar notation for $\check{\delta} = \check{\delta}_{\mathcal{M}}$ and the dimension vectors $d = \underline{\dim}(\mathcal{M})$. Then

$$\delta'(v) = \begin{cases} -\delta(u) & \text{if } v = u \\ \delta(v) - [b_{v,u}]_+ \beta_-(u) + [-b_{v,u}]_+ \beta_+(u) & \text{if } v \neq u. \end{cases}$$
(2.3)

$$\check{\delta}'(v) = \begin{cases} -\check{\delta}(u) & \text{if } v = u\\ \check{\delta}(v) - [b_{u,v}]_+ \check{\beta}_-(u) + [-b_{u,v}]_+ \check{\beta}_+(u) & \text{if } v \neq u. \end{cases}$$
(2.4)

$$d'(u) = d[b_u]_+ - d(u) + \beta_+(u) + \check{\beta}_-(u) = d[-b_u]_+ - d(u) + \beta_-(u) + \check{\beta}_+(u).$$
(2.5)

where b_u is the u-th column of the matrix B_Q .

We remark that the mutated δ -vector δ' is *not* completely determined by δ (we need β_{-} and β_{+}). But see also Remark 3.2.

Lemma 2.6. [6, Proposition 6.1, and Theorem 7.1] Let $\mathcal{M}' = \mu_u(\mathcal{M})$ and $\mathcal{N}' = \mu_u(\mathcal{N})$. We have that

- (1) $\hom_{J'}(\mathcal{M}', \mathcal{N}') \hom_{J}(\mathcal{M}, \mathcal{N}) = \beta_{-,\mathcal{M}}(u)\dot{\beta}_{-,\mathcal{N}}(u) \beta_{+,\mathcal{M}}(u)\dot{\beta}_{+,\mathcal{N}}(u);$
- (2) $e_{J'}(\mathcal{M}', \mathcal{N}') e_J(\mathcal{M}, \mathcal{N}) = \beta_{+,\mathcal{M}}(u)\beta_{-,\mathcal{N}}(u) \beta_{-,\mathcal{M}}(u)\beta_{+,\mathcal{N}}(u);$
- (2*) $\check{\mathrm{e}}_{J'}(\mathcal{M}',\mathcal{N}') \check{\mathrm{e}}_{J}(\mathcal{M},\mathcal{N}) = \check{\beta}_{-,\mathcal{M}}(u)\check{\beta}_{+,\mathcal{N}}(u) \check{\beta}_{+,\mathcal{M}}(u)\check{\beta}_{-,\mathcal{N}}(u).$

In particular, $e_J(\mathcal{M}, \mathcal{M})$ and $\check{e}_J(\mathcal{M}, \mathcal{M})$ are mutation invariant. So any reachable representation is rigid.

Lemma 2.7. [7, Proposition 7.10] The AR-translation τ commutes with the mutation μ_u at any vertex u.

Finally we mention a long-time conjecture of us.

Conjecture 2.8. For a Jacobi-finite QP, any rigid decorated representation can be obtained from a negative representation by a sequence of mutations and some power of τ .

This is equivalent to say that τ acts transitively on the connected components of the cluster complex of (Q, S) introduced in [7].

3. General Presentations

3.1. General Presentations. We shall start our discussion by reviewing some results in [7]. We will consider a more general setting where the algebra A is any basic finite-dimensional k-algebra, which can be presented as kQ/I.

We say that a general presentation in $PHom_A(\delta)$ has property \heartsuit if there is some open (and thus dense) subset U of $PHom_A(\delta)$ such that all presentations in U have property \heartsuit . For example, a general presentation d in $PHom_A(\delta)$ has the following properties: Hom(d, N) has constant dimension for a fixed $N \in \operatorname{rep} A$. Note that E(d, N) has constant dimension on U as well. We denote these two generic values by $hom(\delta, N)$ and $e(\delta, N)$. If $N = A^*$, then $\operatorname{coker}(d)$ has a constant dimension vector, which will be denoted by $\underline{\dim}(\delta)$.

Any $\delta \in \mathbb{Z}^{Q_0}$ can be written as $\delta = \delta_+ - \delta_-$ where $\delta_+ = \max(\delta, 0)$ and $\delta_- = \max(-\delta, 0)$. Here the maximum is taken coordinate-wise. Lemma 3.1 below motivates the following definition.

$$\operatorname{PHom}_A(\delta) := \operatorname{Hom}_A(P(\delta_-), P(\delta_+)).$$

Lemma 3.1 ([15]). A general presentation in $\operatorname{Hom}_A(P(\beta_-), P(\beta_+))$ is homotopy equivalent to a general presentation in $\operatorname{PHom}_A(\beta_+ - \beta_-)$.

Remark 3.2. Due to Lemma 3.1 the δ -vector of a general presentation satisfies $\beta_+ = [\delta]_+$ and $\beta_- = [-\delta]_+$. In particular, for general presentations, Lemma 2.5.(1) reduces to the following rule:

$$\delta'(v) = \begin{cases} -\delta(u) & \text{if } v = u, \\ \delta(v) - b_{v,u}[-\delta(u)]_+ & \text{if } b_{v,u} > 0, \\ \delta(v) - b_{v,u}[\delta(u)]_+ & \text{if } b_{v,u} < 0. \end{cases}$$
(3.1)

If one like, one can combine the last two cases into one $\delta'(v) = \delta(v) + [b_{v,u}]_+ \delta(u) - b_{v,u}[\delta(u)]_+$. We call this the mutation rule for δ -vectors. Later when we write $\mu_u(\delta)$, we refer to this rule.

The presentation space $\operatorname{PHom}_A(\delta)$ comes with a natural group action by

$$\operatorname{Aut}_A(\delta) := \operatorname{Aut}_A(P(\delta_-)) \times \operatorname{Aut}_A(P(\delta_+)).$$

A rigid presentation in $\operatorname{PHom}_A(\delta)$ has a dense $\operatorname{Aut}_A(\delta)$ -orbit [7]. In particular, a rigid presentation is always general.

As explained in [7], the function hom(-,-) and e(-,-) on $PHom_A(\delta_1) \times PHom_A(\delta_2)$ is upper semi-continuous. We will denote its generic value by $hom(\delta_1, \delta_2)$ and $e(\delta_1, \delta_2)$. We say a presentation d_2 is a quotient presentation of d if we have the following diagram

$$0 \longrightarrow P_{-}^{1} \longrightarrow P_{-} \longrightarrow P_{-}^{2} \longrightarrow 0$$

$$d_{1} \downarrow \qquad d_{2} \downarrow \qquad (3.2)$$

$$0 \longrightarrow P_{+}^{1} \longrightarrow P_{+} \longrightarrow P_{+}^{2} \longrightarrow 0$$

We may also write this diagram as the exact sequence:

$$0 \to d_1 \to d \to d_2 \to 0.$$

Theorem 3.3 ([7, Theorem 3.10]). Let $\beta_+ = [\delta_1]_+ + [\delta_2]_+$ and $\beta_- = [-\delta_1]_+ + [-\delta_2]_+$. Then A general presentation in $\operatorname{Hom}_A(P(\beta_-), P(\beta_+))$ has a quotient presentation in $\operatorname{PHom}_A(\delta_2)$ if and only if $e(\delta_1, \delta_2) = 0$.

The following notion is important throughout this paper. Let U be a subset of the product $\prod_{i=1}^{r} X_i$. We say elements in U can be chosen to be general in some X_i if $p_i(U)$ contains an open subset of X_i . We say elements in U can be chosen to be general as a pair in X_i and X_j if $(p_i, p_j)(U)$ contains an open subset of $X_i \times X_j$. Similarly we can talk about generalness in a triple and so on.

8

Recall from [7] that $E(d_2, d_1)$ is a vector space quotient of $Hom_A(P^1_+, P^2_-)$ so any diagram (3.2) gives rises to an element $\eta \in E(d_2, d_1)$.

Lemma 3.4. Keep the notations in Theorem 3.3, and suppose that a general presentation in $\operatorname{Hom}_A(P(\beta_-), P(\beta_+))$ has a quotient presentation d_2 in $\operatorname{PHom}_A(\delta_2)$. Let d_1 be the corresponding subpresentation and η be the corresponding element in $\operatorname{E}(d_2, d_1)$. Then we may assume (d_1, d_2) is a general pair and η is general in $\operatorname{E}(d_2, d_1)$.

Proof. Fix a subrepresentation $P^1_+ \cong P([\delta_1]_+)$ of P_+ such that $P_+/P^1_+ \cong P([\delta_2]_+)$ and let p_+ be the natural projection $P_+ \to P_+/P^1_+$. We also do the similar thing for P_- . We define the variety

$$\operatorname{PHom}_A(\delta_1 \mid \delta_2) := \{ d \in \operatorname{PHom}_A(\delta) \mid p_+ d \mid_{P^1} = 0 \}.$$

Our assumption that a general presentation in $\operatorname{PHom}_A(\delta)$ has a quotient presentation in $\operatorname{PHom}_A(\delta_2)$ implies that the action morphism

 $\operatorname{Aut}_A(\delta) \times \operatorname{PHom}_A(\delta_1 \mid \delta_2) \to \operatorname{PHom}_A(\delta)$

is dominant. It is clear from the definition that

 $\operatorname{PHom}_A(\delta_1 \mid \delta_2) \cong \operatorname{PHom}_A(\delta_1) \times \operatorname{PHom}_A(\delta_2) \times \operatorname{Hom}_A(P^1_+, P^2_-).$

So any open subset of $\operatorname{PHom}_A(\delta)$ pulls back and projects to a nonempty open subset of $\operatorname{PHom}_A(\delta_1) \times \operatorname{PHom}_A(\delta_2) \times \operatorname{Hom}_A(P_+^1, P_-^2)$. Hence, we may assume (d_1, d_2) is a general pair. As $\operatorname{E}(d_2, d_1)$ is a vector space quotient of $\operatorname{Hom}_A(P_+^1, P_-^2)$, the general element in $\operatorname{Hom}_A(P_+^1, P_-^2)$ projects to a general element in $\operatorname{E}(d_2, d_1)$.

We write $\delta = \delta_1 \oplus \delta_2$ if a general presentation of weight δ is a direct sum of a presentation of weight δ_1 and a presentation of weight δ_2 .

3.2. τ **Permutes Principal Components.** In [7, Section 2] we considered the incidence variety $Z(Y, X) = \{(f, \pi, M) \in Y \times \operatorname{Hom}(P_+, k^{\alpha}) \times X \mid \pi \in \operatorname{Hom}_A(P_+, M) \text{ and } P_- \to P_+ \to M \to 0 \text{ is exact}\}$

(3.3) for any $\operatorname{Aut}_A(P_-) \times \operatorname{Aut}_A(P_+)$ -stable subvariety Y of $\operatorname{Hom}_A(P_1, P_0)$ and $\operatorname{GL}_{\alpha}$ -stable subvariety X of $\operatorname{rep}_{\alpha}(A)$. It comes with two projections $p_1: Z \to \operatorname{Hom}_A(P_-, P_+)$ and $p_2: Z \to \operatorname{rep}_{\alpha}(A)$.

Theorem 3.5 ([7, Theorem 2.3, Corollary 2.6]). The projection p_2 is open. Moreover, the map $p_2p_1^{-1}$ gives a bijection between $\operatorname{Aut}_A(P_-) \times \operatorname{Aut}_A(P_+)$ -stable subvariety of $\operatorname{im} p_1$ and $\operatorname{GL}_{\alpha}$ -stable subvariety of $\operatorname{im} p_2$, preserving openness, closure, and irreducibility.

As discussed before, there is an open subset $U \subset \operatorname{PHom}_A(\delta)$ such that $W = p_2(p_1^{-1}(U))$ lies entirely in $\operatorname{rep}_{\alpha}(A)$. In particular, there is an irreducible component C of $\operatorname{rep}_{\alpha}(A)$ such that $p_1(p_2^{-1}(W \cap C))$ is dense in $\operatorname{PHom}_A(\delta)$. Since p_2 is open, W (and thus $W \cap C$) is open. So $p_1(p_2^{-1}(W \cap C))$ is constructible, and thus contains an open subset of $\operatorname{PHom}_A(\delta)$. We conclude that we can shrink U such that $W = p_2(p_1^{-1}(U))$ lies in a single component of $\operatorname{rep}_{\alpha}(A)$.

Definition 3.6. We call this component the *principal component* of δ , denoted by $PC(\delta)$.

As shown in [21], such a component is exactly the *strongly reduced component* introduced in [13]. This fact also follows from our Lemma 3.8 below.

When we say a general representation of weight δ , we mean a general representation in PC(δ). By (2.1) a general representation of weight δ has $\check{\delta}$ -vector $\delta + \underline{\dim}(\delta)B_Q$, which will be denoted by $\check{\delta}$. By the remarks after Definition 2.3, the hom(δ, ϵ) and $e(\delta, \epsilon)$ defined in terms of presentations is nothing but the generic (minimal) values of hom_J(M, N) and $e_J(M, N) + \delta^-(\underline{\dim}(\epsilon))$ on

 $PC(\delta) \times PC(\epsilon)$. Here, δ^- is the nonpositive vector appears in the decomposition $\delta = \delta' \oplus \delta^$ with δ' negative-free, which means that a general presentation in $PHom_J(\delta')$ does not contain a direct summand of form $P \to 0$.

Lemma 3.7. For any $\operatorname{Aut}_A(\delta)$ -stable subvariety $X \subset \operatorname{PHom}_A(\delta)$, there is a nonempty open $\operatorname{Aut}_A(\delta)$ -stable subset U_0 of X such that there is morphism $\pi : U_0 \to \operatorname{PC}(\delta)$ such that $\pi(d)$ is isomorphic to $\operatorname{coker}(d)$.

Proof. This follows from [21, Lemma 2.11].

By a G-variety, we mean a variety with an action of an algebraic group G. By a theorem of Rosenlicht ([22], see also [8, Theorem 6.2]), any irreducible G-variety X contains a nonempty open G-stable subset X_0 which admits a geometric quotient X_0/G . For the definition of geometric quotient, we refer readers to [8, 6.1].

We shall denote by $e_A(\delta)$ the generic value of $e_A(d, d)$ for $d \in \text{PHom}_A(\delta)$. Note the difference between $e_A(\delta)$ and $e_A(\delta, \delta)$. The latter does not require the two arguments d_1 and d_2 to be the same in $e_A(d_1, d_2)$.

Lemma 3.8. There is a non-empty open $\operatorname{Aut}_A(\delta)$ -stable subset U of $\operatorname{PHom}_A(\delta)$ and a nonempty open $\operatorname{GL}_{\underline{\dim}(\delta)}$ -stable subset W of its principal component $\operatorname{PC}(\delta)$ such that there is an isomorphism of geometric quotients

$$U/\operatorname{Aut}_A(\delta) \to W/\operatorname{GL}_{\dim(\delta)}$$
.

Moreover the dimension of the quotient $U/\operatorname{Aut}_A(\delta)$ is equal to $e_A(\delta)$.

Proof. Let U_0 be an open subset of $\operatorname{PHom}_A(\delta)$ as in Lemma 3.7. We get a morphism of varieties $\pi : U_0 \to \operatorname{PC}(\delta)$. Let W be the open subset of $\operatorname{PC}(\delta)$ such that $q : W \to W/\operatorname{GL}_{\operatorname{dim}(\delta)}$ is a geometric quotient. Let U_1 be the open subset of $\operatorname{PHom}_A(\delta)$ such that $U_1 \to U_1/\operatorname{Aut}_A(\delta)$ is a geometric quotient, and set $U = U_0 \cap U_1$. The composition gives $q\pi \mid_U : U \to W/\operatorname{GL}_{\operatorname{dim}(\delta)}$. Recall that a geometric quotient is also a categorical quotient [8]. This map is constant on the $\operatorname{Aut}_A(\delta)$ -orbit of U so it descends to $U/\operatorname{Aut}_A(\delta) \to W/\operatorname{GL}_{\operatorname{dim}(\delta)}$.

For the statement about the dimension, we have that

$$\dim(U/\operatorname{Aut}_{A}(\delta)) = \dim U - \dim \operatorname{Aut}_{A}(\delta) + \min_{d \in \operatorname{PHom}_{A}(\delta)} (\dim \operatorname{Aut}_{A}(\delta)_{d})$$
$$= \dim \operatorname{PHom}_{A}(\delta) - \dim \operatorname{Aut}_{A}(\delta) + \dim(\operatorname{Aut}_{A}(\delta)_{d})$$
(3.4)

where d is a general presentation of weight δ and $\operatorname{Aut}_A(\delta)_d$ is the stabilizer of d in $\operatorname{Aut}_A(\delta)$. It follows from [7, Lemma 3.7] that the Lie algebra of $\operatorname{Aut}_A(\delta)_d$ can be identified with the kernel of the map $f = (d_+, -d_-)$, where d_+, d_- is the induced map by applying $\operatorname{Hom}_A(-, P_+)$ and $\operatorname{Hom}_A(P_-, -)$ to d respectively.

Consider the complex induced from (3.5):

$$0 \to \ker(f) \to \operatorname{Hom}_A(P_-, P_-) \oplus \operatorname{Hom}_A(P_+, P_+) \xrightarrow{J} \operatorname{Hom}_A(P_-, P_+) \to \operatorname{E}_A(d, d) \to 0.$$

We conclude that $(3.4) = e_A(\delta)$ as desired.

Remark 3.9. Slightly modify this proof we can easily show that for any $\operatorname{Aut}_A(\delta)$ -stable subset X of $\operatorname{PHom}_A(\delta)$ which maps onto an open subset of some irreducible component C of $\operatorname{rep}_{\alpha}(A)$, there exist open subsets U and W of X and C respectively such that there is an isomorphism of geometric quotients $U/\operatorname{Aut}_A(\delta) \to W/\operatorname{GL}_{\alpha}$.

We also remark that there is a similar statement for $\operatorname{IHom}(\check{\delta})$. In this case, the dimension formula should read as $\dim(\check{U}/\operatorname{Aut}_A(\check{\delta})) = \check{e}_A(\check{\delta})$.

From now on, we shall assume the algebra A is a Jacobian algebra J as before.

Lemma 3.10. For $\check{\delta} = \delta + \underline{\dim}(\delta)B_Q$, we have that $e_J(\delta) = \check{e}_J(\check{\delta})$.

Proof. Let d be a presentation of weight δ such that $e_J(d, d) = e_J(\delta)$. Let \check{d} be the corresponding injective presentation, that is, $d = d_{\mathcal{M}}$ and $\check{d} = \check{d}_{\mathcal{M}}$. Then \check{d} has weight $\check{\delta}$ by (2.1). We have that $\check{e}_J(\check{\delta}) \leq \check{e}_J(\check{d},\check{d}) = e_J(d,d) = e_J(\delta)$ by Lemma 2.4. Similarly we can get the other inequality. \Box

Due to the relation $\delta_{\tau \mathcal{M}} = -\delta_{\mathcal{M}}$ and (2.1), we have that for a general presentation d of weight δ , the δ -vector of τd is constant. We denote this constant vector by $\tau \delta$.

Theorem 3.11.

- (1) $\operatorname{PHom}_J(\delta)$ and $\operatorname{IHom}_J(\check{\delta})$ have the same principal component where $\check{\delta} = \delta + \underline{\dim}(\delta)B_Q$.
- (2) *M* is a general representation in $PC(\delta)$ if and only if τM is a general representation in $PC(\tau\delta)$.

Proof. (1). It is enough to show that there is an open subset U of $\operatorname{PHom}_J(\delta)$ and an open subset \check{U} of $\operatorname{IHom}_J(\check{\delta})$ such that they correspond to the same open subset W in the principal component $\operatorname{PC}(\delta)$.

Let U and W be the open subset of $\operatorname{PHom}_J(\delta)$ and $\operatorname{PC}(\delta)$ as in Lemma 3.8. Recall the incidence variety Z(Y, X) in (3.3) with projection p_1 and p_2 . Let $\check{U}' = p_1(p_2^{-1}(W))$ be the (constructible) subset of $\operatorname{IHom}_J(\check{\delta})$. By possibly shrinking \check{U}' and W we may assume the isomorphism (see Remark 3.9)

$$\check{U}' / \operatorname{Aut}_A(\check{\delta}) \cong W / \operatorname{GL}_{\operatorname{\underline{dim}}(\delta)}$$

So by Lemma 3.8 we have that

$$\dim(U'/\operatorname{Aut}_A(\delta)) = \dim(W/\operatorname{GL}_{\underline{\dim}(\delta)}) = \dim(U/\operatorname{Aut}_A(\delta)) = e_J(\delta)$$

which is equal to $\check{e}_J(\check{\delta})$ by Lemma 3.10. But $\dim \check{U} / \operatorname{Aut}_A(\check{\delta}) = \check{e}_J(\check{\delta})$ as well, where \check{U} is an open subset of $\operatorname{IHom}_J(\check{\delta})$ as claimed in Remark 3.9. So $\dim \check{U}' = \dim \check{U}$. Hence, \check{U}' is in fact open in $\operatorname{IHom}_J(\check{\delta})$.

(2). Let ι be the isomorphism $U/\operatorname{Aut}_A(\delta) \to W/\operatorname{GL}_{\underline{\dim}(\delta)}$ as in Lemma 3.8. By the part (1), $\check{U}/\operatorname{Aut}_A(-\delta)$ is isomorphic to $\check{W}/\operatorname{GL}_{\underline{\dim}(\tau\delta)}$ for some open subset \check{W} in $\operatorname{PC}(\tau\delta)$. Recall the Nakayama functor ν . It has an obvious algebraic lifting $\operatorname{PHom}(\delta) \to \operatorname{IHom}(-\delta)$ for each δ , which is an isomorphism and sends a linear combination of paths $\sum_i c_i p_i$ to $\sum_i c_i p_i^*$. By properly adjusting the open sets U and \check{U} , this isomorphism descends to the geometric quotients $U/\operatorname{Aut}_A(\delta) \xrightarrow{\nu} \check{U}/\operatorname{Aut}_A(-\delta)$. We thus obtain the following diagram (if necessary we may shrink W and \check{W})

$$\begin{array}{c|c} U/\operatorname{Aut}_{A}(\delta) & \xrightarrow{\iota} & W/\operatorname{GL}_{\operatorname{dim}(\delta)} \\ & \nu & & & \downarrow \\ & & & \downarrow^{\iota\nu\iota^{-1}} \\ \check{U}/\operatorname{Aut}_{A}(-\delta) & \xrightarrow{\check{\iota}} & \check{W}/\operatorname{GL}_{\operatorname{\underline{dim}}(\tau\delta)} \end{array}$$

As W and \dot{W} are open in $PC(\delta)$ and $PC(\tau\delta)$, the isomorphism $W/\operatorname{GL}_{\underline{\dim}(\delta)} \cong \dot{W}/\operatorname{GL}_{\underline{\dim}(\tau\delta)}$ implies (2).

The following lemma is implied in the proof of [21, Theorem 3.21]. It can also be proved in a similar fashion as our Theorem 3.11.

Lemma 3.12. There is an open subset U in $PC(\delta)$ and an open subset U' in $PC(\mu_u(\delta))$ such that the orbit of $\mu_u(M)$ lies in U' if and only if the orbit of M lies in U. In particular, M is a general representation in $PC(\delta)$ if and only if $\mu_u(M)$ is a general representation in $PC(\mu_u(\delta))$.

4. A REVIEW ON 2-CALABI-YAU TRIANGULATED CATEGORIES

4.1. The Cluster Category $C_{Q,S}$. C. Amiot introduced in [1] a triangulated category $C_{Q,S}$ associated to a quiver with potential (Q, S). Let $\Gamma = \Gamma_{Q,S}$ be the complete Ginzburg's dgalgebra attached to (Q, S) [14], and $\mathcal{D}\Gamma$ be its derived category. The perfect derived category per Γ of Γ is the smallest full triangulated subcategory of $\mathcal{D}\Gamma$ containing Γ and closed under taking direct summands. Denote by $\mathcal{D}_{fd}\Gamma$ the full subcategory of $\mathcal{D}\Gamma$ whose objects are those of $\mathcal{D}\Gamma$ with finite-dimensional total homology. As shown in [17, Theorem 2.17], the category $\mathcal{D}_{fd}\Gamma$ is a triangulated subcategory of per Γ . The cluster category $\mathcal{C}_{Q,S}$ of (Q,S) as the idempotent completion of the triangulated quotient $(\text{per }\Gamma)/\mathcal{D}_{fd}\Gamma$.

When (Q, S) is Jacobi-finite, the category $C_{Q,S}$ is Hom-finite and 2-Calabi–Yau, and admits a basic *cluster-tilting* object $\mathbf{T} = \Sigma^{-1}\Gamma$. Its endomorphism algebra is isomorphic to the Jacobian algebra J(Q, S). Here, 2-Calabi-Yau means that there is a bifunctorial isomorphism

$$\mathcal{C}(\mathbf{L}, \Sigma \mathbf{N}) \cong \mathcal{C}(\mathbf{N}, \Sigma \mathbf{L})^*$$

The cluster-tilting object is by definition the object satisfying

- (1) $\mathcal{C}(\mathbf{T}, \Sigma \mathbf{T}) = 0$ and
- (2) for any **M** in \mathcal{C} , if $\mathcal{C}(\mathbf{M}, \Sigma \mathbf{T}) = 0$, then **M** belongs to the full additive subcategory add **T**.

We keep k as an algebraically closed field of characteristic zero. Throughout we will write \mathcal{C} for $\mathcal{C}(Q, \mathcal{S})$ though some of the definitions and results hold in any Hom-finite, 2-Calabi–Yau, Krull–Schmidt k-category which admits a basic cluster-tilting object **T**. Let J be the endomorphism algebra of **T** in \mathcal{C} , and denote by mod J the category of finite-dimensional right J-modules. As shown in [16] the functor $F : \mathcal{C} \to \text{mod } J$ sending **M** to $\mathcal{C}(\mathbf{T}, \mathbf{M})$, induces an equivalence of categories:

$$\mathcal{C}/(\Sigma \mathbf{T}) \cong \mod J,\tag{4.1}$$

where (\mathbf{T}) denotes the ideal of morphisms of \mathcal{C} which factor through an object in add \mathbf{T} . This equivalence restricts to the full subcategories: add $\mathbf{T} \to \text{proj}-J$, which allows us lift *J*-modules to \mathcal{C} using projective presentations.

This equivalence can be slightly extended to incorporate the decorated representations (see [20] for more details). Let \mathbf{M} be an object in \mathcal{C} of the form $\mathbf{M} = \mathbf{M}' \oplus \bigoplus_{u \in Q_0} m_u \Sigma \mathbf{T}_u$ where \mathbf{M} has no direct summands in add $\Sigma \mathbf{T}$. Such an \mathbf{M} will correspond to the decorated representation $(F\mathbf{M}', \bigoplus_{u \in Q_0} m_u(0, S_u))$ where $(0, S_u)$ is the negative simple representation at vertex u. We denote this map by $\widetilde{F} = \widetilde{F}_{Q,S}$, which is denoted by Φ in [20]. If no potential confusion is possible, throughout we will write \mathbf{M} for a lift of M or \mathcal{M} , and denote $F\mathbf{M}$ and $\widetilde{F}\mathbf{M}$ by M and \mathcal{M} respectively.

We also have the following analogue of the space E in \mathcal{C} . Following [21], for $\mathbf{d}' \in \mathcal{C}(\mathbf{T}'_{-}, \mathbf{T}'_{+})$ and $\mathbf{d}'' \in \mathcal{C}(\mathbf{T}''_{-}, \mathbf{T}''_{+})$ we define $E(\mathbf{d}', \mathbf{d}'')$ to be the vector space $\operatorname{Hom}_{K^{b}(\operatorname{add} \mathbf{T})}(\mathbf{d}', \Sigma \mathbf{d}'')$. We denote $F\mathbf{d}'$ and $F\mathbf{d}''$ by d' and d''.

12

Lemma 4.1 ([21, Proposition 3.10]). The space $E(\mathbf{d}', \mathbf{d}'')$ is isomorphic to E(d', d''), and it can be identified with $(\Sigma \mathbf{T})(\mathbf{N}, \Sigma \mathbf{L})$ where \mathbf{N} and \mathbf{L} are the cones of \mathbf{d}' and \mathbf{d}'' respectively. Dually $\check{E}(\check{\mathbf{d}}', \check{\mathbf{d}}'')$ is isomorphic to $\check{E}(\check{d}', \check{d}'')$, and it can be identified with $(\Sigma \mathbf{T})(\Sigma^{-1}\mathbf{N}, \mathbf{L})$ where \mathbf{N} and \mathbf{L} are the fibres of \mathbf{d}' and \mathbf{d}'' respectively.

The following lemma is a reformulation of [19, Lemma 3.3].

Lemma 4.2. $C(\mathbf{N}, \Sigma \mathbf{L}) \cong (\Sigma \mathbf{T})(\mathbf{N}, \Sigma \mathbf{L}) \oplus (\Sigma \mathbf{T})(\mathbf{L}, \Sigma \mathbf{N})^*$.

Corollary 4.3. Let $\mathbf{L} \to \mathbf{M} \to \mathbf{N} \to \Sigma \mathbf{L}$ be a triangle in \mathcal{C} .

- (1) If $e(\mathcal{L}, \mathcal{N}) = 0$, then every $f \in \mathcal{C}(\mathbf{N}, \Sigma \mathbf{L})$ factor through $\Sigma \mathbf{T}$.
- (2) If $\check{\mathbf{e}}(\mathcal{L}, \mathcal{N}) = 0$, then every $f \in \mathcal{C}(\Sigma^{-1}\mathbf{N}, \mathbf{L})$ factor through $\Sigma \mathbf{T}$.
- (3) If $e(\mathcal{L}, \mathcal{N}) = e(\mathcal{N}, \mathcal{L}) = 0$, then $\mathcal{C}(\mathbf{L}, \Sigma \mathbf{N}) = \mathcal{C}(\mathbf{N}, \Sigma \mathbf{L}) = 0$.

Proof. (1). Since $e(\mathcal{L}, \mathcal{N}) = 0$, we have that $(\Sigma \mathbf{T})(\mathbf{L}, \Sigma \mathbf{N}) = 0$. So by Lemma 4.2 every $f \in \mathcal{C}(\mathbf{N}, \Sigma \mathbf{L})$ factor through $\Sigma \mathbf{T}$. The proof of (2) is similar. (3). If $e(\mathcal{L}, \mathcal{N}) = e(\mathcal{N}, \mathcal{L}) = 0$, then $(\Sigma \mathbf{T})(\mathbf{L}, \Sigma \mathbf{N}) = (\Sigma \mathbf{T})(\mathbf{N}, \Sigma \mathbf{L}) = 0$. By Lemma 4.2 $\mathcal{C}(\mathbf{L}, \Sigma \mathbf{N}) = \mathcal{C}(\mathbf{N}, \Sigma \mathbf{L}) = 0$.

In [17] Keller and Yang lifted Derksen-Weyman-Zelevinsky's mutation to the category $\mathcal{D}\Gamma$. The lifted mutation $\tilde{\mu}_u^-$ is in fact a triangle equivalence $\mathcal{D}\Gamma \to \mathcal{D}\Gamma'$, where Γ' the complete Ginzburg dg algebra of $\tilde{\mu}_u(Q, \mathcal{S})$. This equivalence restricts to the subcategories per $\Gamma \to \text{per }\Gamma'$ and $\mathcal{D}_{fd}\Gamma \to \mathcal{D}_{fd}\Gamma'$. There are similar statements for the reduced quivers with potentials (see [17, Theorem 3.2]). In particular, it induces a triangle equivalence $\mathcal{C}_{Q,\mathcal{S}} \to \mathcal{C}_{\mu_u(Q,\mathcal{S})}$, still denoted by $\tilde{\mu}_u^-$. The lifted mutation is compatible with the ordinary one in the following sense.

Theorem 4.4 ([20, Proposition 4.1]). If $u \in Q_0$ is not on any cycle of length 2, then for any object **M** of \mathcal{D} , we have that

$$\widetilde{F}_{\widetilde{\mu}_u(Q,\mathcal{S})}(\widetilde{\mu}_u^-(\mathbf{M})) = \widetilde{\mu}_u(\widetilde{F}_{Q,\mathcal{S}}(\mathbf{M})).$$

Given an admissible sequence of mutations $\mu_{\mathbf{u}} = \mu_{u_r} \cdots \mu_{u_2} \mu_{u_1}$, we have a sequence of triangle equivalences

$$\mu_{\mathbf{u}}^{-}: \mathcal{C}_{Q,\mathcal{S}} \to \mathcal{C}_{\mu_{u_1}(Q,\mathcal{S})} \to \mathcal{C}_{\mu_{u_2}\mu_{u_1}(Q,\mathcal{S})} \to \dots \to \mathcal{C}_{\mu_{\mathbf{u}}(Q,\mathcal{S})}.$$
(4.2)

Let the functor \widetilde{F}' be the composition $\widetilde{F}_{\mu_{\mathbf{u}}(Q,S)} \circ \mu_{\mathbf{u}}^-$. We write J' for the Jacobian algebra of $\mu_{\mathbf{u}}(Q,S)$.

Lemma 4.5. Given an admissible sequence of mutations $\mu_{\mathbf{u}} = \mu_{u_r} \cdots \mu_{u_2} \mu_{u_1}$, the functor \widetilde{F}' sends a general object \mathbf{M} of index δ in \mathcal{C} to a general object of weight $\mu_{\mathbf{u}}(\delta)$ in rep J', and a general morphism \mathbf{g} in $\mathcal{C}(\mathbf{M}, \mathbf{N})$ to a general homomorphism in $\operatorname{Hom}_{J'}(\widetilde{F}'\mathbf{M}, \widetilde{F}'\mathbf{N})$.

Proof. Due to Theorem 4.4, $\widetilde{F}' = \mu_{\mathbf{u}} \circ \widetilde{F}_{Q,\mathcal{S}}$. Note that $\widetilde{F}_{Q,\mathcal{S}}(\mathbf{M})$ is general of weight δ . So it suffices to show that a general representation of weight δ will be mutated to a general representation of weight $\mu_{\mathbf{u}}(\delta)$. But this follows from Lemma 3.12. Since $\mu_{\mathbf{u}}^-$ is a triangle equivalence, $\mu_{\mathbf{u}}^-(\mathbf{g})$ is general in $\mathcal{C}(\mathbf{M}', \mathbf{N}')$ where $\mathbf{M}' = \mu_{\mathbf{u}}^-(\mathbf{M})$ and $\mathbf{N}' = \mu_{\mathbf{u}}^-(\mathbf{N})$. For $\mathbf{T}' = \Sigma^{-1}\Gamma_{\mu_{\mathbf{u}}(Q,\mathcal{S})}$, $\mathcal{C}(\mathbf{M}', \mathbf{N}') \to \mathcal{C}(\mathbf{M}', \mathbf{N}')/(\Sigma \mathbf{T}'(\mathbf{M}', \mathbf{N}')) \cong \operatorname{Hom}_{J'}(M', N')$ is a vector space projection. In particular, it is an open map so a general morphism in $\mathcal{C}(\mathbf{M}', \mathbf{N}')$ descends to a general element in $\operatorname{Hom}_{J'}(M', N')$.

4.2. Palu's Formula on Indices. Let \mathbf{M} be an object of \mathcal{C} . There exist triangles (see [16]) $\mathbf{T}_{-} \rightarrow \mathbf{T}_{+} \rightarrow \mathbf{M} \rightarrow \Sigma \mathbf{T}_{-}$ and $\mathbf{M} \rightarrow \Sigma^{2} \mathbf{T}^{+} \rightarrow \Sigma^{2} \mathbf{T}^{-} \rightarrow \Sigma \mathbf{M}$

with $\mathbf{T}_+, \mathbf{T}_-, \mathbf{T}^+, \mathbf{T}^- \in \operatorname{add} \mathbf{T}$.

Definition 4.6. The *index* and *coindex* of **M** with respect to **T** are the classes in $K_0(\text{add }\mathbf{T})$:

$$\operatorname{ind}_{\mathbf{T}}(\mathbf{M}) = [\mathbf{T}_+] - [\mathbf{T}_-] \text{ and } \operatorname{coind}_{\mathbf{T}}(\mathbf{M}) = [\mathbf{T}^+] - [\mathbf{T}^-].$$

Similar to the projective representations, we denote $\mathbf{T}(\beta) = \bigoplus_{u \in Q_0} \beta(u) \mathbf{T}_u$ where \mathbf{T}_u is the indecomposable direct summands of \mathbf{T} corresponding to u. In this way, the index and coindex can be naturally identified with a vector in \mathbb{Z}^{Q_0} : if $\mathbf{T}_+ = \mathbf{T}(\beta_+)$ and $\mathbf{T}_- = \mathbf{T}(\beta_-)$, then $\operatorname{ind}_{\mathbf{T}}(\mathbf{M}) = \beta_+ - \beta_-$. Under this identification, the following lemma is obvious from the equivalence (4.1) and the fact that $\operatorname{ind}_{\mathbf{T}}(\Sigma \mathbf{T}_u) = e_u$.

Lemma 4.7. As a vector in \mathbb{Z}^{Q_0} , the index of **M** is equal to the weight of $\widetilde{F}(\mathbf{M})$.

We also define the presentation space $\mathbf{T}\mathrm{Hom}(\delta) := \mathcal{C}(\mathbf{T}([\delta]_+), \mathbf{T}([-\delta]_+))$. By a general element **M** of index δ , we mean that **M** is the cone of a general element in $\mathbf{T}\mathrm{Hom}(\delta)$.

Here is Palu's formula on indices in a triangle.

Lemma 4.8 ([19, Proposition 2.2]). Let $\mathbf{L} \xrightarrow{\mathbf{f}} \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{N} \to \Sigma \mathbf{L}$ be a triangle in \mathcal{C} . Take $\mathbf{C} \in \mathcal{C}$ (resp. $\mathbf{K} \in \mathcal{C}$) to be any lift of coker $F\mathbf{g}$ (resp. ker $F\mathbf{f}$). Then

$$\operatorname{ind} \mathbf{M} = \operatorname{ind} \mathbf{L} + \operatorname{ind} \mathbf{N} - \operatorname{ind} \mathbf{C} - \operatorname{ind} \Sigma^{-1} \mathbf{C};$$

coind $\mathbf{M} = \operatorname{coind} \mathbf{L} + \operatorname{coind} \mathbf{N} - \operatorname{coind} \mathbf{K} - \operatorname{coind} \Sigma \mathbf{K}$.

Corollary 4.9. A triangle $\mathbf{L} \to \mathbf{M} \to \mathbf{N} \to \Sigma \mathbf{L}$ in \mathcal{C} gives a long exact sequence in mod J:

$$\cdots \to \tau^{-1}L \to \tau^{-1}M \to \tau^{-1}N \xrightarrow{\theta'} L \to M \to N \xrightarrow{\theta} \tau L \to \tau M \to \tau N \to \tau^2 L \to \cdots .$$
(4.3)

with $\delta_{\mathcal{M}} = \delta_{\mathcal{L}} + \delta_{\mathcal{N}} + \operatorname{rank}(\theta)B$ and $\delta_{\mathcal{M}} = \delta_{\mathcal{L}} + \delta_{\mathcal{N}} - \operatorname{rank}(\theta')B$.

Proof. We apply the functor $F = C(\mathbf{T}, -)$ to the above triangle, and note that $C(\mathbf{T}, \Sigma^i \mathbf{M}) \cong \tau^i M$. So we get the desired long exact sequence. Since $\delta_{\mathcal{M}} = \operatorname{ind} \mathbf{M}$, the last formula then follows from lemma 4.8.

We had a direct explicit proof of this corollary without referring to the category C. We'd like to thank Bernhard Keller for pointing out the connection to the 2-Calabi-Yau categories. This save us at least two pages of space.

4.3. The Lowering and Raising Operators in C.

Definition 4.10. For any object $\mathbf{E} \in C$, we define the operator $\mathbf{r}_{\mathbf{E}}$ on $K_0(\text{add }\mathbf{T})$ as follows. Pick an object \mathbf{M} of index δ , and let \mathbf{g} be a morphism $\mathcal{C}(\mathbf{E}, \Sigma \mathbf{M})$. We complete \mathbf{g} to a triangle

$$\Sigma^{-1}\mathbf{E} \to \mathbf{M} \to \mathbf{R} \to \mathbf{E}.$$
 (4.4)

By Lemma 4.8 the index of **R** is constant if **M** is general and **g** is general in $C(\mathbf{E}, \Sigma \mathbf{M})$. We define $\mathbf{r}_{\mathbf{E}}(\delta)$ to be such a constant index of **R**. Similarly we can take **g** to be a general morphism $C(\mathbf{M}, \mathbf{E})$ and complete it to

$$\Sigma^{-1}\mathbf{E} \to \mathbf{L} \to \mathbf{M} \to \mathbf{E},\tag{4.5}$$

and define $\mathbf{l}_{\mathbf{E}}(\delta)$ to be the index of \mathbf{L} . If working with the coindices, we get another version of $\mathbf{r}_{\mathbf{E}}$ and $\mathbf{l}_{\mathbf{E}}$ which will be denoted by $\mathbf{r}^{\mathbf{E}}$ and $\mathbf{l}^{\mathbf{E}}$. If \mathbf{E} is a general object of index ϵ , then we will denote $\mathbf{r}_{\mathbf{E}}$ and $\mathbf{l}_{\mathbf{E}}$ by \mathbf{r}_{ϵ} and \mathbf{l}_{ϵ} .

It is unclear from the definition if the above **R** or **L** can be chosen as a general element of index \mathbf{r}_{ϵ} or \mathbf{l}_{ϵ} . Later we will see that this indeed can be done in most cases (conjecturally for any nondegenerate (Q, S)). We remark that the general morphism **g** in the above definition can be chosen from other periods of the triangle.

Definition 4.11. We say \mathbf{r}_{ϵ} (resp. \mathbf{l}_{ϵ}) can be generically lifted to a triangle at δ if the above \mathbf{R} (resp. \mathbf{L}) can be chosen as a general element of index $\mathbf{r}_{\epsilon}(\delta)$ (resp. $\mathbf{l}_{\epsilon}(\delta)$). To be more precise, this means that a general object \mathbf{R} (resp. \mathbf{L}) of index $\mathbf{r}_{\epsilon}(\delta)$ (resp. $\mathbf{l}_{\epsilon}(\delta)$) fits into the triangle (4.4) (resp. (4.5)) such that (\mathbf{M}, \mathbf{E}) is a general pair of index (δ, ϵ) and \mathbf{g} is a general morphism in $\mathcal{C}(\mathbf{E}, \Sigma \mathbf{M})$ (resp. $\mathcal{C}(\mathbf{M}, \mathbf{E})$).

We emphasis that in the above definition we do not ask any pair other than (\mathbf{M}, \mathbf{E}) to be general. For example, (\mathbf{M}, \mathbf{R}) or (\mathbf{R}, \mathbf{E}) may not be general as a pair.

Lemma 4.12. If $C(\Sigma^{-1}\mathbf{E}_2, \mathbf{E}_1) = 0$, then we have that $\mathbf{r}_{\mathbf{E}_1 \oplus \mathbf{E}_2} = \mathbf{r}_{\mathbf{E}_2}\mathbf{r}_{\mathbf{E}_1}$ and $\mathbf{l}_{\mathbf{E}_1 \oplus \mathbf{E}_2} = \mathbf{l}_{\mathbf{E}_1}\mathbf{l}_{\mathbf{E}_2}$. If in addition $C(\Sigma^{-1}\mathbf{E}_1, \mathbf{E}_2) = 0$ then $\mathbf{r}_{\mathbf{E}_1}$ and $\mathbf{r}_{\mathbf{E}_2}$ commute and $\mathbf{l}_{\mathbf{E}_1}$ and $\mathbf{l}_{\mathbf{E}_2}$ commute.

Proof. We only prove the statement for $\mathbf{r}_{\mathbf{E}}$ because the argument for $\mathbf{l}_{\mathbf{E}}$ is similar. Let $(\mathbf{f}_1, \mathbf{f}_2)$ be a general morphism in $\mathcal{C}(\Sigma^{-1}(\mathbf{E}_1 \oplus \mathbf{E}_2), \mathbf{M})$. Then \mathbf{f}_1 and \mathbf{f}_2 are general in $\mathcal{C}(\Sigma^{-1}\mathbf{E}_1, \mathbf{M})$ and $\mathcal{C}(\Sigma^{-1}\mathbf{E}_2, \mathbf{M})$. Apply $\mathcal{C}(\Sigma^{-1}\mathbf{E}_2, -)$ to the triangle in the second row, we get

$$\mathcal{C}(\Sigma^{-1}\mathbf{E}_2, \mathbf{M}) \to \mathcal{C}(\Sigma^{-1}\mathbf{E}_2, \mathbf{R}_1) \to \mathcal{C}(\Sigma^{-1}\mathbf{E}_2, \mathbf{E}_1) = 0.$$

So we may choose a morphism **g** general in $C(\Sigma^{-1}\mathbf{E}_2, \mathbf{R}_1)$ making the upper right square commute. Since the triangle of first row is split (in particular contractible), we can complete the first two rows to a nine-diagram below [18]. By construction and definition, the index of $\mathbf{R}_{1\oplus 2}$ is $\mathbf{r}_{\mathbf{E}_1\oplus\mathbf{E}_2}(\delta)$ while the index of $\mathbf{R}_{1,2}$ is $\mathbf{r}_{\mathbf{E}_2}\mathbf{r}_{\mathbf{E}_1}(\delta)$. But these two objects are isomorphic.



If in addition $C(\Sigma^{-1}\mathbf{E}_1, \mathbf{E}_2) = 0$, then $\mathbf{r}_{\mathbf{E}_1 \oplus \mathbf{E}_2} = \mathbf{r}_{\mathbf{E}_1}\mathbf{r}_{\mathbf{E}_2}$ so $\mathbf{r}_{\mathbf{E}_1}$ and $\mathbf{r}_{\mathbf{E}_2}$ commute.

5. The General Ranks

5.1. The Raising and Lowering Operators. Schofield introduced the general rank for quiver representations in his theory of general representations [23]. The following lemma is a straightforward generalization of [23, Lemma 5.1].

Lemma 5.1. Let A be a finite-dimensional algebra. Given any two irreducible closed sets X and Y in representation varieties of A, there is an open subset U of $X \times Y$ and a dimension vector γ such that for $(M, N) \in U$ we have that $\hom_A(M, N)$ is minimal and $\{\phi \in \operatorname{Hom}_A(M, N) \mid \operatorname{rank} \phi = \gamma\}$ is open and non-empty in $\operatorname{Hom}_A(M, N)$.

Below the algebra A will always be the Jacobian algebra of some quiver with potential.

Definition 5.2. If one of X and Y is a single representation, say $Y = \{E\}$, and X is the principal component $PC(\delta)$, then the above dimension vector is denoted by $rank(\delta, E)$. If $X = PC(\delta)$ and $Y = PC(\epsilon)$, then we call this γ is called the *general rank* from δ to ϵ , denoted by $rank(\delta, \epsilon)$. There are obvious variations if we replace δ or ϵ by a $\check{\delta}$ -vector.

Example 5.3. If (δ, ϵ) is an *exchange pair*, that is, δ and ϵ are indecomposable rigid with $e(\delta, \epsilon) = 1$ and $e(\epsilon, \delta) = 0$, then $rank(\epsilon, \tau \delta) \neq 0$ is the *c*-vector associated the exchange pair [9, Section 6]. For the connection to the corresponding concept in cluster algebras, we refer readers to [9, Remark 6.7].

For quivers with potentials, we have that $PC(\delta) = PC(\delta)$ by Theorem 3.11, so $rank(\delta, \epsilon) = rank(\delta, \epsilon)$.

Definition 5.4. For any decorated representation $\mathcal{E} = (E, E^-)$ of weight ϵ , we define the two operators $r_{\mathcal{E}}$ and $l_{\mathcal{E}}$ on the set of δ -vector as follows:

$$r_{\mathcal{E}}(\delta) = \delta + \epsilon + \operatorname{rank}(E, \tau \delta)B;$$
(5.1)

$$l_{\mathcal{E}}(\delta) = \delta - \check{\epsilon} + \operatorname{rank}(\delta, E)B.$$
(5.2)

We also define the two operators $r^{\mathcal{E}}$ and $l^{\mathcal{E}}$ on the set of $\check{\delta}$ -vector

$$r^{\mathcal{E}}(\check{\delta}) = \check{\delta} - \tau^{-1}\epsilon - \operatorname{rank}(\tau^{-1}E,\check{\delta})B;$$
$$l^{\mathcal{E}}(\check{\delta}) = \check{\delta} + \tau^{-1}\check{\epsilon} - \operatorname{rank}(\tau^{-1}\check{\delta},\tau^{-1}E)B.$$

If \mathcal{E} is general of weight ϵ , then we will write ϵ instead of \mathcal{E} in $r_{\mathcal{E}}$ and $l_{\mathcal{E}}$.

We'd like to make it clear as early as possible that the main characters of this paper are r_{ϵ} and l_{ϵ} rather than r_{ε} and l_{ε} . We could have defined r_{ϵ} and l_{ϵ} without mentioning r_{ε} and l_{ε} . It follows directly from the definition that

$$r_{\epsilon}(\delta) = l_{\tau\delta}(\epsilon) \quad \text{and} \quad r^{\epsilon}(\check{\delta}) = l^{\tau\delta}(\check{\epsilon}).$$
 (5.3)

Let **E** be a lift of \mathcal{E} in $\mathcal{C}_{Q,S}$. We pick a general element **M** in $\mathcal{C}_{Q,S}$ of index δ . Let **g** be a general morphism in $\mathcal{C}(\mathbf{E}, \Sigma \mathbf{M})$. We complete **g** to a triangle $\Sigma^{-1}\mathbf{E} \to \mathbf{M} \to \mathbf{R} \to \mathbf{E}$, which gives a long exact sequence as in Corollary 4.9:

$$\cdots \to \tau^{-1} E \xrightarrow{g_{-1}} M \to R \to E \xrightarrow{g} \tau M \cdots$$

By Lemma 4.5 (for void mutation) **g** will descend to a general morphism $g \in \text{Hom}_J(E, \tau M)$. Recall the δ -vector of \mathcal{R} is nothing but the index of **R**. So by Lemma 4.8

$$\delta_{\mathcal{R}} = \delta_{\mathcal{M}} + \delta_{\mathcal{E}} + \operatorname{rank}(g)B = \delta + \epsilon + \operatorname{rank}(E, \tau\delta)B = r_{\mathcal{E}}(\delta).$$

Since $\Sigma^{-1}\mathbf{g}$ is also general in $\mathcal{C}(\Sigma^{-1}\mathbf{E},\mathbf{M})$, similarly we have that

$$\check{\delta}_{\mathcal{R}} = \check{\delta}_{\mathcal{M}} + \check{\delta}_{\mathcal{E}} - \operatorname{rank}(g_{-1})B = \check{\delta} + \check{\epsilon} - \operatorname{rank}(\tau^{-1}E, \delta)B = r^{\mathcal{E}}(\check{\delta}).$$

Note that this $r_{\mathcal{E}}(\delta)$ is equal to $\mathbf{r}_{\mathbf{E}}(\delta)$ from Definition 4.10. If \mathcal{E} is general of weight ϵ , then $r_{\mathcal{E}}(\delta) = r_{\epsilon}(\delta)$ also equals to the $\mathbf{r}_{\epsilon}(\delta)$. We have the similar discussion for $l_{\mathcal{E}}(\delta)$. We summarize in the following lemma.

Lemma 5.5. We have that $\mathbf{r}_{\mathbf{E}}(\delta) = r_{\mathcal{E}}(\delta)$ and $\mathbf{l}_{\mathbf{E}}(\delta) = l_{\mathcal{E}}(\delta)$; $\mathbf{r}_{\epsilon}(\delta) = r_{\epsilon}(\delta)$ and $\mathbf{l}_{\epsilon}(\delta) = l_{\epsilon}(\delta)$.

Recall the direct sum notation for δ -vectors in the end of Section 3.

Proposition 5.6. For any two weight vectors ϵ_1 and ϵ_2 of (Q, S), the following are equivalent:

- (1) $\epsilon = \epsilon_1 \oplus \epsilon_2$;
- $\begin{array}{ll} (2) & r_{\epsilon} = r_{\epsilon_1} r_{\epsilon_2} = r_{\epsilon_2} r_{\epsilon_1}; \\ (3) & l_{\epsilon} = l_{\epsilon_1} l_{\epsilon_2} = l_{\epsilon_2} l_{\epsilon_1}. \end{array}$

Proof. We only show the equivalence of (1) and (2). If $\epsilon = \epsilon_1 \oplus \epsilon_2$, then $e(\epsilon_1, \epsilon_2) = e(\epsilon_2, \epsilon_1) = 0$ by [7, Theorem 4.4]. By Corollary 4.3.(3) we have that $\mathcal{C}(\mathbf{E}_1, \Sigma \mathbf{E}_2) = \mathcal{C}(\mathbf{E}_2, \Sigma \mathbf{E}_1) = 0$ for \mathbf{E}_i general of index ϵ_i (i = 1, 2). By Lemma 4.12 we have that $\mathbf{r}_{\mathbf{E}_1}\mathbf{r}_{\mathbf{E}_2} = \mathbf{r}_{\mathbf{E}_1\oplus\mathbf{E}_2} = \mathbf{r}_{\mathbf{E}_2}\mathbf{r}_{\mathbf{E}_1}$. By Lemma 5.5 we get $r_{\epsilon} = r_{\epsilon_1}r_{\epsilon_2} = r_{\epsilon_2}r_{\epsilon_1}$.

Conversely, if $\epsilon \neq \epsilon_1 \oplus \epsilon_2$, then at least one of $e(\epsilon_1, \epsilon_2)$ and $e(\epsilon_2, \epsilon_1)$ is nonzero, say $e(\epsilon_2, \epsilon_1) \neq 0$, so rank $(\epsilon_1, \tau \epsilon_2) \neq 0$. Let δ be the zero vector, then $r_{\epsilon_2}(\delta) = \epsilon_2$ and $r_{\epsilon}(\delta) = \epsilon_1 + \epsilon_2$. But $r_{\epsilon_1}(\epsilon_2) = \epsilon_1 + \epsilon_2 + \operatorname{rank}(\epsilon_1, \tau \epsilon_2) B$ with $\operatorname{rank}(\epsilon_1, \tau \epsilon_2) \neq 0$. Hence $r_{\epsilon} \neq r_{\epsilon_1} r_{\epsilon_2}$.

This proposition suggest that to study r_{ϵ} and l_{ϵ} it is harmless to assume ϵ is *indecomposable*, that is, a general presentation of weight ϵ is indecomposable.

It is unclear if we can ask \mathcal{R} to be general of weight $r_{\epsilon}(\delta)$. However, this can be done if r_{ϵ} can be generically lifted to $\mathcal{C}_{Q,S}$ at δ in the sense of Definition 4.11. By abuse of language, in this case we also say r_{ϵ} can be generically lifted at δ .

Lemma 5.7. Suppose that the operator r_{ϵ} (or l_{ϵ}) can be generically lifted at δ . Then the operator commutes with any sequence of mutations $\mu_{\mathbf{u}}$ and τ^i :

$$\begin{split} \mu_{\mathbf{u}}(r_{\epsilon}(\delta)) &= r_{\mu_{\mathbf{u}}(\epsilon)}(\mu_{\mathbf{u}}(\delta)) & and & \mu_{\mathbf{u}}(l_{\epsilon}(\delta)) &= l_{\mu_{\mathbf{u}}(\epsilon)}(\mu_{\mathbf{u}}(\delta)); \\ \tau^{i}(r_{\epsilon}(\delta)) &= r_{\tau^{i}\epsilon}(\tau^{i}\delta) & and & \tau^{i}(l_{\epsilon}(\delta)) &= l_{\tau^{i}\epsilon}(\tau^{i}\delta). \end{split}$$

Proof. We only prove the statement for l_{ϵ} . By definition, a general object **L** of index $\mathbf{l}_{\epsilon}(\delta)$ fits into a triangle $\Sigma^{-1}\mathbf{E} \to \mathbf{L} \to \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E}$ with the properties given in Definition 4.11. Then by Lemma 5.5 $\mathbf{l}_{\epsilon}(\delta) = l_{\epsilon}(\delta)$. Let $\mu_{\mathbf{u}}^- : \mathcal{C}_{Q,S} \to \mathcal{C}_{Q',S'} = \mathcal{C}_{\mu_{\mathbf{u}}(Q,S)}$ be the triangle equivalence corresponding to the sequence of mutations. Apply $\mu_{\mathbf{u}}^-$ to the above triangle, and we get another triangle $\Sigma^{-1}\mathbf{E}' \to \mathbf{L}' \to \mathbf{M}' \xrightarrow{\mathbf{g}'} \mathbf{E}'$ in $\mathcal{C}_{Q',S'}$. By Theorem 4.4, $\mathcal{M}' = \mu_{\mathbf{u}}(\mathcal{M})$. Since \mathcal{M} is general of weight δ , $\delta_{\mathcal{M}'} = \mu_{\mathbf{u}}(\delta_{\mathcal{M}}) = \mu_{\mathbf{u}}(\delta)$ by Lemma 3.12. For the same reason, $\delta_{\mathcal{E}'} = \mu_{\mathbf{u}}(\epsilon)$ and $\delta_{\mathcal{L}'} = \mu_{\mathbf{u}}(l_{\epsilon}(\delta))$. By Lemma 4.5, the induced map $g': M' \to E'$ is general, so $\delta_{\mathcal{L}'} = l_{\mu_{\mathbf{u}}(\epsilon)}(\mu_{\mathbf{u}}(\delta))$ by definition. Hence, $\mu_{\mathbf{u}}(l_{\epsilon}(\delta)) = l_{\mu_{\mathbf{u}}(\epsilon)}(\mu_{\mathbf{u}}(\delta)).$

For the statement about τ , let us look at the long exact sequence

$$\cdots \to \tau^{-1}L \to \tau^{-1}M \xrightarrow{\tau^{-1}g} \tau^{-1}E \to L \to M \xrightarrow{g} E \to \tau L \to \tau M \xrightarrow{\tau g} \tau E \to \cdots .$$
(5.4)

As \mathcal{L}, \mathcal{M} and \mathcal{E} are general of weight $l_{\epsilon}(\delta), \delta$ and $\epsilon, \tau^i \mathcal{L}, \tau^i \mathcal{M}$ and $\tau^i \mathcal{E}$ are general of weight $\tau^i l_{\epsilon}(\delta), \tau^i \delta$ and $\tau^i \epsilon$ by Theorem 3.11. Since $\mathbf{g} \in \mathcal{C}(\mathbf{M}, \mathbf{E})$ is general, so is each $\Sigma^i \mathbf{g}$, and so is each $\tau^i g$ by Lemma 4.5. Hence $\tau^i \mathcal{L}$ has weight $l_{\tau^i \epsilon}(\tau^i \delta)$, and the equality $\tau^i(l_{\epsilon}(\delta)) = l_{\tau^i \epsilon}(\tau^i \delta)$ follows.

5.2. Sufficient Conditions for the Generic Lifting. Now we explore some sufficient conditions such that r_{ϵ} and l_{ϵ} can be generically lifted.

Lemma 5.8. Any square given by a quotient presentation



with $\operatorname{coker}(d_M) = M$ and $\operatorname{coker}(d_N) = N$, can be lifted to a nine-diagram



such that each $\mathbf{T}_{\pm}^* \in \operatorname{add} \mathbf{T}$ for $* = \mathbf{L}, \mathbf{M}$ or \mathbf{N} , and $\mathbf{T}_{-}^{\mathrm{L}} \xrightarrow{\mathbf{d}_{\mathrm{L}}} \mathbf{T}_{+}^{\mathrm{L}} \rightarrow \mathbf{L}$ lifts the corresponding subpresentation as well.

Proof. This lemma is a straightforward variation of [20, Lemma 3.2]. The point is that the quotient presentation together with its subpresentation can be lifted to the upper two rows of the diagram. These rows are split triangles so can be completed to the nine diagram (for any \mathbf{d}_{L}) by [18].

Lemma 5.9. If rank $(\delta, \epsilon) = 0$, then l_{ϵ} can be generically lifted at δ . That is, a general object \mathbf{L} of index $\delta + \tau^{-1}\epsilon$ fits into a triangle $\Sigma^{-1}\mathbf{E} \to \mathbf{L} \xrightarrow{\mathbf{f}} \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E}$, where (\mathbf{M}, \mathbf{E}) is a general pair of index (δ, ϵ) and \mathbf{g} is a general morphism in $\mathcal{C}(\mathbf{M}, \mathbf{E})$.

Proof. rank $(\delta, \epsilon) = 0$ is equivalent to hom $(\delta, \epsilon) = 0$. So $e(\tau^{-1}\epsilon, \delta) = 0$. By Theorem 3.3 a general presentation d_L in Hom_J $(P([-\tau^{-1}\epsilon]_+ + [-\delta]_+), P([\tau^{-1}\epsilon]_+ + [\delta]_+]))$ has a quotient presentation d_M of weight δ , and thus a subpresentation d_1 of weight $\tau^{-1}\epsilon$. Note that the presentation d_L is homotopy equivalent to a general presentation of weight $l_{\epsilon}(\delta) = \delta + \tau^{-1}\epsilon$ by Lemma 3.1. Moreover, by Lemma 3.4 we may assume d_M and $d_{\tau^{-1}E}$ are general as a pair, and η is general in $E(d_M, d_{\tau^{-1}E})$. We lift the quotient presentation $d_L \to d_M$ to a nine diagram as in Lemma 5.8:



By Construction **L** is general of index $l_{\epsilon}(\delta)$, (**M**, **E**) is a general pair of index (δ, ϵ) . By Lemma 4.1 the element $\eta \in E(d_2, d_1)$ is now identified with a general element **g** in $(\Sigma \mathbf{T})(\mathbf{M}, \mathbf{E})$. But by Corollary 4.3 every morphism $\mathcal{C}(\mathbf{M}, \mathbf{E})$ factors through $\Sigma \mathbf{T}$. Hence, **g** is in fact a general element in $\mathcal{C}(\mathbf{M}, \mathbf{E})$.

Remark 5.10. (1). Replacing δ and ϵ by ϵ and $\tau\delta$ in Lemma 5.9, we get the following statement: if rank($\epsilon, \tau\delta$) = 0, then \mathbf{r}_{ϵ} can be generically lifted at δ .

(2). With a little more effort, we can show that \mathbf{f} can be assumed to general in $\mathcal{C}(\mathbf{L}, \mathbf{M})$. The proof will start with taking a general morphism $\mathbf{f} : \mathbf{L} \to \mathbf{M}$, then by Corollary 4.9 f is surjective. Then the rest of the proof is similar to that of Lemma 5.11 below. However this by no means says that rank(f) is the general rank from $l_{\epsilon}(\delta)$ to δ because (\mathbf{L}, \mathbf{M}) as a pair may not be in general positions.

Lemma 5.11.

- If rank(δ, ε) = dim(ε), then l_ε can be generically lifted at δ. That is, a general object L of index δ ε fits into a triangle L → M ^g→ E ^h→ ΣL, where (M, E) is a general pair of index (δ, ε) and g is a general morphism in C(M, E).
- (2) If rank $(\delta, \epsilon) = \underline{\dim}(\delta)$, then \mathbf{l}_{ϵ} can be generically lifted at δ .

Proof. (1). Take a general pair (\mathbf{M}, \mathbf{E}) and a general morphism $\mathbf{g} : \mathbf{M} \to \mathbf{E}$, which descends to a general homomorphism $g : M \to E$ by Lemma 4.5. By the rank condition, g is surjective. We complete \mathbf{g} to a triangle $\mathbf{L} \to \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E} \xrightarrow{\mathbf{h}} \Sigma \mathbf{L}$. By Lemma 4.8 the index of \mathbf{L} is $\delta - \epsilon$. We need to show that \mathbf{L} may be assumed to be general of index $\delta - \epsilon$.

Let $\mathbf{T}_{-}^{\mathrm{E}} \xrightarrow{\mathbf{d}_{\mathrm{E}}} \mathbf{T}_{+}^{\mathrm{E}} \to \mathbf{E} \to \Sigma \mathbf{T}_{-}^{\mathrm{E}}$ and $\mathbf{T}_{-}^{\mathrm{L}} \xrightarrow{\mathbf{d}_{\mathrm{L}}} \mathbf{T}_{+}^{\mathrm{L}} \to \mathbf{L} \to \Sigma \mathbf{T}_{-}^{\mathrm{L}}$ be two triangles with $\mathbf{d}_{\mathrm{E}} \in \mathrm{THom}(\epsilon)$ and $\mathbf{d}_{\mathrm{L}} \in \mathrm{THom}(\delta - \epsilon)$. We set $\mathbf{T}_{\pm}^{\mathrm{M}} = \mathbf{T}_{\pm}^{\mathrm{E}} \oplus \mathbf{T}_{\pm}^{\mathrm{L}}$. As h = 0, \mathbf{h} must factor through $\Sigma \mathbf{T}$. Since $\mathcal{C}(\mathbf{T}, \Sigma \mathbf{T}) = 0$, $\mathbf{T}_{+}^{\mathrm{E}} \to \mathbf{E} \xrightarrow{\mathbf{h}} \Sigma \mathbf{L}$ vanishes, so $\mathbf{T}_{+}^{\mathrm{E}} \to \mathbf{E}$ factor through \mathbf{M} . This gives the following square:

$$\begin{array}{c} \mathbf{T}^{\mathrm{M}}_{+} \longrightarrow \mathbf{T}^{\mathrm{E}}_{+} \\ \downarrow & \downarrow \\ \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E} \end{array}$$

which can be completed to a nine-diagram



Apply $\mathcal{C}(\mathbf{T}, -)$ we get the following diagram

$$P_{-}^{L} \longrightarrow P_{-}^{M} \xrightarrow{\pi_{-}} P_{-}^{E}$$

$$d_{L} \downarrow \qquad d_{M} \downarrow \qquad d_{E} \downarrow$$

$$P_{+}^{L} \longrightarrow P_{+}^{M} \xrightarrow{\pi_{+}} P_{+}^{E}$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$L \longrightarrow M \xrightarrow{g} E$$

In particular, we get for a pair of general presentations d_M and d_E of weight δ and ϵ , a general morphism from d_M to d_E is surjective. Finally, a similar argument as in the proof of Lemma 3.4 shows that the kernel d_L of such a morphism is general as well. That is equivalent to say **L** may be assumed to be general of index $\delta - \epsilon$.

(2). Note that if an object **L** has index $\mathbf{l}_{\epsilon}(\delta) = \check{\delta} - \check{\epsilon}$ (by (5.2) and (2.1)), then $\Sigma \mathbf{L}$ has coindex $\check{\epsilon} - \check{\delta}$. So here it is convenient to work with the following obvious variant of Definition 4.11: a general object $\Sigma \mathbf{L}$ of coindex $\check{\epsilon} - \check{\delta}$ fits into a triangle $\mathbf{L} \xrightarrow{\mathbf{f}} \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E} \to \Sigma \mathbf{L}$, where (\mathbf{M}, \mathbf{E}) is

general pair of coindex $(\delta, \check{\epsilon})$ and **g** is a general morphism in $\mathcal{C}(\mathbf{M}, \mathbf{E})$. The proof is then similar to (1).

Remark 5.12. With little more effort, we can show that **h** in Lemma 5.11.(1) is a general morphism in $\mathcal{C}(\mathbf{E}, \Sigma \mathbf{L})$, and **f** in (2) is a general morphism in $\mathcal{C}(\mathbf{L}, \mathbf{M})$. To see this, we take **h** as an example. Note that we do not claim that **g** and **h** are general as a pair, so we can start the proof all over again. Since rank $(\delta, \epsilon) = \underline{\dim}(\epsilon)$, a general representation M in PC (δ) has a quotient representation in PC (ϵ) . So $\mathbf{e}(\delta - \epsilon, \epsilon) = 0$ by Theorem 3.3. Then the same argument as in Lemma 5.9 shows that **h** can be assumed to be general. But rank(h) may not be the general rank from ϵ to $\tau(l_{\epsilon}(\delta))$ because $(\mathbf{E}, \Sigma \mathbf{L})$ may not be general as a pair.

Definition 5.13. We say (δ, ϵ) has completely extremal rank if any of the following occurs:

 $\operatorname{rank}(\delta, \epsilon) = 0$, $\operatorname{rank}(\delta, \epsilon) = \underline{\dim}(\delta)$, $\operatorname{rank}(\delta, \epsilon) = \underline{\dim}(\epsilon)$.

To summarize what Lemmas 5.9 and 5.11 say: if (δ, ϵ) has completely extremal rank, then l_{ϵ} can be generically lifted at δ . So by Lemma 5.7 these operators commutes with mutations and τ^i . The same statement also holds for r_{ϵ} and the other two operators l^{ϵ} and r^{ϵ} . We summarize together in the proposition below.

Proposition 5.14. If the left column has the completely extremal rank, then the right column commutes with any sequence of mutations and τ^i in the sense of Lemma 5.7.

$\operatorname{rank}(\epsilon, \tau \delta)$	$r_{\epsilon}(\delta),$
$\mathrm{rank}(\delta,\epsilon)$	$l_{\epsilon}(\delta),$
$\operatorname{rank}(\tau^{-1}\epsilon,\check{\delta})$	$r^{\epsilon}(\check{\delta}),$
$\operatorname{rank}(\tau^{-1}\check{\delta},\tau^{-1}\epsilon)$	$l^{\epsilon}(\check{\delta}).$

Definition 5.15. An extended mutation sequence is a composition of ordinary mutations μ_u and the AR-translation τ or its inverse τ^{-1} . We also denote τ and τ^{-1} by μ_+ and μ_- respectively, though they are not involutions in general.

As τ commutes with mutations (Lemma 2.7), we can say the right column commutes with any extended sequence of mutations in the conclusion of Proposition 5.14. Moreover, Theorem 4.4 can be naturally extended if we take Σ as the lifted mutation for τ .

5.3. The Main Results.

Theorem 5.16. Let ϵ and δ be two weight vectors for a quiver with potential (Q, \mathcal{S}) .

(1) Suppose that there is an extended sequence of mutation $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\epsilon), \mu_{\mathbf{u}}(\tau\delta))$ has completely extremal rank r. Then there is an exact sequence

$$\cdots \to \tau^{-1}M \xrightarrow{f_{-1}} \tau^{-1}R \xrightarrow{g_{-1}} \tau^{-1}E \xrightarrow{h_{-1}} M \xrightarrow{f_0} R \xrightarrow{g_0} E \xrightarrow{h_0} \tau M \xrightarrow{f_1} \tau R \xrightarrow{g_1} \tau E \xrightarrow{h_1} \tau^2 M \to \cdots,$$
(5.5)

where R is general of weight $r_{\epsilon}(\delta)$, (E, M) is general as a pair of weights δ and ϵ , and h_i is a general homomorphism in $\text{Hom}_J(\tau^i E, \tau^{i+1}M)$.

(2) Suppose that there is an extended sequence of mutation $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ has completely extremal rank r. Then there is an exact sequence

$$\cdots \to \tau^{-1}L \xrightarrow{f_{-1}} \tau^{-1}M \xrightarrow{g_{-1}} \tau^{-1}E \xrightarrow{h_{-1}}L \xrightarrow{f_0} M \xrightarrow{g_0} E \xrightarrow{h_0} \tau L \xrightarrow{f_1} \tau M \xrightarrow{g_1} \tau E \xrightarrow{h_1} \tau^2 L \to \cdots,$$
(5.6)

where L is general of weight $l_{\epsilon}(\delta)$, (M, E) is general as a pair of weight δ and ϵ , and g_i is a general homomorphism in Hom_J $(\tau^i M, \tau^i E)$.

Proof. We will prove the statement (2) only because (1) can be proved in a similar fashion. After application of the extended mutation sequence, we are in one of the three situations of Lemmas 5.9 and 5.11. So we get a triangle $\mathbf{L}' \xrightarrow{\mathbf{f}'} \mathbf{M}' \xrightarrow{\mathbf{g}'} \mathbf{E}' \xrightarrow{\mathbf{h}'} \Sigma \mathbf{L}'$ in $\mathcal{C}_{Q',\mathcal{S}'}$, which generically lift $l_{\epsilon'}$ at δ' where $\delta' = \mu_{\mathbf{u}}(\delta) = \operatorname{ind} \mathbf{M}'$ and $\epsilon' = \mu_{\mathbf{u}}(\epsilon) = \operatorname{ind} \mathbf{E}'$.

By Theorem 4.4 the extended sequence of mutations corresponds to a sequence of triangle equivalence. Under this equivalence, we get a triangle $\mathbf{L} \xrightarrow{\mathbf{f}} \mathbf{M} \xrightarrow{\mathbf{g}} \mathbf{E} \xrightarrow{\mathbf{h}} \Sigma \mathbf{L}$ with \mathbf{L} general of index $l_{\epsilon}(\delta)$, (\mathbf{M}, \mathbf{E}) is a general pair of indices (δ, ϵ) , and \mathbf{g} is general in $\mathcal{C}(\mathbf{M}, \mathbf{E})$. Apply the functor $F = \mathcal{C}(\mathbf{T}, -)$ to this triangle, we get the desired long exact sequence (5.5) with desired generic condition by Lemma 4.5.

Remark 5.17. (1). In fact we can also say something about f_i and h_i in (5.6). If r = 0 or $\underline{\dim}(\mu_{\mathbf{u}}(\delta))$, then we may assume that f_i is general in $\operatorname{Hom}_J(\tau^i L, \tau^i M)$; if $r = \underline{\dim}(\mu_{\mathbf{u}}(\epsilon))$, then we may assume that h_i is general in $\operatorname{Hom}_J(\tau^i E, \tau^{i+1}L)$ (see Remarks 5.10 and 5.12). As remarked there, the ranks of those morphisms may be greater than the general ranks between the corresponding principal components (see Example 5.19 below).

(2). In view of Conjecture 5.21 below, the assumption of existence of such a mutation sequence may not be necessary.

Example 5.18. If $\delta = [\epsilon]_+$, then $l_{\epsilon}(\delta) = [\epsilon]_+ - \check{\epsilon} + \operatorname{rank}([\epsilon]_+, \epsilon)B = [\epsilon]_+ - \epsilon = [-\epsilon]_+$, and the projective presentation of E is a part of the above sequence. Similarly we have that $r_{\epsilon}([-\epsilon]_+) = [\epsilon]_+$.

Example 5.19. In general, we cannot assume (M, R) and (R, E) are general pairs for (1); and cannot assume (L, M) and (E, L) are general pairs for (2). Consider the 3-arrow Kronecker quiver $1 \Longrightarrow 2$. Let $\delta = (0, 1)$ and $\epsilon = (1, -2)$. It is easy to check that $\operatorname{ext}(\delta, \epsilon) = 0$ so $r_{\epsilon}(\delta) = \delta + \epsilon = (1, -1)$. If (R, E) were a general pair, then $\operatorname{rank}(g_0) = \operatorname{rank}(r_{\epsilon}(\delta), \epsilon) = (0, 0)$, and the sequence (5.5) could not be exact at E. We know that a general representation R of weight (1, -1) has a quotient representation E of weight (1, -2) but it is intuitively clear that (R, E) is not a general pair.

Combining Proposition 5.14 and Lemma 5.7, we obtain an algorithm to compute the general rank from δ to ϵ for nondegenerate quivers with potentials. We will see that the algorithm works effectively as long as one of δ and ϵ is reachable, and works to some extent if Conjecture 5.21 is true. Before the invention of Derksen-Weyman-Zelevinsky's mutation, the calculation of rank (δ, ϵ) seems completely out of reach even for acyclic quivers.

Before we state the algorithm, we describe a "frozen-vertex" trick which is needed in the algorithm. An *ice quiver* is a quiver with a set of special vertices, called frozen vertices, which are forbidden to mutate. The extended *B*-matrix of an ice quiver *Q* is the submatrix of the *B*-matrix of *Q* (viewed as an ordinary quiver) given by the rows indexed by the mutable vertices. For any quiver *Q*, we can always add to *Q* some frozen vertices *V* together with some arrows from *V* to Q_0 such that the extended *B*-matrix of the new quiver \widetilde{Q} has full rank. Moreover, for each weight vector δ of (Q, S), a general representation of (\widetilde{Q}, S) of weight $\widetilde{\delta} = (\delta, 0, \ldots, 0)$ is the extension by zeros of a general representation of weight δ . In particular, $\operatorname{rank}(\widetilde{\delta}, \widetilde{\epsilon}) = (\operatorname{rank}(\delta, \epsilon), 0, \ldots, 0)$. A standard way to achieve this is that for each vertex $u \in Q_0$ we add a frozen vertex u' with an arrow from u' to u.

Algorithm 5.20. To find rank (δ, ϵ) we follow the three steps.

Step 0: If B_Q has full rank then go to Step 1. Otherwise we apply the above frozen vertex trick. To ease our notation below, we will denote \tilde{Q} , $\tilde{\delta}$ and $\tilde{\epsilon}$ still by Q, δ and ϵ .

Step 1: Find a sequence of mutations $\mu_{\mathbf{u}}$ such that $r = \operatorname{rank}(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ is completely extremal. Conjecturally such a sequence of mutations always exists.

Step 2. According to Proposition 5.14 and (2.1) we have that

$$l_{\epsilon}(\delta) = \begin{cases} \mu_{\mathbf{u}}^{-1}(\delta' - \check{\epsilon}') & \text{if } r = 0; \\ \mu_{\mathbf{u}}^{-1}(\delta' - \epsilon') & \text{if } r = \underline{\dim}(\epsilon'); \\ \mu_{\mathbf{u}}^{-1}(\check{\delta}' - \check{\epsilon}') & \text{if } r = \underline{\dim}(\delta'), \end{cases}$$

where $\delta' = \mu_{\mathbf{u}}(\delta)$ and $\epsilon' = \mu_{\mathbf{u}}(\epsilon)$. Finally we have that $\operatorname{rank}(\delta, \epsilon)$ is the unique vector $r \in \mathbb{Z}^{Q_0}$ such that

$$rB_Q = l_\epsilon(\delta) - \delta + \check{\epsilon}. \tag{5.7}$$

It is unclear if the sequence in Step 1 always exists. We conjecture the existence of such a sequence (see Conjecture 5.21). This algorithm will be improved in Theorem 6.1 where the formula is division-free.

Conjecture 5.21. Let (Q, S) be a nondegenerate Jacobi-finite QP. For any pair (δ, ϵ) of δ -vectors of (Q, S), there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ has completely extremal rank.

We are not completely confidant about this conjecture. But we are quite confidant about a related conjecture.

Conjecture 5.22. Let (Q, S) be a nondegenerate Jacobi-finite QP. For any pair (δ, ϵ) of δ -vectors of (Q, S), there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\epsilon))$ is either hom-vanishing or e-vanishing.

Clearly rank $(\delta, \epsilon) = 0$ is equivalent to hom $(\delta, \epsilon) = 0$. We suspect that rank $(\delta, \epsilon) = \dim(\epsilon)$ might imply $e(\delta, \epsilon) = 0$ so that Conjecture 5.21 imply Conjecture 5.22. At least the implication holds for acyclic quivers. This follows from the fact that

$$\operatorname{ext}_Q(\alpha,\beta) = \operatorname{ext}_Q(\alpha,\beta-\gamma) = \operatorname{ext}_Q(\alpha-\gamma,\beta),$$

where γ is the general rank from rep_{α}(Q) to rep_{β}(Q). This fact was proved in the proof of [23, Theorem 5.4].

Question 5.23. For a nondegenerate Jacobi-finite QP, do we have that $e(\delta, \epsilon) = e(\delta, N_l)$ where N_l is the cokernel of a general morphism $\delta \to \epsilon$?

It is easy to see that the answer is positive if ϵ is rigid. It is also easy to show that rank $(\delta, \epsilon) = \dim(\epsilon)$ implies $e(\delta, \epsilon) = 0$ if ϵ is rigid.

Another related conjecture is the following. We say that a weight vector δ has projective dimension ≤ 1 if a general presentations in $PHom(\delta)$ is injective.

Conjecture 5.24. Let (Q, S) be a nondegenerate Jacobi-finite QP. For any δ -vector δ of (Q, S), there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $\mu_{\mathbf{u}}(\delta)$ has projective dimension ≤ 1 .

Example 5.25. Let Q be the quiver $1 \Longrightarrow 2 \longrightarrow 3$, and α, β be the dimension vectors (6, 9, 8) and (3, 5, 2). The *B*-matrix is not of full rank so we add a frozen vertex 3' with an arrow from 3' to 3. Then we replace α by (6, 9, 8, 0) and β by (3, 5, 2, 0). By multiplying the Euler matrix

of the quiver, we find that $PC(\delta) = rep_{\alpha}(Q)$ and $PC(\check{\epsilon}) = rep_{\beta}(Q)$ for $\delta = (6, -3, -1, 0)$ and $\check{\epsilon} = (-7, 3, 2, -2)$.

We notice that δ is negative reachable by the sequence $\mathbf{u} = (3, 2, 1, 2, 1, 3)$ of mutations. We find that $\delta' = -e_3$ and $\check{\epsilon}' = (1, 2, -1, 0)$, so $l_{\epsilon}(\delta) = \mu_{\mathbf{u}}^{-1}(-1, -2, 0, 0) = (7, -4, 0, 0)$. Hence the general rank from $\operatorname{rep}_{\alpha}(Q)$ to $\operatorname{rep}_{\beta}(Q)$ is (2, 3, 2).

Example 5.26. The following quiver cannot be mutated to an acyclic quiver



We put the potential S = abc. Let us compute the general rank from $\delta = (-1, 3, -2, 0)$ to $\check{\epsilon} = (1, -9, 1, 4)$. Using an algorithm in [9] we find $\underline{\dim}(\delta) = (3, 3, 4, 5)$ and $\underline{\dim}(\check{\epsilon}) = (5, 3, 2, 5)$. As $\delta(\underline{\dim}(\delta))$ and $\epsilon(\underline{\dim}(\epsilon))$ are not positive, δ and $\check{\epsilon}$ are not rigid (in particular not extended-reachable). However, we can apply the sequence of mutations $\mathbf{u} = (4, 1, 2, 4, 3)$ to them such that $(\delta', \check{\epsilon}') := (\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\check{\epsilon}))$ is hom-vanishing. To see this, we find by Lemma 2.5 that $\delta' = (0, -1, -2, 3), \ \check{\epsilon}' = (-1, 1, -1, -1), \ \text{and } \underline{\dim}(\check{\epsilon}') = (1, 1, 1, 0)$. Since $[\delta']_+(\underline{\dim}(\check{\epsilon}')) = 0$, we see that $\hom(\delta', \check{\epsilon}') = 0$. So $\mu_{\mathbf{u}}(l_{\epsilon}(\delta)) = \delta' - \check{\epsilon}' = (1, -2, -1, 4)$. Apply the mutation backward, and we find $l_{\epsilon}(\delta) = (-3, 5, 1, -1)$. Finally by (5.7) we get $\operatorname{rank}(\delta, \check{\epsilon}) = (2, 2, 1, 4)$.

If ϵ is extended-reachable, the rank condition in Theorem 5.16 is trivially satisfied. In addition, the rigidity of ϵ makes the situation particularly nice. The following two corollaries will play a crucial role in [10].

Corollary 5.27. Assume that ϵ is extended-reachable. Then

(1) we may assume that both (R, E) and (E, M) are general pairs in the sequence

 $\cdots \to \tau^{-1}M \xrightarrow{f_{-1}} \tau^{-1}R \xrightarrow{g_{-1}} \tau^{-1}E \xrightarrow{h_{-1}} M \xrightarrow{f_0} R \xrightarrow{g_0} E \xrightarrow{h_0} \tau M \xrightarrow{f_1} \tau R \xrightarrow{g_1} \tau E \xrightarrow{h_1} \tau^2 M \to \cdots$ In particular, the rank of each g_i is the general rank from $\tau^i(r_{\epsilon}(\delta))$ to $\tau^i \epsilon$. Moreover, we have that

 $r_{\epsilon}(\delta) = \delta + \check{\epsilon} - \operatorname{rank}(g_0)B$ and $\operatorname{hom}_J(r_{\epsilon}(\delta), E) = \operatorname{hom}_J(\delta, E) + \check{\epsilon}(\operatorname{rank}(g_0)),$

 $r_{\epsilon}(\delta) = \delta + \epsilon + \operatorname{rank}(h_0)B$ and $e_J(r_{\epsilon}(\delta), E) = e_J(\delta, E) - \epsilon(\operatorname{rank}(h_0)),$

 $r^{\epsilon}(\check{\delta}) = \check{\delta} - \tau^{-1}\epsilon - \operatorname{rank}(h_{-1})B \quad and \quad \operatorname{hom}_{J}(\tau^{-1}E, r^{\epsilon}(\check{\delta})) = \operatorname{hom}_{J}(\tau^{-1}E, \check{\delta}) - \tau^{-1}\epsilon(\operatorname{rank}(h_{-1})),$ $r^{\epsilon}(\check{\delta}) = \check{\delta} - \tau^{-1}\check{\epsilon} + \operatorname{rank}(g_{-1})B \quad and \quad \check{e}_{J}(\tau^{-1}E, r^{\epsilon}(\check{\delta})) = \check{e}_{J}(\tau^{-1}E, \check{\delta}) + \tau^{-1}\check{\epsilon}(\operatorname{rank}(g_{-1})).$

(2) we may assume that both (M, E) and (E, L) are general pairs in the sequence

$$\cdots \to \tau^{-1}L \xrightarrow{f_{-1}} \tau^{-1}M \xrightarrow{g_{-1}} \tau^{-1}E \xrightarrow{h_{-1}}L \xrightarrow{f_0} M \xrightarrow{g_0} E \xrightarrow{h_0} \tau L \xrightarrow{f_1} \tau M \xrightarrow{g_1} \tau E \xrightarrow{h_1} \tau^2 L \to \cdots$$

In particular, the rank of each h_i is the general rank from $\tau^i \epsilon$ to $\tau^{i+1}(l_{\epsilon}(\delta))$. Moreover, we have that

$$l_{\epsilon}(\delta) = \delta - \check{\epsilon} + \operatorname{rank}(g_0)B$$
 and $\operatorname{hom}_J(l_{\epsilon}(\delta), E) = \operatorname{hom}_J(\delta, E) - \check{\epsilon}(\operatorname{rank}(g_0))$

 $l_{\epsilon}(\delta) = \delta - \epsilon - \operatorname{rank}(h_0)B$ and $e_J(l_{\epsilon}(\delta), E) = e_J(\delta, E) + \epsilon(\operatorname{rank}(h_0)),$

 $l^{\epsilon}(\check{\delta}) = \check{\delta} + \tau^{-1}\epsilon + \operatorname{rank}(h_{-1})B \quad and \quad \hom_{J}(\tau^{-1}E, l^{\epsilon}(\check{\delta})) = \hom_{J}(\tau^{-1}E, \check{\delta}) + \tau^{-1}\epsilon(\operatorname{rank}(h_{-1})),$ $l^{\epsilon}(\check{\delta}) = \check{\delta} + \tau^{-1}\check{\epsilon} - \operatorname{rank}(g_{-1})B \quad and \quad \check{e}_{J}(\tau^{-1}E, l^{\epsilon}(\check{\delta})) = \check{e}_{J}(\tau^{-1}E, \check{\delta}) - \tau^{-1}\check{\epsilon}(\operatorname{rank}(g_{-1})).$

Proof. We will only prove the statement for $l_{\epsilon}(\delta)$. As ϵ is rigid, by Lemma 3.8 E is essentially the only general representation in PC(ϵ). So we may assume that both (M, E) and $(E, \tau L)$ are general pairs. In particular, the rank of each h_i is the general rank from $\tau^i(\epsilon)$ to $\tau^{i+1}(l_{\epsilon}(\delta))$ (see Remark 5.17).

The equalities of the left column follow directly from the definition and the exactness of the sequence. The equalities of the right column are all easy exercises of homological algebras. We prove the first one as an illustration. We apply $\text{Hom}_J(-, E)$ to the exact sequence $0 \rightarrow \text{im}(h_{-1}) \rightarrow L \rightarrow \text{im}(f_0) \rightarrow 0$, and get

$$0 \to \operatorname{Hom}_J(\operatorname{im}(f_0), E) \to \operatorname{Hom}_J(L, E) \to \operatorname{Hom}_J(\operatorname{im}(h_{-1}), E) = 0.$$

Hom_J(im(h_{-1}), E) vanishes because Hom_J($\tau^{-1}E, E$) = $\check{E}_J(E, E) = 0$. Thus Hom_J(im(f_0), E) \cong Hom_J(L, E). Then apply Hom_J(-, E) to the exact sequence $0 \to im(f_0) \to M \to im(g_0) \to 0$. We have

$$0 \to \operatorname{Hom}_J(\operatorname{im}(g_0), E) \to \operatorname{Hom}_J(M, E) \to \operatorname{Hom}_J(\operatorname{im}(f_0), E) \to \dot{\mathrm{E}}_J(\operatorname{im}(g_0), E) = 0.$$

 $\check{\mathbf{E}}_J(\mathrm{im}(g_0), E)$ vanishes because $\check{\mathbf{E}}_J(E, E) = 0$. So $\mathrm{hom}_J(l_{\epsilon}(\delta), E) = \mathrm{hom}_J(\delta, E) - \check{\epsilon}(\mathrm{rank}(g_0))$.

A direct application of Algorithm 5.20 gives the following corollary.

Corollary 5.28. Let **u** be an extended sequence of mutations such that $\mu_{\mathbf{u}}(\epsilon) = -e_v$. The operators l_{ϵ} and r_{ϵ} on the δ -vectors of (Q, S) are given by

$$r_{\epsilon}(\delta) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\delta) - e_{v});$$

$$l_{\epsilon}(\delta) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\delta) + e_{v}).$$

Let **u** be an extended sequence of mutations such that $\mu_{\mathbf{u}}(\check{\epsilon}) = e_v$. The operators l^{ϵ} and r^{ϵ} on the $\check{\delta}$ -vectors of (Q, S) are given by

$$r^{\epsilon}(\check{\delta}) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\check{\delta}) + e_{v});$$
$$l^{\epsilon}(\check{\delta}) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\check{\delta}) - e_{v}).$$

Even in some trivial cases, these formulas produce amusing equalities.

Proposition 5.29. We have the following equalities.

(1) $\tau(\delta + e_v + \operatorname{rank}(P_v, \tau\delta)B) = \tau\delta - e_v;$ (2) $\tau^{-1}(\delta - e_v + \operatorname{rank}(\delta, I_v)B) = \tau^{-1}\delta + e_v.$ (3) $\tau(\check{\delta} - e_v - \operatorname{rank}(P_v, \check{\delta})B) = \tau\check{\delta} + e_v;$ (4) $\tau^{-1}(\check{\delta} + e_v - \operatorname{rank}(\tau^{-1}\check{\delta}, I_v)B) = \tau^{-1}\check{\delta} - e_v.$

Proof. Consider the situation when the extended mutation sequence is just a single $\mu_{+} = \tau$, and $\epsilon = e_v$. By Corollary 5.28 we have $r_{\epsilon}(\delta) = \tau^{-1}(\tau \delta - e_v)$. Comparing with the original definition of r_{ϵ} , we obtain the first equality. Working with $\mu_{-} = \tau^{-1}$ and $\check{\epsilon} = e_v$ for l_{ϵ} , we obtain the second equality. Working with $\mu_{+} = \tau$ and $\epsilon = -e_v$ for r^{ϵ} , we obtain the third equality. Working with $\mu_{-} = \tau^{-1}$ and $\check{\epsilon} = -e_v$ for r^{ϵ} , we obtain the third equality. Working with $\mu_{-} = \tau^{-1}$ and $\check{\epsilon} = -e_v$ for l^{ϵ} , we obtain the third equality. Working with $\mu_{-} = \tau^{-1}$ and $\check{\epsilon} = -e_v$ for l^{ϵ} , we obtain the last equality.

Proposition 5.30. Suppose that we are in the situation of Theorem 5.16. The composition $l_{\epsilon}r_{\epsilon}$ is the identity if and only if $\operatorname{rank}(R, E) = \operatorname{rank}(r_{\epsilon}(\delta), \epsilon)$; and $r_{\epsilon}l_{\epsilon}$ is the identity if and only if $\operatorname{rank}(E, \tau L) = \operatorname{rank}(\epsilon, \tau(l_{\epsilon}(\delta)))$. In particular, when ϵ is rigid, both are identities.

24

Proof. Using the frozen-vertex trick (before Algorithm 5.20), we may assume the *B*-matrix of the quiver has full rank. By Theorem 5.16 we have that $r_{\epsilon}(\delta) = \delta + \check{\epsilon} - \operatorname{rank}(R, E)B$. Then

$$l_{\epsilon}(r_{\epsilon}(\delta)) = r_{\epsilon}(\delta) - \check{\epsilon} + \operatorname{rank}(r_{\epsilon}(\delta), \epsilon)B = \delta + (\operatorname{rank}(r_{\epsilon}(\delta), \epsilon) - \operatorname{rank}(R, E))B.$$

Hence, $l_{\epsilon}r_{\epsilon}$ is the identity if and only if rank $(R, E) = \operatorname{rank}(r_{\epsilon}(\delta), \epsilon)$. The other equality is proved similarly.

As we have seen in Example 5.19, rank(R, E) may be different from rank $(r_{\epsilon}(\delta), \epsilon)$.

5.4. Remarks on Dual Operators. In view of Corollary 5.28 one may wonder the analogous definition of operators on $\check{\delta}$ -vectors such that

$$\check{r}_{\epsilon}(\check{\delta}) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\check{\delta}) - e_{v}); \tag{5.8}$$

$$\hat{l}_{\epsilon}(\hat{\delta}) = \mu_{\mathbf{u}}^{-1}(\mu_{\mathbf{u}}(\hat{\delta}) + e_v).$$
(5.9)

where **u** be a sequence of mutations such that $\mu_{\mathbf{u}}(\check{\epsilon}) = -e_v$.

Definition 5.31. For any δ -vector ϵ , we define the two operators \check{r}_{ϵ} and \check{l}_{ϵ} on the set of δ -vector as follows.

$$\check{r}_{\epsilon}(\check{\delta}) = \check{\delta} + \check{\epsilon} - \operatorname{rank}(\tau^{-1}\check{\delta},\check{\epsilon})B;$$
$$\check{l}_{\epsilon}(\check{\delta}) = \check{\delta} - \epsilon - \operatorname{rank}(\epsilon,\check{\delta})B,$$

and

$$\check{r}^{\epsilon}(\delta) = \delta + \epsilon + \operatorname{rank}(\delta, \tau\check{\epsilon})B;$$
$$\check{l}^{\epsilon}(\delta) = \delta - \check{\epsilon} + \operatorname{rank}(\tau\epsilon, \tau\check{\delta})B.$$

If we compare this definition with Definition 5.4, we immediately get the following lemma.

Lemma 5.32. We have the following equalities

$$\begin{split} \tilde{l}_{-\check{\epsilon}}(\check{\delta}) &= r^{\epsilon}(\check{\delta}) \quad and \quad \check{r}_{-\check{\epsilon}}(\check{\delta}) = l^{\epsilon}(\check{\delta});\\ \tilde{l}^{-\check{\epsilon}}(\delta) &= r_{\epsilon}(\delta) \quad and \quad \check{r}^{-\check{\epsilon}}(\delta) = l_{\epsilon}(\delta). \end{split}$$

where $-\check{\epsilon}$ is viewed as a δ -vector (note that $\ker(\check{\epsilon}) = \tau \operatorname{coker}(-\check{\epsilon})$).

From this lemma, one can easily see that the equalities (5.8) and (5.9) do hold for \check{r}_{ϵ} and \check{l}_{ϵ} in Definition 5.31. We leave it to readers to formulate the analogue of Corollary 5.28 for \check{r}_{ϵ} and \check{l}_{ϵ} .

We also notice that there are more equivalent statements in Proposition 5.6 if we allow \check{r}^{ϵ} and \check{l}^{ϵ} join the game. For example, if $\delta = \delta_1 \oplus \delta_2$, then $\tau \delta = \tau \delta_1 \oplus \tau \delta_2$, so $\check{r}^{\delta} = \check{r}^{\delta_1} \check{r}^{\delta_2}$ by Lemma 5.32.

Corollary 5.33. If $\delta = \delta_1 \oplus \delta_2$, then $r_{\epsilon}(\delta) = r_{r_{\epsilon}(\delta_1)}(\delta_2) = r_{r_{\epsilon}(\delta_2)}(\delta_1)$.

Proof. We have that $r_{\epsilon}(\delta) = \check{r}^{\delta}(\epsilon) = \check{r}^{\delta_1}\check{r}^{\delta_2}(\epsilon) = \check{r}^{\delta_1}(r_{\epsilon}(\delta_2)) = r_{r_{\epsilon}(\delta_2)}(\delta_1)$. The other equality follows from the symmetry.

6. MUTATION OF GENERAL RANKS

Theorem 6.1. Let $\gamma_r = \operatorname{rank}(\epsilon, \tau \delta)$ and $\gamma_l = \operatorname{rank}(\delta, \epsilon)$. We denote $\gamma'_r = \operatorname{rank}(\epsilon', \tau \delta')$ and $\gamma'_l = \operatorname{rank}(\delta', \epsilon')$, where $\delta' = \mu_u(\delta)$ and $\epsilon' = \mu_u(\epsilon)$.

(1) If r_{ϵ} commutes with μ_u , then

$$\gamma_r'(v) = \begin{cases} \gamma_r(v) & \text{for all } v \neq u \\ \gamma_r[b_u]_+ - \gamma_r(u) + [\delta(u)]_+ + [\epsilon(u)]_+ - [r_\epsilon(\delta)(u)]_+ & \text{for all } v = u \end{cases}$$

(2) If l_{ϵ} commutes with μ_u , then

$$\gamma_l'(v) = \begin{cases} \gamma_l(v) & \text{for all } v \neq u, \\ \gamma_l[b_u]_+ - \gamma_l(u) + [\delta(u)]_+ + [-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+ & \text{for all } v = u. \end{cases}$$

Proof. We will only prove the statement for γ_l . The other one can be proved in a similar fashion. If B is not of full rank, then we apply the frozen vertex trick (before Algorithm 5.20). To ease our notation, we will denote $\tilde{B}, \tilde{\delta}$ and $\tilde{\epsilon}$ still by B, δ and $\tilde{\epsilon}$. We know that γ'_l is the unique vector satisfying

$$l_{\epsilon}(\delta)' = l_{\epsilon'}(\delta') = \delta' - \check{\epsilon}' + \gamma'_l B'.$$
(6.1)

Recall the mutation rule for δ -vectors (3.1). For the coordinate u, we have that

$$l_{\epsilon}(\delta)'(u) = -(\delta(u) - \check{\epsilon}(u) + \gamma_l b_u),$$

while

$$\delta'(u) - \check{\epsilon}'(u) + \gamma'_l b'_u = -\delta(u) + \check{\epsilon}(u) + \sum_{v \neq u} \gamma_l(v)(-b_{v,u}) = -\delta(u) + \check{\epsilon}(u) - \gamma_l b_u = l_{\epsilon}(\delta)'(u).$$

For a coordinate $v \neq u$, we have that

$$l_{\epsilon}(\delta)'(v) = l_{\epsilon}(\delta)(v) + [b_{v,u}]_{+}l_{\epsilon}(\delta)(u) - b_{v,u}[l_{\epsilon}(\delta)(u)]_{+},$$

while

$$\begin{split} (\delta' - \check{\epsilon}' + \gamma'_{l}B')(v) \\ = (\delta(v) + [b_{v,u}]_{+}\delta(u) - b_{v,u}[\delta(u)]_{+}) - (\check{\epsilon}(v) + [b_{v,u}]_{+}\check{\epsilon}(u) + b_{v,u}[-\check{\epsilon}(u)]_{+}) + (\gamma'_{l}(u)b'_{u,v} + \sum_{w \neq u} \gamma_{l}(w)b'_{w,v}) \\ = l_{\epsilon}(\delta)(v) + ([b_{v,u}]_{+}\delta(u) - b_{v,u}[\delta(u)]_{+}) - ([b_{v,u}]_{+}\check{\epsilon}(u) + b_{v,u}[-\check{\epsilon}(u)]_{+}) \\ + ((-\gamma_{l}(u) - \gamma'_{l}(u))b_{u,v} + \sum_{w \neq u} \gamma_{l}(w)(b'_{w,v} - b_{w,v})) \\ = l_{\epsilon}(\delta)(v) + [b_{v,u}]_{+}(l_{\epsilon}(\delta)(u) - \gamma_{l}b_{u}) + (-b_{v,u}[\delta(u)]_{+}) - (b_{v,u}[-\check{\epsilon}(u)]_{+}) \\ + ((-\gamma_{l}[b_{u}]_{+} - [\delta(u)]_{+}) - [-\check{\epsilon}(u)]_{+} + -[l_{\epsilon}(\delta)(u)]_{+})b_{u,v} + \sum_{w \neq u} \gamma_{l}(w)(b'_{w,v} - b_{w,v})) \\ = l_{\epsilon}(\delta)(v) + [b_{v,u}]_{+}l_{\epsilon}(\delta)(u) - b_{v,u}[l_{\epsilon}(\delta)(u)]_{+} \\ - \gamma_{l}b_{u}[b_{v,u}]_{+} - \gamma_{l}[b_{u}]_{+}b_{u,v} + \sum_{w \neq u} \gamma_{l}(w)([b_{w,u}]_{+}[b_{u,v}]_{+} - [-b_{w,u}]_{+}[-b_{u,v}]_{+})) \\ = l_{\epsilon}(\delta)'(v) - \gamma_{l}(b_{u}[b_{v,u}]_{+} + [b_{u}]_{+}b_{u,v}) + \gamma_{l}([b_{u}]_{+}[b_{u,v}]_{+} - [-b_{u}]_{+}[-b_{u,v}]_{+}) \\ = l_{\epsilon}(\delta)'(v). \end{split}$$

We thus get the desired equality (6.1).

Remark 6.2. Similarly one can show that

$$\gamma'_r(u) = \gamma_r[-b_u]_+ - \gamma_r(u) + [-\delta(u)]_+ + [-\epsilon(u)]_+ - [-r_\epsilon(\delta)(u)]_+,$$

and

$$\gamma_l'(u) = \gamma_l[-b_u]_+ - \gamma_l(u) + [-\delta(u)]_+ + [\check{\epsilon}(u)]_+ - [-l_{\epsilon}(\delta)(u)]_+.$$

So one can rewrite the formula as

$$\gamma_r'(u) = \begin{cases} \gamma_r[b_u]_+ - \gamma_r(u) + [\delta(u)]_+ + [\epsilon(u)]_+ & \text{if } r_\epsilon(\delta)(u) < 0; \\ \gamma_r[-b_u]_+ - \gamma_r(u) + [-\delta(u)]_+ + [-\epsilon(u)]_+ & \text{if } r_\epsilon(\delta)(u) \ge 0. \end{cases}$$

and

$$\gamma_{l}'(u) = \begin{cases} \gamma_{l}[b_{u}]_{+} - \gamma_{l}(u) + [\delta(u)]_{+} + [-\check{\epsilon}(u)]_{+} & \text{if } l_{\epsilon}(\delta(u)) < 0; \\ \gamma_{l}[-b_{u}]_{+} - \gamma_{l}(u) + [-\delta(u)]_{+} + [\check{\epsilon}(u)]_{+} & \text{if } l_{\epsilon}(\delta)(u) \ge 0, \end{cases}$$

Proposition 6.3. Suppose that l_{ϵ} and r_{ϵ} commutes the mutation μ_u . Let $\gamma_r = \operatorname{rank}(\epsilon, \tau \delta)$ and $\gamma_l = \operatorname{rank}(\delta, \epsilon)$. Then the following quantities are mutation-invariant:

$$h_l(\delta, \epsilon) := \check{\epsilon}(\gamma_l) - \hom(\delta, \epsilon) + \hom(l_{\epsilon}(\delta), \epsilon);$$

$$\check{h}_l(\delta, \epsilon) := \delta(\gamma_l) - \hom(\delta, \epsilon) + \hom(\delta, \check{l}^{\delta}(\epsilon)),$$

and

$$e_r(\delta, \epsilon) := \epsilon(\gamma_r) - e(\delta, \epsilon) + e(r_\epsilon(\delta), \epsilon);$$

$$\check{e}_r(\delta, \epsilon) := -\delta(\gamma_r) - e(\delta, \epsilon) + e(\delta, \check{r}^{\delta}(\epsilon)).$$

Proof. We will only prove the mutation-invariance of $\check{h}_l(\delta, \epsilon)$. The rest can be proved in a similar fashion.

$$\begin{split} \delta'(\gamma') - \delta(\gamma) \\ = \delta'(u)\gamma'(u) - \delta(u)\gamma(u) + \sum_{v \neq u} (\delta'(v)\gamma(v) - \delta(v)\gamma(v)) \\ = \delta'(u)(\gamma[b_u]_+ - \gamma(u) + [\delta(u)]_+ + [-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+) - \delta(u)\gamma(u) + \sum_{v \neq u} \gamma(v)(\delta'(v) - \delta(v)) \quad \text{(Theorem 6.1)} \\ = -\delta(u)(\gamma[b_u]_+ + [\delta(u)]_+ + [-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+) + \sum_{v \neq u} \gamma(v)([b_{v,u}]_+\delta(u) - b_{v,u}[\delta(u)]_+) \quad \text{(by (3.1))} \\ = -\delta(u)([\delta(u)]_+ + [-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+) + \sum_{v \neq u} \gamma(v)(-b_{v,u}[\delta(u)]_+) \\ = - [\delta(u)]_+ (\delta(u) + \gamma b_u) - \delta(u)([-\check{\epsilon}(u)]_+ - [l_{\epsilon}(\delta)(u)]_+) \\ = - [\delta(u)]_+ l_{\epsilon}(\delta)(u) - [\delta(u)]_+ \check{\epsilon}(u) - \delta(u)[-\check{\epsilon}(u)]_+ + \delta(u)[l_{\epsilon}(\delta)(u)]_+ \\ = [\delta(u)]_+ [-l_{\epsilon}(\delta)(u)]_+ - [-\delta(u)]_+ [l_{\epsilon}(\delta)(u)]_+ - [\delta(u)]_+ [\check{\epsilon}(u)]_+ + [-\delta(u)]_+ [-\check{\epsilon}(u)]_+ \\ \text{Comparing this with Lemma 2.6, we find that} \\ \delta'(\gamma') - \delta(\gamma) = - (\hom(\delta', \epsilon') - \hom(\delta, \epsilon)) + (\hom(\delta', \tau l_{\epsilon}(\delta)') - \hom(\delta, \tau l_{\epsilon}(\delta)) \\ = - (\hom(\delta', \epsilon') - \hom(\delta, \epsilon)) + (\hom(\delta', \check{\ell}^{\delta}(\epsilon')) - \hom(\delta, \check{\ell}^{\delta}(\epsilon)) \quad \text{(by (5.3) and Lemma 5.32).} \end{split}$$

27

Remark 6.4. If δ is extended-reachable, then the above invariants always vanish as seen in Corollary 5.28. If δ is not rigid, they may not be zero (but conjecturally always nonnegative).

7. Appendix: Conjecture 5.22 implies the Duality Pairing and the Saturation

Recall from [9] that the tropical *F*-polynomial f_M of a representation M is the function $(\mathbb{Z}^{Q_0})^* \to \mathbb{Z}_{\geq 0}$ defined by

$$\delta \mapsto \max_{L \hookrightarrow M} \delta(\underline{\dim}L);$$

The dual tropical F-polynomial \check{f}_M of a representation M is the function $(\mathbb{Z}^{Q_0})^* \to \mathbb{Z}_{\geq 0}$ defined by

$$\delta \mapsto \max_{M \to N} \delta(\underline{\dim}N),$$

where δ is viewed as an element in $(\mathbb{Z}^{Q_0})^*$ via the usual dot product. Clearly f_M and \check{f}_M are related by $f_M(\delta) - \check{f}_M(-\delta) = \delta(\underline{\dim}M)$. If M is general of weight δ , then we will write f_δ for f_M . Similarly we can define \check{f}_δ and $f_{\check{\delta}}$.

In [9, Section 5] we showed that the Fock-Concharnov's duality pairing conjecture for generic bases of skew-symmetric cluster algebras can be deduced from the following conjecture: for any pair $(\delta, \check{\epsilon})$ we have that

$$f_{\check{\epsilon}}(\delta) = \check{f}_{\delta}(\check{\epsilon})$$

In fact, they are equivalent if the *B*-matrix of *Q* has full rank. We refer readers to [11, Section 12] and [9, Section 5] for the details of the duality pairing conjecture. As mentioned in [9], a more optimistic conjecture is that $f_{\check{\epsilon}}(\delta) = \check{f}_{\delta}(\check{\epsilon}) = \hom(\delta,\check{\epsilon})$. The following theorem generalizes [9, Theorem 2.22].

Theorem 7.1. Assume either of the following two situations:

- (1) there is a sequence of mutations $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\check{\epsilon}))$ is hom-vanishing;
- (2) there are two sequences of mutations $\mu_{\mathbf{u}}$ and $\mu_{\check{\mathbf{u}}}$ such that $(\mu_{\mathbf{u}}(\delta), \mu_{\mathbf{u}}(\check{\epsilon}))$ is e-vanishing and $(\mu_{\check{\mathbf{u}}}(\delta), \mu_{\check{\mathbf{u}}}(\check{\epsilon}))$ is \check{e} -vanishing.

Then
$$f_{\check{\epsilon}}(\delta) = \hom(\delta, \check{\epsilon}) = f_{\delta}(\check{\epsilon}).$$

Proof. Let $\delta' = \mu_{\mathbf{u}}(\delta)$ and $\check{\epsilon}' = \mu_{\mathbf{u}}(\check{\epsilon})$. If $\hom(\delta', \check{\epsilon}') = 0$, then $f_{\check{\epsilon}'}(\delta') \ge \hom(\delta', \check{\epsilon}')$. But $f_{\check{\epsilon}'}(\delta') \le \hom(\delta', \check{\epsilon}')$ by [9, Lemma 2.5]. Hence $f_{\check{\epsilon}'}(\delta') = \hom(\delta', \check{\epsilon}')$. Then by [9, Lemma 2.21] we get $f_{\check{\epsilon}}(\delta) = \hom(\delta, \check{\epsilon})$. Similarly we can show that $\check{f}_{\delta}(\check{\epsilon}) = \hom(\delta, \check{\epsilon})$.

If $e(\delta', \check{\epsilon}') = 0$, then a similar argument shows that $\check{f}_{\check{\epsilon}}(-\delta) = e(\delta, \check{\epsilon})$, which is equivalent to $f_{\check{\epsilon}}(\delta) = \hom(\delta, \check{\epsilon})$. Similarly, from $\check{e}(\delta', \check{\epsilon}') = 0$ we can conclude that $\check{e}(\delta, \check{\epsilon}) = f_{\delta}(-\check{\epsilon})$, which is equivalent to $\hom(\delta, \check{\epsilon}) = \check{f}_{\delta}(\check{\epsilon})$.

Note that if $f_{\check{\epsilon}}(\delta) = \hom(\delta,\check{\epsilon})$, then by [9, Theorem 2.6] $f_{\check{\epsilon}}(m\delta) = \hom(m\delta,\check{\epsilon})$ for any $m \in \mathbb{N}$; similarly if $\check{f}_{\delta}(\check{\epsilon}) = \hom(\delta,\check{\epsilon})$, then $\check{f}_{\delta}(m\check{\epsilon}) = \hom(\delta,m\check{\epsilon})$ for any $m \in \mathbb{N}$. Also note that $f_{\check{\epsilon}}(m\delta) = mf_{\check{\epsilon}}(\delta)$. So the conclusion of Theorem 7.1 implies that

$$\hom(m\delta,\check{\epsilon}) = \hom(\delta,m\check{\epsilon}) = m\hom(\delta,\check{\epsilon}) \quad \text{for any } m \in \mathbb{N}.$$
(7.1)

Conversely, recall from [9, Theorem 2.22] that we always have that

$$f_{\check{\epsilon}}(m\delta) = \hom(m\delta,\check{\epsilon}) \text{ and } f_{\delta}(n\check{\epsilon}) = \hom(\delta,n\check{\epsilon}) \text{ for some } m,n\in\mathbb{N}.$$

So if (7.1) holds for the pair $(\delta, \check{\epsilon})$, then we have that $f_{\check{\epsilon}}(\delta) = \hom(\delta, \check{\epsilon})$ and $\hom(\delta, \check{\epsilon}) = \dot{f}_{\delta}(\check{\epsilon})$.

Definition 7.2. We say that a pair $(\delta, \check{\epsilon})$ has the hom-*fluent property* if (7.1) holds. We say that a pair $(\delta, \check{\epsilon})$ has the *saturation property* if hom $(\delta, \check{\epsilon}) = 0$ whenever hom $(m\delta, n\check{\epsilon}) = 0$ for some $m, n \in \mathbb{N}$. We say a QP is hom-*fluent* (resp. saturated) if the hom-fluent (resp. saturation) property holds for any pair $(\delta, \check{\epsilon})$.

The saturation property generalizes the ordinary saturation property for acyclic quivers [4]. The saturation property seems weaker than the hom-fluent property. But the proof of [9, Theorem 2.22] shows that they are in fact equivalent. We conclude that

Proposition 7.3. The following are equivalent:

- (1) $f_{\check{\epsilon}}(\delta) = \hom(\delta, \check{\epsilon}) = f_{\delta}(\check{\epsilon}).$
- (2) The hom-fluent property holds for $(\delta, \check{\epsilon})$.
- (3) The saturation property holds for $(\delta, \check{\epsilon})$.

Conjecture 7.4 (Saturation for nondegenerate QPs). Every nondegenerate Jacobi-finite QP is saturated.

It is not hard to give a counterexample for degenerate QPs.

Corollary 7.5. Conjecture 5.22 implies both the saturation conjecture and the duality pairing conjecture for nondegenerate QPs.

Proof. By the above discussion, it suffices to show that $f_{\epsilon}(\delta) = \hom(\delta, \epsilon) = f_{\delta}(\epsilon)$ for any pair (δ, ϵ) . For any pair (δ, ϵ) , we also consider the pair $(\tau^{-1}\epsilon, \delta)$. By Conjecture 5.22, there is a sequence of mutation $\mu_{\mathbf{u}}$ such that $(\mu_{\mathbf{u}}(\tau^{-1}\epsilon), \mu_{\mathbf{u}}(\delta)) = (\tau^{-1}\mu_{\mathbf{u}}(\epsilon), \mu_{\mathbf{u}}(\delta))$ is either hom-vanishing or e-vanishing, which is equivalent to say $(\mu_{\mathbf{u}}(\epsilon), \mu_{\mathbf{u}}(\delta))$ is either ϵ -vanishing or hom-vanishing by Lemma 2.4. If it is hom-vanishing, then we are done by Theorem 7.1. Otherwise, there it is ϵ -vanishing and if working with the original pair (δ, ϵ) there is another sequence of mutation $\mu'_{\mathbf{u}}$ such that $(\mu'_{\mathbf{u}}(\epsilon), \mu'_{\mathbf{u}}(\delta))$ is e-vanishing, and we are done as well by Theorem 7.1.

References

- C. Amiot, Cluster categories for algebras of global dimension 2 and quivers with potential, Ann. Inst. Fourier (Grenoble) 59 (2009), no. 6, 2525–2590.
- A. B. Buan, R. Marsh, M. Reineke, I. Reiten, G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. 204 (2006), no. 2, 572–618.
- 3. W. Crawley-Boevey, On homomorphisms from a fixed representation to a general representation of a quiver, Trans. Amer. Math. Soc. 348 (1996), no. 5, 1909–1919.
- H. Derksen, J. Weyman, Semi-invariants of quivers and saturation for Littlewood-Richardson coefficients, J. Amer. Math. Soc. 13 (2000), no. 3, 467–479.
- H. Derksen, J. Weyman, A. Zelevinsky, Quivers with potentials and their representations I, Selecta Math. (N.S.) 14 (2008), no. 1, 59–119.
- H. Derksen, J. Weyman, A. Zelevinsky, Quivers with potentials and their representations II, J. Amer. Math. Soc. 23 (2010), no. 3, 749–790.
- 7. H. Derksen, J. Fei, General presentations of algebras, Adv. Math. 278 (2015), 210-237.
- I. Dolgachev, Lectures on invariant theory, London Mathematical Society Lecture Note Series, 296. Cambridge University Press, Cambridge, 2003. xvi+220 pp.
- 9. J. Fei, Tropical F-polynomials and general presentations, to appear J. London Math. Soc..
- 10. J. Fei, Crystal structures of upper cluster algebras, to appear.
- V. Fock, A. Goncharov, Moduli spaces of local systems and higher Teichmüller theory, Publ. Math. Inst. Hautes Études Sci., 103 (2006) 1–211.
- 12. S. Fomin, A. Zelevinsky, Cluster algebras. IV. Coefficients, Compos. Math. 143 (2007), no. 1, 112–164.
- C. Geiss, B. Leclerc, J. Schröer, Generic bases for cluster algebras and the Chamber Ansatz, J. Amer. Math. Soc. 25 (2012), 21–76.

- 14. V. Ginzburg, Calabi-Yau algebras, arXiv:math/0612139v3.
- K. Igusa, K. Orr, G. Todorov, J. Weyman, *Cluster Complexes via Semi-Invariants*, Compositio Math. 145 (2009), 1001–1034.
- B. Keller, I. Reiten, Cluster-tilted algebras are Gorenstein and stably Calabi-Yau, Adv. Math. 211 (2007), no. 1, 123–151.
- 17. B. Keller, D. Yang, Derived equivalences from mutations of quivers with potential, Adv. Math. 226 (2011), no. 3, 2118–2168.
- 18. A. Neeman, Some new axioms for triangulated categories, J. Algebra 139 (1991), 221–255.
- Y. Palu, Cluster characters for 2-Calabi-Yau triangulated categories, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 6, 2221–2248.
- P. Plamondon, Cluster algebras via cluster categories with infinite-dimensional morphism spaces, Compos. Math. 147 (2011), no. 6, 1921–1954.
- P. Plamondon, Generic bases for cluster algebras from the cluster category, Int. Math. Res. Notices (2013) 2013 (10): 2368–2420.
- 22. M. Rosenlicht, Some basic theorems on algebraic groups, Amer. J. Math. 78 (1956), 401-443.
- 23. A. Schofield, General representations of quivers, Proc. London Math. Soc. (3) 65 (1992), no. 1, 46-64.