

Mean reflected McKean-Vlasov stochastic differential equation

Liu Xiangdong*, Hong Shaopeng

^a*Department of Statistic and Data Science, School of Economics, Jinan University, Guangzhou, 510632, Guangdong, PR China*

Abstract

In this paper, we investigate a class of mean reflected McKean-Vlasov stochastic differential equation, which extends the equation proposed by [1] by allowing the solution's distribution to not only constrain its behavior, but also affect the diffusion and drift coefficients. We establish the existence and uniqueness results of this class, investigate the propagation of chaos, and examine the stability properties with respect to the initial condition, coefficients, and driving process. Moreover, we provide a rigorous proof of a Fredlin-Wentzell type large deviations principle using the weak convergence method. We also demonstrate the existence of an invariant measure and employ the coupling by change of measure method to prove log-Harnack inequality and shift Harnack inequality. Our study sheds light on the properties and behaviors of these reflected McKean-Vlasov stochastic differential equation and contributes to the ongoing research in this field.

Keywords: McKean-Vlasov stochastic differential equation, mean reflected, particle system, stability analysis, large derivation principle, invariant measure, Harnack inequality
2020 MSC: 60H10

1. Introduction

In this paper, we consider the following *mean reflected McKean-Vlasov stochastic differential equation* (MR-MVSDE for short).

$$\begin{cases} X_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dB_s + K_t, & t \geq 0 \\ \mathbb{E}[h(X_t)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s)] dK_s = 0, & t \geq 0. \end{cases} \quad (1.1)$$

Where \mathcal{L}_{X_s} is the law of X_s , $(B_t)_{t \geq 0}$ is a standard Brownian motion defined on some complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We will always assume that h is nondecreasing and that the law of ξ is such that $\mathbb{E}[h(\xi)] \geq 0$. For the precise conditions on b, σ, h , we refer the reader to Section 2. The solution to (1.1) is the couple of continuous processes (X, K) , K being needed to ensure that the constraint is satisfied, in a minimal way according to the last condition namely the Skorokhod condition.

*Corresponding author

Email addresses: tliuxd@jnu.edu.cn (Liu Xiangdong*), hsp1999@stu2021.jnu.edu.cn (Hong Shaopeng)

The McKean-Vlasov stochastic differential equation (McKean-Vlasov SDE for short), also referred to as the mean-field stochastic differential equation or distribution-dependent stochastic differential equation, was first introduced in the literature by [2] and [3]. In recent years, it has garnered significant attention from scholars. The McKean-Vlasov SDE is used to describe the limiting behavior of mean-field interactive particle systems as n goes to infinity. We refer to [4] for a detailed exposition and references. Currently, the McKean-Vlasov SDE plays a crucial role in various fields such as mean field games, financial derivatives pricing, measure value processes, see e.g. [5, 6, 7, 8] and references therein.

The reflected SDE was introduced by [9] and [10]. Since then, it has been widely researched, see e.g. [11, 12, 13, 14]. This type of equation restricts the path of the solution to a given domain. The reflected McKean-Vlasov SDE, first proposed by [15], is well-posed, and its solution satisfies the propagation of chaos. Recently, [16] studied the McKean-Vlasov SDE with self-stabilizing coefficients on unbounded convex regions, replacing the classical Wasserstein Lipschitz condition. Their work proved the existence and uniqueness of the solution, demonstrated chaos propagation, and examined the Freidlin-Wentzell type Large Deviations Principle, as well as an Eyring-Kramer's law for the exit-time from subdomains that lie within the interior of the reflecting domain. In a similar vein, [17] investigated the McKean-Vlasov SDE with oblique reflection on non-smooth time-dependent domains. Their research proved the existence and uniqueness of the solution, demonstrated chaos propagation, and examined the Freidlin-Wentzell type Large Deviations Principle.

In their work, [18] introduced the concept of mean reflective backward stochastic differential equation (MR-BSDE), which places restrictions not on the path of the solution, but on its distribution. They proved that, under certain conditions, MR-BSDE has a unique solution. Further research on MR-BSDE can be found in [19, 20, 21], and their respective references. [1] studied the mean reflected stochastic differential equation (MR-SDE for short), established the well-posedness of the solution, and analyzed the propagation of chaos. They also proposed a numerical simulation method based on particle approximation techniques. [22] extended this work to Lévy driven MR-SDE. [23] examined the MR-SDE with two reflective walls and solved the Skorokhod problem with the mean minimum condition. They also studied the stability of the solution. In a related study, [24] investigated SDE with normal reflection in conditional law and BSDE with normal constraint in law. Using the penalty method, they established the well-posedness of these two methods and proved the forward chaos propagation and backward propagation of chaos [25]. Additionally, they studied a related obstacle problem on the Wasserstein space. Overall, these studies contribute to a better understanding of MR-BSDE and MR-SDE and their applications in various fields.

The large deviation principle (LDP) is primarily concerned with asymptotic estimates of the probability of tail events A , especially whether $\mathbb{P}(X_\varepsilon \in A)$ converges to zero at an exponential rate as $\varepsilon \rightarrow 0$. In this paper, we focus on the Freidlin-Wentzell type of LDP. The study of large deviations for stochastic differential equations originates from [26], who first studied the large deviation principle for diffusion processes. Recently, the weak convergence method [27] based on the variational representation of Brownian motion [28] has become an important and effective tool for proving LDP. Using this method, many LDP for stochastic differential equations and stochastic partial differential equations have been established, see e.g. [29, 30, 31, 32, 33, 34]. For more on the weak convergence method and LDP, see

monograph [35]. [36] proposed sufficient conditions for verifying LDP that are easier to verify and used these conditions to prove LDP for quasilinear stochastic partial differential equations with reflection. Currently, there are relatively few studies on McKean-Vlasov SDE. Using the classical exponential equivalence method, [37] and [38] proved LDP for McKean-Vlasov SDE. For reflected McKean-Vlasov SDE, [16] used the exponential equivalence method to prove the large deviation principle. Notably, [39] proposed a new sufficient condition to verify the criteria of the weak convergence method, and show correct controlled equation for the McKean-Vlasov SDE with jumps, which was used to prove the large deviation principle and moderate deviation principle for the McKean-Vlasov SDE with jumps. [40] using weak convergence method shows the LDP of MR-SDE with jump.

The invariant probability measure is an important tool for studying the long-term behavior of SDE. The principle of invariant probability measure for classical SDE or distribution independent SDE has been extensively studied, as seen in [41, 42, 43, 44, 45]. However, due to the fact that the McKean-Vlasov SDE is not a Markov process, the classical proof method cannot be used. [46] first established the \mathcal{W}_2 -exponential convergence of the solution of the McKean-Vlasov SDE, and subsequently proved the uniqueness and existence of the invariant probability measure. [47] studied the exponential ergodicity of McKean-Vlasov SDE with additive pure jump noise in the relative entropy and (weighted) Wasserstein distances. [48] studied the exponential ergodicity of singular McKean-Vlasov SDE with or without reflection in the relative entropy and (weighted) Wasserstein distances. [49] studied the Donsker-Varadhan large deviation of the invariant measure for path-dependent McKean-Vlasov SPDE.

The Harnack inequality is an important inequality for studying the properties of solutions. [50] first proposed a dimension-free Harnack inequality. If we denote $\{P_t\}_{t \geq 0}$ is the transition semigroup of a Markov process, the dimension-free Harnack inequality states that

$$(P_t f)^\alpha(x) \leq P_t f^\alpha(x+y) \exp\{C_\alpha(y,t)\}, x, y \in \mathbb{R}^d,$$

where $\alpha > 1$, $C_\alpha : \mathbb{R}^d \times (0, \infty) \rightarrow \mathbb{R}_+$ satisfies $C_\alpha(0,t) = 0$, and $x, y \in \mathbb{R}^d$. [51] proved the dimension-free Harnack inequality for a large class of stochastic differential equations with multiplicative noise by constructing a coupling with unbounded time-dependent drift. This coupling method, also known as the variable measure coupling method, has been used to prove the Harnack inequality for many SDE and SPDE, see e.g., [52, 53, 54, 55, 56] and monograph [57]. For the McKean-Vlasov SDE, due to technical difficulties, we can only prove the Harnack inequality when the noise coefficient is independent of the distribution. [46] used the variable measure coupling method to prove the Harnack inequality for the McKean-Vlasov SDE when the noise coefficient is independent of the distribution. [58] studied the Harnack inequality when the drift contains a locally standard integrable term and a Lipschitz continuous term in the spatial variable and is Lipschitz continuous in the distribution variable with respect to a weighted variation distance. [59] studied the Harnack inequality for McKean-Vlasov SDE, where the drift satisfies a standard integrability condition in time-space and may be discontinuous in the distance induced by any Dini function. [60] developed a novel coupling method to effectively prove the shift Harnack inequality. Utilizing this new coupling method, we prove the shift Harnack inequality for the McKean-Vlasov SDE when the noise coefficient is independent of the spatial variable. [61] proved the shift Harnack inequality for

the McKean-Vlasov SPDE in a separable Hilbert space with a Dini continuous drift. [62] used Zvonkin's transform to prove the Harnack inequality and shift Harnack inequality for the McKean-Vlasov SDE when the coefficients are Dini continuous in the spatial variable.

Our contributions.

To the best of our knowledge, there are few results on the MR-MVSDE. The aim of this paper is to investigate this type of MR-MVSDE. In Section 3, we first establish the existence and uniqueness of a strong solution (X, K) for (1.1), followed by a study of its properties, including: (i) the relationship between (1.1) and a PDE with Neumann boundary conditions in the Wasserstein space, i.e., the Feynman-Kac formula (Proposition 3.3); (ii) L^p estimates for the solution (Proposition 3.4); (iii) the absolute continuity of the Stieljes measure dK with respect to the Lebesgue measure, under certain regularity assumptions on h , and an explicit expression for the density (Proposition 3.6). In Section 4, we study propagation of chaos (Theorem 4.1), i.e., we prove the convergence of an interacting particle system (see (4.2)) to (1.1).

In Section 5, we investigate the impact of small perturbations on the system. Specifically, we study the effects of perturbations in the initial condition (Theorem 5.1), the coefficients (Theorem 5.2), and the driving process (Theorem 5.3). We then use Proposition 5.7 and Proposition 5.9 to prove a large deviation principle for (1.1) (Theorem 5.6). In Section 6, we focus on the long-time behavior of solutions and the Harnack inequality. We prove the existence and uniqueness of an invariant measure for (1.1) (Theorem 6.2). Using coupling method, we prove log-Harnack inequality (Theorem 6.4) and shift Harnack inequality (Theorem 6.5) for (1.1).

The paper is organised in the following way: Section 2 introduces the Wasserstein metric and presents some Assumptions and preliminary results. Section 3 shows that the MR-MVSDE admits a unique strong solution (X, K) . Then we study the density of K . Section 4 shows that system (1.1) can be seen as the limit of an interacting particles system with oblique reflection of mean field type. Section 5 studies the effect of small noise on the system. Section 6 proves the existence of invariant probability measure and the Harnack inequality.

Throughout the paper, we adopt the following convention: $\mathcal{B}_b^+(\mathbb{R})$ refers to the space of nonnegative and bounded measurable functions on \mathbb{R} . The symbol C denotes various positive constants whose specific values may vary in different contexts.

2. Preliminary

In this section, we primarily focus on the Wasserstein space and some preliminary results.

2.1. The Wasserstein space

Let $\mathcal{P}(\mathbb{R})$ be the space of probability measures on \mathbb{R} and for any $p > 1$ denote by $\mathcal{P}_p(\mathbb{R})$ the subspace of $\mathcal{P}(\mathbb{R})$ of probability measures with finite moment of order p . Now, we can defined p -Wasserstein distance For $\mu, \nu \in \mathcal{P}_p(\mathbb{R})$, define the p -Wasserstein distance $\mathcal{W}_p(\mu, \nu)$ by

$$\mathcal{W}_p(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \left[\int_{\mathbb{R} \times \mathbb{R}} |x - y|^p d\pi(x, y) \right]^{1/p}$$

where $\Pi(\mu, \nu)$ denotes the set of probability measures on $\mathbb{R} \times \mathbb{R}$ whose first and second marginals are, respectively, μ and ν . An equivalent and more probabilistic definition is

$$\mathcal{W}_p(\mu, \nu) = \left(\inf_{X \sim \mu, Y \sim \nu} \mathbb{E}[|X - Y|^p] \right)^{\frac{1}{p}} \leq \mathbb{E}[|X - Y|^p]^{\frac{1}{p}}$$

In following result, we mainly using 1-Wasserstein distance and 2-Wasserstein distance. More detail about Wasserstein metric can see [63] and references therein.

Then, we introduce the L-derivative (Lions' derivative). A function $f : \mathcal{P}_2(\mathbb{R}) \mapsto \mathbb{R}$ is differential at $\mu \in \mathcal{P}_2(\mathbb{R})$, if for lift map $\tilde{f}(\gamma) := f(\mathcal{L}_\gamma)$, $\gamma \in L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R})$, there exists some $\zeta \in L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R})$ with $\mathcal{L}_\zeta = \mu$ such that \tilde{f} is Fréchet differentiable at ζ , that is, there exists a linear continuous mapping $D_\mu \tilde{f}(\zeta) : L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}) \mapsto \mathbb{R}$ such that for any $\eta \in L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R})$

$$\tilde{f}(\zeta + \eta) - \tilde{f}(\zeta) = D_\mu \tilde{f}(\zeta)(\eta) + o(|\eta|_{L^2}), \quad |\eta|_{L^2} \rightarrow 0$$

In order to obtain the Feynman-Kac formula for (1.1), it is necessary to make certain regularity assumptions on function f and apply a special version of Itô's lemma, as demonstrated in the work of [24].

Defination 2.1. We say f is partially C^2 if the map $y \rightarrow D_\mu f(\mu)(y)$ is continuously differentiable with a derivative $(\mu, y) \rightarrow \partial_y D_\mu f(\mu)(y)$ jointly continuous on $\mathcal{P}_2(\mathbb{R}) \times \mathbb{R}$

Lemma 2.2 (Itô's lemma). Consider

$$dX_t = b_t dt + \sigma_t dB_t + dK_t.$$

where K_t is a deterministic, continuous and nondecreasing process with $K_0 = 0$, (b_t) and (σ_t) are progressively measurable with values in \mathbb{R} and satisfy

$$\mathbb{E} \left[\int_0^T (|b_s|^2 + |\sigma_s|^4) ds \right] < +\infty$$

If f is partially C^2 , we have

$$\begin{aligned} f(\mathcal{L}_{X_t}) &= f(\mathcal{L}_{X_t}) + \int_0^t \mathbb{E} [D_\mu f(\mathcal{L}_{X_t})(X_s) b_s] ds + \int_0^t \mathbb{E} [D_\mu f(\mathcal{L}_{X_t})(X_s)] dK_s \\ &\quad + \frac{1}{2} \int_0^t \mathbb{E} [(\sigma_s^2 \partial_y D_\mu f(\mathcal{L}_{X_t})(X_s))] ds, \end{aligned}$$

2.2. Assumptions and preliminary results

We make following assumptions about the coefficients:

Assumption. (A.1)

- (i) Lipschitz assumption: there exists a constant $K > 0$ such that for all $x, y \in \mathbb{R}$, $\mu, \nu \in \mathcal{P}_2(\mathbb{R})$ we have

$$|b(x, \mu) - b(y, \nu)| + |\sigma(x, \mu) - \sigma(y, \nu)| \leq K(|x - y| + \mathcal{W}_2(\mu, \nu))$$

- (ii) The random variable ξ is square integrable, i.e. $\mathbb{E}[\xi^2] < \infty$.

Remark 2.1. The Assumption (A1) implies that for any $t \in [0, T]$, the coefficients b and σ satisfy

$$|b(x, \mu)| + |\sigma(x, \mu)| \leq C(1 + K)$$

Assumption. (A.2)

- (i) The function $h : \mathbb{R} \rightarrow \mathbb{R}$ is increasing and bi-Lipschitz. There exist $0 < m \leq M$ such that

$$\forall x \in \mathbb{R}, \forall y \in \mathbb{R}, \quad m|x - y| \leq |h(x) - h(y)| \leq M|x - y|$$

- (ii) The initial condition ξ satisfies $\mathbb{E}[h(\xi)] \geq 0$

Assumption. (A.3) There exists $p > 4$ such that $\xi \in L^p$, i.e. $\mathbb{E}[|\xi|^p] < \infty$.

Assumption. (A.4) The function h is twice continuously differentiable with bounded derivatives.

Assumption. (A.5) For some constants $C_2 > C_1 \geq 0$,

$$2(x - y)(b(x, \mu) - b(y, \nu)) + |\sigma(x, \mu) - \sigma(y, \nu)| \leq C_1 \mathcal{W}_2(\mu, \nu)^2 - C_2|x - y|^2$$

We now introduce function H , \bar{G}_0 and G and some properties from [1].

$$H : \mathbb{R} \times \mathcal{P}_1(\mathbb{R}) \ni (x, \nu) \mapsto \int h(x + z)v(dz) \quad (2.1)$$

Let \bar{G}_0 be the inverse function in space of H evaluated at 0 ,

$$\bar{G}_0 : \mathcal{P}_1(\mathbb{R}) \ni \nu \mapsto \inf\{x \in \mathbb{R} : H(x, \nu) \geq 0\} \quad (2.2)$$

and let G_0 denote the positive part of \bar{G}_0 ,

$$G_0 : \mathcal{P}_1(\mathbb{R}) \ni \nu \mapsto \inf\{x \geq 0 : H(x, \nu) \geq 0\} \quad (2.3)$$

Lemma 2.3. *Under Assumption (A.2) we have:*

- (i) *For all ν in $\mathcal{P}_1(\mathbb{R})$, the mapping $H(\cdot, \nu) : \mathbb{R} \ni x \mapsto H(x, \nu)$ is a bi-Lipschitz function, namely:*

$$\forall x, y \in \mathbb{R}, \quad m|x - y| \leq |H(x, \nu) - H(y, \nu)| \leq M|x - y|.$$

- (ii) *For all x in \mathbb{R} , the mapping $H(x, \cdot) : \mathcal{P}_1(\mathbb{R}) \ni \nu \mapsto H(x, \nu)$ satisfies the following Lipschitz estimate:*

$$\forall \nu, \nu' \in \mathcal{P}_1(\mathbb{R}), \quad |H(x, \nu) - H(x, \nu')| \leq \left| \int h(x + \cdot)(d\nu - d\nu') \right|$$

Lemma 2.4. *Under Assumption (A.2), the mapping $G_0 : \mathcal{P}_1(\mathbb{R}) \ni \nu \mapsto G_0(\nu)$ is Lipschitz continuous in the following sense:*

$$|G_0(\nu) - G_0(\nu')| \leq \frac{1}{m} \left| \int h(\bar{G}_0(\nu) + \cdot)(d\nu - d\nu') \right|$$

where we recall that $\bar{G}_0(\nu)$ is the inverse of $H(\cdot, \nu)$ at point 0 . In particular,

$$|G_0(\nu) - G_0(\nu')| \leq \frac{M}{m} \mathcal{W}_1(\nu, \nu').$$

3. Existence, uniqueness and relative properties of the solution

3.1. Existence and Uniqueness of (1.1)

We now introduce the definition of the solution to equation.

Definition 3.1. *A couple of processes (X, K) is said to be a flat deterministic solution to (1.1) if (X, K) satisfy (1.1) with K being a non-decreasing continuous deterministic function with $K_0 = 0$.*

Given this definition, we have the following result.

Theorem 3.2. *Under Assumption (A.1) and (A.2), the MR-MVSDE (1.1) has a unique deterministic flat solution (X, K) . Moreover,*

$$\forall t \geq 0, \quad K_t = \sup_{s \leq t} \inf \{x \geq 0 : \mathbb{E}[h(x + U_s)] \geq 0\} = \sup_{s \leq t} G_0(\mathcal{L}_{U_s}), \quad (3.1)$$

where $(U_t)_{0 \leq t \leq T}$ is the process defined by :

$$U_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dB_s. \quad (3.2)$$

With these notations, denoting by \mathcal{L}_{U_s} the family of marginal laws of $(U_t)_{0 \leq t \leq T}$ we have

$$K_t = \sup_{s \leq t} G_0(\mathcal{L}_{U_s}).$$

PROOF. The proof for the case of MR-SDE is given in [1]. We present here the proof of the MR-MVSDE.

Let us consider the set $\mathcal{C}^2 = \{X \text{ is } \mathcal{F} \text{ - adapted continuous process, } \mathbb{E}(\sup_{0 \leq t \leq T} |X_t|^2) < \infty\}$. And let $\tilde{X} \in \mathcal{C}^2$ be a given process. We set

$$\tilde{U}_t = \xi + \int_0^t b(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s}) ds + \int_0^t \sigma(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s}) dB_s, \quad \text{Law}(\tilde{U}_s) =: \mathcal{L}_{\tilde{U}_s} \quad (3.3)$$

and define the function K by setting

$$K_t = \sup_{s \leq t} \inf \left\{ x \geq 0 : \mathbb{E} \left[h \left(x + \tilde{U}_s \right) \right] \geq 0 \right\} = G_0(\tilde{\mu}_s).$$

The function K being given, let us define the process X by the formula

$$X_t = \xi + \int_0^t b(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s}) ds + \int_0^t \sigma(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s}) dB_s + K_t. \quad (3.4)$$

Let us check (X, K) is the solution of (1.1) with U_t replaced by \tilde{U}_t . First, based on the definition of K , we have $\mathbb{E}[h(X_t)] \geq 0$, $K_t = G_0(\mathcal{L}_{\tilde{U}_t})$, $dK_t - a.e.$ and $G_0(\mathcal{L}_{\tilde{U}_t}) > 0$, $dK_t - a.e.$ Then we obtain

$$\begin{aligned} \int_0^t \mathbb{E}[h(X_s)] dK_s &= \int_0^t \mathbb{E} \left[h \left(\tilde{U}_s + K_s \right) \right] dK_s \\ &= \int_0^t \mathbb{E} \left[h \left(\tilde{U}_s + G_0(\mathcal{L}_{\tilde{U}_s}) \right) \right] dK_s \\ &= \int_0^t \mathbb{E} \left[h \left(\tilde{U}_s + G_0(\mathcal{L}_{\tilde{U}_s}) \right) \right] \mathbb{1}_{\{G_0(\mathcal{L}_{\tilde{U}_s}) > 0\}} dK_s. \end{aligned}$$

Moreover, since h is continuous, we have $\mathbb{E} \left[h \left(\tilde{U}_s + G_0 \left(\mathcal{L}_{\tilde{U}_s} \right) \right) \right] = 0$ as soon as $G_0 \left(\mathcal{L}_{\tilde{U}_s} \right) > 0$, so that

$$\int_0^t \mathbb{E} [h(X_s)] dK_s = 0$$

Second, consider the map Φ which associates to \tilde{X} to the solution X of (1.1). We will show Φ is a contraction. Let $\tilde{X}, \tilde{X}' \in \mathcal{C}^2$ and define \tilde{U}, K and \tilde{U}', K' as above, using the some Brownian motion. We have from Cauchy-Schwarz and Burkholder-Davis-Gundy (B-D-G) inequality

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t| \right] \\ & \leq 3\mathbb{E} \left[\sup_{t \leq T} \left\{ \left| \int_0^t \left(b \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - b \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right) ds \right|^2 + \left| \int_0^t \left(\sigma \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - \sigma \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right) dB_s \right|^2 \right. \right. \\ & \quad \left. \left. + |K_t - K'_t|^2 \right\} \right] \\ & \leq 3\mathbb{E} \left[\sup_{t \leq T} t \int_0^t \left(b \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - b \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right) ds \right]^2 + 3\mathbb{E} \left[\sup_{t \leq T} \left| \int_0^t \left(\sigma \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - \sigma \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right) dB_s \right|^2 \right] \\ & + 3 \sup_{t \leq T} |K_t - K'_t|^2 \\ & := 3(I_1 + I_2 + I_3) \end{aligned} \tag{3.5}$$

From the Lipschitz assumption, we have

$$\begin{aligned} I_1 & \leq T\mathbb{E} \left[\int_0^T \left| b \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - b \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right|^2 ds \right] \\ & \leq CT\mathbb{E} \left[\int_0^T \left| \tilde{X}_s - \tilde{X}'_s \right|^2 + \mathcal{W}_2 \left(\mathcal{L}_{\tilde{X}_s}, \mathcal{L}_{\tilde{X}'_s} \right)^2 ds \right] \\ & \leq CT \left\{ T\mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t| \right] + \mathbb{E} \left[\int_0^T \mathcal{W}_2 \left(\mathcal{L}_{\tilde{X}_s}, \mathcal{L}_{\tilde{X}'_s} \right)^2 ds \right] \right\} \\ & \leq 2CT^2\mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t| \right] \end{aligned} \tag{3.6}$$

$$\begin{aligned} I_2 & \leq C\mathbb{E} \left[\int_0^T \left| \sigma \left(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s} \right) - \sigma \left(\tilde{X}'_s, \mathcal{L}_{\tilde{X}'_s} \right) \right|^2 ds \right] \\ & \leq CT\mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t| \right] + C\mathbb{E} \left[\int_0^T \mathcal{W}_2 \left(\mathcal{L}_{\tilde{X}_s}, \mathcal{L}_{\tilde{X}'_s} \right)^2 ds \right] \\ & \leq 2CT\mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t| \right] \end{aligned} \tag{3.7}$$

$$I_3 \leq \frac{M}{m} \mathbb{E} \left[\sup_{t \leq T} |\tilde{U}_t - \tilde{U}'_t|^2 \right] \leq C(T^2C_1 + TC_2) \mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t|^2 \right]. \tag{3.8}$$

Combining (3.5)-(3.8), we have

$$\mathbb{E} \left[\sup_{t \leq T} |X_t - X'_t|^2 \right] \leq C(1+T)T \mathbb{E} \left[\sup_{t \leq T} |\tilde{X}_t - \tilde{X}'_t|^2 \right].$$

Hence, there exists a positive \tilde{T} , depending on b, σ and h only, such that for all $T < \tilde{T}$, the map Φ is a contraction. We first deduce the existence and uniqueness of the solution on $[0, \tilde{T}]$ and then on \mathbb{R}^+ by iterating the construction.

Remark 3.1. From this construction, we deduce that for all positive r :

$$K_{t+r} - K_t = \sup_{s \in [0, r]} \inf \left\{ x \geq 0 : \mathbb{E} \left[h \left(x + X_t + \int_t^{t+s} b(X_u, \mathcal{L}_{X_u}) du + \int_t^{t+s} \sigma(X_u, \mathcal{L}_{X_u}) dB_u \right) \right] \geq 0 \right\}$$

Then, we connect our MR-MVSDE with a PDE on the Wasserstein space.

Let $\mathcal{O} := \{\mu \in \mathcal{P}_2(\mathbb{R}), \int h(x)\mu(dx) > 0\}$. Given a bounded, continuous map $G : \bar{\mathcal{O}} \rightarrow \mathbb{R}$ we consider the map $u : [0, T] \times \bar{\mathcal{O}} \rightarrow \mathbb{R}$ defined by

$$u(t, \mu) = \mathbb{E}[G(\mathcal{L}_{X_t})] \quad (3.9)$$

Proposition 3.3 (Feynman-Kac formula). *Assume that u is a classical solution to Neumann problem on the Wasserstein space (3.10). Then, u is given by (3.9).*

$$\begin{cases} (i) & (\partial_t + \mathcal{T})u(t, \mu) = 0 & \text{in } (0, T) \times \mathcal{O}, \\ (ii) & \int_{\mathbb{R}} D_\mu u(t, \mu)(y)\mu(dy) = 0, & \text{in } (0, T) \times \partial\mathcal{O}, \\ (iii) & u(T, \mu) = G(\mu), & \text{in } \mathcal{O}, \end{cases} \quad (3.10)$$

where the operator \mathcal{T} is given, for any smooth function $\phi : [0, T] \times \mathcal{P}_2(\mathcal{R}) \rightarrow \mathbb{R}$, by

$$\mathcal{T}\phi(t, \mu) := \int_{\mathbb{R}} b(z, \mu) D_\mu \phi(t, \mu)(z) d\mu(z) + \frac{1}{2} \int_{\mathbb{R}} \sigma(z, \mu)^2 \partial_z D_\mu \phi(t, \mu)(z) d\mu(z) \quad (3.11)$$

PROOF. Let us set $\mu_s := \mathcal{L}_{X_s^{t_0, x_0}}$. Then, by Lemma 2.2:

$$u(T, \mu_T) = u(t_0, \mu_{t_0}) + \int_{t_0}^T (\partial_t + \mathcal{T})u(s, \mu_s) + \int_{t_0}^T \int_{\mathbb{R}} D_\mu u(s, \mu_s)(y)\mu_s(dy) dK_s$$

where, by (3.10)-(ii),

$$\int_{t_0}^T \int_{\mathbb{R}} D_\mu u(s, \mu_s)(y)\mu_s(dy) dK_s = \int_{t_0}^T \mathbb{1}_{\{\mu_s \in \partial\mathcal{O}\}} \left(\int_{\mathbb{R}} D_\mu u(s, \mu_s)(y) \right) dK_s = 0$$

From the equation satisfied by u , we obtain

$$u(t_0, \mu_0) = u(t_0, \mu_{t_0}) = \mathbb{E}[u(T, \mu_T)] = \mathbb{E}[G(\mathcal{L}_{X_T})]$$

Proposition 3.4. *Suppose that Assumption (A.1) and (A.2) hold.*

(i) For all p , there exists a positive C_p , depending on T, b, σ and h such that

$$\mathbb{E} \left[\sup_{t \leq T} |X_t|^p \right] \leq C_p (1 + \mathbb{E} [|\xi|^p])$$

(ii) There exists a positive constant C , depending on T, b, σ and h such that

$$\forall 0 \leq s \leq t \leq T, \quad |K_t - K_s| \leq C |t - s|^{\frac{1}{2}}$$

(iii) For all $p \geq 1$ such that $\xi \in L^p$ we have

$$\forall 0 \leq s \leq t \leq T, \quad \mathbb{E} [|X_t - X_s|^p] \leq C |t - s|^{\frac{p}{2}}$$

PROOF. (i) We have

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq T} |X_t|^p \right] &\leq 4^{p-1} \left\{ \mathbb{E} [|\xi|^p] + \mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |b(X_s, \mathcal{L}_{X_s})| ds \right)^p \right] \right. \\ &\quad \left. + \mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |\sigma(X_s, \mathcal{L}_{X_s})| dB_s \right)^p \right] + K_T^p \right\} \end{aligned} \quad (3.12)$$

We first deal with $K_T^p = \sup_{t \leq T} G_0(\mathcal{L}_{U_t})$. Apply Lipschitz property of G_0 , Lemma 2.4 and the definition of the Wasserstein metric \mathcal{W}_1 , we have

$$\forall t \geq 0, \quad |G_0(\mathcal{L}_{U_t})| \leq \frac{M}{m} \mathbb{E} [|U_t - U_0|],$$

since $G_0(\mathcal{L}_{U_0}) = 0$ as $\mathbb{E} [h(X_0)] \geq 0$, and where U is defined by (3.2). Therefore

$$\begin{aligned} |K_T|^p &= \left| \sup_{t \leq T} G_0(\mathcal{L}_{U_t}) \right|^p \leq 2^{p-1} \left\{ \left(\frac{M}{m} \right)^p \mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |b(X_s, \mathcal{L}_{X_s})| ds \right)^p \right] \right. \\ &\quad \left. + \mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |\sigma(X_s, \mathcal{L}_{X_s})| dB_s \right)^p \right] \right\} \end{aligned} \quad (3.13)$$

Plugging (3.13) into (3.12), we deduce that there exists a constant $C_p \geq 0$ such that

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq T} |X_t|^p \right] &\leq C_p \mathbb{E} \left[|\xi|^p + \sup_{t \leq T} \left(\int_0^t |b(X_s, \mathcal{L}_{X_s})| ds \right)^p + \sup_{t \leq T} \left| \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dB_s \right|^p \right] \\ &\leq C_p \left\{ \mathbb{E} [|\xi|^p] + T^{p-1} \mathbb{E} \left[\int_0^T (1 + |X_s|)^p ds \right] + C \mathbb{E} \left[\int_0^T (1 + |X_s|)^2 ds \right]^{\frac{p}{2}} \right\} \\ &\leq C_p (1 + \mathbb{E} [|\xi|^p]) + C \int_0^T \mathbb{E} \left[\sup_{t \leq r} |X_t|^p \right] dr \end{aligned} \quad (3.14)$$

Apply Gronwall's lemma, we deduce for all $p \geq 2$, there exists a constant $C > 0$ such that

$$\mathbb{E} \left[\sup_{t \leq T} |X_t|^p \right] \leq C (1 + \mathbb{E} [|\xi|^p]).$$

(ii) we observe that $\nu \mapsto G_0(\nu)$ is Lipschitz continuous, we have

$$K_t - K_s = \sup_{r \leq t} G_0(\mathcal{L}_{U_r}) - \sup_{r \leq s} G_0(\mathcal{L}_{U_r}) \leq \frac{M}{m} \mathcal{W}_1(\mathcal{L}_{U_r}, \mathcal{L}_{U_s}) \leq \frac{M}{m} \mathbb{E}[|U_r - U_s|],$$

so the result follows from standard computations.

(iii) we use the decomposition

$$X_t - X_s = U_t - U_s + K_t - K_s$$

and the result of (ii), we complete this proof.

3.2. Density of K_t

Let \mathcal{L}_μ be the partial operator of second order defined by

$$\mathcal{A}_\mu f(x) = b(x, \mu) \frac{\partial}{\partial x} f(x) + \frac{1}{2} \sigma(x, \mu)^2 \frac{\partial^2}{\partial x^2} f(x) \quad (3.15)$$

In order to prove Proposition 3.6, we need following lemma:

Lemma 3.5. *If Assumption (A.1), (A.2) and (A.4) hold*

(i) *If φ is a continuous function such that, for some $C \geq 0$ and $p \geq 1$,*

$$\forall x \in \mathbb{R}, \quad |\varphi(x)| \leq C(1 + |x|^p)$$

then the function $t \mapsto \mathbb{E}[\varphi(X_t)]$ is continuous.

(ii) *The functions $t \mapsto \mathbb{E}[h(X_t)]$ and $t \mapsto \mathbb{E}[\mathcal{A}_{\mathcal{L}_{X_t}} h(X_t)]$ are continuous.*

PROOF. (i) is the [22][Lemma 3].

Now, we prove (ii). Since Assumption (A.1), (A.2) and (A.4) hold, the mapping $x \mapsto \mathcal{A}_{\mathcal{L}_{X_t}} h(X_t)$ is continuous and for all $x \in \mathbb{R}$, there exists constants C_1 and C_2 such that

$$\begin{aligned} |b(x, \mu) h'(x)| &\leq C_1(1 + |x|). \\ |\sigma(x, \mu)^2 h'(x)| &\leq C_2(1 + |x|). \end{aligned}$$

Then, using (i), we deduce that $t \mapsto \mathbb{E} \mathcal{A}_{\mathcal{L}_{X_t}} h(X_t)$ is continuous.

Proposition 3.6. *Suppose Assumptions (A.1), (A.2) and (A.4) hold. Let (X, K) be the unique deterministic flat solution to (1.1). Then the process K is Lipschitz continuous and the Stielje measure dK has the following density w.r.t the Lebesgue measure:*

$$k : \mathbb{R}^+ \ni t \mapsto \frac{(\mathbb{E}[\mathcal{A}_{\mathcal{L}_{X_t}} h(X_t)])^-}{\mathbb{E}[h'(X_t)]} \mathbb{1}_{\mathbb{E}[h(X_t)]=0} \quad (3.16)$$

PROOF. Firstly, we show K is Lipschitz continuous. To do so, we prove that $s \mapsto \bar{G}_0(\mu_s)$ is Lipschitz continuous on $[0, T]$. If $s \leq t$, we have

$$\begin{aligned} |\bar{G}_0(\mathcal{L}_{U_s}) - \bar{G}_0(\mathcal{L}_{U_t})| &\leq \frac{1}{m} |H(\bar{G}_0(\mathcal{L}_{U_s}), \mathcal{L}_{U_t})| \\ &= \frac{1}{m} |\mathbb{E}[h(\bar{G}_0(\mathcal{L}_{U_s}) + U_t)]| \\ &= \frac{1}{m} \left| \mathbb{E} \left[h \left(\bar{G}_0(\mathcal{L}_{U_s}) + U_s + \int_s^t b(X_r, \mathcal{L}_{X_r}) dr + \int_s^t \sigma(X_r, \mathcal{L}_{X_r}) dB_r \right) \right] \right|, \end{aligned}$$

the first equality since $H(\bar{G}_0(\mu_t), \mu_t) = 0$.

Let $\bar{A}_{x,\mu}$ be the partial operator of second order defined by

$$\bar{A}_{y,\mu} f(x) := b(y, \mu) \frac{\partial}{\partial x} f(x) + \frac{1}{2} \sigma(y, \mu)^2 \frac{\partial^2}{\partial x^2} f(x)$$

Apply Itô's lemma, we have

$$\begin{aligned} \mathbb{E}[h(\bar{G}_0(\mathcal{L}_{U_s}) + U_t)] &= \mathbb{E}[h(\bar{G}_0(\mathcal{L}_{U_s}) + U_s)] + \int_s^t \mathbb{E}[\bar{A}_{X_r, \mathcal{L}_{X_r}} h(\bar{G}_0(\mathcal{L}_{U_s}) + U_r)] dr \\ &= H(\bar{G}_0(\mathcal{L}_{U_s}), \mathcal{L}_{U_s}) + \int_s^t \mathbb{E}[\bar{A}_{X_r, \mathcal{L}_{X_r}} h(\bar{G}_0(\mathcal{L}_{U_s}) + U_r)] dr \quad (3.17) \\ &= \int_s^t \mathbb{E}[\bar{A}_{X_r, \mathcal{L}_{X_r}} h(\bar{G}_0(\mathcal{L}_{U_s}) + U_r)] dr. \end{aligned}$$

We get the result follows from that h has bounded derivatives and $\sup_{t \leq T} |X_t| \in L^2$ for each $T > 0$. So from [1], we deduce that K is Lipschitz continuous and has a bounded density on $[0, T]$ for each $T > 0$.

Then we show the density of dK is (3.16). From Assumption (A.4) and apply Itô's lemma, for any $s \leq t$ we can get

$$\begin{aligned} h(X_t) - h(X_s) &= \int_s^t b(X_r, \mathcal{L}_{X_r}) h'(X_r) dr + \int_s^t \sigma(X_r, \mathcal{L}_{X_r}) h'(X_r) dB_r + \frac{1}{2} \int_s^t \sigma(X_r, \mathcal{L}_{X_r})^2 h'(X_r) dr \\ &\quad + \int_s^t h'(X_r) dK_r \\ &= \int_s^t \mathcal{A}_{\mathcal{L}_{X_r}} h(X_r) dr + \int_s^t h'(X_r) dK_r + \int_s^t \sigma(X_r, \mathcal{L}_{X_r}) dB_r \end{aligned} \quad (3.18)$$

Then, we obtain

$$\mathbb{E} \left[\int_s^t h'(X_r) dK_r \right] = \mathbb{E} h(X_t) - \mathbb{E} h(X_s) - \int_s^t \mathbb{E} \mathcal{A}_{\mathcal{L}_{X_r}} h(X_r) dr \quad (3.19)$$

Using (3.19), Lemma 3.5, and the proof of [1][Proposition 2.7], we can get the density of dK w.r.t the Lebesgue measure is

$$k_t = \frac{(\mathbb{E}[\mathcal{A}_{\mathcal{L}_{X_t}} h(X_t)])^-}{\mathbb{E}[h'(X_t)]} \mathbb{1}_{\mathbb{E}[h(X_t)]=0}$$

4. Propagation of Chaos

We can write the unique solution of the MR-MVSDE (1.1) as :

$$X_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dB_s + \sup_{s \leq t} G_0(\mathcal{L}_{U_s}) \quad (4.1)$$

where μ_t stands for the law of

$$U_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dB_s$$

We are here interested in the particle approximation of such a system. Our candidates are the particles

$$X_t^i = \hat{\xi}^i + \int_0^t b\left(X_s, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j}\right) ds + \int_0^t \sigma\left(X_s, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j}\right) dB_s^i + \sup_{s \leq t} G_0(\mathcal{L}_{U_s^N}) \quad (4.2)$$

where B^i are independent Brownian motions, $\hat{\xi}^i = \xi^i + G_0(\mathcal{L}_{U^{\xi, N}})$, $(\xi^i)_i$ are independent copies of ξ , $\mathcal{L}_{U_s^N}$ denotes the empirical distribution at time s of the particles

$$U_s^i = \hat{\xi}^i + \int_0^s b\left(X_r^i, \frac{1}{n} \sum_{j=1}^N \delta_{x_r^j}\right) dr + \int_0^s \sigma\left(X_r^i, \frac{1}{n} \sum_{j=1}^N \delta_{x_r^j}\right) dB_r^i, \quad 1 \leq i \leq N, \quad \mathcal{L}_{U_s^N} = \frac{1}{N} \sum_{i=1}^N \delta_{U_s^i}$$

Remark 4.1. Similar [1] and [22], the previous system of interacting particles can be seen as a multidimensional reflected SDE with oblique reflection. If h is concave, the set

$$\mathcal{S} := \{(x_1, \dots, x_n) \in \mathbb{R}^N : h(x_1) + \dots + h(x_n) \geq 0\}$$

is convex and the system

$$\begin{cases} X_t^i = \hat{\xi}^i + \int_0^t b\left(X_s, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j}\right) ds + \int_0^t \sigma\left(X_s, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j}\right) dB_s^i + K_t^N, & 1 \leq i \leq N \\ \frac{1}{N} \sum_{i=1}^N h(X_t^i) \geq 0, & \frac{1}{N} \sum_{i=1}^N \int_0^t h(X_t^i) dK_s^N = 0 \end{cases} \quad (4.3)$$

is the SDE with oblique reflection in convex \mathcal{S} .

Theorem 4.1 (Propagation of Chaos). *Let $T > 0$, and suppose that Assumptions (A.1) and (A.2) hold.*

- (i) *Under Assumption (A.3) hold, there exists a constant C depending on b, σ, M, m and T such that, for each $j \in \{1, \dots, N\}$,*

$$\mathbb{E} \left[\sup_{t \leq T} \left| X_t^j - \tilde{X}_t^j \right|^2 \right] \leq CN^{-\frac{1}{2}}$$

(ii) Under Assumption (A.4) hold, there exists a constant C depending on b, σ, M, m and T such that, for each $j \in \{1, \dots, N\}$,

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^j - \tilde{X}_t^j|^2 \right] \leq CN^{-1}$$

PROOF. Let $t > 0$, for $r \leq t$, we have

$$\begin{aligned} |X_r^j - \bar{X}_r^j| &\leq \left| \hat{\xi}^j - \xi^j \right| \\ &+ \int_0^r \left| b \left(X_s^j, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j} \right) - b(\bar{X}_s^j, \mathcal{L}_{X_s}) \right| ds \\ &+ \left| \int_0^r \left(\sigma \left(X_s^j, \frac{1}{n} \sum_{j=1}^N \delta_{x_s^j} \right) - \sigma(\bar{X}_s^j, \mathcal{L}_{X_s}) \right) dB_s^j \right| \\ &+ \left| \sup_{s \leq r} G_0(\mathcal{L}_{U_s^N}) - \sup_{s \leq r} G_0(\mathcal{L}_{U_s}) \right|. \end{aligned} \quad (4.4)$$

From the inequality

$$\begin{aligned} \left| \sup_{s \leq r} G_0(\mathcal{L}_{U_s^N}) - \sup_{s \leq r} G_0(\mathcal{L}_{U_s}) \right| &\leq \sup_{s \leq r} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{U_s})| \leq \sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{U_s})| \\ &\leq \sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{\bar{U}_s^N})| + \sup_{s \leq t} |G_0(\mathcal{L}_{\bar{U}_s^N}) - G_0(\mathcal{L}_{U_s})|, \end{aligned}$$

We have

$$\sup_{r \leq t} |X_r^j - \bar{X}_r^j| \leq I_1^j(r) + \sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{\bar{U}_s^N})| + \sup_{s \leq t} |G_0(\mathcal{L}_{\bar{U}_s^N}) - G_0(\mathcal{L}_{U_s})|, \quad (4.5)$$

where, I_1^j is defined by

$$I_1^j = \int_0^t \left| b \left(X_s^j, \frac{1}{n} \sum_{i=1}^N \delta_{X_s^i} \right) - b(\bar{X}_s^j, \mathcal{L}_{\bar{X}_s^j}) \right| ds + \left| \int_0^t \sigma \left(X_s^j, \frac{1}{n} \sum_{i=1}^N \delta_{X_s^i} \right) - \sigma(\bar{X}_s^j, \mathcal{L}_{\bar{X}_s^j}) \right| dB_s^j \quad (4.6)$$

We have

$$\begin{aligned}
\mathbb{E} [|I_1^j(t)|^2] &\leq C \mathbb{E} \left[t \int_0^t \left| b \left(X_s^j, \frac{1}{n} \sum_{i=1}^N \delta_{X_s^i} \right) - b \left(\bar{X}_s^j, \mathcal{L}_{\bar{X}_s^j} \right) \right|^2 ds \right] \\
&\quad + C \mathbb{E} \left[\int_0^t \left| \sigma \left(X_s^j, \frac{1}{n} \sum_{i=1}^N \delta_{X_s^i} \right) - \sigma \left(\bar{X}_s^j, \mathcal{L}_{\bar{X}_s^j} \right) \right|^2 ds \right] \\
&\leq Ct \int_0^t \mathbb{E} \left[|X_s^j - \bar{X}_s^j|^2 + \mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right] ds \\
&\quad + C \int_0^t \mathbb{E} \left[|X_s^j - \bar{X}_s^j|^2 + \mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right] ds \\
&= C(1+t) \int_0^t \mathbb{E} [|X_s^j - \bar{X}_s^j|^2] ds + C(1+t) \int_0^t \mathbb{E} \left[\mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right] ds
\end{aligned} \tag{4.7}$$

We have

$$\sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{\bar{U}_s^N})| \leq \frac{M}{m} \sup_{s \leq t} \frac{1}{N} \sum_{i=1}^N |U_s^i - \bar{U}_s^i| \leq \frac{M}{m} \frac{1}{N} \sum_{i=1}^N \sup_{s \leq t} |U_s^i - \bar{U}_s^i|$$

Moreover, taking into account that the variables are exchangeable, the Cauchy-Schwarz inequality implies

$$\mathbb{E} \left[\sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{\bar{U}_s^N})|^2 \right] \leq \frac{M^2}{m^2} \frac{1}{N} \sum_{i=1}^N \mathbb{E} \left[\sup_{s \leq t} |U_s^i - \bar{U}_s^i|^2 \right] = \frac{M^2}{m^2} \mathbb{E} \left[\sup_{s \leq t} |U_s^j - \bar{U}_s^j|^2 \right]$$

Since

$$U_s^j - \bar{U}_s^j = \int_0^s b \left(X_r^j, \frac{1}{N} \sum_{i=1}^N \delta_{X_r^i} \right) - b \left(\bar{X}_r^j, \mathcal{L}_{\bar{X}_r^j} \right) ds + \int_0^s \sigma \left(X_r^j, \frac{1}{N} \sum_{i=1}^N \delta_{X_r^i} \right) - \sigma \left(\bar{X}_r^j, \mathcal{L}_{\bar{X}_r^j} \right) dB_s^j
\end{aligned} \tag{4.8}$$

We have

$$\mathbb{E} \left[\sup_{s \leq t} |G_0(\mathcal{L}_{U_s^N}) - G_0(\mathcal{L}_{\bar{U}_s^N})| \right] \leq C \frac{M^2}{m^2} (1+t) \left\{ \int_0^t \mathbb{E} [|X_s^j - \bar{X}_s^j|^2] + \mathbb{E} \left[\mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right] ds \right\}$$

Next, dealing with the $\mathbb{E} \left[\mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right]$ terms. By the triangle inequality, we get

$$\mathbb{E} \left[\mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right) \right] \leq \mathbb{E} \left[\left(\frac{1}{N} \sum_{i=1}^N |X_s^i - \bar{X}_s^i|^2 \right)^{\frac{1}{2}} + \mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{\bar{X}_s^i}, \mathcal{L}_{\bar{X}_s^i} \right) \right] \tag{4.9}$$

Noting that the particles are exchangeable and from [5], we have

$$\mathbb{E} \left[\mathcal{W}_2 \left(\frac{1}{n} \sum_{i=1}^N \delta_{X_s^i}, \mathcal{L}_{\bar{X}_s^j} \right)^2 \right] \leq C \mathbb{E} [|X_s^j - \bar{X}_s^j|] + CN^{-\frac{1}{2}}$$

Consequently, we have

$$\mathbb{E} \left[\sup_{r \leq t} |X_r^j - \bar{X}_r^j|^2 \right] \leq 2A \int_0^t \mathbb{E} \left[\sup_{r \leq s} |X_s^j - \bar{X}_s^j|^2 \right] ds + 4 \mathbb{E} \left[\sup_{s \leq t} |G_0(\mathcal{L}_{\bar{U}_s^N}) - G_0(\mathcal{L}_{U_s})|^2 \right] + ATN^{-\frac{1}{2}}, \quad (4.10)$$

where $A = C(1+t)(1+M^2/m^2)$. Apply Gronwall's lemma, we deduce that

$$\mathbb{E} \left[\sup_{r \leq t} |X_r^j - \bar{X}_r^j|^2 \right] \leq 2Ce^A \left\{ \mathbb{E} \left[\sup_{s \leq t} |G_0(\mathcal{L}_{\bar{U}_s^N}) - G_0(\mathcal{L}_{U_s})|^2 \right] + ATN^{-\frac{1}{2}} \right\} \quad (4.11)$$

In view of Lemma 2.4, we have

$$\mathbb{E} \left[\sup_{s \leq t} |G_0(\mathcal{L}_{\bar{U}_s^N}) - G_0(\mathcal{L}_{U_s})|^2 \right] \leq \frac{1}{m^2} \mathbb{E} \left[\sup_{s \leq t} \left| \int h(\bar{G}_0(\mathcal{L}_{U_s}) + \cdot) (d\mathcal{L}_{\bar{U}_s^N} - d\mathcal{L}_{U_s}) \right|^2 \right]$$

which leads to

$$\mathbb{E} \left[\sup_{r \leq t} |X_r^j - \bar{X}_r^j|^2 \right] \leq Ce^A \left\{ \frac{1}{m^2} \mathbb{E} \left[\sup_{s \leq t} \left| \int h(\bar{G}_0(\mathcal{L}_{U_s}) + \cdot) (d\mathcal{L}_{\bar{U}_s^N} - d\mathcal{L}_{U_s}) \right|^2 \right] + ATN^{-\frac{1}{2}} \right\}$$

Proof of (i). As h is Lipschitz continuous, the rate of convergence is determined by the empirical measure convergence of independent and identically distributed diffusion processes. However, obtaining the standard convergence rate is not straightforward since we require uniform convergence in time. If we only suppose that Assumption (A.1) and (A.2) holds, we obtain that

$$\frac{1}{m^2} \mathbb{E} \left[\sup_{s \leq t} \left| \int h(\bar{G}_0(\mathcal{L}_{U_s}) + \cdot) (d\mathcal{L}_{\bar{U}_s^N} - d\mathcal{L}_{U_s}) \right|^2 \right] \leq \frac{M^2}{m^2} \mathbb{E} \left[\sup_{s \leq t} \mathcal{W}_1(\mathcal{L}_{\bar{U}_s^N}, \mathcal{L}_{U_s})^2 \right]$$

In view of the proof of [1][Theorem 3.3 Part (i)], we have

$$\mathbb{E} \left[\sup_{s \leq t} \mathcal{W}_1(\mathcal{L}_{\bar{U}_s^N}, \mathcal{L}_{U_s})^2 \right] \leq CN^{-\frac{1}{2}}$$

So, we have there exists a constant $C > 0$

$$\mathbb{E} \left[\sup_{r \leq t} |X_r^j - \bar{X}_r^j|^2 \right] \leq Ce^A \left(C \frac{M^2}{m^2} + AT \right) N^{-\frac{1}{2}} \leq CN^{-\frac{1}{2}} \quad (4.12)$$

Proof of (ii). We need to estimate

$$\frac{1}{m^2} \mathbb{E} \left[\sup_{s \leq t} \left| \int h(\bar{G}_0(\mathcal{L}_{U_s}) + \cdot) (d\mathcal{L}_{\bar{U}_s^N} - d\mathcal{L}_{U_s}) \right|^2 \right].$$

Let ψ is the Radon-Nikodym derivative of $\bar{G}_0(\mu_s)$, V^i is a semimartingale $s \mapsto \bar{G}_0(\mathcal{L}_{\bar{U}_s^i}) + \bar{U}_s^i$

$$\begin{aligned}
R_N(s) &:= \int h(\bar{G}_0(\mathcal{L}_{U_s}) + \cdot) (d\mathcal{L}_{\bar{U}_s^N} - d\mathcal{L}_{U_s}) \\
&= \frac{1}{N} \sum_{i=1}^N h(\bar{G}_0(\mathcal{L}_{U_s}) + \bar{U}_s^i) - \mathbb{E} [h(\bar{G}_0(\mathcal{L}_{U_s}) + U_s)] \\
&= \frac{1}{N} \sum_{i=1}^N \{h(\bar{G}_0(\mathcal{L}_{U_s}) + \bar{U}_s^i) - \mathbb{E} [h(\bar{G}_0(\mathcal{L}_{U_s}) + \bar{U}_s^i)]\} \\
&= \frac{1}{N} \sum_{i=1}^N \{h(V_s^i) - \mathbb{E} [h(V_s^i)]\}
\end{aligned}$$

Apply Itô's lemma, we have

$$h(V_s^i) = h(V_0^i) + \int_0^s h'(V_r^i) \psi_r + \bar{\mathcal{A}}_{\bar{X}_r, \mathcal{L}_{\bar{X}_r}} h(V_r^i) dr + \int_0^s h'(V_r^i) \sigma(\bar{X}_r^i, \mathcal{L}_{\bar{X}_r^i}) dB_r^i$$

so that

$$\begin{aligned}
\mathbb{E} [h(V_s^i)] &= \mathbb{E} [h(V_0^i)] + \int_0^s \mathbb{E} [h'(V_r^i) \psi_r + \bar{\mathcal{A}}_{\bar{X}_r, \mathcal{L}_{\bar{X}_r}} h(V_r^i)] dr \\
&= 0 + \int_0^s \mathbb{E} [h'(V_r^i) \psi_r + \bar{\mathcal{A}}_{\bar{X}_r, \mathcal{L}_{\bar{X}_r}} h(V_r^i)] dr.
\end{aligned}$$

Then, in view of the proof of [1][Theorem 3.3 Part (ii)] we have there exists a constant $C > 0$ such that

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^j - \tilde{X}_t^j|^2 \right] \leq CN^{-1}.$$

5. Asymptotic analysis

5.1. Stability analysis

In this subsection, We study stability properties of (1.1) with respect to initial condition, coefficients and driven process.

Firstly, we establish the stability of (1.1) with respect to small perturbation of the initial point ξ . For simplicity, We denote by $X_t^{\xi_n}$ as the unique solution of following equation:

$$\begin{cases} X_t^{\xi_n} = \xi_n + \int_0^t b(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) ds + \int_0^t \sigma(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) dB_s + K_t^{\xi_n}, & t \geq 0 \\ \mathbb{E} [h(X_t^{\xi_n})] \geq 0, & \int_0^t \mathbb{E} [h(X_s^{\xi_n})] dK_s^{\xi_n} = 0, & t \geq 0 \end{cases}.$$

Theorem 5.1 (Stability with respect to initial condition). *Suppose that Assumption (A.1) and (A.2) hold. The mapping*

$$\Xi : \mathbb{R} \rightarrow L^2(\Omega, \mathcal{C}([0, T], \mathbb{R}))$$

defined by $\Xi(\xi) \rightarrow X_t^\xi$ is continuous.

PROOF. Let (ξ_n) be a sequence in \mathbb{R} and $\lim_{n \rightarrow \infty} \xi_n = \xi$. We will show that

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \leq T} |X_t^{\xi_n} - X_t^\xi|^2 \right] = 0.$$

That means the mapping Ξ is continuous.

Apply the Lipschitz condition of b and σ , and Cauchy-Schwarz inequality, we have

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq T} |X_t^{\xi_n} - X_t^\xi|^2 \right] &\leq 4|\xi_n - \xi|^2 + 4\mathbb{E} \left[\sup_{t \leq T} \int_0^t |b(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) - b(X_s^\xi, \mathcal{L}_{X_s^\xi})| ds \right]^2 \\ &\quad + 4\mathbb{E} \left[\sup_{t \leq T} \int_0^t |\sigma(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) - \sigma(X_s^\xi, \mathcal{L}_{X_s^\xi})| dB_s \right]^2 + 4 \sup_{t \leq T} |K_t^{\xi_n} - K_t^\xi|^2 \\ &\leq 4|\xi_n - \xi|^2 + 4T\mathbb{E} \left[\int_0^t |b(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) - b(X_s^\xi, \mathcal{L}_{X_s^\xi})|^2 ds \right] \\ &\quad + 4C\mathbb{E} \left[\int_0^t |\sigma(X_s^{\xi_n}, \mathcal{L}_{X_s^{\xi_n}}) - \sigma(X_s^\xi, \mathcal{L}_{X_s^\xi})|^2 ds \right] + 4 \sup_{t \leq T} |K_t^{\xi_n} - K_t^\xi|^2 \\ &\leq 4|\xi_n - \xi| + 4TC\mathbb{E} \left[\int_0^t |X_s^{\xi_n} - X_s^\xi|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^{\xi_n}}, \mathcal{L}_{X_s^\xi})^2 ds \right] \\ &\quad + 4C\mathbb{E} \left[\int_0^t |X_s^{\xi_n} - X_s^\xi|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^{\xi_n}}, \mathcal{L}_{X_s^\xi})^2 ds \right] + 4 \sup_{t \leq T} |K_t^{\xi_n} - K_t^\xi|^2 \end{aligned} \tag{5.1}$$

$$\begin{aligned} \sup_{t \leq T} |K_t^{\xi_n} - K_t^\xi|^2 &\leq \frac{M^2}{m^2} \mathbb{E} \left[\sup_{r \in [0, t]} |U_r^{\xi_n} - U_r^\xi|^2 \right] \\ &\leq \frac{M^2}{m^2} \left\{ 4|\xi_n - \xi| + (4T + 1)C\mathbb{E} \left[\int_0^t |X_s^{\xi_n} - X_s^\xi|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^{\xi_n}}, \mathcal{L}_{X_s^\xi})^2 ds \right] \right\} \end{aligned} \tag{5.2}$$

Apply Gronwall's lemma, we have

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^{\xi_n} - X_t^\xi|^2 \right] \leq 4 \exp \left\{ C \left(1 + \frac{M^2}{m^2} \right) \right\} |\xi_n - \xi|^2 \tag{5.3}$$

Letting n goes to infinity, we complete this proof.

Secondly, we study the stability of (1.1) with respect to small perturbation of the coefficients b and σ . Let X_t^n is the unique solution of following equation:

$$\begin{cases} X_t^n = \xi + \int_0^t b^n(X_s^n, \mathcal{L}_{X_s^n}) ds + \int_0^t \sigma^n(X_s^n, \mathcal{L}_{X_s^n}) dB_s + K_t^n, & t \geq 0 \\ \mathbb{E}[h(X_t^n)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s^n)] dK_s^n = 0, & t \geq 0 \end{cases}.$$

Theorem 5.2 (Stability with respect to coefficients). *Assume that the functions $b(x, \mu)$, $b^n(x, \mu)$, $\sigma^n(x, \mu)$ and $\sigma(x, \mu)$ satisfy Assumption (A.1) and (A.2), Further suppose that for each $T > 0$, and each compact set K there exists $C > 0$ such that*

$$\lim_{n \rightarrow \infty} \sup_{t \leq T} \sup_{x \in K} \sup_{\mu \in \mathcal{P}_2(\mathbb{R})} |b^n(x, \mu) - b(x, \mu)| + |\sigma^n(x, \mu) - \sigma(x, \mu)| = 0$$

then

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] = 0$$

PROOF. Apply the Lipschitz condition of b^n and σ^n , and Cauchy-Schwarz inequality, we have

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] &\leq 4\mathbb{E} \left[\int_0^t b^n(X_s^n, \mathcal{L}_{X_s^n}) - b^n(X_s, \mathcal{L}_{X_s}) ds \right]^2 \\ &\quad + 4\mathbb{E} \left[\int_0^t b^n(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s}) ds \right]^2 \\ &\quad + 4\mathbb{E} \left\{ \sup_{t \leq T} \left[\int_0^t \sigma^n(X_s^n, \mathcal{L}_{X_s^n}) - \sigma^n(X_s, \mathcal{L}_{X_s}) dB_s \right]^2 \right\} \\ &\quad + 4\mathbb{E} \left\{ \sup_{t \leq T} \left[\int_0^t \sigma^n(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s}) dB_s \right]^2 \right\} \\ &\quad + 4 \sup_{t \leq T} |K_t^n - K_t|^2 \\ &\leq 4(CT + C) \int_0^t \mathbb{E} [|X_s^n - X_s|^2] + \mathcal{W}_2(\mathcal{L}_{X_s^n}, \mathcal{L}_{X_s})^2 ds + C\lambda_n + 4 \sup_{t \leq T} |K_t^n - K_t|^2, \end{aligned} \tag{5.4}$$

where

$$\lambda_n = \mathbb{E} \left[\int_0^t |b^n(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s})|^2 + |\sigma^n(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds \right]$$

$$\begin{aligned} \sup_{t \leq T} |K_t^n - K_t|^2 &\leq \frac{M^2}{m^2} \mathbb{E} \left[\sup_{r \in [0, t]} |U_r^n - U_r|^2 \right] \\ &\leq \frac{M^2}{m^2} 4(CT + C) \int_0^t \mathbb{E} [|X_s^n - X_s|^2] + \mathcal{W}_2(\mathcal{L}_{X_s^n}, \mathcal{L}_{X_s})^2 ds + C\lambda_n \end{aligned} \tag{5.5}$$

Apply Gronwall's lemma, we have

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] \leq C \left(1 + \frac{M^2}{m^2} \right) \exp \left\{ 8C(1 + T) \left(1 + \frac{M^2}{m^2} \right) \right\} \lambda_n \tag{5.6}$$

Letting n goes to infinity, we complete this proof.

Thirdly, we study the stability with respect to the driving processes. We consider the MR-MVSDE driving by continuous semimartingales of the following form:

$$\begin{cases} X_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) dA_s + \int_0^t \sigma(X_s, \mathcal{L}_{X_s}) dM_s + K_t, & t \geq 0 \\ \mathbb{E}[h(X_t)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s)] dK_s = 0, & t \geq 0 \end{cases} \tag{5.7}$$

where A_t is an adapted continuous process of bounded variation and M_t is a continuous local martingale.

Let us consider the following sequence $\{X_t^n\}_{n=1}^\infty$ of MR-MVSDE:

$$\begin{cases} X_t^n = \xi + \int_0^t b(X_s^n, \mathcal{L}_{X_s^n}) dA_s^n + \int_0^t \sigma(X_s^n, \mathcal{L}_{X_s^n}) dM_s^n + K_t, & t \geq 0 \\ \mathbb{E}[h(X_t^n)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s^n)] dK_s = 0, & t \geq 0 \end{cases} \quad (5.8)$$

where (A_t^n) is a sequence of \mathcal{F}_t -adapted continuous process of bounded variation and M_t^n is continuous $(\mathcal{F}_t, \mathbb{P})$ -local martingales.

Theorem 5.3 (Stability with respect to driving processes). *Assume that the functions $b(x, \mu)$, $\sigma(x, \mu)$ satisfy Assumption (A.1) and (A.2), Further suppose that (A_t, A_t^n, M_t, M_t^n) satisfy that*

- (i) *The family (A_t, A_t^n, M_t, M_t^n) is bounded in $\mathcal{C}([0, T])^{\otimes 4}$*
- (ii) *$(M_t^n - M_t)$ converges to 0 in probability in $\mathcal{C}([0, T])$*
- (iii) *The total variation $(A_t^n - A_t)$ converges to 0 in probability.*

Then

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t < T} |X_t^n - X_t|^2 \right] = 0$$

where (X_t^n) and (X_t) are, respectively, solutions of (5.8) and (5.7).

PROOF. Apply the Lipschitz condition of b and σ and Cauchy-Schwarz inequality, we have

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] &\leq 4\mathbb{E} \left\{ \sup_{t \leq T} \left[\int_0^t |b(X_s^n, \mathcal{L}_{X_s^n}) - b(X_s, \mathcal{L}_{X_s})| dA_s^n \right]^2 \right\} \\ &+ 4\mathbb{E} \left\{ \sup_{t \leq T} \left[\int_0^t |\sigma(X_s^n, \mathcal{L}_{X_s^n}) - \sigma(X_s, \mathcal{L}_{X_s})| dM_s^n \right]^2 \right\} \\ &+ 4\mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |b(X_s, \mathcal{L}_{X_s})| |dA_s^n - A_s| \right)^2 \right] \\ &+ 4\mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |\sigma(X_s, \mathcal{L}_{X_s})| |dM_s^n - M_s| \right)^2 \right] + 4 \sup_{t \leq T} |K_t^n - K_t|^2 \\ &\leq C \int_0^T \mathbb{E} \left[\sup_{s \leq t} |X_s^n - X_s|^2 \right] + \mathcal{W}_2(\mathcal{L}_{X_s^n}, \mathcal{L}_{X_s})^2 (dA_s^n + d\langle M^n \rangle_s) + e_n \\ &+ 4 \sup_{t \leq T} |K_t^n - K_t|^2 \end{aligned} \quad (5.9)$$

Where

$$e_n = 4\mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |b(X_s, \mathcal{L}_{X_s})| |dA_s^n - A_s| \right)^2 \right] + 4\mathbb{E} \left[\sup_{t \leq T} \left(\int_0^t |\sigma(X_s, \mathcal{L}_{X_s})| |dM_s^n - M_s| \right)^2 \right]$$

$$\begin{aligned}
\sup_{t \leq T} |K_t^n - K_t|^2 &\leq \frac{M^2}{m^2} \mathbb{E} \left[\sup_{r \in [0, t]} |U_r^n - U_r|^2 \right] \\
&\leq \frac{M^2}{m^2} C \int_0^t \mathbb{E} [|X_s^n - X_s|^2] + \mathcal{W}_2 (\mathcal{L}_{X_s^n}, \mathcal{L}_{X_s})^2 ds + C e_n
\end{aligned} \tag{5.10}$$

Combing (5.9) and (5.10), we have

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] \leq C \int_0^T \mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] (dA_s^n + d \langle M^n \rangle_s) + C e_n \tag{5.11}$$

Since $A_s^n + \langle M^n \rangle_s$ is an increasing process, then according to the Stochastic Gronwall lemma, we obtain

$$\mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] \leq C e_n \mathbb{E} [A_T^n + \langle M^n \rangle_T] < \infty,$$

using the assumptions of (A_t, A_t^n, M_t, M_t^n) , we have

$$\lim_{n \rightarrow \infty} e_n = 0$$

Therefore,

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \leq T} |X_t^n - X_t|^2 \right] = 0$$

5.2. Large Deviation Principle

For any $\varepsilon \in (0, 1]$, let $X^\varepsilon = \{X_t^\varepsilon, t \in [0, T]\}$ be the unique strong solution to the following MR-MVSDE:

$$\begin{cases} X_t^\varepsilon = \xi + \int_0^t b(X_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) ds + \sqrt{\varepsilon} \int_0^t \sigma(X_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) dB_s + K_t^\varepsilon, & t \geq 0 \\ \mathbb{E} [h(X_t^\varepsilon)] \geq 0, \quad \int_0^t \mathbb{E} [h(X_s^\varepsilon)] dK_s^\varepsilon = 0, & t \geq 0 \end{cases} \tag{5.12}$$

In this subsection, we mainly consider the LDP for X^ε as ε tending to 0.

Defination 5.4 (Large deviation). *A family $\{X^\varepsilon\}_{\varepsilon > 0}$ of \mathcal{E} -valued random variable is said to satisfy the large deviation principle on \mathcal{E} , with the good rate function I and with the speed function $\lambda(\varepsilon)$ which is a sequence of positive numbers tending to $+\infty$ as $\varepsilon \rightarrow 0$, if the following conditions hold:*

- (i) *for each $M < \infty$, the level set $\{x \in \mathcal{E} : I(x) \leq M\}$ is a compact subset of E ;*
- (ii) *for each closed subset F of \mathcal{E} , $\limsup_{\varepsilon \rightarrow 0} \frac{1}{\lambda(\varepsilon)} \log \mathbb{P}(X^\varepsilon \in F) \leq -\inf_{x \in F} I(x)$;*
- (iii) *for each open subset G of \mathcal{E} , $\liminf_{\varepsilon \rightarrow 0} \frac{1}{\lambda(\varepsilon)} \log \mathbb{P}(X^\varepsilon \in G) \geq -\inf_{x \in G} I(x)$.*

From Theorem 3.2, we know that there exists a unique solution for the following oblique reflected ODE

$$\begin{cases} \psi_t = \xi + \int_0^t b(\psi_s, \delta_{\psi_s}) ds + K_t^\psi, & t \geq 0 \\ h(\psi_t) \geq 0 \quad \int_0^t h(\psi_s) dK_s^\psi = 0 \end{cases}, \tag{5.13}$$

where,

$$K_s^\psi = \sup_{r \leq s} \inf \{x \geq 0, h(x + U_s^{\psi_s}) \geq 0\},$$

and $(U^\psi)_{0 \leq t \leq T}$ is the process defined by

$$U_t^\psi = \xi + \int_0^t b(\psi_s, \delta_{\psi_s}) ds \quad (5.14)$$

We denote ψ is the solution of (5.13). For any $h \in L^2([0, T]; \mathbb{R})$, consider the so called skeleton equation:

$$\begin{cases} Y_t^h = \xi + \int_0^t b(Y_s^h, \delta_{\psi_s}) ds + \int_0^t \sigma(Y_s^h, \delta_{\psi_s}) h(s) ds + K_t^h \\ h(Y_t^h) \geq 0, \quad \int_0^t h(Y_s^h) dK_s^h = 0 \end{cases} \quad (5.15)$$

Proposition 5.5. *Under Assumption (A.1) and (A.2), there exists a unique strong solution to (5.15).*

PROOF. Let $\tilde{b}(y) = b(t, y, \delta_{\psi_t}) + \sigma(t, y, \delta_{\psi_t}) h(t)$, it is easy show $\tilde{b}(y)$ satisfies Assumption (A.1). Then, apply Theorem 3.2, we complete this prove.

Theorem 5.6 (Large deviation principle). *Let Assumption (A.1) and (A.2) hold and X^ε be the unique strong solution to (5.12). Then the family of $\{X^\varepsilon\}_{\varepsilon > 0}$ satisfies a LDP on the space \mathcal{X} with rate function*

$$I(\phi) := \inf_{\{h \in L^2([0, T], \mathbb{R}) : \phi = Y^h\}} \frac{1}{2} \int_0^T |h(t)|^2 dt, \phi \in \mathcal{X},$$

with the convention $\inf\{\emptyset\} = +\infty$, here $Y^h \in \mathcal{X}$ solves equation (5.15).

PROOF. According to Proposition 5.5, there exists a measurable map \mathcal{G}^0 :

$$\mathcal{G}^0 : C([0, T]; \mathbb{R}) \rightarrow \mathcal{X}$$

such that for any $h \in L^2([0, T], \mathbb{R})$

$$Y^h = \mathcal{G}^0 \left(\int_0^\cdot h(s) ds \right).$$

Let

$$\mathcal{H}^M := \left\{ h : [0, T] \rightarrow \mathbb{R}^m, \int_0^T \|h(s)\|^2 ds \leq M \right\},$$

and

$$\tilde{\mathcal{H}}^M := \{h : h \text{ is } \mathbb{R}\text{-valued } \mathcal{F}_t\text{-predictable process such that } h(\omega) \in \mathcal{H}^M, \mathbb{P} - a.s.\}.$$

\mathcal{H}^M is endowed with the weak topology on $L^2([0, T], \mathbb{R})$. Then \mathcal{H}^M is a Polish space. For every $\varepsilon > 0$, there exists a measurable mapping $\mathcal{G}^\varepsilon(\cdot) : C([0, T]; \mathbb{R}) \rightarrow \mathcal{X}$ such that

$$X^\varepsilon = \mathcal{G}^\varepsilon(B(\cdot))$$

and for any $M > 0$ and $h^\varepsilon \in \tilde{\mathcal{H}}^M$,

$$Z^\varepsilon := \mathcal{G}^\varepsilon \left(B(\cdot) + \frac{1}{\sqrt{\varepsilon}} \int_0^\cdot h^\varepsilon(s) ds \right).$$

From Girsanov's Theorem, we know Z^ε is the solution of the following MR-MVSD:

$$\begin{cases} Z_t^\varepsilon = \xi + \int_0^t b(Z_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) ds + \int_0^t \sigma(Y_s^h, \mathcal{L}_{X_s^\varepsilon}) h^\varepsilon(s) ds + K_t^{Z^\varepsilon} \\ \mathbb{E}[h(Z_t^\varepsilon)] \geq 0, \quad \int_0^t h(Z_s^\varepsilon) dK_s^{Z^\varepsilon} = 0 \end{cases} \quad (5.16)$$

Following [39], we only need to verify the two claims:

(LDP1) For every $M < +\infty$ and any family $\{h^n; n \in \mathbb{N}\} \subset \mathcal{H}^M$ converging to some element $h \in \mathcal{H}^M$ as $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \left| \mathcal{G}^0 \left(\int_0^\cdot h^n(s) ds \right) - \mathcal{G}^0 \left(\int_0^\cdot h(s) ds \right) \right| = 0$$

(LDP2) For every $M < +\infty$ and any family $\{h^\varepsilon; \varepsilon > 0\} \subset \tilde{\mathcal{H}}^M$ and any $\theta > 0$,

$$\lim_{\varepsilon \rightarrow 0} \mathbb{P} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}| \geq \theta \right] = 0$$

where Z^ε is the unique solution to (5.16) and $Y^{h^\varepsilon} = \mathcal{G}^0 \left(\int_0^\cdot h^\varepsilon(s) ds \right)$.

The verification of (LDP1) will be given in Proposition 5.7. (LDP2) will be established in Proposition 5.9.

5.2.1. Proof of (LDP1)

Proposition 5.7. *Let Assumption (A.1) and (A.2) hold. For any $M < +\infty$, family $\{h^n\}_{n \geq 1} \in \mathcal{H}^M$ and $h \in \mathcal{H}^M$, suppose that $h^n \rightarrow h$ as $n \rightarrow \infty$ in the weak topology of $L^2([0, T], \mathbb{R})$. Then*

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \left| \mathcal{G}^0 \left(\int_0^\cdot h^n(s) ds \right) - \mathcal{G}^0 \left(\int_0^\cdot h(s) ds \right) \right| = 0$$

PROOF. For simplicity, we denote $Y^{h^n} := \mathcal{G}^0 \left(\int_0^\cdot h^n(s) ds \right)$, $Y^h = \mathcal{G}^0 \left(\int_0^\cdot h(s) ds \right)$.

$$\begin{aligned} Y^{h^n} - Y^h &= \int_0^t b(Y_s^{h^n}, \delta_{\psi_s}) - b(Y_s^h, \delta_{\psi_s}) ds + \int_0^t \sigma(Y_s^{h^n}, \delta_{\psi_s}) h^n(s) - \sigma(Y_s^h, \delta_{\psi_s}) h(s) ds \\ &\quad + K_t^{h^n} - K_t^h \\ &= \int_0^t b(Y_s^{h^n}, \delta_{\psi_s}) - b(Y_s^h, \delta_{\psi_s}) ds + \int_0^t [\sigma(Y_s^{h^n}, \delta_{\psi_s}) - \sigma(Y_s^h, \delta_{\psi_s})] h^n(s) ds \\ &\quad + \int_0^t \sigma(Y_s^h, \delta_{\psi_s}) [h^n(s) - h(s)] ds + K_t^{h^n} - K_t^h \\ &:= I_1^n(t) + I_2^n(t) + I_3^n(t) + I_4^n(t). \end{aligned} \quad (5.17)$$

Let $\kappa^n(t) := \sup_{r \in [0, t]} |Y^{h^n}(r) - Y^h(r)|$. By the Lipschitz condition of b , we have

$$|I_1^n(t)| \leq \int_0^t |b(Y_s^{h^n}, \delta_{\psi_s}) - b(Y_s^h, \delta_{\psi_s})| ds \leq K \int_0^t \kappa^n(s) ds. \quad (5.18)$$

By the Lipschitz condition of σ , we have

$$|I_2^n(t)| \leq K \int_0^t |h^n(s)| \kappa_n(s) ds \quad (5.19)$$

By the linear growth of σ , we have

$$\sup_{t \in [0, T]} |\sigma(x, \mu)|^2 \leq C(1 + K^2) \quad (5.20)$$

Since $h^n(t) \rightarrow h(t)$ weakly in $L^2([0, T], \mathbb{R})$ as $n \rightarrow \infty$, we have

$$\int_0^T |h^n(s)|^2 ds \leq C(h) \quad \int_0^T |h(s)|^2 ds \leq C(h). \quad (5.21)$$

Then, by the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sup_{t \in [0, T]} |I_3^n(t)| &\leq \int_0^T |\sigma(Y_s^h, \delta_{\psi_s}) [h^n(s) - h(s)]| ds \\ &\leq \left(\int_0^T |\sigma(Y_s^h, \psi_s)|^2 ds \right)^{\frac{1}{2}} \left(\int_0^T |h^n(s) - h(s)|^2 ds \right)^{\frac{1}{2}} \\ &\leq \left(T \sup_{t \in [0, T]} |\sigma(Y_s^h, \psi_s)|^2 \right)^{\frac{1}{2}} \left(2 \int_0^T (|h^n(s)|^2 + |h(s)|^2) ds \right)^{\frac{1}{2}} \\ &\leq (CT(1 + K^2)C(h))^{\frac{1}{2}} < \infty. \end{aligned} \quad (5.22)$$

Similarly, we have, for any $0 \leq t_1 \leq t_2 \leq T$,

$$|I_3^n(t_2) - I_3^n(t_1)| \leq [C(t_2 - t_1)(1 + K^2)C(h)]^{\frac{1}{2}},$$

which means that the sequence $\{I_3^n : n \geq 1\}$ is equi-continuous. By the Arzela-Ascoli theorem, we deduce that $\{I_3^n : n \geq 1\}$ is relatively compact in $\mathcal{C}([0, T], \mathbb{R})$.

We can deduce that for any $t \in [0, T]$,

$$I_3^n(t) = \int_0^t (h^n(s) - h(s)) \sigma(Y_s^h, \delta_{\psi_s}) ds \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (5.23)$$

Together with the relative compactness of $\{I_3^n : n \geq 1\}$ in $\mathcal{C}([0, T], \mathbb{R})$, we can get

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} |I_3^n(t)| = 0 \quad (5.24)$$

By lemma 2.4 , we have

$$K_t^{h^n} - K_t^h \leq \frac{M}{m} \sup_{r \in [0, t]} |U_r^{h^n} - U_r^h| = \frac{M}{m} \sup_{r \in [0, t]} [|I_1^n|(r) + |I_2^n|(r)] \quad (5.25)$$

Combing (5.17), (5.18), (5.19), (5.22) and (5.25) together, we have

$$\kappa^n(t) \leq C \left\{ \int_0^t \kappa^n(s) ds + \int_0^t \kappa^n(s) |h^n(s)| ds \right\} + 2 \sup_{t \in [0, T]} |I_3^n(t)|. \quad (5.26)$$

Then, by (5.23) and Gronwall's lemma, we have

$$\begin{aligned} \kappa^n(T) &\leq 2 \sup_{t \in [0, T]} |I_3^n(t)| \exp \left\{ C \int_0^T |h^n(s)| ds \right\} \\ &\leq C \sup_{t \in [0, T]} |I_3^n(t)| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (5.27)$$

The proof is complete.

5.2.2. Proof of (LDP2)

Lemma 5.8. *Suppose Assumption (A.1) and (A.2) hold, let X_t^ε and ψ_t satisfy (5.12) and (5.13), respectively. Then*

$$\lim_{\varepsilon \rightarrow 0} \sup_{t \in [0, T]} \mathcal{W}_2(\mathcal{L}_{X_t^\varepsilon}, \delta_{\psi_t})^2 = 0$$

PROOF.

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |X_t^\varepsilon - \psi_t|^2 \right] &\leq 3 \left\{ \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t b(X_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) - b(\psi_s, \delta_{\psi_s}) ds \right]^2 \right. \\ &\quad \left. + \varepsilon \mathbb{E} \left[\sup_{t \in [0, T]} \int_0^t \sigma(X_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) dB_s \right]^2 + \sup_{t \in [0, T]} |K_t^\varepsilon - K_t^\psi|^2 \right\} \\ &=: I_1 + I_2 + I_3. \end{aligned} \quad (5.28)$$

By the Lipschitz condition of b and σ , we have

$$I_1 \leq TK \int_0^T \left\{ \mathbb{E} [|X_s^\varepsilon - \psi_s|^2] + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right\} ds \quad (5.29)$$

$$I_2 \leq \mathbb{E} \int_0^T \sigma(X_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon})^2 ds \leq \mathbb{E} \int_0^T C(1 + K^2) ds = C(1 + K^2)T < \infty \quad (5.30)$$

$$I_3 = \sup_{t \in [0, T]} |K_t^\varepsilon - K_t^\psi|^2 \leq \frac{M^2}{m^2} \mathbb{E} \left[\sup_{r \in [0, t]} |U_r^\varepsilon - U_r^\psi|^2 \right] \leq 3 \frac{M^2}{m^2} (I_1 + I_2) \quad (5.31)$$

Putting (5.29), (5.30) and (5.31) to (5.28) and applying the Gronwall's lemma, we have

$$\mathbb{E} \left[\sup_{t \in [0, T]} |X_t^\varepsilon - \psi_t|^2 \right] \leq e^C \left\{ T \int_0^T \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 ds + \varepsilon \right\}. \quad (5.32)$$

By the definition of $\mathcal{W}_2(\mathcal{L}_{X_t^\varepsilon}, \delta_{\psi_t})$, we have following estimation

$$\mathcal{W}_2(\mathcal{L}_{X_t^\varepsilon}, \delta_{\psi_t})^2 \leq \mathbb{E} \left[\sup_{t \in [0, T]} |X_t^\varepsilon - \psi_t|^2 \right] \leq e^C \left\{ T \int_0^T \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 ds + \varepsilon \right\} \quad (5.33)$$

Thanking to Gronwall's lemma, we have

$$\lim_{\varepsilon \rightarrow 0} \sup_{t \in [0, T]} \mathcal{W}_2(\mathcal{L}_{X_t^\varepsilon}, \delta_{\psi_t})^2 = 0$$

Proposition 5.9. *Let Assumption (A.1) and (A.2) hold. Then for any $\theta > 0$,*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{P} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}| \geq \theta \right] = 0$$

PROOF. By Chebyshev's inequality, we only need to show that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}|^2 \right] = 0 \\ & \mathbb{E} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}|^2 \right] \leq T \mathbb{E} \left\{ \sup_{t \in [0, T]} \int_0^T |b(Z_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) - b(Y_s^{h^\varepsilon}, \delta_{\psi_s})|^2 ds \right\} \\ & \quad + \mathbb{E} \left\{ \sup_{t \in [0, T]} \left[\int_0^T h^\varepsilon(s) [\sigma(Z_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon}) - \sigma(Y_s^{h^\varepsilon}, \delta_{\psi_s})] ds \right]^2 \right\} \\ & \quad + \varepsilon C_p \int_0^T \sigma(Z_s^\varepsilon, \mathcal{L}_{X_s^\varepsilon})^2 ds + \sup_{t \in [0, T]} |K_t^{Z^\varepsilon} - K_t^{h^\varepsilon}|^2 \\ & \leq KT \sup_{t \in [0, T]} \mathbb{E} \int_0^T \left[|Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \\ & \quad + N \sup_{t \in [0, T]} \mathbb{E} \int_0^T \left[|Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \\ & \quad + \sup_{t \in [0, T]} |K_t^{Z^\varepsilon} - K_t^{h^\varepsilon}|^2 \\ & \sup_{t \in [0, T]} |K_t^{Z^\varepsilon} - K_t^{h^\varepsilon}|^2 \leq \frac{M^2}{m^2} \sup_{r \in [0, t]} |U_r^{Z^\varepsilon} - U_r^{h^\varepsilon}|^2 \\ & \leq \frac{M^2}{m^2} KT \int_0^T \mathbb{E} \left[\sup_{t \in [0, T]} |Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \\ & \quad + \frac{M^2}{m^2} N \int_0^T \mathbb{E} \left[\sup_{t \in [0, T]} |Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \end{aligned} \quad (5.34)$$

$$\begin{aligned} & \leq \frac{M^2}{m^2} KT \int_0^T \mathbb{E} \left[\sup_{t \in [0, T]} |Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \\ & \quad + \frac{M^2}{m^2} N \int_0^T \mathbb{E} \left[\sup_{t \in [0, T]} |Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 + \mathcal{W}_2(\mathcal{L}_{X_s^\varepsilon}, \delta_{\psi_s})^2 \right] ds \end{aligned} \quad (5.35)$$

By (5.34) and (5.35) and apply Gronwall's lemma, we deduce that

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}|^2 \right] &\leq (CT + N) \left(1 + \frac{M^2}{m^2}\right) \int_0^t \mathbb{E} \sup_{s \in [0, s]} |Z_s^\varepsilon - Y_s^{h^\varepsilon}|^2 ds \\ &\quad + (CT + N) \left(1 + \frac{M^2}{m^2}\right) T \sup_{t \in [0, T]} \mathcal{W}(\mathcal{L}_{X_t^\varepsilon}, \delta_{\psi_t}) + C \left(1 + \frac{M^2}{m^2}\right) \varepsilon \end{aligned} \quad (5.36)$$

Apply Gronwall's lemma, Lemma 5.8 and letting $\varepsilon \rightarrow \infty$, we get

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\sup_{t \in [0, T]} |Z_t^\varepsilon - Y_t^{h^\varepsilon}|^2 \right] = 0$$

The proof is complete.

6. Invariant probability measure and Harnack inequality

6.1. Existence of invariant probability measure

For any $X_{s,t}^x$ is the solution of (1.1) with initial condition x from s to t , let us define operator $P_{s,t}$ and its adjoin $P_{s,t}^*$ by

$$P_{s,t} f(x) := \mathbb{E} f(X_{s,t}^x) = \int_{\mathbb{R}} f(y) (P_{s,t}^* \delta_x)(dy)$$

$P_{s,t}$ is not a semigroup, but the dual $P_{s,t}^*$ is a non-linear semigroup on \mathcal{P}_2 .

We first prove the exponential convergence.

Lemma 6.1 (Exponential convergence). *Suppose Assumption (A.1), (A.2) and (A.5) hold, let X_t and Y_t be two solution of (1.1) such that $\mathcal{L}_{\xi^{X_0}} = \mu_0$, $\mathcal{L}_{\xi^{Y_0}} = \nu_0$ and $\mathcal{W}_2(\mu_0, \nu_0)^2 = \mathbb{E} [|X_0 - Y_0|^2]$. We have*

$$\mathcal{W}_2(P_t^* \mu_0, P_t^* \nu_0)^2 \leq e^{-(C_2 - C_1)t} \mathcal{W}_2(\mu_0, \nu_0)^2$$

PROOF. From the definition of P_t^* , we have $P_t^* \mu_0 = \mu_t$, $P_t^* \nu_0 = \nu_t$.

Apply Itô's lemma to $(X_t - Y_t)^2$, we have

$$\begin{aligned} (X_t - Y_t)^2 &= (\xi^{X_0} - \xi^{Y_0})^2 + \int_0^t 2(X_t - Y_t) [b(X_t, \mu_t) - b(Y_t, \nu_t)] dt + \int_0^t \sigma(X_t, \mu_t) - \sigma(Y_t, \nu_t) dt \\ &\quad + \int_0^t 2(X_t - Y_t) [\sigma(X_t, \mu_t) - \sigma(Y_t, \nu_t)] dB_t + \int_0^t 2(X_t - Y_t) d(K_t^X - K_t^Y) \end{aligned} \quad (6.1)$$

Since X_t and Y_t is the solution of (1.1) and h is bi-Lipschitz, we have $\mathbb{E}[X_t] \geq 0$ and $\mathbb{E}[Y_t] \geq 0$. Combining K_t^X and K_t^Y are non-decrease processes, we have

$$\int_0^t \mathbb{E}[X_t] dK_t^Y \geq 0 \quad \int_0^t \mathbb{E}[Y_t] dK_t^X \geq 0$$

By the definition of solution (1.1), we obtain

$$\int_0^t \mathbb{E} [h(X_t)] dK_t^X = \int_0^t \mathbb{E} [h(Y_t)] dK_t^Y = 0$$

Taking expectation of (6.1) and applying stochastic Fubini's Theorem, we have following estimation

$$\mathbb{E} (X_t - Y_t)^2 \leq (\xi^{X_0} - \xi^{Y_0})^2 + \mathbb{E} \int_0^t 2 (X_t - Y_t) [b(X_t, \mu_t) - b(Y_t, \nu_t)] dt + \mathbb{E} \int_0^t \sigma(X_t, \mu_t) - \sigma(Y_t, \nu_t) dt$$

Apply Assumption 5 and Gronwall's lemma, we obtain

$$\mathcal{W}_2 (P_t^* \mu_0, P_t^* \nu_0)^2 \leq \mathbb{E} [|X_t - Y_t|^2] \leq e^{-(C_2 - C_1)t} \mathbb{E} [|\xi^{X_0} - \xi^{Y_0}|^2] \leq e^{-(C_2 - C_1)t} \mathcal{W}_2 (\mu_0, \nu_0)^2$$

Theorem 6.2 (Existence of invariant probability measure). *Suppose Assumption (A.1), (A.2) and (A.5) hold. P_t has a unique invariant probability measure $\mu \in \mathcal{P}_2$ i.e. $P_t^* \mu = \mu, t > 0$ such that*

$$\mathcal{W}_2 (P_t^* \nu_0, \mu)^2 \leq \mathcal{W}_2 (\nu_0, \mu)^2 e^{-(C_2 - C_1)t}, \quad t \geq 0, \nu_0 \in \mathcal{P}_2.$$

PROOF. We first show that for any $\mu \in \mathcal{P}_2(\mathbb{R})$ we have

$$\lim_{t \rightarrow \infty} \mathcal{W}_2 (P_t^* \delta_0, \mu) = 0 \tag{6.2}$$

To this end, it suffices to show that $\{P_t^* \delta_0\}_{t \geq 0}$ is a \mathcal{W}_2 -Cauchy family when $t \rightarrow \infty$. Using the semigroup property of P_{t+s}^* and the exponential convergence, we have

$$\mathcal{W}_2 (P_{t+s}^* \delta_0, P_t^* \delta_0)^2 \leq \mathcal{W}_2 (P_s^* \delta_0, \delta_0)^2 e^{-(C_2 - C_1)t} = e^{-(C_2 - C_1)t} \mathbb{E} |X_s^0|^2, \quad s, t \geq 0.$$

where X_s^0 is a solution of MR-MVSDE which initial condition is $\xi = 0$. Then, we only need to show

$$\sup_{s \geq 0} \mathbb{E} [|X_s^0|^2] \leq \infty$$

It is true from Proposition 3.4. By (6.2), $P_{t+s}^* = P_t^* P_s^*$ and

$$\mathcal{W}_2 (P_t^* \mu_1, P_t^* \mu_2) \leq e^{-(C_2 - C_1)t/2} \mathcal{W}_2 (\mu_1, \mu_2),$$

we obtain that

$$\mathcal{W}_2 (P_t^* \mu, \mu) = \lim_{s \rightarrow \infty} \mathcal{W}_2 (P_t^* \mu, P_{t+s}^* \delta_0) \leq e^{-(C_2 - C_1)t/2} \lim_{s \rightarrow \infty} \mathcal{W}_2 (\mu, P_s^* \delta_0) = 0$$

That means $P_t^* \mu = \mu$.

6.2. Log-Harnack inequality and Shift Harnack inequality

In this subsection, we investigate the dimension-free log-Harnack inequality and Shift Harnack inequality. Using the coupling by change of measure method, we establish log-Harnack inequalities for $P_t f$, as detailed in [51, 57]. However, for this purpose, we need to assume that the noise term is distribution-free and that $|\sigma(x)^{-1}| \leq \lambda$. This implies that we consider a particular version of (1.1):

$$\begin{cases} X_t = \xi^{X_0} + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(X_s) dB_s + K_t, & t \geq 0 \\ \mathbb{E}[h(X_t)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s)] dK_s = 0, & t \geq 0 \end{cases} \quad (6.3)$$

For any $\mu_0 \in \mathcal{P}_2(\mathbb{R})$ and $r \geq 0$, let $B(\mu_0, r) = \{\nu \in \mathcal{P}_2(\mathbb{R}) : \mathcal{W}_2(\mu_0, \nu) \leq r\}$. Let

$$\phi(s, t) = \lambda^2 \left(\frac{C_1}{1 - e^{-C_1(t-s)}} + \frac{tC_2^2 \exp(2(C_1 + C_2)(t-s))}{2} \right), \quad 0 \leq s < t.$$

Let $\xi_t = \frac{1}{C_1}(1 - e^{C_1(t-T)})$, $\nu_t = P_t^* \nu_0$ and let ξ^{Y_0} be \mathcal{F}_0 -measurable with $\mathcal{L}_{\xi^{Y_0}} = \nu_0$. Consider the following MR-MVSDE:

$$\begin{cases} Y_t = \xi^{Y_0} + \int_0^t b(Y_s, \mathcal{L}_{Y_s}) ds + \int_0^t \frac{1}{\xi_s} \sigma(Y_s) \sigma(X_s)^{-1} (X_s - Y_s) ds + \int_0^t \sigma(Y_s) dB_s + K_t^Y, & t \geq 0 \\ \mathbb{E}[h(Y_t)] \geq 0, \quad \int_0^t \mathbb{E}[h(Y_s)] dK_s^Y = 0, & t \geq 0 \end{cases} \quad (6.4)$$

By Theorem 3.2, this MR-MVSDE has unique solution $(Y_t)_{t \in [0, T]}$. Let

$$\tau_n = T \wedge \inf \{t \in [0, T] : |X_t| + |Y_t| \geq n\},$$

we have $\tau_n \rightarrow T$ as $n \rightarrow \infty$. We first show

$$R_t := \exp \left\{ \int_0^t \frac{1}{\xi_s} \sigma(X_s)^{-1} (Y_s - X_s) dB_s - \frac{1}{2} \int_0^t \frac{|\sigma(X_s)^{-1} (Y_s - X_s)|^2}{\xi_s^2} ds \right\} \quad (6.5)$$

is a uniformly integrable martingale for $t \in [0, T]$.

Lemma 6.3. *Suppose Assumption (A.1) and (A.2) hold, and $\sigma(x)^{-1} < \infty$. Let X_0, Y_0 be two \mathcal{F}_0 -measurable random variables such that $\mathcal{L}_{\xi^{X_0}} = \mu_0$, $\mathcal{L}_{\xi^{Y_0}} = \nu_0$ and*

$$\mathbb{E} \left[|\xi^{X_0} - \xi^{Y_0}|^2 \right] = \mathcal{W}_2(\mu_0, \nu_0)^2 \quad (6.6)$$

Then $(R_t)_{t \in [0, T]}$ is a uniformly integrable martingale with

$$\sup_{t \in [0, T]} \mathbb{E}[R_t \log R_t] \leq \phi(0, T) \mathcal{W}_2(\mu_0, \nu_0)^2 \quad (6.7)$$

PROOF. By Assumption (A.1), for any $n \geq 1$, $(R_{t \wedge \tau_n})_{t \in [0, T]}$ is a uniformly integrable continuous martingale. Since $\tau_n \rightarrow T$ as $n \rightarrow \infty$, by the martingale convergence Theorem, we only need to prove

$$\sup_{t \in [0, T], n \geq 1} \mathbb{E}[R_{t \wedge \tau_n} \log R_{t \wedge \tau_n}] \leq \phi(0, T) \mathcal{W}_2(\mu_0, \nu_0)^2 \quad (6.8)$$

We fix $t \in [0, T)$ and $n \geq 1$. By Girsanov's Theorem,

$$\tilde{B}_s := B_s - \frac{1}{\xi_s} \sigma(X_s)^{-1} (Y_s - X_s), s \in [0, t \wedge \tau_n]$$

is a Brownian motion under the weighted probability $\mathbb{Q}_{t,n} := R_{t \wedge \tau_n} \mathbb{P}$.

Under $\mathbb{Q}_{t,n}$, reformulating (6.3) and (6.4)

$$\begin{cases} X_{t \wedge \tau_n} = \xi^{X_0} + \int_0^{t \wedge \tau_n} b(X_s, \mathcal{L}_{X_s}) - \frac{X_s - Y_s}{\xi_s} ds + \int_0^{t \wedge \tau_n} \sigma(X_s) d\tilde{B}_s + \tilde{K}_{t \wedge \tau_n}^X \\ \mathbb{E}^{\mathbb{Q}_{t,n}} [h(X_t)] \geq 0, \quad \int_0^t \mathbb{E}^{\mathbb{Q}_{t,n}} [h(X_s)] d\tilde{K}_s^X = 0 \end{cases} \quad (6.9)$$

$$\begin{cases} Y_{t \wedge \tau_n} = \xi^{Y_0} + \int_0^{t \wedge \tau_n} b(Y_s, \mathcal{L}_{Y_s}) ds + \int_0^{t \wedge \tau_n} \sigma(Y_s) d\tilde{B}_s + \tilde{K}_{t \wedge \tau_n}^Y \\ \mathbb{E} [h(Y_{t \wedge \tau_n})] \geq 0, \quad \int_0^{t \wedge \tau_n} \mathbb{E}^{\mathbb{Q}_{t,n}} [h(Y_s)] d\tilde{K}_s^Y = 0 \end{cases} \quad (6.10)$$

Apply Itô's Lemma, we have

$$\begin{aligned} (X_{t \wedge \tau_n} - Y_{t \wedge \tau_n})^2 &\leq \int_0^{t \wedge \tau_n} C_1 |X_s - Y_s|^2 + C_2 \mathcal{W}_2(\mu_s, \nu_s) - \frac{2|X_s - Y_s|^2}{\xi_s} ds \\ &\quad + \int_0^{t \wedge \tau_n} (X_s - Y_s) (\sigma(X_s) - \sigma(Y_s)) d\tilde{B}_s + \int_0^{t \wedge \tau_n} (X_s - Y_s) d(\tilde{K}_s^X - \tilde{K}_s^Y) \end{aligned} \quad (6.11)$$

Using the some argument of proving Lemma 6.1, we have

$$\int_0^{t \wedge \tau_n} (X_s - Y_s) d(\tilde{K}_s^X - \tilde{K}_s^Y) \leq 0$$

Then, apply Young's inequality, we have

$$\begin{aligned} \frac{(X_{t \wedge \tau_n} - Y_{t \wedge \tau_n})^2}{\xi_{t \wedge \tau_n}} &\leq \frac{C^2}{2} \int_0^{t \wedge \tau_n} \mathcal{W}_2(\mu_s, \nu_s)^2 ds - \int_0^{t \wedge \tau_n} \left(\frac{3}{2} - C_1 \xi_s \right) \frac{|X_s - Y_s|^2}{\xi_s^2} ds \\ &\quad + \int_0^{t \wedge \tau_n} \frac{(X_s - Y_s)}{\xi_s} (\sigma(X_s) - \sigma(Y_s)) d\tilde{B}_s \end{aligned} \quad (6.12)$$

By the exponential convergence (Lemma 6.1), we have

$$\mathcal{W}_2(\mu_s, \nu_s) \leq \mathcal{W}_2(\mu_0, \nu_0) e^{(C_1 + C_2)s}$$

Then, we have

$$\begin{aligned} \mathbb{E}^{\mathbb{Q}_{t,n}} \int_0^{t \wedge \tau_n} \frac{|X_s - Y_s|^2}{\xi_s^2} ds &\leq \frac{2}{\xi_0} + \mathbb{E}^{\mathbb{Q}_{t,n}} \int_0^{t \wedge \tau_n} C_2^2 \mathcal{W}_2(\mu_s, \nu_s)^2 ds \\ &\leq \frac{2}{\xi_0} + T C_2^2 e^{2(C_1 + C_2)T} \mathcal{W}_2(\mu_0, \nu_0)^2. \end{aligned} \quad (6.13)$$

Therefore, (6.8) holds since $t \in [0, T)$ and $n \geq 1$ are arbitrary.

Theorem 6.4 (Log-Harnack inequality). *Suppose Assumption (A.1) and (A.2) hold and $\sigma(x)^{-1} < \infty$. Let $0 \leq s < t \leq T$*

(i) For any $\mu_0, \nu_0 \in \mathcal{P}_2(\mathbb{R})$,

$$(P_{s,t} \log f)(\nu_0) \leq \log(P_{s,t} f)(\mu_0) + \phi(s, t) \mathcal{W}_2(\mu_0, \nu_0)^2, \quad f \in \mathcal{B}_b^+(\mathbb{R}).$$

Consequently,

$$|\nabla P_{s,t} f|^2 \leq 2\phi(s, t) \{P_{s,t} f^2 - (P_{s,t} f)^2\}, \quad f \in \mathcal{B}_b(\mathbb{R})$$

(ii) For any different $\mu_0, \nu_0 \in \mathcal{P}_2(\mathbb{R})$ and $f \in \mathcal{B}_b^+(\mathbb{R})$,

$$\frac{|(P_{s,t} f)(\mu_0) - (P_{s,t} f)(\nu_0)|^2}{\mathcal{W}_2(\mu_0, \nu_0)^2} \leq 2\phi(s, t) \sup_{\nu \in B(\mu_0, \mathcal{W}_2(\mu_0, \nu_0))} \{(P_{s,t} f^2)(\nu) - (P_{s,t} f)^2(\nu)\}.$$

Consequently,

$$\|P_{s,t}^* \mu_0 - P_{s,t}^* \nu_0\|_{var} := 2 \sup_{A \in \mathcal{B}_b(\mathbb{R})} |(P_{s,t}^* \mu_0)(A) - (P_{s,t}^* \nu_0)(A)| \leq \sqrt{2\phi(s, t)} \mathcal{W}_2(\mu_0, \nu_0).$$

PROOF. By Lemma 6.3 and Girsanov's Theorem, $d\mathbb{Q} := R_T d\mathbb{P}$ is a probability measure such that

$$\tilde{B}_t := B_t - \int_0^t \frac{\sigma(X_s)^{-1}(Y_s - X_s)}{\xi_s} ds \quad t \in [0, T]$$

is a Brownian motion. Then, (6.4) can be written to

$$\begin{cases} Y_t = \xi^{Y_0} + \int_0^t b(Y_s, \mathcal{L}_{Y_s}) ds + \int_0^t \sigma(Y_s) d\tilde{B}_s + \tilde{K}_t^Y \\ \mathbb{E}^{\mathbb{Q}}[h(Y_t)] \geq 0, \quad \int_0^t \mathbb{E}^{\mathbb{Q}}[h(Y_s)] d\tilde{K}_s^Y = 0 \end{cases} \quad (6.14)$$

Consider MR-MVSDE under \mathbb{Q} measure

$$\begin{cases} \tilde{X}_t = \xi^{Y_0} + \int_0^t b(\tilde{X}_s, \mathcal{L}_{\tilde{X}_s}) ds + \int_0^t \sigma(\tilde{X}_s) d\tilde{B}_s + \tilde{K}_t^{\tilde{X}} \\ \mathbb{E}^{\mathbb{Q}}[h(\tilde{X}_t)] \geq 0, \quad \int_0^t \mathbb{E}^{\mathbb{Q}}[h(\tilde{X}_s)] d\tilde{K}_s^{\tilde{X}} = 0 \end{cases} \quad (6.15)$$

By Theorem 3.2, we have $\tilde{X}_t = Y_t$ for $t \in [0, T]$. So we deduce that

$$(P_T \log f)(\nu_0) \leq \log(P_T f)(\mu_0) + \phi(0, T) \mathcal{W}_2(\mu_0, \nu_0)^2.$$

In particular,

$$P_T \log f(x) \leq (\log P_T f)(y) + \phi(0, T) |x - y|^2.$$

Following [64] We can get for any $f \in \mathcal{B}_b^+(\mathbb{R})$

$$|\nabla P_{s,t} f|^2 \leq 2\phi(s, t) [P_{s,t} f^2 - (P_{s,t} f)^2]$$

Repeating the proof of [46][Theorem 4.1 (ii)] we can obtain for any different μ_0 and $\nu_0 \in \mathcal{P}_2(\mathbb{R})$ and any $f \in \mathcal{B}_b^+(\mathbb{R})$,

$$\begin{aligned} & \frac{|(P_{s,t} f)(\mu_0) - (P_{s,t} f)(\nu_0)|^2}{\mathcal{W}_2(\mu_0, \nu_0)^2} \\ & \leq 2\phi(s, t) \sup_{\nu \in B(\mu_0, \mathcal{W}_2(\mu_0, \nu_0))} \{(P_{s,t} f^2)(\nu) - (P_{s,t} f)^2(\nu)\}. \end{aligned} \quad (6.16)$$

and

$$\begin{aligned} \|P_{s,t}^* \mu_0 - P_{s,t}^* \nu_0\|_{var} &:= 2 \sup_{A \in \mathcal{B}_b(\mathbb{R}^d)} |(P_{s,t}^* \mu_0)(A) - (P_{s,t}^* \nu_0)(A)| \\ &\leq \sqrt{2\phi(s,t)} \mathcal{W}_2(\mu_0, \nu_0). \end{aligned} \quad (6.17)$$

Then we establish Shift Harnack inequality for P_t introduced in [60]. We should assume that $\sigma(x, \mu)$ does not depend on x . Then, MR-MVSDE (1.1) becomes

$$\begin{cases} X_t = \xi + \int_0^t b(X_s, \mathcal{L}_{X_s}) ds + \int_0^t \sigma(\mathcal{L}_{X_s}) dB_s + K_t, & t \geq 0 \\ \mathbb{E}[h(X_t)] \geq 0, \quad \int_0^t \mathbb{E}[h(X_s)] dK_s = 0, & t \geq 0 \end{cases} \quad (6.18)$$

Theorem 6.5 (Shift Harnack inequality). *Suppose Assumption (A.1) and (A.2) hold and $\sigma(\mu)$ is invertible with $\sigma(\mu) + \sigma(\mu)^{-1}$ is bounded. For any $p > 1$, $t \in [0, T]$, $\mu \in \mathcal{P}_2(\mathbb{R})$, $\nu \in \mathbb{R}$ and $f \in \mathcal{B}_b^+(\mathbb{R})$.*

$$(P_t f)^p(\mu_0) \leq (P_t f(\nu + \cdot))^p(\mu_0) \exp \frac{p \int_0^t \sigma(\mu_s)^{-2} \{|v|/t + s|v|/t\}^2 ds}{2(p-1)}$$

Moreover, for any $f \in \mathcal{B}_b^+(\mathbb{R})$ with $f > 1$,

$$(P_t \log f)(\mu_0) \leq \log(P_t f(\nu + \cdot))(\mu_0) + \frac{1}{2} \int_0^t \sigma(\mu_s)^{-2} \{|v|/t + s|v|/t\}^2 ds$$

PROOF. Let $Y_t = X_t + \frac{tv}{T}$, then

$$dY_t = b(Y_t, \mu_t) dt + \sigma(\mu_t) d\tilde{B}_t + dK_t$$

where

$$\begin{aligned} \tilde{B}_t &:= B_t + \int_0^t \eta_s ds \\ \eta_t &:= \sigma(\mu_t)^{-1} \left\{ \frac{v}{T} + b(X_t, \mu_t) - b\left(X_t + \frac{tv}{T}, \mu_t\right) \right\}. \end{aligned}$$

Let $R_T = \exp \left\{ \int_0^T \eta_s dW_s - \frac{1}{2} \int_0^T |\eta_s|^2 ds \right\}$. Since $\mathbb{E} \left[\exp \left(\frac{1}{2} \int_0^T |\eta_s|^2 ds \right) \right] < \infty$, from Novikov's condition, we deduce that R_t is a martingale and $\mathbb{E} R_T = 1$. By the Girsanov's Theorem we can get

$$(P_T f)(\mu_0) = \mathbb{E} [R_T f(Y_T)] = \mathbb{E} [R_T f(X_T + v)] \leq (P_T f(\nu + \cdot))^{\frac{1}{p}}(\mu_0) \left(\mathbb{E} R_T^{\frac{p}{p-1}} \right)^{\frac{p-1}{p}}$$

apply Young's inequality, we obtain

$$\begin{aligned} (P_T \log f)(\mu_0) &= \mathbb{E} [R_T \log f(Y_T)] = \mathbb{E} [R_T \log f(X_T + v)] \\ &\leq \log P_T f(\nu + \cdot)(\mu_0) + \mathbb{E} [R_T \log R_T] \end{aligned}$$

Then we have

$$\mathbb{E} R_T^{\frac{p}{p-1}} \leq \exp \left\{ \frac{p}{2(p-1)^2} \int_0^T |\eta_s|^2 ds \right\} \leq \exp \left\{ \frac{p \int_0^T \sigma(\mu_t)^{-2} \{|v|/T + t|v|/T\}^2 dt}{2(p-1)^2} \right\}$$

and

$$\mathbb{E} [R_T \log R_T] = \mathbb{E}^{\mathbb{Q}} \log R_T \leq \frac{1}{2} \mathbb{E}^{\mathbb{Q}} \int_0^T |\eta_s|^2 ds \leq \frac{1}{2} \int_0^T \sigma(\mu_t)^{-2} \{|v|/T + t|v|/T\}^2 dt$$

Acknowledgements

The research is supported by NSFC (No. 71471075)

References

- [1] P. Briand, P.-E. C. De Raynal, A. Guillin, C. Labart, Particles systems and numerical schemes for mean reflected stochastic differential equations, *Ann. Appl. Probab.* 30 (4) (2020) 1884–1909.
- [2] H. P. McKean Jr, A class of Markov processes associated with nonlinear parabolic equations, *Proc. Natl. Acad. Sci.* 56 (6) (1966) 1907–1911.
- [3] A. A. Vlasov, The vibrational properties of an electron gas, *Sov. Phys. Uspekhi* 10 (6) (1968) 721.
- [4] A.-S. Sznitman, Topics in propagation of chaos, in: *Ecole d’Eté de Probabilités de Saint-Flour XIX—1989*, Vol. 1464, Springer, Berlin, Heidelberg, 2006, pp. 165–251.
- [5] R. Carmona, F. Delarue, Probabilistic theory of mean field games with applications I: Mean field FBSDEs, control, and games, *Probability theory and stochastic modelling*, Springer Nature, United States, 2018.
- [6] N. Bush, B. M. Hambly, H. Haworth, L. Jin, C. Reisinger, Stochastic evolution equations in portfolio credit modelling, *SIAM J. Financ. Math.* 2 (1) (2011) 627–664.
- [7] B. Hambly, A. Søjmark, An SPDE model for systemic risk with endogenous contagion, *Finance Stoch.* 23 (3) (2019) 535–594.
- [8] D. Dawson, J. Vaillancourt, Stochastic McKean-Vlasov equations, *Nonlinear Differ. Equ. Appl.* 2 (1995) 199–229.
- [9] A. V. Skorokhod, Stochastic equations for diffusion processes in a bounded region, *Theory Probab. Appl.* 6 (3) (1961) 264–274.
- [10] A. V. Skorokhod, Stochastic equations for diffusion processes in a bounded region. II, *Theory Probab. Appl.* 7 (1) (1962) 3–23.
- [11] P.-L. Lions, A.-S. Sznitman, Stochastic differential equations with reflecting boundary conditions, *Commun. Pure Appl. Math.* 37 (4) (1984) 511–537.
- [12] N. L. Lundström, T. Önskog, Stochastic and partial differential equations on non-smooth time-dependent domains, *Stoch. Process. Their Appl.* 129 (4) (2019) 1097–1131.
- [13] K. Burdzy, Z.-Q. Chen, J. Sylvester, The heat equation and reflected Brownian motion in time-dependent domains.: II. Singularities of solutions, *J. Funct. Anal.* 204 (1) (2003) 1–34.

- [14] K. Burdzy, Z.-Q. Chen, J. Sylvester, The heat equation and reflected Brownian motion in time-dependent domains, *Ann. Probab.* 32 (1B) (2004) 775–804.
- [15] A.-S. Sznitman, Nonlinear reflecting diffusion process, and the propagation of chaos and fluctuations associated, *J. Funct. Anal.* 56 (3) (1984) 311–336.
- [16] D. Adams, G. Dos Reis, R. Ravaille, W. Salkeld, J. Tugaut, Large deviations and exit-times for reflected McKean-Vlasov equations with self-stabilising terms and superlinear drifts, *Stoch. Process. Their Appl.* 146 (2022) 264–310.
- [17] R. Wei, S. Yang, J. Zhai, McKean-Vlasov stochastic differential equations with oblique reflection on non-smooth time dependent domains, *arXiv preprint arXiv:2208.10702* (2022).
- [18] P. Briand, R. Elie, Y. Hu, BSDEs with mean reflection, *Ann. Appl. Probab.* 28 (1) (2018) 482–510.
- [19] P. Briand, H. Hibon, Particles systems for mean reflected BSDEs, *Stoch. Process. Their Appl.* 131 (2021) 253–275.
- [20] G. Liu, F. Wang, BSDEs with mean reflection driven by G-Brownian motion, *J. Math. Anal. Appl.* 470 (1) (2019) 599–618.
- [21] F. Cui, W. Zhao, Well-posedness of mean reflected BSDEs with non-Lipschitz coefficients, *Stat. Probab. Lett.* 193 (2023) 109718.
- [22] P. Briand, A. Ghannoum, C. Labart, Mean reflected stochastic differential equations with jumps, *Adv. Appl. Probab.* 52 (2) (2020) 523–562.
- [23] A. Falkowski, L. Słomiński, Mean reflected stochastic differential equations with two constraints, *Stoch. Process. Their Appl.* 141 (2021) 172–196.
- [24] P. Briand, P. Cardaliaguet, P.-E. C. De Raynal, Y. Hu, Forward and backward stochastic differential equations with normal constraints in law, *Stoch. Process. Their Appl.* 130 (12) (2020) 7021–7097.
- [25] M. Laurière, L. Tangpi, Backward propagation of chaos, *Electron. J. Probab.* 27 (2022) 1–30.
- [26] M. I. Freidlin, A. D. Wentzell, *Random perturbations of dynamical systems*, Springer, Berlin, Heidelberg, 1984.
- [27] A. Budhiraja, P. Dupuis, V. Maroulas, Large deviations for infinite dimensional stochastic dynamical systems, *Ann. Probab.* 36 (4) (2008) 1390–1420.
- [28] M. Boué, P. Dupuis, A variational representation for certain functionals of Brownian motion, *Ann. Probab.* 26 (4) (1998) 1641–1659.

- [29] A. Jacquier, A. Pannier, Large and moderate deviations for stochastic Volterra systems, *Stoch. Process. Their Appl.* 149 (2022) 142–187.
- [30] M. Róckner, T. Zhang, X. Zhang, Large deviations for stochastic tamed 3D Navier-Stokes equations, *Appl. Math. Optim.* 61 (2010) 267–285.
- [31] A. Chiarini, M. Fischer, On large deviations for small noise itô processes, *Adv. Appl. Probab.* 46 (4) (2014) 1126–1147.
- [32] J. Zhai, T. Zhang, Large deviations for 2-D stochastic Navier-Stokes equations driven by multiplicative Lévy noises, *Bernoulli* 21 (4) (2015) 2351–2392.
- [33] J. Ren, S. Xu, X. Zhang, Large deviations for multivalued stochastic differential equations, *J. Theor. Probab.* 23 (2010) 1142–1156.
- [34] P. Fatheddin, J. Xiong, Large deviation principle for some measure-valued processes, *Stoch. Process. Their. Appl.* 125 (3) (2015) 970–993.
- [35] A. Budhiraja, P. Dupuis, Analysis and approximation of rare events: Representations and weak convergence methods, Springer, New York, NY, 2019.
- [36] A. Matoussi, W. Sabbagh, T. Zhang, Large deviation principles of obstacle problems for quasilinear stochastic PDEs, *Appl. Math. Optim.* 83 (2021) 849–879.
- [37] G. Dos Reis, W. Salkeld, J. Tugaut, Freidlin-Wentzell LDP in path space for McKean-Vlasov equations and the functional iterated logarithm law, *Ann. Appl. Probab.* 29 (3) (2019) 1487–1540.
- [38] S. Herrmann, P. Imkeller, D. Peithmann, Large deviations and a Kramers’ type law for self-stabilizing diffusions, *Ann. Appl. Probab.* 18 (4) (2008) 1379–1423.
- [39] W. Liu, Y. Song, J. Zhai, T. Zhang, Large and moderate deviation principles for McKean-Vlasov SDEs with jumps, *Potential Anal.* (2022) 1–50.
- [40] Y. Li, Large deviation principle for the mean reflected stochastic differential equation with jumps, *J. Inequal. Appl.* 2018 (2018) 1–15.
- [41] G. Da Prato, J. Zabczyk, Ergodicity for infinite dimensional systems, Vol. 229, Cambridge University Press, Cambridge, England, 1996.
- [42] H. Qiao, Exponential ergodicity for SDEs with jumps and non-Lipschitz coefficients, *J. Theor. Probab.* 27 (2014) 137–152.
- [43] X. Zhang, Exponential ergodicity of non-Lipschitz stochastic differential equations, *Proc. Am. Math. Soc.* 137 (1) (2009) 329–337.
- [44] J. Ren, J. Wu, X. Zhang, Exponential ergodicity of non-Lipschitz multivalued stochastic differential equations, *Bull. Sci. Math.* 134 (4) (2010) 391–404.

- [45] A. M. Kulik, Exponential ergodicity of the solutions to SDE's with a jump noise, *Stoch. Process. Their Appl.* 119 (2) (2009) 602–632.
- [46] F.-Y. Wang, Distribution dependent SDEs for landau type equations, *Stoch. Process. Their. Appl.* 128 (2) (2018) 595–621.
- [47] M. Liang, M. B. Majka, J. Wang, Exponential ergodicity for SDEs and McKean-Vlasov processes with lévy noise, *Ann. Inst. H. Poincaré Probab. Statist.* 57 (3) (2021) 1665–1701.
- [48] F.-Y. Wang, Exponential ergodicity for singular reflecting McKean-Vlasov SDEs, arXiv preprint arXiv:2108.03908 (2021).
- [49] P. Ren, F.-Y. Wang, Donsker-Varadhan large deviations for path-distribution dependent SPDEs, *J. Math. Anal. Appl.* 499 (1) (2021) 125000.
- [50] F.-Y. Wang, On estimation of the logarithmic sobolev constant and gradient estimates of heat semigroups, *Probab. Theory Relat. Fields* 108 (1997) 87–101.
- [51] F.-Y. Wang, Harnack inequality for SDE with multiplicative noise and extension to neumann semigroup on nonconvex manifolds, *Ann. Probab.* 39 (4) (2011) 1449–1467.
- [52] F.-Y. Wang, Harnack inequality and applications for stochastic generalized porous media equations, *Ann. Probab.* 35 (4) (2007) 1333–1350.
- [53] M. Niu, B. Xie, Wang's Harnack inequalities for space-time white noises driven SPDEs with two reflecting walls and their applications, *J. Math. Anal. Appl.* 469 (2) (2019) 568–593.
- [54] F.-Y. Wang, J. Wang, Harnack inequalities for stochastic equations driven by lévy noise, *J. Math. Anal. Appl.* 410 (1) (2014) 513–523.
- [55] L. Wang, X. Zhang, Harnack inequalities for SDEs driven by cylindrical α -stable processes, *Potential Anal.* 42 (2015) 657–669.
- [56] F.-Y. Wang, Gradient estimates and applications for SDEs in Hilbert space with multiplicative noise and Dini continuous drift, *J. Differ. Equ.* 260 (3) (2016) 2792–2829.
- [57] F.-Y. Wang, Harnack inequalities for stochastic partial differential equations, Vol. 1332, Springer, Heidelberg, Germany, 2013.
- [58] P. Ren, Singular McKean-Vlasov SDEs: Well-posedness, regularities and Wang's Harnack inequality, *Stoch. Process. Their Appl.* 156 (2023) 291–311.
- [59] X. Huang, F.-Y. Wang, Log-Harnack inequality and Bismut formula for singular McKean-Vlasov SDEs, arXiv preprint arXiv:2207.11536 (2022).
- [60] F.-Y. Wang, Integration by parts formula and shift Harnack inequality for stochastic equations, *Ann. Probab.* 42 (3) (2014) 994–1019.

- [61] X. Huang, Y. Song, Well-posedness and regularity for distribution dependent SPDEs with singular drifts, *Nonlinear Anal.* 203 (2021) 112167.
- [62] X. Huang, F.-Y. Wang, Distribution dependent SDEs with singular coefficients, *Stoch. Process. Their Appl.* 129 (11) (2019) 4747–4770.
- [63] C. Villani, *Topics in optimal transportation*, Vol. 58, American Mathematical Soc., Providence, Rhode Island, 2021.
- [64] M. Arnaudon, A. Thalmaier, F.-Y. Wang, Equivalent Harnack and gradient inequalities for pointwise curvature lower bound, *Bull. Sci. Math.* 138 (5) (2014) 643–655.