

WAVEFRONT SETS OF UNIPOTENT REPRESENTATIONS OF REDUCTIVE p -ADIC GROUPS II

DAN CIUBOTARU, LUCAS MASON-BROWN, AND EMILE OKADA

ABSTRACT. The wavefront set is a fundamental invariant of an admissible representation arising from the Harish-Chandra-Howe local character expansion. In this paper, we give a precise formula for the wavefront set of an irreducible representation of real infinitesimal character in Lusztig's category of unipotent representations in terms of the Deligne-Langlands-Lusztig correspondence. Our formula generalizes the main result of [CMO21], where this formula was obtained in the Iwahori-spherical case. We deduce that for any irreducible unipotent representation with real infinitesimal character, the algebraic wavefront set is a singleton, verifying a conjecture of Mœglin and Waldspurger. In the process, we establish new properties of the generalized Springer correspondence in relation to Lusztig's families of unipotent representations of finite reductive groups.

CONTENTS

| | |
|---|----|
| 1. Introduction | 2 |
| 1.1. Wavefront sets | 2 |
| 1.2. Representations with unipotent cuspidal support | 2 |
| 1.3. Aubert-Zelevinsky duality | 3 |
| 1.4. Main results | 3 |
| 1.5. Structure of paper | 4 |
| 1.6. Acknowledgments | 4 |
| 2. Preliminaries | 4 |
| 2.1. Notation | 4 |
| 2.2. The Bruhat-Tits Building | 5 |
| 2.3. Nilpotent orbits | 6 |
| 2.4. Wavefront sets | 9 |
| 3. Affine Hecke algebras and graded affine Hecke algebras | 9 |
| 3.1. Generic affine Hecke algebras | 10 |
| 3.2. Graded Hecke algebras | 11 |
| 3.3. Reduction theorems | 12 |
| 3.4. Tempered modules | 14 |
| 3.5. Deformations of standard modules | 14 |
| 3.6. Parabolic subalgebras | 15 |
| 3.7. Geometric graded Hecke algebras | 16 |
| 4. Lusztig's arithmetic-geometric correspondence | 18 |
| 4.1. Representations with unipotent cuspidal support | 18 |
| 4.2. The arithmetic side | 19 |
| 4.3. The geometric side | 20 |
| 4.4. Multiplicities | 21 |
| 5. Faithfulness for the generalised Springer correspondence | 22 |
| 5.1. Preliminaries on partitions | 22 |
| 5.2. Preliminaries on nilpotent orbits in classical types | 23 |
| 5.3. Preliminaries on symbols | 23 |
| 5.4. Combinatorics for Spin groups | 26 |

| | |
|---|----|
| 5.5. Unipotent representations of finite groups of Lie type | 31 |
| 5.6. The generalised Springer correspondence | 33 |
| 5.7. The Arithmetic-Geometric correspondence | 35 |
| 5.8. Faithful nilpotent orbits | 36 |
| 5.9. Proof of faithfulness in classical types | 37 |
| 5.10. Proof of faithfulness in exceptional types | 48 |
| 6. Main results | 49 |
| References | 52 |

1. INTRODUCTION

Let k be a nonarchimedean local field of characteristic 0 and residue field \mathbb{F}_q of sufficiently large characteristic, and let \mathbf{G} be a connected reductive algebraic group defined over k , inner to split. Fix an algebraic closure \bar{k} of k and let $K \subset \bar{k}$ be the maximal unramified extension of k in \bar{k} . The main result in this paper is a formula for the wavefront set of an arbitrary irreducible unipotent representation of $\mathbf{G}(k)$ of real infinitesimal character. To formulate our result, we will need to review some preliminary ideas.

1.1. Wavefront sets. Let X be an irreducible admissible representation of $\mathbf{G}(k)$. The *wavefront set* of X is a fundamental invariant related to the Harish-Chandra-Howe local character expansion. In its classical form, the wavefront set of X is a collection of nilpotent $\mathbf{G}(k)$ -orbits in the Lie algebra $\mathfrak{g}(k)$, denoted $\mathrm{WF}(X)$. We will recall the definition in Section 2.4. In this paper, we consider two coarser invariants. The first of these invariants is the *algebraic wavefront set*, denoted ${}^{\bar{k}}\mathrm{WF}(X)$. This is a collection of nilpotent orbits in $\mathfrak{g}(\bar{k})$, see for example [Wal18, p. 1108]. The second invariant is a natural refinement ${}^K\mathrm{WF}(X)$ of ${}^{\bar{k}}\mathrm{WF}(X)$ called the *canonical unramified wavefront set*, defined recently in [Oka21]. This is a collection of nilpotent orbits $\mathfrak{g}(K)$ (modulo a certain equivalence relation \sim_A). The relationship between these three invariants is as follows: the algebraic wavefront set ${}^{\bar{k}}\mathrm{WF}(X)$ is deducible from the usual wavefront set $\mathrm{WF}(X)$ as well as the canonical unramified wavefront set ${}^K\mathrm{WF}(X)$. It is not known whether the canonical unramified wavefront set is deducible from the usual wavefront set, but we expect this to be the case.

1.2. Representations with unipotent cuspidal support. In [Lus95a], Lusztig defines the notion of an irreducible *unipotent representation* of $\mathbf{G}(k)$. These are the quotients of the representations which are compactly induced from Deligne-Lusztig unipotent cuspidal representations of the reductive quotients of parahoric subgroups. We denote the set of such $\mathbf{G}(k)$ -representations by $\Pi^{\mathrm{Lus}}(\mathbf{G}(k))$. We will now briefly recall the classification of $\Pi^{\mathrm{Lus}}(\mathbf{G}(k))$ (see Section 4.1 for a more detailed statement). While our main results are independent of isogeny (see in particular Remark 4.1.2), we will assume in the introduction that \mathbf{G} is of adjoint type. Let G^\vee denote the complex points of the Langlands dual group of \mathbf{G} . Then $\Pi^{\mathrm{Lus}}(\mathbf{G}(k))$ is in one-to-one correspondence with G^\vee -conjugacy class of triples (s, n, ρ) consisting of

- a semisimple element $s \in G^\vee$,
- a nilpotent element $n \in \mathfrak{g}^\vee$ such that $\mathrm{Ad}(s)n = qn$, and
- an irreducible representation ρ of the component group of the centralizer of s and n in G^\vee .

We write $X(s, n, \rho)$ for the unipotent representation associated to the triple (s, n, ρ) . Conjugating if necessary, we can assume that s belongs to a fixed maximal torus $T^\vee \subset G^\vee$. There is a canonical decomposition $T^\vee = T_c^\vee T_r^\vee$, where T_c^\vee is the maximal compact subgroup of T^\vee and T_r^\vee is a real vector group. The representation $X(s, n, \rho)$ is said to have *real infinitesimal character* if s belongs to T_r^\vee .

1.3. Aubert-Zelevinsky duality. In [Aub95], Aubert defines an involution $X \mapsto \text{AZ}(X)$ on the Grothendieck group of finite-length smooth $\mathbf{G}(\mathbf{k})$ -representations (generalizing an involution defined by Zelevinsky in [Zel80]). For details, we refer the reader to [CMO21, Section 2.7]. After a sign normalization depending only on the Bernstein component, this involution takes irreducibles to irreducibles and preserves the set $\Pi^{\text{Lus}}(\mathbf{G}(\mathbf{k}))$ of unipotent representations. Moreover, if (s, n, ρ) is the parameter of a unipotent representation, then $\text{AZ}(X(s, n, \rho)) = X(s, n', \rho')$.

1.4. Main results. Let G be the complex connected reductive algebraic group associated to \mathbf{G} . Let \mathcal{N}_o be the set of nilpotent orbits in the Lie algebra \mathfrak{g} and let $\mathcal{N}_{o, \bar{c}}$ be the set of pairs (\mathbb{O}, \bar{C}) consisting of a nilpotent orbit $\mathbb{O} \in \mathcal{N}_o$ and a conjugacy class \bar{C} in Lusztig's canonical quotient $\bar{A}(\mathbb{O})$ of the G -equivariant fundamental group $A(\mathbb{O})$ of \mathbb{O} . Write \mathcal{N}_o^\vee and $\mathcal{N}_{o, \bar{c}}^\vee$ for the corresponding sets for G^\vee . In [Spa82, Proposition 10.3], Spaltenstein defines a duality map

$$d : \mathcal{N}_o^\vee \rightarrow \mathcal{N}_o$$

An orbit $\mathbb{O} \in \mathcal{N}_o$ is in the image of d if and only if it is special in the sense of Lusztig. In [Ach03], Achar defines a related map

$$d_A : \mathcal{N}_{o, \bar{c}}^\vee \rightarrow \mathcal{N}_{o, \bar{c}}$$

This map generalizes d in the following sense: for every orbit $\mathbb{O}^\vee \in \mathcal{N}_o^\vee$, there is a conjugacy class \bar{C}' in $\bar{A}(d(\mathbb{O}^\vee))$ such that

$$d_A(\mathbb{O}^\vee, 1) = (d(\mathbb{O}^\vee), \bar{C}')$$

Furthermore, $\bar{C}' = 1$ if \mathbb{O}^\vee is special.

There is a natural bijection between $\mathcal{N}_{o, \bar{c}}$ and the set of \sim_A -equivalence classes of unramified nilpotent orbits $\mathcal{N}_o(K)$ of $\mathbf{G}(K)$, see [Oka21, Section 5.1]. Thus, we can regard $d_A(\mathbb{O}^\vee, \bar{C})$ as an element of $\mathcal{N}_o(K) / \sim_A$.

Theorem 1.4.1 (See Theorem 6.0.3 and Corollary 6.0.5 below). *Let $X = X(s, n, \rho)$ be a unipotent representation of $\mathbf{G}(\mathbf{k})$ with real infinitesimal character and let $\text{AZ}(X) = X(s, n', \rho')$.*

(1) *The canonical unramified wavefront set ${}^K\text{WF}(X)$ is a singleton, and*

$${}^K\text{WF}(X) = d_A(\mathbb{O}_{\text{AZ}(X)}^\vee, 1),$$

where $\mathbb{O}_{\text{AZ}(X)}^\vee$ is the G^\vee -orbit of n' and d_A is the duality map defined by Achar. In particular,

$$\bar{k}\text{WF}(X) = d(\mathbb{O}_{\text{AZ}(X)}^\vee).$$

(2) *Suppose $X = X(q^{\frac{1}{2}h^\vee}, n, \rho)$ where h^\vee is the neutral element of a Lie triple attached to a nilpotent orbit $\mathbb{O}^\vee \subset \mathfrak{g}^\vee$. Then*

$$d_A(\mathbb{O}^\vee, 1) \leq_A {}^K\text{WF}(X),$$

where \leq_A is the partial order defined by Achar. In particular,

$$d(\mathbb{O}^\vee) \leq \bar{k}\text{WF}(X).$$

This result is a vast generalization of [CMO21, Theorem 1.2.1] where a similar result was proved in the case when \mathbf{G} is split and X is Iwahori-spherical. When \mathbf{G} is inner to $\text{SO}(2n+1)$ and $\text{AZ}(X)$ is tempered, a formula for ${}^k\text{WF}(X)$ was previously obtained by Waldspurger in [Wal18].

We note that the class of representations considered in Theorem 1.4.1 is probably the most general setting for a result of this form. If X is unipotent of *non-real* infinitesimal character, then both formulas in (1) may not hold.

Example 1.4.2. *Let $G^\vee = E_7$, let T^\vee be a maximal torus, and let $s \in T^\vee$ be a semisimple element such that the centralizer $C_{G^\vee}(s)$ of s is a subgroup of type $A_1 + D_6$ (there are two such s up to W -conjugacy). Let u be a unipotent element in the orbit $(2) \times (5, 3^2, 1)$ of $C_{G^\vee}(s)$. Then $A(s, u)$ is a cyclic group of order 2 and u belongs to the orbit $A_3 + A_2 + A_1$ of G^\vee . Let ρ be the sign*

representation of $A(s, u)$. We have that $d(A_3 + A_2 + A_1) = A_4 + A_2$. However if $AZ(X)$ has parameters $(sq^{\frac{1}{2}h^\vee}, u, \rho)$ then X has wavefront set $A_4 + A_1$ which is the special orbit directly below $d(A_3 + A_2 + A_1)$. This is an example where both equalities in (1) fail (it suffices to show that second fails since the first implies the second).

We note that there is a strictly weaker result which appears to hold for all unipotent representations (including the example above). Let d_S be the duality map (2.3.3) defined by Sommers [Som01]. The following conjecture is due to the third named author.

Conjecture 1.4.3. *Let X be a unipotent representation of $\mathbf{G}(\mathfrak{k})$. Then ${}^K\mathrm{WF}(X)$ is a singleton and*

$$d_S({}^K\mathrm{WF}(X)) = \mathbb{O}_{AZ(X)}^\vee.$$

This is a weakening of our main theorem in the following sense: for a representation X

$${}^K\mathrm{WF}(X) = d_A(\mathbb{O}_{AZ(X)}^\vee, 1) \Leftrightarrow d_S({}^K\mathrm{WF}(X)) = \mathbb{O}_{AZ(X)}^\vee \text{ and } \bar{\mathfrak{k}}\mathrm{WF}(X) = d(\mathbb{O}_{AZ(X)}^\vee).$$

If we leave the world of unipotent representations, further difficulties arise. For one, there is no complete classification of irreducible representations in this more general setting. Moreover, it was recently announced in [Tsa22] that $\bar{\mathfrak{k}}\mathrm{WF}(X)$ may not always be a singleton for positive depth representations.

We conclude with the following remark. The study of wavefront sets has traditionally been guided by the question ‘How can we determine the wavefront set from the L-parameter?’. However since d_S is a many to one map, Conjecture 1.4.3 suggests it is more natural to ask ‘How can we determine the nilpotent part of the L-parameter from the wavefront set of the AZ dual?’. Moreover, since ${}^K\mathrm{WF}(X)$ is a coarsening of $\mathrm{WF}(X)$ one might further wonder how much arithmetic information exactly is encoded in $\mathrm{WF}(X)$?

1.5. Structure of paper. In Section 2 we collect some preliminaries regarding the structure theory of p -adic groups, nilpotent orbits, and wavefront sets. In Section 3, we recall some facts regarding (graded) affine Hecke algebras, including Lusztig’s reduction theorems, the classification of simple modules, analytic deformations, and branching to parahoric subalgebras. In Section 4, we recall the details of Lusztig’s arithmetic-geometric correspondence, and use it to obtain a formula for the structure of the parahoric restrictions of a unipotent representation. Our main results, stated and proved in Section 6, depend on verifying a certain technical property of nilpotent orbits related to the generalized Springer correspondence. We call this property *x-faithfulness*; it can be viewed as a generalization of the notion of an *adapted pair* from [Lus20]. In Section 5 we check this property in all types. In the process, we establish a new combinatorial characterization of the generalized Springer correspondence for spin groups. In Section 6, we use *x-faithfulness* to prove our main results.

We mention that the motivation of [Lus20] is to give a new, purely algebraic, description of the Springer correspondence, via families of unipotent representations for (maximal) parahoric subgroups. We expect that our results in Section 5 will play a similar role for the generalized Springer correspondence.

1.6. Acknowledgments. This research was partially supported by EP/V046713/1. The authors thank Ruben La for helpful discussions about the generalized Springer correspondence.

2. PRELIMINARIES

2.1. Notation. Let \mathfrak{k} be a nonarchimedean local field of characteristic 0 with residue field \mathbb{F}_q of sufficiently large characteristic, ring of integers $\mathfrak{o} \subset \mathfrak{k}$, and valuation $\mathrm{val}_{\mathfrak{k}}$. Fix an algebraic closure $\bar{\mathfrak{k}}$ of \mathfrak{k} with Galois group Γ , and let $K \subset \bar{\mathfrak{k}}$ be the maximal unramified extension of \mathfrak{k} in $\bar{\mathfrak{k}}$. Let

\mathfrak{O} be the ring of integers of K . Let Frob be the geometric Frobenius element of $\text{Gal}(K/k)$, the topological generator which induces the inverse of the automorphism $x \rightarrow x^q$ of \mathbb{F}_q .

Let \mathbf{G} be a connected reductive algebraic group defined over \mathfrak{k} , inner to split, and let $\mathbf{T} \subset \mathbf{G}$ be a maximal torus. For any field F containing \mathfrak{k} , we write $\mathbf{G}(F)$, $\mathbf{T}(F)$, etc. for the groups of F -rational points. Let $\mathbf{G}_{\text{ad}} = \mathbf{G}/Z(\mathbf{G})$ denote the adjoint group of \mathbf{G} .

Write $X^*(\mathbf{T}, \bar{\mathfrak{k}})$ (resp. $X_*(\mathbf{T}, \bar{\mathfrak{k}})$) for the lattice of algebraic characters (resp. co-characters) of $\mathbf{T}(\bar{\mathfrak{k}})$, and write $\Phi(\mathbf{T}, \bar{\mathfrak{k}})$ (resp. $\Phi^\vee(\mathbf{T}, \bar{\mathfrak{k}})$) for the set of roots (resp. co-roots). Let

$$\mathcal{R} = (X^*(\mathbf{T}, \bar{\mathfrak{k}}), \Phi(\mathbf{T}, \bar{\mathfrak{k}}), X_*(\mathbf{T}, \bar{\mathfrak{k}}), \Phi^\vee(\mathbf{T}, \bar{\mathfrak{k}}), \langle , \rangle)$$

be the root datum corresponding to \mathbf{G} , and let W the associated (finite) Weyl group. Let G be the complex reductive group with the same absolute root datum as \mathbf{G} and let \mathbf{G}^\vee be the Langlands dual group of \mathbf{G} , i.e. the connected reductive algebraic group defined and split over \mathbb{Z} corresponding to the dual root datum

$$\mathcal{R}^\vee = (X_*(\mathbf{T}, \bar{\mathfrak{k}}), \Phi^\vee(\mathbf{T}, \bar{\mathfrak{k}}), X^*(\mathbf{T}, \bar{\mathfrak{k}}), \Phi(\mathbf{T}, \bar{\mathfrak{k}}), \langle , \rangle).$$

Set $\Omega = X_*(\mathbf{T}, \bar{\mathfrak{k}})/\mathbb{Z}\Phi^\vee(\mathbf{T}, \bar{\mathfrak{k}})$. The center $Z(\mathbf{G}^\vee)$ can be naturally identified with the irreducible characters $\text{Irr}\Omega$, and dually, $\Omega \cong X^*(Z(\mathbf{G}^\vee))$. For $\omega \in \Omega$, let ζ_ω denote the corresponding irreducible character of $Z(\mathbf{G}^\vee)$.

For details regarding the parametrization of inner twists of \mathbf{G} , see [Vog93, §2], [Kot84], [Kal16, §2]. For our purposes, it suffices to note that there is a natural bijection between equivalence classes of inner twists of \mathbf{G} and elements of the Galois cohomology group

$$H^1(\Gamma, \mathbf{G}_{\text{ad}}) \cong H^1(F, \mathbf{G}_{\text{ad}}(K)) \cong \Omega_{\text{ad}} \cong \text{Irr } Z(\mathbf{G}_{\text{sc}}^\vee),$$

Here $\mathbf{G}_{\text{sc}}^\vee$ denotes the Langlands dual group of \mathbf{G}_{ad} and F denotes the action of Frob on $\mathbf{G}(K)$. The isomorphism above is defined as follows: for a cohomology class h in $H^1(F, \mathbf{G}_{\text{ad}}(K))$, let z be a representative cocycle. Let $u \in \mathbf{G}_{\text{ad}}(K)$ be the image of F under z , and let ω denote the image of u in Ω_{ad} . Set $F_\omega = \text{Ad}(u) \circ F$. Then the rational structure of \mathbf{G} corresponding to h is given by F_ω . Write \mathbf{G}^ω for the connected reductive group defined over \mathfrak{k} such that $\mathbf{G}(K)^{F_\omega} = \mathbf{G}^\omega(K)$.

If H is a complex reductive group and x is an element of H or \mathfrak{h} , we write $H(x) = C_H(x)$ for the centralizer of x in H , and $A_H(x)$ for the group of connected components of $H(x)$. If S is a subset of H or \mathfrak{h} (or indeed, of $H \cup \mathfrak{h}$), we can similarly define $H(S)$ and $A_H(S)$. We will sometimes write $A(x)$, $A(S)$ when the group H is implicit. The subgroups of H of the form $H(x)$ where x is a semisimple element of H are called *pseudo-Levi* subgroups of H .

Let $\mathcal{C}(\mathbf{G}(\mathfrak{k}))$ be the category of finite-length smooth complex $\mathbf{G}(\mathfrak{k})$ -representations and let $\Pi(\mathbf{G}(\mathfrak{k})) \subset \mathcal{C}(\mathbf{G}(\mathfrak{k}))$ be the set of irreducible objects. Let $R(\mathbf{G}(\mathfrak{k}))$ denote the Grothendieck group of $\mathcal{C}(\mathbf{G}(\mathfrak{k}))$.

2.2. The Bruhat-Tits Building. In this section we will recall some standard facts about the Bruhat-Tits building.

Fix a $\omega \in \Omega$ and let \mathbf{G}^ω be the inner twist of \mathbf{G} corresponding ω as defined in the previous section. Let $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ denote the (enlarged) Bruhat-Tits building for $\mathbf{G}^\omega(\mathfrak{k})$. Let $\mathcal{B}(\mathbf{G}, K)$ denote the (enlarged) Bruhat-Tits building for $\mathbf{G}(K)$. For an apartment \mathcal{A} of $\mathcal{B}(\mathbf{G}, K)$ and $\Omega \subseteq \mathcal{A}$ we write $\mathcal{A}(\Omega, \mathcal{A})$ for the smallest affine subspace of \mathcal{A} containing Ω . The inner twist \mathbf{G}^ω of \mathbf{G} gives rise to an action of the Galois group $\text{Gal}(K/k)$ on $\mathcal{B}(\mathbf{G}, K)$ and we can (and will) identify $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ with the fixed points of this action. We use the notation $c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ to indicate that c is a face of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$. Given a maximal \mathfrak{k} -split torus \mathbf{T} of \mathbf{G}^ω , write $\mathcal{A}(\mathbf{T}, \mathfrak{k})$ for the corresponding apartment in $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$. For a face $c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ there is a group \mathbf{P}_c^\dagger defined over \mathfrak{o} such that $\mathbf{P}_c^\dagger(\mathfrak{o})$ identifies with the stabiliser of c in $\mathbf{G}(\mathfrak{k})$. There is an exact sequence

$$(2.2.1) \quad 1 \rightarrow \mathbf{U}_c(\mathfrak{o}) \rightarrow \mathbf{P}_c^\dagger(\mathfrak{o}) \rightarrow \mathbf{L}_c^\dagger(\mathbb{F}_q) \rightarrow 1,$$

where $\mathbf{U}_c(\mathfrak{o})$ is the pro-unipotent radical of $\mathbf{P}_c^\dagger(\mathfrak{o})$ and \mathbf{L}_c^\dagger is the reductive quotient of the special fibre of \mathbf{P}_c^\dagger . Let \mathbf{L}_c denote the identity component of \mathbf{L}_c^\dagger , and let \mathbf{P}_c be the subgroup of \mathbf{P}_c^\dagger defined over \mathfrak{o} such that $\mathbf{P}_c(\mathfrak{o})$ is the inverse image of $\mathbf{L}_c(\overline{\mathbb{F}}_q)$ in $\mathbf{P}_c^\dagger(\mathfrak{o})$. We also write \mathbf{T} for the well defined split torus scheme over \mathfrak{o} with generic fibre \mathbf{T} . This scheme \mathbf{T} defined over \mathfrak{o} is a subgroup of \mathbf{P}_c and the special fibre of \mathbf{T} , denoted $\overline{\mathbf{T}}$, is a maximal torus of \mathbf{L}_c . For c viewed as a face of $\mathcal{B}(\mathbf{G}, K)$, the stabiliser of c in $\mathbf{G}(K)$ identifies with $\mathbf{P}_c^\dagger(\mathfrak{O})$. It has pro-unipotent radical $\mathbf{U}_c(\mathfrak{O})$ and $\mathbf{P}_c^\dagger(\mathfrak{O})/\mathbf{U}_c(\mathfrak{O}) = \mathbf{L}_c^\dagger(\overline{\mathbb{F}}_q)$. For c a face lying in $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k}) \subseteq \mathcal{B}(\mathbf{G}, K)$, F_ω stabilises $\mathbf{P}_c(\mathfrak{O})$ and induces a Frobenius on $\mathbf{L}_c(\overline{\mathbb{F}}_q)$. The group $\mathbf{L}_c(\overline{\mathbb{F}}_q)$ consists of the fixed points of this Frobenius. The groups $\mathbf{P}_c(\mathfrak{o})$ obtained in this manner are called (k -)parahoric subgroups of \mathbf{G}^ω . When c is a chamber, then we call $\mathbf{P}_c(\mathfrak{o})$ an *Iwahori subgroup* of \mathbf{G} .

For the purposes of this paper, it will be convenient to fix a maximal k -split torus \mathbf{T} of \mathbf{G}^ω and a maximal K -split torus \mathbf{T}_1 of \mathbf{G}^ω containing \mathbf{T} and defined over k . We have that $\mathcal{A}(\mathbf{T}, k) = \mathcal{A}(\mathbf{T}_1, K)^{\text{Gal}(K/k)}$. Write $\Phi(\mathbf{T}_1, K)$ (resp. $\Psi(\mathbf{T}_1, K)$) for the set of roots of $\mathbf{G}(K)$ (resp. affine roots) with respect to $\mathbf{T}_1(K)$. For $\psi \in \Psi(\mathbf{T}, k)$ write $\dot{\psi} \in \Phi(\mathbf{T}, k)$ for the gradient of ψ , and $W = W(\mathbf{T}_1, K)$ for the Weyl group of $\mathbf{G}(K)$ with respect to $\mathbf{T}_1(K)$. We will also fix a $\text{Gal}(K/k)$ -stable chamber c_0 of $\mathcal{A}(\mathbf{T}_1, K)$ and a special point $x_0 \in c_0$. Let $\widetilde{W} = W \rtimes X_*(\mathbf{T}_1, K)$ be the (extended) affine Weyl group. The choice of special point x_0 of $\mathcal{B}(\mathbf{G}, K)$ fixes an inclusion $\Phi(\mathbf{T}_1, K) \rightarrow \Psi(\mathbf{T}_1, K)$ and an isomorphism between \widetilde{W} and $N_{\mathbf{G}(K)}(\mathbf{T}_1(K))/\mathbf{T}_1(\mathfrak{O}^\times)$. Write

$$(2.2.2) \quad \widetilde{W} \rightarrow W, \quad w \mapsto \dot{w}$$

for the natural projection map. For a face $c \subseteq \mathcal{A}(\mathbf{T}_1, K)$ let W_c be the subgroup of \widetilde{W} generated by reflections in the hyperplanes through c . The special fibre of \mathbf{T}_1 (as a scheme over \mathfrak{O}) which we denote by $\overline{\mathbf{T}}_1$, is a split maximal torus of $\mathbf{L}_c(\overline{\mathbb{F}}_q)$. Write $\Phi_c(\overline{\mathbf{T}}_1, \overline{\mathbb{F}}_q)$ for the root system of \mathbf{L}_c with respect to $\overline{\mathbf{T}}_1$. Then $\Phi_c(\overline{\mathbf{T}}_1, \overline{\mathbb{F}}_q)$ naturally identifies with the set of $\psi \in \Psi(\mathbf{T}_1, K)$ that vanish on c , and the Weyl group of $\overline{\mathbf{T}}_1$ in \mathbf{L}_c is isomorphic to W_c .

Recall that a choice of x_0 fixes an embedding $\Phi(\mathbf{T}_1, K) \rightarrow \Psi(\mathbf{T}_1, L)$. If we fix a set of simple roots $\Delta \subset \Phi(\mathbf{T}_1, K)$, this embedding determines a set of extended simple roots $\tilde{\Delta} \subseteq \Psi(\mathbf{T}_1, K)$. When $\Phi(\mathbf{T}_1, K)$ is irreducible, $\tilde{\Delta}$ is just the set $\Delta \cup \{1 - \alpha_0\}$ where α_0 is the highest root of $\Phi(\mathbf{T}_1, K)$ with respect to Δ . When $\Phi(\mathbf{T}_1, K)$ is reducible, say $\Phi(\mathbf{T}_1, K) = \cup_i \Phi_i$ where each Φ_i is irreducible, then $\tilde{\Delta} = \cup_i \tilde{\Delta}_i$ where $\tilde{\Delta}_i = \Phi_i \cap \Delta$. Fix Δ so that the chamber cut out by $\tilde{\Delta}$ is c_0 . Let

$$\mathbf{P}(\tilde{\Delta}) := \{J \subsetneq \tilde{\Delta} : J \cap \tilde{\Delta}_i \subsetneq \tilde{\Delta}_i, \forall i\}.$$

Each $J \in \mathbf{P}(\tilde{\Delta})$ cuts out a face of c_0 which we denote by $c(J)$. In particular $c(\Delta) = x_0$. Note that since $\Omega \simeq \widetilde{W}/W \rtimes \mathbb{Z}\Phi(\mathbf{T}_1, K)$ (recall \mathbf{G} is semisimple), and $W \rtimes \mathbb{Z}\Phi(\mathbf{T}_1, K)$ acts simply transitively on the chambers of $\mathcal{A}(\mathbf{T}_1, K)$, the action of \widetilde{W} on $\mathcal{A}(\mathbf{T}_1, K)$ induces an action of Ω on the faces of c_0 and hence on $\tilde{\Delta}$ (and $\mathbf{P}(\tilde{\Delta})$). For $\omega \in \Omega$ let σ_ω denote the corresponding permutation of $\tilde{\Delta}$. Let

$$\mathbf{P}^\omega(\tilde{\Delta}) := \{J \in \mathbf{P}(\tilde{\Delta}) \mid \sigma_\omega(J) = J\}$$

and let c_0^ω be the chamber of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ lying in c_0 . The set $\mathbf{P}^\omega(\tilde{\Delta})$ is an indexing set for the faces of c_0^ω . For $J \in \mathbf{P}^\omega(\tilde{\Delta})$ write $c^\omega(J)$ for the face of c_0^ω corresponding to J . The face $c^\omega(J)$ lies in $c(J)$. Moreover for $J, J' \in \mathbf{P}^\omega(\tilde{\Delta})$ (resp. $\mathbf{P}(\tilde{\Delta})$) we have $J \subseteq J'$ if and only if $\overline{c^\omega(J)} \supseteq \overline{c^\omega(J')}$ (resp. $\overline{c(J)} \supseteq \overline{c(J')}$). After this section we will write I for $\tilde{\Delta}$ and I_0 for Δ .

2.3. Nilpotent orbits. Let \mathcal{N} be the functor which takes a field F to the set of nilpotent elements in $\mathfrak{g}(F)$, and let \mathcal{N}_o be the functor which takes F to the set of adjoint $\mathbf{G}(F)$ -orbits on $\mathcal{N}(F)$. When F is k or K , we view $\mathcal{N}_o(F)$ as a partially ordered set with respect to the closure ordering in the topology induced by the topology on F . When F is algebraically closed, we view $\mathcal{N}_o(F)$ as a partially ordered set with respect to the closure ordering in the Zariski topology. To simplify the notation, we will write $\mathcal{N}(F'/F)$ (resp. $\mathcal{N}_o(F'/F)$) for $\mathcal{N}(F \rightarrow F')$ (resp. $\mathcal{N}_o(F \rightarrow F')$) where

$F \rightarrow F'$ is a morphism of fields. For $(F, F') = (k, K)$ (resp. (\bar{k}, \bar{K})), the map $\mathcal{N}_o(F'/F)$ is strictly increasing (resp. strictly increasing, non-decreasing). We will simply write \mathcal{N} for $\mathcal{N}(\mathbb{C})$ and \mathcal{N}_o for $\mathcal{N}_o(\mathbb{C})$. In this case we also define $\mathcal{N}_{o,c}$ (resp. $\mathcal{N}_{o,\bar{c}}$) to be the set of all pairs (\mathbb{O}, C) such that $\mathbb{O} \in \mathcal{N}_o$ and C is a conjugacy class in the fundamental group $A(\mathbb{O})$ of \mathbb{O} (resp. Lusztig's canonical quotient $\bar{A}(\mathbb{O})$ of $A(\mathbb{O})$, see [Som01, Section 5]). There is a natural map

$$(2.3.1) \quad \mathfrak{Q} : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_{o,\bar{c}}, \quad (\mathbb{O}, C) \mapsto (\mathbb{O}, \bar{C})$$

where \bar{C} is the image of C in $\bar{A}(\mathbb{O})$ under the natural homomorphism $A(\mathbb{O}) \twoheadrightarrow \bar{A}(\mathbb{O})$. There are also projection maps $\text{pr}_1 : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_o$, $\text{pr}_1 : \mathcal{N}_{o,\bar{c}} \rightarrow \mathcal{N}_o$. We will typically write \mathcal{N}^\vee , \mathcal{N}_o^\vee , $\mathcal{N}_{o,c}^\vee$, and $\mathcal{N}_{o,\bar{c}}^\vee$ for the sets \mathcal{N} , \mathcal{N}_o , $\mathcal{N}_{o,c}$, and $\mathcal{N}_{o,\bar{c}}$ associated to the Langlands dual group G^\vee . When we wish to emphasise the group we are working with we include it as a superscript e.g. $\mathcal{N}_o^{\mathbf{G}^\omega}$. Note that since \mathbf{G}^ω splits over K we have that $\mathcal{N}_o^{\mathbf{G}}(F) = \mathcal{N}_o^{\mathbf{G}^\omega}(F)$ for field extensions F of K . The following well-known result is due to Pommerening, Spaltenstein, Gerstenhaber, and Hesselink.

Lemma 2.3.1. *Let F be algebraically closed of sufficiently large characteristic. Then there is canonical isomorphism of partially ordered sets $\Lambda^F : \mathcal{N}_o^{\mathbf{G}}(F) \xrightarrow{\sim} \mathcal{N}_o$.*

Proof. Pommerening establishes a canonical bijection $\Lambda^F : \mathcal{N}_o^{\mathbf{G}}(F) \xrightarrow{\sim} \mathcal{N}_o$ in [Pom77, Corollary 3.5] and [Pom80, Theorem 1.5]. In classical types, it is well-known that this bijection is order-preserving (the partial order is given by the dominance order on partitions on both sides, see [Ger61] and [Hes76]). In exceptional types, one sees that Λ^F is order-preserving by examining the diagrams in [Spa82, Chapter 4]. \square

There are several duality maps relating the sets \mathcal{N}_o , $\mathcal{N}_{o,c}$, $\mathcal{N}_{o,\bar{c}}$ and their dual group counterparts.

(1) Write

$$(2.3.2) \quad d : \mathcal{N}_o \rightarrow \mathcal{N}_o^\vee, \quad d : \mathcal{N}_o^\vee \rightarrow \mathcal{N}_o.$$

for the duality maps defined by Spaltenstein ([Spa82, Proposition 10.3]), Lusztig ([Lus84c, §13.3]), and Barbasch-Vogan ([BV85, Appendix A]).

(2) Write

$$(2.3.3) \quad d_S : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_o^\vee, \quad d_S : \mathcal{N}_{o,c}^\vee \rightarrow \mathcal{N}_o$$

for the duality maps defined by Sommers ([Som01, Section 6]).

(3) Write

$$(2.3.4) \quad d_A : \mathcal{N}_{o,\bar{c}} \rightarrow \mathcal{N}_{o,\bar{c}}^\vee, \quad d_A : \mathcal{N}_{o,\bar{c}}^\vee \rightarrow \mathcal{N}_{o,\bar{c}}$$

for the duality maps defined by Achar ([Ach03, Section 1]).

Write \leq_A for Achar's preorder on $\mathcal{N}_{o,c}$ [Ach03, Introduction] and \sim_A for the equivalence relation on $\mathcal{N}_{o,c}$ induced by this pre-order, i.e.

$$(\mathbb{O}_1, C_1) \sim_A (\mathbb{O}_2, C_2) \iff (\mathbb{O}_1, C_1) \leq_A (\mathbb{O}_2, C_2) \text{ and } (\mathbb{O}_2, C_2) \leq_A (\mathbb{O}_1, C_1).$$

Write $[(\mathbb{O}, C)]$ for the equivalence class of $(\mathbb{O}, C) \in \mathcal{N}_{o,c}$. The \sim_A -equivalence classes in $\mathcal{N}_{o,c}$ coincide with the fibres of the projection map $\mathfrak{Q} : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_{o,\bar{c}}$ [Ach03, Theorem 1]. So \leq_A descends to a partial order on $\mathcal{N}_{o,\bar{c}}$, which we simply denote by \leq_A . The maps d, d_S, d_A are all order reversing with respect to the relevant pre/partial orders.

2.3.1. Structure of $\mathcal{N}_o^{\mathbf{G}}(K)$. Let \mathbf{T} be a maximal k -split torus of \mathbf{G}^ω , \mathbf{T}_1 be a maximal K -split torus of \mathbf{G}^ω defined over k and containing \mathbf{T} , and let x_0 be a special point in $\mathcal{A}(\mathbf{T}_1, K)$. In [Oka21, Section 2.1.5] the third-named author constructs a bijection

$$\theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o^{\mathbf{G}^\omega}(K) \xrightarrow{\sim} \mathcal{N}_{o,c}.$$

The composition

$$d^{un} := d_S \circ \theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o(K) \rightarrow \mathcal{N}_o^\vee$$

is independent of the choice of x_0 and natural in \mathbf{T}_1 [Oka21, Proposition 2.32].

For $\mathbb{O}_1, \mathbb{O}_2 \in \mathcal{N}_o(K)$ define $\mathbb{O}_1 \leq_A \mathbb{O}_2$ by

$$\mathbb{O}_1 \leq_A \mathbb{O}_2 \iff \mathcal{N}_o(\bar{k}/K)(\mathbb{O}_1) \leq \mathcal{N}_o(\bar{k}/K)(\mathbb{O}_2), \text{ and } d^{un}(\mathbb{O}_1) \geq d^{un}(\mathbb{O}_2)$$

and let \sim_A denote the equivalence classes of this pre-order.

Theorem 2.3.2. [Oka21, Theorem 2.33] *The composition $\Omega \circ \theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o(K) \rightarrow \mathcal{N}_{o, \bar{c}}$ descends to a (natural in \mathbf{T}_1) isomorphism of partial orders*

$$\bar{\theta} : \mathcal{N}_o(K) / \sim_A \rightarrow \mathcal{N}_{o, \bar{c}}$$

which does not depend on x_0 .

2.3.2. *Lifting nilpotent orbits.* Define

$$(2.3.5) \quad \mathcal{I}_o = \{(c, \mathbb{O}) \mid c \subseteq \mathcal{B}(\mathbf{G}, K), \mathbb{O} \in \mathcal{N}_o^{\mathbf{L}_c}(\bar{\mathbb{F}}_q)\}.$$

There is a partial order on \mathcal{I}_o , defined by

$$(c_1, \mathbb{O}_1) \leq (c_2, \mathbb{O}_2) \iff c_1 = c_2 \text{ and } \mathbb{O}_1 \leq \mathbb{O}_2,$$

and a strictly increasing surjective map [Oka21, Section 1.1.2]

$$(2.3.6) \quad \mathcal{L} : (\mathcal{I}_o, \leq) \rightarrow (\mathcal{N}_o(K), \leq).$$

Recall the groups $G = \mathbf{G}(\mathbb{C}), T = \mathbf{T}(\mathbb{C})$ from section 2.1. We call a pseudo-Levi subgroup L of G *standard* if it contains T , and we write Z_L for its center. Let $\mathcal{A} = \mathcal{A}(\mathbf{T}_1, K)$.

Recall from the end of section 2.2 the definitions of $\Delta, \tilde{\Delta}, c_0, \mathbf{P}(\tilde{\Delta})$ and $c(J)$. Note that the definitions of c_0 and $c(J)$ depend on a choice of x_0 and Δ . Let L_J denote the pseudo-Levi subgroup of G generated by T and the root groups corresponding to $\dot{\alpha}$ for $\alpha \in J$. Then L_J is the complex connected reductive group with the same absolute root datum as $\mathbf{L}_{c(J)}$. Thus Lemma ?? gives rise to an isomorphism

$$\Lambda_J^{\bar{\mathbb{F}}_q} : \mathcal{N}_o^{\mathbf{L}_{c(J)}(o)}(\bar{\mathbb{F}}_q) \rightarrow \mathcal{N}_o^{L_J}(\mathbb{C}).$$

Define

$$(2.3.7) \quad \mathcal{I}_{x_0, \tilde{\Delta}} = \{(J, \mathbb{O}) \mid J \in \mathbf{P}(\tilde{\Delta}), \mathbb{O} \in \mathcal{N}_o^{\mathbf{L}_{c(J)}}(\bar{\mathbb{F}}_q)\},$$

$$(2.3.8) \quad \mathcal{K}_{\tilde{\Delta}} = \{(J, \mathbb{O}) \mid J \in \mathbf{P}(\tilde{\Delta}), \mathbb{O} \in \mathcal{N}_o^{L_J}(\mathbb{C})\}.$$

The map

$$\iota_{x_0} : \mathcal{I}_{x_0, \tilde{\Delta}} \rightarrow \mathcal{K}_{\tilde{\Delta}}, \quad (J, \mathbb{O}) \mapsto (J, \Lambda_J^{\bar{\mathbb{F}}_q}(\mathbb{O}))$$

is a bijection. Let

$$(2.3.9) \quad \mathbb{L} : \mathcal{K}_{\tilde{\Delta}} \rightarrow \mathcal{N}_{c, o}$$

be the map [Oka21, Corollary 2.19]. By [CMO21, Theorem 2.1.7] the diagram

$$(2.3.10) \quad \begin{array}{ccc} \mathcal{I}_{x_0, \tilde{\Delta}} & \xrightarrow{\sim \iota_{x_0}} & \mathcal{K}_{\tilde{\Delta}} \\ \downarrow \mathcal{L} & & \downarrow \mathbb{L} \\ \mathcal{N}_o(K) & \xrightarrow{\sim \theta_{x_0, \mathbf{T}_1}} & \mathcal{N}_{o, c} \end{array}$$

commutes. Define

$$\bar{\mathbb{L}} = \Omega \circ \mathbb{L}.$$

This map can be computed using the algorithms in [Ach03, Section 3.4].

2.4. Wavefront sets. Let X be an admissible smooth representation of $\mathbf{G}^\omega(\mathbf{k})$ and let Θ_X be the character of X . Recall that for each nilpotent orbit $\mathbb{O} \in \mathcal{N}_o^{\mathbf{G}^\omega}(\mathbf{k})$ there is an associated distribution $\mu_{\mathbb{O}}$ on $C_c^\infty(\mathfrak{g}^\omega(\mathbf{k}))$ called the *nilpotent orbital integral* of \mathbb{O} ([Ran72]). Write $\hat{\mu}_{\mathbb{O}}$ for its Fourier transform. Generalizing a result of Howe ([How74]), Harish-Chandra in [Har99] shows that there are complex numbers $c_{\mathbb{O}}(X) \in \mathbb{C}$ such that

$$(2.4.1) \quad \Theta_X(\exp(\xi)) = \sum_{\mathbb{O}} c_{\mathbb{O}}(X) \hat{\mu}_{\mathbb{O}}(\xi)$$

for $\xi \in \mathfrak{g}^\omega(\mathbf{k})$ a regular element in a small neighborhood of 0. The formula (2.4.1) is called the *local character expansion* of X . The (p -adic) *wavefront set* of X is

$$\mathrm{WF}(X) := \max\{\mathbb{O} \mid c_{\mathbb{O}}(X) \neq 0\} \subseteq \mathcal{N}_o(\mathbf{k}).$$

The *algebraic wavefront set* of X is

$$\bar{\mathbf{k}}\mathrm{WF}(X) := \max\{\mathcal{N}_o(\bar{\mathbf{k}}/\mathbf{k})(\mathbb{O}) \mid c_{\mathbb{O}}(X) \neq 0\} \subseteq \mathcal{N}_o(\bar{\mathbf{k}}),$$

see [Wal18, p. 1108] (warning: in [Wal18], the invariant $\bar{\mathbf{k}}\mathrm{WF}(X)$ is called simply the ‘wavefront set’ of X).

In [Oka21, Section 2.2.3] the third author has introduced a third type of wavefront set for depth-0 representations, called the *canonical unramified wavefront set*. This invariant is a natural refinement of $\bar{\mathbf{k}}\mathrm{WF}(X)$. We will now define ${}^K\mathrm{WF}(X)$ and explain how to compute it.

Recall from Equation 2.3.6 the lifting map \mathcal{L} . For every face $c \subseteq \mathcal{B}(\mathbf{G})$, the space of invariants $X^{\mathbf{U}_c(o)}$ is a (finite-dimensional) $\mathbf{L}_c(\mathbb{F}_q)$ -representation. Let $\mathrm{WF}(X^{\mathbf{U}_c(o)}) \subseteq \mathcal{N}_o^{\mathbf{L}_c}(\bar{\mathbb{F}}_q)$ denote the Kawanaka wavefront set [Kaw87] and let

$$(2.4.2) \quad {}^K\mathrm{WF}_c(X) := \{[\mathcal{L}(c, \mathbb{O})] \mid \mathbb{O} \in \mathrm{WF}(X^{\mathbf{U}_c(o)})\} \subseteq \mathcal{N}_o(K).$$

Definition 2.4.1. *Let X be a depth-0 representation of $\mathbf{G}^\omega(\mathbf{k})$. The canonical unramified wavefront set of X is*

$$(2.4.3) \quad {}^K\mathrm{WF}(X) := \max\{{}^K\mathrm{WF}_c(X) \mid c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathbf{k})\} \subseteq \mathcal{N}_o(K) / \sim_A.$$

Fix $\mathbf{T}, \mathbf{T}_1, c_0, x_0, \Delta$ as at the end of section 2.2. By [Oka21, Lemma 2.36] we have that

$$(2.4.4) \quad {}^K\mathrm{WF}(X) = \max\{{}^K\mathrm{WF}_{c^\omega(J)}(X) \mid J \in \mathbf{P}^\omega(\tilde{\Delta})\}.$$

We will often want to view ${}^K\mathrm{WF}(X)$ and ${}^K\mathrm{WF}_c(X)$ as subsets of $\mathcal{N}_{o, \bar{c}}$ using the identification $\bar{\theta}_{\mathbf{T}_1}$ from Theorem 2.3.2. We will write

$${}^K\mathrm{WF}(X, \mathbb{C}) := \bar{\theta}_{\mathbf{T}_1}({}^K\mathrm{WF}(X)), \quad {}^K\mathrm{WF}_c(X, \mathbb{C}) := \bar{\theta}_{\mathbf{T}_1}({}^K\mathrm{WF}_c(X)).$$

We will also want to view $\bar{\mathbf{k}}\mathrm{WF}(X)$, which is naturally contained in $\mathcal{N}_o(\bar{\mathbf{k}})$, as a subset of \mathcal{N}_o . Thus, we will write

$$\bar{\mathbf{k}}\mathrm{WF}(X, \mathbb{C}) := \Lambda^{\bar{\mathbf{k}}}(\bar{\mathbf{k}}\mathrm{WF}(X)).$$

By [Oka21, Theorem 2.37], if ${}^K\mathrm{WF}(X)$ is a singleton, then $\bar{\mathbf{k}}\mathrm{WF}(X)$ is also a singleton and

$$(2.4.5) \quad {}^K\mathrm{WF}(X, \mathbb{C}) = (\bar{\mathbf{k}}\mathrm{WF}(X, \mathbb{C}), \bar{C}).$$

for some conjugacy class \bar{C} in $\bar{A}(\bar{\mathbf{k}}\mathrm{WF}(X, \mathbb{C}))$.

3. AFFINE HECKE ALGEBRAS AND GRADED AFFINE HECKE ALGEBRAS

In this section we will collect some preliminaries regarding affine Hecke algebras and graded affine Hecke algebras. We follow [Lus89; Lus88; Lus95b; Lus02b], [BM93], [BC13], and [Sol12].

3.1. Generic affine Hecke algebras. Let $\Psi = (X, X^\vee, R, R^\vee)$ be a reduced root datum. Let W be the finite Weyl group associated to Ψ . Fix a choice of positive roots $R^+ \subset R$, with basis Π of simple roots, and let $R^{\vee,+}$, Π^\vee be the corresponding subsets of R^\vee . Let $\ell : W \rightarrow \mathbb{Z}$ be the length function corresponding to Π . Let $\widetilde{W} = W \ltimes X$ be the extended affine Weyl group.

We write $G(\Psi)$ (or just G if there is no ambiguity) for the complex connected reductive algebraic group associated to Ψ . Then $T := X^\vee \otimes_{\mathbb{Z}} \mathbb{C}^\times$ is a maximal torus in G . Let $B \supset T$ be the Borel subgroup corresponding to R^+ . We may identify:

$$X = \text{Hom}(T, \mathbb{C}^\times), \quad X^\vee = \text{Hom}(\mathbb{C}^\times, T).$$

A parameter set for Ψ is a pair of functions (λ, λ^*) ,

$$\lambda : \Pi \rightarrow \mathbb{Z}_{\geq 0}, \quad \lambda^* : \{\alpha \in \Pi : \check{\alpha} \in 2X^\vee\} \rightarrow \mathbb{Z}_{\geq 0},$$

such that $\lambda(\alpha) = \lambda(\alpha')$ and $\lambda^*(\alpha) = \lambda^*(\alpha')$ whenever α, α' are W -conjugate.

Definition 3.1.1 (Bernstein presentation, [Lus89, Section 3]). *The affine Hecke algebra $\mathcal{H}^{\lambda, \lambda^*}(\Psi, z)$, or just $\mathcal{H}(\Psi, z)$, associated to the root datum Ψ with parameter set (λ, λ^*) is the associative algebra over $\mathbb{C}[z, z^{-1}]$ with unit (z is an indeterminate), generated by the elements T_w , $w \in W$, and θ_x , $x \in X$ subject to the relations:*

$$(3.1.1) \quad (T_{s_\alpha} + 1)(T_{s_\alpha} - z^{2\lambda(\alpha)}) = 0, \text{ for all } \alpha \in \Pi,$$

$$T_w T_{w'} = T_{ww'}, \text{ for all } w, w' \in W \text{ such that } \ell(ww') = \ell(w) + \ell(w'),$$

$$(3.1.2) \quad \theta_x \theta_{x'} = \theta_{x+x'}, \text{ for all } x, x' \in X,$$

$$\theta_x T_{s_\alpha} - T_{s_\alpha} \theta_{s_\alpha(x)} = (\theta_x - \theta_{s_\alpha(x)})(\mathcal{G}(\alpha) - 1), \text{ where } x \in X, \alpha \in \Pi, \text{ and}$$

$$(3.1.3) \quad \mathcal{G}(\alpha) = \begin{cases} \frac{\theta_\alpha z^{2\lambda(\alpha)} - 1}{\theta_\alpha - 1}, & \text{if } \check{\alpha} \notin 2X^\vee, \\ \frac{(\theta_\alpha z^{\lambda(\alpha) + \lambda^*(\alpha)} - 1)(\theta_\alpha z^{\lambda(\alpha) - \lambda^*(\alpha)} + 1)}{\theta_{2\alpha} - 1}, & \text{if } \check{\alpha} \in 2X^\vee. \end{cases}$$

Let $\mathcal{A} = \mathcal{A}(\Psi, z)$ be the algebra of regular functions on $\mathbb{C}^\times \times T$. Note that \mathcal{A} can be identified with the abelian $\mathbb{C}[z, z^{-1}]$ -subalgebra of $\mathcal{H}(\Psi, z)$ generated by $\{\theta_x : x \in X\}$, where for every $x \in X$, $\theta_x : T \rightarrow \mathbb{C}^\times$ is defined by

$$(3.1.4) \quad \theta_x(y \otimes \zeta) = \zeta^{\langle x, y \rangle}, \quad y \in X^\vee, \quad \zeta \in \mathbb{C}^\times.$$

If we denote by $\mathcal{H}_W(z)$ the $\mathbb{C}[z, z^{-1}]$ -subalgebra generated by $\{T_w : w \in W\}$, then

$$(3.1.5) \quad \mathcal{H}(\Psi, z) = \mathcal{H}_W(z) \otimes_{\mathbb{C}[z, z^{-1}]} \mathcal{A}(\Psi, z),$$

as a $\mathbb{C}[z, z^{-1}]$ -module.

Theorem 3.1.2 ([Lus89, Proposition 3.11]). *The center of $\mathcal{H}(\Psi, z)$ is $\mathcal{Z} = \mathcal{A}(\Psi, z)^W$, i.e. the W -invariants in $\mathcal{A}(\Psi, z)$.*

Let $\mathcal{H}(\Psi)\text{-mod}$ denote the category of finite dimensional $\mathcal{H}(\Psi, z)$ -modules. By Schur's lemma, every irreducible module (π, V) has a *central character*, i.e. there is a homomorphism $\chi : \mathcal{Z} \rightarrow \mathbb{C}$ such that $\pi(z)v = \chi(z)v$ for every $v \in V$ and $z \in \mathcal{Z}$. By Theorem 3.1.2, there is a one-to-one correspondence between central characters for $\mathcal{H}(\Psi, z)$ and W -conjugacy classes $(z_0, s) \in \mathbb{C}^\times \times T$. There is a block decomposition:

$$(3.1.6) \quad \mathcal{H}(\Psi, z)\text{-mod} = \coprod_{(z_0, s) \in \mathbb{C}^\times \times W \backslash T} \mathcal{H}(\Psi, z)\text{-mod}_{(z_0, s)},$$

where $\mathcal{H}(\Psi, z)\text{-mod}_{(z_0, s)}$ is the subcategory of modules with central character (z_0, s) . Let $\text{Irr}_{(z_0, s)} \mathcal{H}(\Psi, z)$ be the set of isomorphism classes of simple objects in this category. For the remainder of this section, we will assume that z_0 is a fixed element of $\mathbb{R}_{>1}$.

A second presentation of the affine Hecke algebra is due to Iwahori and Matsumoto [IM65] (and, in fact, predates the Bernstein presentation). Let $W^a = W \rtimes \mathbb{Z}R \subseteq \widetilde{W}$ be the affine Weyl group and let S^a denote a set of reflections of S^a containing in particular all s_α , $\alpha \in \Pi$. The length function on W^a , still denoted by ℓ , can be extended to a length function on \widetilde{W} . Let $L : S^a \rightarrow \mathbb{Z}_{\geq 0}$ be a \widetilde{W} -invariant function.

Definition 3.1.3 (Iwahori presentation). *The affine Hecke algebra $\mathcal{H}^L(\Psi, z)$ associated to the root datum Ψ with parameter function L is the associative algebra over $\mathbb{C}[z, z^{-1}]$ with unit (z is an indeterminate) generated by the elements T_w , $w \in \widetilde{W}$ subject to the relations:*

$$(3.1.7) \quad \begin{aligned} (T_s + 1)(T_s - z^{L(s)}) &= 0, \text{ for all } s \in S^a, \\ T_w T_{w'} &= T_{ww'}, \text{ for all } w, w' \in \widetilde{W} \text{ such that } \ell(ww') = \ell(w) + \ell(w'). \end{aligned}$$

For the dictionary between the parameter functions L and (λ, λ^*) as well as the relations between θ_x and T_w , we refer to [Lus89, Section 3].

If $J \subsetneq S^a$, denote by \widetilde{W}_J the stabiliser in \widetilde{W} of $S^a - J$. Let $\mathcal{H}_J(\Psi, z)$ the subalgebra of $\mathcal{H}^L(\Psi, z)$ generated by T_w , $w \in \widetilde{W}_J$. This is a finite Hecke algebra with parameters (possibly extended by diagram automorphisms).

3.2. Graded Hecke algebras. Fix a W -orbit $\mathcal{O} \subset T$. Following [Lus89, Section 4] and [BM93, Section 3], we will define a decreasing chain of ideals $\mathcal{I}^1, \mathcal{I}^2, \dots$ in \mathcal{A} . Let

$$(3.2.1) \quad \mathcal{I} := \{f \text{ regular function on } \mathbb{C}^\times \times T : f(1, \sigma) = 0, \text{ for all } \sigma \in \mathcal{O}\},$$

and let \mathcal{I}^k be the ideal of functions vanishing on $(1, \mathcal{O})$ to at least order k . Let $\widetilde{\mathcal{I}}^k := \mathcal{I}^k \mathcal{H} = \mathcal{H} \mathcal{I}^k$, $k \geq 1$, be the chain of ideals in $\mathcal{H}(\Psi)$, generated by the \mathcal{I}^k 's.

Definition 3.2.1. *The graded affine Hecke algebra $\mathbf{H}_{\mathcal{O}}(\Psi)$ is the associated graded of the chain of ideals $\dots \supset \widetilde{\mathcal{I}}^k \supset \dots$ in $\mathcal{H}(\Psi)$.*

Let $\mathfrak{t} = X^\vee \otimes_{\mathbb{Z}} \mathbb{C}$ be the Lie algebra of T , and let $\mathfrak{t}^* = X \otimes_{\mathbb{Z}} \mathbb{C}$ be the dual space. Extend the pairing $\langle \cdot, \cdot \rangle$ to $\mathfrak{t}^* \times \mathfrak{t}$. Let \mathbf{A} be the algebra of regular functions on $\mathbb{C} \oplus \mathfrak{t}$. Note that \mathbf{A} can be identified with $\mathbb{C}[r] \otimes_{\mathbb{C}} S(\mathfrak{t}^*)$, where $S(\cdot)$ denotes the symmetric algebra, and r is an indeterminate. In the following δ denotes the delta function.

Theorem 3.2.2 ([Lus89, Proposition 4.4 and §8.7],[BM93, Proposition 3.2]). *The graded Hecke algebra $\mathbf{H}_{\mathcal{O}}(\Psi)$ is a $\mathbb{C}[r]$ -algebra generated by $\{t_w : w \in W\}$, $S(\mathfrak{t}^*)$, and a set of orthogonal idempotents $\{E_\sigma : \sigma \in \mathcal{O}\}$ subject to the relations*

$$(3.2.2) \quad \begin{aligned} t_w \cdot t_{w'} &= t_{ww'}, \quad w, w' \in W, \quad \sum_{\sigma \in \mathcal{O}} E_\sigma = 1, \quad E_{\sigma'} E_{\sigma''} = \delta_{\sigma', \sigma''} E_{\sigma'}, \\ E_\sigma t_{s_\alpha} &= t_{s_\alpha} E_{s_\alpha \sigma}, \quad E_\sigma \omega = \omega E_\sigma, \\ \omega \cdot t_{s_\alpha} - t_{s_\alpha} \cdot s_\alpha(\omega) &= rg(\alpha) \langle \omega, \check{\alpha} \rangle, \quad \text{where } \alpha \in \Pi, \quad \omega \in \mathfrak{t}^*, \\ g(\alpha) &= \sum_{\sigma \in \mathcal{O}} E_\sigma \mu_\sigma(\alpha), \quad \text{and } \mu_\sigma(\alpha) = \begin{cases} 0, & \text{if } s_\alpha \sigma \neq \sigma, \\ 2\lambda(\alpha), & \text{if } s_\alpha \sigma = \sigma, \quad \check{\alpha} \notin 2X^\vee, \\ \lambda(\alpha) + \lambda^*(\alpha) \theta_{-\alpha}(\sigma), & \text{if } s_\alpha \sigma = \sigma, \quad \check{\alpha} \in 2X^\vee. \end{cases} \end{aligned}$$

Remark 3.2.3. *An important special case is when \mathcal{O} is formed of a single (W -invariant) element σ . Then there is only one idempotent generator $E_\sigma = 1$, so it is suppressed from the notation. The algebra $\mathbf{H}_{\mathcal{O}}(\Psi)$ is generated by $\{t_w : w \in W\}$ and $S(\mathfrak{t}^*)$ subject to the commutation relation*

$$(3.2.3) \quad \begin{aligned} \omega \cdot t_{s_\alpha} - t_{s_\alpha} \cdot s_\alpha(\omega) &= r \mu_\sigma(\alpha) \langle \omega, \check{\alpha} \rangle, \quad \alpha \in \Pi, \quad \omega \in \mathfrak{t}^*, \quad \text{where} \\ \mu_\sigma(\alpha) &= \begin{cases} 2\lambda(\alpha), & \text{if } \check{\alpha} \notin 2X^\vee, \\ \lambda(\alpha) + \lambda^*(\alpha) \theta_{-\alpha}(\sigma), & \text{if } \check{\alpha} \in 2X^\vee. \end{cases} \end{aligned}$$

Still assuming that σ is W -invariant, we have $\theta_\alpha(\sigma) \in \{\pm 1\}$, for all $\alpha \in \Pi$. If σ is in the center of group $G(\Psi)$, then $\theta_\alpha(\sigma) = 1$, for all $\alpha \in \Pi$.

We will use the notation $\mathbf{H}_{\mu_\sigma}(r)$ for the graded affine Hecke algebra in the particular case defined by equations (3.2.3). Define

$$(3.2.4) \quad W(\sigma) := \{w \in W : w\sigma = \sigma\}.$$

Then $W \cdot \sigma = \{w_j \cdot \sigma : 1 \leq j \leq n\}$, where $\{w_1 = 1, w_2, \dots, w_n\}$ are coset representatives for $W/W(\sigma)$. Then $\{E_\tau\} = \{E_{w_j\sigma} : 1 \leq j \leq n\}$, and from Proposition 3.2.2, $\mathbf{H}_\mathcal{O} = \mathbb{C}[W] \otimes (\mathcal{E} \otimes \mathbf{A})$, as a $\mathbb{C}[r]$ -vector space, where \mathcal{E} is the algebra generated by the $E_{w_j\sigma}$'s with $1 \leq j \leq n$.

Proposition 3.2.4 ([Lus89, Proposition 4.5]). *The center of $\mathbf{H}_\mathcal{O}$ is $Z = (\mathcal{E} \otimes \mathbf{A})^W$.*

It follows that the central characters of $\mathbf{H}_\mathcal{O}$ are parameterized by $W(\sigma)$ -orbits in $\mathbb{C} \oplus \mathfrak{t}$. Define the category $\text{mod}_{(r_0, x)} \mathbf{H}_\sigma$, where r acts by $r_0 > 0$ and $x \in \mathfrak{t}$.

We describe the structure of $\mathbf{H}_\mathcal{O}(\Psi)$ in more detail. Fix a $\sigma \in \mathcal{O} \subset T$. We define a root datum $\Psi_\sigma = (X, R_\sigma, X^\vee, R_\sigma^\vee)$ with positive roots R_σ^+ , defined as follows:

$$(3.2.5) \quad R_\sigma = \left\{ \alpha \in R : \theta_\alpha(\sigma) = \begin{cases} 1, & \text{if } \check{\alpha} \notin 2X^\vee, \\ \pm 1, & \text{if } \check{\alpha} \in 2X^\vee \end{cases} \right\},$$

$$R_\sigma^+ = R_\sigma \cap R^+, \quad R_\sigma^\vee = \{\check{\alpha} \in R^\vee : \alpha \in R_\sigma\}.$$

Note that Ψ_σ is the root datum for $(G(\sigma)_0, T)$, with positive roots R_σ^+ with respect to the Borel subgroup $G(\sigma)_0 \cap B$.

Define

$$(3.2.6) \quad \Gamma_\sigma := \{w \in W(\sigma) : w(R_\sigma^+) = R_\sigma^+\}.$$

Note that Γ_σ acts on the root datum Ψ_σ and preserves the set of positive roots. Thus, it defines an extended graded Hecke algebra

$$(3.2.7) \quad \mathbf{H}'_{\mu_\sigma}(\Psi_\sigma) := \mathbf{H}_{\mu_\sigma}(\Psi_\sigma) \rtimes \Gamma_\sigma.$$

Recall the coset representatives $\{w_1, \dots, w_n\}$ for $W/W(\sigma)$ from the paragraph after (3.2.4). Set

$$(3.2.8) \quad E_{i,j} = t_{w_i^{-1}w_j} E_{w_j\sigma} = E_{w_i\sigma} t_{w_i^{-1}w_j}, \text{ for all } 1 \leq i, j \leq n,$$

and let \mathcal{M}_n be the matrix algebra with basis $\{E_{i,j}\}$.

Theorem 3.2.5 ([Lus89, Section 8], [BM93, Theorem 3.3]). *There is an algebra isomorphism*

$$\mathbf{H}_\mathcal{O}(\Psi) \cong \mathcal{M}_n \otimes_{\mathbb{C}} \mathbf{H}'_{\mu_\sigma}(\Psi_\sigma) = \mathcal{M}_n \otimes_{\mathbb{C}} (\mathbf{H}_{\mu_\sigma}(\Psi_\sigma) \rtimes \Gamma_\sigma).$$

Since the only irreducible representation of \mathcal{M}_n is the n -dimensional standard representation, one obtains immediately the equivalences of categories:

$$(3.2.9) \quad \text{mod}_{(r_0, x)} \mathbf{H}_\mathcal{O}(\Psi) \cong \text{mod}_{(r_0, x)} \mathbf{H}'_{\mu_\sigma}(\Psi_\sigma).$$

3.3. Reduction theorems. In this section, we recall the relation between $\mathcal{H}(\Psi)$ and $\mathbf{H}_\mathcal{O}(\Psi)$. The torus $T = X^\vee \otimes_{\mathbb{Z}} \mathbb{C}^\times$ admits a polar decomposition $T = T_c \times T_r$, where $T_c = X^\vee \otimes_{\mathbb{Z}} S^1$, and $T_r = X^\vee \otimes_{\mathbb{Z}} \mathbb{R}_{>0}$. Consequently, every $s \in T$ decomposes uniquely as $s = s_c \cdot s_h$, with $s_c \in T_c$ and $s_h \in T_r$. We call an element $s_c \in T_c$ elliptic, and an element $s_h \in T_r$ hyperbolic. Similarly, $\mathfrak{t} = X^\vee \otimes_{\mathbb{Z}} \mathbb{C}$ admits a decomposition $\mathfrak{t} = \mathfrak{t}_{i\mathbb{R}} \oplus \mathfrak{t}_{\mathbb{R}}$, where $\mathfrak{t}_{i\mathbb{R}} = X^\vee \otimes_{\mathbb{Z}} i\mathbb{R}$ and $\mathfrak{t}_{\mathbb{R}} = X^\vee \otimes_{\mathbb{Z}} \mathbb{R}$.

There are certain completions of the Hecke algebras which appear in the reduction theorems. The algebras $\mathbb{C}[r]$, \mathcal{S} , and $\mathbb{C}[r] \otimes \mathcal{S}$ consist of polynomial functions on \mathbb{C} , \mathfrak{t} and $\mathcal{M} := \mathbb{C} \oplus \mathfrak{t}$,

respectively. Let $\widehat{\mathbb{C}[r]}$ and $\widehat{\mathcal{S}}$ be the corresponding algebras of holomorphic functions. Let \mathcal{K} and $\widehat{\mathcal{K}}$ be the fields of rational and meromorphic functions on \mathcal{M} . Finally set $\widehat{\mathbf{A}} := \mathbf{A} \otimes_{\mathcal{S}} \widehat{\mathcal{S}} \subset \widehat{\mathcal{K}}$, and

$$\begin{aligned} \widehat{\mathbf{H}}_{\mathcal{O}} &:= \mathbb{C}[W] \otimes (\widehat{\mathbb{C}[r]} \otimes \widehat{\mathbf{A}}), & \widehat{\mathbf{H}}_{\mathcal{O}}(\mathcal{K}) &:= \mathbb{C}[W] \otimes (\mathcal{E} \otimes \mathcal{K}) \supset \widehat{\mathbf{H}}_{\mathcal{O}}, \\ \widehat{\mathbf{H}}_{\mathcal{O}}(\widehat{\mathcal{K}}) &:= \mathbb{C}[W] \otimes (\mathcal{E} \otimes \widehat{\mathcal{K}}) \supset \mathbf{H}_{\mathcal{O}}(\mathcal{K}), \widehat{\mathbf{H}}_{\mathcal{O}}. \end{aligned}$$

Theorem 3.3.1 ([Lus89, Proposition 5.2] and [BM93, Theorem 3.5]). *The map $\iota : \mathbb{C}[W] \times (\mathcal{E} \otimes \widehat{\mathcal{K}}) \rightarrow \widehat{\mathbf{H}}_{\mathcal{O}}(\widehat{\mathcal{K}})$, defined by*

$$\iota(E_{\sigma}) = E_{\sigma}, \quad \iota(f) = f, \quad f \in \widehat{\mathcal{K}}, \quad \iota(t_{\alpha}) = (t_{\alpha} + 1) \left(\sum_{\sigma \in \mathcal{O}} g_{\sigma}(\alpha)^{-1} E_{\sigma} \right) - 1,$$

where

$$(3.3.1) \quad g_{\sigma}(\alpha) = 1 + \mu_{\sigma}(\alpha) \alpha^{-1} \in \widehat{\mathcal{K}},$$

is an algebra isomorphism.

For every character χ of \mathcal{Z} (the center of $\mathcal{H} = \mathcal{H}_W \otimes \mathcal{A}$), there is a maximal ideal $\mathcal{J}_{\chi} = \{z \in \mathcal{Z} : \chi(z) = 0\}$ of \mathcal{Z} . Consider the quotients

$$\mathcal{A}_{\chi} = \mathcal{A}/\mathcal{A} \cdot \mathcal{J}_{\chi}, \quad \mathcal{H}(\Psi)_{\chi} = \mathcal{H}(\Psi)/\mathcal{H}(\Psi) \cdot \mathcal{J}_{\chi}.$$

Similarly, consider the ideal $\mathcal{I}_{\bar{\chi}}$ for every character $\bar{\chi}$ of Z in $\mathbf{H}_{\mathcal{O}}$, and define the analogous quotients. Then

$$\mathbf{A}_{\bar{\chi}} := \mathbf{A}/(\mathbf{A} \cdot \mathcal{I}_{\bar{\chi}}) = \widehat{\mathbf{A}}/(\widehat{\mathbf{A}} \cdot \mathcal{I}_{\bar{\chi}}) = \widehat{\mathbf{A}}_{\bar{\chi}}, \quad \mathbf{H}_{\bar{\chi}} := \mathbf{H}_{\mathcal{O}}/(\mathbf{H}_{\mathcal{O}} \cdot \mathcal{I}_{\bar{\chi}}) = \widehat{\mathbf{H}}_{\mathcal{O}}/(\widehat{\mathbf{H}}_{\mathcal{O}} \cdot \mathcal{I}_{\bar{\chi}}) = \widehat{\mathbf{H}}_{\bar{\chi}}.$$

The map

$$\tau : \mathbb{C} \oplus \mathfrak{t}_{\mathbb{R}} \rightarrow \mathbb{C}^{\times} \times T, \quad (r_0, \nu) \mapsto (z_0, s) = (e^{r_0}, \sigma \cdot e^{\nu})$$

is $W(\sigma)$ -invariant. It matches the central characters

$$(3.3.2) \quad \tau : \bar{\chi} = W(\sigma) \cdot (r_0, \nu) \longleftrightarrow \chi = W \cdot (z_0, s).$$

Moreover, τ is a bijection onto the central characters of \mathcal{H} with elliptic part in \mathcal{O} .

Theorem 3.3.2 ([BM93, Proposition 4.1, Theorem 4.3]).

(1) *The map $\phi : \mathcal{A}[z, z^{-1}] \rightarrow \widehat{\mathbb{C}[r]} \otimes \widehat{\mathbf{A}}$ defined by*

$$(3.3.3) \quad \phi(z) = e^r, \quad \phi(\theta_x) = \sum_{\sigma \in \mathcal{O}} \theta_x(\sigma) E_{\sigma} e^x, \quad x \in X,$$

is a \mathbb{C} -algebra homomorphism which maps \mathcal{J}_{χ} to $\mathcal{I}_{\bar{\chi}}$ and defines by passage to the quotients an isomorphism $\mathcal{A}_{\chi} \cong \mathbf{A}_{\bar{\chi}}$.

(2) *The map $\Phi : \mathcal{H} \rightarrow \widehat{\mathbf{H}}_{\mathcal{O}}(\widehat{\mathcal{K}})$ defined by*

$$(3.3.4) \quad \Phi(a) = \phi(a), \quad \Phi(T_{\alpha} + 1) = \sum_{\sigma \in \mathcal{O}} E_{\sigma} (t_{\alpha} + 1) \phi(\mathcal{G}_{\alpha}) g_{\sigma}(\alpha)^{-1}$$

with \mathcal{G}_{α} as in Definition 3.1.1, and $g_{\sigma}(\alpha)$ as in (3.3.1), induces an isomorphism $\mathcal{H}(\Psi)_{\chi} \cong (\mathbf{H}_{\mathcal{O}}(\Psi))_{\bar{\chi}}$.

(3) *Assume $\sigma \in T_c$. Let $\chi = W \cdot (e^{r_0}, \sigma \cdot e^{\nu})$, $\bar{\chi} = W(\sigma) \cdot (r_0, \nu)$ be as in (3.3.2), with $(r_0, \nu) \in \mathbb{C} \oplus \mathfrak{t}_{\mathbb{R}}$. The map (3.3.4)*

$$\Phi : \mathcal{H}(\Psi)_{\chi} \xrightarrow{\cong} \mathbf{H}_{\mathcal{O}}(\Psi)_{\bar{\chi}}$$

is an analytic isomorphism in $(r_0, \nu) \in \mathbb{C} \oplus \mathfrak{t}_{\mathbb{R}}$ between two families of analytic \mathbb{C} -algebras, in the sense of [BM93, Definition 4.2].

If $\sigma \in T_c$, denote by r_{σ} the bijection defined by Φ between simple $\mathbf{H}_{\mu_{\sigma}}(\Psi_{\sigma}) \rtimes \Gamma_{\sigma}$ -modules with real central character and simple $\mathcal{H}(\Psi, z_0)$ -modules with central character having compact part in $W \cdot \sigma$.

3.4. Tempered modules. Let V be a $\mathcal{H}(\Psi, z)$ -module on which z acts by $z_0 \in \mathbb{R}_{>1}$. For every $t \in T$, define a \mathcal{A} -generalized eigenspace of V :

$$(3.4.1) \quad V_t = \{v \in V : \text{for all } x \in X, (x(t) - \theta_x)^k v = 0, \text{ for some } k \geq 0\}.$$

We say that t is a weight of V if $V_t \neq \{0\}$. Let $\Phi(V)$ denote the set of weights of V . We have $V = \bigoplus_{t \in \Phi(V)} V_t$ as \mathcal{A} -modules.

Definition 3.4.1. *We say that V is tempered if, for all $x \in X^+ := \{x \in X : \langle x, \check{\alpha} \rangle \geq 0, \text{ for all } \alpha \in R^+\}$, and all $t \in \Phi(V)$, $|x(t)| \leq 1$ holds.*

Let \bar{V} be a $\mathbf{H}'_{\mu_\sigma}(\Psi_\sigma)$ -module on which r acts by $r_0 > 0$. For every $\nu \in \mathfrak{t}$, define a \mathbf{A} -generalized eigenspace of \bar{V} ,

$$(3.4.2) \quad \bar{V}_\nu = \{v \in \bar{V} : \text{for all } \omega \in S(\mathfrak{t}^*), (\omega(\nu) - \omega)^k v = 0, \text{ for some } k \geq 0\}.$$

We say that ν is a weight of \bar{V} if $\bar{V}_\nu \neq \{0\}$. Let $\Phi(\bar{V})$ denote the set of weights of \bar{V} . We have $\bar{V} = \bigoplus_{\nu \in \Phi(\bar{V})} \bar{V}_\nu$, as \mathbf{A} -modules.

Definition 3.4.2. *We say that \bar{V} is tempered if, for all $\omega \in \mathbf{A}^+ := \{\omega \in S(\mathfrak{t}^*) : \langle \omega, \check{\alpha} \rangle \geq 0, \text{ for all } \alpha \in R^+\}$, and all $\nu \in \Phi(\bar{V})$, we have $\langle \omega, \nu \rangle \leq 0$.*

The two notions of tempered are naturally related, see for example [BM93, Sections 6, 7]: under the bijection r_σ from Theorem 3.3.2, the tempered modules correspond.

3.5. Deformations of standard modules. We make use of certain continuous deformations of affine Hecke algebras and their Schwarz completions due to Solleveld [Sol12, §4], based also on an idea of Opdam [Opd04]. Let $\mathcal{H}(\Psi, z_0)$ be the specialisation of the generic affine Hecke algebra $\mathcal{H}(\Psi, z_0)$ at $z = z_0 > 1$. There is a family of additive functors

$$(3.5.1) \quad \tilde{\sigma}_\epsilon : \text{mod}\mathcal{H}(\Psi, z_0) \rightarrow \text{mod}\mathcal{H}(\Psi, z_0^\epsilon), \quad \tilde{\sigma}_\epsilon(\pi, V) = (\pi_\epsilon, V), \quad \epsilon \in [-1, 1]$$

such that for each $w \in \widetilde{W}$, the map

$$[-1, 1] \rightarrow \text{End}(V), \quad \epsilon \mapsto \pi_\epsilon(T_w)$$

is analytic in ϵ . The functors $\tilde{\sigma}_\epsilon$ are defined by

$$\pi_\epsilon = \pi \circ \rho_\epsilon,$$

see [Sol12, Proposition 4.1.2], where ρ_ϵ is a scaling map (which is an isomorphism of topological algebras when $\epsilon \neq 0$) between certain analytic completions of $\mathcal{H}(\Psi, z_0^\epsilon)$ and $\mathcal{H}(\Psi, z_0)$, such that the maps $\epsilon \mapsto \rho_\epsilon(T_w)$ are analytic for all $w \in \widetilde{W}$. We summarize the properties of $\tilde{\sigma}_\epsilon$.

Theorem 3.5.1 ([Sol12, Corollary 4.2.2]). *The functors $\tilde{\sigma}_\epsilon$:*

- (1) *map modules with central character $s \in W \setminus T$ to modules with the same central character;*
- (2) *are equivalences of categories if $\epsilon \neq 0$;*
- (3) *map tempered modules to tempered modules when $\epsilon \geq 0$.*

We are particularly interested in the functor $\tilde{\sigma}_0$. For finite-dimensional tempered modules, a precise description is available. Let

$$(3.5.2) \quad \zeta^* : \{\text{simple tempered } \mathcal{H}(\Psi, z_0)\text{-modules}\} \longrightarrow \{\text{tempered } \mathbb{C}[X \rtimes W]\text{-modules}\}$$

denote the restriction of $\tilde{\sigma}_0$ to the set of tempered modules. Fix a compact semisimple element $\sigma \in T_c$, and let r_σ denote the corresponding bijection given after Theorem 3.3.2.

Theorem 3.5.2 ([Sol12, Theorem 2.3.1, Corollary 4.4.3]). *If $\tilde{\pi}$ is any simple tempered $\mathbf{H}_{\mu_\sigma}(\Psi_\sigma) \rtimes \Gamma_\sigma$ -module with real central character and $\tilde{\pi} \circ r_\sigma$ the corresponding simple tempered $\mathcal{H}(\Psi, z_0)$ -module, then*

$$\zeta^*(\tilde{\pi} \circ r_\sigma) = \text{Ind}_{X \rtimes W(\sigma)}^{X \rtimes W}(\mathbb{C}_\sigma \otimes \tilde{\pi}|_{W(\sigma)}).$$

Let $J \subsetneq S^a$ be given. The finite Hecke algebra $\mathcal{H}_J(\Psi, z_0^\epsilon)$ is semisimple and finite dimensional with the underlying vector space $\mathbb{C}[\widetilde{W}_J]$. Since the multiplication varies continuously with ϵ , it follows by Tits' deformation theorem, that the algebras $\mathcal{H}_J(\Psi, z_0^\epsilon)$ are isomorphic for all ϵ . Moreover since the maps $\epsilon \rightarrow \pi_\epsilon(T_w)$ are continuous in ϵ , it follows that the multiplicities of the restrictions of π_ϵ to $\mathcal{H}_J(\Psi, z_0^\epsilon)$ are constant with respect to ϵ . We record these facts in the following corollary.

Corollary 3.5.3. *Let τ be a simple $\mathcal{H}_J(\Psi, z_0)$ -module and let $\tau|_{z_0 \rightarrow 1}$ be the corresponding simple $\mathbb{C}[\widetilde{W}_J]$ -module. Then, for every finite-dimensional $\mathcal{H}(\Psi, z_0)$ -module π ,*

$$\text{Hom}_{\mathcal{H}_J(\Psi, z_0)}[\tau, \pi] = \text{Hom}_{\widetilde{W}_J}[\tau|_{z_0 \rightarrow 1}, \tilde{\sigma}_0(\pi)].$$

In particular, for every simple tempered $\pi = \tilde{\pi} \circ r_\sigma$ -module as in Theorem 3.5.2,

$$(3.5.3) \quad \text{Hom}_{\mathcal{H}_J(\Psi, z_0)}[\tau, \pi] = \text{Hom}_{\widetilde{W}_J}[\tau|_{z_0 \rightarrow 1}, \zeta^*(\pi)] = \text{Hom}_{\widetilde{W}_J}[\tau|_{z_0 \rightarrow 1}, \text{Ind}_{X \rtimes W(\sigma)}^{X \rtimes W}(\mathbb{C}_\sigma \otimes \tilde{\pi}|_{W(\sigma)})].$$

3.6. Parabolic subalgebras. For $P \subseteq \Pi$, define

$$R_P = \Pi \cap \mathbb{Q}P, \quad R_P^\vee = R^\vee \cap \mathbb{Q}P^\vee, \quad X_P = X/X \cap (P^\vee)^\perp, \quad X_P^\vee = X^\vee \cap \mathbb{Q}P^\vee.$$

Let $\Psi^P = (X, X^\vee, R_P, R_P^\vee)$ and $\Psi_P = (X_P, X_P^\vee, R_P, R_P^\vee)$ be the corresponding root data. The parameters λ, λ^* of Ψ give rise by restriction to certain parameters for Ψ^P, Ψ_P , see [Sol12, §1.4, (1.3)]. Denote the resulting affine Hecke algebras by $\mathcal{H}_P = \mathcal{H}(\Psi_P, z_0)$ and $\mathcal{H}^P = \mathcal{H}(\Psi^P, z_0)$. The algebra \mathcal{H}^P is a subalgebra of $\mathcal{H}(\Psi, z_0)$ and there is a natural surjective algebra homomorphism

$$\mathcal{H}^P \rightarrow \mathcal{H}_P, \quad \theta_x \mapsto \theta_{x_P}, \quad T_w \mapsto T_w, \quad x \in X, \quad w \in W,$$

arising from the natural projection $X \rightarrow X_P, x \mapsto x_P$.

For every $t \in T^P := \text{Hom}[X^P, \mathbb{C}^\times]$, we may define an algebra automorphism

$$\chi_t : \mathcal{H}^P \rightarrow \mathcal{H}^P, \quad \chi_t(\theta_x) = t(x)\theta_x, \quad \chi_t(T_w) = T_w.$$

Every finite-dimensional \mathcal{H}_P -module π_P can be regarded as an \mathcal{H}^P -module by pullback and we may define the parabolically induced representations

$$(3.6.1) \quad I(P, \pi_P, t) = \text{Ind}_{\mathcal{H}^P}^{\mathcal{H}_P}(\pi_P \circ \chi_t), \quad t \in T^P.$$

Since the action varies continuously in t , the restriction of $I(P, \pi_P, t)$ to any finite Hecke algebra $\mathcal{H}_J(\Psi, z_0)$ is independent of t . It is clear from the definitions that if π_P is a simple tempered module for \mathcal{H}_P , then $I(P, \pi_P, 1) = \text{Ind}_{\mathcal{H}^P}^{\mathcal{H}_P}(\pi_P)$ is a (not necessarily simple) tempered module for \mathcal{H} . Hence Corollary 3.5.3 implies at once:

Proposition 3.6.1. *Let π_P be a simple tempered \mathcal{H}_P -module, $t \in T^P$, and $J \subsetneq S^a$. Then for every simple $\mathcal{H}_J(\Psi, z_0)$ -module τ ,*

$$(3.6.2) \quad \text{Hom}_{\mathcal{H}_J(\Psi, z_0)}[\tau, I(P, \pi_P, t)] = \text{Hom}_{\widetilde{W}_J}[\tau|_{z_0 \rightarrow 1}, \zeta^*(I(P, \pi_P, 1))],$$

which can be computed using Corollary 3.5.3.

3.7. Geometric graded Hecke algebras. We analyze the graded Hecke algebra $\mathbf{H} = \mathbf{H}_{\mu_\sigma}(\Psi_\sigma)$ defined by the relations in (3.2.3) for the case $\sigma = 1$. We will consider only the Hecke algebras arising by the constructions of [Lus88; Lus95b]. Let G be a complex connected reductive group, with a fixed Borel subgroup B , and maximal torus $A \subset B$. The Lie algebras will be denoted by the corresponding Gothic letters.

Definition 3.7.1. *A cuspidal triple for G is a triple $(L, \mathbb{O}, \mathcal{E})$, where L is a Levi subgroup of G , \mathbb{O} is a nilpotent adjoint L -orbit on the Lie algebra \mathfrak{l} , and \mathcal{E} is an irreducible G -equivariant local system on \mathbb{O} which is cuspidal in the sense of [Lus84b, Definition 2.4].*

Let $\mathfrak{L}(G)$ denote the set of G -conjugacy classes of cuspidal triples for G . Let us fix $(L, \mathbb{O}, \mathcal{E}) \in \mathfrak{L}(G)$, such that $A \subset L$, and $P = LU \supset B$ is a parabolic subgroup. Let T denote the identity component of the center of L . Set $W = N_G(L)/L$, a (finite) Coxeter group.

Let $\mathbf{H} = \mathbf{H}(L, \mathbb{O}, \mathcal{E})$ denote the graded affine Hecke algebra attached to $(L, \mathbb{O}, \mathcal{E})$ in [Lus88, section 2]. The explicit classification of cuspidal triples when G is simple, along with the corresponding values for the parameters of the graded Hecke algebra can be found in the tables of [Lus88, §2.13].

Consider the algebraic varieties

$$(3.7.1) \quad \tilde{\mathfrak{g}} = \{(x, g) \in \mathfrak{g} \times G : \text{Ad}(g^{-1})x \in \mathbb{O} + \mathfrak{t} + \mathfrak{u}\}, \quad \dot{\mathfrak{g}} = \{(x, gP) \in \mathfrak{g} \times G/P : (x, g) \in \tilde{\mathfrak{g}}\},$$

on which $G \times \mathbb{C}^\times$ acts via $(g_1, \lambda): x \mapsto \lambda^{-2}\text{Ad}(g_1)x$, $x \in \mathfrak{g}$, and $gP \mapsto g_1gP$, $g \in G$. Let $\text{pr}_P : \tilde{\mathfrak{g}} \rightarrow \dot{\mathfrak{g}}$ denote the obvious $G \times \mathbb{C}^\times$ -equivariant map.

If \mathcal{V} is any $G \times \mathbb{C}^\times$ -stable subvariety of \mathfrak{g} , denote by $\dot{\mathcal{V}}$ the preimage under the first projection $\text{pr}_1 : \dot{\mathfrak{g}} \rightarrow \mathfrak{g}$. Let \mathcal{N} denote the variety of nilpotent elements in \mathfrak{g} . Define

$$\mathcal{P}_n = \text{pr}_2(\dot{\mathcal{N}}) = \{gP : \text{Ad}(g^{-1})n \in \mathbb{O} + \mathfrak{u}\},$$

for any $n \in \mathcal{N}$, where $\text{pr}_2 : \dot{\mathfrak{g}} \rightarrow G/P$ is the second projection. Define also $\mathcal{P}_n^s = \{gP \in \mathcal{P}_n : \text{Ad}(g^{-1})s \in \mathfrak{p}\}$, for any semisimple $s \in \mathfrak{g}$.

Consider the $G \times \mathbb{C}^\times$ -equivariant projection $\text{pr}_\mathbb{O} : \tilde{\mathfrak{g}} \rightarrow \mathbb{O}$, $\text{pr}_\mathbb{O}(x, g) = \text{pr}_\mathbb{O}(\text{Ad}(g^{-1})x)$. Let $\dot{\mathcal{E}}$ be the $G \times \mathbb{C}^\times$ -equivariant local system on $\dot{\mathfrak{g}}$ defined by the condition $\text{pr}_\mathbb{O}^*(\mathcal{E}) = \text{pr}_P^*(\dot{\mathcal{E}})$.

The classification of simple modules for \mathbf{H} is in [Lus88; Lus95b; Lus02b]. Fix a semisimple element $s \in \mathfrak{g}$ and $r_0 \in \mathbb{C}^\times$, and let $\mathcal{T} = \mathcal{T}_{s, r_0}$ be the smallest torus in $G \times \mathbb{C}^\times$ whose Lie algebra contains (s, r_0) . Let \mathfrak{g}_{2r_0} be the set of \mathcal{T} -fixed vectors in \mathfrak{g} , namely

$$(3.7.2) \quad \mathfrak{g}_{2r_0} = \{x \in \mathfrak{g} : [s, x] = 2r_0x\}.$$

Let $G(s) \times \mathbb{C}^\times$ be the centralizer of (s, r_0) in $G \times \mathbb{C}^\times$. The group $G(s)$ is connected (in fact, a Levi subgroup of G). Denote

$$M(n) = C_{G \times \mathbb{C}^\times}(n) = \{(x, \lambda) : \text{Ad}(x)n = \lambda^2n\},$$

and let $M(n)^\circ$ be its identity component.

If Q is an algebraic group, X a Q -variety, and \mathcal{L} a Q -equivariant local system on X , let $H_\bullet^Q(X, \mathcal{L})$ denote the equivariant homology groups as in [Lus88, §1]. In particular, if $Q = \{1\}$, then

$$(3.7.3) \quad H_j(X, \mathcal{L}) := H_j^{\{1\}}(X, \mathcal{L}) = H_c^{2 \dim X - j}(X, \mathcal{L}^*)^*.$$

The construction of standard modules using equivariant homology $H_\bullet^{M(n)^\circ}$ is in §8 in *loc. cit.*, see also [Lus95b, pp. 10.7, 10.12]). It is shown that for $n \in \mathfrak{g}_{2r_0}$, the space

$$Y_{\mathbf{H}}(s, n) = \mathbb{C}_{(s, r_0)} \otimes_{H_\bullet^{M(n)^\circ}} H_\bullet^{M(n)^\circ}(\mathcal{P}_n, \dot{\mathcal{E}}) \cong H_\bullet^{\{1\}}(\mathcal{P}_n^s, \dot{\mathcal{E}})$$

carries an action of $A(s, n) \times \mathbf{H}$. In particular, the groups $H_\bullet(\mathcal{P}_n, \dot{\mathcal{E}})$ carry an $A(n) \times W$ -action which is compatible via the identification (3.7.3), and with the appropriate normalization, with the

action defined in [Lus84b]. We normalize the W -action so that $H_0(\mathcal{P}_n, \dot{\mathcal{E}}) = \text{triv}$ when n is the *smallest* nilpotent orbit that occurs in this setting.

Let $\widehat{A(n)}_{\mathcal{E}}$ denote the irreducible representations that occur in the action of $A(n)$ on $H^{\{1\}}(\mathcal{P}_n, \dot{\mathcal{E}})$. We have a natural injection $A(s, n) \rightarrow A(n)$ and let $\widehat{A(s, n)}_{\mathcal{E}}$ denote the irreducible $A(s, n)$ -representations that occur in the restrictions of $\widehat{A(n)}_{\mathcal{E}}$. For every $\rho \in \widehat{A(s, n)}$, define the geometric *standard module*

$$(3.7.4) \quad Y_{\mathbf{H}}(s, n, \rho) = \text{Hom}_{A(s, n)}[\rho, Y_{\mathbf{H}}(s, n)].$$

By [Lus88, Proposition 8.10], $Y_{\mathbf{H}}(s, n, \rho) \neq 0$ if and only if $\rho \in \widehat{A(s, n)}_{\mathcal{E}}$. One can phrase the classification as follows.

Theorem 3.7.2 ([Lus95b, §8.10, 8.14, 8.17], [Lus02b, Theorem 1.15]). *The standard module $Y_{\mathbf{H}}(s, n, \rho)$, $s \in \mathfrak{g}$ semisimple, $n \in \mathfrak{g}_{2r_0}$, $\rho \in \widehat{A(s, n)}_{\mathcal{E}}$ has a unique irreducible quotient $X_{\mathbf{H}}(s, n, \rho)$. This induces a natural one-to-one correspondence*

$$(3.7.5) \quad \text{Irr}_{r_0} \mathbf{H}(L, \mathbb{O}, \mathcal{E}) \leftrightarrow \{(s, n, \rho) : [s, e] = 2r_0e, s \in \mathfrak{g} \text{ semisimple}, n \in \mathcal{N}, \widehat{A(s, n)}_{\mathcal{E}}\}/G.$$

We are interested in the W -structure of standard modules.

Lemma 3.7.3 ([Lus88], 10.13). *There is an isomorphism of W -representations*

$$(3.7.6) \quad Y_{\mathbf{H}}(s, n, \rho) = \text{Hom}_{A(s, n)}[\rho, H_{\bullet}(\mathcal{P}_n^s, \dot{\mathcal{E}})] = \text{Hom}_{A(s, n)}[\rho, H_{\bullet}(\mathcal{P}_n, \dot{\mathcal{E}})].$$

Denote

$$(3.7.7) \quad \mu(n, \tilde{\rho}) := \text{Hom}_{A(n)}[\tilde{\rho}, H_0(\mathcal{P}_n, \dot{\mathcal{E}})], \quad \tilde{\rho} \in \widehat{A(n)}_{\mathcal{E}},$$

an irreducible W -representation. The correspondence $\widehat{A(n)}_{\mathcal{E}} \rightarrow \widehat{W}$, $(n, \tilde{\rho}) \rightarrow \mu(n, \tilde{\rho})$ is a bijection.

The results from [Lus84b, §6.5], [Lus86, §24.4], see also the summary in [BC13, Theorem 3.4.1], imply the following W -character formula:

$$(3.7.8) \quad \text{Hom}_{A(n)}[\tilde{\rho}, H_{\bullet}(\mathcal{P}_n, \dot{\mathcal{E}})] = \mu(n, \tilde{\rho}) + \sum_{(n', \tilde{\rho}')} m_{(n, \tilde{\rho}), (n', \tilde{\rho}')} \mu(n', \tilde{\rho}'),$$

where $m_{(n, \tilde{\rho}), (n', \tilde{\rho}')} \in \mathbb{Z}_{\geq 0}$, $\tilde{\rho}' \in \widehat{A(n')}_{\mathcal{E}}$, and n' ranges over a set of representatives of G -orbits in \mathcal{N} such that $n \in \overline{G \cdot n'} \setminus G \cdot n'$. Combining this with Lemma 3.7.3, we arrive at a character formula for the standard modules.

Proposition 3.7.4. *Let (s, n, ρ) be as in Theorem 3.7.2. As W -representations,*

$$Y_{\mathbf{H}}(s, n, \rho) = \sum_{\tilde{\rho} \in \widehat{A(n)}_{\mathcal{E}}} m(\rho, \tilde{\rho}) \mu(n, \tilde{\rho}) + \sum_{(n', \tilde{\rho}')} m_{(n, \tilde{\rho}), (n', \tilde{\rho}')} \mu(n', \tilde{\rho}'),$$

where $m(\rho, \tilde{\rho}) = \dim \text{Hom}_{A(s, n)}[\rho, \tilde{\rho}|_{A(s, n)}]$, $m_{(n, \tilde{\rho}), (n', \tilde{\rho}')} \in \mathbb{Z}_{\geq 0}$, $\tilde{\rho}' \in \widehat{A(n')}_{\mathcal{E}}$, and n' ranges over a set of representatives of G -orbits in \mathcal{N} such that $n \in \overline{G \cdot n'} \setminus G \cdot n'$.

Finally, we record the classification of tempered modules with real central character, see [BC13, §3.5] for a summary of Lusztig's results [Lus02b]. If $n \in \mathcal{N}_G$, let $\{\bar{n}, h, n\}$ be a Lie triple for n . Then $A(h, n) = A(n)$, so we may identify $\widehat{A(n)}_{\mathcal{E}} = \widehat{A(h, n)}_{\mathcal{E}}$.

Theorem 3.7.5. *The map*

$$(3.7.9) \quad \{(n, \rho) : e \in \mathcal{N}, \rho \in \widehat{A(n)}_{\mathcal{E}}\}/G \longleftrightarrow X_{\mathbf{H}}(r_0h, n, \rho) = Y_{\mathbf{H}}(r_0h, n, \rho),$$

is a one-to-one correspondence onto the set of isomorphism classes of tempered irreducible $\mathbf{H}(L, \mathbb{O}, \mathcal{E})$ -modules with real central character, on which r acts by r_0 .

4. LUSZTIG'S ARITHMETIC-GEOMETRIC CORRESPONDENCE

We assume in this section that \mathbf{G} is adjoint simple.

4.1. Representations with unipotent cuspidal support. We briefly recall the classification of irreducible representations with unipotent cuspidal support defined in [Lus95a, §1.6, §1.21]. With the notation as in section 2.2, let (π, V) be a $\mathbf{G}^\omega(\mathbf{k})$ -representation. We say V has *unipotent cuspidal support* if there exists a parahoric subgroup $\mathbf{P}_c(\mathfrak{o}) \subset \mathbf{G}^\omega(\mathbf{k})$ and an irreducible unipotent cuspidal representation $(\sigma_{\mathbf{E}}, \mathbf{E})$ of $\mathbf{L}_c(\mathbb{F}_q)$ such that the \mathbf{E} -isotypic component of $V|_{\mathbf{P}_c(\mathfrak{o})}$ generates V . (By abuse of notation, we denote by \mathbf{E} also the pullback of \mathbf{E} to $\mathbf{P}_c(\mathfrak{o})$). Let $\mathcal{C}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E})$ denote the corresponding full subcategory of smooth $\mathbf{G}^\omega(\mathbf{k})$ -representations. The assignment

$$m_{\mathbf{E}} : V \mapsto \text{Hom}_{\mathbf{P}_c(\mathfrak{o})}(\mathbf{E}, V)$$

defines an equivalence of categories between $\mathcal{C}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E})$ and the category of modules for the convolution algebra

$$\mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E}) = \{f : \mathbf{G}^\omega(\mathbf{k}) \rightarrow \text{End}_{\mathbb{C}}(\mathbf{E}) \mid f(p_1 g p_2) = \sigma_{\mathbf{E}}(p_1) \circ f(g) \circ \sigma_{\mathbf{E}}(p_2), g \in \mathbf{G}^\omega(\mathbf{k}), p_1, p_2 \in \mathbf{P}_c(\mathfrak{o})\}.$$

Let $G^\vee = \mathbf{G}^\vee(\mathbb{C})$ and let $T^\vee = X^*(\mathbf{T}, \bar{\mathbf{k}}) \otimes_{\mathbb{Z}} \mathbb{C}^\times$. The polar decomposition of T^\vee is $T^\vee = T_c^\vee T_r^\vee$, where $T_c^\vee = X^*(\mathbf{T}, \bar{\mathbf{k}}) \otimes_{\mathbb{Z}} S^1$ and $T_r^\vee = X^*(\mathbf{T}, \bar{\mathbf{k}}) \otimes_{\mathbb{Z}} \mathbb{R}_{>0}$.

Write $\Phi(G^\vee)$ for the set of G^\vee -orbits (under conjugation) of triples (s, n, ρ) where

- $s \in G^\vee$ is semisimple,
- $n \in \mathfrak{g}^\vee$ such that $\text{Ad}(s)n = qn$,
- $\rho \in \text{Irr}(A_{G^\vee}(s, n))$.

Without loss of generality, we may assume that $s \in T^\vee$. Note that $n \in \mathfrak{g}^\vee$ is necessarily nilpotent. The group $G^\vee(s)$ acts with finitely many orbits on the q -eigenspace of $\text{Ad}(s)$

$$\mathfrak{g}_q^\vee = \{x \in \mathfrak{g}^\vee \mid \text{Ad}(s)x = qx\}.$$

In particular, there is a unique open $G^\vee(s)$ -orbit in \mathfrak{g}_q^\vee .

Fix an $\mathfrak{sl}(2)$ -triple $\{n^-, h^\vee, n\} \subset \mathfrak{g}^\vee$ with $h^\vee \in \mathfrak{t}_{\mathbb{R}}^\vee$ and set

$$s_0 := sq^{-\frac{h^\vee}{2}}.$$

Then $\text{Ad}(s_0)n = n$.

A parameter $(s, n, \rho) \in \Phi(G^\vee)$ is called *discrete* if $G^\vee(s, n)$ does not contain a nontrivial torus. A discrete parameter is called *cuspidal* if (n, ρ) is a cuspidal pair in $G^\vee(s)$ in the sense of section 3.7. Let $\Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathbf{k}))$ denote the set of equivalence classes of irreducible $\mathbf{G}^\omega(\mathbf{k})$ -representations with unipotent cuspidal support and let

$$\Pi^{\text{Lus}}(\mathbf{G}) = \bigsqcup_{\omega \in \Omega_{\text{ad}}} \Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathbf{k})).$$

(In this subsection, $\mathbf{G}^1(\mathbf{k})$ is the split form.) The following theorem is a combination of several results, namely [KL87, Theorems 7.12, 8.2, 8.3] for Iwahori-spherical representations, and [Lus95a, Corollary 6.5] and [Lus02a, Theorem 10.5] for representations with unipotent cuspidal support. See [AMS21, §2.3] for a discussion of the compatibility between these classifications.

Theorem 4.1.1 (Deligne-Langlands-Lusztig correspondence). *There is a bijection*

$$\Phi(G^\vee) \xrightarrow{\sim} \Pi^{\text{Lus}}(\mathbf{G}), \quad (s, n, \rho) \mapsto X(s, n, \rho),$$

with the following properties:

- (1) $X(s, n, \rho)$ is tempered if and only if $s_0 \in T_c^\vee$ (in particular, $\overline{G^\vee(s)n} = \mathfrak{g}_q^\vee$),
- (2) $X(s, n, \rho)$ is square integrable (modulo center) if and only if it is tempered and (s, n, ρ) is discrete.
- (3) $X(s, n, \rho)$ is supercuspidal if and only if (s, n, ρ) is a cuspidal parameter.

(4) $X(s, n, \rho) \in \Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathbf{k}))$ if and only if $\rho|_{Z(G^\vee)}$ is a multiple of ζ_ω .

For each parameter (s, n, ρ) , there is an associated admissible *standard module* $Y(s, n, \rho)$ with the following properties, see [Lus02a, Section 10], which relies on the graded affine Hecke algebra results in [Lus02b, Theorems 1.15, 1.21, 1.22]):

- (1) $X(s, n, \rho)$ is the unique simple quotient of $Y(s, n, \rho)$;
- (2) If $G^\vee(s)\bar{n} = \mathfrak{g}_q^\vee$, then $X(s, n, \rho) = Y(s, n, \rho)$; in particular, this is the case when $X(s, n, \rho)$ is tempered. More generally:
- (3) There is an identity of characters ([Lus95b, Corollary 10.7])

$$(4.1.1) \quad Y(s, n, \rho) = X(s, n, \rho) + \sum_{(n', \rho')} m((n, \rho), (n', \rho')) X(s, n', \rho'),$$

where $m((n, \rho), (n', \rho')) \in \mathbb{Z}_{\geq 0}$, and the sum is over all $G^\vee(s)$ -orbits (n', ρ') , such that $n \in \overline{G^\vee(s)n'}$.

We denote by $\Pi_s^{\text{Lus}}(\mathbf{G}(\mathbf{k}))$ the set of irreducible $\mathbf{G}(\mathbf{k})$ -representations $X(s, n, \rho)$ for a fixed $s \in T^\vee$. We say that $X(s, n, \rho)$ has *real infinitesimal character* if $s \in T_r^\vee$.

Remark 4.1.2. *When \mathbf{G} is not adjoint, the parameters of the unipotent representations take a similar form (s, n, ρ) , see [Sol18], except ρ is an irreducible representation of the group $A_{G^\vee}^1(s, n)$, rather than $A_{G^\vee}(s, n)$. To define $A_{G^\vee}^1(s, n)$, let $G_{\text{sc}}^\vee \rightarrow G^\vee \rightarrow G_{\text{ad}}^\vee$ be the chain of isogenies, and let $Z_{G^\vee}^1(s)$ be the preimage in G^\vee of the image of $G^\vee(s)$ in G_{ad}^\vee . Then $A_{G^\vee}^1(s, n)$ is the group of components of the centralizer of n in $Z_{G^\vee}^1(s)$.*

If $s \in T_r^\vee$, let \bar{s} be the unique element of the Lie algebra $\mathfrak{t}_r^\vee \subset \mathfrak{g}^\vee$ that exponentiates to s . Then $G^\vee(s) = C_{G^\vee}(\bar{s})$, and $Z_{G^\vee}^1(s) = C_{G_{\text{sc}}^\vee}(\bar{s})$, and therefore $A^1(s, n) = A_{G_{\text{sc}}^\vee}(\bar{s}, n)$. In other words, the classification on unipotent representations with real infinitesimal character is independent of isogeny. We will use this implicitly in Section 6.

4.2. The arithmetic side. We follow [Lus92]. Let $\mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E})$ be a Hecke algebra as in Section 4.1. Let $J \in \mathbf{P}^\omega(I)$ be the subset of the affine simple roots corresponding to $\mathbf{P}_c(\mathfrak{o})$, and let \widetilde{W}_J be the finite subgroup of \widetilde{W} generated by the simple reflections in J . Write $\mathcal{W} = N_{\widetilde{W}}(\widetilde{W}_J)/\widetilde{W}_J$. As a set, \mathcal{W} can be identified with the set of minimal-length $(\widetilde{W}_J, \widetilde{W}_J)$ -double cosets in \widetilde{W} which are contained in $N_{\widetilde{W}}(\widetilde{W}_J)$. Let \mathcal{W}' (resp. $\overline{\Omega}$) denote the subgroup of \mathcal{W} given by the double cosets contained in W^a (resp. Ω).

The element $\omega \in \Omega$ acts on I and preserves J . Let $\mathcal{W}^\omega, \mathcal{W}'^\omega, \overline{\Omega}^\omega$ denote the fixed points. Let $(I - J)/\omega$ denote the orbits of the action of ω on $I - J$. The group $\overline{\Omega}^\omega$ acts on $(I - J)/\omega$ by permutations. Let $\overline{\Omega}_1^\omega$ (resp. $\overline{\Omega}_2^\omega$) denote the kernel (resp. the image) of the resulting homomorphism $\overline{\Omega}^\omega \rightarrow \text{Perm}(I - J/\omega)$, see [Lus92, §1.20].

The affine Coxeter group \mathcal{W}'^ω has generators $s_{\underline{k}}$, where \underline{k} ranges over $(I - J)/\omega$ if this set has cardinality at least two. Otherwise $\mathcal{W}'^\omega = 1$. Let $\mathcal{H}'(I, J, \omega, \mathbf{E})$ be the affine Hecke algebra with parameters $L(\underline{k})$ constructed in [Lus92, §1.18] for the extended affine Weyl group

$$(4.2.1) \quad \mathcal{W}(J, \omega, \mathbf{E}) := \mathcal{W}'^\omega \rtimes \overline{\Omega}_2^\omega.$$

Then

$$(4.2.2) \quad \mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E}) = \mathcal{H}'(I, J, \omega, \mathbf{E}) \otimes \mathcal{C}[\overline{\Omega}_1^\omega].$$

This can be rewritten as $\mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E}) = \bigoplus_{\psi \in \overline{\Omega}_1^\omega} \mathcal{H}'(I, J, \omega, \mathbf{E})$. Each (conjugacy class of) triple (J, \mathbf{E}, ψ) as above is called an *arithmetic diagram* for \mathbf{G}^ω with associated affine Hecke algebra $\mathcal{H}'(I, J, \omega, \mathbf{E})$ (independent of ψ). Let $\overline{\mathfrak{A}}_\omega$ denote the set of arithmetic diagrams (J, \mathbf{E}, ψ) of \mathbf{G}^ω and set $\overline{\mathfrak{A}} = \sqcup_{\omega \in \Omega} \overline{\mathfrak{A}}_\omega$.

There is a natural one-to-one correspondence (coming from an equivalence of categories)

$$(4.2.3) \quad \Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathbf{k})) \longleftrightarrow \bigsqcup_{(J, \mathbf{E}, \psi) \in \bar{\mathfrak{A}}_\omega} \text{Irr } \mathcal{H}'(I, J, \omega, \mathbf{E}).$$

Denote by $\text{Acusp}(V) = (J, \omega, \mathbf{E})$, the supercuspidal support of a simple $\mathbf{G}^\omega(\mathbf{k})$ -representation V .

4.3. The geometric side. Let I^\vee denote the affine Dynkin diagram of the simply-connected simple group G^\vee . If $J^\vee \subsetneq I^\vee$, denote by $G_{J^\vee}^\vee$ the corresponding pseudo-Levi subgroup of G^\vee . Let $\bar{\mathfrak{S}}$ denote the set of triples (*geometric diagrams*) $(J^\vee, \mathcal{C}^\vee, \mathcal{F}^\vee)$, where $J^\vee \subsetneq I^\vee$, \mathcal{C}^\vee is a nilpotent $G_{J^\vee}^\vee$ -orbit on the Lie algebra $\mathfrak{g}_{J^\vee}^\vee$ and \mathcal{E}^\vee is an isomorphism class of $G_{J^\vee}^\vee$ -equivariant cuspidal local systems on \mathcal{C}^\vee . To such a triple, in [Lus92, §5.12], Lusztig associates a generic affine Hecke algebra $\mathcal{H}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) = \mathcal{H}^{\lambda, \lambda^*}(\Psi, z)$, for some root datum Ψ and parameters λ, λ^* as in *loc.cit.*. Denote by $\mathcal{W}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) = \bar{W} \ltimes \mathcal{X}$ the extended affine Weyl group underlying this affine Hecke algebra. The classification of the simple modules of $\mathcal{H}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ goes via the reduction to graded affine Hecke algebras. More precisely, if $\mathcal{T} = \mathcal{T}_c \cdot \mathcal{T}_r = \text{Hom}_{\mathbb{Z}}(\mathcal{X}, \mathbb{C}^\times)$, then fixing a semisimple conjugacy class $\bar{W} \cdot t \subset \mathcal{T}$, $t = t_c t_h$ we have

$$\text{Irr}_{\bar{W}t, z_0} \mathcal{H}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \longleftrightarrow \text{Irr}_{\log t_h, r_0} \mathbf{H}(G^\vee(s_c), G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee),$$

where s_c (also denoted by t_c^ψ in *loc.cit.*) is the element of T_c^\vee associated to t_c in [Lus92, §5.13]. The group $G_{J^\vee}^\vee$ is a Levi subgroup of $Z_{G^\vee}(s_c) = G_{I^\vee - S^\vee}^\vee$, for a nonempty subset $S^\vee \subset I^\vee - J^\vee$. Therefore, the graded affine Hecke algebra in the right hand side is of the type defined in section 3.7, with $G = G_{I^\vee - S^\vee}^\vee$, $L = G_{J^\vee}^\vee$, and $(\mathcal{C}, \mathcal{L}) = (\mathcal{C}^\vee, \mathcal{E}^\vee)$.

For every $\zeta \in \text{Irr} Z(G^\vee)$, let $\bar{\mathfrak{S}}_\zeta$ denote subset of the triples $(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ such that $Z(G^\vee) \subset G_{J^\vee}^\vee$ acts on each stalk of \mathcal{E}^\vee by ζ .

For every $z_0 > 0$, denote by $\mathfrak{S}(z_0)$ the set of G^\vee -conjugacy classes of triples (s, n, ρ) , where $s \in T^\vee$, $n \in \mathfrak{g}^\vee$ nilpotent such that $\text{Ad}(s)n = z_0^2 n$, and $\rho \in \text{Irr } A(s, n)$. Denote by $\mathfrak{S}(z_0)_\zeta$ the subset of $\mathfrak{S}(z_0)$ with the property that $Z(G^\vee)$ acts in ρ by the character ζ .

Via the classification of simple modules for graded Hecke algebras recalled in Theorem 3.7.2, we have (recall s_c is determined by t_c):

$$(4.3.1) \quad \text{Irr}_{\bar{W}t, z_0} \mathcal{H}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \leftrightarrow \{(s, n, \rho) \in \mathfrak{S}(z_0) \mid s = s_c s_h, s_h \in T_h^\vee, \rho \in \widehat{A(s, n)}_{\mathcal{E}^\vee}\}.$$

Then, by combining all such correspondences for all geometric diagrams, [Lus92, Theorem 5.21]] says that there is a natural bijection

$$(4.3.2) \quad \mathfrak{S}(z_0)_\zeta \longleftrightarrow \bigsqcup_{(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \in \bar{\mathfrak{S}}_\zeta} \text{Irr } \mathcal{H}_{z_0}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee).$$

If $(s, n, \rho) \in \mathfrak{S}(z_0)$ parametrizes a module for $\mathcal{H}_{z_0}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$, denote by $\text{Gcusp}(s, n, \rho) = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ and call it the geometric cuspidal support of (s, n, ρ) (or $X(s, n, \rho), Y(s, n, \rho)$).

Remark 4.3.1. *If X has real infinitesimal character and $\text{Gcusp}(X) = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$, then $J^\vee \subseteq I_0^\vee$.*

To complete the correspondence and the proof of Theorem 4.1.1, Lusztig matched the arithmetic and geometric diagrams. More precisely, [Lus92, Theorem 6.3] gives an explicit bijection, for each $\omega \in \Omega$,

$$(4.3.3) \quad \theta : \bar{\mathfrak{S}}_{\zeta_\omega} \longleftrightarrow \bar{\mathfrak{A}}_\omega,$$

such that if $\theta((J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)) = (J, \mathbf{E}, \psi)$, then

$$\mathcal{H}_{\sqrt{q}}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \cong \mathcal{H}'(I, J, \omega, \mathbf{E}).$$

In particular, the underlying (extended) affine Weyl groups are isomorphic

$$\mathcal{W}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \cong \mathcal{W}(J, \omega, \mathbf{E})$$

Denote this isomorphism of Hecke algebras also by θ and let θ^* the dual module correspondence. We also write θ for the bijection between $\overline{\mathfrak{S}}$ and $\overline{\mathfrak{U}}$. In this case write $\theta(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) = (J, \mathbf{E}, \psi, \omega)$.

4.4. Multiplicities. Let $(\pi, V) \in \mathcal{C}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E})$ be an irreducible or standard representation with unipotent cuspidal support $(\mathbf{P}_c(\mathfrak{o}), \mathbf{E})$, and $\mathbf{P}_{c'}(\mathfrak{o}) \supseteq \mathbf{P}_c(\mathfrak{o})$ another parahoric subgroup. Define the convolution algebra

$$\mathcal{H}(\mathbf{P}_{c'}(\mathfrak{o}), \mathbf{E}) = \{f \in \mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E}) \mid \text{supp } f \subseteq \mathbf{P}_{c'}(\mathfrak{o})\}.$$

If τ is irreducible representation of $\mathbf{L}_{c'}(\mathbb{F}_q)$, denote by $\mathbf{m}_{\mathbf{E}}(\tau)$ the representation of $\mathcal{H}(\mathbf{P}_{c'}(\mathfrak{o}), \mathbf{E})$ on $\text{Hom}_{\mathbf{L}_c(\mathbb{F}_q)}(\mathbf{E}, \tau|_{\mathbf{L}_c(\mathbb{F}_q)})$. Then we have

$$(4.4.1) \quad \text{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, V) = \text{Hom}_{\mathcal{H}(\mathbf{P}_{c'}(\mathfrak{o}), \mathbf{E})}(\mathbf{m}_{\mathbf{E}}(\tau), \mathbf{m}_{\mathbf{E}}(V)|_{\mathcal{H}(\mathbf{P}_{c'}(\mathfrak{o}), \mathbf{E})}).$$

Next, let $J \subseteq J' \in \mathbf{P}^\omega(I)$ be the subsets defining $\mathbf{P}_c(\mathfrak{o}), \mathbf{P}_{c'}(\mathfrak{o})$, respectively. Then using (4.2.2), we find

$$(4.4.2) \quad \text{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, V) = \text{Hom}_{\mathcal{H}(J', J, \omega, \mathbf{E})}(\mathbf{m}_{\mathbf{E}}(\tau), \mathbf{m}_{\mathbf{E}}(V)).$$

Here, $\mathbf{m}_{\mathbf{E}}(V)$ is a module an affine Hecke algebras $\mathcal{H}'(I, J, \omega, \mathbf{E})$ (for one ψ), and $\mathcal{H}(J', J, \omega, \mathbf{E})$ is the parahoric (finite) Hecke algebra with support in J' .

Let $(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) = \theta^{-1}(J, \mathbf{E}, \psi)$ be the corresponding geometric diagram in $\tilde{\mathfrak{S}}_{\zeta_\omega}$. Denote by $J'^\vee = \theta^{-1}(J')$ the corresponding subset of I^\vee such that there is an isomorphism of finite Hecke algebras

$$\mathcal{H}_{\sqrt{q}}(G_{J'^\vee}^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \cong \mathcal{H}'(J', J, \omega, \mathbf{E}).$$

Let $\mathcal{W}_{J'^\vee}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \cong \mathcal{W}_{J'}(J, \omega, \mathbf{E})$ denote the finite parahoric subgroups defined by $J'^\vee \supseteq J^\vee$ and $J' \supseteq J$, respectively. It follows that

$$\text{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, V) = \text{Hom}_{\mathcal{H}_{\sqrt{q}}(G_{J'^\vee}^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)}(\theta^*(\mathbf{m}_{\mathbf{E}}(\tau)), \theta^*(\mathbf{m}_{\mathbf{E}}(V))),$$

where $\theta^*(\mathbf{m}_{\mathbf{E}}(V))$ is an $\mathcal{H}_{\sqrt{q}}(G_{J'^\vee}^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ -module. We are now in position to apply the deformation argument from section 3.5. From the first part of Corollary 3.5.3, it follows at once that, when V is irreducible,

$$(4.4.3) \quad \text{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, V) = \text{Hom}_{\mathcal{W}_{J'^\vee}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)}(\theta^*(\mathbf{m}_{\mathbf{E}}(\tau))_{q \rightarrow 1}, \tilde{\sigma}_0(\theta^*(\mathbf{m}_{\mathbf{E}}(V))),$$

where $\tilde{\sigma}_0$ is the deformation functor from (3.5.1). Here we use the notation $q \rightarrow 1$ in place of $z_0 \rightarrow 1$ as in section 3.5. Recall that $\tilde{\sigma}_0(\theta^*(\mathbf{m}_{\mathbf{E}}(V)))$ is a module for the extended affine Weyl group $\mathcal{W}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$.

For standard modules, we can explicitate this formula further using Proposition 3.6.1. Let $Y(s, n, \rho)$ be a standard $\mathbf{G}^\omega(\mathbf{k})$ -module with unipotent cuspidal support. Let (J, \mathbf{E}) be the supercuspidal support of $X(s, n, \rho)$, and $(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) = \text{Gcusp}(s, n, \rho)$. Let $G_{J^\vee - S^\vee}^\vee = Z_{G^\vee}(s_c)$. Let $\mathcal{P}_{s_c, n}$ denote the generalized Springer fiber \mathcal{P}_n defined as in section 3.7 for the pair $(\mathcal{C}^\vee, \mathcal{E}^\vee)$ in $\mathfrak{g}_{J^\vee - S^\vee}^\vee$. Let s_c be the compact part of s and t_c the corresponding element in \mathcal{T}_c for the geometric Hecke algebra $\mathcal{H}(G_{J^\vee}^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$.

Theorem 4.4.1. *Let $\mathbf{P}_c(\mathfrak{o})$ be a parahoric subgroup for $\mathbf{G}^\omega(\mathbf{k})$ corresponding to J , $J \subset J' \in \mathbf{P}^\omega(I)$, and τ an irreducible $\mathbf{L}_{c'}(\mathbb{F}_q)$ -representation. Write $\mathcal{W}^\vee := \mathcal{W}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ and $\mathcal{W}_{J'^\vee}^\vee := \mathcal{W}_{J'^\vee}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ for simplicity, and let $\mathcal{W}^\vee(t_c)$ be the centralizer of t_c in \mathcal{W}^\vee . Then*

$$(4.4.4) \quad \begin{aligned} \text{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, Y(s, n, \rho)) &= \text{Hom}_{\mathcal{W}_{J'^\vee}^\vee}(\theta^*(\mathbf{m}_{\mathbf{E}}(\tau))_{q \rightarrow 1}, \text{Ind}_{\mathcal{W}^\vee(t_c)}^{\mathcal{W}^\vee}(\mathbb{C}_{t_c} \otimes Y_{\mathbf{H}}(s, n, \rho))) \\ &= \text{Hom}_{\mathcal{W}_{J'^\vee}^\vee}(\theta^*(\mathbf{m}_{\mathbf{E}}(\tau))_{q \rightarrow 1}, \text{Ind}_{\mathcal{W}^\vee(t_c)}^{\mathcal{W}^\vee}(\mathbb{C}_{t_c} \otimes H_\bullet(\mathcal{P}_{s_c, n}, \dot{\mathcal{E}}^\vee)^\rho)), \end{aligned}$$

where $Y_{\mathbf{H}}(s, n, \rho)$ is the standard module for the graded affine Hecke algebra $\mathbf{H}(G^\vee(s_c), G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$, and $H_\bullet(\mathcal{P}_{s_c, n}, \dot{\mathcal{E}}^\vee)^\rho$ is a (reducible) generalized Springer representation of the finite Weyl group $N_{G^\vee(s_c)}(G_{J^\vee}^\vee)/G_{J^\vee}^\vee$, as expositied in section 3.7.

Proof. The first equality follows immediately from (4.4.3), Proposition 3.6.1 and the second part of Corollary 3.5.3. The second equality is implied directly by Lemma 3.7.3. \square

In the case of real infinitesimal character in this paper, this result takes the following simpler form.

Corollary 4.4.2. *Retain the notation from Theorem 4.4.1. Suppose $Y(s, n, \rho)$ has real infinitesimal character, i.e. $s_c = 1$. Then*

$$(4.4.5) \quad \begin{aligned} \mathrm{Hom}_{\mathbf{P}_{c'}(\mathfrak{o})}(\tau, Y(s, n, \rho)) &= \mathrm{Hom}_{\mathcal{W}_{j^\vee}^\vee}(\theta^*(m_{\mathbf{E}}(\tau))_{q \rightarrow 1}, Y_{\mathbf{H}}(s, n, \rho)) \\ &= \mathrm{Hom}_{\mathcal{W}_{j^\vee}^\vee}(\theta^*(m_{\mathbf{E}}(\tau))_{q \rightarrow 1}, H_\bullet(\mathcal{P}_n, \dot{\mathcal{E}}^\vee)^\rho), \end{aligned}$$

where $Y_{\mathbf{H}}(s, n, \rho)$ is the standard module for the graded affine Hecke algebra $\mathbf{H}(G^\vee, G_{j^\vee}^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$, $\mathcal{W}_{j^\vee}^\vee = N_{G^\vee}(G_{j^\vee}^\vee)/G_{j^\vee}^\vee$, and $H_\bullet(\mathcal{P}_n, \dot{\mathcal{E}}^\vee)$ are the homology groups carrying a $\mathcal{W}_{j^\vee}^\vee \times A(n)$ -representation that occur in the generalized Springer correspondence for $G_{j^\vee}^\vee \subset G^\vee$ (see 3.7).

5. FAITHFULNESS FOR THE GENERALISED SPRINGER CORRESPONDENCE

5.1. Preliminaries on partitions. In this section, we will recall some of the basic notation for partitions, and some related objects. A *partition of n* is a non-increasing sequence of positive integers $(\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m)$ such that $\lambda_1 + \lambda_2 + \dots + \lambda_m = n$ (occasionally, it will be more convenient to denote a partition as a non-decreasing sequence, but we will clearly indicate when this is the case). Let $\mathcal{P}(n)$ denote the set of partitions of n . We write $\#\lambda := m$ and $|\lambda| := n$. If $x \in \mathbb{Z}_{\geq 0}$, we write $m_\lambda(x)$ for the multiplicity of x in λ and $\mathrm{ht}_\lambda(x)$ for its *height*, i.e. $\mathrm{ht}_\lambda(x) = \sum_{y \geq x} m_\lambda(y)$. A partition is *very even* if all parts are even, occurring with even multiplicity. A partition is *rather odd* if every even part occurs with even multiplicity and every odd part occurs with multiplicity ≤ 1 . We write $\mathcal{P}_{ve}(n)$ (resp. $\mathcal{P}_{ro}(n)$) for the set of very even (resp. rather odd) partitions of n .

There is a partial order on integer sequences (and in particular, on $\mathcal{P}(n)$), defined by the formula

$$(5.1.1) \quad \lambda \leq \mu \iff \sum_{i \leq j} \lambda_i \leq \sum_{i \leq j} \mu_i.$$

Given an arbitrary partition $\lambda = (\lambda_1, \dots, \lambda_m)$, we define two new partitions (of $|\lambda| + 1$ and $|\lambda| - 1$)

$$\lambda^+ := (\lambda_1 + 1, \lambda_2, \dots, \lambda_m), \quad \lambda^- := (\lambda_1, \lambda_2, \dots, \lambda_m - 1)$$

and we write λ^t for the transpose of λ .

Given $\lambda \in \mathcal{P}(n)$ and $\mu \in \mathcal{P}(m)$, we define $\lambda \cup \mu \in \mathcal{P}(m+n)$ by ‘adding multiplicities’

$$m_{\lambda \cup \mu}(x) = m_\lambda(x) + m_\mu(x), \quad \forall x \in \mathbb{Z}_{\geq 0}.$$

We will need the following elementary fact.

Lemma 5.1.1. *Let $\alpha_1, \alpha_2, \beta_1, \beta_2$ be partitions such that $\alpha_1 \leq \alpha_2$ and $\beta_1 \leq \beta_2$. Then $\alpha_1 \cup \beta_1 \leq \alpha_2 \cup \beta_2$.*

Proof. We begin with the following observation. If λ is an integer sequence and λ' is the partition obtained by permuting the elements of λ , then $\lambda \leq \lambda'$. Indeed, λ' is obtained from λ through the following algorithm: if $\lambda_i > \lambda_j$ for $i > j$, swap λ_i and λ_j , and swaps of this type are strictly increasing with respect to \leq .

Add 0’s to α_i, β_i so that $\#\alpha_1 + \#\beta_1 = \#\alpha_2 + \#\beta_2$, and choose a function $f : \{1, 2, \dots, \#\alpha_1 + \#\beta_1\} \rightarrow \{\alpha, \beta\}$ such that $\alpha_1 = ((\alpha_1 \cup \beta_1)_i \mid f(i) = \alpha)$ and $\beta_1 = ((\alpha_1 \cup \beta_1)_i \mid f(i) = \beta)$. Define a sequence λ by replacing each element of $\alpha_1 \cup \beta_2$ with the corresponding element in α_2 or β_2 according to f , i.e.

$$\lambda_i = \begin{cases} (\alpha_2)_{\#\{j \leq i \mid f(j) = \alpha\}} & \text{if } f(i) = \alpha \\ (\beta_2)_{\#\{j \leq i \mid f(j) = \beta\}} & \text{if } f(i) = \beta \end{cases}$$

Since $\alpha_1 \leq \alpha_2$ and $\beta_1 \leq \beta_2$, we have $\alpha_1 \cup \beta_1 \leq \lambda$. On the other hand, $\alpha_2 \cup \beta_2$ is the partition obtained by permuting the entries of λ . So by the discussion of the previous paragraph, $\lambda \leq \alpha_2 \cup \beta_2$. So $\alpha_1 \cup \beta_1 \leq \alpha_2 \cup \beta_2$, as asserted. \square

If $r \geq \lambda_1$, we write $(r, \lambda) = \lambda \cup \{r\}$. We write $\alpha = \lambda \cup_{\geq} \mu$ if $\alpha = \lambda \cup \mu$ and the smallest part of λ is at least as large as the largest part of μ . We write $\mu \subseteq \lambda$ if $m_\mu(x) \leq m_\lambda(x)$ for every $x \in \mathbb{Z}_{\geq 0}$. In this case, there is a unique subpartition $\lambda \setminus \mu \subseteq \lambda$ such that $\lambda = \mu \cup (\lambda \setminus \mu)$.

A *bipartition* of n is a pair of partitions (λ, μ) such that $|\lambda| + |\mu| = n$. Write $\mathcal{B}(n)$ for the set of bipartitions of n . \mathbb{Z}_2 acts by flipping on $\mathcal{B}(n)$, and we write $\mathcal{D}(n)$ for the set of \mathbb{Z}_2 -orbits on $\mathcal{B}(n)$.

5.2. Preliminaries on nilpotent orbits in classical types. For $X \in \{B, C, D\}$ and n a positive integer, let $\mathfrak{g}_X(n)$ denote the simple complex Lie algebra of type X and rank n . Define the sets

$$\begin{aligned} \mathcal{P}_B(2n+1) &= \{\lambda \in \mathcal{P}(2n+1) \mid x \text{ even} \implies m_\lambda(x) \text{ even}\} \\ \mathcal{P}_C(2n) &= \{\lambda \in \mathcal{P}(2n) \mid x \text{ odd} \implies m_\lambda(x) \text{ odd (or 0)}\} \\ \mathcal{P}_D(2n) &= \{\lambda \in \mathcal{P}(2n) \mid x \text{ even} \implies m_\lambda(x) \text{ even}\} \end{aligned}$$

The elements of $\mathcal{P}_B(2n+1)$ (resp. $\mathcal{P}_C(2n)$, resp. $\mathcal{P}_D(2n)$) are called *B-partitions* (resp. *C-partitions*, resp. *D-partitions*). There are well-known bijections

$$\mathcal{N}_o^{\mathfrak{g}_B} \xrightarrow{\sim} \mathcal{P}_B(2n+1), \quad \mathcal{N}_o^{\mathfrak{g}_C} \xrightarrow{\sim} \mathcal{P}_C(2n)$$

For $X = D$, there is a surjection

$$\mathcal{N}_o^{\mathfrak{g}_D} \twoheadrightarrow \mathcal{P}_D(2n)$$

which is a bijection over partitions in $\mathcal{P}_D(2n)$, except the very even partitions, over which is two-to-one, see [CM93, Section 5.1].

For $X \in \{B, C, D\}$, the X -collapse of λ is the unique largest X -partition λ_X such that $\lambda_X \leq \lambda$. A formula for λ_X is provided in [CM93, Lem 6.3.8].

If λ is an X -partition, define the *dual* partition $d_{BV}(\lambda)$ of λ as follows:

- If $\lambda \in \mathcal{P}_B(2n+1)$, then $d_{BV}(\lambda) = ((\lambda^t)^-)_C \in \mathcal{P}_C(2n)$.
- If $\lambda \in \mathcal{P}_C(2n)$, then $d_{BV}(\lambda) = ((\lambda^t)^+)_B \in \mathcal{P}_B(2n+1)$.
- If $\lambda \in \mathcal{P}_D(2n)$, then $d_{BV}(\lambda) = (\lambda^t)_D \in \mathcal{P}_D(2n)$.

5.3. Preliminaries on symbols. In this section, we will recall some of the basic notation for symbols, following [GM20, Section 4.4].

For a sequence $X = (x_1, x_2, \dots, x_n)$ and integer $a \in \mathbb{Z}$ write $X + a$ for the sequence $(x_1 + a, x_2 + a, \dots, x_n + a)$. We write z for the sequence $(0, 1, \dots, n-1)$ (n will always be clear from the context and therefore omitted from the notation).

A β -set is a strictly increasing sequence $x = (x_1 < x_2 < \dots < x_m)$. The partition $\mathcal{P}(x)$ associated to x is $(x_1 - 0, x_2 - 1, \dots, x_m - (m-1))$. We say that x is associated to a partition λ if $\mathcal{P}(x) = \lambda$. For example, the β -sets associated to $\lambda = (1, 2, 4)$ are those of the form $(1, 3, 6)$ or $(0, 1, 2, \dots, k-1, k+1, k+3, k+6)$ for some k .

For integers $a, b \geq 0$ and d , let $\tilde{\mathcal{X}}_d^{a,b}$ denote the set of pairs (X, Y) of finite sequences $X = (x_1, \dots, x_r)$ and $Y = (y_1, \dots, y_s)$ of nonnegative integers, such that

$$\begin{aligned} x_j &\geq x_{j-1} + a + b & 2 \leq j \leq r, \\ y_j &\geq y_{j-1} + a + b & 2 \leq j \leq s, \\ y_1 &\geq b, \\ r - s &= d. \end{aligned}$$

The elements of $\tilde{\mathcal{X}}_d^{a,b}$ are called *symbols*, and are often denoted as follows

$$(X, Y) = \begin{pmatrix} x_1 & x_2 & \dots & x_r \\ & y_1 & \dots & y_s \end{pmatrix}$$

The *defect* of $S = (X, Y) \in \tilde{\mathcal{X}}_d^{a,b}$ is the integer $d(S) = d = |X| - |Y|$. The *content* is the integer $|S| := |X| + |Y|$. Define

$$r(S) := \sum_{x \in X \cup Y} x - (a+b) \lfloor \frac{(|S|-1)^2}{4} \rfloor - b \lfloor \frac{|S|}{2} \rfloor$$

$$m(S) := \sum_{x \in X \cup Y} x - (a+b) \left(\binom{|X|}{2} + \binom{|Y|}{2} \right) - b|Y|.$$

where the sums run over the multiset $X \cup Y$. We call $r(S)$ (resp. $m(S)$) the *rank* (resp. *relative rank*) of S . It is easy to check that both $r(S)$ and $m(S)$ are nonnegative integers. Define

$$\Delta_d^{a,b} := (a+b) \left\lfloor \frac{d^2}{4} \right\rfloor - b \left\lfloor \frac{d}{2} \right\rfloor = \begin{cases} (a+b) \left(\frac{d}{2}\right)^2 - b \left(\frac{d}{2}\right) & d \text{ even} \\ (a+b) \left(\frac{d-1}{2}\right) \left(\frac{d-1}{2} + 1\right) - b \left(\frac{d-1}{2}\right) & d \text{ odd.} \end{cases}$$

Then for $S \in \tilde{\mathcal{X}}_d^{a,b}$

$$m(S) = r(S) - \Delta_d^{a,b}.$$

The *shift* of a symbol $S = (X, Y) \in \tilde{\mathcal{X}}_d^{a,b}$ is the symbol

$$S' = (\{0\} \cup (X + a + b), \{b\} \cup (Y + a + b)) \in \tilde{\mathcal{X}}_d^{a,b}$$

Note that both defect and rank are invariant under shift. Write $\mathcal{X}_d^{a,b}$ for the set of equivalence classes under the equivalence relation generated by the shift operation. Write $\mathcal{X}_{d,n}^{a,b}$ for the set of classes in $\mathcal{X}_d^{a,b}$ of rank n and write

$$\mathcal{X}_{\bullet,n}^{a,b} = \bigsqcup_{d \in \mathbb{Z}} \mathcal{X}_{d,n}^{a,b}, \quad \mathcal{X}_{\text{even},n}^{a,b} = \bigsqcup_{d \in \mathbb{Z}, d \text{ even}} \mathcal{X}_{d,n}^{a,b}, \quad \mathcal{X}_{\text{odd},n}^{a,b} = \bigsqcup_{d \in \mathbb{Z}, d \text{ odd}} \mathcal{X}_{d,n}^{a,b}.$$

There is an equivalence relation \sim on $\mathcal{X}_{\bullet,n}^{a,b}$ defined as follows: $S_1 \sim S_2$ if and only if there are representatives (X_1, Y_1) and (X_2, Y_2) of S_1 and S_2 such that the multisets $X_1 \cup Y_1$ and $X_2 \cup Y_2$ coincide.

There is a bijection $\mathcal{B} : \mathcal{X}_{d,n}^{a,b} \rightarrow \mathcal{B}(n - \Delta_d^{a,b})$ given by

$$(5.3.1) \quad (X, Y) \mapsto (X - (0, (a+b), 2(a+b), \dots), Y - (b, b + (a+b), b + 2(a+b), \dots)).$$

For $a = b = 0$ this specialises to a bijection $\mathcal{B} : \mathcal{X}_{d,n}^{0,0} \rightarrow \mathcal{B}(n)$.

If $b = 0$, then \mathbb{Z}_2 acts on $\mathcal{X}_d^{a,0} \cup \mathcal{X}_{-d}^{a,0}$ by interchanging the rows. Let \mathcal{Y}_d^a denote the set of \mathbb{Z}_2 -orbits on $\mathcal{X}_d^{a,0} \cup \mathcal{X}_{-d}^{a,0}$. Note that the rank of a symbol is invariant under flipping. We call $S \in \mathcal{Y}_d^a$ degenerate if its \mathbb{Z}_2 -orbit is a singleton. We write $\mathcal{Y}_{d,n}^a$ for the set of orbits of rank n and

$$\mathcal{Y}_{\bullet,n}^a = \bigsqcup_{d \geq 0} \mathcal{Y}_{d,n}^a, \quad \mathcal{Y}_{\text{even},n}^a = \bigsqcup_{d \geq 0, d \text{ even}} \mathcal{Y}_{d,n}^a, \quad \mathcal{Y}_{\text{odd},n}^a = \bigsqcup_{d \geq 0, d \text{ odd}} \mathcal{Y}_{d,n}^a.$$

It is clear that \sim induces an equivalence relation on $\mathcal{Y}_{\bullet,n}^a$.

Given a symbol $S = (X, Y) \in \mathcal{Y}_{d,n}^a$ define $\underline{S} \in \mathcal{X}_{|d|,n}^{a,0}$ as follows: if $d \neq 0$

$$(5.3.2) \quad \underline{S} := \begin{cases} (X, Y) & \text{if } |X| > |Y| \\ (Y, X) & \text{if } |Y| > |X| \end{cases}$$

and if $d = 0$

$$(5.3.3) \quad \underline{S} := \begin{cases} (X, Y) & \text{if } \sum_{x \in X} x \geq \sum_{y \in Y} y \\ (Y, X) & \text{if } \sum_{x \in X} x < \sum_{y \in Y} y \end{cases}$$

There is a bijection

$$(5.3.4) \quad \mathcal{D} : \mathcal{Y}_{d,n}^a \rightarrow \begin{cases} \mathcal{D}(n) & d = 0 \\ \mathcal{B}_{n-\Delta_d^{a,0}} & d \neq 0. \end{cases}$$

given by

$$(5.3.5) \quad S \mapsto (X - (0, (a+b), 2(a+b), \dots), Y - (0, (a+b), 2(a+b), \dots))$$

where $(X, Y) = \underline{S}$.

For $d \in \mathbb{Z}$ there is a bijection $\mathcal{X}_{d,n}^{1,0} \rightarrow \mathcal{X}_{d+1,n}^{0,1}$ given by $S = (X, Y) \mapsto S^!$ where

$$S^! = (\{0\} \cup (X+1), Y+1).$$

We have that $\mathcal{B}(S) = \mathcal{B}(S^!)$.

A symbol $S = (X, Y) \in \mathcal{X}_{d,n}^{a,b}$ is called *special* if $d \in \{0, 1\}$ and $x_1 \leq y_1 \leq x_2 \leq \dots$. Note that this definition is independent of the choice of representative of S . Every \sim -equivalence class in $\mathcal{X}^{a,b}(n)$ contains a unique special element. A symbol $S \in \mathcal{Y}_n^{a,b}$ is called *special* if the corresponding \mathbb{Z}_2 -orbit in $\mathcal{X}_n^{a,b}$ contains a special symbol. Given a symbol S in $\mathcal{X}_n^{a,b}$ or \mathcal{Y}_n^a , we write S^{sp} for the (unique) special symbol in the \sim -equivalence class of S .

For $n \geq 0$, write W_n (resp. W'_n) for the Weyl group of type B_n (resp. D_n). Recall that $\text{Irr}(W_n) \leftrightarrow \mathcal{B}_n$ and $\text{Irr}(W'_n) \leftrightarrow \mathcal{D}_n$ except to each degenerate orbit in \mathcal{D}_n there are two irreducible representations of W'_n .

To make our terminology consistent with that of [CMO21] we say that S is an *a-symbol* if $a+b = 1$ and an *s-symbol* if $a+b = 2$.

We conclude this subsection with two elementary lemmas. A *linear presentation* of a symbol $S = (X, Y) \in \tilde{\mathcal{X}}_{\bullet}^{a,0}$ is a pair (Z, ϵ) consisting of sequences $Z \in \mathbb{Z}_{\geq 0}^{|S|}$ and $\epsilon \in \{0, 1\}^{|S|}$ such that $X = (Z_i \mid \epsilon_i = 0)$ and $Y = (Z_i \mid \epsilon_i = 1)$. We say that (Z, ϵ) is *canonical* if Z is non-decreasing and $\epsilon_i = 0$ whenever $Z_i = Z_{i+1}$. Note that if $a > 0$, then S has a unique canonical linear presentation, which we will denote by $(Z^*(S), \epsilon^*(S))$ or simply (Z^*, ϵ^*) when there is no risk for confusion.

Lemma 5.3.1. *Suppose $S = (X, Y) \in \tilde{\mathcal{X}}_{\bullet}^{1,0}$. Let (Z^*, ϵ^*) be the canonical linear presentation of S and let (Z, ϵ) be any linear presentation of S . Then*

$$Z^* + \epsilon^* \geq Z + \epsilon$$

Proof. Note that $Z^* + \epsilon^* = Z + \epsilon$ as multisets. So by the argument in the proof of Lemma 5.1.1, it suffices to show that the sequence $Z^* + \epsilon^*$ is non-decreasing. If $i < j$, then either

- (1) $Z_i^* < Z_j^*$, or
- (2) $Z_i^* = Z_j^*$, $\epsilon_i^* = 0$, and $\epsilon_j^* = 1$.

In the first case, we have $(Z^* + \epsilon^*)_i \leq Z_i^* + 1 \leq Z_j^* \leq (Z^* + \epsilon^*)_j$. In the second case, $(Z^* + \epsilon^*)_i = Z_i^* < Z_i^* + 1 = (Z^* + \epsilon^*)_j$. Hence $Z^* + \epsilon^*$ is non-decreasing. \square

For a symbol $S = (X, Y)$ let $\tilde{S} = (2X, 2Y + 1)^{sp}$.

Lemma 5.3.2. *Let $S = (X, Y)$ be an *a-symbol* and let (Z, ϵ) be the canonical linear presentation for S . Then $Z^*(\tilde{S}) = 2Z + \epsilon$.*

Proof. Clearly \tilde{S} and $2Z + \epsilon$ are the same as multisets. Thus it suffices to prove that $2Z + \epsilon$ is non-decreasing. Let $i < j$. Then either

- (1) $Z_i < Z_j$, or
- (2) $Z_i = Z_j$, $\epsilon_i = 0$, and $\epsilon_j = 1$.

In the first case $2Z_i + 2 \leq 2Z_j$ and so $2Z_i + \epsilon_i < 2Z_i + 2 \leq 2Z_j \leq 2Z_j + \epsilon_j$. In the second case $2Z_i + \epsilon_i = 2Z_i < 2Z_i + 1 = 2Z_j + \epsilon_j$. Hence $2Z + \epsilon$ is non-decreasing (in fact increasing). \square

5.4. Combinatorics for Spin groups. In this section, we will recall, and reinterpret, some of the combinatorics from [LS85, Section 4] required to formulate the generalized Springer correspondence for $Spin(N)$.

If $m \in \mathbb{Z}$, define $d(m) \in \{0, \pm 1\}$ by

$$d(m) = \begin{cases} 0 & \text{if } m \text{ is even} \\ 1 & \text{if } m \equiv 1 \pmod{4} \\ -1 & \text{if } m \equiv 3 \pmod{4} \end{cases}$$

If $\lambda \in \mathcal{P}(n)$, define

$$(5.4.1) \quad d(\lambda) = \sum_i d(\lambda_i).$$

Now suppose $\lambda = (\lambda_1 \leq \dots \leq \lambda_m) \in \mathcal{P}_{ro}(N)$. Following [LS85, Section 4.11], we will define a bipartition $\rho(\lambda)$. If $m = 0$, we put $\rho(\lambda) = ((0), (0))$. Suppose $m \neq 0$. Let

$$(5.4.2) \quad \mu = \begin{cases} (\lambda_1 \leq \dots \leq \lambda_{m-1}) & \text{if } \lambda_m \text{ is odd} \\ (\lambda_1 \leq \dots \leq \lambda_{m-2}) & \text{if } \lambda_m \text{ is even} \end{cases}$$

We will now define $\rho(\lambda) = (\alpha, \beta)$ in terms of $\rho(\mu) = (\gamma, \delta)$. There are eight cases:

- (a) If $d(\lambda_m) = 0$, set $r = \lfloor \frac{1}{4}(\lambda_m + 2) \rfloor - d(\mu)$, $d = \lfloor \frac{1}{4}\lambda_m \rfloor + d(\mu)$.
 - (a₁) If $d(\mu) < 0$, then $\alpha = (\gamma, r)$ and $\beta = (\delta, s)$.
 - (a₂) If $d(\mu) \leq 0$, then $\alpha = (\gamma, s)$ and $\beta = (\gamma, r)$.
- (b) If $d(\lambda_m) = 1$, set $r = \frac{1}{4}(\lambda_m - 1) - d(\mu)$.
 - (b₁) If $d(\mu) > 0$, then $\alpha = (\gamma, r)$ and $\beta = \delta$.
 - (b₂) If $d(\mu) = 0$, then $\alpha = (\delta, r)$ and $\beta = \gamma$.
 - (b₃) If $d(\mu) < 0$, then $\alpha = \gamma$ and $\beta = (\delta, r)$.
- (c) If $d(\lambda_m) = -1$, set $r = \frac{1}{4}(\lambda_m - 3) + d(\mu)$.
 - (c₁) If $d(\mu) > 1$, then $\alpha = \gamma$ and $\beta = (\delta, r)$.
 - (c₂) If $d(\mu) = 1$, then $\alpha = (\delta, r)$ and $\beta = \gamma$.
 - (c₃) If $d(\mu) < 1$, then $\alpha = (\gamma, r)$ and $\beta = \delta$.

It is shown in [LS85, Lemma 4.12] that $\rho(\lambda) \in \mathcal{B}(\frac{1}{4}(N - d(\lambda)(2d(\lambda) - 1)))$.

Below, it will be convenient to have a non-recursive formula for $\rho(\lambda)$. This will require some additional notation. For $n \in \mathbb{Z}$, let $r(n) \in \{0, 1, 2, 3\}$ denote the residue of $n \pmod{4}$ and let $q(n) \in \mathbb{Z}$ be such that $n = 4q(n) + r(n)$. For $\lambda \in \mathcal{P}_{ro}(n)$ and $x \in \{0, 1, 2, 3\}$, define the partition

$$\lambda^x = (q(\lambda_i) \mid r(\lambda_i) = x)$$

For any sequence $\mu = (\mu_1, \dots, \mu_m)$ of the same length as λ define

$$\mu^x = (\mu_i \mid r(\lambda_i) = x)$$

Also let

$$D(\lambda) = (\sum_{j < i} d(\lambda_j))_i$$

Since every even part of λ has even multiplicity, every part of λ^x , for $x \in \{0, 2\}$, has even multiplicity. So, for $x \in \{0, 2\}$, there is a partition $\underline{\lambda}^x$ such that

$$\lambda^x = \underline{\lambda}^x \cup \underline{\lambda}^x$$

Similarly, for $x \in \{0, 2\}$, there is a partition $\underline{D}(\lambda)^x$ such that

$$D(\lambda)^x = \underline{D}(\lambda)^x \cup \underline{D}(\lambda)^x$$

Proposition 5.4.1. *Let $\lambda \in \mathcal{P}_{ro}(n)$ be rather odd and put $D = D(\lambda)$. Then*

$$\rho(\lambda) = \begin{cases} (\underline{\lambda}^0 - \underline{D}^0) \cup (\lambda^1 - D^1) \cup (\underline{\lambda}^2 + 1 - \underline{D}^2) \times (\underline{\lambda}^0 + \underline{D}^0) \cup (\lambda^3 + D^3) \cup (\underline{\lambda}^2 + \underline{D}^2) & \text{if } d(\lambda) > 0 \\ (\underline{\lambda}^0 + \underline{D}^0) \cup (\lambda^3 + D^3) \cup (\underline{\lambda}^2 + \underline{D}^2) \times (\underline{\lambda}^0 - \underline{D}^0) \cup (\lambda^1 - D^1) \cup (\underline{\lambda}^2 + 1 - \underline{D}^2) & \text{if } d(\lambda) \leq 0 \end{cases}$$

Proof. Let $\rho^*(\lambda)$ denote the right hand side of the equation in the statement of the proposition. We will show by induction on the number of parts in λ that $\rho^*(\lambda) = \rho(\lambda)$. Fix μ as in (5.4.2) and let $(\gamma, \delta) = \rho(\mu) = \rho^*(\mu)$. There are eight cases to consider:

(a₁) $d(\lambda_m) = 0$ and $d(\mu) > 0$. Let $r = \lfloor \frac{1}{4}(\lambda_m + 2) \rfloor - d(\mu)$ and $s = \lfloor \frac{1}{4}\lambda_m \rfloor + d(\mu)$. Then by the definition of ρ

$$\rho(\lambda) = (\gamma, r) \times (\delta, s)$$

On the other hand $D(\lambda) = (D(\mu), d(\mu), d(\mu))$, $D(\lambda)^1 = D(\mu)^1$, $D(\lambda)^3 = D(\mu)^3$, $\lambda^1 = \mu^1$, and $\lambda^3 = \mu^3$. To compute $\rho^*(\lambda)$, there are two subcases to consider. First assume that $\lambda_m \equiv 0 \pmod{4}$, then $\underline{D}(\lambda)^0 = (\underline{D}(\mu)^0, d(\mu))$, $\underline{D}(\lambda)^2 = \underline{D}(\mu)^2$, $\underline{\lambda}^0 = (\underline{\mu}^0, \frac{1}{4}\lambda_m)$, and $\underline{\lambda}^2 = \underline{\mu}^2$. So

$$\rho^*(\lambda) = (\gamma, \frac{1}{4}\lambda_m - d(\mu)) \times (\delta, \frac{1}{4}\lambda_m + d(\mu)) = (\gamma, r) \times (\gamma, s)$$

If instead $\lambda_m \equiv 2 \pmod{4}$, then $\underline{D}(\lambda)^0 = \underline{D}(\mu)^0$, $\underline{D}(\lambda)^2 = (\underline{D}(\mu), d(\mu))$, $\underline{\lambda}^0 = \underline{\mu}^0$, and $\underline{\lambda}^2 = (\underline{\mu}^2, \frac{1}{4}(\lambda_m - 2))$. So

$$\rho^*(\lambda) = (\gamma, \frac{1}{4}(\lambda_m - 2) + 1 - d(\mu)) \times (\delta, \frac{1}{4}(\lambda_m - 2) + d(\mu)) = (\gamma, r) \times (\gamma, s)$$

(a₂) $d(\lambda_m) = 0$ and $d(\mu) \leq 0$. This is identical to case (a₁) except with $\rho(\lambda)$ and $\rho^*(\lambda)$ flipped.

(b₁) $d(\lambda_m) = 1$ and $d(\mu) > 0$. Let $r = \frac{1}{4}(\lambda_m - 1) - d(\mu)$. Then by the definition of ρ

$$\rho(\lambda) = (\gamma, r) \times \delta.$$

On the other hand, $D(\lambda) = (D(\mu), d(\mu))$, $\underline{D}(\lambda)^0 = \underline{D}(\mu)^0$, $D(\lambda)^1 = (D(\mu)^1, d(\mu))$, $\underline{D}(\lambda)^2 = \underline{D}(\mu)^2$, $D(\lambda)^3 = D(\mu)^3$, $\underline{\lambda}^0 = \underline{\mu}^0$, $\lambda^1 = (\mu^1, \frac{1}{4}(\lambda_m - 1))$, $\underline{\lambda}^2 = \underline{\mu}^2$, and $\lambda^3 = \mu^3$. So

$$\rho^*(\lambda) = (\gamma, \frac{1}{4}(\lambda_m - 1) - d(\mu)) \times \delta = (\gamma, r) \times \delta$$

(b₂) $d(\lambda_m) = 1$ and $d(\mu) = 0$. Let $r = \frac{1}{4}(\lambda_m - 1) - d(\mu)$. Then by the definition of ρ

$$\rho(\lambda) = (\delta, r) \times \gamma.$$

On the other hand, $D(\lambda) = (D(\mu), d(\mu))$, $\underline{D}(\lambda)^0 = \underline{D}(\mu)^0$, $D(\lambda)^1 = (D(\mu)^1, d(\mu))$, $\underline{D}(\lambda)^2 = \underline{D}(\mu)^2$, $D(\lambda)^3 = D(\mu)^3$, $\underline{\lambda}^0 = \underline{\mu}^0$, $\lambda^1 = (\mu^1, \frac{1}{4}(\lambda_m - 1))$, $\underline{\lambda}^2 = \underline{\mu}^2$, and $\lambda^3 = \mu^3$. So

$$\rho^*(\lambda) = (\delta, \frac{1}{4}(\lambda_m - 1) - d(\mu)) \times \gamma = (\delta, r) \times \gamma$$

(b₃) $d(\lambda_m) = 1$ and $d(\mu) < 0$. This is identical to case (b₁) except with $\rho(\lambda)$ and $\rho^*(\lambda)$ flipped.

(c₁) $d(\lambda_m) = -1$ and $d(\mu) > 1$. Let $r = \frac{1}{4}(\lambda_m - 3) + d(\mu)$. Then by the definition of ρ

$$\rho(\lambda) = \gamma \times (\delta, r).$$

On the other hand, $D(\lambda) = (D(\mu), d(\mu) - 1)$, $\underline{D}(\lambda)^0 = \underline{D}(\mu)^0$, $D(\lambda)^1 = D(\mu)^1$, $\underline{D}(\lambda)^2 = \underline{D}(\mu)^2$, $D(\lambda)^3 = (D(\mu)^3, d(\mu))$, $\underline{\lambda}^0 = \underline{\mu}^0$, $\lambda^1 = \mu^1$, $\underline{\lambda}^2 = \underline{\mu}^2$, and $\lambda^3 = (\mu^3, \frac{1}{4}(\lambda_m - 3))$. So

$$\rho^*(\lambda) = \gamma \times (\delta, \frac{1}{4}(\lambda_m - 3) + d(\mu)) = \gamma \times (\delta, r)$$

(c₂) $d(\lambda_m) = 1$ and $d(\mu) = 1$. This is identical to case (c₁) except with $\rho(\lambda)$ and $\rho^*(\lambda)$ flipped.

(c₃) $\underline{d(\lambda_m)} = -1$ and $d(\mu) < 1$. Let $r = \frac{1}{4}(\lambda_m - 3) + d(\mu)$. Then by the definition of ρ

$$\rho(\lambda) = (\gamma, r) \times \delta.$$

On the other hand, $D(\lambda) = (D(\mu), d(\mu))$, $\underline{D}(\lambda)^0 = \underline{D}(\mu)^0$, $D(\lambda)^1 = D(\mu)^1$, $\underline{D}(\lambda)^2 = \underline{D}(\mu)^2$, $D(\lambda)^3 = (D(\mu)^3, d(\mu) - 1)$, $\underline{\lambda}^0 = \underline{\mu}^0$, $\lambda^1 = \mu^1$, $\underline{\lambda}^2 = \underline{\mu}^2$, and $\lambda^3 = (\mu^3, \frac{1}{4}(\lambda_m - 3))$. So

$$\rho^*(\lambda) = (\gamma, \frac{1}{4}(\lambda_m - 3) + d(\mu)) \times \delta = (\gamma, r) \times \delta$$

□

Proposition 5.4.1 motivates the following much simpler recursive definition for $\rho(\lambda)$ which is obvious from Proposition 5.4.1.

Let $\lambda = (\lambda_1 \leq \dots \leq \lambda_m) \in \mathcal{P}_{ro}(n)$. If $m = 0$ set $\tilde{\rho}(\lambda) = (0) \times (0)$. Otherwise, define μ as in (5.4.2). Suppose $\tilde{\rho}(\mu) = \gamma \times \delta$. We define $\tilde{\rho}(\lambda) = \alpha \times \beta$ where α, β are defined as follows: set $q = q(\lambda_m), d = d(\mu)$

- (1) if $\lambda_m \equiv 0 \pmod{4}$ then set $\alpha = (\gamma, q - d), \beta = (\delta, q + d)$;
- (2) if $\lambda_m \equiv 1 \pmod{4}$ then set $\alpha = (\gamma, q - d), \beta = \delta$;
- (3) if $\lambda_m \equiv 2 \pmod{4}$ then set $\alpha = (\gamma, q - d + 1), \beta = (\delta, q + d)$;
- (4) if $\lambda_m \equiv 3 \pmod{4}$ then set $\alpha = \gamma, \beta = (\delta, q + d)$.

Then

$$\rho(\lambda) := \begin{cases} \alpha \times \beta & \text{if } d(\lambda) > 0 \\ \beta \times \alpha & \text{if } d(\lambda) \leq 0. \end{cases}$$

Finally we remark that from the above recursive description of $\tilde{\rho}(\lambda)$ one can deduce the following explicit description of $\tilde{\rho}(\lambda)$.

For $r \in \{0, 1, 2, 3\}$ and $a \geq 0$, define $\epsilon(r, a)$ to be the sequence of length a of the form

$$(5.4.3) \quad \begin{cases} (0, 1, 0, 1, \dots) & \text{if } r = 0 \\ (0, 0, 0, 0, \dots) & \text{if } r = 1 \\ (1, 0, 1, 0, \dots) & \text{if } r = 2 \\ (1, 1, 1, 1, \dots) & \text{if } r = 3. \end{cases}$$

Similarly for $r \in \{0, 1, 2, 3\}$ and $a \geq 0$, define $\delta(r, a)$ to be the sequence of length a of the form

$$(5.4.4) \quad \begin{cases} (0, 1, 0, 1, \dots) & \text{if } r = 0 \\ (0, 0, 0, 0, \dots) & \text{if } r = 1, 2, 3. \end{cases}$$

Now for $\lambda \in \mathcal{N}_{ro}$ write λ as $(\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_k^{a_k})$. Define

$$\begin{aligned} q(\lambda) &= (q(\lambda_1)^{a_1}, q(\lambda_2)^{a_2}, \dots, q(\lambda_k)^{a_k}), \\ r(\lambda) &= (r(\lambda_1)^{a_1}, r(\lambda_2)^{a_2}, \dots, r(\lambda_k)^{a_k}), \\ \epsilon(\lambda) &= (\epsilon(r_1, a_1), \epsilon(r_2, a_2), \dots, \epsilon(r_k, a_k)), \\ \delta(\lambda) &= (\delta(r_1, a_1), \delta(r_2, a_2), \dots, \delta(r_k, a_k)). \end{aligned}$$

When λ is clear from the context, we will write q_i (resp. r_i) for $q(\lambda_i)$ (resp. $r(\lambda_i)$).

Recall from Section 5.3 the sequence $z = (0, 1, \dots)$.

Proposition 5.4.2. *Let $\lambda \in \mathcal{N}_{ro}$, $q = q(\lambda), \epsilon = \epsilon(\lambda), \delta = \delta(\lambda)$. Let $\Lambda \in \mathcal{X}_{d(\lambda)}^{2,0}$ be the symbol with linear presentation $(q + z - \delta, \epsilon)$. Then $\tilde{\rho}(\lambda) = \mathcal{B}(\Lambda)$ and $(q + z - \delta, \epsilon)$ is the canonical linear presentation for Λ .*

Proof. Let $d_i = \sum_{j<i} d(\lambda_j)$, $A_i = \sum_{j<i} a_j$, and

$$x_i = \sum_{j<i} [r_j = 1] + [2 \mid \lambda_j] a_j / 2, \quad y_i = \sum_{j<i} [r_j = 3] + [2 \mid \lambda_j] a_j / 2.$$

Clearly $d_i = x_i - y_i$ and $x_i + y_i = A_i$. Write $\tilde{\rho}(\lambda) = (B_1, B_2)$. By the recursive formula for $\tilde{\rho}(\lambda)$ we have that B_1 consists of blocks of sequences of the form

$$(5.4.5) \quad \begin{cases} q_i - d_i + (0, 0, \dots, 0) & \text{if } r_i = 0 \\ q_i - d_i + (0) & \text{if } r_i = 1 \\ q_i - d_i + 1 + (0, 0, \dots, 0) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3 \end{cases}$$

and B_2 consists of blocks of sequences of the form

$$(5.4.6) \quad \begin{cases} q_i + d_i + (0, 0, \dots, 0) & \text{if } r_i = 0 \\ \emptyset & \text{if } r_i = 1 \\ q_i + d_i + (0, 0, \dots, 0) & \text{if } r_i = 2 \\ q_i + d_i + (0) & \text{if } r_i = 3. \end{cases}$$

Here the sequences of the form $(0, 0, \dots, 0)$ have length $a_i/2$. We wish to compute $\Lambda = (B_1, B_2) + 2((0, 1, 2, \dots), (0, 1, 2, \dots))$. To compute $B_1 + 2(0, 1, 2, \dots)$ we compute along blocks. The sequence $(0, 1, 2, \dots)$ can be written as a concatenation of sequences of the form

$$(5.4.7) \quad \begin{cases} x_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0 \\ x_i + (0) & \text{if } r_i = 1 \\ x_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3. \end{cases}$$

These blocks have the same form as the blocks for B_1 so we can add block-wise. Noting that $2x_i - d_i = x_i + y_i = A_i$, we get that $B_1 + 2(0, 1, 2, \dots)$ consists of blocks of the form

$$(5.4.8) \quad \begin{cases} q_i + A_i + (0, 2, \dots, a_i - 2) & \text{if } r_i = 0 \\ q_i + A_i + (0) & \text{if } r_i = 1 \\ q_i + A_i + (1, 3, \dots, a_i - 1) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3. \end{cases}$$

Similarly, using that $d_i + 2y_i = x_i + y_i = A_i$, we get that $B_2 + 2(0, 1, 2, \dots)$ consists of blocks of the form

$$(5.4.9) \quad \begin{cases} q_i + A_i + (0, 2, \dots, a_i - 2) & \text{if } r_i = 0 \\ \emptyset & \text{if } r_i = 1 \\ q_i + A_i + (0, 2, \dots, a_i - 2) & \text{if } r_i = 2 \\ q_i + A_i + (0) & \text{if } r_i = 3. \end{cases}$$

Thus we see that $(q + z - \delta, \epsilon)$ is a linear presentation of Λ . To see that it is the canonical presentation for Λ note that q is non-decreasing and z is strictly increasing. Thus $q + z$ is strictly increasing. Thus $q + z - \delta$ is strictly increasing except for in blocks corresponding to $r_i = 0$ where the block takes the form $(x, x, x + 2, x + 2, x + 4, x + 4, \dots)$. But for such blocks ϵ is of the form $(0, 1, 0, 1, 0, 1, \dots)$ and so $(q + z - \delta, \epsilon)$ is indeed the canonical linear presentation for Λ . \square

Example 5.4.3. Let $\lambda = (2^4, 5, 7, 12^6, 15)$. Then

$$\begin{aligned} q(\lambda) &= (0, 0, 0, 0, 1, 1, 3, 3, 3, 3, 3, 3, 3) \\ r(\lambda) &= (2, 2, 2, 2, 1, 3, 0, 0, 0, 0, 0, 0, 3) \\ \epsilon(\lambda) &= (1, 0, 1, 0, 0, 1, 0, 1, 0, 1, 0, 1, 1) \\ \delta(\lambda) &= (0, 0, 0, 0, 0, 0, 0, 1, 0, 1, 0, 1, 0) \\ q(\lambda) + z - \delta(\lambda) &= (0, 1, 2, 3, 5, 6, 9, 9, 11, 11, 13, 13, 15) \\ \Lambda &= \begin{pmatrix} 1 & 3 & 5 & 9 & 11 & 13 \\ 0 & 2 & 6 & 9 & 11 & 13 & 15 \end{pmatrix} \\ \tilde{\rho}(\lambda) &= \begin{pmatrix} 1 & 1 & 1 & 3 & 3 & 3 \\ 0 & 0 & 2 & 3 & 3 & 3 & 3 \end{pmatrix} \\ \rho(\lambda) &= \begin{pmatrix} 0 & 0 & 2 & 3 & 3 & 3 & 3 \\ 1 & 1 & 1 & 3 & 3 & 3 \end{pmatrix}. \end{aligned}$$

We now give an explicit description of $d_{BV}(\lambda)$ for $\lambda \in \mathcal{P}_{ro}(N)$.

Let $\lambda \in \mathcal{P}_{ro}(N)$. Write $\lambda = (\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_k^{a_k})$ for $\lambda_1 < \dots < \lambda_k$. Let

$$f : \{1, \dots, \tilde{l}\} \rightarrow \{1, \dots, k\}$$

be the increasing function which enumerates the odd parts of λ . For example, if $\lambda = (1, 2^2, 5, 8^4, 9)$, then $\tilde{l} = 3$ and f is given by $f(1) = 1$, $f(2) = 3$, $f(3) = 5$. Note that N and \tilde{l} have the same parity. Extend the domain of f to a set of even size, as follows. If N is odd, write \tilde{l} as $\tilde{l} = 2l - 1$ and define $f(2l) = k$. In this manner we obtain a function

$$f : \{1, \dots, 2l\} \rightarrow \{1, \dots, k\}.$$

Define δ^i to be the sequence of length k of the form

$$(5.4.10) \quad \delta^i = \begin{cases} (0, \dots, 0, 1, -1, 0, \dots, 0) & \text{if } i \text{ is even} \\ (0, \dots, 0, -1, 1, 0, \dots, 0) & \text{if } i \text{ is odd.} \end{cases}$$

where the first non-zero entry arises at position $f(i)$. If $f(i) = k$ then we interpret δ^i as the sequence

$$(5.4.11) \quad \delta^i = \begin{cases} (0, \dots, 0, 1) & \text{if } i \text{ is even} \\ (0, \dots, 0, -1) & \text{if } i \text{ is odd.} \end{cases}$$

Lemma 5.4.4. Let $\lambda \in \mathcal{P}_{ro}(N)$. Let $\delta = \sum_{i=1}^{\tilde{l}} \delta^i$, $A_i = \sum_{j \geq i} a_j$, and $\bar{\lambda}_i = \lambda_i - \lambda_{i-1}$ (with the convention $\lambda_0 = 0$). Then

$$(5.4.12) \quad d_{BV}(\lambda) = (A_i^{\bar{\lambda}_i + \delta_i})_i.$$

Proof. If a is even, then $d_{BV}(\lambda \cup (a, a)) = d_{BV}(\lambda) + (2^a)$. So if (5.4.12) holds for λ , it also holds for $\lambda \cup (a, a)$. Thus, we can reduce to the case when λ consists of only odd parts, i.e. $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ with all λ_i odd. In this case,

$$\lambda^t = (k^{\bar{\lambda}_1}, (k-1)^{\bar{\lambda}_2}, \dots, 2^{\bar{\lambda}_{k-1}}, 1^{\bar{\lambda}_k}).$$

By the standard algorithm for the collapse of a partition (cf. [CM93, Lemma 6.3.3])

$$d_{BV}(\lambda) = (k^{\bar{\lambda}_1 + \delta_1}, (k-1)^{\bar{\lambda}_2 + \delta_2}, \dots, 2^{\bar{\lambda}_{k-1} + \delta_{k-1}}, 1^{\bar{\lambda}_k + \delta_k})$$

where

$$\delta = \begin{cases} (-1, 2, -2, 2, \dots, -2, 2) & \text{if } N \text{ is even} \\ (-1, 2, -2, 2, \dots, 2, -2) & \text{if } N \text{ is odd.} \end{cases}$$

But in both cases, $\delta = \sum_{i=1}^{\tilde{l}} \delta^i$. This completes the proof. \square

Corollary 5.4.5. *There exists a partition $\underline{d}_{BV}(\lambda)$ such that $d_{BV}(\lambda) = \underline{d}_{BV}(\lambda) \cup \underline{d}_{BV}(\lambda)$.*

Proof. It suffices to show that $\bar{\lambda}_i + \delta_i$ is even for all i . Suppose λ_i, λ_{i-1} are both even. Then $\delta_i = 0$ and $\bar{\lambda}_i + \delta_i = \lambda_i - \lambda_{i-1}$ is even. If λ_{i-1} is odd and λ_i is even, or λ_{i-1} is even and λ_i is odd, then $\delta_i \equiv 1 \pmod{2}$ so $\bar{\lambda}_i + \delta_i$ is even. Finally, if both λ_{i-1}, λ_i are odd then $\delta_i \equiv 1 + 1 \pmod{2}$ and so $\bar{\lambda}_i + \delta_i \equiv 0 \pmod{2}$. \square

5.5. Unipotent representations of finite groups of Lie type. In this section, we will collect various facts about unipotent representations of finite classical groups of Lie type (e.g. classification, decomposition into Harish-Chandra series, decomposition into families, computation of Kawanaka wavefront sets). Essentially all of this information appears in [Lus84c]. However, the presentation below is from [GM20].

Let \mathbf{G} be a connected reductive group defined over \mathbb{F}_q and write $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$ for the set of unipotent representations of $\mathbf{G}(\mathbb{F}_q)$. The set $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$ decomposes into *Harish-Chandra series*

$$\text{Unip}(\mathbf{G}(\mathbb{F}_q)) = \coprod_{(\mathbf{L}, \mathbf{E})} \text{Unip}_{\mathbf{L}, \mathbf{E}}(\mathbf{G}(\mathbb{F}_q))$$

where (\mathbf{L}, \mathbf{E}) ranges over $\mathbf{G}(\mathbb{F}_q)$ -conjugacy classes of pairs consisting of a \mathbf{G} -split Levi subgroup \mathbf{L} of \mathbf{G} and a cuspidal unipotent representation \mathbf{E} of $\mathbf{L}(\mathbb{F}_q)$, and $\text{Unip}_{\mathbf{L}, \mathbf{E}}(\mathbf{G}(\mathbb{F}_q))$ denotes the subset of $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$ consisting of unipotent representations with cuspidal support (\mathbf{L}, \mathbf{E}) . Moreover $\text{Unip}_{\mathbf{L}, \mathbf{E}}(\mathbf{G}(\mathbb{F}_q))$ is parameterised by $\text{Irr}(W_{\mathbf{L}})$ where $W_{\mathbf{L}} := N_{\mathbf{G}(\mathbb{F}_q)}(\mathbf{L}(\mathbb{F}_q))/\mathbf{L}(\mathbb{F}_q)$. Thus there is a bijection

$$(5.5.1) \quad \text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \coprod_{(\mathbf{L}, \mathbf{E})} \text{Irr}(W_{\mathbf{L}}).$$

where, again, the union runs over all $\mathbf{G}(\mathbb{F}_q)$ -conjugacy classes of pairs (\mathbf{L}, \mathbf{E}) consisting of a \mathbf{G} -split Levi subgroup \mathbf{L} of \mathbf{G} and a cuspidal unipotent representation \mathbf{E} of $\mathbf{L}(\mathbb{F}_q)$.

There is also a decomposition of $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$ into families. It is clear from [Lus84c, Section 13.4] that WF is constant on families, so for a family ϕ , the notation $\text{WF}(\phi)$ makes sense. In fact WF induces a bijection between families and special nilpotent orbits of \mathbf{G} defined over \mathbb{F}_q . It follows, and will be useful to note, that a unipotent representation X lies in a family ϕ if and only if $\text{WF}(X) = \text{WF}(\phi)$.

Below, we list all simple finite groups of Lie type which arise in reductive quotients of classical p -adic groups, inner to split. For each group, we give

- (1) A parameterization of the set $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$. See [GM20, Theorem 4.4.13].
- (2) A description of the families in $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$. See [GM20, Proposition 4.4.20].
- (3) A formula for the Kawanaka wavefront set of the elements of $\text{Unip}(\mathbf{G}(\mathbb{F}_q))$, in terms of their parameters.
- (4) A list of all Levi subgroups (up to conjugation) which admit cuspidal representations. In all the cases we consider, there is at most one cuspidal per Levi. See [GM20, Theorem 4.4.28].
- (5) A description of the relative Weyl groups corresponding to the Levi subgroups in (4). See [GM20, Proposition 4.4.29].
- (6) A description of the matching (5.5.1). See [GM20, Proposition 4.4.29].

5.5.1. $A_{n-1}(q)$. Let $\mathbf{G}(\mathbb{F}_q) = PGL(n, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \mathcal{P}(n)$
- (2) Every family is a singleton.
- (3) $\text{WF}(\lambda) = \lambda^t$.
- (4) The only Levi supporting a cuspidal representation is the maximal torus.
- (5) The relative Weyl is S_n .
- (6) The matching is $\lambda \mapsto \lambda$.

5.5.2. ${}^2A_{n-1}(q^2)$. Let $\mathbf{G}(\mathbb{F}_q) = PU(n, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \mathcal{P}(n)$
- (2) Every family is a singleton.
- (3) $WF(\lambda) = \lambda^t$.
- (4) \mathbf{L} is of type $A_{r(r+1)/2-1}$ for $r \geq 0$ such that $r(r+1)/2 \leq n$ and $2 \mid n - r(r+1)/2$.
- (5) $W_{\mathbf{L}} = W_{\frac{1}{2}(n - \frac{1}{2}r(r+1))}$, where W_k denotes the Weyl group of type B_k .
- (6) Let $\lambda \in \mathcal{P}(n)$. The number of boxes in the young diagram of λ with odd hook length minus the number of boxes with even hook length is of the form $\frac{1}{2}s(s+1)$ for some $s \geq 0$. The partition λ belongs to the Harish-Chandra series corresponding to $r = s$. Now let Z be a β -set for λ of odd length. Enumerate the even and odd parts of Z as

$$2x_1 < 2x_2 < \cdots < 2x_k, 2y_1 + 1 < 2y_2 + 1 < \cdots < 2y_l + 1$$

and let $X = \{x_1, x_2, \dots, x_k\}, Y = \{y_1, y_2, \dots, y_l\}$. If s is even then λ corresponds to the bipartition $(\mathcal{P}(X), \mathcal{P}(Y))$, and if s is odd it corresponds to $(\mathcal{P}(Y), \mathcal{P}(X))$.

5.5.3. $B_n(q)$. Let $\mathbf{G}(\mathbb{F}_q) = PSO(2n+1, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \mathcal{Y}_{odd, n}^1$.
- (2) The families are the \sim -classes in $\mathcal{Y}_{odd, n}^1$.
- (3) $WF(S) = \mathcal{P}(2X + 1 \cup 2Y)^t$, where $(X, Y) = S^{sp}$.
- (4) \mathbf{L} is of type B_{r^2+r} for $r \geq 0$ such that $r^2 + r \leq n$.
- (5) $W_{\mathbf{L}} = W_{n-r^2-r}$.
- (6) Let $S \in \mathcal{Y}_{d, n}^1$ for $d \geq 1$ odd. Then S belongs to the Harish-Chandra series corresponding to $r = (d-1)/2$. The corresponding bipartition is $\mathcal{D}(S)$.

5.5.4. $C_n(q)$. Let $\mathbf{G}(\mathbb{F}_q) = PSp(2n, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \mathcal{Y}_{odd, n}^1$.
- (2) The families are the \sim -classes in $\mathcal{Y}_{odd, n}^1$.
- (3) $WF(S) = \mathcal{P}(2X \cup 2Y + 1)^t$, where $(X, Y) = S^{sp}$.
- (4) \mathbf{L} is of type C_{r^2+r} for $r \geq 0$ such that $r^2 + r \leq n$.
- (5) $W_{\mathbf{L}} = W_{n-r^2-r}$.
- (6) Let $S \in \mathcal{Y}_{d, n}^1$ for $d \geq 1$ odd. Then S belongs to the Harish-Chandra series corresponding to $r = (d-1)/2$. The corresponding bipartition is $\mathcal{D}(S)$.

5.5.5. $D_n(q)$. Let $\mathbf{G}(\mathbb{F}_q) = PCO^+(2n, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \sqcup_{d \geq 0, 4 \mid d} \mathcal{Y}_{d, n}^1$ except to each degenerate symbol in $\mathcal{Y}_{0, n}^1$ there are two unipotent representations.
- (2) The families are the \sim -classes in $\sqcup_{d \geq 0, 4 \mid d} \mathcal{Y}_{d, n}^1$.
- (3) $WF(S) = (\mathcal{P}(2X + 1 \cup 2Y)^t)_D$, where $(X, Y) = S^{sp}$.
- (4) \mathbf{L} is of type D_{r^2} for $r \geq 0$ even such that $r^2 \leq n$.
- (5) $W_{\mathbf{L}} = W_{n-r^2}$ if $r \geq 2$ and W'_n if $r = 0$.
- (6) Let $S \in \mathcal{Y}_{d, n}^1$ for $d \in \mathbb{Z}, 4 \mid d$. Then S belongs to the Harish-Chandra series corresponding to $r = d/2$. The corresponding bipartition is $\mathcal{D}(S)$. In the case where S is degenerate there are two unipotent representations parameterised by S , but also two irreducible representations of W'_n parameterised by $\mathcal{D}(S)$ and these are matched in some manner (the precise nature of which is not important to us).

5.5.6. ${}^2D_n(q^2)$. Let $\mathbf{G}(\mathbb{F}_q) = PCO^-(2n, \mathbb{F}_q)$.

- (1) $\text{Unip}(\mathbf{G}(\mathbb{F}_q)) \leftrightarrow \sqcup_{d \geq 0, 4 \mid d-2} \mathcal{Y}_{d, n}^1$.
- (2) The families are the \sim -classes in $\sqcup_{d \geq 0, 4 \mid d-2} \mathcal{Y}_{d, n}^1$.

- (3) $\text{WF}(S) = (\mathcal{P}(2X + 1 \cup 2Y)^t)_D$, where $(X, Y) = S^{sp}$.
- (4) \mathbf{L} is of type ${}^2D_{r,2}$ for $r \geq 1$ odd such that $r^2 \leq n$. Here 2D_1 is regarded as the empty graph.
- (5) $W_{\mathbf{L}} = W_{n-r^2}$.
- (6) Let $S \in \mathcal{Y}_{d,n}^1$ for $d \geq 0, 4 \mid d-2$. Then S belongs to the Harish-Chandra series corresponding to $r = d/2$. The corresponding bipartition is $\mathcal{D}(S)$.

5.6. The generalised Springer correspondence. In this section we will recall the statement of the generalised Springer correspondence and give a description of this correspondence in classical types. The main ideas are from [Lus84a] and [LS85].

Let G be a complex reductive group, T a maximal torus of G , B a Borel subgroup of G containing T , and I_0 the set of simple roots determined by T and B . For a subset $J \subseteq I_0$ write G_J for the Levi subgroup of G determined by T and J . For a Levi subgroup L of G write $W_L = N_G(L)/L$. Let $\mathcal{N}_{o,l}$ be the set of pairs $(\mathbb{O}, \mathcal{L})$ consisting of a nilpotent orbit $\mathbb{O} \in \mathcal{N}_o$ and a G -equivariant irreducible local system \mathcal{L} on \mathbb{O} . The generalised Springer correspondence is a bijection

$$(5.6.1) \quad \mathcal{N}_{o,l} \leftrightarrow \coprod_{(L, \mathcal{C}, \mathcal{E}) \in \mathfrak{C}(G)} \text{Irr}(W_L), \quad (\mathbb{O}, \rho) \mapsto E(\mathbb{O}, \rho)$$

where $\mathfrak{C}(G)$ is the set of G -conjugacy classes of triples $(L, \mathcal{C}, \mathcal{E})$ where L is a Levi subgroup of G , \mathcal{C} is a nilpotent orbit of the Lie algebras \mathfrak{l} of L , and \mathcal{E} is a L -equivariant cuspidal local system on \mathcal{C} . Let $\bar{\mathfrak{C}}(G)$ be the set of all triples $(J, \mathcal{C}, \mathcal{E})$ where $J \subseteq I_0$, \mathcal{C} is a nilpotent orbit of the Lie algebra \mathfrak{g}_J of G_J , and \mathcal{E} is a G_J -equivariant cuspidal local system on \mathcal{C} . There is the obvious map $\bar{\mathfrak{C}}(G) \rightarrow \mathfrak{C}(G)$, given by $(J, \mathcal{C}, \mathcal{E}) \mapsto (G_J, \mathcal{C}, \mathcal{E})$. It is clear from form of the Levis that admit cuspidal local systems that this map is a bijection. For $x = (J, \mathcal{C}, \mathcal{E}) \in \bar{\mathfrak{C}}(G)$ we will write L_x for G_J and W_x for W_{L_x} . We write $(\mathcal{N}_{o,l})_x$ for the pairs lying in the block parameterised by $x \in \bar{\mathfrak{C}}(G)$ and $(\mathcal{N}_o)_x$ for $p_1((\mathcal{N}_{o,l})_x)$. So if $x = (J, \mathcal{C}, \mathcal{E})$ then

$$(\mathcal{N}_{o,l})_x \leftrightarrow \text{Irr}(W_x).$$

Note that for a Levi subgroup $L \subset G$ and a character χ of $Z(L)$, there is at most one pair $(\mathcal{C}, \mathcal{E})$ such that $(L, \mathcal{C}, \mathcal{E}) \in \mathfrak{C}(G)$ and $Z(L)$ acts via (a multiple of) χ on \mathcal{E} .

Below, we describe this bijection for all simple simply connected groups of classical type. For each group, we give

- (1) A parameterization of $\mathcal{N}_{o,l}$.
- (2) A description of the projection $p_1 : \mathcal{N}_{o,l} \rightarrow \mathcal{N}_o$ in terms of their parameter.
- (3) A list of all pairs (L, χ) consisting of a Levi subgroup $L \subset G$ and a character χ of $Z(L)$ such that there is a triple $(L, \mathbb{O}, \mathcal{E}) \in \mathfrak{C}$ and $Z(L)$ acts via (a multiple of) χ on \mathcal{E} . In most cases, χ is uniquely determined by L ; in such cases we omit it.
- (4) A description of the relative Weyl groups corresponding to the Levi subgroups in (3).
- (5) A description of the matching (5.6.1).

The references are: [LS85, Section 5.1] for $SL(N)$, [Lus84a, Section 10.6] for $SO(N)$, [Lus84a, Section 10.4] and $Sp(2n)$, and [LS85, Section 0.10, Section 4] for $Spin(N)$.

For $G = Spin(N)$, we present (1)-(5) in a slightly different manner. There are natural inclusions

$$\mathcal{N}_{o,l}^{SO(N)} \rightarrow \mathcal{N}_{o,l}^{Spin(N)}, \quad \bar{\mathfrak{C}}(SO(N)) \rightarrow \bar{\mathfrak{C}}(Spin(N)).$$

With respect to these inclusions there are decompositions

$$\mathcal{N}_{o,l}^{Spin(N)} = \mathcal{N}_{o,l}^{SO(N)} \sqcup \mathcal{N}_{o,l}^*$$

where $\mathcal{N}_{o,l}^*$ is the set of pairs $(\mathbb{O}, \mathcal{L}) \in \mathcal{N}_{o,l}^{Spin(n)}$ consisting of a nilpotent orbit $\mathbb{O} \in \mathcal{N}_o$ and a $Spin(N)$ -equivariant local system \mathcal{L} on \mathbb{O} which is not $SO(N)$ -equivariant, and

$$\bar{\mathfrak{C}}(Spin(N)) = \bar{\mathfrak{C}}(SO(N)) \sqcup \bar{\mathfrak{C}}^*(Spin(N))$$

where $\overline{\mathcal{C}}^*$ is the set of triples $(J, \mathcal{C}, \mathcal{E}) \in \overline{\mathcal{C}}(\text{Spin}(N))$ where \mathcal{E} is not equivariant for the associated Levi subgroup of $SO(N)$. The matching (5.6.1) takes $\mathcal{N}_{o,l}^{SO(N)}$ to $\overline{\mathcal{C}}(SO(N))$ and $\mathcal{N}_{o,l}^*$ to $\overline{\mathcal{C}}^*(\text{Spin}(N))$. In (5.6.2) and (5.6.3), we describe the correspondence for $\mathcal{N}_{o,l}^{SO(N)}$, and in (5.6.5) and (5.6.6), we describe the correspondence for $\mathcal{N}_{o,l}^*$.

5.6.1. $SL(N)$.

- (1) $\mathcal{N}_{o,l}$ is parameterized by pairs (λ, ρ) , where $\lambda = (\lambda_1, \dots, \lambda_k) \in \mathcal{P}(N)$ and ρ is a character of \mathbb{Z}_d , where $d = \gcd(\lambda_1, \dots, \lambda_k)$.
- (2) $p_1(\lambda, \rho) = \lambda$.
- (3) L is of type $(N/r)A_{r-1}$ for $r \mid N$, and χ is a character (trivial on connected center) of order r .
- (4) $W_L = S_{N/r}$.
- (5) Let $\lambda = (\lambda_1, \dots, \lambda_k)$, $d = \gcd(\lambda_1, \dots, \lambda_k)$, and let ρ be a character of \mathbb{Z}_d of degree m . Then (λ, ρ) belongs to a block with $r = m$. The specific χ it corresponds to depends on ρ , but is not important for our purposes. The corresponding $S_{N/r}$ -representation is given by the partition $(\lambda_1/r, \dots, \lambda_k/r)$ of N/r .

5.6.2. $SO(N)$, N odd. Let $n = \lfloor N/2 \rfloor$.

- (1) $\mathcal{N}_{o,l} \leftrightarrow \mathcal{Y}_{odd,n}^2$.
- (2) Let $S \in \mathcal{Y}_{odd,n}^2$. Then $p_1(S) = \mathcal{P}(2(X - z) + 1 \cup 2(Y - z))$ where $(X, Y) = S^{sp}$.
- (3) L is of type $B_{(r^2-1)/2}$ for $r \geq 0$ odd such that $r^2 \leq N$.
- (4) $W_L = W_{\frac{1}{2}(N-r^2)}$.
- (5) Let $S \in \mathcal{Y}_{d,n}^2$ for $d \geq 0$ odd. Then S belongs to the block corresponding to $r = d$. The corresponding bipartition is $\mathcal{D}(S)$.

5.6.3. $SO(N)$, N even. Let $n = \lfloor N/2 \rfloor$.

- (1) $\mathcal{N}_{o,l} \leftrightarrow \mathcal{Y}_{even,n}^2$.
- (2) Let $S \in \mathcal{Y}_{even,n}^2$. Then $p_1(S) = \mathcal{P}(2(X - z) + 1 \cup 2(Y - z))$ where $(X, Y) = S^{sp}$.
- (3) L is of type $D_{r^2/2}$ for $r \geq 0$ even such that $r^2 \leq N$.
- (4) $W_L = W_{\frac{1}{2}(N-r^2)}$ if $r > 0$ and W'_n if $r = 0$.
- (5) Let $S \in \mathcal{Y}_{d,n}^2$ for $d \geq 0$ even. Then S belongs to the block corresponding to $r = d$. The corresponding bipartition is $\mathcal{D}(S)$. The label of $\mathcal{D}(S)$ when S is degenerate is not important for our purposes.

5.6.4. $Sp(2n)$.

- (1) $\mathcal{N}_{o,l} \leftrightarrow \mathcal{X}_{odd,n}^{1,1}$.
- (2) Let $S \in \mathcal{X}_{odd,n}^{1,1}$. Then $p_1(S) = \mathcal{P}(2(X - z) \cup 2(Y - z) + 1)$ where $(X, Y) = S^{sp}$.
- (3) L is of type $C_{\frac{1}{2}r(r+1)}$ for $r \geq 0$ such that $\frac{1}{2}r(r+1) \leq n$.
- (4) $W_L = W_{n - \frac{1}{2}r(r+1)}$.
- (5) Let $S \in \mathcal{X}_{d,n}^{1,1}$ for d odd. Then S belongs to the block corresponding to $r = d - 1$ if $d \geq 0$ and $r = -d$ if $d < 0$. The corresponding bipartition is $\mathcal{B}(S)$ if $d \geq 0$ and the flip of $\mathcal{B}(S)$ otherwise.

5.6.5. $Spin(N)$, N odd.

- (1) $\mathcal{N}_{o,l}^* \leftrightarrow \mathcal{P}_{ro}(N)$.
- (2) $p_1(\lambda) = \lambda$.
- (3) L is of type $B_{(d-1)(2d+1)/2} + \frac{1}{4}(N - d(2d-1))A_1$ for $d \in \mathbb{Z}$, $d \equiv N \pmod{4}$ and $d(2d-1) \leq N$.
- (4) $W_L = W_{(N-d(2d-1))/4}$.

- (5) Let $\lambda \in \mathcal{P}_{ro}(N)$. Then λ belongs to the block corresponding to $d = d(\lambda)$, cf. (5.4.1), and the corresponding bipartition is $\rho(\lambda)$, cf. Proposition 5.4.1.

5.6.6. *Spin(N), N even.*

- (1) $\mathcal{N}_{o,l}^* \leftrightarrow \mathcal{P}_{ro}(N) \sqcup \mathcal{P}_{ro}(N)$.
- (2) $p_1(\lambda) = \lambda$.
- (3) L is of type $D_{d(2d-1)/2} + \frac{1}{4}(N - d(2d-1))A_1$ for $d \in \mathbb{Z}$, $d \equiv N \pmod{4}$ and $d(2d-1) \leq N$.
- (4) $W_L = W_{(N-d(2d-1))/4}$.
- (5) Let $\lambda \in \mathcal{P}_{ro}(N)$. Then λ belongs to the block corresponding to $d = d(\lambda)$, cf. (5.4.1), and the corresponding bipartition is $\rho(\lambda)$, cf. Proposition 5.4.1.

5.7. The Arithmetic-Geometric correspondence. Recall from Section 4 the definitions of $\mathbf{G}, G^\vee, \overline{\mathcal{C}}, \overline{\mathcal{M}}, \theta$ etc. Recall from the previous section the definition of the set $\overline{\mathcal{C}}(G^\vee)$ and note that there is a natural inclusion $\overline{\mathcal{C}}(G^\vee) \rightarrow \overline{\mathcal{C}}$. In this section, we will give an explicit description of Lusztig's arithmetic-geometric correspondence (cf. Section 4) for the geometric diagrams $\overline{\mathcal{C}}(G^\vee)$ for the classical groups.

Below we record

- (1) A list of all pairs (L^\vee, χ) consisting of a Levi subgroup $L^\vee \subset G^\vee$ and a character χ of $Z(L^\vee)$ such that there is a triple $(L^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \in \overline{\mathcal{C}}(G^\vee)$ and $Z(L^\vee)$ acts on the stalks of \mathcal{E}^\vee by (a multiple of) χ . In most cases, χ is uniquely determined by L^\vee ; in such cases we omit it.
- (2) The type of the affine Weyl group $\mathcal{W}^\vee(J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$.
- (3) A description of J and ω in $\theta(x) = (J, \mathbf{E}, \psi, \omega)$ for each $x \in \overline{\mathcal{C}}(G^\vee)$. In all the cases we consider there is a unique choice of \mathbf{E} and so we omit it.

This information is taken from [Lus95a, Section 7].

For $G^\vee = \text{Spin}(N)$ we present this information in a slightly different manner. As discussed in Section 5.6, there is a decomposition

$$\overline{\mathcal{C}}(\text{Spin}(N)) = \overline{\mathcal{C}}(\text{SO}(N)) \sqcup \overline{\mathcal{C}}^*(\text{Spin}(N)).$$

In (5.7.2) and (5.7.3), we describe the restriction of θ to $\overline{\mathcal{C}}(\text{SO}(N))$, and in (5.7.5) and (5.7.6), we describe the restriction of θ to $\overline{\mathcal{C}}^*(\text{Spin}(N))$.

5.7.1. $G^\vee = \text{SL}(N)$.

- (1) L^\vee is of type $(N/r)A_{r-1}$ for $r \mid N$, χ is of order r .
- (2) $\tilde{A}_{N/r-1}$
- (3) J is \emptyset and ω is of order r .

5.7.2. $G^\vee = \text{SO}(N)$, N odd.

- (1) L^\vee is of type $B_{\frac{1}{2}(r^2-1)}$ for $r \geq 0$ odd and $\frac{1}{2}(r^2-1) \leq n$.
- (2) $\tilde{C}_{\frac{1}{2}(N-r^2)}$
- (3) J is $C_{a^2+a} \times C_{a^2+a}$ where $a := (r-1)/2$. $\omega = 1$.

5.7.3. $G^\vee = \text{SO}(N)$, N even.

- (1) L^\vee is of type $D_{r^2/2}$ for $r \geq 0$ even and $r^2/2 \leq n$.
- (2) $\tilde{C}_{\frac{1}{2}(N-r^2)}$ if $r > 0$, \tilde{D}_n if $r = 0$.
- (3) J is $D_{a^2} \times D_{a^2}$ where $a := r/2$. $\omega = 1$ if a is even and ω of type I if a is odd.

5.7.4. $G^\vee = Sp(2n)$.

- (1) L^\vee is of type $C_{\frac{1}{2}r(r+1)}$ for $r \geq 0$ and $\frac{1}{2}r(r+1) \leq n$.
- (2) $\tilde{C}_{n-\frac{1}{2}r(r+1)}$ if $r > 0$, \tilde{B}_n if $r = 0$.
- (3) J is $D_{a^2} \times B_{a^2+a}$ if r is even and $D_{a^2} \times B_{a^2-a}$ if r is odd where $a := \lfloor (r+1)/2 \rfloor$. $\omega = 1$ if a is even and $\omega \neq 1$ if a is odd.

5.7.5. $G^\vee = Spin(N)$, N odd.

- (1) L^\vee is of type $B_{(d-1)(2d+1)/2} + \frac{1}{4}(N - d(2d-1))A_1$ for $d \equiv N \pmod{4}$ and $d(2d-1) \leq N$.
- (2) $\tilde{C}_{(N-d(2d-1))/4}$
- (3) J is $C_{(d^2-1)/4} \times A_{d(d-1)/2-1} \times C_{(d^2-1)/4}$, $\omega \neq 1$.

5.7.6. $G^\vee = Spin(N)$, N even.

- (1) L^\vee is of type $D_{d(2d-1)/2} + \frac{1}{4}(N - d(2d-1))A_1$ for $d \equiv N \pmod{4}$ and $d(2d-1) \leq N$. There are two possible χ .
- (2) $\tilde{C}_{(N-d(2d-1))/4}$
- (3) J is $D_{d^2/4} \times A_{d(d-1)/2-1} \times D_{d^2/4}$, ω of type II.

5.8. **Faithful nilpotent orbits.** Recall the notation $\mathbf{G}, G^\vee, \overline{\mathfrak{S}}, \overline{\mathfrak{U}}, \theta$ from Section 4, $\mathcal{N}_{o,l}, (\mathcal{N}_{o,l})_x, W_x, \overline{\mathfrak{C}}$ from Section 4.3, $d_A, \mathcal{K}_I, \overline{\mathfrak{L}}$ from Section 2.3, and $\text{Unip}, \text{Unip}_{\mathbf{L}, \mathbf{E}}, \text{WF}(\phi)$ from Section 5.5.

- If $x = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \in \overline{\mathfrak{S}}$, and $\theta(x) = (K, \mathbf{E}, \psi, \omega)$, write \mathcal{W}_x for the affine Weyl group $\mathcal{W}(K, \omega, \mathbf{E})$ and S_x for the Coxeter generators for \mathcal{W}_x . Recall from Section 4.2 that $S_x \leftrightarrow (I - K)/\omega$. Write W_x for the finite part of \mathcal{W}_x (when $x \in \overline{\mathfrak{C}}(G^\vee)$ this agrees with the definition of W_x given in Section 5.6). Write $\pi : \mathcal{W}_x \rightarrow W_x$ for the projection map and $\mathcal{I} : (I - K)/\omega \xrightarrow{\sim} S_x$ for the bijection.
- Let $\mathcal{J}_x = \{J \mid K \subseteq J \subsetneq I, \omega J = J\}$.
- Write $I_{0,K}$ for the element of \mathcal{J}_x with $\mathcal{I}(I_{0,K} - K)$ equal to the Coxeter generators of W_x .
- Let $\mathcal{F}_x = \{(J, \phi) \mid J \in \mathcal{J}_x, \phi \text{ a family of } \text{Unip}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))\}$.
- For $J \in \mathcal{J}_x$ write $\mathcal{W}_{x,J}$ for $\langle y \mid y \in \mathcal{I}((J - K)/\omega) \rangle$ and $W_{x,J}$ for $\pi(\mathcal{W}_{x,J})$.
- Let $J \in \mathcal{J}_x$. Then $\mathbf{L}_{c^\omega(K)}(\mathbb{F}_q)$ is a $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ -split Levi subgroup of $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ affording the cuspidal representation \mathbf{E} , and $\mathcal{H}(\mathbf{P}_{c^\omega(J)(\mathfrak{o})}, \mathbf{E})$ (c.f. Section 4.4) has underlying coxeter system $\mathcal{W}_{x,J} \cong W_{x,J}$. Thus $\text{Irr}(W_{x,J})$ (resp. $\text{Irr}(\mathcal{W}_{x,J})$) parameterizes the irreducible representations in the Harish-Chandra series of $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ determined by the cuspidal data $(\mathbf{L}_{c^\omega(K)}, \mathbf{E})$. Let $\mathcal{U}_{x,J} : \text{Irr}(W_{x,J}) \rightarrow \text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$ (resp. $\tilde{\mathcal{U}}_{x,J} : \text{Irr}(\mathcal{W}_{x,J}) \rightarrow \text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$) denote this parameterisation. We also write $\mathcal{U}_{x,J}$ for the composition

$$\text{Irr}(W_{x,J}) \xrightarrow{\sim} \text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)) \hookrightarrow \text{Unip}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)).$$

Definition 5.8.1. Let $x = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee) \in \overline{\mathfrak{C}}(G^\vee)$, $\theta(x) = (K, \mathbf{E}, \psi, \omega)$ and $\mathbb{O}^\vee \in (\mathcal{N}_o^\vee)_x$. We say that \mathbb{O}^\vee is x -faithful if there exists a pair $(J, \phi) \in \mathcal{F}_x$ such that

$$(i) \quad \overline{\mathfrak{L}}(J, \text{WF}(\phi, \mathbb{C})) = d_A(\mathbb{O}^\vee, 1),$$

and for every irreducible representation ρ of $A(\mathbb{O}^\vee)$ such that $(\mathbb{O}, \rho) \in (\mathcal{N}_{o,l}^\vee)_x$,

$$(ii) \quad \text{there is a representation } F \in \text{Irr}(W_{x,J}) \text{ such that } \mathcal{U}_{x,J}(F) \in \phi \text{ and}$$

$$\text{Hom}_{W_{x,J}}(F, E(\mathbb{O}^\vee, \rho)|_{W_{x,J}}) \neq 0;$$

(iii) for every $J' \in \mathcal{J}_x$ and $F' \in \text{Irr}(W_{x,J'})$, if

$$\text{Hom}_{W_{x,J'}}(F', E(\mathbb{O}^\vee, \rho)|_{W_{x,J'}}) \neq 0$$

then

$$(5.8.1) \quad \overline{\mathbb{L}}(J', \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C})) \leq d_A(\mathbb{O}^\vee, 1).$$

Note that Equation 5.8.1 is equivalent to the condition

$$d_S(J', \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C})) \geq \mathbb{O}^\vee.$$

Lemma 5.8.2. *It suffices to check (iii) in Definition 5.8.1 for J' maximal in \mathcal{J}_x with respect to inclusion.*

Proof. Suppose condition (iii) of Definition 5.8.1 holds for all J' maximal. Let $J' \in \mathcal{J}_x$ and $F' \in \text{Irr}(W_{x,J'})$ be such that

$$\text{Hom}_{W_{x,J'}}(F', E(\mathbb{O}^\vee, \rho)|_{W_{x,J'}}) \neq 0.$$

Let $J'' \in \mathcal{J}_x$ be maximal among elements of \mathcal{J}_x containing J' . Then there exists an $F'' \in \text{Irr}(W_{x,J''})$ such that

$$\text{Hom}_{W_{x,J''}}(F'', E(\mathbb{O}^\vee, \rho)|_{W_{x,J''}}) \neq 0$$

and

$$\text{Hom}_{W_{x,J'}}(F', F''|_{W_{x,J'}}) \neq 0.$$

By assumption

$$d_S(J'', \text{WF}(\mathcal{U}_{x,J''}(F''), \mathbb{C})) \geq \mathbb{O}^\vee.$$

Thus it suffices to prove

$$d_S(J', \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C})) \geq d_S(J'', \text{WF}(\mathcal{U}_{x,J''}(F''), \mathbb{C})).$$

By [Oka21, Proposition 2.31] and the remarks in the proof we have

$$d_S(J', \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C})) = d_S(J'', \text{Sat}_{L_{J''}}^{L_{J''}} \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C})).$$

Thus it suffices to prove that for $F' \in \text{Irr}(W_{x,J'})$, $F'' \in \text{Irr}(W_{x,J''})$

$$(5.8.2) \quad \text{Hom}_{W_{x,J'}}(F', F''|_{W_{x,J'}}) \neq 0 \implies \text{Sat}_{L_{J''}}^{L_{J''}} \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C}) \leq \text{WF}(\mathcal{U}_{x,J''}(F'')).$$

By [HL83, Theorem 5.9],

$$\mathcal{U}_{x,J''} \left(\text{Ind}_{W_{x,J'}}^{W_{x,J''}} F' \right) = \text{Ind}_{L_{J''}}^{L_{J''}} \mathcal{U}_{x,J'}(F')$$

where Ind applied to unipotent characters means Harish-Chandra induction. Thus by Frobenius reciprocity we have that

$$(5.8.3) \quad \text{Hom}_{W_{x,J'}}(F', F''|_{W_{x,J'}}) \neq 0 \Leftrightarrow \mathcal{U}_{x,J''}(F'') \subseteq \text{Ind}_{L_{J''}}^{L_{J''}} \mathcal{U}_{x,J'}(F').$$

By [TT20, Lemma 10.3] (and noting that we may replace the Deligne-Lusztig induction in the lemma by Harish-Chandra induction since $\mathbf{L}_{c^\omega(J')}$ is a $\mathbf{L}_{c^\omega(J'')}$ -split Levi of $\mathbf{L}_{c^\omega(J'')}$ by [TT20, Section 2.1]) we have that

$$(5.8.4) \quad \mathcal{U}_{x,J''}(F'') \subseteq \text{Ind}_{L_{J''}}^{L_{J''}} \mathcal{U}_{x,J'}(F') \implies \text{Sat}_{L_{J''}}^{L_{J''}} \text{WF}(\mathcal{U}_{x,J'}(F'), \mathbb{C}) \leq \text{WF}(\mathcal{U}_{x,J''}(F'')).$$

Stringing together Equation 5.8.3 and Equation 5.8.4 gives Equation 5.8.2 as required. \square

Theorem 5.8.3. *Suppose \mathbf{G} is a simple group of adjoint type and let $\mathbb{O}^\vee \in \mathcal{N}_o^\vee$. Then for every $x \in \overline{\mathfrak{C}}(G^\vee)$ such that $\mathbb{O}^\vee \in (\mathcal{N}_o)_x$, \mathbb{O}^\vee is x -faithful.*

Proof. We consider separately the cases of classical and exceptional groups. If \mathbf{G} is classical, the statement is proved in Section 5.9. If \mathbf{G} is exceptional, it is proved in Section 5.10. \square

5.9. Proof of faithfulness in classical types.

5.9.1. $G^\vee = SL(N)$. Let $n = N - 1$. Let $x \in \overline{\mathfrak{C}}(G^\vee)$ correspond to $\chi \in \text{Irr}(\mathbb{Z}_N)$ of order $r \mid N$. Then $\theta(x) = (\emptyset, 1, \psi, \chi)$ and \mathcal{W}_x is of type \tilde{A}_{k-1} where $k = N/r$. A nilpotent orbit $\mathbb{O}^\vee \in \mathcal{N}_o^\vee \leftrightarrow \lambda = (\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_e)$ lies in $(\mathcal{N}_o^\vee)_x$ iff $r \mid \lambda_i$ for all i .

We wish to show that λ is x -faithful. Let $J = rA_{k-1}$ and $\mu = (\lambda_1/r \leq \lambda_2/r \leq \dots \leq \lambda_e/r)$. Then $\mathbf{L}_{c^\chi(J)}$ is isomorphic to $A_{k-1}(q)$ and so the families are parameterised by $\mathcal{P}(k)$. Let ϕ be the family parameterised by the partition μ . Then $\overline{\mathbf{L}}(J, \text{WF}(\phi, \mathbb{C})) = \mu^t \cup \mu^t \cup \dots \cup \mu^t$ where there are r terms in this expression. But $\mu^t \cup \mu^t \cup \dots \cup \mu^t = (\mu + \mu + \dots + \mu)^t = \lambda^t = d(\mathbb{O}^\vee, 1)$. Thus condition (i) holds. Condition (ii) holds because $W_{x,J} = W_x$ and $E(\mathbb{O}^\vee, \rho) = \mu$. Finally condition (iii) holds by using Lemma 5.8.2 to reduce to the case where $J \in \mathcal{J}_x$ is maximal and noting that for such J , $W_{x,J} = W_x$. Condition (iii) then holds as a consequence of condition (ii) (the only possible F' that can arise is $F' = F$).

5.9.2. $SO(N)$, N odd. Let $n = \lfloor N/2 \rfloor$, $G^\vee = Spin(N)$. Fix $\mathbb{O}^\vee \in \mathcal{N}_o^\vee$. We show that \mathbb{O}^\vee is x -faithful for all $x \in \overline{\mathfrak{C}}(SO(N))$ such that $\mathbb{O}^\vee \in (\mathcal{N}_o^\vee)_x$ by exhibiting a single pair (J, ϕ) that works for all such blocks. Let $\lambda = (\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_e)$ be the orthogonal partition of N corresponding to \mathbb{O}^\vee . Define $\mu(\lambda) \subseteq \lambda^t$ as follows

$$m_{\mu(\lambda)}(x) = \begin{cases} 2 & \text{if } x \text{ is even and } m_{\lambda^t}(x) \text{ is} \\ & \text{even and nonzero} \\ 1 & \text{if } x \text{ is even and } m_{\lambda^t}(x) \text{ is} \\ & \text{odd} \\ 0 & \text{otherwise} \end{cases}$$

and define $\nu(\lambda) = d_{BV}(\lambda) \setminus \mu(\lambda)$. By [CMO21, Lemma 4.8.4] $\mu(\lambda)$ and $\nu(\lambda)$ are both special symplectic partitions. We set J to be the set $C_m \times C_{n-m}$ where $m = \lfloor \mu \rfloor / 2$, and ϕ to be the family characterised by $\text{WF}(\phi, \mathbb{C}) = (\mu(\lambda), \nu(\lambda))$. Since $J = C_m \times C_{n-m}$, the family ϕ decomposes as a product $\phi_1 \times \phi_2$ where ϕ_1 is a family of C_m , ϕ_2 is a family of C_{n-m} and

$$\text{WF}(\phi_1) = \mu(\lambda), \quad \text{WF}(\phi_2) = \nu(\lambda).$$

It follows from [CMO21, Lemma 4.8.6] that condition (i) holds for this choice of (J, ϕ) . For condition (ii), let $x \in \overline{\mathfrak{C}}(SO(N))$ and suppose $(\mathbb{O}^\vee, \rho) \in (\mathcal{N}_{o,l})_x$. As described in Section 5.6.2 (3) let r be the odd integer corresponding to x . Then $\theta(x) = (C_{a^2+a} \times C_{a^2+a}, \mathbf{E}, \psi, \mathbf{1})$, $a = (r-1)/2$, \mathbf{E} is a uniquely determined cuspidal unipotent representation, and \mathcal{W}_x is of type $\tilde{C}_{\frac{1}{2}(r^2-1)}$. By Section 5.5.4, $\text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)) \leftrightarrow \mathcal{Y}_{r,m}^1 \times \mathcal{Y}_{r,n-m}^1$. Note that $K \subseteq J$ (and hence $J \in \mathcal{J}_x$) if and only if $\mathcal{Y}_{r,m}^1 \times \mathcal{Y}_{r,n-m}^1$ is non-empty. Under the bijection in Section 5.6.2 (1), (\mathbb{O}^\vee, ρ) corresponds to a $S \in \mathcal{Y}_{r,n}^2$ with $p_1(S) = \lambda$. By [CMO21, Lemma 4.4.2], S has the following description in terms of $S^{sp} = (X, Y)$. Let $X = (X_1, \dots, X_c), Y = (Y_1, \dots, Y_c)$ be the refinements of X and Y (see paragraph preceding [CMO21, Example 4.4.1]). Then $S = (A, B)$ where $A = (A_1, \dots, A_c), B = (B_1, \dots, B_c)$ and $\{A_i, B_i\} = \{X_i, Y_i\}$ for all i . Let $S^{1,sp} = (X^1, Y^1) \in \mathcal{Y}_{1,m}^1, S^{2,sp} = (X^2, Y^2) \in \mathcal{Y}_{1,n-m}^1$ be the special symbols in the families ϕ_1 and ϕ_2 respectively. Taking sufficiently large representatives in the relevant shift equivalence classes we have by condition (i) and [CMO21, Proposition 4.7.1] that $S^{sp} = S^{1,sp} + S^{2,sp}$. Let $S^{i,sp} = (X^i, Y^i) = ((X_1^i, \dots, X_c^i), (Y_1^i, \dots, Y_c^i))$ be the decomposition into subsequences along the same indices as the decomposition of X, Y . Define $S^i = ((A_1^i, \dots, A_c^i), (B_1^i, \dots, B_c^i))$ where

$$(5.9.1) \quad (A_j^i, B_j^i) = \begin{cases} (X_j^i, Y_j^i) & \text{if } (A_j, B_j) = (X_j, Y_j) \\ (Y_j^i, X_j^i) & \text{if } (A_j, B_j) = (Y_j, X_j). \end{cases}$$

By construction $S^{i,sp} \sim S^i$ and $S = S^1 + S^2$. Moreover, by [CMO21, Lemma 4.5.2] and [CMO21, Lemma 4.9.4] we have that $S^1 \in \mathcal{Y}_{r,m}^1, S^2 \in \mathcal{Y}_{r,n-m}^1$. Thus $\text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)) \neq \emptyset$, so $J \in \mathcal{J}_x$,

and

$$(5.9.2) \quad \mathcal{D}(S) = \mathcal{D}(S^1) + \mathcal{D}(S^2).$$

Note that $\mathcal{D}(S)$ parameterises an irreducible representation of W_x and $(\mathcal{D}(S^1), \mathcal{D}(S^2))$ parameterises an irreducible representation of $W_{x,J}$. By the Littlewood-Richardson rule (c.f. the proof of [CMO21, Proposition 4.7.2]) Equation 5.9.2 implies that

$$\mathrm{Hom}_{W_{x,J}}(F, E(\mathbb{O}^\vee, \rho)|_{W_{x,J}}) \neq 0.$$

Finally for condition (iii), let $S \in \mathcal{Y}_{r,n}^2$ of defect $r \geq 1$ correspond to (\mathbb{O}^\vee, ρ) as above and suppose $J \in \mathcal{J}_x$ is maximal and $F \in \mathrm{Irr}(W_{x,J})$ satisfies

$$\mathrm{Hom}_{W_{x,J}}(F, E(\mathbb{O}^\vee, \rho)|_{W_{x,J}}) \neq 0.$$

Since J is maximal it is of the form $C_m \times C_{n-m}$ and $W_{x,J}$ is of the form $W_k \times W_{l-k}$. Let the pair of bipartitions (B^1, B^2) correspond to F . Let (S^1, S^2) be the corresponding symbols of defect r . By the Littlewood-Richardson rule (cf. [CMO21, Proposition 4.7.2]) we have that $S^1 + S^2 \geq S$. We wish to show that

$$d_S(J, \mathrm{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) \geq \mathbb{O}^\vee.$$

By [AA07, Lemma 3.2] it suffices to show that the inequality holds with respect to the dominance order for the respective s -symbols viewed as non-decreasing sequences i.e. partitions. The s -symbol for $E(\mathbb{O}^\vee, 1)$ is just S^{sp} . Let us compute the s -symbol for the left hand side. Let $\lambda_1 = \mathcal{P}(2S_1^{1,sp} \cup 2S_2^{1,sp} + 1)$ and $\lambda_2 = \mathcal{P}(2S_2^{2,sp} \cup 2S_2^{2,sp} + 1)$ (these are both special symplectic partitions). Then

$$\mathrm{WF}(\mathcal{U}_{x,J}(F)) = \lambda_1^t \times \lambda_2^t.$$

Thus by [Oka21, Proposition 2.31]

$$d_S(J, \mathrm{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) = j_{W_J}^{W_J^\vee}(E(\lambda_1, 1) \otimes E(\lambda_2, 1)).$$

where j is j -induction, see [Lus84c, Chapter 4]. By [Lus09, 4.5 (a)], W_J is the parahoric of W (the Weyl group of \mathbf{G}) determined by J , and the s -symbol for the j -induced representation is

$$(5.9.3) \quad S^{1,sp} + S^{2,sp}.$$

Since $S^1 + S^2 \geq S$, by Lemma 5.1.1 we have that $S^{1,sp} + S^{2,sp} \geq S$. This is the required inequality.

5.9.3. $SO(N)$, N even. Suppose λ is not one of the edge cases listed in [CMO21, Proposition 4.8.2]. Then the proof is identical to the N odd case except for a few minor technical differences which we note.

- The partition $\mu(\lambda)$ is defined as

$$m_{\mu(\lambda)}(x) = \begin{cases} 2 & \text{if } x \text{ is odd and } m_{\lambda^t}(x) \text{ is} \\ & \text{even and nonzero} \\ 1 & \text{if } x \text{ is odd and } m_{\lambda^t}(x) \text{ is} \\ & \text{odd} \\ 0 & \text{otherwise.} \end{cases}$$

- $J = D_m \times D_{n-m}$.
- r is even. $\omega = 1$ if $4 \mid r$, and ω is type I if $4 \mid r - 2$. $K = D_{a^2} \times D_{a^2}$ where $a = r/2$. Despite these differences, we still have $\mathrm{Unip}_{\mathbf{L}^{c\omega(K)}, \mathbf{E}}(\mathbf{L}^{c\omega(J)}(\mathbb{F}_q)) \leftrightarrow \mathcal{Y}_{r,m}^1 \times \mathcal{Y}_{r,n-m}^1$.
- The rest of the argument follows through identically, except for the case $r = 0$ (see Equation 5.3.4 for why this case is different) which is treated in [CMO21, Lemma 4.9.5].

Now suppose λ is one of the edge cases listed in [CMO21, Porposition 4.8.2]. Recall that for these λ , $\mu(\lambda) = (1^2)$ which is a partition of 2. The issue with these edge cases can be traced down to the fact that $m = 1$, but $D_1 \times D_{n-1}$ is not a subgraph of \tilde{D}_n if we interpret D_1 as A_1 . If we interpret D_1 as the empty graph however, then as long as we interpret things correctly, the same argument as the one given for λ not an edge case works. We briefly give the details.

Suppose λ is one of said edge cases. Then λ is special, r takes values in $\{0, 2\}$, $\mu(\lambda) = (1^2)$ and $\nu(\lambda) = d_{BV}(\lambda) \setminus \mu(\lambda)$. For both $r = 0, 2$ we have that $K = \emptyset$ (since [Lus95a] uses the convention $D_1 = \emptyset$). Thus we can take J to be D_{n-1} . Take ϕ to be the family characterised by $\text{WF}(\phi, \mathbb{C}) = \nu(\lambda)$. Since $d_{BV}(\lambda) = \nu(\lambda) \cup \mu(\lambda) = \nu(\lambda) \cup (1^2)$, we have that $\nu(\lambda)$ saturates to $d_{BV}(\lambda)$ in D_n and so since λ is special we get that

$$\bar{\mathbb{L}}(D_{n-1}, \nu(\lambda)) = \bar{\mathbb{L}}(D_n, d_{BV}(\lambda)) = d_A(\mathbb{O}^\vee, 1).$$

This establishes property (i). For property (ii) we can construct as usual the a-symbols S_1, S_2 so that Equation 5.9.2 holds. Supposing $r \neq 0$ (the $r = 0$ case has already been treated in [CMO21, Proposition 4.8.2]), then $W_x = W_l$ where $l = \frac{1}{2}(N - r^2)$, and $W_{x,J} = W_{l-1}$. Since $W_{l-1} \leq W_1 \times W_{l-1} \leq W_l$ and $\mathcal{D}(S^2) \subseteq \mathcal{D}(S^1) \otimes \mathcal{D}(S^2)|_{W_{l-1}}$, we get from Equation 5.9.2 that

$$\text{Hom}_{W_{x,J}}(F, E(\mathbb{O}^\vee, \rho)|_{W_{x,J}}) \neq 0$$

as required. The proof for condition (iii) requires no modifications.

5.9.4. $Sp(2n)$. Suppose λ is not one of the edge cases listed in [CMO21, Proposition 4.8.2]. The proof for this case is similar to the $SO(N)$, N odd case with a few (more substantial) technical differences which we note.

- λ is a symplectic partition of $2n$. The partition $\mu(\lambda)$ is defined as

$$m_{\mu(\lambda)}(x) = \begin{cases} 2 & \text{if } x \text{ is odd and } m_{\lambda^t}(x) \text{ is} \\ & \text{even and nonzero} \\ 1 & \text{if } x \text{ is odd and } m_{\lambda^t}(x) \text{ is} \\ & \text{odd} \\ 0 & \text{otherwise.} \end{cases}$$

- $J = D_m \times B_{n-m}$.
- r has no parity restriction. $K = D_{a^2} \times B_{a^2+a}$ if r is even and $K = D_{a^2} \times B_{a^2-a}$ if r is odd where $a = \lfloor (r+1)/2 \rfloor$. $\omega = 1$ if a is even, and $\omega = -1$ if a is odd.
- $S \in \mathcal{X}_{d,n}^{1,1}$ where $d = r+1$ if r is even and $d = -r$ if r is odd.
- We have $\text{Unip}_{\mathbf{L}^{c\omega(K)}, \mathbf{E}}(\mathbf{L}^{c\omega(J)}(\mathbb{F}_q)) \leftrightarrow \mathcal{Y}_{d-1,m}^1 \times \mathcal{Y}_{d,n-m}^1$.
- $S^{1,sp}$ lies in $\mathcal{Y}_{0,m}^1$. The additive identity is $S^{sp} = (S^{1,sp})^\bullet + S^{2,sp}$.
- $(S^1)^\bullet$ is constructed from $(S^{1,sp})^\bullet$. We have $S^1 \in \mathcal{X}_{d-1,m}^{1,0}$, $(S^1)^\bullet \in \mathcal{X}_{d,m}^{0,1}$, $S^2 \in \mathcal{X}_{d,n-m}^{1,0}$ and $S = (S^1)^\bullet + S^2$. The \bullet map reflects similarity classes and so $S^1 \sim S^{1,sp}$.
- The rest of the argument follows through identically.

If λ is one of the edge cases listed in [CMO21, Proposition 4.8.2] then if we interpret D_1 as the empty graph, the same argument as in Section 5.9.3 follows through.

5.9.5. $Spin(N)$, N odd. Let $n = \lfloor N/2 \rfloor$, $G^\vee = Spin(N)$. Fix $(\mathbb{O}^\vee, \rho) \in \mathcal{N}_{o,l}^*$ and let $\lambda = (\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_k^{a_k}) \in \mathcal{N}_{ro}$ for $\lambda_1 < \dots < \lambda_k$ and $a_i \geq 1, i = 1, \dots, k$ be the corresponding partition under the bijection in Section 5.6.5 (1). Define $\mu(\lambda) \subseteq \lambda^t$ as follows

$$m_{\mu(\lambda)}(x) = \begin{cases} 2 & \text{if } x \text{ is even and } 4 \mid m_{\lambda^t}(x) \\ & \text{and } m_{\lambda^t}(x) \neq 0 \\ 1 & \text{if } x \text{ is even and } m_{\lambda^t}(x) \text{ is} \\ & \text{odd} \\ 0 & \text{otherwise.} \end{cases}$$

Recall from Corollary 5.4.5 that we can write $d_{BV}(\lambda) = \underline{d_{BV}(\lambda)} \cup \underline{d_{BV}(\lambda)}$.

Lemma 5.9.1. $\mu(\lambda) \subseteq \underline{d_{BV}(\lambda)}$.

Proof. Arguing as in the proof of Lemma 5.4.4, we can reduce to the case when λ consists of only odd parts, i.e. $\lambda = (\lambda_1 < \lambda_2 < \dots < \lambda_k)$. Recall that

$$\lambda^t = (k\bar{\lambda}_1, (k-1)\bar{\lambda}_2, \dots, 1\bar{\lambda}_k)$$

and

$$d_{BV}(\lambda) = (k\bar{\lambda}_1 + \delta_1, (k-1)\bar{\lambda}_2 + \delta_2, \dots, 1\bar{\lambda}_k + \delta_k)$$

where $\delta = (-1, 2, -2, \dots, 2, -2)$. If x is even, then $m_{\lambda^t}(x) \geq 2$. So $m_{d_{BV}(\lambda)}(x) = m_{\lambda^t}(x) + 2 \geq 4$. Hence, $m_{\underline{d_{BV}(\lambda)}}(x) = \frac{1}{2}m_{d_{BV}(\lambda)}(x) \geq 2$. \square

Define $\nu(\lambda) = \underline{d_{BV}(\lambda)} \setminus \mu(\lambda)$.

Let $x \in \bar{\mathfrak{C}}^*(Spin(N))$ be such that $(\mathbb{O}, \rho) \in (\mathcal{N}_{o,l}^*)_x$. x is uniquely determined and corresponds to the block parameterised by $d = d(\lambda)$ in the description of blocks given in 5.6.5 (3). We have $\theta(x) = (C_{(d^2-1)/4} \times A_{d(d-1)/2-1} \times C_{(d^2-1)/4}, \mathbf{E}, \psi, \omega)$ where $\omega = -1$, and the affine Weyl group is of type \tilde{C} . Note that since N is odd and $d(\lambda) \equiv N \pmod{2}$ we have that d is also odd. We set J to be the ω -stable set $C_m \times A_{n-2m-1} \times C_m$ where $m = \lfloor \mu \rfloor / 2$. The reductive quotient has type $C_m(q) \times {}^2A_{n-2m-1}(q^2)$. By Section 5.5,

$$\text{Irr}(W_{m-(d^2-1)/4}) \times \text{Irr}(W_{\frac{1}{2}(n-2m-\frac{1}{2}r(r+1))}) \leftrightarrow \text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$$

where $r = d - 1$ if $d > 0$ and $r = -d$ if $d \leq 0$. It will be useful to note that

$$\text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)) \subseteq \text{Unip}(C_m(q)) \times \text{Unip}({}^2A_{n-2m-1}).$$

and that $\mathcal{U}_{x,J}$ is a product of maps

$$\text{Irr}(W_{m-(d^2-1)/4}) \rightarrow \text{Unip}(C_m(q)), \quad \text{Irr}(W_{\frac{1}{2}(n-2m-\frac{1}{2}r(r+1))}) \rightarrow \text{Unip}({}^2A_{n-2m-1}).$$

We call these maps $\mathcal{U}_{x,J,1}$ and $\mathcal{U}_{x,J,2}$ respectively.

We set ϕ to be the family characterised by $\text{WF}(\phi, \mathbb{C}) = (\mu(\lambda), \nu(\lambda))$. Note the family ϕ decomposes as a product $\phi_1 \times \phi_2$ where ϕ_1 is a family of $C_m(q)$, ϕ_2 is a family of ${}^2A_{n-2m-1}(q^2)$ and

$$\text{WF}(\phi_1) = \mu(\lambda), \quad \text{WF}(\phi_2) = \nu(\lambda).$$

The following Lemma establishes condition (i) for the pair (J, ϕ) .

Lemma 5.9.2. $\bar{\mathbb{L}}(J, \text{WF}(\phi, \mathbb{C})) = d_A(\mathbb{O}^\vee, 1)$.

Proof. We have

$$\begin{aligned}\bar{\mathbb{L}}(J, \text{WF}(\phi, \mathbb{C})) &= \bar{\mathbb{L}}(C_m \times A_{n-2m-1} \times C_m, (\mu(\lambda), \nu(\lambda), \mu(\lambda))) \\ &= \bar{\mathbb{L}}(C_m \times C_{n-m}, (\mu(\lambda), \nu(\lambda) \cup \nu(\lambda) \cup \mu(\lambda))) \\ &= \bar{\mathbb{L}}(C_m \times C_{n-m}, (\mu(\lambda), d_{BV}(\lambda) \setminus \mu(\lambda))) \\ &= d_A(\mathbb{O}^\vee, 1)\end{aligned}$$

where the last equality follows from essentially the same argument given in [CMO21, Lemma 4.8.6]. \square

Recall the definition of ρ from Section 5.4 and let $B = (B_1, B_2) = \rho(\lambda)$. Note that there is a case distinction involving a flip depending on if $d(\lambda) > 0$ or ≤ 0 . Let us assume for now that $d(\lambda) > 0$. We will discuss the case $d(\lambda) \leq 0$ separately at the end.

To verify condition (ii) it suffices to find $B^1 \in \mathcal{B}(m - (d^2 - 1)/4)$, $B^2 \in \mathcal{B}(\frac{1}{2}(n - 2m - \frac{1}{2}r(r + 1)))$ such that

$$B = B^1 + B^2$$

(c.f. the proof of [CMO21, Proposition 4.7.2]) and $\mathcal{U}_{x,J,1}(B^1) \in \phi_1, \mathcal{U}_{x,J,2}(B^2) \in \phi_2$. Note that since the families in ${}^2A_{n-2m-1}$ are all singletons, the condition $\text{WF}(\mathcal{U}_{x,J,2}(B^2)) = \nu(\lambda)$ in fact completely determines B^2 provided such a B^2 exists. For this reason it is convenient to determine B^2 first.

Recall the definitions of $f, \delta, \bar{\lambda}_i, A_i$ from the end of Section 5.4. Let $(a, b]$ denote the set $\{a + 1, \dots, b\}$ when $a < b$ and \emptyset otherwise. Let (a, b) denote the set $\{a + 1, \dots, b - 1\}$ if $a + 1 < b$ and \emptyset otherwise. Let $[P]$ denote the function that takes value 1 if P is true and 0 if P is false. Define $\xi_i = [4 \mid \bar{\lambda}_i]$, $\zeta_i = [2 \nmid \bar{\lambda}_i]$ and $\mathcal{S} = \bigcup_{i=1}^l (f(2i - 1), f(2i)]$. Since $a_i = 1$ whenever λ_i is odd, we have that A_i is even if and only if $i \in \mathcal{S}$. Therefore, we can write μ as $(A_i^{\zeta_i + 2\xi_i})_{i \in \mathcal{S}}$ (this is a decreasing sequence in i). It follows that

$$\nu = (A_i^{\frac{1}{2}(\bar{\lambda}_i + \delta_i) - [i \in \mathcal{S}](\zeta_i + 2\xi_i)}).$$

A direct computation then shows that

$$\nu^t = (\eta_1^{a_1}, \eta_2^{a_2}, \dots, \eta_k^{a_k})$$

where $\eta_1 \leq \eta_2 \leq \dots \leq \eta_k$,

$$\eta_i = \frac{1}{2}(\lambda_i + [i \in \mathcal{O}_0] - [i \in \mathcal{O}_1]) - E_i, \quad E_i = \sum_{j \leq i, j \in \mathcal{S}} \zeta_j + 2\xi_j$$

and

$$\mathcal{O}_i = \{f(j) \mid 1 \leq j \leq 2l, j \equiv i \pmod{2}\}.$$

By Section 5.5.2 (2), the partition ν^t parameterises $\mathcal{U}_{x,J,2}(B^2)$ as an element of $\text{Unip}({}^2A_{n-2m-1}(q^2))$. Let us now compute $B_2 = (B_1^2, B_2^2)$ using the algorithm in Section 5.5.2 (6). Let $z_i = \lfloor \lambda_i/2 \rfloor$ so z_i is even if $r_i = 0, 1$ and is odd if $r_i = 2, 3$. In terms of z_i we have

$$\eta_i = z_i + [i \in \mathcal{O}_0] - E_i.$$

The β -set Z for ν^t of length $\#\lambda$ consists of blocks of sequences of the form

$$z_i + [i \in \mathcal{O}_0] - E_i + C_i + (0, 1, \dots, a_i - 1)$$

where $C_i = \sum_{j < i} a_j$. Note that

$$\begin{aligned}& z_i + [i \in \mathcal{O}_0] - E_i + C_i \\ & \equiv [r_i = 2, 3] + [i \in \mathcal{S}, \lambda_i \text{ is odd}] + [i \in \mathcal{S}, \lambda_i \text{ is even}] + [i \in \mathcal{S}] \pmod{2} \\ & \equiv [r_i = 2, 3] \pmod{2}.\end{aligned}$$

Thus the β -set for B_1^2 , which we denote S_1^2 , consists of blocks of sequences of the form

$$(5.9.4) \quad \begin{cases} q_i + \frac{1}{2}\Delta_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0 \\ q_i + \frac{1}{2}\Delta_i + (0) & \text{if } r_i = 1 \\ q_i + \frac{1}{2}\Delta_i + 1 + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3 \end{cases}$$

and the β -set for B_2^2 , which we denote S_2^2 , consists of blocks of sequences of the form

$$(5.9.5) \quad \begin{cases} q_i + \frac{1}{2}\Delta_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0 \\ \emptyset & \text{if } r_i = 1 \\ q_i + \frac{1}{2}\Delta_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 2 \\ q_i + \frac{1}{2}\Delta_i + (0) & \text{if } r_i = 3. \end{cases}$$

where $\Delta_i = [i \in \mathcal{O}_0] + C_i - E_i$ and $1 \leq i \leq k$. The partitions B_1^2, B_2^2 can then be obtained by applying \mathcal{P} to the β -sets. Note that $|S_1^2| - |S_2^2| = d$.

Let us now produce a bipartition B_1 satisfying $\mathcal{U}_{x,J,1}(B_1) \in \phi_1$ which, we will see, satisfies $B = B^1 + B^2$. First suppose such a B^1 exists. Let $S = \mathcal{D}^{-1}(B^1) \in \mathcal{Y}_{d,m}^1$. Using Section 5.5.4 (3), we can compute S^{sp} from the condition $\text{WF}(\mathcal{U}_{x,J,1}(B^1)) = \mu(\lambda)$. A direct computation shows that

$$\mu^t = (E_1^{a_1}, E_2^{a_2}, \dots, E_k^{a_k}).$$

Taking a β -set of length $\#\lambda$ one computes that $S^{sp} = (X, Y)$ where X consists of blocks of sequences of the form

$$(5.9.6) \quad \begin{cases} \emptyset & \text{if } i \in \mathcal{O}_0 \\ (C_i + E_i)/2 + (0) & \text{if } i \in \mathcal{O}_1 \\ (C_i + E_i)/2 + (0, 1, \dots, a_i/2 - 1) & \text{if } \lambda_i \text{ is even} \end{cases}$$

and Y consists of blocks of sequences of the form

$$(5.9.7) \quad \begin{cases} (C_i + E_i - 1)/2 + (0) & \text{if } i \in \mathcal{O}_0 \\ \emptyset & \text{if } i \in \mathcal{O}_1 \\ (C_i + E_i)/2 + (0, 1, \dots, a_i/2 - 1) & \text{if } \lambda_i \text{ is even.} \end{cases}$$

Thus we know that if such a S exists, it must have entries coming from the above special symbol. We now define such an S . Let $S = (S_1^1, S_2^1)$ where S_1^1 consists of blocks of sequences of the form

$$(5.9.8) \quad \begin{cases} (C_i + E_i - [i \in \mathcal{O}_0])/2 + (0) & \text{if } r_i = 1 \\ \emptyset & \text{if } r_i = 3 \\ (C_i + E_i)/2 + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0, 2 \end{cases}$$

and S_2^1 consists of blocks of sequences of the form

$$(5.9.9) \quad \begin{cases} \emptyset & \text{if } r_i = 1 \\ (C_i + E_i - [i \in \mathcal{O}_0])/2 + (0) & \text{if } r_i = 3 \\ (C_i + E_i)/2 + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0, 2. \end{cases}$$

Note as a sanity check that $C_i + E_i \equiv [i \in \mathcal{S}] + [i \in \mathcal{S}, \lambda_i \text{ even}] \equiv [i \in \mathcal{O}_0] \pmod{2}$ since $\mathcal{O}_0 = \{i \in \mathcal{S} \mid \lambda_i \text{ is odd}\}$ and so all the entries are integers. It is clear that S has the same entries as S^{sp} and has defect d . To show that S lies in $\mathcal{Y}_{d,m}^1$ it suffices to check that X', Y' are both strictly increasing. Clearly within each block the sequence is strictly increasing so we only need to compare endpoints of blocks. We prove this for S_1^1 . The proof for S_2^1 is analogous. Let $o_i = \lfloor (C_i + E_i)/2 \rfloor = (C_i + E_i - [i \in \mathcal{O}_0])/2$, $e_i = (C_i + E_i)/2$ and $e'_i = (C_{i+1} + E_i)/2 - 1$. Here e_i and e'_i are the entries

at the start and at the end of the i th block when λ_i is even. Let i and j index consecutive blocks of S_1^1 (so $i < j$ and $r_l = 3$ for all $i < l < j$). There are a number of cases to consider:

- $r_i = 1, r_j = 1 \implies o_i < o_j$,
- $r_i = 1, r_j \in \{0, 2\} \implies o_i < e_j$,
- $r_i \in \{0, 2\}, r_j = 1 \implies e'_i < o_j$,
- $r_i \in \{0, 2\}, r_j \in \{0, 2\} \implies e'_i < e_j$.

We prove each case as follows:

- Note that $C_j = C_i + (j - i)$ and so $o_i < o_j$ certainly holds if $j - i \geq 2$. Thus it remains to verify the case when $j = i + 1$. Suppose $j = i + 1$. Then $C_j = C_i + 1$. Moreover, since $r_{j-1} = r_i = r_j$ we have $\xi_j = 1$ and so $E_j = E_i + 2[j \in \mathcal{S}]$. Thus

$$o_j = o_i + (1 + 2[j \in \mathcal{S}] + [j - 1 \in \mathcal{O}_0] - [j \in \mathcal{O}_0])/2 = o_i + 1 > o_i$$

as required.

- By a similar reduction to the one performed in the previous case we may assume that $j = i + 1$. Then $C_j = C_i + 1$ and

$$E_j = E_i + [j \in \mathcal{S}] = E_i + 1 - [j - 1 \in \mathcal{O}_0].$$

Thus

$$e_j = (C_j + E_j)/2 = (C_i + E_i - [i \in \mathcal{O}_0])/2 + 1 > o_i$$

as required.

- Suppose $j - i \geq 2$. Then $C_j = C_{i+1} + (j - i - 1) \geq C_{i+1} + 1$ and $E_j \geq E_i$ so

$$o_j = (C_j + E_j - [j \in \mathcal{O}_0])/2 \geq (C_{i+1} + E_i + 1 - [j \in \mathcal{O}_0])/2 > e'_i.$$

If $j = i + 1$ then $E_j = E_i + [j \in \mathcal{O}_0]$ so $o_j = e'_i + 1 > e'_i$ as required.

- Since $j > i$ we have $C_j \geq C_{i+1}$ and $E_j \geq E_i$ and so

$$e'_i \leq (C_j + E_j)/2 - 1 < e_j.$$

It thus now remains to check that $\mathcal{P}(S_y^1) + \mathcal{P}(S_y^2) = B_y$ for $y = 1, 2$. For this we prove the equivalent statement that

$$(5.9.10) \quad S_y^1 + S_y^2 = B_y + 2 \cdot (0, 1, 2, \dots).$$

Let λ^i denote the partition $(\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_{i-1}^{a_{i-1}})$. We note that by Proposition 5.4.1, B_1 consists of blocks of the form

$$(5.9.11) \quad \begin{cases} q_i - d(\lambda^i) + (0, 0, \dots, 0) & \text{if } r_i = 0 \\ q_i - d(\lambda^i) + (0) & \text{if } r_i = 1 \\ q_i - d(\lambda^i) + 1 + (0, 0, \dots, 0) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3 \end{cases}$$

where in the $r_i = 0, 2$ cases the blocks have length $a_i/2$, and B_2 consists of blocks of sequences of the form

$$(5.9.12) \quad \begin{cases} q_i + d(\lambda^i) + (0, 0, \dots, 0) & \text{if } r_i = 0 \\ \emptyset & \text{if } r_i = 1 \\ q_i + d(\lambda^i) + (0, 0, \dots, 0) & \text{if } r_i = 2 \\ q_i + d(\lambda^i) + (0) & \text{if } r_i = 3 \end{cases}$$

where in the $r_i = 0, 2$ cases the blocks have length $a_i/2$. We prove Equation 5.9.10 for $y = 1$. The case $y = 2$ is analogous. Note that the block structures of S_i^1, S_i^2 , and B_i all coincide (by which we mean they have the same number of blocks and the blocks have the same size) and so we prove the

equality on an arbitrary block. For this it will be necessary to decompose the sequence $(0, 1, \dots)$ into blocks of the same size. We have for $y = 1$, the sequence $(0, 1, \dots)$ decomposes into the blocks

$$(5.9.13) \quad \begin{cases} F_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 0 \\ F_i + (0) & \text{if } r_i = 1 \\ F_i + (0, 1, \dots, a_i/2 - 1) & \text{if } r_i = 2 \\ \emptyset & \text{if } r_i = 3 \end{cases}$$

where $F_i = \sum_{j < i} ([2 \mid \lambda_j] a_j / 2 + [r(\lambda_j) = 1])$. Unpacking Equation 5.9.10 for a block with $r_i = 0, 2$ we see that we must show that

$$\begin{aligned} & q_i + \frac{1}{2} \Delta_i + [r_i = 2] + (0, 1, \dots, a_i/2 - 1) + (C_i + E_i)/2 + (0, 1, \dots, a_i/2 - 1) \\ &= q_i - d(\lambda^i) + [r_i = 2] + 2F_i + 2(0, 1, \dots, a_i/2 - 1) \end{aligned}$$

After cancelling equal terms we get that we must show

$$(5.9.14) \quad \frac{1}{2} \Delta_i + (C_i + E_i)/2 = 2F_i - d(\lambda^i).$$

Now the left hand side simplifies to C_i , whereas the right hand side simplifies to

$$\begin{aligned} & \sum_{j < i} ([2 \mid a_j] a_j + 2[r(\lambda_j) = 1]) - \sum_{j < i} ([r(\lambda_j) = 1] - [r(\lambda_j) = 3]) \\ &= \sum_{j < i} ([2 \mid a_j] a_j + [r(\lambda_j) = 1] + [r(\lambda_j) = 3]) = C_i. \end{aligned}$$

Thus we have the equality that we desire. For blocks with $r_i = 1$ we must show that

$$\begin{aligned} & q_i + \frac{1}{2} \Delta_i + (0) + (C_i + E_i - [i \in \mathcal{O}_0])/2 + (0) \\ &= q_i - d(\lambda^i) + 2F_i + 2(0). \end{aligned}$$

Cancelling equal terms yields exactly Equation 5.9.14 which we have just established. Finally, the $r_i = 3$ case is vacuous. This concludes our proof that condition (ii) holds when $d(\lambda) > 0$. The case $d(\lambda) \leq 0$ is analogous after making the following modifications

- In terms of B_1, B_2 as defined above, $B = (B_2, B_1)$;
- In terms of B_1^2, B_2^2 as defined above, $B_2 = (B_2^2, B_1^2)$;
- In terms of S_1^1, S_2^1 as defined above, we define $S = (S_2^1, S_1^1)$.

Since all the terms are just flipped the proof follows through identically.

Finally we show that condition (iii) holds. We first need a preparatory lemma. For $r \in \{0, 1, 2, 3\}$ and $a \geq 0$, define $\gamma(r, a)$ to be the sequence of length a of the form

$$(5.9.15) \quad \begin{cases} (0, 1, 0, 1, \dots) & \text{if } r = 0, 2 \\ (0, 0, 0, 0, \dots) & \text{if } r = 1, 3. \end{cases}$$

Define

$$\gamma(\lambda) = (\gamma(r_1, a_1), \gamma(r_2, a_2), \dots, \gamma(r_k, a_k)).$$

Lemma 5.9.3. *Let $\lambda \in \mathcal{N}_{ro}$. Let $\Lambda \in \mathcal{Y}_{\bullet, n}^2$ be the symbol with $\mathcal{D}(\Lambda) = E(\lambda, 1)$. Then*

$$Z^*(\Lambda) = 2q(\lambda) + z + \epsilon(\lambda) - 2\delta(\lambda).$$

Proof. Write $\lambda = (\lambda_1^{a_1}, \dots, \lambda_k^{a_k})$ with $\lambda_1 < \lambda_2 < \dots < \lambda_k$. Let $1 \leq n_1 < \dots < n_l \leq k$ denote the indicies for which λ_i is odd. Let $n_0 = 0$, $\lambda_0 = 0$, and $a_0 = 0$. We can break λ up into subsequences

of the form $(\lambda_{n_i}^{a_{n_i}}, \dots, \lambda_{n_{i+1}-1}^{a_{n_{i+1}-1}})$. Note that for $i > 0$, $a_{n_i} = 1$. Let $A_i = \sum_{j < i} a_j$. Then $\lambda + z$ is block-wise of the form

$$(5.9.16) \quad A_{n_i} + (\lambda_{n_i}^{a_{n_i}}, \dots, \lambda_{n_{i+1}-1}^{a_{n_{i+1}-1}}) + (0, 1, \dots).$$

Since $A_{n_i} \equiv i - 1 \pmod{2}$ for $i > 0$ and $A_{n_0} = 0$, considering this sequence in Equation 5.9.16 modulo 2 we see that it is of the form $i + (0, 0, 1, 0, 1, \dots) \pmod{2}$ for $i > 0$ and $(0, 1, 0, 1, \dots) \pmod{2}$ for $i = 0$. Let Thus the a-symbol consists of blocks of the form

$$\begin{pmatrix} \lambda_{n_i}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lambda_{n_i}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 0, & 1, & \dots \\ 0, & 1, & \dots \end{pmatrix}$$

for $i = 0$,

$$A_{n_i}/2 + \begin{pmatrix} \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 0, & 0, & 1, & \dots \\ 1, & 2, & \dots \end{pmatrix}$$

for $i > 0$ and odd, and

$$(A_{n_i} - 1)/2 + \begin{pmatrix} \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 1, & 2, & \dots \\ 1, & 2, & \dots \end{pmatrix}$$

for $i > 0$ and even. The s-symbol thus consists of blocks of the form

$$\begin{pmatrix} \lambda_{n_i}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lambda_{n_i}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 0, & 2, & \dots \\ 0, & 2, & \dots \end{pmatrix}$$

for $i = 0$,

$$A_{n_i} + \begin{pmatrix} \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 0, & 1, & 3, & 5, & \dots \\ 1, & 3, & 5, & \dots \end{pmatrix}$$

for $i > 0$ and odd, and

$$A_{n_i} + \begin{pmatrix} \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \\ \lfloor \lambda_{n_i}/2 \rfloor, & \lambda_{n_{i+1}}/2, & \dots, & \lambda_{n_{i+1}-1}/2, & \dots \end{pmatrix} + \begin{pmatrix} 1, & 3, & 5, & \dots \\ 0, & 1, & 3, & 5, & \dots \end{pmatrix}$$

for $i > 0$ and even. Thus as a non-decreasing sequence, the s-symbol is equal to

$$\lfloor \lambda/2 \rfloor + z - \gamma(\lambda).$$

But $\lfloor \lambda/2 \rfloor = 2\lfloor \lambda/4 \rfloor + [r = 2, 3]$. Since $[r = 2, 3] - \gamma(\lambda) = \epsilon(\lambda) - 2\delta(\lambda)$ we get that

$$\lfloor \lambda/2 \rfloor + z - \gamma(\lambda) = 2q(\lambda) + z + \epsilon(\lambda) - 2\delta(\lambda).$$

□

We handle the case $d(\lambda) > 0$ first. Suppose $J \in \mathcal{J}_x$ is maximal and $F \in \text{Irr}(W_{x,J})$ is such that

$$(5.9.17) \quad \text{Hom}_{W_{x,J}}(F, E(\mathbb{O}^\vee, \rho)|_{W_{x,J}}) \neq 0.$$

Since J is maximal $W_{x,J} = W_i \times W_{l-i}$ for $l = (N - d(2d - 1))/4$ and some $0 \leq i \leq l$. Let (B^1, B^2) be the pair of bipartitions parameterising F (so B^1 is a bipartition of i , and B^2 is a bipartition of $l - i$). We wish to show that

$$d_S(J, \text{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) \geq \mathbb{O}^\vee.$$

By [AA07, Lemma 3.2] it suffices to show that the inequality holds for the respective s-symbols viewed as non-decreasing sequences. By Lemma 5.9.3 the multiset of entries of the s-symbol for $E(\mathbb{O}^\vee, 1)$ viewed as a non-decreasing sequence is

$$2q(\lambda) + z + \epsilon(\lambda) - 2\delta(\lambda).$$

Let us compute the s -symbol attached to $d_S(J, \text{WF}(\mathcal{U}_{x,J}(F), \mathbb{C}))$. Let $S^i \in \mathcal{X}_{d(\lambda)}^{1,0}$ be the symbol associated to B^i . Write $S^i = (S_1^i, S_2^i)$. Let $\lambda_1 = \mathcal{P}(2S_1^{1,sp} \cup 2S_2^{1,sp} + 1)$ (a special symplectic partition) and $\lambda_2 = \mathcal{P}(2S_1^2 \cup 2S_2^2 + 1)$. Then

$$\text{WF}(\mathcal{U}_{x,J,1})(B^1) = \lambda_1^t, \quad \text{WF}(\mathcal{U}_{x,J,2}(B^2)) = \lambda_2^t.$$

Thus by [Oka21, Proposition 2.31]

$$d_S(J, \text{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) = j_{W_J}^{W^\vee}(E(\lambda_1, 1) \otimes E(\lambda_2, 1) \otimes E(\lambda_1, 1))$$

where W_J is the parahoric of W (the Weyl group of \mathbf{G}) determined by J . By [Lus09, 4.5 (a)], Z^* of the s -symbol for the j -induced representation is

$$(5.9.18) \quad 2S^{1,sp} + \widetilde{S}^2 - z.$$

Thus we must show that

$$(5.9.19) \quad 2S^{1,sp} + \widetilde{S}^2 - z \geq 2q(\lambda) + z + \epsilon(\lambda) - 2\delta(\lambda).$$

For this we first observe that by the Littlewood-Richardson rule (c.f. [CMO21, Proposition 4.7.2]), Equation 5.9.17 implies that $B^1 + B^2 \geq B$ where $B = \rho(\lambda)$. Let $q = q(\lambda), \epsilon = \epsilon(\lambda), \delta = \delta(\lambda)$ and Λ be the symbol with linear presentation $(q + z - \delta, \epsilon)$. By Proposition 5.4.2, $\Lambda = \rho(B) + 2(z, z)$. Thus

$$B^1 + B^2 \geq B \Rightarrow S^1 + S^2 \geq \Lambda.$$

Since ϵ has exactly $d(\lambda)$ many 1's, there is a unique sequence $S^{i,r}$ such that $(S^{i,r}, \epsilon)$ is a linear presentation for S^i . Write Λ^r for $q + z - \delta$. We thus get that

$$S^{1,r} + S^{2,r} - \Lambda^r \geq 0.$$

By Lemma 5.3.1 we have that

$$S^{2,sp} + \epsilon^*(S^2) \geq S^{2,r} + \epsilon.$$

In particular

$$S^{2,sp} + \epsilon^*(S^2) - S^{2,r} \geq \epsilon.$$

Finally we have by Lemma 5.3.2 that

$$2S^{1,sp} + \widetilde{S}^2 - z = 2(S^{1,sp} + S^{2,sp}) + \epsilon^*(S^2) - z$$

Therefore

$$\begin{aligned} & 2S^{1,sp} + \widetilde{S}^2 - z \\ &= 2(S^{1,sp} + S^{2,sp}) + \epsilon^*(S^2) - z \\ &= 2(S^{1,sp} + S^{2,sp} - S^{1,r} - S^{2,r} + S^{1,r} + S^{2,r} - \Lambda^r + \Lambda^r) + \epsilon^*(S^2) - z \\ &= 2(S^{1,sp} - S^{1,r}) + (S^{2,sp} - S^{2,r}) + (S^{2,sp} + \epsilon^*(S^2) - S^{2,r}) + 2(S^{1,r} + S^{2,r} - \Lambda) + 2q + z - 2\delta \\ &\geq 2 \cdot 0 + 0 + \epsilon + 2 \cdot 0 + 2q + z - 2\delta \\ &= 2q + z + \epsilon - 2\delta. \end{aligned}$$

where we use Lemma 5.1.1 to deduce that $S^{i,sp} - S^{i,r} \geq 0$. This proves Equation 5.9.19 as required. For the case $d(\lambda) \leq 0$ we note that Λ and S^2 get replaced by their flips, Λ^f and $(S^2)^f$. The linear presentation for Λ^f is just $(q + z - \delta, 1 - \epsilon)$. Also $\widetilde{(S^2)^f} = 2S^{2,sp} + 1 - \epsilon^*(S^2)$. In particular

$$S^{2,sp} + 1 - \epsilon^*(S^2) - S^{2,r} \geq 1 - \epsilon$$

where $S^{2,r}$ is now the linear presentation with respect to $1 - \epsilon$, and so the rest of the argument now follows through identically.

5.9.6. *Spin(N), N even.* When N is even, $\mathcal{N}_{o,l}^*$ consists of two copies of $\mathcal{P}_{ro}(N)$ instead of one. However the associated Hecke algebras are the same for both copies and so we treat both cases the same. Otherwise the proof is essentially the same as the N odd case and all the relevant lemmas have been proved in sufficient generality for the even case. We do not repeat the proof, but mention the main technical differences.

- The partition $\mu(\lambda)$ is defined as

$$m_{\mu(\lambda)}(x) = \begin{cases} 2 & \text{if } x \text{ is odd and } 4 \mid m_{\lambda^t}(x) \\ & \text{and } m_{\lambda^t}(x) \neq 0 \\ 1 & \text{if } x \text{ is odd and } m_{\lambda^t}(x) \text{ is} \\ & \text{odd} \\ 0 & \text{otherwise.} \end{cases}$$

- $J = D_m \times A_{n-2m-1} \times D_m$.
- d is even. ω is type II. $K = D_{d^2} \times A_{d(d-1)/2-1} \times D_{d^2}$.

5.10. **Proof of faithfulness in exceptional types.** Let G^\vee be a complex exceptional group and $x \in \overline{\mathfrak{C}}(G^\vee)$. Let $\theta(x) = (K, \mathbf{E}, \psi, \omega)$. The block x takes one of the following three forms:

- (1) $L^\vee = T$ and \mathcal{E}^\vee is the trivial local system;
- (2) $L^\vee = G^\vee$ and \mathcal{E}^\vee is a cuspidal local system;
- (3) $L^\vee \neq G^\vee, T^\vee$.

The first case is covered in [CMO21, Section 4.10, Lemma 3.0.2], and the second case in [CMO22]. We now cover the third case. There are very few possibilities for this case: we must have G^\vee is either of type E_6 or E_7 . If $G^\vee = E_6$ there are two possibilities for x . In both cases however the relative Weyl group is of type G_2 and \mathcal{J}_x is the same set for both. We can treat these two cases simultaneously. If $G^\vee = E_7$ there is only one choice for x and the relative Weyl group is of type F_4 . We use GAP to exhibit explicit pairs $(J, \phi) \in \mathcal{F}_x$ that realise faithfulness. We summarise the results in the next paragraph.

We treat the cases E_6, E_7 simultaneously. Suppose $\mathbb{O}^\vee \in (\mathcal{N}_o^\vee)_x$. If \mathbb{O}^\vee appears in Table 1 or Table 2 take the indicated (J, ϕ) (recall that $\text{WF}(\phi, \mathbb{C})$ completely characterises ϕ). If \mathbb{O}^\vee does not appear then \mathbb{O}^\vee has the following property: \mathbb{O}^\vee is special and $d(\mathbb{O}^\vee)$ intersects $\mathbb{L}_{c^\omega(I_{0,K})}(\overline{\mathbb{F}}_q)$ in a single F_ω -stable special orbit which we denote \mathbb{O}_0 . We take (J, ϕ) to be such that $J = I_{0,K}$ and $\text{WF}(\phi, \mathbb{C}) = \mathbb{O}_0$.


| \mathbb{O}^\vee | J | Type of $\mathbf{L}_{c^\omega(J)}(\overline{\mathbb{F}}_q)$ | $\text{WF}(\phi, \mathbb{C})$ |
|-------------------|-----|---|-------------------------------|
| A_5 | | $A_1 \times A_1$ | $(2) \times (2)$ |
| $2A_2 + A_1$ | | A_2 | (3) |

TABLE 1. (J, ϕ) for $G^\vee = E_6$

| \mathbb{O}^\vee | J | Type of $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ | WF(ϕ, \mathbb{C}) |
|-------------------|-----|--|--------------------------|
| D_6 | | 2A_7 | (2^4) |
| $E_7(a_4)$ | | $A_1 \times {}^2D_4$ | $(2) \times (3^2 1^2)$ |
| $D_6(a_2)$ | | 2A_7 | (42^2) |
| $D_5(a_1) + A_1$ | | 2A_7 | (431) |
| $A_5 + A_1$ | | $A_2 \times A_2$ | $(3) \times (3)$ |
| $D_4 + A_1$ | | 2A_7 | (4^2) |
| $A_3 + A_2 + A_1$ | | 2A_7 | (53) |
| $A_3 + 2A_1$ | | 2A_7 | (62) |
| $A_2 + 3A_1$ | | 2A_7 | (71) |
| $4A_1$ | | 2A_7 | (8) |

 TABLE 2. (J, ϕ) for $G^\vee = E_7$

Remark 5.10.1. We make a brief remark on how to interpret the table. Take for example $G^\vee = E_7, \mathbb{O}^\vee = E_7(a_4)$. J takes the form



The reductive quotient $\mathbf{L}_{c^\omega(J)}(\overline{\mathbb{F}_q})$ is of type $A_1 \times D_4 \times A_1$ and comes with a Frobenius F_ω that identifies the two A_1 factors and folds the D_4 factor. Thus the \mathbb{F}_q points $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ has type $A_1 \times {}^2D_4$. When we say $\text{WF}(\phi, \mathbb{C}) = (2) \times (3^2 1^2)$ we are listing the geometric orbit in $A_1 \times D_4$. The actual geometric orbit in $\mathbf{L}_{c^\omega(J)}(\overline{\mathbb{F}_q})$ is $(2) \times (3^2 1^2) \times (2)$.

Remark 5.10.2. We remark that there appears to be a mistake in table for the generalised Springer correspondence for E_7 in [Spa85] (and hence in GAP). It is listed that

$$E(E_7(a_4), \epsilon'') = \chi_{8,3}, \quad E(D_5 + A_1, -1) = \chi_{2,3}.$$

However, it should be

$$E(E_7(a_4), \epsilon'') = \chi_{2,3}, \quad E(D_5 + A_1, -1) = \chi_{8,3}.$$

This was verified in [Ciu08, Table 5], where the $W(F_4)$ -structure of the simple tempered modules with real infinitesimal character (and as a byproduct the generalized Springer correspondence) was computed for the graded affine Hecke algebra attached to the cuspidal local system on the principal nilpotent orbit in the Levi subgroup $(3A_1)' \subset E_7$.

6. MAIN RESULTS

Our main result is a general formula for the (canonical unramified) wavefront set of an irreducible unipotent representation with real infinitesimal character.

For this we first need an analogue of [Oka21, Theorem 3.3]. Let X be a unipotent representation of $\mathbf{G}^\omega(\mathbf{k})$ and let $x = \text{Gcusp}(X) = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ and $\theta(x) = \text{Acusp}(X) = (K, \mathbf{E}, \psi, \omega)$ (c.f. Section 4.2 and Section 4.3]). Recall from Section 4.1 that we can associate to X a module $m_{\mathbf{E}}(X)$ of the Hecke algebra $\mathcal{H} = \mathcal{H}(\mathbf{G}^\omega(\mathbf{k}), \mathbf{E})$. Our first order of business is to describe the structure of $X^{\mathbf{U}_c(\mathfrak{o})}$ in terms of restrictions of $m_{\mathbf{E}}(X)$ to subalgebras of \mathcal{H} .

Recall from Section 2.4 the definition of the local wavefront sets ${}^K\text{WF}_c(X)$ for $c \in \mathcal{B}(\mathbf{G}^\omega, \mathbf{k})$.

Lemma 6.0.1. *We have*

$${}^K\text{WF}(X) = \max_J {}^K\text{WF}_{c^\omega(J)}(X)$$

where J ranges over ω -stable subsets of I containing K .

Proof. By [Ree00, Lemma 5.1], if $X^{\mathbf{U}_{c^\omega(J)}(\mathfrak{o})} \neq 0$ then there exists a $\sigma \in \Omega$ such that $\sigma(J) \supseteq K$. But since ${}^K\text{WF}_{c^\omega(J)}$ is constant on Ω -orbits the result follows from the analogue of [Oka21, Theorem 1.18] for canonical unramified wavefront sets (this is essentially the content of [Oka21, Lemma 2.36]). \square

Recall from Section 5.8 the definition of $\mathcal{J}_x, \tilde{\mathcal{U}}_{x,J}, \mathcal{W}_x, W_x, \mathcal{W}_{x,J}$ and $W_{x,J}$. Recall from Section 4.4 the definition of the \mathcal{W}_x representation $\tilde{\sigma}_0(\theta^*(m_{\mathbf{E}}(V)))$. Write E for this representation.

Lemma 6.0.2. *We have*

$${}^K\text{WF}(X) = \max\{\overline{\mathbb{L}}(J, \text{WF}(\tilde{\mathcal{U}}_{x,J}(F), \mathbb{C})) \mid J \in \mathcal{J}_x, F \in \text{Irr}(\mathcal{W}_{x,J}), \text{Hom}_{\mathcal{W}_J}(F, E) \neq 0\}.$$

Proof. Let $J \in \mathcal{J}_x$. By [MP94] the representation $X^{\mathbf{U}_{c^\omega(J)}(\mathfrak{o})}$ of $\mathbf{L}_{c^\omega(J)}(\mathfrak{o})$ is a direct sum of representations in $\text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$. Moreover $(\theta^* \circ m_{\mathbf{E}})_{q \rightarrow 1}$ furnishes a bijection between $\text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$ and $\text{Irr}(\mathcal{W}_J)$ with inverse $\tilde{\mathcal{U}}_{x,J}$. The result then follows from Equation 4.4.3 and Lemma 6.0.1. \square

Theorem 6.0.3. *Let $X = X(s, n, \rho)$ be an irreducible unipotent $\mathbf{G}^\omega(\mathbf{k})$ -representation with $s \in T_{\mathbb{R}}^\vee$. Write $\mathbb{O}_X^\vee = G^\vee n \subset \mathfrak{g}^\vee$. Then ${}^K\text{WF}(\text{AZ}(X)), \bar{k}\text{WF}(\text{AZ}(X))$ are singletons, and*

$$\begin{aligned} {}^K\text{WF}(\text{AZ}(X)) &= d_A(\mathbb{O}_X^\vee, 1) \\ \bar{k}\text{WF}(\text{AZ}(X)) &= d(\mathbb{O}_X^\vee). \end{aligned}$$

Proof. Arguing as in the proof of [CMO22, Theorem 3.0.2], we can reduce to the case when \mathbf{G} is simple and adjoint.

By Equation (2.4.5) and the identity

$$(6.0.1) \quad \text{pr}_1(d_A(\mathbb{O}_X^\vee, 1)) = d(\mathbb{O}_X^\vee)$$

the formula for $\bar{k}\text{WF}(\text{AZ}(X))$ follows from the formula for ${}^K\text{WF}(\text{AZ}(X))$. So it suffices to show that ${}^K\text{WF}(\text{AZ}(X)) = d_A(\mathbb{O}_X^\vee, 1)$.

By inverting the character identities (4.1.1), we see that there is an equality in $R(\mathbf{G}^\omega(\mathbf{k}))$

$$X = Y(s, n, \rho) + \sum_{n \in \partial(G^\vee n')} m_{s, n', \rho'} Y(s, n', \rho')$$

for standard modules $Y(s, n', \rho')$ and $m_{s, n', \rho'} \in \mathbb{Z}$. Let $x = \text{Gcusp}(X(s, n, \rho)) = (J^\vee, \mathcal{C}^\vee, \mathcal{E}^\vee)$ and $\theta(x) = \text{Acusp}(X(s, n, \rho)) = (K, \mathbf{E}, \psi, \omega)$. By Remark 4.3.1 x lies in $\bar{\mathcal{C}}(G^\vee)$.

If $J \in \mathcal{J}_x$ and $\tau \in \text{Unip}_{\mathbf{L}_{c^\omega(K)}, \mathbf{E}}(\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q))$, let $\tau|_{q \rightarrow 1}$ be the corresponding irreducible $\mathcal{W}_{x,J}$ -representation (this was denoted $\theta^*(m_{\mathbf{E}}(\tau))_{q \rightarrow 1}$ in section 4.4). By Corollary 4.4.2,

$$(6.0.2) \quad \text{Hom}_{\mathbf{P}_{c^\omega(J)}(\mathfrak{o})}(\tau, Y(s, n', \rho')) = \text{Hom}_{W_{x,J}}(\tau_{q \rightarrow 1}, Y_{\mathbf{H}}(s, n', \rho')|_{W_x}),$$

where $Y_{\mathbf{H}}(s, n, \rho)$ is the standard module for the graded affine Hecke algebra $\mathbf{H}(G^\vee, G_{J^\vee}^\vee, \mathcal{C}^\vee, \mathcal{F}^\vee)$. To simplify the notation, let us denote by $\text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(X)$ the character of the parahoric restriction of an admissible representation V . Then we can rewrite (6.0.2) in the following form:

$$(6.0.3) \quad \text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(Y(s, n', \rho')) = \text{Res}_{W_{x,J}}^{W_x}(Y_{\mathbf{H}}(s, n', \rho')),$$

by which we understand that the multiplicity of τ in the left hand side equals the multiplicity of $\tau_{q \rightarrow 1}$ in the right hand side.

By Proposition 3.7.4, we have the following identity of W_x -characters:

$$Y_{\mathbf{H}}(s, n', \rho')|_{W_x} = \sum_{\tilde{\rho} \in \overline{A(n')_{\mathcal{L}}}} m(\rho', \tilde{\rho}) \mu(n', \tilde{\rho}) + \sum_{(n'', \tilde{\rho}'')} m_{(n', \tilde{\rho}), (n'', \tilde{\rho}')} \mu(n'', \tilde{\rho}'),$$

where

$$m(\rho', \tilde{\rho}) = \dim \text{Hom}_{A(s, n')}[\rho', \tilde{\rho}|_{A(s, n')}],$$

$m_{(n', \tilde{\rho}), (n'', \tilde{\rho}')} \in \mathbb{Z}_{\geq 0}$, $\tilde{\rho}' \in \overline{A(n'')_{\mathcal{L}}}$, and n'' ranges over a set of representatives of G^\vee -orbits in \mathcal{N}^\vee such that $n' \in \partial G^\vee n''$.

This means that

$$(6.0.4) \quad \text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(X) = \sum_{\tilde{\rho} \in \overline{A(n)_{\mathcal{L}}}} m(\rho, \tilde{\rho}) \mu(n, \tilde{\rho})|_{W_{x,J}} + \sum_{\substack{\mu' \in \text{Irr}(W_x) \\ n \in \partial \mathbb{O}^\vee(\mu', \mathbb{C})}} m_{\mu'} \mu'|_{W_{x,J}},$$

for some integers $m_{\mu'}$. Consequently (see Remark 6.0.4),

$$(6.0.5) \quad \text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(\text{AZ}(X)) = \sum_{\tilde{\rho} \in \overline{A(n)_{\mathcal{L}}}} m(\rho, \tilde{\rho}) (\mu(n, \tilde{\rho}) \otimes \text{sgn})|_{W_{x,J}} + \sum_{\substack{\mu' \in \text{Irr}(W_x) \\ n \in \partial \mathbb{O}^\vee(\mu', \mathbb{C})}} m_{\mu'} (\mu' \otimes \text{sgn})|_{W_{x,J}},$$

Notice that our normalization is such that $\mu(n, \tilde{\rho}) \otimes \text{sgn}$ is precisely the irreducible generalized Springer representation attached to the pair $(n, \tilde{\rho})$ i.e. $E(\mathbb{O}_n^\vee, \tilde{\rho})$. Thus

$$(6.0.6) \quad \text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(\text{AZ}(X)) = \sum_{\tilde{\rho} \in \overline{A(n)_{\mathcal{L}}}} m(\rho, \tilde{\rho}) E(\mathbb{O}_X^\vee, \tilde{\rho})|_{W_{x,J}} + \sum_{\substack{\mathbb{O}^\vee < \mathbb{O}'^\vee \\ (\mathbb{O}'^\vee, \tilde{\rho}') \in (\mathcal{N}_{o,l})_x}} m_{\mathbb{O}'^\vee, \tilde{\rho}'} E(\mathbb{O}_1^\vee, \tilde{\rho}')|_{W_{x,J}}.$$

Let F be an irreducible constituent of $\text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(\text{AZ}(X))$. Then $\text{Hom}_{W_{x,J}}(F, E(\mathbb{O}'^\vee, \tilde{\rho}')) \neq 0$ for some $\mathbb{O}'^\vee \geq \mathbb{O}_X^\vee$. By condition (iii) of faithfulness, we have

$$\overline{\mathbb{L}}(J, \text{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) \leq d_A(\mathbb{O}'^\vee, 1) \leq d_A(\mathbb{O}_X^\vee, 1).$$

Thus by Lemma 6.0.2 ${}^K\text{WF}(X) \leq d_A(\mathbb{O}_X^\vee, 1)$. To establish equality note that there exists at least one $\tilde{\rho}$ such that $m(\rho, \tilde{\rho}) \neq 0$. By conditions (i) and (ii) of faithfulness there exists a J and $F \in \text{Irr}(W_{x,J})$ such that $\text{Hom}_{W_{x,J}}(F, E(\mathbb{O}_X^\vee, \tilde{\rho})) \neq 0$ and

$$\overline{\mathbb{L}}(J, \text{WF}(\mathcal{U}_{x,J}(F), \mathbb{C})) = d_A(\mathbb{O}_X^\vee, 1).$$

But

$$\text{Hom}_{W_{x,J}}(F, E(\mathbb{O}_X^\vee, \tilde{\rho})) \neq 0 \implies \text{Hom}_{W_{x,J}}(F, \text{Res}_{\mathbf{L}_{c\omega(J)}(\mathbb{F}_q)}(\text{AZ}(X))) \neq 0$$

since $m(\rho, \tilde{\rho}) \neq 0$ and so by Lemma 6.0.2 we get ${}^K\text{WF}(\text{AZ}(X)) = d_A(\mathbb{O}_X^\vee, 1)$ as desired. □

Remark 6.0.4. *In the above proof, we have used implicitly (see (6.0.5)) that the restrictions of $\text{AZ}(X)$ are obtained from those of X by tensoring with sgn . In the Iwahori-spherical case, we presented the details in [CMO21, Section 2.7]. In the general case, the argument is identical using Kato's duality for affine Hecke algebras with unequal parameters [Kat93] (see also [Cha16, Section 6] for graded Hecke algebras), and the known compatibility between the parabolic induction*

(respectively, Jacquet restriction) functors for $\mathbf{G}^\omega(\mathbf{k})$ -representations in a Bernstein component and the parabolic induction (respectively, restriction) functors for the modules of the corresponding Hecke algebras (see [BK98]).

Now let $\mathbb{O}^\vee \subset \mathfrak{g}^\vee$ be a nilpotent orbit. Choose an $\mathfrak{sl}(2)$ -triple (e^\vee, f^\vee, h^\vee) with $e^\vee \in \mathbb{O}^\vee$.

Corollary 6.0.5. *Let $X = X(\frac{1}{2}h^\vee, n, \rho)$ be an irreducible unipotent $\mathbf{G}^\omega(\mathbf{k})$ -representation. Then*

$$(6.0.7) \quad \begin{aligned} d_A(\mathbb{O}^\vee, 1) &\leq_A {}^K\mathrm{WF}(X) \\ d(\mathbb{O}^\vee) &\leq {}^{\bar{k}}\mathrm{WF}(X) \end{aligned}$$

Proof. The second inequality follows from the first by applying pr_1 to both sides. So it suffices to show that $d_A(\mathbb{O}^\vee, 1) \leq_A {}^K\mathrm{WF}(X)$.

Let $X' = \mathrm{AZ}(X)$. Note that $X' = X(\frac{1}{2}h^\vee, n', \rho')$, for some n' and ρ' . By [CMO21, Lemma 3.0.9], we have $\mathbb{O}_{X'}^\vee = G^\vee n' \leq \mathbb{O}^\vee$. In particular, $(\mathbb{O}_{X'}^\vee, 1) \leq_A (\mathbb{O}^\vee, 1)$ and so $d_A(\mathbb{O}^\vee, 1) \leq_A d_A(\mathbb{O}_{X'}^\vee, 1)$ since d_A is order-reversing. But by Theorem 6.0.3, $d_A(\mathbb{O}_{X'}^\vee, 1) = {}^K\mathrm{WF}(X)$. This completes the proof. \square

REFERENCES

- [Ach03] P. Achar. “An order-reversing duality map for conjugacy classes in Lusztig’s canonical quotient”. In: *Transform. Groups* 8 (2003), pp. 107–145.
- [AA07] Pramod N. Achar and Anne-Marie Aubert. “Supports unipotents de faisceaux caractères”. In: *J. Inst. Math. Jussieu* 6.2 (2007), pp. 173–207.
- [Aub95] A. M. Aubert. “Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d’un groupe réductif p -adique”. In: *Trans. Amer. Math. Soc.* 347.6 (1995), pp. 2179–2189.
- [AMS21] A.-M. Aubert, A. Moussaoui, and M. Solleveld. “Affine Hecke algebras for Langlands parameters”. In: *arXiv:1701.03593v4* (2021).
- [BC13] D. Barbasch and D. Ciubotaru. “Unitary equivalences for reductive p -adic groups”. In: *Amer. J. Math.* 135 (6 2013), pp. 1633–1674.
- [BM93] D. Barbasch and A. Moy. “Reduction to real infinitesimal character in affine Hecke algebras”. In: *J. Amer. Math. Soc.* 6.3 (1993), pp. 611–635.
- [BV85] D. Barbasch and D. Vogan. “Unipotent representations of complex semisimple groups”. In: *Ann. of Math. (2)* 121.1 (1985), pp. 41–110.
- [BK98] C. Bushnell and P. Kutzko. “Smooth representations of reductive p -adic groups: structure theory via types”. In: *Proc. London Math. Soc.* 77 (3 1998), pp. 582–634.
- [Cha16] K.-Y. Chan. “Duality for Ext-groups and extensions of discrete series for graded Hecke algebras”. In: *Adv. Math.* 294 (2016), pp. 410–453.
- [Ciu08] Dan Ciubotaru. *On unitary unipotent representations of p -adic groups and affine Hecke algebras with unequal parameters*. 2008. DOI: 10.48550/ARXIV.0810.5366. URL: <https://arxiv.org/abs/0810.5366>.
- [CMO21] Dan Ciubotaru, Lucas Mason-Brown, and Emile Okada. *The Wavefront Sets of Unipotent Representations of Reductive p -adic Groups I*. 2021. DOI: 10.48550/ARXIV.2112.14354. URL: <https://arxiv.org/abs/2112.14354>.
- [CMO22] Dan Ciubotaru, Lucas Mason-Brown, and Emile Okada. *The Wavefront Sets of Unipotent Supercuspidal Representations*. 2022. DOI: 10.48550/ARXIV.2206.08628. URL: <https://arxiv.org/abs/2206.08628>.
- [CM93] D. Collingwood and W. McGovern. *Nilpotent Orbits In Semisimple Lie Algebra: An Introduction*. Chapman and Hall/CRC, 1993.
- [GM20] M. Geck and G. Malle. *The character theory of finite groups of Lie type*. Vol. 187. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2020.

- [Ger61] M. Gerstenhaber. “Dominance Over The Classical Groups”. In: *Annals of Mathematics* 74.3 (1961).
- [Har99] Harish-Chandra. “Harish-Chandra Admissible invariant distributions on reductive p -adic groups. With a preface and notes by Stephen DeBacker and Paul J. Sally, Jr.” In: vol. 16. University Lecture Series. American Mathematical Society, Providence, RI, 1999, xiv+97 pp.
- [Hes76] W. Hesselink. “Singularities in the nilpotent scheme of a classical group”. In: *Trans. Amer. Math. Soc.* (1976).
- [How74] R. Howe. “The Fourier transform and germs of characters (case of Gl_n over a p -adic field)”. In: *Math. Ann.* 208 (1974), pp. 305–322.
- [HL83] R. B. Howlett and G. I. Lehrer. “Representations of generic algebras and finite groups of Lie type”. In: *Trans. Amer. Math. Soc.* 280.2 (1983), pp. 753–779. ISSN: 0002-9947. DOI: 10.2307/1999645. URL: <https://doi-org.ezproxy-prd.bodleian.ox.ac.uk/10.2307/1999645>.
- [IM65] N. Iwahori and H. Matsumoto. “On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups”. In: *Inst. Hautes Études Sci. Publ. Math.* 25 (1965), pp. 5–48.
- [Kal16] T. Kaletha. “The local Langlands conjecture for non-quasi-split groups”. In: *Families of automorphic forms and the trace formula*. Simons Symp. Springer, 2016, pp. 217–257.
- [Kat93] S. Kato. “Duality for representations of a Hecke algebra”. In: *Proc. Amer. Math. Soc.* 119.3 (1993), pp. 941–946.
- [Kaw87] Noriaki Kawanaka. “Shintani lifting and Gel’fand-Graev representations”. In: *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*. Vol. 47. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1987, pp. 147–163.
- [KL87] D. Kazhdan and G. Lusztig. “Proof of the Deligne-Langlands conjecture for Hecke algebras”. In: *Invent. Math.* 1.2 (1987), pp. 153–215.
- [Kot84] R. Kottwitz. “Stable trace formula: cuspidal tempered terms”. In: *Duke Math. J.* 51 (3 1984), pp. 611–650.
- [Lus84a] G. Lusztig. “Intersection Cohomology Complexes on a Reductive Group”. In: *Invent. math.* 75 (1984).
- [Lus84b] G. Lusztig. “Intersection cohomology complexes on a reductive group”. In: *Invent. Math.* 75 (2 1984), pp. 205–272.
- [Lus86] G. Lusztig. “Character sheaves. V.” In: *Adv. Math.* 61 (2 1986), pp. 103–155.
- [Lus88] G. Lusztig. “Cuspidal local systems and graded Hecke algebras. I.” In: *Inst. Hautes Études Sci. Publ. Math.* 67 (1988), pp. 145–202.
- [Lus89] G. Lusztig. “Affine Hecke algebras and their graded version”. In: *J. Amer. Math. Soc.* 2 (1989), pp. 599–635.
- [Lus95a] G. Lusztig. “Classification of unipotent representations of simple p -adic groups”. In: *Internat. Math. Res. Notices* 11 (1995), pp. 517–589.
- [Lus95b] G. Lusztig. “Cuspidal local systems and graded Hecke algebras. II.” In: *Representations of groups (Banff, AB, 1994)*. Vol. 16. CMS Conf. Proc. Amer. Math. Soc., Providence, RI, 1995, pp. 217–275.
- [Lus02a] G. Lusztig. “Classification of unipotent representations of simple p -adic groups. II”. In: *Represent. Theory* 6 (2002), pp. 243–289.
- [Lus02b] G. Lusztig. “Cuspidal local systems and graded Hecke algebras. III.” In: *Represent. Theory* 6 (2002), pp. 202–242.
- [Lus09] G. Lusztig. “Unipotent classes and special Weyl group representations”. In: *J. Algebra* 321 (2009), pp. 3418–3449.
- [Lus20] G. Lusztig. *From families in Weyl groups to Springer representations*. 2020. arXiv: 2006.16159 [math.RT].

- [LS85] G. Lusztig and N. Spaltenstein. “On the Generalized Springer Correspondence for Classical Groups”. In: *Advanced Studies in Pure Mathematics* 6 (1985).
- [Lus84c] George Lusztig. *Characters of Reductive Groups over a Finite Field. (AM-107)*. Princeton University Press, 1984.
- [Lus92] George Lusztig. “A unipotent support for irreducible representations”. In: *Adv. Math.* 94.2 (1992), pp. 139–179. ISSN: 0001-8708. DOI: 10.1016/0001-8708(92)90035-J. URL: [https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.1016/0001-8708\(92\)90035-J](https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.1016/0001-8708(92)90035-J).
- [MP94] A. Moy and G. Prasad. “Unrefined minimal K -types for p -adic groups”. In: *Invent. Math.* 116.1-3 (1994), pp. 393–408.
- [Oka21] Emile Takahiro Okada. *The wavefront set of spherical Arthur representations*. 2021. arXiv: 2107.10591 [math.RT].
- [Opd04] E.M. Opdam. “On the spectral decomposition of affine Hecke algebras”. In: *J. Inst. Math. Jussieu* 4 (2004), pp. 531–648.
- [Pom77] K. Pommerening. “Über die unipotenten Klassen reductiver Gruppen”. In: *J. Algebra* (1977), pp. 373–398.
- [Pom80] K. Pommerening. “Über die unipotenten Klassen reductiver Gruppen II”. In: *J. Algebra* 65 (1980), pp. 525–536.
- [Ran72] R. Ranga Rao. “Orbital integrals in reductive groups”. In: *Ann. of Math. (2)* 96 (1972), pp. 505–510. ISSN: 0003-486X. DOI: 10.2307/1970822. URL: <https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.2307/1970822>.
- [Ree00] Mark Reeder. “Formal degrees and L -packets of unipotent discrete series representations of exceptional p -adic groups”. In: *J. Reine Angew. Math.* 520 (2000). With an appendix by Frank Lübeck, pp. 37–93.
- [Sol12] M. Solleveld. “On the classification of irreducible representations of affine Hecke algebras with unequal parameters”. In: *Represent. Theory* 16 (2012), pp. 1–87.
- [Sol18] M. Solleveld. *A local Langlands correspondence for unipotent representations*. 2018. eprint: 1806.11357 (math.RT).
- [Som01] E. Sommers. “Lusztig’s canonical quotient and generalized duality”. In: *J. Algebra* 243 (2001), pp. 790–812.
- [Spa82] N. Spaltenstein. *Classes Unipotentes et Sous-groupes de Borel*. Springer Berlin Heidelberg, 1982.
- [Spa85] N. Spaltenstein. “On the generalized Springer correspondence for exceptional groups”. In: *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*. Vol. 6. Adv. Stud. Pure Math. North-Holland, Amsterdam, 1985, pp. 317–338. DOI: 10.2969/aspm/00610317. URL: <https://doi-org.ezproxy-prd.bodleian.ox.ac.uk/10.2969/aspm/00610317>.
- [TT20] Jay Taylor and Pham Huu Tiep. “Lusztig induction, unipotent supports, and character bounds”. In: *Trans. Amer. Math. Soc.* 373.12 (2020), pp. 8637–8676. ISSN: 0002-9947. DOI: 10.1090/tran/8188. URL: <https://doi-org.ezproxy-prd.bodleian.ox.ac.uk/10.1090/tran/8188>.
- [Tsa22] Cheng-Chiang Tsai. *Geometric wave-front set may not be a singleton*. 2022. arXiv: 2207.13445 [math.RT].
- [Vog93] D.A. Vogan. “The local Langlands conjecture”. In: *Representation theory of groups and algebras*. Vol. 145. Contemp. Math. Amer. Math. Soc., Providence, RI, 1993, pp. 305–379.
- [Wal18] J.-L. Waldspurger. “Représentations de réduction unipotente pour $SO(2n + 1)$, III: exemples de fronts d’onde”. In: *Algebra Number Theory* 12.5 (2018), pp. 1107–1171.
- [Zel80] A. Zelevinsky. “Induced representations of reductive p -adic groups II”. In: *Ann. Sci. École Norm. Sup.* 13 (1980), pp. 154–210.

(D. Ciubotaru) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD OX2 6GG, UK
Email address: `dan.ciubotaru@maths.ox.ac.uk`

(L. Mason-Brown) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD OX2 6GG, UK
Email address: `lucas.mason-brown@maths.ox.ac.uk`

(E. Okada) DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076
Email address: `okada@nus.edu.sg`