

# REAL ALGEBRAIC MAPS OF NON-POSITIVE CODIMENSIONS OF CERTAIN GENERAL TYPES

NAOKI KITAZAWA

ABSTRACT. We present new real algebraic maps of non-positive codimensions with images of some general types. We also investigate the compositions of the maps with canonical projections.

This gives a new answer to construction of explicit real algebraic maps of negative codimensions with explicit information on preimages. Thanks to some celebrated theory of Nash, smooth closed manifolds are smooth real algebraic manifolds and the zero sets of some real polynomials. In considerable cases, we can approximate smooth functions or more generally, maps, by real algebraic ones. It is in general difficult to have explicit examples. We have constructed some previously and we have ones with new topological structures.

## 1. INTRODUCTION.

Real algebraic geometry has a long nice history. This studies real algebraic manifolds or more generally, varieties. For related topics, see [2, 3, 16, 17, 22, 31] for example. We only consider smooth real algebraic manifolds essentially here.

A part of Nash's theory says that smooth closed manifolds are smooth real algebraic manifolds and the zero sets of some real polynomials. In considerable cases, we can approximate smooth functions on smooth real algebraic manifolds or more generally, maps between smooth algebraic manifolds, by real algebraic ones.

On the other hand, our study is on construction of explicit real algebraic maps of negative codimensions and knowing global topological properties. We respect differentiable topology and singularity theory of differentiable maps. Explicit examples of such functions and maps have been well-known. Canonical projections of spheres embedded naturally in Euclidean spaces are simplest ones. Some natural functions on so-called projective spaces, Lie groups and their quotient spaces are also well-known. See [18, 23, 30] for example. We apply some theory of smooth manifolds and maps and some elementary real algebraic arguments. We add to our related pioneering studies [10, 11]. [4, 28, 29] have also motivated us.

**1.1. Polyhedra, graphs and smooth manifolds and maps.** Let  $X$  be a topological space having the structure of some cell complex whose dimension is finite. We can define the dimension  $\dim X$  uniquely.  $\dim X$  is an integer of course. A topological manifold is well-known to have the structure of a CW complex. A smooth manifold is well-known to have the structure of a polyhedron. We can define the structure of a certain polyhedron for a smooth manifold canonically. This

---

*Key words and phrases.* (Smooth) real algebraic manifolds and maps. Smooth maps. Graphs. Reeb graphs.

2020 *Mathematics Subject Classification:* Primary 14P05, 14P25, 57R45, 58C05. Secondary 57R19.

is a so-called PL manifold. It is also well-known that a topological space having the structure of a polyhedron whose dimension is at most 2 has the structure of a polyhedron uniquely. For a topological manifold, we can also say in the case where the dimension is at most 3. This respects the well-known study [21] for example.

Let  $\mathbb{R}^k$  denote the  $k$ -dimensional Euclidean space, which is a simplest  $k$ -dimensional smooth manifold and also regarded as the Riemannian manifold endowed with the standard Euclidean metric. Let  $\mathbb{R} := \mathbb{R}^1$  and  $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{R}$  be the set of all positive integers and that of all integers respectively. For each point  $x \in \mathbb{R}^k$ , we can define  $\|x\| \geq 0$  as the distance between  $x$  and the origin 0 under this metric. This is also naturally a smooth real algebraic manifold and we call this the  $k$ -dimensional real affine space.  $S^k := \{x \in \mathbb{R}^{k+1} \mid \|x\| = 1\}$  denotes the  $k$ -dimensional unit sphere, which is a  $k$ -dimensional smooth compact submanifold of  $\mathbb{R}^{k+1}$  and has no boundary. It is connected for any positive integer  $k \geq 1$ . It is a discrete set with exactly two points for  $k = 0$ . It is the zero set of the real polynomial  $\|x\|^2 = \sum_{j=1}^{k+1} x_j^2$  with  $x := (x_1, \dots, x_{k+1})$  and a smooth real algebraic (sub)manifold (hypersurface).  $D^k := \{x \in \mathbb{R}^k \mid \|x\| \leq 1\}$  is the  $k$ -dimensional unit disk, which is a  $k$ -dimensional smooth compact and connected submanifold of  $\mathbb{R}^k$  for any non-negative integer  $k \geq 0$ . We can also easily understand the topologies of these fundamental spaces.

Let  $c : X \rightarrow Y$  be a differentiable map from a differentiable manifold  $X$  to another one  $Y$ .  $x \in X$  is a *singular* point of  $c$  if the rank of the differential at  $x$  is smaller than the minimum between  $\dim X$  and  $\dim Y$ .  $c(x)$  is a *singular value* of  $c$ . Let  $S(c)$  denote the *singular set* of  $c$ , which is the set of all singular points of  $c$ . In our paper, we consider smooth maps, defined as maps of the class  $C^\infty$ , as differentiable maps, unless otherwise stated.

A canonical projection of a Euclidean space  $\mathbb{R}^k$  is denoted by  $\pi_{k,k_1} : \mathbb{R}^k \rightarrow \mathbb{R}^{k_1}$ , mapping each point  $x = (x_1, x_2) \in \mathbb{R}^{k_1} \times \mathbb{R}^{k_2} = \mathbb{R}^k$  to the first component  $x_1 \in \mathbb{R}^{k_1}$  with the conditions on the dimensions given by  $k_1, k_2 > 0$  and  $k = k_1 + k_2$ . A canonical projection of the unit sphere  $S^{k-1}$  is the restriction of it.

**1.2. Graphs and Reeb graphs.** (Reeb) graphs are our fundamental tools.

A graph can be defined as a 1-dimensional CW complex with the *vertex set* and the *edge set*. They are the set of all 0-dimensional cells and the set of all 1-dimensional cells, respectively. A vertex is an element of the vertex set. An edge is an element of the edge set. The closure of an edge homeomorphic to a circle is a *loop*. We do not consider graphs with having loops. In other words, a graph is always a 1-dimensional polyhedron. Furthermore, it is *finite*. In a word, its vertex set and edge set are finite. On the other hand, a graph may be a so-called multi-graph, a graph with more than one edge connecting some distinct vertices. An *isomorphism* from a graph  $K_1$  onto another graph  $K_2$  means a PL or a piecewise smooth homeomorphism mapping the edge set and the vertex set of  $K_1$  onto those of  $K_2$ . This defines a natural equivalence relation on the family of all graphs. Two graphs  $K_1$  and  $K_2$  are *isomorphic* if an isomorphism from  $K_1$  onto  $K_2$  exists. For a smooth function  $c : X \rightarrow \mathbb{R}$ , we can define an equivalence relation by the rule that two points  $x_1$  and  $x_2$  in  $X$  are equivalent if and only if they are in some same connected component of some preimage  $c^{-1}(y)$ . Let  $W_c$  denote the quotient space.

**Definition 1.** If  $W_c$  has the structure of a graph where a point  $p$  is a vertex if and only if  $q_c^{-1}(p)$  has some singular points, then  $W_c$  is the *Reeb graph* of  $c$ .

We explain about a main result of [26]. For a smooth function on a compact manifold having finitely many singular values, the quotient space  $W_c$  has been shown to be homeomorphic to a graph. If the manifold of the domain is closed, then we can define the Reeb graph  $W_c$  of  $c$ . Morse(-Bott) functions and functions of some considerably wide classes satisfy such conditions.

The Reeb graph of a smooth function has been already defined in [24] for example. The Reeb graphs of nice smooth functions have been important tools and objects in the theory on singularities and applications to geometry. These graphs have some information on the manifolds.

**1.3. Our main results.** Hereafter, let  $\mathbb{N}_a$  be the set of all elements of  $\mathbb{N}$  smaller than or equal to a given real number  $a$ .

**Main Theorem 1.** *Let  $l_1, l_2$  and  $n$  be positive integers. Let  $\{S_j\}_{j=1}^{l_1}$  be a family of smooth connected algebraic hypersurfaces with no boundaries each  $S_j$  of which is the zero set of a real polynomial  $f_j$ . Suppose the existence of some connected open set  $D$  of  $\mathbb{R}^n$  whose closure  $\overline{D}$  is compact and connected and suppose that for an arbitrary sufficiently small connected open neighborhood  $U_D$  of  $\overline{D}$  the following are enjoyed.*

- $S_j \cap D$  is the empty set and the intersection  $S_j \cap \overline{D}$  of  $S_j$  and  $\overline{D}$  is connected and is not empty.  $\overline{D} - D \subset \bigcup_j S_j$  holds. We can also have  $D = U_D \cap \bigcap_{j=1}^{l_1} \{x \mid f_j(x) > 0\}$ .
- A surjective map  $m_{l_1, l_2} : \mathbb{N}_{l_1} \rightarrow \mathbb{N}_{l_2}$  enjoying the following properties exists.
  - For the intersection  $\bigcap_{i=1}^{i_0} S_{j_i}$  which is not empty for some positive integer  $i_0$  and some increasing sequence  $\{j_i\}_{i=1}^{i_0}$ , the restriction of  $m_{l_1, l_2}$  to the set  $\{j_i\}_{i=1}^{i_0}$  is injective and the intersection  $\bigcap_{i=1}^{i_0} S_{j_i}$  is a real algebraic submanifold of dimension  $n - i_0$  with no boundary.
  - The intersection of  $\bigcap_{i=1}^{i_0} S_{j_i}$  and the closure  $\overline{D}$  is a smooth compact manifold and has its corner, which is non-empty in general.
  - A kind of "transversality", holds: for each point  $p$  in the intersection  $\bigcap_{i=1}^{i_0} S_{j_i}$ , the intersection of the images of the differentials at points  $e_{j_i}^{-1}(p)$  of the canonical embeddings  $e_{j_i}$  for all  $S_{j_i}$  and the image of the differential of the canonical embedding of the intersection  $\bigcap_{i=1}^{i_0} S_{j_i}$  at one point  $e_{j_i}^{-1}(p)$  agree. See [7] for related notions.

Let  $m_{l_2} : \mathbb{N}_{l_2} \rightarrow \mathbb{N} \sqcup \{0\}$  be a map. Let  $m := n + \sum_{j=1}^{l_2} m_{l_2}(j)$ . Then we have an  $m$ -dimensional smooth connected real algebraic closed manifold  $M$  and a smooth real algebraic map  $f : M \rightarrow \mathbb{R}^n$  enjoying the properties that the image  $f(M)$  is the closure  $\overline{D}$  and that the dimension of each preimage  $f^{-1}(p)$  ( $p \in \overline{D}$ ) is at most  $m - n$ .

This is an extension or a variant of main results of the author, presented in [10, 11]. So is the following. The next section explains about our Main Theorems.

**Main Theorem 2.** *Let  $m$  be a sufficiently large integer. For each graph in Example 1 of [10], we have a family of countably many smooth real algebraic functions on some  $m$ -dimensional smooth closed and connected manifolds such that the Reeb graphs collapse to the graph and that the Reeb graphs of distinct functions are mutually not isomorphic.*

**Conflict of interest.**

The author is a member of the project JSPS KAKENHI Grant Number JP22K18267 "Visualizing twists in data through monodromy" (Principal Investigator: Osamu Saeki). Under its support we do this study.

**Data availability.**

Data essentially supporting our present study are all in the paper.

## 2. ON MAIN THEOREMS.

**Theorem 1.** *In Main Theorem 1, each preimage  $f^{-1}(p)$  is as follows.*

- (1) For  $p \in D$ , it is diffeomorphic to  $\prod_{j=1}^{l_2} S^{m_{l_2}(j)}$ .
- (2) For  $p \in \bigcap_{i=1}^{i_0} S_{j_i} \cap \overline{D}$  in the increasing sequence  $\{j_i\}_{i=1}^{i_0}$  such that  $p \notin \bigcap_{i=1}^{i_0+1} S_{j'_i}$  for any increasing sequence  $\{j'_i\}_{i=1}^{i_0+1}$  containing the sequence  $\{j_i\}_{i=1}^{i_0}$  as a subsequence, it is diffeomorphic to  $\prod_{j \in \mathbb{N}_{l_2} - \{m_{l_2}(j_i)\}_{i=1}^{i_0}} S^{m_{l_2}(j)}$ .

This is an extension of a main result of the author of [10]. The result is also presented as a part of Theorem 1 of [11]. Some essential steps are same as those of their original proofs.

*A proof of Main Theorem 1 with Theorem 1.* In the proof, we use "  $\prod$  " for the products of numbers or functions. We use the notation in the forms  $\prod_{j=1}^l$  for a positive integer  $l$  and  $\prod_X$  for a finite set  $X$  for example. For coordinates and points for example, we use the notation like  $x = (x_1, \dots, x_l)$  with a positive integer  $l$  for example.

We define a set  $S_0 \subset \mathbb{R}^{m+l_2}$  by  $S_0 := \{(x, y_1, \dots, y_{l_2}) \in U_D \times \prod_{i=1}^{l_2} \mathbb{R}^{m_{l_2}(i)+1} \subset \mathbb{R}^n \times \prod_{i=1}^{l_2} \mathbb{R}^{m_{l_2}(i)+1} = \mathbb{R}^{m+l_2} \mid \prod_{j \in m_{l_1, l_2}^{-1}(i)} (f_j(x_1, \dots, x_n)) - \|y_i\|^2 = 0, i \in \mathbb{N}_{l_2}\}$ .

As important tools, we consider the partial derivatives of the real polynomial function defined by the real polynomial  $\prod_{j \in m_{l_1, l_2}^{-1}(i)} (f_j(x_1, \dots, x_n)) - \|y_i\|^2 = \prod_{j \in m_{l_1, l_2}^{-1}(i)} (f_j(x_1, \dots, x_n)) - \sum_{j'=1}^{m_{l_2}(i)} (y_{i, j'})^2$  for variants  $x_j$  and  $y_{i, j'}$ . Here  $y_i := (y_{i,1}, \dots, y_{i, m_{l_2}(i)})$ .

First, we take a point  $(x_0, y_0) \in S_0 \subset U_D \times \mathbb{R}^{m-n+l_2}$  such that  $y_0 := (y_{0,1}, \dots, y_{0, l_2})$  is not the origin. We have  $\prod_{j \in m_{l_1, l_2}^{-1}(i)} (f_j(x_0)) > 0$ . We have  $2y_{i, i_{j_0'}} = 2y_{0, i, i_{j_0'}} \neq 0$  as the derivative for some variant  $y_{i, i_{j_0'}}$ , where  $y_{0, i, i_{j_0'}}$  is from  $y_{0, i} := (y_{0, i, 1}, \dots, y_{0, i, m_{l_2}(i)})$  with  $y_0 := (y_{0,1}, \dots, y_{0, l_2})$ .

The differential of the restriction of the map into  $\mathbb{R}^{l_2}$  defined canonically from the  $l_2 > 0$  real polynomials  $\prod_{j \in m_{l_1, l_2}^{-1}(i)} (f_j(x)) - \|y_i\|^2$  at  $(x_0, y_0)$  is of rank  $l_2$ . The point is not a singular point of the map.

Second, we take a point  $(x_O, y_O) \in S_0 \subset U_D \times \mathbb{R}^{m-n+l_2}$  such that  $y_O \in \mathbb{R}^{m-n+l_2}$  is the origin. Assume also that  $x_O \in \bigcap_{i=1}^{i_0} S_{j_i} \cap \overline{D}$  holds for some increasing sequence  $\{j_i\}_{i=1}^{i_0}$  and that  $x_O \notin \bigcap_{i=1}^{i_0+1} S_{j'_i}$  holds for any increasing sequence  $\{j'_i\}_{i=1}^{i_0+1}$  containing the sequence  $\{j_i\}_{i=1}^{i_0}$  as a subsequence.

Here we consider the partial derivative of the function defined canonically from the real polynomial  $\prod_{j \in m_{l_1, l_2}^{-1}(i_x)} (f_j(x_1, \dots, x_n)) - \|y_{i_x}\|^2 = \prod_{j \in m_{l_1, l_2}^{-1}(i_x)} (f_j(x_1, \dots, x_n)) - \sum_{j'=1}^{m_{l_2}(i_x)} (y_{i_x, j'})^2$  for each variant  $x_j$  and for each  $i_x \in m_{l_1, l_2}(\{j_i\}_{i=1}^{i_0})$ . For the integer  $i_x$ , we have the value represented as the product of the partial derivative of the function for the unique number  $j_{i_x}$  satisfying  $i_x = (m_{l_1, l_2})(j_{i_x})$  at  $(x_O, y_O)$

and the product of the real numbers obtained as the values of the canonically defined real polynomial functions in the family  $\{f_j(x_1, \dots, x_n)\}_{j \in m_{l_1, l_2}^{-1}(i_x) - \{j_{i_x}\}}$  at  $x_0$ . As an important part on our arguments, we apply the assumption that certain important restrictions of  $m_{l_1, l_2}$  related to our arguments are injective. We have a real number which is not 0 by the fact that the functions defined canonically from the real polynomials  $\prod_{j \in m_{l_1, l_2}^{-1}(i_x)} (f_j(x_1, \dots, x_n)) - \|y_{i_x}\|^2 = \prod_{j \in m_{l_1, l_2}^{-1}(i_x)} (f_j(x_1, \dots, x_n)) - \sum_{j'=1}^{m_{l_2}(i_x)} (y_{i_x, j'})^2$  are assumed to have no singular points on the corresponding hypersurfaces  $S_j$  and the properties on the zero sets of the real polynomials with the nicely chosen sets  $D$  and  $U_D$ .

Let  $i_y \in \mathbb{N}_{l_2} - m_{l_1, l_2}(\{j_i\}_{j=1}^{i_0})$ . We consider the partial derivative of the function defined by the real polynomials  $\prod_{j \in m_{l_1, l_2}^{-1}(i_y)} (f_j(x_1, \dots, x_n)) - \|y_{i_y}\|^2 = \prod_{j \in m_{l_1, l_2}^{-1}(i_y)} (f_j(x_1, \dots, x_n)) - \sum_{j'=1}^{m_{l_2}(i_y)} (y_{i_y, j'})^2$ . We have  $2y_{i_y, i_y j_0'} = 2y_{0, i_y, i_y j_0'} \neq 0$  for some variant  $y_{i_y, i_y j_0'}$ , where we abuse the notation in the first case.

Remember the assumption on the transversality on intersections for hypersurfaces  $S_j$ . The differential of the restriction of the map into  $\mathbb{R}^{l_2}$  defined canonically from the  $l_2 > 0$  real polynomials same as the one before is of rank  $l_2 > 0$ . We have a result same as one in the case  $(x_0, y_0)$ .

By the assumption on  $U_D$ , which is an arbitrary sufficiently small connected open neighborhood of a nicely given connected open set  $D \subset \mathbb{R}^n$ , we cannot take a point  $x \in U_D - \overline{D}$  as the component of  $(x, y) \in S_0$ .

More precisely, by the condition that the important restrictions of  $m_{l_1, l_2}$  are always injective, at any point in  $x_U \in U_D - \overline{D}$ , for some integer  $1 \leq i_U \leq l_2$ , the value  $\prod_{j \in m_{l_1, l_2}^{-1}(i_U)} (f_j(x_{i_U}))$  is smaller than 0.

By the construction and implicit function theorem,  $S_0$  is a smooth compact submanifold in  $U_D \times \mathbb{R}^{m-n+l_2}$  and has no boundary. This is also regarded as a smooth real algebraic manifold with no boundary in the real affine space  $\mathbb{R}^{m+l_2}$ .

By considering the canonical projection into  $\mathbb{R}^n$ , we have a smooth real algebraic map  $f : M := S_0 \rightarrow \mathbb{R}^n$ . We show that  $f$  is a desired map. The properties on connectedness are enjoyed by our construction. This smooth real algebraic manifold  $M = S_0$  is a connected component of the zero set of some real polynomial map in the real affine space. The property on the image and the property on the preimages with Theorem 1 are also enjoyed. We can see this easily by our construction. We can see that  $f$  is our desired map.

This completes the proof. □

More precisely, the obtained map is, *at least topologically*, a so-called *special generic* map or a smooth map locally regarded as a so-called *moment* map. In other words, there exists such a nice map  $f_0$  and a homeomorphism  $\Phi$  enjoying the relation  $f_0 \circ \Phi = f$ . Here we adopt methods for some presentations in [11].

For special generic maps, see [25] and see also recent preprints [12, 14] of the author. See [1, 6] for moment maps on so-called *symplectic toric* manifolds. [13] is a preprint introducing a certain class of smooth maps generalizing the class of special generic maps first as the class of *simply generalized special generic* maps and investigating their topological properties, especially the cohomology rings, respecting and extending the results in [14]. These maps are locally moment maps.

We can have maps at least topologically simply generalized special generic maps in cases where hypersurfaces  $S_j$  do not intersect.

This is also related to the singularity theory of differentiable maps. See [7] for the theory. As presented in the assumption of Main Theorem 1, the notion "transversality" is also from this, for example.

For here, see also [15]. This is on explicit classifications of regions surrounded by smooth real algebraic hypersurfaces intersecting satisfying the condition on the "transversality" as in our case in explicit situations. Related to this, [4, 28, 29] are on regions surrounded by smooth real algebraic hypersurfaces with no intersections. This is explained explicitly in Example 1 and FIGURE 1 of [10], which are presented in the following proof.

*A proof of Main Theorem 2.* In Example 1 and FIGURE 1 in [10], we can take a sufficiently large outermost circle. We choose the circle as the one centered at the origin  $(0, 0) \in \mathbb{R}^2$  and of a sufficiently large radius  $R > 0$ . For any integer  $k > 0$ , we can take  $k > 0$  disks of suitable chosen radii enjoying the following properties.

- They are located sufficiently close to the point  $(-R, 0)$ .
- They are centered at points in the given sufficiently large circle centered at the origin  $(0, 0) \in \mathbb{R}^2$  and of radius  $R > 0$ .
- They are mutually disjoint.

To have our desired family of Reeb graphs, we construct maps into  $\mathbb{R}^2$  by applying Main Theorem 1 respecting the situations and compose the canonical projection to the first component.  $\square$

We close our paper by giving Remarks on Main Theorem 2.

Remark 1. This is regarded as a variant of results in [10, 11]. Moreover, these studies are originally motivated by Sharko's question in [27]. It asks whether we can have an explicit nice smooth function whose Reeb graph is isomorphic to a given graph on some closed surface or more generally, some smooth closed manifold. More precisely, these studies are also motivated by a revised question by the author. It asks whether we can respect singularities of the functions and preimages in addition. For related studies, see [8, 9] as pioneering studies by the author for example. [26] is also a related study based on our informal discussions on [8]. Note that for the case of Morse functions, [19, 20] are pioneering works and have motivated the author.

It is also remarkable that we construct smooth functions which are not real algebraic or real analytic functions in several scenes in the study. Important functions in [9, 26] show explicit cases.

Remark 2. As presented in Remark 1 of [11], in several cases such as so-called "generic" cases, we can have a similar result by considering "approximations". Approximations are presented shortly in our first section or our introduction.

We have such a result by avoiding approximations in addition to our studies [10, 11].

#### REFERENCES

- [1] V. M. Buchstaber and T. E. Panov, *Toric topology*, Mathematical Surveys and Monographs, Vol. 204, American Mathematical Society, Providence, RI, 2015.
- [2] J. Bochnak, M. Coste and M.-F. Roy, *Real algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 36, Springer-Verlag, Berlin, 1998. Translated from the 1987 French original; Revised by the authors.

- [3] J. Bochnak and W. Kucharz, *Algebraic approximation of mappings into spheres*, Michigan Mathematical Journal, vol. 34, no. 1, 1987.
- [4] A. Bodin, P. Popescu-Pampu and M. S. Sorea, *Poincaré-Reeb graphs of real algebraic domains*, arXiv:2207.06871.
- [5] R. Bott, *Nondegenerate critical manifolds*, Ann. of Math. 60 (1954), 248–261.
- [6] T. Delzant, *Hamiltoniens périodiques et images convexes de l'application moment*, Bull. Soc. Math. France 116 (1988), No. 3, 315–339.
- [7] M. Golubitsky and V. Guillemin, *Stable Mappings and Their Singularities*, Graduate Texts in Mathematics (14), Springer-Verlag(1974).
- [8] N. Kitazawa, *On Reeb graphs induced from smooth functions on 3-dimensional closed orientable manifolds with finitely many singular values*, Topol. Methods in Nonlinear Anal. Vol. 59 No. 2B, 897–912, arXiv:1902.08841.
- [9] N. Kitazawa, *On Reeb graphs induced from smooth functions on closed or open surfaces*, Methods of Functional Analysis and Topology Vol. 28 No. 2 (2022), 127–143, arXiv:1908.04340.
- [10] N. Kitazawa, *Real algebraic functions on closed manifolds whose Reeb graphs are given graphs*, a positive report for publication has been announced to have been sent and this will be published in Methods of Functional Analysis and Topology, arXiv:2302.02339.
- [11] N. Kitazawa, *Construction of real algebraic functions with prescribed preimages*, submitted to a refereed journal, arXiv:2303.00953.
- [12] N. Kitazawa, *Notes on explicit special generic maps into Euclidean spaces whose dimensions are greater than 4*, a revised version is submitted based on positive comments (major revision) by referees and editors after the first submission to a refereed journal, arxiv:2010.10078.
- [13] N. Kitazawa, *A class of naturally generalized special generic maps*, submitted to a refereed journal, arXiv:2212.03174.
- [14] N. Kitazawa, *A note on cohomological structures of special generic maps*, a revised version is submitted based on positive comments by referees and editors after the second submission to a refereed journal.
- [15] K. Kohn, R. Piene, K. Ranestad, F. Rydell, B. Shapiro, R. Sinn, M-S. Sorea and S. Telen, *Adjoint and Canonical Forms of Polypols*, arXiv:2108.11747.
- [16] J. Kollár, *Nash's work in algebraic geometry*, Bulletin (New Series) of the American Mathematical Society (2) 54, 2017, 307–324.
- [17] W. Kucharz, *Some open questions in real algebraic geometry*, Proyecciones Journal of Mathematics, Vol. 41 No. 2 (2022), Universidad Católica del Norte Antofagasta, Chile, 437–448.
- [18] E. Macías-Virgós and M. J. Pereira-Sáez, *Height functions on compact symmetric spaces*, Monatshefte für Mathematik 177 (2015), 119–140.
- [19] L. P. Michalak, *Realization of a graph as the Reeb graph of a Morse function on a manifold*, Topol. Methods in Nonlinear Anal. 52 (2) (2018), 749–762, arXiv:1805.06727.
- [20] L. P. Michalak, *Combinatorial modifications of Reeb graphs and the realization problem*, arxiv:1811.08031.
- [21] E. E. Moise, *Affine Structures in 3-Manifold: V. The Triangulation Theorem and Hauptvermutung*, Ann. of Math., Second Series, Vol. 56, No. 1 (1952), 96–114.
- [22] J. Nash, *Real algebraic manifolds*, Ann. of Math. (2) 56 (1952), 405–421.
- [23] S. Ramanujam, *Morse theory of certain symmetric spaces*, J. Diff. Geom. 3 (1969), 213–229.
- [24] G. Reeb, *Sur les points singuliers d'une forme de Pfaff complètement intégrable ou d'une fonction numérique*, Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences 222 (1946), 847–849.
- [25] O. Saeki, *Topology of special generic maps of manifolds into Euclidean spaces*, Topology Appl. 49 (1993), 265–293.
- [26] O. Saeki, *Reeb spaces of smooth functions on manifolds*, International Mathematics Research Notices, maa301, Volume 2022, Issue 11, June 2022, <https://doi.org/10.1093/imrn/maa301>, arXiv:2006.01689.
- [27] V. Sharko, *About Kronrod-Reeb graph of a function on a manifold*, Methods of Functional Analysis and Topology 12 (2006), 389–396.
- [28] M. S. Sorea, *The shapes of level curves of real polynomials near strict local maxima*, Ph. D. Thesis, Université de Lille, Laboratoire Paul Painlevé, 2018.
- [29] M. S. Sorea, *Measuring the local non-convexity of real algebraic curves*, J. Symbolic Compute. 109 (2022), 482–509.

- [30] M. Takeuchi, *Nice functions on symmetric spaces*, Osaka. J. Mat. (2) Vol. 6 (1969), 283–289.
- [31] A. Tognoli, *Su una congettura di Nash*, Ann. Scuola Norm. Sup. Pisa (3) 27 (1973), 167–185.

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU  
FUKUOKA 819-0395, JAPAN, TEL (OFFICE): +81-92-802-4402, FAX (OFFICE): +81-92-802-  
4405,

*Email address:* `n-kitazawa@imi.kyushu-u.ac.jp`

*Webpage:* <https://naokikitazawa.github.io/NaokiKitazawa.html>